FINITE ENERGY GEODESIC RAYS IN BIG COHOMOLOGY CLASSES

PRAKHAR GUPTA

ABSTRACT. For a big class represented by θ , we show that the metric space $(\mathcal{E}^p(X,\theta), d_p)$ for $p \ge 1$ is Buseman convex. This allows us to construct a chordal metric d_p^c on the space of geodesic rays in $\mathcal{E}^p(X,\theta)$. We also prove that the space of finite *p*-energy geodesic rays with the chordal metric d_p^c is a complete geodesic metric space.

With the help of the metric d_p , we find a characterization of geodesic rays lying in $\mathcal{E}^p(X,\theta)$ in terms of the corresponding test curves via the Ross-Witt Nyström correspondence. This result is new even in the Kähler setting.

CONTENTS

1.	Introduction	1
2.	Preliminaries	3
3.	Lidskii type inequality	7
4.	Buseman Convexity of $\mathcal{E}^p(X,\theta)$	8
5.	Space of geodesic rays	11
6.	The Ross-Witt Nyström correspondence	15
References		17

1. INTRODUCTION

Finding canonical metrics on a compact Kähler manifold (X, ω) is a problem that has guided the field of Kähler geometry for many decades now. To tackle this problem, [Mab87], [Sem92], and [Don99] constructed a Riemannian structure on

$$\mathcal{H}_{\omega} = \{ u \in C^{\infty}(X) : \omega + dd^{c}u > 0 \},\$$

the space of smooth Kähler potentials in ω . To find the canonical metric cohomologous to ω , we needed to understand the geometry of \mathcal{H}_{ω} endowed with this Riemannian structure. In [Che00], Chen proved that this Riemannian structure gives rise to a metric d_2 on \mathcal{H}_{ω} . In [Dar17], Darvas showed that the completion $\overline{(\mathcal{H}_{\omega}, d_2)}$ can be identified with $(\mathcal{E}^2(X, \omega), d_2)$, the space of finite energy potentials, confirming a conjecture of Guedj [Gue14]. Moreover, any two points in $\mathcal{E}^2(X, \omega)$ can be joined by a metric geodesic lying in $\mathcal{E}^2(X, \omega)$. Using Finsler metric structure on \mathcal{H}_{ω} for $p \geq 1$, in [Dar15], Darvas constructed complete geodesic metrics $(\mathcal{E}^p(X, \omega), d_p)$. These metrics proved useful in finding Kähler-Einstein [DR17], and cscK metrics in (X, ω) [CC21a], [CC21b], [CC18], [BDL17].

Further properties of the metric structure of $(\mathcal{E}^p(X,\omega), d_p)$ were studied to improve our understanding of canonical metrics. In [CC18, Theorem 1.5], Chen-Cheng proved that the metric space $(\mathcal{E}^p(X,\omega), d_p)$ is Buseman convex, with the case p = 1 being proved in [BDL17]. They use the Buseman convexity property for p = 1, to prove that the L^1 geodesic stability of (X,ω) is equivalent to the existence of cscK metric cohomologous to ω [CC18]. In [DL20a], Darvas-Lu proved a uniform convexity property for $(\mathcal{E}^p(X,\omega), d_p)$ for p > 1, that allowed them to prove $C^{1,\bar{1}}$ -geodesic stability for the existence of cscK metric. In op. cit., the authors used Buseman convexity and uniform convexity to prove that the space of p-finite energy geodesic rays \mathcal{R}^p_{ω} can be endowed with a complete geodesic metric d^p_p .

In the author's previous paper [Gup24], he showed that for a big cohomology class $\{\theta\}$, the space of finite energy potentials $\mathcal{E}^p(X,\theta)$ can be endowed with a complete geodesic metric d_p . Moreover, he showed that the metric space $(\mathcal{E}^p(X,\theta), d_p)$ is uniformly convex if p > 1. With the prospects of studying

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MARYLAND, COLLEGE PARK, MARYLAND, USA

 $E\text{-}mail\ address: \texttt{pgupta8@umd.edu}.$

²⁰⁰⁰ Mathematics Subject Classification. Primary: 32U05; Secondary: 32Q15, 53C55.

Key words and phrases. Kähler Manifolds, Pluripotential Theory, Monge-Ampère Measures, Finite Energy Classes.

stability in the big case in mind (c.f. [DXZ23]), in this paper, we will explore the space of geodesic rays in the metric space $(\mathcal{E}^p(X,\theta), d_p)$.

In [Gup24], the author used potentials of analytic singularity type to approximate the minimal singularity type. He used this approximation to construct the complete geodesic metric space $(\mathcal{E}^p(X,\theta), d_p)$ for $p \geq 1$. In this paper, we show that the same approximation scheme can be used to prove various properties of the metric space $(\mathcal{E}^p(X,\theta), d_p)$.

As a first application of the approximation scheme, we prove a Lidskii-type inequality, the analog of a well known inequality for matrices (for a survey, see [DLR20, Theorem 2.7]). This result was first proved in [DLR20, Theorem 5.1] for Kähler classes $\omega = c_1(L)$ induced by an ample line bundle L. In [DL20a, Corollary 3.2], Darvas-Lu extended this result to an arbitrary Kähler class. In this paper, we use the approximation scheme to extend the result to arbitrary big classes. In particular, we prove

Theorem 1.1. If $u, v, w \in \mathcal{E}^p(X, \theta)$ satisfy $u \ge v \ge w$, then

$$d_p^p(v,w) \le d_p^p(u,w) - d_p^p(u,v)$$

Next, we extend [CC18, Theorem 1.5] of Chen-Cheng to the big cohomology classes. In the Kähler setting, they prove a Buseman type convexity of the metric space $(\mathcal{E}^p(X, \omega), d_p)$. We prove

Theorem 1.2. Let θ be a real, smooth, closed (1,1)-form representing a big cohomology class. If $p \ge 1$, then the metric space $(\mathcal{E}^p(X,\theta), d_p)$ is Buseman convex. This means if $u_0, u_1, v_0, v_1 \in \mathcal{E}^p(X,\theta)$, and $[0,1] \ni t \mapsto u_t$ and $[0,1] \ni t \mapsto v_t$ are the weak geodesics joining u_0, u_1 , and v_0, v_1 respectively, then

$$d_p(u_t, v_t) \le (1-t)d_p(u_0, v_0) + td_p(u_1, v_1).$$

These results allow us to study the metric geometry of the space of geodesic rays in $\mathcal{E}^p(X,\theta)$ in more depth. Given $u \in \mathcal{E}^p(X,\theta)$, we say the geodesic ray $[0,\infty) \ni t \mapsto u_t \in \mathcal{E}^p(X,\theta)$ is a finite *p*-energy geodesic ray starting at $u_0 = u$. We denote such a ray by $\{u_t\}_t \in \mathcal{R}^p_u$. We can endow \mathcal{R}^p_u with a metric, as follows. Given geodesic rays $\{u_t^0\}_t, \{u_t^1\}_t \in \mathcal{R}^p_u$ we define the chordal distance between them by

(1)
$$d_p^c(\{u_t^0\}_t, \{u_t^1\}_t) = \lim_{t \to \infty} \frac{d_p(u_t^0, u_t^1)}{t}.$$

Let θ represent a big cohomology class. By \mathcal{R}^p_u we denote the space of finite *p*-energy geodesic rays emanating from $u \in \mathcal{E}^p(X, \theta)$.

Theorem 1.3. (\mathcal{R}^p_u, d^c_n) is a complete geodesic metric space.

1.1. Ross-Witt Nyström correspondence. The notion of test curves was introduced in [RW14] by Ross-Witt Nyström in the Kähler setting, giving a potential theoretic framework to study stability.

Test curves are dual to (sub)geodesic rays via the Legendre transform. Ross-Witt Nyströms's work generalized the work of Phong-Sturm [PS07] where they associated geodesic rays to test configurations. The study of test curves was extended to the big setting in [DDL18a] by Darvas-Di Nezza-Lu. Furthermore, the notion of finite energy test curves was introduced in [DX22] (in the Kähler setting) and [DZ23] (in the big setting). Test curves can detect several properties of the geodesic rays. Using the metric d_p on $\mathcal{E}^p(X, \theta)$, the following theorem proves that the geodesic rays lying in $\mathcal{E}^p(X, \theta)$ can be detected from the corresponding test curve.

If $(0, \infty) \ni t \mapsto u_t \in PSH(X, \theta)$ is a θ -geodesic ray starting at V_{θ} , then its Legendre dual $\mathbb{R} \ni \tau \mapsto \hat{u}_{\tau} \in PSH(X, \theta)$ is defined by

$$\hat{u}_{\tau} = \inf_{t>0} (u_t - t\tau).$$

For τ large enough, $\hat{u}_{\tau} \equiv -\infty$. Thus we define,

$$au_{\hat{u}}^+ := \inf\{ au \in \mathbb{R} \, : \, \hat{u}_{ au} \equiv -\infty\} < \infty.$$

Theorem 1.4. Let $\{u_t\}_t$ be a θ -geodesic ray starting from V_{θ} . Let $\{\hat{u}_{\tau}\}_{\tau}$ be the Legendre dual of $\{u_t\}_t$. Then $u_t \in \mathcal{E}^p(X, \theta)$ for all $t \ge 0$ iff

(2)
$$\int_{-\infty}^{\tau_{\hat{u}}^+} (-\tau + \tau_{\hat{u}}^+)^{p-1} \left(\int_X \theta_{V_\theta}^n - \int_X \theta_{\hat{u}_\tau}^n \right) d\tau < \infty$$

When $\tau_{\hat{u}}^+ = 0$, the expression in Equation (2) equals $\frac{1}{p} d_p^p(V_{\theta}, u_1) = \frac{1}{p} \int_X |\dot{u}_0|^p \theta_{V_{\theta}}^n$. Thus this expression is closely related to the speed of the geodesic ray in $\mathcal{E}^p(X, \theta)$. When p = 1, such characterization of finite energy geodesic rays was obtained in [DX22, Theorem 3.7] (in the Kähler setting) and [DZ23, Theorem 3.9] (in the big setting). For p > 1, this is new in the Kähler case as well.

Organization. In Section 2, we recall some background material and in particular the approximation scheme that is used in the construction of the metric d_p on $\mathcal{E}^p(X,\theta)$. In Section 3, we prove Theorem 1.1. In Section 4, we prove Theorem 1.2. In Section 5, we prove Theorem 1.3. Lastly, in Section 6, we prove theorem 1.4.

Acknowledgement. I want to thank my advisor Tamás Darvas for his constant support and valuable suggestions on an early draft that improved the paper. I am also grateful to Mingchen Xia and Antonio Trusiani for carefully reading the preliminary version of this paper and suggesting helpful improvements. Research is partially supported by NSF CAREER grant DMS-1846942.

2. Preliminaries

2.1. Pluripotential theory. In this section, we will recall the notions from the pluripotential theory developed in [BEGZ10]. Throughout, we will work on a compact Kähler manifold (X, ω) .

Let θ be a smooth closed real (1,1)-from representing the cohomology class $\{\theta\} \in H^{1,1}(X,\mathbb{R})$. A function $u : X \to \mathbb{R} \cup \{-\infty\}$ is called a θ -psh function if locally u + g is plurisubharmonic, where $\theta = dd^c g$. By dd^c here we mean $\sqrt{-1}\partial\bar{\partial}/2\pi$. We use $PSH(X,\theta)$ to denote the set of all θ -psh functions. If u is θ -psh, then the (1,1)-current $\theta_u := \theta + dd^c u$ is a closed positive current.

We say θ represents a pseudoeffective class if $PSH(X, \theta) \neq \emptyset$. We say θ represents a big cohomology class if there exists $u \in PSH(X, \theta)$ such that $\theta_u \geq \varepsilon \omega$ for some small enough $\varepsilon > 0$. We say that θ represents a Kähler class if there is $u \in PSH(X, \theta) \cap C^{\infty}(X)$ such that $\theta_u \geq \varepsilon \omega$ for some $\varepsilon > 0$. We say that θ represents a nef cohomology class $\{\theta + \varepsilon \omega\}$ is a Kähler class. In a Kähler class, there are plenty of smooth potentials, but in an arbitrary big class, we do not expect any smooth potentials.

Let θ represent a big cohomology class. If $u, v \in PSH(X, \theta)$ we say that u is more singular than v, denoted by $u \leq v$ if $u \leq v + C$ for some constant C. We say that $u, v \in PSH(X, \theta)$ have the same singularity type if $u \leq v$ and $v \leq u$. The potential

(3)
$$V_{\theta} = \sup\{u \in PSH(X, \theta) : u \le 0\}$$

has the least singularity among all θ -psh functions. Any potential with the same singularity type as V_{θ} is called a minimal singularity potential.

In [BEGZ10], the authors introduced a non-pluripolar measure $\theta_u^n := (\theta + dd^c u)^n$ associated to any θ -psh function u. Witt Nyström proved [Wit19] proved that if $u \leq v$, then $\int_X \theta_u^n \leq \int_X \theta_v^n$. Thus for any $u \in \text{PSH}(X, \theta), \int_X \theta_u^n \leq \int_X \theta_{V_\theta}^n := \text{Vol}(\theta)$.

We denote the potentials of full mass by $\mathcal{E}(X, \theta)$. In particular

$$\mathcal{E}(X,\theta) := \left\{ u \in \mathrm{PSH}(X,\theta) : \int_X \theta_u^n = \int_X \theta_{V_\theta}^n \right\}.$$

For $p \ge 1$, we say that $u \in \mathcal{E}(X, \theta)$ has finite *p*-energy if $\int_X |u - V_\theta|^p \theta_u^n < \infty$. We denote

$$\mathcal{E}^p(X,\theta) := \left\{ u \in \mathcal{E}(X,\theta) : \int_X |u - V_\theta|^p \theta_u^n < \infty \right\}.$$

We can do the same construction in the prescribed singularity setting as well. See [DDL23] for more details. If ϕ is a model potential, meaning $P_{\theta}[\phi](0) = \phi$ (see Equation (6)), then we define the space of relative full mass as

$$\mathcal{E}(X,\theta,\phi) = \left\{ u \in \mathrm{PSH}(X,\theta) \, : \, u \preceq \phi, \int_X \theta_u^n = \int_X \theta_\phi^n \right\}$$

and the space of relatively finite p-energy as

$$\mathcal{E}^p(X,\theta,\phi) = \left\{ u \in \mathcal{E}(X,\theta,\phi) : \int_X |u-\phi|^p \theta_\phi^n < \infty \right\}.$$

2.2. Geodesic rays. Following Berndtsson [Ber15] and Darvas-Di Nezza-Lu [DDL18b] we define the geodesic segments by an envelope construction. Let $S \subset \mathbb{C}$ be a vertical strip given by $S = \{z \in \mathbb{C} \mid 0 \leq \text{Re}(z) \leq 1\}$. Let $\pi : S \times X \to X$ be the projection. Let $u_0, u_1 \in \text{PSH}(X, \theta)$, then we say that a path $(0,1) \ni t \mapsto \text{PSH}(X,\theta)$ is a subgeodesic between u_0 and u_1 if the map $S \times X \ni (z,x) \mapsto V(z,x) := v_{\text{Re}(z)}(x)$ is $\pi^*\theta$ -psh and $\lim_{t\to 0,1} v_t \leq u_{0,1}$. Let $S(u_0, u_1)$ to be the collection of all subgeodesics between u_0 and u_1 . We define the geodesic joining u_0 and u_1 to be a path $(0,1) \ni t \mapsto u_t \in \text{PSH}(X,\theta)$ given by

(4)
$$u_t(x) = \sup_{\mathcal{S}(u_0, u_1)} v_t(x)$$

In the Kähler setting, the geodesics joining "smooth" points have some regularity properties. If η represents a Kähler class (and not necessarily be a Kähler form), then we denote

$$\mathcal{H}_n^{1,1} = \mathrm{PSH}(X,\eta) \cap C^{1,1}(X)$$

which will act as the space of "smooth" potentials for us.

We recall that in [He15], He proved

Lemma 2.1. If ω is a Kähler form and $u_0, u_1 \in \mathcal{H}^{1,\bar{1}}_{\omega}$, then the geodesic u_t joining u_0 and u_1 is also in $\mathcal{H}^{1,\bar{1}}_{\omega}$.

We can modify He's result to prove

Lemma 2.2. If η represents a Kähler class and $u_0, u_1 \in \mathcal{H}^{1,\bar{1}}_{\eta}$, then the geodesic u_t joining them is in $\mathcal{H}^{1,\bar{1}}_{\eta}$ as well.

Proof. Let $g \in C^{\infty}(X)$ such that $\eta_g := \eta + dd^c g$ is Kähler. Then $u_0 - g, u_1 - g \in C^{1,\bar{1}}(X)$ are η_g -psh. Lemma 2.1 says that the goedesic $u_{t,g}$ joining them is in $C^{1,\bar{1}}(X)$. Since $u_{t,g} + g$ is the η -geodesic joining u_0 and u_1 , we get that $u_t = u_{t,g} + g \in C^{1,\bar{1}}(X)$ as well.

Now we define geodesic rays. Again we assume that θ represents a big cohomology class. A path $(0,\infty) \ni t \mapsto v_t \in PSH(X,\theta)$ is a sublinear subgeodesic ray starting at $u_0 \in PSH(X,\theta)$ if $u_t \to u_0$ in $L^1(X)$ as $t \to 0$ and for any $0 < a < b < \infty$, the path $(a,b) \ni t \mapsto u_t$ is a subgeodesic between u_a and u_b and $u_t \leq u_0 + Ct$ for some constant C.

A sublinear subgeodesic ray $(0, \infty) \ni t \mapsto u_t \in PSH(X, \theta)$ is a geodesic ray starting at $u_0 \in PSH(X, \theta)$ if $u_t \to u_0$ in $L^1(X)$ as $t \to 0$ and for any $0 < a < b < \infty$, the path $(a, b) \ni t \mapsto u_t$ is a geodesic ray joining u_a and u_b . We recall a few useful results about the geodesic rays.

Lemma 2.3 ([DZ23, Remark 3.3]). If $[0, \infty) \ni t \mapsto u_t \in \mathcal{E}^p(X, \theta)$ is a geodesic ray starting from $u_0 = V_{\theta}$, then $t \mapsto \sup_X u_t = \sup_{Amp(\theta)} (u_t - V_{\theta})$ is linear.

Theorem 2.4 ([DZ23, Proposition 3.8]). If $[0, \infty) \ni t \mapsto u_t \in \mathcal{E}^p(X, \theta)$ is a geodesic ray starting at $u_0 = V_{\theta}$, then one can find geodesic rays of minimal singularity type $[0, \infty) \ni t \mapsto u_t^j \in \mathcal{E}^p(X, \theta)$ such that each u_t^j has minimal singularity and $u_t^j \searrow u_t$ as $j \to \infty$ for all $t \ge 0$.

2.3. Plurisubharmonic envelopes. The envelope construction has several applications in pluripotential theory. We have already seen two examples of such construction in Equations 3 and 4. In this section we will see more such examples are recall some theorems about them.

Assume that θ is a smooth closed real (1,1)-from that represents a big cohomology class. Given a measurable function $f: X \to \mathbb{R} \cup \{\pm \infty\}$, we define

(5)
$$P_{\theta}(f) := (\sup\{u \in PSH(X, \theta) : u \le f\})^*,$$

where φ^* is the upper semicontinuous regularization of φ . $P_{\theta}(f) \equiv -\infty$ if there is no θ -psh function such that $u \leq f$.

If $\phi \in PSH(X, \theta)$, then the envelope with respect to the singularity type of ϕ is constructed by

(6)
$$P_{\theta}[\phi](f) := (\sup\{u \in \mathrm{PSH}(X,\theta) : u \preceq \phi, u \leq f\})^* = \left(\lim_{C \to \infty} P_{\theta}(\phi + C, f)\right)^*.$$

In the Kähler setting, we have the following regularity result.

(

Lemma 2.5. If η represents a Kähler class (and not necessarily be a Kähler form) and $f \in C^{1,\bar{1}}(X)$, then $P_{\eta}(f) \in C^{1,\bar{1}}(X)$.

Proof. Let $g \in C^{\infty}(X)$ such that $\eta_g = \eta + dd^c g$ is a Kähler form. We can see that $P_{\eta}(f) = P_{\eta_g}(f-g) + g$. Now due to [Ber19] (for a survey see [Dar19, Appendix A.3]), $P_{\eta_g}(f-g) \in C^{1,\bar{1}}(X)$ as $f-g \in C^{1,\bar{1}}(X)$. Therefore, $P_{\eta_g}(f-g) + g \in C^{1,\bar{1}}(X)$. Thus $P_{\eta}(f) \in C^{1,\bar{1}}(X)$.

In general, we have the following formula for the non-pluripolar measures of potentials obtained by taking an envelope :

Lemma 2.6 ([DT21]). If θ represents a big cohomology class, $\phi \in PSH(X, \theta)$, and $f \in C^{1,\overline{1}}(X)$, then

$$\theta_{P_{\theta}[\phi](f)}^{n} = \mathbb{1}_{\{P_{\theta}[\phi](f)=f\}} \theta_{f}^{n}.$$

2.4. The complete geodesic metric d_p . If ω is a Kähler metric, a complete geodesic metric d_p on $\mathcal{E}^p(X,\omega)$ was constructed by Darvas in [Dar15]. If β represents a big and nef cohomology class, then by approximating from the Kähler case, Di Nezza-Lu constructed such a complete geodesic metric d_p on $\mathcal{E}^p(X,\beta)$ in [DL20b]. If θ represents a big cohomology class, then in [Gup24], the author constructed an approximation scheme via analytic singularity types to construct a complete geodesic metric d_p on $\mathcal{E}^p(X,\theta)$. In this subsection, we will describe the approximation method used to construct d_p on big and nef, and big classes.

2.4.1. d_p metric in big and nef classes. Let β represent a big and nef cohomology class. Here we explain the construction of the complete geodesic metric d_p on $\mathcal{E}^p(X,\beta)$ by Di Nezza-Lu in [DL20b]. Let ω be a Kähler form. Denote by $\omega_{\varepsilon} := \beta + \varepsilon \omega$, a smooth form which represents a Kähler class. There is a complete geodesic metric d_p on $\mathcal{E}^p(X, \omega_{\varepsilon})$. Let

$$\mathcal{H}_{\beta} = \{ u \in \text{PSH}(X, \beta) : u = P_{\beta}(f) \text{ where } f \in C^{1,1}(X) \}.$$

 \mathcal{H}_{β} is the analog of the space of smooth potentials from the Kähler case, and it was first used in [DL20b]. If $u_0, u_1 \in \mathcal{H}_{\beta}$, then there are $f_0, f_1 \in C^{1,\bar{1}}(X)$ such that $u_0 = P_{\beta}(f_0)$ and $u_1 = P_{\beta}(f_1)$. We define $u_{0,\varepsilon} = P_{\omega_{\varepsilon}}(f_0)$ and $u_{1,\varepsilon} = P_{\omega_{\varepsilon}}(f_1)$. We define

$$d_p(u_0, u_1) = \lim_{\varepsilon \to 0} d_p(u_{0,\varepsilon}, u_{1,\varepsilon})$$

For $u_0, u_1 \in \mathcal{E}^p(X, \beta)$, we can find $u_0^j, u_1^j \in \mathcal{H}_\beta$ such that $u_0^j \searrow u_0$ and $u_1^j \searrow u_1$. We define

$$d_p(u_0, u_1) = \lim_{j \to \infty} d_p(u_0^j, u_1^j)$$

2.4.2. d_p metric in the analytic singularity setting. Let ψ be a θ -psh function with analytic singularities. By Hironaka's resolution, we can find a modification $\mu : \tilde{X} \to X$ that resolves the singularities of ψ . See [Gup24, Section 3] for more details. There is a smooth closed real (1, 1)-form $\tilde{\theta}$ and a bounded $\tilde{\theta}$ -psh function g on \tilde{X} such that the map

$$\operatorname{PSH}(X, \theta, \psi) \ni u \mapsto \tilde{u} := (u - \psi) \circ \mu + g \in \operatorname{PSH}(X, \theta)$$

is an order-preserving bijection. Moreover, this mapping is a bijection between $\mathcal{E}^p(X, \theta, \psi)$ and $\mathcal{E}^p(\tilde{X}, \tilde{\theta})$ as well. Since $\tilde{\theta}$ represents a big and nef class, there is a complete geodesic metric d_p on $\mathcal{E}^p(\tilde{X}, \tilde{\theta})$. Using this correspondence, we can define the metric on $\mathcal{E}^p(X, \theta, \psi)$ as follows. Let $u_0, u_1 \in \mathcal{E}^p(X, \theta, \psi)$, then

(7)
$$d_p(u_0, u_1) := d_p(\tilde{u}_0, \tilde{u}_1).$$

2.4.3. d_p metric in big classes. Let θ represent a big cohomology class. By Demailly's regularization, we can find an increasing sequence ψ_k of θ -psh functions with analytic singularities such that $\psi_k \nearrow V_{\theta}$.

By [Gup24, Theorem 7.4], we can define the complete geodesic metric d_p on $\mathcal{E}^p(X, \theta)$ by the following. If $u_0, u_1 \in \mathcal{E}^p(X, \theta)$, then

(8)
$$d_p(u_0, u_1) = \lim_{k \to \infty} d_p(P_{\theta}[\psi_k](u_0), P_{\theta}[\psi_k](u_1)).$$

2.4.4. Properties of the metric d_p . Let θ represent a big cohomology class. We denote by

$$\mathcal{H}_{\theta} = \{ u \in PSH(X, \theta) : u = P_{\theta}(f) \text{ for some } f \in C^{1,1}(X) \}$$

Notice that if θ represents a Kähler class, then due to Lemma 2.5 \mathcal{H}_{θ} consists of $C^{1,\bar{1}}$ potentials.

Theorem 2.7 ([Gup24, Theorem 4.7]). Let $u_0, u_1 \in \mathcal{H}_{\theta}$, and let u_t be the weak geodesic joining u_0 and u_1 . Then

$$d_p^p(u_0, u_1) = \int_X |\dot{u}_0|^p \theta_{u_0}^n = \int_X |\dot{u}_1|^p \theta_{u_1}^n.$$

Lemma 2.8 (Pythagorean identity, [Gup24, Theorem 5.5]). The metric d_p satisfies the following Pythagorean identity. If $u_0, u_1 \in \mathcal{E}^p(X, \theta)$, then

$$d_p^p(u_0, u_1) = d_p^p(u_0, P_\theta(u_0, u_1)) + d_p^p(u_1, P_\theta(u_0, u_1)).$$

Lemma 2.9 ([Gup24, Lemma 5.2]). If $u_0^j, u_1^j, u_0, u_1 \in \mathcal{E}^p(X, \theta)$ satisfy $u_0^j \searrow u_0$ and $u_1^j \searrow u_1$, then

$$\lim_{j \to \infty} d_p(u_0^j, u_1^j) = d_p(u_0, u_1)$$

Theorem 2.10 (Uniform Convexity, [Gup24, Theorem 8.3]). Let p > 1. Let $u, v_0, v_1 \in \mathcal{E}^p(X, \theta)$ and v_λ be the weak geodesic joining v_0 and v_1 . Then

$$d_p(u, v_{\lambda})^2 \leq (1 - \lambda)d_p(u, v_0)^2 + \lambda d_p(u, v_1)^2 - (p - 1)\lambda(1 - \lambda)d_p(v_0, v_1)^2, \text{ if } 1$$

A corollary of Theorem 2.10 is

Corollary 2.11 ([Gup24, Corollary 8.5]). For p > 1, if $u, v_0, v_1 \in \mathcal{E}^p(X, \theta)$ such that for some $\lambda \in [0, 1]$ and $\varepsilon > 0$, $d_p(u, v_0) \le (\lambda + \varepsilon)d_p(v_0, v_1)$ and $d_p(u, v_1) \le (1 - \lambda + \varepsilon)d_p(v_0, v_1)$, then there is a constant C > 0 such that

$$d_p(u, v_{\lambda}) \le C\varepsilon^{\frac{1}{r}} d_p(v_0, v_1)$$

where $r = \max\{2, p\}$ and v_t is the geodesic joining v_0 and v_1 .

A consequence of these properties is the following

Lemma 2.12. If $1 \le p' \le p < \infty$ then for any $u_0, u_1 \in \mathcal{E}^p(X, \theta)$,

$$d_{p'}(u_0, u_1) \le d_p(u_0, u_1) \operatorname{Vol}(\theta)^{\frac{1}{p'} - \frac{1}{p}}.$$

Proof. First we assume that $u_0, u_1 \in \mathcal{H}_{\theta}$ and u_t is the weak geodesic joining u_0 and u_1 . Then by Theorem 2.7,

$$d_p^p(u_0, u_1) = \int_X |\dot{u}_0|^p \theta_{u_0}^n \quad \text{and} \quad d_{p'}^{p'}(u_0, u_1) = \int_X |\dot{u}_0|^{p'} \theta_{u_0}^n.$$

Applying Hölder inequality with factors p/p' and p/(p - p') we can write,

$$\int_{X} |\dot{u}_{0}|^{p'} \theta_{u_{0}}^{n} \leq \left(\int_{X} |\dot{u}_{0}|^{p} \theta_{u_{0}}^{n}\right)^{p'/p} \left(\int_{X} \theta_{u_{0}}^{n}\right)^{\frac{p-p'}{p}}.$$

Raising both sides to 1/p' and noticing that $\int_X \theta_{u_0}^n = \operatorname{Vol}(\theta)$, we get that

$$d_{p'}(u_0, u_1) \le d_p(u_0, u_1) \operatorname{Vol}(\theta)^{\frac{1}{p'} - \frac{1}{p}}.$$

More generally, if $u_0, u_1 \in \mathcal{E}^p(X, \theta)$, then we can find $u_0^j, u_1^j \in \mathcal{H}_{\theta}$ such that $u_0^j \searrow u_0$ and $u_1^j \searrow u_1$. By Lemma 2.9,

$$\lim_{j \to \infty} d_{p'}(u_0^j, u_1^j) = d_{p'}(u_0, u_1) \quad \text{and} \quad \lim_{j \to \infty} d_p(u_0^j, u_1^j) = d_p(u_0, u_1).$$

From the proof above we can write,

$$d_{p'}(u_0, u_1) = \lim_{j \to \infty} d_{p'}(u_0^j, u_1^j) \le \lim_{j \to \infty} d_p(u_0^j, u_1^j) \operatorname{Vol}(\theta)^{\frac{1}{p'} - \frac{1}{p}} = d_p(u_0, u_1) \operatorname{Vol}(\theta)^{\frac{1}{p'} - \frac{1}{p}}.$$

The following Lemma from [BDL17] shows that the weak geodesics are stable under perturbations of the endpoints.

Lemma 2.13. If $u_0^j, u_1^j, u_0, u_1 \in \mathcal{E}^p(X, \theta)$ satisfy $d_p(u_0^j, u_0) \to 0$ and $d_p(u_1^j, u_1) \to 0$ as $j \to \infty$, then $d_p(u_t^j, u_t) \to 0$ as $j \to \infty$ where u_t^j, u_t are the weak geodesics joining u_0^j, u_1^j and u_0, u_1 respectively.

Proof. The proof is the same as in [BDL17, Proposition 4.3]. We just need the corresponding results used in the proof in [BDL17, Proposition 4.3] for the big setting. For that we can obtain the quasimonotonicity for the d_p metric from [Gup23, Theorem 5.2] and [Gup24, Lemma 4.12]. The fact that for $u, v \in \mathcal{E}^1(X, \theta)$ satisfying $u \leq v$ and I(u) = I(v) imply u = v follows by combining [DDL18a, Theorem 2.4, Proposition 2.5, and Proposition 2.8]. Moreover, the fact that $u \leq v \leq w$ implies $d_p(u, v) \leq d_p(u, w)$ can be obtained from Theorem 1.1 (whose proof does not rely on this lemma).

Lemma 2.14. If $u_j \in \mathcal{E}^p(X, \theta)$ is a d_p -bounded decreasing sequence, then $\lim_{j\to\infty} u_j =: u \in \mathcal{E}^p(X, \theta)$.

Proof. By [Gup24, Lemma 4.12], the metric d_p is comparable to the quasi-metric I_p i.e., there exists C > 1, such that

$$\frac{1}{C}I_p(u_0, u_1) \le d_p^p(u_0, u_1) \le CI_p(u_0, u_1).$$

Here the functional I_p is given by

$$I_p(u_0, u_1) = \int_X |u_0 - u_1|^p (\theta_{u_0}^n + \theta_{u_1}^n)$$

for $u_0, u_1 \in \mathcal{E}^p(X, \theta)$. So, if the sequence u_j is d_p -bounded, then it is I_p -bounded as well. This means that

$$\sup_{n \in \mathbb{N}} \int_{X} |u_0 - u_j|^p \theta_{u_j}^n < \infty.$$

$$=: u \in \mathcal{E}^p(X, \theta).$$

Thus by [GLZ19], the limit $\lim_{j\to\infty} u_j =: u \in \mathcal{E}^p(X,\theta)$

 s^{j}

We will end this section by proving a d_p -contraction property for the projection into different cohomology classes. This result will play an important role in obtaining our main results. This is an analog of the contraction property in [Gup24, Theorem 7.3], and [Tru22, Theorem 4.3] for p = 1, where the projection operator maps to a different singularity type in the same cohomology class.

Lemma 2.15 (Contraction property). If θ and η are closed smooth real (1,1)-forms representing big cohomology classes such that $\theta \leq \eta$. Then the map $P_{\theta} : \mathcal{H}_{\eta} \mapsto \mathcal{H}_{\theta}$ is a contraction. This means that $v_0, v_1 \in \mathcal{H}_{\eta}$, then

$$d_p(P_\theta(v_0), P_\theta(v_1)) \le d_p(v_0, v_1)$$

Proof. First we will show that P_{θ} maps \mathcal{H}_{η} to \mathcal{H}_{θ} . If $v \in \mathcal{H}_{\eta}$, then $v = P_{\eta}(f)$ for some $f \in C^{1,\overline{1}}$. By a standard argument, we can show that $P_{\theta}(v) = P_{\theta}(f)$. Thus showing that $P_{\theta}(v) \in \mathcal{H}_{\theta}$.

Now, assume that $f_0, f_1 \in C^{1,\overline{1}}$ be such that $v_0 = P_{\eta}(f_0)$ and $v_1 = P_{\eta}(f_1)$. Also call $u_0 = P_{\theta}(f_0) = P_{\theta}(v_0)$ and $u_1 = P_{\theta}(f_1) = P_{\theta}(v_1)$. For now, we also assume that $f_0 \leq f_1$. By Theorem 2.7 and Lemma 2.6 the distances are given by

$$d_p^p(v_0, v_1) = \int_X |\dot{v}_0|^p \eta_{v_0}^n = \int_{\{v_0 = f_0\}} |\dot{v}_0|^p \eta_{f_0}^n$$

and

$$d_p^p(u_0, u_1) = \int_X |\dot{u}_0|^p \theta_{u_0}^n = \int_{\{u_0=f_0\}} |\dot{u}_0|^p \theta_{f_0}^n.$$

where v_t is the η -geodesic joining v_0 and v_1 , and u_t is the θ -geodesic joining u_0 and u_1 . Since $\theta \leq \eta$, $u_0 \leq v_0$ and $u_1 \leq v_1$, u_t is the η -subgeodesic joining v_0 and v_1 , therefore, $u_t \leq v_t$.

Since $u_0 \le v_0 \le f_0$, the set $\{u_0 = f_0\} \subset \{v_0 = f_0\}$. Moreover, when $x \in \{u_0 = f_0\}$,

$$\dot{u}_0(x) = \lim_{t \to 0} \frac{u_t(x) - u_0(x)}{t} = \lim_{t \to 0} \frac{u_t(x) - f_0(x)}{t} \le \lim_{t \to 0} \frac{v_t(x) - f_0(x)}{t} = \lim_{t \to 0} \frac{v_t(x) - v_0(x)}{t} = \dot{v}_0(x).$$

Due to the assumption that $f_0 \leq f_1$, $0 \leq \dot{u}_0$, \dot{v}_0 , thus we obtain that $|\dot{u}_0| \leq |\dot{v}_0|$. Again due to $\theta \leq \eta$, we have $\theta_{f_0}^n \leq \eta_{f_0}^n$. Combining these we get

$$\int_{\{u_0=f_0\}} |\dot{u}_0|^p \theta_{f_0}^n \le \int_{\{v_0=f_0\}} |\dot{v}_0|^p \eta_{f_0}^n$$

Therefore, $d_p(u_0, u_1) \le d_p(v_0, v_1)$.

More generally, if f_0 and f_1 are unrelated, we can use the Pythagorean identity (see Lemma 2.8) to prove the general result. We can do this because of [Gup24, Lemma 4.1] implies that $P_{\theta}(u_0, u_1) \in \mathcal{H}_{\theta}$ and the fact that $P_{\theta}(P_{\eta}(v_0, v_1)) = P_{\theta}(u_0, u_1)$.

$$d_p^p(u_0, u_1) = d_p^p(u_0, P_{\theta}(u_0, u_1)) + d_p^p(u_1, P_{\theta}(u_0, u_1))$$

$$\leq d_p^p(v_0, P_{\eta}(v_0, v_1)) + d_p^p(v_1, P_{\eta}(v_0, v_1))$$

$$= d_p^p(v_0, v_1).$$

3. Lidskii type inequality

In this section we will prove Theorem 1.1. The proof will follow the approximation process used to construct the metric d_p on $\mathcal{E}^p(X, \theta)$. First, we will prove the Lidskii-type inequality for the big and nef cohomology classes and then prove it for the big cohomology classes.

Lemma 3.1. If β represents a big and nef cohomology class and $u, v, w \in \mathcal{E}^p(X, \beta)$ satisfy $u \ge v \ge w$, then

$$d_p^p(v,w) \le d_p^p(u,w) - d_p^p(u,v).$$

Proof. First, we assume that $u, v, w \in \mathcal{H}_{\beta}$. Let $\omega_{\varepsilon} = \beta + \varepsilon \omega$ represent a Kähler class. Let $u = P_{\beta}(f)$, $v = P_{\beta}(g)$, and $w = P_{\beta}(h)$ where $f, g, h \in C^{1,\overline{1}}(X)$ satisfy $f \geq g \geq h$. We denote by $u_{\varepsilon} = P_{\omega_{\varepsilon}}(f)$, $v_{\varepsilon} = P_{\omega_{\varepsilon}}(g)$, and $w_{\varepsilon} = P_{\omega_{\varepsilon}}(h)$. Since $u_{\varepsilon} \geq v_{\varepsilon} \geq w_{\varepsilon}$ are in $\mathcal{E}^{p}(X, \omega_{\varepsilon})$, by [DL20a, Corollary 3.2],

$$d_p^p(v_{\varepsilon}, w_{\varepsilon}) \le d_p^p(u_{\varepsilon}, w_{\varepsilon}) - d_p^p(u_{\varepsilon}, v_{\varepsilon}).$$

By the definition of d_p on \mathcal{H}_{β} (see Section 2.4.1), we know that $\lim_{\varepsilon \to 0} d_p(u_{\varepsilon}, v_{\varepsilon}) = d_p(u, v)$, $\lim_{\varepsilon \to 0} d_p(u_{\varepsilon}, w_{\varepsilon}) = d_p(u, w)$, and $\lim_{\varepsilon \to 0} d_p(v_{\varepsilon}, w_{\varepsilon}) = d_p(v, w)$. Thus taking the limit $\varepsilon \to 0$ in the above equation we get

$$d_p^p(v,w) \le d_p^p(u,w) - d_p^p(u,v).$$

More generally, if $u, v, w \in \mathcal{E}^p(X, \beta)$, then we can find $u^j, v^j, w^j \in \mathcal{H}_\beta$ such that $u^j \ge v^j \ge w^j$ and $u^j \searrow u, v^j \searrow v$, and $w^j \searrow w$ as $j \to \infty$. We just proved that

$$d_p^p(v^j, w^j) \le d_p^p(u^j, w^j) - d_p^p(u^j, v^j).$$

Taking the limit $j \to \infty$ and using Lemma 2.9, we get

$$d_p^p(v,w) \le d_p^p(u,w) - d_p^p(u,v).$$

Before we can prove the Lidskii-type inequality for the big cohomology class, we need to show the result for the analytic singularity type.

Lemma 3.2. If θ represents a big cohomology class and ψ is a θ -psh function with analytic singularity type, then for $u, v, w \in \mathcal{E}^p(X, \theta, \psi)$ satisfying $u \ge v \ge w$, we have

$$d_p^p(v,w) \le d_p^p(u,w) - d_p^p(u,v).$$

Proof. Let $\mu : \tilde{X} \to X$ be the modification that resolves the singularities of ψ . As discussed in Section 2.4.2 there is a smooth closed real (1, 1)-form $\tilde{\theta}$ representing a big and nef cohomology class, and a bounded $\tilde{\theta}$ -psh function g on \tilde{X} such that $\mathcal{E}^p(X, \theta, \psi) \ni u \mapsto \tilde{u} = (u - \psi) \circ \mu + g \in \mathcal{E}^p(\tilde{X}, \tilde{\theta})$ is a bijection.

Since $\tilde{u}, \tilde{v}, \tilde{w} \in \mathcal{E}^p(\tilde{X}, \tilde{\theta})$ satisfy $\tilde{u} \geq \tilde{v} \geq \tilde{w}$, by Lemma 3.1 we have

$$d_p^p(\tilde{v}, \tilde{w}) \le d_p^p(\tilde{u}, \tilde{w}) - d_p^p(\tilde{u}, \tilde{v})$$

By Equation (7), we can write

$$d_p^p(v,w) \le d_p^p(u,w) - d_p^p(u,v)$$

Now we can prove the Lidskii-type inequality for the big classes.

Theorem 3.3. Let θ represent a big cohomology class. If $u, v, w \in \mathcal{E}^p(X, \theta)$ satisfy $u \geq v \geq w$, then

$$d_p^p(v,w) \le d_p^p(u,w) - d_p^p(u,v).$$

Proof. As discussed in Section 2.4.3, let ψ_k be an increasing sequence of θ -psh functions with analytic singularities such that $\psi_k \nearrow V_{\theta}$ a.e. We denote by $u^k = P_{\theta}[\psi_k](u)$, $v^k = P_{\theta}[\psi_k](v)$, and $w^k = P_{\theta}[\psi_k](w)$. Since $u \ge v \ge w$, we have $u^k \ge v^k \ge w^k$. By Lemma 3.2,

$$d_{p}^{p}(v^{k}, w^{k}) \leq d_{p}^{p}(u^{k}, w^{k}) - d_{p}^{p}(u^{k}, v^{k}).$$

Notice that the metric d_p applied to u^k, v^k, w^k is the metric on the space $\mathcal{E}^p(X, \theta, \psi_k)$, whereas the metric applied to u, v, w is the metric on the space $\mathcal{E}^p(X, \theta)$. Taking the limit $k \to \infty$ and using Equation (8) we get

$$d_p^p(v,w) \le d_p^p(u,w) - d_p^p(u,v).$$

4. BUSEMAN CONVEXITY OF $\mathcal{E}^p(X,\theta)$

In this section, we will prove Buseman convexity of the metric space $(\mathcal{E}^p(X,\theta), d_p)$ for $p \ge 1$ where θ represents a big cohomology class. This means that we will prove that if $u_0, u_1, v_0, v_1 \in \mathcal{E}^p(X,\theta)$, and u_t and v_t are geodesic joining u_0, u_1 , and v_0, v_1 respectively, then

$$d_p(u_t, v_t) \le (1-t)d_p(u_0, v_0) + td_p(u_1, v_1).$$

4.1. Buseman convexity for p = 1. The Buseman convexity for p = 1 follows from the arguments in [BDL17, Proposition 5.1]. We reproduce the arguments here for completeness.

Theorem 4.1. If θ represents a big cohomology class, $u_0, u_1, v_0, v_1 \in \mathcal{E}^p(X, \theta)$, and u_t and v_t are the weak geodesics joining u_0, u_1 , and v_0, v_1 respectively, then

$$d_1(u_t, v_t) \le (1-t)d_1(u_0, v_0) + td_1(u_1, v_1).$$

Proof. Fix $a \leq b \in [0, 1]$. Let $w_a := P_{\theta}(u_a, v_a)$ and $w_b = P_{\theta}(u_b, v_b)$. Let $[a, b] \ni t \mapsto w_t \in \mathcal{E}^p(X, \theta)$ be the geodesic joining w_a and w_b . Since $w_a \leq u_a, v_a$ and $w_b \leq u_b, v_b$, the comparison principle for geodesics implies that $w_t \leq u_t, v_t$ for all $t \in [a, b]$. Thus $w_t \leq P_{\theta}(u_t, v_t)$. By monotonicity of I (see [DDL18a, Theorem 2.4 and Proposition 2.5], we get that $I(w_t) \leq I(P_{\theta}(u_t, v_t))$. As I is linear along geodesics, we have $I(P_{\theta}(u_t, v_t)) \geq I(w_t) = \frac{b-t}{b-a}I(P_{\theta}(u_a, v_a)) + \frac{t-a}{b-a}I(P_{\theta}(u_b, v_b))$ for all $t \in [a, b]$. Thus $t \mapsto I(P_{\theta}(u_t, v_t))$ is a concave map.

By [Gup24, Theorem 5.7], $d_1(u_t, v_t) = I(u_t) + I(v_t) - 2I(P_{\theta}(u_t, v_t))$, thus it agrees with the metric introduced in [DDL18a]. As $t \mapsto I(u_t)$ and $t \mapsto I(v_t)$ are linear and $t \mapsto I(P_{\theta}(u_t, v_t))$ is concave, we obtain that $t \mapsto d_1(u_t, v_t)$ is convex.

Thus, we only need to prove Buseman convexity for p > 1. This is great because now we can use uniform convexity of $(\mathcal{E}^p(X,\theta), d_p)$ from Theorem 2.10. First, we will prove Buseman convexity for the big and nef case, and then prove the Buseman convexity for the big classes.

4.2. Buseman convexity for big and nef classes when p > 1. Fix a smooth closed real (1, 1)-form β representing a big and nef cohomology class and fix a p > 1. Recall from Section 2.4.1 that $\omega_{\varepsilon} = \beta + \varepsilon \omega$ represents a Kähler class. Let $u_0, u_1, v_0, v_1 \in \mathcal{H}_{\beta}$ be given by $u_0 = P_{\beta}(f_0), u_1 = P_{\beta}(f_1), v_0 = P_{\beta}(g_0)$, and $v_1 = P_{\beta}(g_1)$ where $f_0, f_1, g_0, g_1 \in C^{1,\overline{1}}(X)$. Let $u_{0,\varepsilon} = P_{\omega_{\varepsilon}}(f_0), u_{1,\varepsilon} = P_{\omega_{\varepsilon}}(f_1), v_{0,\varepsilon} = P_{\omega_{\varepsilon}}(g_0)$, and $v_{1,\varepsilon} = P_{\omega_{\varepsilon}}(g_1)$. Let $u_{t,\varepsilon}$ and $v_{t,\varepsilon}$ be the ω_{ε} -geodesic joining $u_{0,\varepsilon}, u_{1,\varepsilon}$, and $v_{0,\varepsilon}, v_{1,\varepsilon}$ respectively.

By the Buseman convexity in the Kähler case, we obtain that

$$d_p(u_{t,\varepsilon}, v_{t,\varepsilon}) \le (1-t)d_p(u_{0,\varepsilon}, v_{0,\varepsilon}) + td_p(u_{1,\varepsilon}, v_{1,\varepsilon}).$$

By construction of d_p on $\mathcal{E}^p(X,\beta)$, we know that $\lim_{\varepsilon \to 0} d_p(u_{0,\varepsilon}, v_{0,\varepsilon}) = d_p(u_0, v_0)$ and $\lim_{\varepsilon \to 0} d_p(u_{1,\varepsilon}, v_{1,\varepsilon}) = d_p(u_1, v_1)$. Thus to show Buseman convexity for potentials in \mathcal{H}_β , it is enough to show that

$$d_p(u_t, v_t) \leq \lim_{t \to 0} d_p(u_{t,\varepsilon}, v_{t,\varepsilon}),$$

where u_t and v_t are the β -geodesics joining u_0, u_1 and v_0, v_1 respectively.

Lemma 4.2. Let $u_0, u_1 \in \mathcal{H}_\beta$ and u_t be the geodesic joining them. Let $u_{0,\varepsilon}, u_{1,\varepsilon}$ be as above and $u_{t,\varepsilon}$ be the geodesic joining $u_{0,\varepsilon}$ and $u_{1,\varepsilon}$. Then for p > 1,

$$\lim_{\varepsilon \to 0} d_p(u_t, P_\beta(u_{t,\varepsilon})) = 0.$$

Proof. We claim that $P_{\beta}(u_{t,\varepsilon}) \searrow u_t$.

First notice that $u_{0,\varepsilon} \searrow u_0$ and $u_{1,\varepsilon} \searrow u_1$. Recall that by definition, $u_{0,\varepsilon} = P_{\omega_{\varepsilon}}(f_0)$ and $u_0 = P_{\beta}(f_0)$ for $f_0 \in C^{1,\overline{1}}$. If $\varepsilon_1 < \varepsilon_2$, then u_{0,ε_1} is ω_{ε_2} -psh as well and $u_{0,\varepsilon_1} \leq f_0$, therefore, $u_{0,\varepsilon_1} \leq u_{0,\varepsilon_2}$. By a similar argument $u_{0,\varepsilon} \geq u_0$ as well. Therefore, $u_{0,\varepsilon}$ are increasing sequence of functions, and $\lim_{\varepsilon \to 0} u_{0,\varepsilon} \geq u_0$. The limit $\lim_{\varepsilon \to 0} u_{0,\varepsilon}$ is a $\beta + \varepsilon \omega$ -psh for every $\varepsilon > 0$, therefore, $\lim_{\varepsilon \to 0} u_{0,\varepsilon}$ is a β -psh function that satisfies $\lim_{\varepsilon \to 0} u_{0,\varepsilon} \leq f_0$, thus $\lim_{\varepsilon \to 0} u_{0,\varepsilon} \leq u_0$. Thus $\lim_{\varepsilon \to 0} u_{0,\varepsilon} = u_0$. Similarly, $u_{1,\varepsilon} \searrow u_1$.

We can make the same argument for the ω_{ε} -geodesics $u_{t,\varepsilon}$ joining $u_{0,\varepsilon}$ and $u_{1,\varepsilon}$. If $\varepsilon_1 \leq \varepsilon_2$, then the path $(0,1) \ni t \mapsto u_{t,\varepsilon_1}$ is an ω_{ε_2} -subgeodesic joining u_{0,ε_2} and u_{1,ε_2} . Therefore, $u_{t,\varepsilon_1} \leq u_{t,\varepsilon_2}$. Similarly, $u_{t,\varepsilon} \geq u_t$. Therefore, $(0,1) \ni t \mapsto u_{t,\varepsilon}$ is a decreasing sequence of paths and $\lim_{\varepsilon \to 0} u_{t,\varepsilon} \geq u_t$. The limit path $(0,1) \ni t \mapsto \lim_{\varepsilon \to 0} u_{t,\varepsilon}$ is a $\beta + \varepsilon \omega$ -subgeodesic joining u_0 and u_1 for each $\varepsilon > 0$, therefore it is a β -subgeodesic as well. Hence $\lim_{\varepsilon \to 0} u_{t,\varepsilon} \leq u_t$. So we get $\lim_{\varepsilon \to 0} u_{t,\varepsilon} = u_t$.

Since $u_{t,\varepsilon} \searrow u_t$ and $u_{t,\varepsilon} \ge P_{\beta}(u_{t,\varepsilon}) \ge u_t$, we have that $P_{\beta}(u_{t,\varepsilon}) \searrow u_t$ as well. Thus proving the claim. Now the proof of the lemma follows from Lemma 2.9.

Similarly for p > 1, $\lim_{\varepsilon \to 0} d_p(v_t, P_\beta(v_{t,\varepsilon})) = 0$. By the triangle inequality,

$$\lim_{\epsilon \to 0} d_p(P_\beta(u_{t,\varepsilon}), P_\beta(v_{t,\varepsilon})) = d_p(u_t, v_t).$$

Since $f_0, f_1 \in C^{1,\bar{1}}(X)$, Lemma 2.5 implies that $u_{0,\varepsilon}, u_{1,\varepsilon} \in C^{1,\bar{1}}(X)$. Thus Lemma 2.2 implies that the geodesic $u_{t,\varepsilon} \in C^{1,\bar{1}}(X)$ as well. Similarly, the geodesic $v_{t,\varepsilon} \in C^{1,\bar{1}}(X)$. Thus we can apply Lemma 2.15,

to get $d_p(P_\beta(u_{t,\varepsilon}), P_\beta(v_{t,\varepsilon})) \leq d_p(u_{t,\varepsilon}, v_{t,\varepsilon})$. Combining this with the above equation we get

$$d_p(u_t, v_t) = \lim_{\varepsilon \to 0} d_p(P_{\beta}(u_{t,\varepsilon}), P_{\beta}(v_{t,\varepsilon})) \le \lim_{\varepsilon \to 0} d_p(u_{t,\varepsilon}, v_{t,\varepsilon})$$

as we needed to show. Thus we have proved

Lemma 4.3. Let β represent a big and nef cohomology class and p > 1. Let $u_0, u_1, v_0, v_1 \in \mathcal{H}_{\beta}$. If u_t and v_t are the geodesics joining u_0, u_1 and v_0, v_1 respectively, then

$$d_p(u_t, v_t) \le (1-t)d_p(u_0, v_0) + td_p(u_1, v_1).$$

Now we will extend this result to $\mathcal{E}^p(X,\beta)$ for p > 1.

Theorem 4.4. Let β represent a big and nef cohomology class and p > 1. Let $u_0, u_1, v_0, v_1 \in \mathcal{E}^p(X, \beta)$. If u_t and v_t are the geodesics joining u_0, u_1 and v_0, v_1 respectively, then

$$l_p(u_t, v_t) \le (1-t)d_p(u_0, v_0) + td_p(u_1, v_1).$$

Proof. As standard, we can find $u_0^j, u_1^j, v_0^j, v_1^j \in \mathcal{H}_\beta$ such that $u_0^j \searrow u_0, u_1^j \searrow u_1, v_0^j \searrow v_0$, and $v_1^j \searrow v_1$. If u_t^j and v_t^j are the geodesics joining u_0^j, u_1^j , and v_0^j, v_1^j respectively, then $u_t^j \searrow u_t$ and $v_t^j \searrow v_t$.

By the definition of d_p for $\mathcal{E}^p(X,\beta)$ in Section 2.4.1, we know that $\lim_{j\to\infty} d_p(u_0^j, v_0^j) = d_p(u_0, v_0)$ and that $\lim_{j\to\infty} d_p(u_1^j, v_1^j) = d_p(u_1, v_1)$. Moreover, due to Lemma 2.9, $\lim_{j\to\infty} d_p(u_t^j, v_t^j) = d_p(u_t, v_t)$. By Lemma 4.3, we have

$$d_p(u_t^j, v_t^j) \le (1 - t)d_p(u_0^j, v_0^j) + td_p(u_1^j, v_1^j).$$

Taking limit $j \to \infty$, we get that

$$d_p(u_t, v_t) \le (1-t)d_p(u_0, v_0) + td_p(u_1, v_1).$$

4.3. Buseman Convexity for big classes when p > 1. As in the case of Lidskii-type inequality in Section 3, we will first show the Buseman convexity for the analytic singularity case, and then for the minimal singularity case for the big classes.

Assume that θ is a smooth closed real (1, 1)-form representing a big cohomology class. Let ψ be a θ -psh function with analytic singularities. We will first show that $(\mathcal{E}^p(X, \theta, \psi), d_p)$ is Buseman convex.

Lemma 4.5. For p > 1, the metric space $(\mathcal{E}^p(X, \theta, \psi), d_p)$ is a Buseman convex metric space. This means that if $u_0, u_1, v_0, v_1 \in \mathcal{E}^p(X, \theta, \psi)$, and u_t and v_t are the weak geodesics joining u_0, u_1 , and v_0, v_1 respectively, then

$$d_p(u_t, v_t) \le (1-t)d_p(u_0, v_0) + td_p(u_1, v_1)$$

Proof. Recall from Section 2.4.2 that there exists a modification $\mu : \tilde{X} \to X$, a big and nef cohomology class $\tilde{\theta}$ and a bounded $\tilde{\theta}$ -psh function g on \tilde{X} such that the metric d_p on $\mathcal{E}^p(X, \theta, \psi)$ is given by

$$d_p(u_0, u_1) = d_p(\tilde{u}_0, \tilde{u}_1)$$

where $\mathcal{E}^p(X, \theta, \psi) \ni u \mapsto \tilde{u} := (u - \psi) \circ \mu + g \in \mathcal{E}^p(\tilde{X}, \tilde{\theta})$ is a bijection.

By Theorem 4.4, the metric space $(\mathcal{E}^p(\tilde{X}, \tilde{\theta}), d_p)$ is Buseman convex. Therefore,

$$d_p(u_t, v_t) = d_p(\tilde{u}_t, \tilde{v}_t) \le (1 - t)d_p(\tilde{u}_0, \tilde{v}_0) + td_p(\tilde{u}_1, \tilde{v}_1) = (1 - t)d_p(u_0, v_0) + td_p(u_1, v_1).$$

Now we can show that the metric space $(\mathcal{E}^p(X,\theta), d_p)$ is Buseman convex for p > 1.

Theorem 4.6. For p > 1, the metric space $(\mathcal{E}^p(X,\theta), d_p)$ is a Buseman convex metric space. This means that if $u_0, u_1, v_0, v_1 \in \mathcal{E}^p(X,\theta)$, and u_t and v_t are the weak geodesics joining u_0, u_1 , and v_0, v_1 respectively, then

$$d_p(u_t, v_t) \le (1-t)d_p(u_0, v_0) + td_p(u_1, v_1).$$

Proof. As described in Section 2.4.3, let ψ_k be an increasing sequence of θ -psh functions with analytic singularities such $\psi_k \nearrow V_{\theta}$ a.e.

We denote the projections of u_0, u_1, v_0 , and v_1 to $\mathcal{E}^p(X, \theta, \psi_k)$ by $u_0^k = P_\theta[\psi_k](u_0), u_1^k = P_\theta[\psi_k](u_1), v_0^k = P_\theta[\psi_k](v_0)$, and $v_1^k = P_\theta[\psi_k](v_1)$ respectively. We denote by u_t^k and v_t^k the weak geodesics joining u_0^k, u_1^k , and v_0^k, v_1^k respectively.

By a proof similar to Lemma 4.2 we can show that

(9)
$$\lim_{k \to \infty} d_p(u_t^k, P_{\theta}[\psi_k](u_t)) = 0 \quad \text{and} \quad \lim_{k \to \infty} d_p(v_t^k, P_{\theta}[\psi_k](v_t)) = 0.$$

See the proof of [Gup24, Theorem 8.3, Equation 26] for more details.

From Equation (8) and 9, we get

$$d_p(u_t, v_t) = \lim_{k \to \infty} d_p(P_{\theta}[\psi_k](u_t), P_{\theta}[\psi_k](v_t)) = \lim_{k \to \infty} d_p(u_t^k, v_t^k).$$

By Lemma 4.5, we can write

$$d_p(u_t^k, v_t^k) \le (1-t)d_p(u_0^k, v_0^k) + td_p(u_1^k, v_1^k).$$

Combining the last two equations we get

$$d_p(u_t, v_t) = \lim_{k \to \infty} d_p(u_t^k, v_t^k) \le \lim_{k \to \infty} (1 - t) d_p(u_0^k, v_0^k) + t d_p(u_1^k, v_1^k) = (1 - t) d_p(u_0, v_0) + t d_p(u_1, v_1).$$

Combined with Theorem 4.1, this finishes the proof of Theorem 1.2.

5. Space of geodesic rays

In this section, we will study the space of geodesic rays and prove Theorem 1.3. Let $u \in \mathcal{E}^p(X, \theta)$. A geodesic ray $\{u_t\}_t$ is a finite *p*-energy geodesic ray emanating from *u* if $u_t \to u$ as $t \to 0$ in $L^1(X)$ and $u_t \in \mathcal{E}^p(X, \theta)$ for all t > 0. We denote by \mathcal{R}^p_u the set of all finite *p*-energy geodesic rays emanating from *u*.

We construct a metric d_p^c on \mathcal{R}_u^p as follows. If $\{u_t\}_t, \{v_t\}_t \in \mathcal{R}_u^p$ are two finite energy geodesic rays, then we define

(10)
$$d_p^c(\{u_t\}_t, \{v_t\}_t) = \lim_{t \to \infty} \frac{d_p(u_t, v_t)}{t}$$

Due to Buseman convexity of d_p from Theorem 1.2, we know that if $t_1 \ge t_2$, then

$$d_p(u_{t_2}, v_{t_2}) \le \left(1 - \frac{t_2}{t_1}\right) d_p(u_0, v_0) + \frac{t_2}{t_1} d_p(u_{t_1}, v_{t_1}).$$

Since $u_0 = v_0 = u$, we have

$$\frac{d_p(u_{t_2}, v_{t_2})}{t_2} \le \frac{d_p(u_{t_1}, v_{t_1})}{t_1}.$$

Thus the limit in Equation (10) is increasing and is bounded from above due to the triangle inequality. This is because $d_p(u_t, v_t) \leq d_p(u_t, u) + d_p(v_t, u) = td_p(u_1, u) + td_p(v_1, u)$. Thus

$$\frac{d_p(u_t, v_t)}{t} \le d_p(u_1, u) + d_p(v_1, u).$$

Thus the limit in Equation (10) is well defined.

Lemma 5.1. Equation (10) defines a metric on \mathcal{R}_{u}^{p} .

Proof. Let $\{u_t\}_t, \{v_t\}_t, \{w_t\}_t \in \mathcal{R}^p_u$ be three finite *p*-energy geodesic rays. $d_p^c(\{u_t\}_t, \{v_t\}_t) \ge 0$ because $d_p(u_t, v_t) \ge 0$.

If $d_p^c(\{u_t\}_t, \{v_t\}_t) = 0$, then the function $f(t) = d_p(u_t, v_t)/t$ is an increasing function that satisfies f(0) = 0 and $\lim_{t\to\infty} f(t) = 0$. Therefore f(t) = 0 for all $t \ge 0$. Thus $d_p(u_t, v_t) = 0$ for all $t \ge 0$. Therefore $u_t = v_t$ for all $t \ge 0$. So $\{u_t\}_t = \{v_t\}_t$. This proves non-degeneracy of d_p^c .

The triangle inequality for d_p^c follows from the triangle inequality for d_p .

$$d_{p}^{c}(\{u_{t}\}_{t},\{v_{t}\}_{t}) = \lim_{t \to \infty} \frac{d_{p}(u_{t},v_{t})}{t}$$
$$\leq \lim_{t \to \infty} \frac{d_{p}(u_{t},w_{t})}{t} + \lim_{t \to \infty} \frac{d_{p}(w_{t},v_{t})}{t}$$
$$= d_{p}^{c}(\{u_{t}\}_{t},\{w_{t}\}_{t}) + d_{p}^{c}(\{w_{t}\}_{t},\{v_{t}\}_{t}).$$

The next theorem shows that the metric space (\mathcal{R}^p_u, d^c_p) are all isometric for all $u \in \mathcal{E}^p(X, \theta)$.

Theorem 5.2. Given $u, v \in \mathcal{E}^p(X, \theta)$, there exists a map $P_{uv} : \mathcal{R}^p_u \to \mathcal{R}^p_v$ that induces a bijective isometry $P_{uv} : (\mathcal{R}^p_u, d^c_p) \to (\mathcal{R}^p_v, d^c_p)$.

Proof. Let $\{u_t\}_t \in \mathcal{R}^p_u$. We will construct a finite energy geodesic ray $\{v_t\}_t$ starting from v.

First, assume that $u \ge v$. We can construct a finite energy geodesic $[0, t] \ni l \mapsto v_l^t \in \mathcal{E}^p(X, \theta)$ joining $v = v_0^t$ and $u_t = v_t^t$. Consider $0 \le t \le t'$. Since $[0, t'] \ni l \mapsto u_l$ is a finite energy geodesic joining u and $u_{t'}$, whereas $[0, t'] \ni l \mapsto v_l^{t'}$ is a finite energy geodesic joining v and $u_{t'}$. Since $u \ge v$, by the comparison principle, $v_t^{t'} \le u_t = v_t^t$. Again applying the comparison principle, we get that for $l \le t$, $v_l^{t'} \le v_l^t$.

Moreover, by Theorem 1.2,

$$d_p(v_l^t, u_l) \le \left(1 - \frac{l}{t}\right) d_p(u, v).$$

By Lemma 2.13, the map $[0, \infty) \ni l \mapsto v_l := \lim_{t\to\infty} v_l^t \in \mathcal{E}^p(X, \theta)$ is a finite energy geodesic ray. Note that the limit geodesic v_l still emanates from v. This is because by Lemma 2.13, the path $[0, t] \ni l \mapsto v_l \in \mathcal{E}^p(X, \theta)$ is a geodesic joining v and v_t . Because $v, v_t \in E^p(X, \theta)$ the result follows from the following argument. As $d_p(v_l, v_0) = \frac{l}{t}$, we get that $\lim_{l\to 0} d_p(v_l, v_0) = 0$. Thus by [Gup24, Lemma 4.12] and [Gup23, Theorem 1.2], we get that $v_l \to v_0$ in $L^1(X)$. See [Xia24, Proposition 4.2.1] for a more general result. Now, taking the limit $t \to \infty$ in the equation above we get that $d_p(v_l, u_l) \leq d_p(u, v)$.

A similar proof works when $u \leq v$.

More generally, we can consider $h = P_{\theta}(u, v) \in \mathcal{E}^p(X, \theta)$. Now $h \leq u, v$. Thus from above, we can construct a finite energy geodesic ray $\{h_t\}_t \in \mathcal{R}^p_h$ that satisfies $d_p(h_t, u_t) \leq d_p(h, u)$. We can now also construct a finite energy geodesic ray $\{v_t\}_t \in \mathcal{R}^p_v$ such that $d_p(h_t, v_t) \leq d_p(h, v)$. Thus by the triangle inequality $d_p(u_t, v_t) \leq d_p(u_t, h_t) + d_p(h_t, v_t) \leq d_p(u, h) + d_p(h, v)$. Thus $\{u_t\}_t, \{v_t\}_t$ are parallel geodesic rays.

To see that $P_{uv}: (\mathcal{R}^p_u, d^c_p) \to (\mathcal{R}^p_v, d^c_p)$ is an isometry consider $\{u^0_t\}_t, \{u^1_t\}_t \in \mathcal{R}^p_u$. Let $P_{uv}(\{u^i_t\}_t) = \{v^i_t\}_t$ for i = 0, 1. By the triangle inequality we have

$$\begin{split} d_p^c(\{v_t^0\}_t, \{v_t^1\}_t) &= \lim_{t \to \infty} \frac{d_p(v_t^0, v_t^1)}{t} \\ &\leq \lim_{t \to \infty} \frac{d_p(v_t^0, u_t^0) + d_p(u_t^0, u_t^1) + d_p(u_t^1, v_t^1)}{t} \\ &= \lim_{t \to \infty} \frac{d_p(u_t^0, u_t^1)}{t} \\ &= d_p^c(\{u_t^0\}_t, \{u_t^1\}_t). \end{split}$$

In the second line, we used that fact that $d_p(u_t^i, v_t^i)$ remain bounded as $t \to \infty$ for i = 0, 1. The other side inequality is obtained similarly. Thus $d_p^c(\{u_t^0\}_t, \{u_t^1\}_t) = d_p^c(\{v_t^0\}_t, \{v_t^1\}_t)$.

Since the spaces (\mathcal{R}^p_u, d^c_p) are all isometric to each other, it is good enough to study just one of them. We will fix $u = V_{\theta}$ and use \mathcal{R}^p_{θ} to denote $\mathcal{R}^p_{V_{\theta}}$. The goal of the rest of the section is to prove that $(\mathcal{R}^p_{\theta}, d^c_p)$ is a complete geodesic metric space. First, we prove

Lemma 5.3. The metric space $(\mathcal{R}^p_{\theta}, d^c_p)$ is a complete metric space.

Proof. Let $\{u_t^j\}_t \in \mathcal{R}^p_{\theta}$ be a Cauchy sequence of finite energy geodesic rays. This means that $d_p^c(\{u_t^j\}_t, \{u_t^k\}_t) \to 0$ as $j, k \to \infty$. For a fixed $t \ge 0$, we know that $d_p(u_t^j, u_t^k) \le t d_p^c(\{u_t^j\}_t, \{u_t^k\}_t)$. Thus u_t^j is a Cauchy sequence in $(\mathcal{E}^p(X, \theta), d_p)$ with limit point $u_t \in \mathcal{E}^p(X, \theta)$. By the endpoint stability Lemma 2.13, u_t is a finite energy geodesic ray. Now we observe that

$$d_p^c(\{u_t^j\}_t, \{u_t\}_t) = \lim_{t \to \infty} \frac{d_p(u_t^j, u_t)}{t}$$
$$= \lim_{t \to \infty} \lim_{k \to \infty} \frac{d_p(u_t^j, u_t^k)}{t}$$
$$\leq d_p^c(\{u_t^j\}_t, \{u_t^k\}_t)$$

which can be arbitrarily small. Therefore, $d_p^c(\{u_t^j\}_t, \{u_t\}_t) \to 0$ as $j \to \infty$.

First, we will show that for p > 1, $(\mathcal{R}^p_{\theta}, d^c_p)$ is a complete geodesic metric space. We have already seen that it is a complete metric space. We just need to show that there are geodesics in this space.

Theorem 5.4. For p > 1, $(\mathcal{R}^p_{\theta}, d^c_p)$ is a complete geodesic metric space.

Proof. Let $\{u_t^0\}_t, \{u_t^1\}_t \in \mathcal{R}^p_\theta$ be two finite energy geodesic rays. We will construct finite energy geodesic rays $\{u_t^\alpha\}_t \in \mathcal{R}^p_\theta$ for $0 \le \alpha \le 1$, such that the path $[0,1] \ni \alpha \mapsto \{u_t^\alpha\}_t \in \mathcal{R}^p_\theta$ is a d_p^c -geodesic.

First, consider the finite energy geodesic $[0,1] \ni \alpha \mapsto v_t^{\alpha} \in \mathcal{E}^p(X,\theta)$ joining u_t^0 and u_t^1 . Let $[0,t] \ni l \mapsto w_l^{\alpha,t} \in \mathcal{E}^p(X,\theta)$ be the finite energy geodesic joining $w_0^{\alpha,t} := V_{\theta}$ and $w_t^{\alpha,t} = v_t^{\alpha}$. We claim that for a fixed $l \ge 0$, $w_l^{\alpha,t} \in \mathcal{E}^p(X,\theta)$ is d_p -Cauchy as $t \to \infty$. Thus there exist $u_l^{\alpha} \in \mathcal{E}^p(X,\theta)$ such that $\lim_{t\to\infty} d_p(w_l^{\alpha,t}, u_l^{\alpha}) = 0$. By the endpoint stability Lemma 2.13, $\{u_t^{\alpha}\}_t$ is a finite energy geodesic ray.

Now we will prove the claim. Let $s \leq t$. By Theorem 1.2,

(11)
$$\frac{d_p(u_s^0, w_s^{\alpha, t})}{s} \le \frac{d_p(u_t^0, v_t^{\alpha})}{t} = \alpha \frac{d_p(u_t^0, u_t^1)}{t} \le \alpha d_p^c(\{u_t^0\}_t, \{u_t^1\}_t)$$

Similarly, we have

(12)
$$\frac{d_p(u_s^1, w_s^{\alpha, t})}{s} \le \frac{d_p(u_t^1, v_t^{\alpha})}{t} = (1 - \alpha) \frac{d_p(u_t^0, u_t^1)}{t} \le (1 - \alpha) d_p^c(\{u_t^0\}_t, \{u_t^1\}_t)$$

Since $\frac{d_p(u_s^0, u_s^1)}{s} \nearrow d_p^c(\{u_t^0\}_t, \{u_t^1\}_t)$. We can write $d_p^c(\{u_t^0\}_t, \{u_t^1\}_t) \le (1 + \varepsilon(s)) \frac{d_p(u_s^0, u_s^1)}{s}$ where $\varepsilon(s) \to 0$ as $s \to \infty$. Combining with Equations (11) and (12), we get

$$d_p(u_s^0, w_s^{\alpha, t}) \le (\alpha + \varepsilon(s))d_p(u_s^0, u_s^1)$$

and

$$d_p(u_s^1, w_s^{\alpha, t}) \le (1 - \alpha + \varepsilon(s))d_p(u_s^0, u_s^1)$$

By Corollary 2.11 of the uniform convexity, we can write

$$d_p(v_s^{\alpha}, w_s^{\alpha, t}) \le C\varepsilon(s)^{\frac{1}{r}} d_p(u_s^0, u_s^1).$$

This is the only place where we need the condition that p > 1. For $l \leq s$, using Theorem 1.2,

$$\frac{d_p(w_l^{\alpha,s}, w_l^{\alpha,t})}{l} \le \frac{d_p(v_s^{\alpha}, w_s^{\alpha,t})}{s} \le C(\varepsilon(s))^{\frac{1}{r}} d_p^c(\{u_t^0\}_t, \{u_t^1\}_t)$$

As $\varepsilon(s)$ can be made arbitrarily small, we get that $w_l^{\alpha,t}$ is a d_p -Cauchy sequence as $t \to \infty$. By [Gup24, Main Theorem], the metric d_p is complete, so there exists $u_l^{\alpha} \in \mathcal{E}^p(X, \theta)$ such that $\lim_{t\to\infty} d_p(w_l^{\alpha,t}, u_l^{\alpha}) = 0$.

Now we will prove that the map $[0,1] \ni \alpha \mapsto \{u_t^{\alpha}\}_t \in \mathcal{R}^p_{\theta}$ is a d_p^c -geodesic. Taking the limit $t \to \infty$ in Equations (11) and (12), we get that

$$\frac{d_p(u_s^0, u_s^\alpha)}{s} \le \alpha d_p^c(\{u_t^0\}_t, \{u_t^1\}_t).$$

and

$$\frac{d_p(u_s^1, u_s^\alpha)}{s} \le (1 - \alpha) d_p^c(\{u_t^0\}_t, \{u_t^1\}_t).$$

Taking the limit $s \to \infty$, we get that

 $d_p^c(\{u_t^0\}_t, \{u_t^{\alpha}\}_t) \le \alpha d_p^c(\{u_t^0\}_t, \{u_t^1\}_t) \quad \text{and} \quad d_p^c(\{u_t^1\}_t, \{u_t^{\alpha}\}_t) \le (1-\alpha) d_p^c(\{u_t^0\}_t, \{u_t^1\}_t)$ Combining with the triangle inequality, we get

$$d_p^c(\{u_t^0\}_t, \{u_t^\alpha\}_t) = \alpha d_p^c(\{u_t^0\}_t, \{u_t^1\}_t) \quad \text{and} \quad d_p^c(\{u_t^1\}_t, \{u_t^\alpha\}_t) = (1 - \alpha) d_p^c(\{u_t^0\}_t, \{u_t^1\}_t)$$

For $0 \le \alpha \le \beta \le 1$, the triangle inequality implies

$$d_p^c(\{u_t^{\alpha}\}_t, \{u_t^{\beta}\}_t) \ge d_p^c(\{u_t^{0}\}_t, \{u_t^{\beta}\}_t) - d_p^c(\{u_t^{0}\}_t, \{u_t^{\alpha}\}_t) = (\beta - \alpha)d_p^c(\{u_t^{0}\}_t, \{u_t^{1}\}_t).$$

For the other side, we again use Theorem 1.2. We notice that

$$\frac{d_p(w_l^{\alpha,t}, w_l^{\beta,t})}{l} \le \frac{d_p(v_t^{\alpha}, v_t^{\beta})}{t} = (\beta - \alpha) \frac{d_p(u_t^0, u_t^1)}{t}$$

Taking limit $t \to \infty$, we get

$$\frac{d_p(u_l^{\alpha}, u_l^{\beta})}{l} \le (\beta - \alpha) d_p^c(\{u_t^0\}_t, \{u_t^1\}_t)$$

Now taking limit $l \to \infty$, we get

$$d_p^c(\{u_t^{\alpha}\}_t, \{u_t^{\beta}\}_t) \le (\beta - \alpha) d_p^c(\{u_t^{0}\}_t, \{u_t^{1}\}_t).$$

Thus

$$d_p^c(\{u_t^{\alpha}\}_t, \{u_t^{\beta}\}_t) = (\beta - \alpha)d_p^c(\{u_t^{0}\}_t, \{u_t^{1}\}_t)$$

proving that the map $[0,1] \ni \alpha \mapsto \{u_t^{\alpha}\}_t \in \mathcal{R}^p_{\theta}$ is a d_p^c -geodesic.

Now we need to extend this proof to p = 1 by approximating it from p > 1. First, we will show that the d_p^c -geodesics depend only on the end-points, and not the value of p, just like in the metric space $(\mathcal{E}^p(X,\theta), d_p)$, where the metric geodesics are described by the weak geodesics whose construction does not depend on p.

Lemma 5.5. If $1 \leq p' \leq p$, then $\mathcal{R}^p_{\theta} \subset \mathcal{R}^{p'}_{\theta}$. Moreover, if $[0,1] \ni \alpha \mapsto \{u^{\alpha}_t\}_t \in \mathcal{R}^p_{\theta}$ is the d^c_p -geodesic joining $\{u^0_t\}_t$ and $\{u^1_t\}_t$, then $[0,1] \ni \alpha \mapsto \{u^{\alpha}_t\}_t$ is also the $d^c_{p'}$ -geodesic.

Proof. Recall from the proof of Theorem 5.4 that the geodesics $[0,1] \ni \alpha \mapsto v_t^{\alpha} \in \mathcal{E}^p(X,\theta)$ joining u_t^0 and u_t^1 do not depend on the value of p. Similarly, the geodesics $[0,t] \ni l \mapsto w_l^{\alpha,t} \in \mathcal{E}^p(X,\theta)$ joining V_{θ} and v_t^{α} also do not depend on the value of p. If we can show that the d_p limit u_l^{α} of $w_l^{\alpha,t}$ as $t \to \infty$ does not depend on p as well, we will be done.

If we can show that the d_p limit u_l^{α} of $w_l^{\alpha, c}$ as $t \to \infty$ does not depend on p as well, we will be done. Recall from Lemma 2.12, that $d_{p'}(\cdot, \cdot) \leq d_p(\cdot, \cdot) \operatorname{Vol}(\theta)^{\frac{1}{p'} - \frac{1}{p}}$ on $\mathcal{E}^p(X, \theta)$. Thus if $\lim_{t\to\infty} d_p(w_l^{\alpha, t}, u_l^{\alpha}) = 0$, then

$$\lim_{t \to \infty} d_{p'}(w_l^{\alpha,t}, u_l^{\alpha}) \le \lim_{t \to \infty} d_p(w_l^{\alpha,t}, u_l^{\alpha}) \operatorname{Vol}(\theta)^{\frac{1}{p'} - \frac{1}{p}} = 0$$

Knowing that the $d_{p'}$ limit of $w_l^{\alpha,t}$ as $t \to \infty$ is u_l^{α} irrespective of the value of p', the proof of the fact that $[0,1] \ni \alpha \mapsto \{u_t^{\alpha}\}_t \in \mathcal{R}_{\theta}^{p'}$ is a $d_{p'}^c$ -geodesic goes through as in Theorem 5.4.

Now we will establish the version of Theorem 5.4 for p = 1. Before that, we need the following

Lemma 5.6. For $p \ge 1$, if $\{u_t^j\}_t, \{u_t\} \in \mathcal{R}^p_\theta$ are finite energy geodesic rays such that $u_t^j \searrow u_t$ then

$$\lim_{j \to \infty} d_p^c(\{u_t^j\}_t, \{u_t\}_t) = 0$$

Proof. By Lemma 2.3, the map $t \mapsto \sup_X u_t$ and $t \mapsto \sup_X u_t^j$ is linear. Therefore, by replacing u_t, u_t^j by $u_t - Ct, u_t^j - Ct$, we can assume that $0 \ge u_t^j \ge u_t$.

Applying the Lidskii-type inequality from Theorem 3.3, we get that

$$\frac{d_p^p(u_t^j, u_t)}{t^p} \le \frac{d_p^p(u_t, 0) - d_p^p(u_t^j, 0)}{t^p} = d_p^p(u_1, 0) - d_p^p(u_1^j, 0).$$

Since $u_1^j \searrow u_1$, by Lemma 2.9, $\lim_{j\to\infty} d_p(u_1^j, 0) = d_p(u_1, 0)$. Thus

$$\lim_{j \to \infty} \frac{d_p(u_t^j, u_t)}{t} = 0$$

uniformly in t. Therefore, $\lim_{j\to\infty} d_p^c(\{u_t^j\}_t, \{u_t\}_t) = 0.$

With the help of Lemma 5.6 and Theorem 2.4, we can prove the existence of geodesics in \mathcal{R}^1_{θ} .

Theorem 5.7. For $p \ge 1$, the space $(\mathcal{R}^p_{\theta}, d^c_p)$ is a complete geodesic metric.

Proof. We have already proved the theorem for p > 1. We just need to prove it for p = 1 now. Let $\{u_t^0\}_t, \{u_t^1\}_t \in \mathcal{R}^1_{\theta}$ be two finite energy geodelisc rays. As in the proof of Theorem 5.4, we can construct geodesics $[0,1] \ni \alpha \mapsto v_t^{\alpha} \in \mathcal{E}^1(X,\theta)$ joining u_t^0 and u_t^1 . We can also construct geodesics $[0,t] \ni l \mapsto w_l^{\alpha,t} \in \mathcal{E}^1(X,\theta)$ joining V_{θ} and v_t^{α} . We used p > 1 to construct u_l^{α} as the d_p limit of $w_l^{\alpha,t}$ as $t \to \infty$. In the case p = 1, we will use approximation to construct u_l^{α} .

Fix any p > 1. From Theorem 2.4 we can construct geodesic rays $\{u_t^{0,j}\}_t, \{u_t^{1,j}\}_t \in \mathcal{R}^p_{\theta}$, such that $u_t^{0,j} \searrow u_t^0$ and $u_t^{1,j} \searrow u_t^1$ as $j \to \infty$ and for all $t \ge 0$.

Let $[0,1] \ni \alpha \mapsto \{u_t^{\alpha,j}\}_t \in \mathcal{R}^p_{\theta} \subset \mathcal{R}^1_{\theta}$ be the d_p^c -geodesic constructed in Theorem 5.4. By Lemma 5.5 it is also the d_1^c -geodesic. Recall that we constructed $u_l^{\alpha,j}$ as d_p -limit (hence d_1 -limit) of $w_l^{\alpha,t,j}$ as $t \to \infty$. Since $w_l^{\alpha,t,j} \ge w_l^{\alpha,t}$, we have

$$d_1(w_l^{\alpha,t,j}, w_l^{\alpha,t}) = I(w_l^{\alpha,t,j}) - I(w_l^{\alpha,t})$$

= $\frac{l}{t}(I(v_t^{\alpha,j}) - I(v_t^{\alpha}))$
= $\frac{l}{t}((1 - \alpha)I(u_t^{0,j}) + I(u_t^{1,j}) - (1 - \alpha)I(u_t^0) - \alpha I(u_t^1))$
= $l(1 - \alpha)(I(u_1^{0,j}) - I(u_1^0)) + l\alpha(I(u_1^{1,j}) - I(u_1^1)).$

As $j \to \infty$, $u_1^{0,j} \searrow u_1^0$ and $u_1^{1,j} \searrow u_1^1$. Therefore $I(u_1^{0,j}) \to I(u_1^0)$ and $I(u_1^{1,j}) \to I(u_1^1)$. Therefore, $d_1(w_l^{\alpha,t,j}, w_l^{\alpha,t}) \to 0$ as $j \to \infty$ uniformly over t.

Now since $u_l^{\alpha,j}$ is a decreasing sequence and is bounded from below because $I(u_l^{\alpha,j})$ is bounded. Therefore, the limit $\lim_{j\to\infty} u_l^{\alpha,j} =: u_l^{\alpha} \in \mathcal{E}^1(X,\theta)$. By the proof of Theorem 5.4, we will be done, if we can show that $\lim_{t\to\infty} d_1(w_l^{\alpha,t}, u_l^{\alpha}) = 0$. To see this notice that

$$d_1(w_l^{\alpha,t}, u_l^{\alpha}) \le d_1(w_l^{\alpha,t}, w_l^{\alpha,t,j}) + d_1(w_l^{\alpha,t,j}, u_l^{\alpha,j}) + d_1(u_l^{\alpha,j}, u_l^{\alpha}).$$

We have seen above that the first term goes to 0 as $j \to \infty$ uniformly in t. The last term also goes to 0 as $j \to \infty$. For a fixed j, the second term gets arbitrarily small for large t. Therefore, $\lim_{t\to\infty} d_1(w_l^{\alpha,t}, u_l^{\alpha}) \to 0$.

6. The Ross-Witt Nyström correspondence

The notion of test curves, introduced by Ross-Witt Nyström in [RW14] in the Kähler setting, and further studied by Darvas-Di Nezza-Lu [DDL18a], Darvas-Zhang [DZ23], Darvas-Xia [DX22] and Darvas-Xia-Zhang [DXZ23], is a powerful tool to study geodesic rays in connection with K-stability.

Definition 6.1. A map $\mathbb{R} \ni \tau \mapsto \psi_{\tau} \in PSH(X, \theta)$ is a test curve, denoted by $\{\psi_{\tau}\}_{\tau}$, if

- (1) The map $\mathbb{R} \ni \tau \mapsto \psi_{\tau}(x)$ is decreasing, usc, concave for all $x \in X$, and
 - (2) $\psi_{\tau} = -\infty$ for all τ big enough, and
 - (3) $\psi_{\tau} \nearrow V_{\theta}$ a.e. as $\tau \to -\infty$.

From condition (2) above, we can define

$$\tau_{\psi}^{+} = \inf\{\tau \in \mathbb{R} : \psi_{\tau} \equiv -\infty\}$$

Test curves are Legendre dual of sublinear subgeodesic rays starting at V_{θ} [DZ23]. Several properties of the geodesic rays can be detected by the corresponding test curves. Based on that we define several subclasses of test curves.

Definition 6.2. A test curve $\{\psi_{\tau}\}_{\tau}$ can have the following properties.

- (1) It is maximal if $P_{\theta}[\psi_{\tau}](0) = \psi_{\tau}$ for all $\tau \in \mathbb{R}$.
- (2) It has finite p-energy for $p \ge 1$, if

$$\int_{-\infty}^{\tau_{\psi}^+} (-\tau + \tau_{\psi}^+)^{p-1} \left(\int_X \theta_{V_{\theta}}^n - \int_X \theta_{\psi_{\tau}}^n \right) d\tau < \infty$$

(3) It is *bounded* if for all τ negative enough, $\psi_{\tau} = V_{\theta}$. In this case, we define

$$\tau_{\psi}^{-} = \sup\{\tau \in \mathbb{R} : \psi_{\tau} = V_{\theta}\}.$$

When p = 1, the notion of finite *p*-energy test curves was introduced by Darvas-Xia [DX22] in the Kähler setting, and by Darvas-Zhang [DZ23] in the big setting.

The Ross-Witt Nyström correspondence is the observation of Ross-Witt Nyström [RW14] that test curves and subgeodesic rays are dual to each other via Legendre transform. If $\{u_t\}_t$ is a sublinear sub geodesic ray starting at V_{θ} , then its Legendre transform is given by

$$\hat{u}_{\tau}(x) = \inf_{t>0} (u_t(x) - t\tau).$$

If $\{\psi_{\tau}\}_{\tau}$ is a test curve, then its inverse Legendre transform is given by

$$\check{\psi}_t(x) = \sup_{\tau \in \mathbb{R}} (\psi_\tau(x) - t\tau).$$

In particular, we have

Theorem 6.3 ([DZ23, Theorem 3.7]). The Legendre transform $\{\psi_{\tau}\}_{\tau} \mapsto \{\check{\psi}_t\}_t$ is a bijection with inverse $\{u_t\}_t \mapsto \{\hat{u}_{\tau}\}_{\tau}$ between

- (1) test curves and sublinear subgeodesic rays starting at V_{θ} ;
- (2) maximal test curves and geodesic rays starting at V_{θ} ;
- (3) bounded maximal test curves and geodesic rays with minimal singularity type. Moreover, in this case, $V_{\theta} + \tau_{\psi}^{-} t \leq \check{\psi}_{t} \leq V_{\theta} + \tau_{\psi}^{+} t$.

Darvas-Xia [DX22, Theorem 3.7] in the Kähler setting, and Darvas-Zhang [DZ23, Theorem 3.9] in the big setting proved that the Ross-Witt Nyström correspondence as described above is a bijection between maximal finite 1-energy test curves and finite 1-energy geodesic rays. In this section, we will generalize this statement to finite p-energy test curves and geodesic rays.

For that, first, we need a lemma about the speed of the geodesic segments, which can be seen as a converse of [Gup24, Lemma 5.3].

Lemma 6.4. If $u_0 \in \mathcal{H}_{\theta}$, $u_1 \in PSH(X, \theta)$, and $[0, 1] \ni t \mapsto u_t \in PSH(X, \theta)$ is the weak geodesic joining u_0 and u_1 such that

$$\int_X |\dot{u}_0|^p \theta_{u_0}^n < \infty$$

then $u_1 \in \mathcal{E}^p(X, \theta)$.

Proof. First, assume that $u_0 \ge u_1 + 1$, so that we can find $u_1^j \in \mathcal{H}_{\theta}$ such that $u_1^j \searrow u_1$ and $u_0 \ge u_1^j$. Let $[0,1] \ni t \mapsto u_t^j \in \mathcal{E}^p(X,\theta)$ be the weak geodesic joining u_0 and u_1^j . The geodesics are decreasing, therefore $\dot{u}_0, \dot{u}_0^j \le 0$. By Theorem 2.7,

$$d_p^p(u_0, u_1^j) = \int_X |\dot{u}_0^j|^p \theta_{u_0}^n = \int_X (-\dot{u}_0^j)^p \theta_{u_0}^n.$$

Since $u_1^j \ge u_1$, the weak geodesics satisfy $u_t^j \ge u_t$. Therefore,

$$\dot{u}_0^j = \lim_{t \to 0} \frac{u_t^j - u_0}{t} \ge \lim_{t \to 0} \frac{u_t - u_0}{t} = \dot{u}_0.$$

Hence, $(-\dot{u}_0^j)^p \le (-\dot{u}_0)^p$. Now,

$$d_p^p(u_0, u_1^j) = \int_X (-\dot{u}_0^j)^p \theta_{u_0}^n \le \int_X |\dot{u}_0|^p \theta_{u_0}^n < \infty.$$

Therefore, u_1^j is a d_p -bounded decreasing sequence such that $u_1^j \searrow u_1$. Therefore by Lemma 2.14 $u_1 \in \mathcal{E}^p(X, \theta)$.

Now we drop the assumption that $u_0 \ge u_1 + 1$. Let C > 0 such that $u_1 - C + 1 \le u_0$. Let $w_1 = u_1 - C$, so that $u_0 \ge w_1 + 1$. If w_t is the geodesic joining u_0 and w_1 , then $w_t = u_t - Ct$. Therefore, $\dot{w}_0 = \dot{u}_0 - C$. Hence,

$$\int_X |\dot{w}_0|^p \theta_{u_0}^n = \int_X |\dot{u}_0 - C|^p \theta_{u_0}^n \le 2^{p-1} \left(\int_X |\dot{u}_0|^p \theta_{u_0}^n + C^p \operatorname{Vol}(\theta) \right) < \infty.$$

By the argument above, we find that $w_1 \in \mathcal{E}^p(X, \theta)$, therefore, $w_1 + C = u_1 \in \mathcal{E}^p(X, \theta)$ as well.

This Lemma allows us to prove

Theorem 6.5. For $p \ge 1$, the Legendre Transform $\{u_t\}_t \mapsto \{\hat{u}_\tau\}_\tau$ is the bijective map between finite *p*-energy geodesic rays in \mathcal{R}^p_{θ} and maximal finite *p*-energy test curves.

Proof. First, we assume that $\sup_X u_t = 0$ for all $t \ge 0$. In this case $\tau_{\hat{u}}^+ = 0$. To see this we observe that u_t decreases as $t \to \infty$. As $\sup_X u_t = 0$, by Hartog's Lemma (see [GZBook, Proposition 8.4]) $\sup_X (\lim_{t\to\infty} u_t) = 0$. Therefore, when $\tau = 0$,

$$\hat{u}_0 = \inf_{t \ge 0} u_t = \lim_{t \to \infty} u_t \not\equiv -\infty$$

Whereas for $\tau > 0$,

$$\hat{u}_{\tau} = \inf_{t \ge 0} (u_t - t\tau) \equiv -\infty.$$

Thus $\tau_{\hat{u}}^+ = 0$. Again since u_t is decreasing $\dot{u}_0 \leq 0$. We will show that

(13)
$$\int_X |\dot{u}_0|^p \theta_{u_0}^n = p \int_{-\infty}^0 (-\tau)^{p-1} \left(\int_X \theta_{V_\theta}^n - \int_X \theta_{\dot{u}_\tau}^n \right) d\tau.$$

To see this we start with the left-hand side

$$\int_{X} |\dot{u}_{0}|^{p} \theta_{u_{0}}^{n} = p \int_{X} (-\dot{u}_{0})^{p} \theta_{V_{\theta}}^{n}$$
$$= p \int_{0}^{\infty} \tau^{p-1} \theta_{V_{\theta}}^{n} (\{\dot{u}_{0} < -\tau\}) d\tau$$

Changing the variable from τ to $-\tau$, we can continue

$$= p \int_{-\infty}^{0} (-\tau)^{p-1} \theta_{V_{\theta}}^{n} (\{ \dot{u}_{0} < \tau \}) d\tau$$

Recall that $\hat{u}_{\tau} = \inf_{t \ge 0} (u_t - t\tau)$. Thus the set $\{\dot{u}_0 \ge \tau\} = \{\hat{u}_{\tau} = u_0 = V_{\theta}\}$. Therefore,

$$= p \int_{-\infty}^{0} (-\tau)^{p-1} \left(\operatorname{Vol}(\theta) - \theta_{V_{\theta}}^{n} (\{ \hat{u}_{\tau} = V_{\theta} \}) \right) d\tau.$$

Since $\{\hat{u}_{\tau}\}$ is a maximal test curve, we know that $P_{\theta}[\hat{u}_{\tau}] = \hat{u}_{\tau}$. Therefore, by Lemma 2.6, $\theta_{P_{\theta}[\hat{u}_{\tau}]}^n = \mathbb{1}_{\{P_{\theta}[\hat{u}_{\tau}]=0\}}\theta^n$. Moreover, $\theta_{V_{\theta}}^n = \mathbb{1}_{\{V_{\theta}=0\}}\theta^n$. As $P_{\theta}[\hat{u}_{\tau}] = \hat{u}_{\tau}$, and $\{\hat{u}_{\tau}=0\} \subset \{V_{\theta}=0\}$, we get that $\theta_{\hat{u}_{\tau}}^n = \mathbb{1}_{\{\hat{u}_{\tau}=0\}}\theta_{V_{\theta}}^n$. Thus we can write, $\theta_{V_{\theta}}^n(\{\hat{u}_{\tau}=V_{\theta}\}) = \int_X \theta_{\hat{u}_{\tau}}^n$. So we get

$$= p \int_{-\infty}^{0} (-\tau)^{p-1} \left(\int_{X} \theta_{V_{\theta}}^{n} - \int_{X} \theta_{\hat{u}_{\tau}}^{n} \right) d\tau.$$

This proves (13).

If the test curve $\{\hat{u}_{\tau}\}_{\tau}$ has finite *p*-energy, then by (13), $\int_{X} |\dot{u}_{0}|^{p} \theta_{u_{0}}^{n} < \infty$. Since $[0, 1] \ni s \mapsto w_{s} := u_{ts}$ is the geodesic joining u_{0} and u_{t} . Thus, $\dot{w}_{0} = t\dot{u}_{0}$ and

$$\int_X |\dot{w}_0|^p \theta_{u_0}^n = t^p \int_X |\dot{u}_0|^p \theta_{u_0}^n < \infty$$

Therefore, by Lemma 6.4, $u_t \in \mathcal{E}^p(X, \theta)$. Thus the geodesic ray $\{u_t\}_t$ is a finite *p*-energy geodesic ray.

Similarly, if $\{u_t\}_t$ is a finite *p*-energy geodesic ray then $\int_X |\dot{u}_0|^p \theta_{u_0}^n = d_p^p(u_0, u_1) < \infty$, thus by (13), $\{\hat{u}_\tau\}_\tau$ is a finite *p*-energy test curve.

More generally, by Lemma 2.3, $\sup_X u_t = Ct$ for some C. Moreover, by the same argument as earlier $C = \tau_{\hat{u}}^+$. If $w_t = u_t - Ct = u_t - \tau_{\hat{u}}^+ t$, then $\{w_t\}_t$ is a geodesic whose Legendre dual $\{\hat{w}_{\tau}\}_{\tau}$ is given by $\hat{w}_{\tau} = \hat{u}_{\tau+\tau_c} = \hat{u}_{\tau+\tau_c}^+$.

From (13), we can say that

$$\begin{split} \int_X |\dot{w}_0|^p \theta_{V_\theta}^n &= p \int_{-\infty}^0 (-\tau)^{p-1} \left(\int_X \theta_{V_\theta}^n - \int_X \theta_{\dot{w}_\tau}^n \right) d\tau \\ &= p \int_{-\infty}^0 (-\tau)^{p-1} \left(\int_X \theta_{V_\theta}^n - \int_X \theta_{\dot{u}_{\tau+\tau_{\dot{u}}}^+}^n \right) d\tau \\ &= p \int_{-\infty}^{\tau_{\dot{u}}^+} (-\tau + \tau_{\dot{u}}^+)^{p-1} \left(\int_X \theta_{V_\theta}^n - \int_X \theta_{\dot{u}_\tau}^n \right) d\tau \end{split}$$

As $\dot{w}_0 = \dot{u}_0 - \tau_{\hat{u}}^+$, we get that

$$\int_X |\dot{u}_0|^p \theta_{V_\theta}^n < \infty \iff \int_X |\dot{w}_0|^p \theta_{V_\theta}^n < \infty.$$

Combining the two equations above, we get that $\{u_t\}_t \in \mathcal{R}^p_{\theta}$ is the finite *p*-energy geodesic ray iff the Legendre transform $\{\hat{u}_{\tau}\}_{\tau}$ is a maximal finite *p*-energy test curve.

References

- [Ber19] Robert J. Berman. "From Monge-Ampère equations to envelopes and geodesic rays in the zero temperature limit". In: *Math. Z.* 291.1-2 (2019), pp. 365–394. ISSN: 0025-5874,1432-1823. DOI: 10.1007/s00209-018-2087-0.
- [BDL17] Robert J. Berman, Tamás Darvas, and Chinh H. Lu. "Convexity of the extended K-energy and the large time behavior of the weak Calabi flow". In: *Geom. Topol.* 21.5 (2017), pp. 2945– 2988. ISSN: 1465-3060. DOI: 10.2140/gt.2017.21.2945.
- [Ber15] Bo Berndtsson. "A Brunn-Minkowski type inequality for Fano manifolds and some uniqueness theorems in Kähler geometry". In: *Invent. Math.* 200.1 (2015), pp. 149–200. ISSN: 0020-9910,1432-1297. DOI: 10.1007/s00222-014-0532-1.
- [BEGZ10] Sébastien Boucksom et al. "Monge-Ampère equations in big cohomology classes". In: Acta Math. 205.2 (2010), pp. 199–262. ISSN: 0001-5962. DOI: 10.1007/s11511-010-0054-7.
- [Che00] Xiuxiong Chen. "The space of Kähler metrics". In: J. Differential Geom. 56.2 (2000), pp. 189– 234. ISSN: 0022-040X. URL: http://projecteuclid.org.proxy-um.researchport.umd. edu/euclid.jdg/1090347643.
- [CC18] Xiuxiong Chen and Jingrui Cheng. On the constant scalar curvature Kähler metrics, general automorphism group. 2018. arXiv: 1801.05907 [math.DG].
- [CC21a] Xiuxiong Chen and Jingrui Cheng. "On the constant scalar curvature Kähler metrics (I)—A priori estimates". In: J. Amer. Math. Soc. 34.4 (2021), pp. 909–936. ISSN: 0894-0347. DOI: 10.1090/jams/967.
- [CC21b] Xiuxiong Chen and Jingrui Cheng. "On the constant scalar curvature Kähler metrics (II)— Existence results". In: J. Amer. Math. Soc. 34.4 (2021), pp. 937–1009. ISSN: 0894-0347. DOI: 10.1090/jams/966.

[Dar15]	Tamás Darvas. "The Mabuchi geometry of finite energy classes". In: <i>Adv. Math.</i> 285 (2015), pp. 182–219. ISSN: 0001-8708. DOI: 10.1016/j.aim.2015.08.005.
[Dar17]	Tamás Darvas. "The Mabuchi completion of the space of Kähler potentials". In: Amer. J. Math. 139.5 (2017), pp. 1275–1313. ISSN: 0002-9327. DOI: 10.1353/ajm.2017.0032.
[Dar19]	Tamás Darvas. "Geometric pluripotential theory on Kähler manifolds". In: Contemp. Math. 735 (2019), pp. 1–104. DOI: 10.1090/conm/735/14822.
[DDL18a]	Tamás Darvas, Eleonora Di Nezza, and Chinh H. Lu. "L ¹ metric geometry of big cohomology classes". In: Ann. Inst. Fourier (Grenoble) 68.7 (2018), pp. 3053–3086. ISSN: 0373-0956,1777-5310. URL: http://aif.cedram.org/item?id=AIF_201868_7_3053_0.
[DDL18b]	Tamás Darvas, Eleonora Di Nezza, and Chinh H. Lu. "On the singularity type of full mass currents in big cohomology classes". In: <i>Compos. Math.</i> 154.2 (2018), pp. 380–409. ISSN: 0010-437X. DOI: 10.1112/S0010437X1700759X.
[DDL23]	Tamás Darvas, Eleonora Di Nezza, and Chinh H. Lu. "Relative pluripotential theory on compact Kähler manifolds". In: (2023). arXiv: 2303.11584 [math.CV].
[DL20a]	Tamás Darvas and Chinh H. Lu. "Geodesic stability, the space of rays and uniform convexity in Mabuchi geometry". In: <i>Geom. Topol.</i> 24.4 (2020), pp. 1907–1967. ISSN: 1465-3060,1364-0380. DOI: 10.2140/gt.2020.24.1907.
[DLR20]	Tamás Darvas, Chinh H. Lu, and Yanir A. Rubinstein. "Quantization in geometric pluripo- tential theory". In: <i>Comm. Pure Appl. Math.</i> 73.5 (2020), pp. 1100–1138. ISSN: 0010- 3640,1097-0312. DOI: 10.1002/cpa.21857.
[DR17]	Tamás Darvas and Yanir A. Rubinstein. "Tian's properness conjectures and Finsler geometry of the space of Kähler metrics". In: J. Amer. Math. Soc. 30.2 (2017), pp. 347–387. ISSN: 0894-0347. DOI: 10.1090/jams/873.
[DX22]	Tamás Darvas and Mingchen Xia. "The closures of test configurations and algebraic sin- gularity types". In: <i>Advances in Mathematics</i> 397 (2022), p. 108198. ISSN: 0001-8708. DOI: https://doi.org/10.1016/j.aim.2022.108198.
[DXZ23]	Tamás Darvas, Mingchen Xia, and Kewei Zhang. A transcendental approach to non- Archimedean metrics of pseudoeffective classes. 2023. arXiv: 2302.02541 [math.AG].
[DZ23]	Tamás Darvas and Kewei Zhang. "Twisted Kähler-Einstein metrics in big classes". In: (2023). arXiv: 2208.08324 [math.DG].
[DL20b]	Eleonora Di Nezza and Chinh H. Lu. " L^p metric geometry of big and nef cohomology classes". In: Acta Math. Vietnam. 45.1 (2020), pp. 53–69. ISSN: 0251-4184. DOI: 10.1007/s40306-019-00343-4.
[DT21]	Eleonora Di Nezza and Stefano Trapani. "Monge-Ampère measures on contact sets". In: <i>Math. Res. Lett.</i> 28.5 (2021), pp. 1337–1352. ISSN: 1073-2780,1945-001X.
[Don99]	 S. K. Donaldson. "Symmetric spaces, Kähler geometry and Hamiltonian dynamics". In: Northern California Symplectic Geometry Seminar. Vol. 196. Amer. Math. Soc. Transl. Ser. 2. Amer. Math. Soc., Providence, RI, 1999, pp. 13–33. DOI: 10.1090/trans2/196/02.
[Gue14]	Vincent Guedj. The metric completion of the Riemannian space of Kähler metrics. 2014. arXiv: 1401.7857 [math.DG].
[GLZ19]	Vincent Guedj, Chinh H. Lu, and Ahmed Zeriahi. "Plurisubharmonic envelopes and supersolutions". In: J. Differential Geom. 113.2 (2019), pp. 273–313. ISSN: 0022-040X. DOI: 10.4310/jdg/1571882428.
[GZBook]	Vincent Guedj and Ahmed Zeriahi. <i>Degenerate complex Monge-Ampère equations</i> . Vol. 26. EMS Tracts in Mathematics. European Mathematical Society (EMS), Zürich, 2017, pp. xxiv+472. ISBN: 978-3-03719-167-5. DOI: 10.4171/167.
[Gup23]	Prakhar Gupta. "A complete metric topology on relative low energy spaces". In: <i>Math. Z.</i> 303.3 (2023), Paper No. 56, 27. ISSN: 0025-5874. DOI: 10.1007/s00209-023-03218-5.
[Gup24]	Prakhar Gupta. Complete Geodesic Metrics in Big Classes. 2024. arXiv: 2401.01688 [math.DG].
[He15]	Weiyong He. "On the Space of Kähler Potentials". In: <i>Communications on Pure and Applied Mathematics</i> 68.2 (2015), pp. 332–343. DOI: https://doi.org/10.1002/cpa.21515.
[Mab87]	Toshiki Mabuchi. "Some symplectic geometry on compact Kähler manifolds. I". In: Osaka J. Math. 24.2 (1987), pp. 227–252. ISSN: 0030-6126. URL: http://projecteuclid.org.proxy-um.researchport.umd.edu/euclid.ojm/1200780161.
[PS07]	Duong H. Phong and Jacob Sturm. "Test configurations for K-stability and geodesic rays". In: J. Symplectic Geom. 5.2 (2007), pp. 221–247. ISSN: 1527-5256,1540-2347. URL: http://projecteuclid.org/euclid.jsg/1202004456.

REFERENCES

- [RW14] Julius Ross and David Witt Nyström. "Analytic test configurations and geodesic rays". In: J. Symplectic Geom. 12.1 (2014), pp. 125–169. ISSN: 1527-5256,1540-2347. DOI: 10.4310/JSG.2014.v12.n1.a5.
- [Sem92] Stephen Semmes. "Complex Monge-Ampère and symplectic manifolds". In: Amer. J. Math. 114.3 (1992), pp. 495–550. ISSN: 0002-9327. DOI: 10.2307/2374768.
- [Tru22] Antonio Trusiani. "L¹ Metric Geometry of Potentials with Prescribed Singularities on Compact Kähler Manifolds". In: *The Journal of Geometric Analysis* 32 (2022), pp. 1–37.
- [Wit19] David Witt Nyström. "Monotonicity of non-pluripolar Monge-Ampère masses". In: *Indiana Univ. Math. J.* 68.2 (2019), pp. 579–591. ISSN: 0022-2518,1943-5258. DOI: 10.1512/iumj.2019.68.7630.
- [Xia24] Mingchen Xia. Singularities in global pluripotential theory. 2024. URL: https://mingchenxia.github.io/home/Lectures/SGPT.pdf.