# DEGENERATE TWISTED TRACES ON QUANTIZED KLEINIAN SINGULARITIES OF TYPE A

ZEV FRIEDMAN AND BEN WEBSTER

ABSTRACT. We study the space of non-degenerate traces on quantized Kleinian singularities of type A by studying their complement, the degenerate traces. In particular, we find the dimension of the space of twisted traces as a function of the corresponding automorphism and the quantization parameters, encoded in a polynomial P.

## 1. INTRODUCTION

Given a  $\mathbb{C}$ -algebra  $\mathcal{A}$  and automorphism  $\omega$ , a **twisted trace** on  $\mathcal{A}$  is a  $\mathbb{C}$ -linear map  $T: \mathcal{A} \to \mathbb{C}$ such that  $T(ab) = T(\omega(b)a)$ . The topic of twisted traces on noncommutative algebras, especially quantizations of symplectic singularities, has attracted considerable mathematical attention in recent years. The traces of interest to use have arisen in the study of conformal field theory [BPR17], but have raised mathematical questions of independent interest. As shown in [ES20], a particularly important role in the construction of short star products is played by twisted traces which are strongly nondegenerate, that is, where for all a, there is a b with deg  $b \leq \deg a$  such that  $T(ab) \neq 0$  (see Definition 1). In this paper, we study a weaker form of nondegeneracy, where we remove the condition deg  $b \leq \deg a$ which is easier to study. Throughout, we'll use **nondegenerate** to mean a trace satisfying this weaker condition.

In this paper, we consider the quantizations  $\mathcal{A}_P$  of Kleinian singularities of type A and study their space of twisted traces for a natural family of automorphisms  $g_t$  depending on a parameter  $t \in \mathbb{C} \setminus \{0\}$ . This algebra depends on a choice of monic polynomial P, whose degree  $d = \deg P$  fixes the singularity that we quantize. These algebras have already been considered in [Eti+21], but primarily with a focus on twisted traces on even quantizations and the question of positive definiteness. We will consider these questions from a more elementary, algebraic perspective, focusing on the question: What is the set of non-degenerate twisted traces of  $\mathcal{A}_P$  as a function of P?

As is often the case, we can answer this question by studying the complement of the non-degenerate traces, which are naturally the **degenerate traces**. Since the space of all twisted traces is known to be a d-1 or d-dimensional  $\mathbb{C}$ -vector space (depending on whether t = 1 or not) and degenerate traces form a subspace, the only interesting invariant determining this structure is the dimension of the space of degenerate traces.

**Theorem A.** The space of degenerate traces is a subspace whose dimension is  $0 \le \delta(P) \le d-1$  as defined in (9), with every integer in this range being realized for every t. In particular, we have:

- $\delta(P) = 0$  if and only if P has no pair of distinct roots whose difference is an integer.
- $\delta(P) = d 1$  if and only if all roots of P are distinct and belong to a single coset  $a + \mathbb{Z}$  for  $a \in \mathbb{C}$ .

In particular, the algebra  $\mathcal{A}_P$  has a non-degenerate trace if  $t \neq 1$ . On the other hand, when t = 1, a non-degenerate trace exists if and only if  $\delta(P) < d - 1$ , that is, P has a multiple root or two roots whose difference is not integral.

We prove this by studying the **formal Stieltjes transform**  $F_T$  of a trace T. This is a power series which encodes the information of a twisted trace. We show below that this power series is the Taylor expansion at  $z = \infty$  of a rational function if and only if T is degenerate (Theorem 6). Thus, we prove Theorem A by characterizing the rational functions which appear in this way (Lemma 9). This allows us to prove a general characterization of the dimension of the space of degenerate traces (Theorem 10). We'll also discuss how similar tools can be applied to study strong non-degeneracy; this is a nonlinear condition, so it is a much more complicated to study, but it can be characterized by the Padé approximants of  $F_T(x)$ . Nondegeneracy will fail on polynomials of degree  $\leq n$  if the *n*th Padé approximant to  $F_T(x)$  has denominator of lower than expected degree (Lemma 13).

There are many possible extensions of this work, for example, considering general hypertoric enveloping algebras. The space of twisted traces for these algebras is characterized by the "quantum Hikita conjecture" [KMP21] of Kamnitzer, McBreen, and Proudfoot (a theorem in the hypertoric case). These have been studied in more generality by the second author, Gaiotto, Hilburn, Redondo-Yuste, and Zhou [Gai+24]. However, no consideration has been given there to understanding which of these traces are degenerate.

In addition to the proof of Theorem A, we discuss a few other interesting details about these traces. In particular, we show how to write any degenerate trace as a sum of pullbacks of degenerate traces from algebras where P has two roots and characterize the degenerate traces that appear concretely as twisted traces of finite-dimensional representations (Lemma 15). Finally, we point out the existence of a meromorphic function satisfying the recursion (17) expected of the formal Stieltjes transform of a trace, though the transform is a power series with radius of convergence 0 and thus does not define a meromorphic function.

Acknowledgements. The authors appreciate useful conversations with Pavel Etingof, Davide Gaiotto, and Daniil Klyuev about twisted traces. B. W. is supported by an NSERC Discovery Grant. This research was supported in part by a Mathematics Undergraduate Research Award from the Faculty of Mathematics, University of Waterloo, and in part by Perimeter Institute for Theoretical Physics. Research at Perimeter Institute is supported by the Government of Canada through the Department of Innovation, Science and Economic Development Canada and by the Province of Ontario through the Ministry of Research, Innovation and Science.

### 2. Background

Given a nonconstant polynomial  $P \in \mathbb{C}[x]$ , the quantized Kleinian singularity  $\mathcal{A}_P$  is the free algebra on  $\mathbb{C}$  in three variables u, v, z, with the following relations:

$$zu = uz - u, \quad zv = vz + v, \quad uv = P(z - \frac{1}{2}), \quad vu = P(z + \frac{1}{2}).$$

We will write  $\mathcal{A} = \mathcal{A}_P$  for brevity unless the identity of P is crucial for a given result. This algebra makes a number of appearances in different aspects of the theory of symplectic singularities. It's the W-algebra associated to a subregular nilpotent of  $\mathfrak{sl}_d$ , a hypertoric enveloping algebra for a rank d-1torus acting on a d-dimensional vector space, and the spherical Cherednik algebra for  $\mathbb{Z}/d\mathbb{Z}$  acting on  $\mathbb{C}$  by the dth roots of unity.

Notice that  $\mathcal{A}$  decomposes as a sum of eigenspaces of adz, where adz maps  $a \in \mathcal{A}$  to za - az, and that the eigenvalues are exactly the integers. For each  $k \in \mathbb{Z}$ , let  $\mathcal{A}_k$  be the eigenspace of adz for the eigenvalue k. Note that  $\mathcal{A}_0 = \mathbb{C}[z]$  and that if  $a \in \mathcal{A}_j, b \in \mathcal{A}_k$  then  $ab \in \mathcal{A}_{j+k}$ .

For all  $t \in \mathbb{C} \setminus \{0\}$ , let  $g_t : \mathcal{A} \to \mathcal{A}$  be the automorphism given by  $u \mapsto t^{-1}u, v \mapsto tv$ , and  $z \mapsto z$ .

**Definition 1.** A twisted trace on  $\mathcal{A}$  with respect to  $g_t$  is a linear functional  $T : \mathcal{A} \to \mathbb{C}$  with the property  $T(ab) = T(g_t(b)a)$  for all  $a, b \in \mathcal{A}$ .

Consider the radical  $\operatorname{rad} T$  of a twisted trace T

$$rad(T) = \{ a \in \mathcal{A} \mid T(ab) = 0 \text{ for all } b \in \mathcal{A} \}$$

The trace T is degenerate if  $rad(T) \neq 0$ , that is, if there exists  $a \in A$  such that T(ab) = 0 for all  $b \in A$ . Otherwise, we have rad(T) = 0 and we call T nondegenerate.

We'll also be interested in a stronger version of non-degeneracy: we call T n-degenerate if for some p(z) of degree  $\leq n$ , we have T(p(z)q(z)) = 0 for all  $q \in \mathbb{C}[x]$  satisfying deg  $q \leq n$ . We can also think about this nondegeneracy in terms of the infinite Hankel matrix

(1) 
$$H = \begin{bmatrix} T(1) & T(z) & T(z^2) & \cdots \\ T(z) & T(z^2) & T(z^3) & \cdots \\ T(z^2) & T(z^3) & T(z^4) & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

The trace T is nondegenerate if the matrix H is nondegenerate and n-degenerate if its principal  $n \times n$  minor is degenerate.

Let  $\mathcal{C}_{P,t}$  denote the space of twisted traces on  $\mathcal{A}_P$  with respect to  $g_t$ , and let  $\mathcal{D}_{P,t}$  denote the set of degenerate twisted traces on  $\mathcal{A}_P$  with respect to  $g_t$ . One can readily verify that  $\mathcal{C}_{P,t}$  has a natural vector space structure and that  $\mathcal{D}_{P,t}$  is a vector subspace, since if T(ab) = 0 and T'(a'b) = 0 for all b, then (T + T')(aa'b) = 0 as well.

Note that the twisted trace property shows that rad T is a 2-sided ideal, since if  $a \in \operatorname{rad} T, a', b \in \mathcal{A}$ , we have

$$T((aa')b) = T(a(a'b)) = 0 T((a'a)b) = T(a(bg_t^{-1}(a'))) = 0$$

If  $P_1(x) | P_2(x) \in \mathbb{C}[x]$  and we choose  $Q_1, Q_2$  so that  $P_1(x)Q_1(x)Q_2(x) = P_2(x)$ , then there is a natural homomorphism  $\varphi_{Q_1,Q_2} \colon \mathcal{A}_{P_2} \to \mathcal{A}_{P_1}$  defined by

(2) 
$$\varphi_{Q_1,Q_2}(u) = Q_1(z - \frac{1}{2})u \qquad \varphi_{Q_1,Q_2}(v) = Q_2(z + \frac{1}{2})v \qquad \varphi_{Q_1,Q_2}(z) = z.$$

Note that if  $aP_1(x) = P_2(x)$  for some  $0 \neq a \in \mathbb{C}$ , then the map  $\varphi_{a,1}$  is an isomorphism. So for the remainder of the paper it suffices to consider  $\mathcal{A}_P$  where P is monic. This map commutes with the action of  $g_t$ , so the pullback by  $\varphi_{Q_1,Q_2}$  preserves twisted traces.

It will also be occasionally useful to note that  $\mathcal{A}_{P(x)} \cong \mathcal{A}_{P(-x)}$  via the isomorphism

$$(3) u \mapsto v v \mapsto u z \mapsto -z$$

Note that this isomorphism intertwines  $g_t$  and  $g_{t^{-1}}$ , so pullback by it swaps twisted traces for these automorphisms. Similarly,  $\mathcal{A}_{P(x)} \cong \mathcal{A}_{P(x-r)}$  via

$$(4) u \mapsto u v \mapsto v z \mapsto z + r$$

In their paper [Eti+21], Etingof et al. showed that:

**Lemma 2** ([Eti+21, Prop. 2.3]). A linear functional T on  $\mathcal{A}$  is a twisted trace with respect to  $g_t$  if and only if

(1) 
$$T(\mathcal{A}_k) = 0$$
 for all  $k \neq 0$ , and

(2)  $T(S(z-\frac{1}{2})P(z-\frac{1}{2})) = tT(S(z+\frac{1}{2})P(z+\frac{1}{2}))$  for all  $S \in \mathbb{C}[x]$ .

This means that a twisted trace is uniquely determined by its behaviour on  $\mathcal{A}_0$ .

**Definition 3.** If T is a linear functional on  $\mathbb{C}[z]$ , then the formal Stieljes transform of T is the formal power series  $F_T \in x^{-1}\mathbb{C}[[x^{-1}]]$  given by

$$F_T(x) = \sum_{n=0}^{\infty} x^{-n-1} T(z^n) = T(\frac{1}{x-z}).$$

For  $n \in \mathbb{Z}$ , let  $\mathbb{C}_n[x]$  denote the vector space of polynomials of degree at most n (in particular,  $\mathbb{C}_n[x] = \{0\}$  for n < 0).

**Lemma 4** ([Eti+21, Prop. 2.5]). The map  $T \mapsto P(x)(F_T(x+\frac{1}{2})-tF_T(x-\frac{1}{2}))$  is a vector space isomorphism  $\mathcal{C}_{P,t} \xrightarrow{\sim} \mathbb{C}_{\deg P-1}[x]$  if  $t \neq 1$ , and from  $\mathcal{C}_{P,t} \xrightarrow{\sim} \mathbb{C}_{\deg P-2}[x]$  if t = 1. In particular, if  $P(x)(F(x+\frac{1}{2})-tF(x-\frac{1}{2}))$  is a polynomial, then F(x) must be the formal Stieljes transform of some twisted trace on  $\mathcal{A}$ . Note that this implies that  $F_T(x + \frac{1}{2}) - tF_T(x - \frac{1}{2})$  is a rational function which has a pole at a of order at most the vanishing order  $p_a$  of P at x = a. Thus, we have the principal parts expansion

(5) 
$$F_T(x+\frac{1}{2}) - tF_T(x-\frac{1}{2}) = \sum_{a \in \mathbb{C}} \frac{D_a^{(1)}}{x-a} + \dots + \frac{D_a^{(p_a)}}{(x-a)^{p_a}}.$$

# 3. Degenerate traces

**Lemma 5.** A twisted trace T is degenerate on  $\mathcal{A}$  if and only if it is degenerate on  $\mathbb{C}[z] \subset \mathcal{A}$ .

Proof. ( $\Rightarrow$ ) Suppose that T is degenerate on  $\mathcal{A}$ . Then there exists some nonzero  $a \in A$  such that T(ab) = 0 for all  $b \in \mathcal{A}$ . Since  $A = \bigoplus_{k \in \mathbb{Z}} \mathcal{A}_k$ , we may write  $a = \sum_{k=-K}^{K} a_k$  for  $a_k \in \mathcal{A}_k$ , and  $K \in \mathbb{N}$ . Then since  $a \neq 0$ , there exists  $j \in \mathbb{Z}$  such that  $a_j \neq 0$ . For all  $b \in A_{-j}$ , we have  $T(ab) = \sum_{k=-K}^{K} T(a_k b) = T(a_j b)$  since  $a_k b \in \mathcal{A}_{k-j}$ . Choose some nonzero  $c \in A_{-j}$  and note that  $0 \neq a_k c \in \mathcal{A}_0 = \mathbb{C}[z]$ . Then for all  $R(z) \in \mathbb{C}[z]$ , we have that  $cR(z) \in \mathcal{A}_{-j}$  and so  $T(a_k cR(z)) = T(acR(z)) = 0$ , so T is degenerate on  $\mathbb{C}[z]$ .

( $\Leftarrow$ ) Suppose T is degenerate on  $\mathbb{C}[z]$ . Then there exists some nonzero  $S(z) \in \mathbb{C}[z]$  such that T(S(z)R(z)) = 0 for all  $R \in \mathbb{C}[x]$ . Then for all  $b \in \mathcal{A}_0$  we have T(S(z)b) = 0, and for all  $b \in \mathcal{A}_k$  with  $k \neq 0$ , we have  $S(z)b \in \mathcal{A}_k$  and so T(S(z)b) = 0. Since  $\mathcal{A}$  is the direct sum of these spaces, we have T(S(z)b) = 0 for all  $b \in \mathcal{A}$ .

Note that  $I_T = \operatorname{rad} T \cap \mathbb{C}[z]$  is an ideal, so it is generated by a unique monic polynomial S(x), which is characterized as the unique monic polynomial of minimal degree in  $I_T$ .

**Theorem 6.** We have an equality

$$I_T = \{ S(z) \mid F_T(z)S(z) \in \mathbb{C}[z] \}$$

In particular, a twisted trace T on  $\mathcal{A}$  is degenerate if and only if  $F_T(x)$  is a rational function of x. In this case, if we write  $F_T(x) = \frac{R(x)}{S(x)}$  with S monic and gcd(R, S) = 1, then S generates  $I_T$ .

*Proof.* Suppose that  $S(z) \in I_T$ . Then we have

$$S(x)F_T(x) = S(x)F_T(x) - \sum_{n=0}^{\infty} x^{-n-1}T(S(z)z^n) = T(\frac{S(x) - S(z)}{x - z})$$

Let  $R(x) = T(\frac{S(x) - S(z)}{x - z})$ , and note that R(x) is a polynomial. Then  $S(x)F_T(x) = R(x)$  as desired. Now suppose that  $R, S \in \mathbb{C}[x]$  satisfy  $S(x)F_T(x) = R(x)$ . Then

$$R(x) = S(x)F_T(x) = T(\frac{S(x)}{x-z}) = T(\frac{S(x)-S(z)}{x-z}) + T(\frac{S(z)}{x-z})$$

Since every term in R(x) and  $T(\frac{S(x)-S(z)}{x-z})$  has a nonnegative exponent on x and every term in  $T(\frac{S(z)}{x-z})$  has a negative exponent on x, in order for the two sides to be equal we must have

$$0 = T(\frac{S(z)}{x-z}) = \sum_{n=0}^{\infty} x^{-n-1} T(S(z)z^n),$$

and so T(S(z)b) = 0 for all  $b \in \mathbb{C}[z]$  since  $\{z^n\}_{n=0}^{\infty}$  is a basis for  $\mathbb{C}[z]$ . Thus, we have  $S(z) \in I_T$ . This completes the proof.

Note that since the Laurent expansion of  $F_T(x)$  at  $x = \infty$  (that is, its expansion as a power series in  $\mathbb{C}[[x^{-1}]]$ ) contains no nonnegative powers of x, we must have deg  $R < \deg S$ . Thus, we can expand R/S in its principal parts expansion—if we let  $m_a$  be the order of vanishing of S at x = a for  $a \in \mathbb{C}$ , then we have

$$\frac{R(x)}{S(x)} = \sum_{a \in \mathbb{C}} \left( \frac{C_a^{(1)}}{x-a} + \dots + \frac{C_a^{(m_a)}}{(x-a)^{m_a}} \right) \qquad C_a^{(k)} = \operatorname{res}_{x=a} \left( (x-a)^{k-1} \frac{R(x)}{S(x)} \right)$$

Note that this principal part decomposition is equivalent to the Vandermonde decomposition of a Hankel matrix H defined in (1). For simplicity, assume  $F_T(x) = \prod_a(x)$  for a single  $a \in \mathbb{C}$ . In this case,  $T(z^k) = \sum_{r=1}^{\infty} C_a^{(r)} {r-1 \choose k} a^{k-r+1}$  by the usual Taylor expansion

$$\frac{1}{(x-a)^r} = \frac{1}{x^r} + a\binom{r-1}{r} \frac{1}{x^{r+1}} + a^2\binom{r-1}{r+1} \frac{1}{x^{r+2}} + \cdots$$

at  $x = \infty$ . Let r be the order of the pole of  $\Pi_a(x)$ . By [BLV97, Th. 2], we can factor the Hankel matrix as a product  $H = (V^{(a)})^T D^{(a)} V^{(a)}$  where V is an  $r \times \infty$  matrix with  $V_{ij}^{(a)} = (\binom{j-1}{i-1}a^{j-i})$  and D is the matrix  $D_{ij}^{(a)} = (C_a^{(i+j-1)})$ . One can readily calculate that the i, j entry of  $(V^{(a)})^T D^{(a)} V^{(a)}$  is

$$\sum_{k,\ell \ge 1} a^{i+j-k-\ell} C_a^{(k+\ell-1)} \binom{k-1}{i-1} \binom{\ell-1}{j-1} = \sum_{s \ge 1} a^{i+j-s-1} C^{(s)} \binom{s-1}{i+j-2} = T(z^{i+j-2})$$

using the reindexing  $s = k + \ell - 1$  and the usual relation on binomial coefficients. The general case is found by stacking the matrices  $V^{(a)}$  for different  $a \in \mathbb{C}$  and taking the matrices  $D^{(a)}$  as diagonal blocks, as in [BLV97, Th. 2].

By (5), the coefficients  $C_a^{(k)}$  satisfy the relations

(6) 
$$C_{a-1/2}^{(k)} - tC_{a+1/2}^{(k)} = D_a^{(k)}.$$

If  $r \in \mathbb{C}$  and  $P \in \mathbb{C}[x]$ , if r is a root of P of order n, we will say that r is an n-fold root of P. For each complex number a, we let  $n_a^+$  be the largest integer such that there is a root of order  $\geq n_a^+$  of P in the set  $a + \frac{1}{2} + \mathbb{Z}_{\geq 0}$  and similarly with  $\geq n_a^-$  and the set  $a - \frac{1}{2} - \mathbb{Z}_{\geq 0}$ . Let  $n_a = \min(n_a^+, n_a^-)$ .

**Lemma 7.** For a fixed polynomial P and any degenerate trace T, we have  $m_a \leq n_a$  for all  $a \in \mathbb{C}$ .

*Proof.* We need only prove that  $n_a^+ \ge m_a$ . Applying the same argument after the isomorphism (3) will imply that  $n_a^- \ge m_a$  as well. Restrict consideration to the set  $a + \mathbb{Z}$ . Since P and S have finitely many zeros, we have  $m_{a'} = n_{a'}^+ = 0$  for all but finitely many  $a' \in a + \mathbb{Z}_{\ge 0}$ . Thus, we can apply induction with base case being some a such that  $m_{a'} = n_{a'}^+ = 0$  for all  $a' \ge a$ . For the inductive step, assume that  $m_{a+1} \le n_{a+1}^+$ .

Recall that  $P(x)(F_T(x+\frac{1}{2})-tF_T(x-\frac{1}{2})) \in \mathbb{C}[x]$ , and observe that

$$F_T(x+\frac{1}{2}) - tF_T(x-\frac{1}{2}) = \frac{R(x+\frac{1}{2})}{S(x+\frac{1}{2})} - t\frac{R(x-\frac{1}{2})}{S(x-\frac{1}{2})}$$

Consider this rational function at  $x = a + \frac{1}{2}$ . It can only have a pole if  $P(a + \frac{1}{2}) = 0$ , and the order p of its pole is at most the vanishing order of P at this point. Let us compare this with the vanishing orders  $m_{a+1}$  and  $m_a$ ; since R and S are coprime, these are also the order of the pole of R/S at these points.

If  $p < m_a$ , then by the non-archimedean triangle inequality for vanishing orders, we must have that  $m_{a+1} = m_a$ . By induction, we know that  $n_a^+ \ge n_{a+1}^+ \ge m_{a+1} = m_a$ .

On the other hand, if  $p \ge m_a$ , then  $b = a + \frac{1}{2}$  is the desired root of order  $\ge m_a$ . This completes the proof.

Thus, we can write the formal Stieltjes transform of any degenerate twisted trace as:

$$\frac{R(x)}{S(x)} = \sum_{a \in \mathbb{C}} \Pi_a(x) \qquad \Pi_a(x) = \left(\frac{C_a^{(1)}}{x-a} + \dots + \frac{C_a^{(n_a)}}{(x-a)^{n_a}}\right)$$

Furthermore, since  $P(x)(F_T(x+\frac{1}{2})-tF_T(x-\frac{1}{2})) \in \mathbb{C}[x]$ , it has vanishing principal parts at all  $a \in \mathbb{C}$ .

**Lemma 8.** The rational function  $\frac{R(x)}{S(x)}$  is the formal Stieltjes transform of a degenerate twisted trace if and only if for all  $a \in \mathbb{C}$  which are not roots of P

(7) 
$$C_{a-1/2}^{(\ell)} = t C_{a+1/2}^{(\ell)}$$

for all  $\ell \leq n_{a\pm 1/2}$ , and

$$C_{a-1/2}^{(\ell)} = C_{a+1/2}^{(\ell)} = 0$$

for all  $\ell > n_{a\pm 1/2}$ .

We can also write this condition in terms of the coefficients  $D_a^{(\ell)}$  defined by (5). Applying (6), we have

(8) 
$$C_a^{(k)} = (C_a^{(k)} - tC_{a+1}^{(k)}) + t(C_{a+1}^{(k)} - tC_{a+2}^{(k)}) + \dots = \sum_{j=0}^{\infty} t^j D_{a+j+1/2}^{(k)}$$

Since  $C_a^{(k)} = 0$  for all but finitely many a, we find that:

**Lemma 9.** The twisted trace with the formal Stieltjes transform  $F_T(x)$  is degenerate if and only if

$$\Delta_a^{(k)} = \sum_{j=-\infty}^{\infty} t^j D_{a+j+1/2}^{(k)} = 0$$

for all  $a \in \mathbb{C}$  and k.

Note that  $\Delta_a^{(k)} = t \Delta_{a-1}^{(k)}$ , so the vanishing of this quantity is really a condition on the coset of a modulo  $\mathbb{Z}$ . This description makes it clear how large the space of degenerate traces is. For each coset  $[a] \in \mathbb{C}/\mathbb{Z}$ , let  $\delta_{[a]}(P)$  be the number of roots in [a], counted with multiplicity, minus the maximum multiplicity of a root in this set. Let  $\delta(P)$  be the sum of these statistics over  $\mathbb{C}/\mathbb{Z}$ . That is:

(9) 
$$\delta(P) = \sum_{[a] \in \mathbb{C}/\mathbb{Z}} \delta_{[a]}(P) = \sum_{a' \in [a]} p_{a'} - \max_{a' \in [a]} p_a$$

**Theorem 10.** dim  $\mathcal{D}_{P,t} = \delta(P)$ .

Proof. For simplicity, we can assume that all the roots of P lie in  $\mathbb{Z} + \frac{1}{2}$  and prove that dim  $\mathcal{D}_{P,t} = \delta_{[1/2]}(P)$ . Let  $\pi = \max_{a' \in \mathbb{Z} + 1/2} p_{a'}$ . The space of all possible choices of  $D_a^{(\ell)}$  for  $\ell = 1, \ldots, p_a$  and all  $a \in \mathbb{Z} + \frac{1}{2}$  is exactly  $\sum_{a \in \mathbb{Z} + 1/2} p_a$ , and the statistics  $\Delta_0^{(\ell)} = \sum_{j=-\infty}^{\infty} t^j D_{j+1/2}^{(\ell)}$  for  $\ell = 1, \ldots, \pi$  are linearly independent functions. Thus, the dimension of their common kernel, which is the space of degenerate traces, is exactly  $\delta_{[1/2]}(P) = \deg P - \pi$ . This completes the proof.

This shows that the dimension of  $\mathcal{D}_{P,t}$  can range anywhere from 0 (if no distinct roots of P differ by an element of  $\mathbb{Z}$ ) to deg P-1 (if all roots are simple and lie in the same coset of  $\mathbb{Z}$ ).

More explicitly, we see that we have coordinates on  $\mathcal{D}_{P,t}$  given by the functions  $D_a^{(k)}$  where k ranges over integers such that P vanishes to order  $\geq k$  at a and at a + j for some  $j \in \mathbb{Z}_{>0}$ .

Now we discuss how we can compare the traces for different values of P. We have introduced an algebra homomorphism  $\varphi_{Q_1,Q_2} : \mathcal{A}_{PQ_1Q_2} \to \mathcal{A}_P$  and considered pullback of twisted traces under it. Since this map is the identity on  $\mathbb{C}[z]$ , it preserves the formal Stieltjes transform. Since we can easily tell if a trace is 0 or degenerate from its formal Stieltjes transform, this shows:

**Theorem 11.** If  $P_1(x) | P_2(x) \in \mathbb{C}[x]$ , then the pullback by the homomorphism  $\varphi_{Q_1,Q_2}$  introduced in (2) induces an injective map  $\mathcal{C}_{P_1,t} \hookrightarrow \mathcal{C}_{P_2,t}$ . This homomorphism preserves the formal Stieltjes transform. The subspace  $\mathcal{D}_{P_1,t}$  is the preimage of  $\mathcal{D}_{P_2,t}$ .

*Proof.* Since the map  $\varphi = \varphi_{Q_1,Q_2}$  acts by the identity on  $\mathbb{C}[z]$ , it preserves formal Stieltjes transforms:  $F_T = F_{T \circ \varphi}$ . Thus, by Lemma 4, pullback sends non-zero traces to non-zero traces, and thus is injective. Similarly, by Theorem 6, the pullback of a trace is degenerate if and only if the original trace is.  $\Box$ 

Note that if  $P_1 \neq P_2$  then this map is never an isomorphism on the space of all traces, but it may be on the space of degenerate traces, which we can check by seeing that the dimensions on the same. This happens when  $\delta(P_1) = \delta(P_2)$ . This condition can be stated in a number of ways, but one way to say it is that if a is a zero of different orders p < p' for  $P_1$  and  $P_2$ , then all other zeros in  $a + \mathbb{Z}$  must have the same vanishing order  $\leq p$  for  $P_1$  and  $P_2$ .

Employing different decompositions of the rational function  $F_T(x+\frac{1}{2})-tF_t(x-\frac{1}{2})$ , we obtain different decompositions of our twisted trace, which are often interestingly compatible with this pullback.

For example, a rational function with a single pole of order s at a will give a twisted trace for  $P(x) = (x - a)^s$ , and as long as  $t \neq 1$ , we can write any twisted trace canonically as the sum of pullbacks of these traces, corresponding to the principal part decomposition of  $F_T(x + \frac{1}{2}) - tF_t(x - \frac{1}{2})$ . However, this process is badly incompatible with degeneracy, since when P has a single root, all traces will be non-degenerate.

More natural for us is to decompose according to the order of poles. That is, we let:

$$D^{(k)}(x) = \sum_{a \in \mathbb{C}} \frac{D_a^{(k)}}{(x-a)^k}.$$

Of course,  $F_T(x + \frac{1}{2}) - tF_T(x - \frac{1}{2}) = \sum_{k=1}^{\infty} D^{(k)}(x).$ 

**Lemma 12.** We have a unique decomposition  $T = \sum_{k=1}^{\infty} T^{(k)}$  into twisted traces pulled back from  $\mathcal{A}_{P^{(k)}}$ where  $P^{(k)} = \prod (x-a)^{\min(p_a,k)}$ , with the property that every term in the partial fraction expansion of  $F_{T^{(k)}}$  has only poles of order k. The trace T is degenerate if and only if  $T^{(k)}$  is degenerate for all k.

Proof. We can characterize these traces uniquely by  $F_{T^{(k)}}(x+\frac{1}{2}) - tF_{T^{(k)}}(x-\frac{1}{2}) = D^{(k)}(x)$ . If  $t \neq 1$ , any rational function which only has poles at the right points corresponds to a twisted trace. If t = 1, we need the additional condition  $\sum_{a \in \mathbb{C}} D_a^{(1)} = 0$ . This holds for  $D^{(1)}$  if and only if it holds for  $F_T(x+\frac{1}{2}) - tF_T(x-\frac{1}{2})$ , and is automatic for  $D^{(k)}$  for k > 1.

The condition  $\Delta_a^{(k)} = 0$  holds for  $D^{(k)}(x)$  if and only if it does for  $F_T(x + \frac{1}{2}) - tF_T(x - \frac{1}{2})$ , so Lemma 9 shows that T is degenerate if and only if  $T^{(k)}$  is degenerate for all k.

Notice that we can also uniquely characterize the twisted traces in the previous lemma by simply taking the partial fraction expansion of  $F_T$  and extracting the terms whose poles are of order k to be  $F_{T^{(k)}}$ , and this implies that  $T^{(k)}$  is a degenerate twisted trace on  $P^{(k)}$ .

There are various possibilities for decomposing further: for example, for  $a \in \mathbb{C}$  and  $j \in \mathbb{Z}_{\geq 0}$ , if a, a + j are roots of order  $\geq k$  of P with no roots of order  $\geq k$  in  $a + 1, \ldots, a + j - 1$ , the sum

$$\frac{C_{a+1/2}^{(k)}}{\left(x-a-1/2\right)^{k}} + \dots + \frac{C_{a-j-1/2}^{(k)}}{\left(x-a-j+1/2\right)^{k}}$$

is the Stieltjes transform of a degenerate twisted trace for  $P = (x - a)^k (x - a - j)^k$ , and  $T^{(k)}$  can be written uniquely as the sum of these traces.

Alternatively, if a and a + j are as above, then we have a degenerate trace T' with  $F_{T'}(x + \frac{1}{2}) - tF_{T'}(x - \frac{1}{2}) = \sum_{i=1}^{k} D_a^{(k)} \left(\frac{1}{(x-a)^i} - \frac{t^{-j}}{(x-a-j)^i}\right)$  such that T'' = T - T' is a degenerate twisted trace such that  $F_{T''}(x + \frac{1}{2}) - tF_{T''}(x - \frac{1}{2})$  has trivial principal part at x = a. Applying this inductively, we can write any degenerate trace as a sum of traces pulled back from  $\mathcal{A}_{P'}$  where P' has two roots with integer difference.

# 4. Strong non-degeneracy

We now consider how similar ideas can be applied to study *n*-degeneracy. A useful fact to note is that any formal series  $F_T(x) \in x^{-1}\mathbb{C}[\![x^{-1}]\!]$  has a unique [n-1/n]-Padé approximant at  $x = \infty$ 

$$\frac{R_{n-1}(x)}{S_n(x)} \approx F_T(x)$$

for  $S_n$  a monic polynomial of degree  $\leq n$  and deg  $R_{n-1} < \deg S_n$ . This rational function is uniquely characterized by the equivalent conditions

$$F_T(x) - \frac{R_{n-1}(x)}{S_n(x)} \in x^{-2n-1} \mathbb{C}[x^{-1}] \qquad S_n(x) F_T(x) - R_{n-1}(x) \in x^{-n-1} \mathbb{C}[x^{-1}]$$

These conditions show that  $R_{n-1}(x)$  is determined by  $S_n(x)F_T(x)$ , so the polynomial  $S_n(x)$  is determined by the condition that the  $x^{-1-k}$  coefficients of  $S_n(x)F_T(x)$  vanish for  $k = 0, \ldots, n-1$ . These coefficients can also be interpreted as the traces  $T(S_n(z)z^k)$ . That is,  $S_n(x)$  is the unique polynomial of minimal degree such that  $S_n(x)$  is perpendicular to all polynomials of degree < n. There must be such a polynomial of degree n for dimension reasons, but it might be that in some cases, there is one of lower degree, in which case T is (n-1)-degenerate. That is:

**Lemma 13.** The trace is n-degenerate if and only if deg  $S_{n+1} \leq n$ , that is, if  $S_{n+1} = S_n$  or equivalently  $R_n/S_{n+1} = R_{n-1}/S_n.$ 

For any value of n, it is easy to construct examples of non-degenerate traces which are nonetheless *n*-degenerate: For any fixed  $\frac{R}{S}$  with deg  $R < \deg S < n$ , we can consider the twisted trace with  $F_T(x) = \frac{R(x+\frac{1}{2})}{S(x+\frac{1}{2})} - t\frac{R(x-\frac{1}{2})}{S(x-\frac{1}{2})} + \frac{1}{(x-a)^k} \text{ for some } k \gg n, \text{ which has } S_n = S_{n+1} = \dots = S_{k-2} = S, \text{ but}$ this trace is non-degenerate.

It seems to be a very complicated problem to compute exactly which traces are *n*-degenerate for a fixed n. We hope that some progress can be made from the observation that:

(10) 
$$\frac{R_{n-1}(x+\frac{1}{2})}{S_n(x+\frac{1}{2})} - t\frac{R_{n-1}(x-\frac{1}{2})}{S_n(x-\frac{1}{2})} \in \frac{Q(x)}{P(x)} + x^{-2n-1}\mathbb{C}[\![x^{-1}]\!] \qquad t \neq 1$$

(11) 
$$\frac{R_{n-1}(x+\frac{1}{2})}{S_n(x+\frac{1}{2})} - t\frac{R_{n-1}(x-\frac{1}{2})}{S_n(x-\frac{1}{2})} \in \frac{Q(x)}{P(x)} + x^{-2n-2}\mathbb{C}[x^{-1}] \qquad t = 1$$

where  $Q(x) = P(x)(F_T(x + \frac{1}{2}) - tF_T(x - \frac{1}{2})) \in \mathbb{C}[x].$ 

**Theorem 14.** The polynomials  $R_{n-1}(x)$  and  $S_n(x)$  are the unique polynomials such that deg  $R_{n-1} <$  $\deg S_n \leq n, \gcd(R_{n-1}, S_n) = 1, S_n$  is monic and (10) or (11) holds, depending on whether  $t \neq 1$  or t = 1.

*Proof.* If all these conditions hold for some polynomials r, s, then  $\overline{F}(x) = F_T(x) - r(x)/s(x)$  is a power series that satisfies

$$\bar{F}(x+\frac{1}{2}) - t\bar{F}(x-\frac{1}{2}) = \frac{Q(x)}{P(x)} - \frac{r(x+\frac{1}{2})}{s(x+\frac{1}{2})} - t\frac{r(x-\frac{1}{2})}{s(x-\frac{1}{2})} \in x^{-2n-1}\mathbb{C}[\![x^{-1}]\!].$$

If t = 1, we can further conclude that  $\overline{F}(x + \frac{1}{2}) - t\overline{F}(x - \frac{1}{2}) \in x^{-2n-2}\mathbb{C}[x^{-1}]$ . Let k + 1 for  $k \ge 0$  be the order of vanishing of  $\overline{F}(x) = a_k x^{-k-1}$  at  $x = \infty$ . Taylor expanding  $(x \pm \frac{1}{2})^{-k} = x^{-k} \pm \frac{k+1}{2} x^{-k-1} + \cdots$  at  $x = \infty$ , we find that

$$\bar{F}(x+\frac{1}{2}) - t\bar{F}(x-\frac{1}{2}) = (t-1)a_k x^{-k-1} + ((1-t)a_{k+1} - (k+1)a_k)x^{-k-2}$$

vanishes to order k+1 and order k+2 if t=1, so we must have  $k \ge 2n$ . This proves that r(x)/s(x) = $R_{n-1}(x)/S_n(x)$  is the expected Padé approximant, and the conditions on these polynomials guarantee that  $r(x) = R_{n-1}(x)$  and  $s(x) = S_n(x)$ . 

It is a very interesting question how we might be able to find the principal parts expansion of  $R_{n-1}(x)/S_n(x)$  from that of  $\frac{Q(x)}{P(x)}$ . At the moment, we see no practical way of doing this.

#### DEGENERATE TWISTED TRACES

# 5. FINITE-DIMENSIONAL MODULES

One interesting source of degenerate twisted traces is finite-dimensional modules over  $\mathcal{A}$ . Assume M is such a finite-dimensional module and  $\alpha \colon M \to M$  is a map satisfying

(12) 
$$a \cdot \alpha(m) = \alpha(g_t(a) \cdot m)$$
 for all  $a \in \mathcal{A}, m \in M$ .

Such an  $\alpha$  is called a **twisting map**. We call the map  $\mathcal{A} \to \mathbb{C}$  defined by  $a \mapsto \operatorname{tr}(M, \alpha \circ a)$  the **twisted trace of the module** M and map  $\alpha$ . This is manifestly a twisted trace in the sense of Definition 1 since  $\operatorname{tr}(M, \alpha \circ ab) = \operatorname{tr}(M, b \circ \alpha \circ a) = \operatorname{tr}(M, \alpha \circ g_t(b)a)$ .

**Lemma 15.** A degenerate twisted trace is a twisted trace of a finite-dimensional module if and only  $C_a^{(k)} = 0$  for all a and all k > 1, or equivalently  $D_a^{(k)} = 0$  for all a and all k > 1.

In particular, every degenerate trace is of this form if and only if P has no multiple roots.

*Proof.* The finite-dimensional representations of this algebra are well-known: if a, a + j are roots of P with no roots in  $a + 1, \ldots, a + j - 1$ , then there is a unique irreducible representation  $S_{a,a+j}$  where the element z has simple spectrum  $\{a + \frac{1}{2}, \ldots, a + j - \frac{1}{2}\}$ , and all irreducible representations are of this form.

Given a finite-dimensional module M over  $\mathcal{A}$  and a twisting map  $\alpha: M \to M$ , since  $\alpha$  and z commute, they must have a simultaneous eigenvector x with eigenvalues b and  $\lambda$ . Note that ux, vx are also eigenvectors with eigenvalues  $b \mp 1$  and  $t^{\pm 1}\lambda$ . Since z has finite spectrum, we must have  $v^m x = u^m x = 0$  for some m. If for some q, we have  $uv^q x \neq 0$ , we replace x by  $v^q x$  for the largest such q. If no such q exists, we replace x by  $u^s x$  where s is maximal amongst integers such that  $u^s x \neq 0$ . Thus, we assume that ux = 0, and the assumptions above guarantee that x generates a copy of  $S_{b-1/2,b+j+1/2}$  for some j. Furthermore, this submodule has a basis given by  $\{x, vx, \ldots, v^{j-1}x\}$ , so it is invariant under  $\alpha$ . Thus

$$\operatorname{tr}(M, \alpha \circ a) = \operatorname{tr}(M/\mathcal{A}x, \alpha \circ a) + \operatorname{tr}(\mathcal{A}x, \alpha \circ a)$$

and we can reduce to assuming M is irreducible, and  $\alpha$  and z have a basis of simultaneous eigenvectors. We can explicitly calculate that

$$F_T(x) = \sum_{n=0}^{\infty} x^{-n-1} \sum_{s=0}^{j-1} \lambda t^{-s} (b+s)^n = \sum_{s=0}^{j-1} \frac{\lambda t^{-s}}{x-a}.$$

In the notation introduced earlier, this is the twisted trace corresponding to  $C_b^{(1)} = tC_{b+1}^{(1)} = \cdots = t^j C_{b+j}^{(1)} = \lambda$  and all other coefficients 0. This shows that a twisted trace of a finite-dimensional module must have this form and, in particular, has  $C_a^{(k)} = 0$  for k > 1.

On the other hand, we have shown that a degenerate twisted trace with  $C_a^{(k)} = 0$  whenever k > 1 is a sum of the traces above, and thus it can be written as a twisted trace on a sum of simples with an appropriate choice of  $\alpha$ .

It is possible to deal with the higher order poles by loosening the restriction (12). Consider the case where

(13) 
$$F_T(x) = C \sum_{j=0}^{n-1} \frac{t^j}{(x-a-j)^k}$$
$$F_T(x+\frac{1}{2}) - tF_T(x-\frac{1}{2}) = \frac{C}{(x-a+\frac{1}{2})^k} - \frac{Ct^n}{(x-a-n+\frac{1}{2})^k},$$

so  $a - \frac{1}{2}$  and  $a + n - \frac{1}{2}$  are both k-fold roots of P. We can define a finite-dimensional representation M with a basis  $e_j^{(p)}$  for  $j = 0, \ldots, n-1$  and  $p = 1, \ldots, k$  (in all equations below, whenever an index is outside these ranges,  $e_j^{(p)} = 0$  by convention), with the action:

$$z \cdot e_j^{(p)} = (a+j)e_j^{(p)} + e_j^{(p+1)} \qquad u \cdot e_j^{(p)} = P(z-\frac{1}{2})e_{j-1}^{(p)} \qquad v \cdot e_j^{(p)} = e_{j+1}^{(p)}$$

This is the same as the representation generated by the element  $e_0^{(1)}$  subject to the relations

$$(z-a)^k \cdot e_0^{(1)} = 0$$
  $u \cdot e_0^{(1)} = 0$   $v^n \cdot e_0^{(1)} = 0.$ 

We can define a map  $\alpha: V \to V$  sending  $e_j^{(k)} \mapsto Ct^j e_j^{(k-p+1)}$ ; note that  $z \cdot \alpha(m) \neq \alpha(z \cdot m)$ , so this does not satisfy (12). Despite this, one can calculate that  $T(a) = \operatorname{tr}(M, \alpha \circ a)$  is a twisted trace with  $F_T(z)$  given by (13). Summing over several representations, we can get any degenerate trace as  $T(a) = \operatorname{tr}(M, \alpha \circ a)$  for some  $(M, \alpha)$ . It's not clear to us in general how to test if a map  $\alpha$  gives a twisted trace.

# 6. Lerch transcendents

A natural question raised by our calculations with Stieltjes transforms is whether we can interpret  $F_T(x)$  as an analytic function for any non-degenerate traces. In a certain sense, the answer is obviously not:

**Lemma 16.** If  $F_T(x)$  is not the Taylor expansion of a rational function at  $x = \infty$ , then the radius convergence (in terms of  $x^{-1}$ ) of this power series is 0.

Proof. Assume not. Then for some real number  $\epsilon$ , we have that  $F_T(x)$  converges absolutely to a holomorphic function on  $\{x \in \mathbb{C} \mid |x| > \epsilon\}$ , and in fact on its closure in the Riemann sphere. We can use (5) to meromorphically extend this function to the entirety of the Riemann sphere. The argument of Lemma 8 still applies and shows that this function can only have poles at points of the form  $a + \frac{1}{2} + \mathbb{Z}$  for a a root of P which lie inside  $\{x \in \mathbb{C} \mid |x| \le \epsilon\}$ , with the order of these poles bounded by the vanishing order of p at x = a. There are only finitely many such points, which shows that  $F_T(x)$  is a meromorphic function on the Riemann sphere with finitely many poles, all of finite order. This implies that  $F_T(x)$  is a rational function. Thus, the twisted trace T is degenerate, contradicting our assumption.

Interestingly, if we focus solely on the recursion

$$F(x + \frac{1}{2}) - tF(x - \frac{1}{2}) = \frac{Q(x)}{P(x)}$$

for deg  $Q(x) < \deg P(x)$ , there is a solution to this recursion which is meromorphic in the complex plane, though not at  $x = \infty$ . It's unclear whether we can learn anything interesting from this function, but we note it here, since it is an intriguing coincidence.

Consider the Lerch transcendent

$$\Phi(t,n,x) = \sum_{j=0}^{\infty} \frac{t^j}{(x+j)^n}.$$

While frequently in the literature, this is considered as a meromorphic function of n, we will only consider the case where n is a positive integer; this sum absolutely converges and thus defines a meromorphic function on the open domain  $\mathbb{C} \setminus \mathbb{Z}_{\leq 0}$  if  $|t| \leq 1$  and  $n \geq 1$  and one of these inequalities is strict. If  $t \notin [1, \infty)$ , then the integral formula

(14) 
$$\Phi(t,n,x) = \frac{1}{n!} \int_0^\infty \frac{z^{n-1}e^{-xz}}{1 - te^{-z}} dz$$

defines this function on the half-plane  $\operatorname{Re}(x) > 0$ . This function meromorphically extends to the rest of the complex plane with poles at  $x \in \mathbb{Z}_{\leq 0}$  via the equality  $\Phi(t, n, x) - t\Phi(t, n, x+1) = \frac{1}{x^n}$ . Consider the function  $\Psi_a^{(n)}(x) = \Phi(t, n, a - x + \frac{1}{2})$ . This satisfies

(15) 
$$\Psi_a^{(n)}(x+\frac{1}{2}) - t\Psi_a^{(n)}(x-\frac{1}{2}) = \frac{1}{(a-x)^n}$$

Thus, we find that:

**Theorem 17.** If  $|t| \leq 1$  and  $t \neq 1$  and (5) gives the principal part decomposition of  $\frac{Q(z)}{P(z)}$  then:

(16) 
$$\tilde{F}(x) = \sum_{a \in \mathbb{C}} \sum_{\ell=1}^{p_a} (-1)^{\ell} D_a^{(\ell)} \Psi_a^{(\ell)}(x)$$

satisfies the recursion

(17) 
$$\tilde{F}(x+\frac{1}{2}) - t\tilde{F}(x-\frac{1}{2}) = \frac{Q(z)}{P(z)} = \sum_{k=1}^{\infty} D^{(k)}(x)$$

If t = 1, then the situation is more subtle. One can write the formal sum as in the RHS of (16), but if  $D_a^{(1)} \neq 0$  for any  $a \in \mathbb{C}$ , then this sum is not absolutely convergent. Thus, constructing a limit for it is considerably more subtle. However, in this case, we can use the formulas (14–16), but move the sum inside the integral and obtain the formula:

(18) 
$$\tilde{F}(x) = \int_0^\infty \sum_{a \in \mathbb{C}} \sum_{\ell=1}^{p_a} D_a^{(\ell)} (-1)^\ell \frac{z^{\ell-1} e^{(x-a-1/2)z}}{1-e^{-z}} dz$$

If we have  $\sum_{a \in \mathbb{C}} D_a^{(1)} = 0$ , then the numerator and denominator both have simple zeros at z = 0, so the ratio is holomorphic and decays exponentially if  $\operatorname{Re}(x) \ll 0$ . Thus, the improper integral in (18) is well-defined and holomorphic as a function of x on some half-plane, as well as satisfying (17) on this half-plane. This shows that we can use (17) to meromorphically extend this function to the whole of the complex plane.

# References

- [BLV97] DL Boley, FT Luk, and David Vandevoorde. "A general Vandermonde factorization of a Hankel matrix". Int'l Lin. Alg. Soc. (ILAS) Symp. on Fast Algorithms for Control, Signals and Image Processing. accessed from https://www-users.cselabs.umn.edu/~boley/publications/papers/ 1997.
- [BPR17] Christopher Beem, Wolfger Peelaers, and Leonardo Rastelli. "Deformation quantization and superconformal symmetry in three dimensions". Communications in Mathematical Physics 354.1 (2017), pp. 345–392. arXiv: 1601.05378 [hep-th].
- [ES20] Pavel Etingof and Douglas Stryker. "Short Star-Products for Filtered Quantizations, I". SIGMA. Symmetry, Integrability and Geometry: Methods and Applications 16 (2020), p. 014.
- [Eti+21] Pavel Etingof, Daniil Klyuev, Eric Rains, and Douglas Stryker. "Twisted Traces and Positive Forms on Quantized Kleinian Singularities of Type A". SIGMA. Symmetry, Integrability and Geometry: Methods and Applications 17 (2021), p. 029.
- [Gai+24] Davide Gaiotto, Justin Hilburn, Jaime Redondo-Yuste, Ben Webster, and Zheng Zhou. Twisted traces on abelian quantum Higgs and Coulomb branches. 2024. arXiv: 2308.15198 [hep-th, physics:n URL: http://arxiv.org/abs/2308.15198. preprint.
- [KMP21] Joel Kamnitzer, Michael McBreen, and Nicholas Proudfoot. "The quantum Hikita conjecture". Advances in Mathematics 390 (2021), Paper No. 107947, 53.

Email address: zfriedma@uwaterloo.ca

DEPARTMENT OF PURE MATHEMATICS, UNIVERSITY OF WATERLOO, WATERLOO, ON, CANADA

# Email address: ben.webster@uwaterloo.ca

DEPARTMENT OF PURE MATHEMATICS, UNIVERSITY OF WATERLOO & PERIMETER INSTITUTE FOR THEORETICAL PHYSICS, WATERLOO, CANADA, ON