

Quantum fluctuation on the worldsheet of probe string in BTZ black hole

Yu-Ting Zhou*

*College of Physics, Nanjing University of Aeronautics
and Astronautics, Nanjing 210016, China and
Key Laboratory of Aerospace Information Materials
and Physics (NUAA), MIIT, Nanjing 211106, China*

Xiao-Mei Kuang†

*Center for Gravitation and Cosmology,
College of Physical Science and Technology,
Yangzhou University, Yangzhou 225009, China.*

Abstract

In this paper, we investigate the second-order normal quantum fluctuation on the worldsheet of a probe string in the Bañados-Teitelboim-Zanelli (BTZ) black hole. These fluctuations is treated as the projection of Hawking radiation on the worldsheet and indeed modify the action growth of the string. Then in the string field theory/boundary conformal field theory framework, via the boundary vertex operator we study the correlation function of the Schrödinger functional of excited fields on the worldsheet and further extract the field's formula. Our study could shed light on the potential connection between complexity growth and correlation function.

*Electronic address: yu-tingzhou@nuaa.edu.cn

†Electronic address: xmeikuang@yzu.edu.cn

Contents

I. Introduction	2
II. Action growth of probe string in BTZ black hole	4
III. Quantum fluctuation on the worldsheet of probe string	5
IV. Further study from the perspective of String Field Theory	8
V. Conclusion	11
Acknowledgments	12
References	13

I. INTRODUCTION

Black holes have become a natural platform for us to understand spacetime and gravity at a semi-classical level after Stephen Hawking proposed the exciting theory of black hole radiation. Then, drawing inspiration from black hole thermodynamics, physicists proposed the holographic nature of gravity [1, 2], which then gives rise to the gauge/gravity duality [3-5]. The extensive applications of this duality provide novel perspectives for exploring gravity and strongly coupled system.

Recently, quantum information theory has begun to transcend its traditional framework, providing more and more insights in the field of quantum gravity such as computational complexity [6-12]. The complexity essentially measures the difficulty of changing a quantum state into another state, however, it is still not clear to define the initial states and target states when one applies complexity into quantum field theory (QFT). Though considerable attempts have been made in this field [13-26], a widely acceptable understanding of the complexity still poses an unresolved query. Thanks to holography, two elegant descriptions of complexity have been proposed from gravity side. One is the “complexity=volume (CV)” conjecture, where V represents the volume of the Einstein-Rosen (ER) bridge linking the two sides of the AdS black hole’s boundary. The other is the “complexity=action (CA)” conjecture, where A denotes the classical action of a space-time region enclosed by the bulk

Cauchy slice anchored at the boundaries, also known as the “Wheeler-Dewitt (WdW)” patch [27–29].

Based on the CA conjecture, there are many studies on the stationary systems, see for examples [30–49] and references therein. Similar efforts are evident in dynamic systems, such as the investigation of complexity growth with probe branes [50] and the exploration of non-local operator effects in the BTZ black hole [51–59]. Remarkable progress has been made in holographic complexity, but a well-defined reference state in holography is also open. Therefore, the study of quantum fluctuation or perturbation of complexity could circumvent the state puzzle and help to step forward to investigate the complexity.

Thus, the aim of this paper is to study the second-order normal fluctuation on a probe string whose two points end in the dual boundary of the BTZ black hole. As a first attempt, our goal is to build a possible connection between the complexity growth and correlation function. The purpose we consider the normal fluctuations on the probe string mainly consists of two aspects. On one hand, the probe string in BTZ black hole can be treated as a 2-dimensional quantum field theory in the curved space-time background, similar as the handling with holographic Brownian motion proposed in [60, 61]. Then in this scenario, various fluctuation modes could be excited on the string’s worldsheet due to Hawking radiation in the black hole environment [62]. Here for convenience, we shall fix the end points of the string onto the boundary and focus on the normal direction fluctuation of the string. This operation allows us to obtain the normal operator of those excited normal modes along the string, as we will show soon. On the other hand, this consideration allows us to work with the string field theory/boundary conformal field theory (SFT/BCFT) correspondence, stating that every classical field in worldsheet of the probe string can be described by a BCFT of an open string attached to the boundary [63–68]. In this framework, we can treat the excited fields as a single object and construct the corresponding vertex operator on the boundary. This then further helps us to calculate the correlation function of the bulk fields in the Schrödinger functional representation and extract the curved-dependent worldsheet excited field.

The remaining of this paper is organized as follows. In Sec. II, we briefly review the BTZ black hole and the action growth of probe string model. In Sec. III, we consider second order normal fluctuation on the Nambu-Goto action of the probe string in the black hole and then we get the normal fluctuation operator. In Sec. IV, we explore the two-point

correlation function of these excited modes and extract the field function on the worldsheet. We summarize our work in the last section.

II. ACTION GROWTH OF PROBE STRING IN BTZ BLACK HOLE

We start with the 3-dimensional BTZ black hole with a negative cosmological constant of which the metric is [69, 70]

$$ds^2 = -\left(-M + \frac{r^2}{l^2}\right)dt^2 + \frac{1}{-M + \frac{r^2}{l^2}}dr^2 + r^2d\phi^2, \quad (1)$$

where $-\infty \leq t \leq \infty$ and $0 \leq \phi \leq 2\pi$, M is the black hole mass and l is AdS radius. Then for convenience, we shall set $l = 1$ and rewrite metric (1) under Poincare coordinate as,

$$ds^2 = \frac{1}{z^2} \left[-f(z)dt^2 + \frac{1}{f(z)}dz^2 + d\phi^2 \right], \quad f(z) = 1 - Mz^2, \quad (2)$$

where we have $z = \frac{1}{r}$, the horizon location $z_h = \frac{1}{\sqrt{M}}$ and the Hawking temperature $T = \frac{1}{2\pi\sqrt{M}}$.

We proceed to consider a probe string in this background spacetime and its two endpoints are attached in the boundary subspace. The configuration is shown in FIG. 1 where we have omitted the time direction. Subsequently, we can work in the worldsheet coordinates $\tau = t$




FIG. 1: The geometric description of a probe string ending on the boundary of BTZ black hole background.

and $\sigma = \phi$, and further perform $X^\mu(t, \phi)$ as

$$X^\mu(t, \phi) = \begin{pmatrix} t \\ z(\phi) \\ \phi \end{pmatrix}. \quad (3)$$

Then the Nambu-Goto action of this string is

$$S_{NG} = T_s \int dt d\phi \sqrt{-\det [h_{ab}]}, \quad (4)$$

where T_s is the tension of string. h_{ab} is the induced metric of worldsheet, which can be written in the form

$$h_{ab} = g_{\mu\nu} \partial_a X^\mu \partial_b X^\nu, \quad (5)$$

with $g_{\mu\nu}$ the metric defined in (2). Then directly from (4), we can define the action growth of the probe string as

$$\frac{1}{T_s} \frac{dS_{NG}}{dt} = \int_{-\frac{\phi_c}{2}}^{\frac{\phi_c}{2}} d\phi \sqrt{-\det [h_{ab}]}. \quad (6)$$

Further calculations on the above action growth has been addressed in [52–54], in which the authors discussed the significant effects of the graviton mass and black hole mass. In the following studies, we shall concern the quadratic fluctuation on this action growth.

III. QUANTUM FLUCTUATION ON THE WORLDSHEET OF PROBE STRING

We shall then adopt the normal coordinate gauge, under which the fluctuation is normal to the probe string everywhere, to expand the action growth into the quadratic order. The reason exists in that the normal coordinate gauge could be the safest choice comparing to other coordinate gauges, as addressed in [71]. The explicit geometric description of this fluctuation is shown in FIG.2, in which the dashed curve describes a fluctuated string on the original probe string.




FIG. 2: The geometric description of the fluctuation on the string.

To proceed, we expand the action growth (6) up to the quadratic term as

$$\frac{1}{T_s} \frac{dS_{NG}}{dt} \simeq \frac{1}{T_s} \frac{dS_{NG}^{(0)}}{dt} + \frac{1}{T_s} \frac{dS_{NG}^{(2)}}{dt} + \dots, \quad (7)$$

where the first term in the right side denotes the initial classical action growth. It is noticed that in general we can have infinite higher order terms but in our following study we consider till quadratic term. Then we shall assume that the fluctuation field is a scalar field ξ_n and account for the contributions from the quantum fluctuations. To this end, we first expand the string coordinates around (3) as

$$X^\mu(t, \phi) = X_0^\mu(t, \phi) + n_\mu \xi_n(t, \phi). \quad (8)$$

Here we introduce the unit normal vector $n_\mu = (n_z, n_\phi)$ and consider the normal fluctuation on the coordinate. Subsequently, (8) can be rewritten as

$$z = z + n_z \xi_n(t, \phi), \quad (9)$$

$$\phi = \phi + n_\phi \xi_n(t, \phi). \quad (10)$$

Noted that there is no fluctuation on the time direction, which means that we keep the time coordinate fixed. The normal vector should satisfy the condition [71]

$$n_\phi \lambda_\phi g_{\phi\phi} + n_z \lambda_z g_{zz} = 0, \quad (11)$$

$$n_\phi^2 g_{\phi\phi} + n_z^2 g_{zz} = 1, \quad (12)$$

where λ_ϕ and λ_z are the parameters satisfying the tangent vector equation of string

$$z' = \frac{\partial z}{\partial \phi} = \frac{\lambda_z}{\lambda_\phi}. \quad (13)$$

Now we have to fix the normal vector. The Hamiltonian of the string can be extracted from (4) as

$$H = \frac{Mz^2 - 1}{z^2 \sqrt{1 - Mz^2 + z'^2}}, \quad (14)$$

which does not explicitly depend on ϕ , so it is conserved. Then z' could be written as

$$z' = \frac{z_*^2 \sqrt{Mz^2 - 1} \sqrt{Mz^2 - 1 + \frac{z^4(1 - Mz_*^2)}{z_*^4}}}{z^2 \sqrt{1 - Mz^2}}, \quad (15)$$

where z_* is the turning point at the ϕ_0 (see FIG.1). Then combining (11)-(13) and (15), we can solve out n_z and n_ϕ as

$$n_z = \frac{z^3 \sqrt{1 - Mz_*^2}}{z_*^2},$$

$$n_\phi = \frac{\sqrt{z^6 - Mz^6 z_*^2 - z^2 z_*^4 + Mz^4 z_*^4}}{\sqrt{Mz^2 z_*^4 - z_*^4}}. \quad (16)$$

Under the fluctuated string coordinates

$$\begin{aligned}
\tilde{X}^t &= t, \\
\tilde{X}^z &= z + n_z \xi_n(t, \phi), \\
\tilde{X}^\phi &= \phi + n_\phi \xi_n(t, \phi),
\end{aligned} \tag{17}$$

the induced metric (5) can be expanded as

$$\begin{aligned}
\tilde{h}_{tt} &= \partial_t \tilde{X}^t \partial_t \tilde{X}^t g_{tt} + \partial_t \tilde{X}^z \partial_t \tilde{X}^z g_{zz} + \partial_t \tilde{X}^\phi \partial_t \tilde{X}^\phi g_{\phi\phi}, \\
&= g_{tt} + (n_z \partial_t \xi_n)^2 g_{zz} + (n_\phi \partial_t \xi_n)^2 g_{\phi\phi} \\
\tilde{h}_{\phi\phi} &= \partial_\phi \tilde{X}^z \partial_\phi \tilde{X}^z g_{zz} + \partial_\phi \tilde{X}^\phi \partial_\phi \tilde{X}^\phi g_{\phi\phi} \\
&= [z' + \partial_\phi(n_z \xi_n)]^2 g_{zz} + [1 + \partial_\phi(n_\phi \xi_n)]^2 g_{\phi\phi} \\
&= (z')^2 g_{zz} + 2z' \xi_n \partial_\phi n_z \cdot g_{zz} + 2z' n_z \partial_\phi \xi_n \cdot g_{zz} + \xi_n^2 (\partial_\phi n_z)^2 g_{zz} \\
&\quad + 2\xi_n n_z \partial_\phi n_z \cdot \partial_\phi \xi_n \cdot g_{zz} + n_z^2 (\partial_\phi \xi_n)^2 g_{zz} + g_{\phi\phi} + 2\xi_n \partial_\phi n_\phi \cdot g_{\phi\phi} \\
&\quad + 2n_\phi \partial_\phi \xi_n \cdot g_{\phi\phi} + \xi_n^2 (\partial_\phi n_\phi)^2 g_{\phi\phi} + 2\xi_n n_\phi \partial_\phi n_\phi \cdot \partial_\phi \xi_n \cdot g_{\phi\phi} + n_\phi^2 (\partial_\phi \xi_n)^2 g_{\phi\phi}.
\end{aligned} \tag{18}$$

After straightforward calculation, we can expand the $\det[\tilde{h}_{ab}]$ up to the second order and obtain the quadratic term

$$\frac{1}{T_s} \frac{dS_{NG}^{(2)}}{dt} = \int d\phi \xi_n^\dagger [(z'^2 g_{zz} + g_{\phi\phi}) \partial_t^2 + \partial_\phi (g_{tt} \partial_\phi) - [(\partial_\phi n_z)^2 g_{zz} g_{tt} + (\partial_\phi n_\phi^2) g_{\phi\phi} g_{tt}]] \xi_n, \tag{19}$$

where we have used the relation of the normal vector in (13). By defining a normal fluctuation operator as

$$O_n = (z'^2 g_{zz} + g_{\phi\phi}) \partial_t^2 + \partial_\phi (g_{tt} \partial_\phi) - [(\partial_\phi n_z)^2 g_{zz} g_{tt} + (\partial_\phi n_\phi)^2 g_{\phi\phi} g_{tt}], \tag{20}$$

we can rewrite the quadratic terms (19) as

$$\frac{1}{T_s} \frac{dS_{NG}^{(2)}}{dt} = \sum_n^N \int d\phi \xi_n^\dagger O_n \xi_n, \tag{21}$$

which was proposed in [71].

In principle, we can solve out ξ_n from (21) by the equation of motion

$$O_n \xi_n = 0, \tag{22}$$

however, it is a very difficult mission for us to go ahead. The difficulties come from two aspects: the first is that the equation $O_n \xi_n$ is highly nonlinear, so analytical solution is a

big challenge and numerical method is called for. The second is that ξ_n can be any field, which further reinforces the complex of this issue. Therefore, we shall shift our strategy and adopt alternative approach to find ξ_n , which will be elaborated in the next section.

IV. FURTHER STUDY FROM THE PERSPECTIVE OF STRING FIELD THEORY

In this section, we shall endow the quantum fluctuation on the probe string with a potential physical process and discuss what we can construct. The probe string living in BTZ black hole environment should be inevitably affected by the Hawking radiation [60–62], besides, combining (8) and (21) further inspires us to interpret ξ_n as the fluctuation fields caused by the Hawking radiation. It is noted that random fluctuations on the worldsheet caused by Hawking radiation can also lead to a random motion of the endpoint of the string on the boundary, which is dubbed holographic Brownian motion [60]. But here our case is different because we fix the two endpoints of the string on the boundary and only consider the fluctuation on the worldsheet. Then one interesting question we shall ask is which kind of physical quantity in the fluctuations can correspond to complexity/action growth caused by this process. We shall do some analysis and expect to give insight in this issue.

As we aforementioned that the quantum fluctuations could be excited by any kind of field so that the sum in (21) should collect all possible fields. Fortunately, inspired by the strategy in string field theory (SFT) [63, 64, 72], we can treat this set of fluctuations as a single object, called excited string fields (ESF) $\Xi[\phi_i, A_\mu, \dots]$, where ϕ_i and A_μ are the scalar fields and $U(1)$ gauge field, while the ellipsis denote other possible excited fields. Moreover, in our setup the worldline of the string is swept out by the endpoints attached to the boundary, so all of these components collectively form boundary conformal field theory. Thus, we shall apply SFT/BCFT correspondence [63–68] to proceed.

We define a worldsheet’s excited space , \mathcal{H}_{BCFT} , which can be viewed as a tensor product of “matter ” and “ghost” sectors [73]

$$\mathcal{H}_{BCFT} = \mathcal{H}_{BCFT}^m \otimes \mathcal{H}_{BCFT}^{gh}, \quad (23)$$

where m and gh denote matter and ghost sectors, respectively. The ghost sector is a bc system that is characterized by anticommuting, holomorphic, and anti-holomorphic worldsheet

fields b, \bar{b}, c, \bar{c} . Since the ghost sector comes from the gauge fixing process and its physics is usually not well understood, so here we do not consider the ghost sector for the fluctuations. For a state $|\xi_n\rangle \in \mathcal{H}_{BCFT}$, there exists a corresponding boundary operator V_{ξ_n} , called vertex operator, so that we have

$$|\xi_n\rangle = V_{\xi_n} |0\rangle. \quad (24)$$

Here $|0\rangle$ is the $SL(2, \mathbb{R})$ vacuum in the boundary without vertex operator. Then we extend the vertex operator at the point ϕ_0 as $\mathcal{V}(\phi_0) =: e^{i\beta_n X^\mu(\phi_0)}$: of which the integral form is

$$V_{\xi_n}(\phi_0) = \int dt d\phi \sqrt{\gamma} \mathcal{V}(\phi_0). \quad (25)$$

Here, $::$ means the normal ordering, and β_n is the charge of ξ_n which plays the role of space-time momentum along the X^μ . γ is the induced metric in the boundary. For a physical state $|\xi_n\rangle$, it must be a BRST invariant state in BCFT at ghost number 1, satisfying [72, 73]

$$Q|\xi_n\rangle = 0, \quad |\xi_n\rangle \sim |\xi_n\rangle + Q|\Lambda\rangle, \quad |\xi_n\rangle = \text{ghost number } 1, \quad (26)$$

where Q is the BRST operator and $|\Lambda\rangle$ is ghost number 0 state. Recalling the equation of motion (22), we find that it is natural to treat the normal fluctuation operator O_n as the BRST operator Q .

Next, we use the *Schrödinger* representation for the excited worldsheet fields and consider Dirichlet boundary conditions of the probe string on both endpoints

$$X^\mu(\phi = 0) = x^\mu, \quad X^\mu(\phi = \pi) = x^\mu. \quad (27)$$

Subsequently, we can expand $X^\mu(t, \phi)$ in (3) as [74]

$$X^\mu(\phi) = x^\mu(\phi) + \frac{(x^\mu_{\frac{\phi_c}{2}} - x^\mu_{-\frac{\phi_c}{2}})}{\pi} \phi + \sqrt{2\alpha'} \sum_{n \neq 0} \frac{1}{n} \alpha_n^\mu \sin(n\phi), \quad (28)$$

or more explicitly in the form

$$\begin{aligned} z(\phi) &= z(\phi) + \sqrt{2\alpha'} \sum_{n \neq 0} \frac{1}{n} \alpha_n^\mu \sin[n\phi], \\ \phi &= -\frac{\phi_c}{2} + \frac{\phi_c}{\pi} \phi + \sqrt{2\alpha'} \sum_{n \neq 0} \frac{1}{n} \alpha_n^\mu \sin[n\phi], \end{aligned} \quad (29)$$

and the vertex operator (24) could be

$$\begin{aligned} V_{\xi_n(z)}(\phi_0) &= e^{i\beta_n z^*} \times \exp\left[i\sqrt{2\alpha'} \beta_n \sum_{n \neq 0} \frac{\alpha_n^\mu}{n} \sin[n\phi_0]\right], \\ V_{\xi_n(\phi)}(\phi_0) &= e^{i\beta_n (\frac{\phi_0}{\pi} - \frac{1}{2}) \phi_c} \times \exp\left[i\sqrt{2\alpha'} \beta_n \sum_{n \neq 0} \frac{\alpha_n^\mu}{n} \sin[n\phi_0]\right]. \end{aligned} \quad (30)$$

Then we define the overlap

$$\Xi_n[X^\mu(\phi)] = \langle X^\mu(\phi) | \Xi_n \rangle, \quad (31)$$

where Ξ_n denotes the collection of excited fields on the worldsheet, depending on the curve X^μ in the background. It is noticed that Ξ_n could also include the $b(\phi)$ and $c(\phi)$ fields, but here we do not consider this ghost sector. For the chosen point ϕ_0 in the boundary, we calculate the correlation of the vertex operator in the worldsheet as

$$\langle \Xi_n, \Xi'_n \rangle = \langle (I \circ V_{\xi_n}(\phi_0)) V_{\xi'_n}(\phi_0) \rangle_{worldsheet}. \quad (32)$$

where I denotes the Belavin-Polyakov-Zamolodchikov (BPZ) conjugation, $I(a) = -\frac{1}{a}$. That is to say, the $I \circ V_{\xi_n}(\phi_0)$ corresponds to another point in the boundary. Thus, (32) contains two vertex operators, which are at ϕ_0 and $-\frac{1}{\phi_0}$, respectively, and the formula of the path integral over the worldsheet is

$$\langle \Xi_n, \Xi'_n \rangle = \int [dX^\mu] (I \circ V_{\xi_n}(\phi_0)) V_{\xi'_n}(\phi_0) e^{-(S_{NG}^{(0)} + S_{NG}^{(2)})}. \quad (33)$$

Then we factorize this integration into three parts, saying outside ($z > |z(\phi)|$), on ($z = |z(\phi)|$), and below ($z < |z(\phi)|$) the probe string,

$$\begin{aligned} \langle \Xi_n, \Xi'_n \rangle &= \int_{-\frac{\phi_c}{2}}^{\frac{\phi_c}{2}} [dX^\mu]_{z=|z(\phi)|} \int_{-\frac{\phi_c}{2}}^{\frac{\phi_c}{2}} [dX^\mu]_{z>|z(\phi)|} \\ &\times \int_{-\frac{\phi_c}{2}}^{\frac{\phi_c}{2}} [dX^\mu]_{z<|z(\phi)|} (I \circ V_{\xi_n}(\phi_0)) V_{\xi'_n}(\phi_0) e^{-(S_{NG}^{(0)} + S_{NG}^{(2)})}. \end{aligned} \quad (34)$$

by using the BPZ inverse, which takes the form

$$\begin{aligned} \langle \Xi_n, \Xi'_n \rangle &= \int_{-\frac{\phi_c}{2}}^{\frac{\phi_c}{2}} [dX^\mu]_{z=|z(\phi)|} \left(\int_{-\frac{\phi_c}{2}}^{\frac{\phi_c}{2}} [dX^\mu]_{z<|z(\phi)|} V_{\xi_n}(\phi_0) e^{-S_{NG}^{(0)} - S_{NG}^{(2)}} \right) \\ &\times \left(\int_{-\frac{\phi_c}{2}}^{\frac{\phi_c}{2}} [dX^\mu]_{z<|z(\phi)|} V_{\xi'_n}(\phi_0) e^{-S_{NG}^{(0)} - S_{NG}^{(2)}} \right). \end{aligned} \quad (35)$$

From (35), we can extract the induced string field on the worldsheet or the projection of excited fields of Hawking radiation on the worldsheet as

$$\begin{aligned} \Xi_n[X^\mu(\phi)] &= \int [dX^\mu] V_{\xi_n}(\phi_0) e^{-S_{NG}^{(0)} - S_{NG}^{(2)}} \\ &= \int d(\delta X^\mu) V_{\xi_n}(\phi_0) e^{-S_{NG}^{(0)} - S_{NG}^{(2)}} \\ &= \int n_\mu d\xi_n V_{\xi_n}(\phi_0) e^{-S_{NG}^{(0)} - S_{NG}^{(2)}}. \end{aligned} \quad (36)$$

In the second equality, we change the integration variable in term of the facts that $dX^\mu = dX^0 + d(\delta X)$ and dX^0 is fixed. In the third equality, we recall (8) and take n_μ as a constant vector along the worldsheet. Then considering that $\int d\xi e^{-S_{NG}^{(2)}}$ is a Gaussian integral, we shall further reduce the above formula into

$$\Xi_n[X^\mu] = n_\mu \int dt d\phi \sqrt{\frac{\pi}{T_s O_n}} \cdot V_{\xi_n}(\phi_0) e^{-S_{NG}^0}, \quad (37)$$

which could be defined as the induced fields excited by Hawking radiation under the Schrödinger functional representation. As expected, n_μ appears here indicates that these fields are polarized and normal to the worldsheet. One interesting property we can read off from (37) is when the string tension T_s approaches to zero, the fields will blow up. This could be understood as that the string is melted by the high Hawking temperature when it goes near the horizon.

Before closing this section, we shall present some discussions on our results. Firstly, we argue that $\Xi_n[X^\mu]$ can be defined as the induced field excited by Hawking radiation based on the fact that the fluctuation of worldsheet can be caused by Hawking radiation as the probe string is in black hole environment, even though the physical quantities related to Hawking radiation are not reflected in (37). It is noticed that besides Hawking radiation, the fluctuations may also be triggered by other mechanisms, such as a scalar field on the worldsheet as a defect [75], but here we take more care of the worldsheet fluctuation itself, which means that fields $\Xi_n[X^\mu]$ are only curve-dependent. Secondly, in our analysis, we see that under the second-order fluctuation on the action/complexity growth, the two-point correlation emerge from the worldsheet's perturbation. Moreover, (35) shows that the second order fluctuation of action growth of probe string indeed contributes to the correlation of vertex operators in the boundary conformal field theory in the form of functional path integrals. We argue that this may indicate a profound connection between complexity and the correlation function, which deserves further clarification.

V. CONCLUSION

There are plenty of schemes to define complexity in quantum field theory and conformal field theory. Relating with gravity, one also has two holographic versions of complexity. So further studies on complexity in a suitable framework will help us to further understand

complexity in physics. In this work, we handle with complexity in string field theory and expect to shed light on connecting complexity with correlation function in quantum field theory and holography.

We investigated the second-order quantum fluctuation of a probe string ending on the boundary of the BTZ black hole background. Since the string is in black hole environment, we assumed that the unique normal fluctuation on the string is introduced by the Hawking radiation. Considering that the fluctuation can also affect the complexity or action growth of the string and will bring in a high-order correction on the Nambu-Goto action. So we calculate the second-order correction on the Nambu-Goto action and obtain the normal fluctuation operator.

Moreover, in the SFT/BCFT framework, by treating the excited fields on the worldsheet as a single object, and further defining them as excited string field, we found that the normal fluctuation operator also can be viewed as a BRST operator. Then we calculated the correlation function of vertex operators which is given in the form of functional path integrals. This corresponds to the correlation function of the excited fields on the worldsheet. We then extracted the induced field excited by Hawking radiation under the Schrödinger functional representation. Our study shows that under the second-order fluctuation on the action/complexity growth, the two-point correlation can emerge from the worldsheet's perturbation. This indicates that somehow we may explain the complexity as scattering amplitude, which could be an interesting direct, for example, one should clarify the definition of the minimal path in the scattering amplitude, and so on. We hope to further study the related topics in the future.

Acknowledgments

We appreciate Profs. Rui-Hong Yue, Ya-Peng Hu and Jian-Pin Wu for helpful and intriguing discussions. Moreover, We are grateful to Dr. Guo-Yang Fu and Dr. Kang Zhou for their constructive advice. This work is supported by National Natural Science Foundation of China (NSFC) under grant Nos. 12247170, 12175105, Top-notch Academic

- [1] L. Susskind, “The World as a hologram,” [J. Math. Phys.](#) **36** (1995) 6377–6396, [arXiv:hep-th/9409089](#).
- [2] G. 't Hooft, “The Holographic principle: Opening lecture,” [Subnucl. Ser.](#) **37** (2001) 72–100, [arXiv:hep-th/0003004](#).
- [3] J. M. Maldacena, “The Large N limit of superconformal field theories and supergravity,” [Adv. Theor. Math. Phys.](#) **2** (1998) 231–252, [arXiv:hep-th/9711200](#).
- [4] S. S. Gubser, I. R. Klebanov, and A. M. Polyakov, “Gauge theory correlators from noncritical string theory,” [Phys. Lett. B](#) **428** (1998) 105–114, [arXiv:hep-th/9802109](#).
- [5] E. Witten, “Anti-de Sitter space and holography,” [Adv. Theor. Math. Phys.](#) **2** (1998) 253–291, [arXiv:hep-th/9802150](#).
- [6] T. J. Osborne, “Hamiltonian complexity,” [Reports on Progress in Physics](#) **75** no. 2, (Jan, 2012) 022001.
- [7] S. Gharibian, Y. Huang, Z. Landau, and S. W. Shin, “Quantum hamiltonian complexity,” [Foundations and Trends in Theoretical Computer Science](#) **10** no. 3, (2015) 159–282.
- [8] G. Dvali, C. Gomez, D. Lst, Y. Omar, and B. Richter, “Universality of black hole quantum computing,” [Fortschritte der Physik](#) **65** no. 1, (2017) 1600111.
- [9] B. Swingle, G. Bentsen, M. Schleier-Smith, and P. Hayden, “Measuring the scrambling of quantum information,” [Phys. Rev. A](#) **94** (Oct, 2016) 040302.
- [10] K. Hashimoto, N. Iizuka, and S. Sugishita, “Time evolution of complexity in abelian gauge theories,” [Phys. Rev. D](#) **96** (Dec, 2017) 126001.
- [11] J. Watrous, “Quantum computational complexity,” [arXiv:0804.3401 \[quant-ph\]](#).
- [12] N. Bao and J. Liu, “Quantum complexity and the virial theorem,” [JHEP](#) **08** (2018) 144, [arXiv:1804.03242 \[hep-th\]](#).
- [13] V. Vanchurin, “Dual Field Theories of Quantum Computation,” [JHEP](#) **06** (2016) 001, [arXiv:1603.07982 \[hep-th\]](#).
- [14] S. Chapman, M. P. Heller, H. Marrochio, and F. Pastawski, “Toward a Definition of Complexity for Quantum Field Theory States,” [Phys. Rev. Lett.](#) **120** no. 12, (2018) 121602, [arXiv:1707.08582 \[hep-th\]](#).

- [15] J. Molina-Vilaplana and A. Del Campo, “Complexity Functionals and Complexity Growth Limits in Continuous MERA Circuits,” [JHEP](#) **08** (2018) 012, [arXiv:1803.02356 \[hep-th\]](#).
- [16] A. Bhattacharyya, P. Caputa, S. R. Das, N. Kundu, M. Miyaji, and T. Takayanagi, “Path-Integral Complexity for Perturbed CFTs,” [JHEP](#) **07** (2018) 086, [arXiv:1804.01999 \[hep-th\]](#).
- [17] M. A. Nielsen, “A geometric approach to quantum circuit lower bounds,” [Quant. Inf. Comput.](#) **6** no. 3, (2006) 213–262, [arXiv:quant-ph/0502070](#).
- [18] M. A. Nielsen, M. R. Dowling, M. Gu, and A. C. Doherty, “Quantum computation as geometry,” [Science](#) **311** no. 5764, (2006) 1133–1135.
- [19] R. Jefferson and R. C. Myers, “Circuit complexity in quantum field theory,” [JHEP](#) **10** (2017) 107, [arXiv:1707.08570 \[hep-th\]](#).
- [20] R.-Q. Yang, Y.-S. An, C. Niu, C.-Y. Zhang, and K.-Y. Kim, “Principles and symmetries of complexity in quantum field theory,” [Eur. Phys. J. C](#) **79** no. 2, (2019) 109, [arXiv:1803.01797 \[hep-th\]](#).
- [21] A. Bhattacharyya, A. Shekar, and A. Sinha, “Circuit complexity in interacting QFTs and RG flows,” [JHEP](#) **10** (2018) 140, [arXiv:1808.03105 \[hep-th\]](#).
- [22] A. Bhattacharyya, P. Nandy, and A. Sinha, “Renormalized Circuit Complexity,” [Phys. Rev. Lett.](#) **124** no. 10, (2020) 101602, [arXiv:1907.08223 \[hep-th\]](#).
- [23] H. A. Camargo, P. Caputa, and P. Nandy, “Q-curvature and path integral complexity,” [JHEP](#) **04** (2022) 081, [arXiv:2201.00562 \[hep-th\]](#). [Erratum: [JHEP](#) 10, 038 (2023)].
- [24] K. Adhikari, S. Choudhury, and A. Roy, “Krylov Complexity in Quantum Field Theory,” [Nucl. Phys. B](#) **993** (2023) 116263, [arXiv:2204.02250 \[hep-th\]](#).
- [25] K. Adhikari, S. Choudhury, S. Chowdhury, K. Shirish, and A. Swain, “Circuit complexity as a novel probe of quantum entanglement: A study with black hole gas in arbitrary dimensions,” [Phys. Rev. D](#) **104** no. 6, (2021) 065002, [arXiv:2104.13940 \[hep-th\]](#).
- [26] K. Adhikari and S. Choudhury, “Cosmological Krylov Complexity,” [Fortsch. Phys.](#) **70** no. 12, (2022) 2200126, [arXiv:2203.14330 \[hep-th\]](#).
- [27] S. Chapman, H. Marrochio, and R. C. Myers, “Complexity of Formation in Holography,” [JHEP](#) **01** (2017) 062, [arXiv:1610.08063 \[hep-th\]](#).
- [28] A. R. Brown, D. A. Roberts, L. Susskind, B. Swingle, and Y. Zhao, “Holographic Complexity Equals Bulk Action?,” [Phys. Rev. Lett.](#) **116** no. 19, (2016) 191301,

- [arXiv:1509.07876 \[hep-th\]](#).
- [29] A. R. Brown, D. A. Roberts, L. Susskind, B. Swingle, and Y. Zhao, “Complexity, action, and black holes,” [Phys. Rev. D](#) **93** no. 8, (2016) 086006, [arXiv:1512.04993 \[hep-th\]](#).
- [30] W.-J. Pan and Y.-C. Huang, “Holographic complexity and action growth in massive gravities,” [Phys. Rev. D](#) **95** no. 12, (2017) 126013, [arXiv:1612.03627 \[hep-th\]](#).
- [31] W.-D. Guo, S.-W. Wei, Y.-Y. Li, and Y.-X. Liu, “Complexity growth rates for AdS black holes in massive gravity and $f(R)$ gravity,” [Eur. Phys. J. C](#) **77** no. 12, (2017) 904, [arXiv:1703.10468 \[gr-qc\]](#).
- [32] D. Momeni, S. A. Hosseini Mansoori, and R. Myrzakulov, “Holographic Complexity in Gauge/String Superconductors,” [Phys. Lett. B](#) **756** (2016) 354–357, [arXiv:1601.03011 \[hep-th\]](#).
- [33] J. Tao, P. Wang, and H. Yang, “Testing holographic conjectures of complexity with Born–Infeld black holes,” [Eur. Phys. J. C](#) **77** no. 12, (2017) 817, [arXiv:1703.06297 \[hep-th\]](#).
- [34] M. Alishahiha, A. Faraji Astaneh, A. Naseh, and M. H. Vahidinia, “On complexity for F(R) and critical gravity,” [JHEP](#) **05** (2017) 009, [arXiv:1702.06796 \[hep-th\]](#).
- [35] A. Reynolds and S. F. Ross, “Complexity in de Sitter Space,” [Class. Quant. Grav.](#) **34** no. 17, (2017) 175013, [arXiv:1706.03788 \[hep-th\]](#).
- [36] M. M. Qaemmaqami, “Complexity growth in minimal massive 3D gravity,” [Phys. Rev. D](#) **97** no. 2, (2018) 026006, [arXiv:1709.05894 \[hep-th\]](#).
- [37] L. Sebastiani, L. Vanzo, and S. Zerbini, “Action growth for black holes in modified gravity,” [Phys. Rev. D](#) **97** no. 4, (2018) 044009, [arXiv:1710.05686 \[hep-th\]](#).
- [38] J. Couch, S. Eccles, W. Fischler, and M.-L. Xiao, “Holographic complexity and noncommutative gauge theory,” [JHEP](#) **03** (2018) 108, [arXiv:1710.07833 \[hep-th\]](#).
- [39] B. Swingle and Y. Wang, “Holographic Complexity of Einstein-Maxwell-Dilaton Gravity,” [JHEP](#) **09** (2018) 106, [arXiv:1712.09826 \[hep-th\]](#).
- [40] P. A. Cano, R. A. Hennigar, and H. Marrochio, “Complexity Growth Rate in Lovelock Gravity,” [Phys. Rev. Lett.](#) **121** no. 12, (2018) 121602, [arXiv:1803.02795 \[hep-th\]](#).
- [41] S. Chapman, H. Marrochio, and R. C. Myers, “Holographic complexity in Vaidya spacetimes. Part I,” [JHEP](#) **06** (2018) 046, [arXiv:1804.07410 \[hep-th\]](#).
- [42] S. Chapman, H. Marrochio, and R. C. Myers, “Holographic complexity in Vaidya

- spacetimes. Part II,” [JHEP](#) **06** (2018) 114, [arXiv:1805.07262 \[hep-th\]](#).
- [43] R. Auzzi, S. Baiguera, M. Grassi, G. Nardelli, and N. Zenoni, “Complexity and action for warped AdS black holes,” [JHEP](#) **09** (2018) 013, [arXiv:1806.06216 \[hep-th\]](#).
- [44] E. Yaraie, H. Ghaffarnejad, and M. Farsam, “Complexity growth and shock wave geometry in AdS-Maxwell-power-Yang–Mills theory,” [Eur. Phys. J. C](#) **78** no. 11, (2018) 967, [arXiv:1806.07242 \[gr-qc\]](#).
- [45] M. Alishahiha, A. Faraji Astaneh, M. R. Mohammadi Mozaffar, and A. Mollabashi, “Complexity Growth with Lifshitz Scaling and Hyperscaling Violation,” [JHEP](#) **07** (2018) 042, [arXiv:1802.06740 \[hep-th\]](#).
- [46] Y.-S. An and R.-H. Peng, “Effect of the dilaton on holographic complexity growth,” [Phys. Rev. D](#) **97** no. 6, (2018) 066022, [arXiv:1801.03638 \[hep-th\]](#).
- [47] R.-G. Cai, S.-M. Ruan, S.-J. Wang, R.-Q. Yang, and R.-H. Peng, “Action growth for AdS black holes,” [JHEP](#) **09** (2016) 161, [arXiv:1606.08307 \[gr-qc\]](#).
- [48] A. Ghodsi, S. Qolibikloo, and S. Karimi, “Holographic complexity in general quadratic curvature theory of gravity,” [Eur. Phys. J. C](#) **80** no. 10, (2020) 920, [arXiv:2005.08989 \[hep-th\]](#).
- [49] A. M. Frassino, R. B. Mann, and J. R. Mureika, “Extended Thermodynamics and Complexity in Gravitational Chern-Simons Theory,” [JHEP](#) **11** (2019) 112, [arXiv:1906.07190 \[gr-qc\]](#).
- [50] F. J. G. Abad, M. Kulaxizi, and A. Parnachev, “On Complexity of Holographic Flavors,” [JHEP](#) **01** (2018) 127, [arXiv:1705.08424 \[hep-th\]](#).
- [51] D. S. Ageev and I. Y. Aref’eva, “Holography and nonlocal operators for the BTZ black hole with nonzero angular momentum,” [Theor. Math. Phys.](#) **180** (2014) 881–893, [arXiv:1402.6937 \[hep-th\]](#).
- [52] Y.-T. Zhou, X.-M. Kuang, and J.-P. Wu, “Complexity growth of massive black hole with a probe string,” [Eur. Phys. J. C](#) **81** no. 8, (2021) 768, [arXiv:2104.12998 \[hep-th\]](#).
- [53] Y.-T. Zhou, “Complexity growth of BTZ black hole in massive gravity with a null string,” [Eur. Phys. J. C](#) **83** no. 12, (2023) 1109, [arXiv:2302.10565 \[hep-th\]](#).
- [54] K. Nagasaki, “Complexity growth of rotating black holes with a probe string,” [Phys. Rev. D](#) **98** no. 12, (2018) 126014, [arXiv:1807.01088 \[hep-th\]](#).
- [55] K. Nagasaki, “Complexity growth for topological black holes by holographic method,” [Int. J.](#)

- [Mod. Phys. A](#) **35** no. 25, (2020) 2050152, [arXiv:1912.03567 \[hep-th\]](#).
- [56] M. Bravo-Gaete and F. F. Santos, “Complexity of four-dimensional hairy anti-de-Sitter black holes with a rotating string and shear viscosity in generalized scalar–tensor theories,” [Eur. Phys. J. C](#) **82** no. 2, (2022) 101, [arXiv:2010.10942 \[hep-th\]](#).
- [57] F. F. Santos, “Rotating black hole with a probe string in Horndeski Gravity,” [Eur. Phys. J. Plus](#) **135** no. 10, (2020) 810, [arXiv:2005.10983 \[hep-th\]](#).
- [58] K. Nagasaki, “Probe strings on anti-de Sitter accelerating black holes,” [PTEP](#) **2022** no. 4, (2022) 043B02, [arXiv:2108.05429 \[hep-th\]](#).
- [59] K. Nagasaki, “Effects of the acceleration on holographic complexity,” [Int. J. Mod. Phys. A](#) **38** no. 04n05, (2023) 2350027, [arXiv:2205.00196 \[hep-th\]](#).
- [60] J. de Boer, V. E. Hubeny, M. Rangamani, and M. Shigemori, “Brownian motion in AdS/CFT,” [JHEP](#) **07** (2009) 094, [arXiv:0812.5112 \[hep-th\]](#).
- [61] A. N. Atmaja, J. de Boer, and M. Shigemori, “Holographic Brownian Motion and Time Scales in Strongly Coupled Plasmas,” [Nucl. Phys. B](#) **880** (2014) 23–75, [arXiv:1002.2429 \[hep-th\]](#).
- [62] A. E. Lawrence and E. J. Martinec, “Black hole evaporation along macroscopic strings,” [Phys. Rev. D](#) **50** (1994) 2680–2691, [arXiv:hep-th/9312127](#).
- [63] E. Fuchs and M. Kroyter, “Analytical Solutions of Open String Field Theory,” [Phys. Rept.](#) **502** (2011) 89–149, [arXiv:0807.4722 \[hep-th\]](#).
- [64] W. Taylor and B. Zwiebach, “D-branes, tachyons, and string field theory,” pp. 641–759. 10, 2003. [arXiv:hep-th/0311017](#).
- [65] L. Rastelli, “String field theory,” 9, 2005. [arXiv:hep-th/0509129](#).
- [66] W. Taylor, “String field theory,” [arXiv:hep-th/0605202](#).
- [67] M. Kiermaier, Y. Okawa, and B. Zwiebach, “The boundary state from open string fields,” [arXiv:0810.1737 \[hep-th\]](#).
- [68] M. Kudrna, C. Maccaferri, and M. Schnabl, “Boundary State from Ellwood Invariants,” [JHEP](#) **07** (2013) 033, [arXiv:1207.4785 \[hep-th\]](#).
- [69] M. Bañados, C. Teitelboim, and J. Zanelli, “Black hole in three-dimensional spacetime,” [Phys. Rev. Lett.](#) **69** (Sep, 1992) 1849–1851.
- [70] M. Bañados, M. Henneaux, C. Teitelboim, and J. Zanelli, “Geometry of the 2+1 black hole,” [Phys. Rev. D](#) **48** (Aug, 1993) 1506–1525.

- [71] Y. Kinar, E. Schreiber, J. Sonnenschein, and N. Weiss, “Quantum fluctuations of Wilson loops from string models,” [Nucl. Phys. B](#) **583** (2000) 76–104, [arXiv:hep-th/9911123](#).
- [72] T. Erler, “Four lectures on analytic solutions in open string field theory,” [Phys. Rept.](#) **980** (2022) 1–95, [arXiv:1912.00521 \[hep-th\]](#).
- [73] J. Polchinski, [String theory, Vol. 1 & 2](#), Cambridge University Press.
- [74] C. Maccaferri, F. Marino, and B. Valsesia, “Introduction to String Theory,” [arXiv:2311.18111 \[hep-th\]](#).
- [75] J. Garriga and A. Vilenkin, “Quantum fluctuations on domain walls, strings and vacuum bubbles,” [Phys. Rev. D](#) **45** (1992) 3469–3486.