

# McCormick envelopes in mixed-integer PDE-constrained optimization

Sven Leyffer<sup>1</sup> and Paul Manns<sup>2</sup>

<sup>1</sup>Mathematics and Computer Science Division, Argonne National Laboratory,  
Lemont, IL 60439, USA, *leyffer@anl.gov*

<sup>2</sup>Faculty of Mathematics, TU Dortmund University, 44227 Dortmund, Germany,  
*paul.manns@tu-dortmund.de*

June 13, 2024

## Abstract

McCormick envelopes are a standard tool for deriving convex relaxations of optimization problems that involve polynomial terms. McCormick envelopes provide lower bounds, for example, in branch-and-bound procedures for mixed-integer nonlinear programs but have not gained much attention in infinite-dimensional optimization so far. This lack of attention may be due to the distributed nature of infinite-dimensional problems, which on the one hand leads to infinitely many constraints (generally state constraints that may be difficult to handle) that are added to the problem under consideration and on the other hand renders bound-tightening procedures that successively improve these convex envelopes computationally intractable.

We take on the challenge and analyze McCormick envelopes for a model problem class that is governed by a semilinear PDE involving a bilinearity. We approximate the McCormick envelopes by averaging them over the cells of a discretization of the computational domain. This process yields convex problems that underestimate the original problem up to an a priori error estimate that depends on the mesh size of the discretization. These approximate McCormick envelopes can be improved by means of an optimization-based bound-tightening procedure and can be shown to  $\Gamma$ -converge to a limit problem with a pointwise formulation of the McCormick envelopes.

We provide a computational example, for which we certify all of our imposed assumptions. The results point to both the potential of the methodology and the gaps in the research that need to be closed.

Our methodology provides a framework first for obtaining pointwise underestimators for nonconvexities and second for approximating them with finitely many linear inequalities in a superordinate infinite-dimensional setting.

## 1 Introduction

We are interested in the global optimization of mixed-integer PDE-constrained optimization problems (MIPDECOS) that feature nonconvex terms. MIPDECOS arise in many

real-world applications such as topology optimization [2, 19, 22] and supply network optimization [16, 32]. A prototypical problem class is

$$\begin{aligned} \min_{u, w} \quad & j(u, w) + \alpha R(w) \\ \text{s.t.} \quad & Au + N(u, w) = 0 \\ & w \in C \text{ and } w(x) \in \mathbb{Z} \text{ pointwise almost everywhere (a.e.) in } \Omega, \end{aligned} \tag{MIPDECO}$$

where  $w$  is a measurable input function (generally also called *control*) on a computational domain  $\Omega \subset \mathbb{R}^d$ ,  $d \in \mathbb{N}$ , which enters the problem as an input of the state equation (PDE)  $Au + N(u, w) = 0$ , which we assume to have a unique solution  $u$  for a given  $w$ . The operator  $A$  could be an elliptic differential operator and  $N(u, w)$  a lower-order term that may involve a nonlinearity. The function  $w$  needs to lie in a convex set  $C$ , for example, abide by bound constraints, and additionally satisfy the pointwise integrality restriction  $w(x) \in \mathbb{Z}$  in  $\Omega$ . The pair of  $w$  and the implied solution  $u$  to the state equation will minimize an objective term  $j$ , which is often a quadratic fidelity term on  $u$ . Moreover, desired structures on  $w$  can be promoted by means of the regularization term  $\alpha R(w)$  for some  $\alpha > 0$ .

A beneficial choice for  $R$  has been the regularization  $R = \text{TV}$ , where TV denotes the total variation seminorm on the space of integrable functions [1]. The total variation of an integer-valued function is the  $d - 1$ -dimensional volume of the interfaces between its level sets weighted by the jump heights over the respective interfaces. This choice implies that instances of (MIPDECO) admit solutions under mild assumptions; see, for example, [8, 27]. Recent work on solving such problems to the satisfaction of the stationarity concept has been studied in [21, 25, 27, 37]. The computation of lower bounds for a linear PDE, thereby yielding a convex set after omitting the integrality constraints, and their integration into a branch-and-bound algorithm have recently been studied by Buchheim et al. in [3–7] for  $d = 1$ .

In this work we strive to obtain convex relaxations and thus lower bounds on (MIPDECO) in the presence of a structured nonlinearity of the feasible set induced by a specific choice of  $N$ . To be precise, we restrict our work to the case that  $A$  is an elliptic operator,  $R = \text{TV}$ , and the nonlinearity  $N$  in the state equation is in fact a bilinearity, specifically

$$N(u, w)(x) := u(x)w(x) \text{ for } x \in \Omega.$$

We replace the nonlinear state equation by a linear state equation and linear inequalities by generalizing McCormick envelopes that have been introduced for finite-dimensional nonconvex optimization problems in [28]. Our aim is to obtain valid and (ideally) tight lower bounds for the superordinate instances of (MIPDECO). They have already been studied in the context of global optimization with ODEs and finite-dimensional control inputs in [30, 36, 38].

In order to improve the tightness, our goal is to combine the generalized McCormick envelopes with an optimality-based bound-tightening procedure in order to improve the lower bounds provided by the McCormick envelope. This bound-tightening procedure can become very compute-intensive, however. Moreover, state-constrained optimal control problems, in particular with more than finitely many state constraints, are notoriously

hard to analyze and solve.

Therefore, we introduce an approximation scheme that allows us to work with a finite number of linear inequalities and variables that is substantially smaller than the number of variables used for the PDE ansatz. Using beneficial regularity or continuity properties of the solution to the underlying (discretized) PDE, specifically that its solution attains relatively close values in grid cells that are spatially close to each other, we expect that a moderate mesh size may already yield a good approximation of the lower bound. In this way we strive for a much faster computation of the approximate McCormick envelopes in order to obtain approximate lower bounds on (MIPDECO), which are required for our superordinate goal of solving instances of (MIPDECO) to global optimality. We highlight that the computation of such lower bounds is also interesting for global optimization in its own right even without the context of integer-valued input functions in the superordinate problem.

We believe that our restrictions on the setting are sensible. First,  $R = \text{TV}$  is a useful regularization for MIPDECOs because it results in crisp designs (level sets of the input function). Moreover, the analytical properties of the TV-regularization term also help streamline our analysis although the general strategy can also be carried out for many other choices of  $R$ . Similarly, higher-order multilinear terms for  $N(u, w)$  involving  $u$ , for example,  $u^k w$ ,  $k \geq 2$ , would make the PDE analysis, namely, the existence of solutions and their regularity, considerably more complicated. In this regard many of our arguments are streamlined and do not distract from the analysis and approximation of the McCormick envelopes. Consequently, this article will be read as a recipe to first obtain a pointwise generalization of underestimators for nonconvexities as are known from finite-dimensional nonconvex optimization and then approximate this pointwise generalization with finitely many linear inequalities using approximation properties that are due to the PDE setting.

## Contribution

We provide a formal definition of an optimal control problem with the aforementioned features and derive McCormick envelopes using pointwise constraints. We then introduce a grid that discretizes the computational domain  $\Omega$ , and we approximate the inequalities defining the McCormick envelope by a local averaging over the grid cells. In a second step, we discretize the control input function on the same grid, thereby reducing the complexity of the problem further. We prove the existence of solutions to the approximate convex relaxations and an estimate on the lower bound for the optimal control problem depending on the mesh size of the grid. Moreover, we prove that minimizers of the approximate convex relaxations converge to minimizers of the convex relaxation obtained by imposing the McCormick envelopes by means of pointwise constraints when driving the mesh size to zero.

We introduce and analyze the aforementioned optimality-based bound-tightening procedure that allows us to tighten given bounds on the state variable  $u$  and in turn to increase the objective value of the (approximate) McCormick envelope and thus to tighten the induced lower bounds on the optimal objective value of the superordinate optimal control problem.

We then verify all assumptions imposed on our analysis for an exemplary optimal

control problem with  $\Omega \subset \mathbb{R}$ . In particular, this example allows us to provide tight values for all constants in the estimates we impose. We then give details on how we discretize the PDE and set up a computational experiment where we insert the aforementioned constants. We compute and compare the approximate lower bounds with and without the bound-tightening procedure applied with each other to upper bounds with and without an integrality restriction on the controls in order to assess the quality of the analyzed methodology.

## Structure of the remainder

We continue with a brief introduction to our notation. In [section 2](#) we introduce McCormick envelopes for optimal control problems with semilinear state equations that feature a bilinear term. The McCormick envelopes are first introduced and analyzed in a pointwise fashion. Then a local averaging is introduced in two steps to approximate the pointwise inequalities by finitely many linear inequalities and the control discretization by a piecewise constant control ansatz, as is desired in our motivating application of integer optimal control. In [section 3](#) we introduce an elliptic PDE that we use as a showcase to verify all of the assumptions that are imposed for the well-definedness and approximation properties of the different McCormick settings from [section 2](#). We perform our computational experiments on this example in [section 4](#) and draw a conclusion in [section 5](#).

## Notation

Let  $\Omega \subset \mathbb{R}^d$ ,  $d \in \mathbb{N}$ , be a bounded domain. The projection in the space of integrable functions  $L^1(\Omega)$  to piecewise constant functions on a partition  $\mathcal{Q}_h = \{Q_h^1, \dots, Q_h^{N_h}\}$  of  $\Omega$  is denoted by  $P_h$ ; specifically

$$P_h : L^1(\Omega) \ni f \mapsto \sum_{i=1}^{N_h} \frac{1}{|Q_h^i|} \int_{Q_h^i} f(x) \, dx \chi_{Q_h^i} \in L^1(\Omega). \quad (1)$$

For a measurable set  $A \subset \mathbb{R}^d$ ,  $d \in \mathbb{N}$ , we denote its  $d$ -dimensional Lebesgue measure by  $|A|$ . We say that a function  $w \in L^1(\Omega)$  is of bounded variation if its total variation seminorm  $\text{TV}(w)$  is finite. The Banach space of functions in  $L^1(\Omega)$  of bounded variation with norm  $\|\cdot\|_{\text{BV}} = \|\cdot\|_{L^1} + \text{TV}(\cdot)$  is then denoted by  $\text{BV}(\Omega)$ ; see [\[1\]](#) for details on functions of bounded variation. We recall that the total variation seminorm for  $w \in L^1(\Omega)$  is defined as

$$\text{TV}(w) := \sup \left\{ \int_{\Omega} w(x) \operatorname{div} \phi(x) \, dx \mid \phi \in C_c^1(\Omega, \mathbb{R}^d), \max_{x \in \Omega} \|\phi(x)\| \leq 1 \right\},$$

where  $\|\cdot\|$  denotes the Euclidean norm on  $\mathbb{R}^d$ . On one-dimensional domains  $\Omega = (a, b)$  for  $a, b \in \mathbb{R}$ , every element of  $\text{BV}(\Omega)$  has a left-continuous representative, and the total variation is the sum of the jump heights of the left-continuous representative (this interpretation also works for the right-continuous representative or other so-called good representatives; see [\[1\]](#)).

## 2 McCormick envelopes for optimal control problems with state equations that have bilinear terms

In this section we derive and analyze McCormick envelopes for a class of nonconvex optimal control problems where the nonconvexity stems from bilinear terms in the state equation. We first introduce a prototypical optimal control problem in [section 2.1](#). We then derive pointwise a.e. McCormick envelopes in [section 2.2](#) and, subsequently, provide a local averaging in two stages that yields approximate lower bounds but implies problems that feature only finitely many additional linear inequalities in [section 2.3](#). In [section 2.4](#) we prove the validity of a bound-tightening procedure that may improve (increase) the (approximate) lower bounds.

### 2.1 Optimal control problem with nonconvex bilinearity

We consider the following prototypical setting with a nonconvexity induced by a bilinear term in the state equation, but note that the strategy we develop can be transferred to many other settings with lower or slightly different regularity assumptions with a small amount of modifications.

Let  $\Omega$  be a bounded domain. Let  $U$  be a reflexive Banach space that is compactly embedded into  $H := L^2(\Omega)$ , that is,  $U \hookrightarrow^c H$ , so that we may work with the triple  $U \hookrightarrow^c H \cong H^* \hookrightarrow^c U^*$  of compact embeddings and the isometric isomorphic identification  $H \cong H^*$ . In optimal control terminology,  $U$  is the *state space*. For the *control space*, we consider the space  $W := \text{BV}(\Omega)$  of functions of bounded variation, which is motivated by the compactness and approximation properties it provides in the control space and our intended application in the context of integer optimal control. Specifically,  $\text{BV}(\Omega)$ -regularity allows us to approximate a function by its average on a partition of the domain with an error that is bounded by a scalar multiple of the mesh size. Note that other spaces like the Sobolev space  $H^1(\Omega)$  also provide this property.

The optimal control problem is

$$\begin{aligned} \min_{u,w} \quad & j(u, w) + \alpha \text{TV}(w) \\ \text{s.t.} \quad & Au + uw = f \quad \text{in } U^*, \\ & w \in C, u \in U \end{aligned} \tag{OCP}$$

for a nonempty, bounded, closed, and convex set  $C \subset H$ , an objective  $j : U \times H \rightarrow \mathbb{R}$  that is Lipschitz continuous on bounded sets and a linear and bounded differential operator  $A : U \rightarrow U^*$  with bounded inverse  $A^{-1} : U^* \rightarrow U$ . The nonconvexity of the problem stems from the bilinear term  $uw$  in the state equation.

Note that the assumptions on  $C$  imply that it is a weakly sequentially compact subset of  $H$ . The vector  $f \in U^*$  is a fixed datum that parameterizes the optimization problem, and we assume that the product  $uw$  is a measurable function that is (also) an element of  $U^*$  for all tuples  $(u, w) \in U \times W$ . This follows, for example, if  $C$  is bounded in  $L^\infty(\Omega)$ , too. The term  $\text{TV}(w)$  denotes the total variation seminorm of  $w$  that enforces its required  $\text{BV}(\Omega)$ -regularity, and  $\alpha > 0$  is a positive scalar. Assuming that the state equation and the original optimal control problem admit unique solutions, the presence of the term  $uw$

implies a nonlinear dependence of the solution to the state equation on  $w$ , which in turn implies that (OCP) is generally nonconvex even if  $j$  is convex.

## 2.2 Pointwise McCormick envelopes

McCormick envelopes have been introduced for finite-dimensional nonconvex optimization problems in [28]. The key idea for the case of (OCP) is to derive a new optimal control problem, the so-called McCormick envelope, by relaxing the state constraints. Specifically, the bilinear term is replaced by a new variable  $z$ , which is a measurable function that is also an element of  $U^*$ , and a set of linear inequalities on  $z$ ,  $u$ , and  $w$  that preserve the feasibility of the state equation; that is, if  $u$  and  $w$  satisfy  $Au = uw + f$ , then the choice  $z = uw$  satisfies all of the newly introduced linear inequalities. Here, we make the following assumption.

**Assumption 2.1.** Let there be bounds  $u_\ell, u_u \in H$  and  $w_\ell, w_u \in L^\infty(\Omega)$  so that  $u_\ell \leq u \leq u_u$  and  $w_\ell \leq w \leq w_u$  hold pointwise a.e. for all  $(u, w) \in U \times W$  that solve  $Au + uw = f$  and satisfy  $w \in C$ .

Clearly, the multiplication of these bounds,  $u_\ell w_\ell, u_\ell w_u, u_u w_\ell, u_u w_u \in H$  by Hölder's inequality. We note that the assumption of bounds on the state is quite strong, but we emphasize that such bounds are generally required only in the subset of the computational domain  $\Omega$  on which the nonlinearity is defined so that higher interior regularity of the PDE solutions can help establish implementable (i.e., generally uniform) bounds.

We note that the different parts of the proofs do not require the full assumed regularity here, but again we prefer this slightly more restrictive setting in the interest of a cleaner presentation. Under **assumption 2.1**, the McCormick envelope of (OCP) can be derived pointwise a.e. in a similar way as for finite-dimensional problems; see [28]. It is as follows.

$$\begin{aligned}
& \min_{u, w, z} j(u, w) + \alpha \text{TV}(w) \\
& \text{s.t. } Au + z = f \quad \text{in } U^*, \\
& \quad z \geq u_\ell w + uw_\ell - u_\ell w_\ell \quad \text{a.e.}, \\
& \quad z \geq u_u w + uw_u - u_u w_u \quad \text{a.e.}, \\
& \quad z \leq u_u w + uw_\ell - u_u w_\ell \quad \text{a.e.}, \\
& \quad z \leq u_\ell w + uw_u - u_\ell w_u \quad \text{a.e.}, \\
& \quad u \in U, \quad u_\ell \leq u \leq u_u \quad \text{a.e.}, \\
& \quad w \in C, \quad w_\ell \leq w \leq w_u \quad \text{a.e.}
\end{aligned} \tag{McC}$$

We immediately obtain that (McC) is a relaxation of (OCP) with a convex feasible set.

**Proposition 2.2.** Let **assumption 2.1** hold.

1. If  $(u, w)$  is feasible for (OCP), then  $(u, w, z)$  with  $z = uw$  is feasible for (McC). In particular, the infimum (McC) is a lower bound on the infimum of (OCP).
2. The feasible set of (McC) is convex and bounded in  $U \times H \times H$ .

*Proof.* Regarding feasibility, the state equation, the inclusion in  $C$ , and the two last pointwise inequalities are immediate. The satisfaction of the four additional pointwise inequalities for the choice  $z = uw$  follows as in the finite-dimensional case by rearranging them.

For example, we obtain  $(u - u_\ell)w \geq (u - u_\ell)w_\ell$  a.e. for the first inequality. The state equation in (McC) and the additional inequalities are affine in  $(u, w, z)$ . Moreover,  $C$  is assumed to be convex so that (McC) has a convex feasible set. The feasible  $w$  are bounded in  $H$ . In combination with the pointwise bounds on  $u$ , we obtain boundedness of the feasible  $z$  in  $H$  from the McCormick inequalities. In turn, the continuous invertibility of  $A$  and the embedding  $H \cong H^* \hookrightarrow^c U^*$  imply the boundedness of the feasible  $u$  in  $U$ .  $\square$

*Remark 2.3.* The pointwise a.e. bound constraints  $u_\ell \leq u \leq u_u$  are unattractive in optimal control settings because they involve only the state variable  $u$ , which may imply that the regularity of corresponding Lagrange multipliers is low (a measure); see, for example, [9, 10]. This may necessitate additional regularization (see, e.g., [29]) and cause difficulties in computational approaches. Because PDE-constrained optimization problems generally feature a large number of variables after discretization due to their distributed nature, the McCormick inequalities add many linear constraints—on the order of six times the number of discretization cells—to the fully discretized problem.

Without any additional constraints, we do not know whether we can actually bound  $z$  uniformly in  $H$ , however, so we add these bound constraints on  $u$ , which are again satisfied by all feasible points for (OCP).

**Proposition 2.4.** *In the setting of section 2.1, the problems (OCP) and (McC) admit solutions.*

*Proof.* The existence of solutions for (OCP) follow with standard arguments from the direct method of calculus of variations.

By construction of (McC) as a relaxation of (OCP), its feasible set is nonempty, and there exists a minimizing sequence  $(u^n, w^n, z^n)_n \subset U \times H \times H$  of feasible points, which has at least one accumulation point  $(u, w, z) \in H \times H \times H$  with respect to weak convergence in  $H \times H \times H$ , which follows directly from proposition 2.2. Because  $A^{-1}$  is a bounded, linear operator,  $u$  satisfies the state equation and is an element of  $U$ , too. Moreover, because  $H$  is compactly embedded in  $U^*$  and thus  $z^n \rightarrow z$  holds in  $U^*$ , we even obtain

$$u^n = A^{-1}(f - z^n) \rightarrow A^{-1}(f - z) = u \text{ in } U$$

and in turn  $u^n \rightarrow u$  in  $H$  and pointwise a.e. after restricting to a suitable subsubsequence. The pointwise a.e. convergence yields  $u_\ell \leq u \leq u_u$ .

The assumptions on  $C$  imply  $w \in C$ . Because  $j$  is bounded below,  $(\text{TV}(w^n))_n$  is bounded; and again after restricting to a suitable subsubsequence, we obtain  $w^n \rightarrow w$  in  $L^1(\Omega)$  and pointwise a.e. from the weak-\* sequential compactness properties of the TV-seminorm; see Theorem 3.23 in [1]. We obtain  $w_\ell \leq w \leq w_u$ .

It remains to assert that the weak limit  $z$  satisfies the McCormick inequalities before proving the optimality. This follows from lemma A.1.

The continuity properties of  $j$  and the lower semi-continuity of the TV-seminorm with respect to convergence in  $L^1(\Omega)$  and in turn also in  $H$  yield that  $(u, w, z)$  is a minimizer.  $\square$

*Remark 2.5.* In our examples, the space  $W$  will be (a subspace of) the space of functions of bounded variation  $\text{BV}(0, 1)$ , which is not reflexive. There are important sets  $C$  that are subsets of such spaces and also convex, bounded, and weakly sequentially closed in  $H$ . An




Figure 1: Two-stage approximation of  $(\text{McC})$  by reduced problems  $(\text{McC}_h)$  and  $(\text{McC}_{hh})$  in order to obtain approximate lower bounds on  $(\text{OCP})$ .

example is  $\{w \in \text{BV}(0,1) \mid \text{TV}(w) \leq \kappa\}$  for some  $\kappa > 0$ ; see also [4] for such a constraint set in the context of integer optimal control.

The challenges of the pointwise McCormick envelopes for  $(\text{McC})$  have already been sketched in remark 2.3. The situation is further complicated if one wants to improve the lower bound induced by  $(\text{McC})$ . A straightforward application of a bound-tightening procedure implies that one alternately picks  $\tilde{x} \in \Omega$  and solves (one of) the optimization problems

$$\inf_{u,w,z} u(\tilde{x}) \text{ s.t. } (u, w, z) \text{ is feasible for } (\text{McC})$$

and

$$\sup_{u,w,z} u(\tilde{x}) \text{ s.t. } (u, w, z) \text{ is feasible for } (\text{McC})$$

in order to then update  $u_\ell(\tilde{x})$  and  $u_u(\tilde{x})$  to the computed values. Of course, this is sensible only if  $u \in C(\bar{\Omega})$  or has a meaningful pointwise interpretation. In a practical implementation, such optimization problems may, for example, be solved for nodes  $\tilde{x}$  of a finite-element discretization with a nodal basis. Since improving the bounds yields further possible improvements, it makes sense to repeatedly solve such problems until no further progress is made. In total, the bound-tightening procedure can become computationally very expensive. Therefore, in the next section we analyze a local averaging of the McCormick envelopes that leads to approximate lower bounds.

### 2.3 Locally averaged McCormick envelopes

We propose, analyze, and assess a local averaging that modifies the problem before introducing the variable  $z$ . Thus, the variable  $z$  and in a second approximation step  $w$ , which only enters the relaxed PDE implicitly through  $z$ , and the bounds on the product  $uw$  are discretized on a coarser grid that allows us to reduce the size of the resulting optimization problem. We give an overview on the approximation arguments of this subsection in fig. 1.

Let  $\mathcal{Q}_h = \{Q_h^1, \dots, Q_h^N\}$  be a partition of the domain  $\Omega$  into intervals / squares / cubes



/ ... with mesh size  $h$ . Locally averaging the nonlinearity of the state equation from (McC) means that  $Au + uw = f$  is replaced by the variant

$$Au + (P_h u)(P_h w) = f \quad (2)$$

with  $P_h$  being defined in (1). We define the locally averaged McCormick envelope with respect to the partition  $\mathcal{Q}_h$  as

$$\begin{aligned} \min_{\substack{u, w, \\ z_1, \dots, z_{N_h}}} \quad & j(u, w) + \alpha \text{TV}(w) \\ \text{s.t.} \quad & Au + z = f \quad \text{in } U^*, \\ & z = \sum_{i=1}^{N_h} z_i \chi_{Q_h^i}, \\ & z_i \geq u_\ell^i (P_h w) + (P_h u) w_\ell^i - u_\ell^i w_\ell^i \quad \text{on } Q_h^i \text{ for all } i \in \{1, \dots, N_h\}, \\ & z_i \geq u_u^i (P_h w) + (P_h u) w_u^i - u_u^i w_u^i \quad \text{on } Q_h^i \text{ for all } i \in \{1, \dots, N_h\}, \\ & z_i \leq u_\ell^i (P_h w) + (P_h u) w_\ell^i - u_\ell^i w_\ell^i \quad \text{on } Q_h^i \text{ for all } i \in \{1, \dots, N_h\}, \\ & z_i \leq u_u^i (P_h w) + (P_h u) w_u^i - u_u^i w_u^i \quad \text{on } Q_h^i \text{ for all } i \in \{1, \dots, N_h\}, \\ & u \in U, u_\ell^i \leq P_h u \leq u_u^i \quad \text{on } Q_h^i \text{ for all } i \in \{1, \dots, N_h\}, \\ & w \in C, w_\ell \leq w \leq w_u \text{ a.e.}, \\ & z_1, \dots, z_{N_h} \in \mathbb{R}. \end{aligned} \quad (\text{McC}_h)$$

In (McC<sub>h</sub>) the statement that the four McCormick inequalities on  $z_i$  hold on  $Q_h^i$  might be misunderstood. We therefore highlight that  $P_h u$  and  $P_h w$  are constant on the cells  $Q_h^i$  so that there are exactly four McCormick inequalities per cell  $Q_h^i \in \mathcal{Q}_h$ . The same argument applies for the bounds on  $P_h u$ . The bounds  $u_\ell^i \in \mathbb{R}$  and  $u_u^i \in \mathbb{R}$  assume the roles of the functions  $u_\ell$  and  $u_u$  per grid cell for the locally averaged function  $P_h u$ . Specifically, the function  $P_h u$  is bounded from below by  $u_\ell^i$  and from above by  $u_u^i$  on the  $i$ -th grid cell  $Q_h^i$ . Similarly, the bounds  $w_\ell^i$  and  $w_u^i$  assume the roles of  $w_\ell$  and  $w_u$ . Note that the problem is still infinite-dimensional due to the variables  $u$  and  $w$ .

By employing only finitely many bounds on  $P_h u$ , we can circumvent the aforementioned regularity problems if the number of constraints is not too large; that is, its Lagrange multiplier may sensibly be interpreted as a vector in  $\mathbb{R}^n$ . Moreover, the local averaging opens up possibilities for adaptive McCormick envelopes when using a corresponding adaptive refinement of the partition  $\mathcal{Q}_h$  and will also reduce the computational effort for improving the (approximate) lower bound. We can now impose inequalities on the locally averaged state  $P_h u$  and tighten them algorithmically in order to reduce the gap between the relaxation and the original problem. We now impose an assumption that implies that (McC<sub>h</sub>) admits a solution and is an approximate relaxation of (OCP). This means that the optimal objective of (OCP) is bounded from below by the objective of (McC<sub>h</sub>) minus a bound on the approximation error.

**Assumption 2.6.** In addition to the setting from section 2.1, we assume the following.

1. Let  $\{\mathcal{Q}_h\}_h$  be a sequence of partitions of the domain  $\Omega$  with mesh sizes  $h$ .
2. Let the equation (2) have a unique solution for all  $w \in \text{BV}(\Omega)$ .

3. For all  $h$  and all  $Q_i \in \mathcal{Q}_h$ , let the bounds  $u_\ell^i, u_u^i, w_\ell^i, w_u^i \in \mathbb{R}$  be such that  $u_\ell^i \leq (P_h u)|_{Q_i} \leq u_u^i$  and  $w_\ell^i \leq (P_h w)|_{Q_i} \leq w_u^i$  hold for all  $w \in C \cap [w_\ell, w_u]$  and  $u \in U$  that solve  $Au + (P_h u)(P_h w) = f$ .

We will verify all parts of [assumption 2.6](#) for an example in [section 3](#).

**Theorem 2.7.** *Let [assumption 2.6](#) hold. Then  $(\text{McC}_h)$  has a convex feasible set that is bounded in  $U \times H \times \mathbb{R}^{N_h}$  and admits a minimizer. Let  $(u, w)$  be feasible for  $(\text{OCP})$ . Then the tuple  $(u_h, w_h, z_h)$  with  $w_h := w$ ,  $u_h$  being the unique solution to (2), and  $z_h := (P_h u_h)(P_h w_h)$  is feasible for  $(\text{McC}_h)$ , and the objective value satisfies*

$$|j(u_h, w_h) + \alpha \text{TV}(w_h) - j(u, w) - \alpha \text{TV}(w)| \leq L_u \|u - u_h\|_U, \quad (3)$$

where  $L_u$  is the Lipschitz constant of  $j$  on the feasible set of  $(\text{McC}_h)$  for the first argument and fixed  $w$ . Moreover,

$$m_{(\text{OCP})} \geq m_{(\text{McC}_h)} - L_u \|\bar{u} - \bar{u}_h\|_U \quad (4)$$

if  $m_{(\text{McC}_h)}$  denotes the infimum of  $(\text{McC}_h)$ ,  $m_{(\text{OCP})}$  denotes the infimum of  $(\text{OCP})$ ,  $(\bar{u}, \bar{w})$  denotes the minimizer of  $(\text{OCP})$ , and  $\bar{u}_h$  the corresponding solution to (2).

*Proof.* The convexity and boundedness of the feasible set follow as in [proposition 2.2](#). The feasibility of  $(u_h, w_h, z_h)$  for  $(\text{McC}_h)$  follows by construction. Because the feasible set of  $(\text{McC})$  is nonempty (see [proposition 2.4](#)), there exist feasible points of  $(\text{McC}_h)$ . Because the feasible set of  $(\text{McC}_h)$  is bounded, the desired constant  $L_u$  exists by assumption on  $j$ .

Consequently, we can consider a minimizing sequence  $\{(u_h^k, w_h^k, z_h^k)\}_k$  for  $(\text{McC}_h)$  and apply similar arguments as in the proof of [proposition 2.4](#) in order to prove existence of a minimizer. The  $z_h^k$  have a finite-dimensional piecewise constant ansatz and are bounded because of the boundedness of  $w$  implied by the properties of  $C$  so that they admit a (norm-)convergent subsequence in  $H$  with limit  $\bar{z}_h$ . Because of the continuous invertibility of  $A$ , there is a corresponding subsequence of  $\{u_h^k\}_k$  that converges to  $\bar{u}_h = A^{-1}(-\bar{z}_h + f)$ . By possibly passing to a further subsequence, the corresponding subsequence of  $\{w_h^k\}_k$  converges weakly\* to  $\bar{w}_h$  in  $W$ . This implies  $\bar{w}_h \in C$  and  $w_\ell \leq \bar{w}_h \leq w_u$ . Passing to a further subsequence, we obtain pointwise a.e. convergence for all three subsequences and in turn feasibility of the limit triple  $(\bar{u}_h, \bar{w}_h, \bar{z}_h)$  for  $(\text{McC}_h)$ , where we use that the projection to the piecewise constant functions is continuous with respect to all  $L^p$ -norms. For the objective, we obtain that the first term converges because of its assumed continuity properties. Moreover, the TV-term is lower semicontinuous with respect to convergence in  $H$ ; and, in turn, we obtain that  $(\bar{u}_h, \bar{w}_h, \bar{z}_h)$  realizes the infimal objective value and is thus a minimizer.

The approximation bound (3) follows from the Lipschitz continuity of  $j$  and the assumed choice  $(u_h, w_h, z_h)$ .

To prove the approximate lower bound (4), let  $(\bar{u}, \bar{w})$  be a minimizer to  $(\text{OCP})$ , and let  $\bar{w}_h = \bar{w}$ . Then [assumption 2.6 2](#) gives a unique solution  $\bar{u}_h$  to (2), and we choose  $\bar{z}_h = (P_h \bar{u}_h)(P_h \bar{w})$ . [Assumption 2.6 3](#) yields that the triple  $(\bar{u}_h, \bar{w}_h, \bar{z}_h)$  is feasible for

(**McC<sub>h</sub>**). We deduce by means of  $\bar{w}_h = \bar{w}$  and the optimality of  $(\bar{u}, \bar{w}, \bar{z})$  that

$$\begin{aligned} m_{(\text{OCP})} &= j(\bar{u}, \bar{w}) + \alpha \text{TV}(\bar{w}) \geq j(\bar{u}_h, \bar{w}_h) + \alpha \text{TV}(\bar{w}_h) - L_u \|u - u_h\|_U \\ &\geq m_{(\text{McC}_h)} - L_u \|u - u_h\|_U. \end{aligned}$$

□

We note that approximation results on the difference  $\|u - u_h\|_U$  that are required to obtain a meaningful bound can generally be obtained for broad classes of elliptic and parabolic PDEs.

The locally averaged McCormick envelope (**McC<sub>h</sub>**) has the drawback that while only a local average of  $w$  is required for the McCormick inequalities, the full information is required in order to represent the total variation term correctly. To reduce the number of optimization variables further, one is tempted to replace  $\text{TV}(w)$  by  $\text{TV}(P_h w)$ . Then one could use  $P_h w = \sum_{i=1}^{N_h} w_i \chi_{Q_h^i}$  for real coefficients  $w_i$ ,  $i \in \{1, \dots, N_h\}$ , in order to replace the optimization over  $w$  in  $\text{BV}(\Omega)$  by an optimization over  $\mathbb{R}^N$ . The lower semicontinuity inequality  $\text{TV}(w) \leq \liminf_{h \searrow 0} \text{TV}(P_h w)$  is strict in general if the dimension of  $\Omega$  is larger than one. To make things worse, even a  $\Gamma$ -convergence result cannot be obtained directly because the geometries of the functions in the limit  $h \searrow 0$  are restricted due to the restrictions of the geometry of the elements of the partitions  $\mathcal{Q}_h$ . This is noted at the end of section 1 in [12] with references to [20] and [11] that employ discrete (anisotropic) approximations of the total variation. Specifically, the isotropic unit ball that is due to the Euclidean norm in the definition of the total variation is not recovered in the limit process for  $h \searrow 0$ , and other limit functionals that have nonisotropic unit balls may arise for periodic discretizations; this situation is analyzed and reported in [15].

We introduce a variant of the McCormick envelope where both the McCormick inequalities and the control discretization employ a local averaging with respect to the same grid based on the following assumption. To be able to obtain the desired approximation bound and a  $\Gamma$ -convergence to the limit problem (**McC**), we assume that a regularization of the total variation penalty, such as the  $L^2$ -regularization of the dual formulation presented in [13], is coupled to the discretization.

**Assumption 2.8.** In addition to **assumption 2.6**, we assume that there is a sequence of approximations  $(\text{TV}_h)_h$  of  $\text{TV}$  that is consistent with local averaging. Specifically, we assume for all  $h$  that  $\text{TV}_h : L^1(\Omega) \rightarrow [0, \infty]$  is lower semicontinuous,  $\text{TV}_h \circ P_h \leq \text{TV}$ , and  $\text{TV}_h \circ P_h \leq \text{TV}_h$ .

*Remark 2.9.* In one-dimensional domains, the choice  $\text{TV} = \text{TV}_h$  immediately satisfies **assumption 2.8**, which follows from **lemma 2.14** below. On multidimensional domains, one can, for example, choose suitable finite-element discretizations of the total variation seminorm like a approximation with lowest-order Raviart–Thomas elements; see §4.1 in [12] and §2 in [34].

The variant of the McCormick envelope that replaces  $\text{TV}(w)$  by  $\text{TV}_h(w)$  is:

$$\begin{aligned}
& \min_{\substack{u, w_1, \dots, w_{N_h}, \\ z_1, \dots, z_{N_h}}} j(u, w) + \alpha \text{TV}_h(w) \\
& \text{s.t.} \quad Au + z = f \quad \text{in } U^*, \\
& \quad w = \sum_{i=1}^{N_h} w_i \chi_{Q_h^i}, z = \sum_{i=1}^{N_h} z_i \chi_{Q_h^i}, \\
& \quad z_i \geq u_\ell^i w_i + (P_h u) w_\ell^i - u_\ell^i w_\ell^i \quad \text{on } Q_h^i \text{ for all } i \in \{1, \dots, N_h\}, \\
& \quad z_i \geq u_u^i w_i + (P_h u) w_u^i - u_u^i w_u^i \quad \text{on } Q_h^i \text{ for all } i \in \{1, \dots, N_h\}, \\
& \quad z_i \leq u_u^i w_i + (P_h u) w_\ell^i - u_u^i w_\ell^i \quad \text{on } Q_h^i \text{ for all } i \in \{1, \dots, N_h\}, \\
& \quad z_i \leq u_\ell^i w_i + (P_h u) w_u^i - u_\ell^i w_u^i \quad \text{on } Q_h^i \text{ for all } i \in \{1, \dots, N_h\}, \\
& \quad u \in U, u_\ell^i \leq P_h u \leq u_u^i \quad \text{on } Q_h^i \text{ for all } i \in \{1, \dots, N_h\}, \\
& \quad w \in C, w_\ell \leq w \leq w_u \text{ a.e.}, \\
& \quad z_1, \dots, z_{N_h} \in \mathbb{R}
\end{aligned} \tag{McC}_{hh}$$

With the help of [assumption 2.8](#), we can show that  $(\text{McC}_{hh})$  is an approximate lower bound on  $(\text{McC}_h)$  and in turn on  $(\text{OCP})$ . We note that the inequalities in [assumption 2.8](#) are implementable in practice and can be relaxed to approximate inequalities with an  $h$ -dependent error that tends to zero for  $h \searrow 0$ .

**Theorem 2.10.** *Let [assumption 2.8](#) hold. Then  $(\text{McC}_{hh})$  has a convex feasible set that is bounded in  $U \times \mathbb{R}^{N_h} \times \mathbb{R}^{N_h}$  and admits a minimizer. We define  $w_h := \sum_{i=1}^{N_h} w_i^h \chi_{Q_h^i}$  for a feasible point  $(u_{hh}, w_h^1, \dots, w_h^{N_h}, z_h^1, \dots, z_h^{N_h})$  of  $(\text{McC}_{hh})$ . Then the tuple  $(u_{hh}, w_h, z_h^1, \dots, z_h^{N_h})$  is feasible for  $(\text{McC}_h)$ .*

*Let  $(u_h, w_h, z_h^1, \dots, z_h^{N_h})$  be feasible for  $(\text{McC}_h)$ . Then  $(u_h, w_h^1, \dots, w_h^{N_h}, z_h^1, \dots, z_h^{N_h})$  with  $w_h^i := (P_h w_h)|_{Q_h^i}$ ,  $i \in \{1, \dots, N_h\}$ , is feasible for  $(\text{McC}_{hh})$ . Moreover,*

$$j(\bar{u}_h, w_h) + \alpha \text{TV}(w_h) \geq m_{(\text{McC}_{hh})} - L_w \|w_u - w_\ell\|_{L^\infty}^{\frac{1}{2}} \sqrt{d}^{\frac{1}{2}} \text{TV}(\bar{w}_h)^{\frac{1}{2}} h^{\frac{1}{2}}, \tag{5}$$

where  $m_{(\text{McC}_{hh})}$  is the infimum of  $(\text{McC}_{hh})$  and  $\bar{w}_h$  is a minimizer of  $(\text{McC}_h)$ ,  $\bar{u}_h$  the corresponding solution to [\(2\)](#), and  $L_w$  is the Lipschitz constant of  $j$  with respect to its second argument on  $C$ .

Moreover,

$$m_{(\text{OCP})} \geq m_{(\text{McC}_{hh})} - L_u \|\bar{u} - \bar{u}_h\|_U - L_w \|w_u - w_\ell\|_{L^\infty}^{\frac{1}{2}} \sqrt{d}^{\frac{1}{2}} \text{TV}(\bar{w})^{\frac{1}{2}} h^{\frac{1}{2}}, \tag{6}$$

where  $m_{(\text{OCP})}$  is the infimum of  $(\text{OCP})$ ,  $(\bar{u}, \bar{w})$  is a minimizer of  $(\text{OCP})$ , and  $L_u$  is the Lipschitz constant of  $j$  with respect to its first argument on the feasible set of  $(\text{McC}_{hh})$ .

*Proof.* The convexity and boundedness of the feasible set follow as in [proposition 2.2](#). The respective feasibility relations follow by construction, the definitions of  $(\text{McC}_h)$  and  $(\text{McC}_{hh})$ , and the assumed properties of  $C$ . The existence of minimizers for  $(\text{McC}_{hh})$  follows with the same arguments as for  $(\text{McC}_h)$  with the only difference being that the existence of a convergent subsequence in  $H$  of the control inputs  $w$  as part of a minimizing sequence can already be deduced from the finite-dimensional ansatz. Because the feasible

set of  $(\text{McC}_h)$  is bounded, the desired constant  $L_u$  exists. To prove the approximate lower bound (5), we deduce that, for all  $(u_h, w_h, z_1^h, \dots, z_{N_h}^h)$  that are feasible for  $(\text{McC}_h)$ , the estimates

$$\begin{aligned} m_{(\text{McC}_{hh})} &\leq j(u_h, P_h w_h) + \alpha \text{TV}_h(P_h w_h) \\ &\leq j(u_h, w_h) + \alpha \text{TV}_h(P_h w_h) + L_w \|w_h - P_h w_h\|_H \\ &\leq j(u_h, w_h) + \alpha \text{TV}(w_h) + L_w \|w_u - w_\ell\|_{L^\infty}^{\frac{1}{2}} \|w_h - P_h w_h\|_{L^1(\Omega)}^{\frac{1}{2}} \\ &\leq j(u_h, w_h) + \alpha \text{TV}(w_h) + L_w \|w_u - w_\ell\|_{L^\infty}^{\frac{1}{2}} \sqrt{d}^{\frac{1}{2}} \text{TV}(w_h)^{\frac{1}{2}} h^{\frac{1}{2}}, \end{aligned}$$

hold, where we have used [assumption 2.8](#) and the Lipschitz continuity of  $j$  with respect to the second argument for the second inequality and (the proof of) (12.24) in Theorem 12.26 in [23] for the third inequality.

To prove the approximate lower bound (6), we chain this estimate with the arguments from [theorem 2.7](#). Let  $(\bar{u}, \bar{w})$  be a minimizer to  $(\text{OCP})$ , and let  $\bar{w}_h = \bar{w}$ . Then [assumption 2.6 2](#) gives a unique solution  $\bar{u}_h$  to (2), and we choose  $\bar{z}_h = (P_h \bar{u}_h)(P_h \bar{w})$ . [Assumption 2.6 3](#) gives the required feasibility, and using  $P_h \bar{w}$  as well as  $(\bar{u}_h, P_h \bar{w}, \bar{z}_1^h, \dots, \bar{z}_{N_h}^h)$  in the estimate above gives

$$\begin{aligned} m_{(\text{McC}_{hh})} &\leq j(\bar{u}_h, \bar{w}) + \alpha \text{TV}(\bar{w}) + L_w \|w_u - w_\ell\|_{L^\infty}^{\frac{1}{2}} \sqrt{d}^{\frac{1}{2}} \text{TV}(\bar{w})^{\frac{1}{2}} h^{\frac{1}{2}} \\ &\leq j(\bar{u}, \bar{w}) + \alpha \text{TV}(\bar{w}) + L_u \|\bar{u} - \bar{u}_h\|_U + L_w \|w_u - w_\ell\|_{L^\infty}^{\frac{1}{2}} \sqrt{d}^{\frac{1}{2}} \text{TV}(\bar{w})^{\frac{1}{2}} h^{\frac{1}{2}}, \quad (7) \end{aligned}$$

where the second inequality follows from the Lipschitz continuity of  $j$  in the first argument.  $\square$

We now employ a  $\Gamma$ -convergence argument to prove that the locally averaged McCormick envelopes  $(\text{McC}_{hh})$  approximate  $(\text{McC})$  for  $h \searrow 0$ . To this end, we need further compatibility conditions for the bounds on the control and state variables.

**Assumption 2.11.** In addition to [assumption 2.8](#), we assume the following.

1. Let  $\{\mathcal{Q}_h\}_h$  satisfy a uniform bounded eccentricity condition; that is, there exists  $C > 0$  such that for each  $Q_i^h$  there is a ball  $B_i^h$  such that  $Q_i^h \subset B_i^h$  and  $|B_i^h| \leq C|Q_i^h|$ . See, for example, Definition 4.3 4. in [24].
2. Let  $\sup_{h \searrow 0} \text{TV}_h(w_h) \leq C$  and  $\sup_{h \searrow 0} \|w_h\|_{L^1} \leq C$  for some  $C > 0$  imply that there exists a subsequence  $\{w_h\}_h$  and  $w \in W$  such that  $w_h \rightarrow w$  in  $L^1(\Omega)$ .
3. Let  $\text{TV}_h \circ P_h$   $\Gamma$ -converge to  $\text{TV}$  for  $h \searrow 0$ .
4. Let  $u_{\ell,h} := \sum_{i=1}^{N_h} u_\ell^i \chi_{Q_i^h}$ ,  $u_{u,h} := \sum_{i=1}^{N_h} u_u^i \chi_{Q_i^h}$  satisfy  $u_{\ell,h} \rightarrow u_\ell$  in  $H$  and  $u_{u,h} \rightarrow u_u$  in  $H$ .
5. Let  $w_{\ell,h} := \sum_{i=1}^{N_h} w_\ell^i \chi_{Q_i^h}$ ,  $w_{u,h} := \sum_{i=1}^{N_h} w_u^i \chi_{Q_i^h}$  satisfy  $w_{\ell,h} \rightarrow w_\ell$  in  $H$  and  $w_{u,h} \rightarrow w_u$  in  $H$ .
6. Let  $u$  solve  $Au + uw = f$  for some  $w \in H$ . Then  $u_h \rightarrow u$  in  $U$  holds for the solutions  $u_h$  to (2) for  $h \searrow 0$ .

*Remark 2.12.* We note that Assumptions 2.11, 4, and 5 can generally be ensured by inferring suitable bounds  $w_{\ell,h}$ ,  $w_{u,h}$ ,  $u_{\ell,h}$ ,  $u_{u,h}$  from valid bounds  $w_\ell$ ,  $w_u$ ,  $u_\ell$ ,  $u_u$ . The other assumptions are quite typical when discretizing such problems.

**Proposition 2.13.** *Let assumption 2.11 hold. Let the  $\{(\bar{u}_{hh}, \bar{w}_h^1, \dots, \bar{w}_h^{N_h}, \bar{z}_h^1, \dots, \bar{z}_h^{N_h})\}_h$  be solutions to the problems (McC<sub>hh</sub>) for  $h \searrow 0$ . Then the sequence  $\{(\bar{u}_{hh}, \bar{w}_h, \bar{z}_h)\}_h$  with  $\bar{w}_h := \sum_{i=1}^{N_h} \bar{w}_h^i \chi_{Q_h^i}$  and  $\bar{z}_h := \sum_{i=1}^{N_h} \bar{z}_h^i \chi_{Q_h^i}$  admits an accumulation point  $(\bar{u}, \bar{w}, \bar{z}) \in U \times W \times H$  such that for a subsequence (for ease of notation denoted by the same symbol)*

$$\bar{u}_{hh} \rightarrow \bar{u} \text{ in } U \text{ and } \bar{w}_h \xrightarrow{*} \bar{w} \text{ in } W \text{ and } \bar{z}_h \rightarrow \bar{z} \text{ in } H.$$

Moreover, the point  $(\bar{u}, \bar{w}, \bar{z})$  minimizes (McC).

*Proof.* The sequence  $\{\bar{z}_h\}_h$  is bounded in  $H$  by assumption 2.11 4 and 5 and the McCormick inequalities. Consequently,  $\{\bar{u}_{hh}\}_h$  is also bounded in  $U$  because  $A$  is continuously invertible. In turn, after possibly passing to a subsequence,  $\bar{z}_h \rightarrow \bar{z}$  holds for some  $\bar{z} \in H$  and  $\bar{u}_{hh} \rightarrow \bar{u}$  in  $U$  for some  $\bar{u} \in U$  because of the compact embedding  $H \hookrightarrow^c U^*$  and the continuous invertibility of  $A$ . This argument also implies that the original state equation  $A\bar{u} + \bar{z} = f$  holds for the limit.

Let  $u_{hh}$  solve (2) for  $\bar{w}$ . Then  $(u_{hh}, P_h \bar{w}, (P_h \bar{w})(P_h u_{hh}))$  is feasible for (McC<sub>hh</sub>) by assumption 2.6 3. The optimality of the tuples  $(\bar{u}_{hh}, \bar{w}_h^1, \dots, \bar{w}_h^{N_h}, \bar{z}_h^1, \dots, \bar{z}_h^{N_h})$  for the problems (McC<sub>hh</sub>) and assumption 2.8 give

$$j(\bar{u}_{hh}, \bar{w}_h) + \alpha \text{TV}_h(\bar{w}_h) \leq j(u_{hh}, P_h \bar{w}) + \alpha \text{TV}_h(P_h \bar{w}) \leq j(u_{hh}, P_h \bar{w}) + \alpha \text{TV}(\bar{w}).$$

Because  $\{\bar{w}_h\}_h$ ,  $\{P_h \bar{w}\}_h$ , and in turn  $\{u_{hh}\}_h$  are bounded and  $j$  is Lipschitz and thus bounded on bounded sets, this implies that  $\{\text{TV}_h(\bar{w}_h)\}_h$  is bounded in  $[0, \infty)$ .

The properties of the set  $C$  imply that  $\{\|\bar{w}_h\|_H\}_h$  is bounded so that  $\{\|\bar{w}_h\|_{L^1}\}_h$  is bounded in  $\mathbb{R}$ , too. Thus, after passing to a subsequence (for ease of notation denoted by the same symbol), assumption 2.11 2 implies  $\bar{w}_h \rightarrow \bar{w}$  in  $L^1(\Omega)$  for some  $\bar{w} \in W$ .

The pointwise a.e. bounds on  $P_h \bar{w}_h$  and  $P_h \bar{u}_{hh}$  in the McCormick inequalities together with assumption 2.11 4 and 5 imply boundedness of the  $\bar{z}_h$  in  $H$ .

We are now concerned with feasibility of  $(\bar{u}, \bar{w}, \bar{z})$  for (McC). The inclusion  $\bar{w} \in C$  follows from the assumed properties of  $C$ . For the bounds on the state variable, we first observe

$$\|\bar{u} - P_h \bar{u}_{hh}\|_H \leq \|\bar{u} - P_h \bar{u}\|_H + \|P_h\|_{H,H} \|\bar{u} - \bar{u}_{hh}\|_H \rightarrow 0$$

from the triangle inequality, the nonexpansiveness of  $P_h$ , and Lebesgue's differentiation theorem, which requires assumption 2.11 1. Because, in addition,  $u_{\ell,h} \rightarrow u_\ell$  and  $u_{u,h} \rightarrow u_u$  in  $H$  hold, we can choose a further (subsubsub)sequence such that all three parts of the inequalities  $u_{\ell,h} \leq P_h \bar{u}_{hh} \leq u_{u,h}$  converge pointwise a.e. In turn, we obtain the pointwise a.e. inequalities  $u_\ell \leq \bar{u} \leq u_u$  for the limit.

It remains to show the feasibility of  $\bar{z}$  for the four pointwise McCormick inequalities in (McC) and the state equation. We prove the claim only for the first one since the others follow with an analogous argument. The feasibility of the solutions for (McC<sub>hh</sub>) gives

$$\bar{z}_h \geq u_{\ell,h} \bar{w}_h + \bar{u}_{hh} w_{\ell,h} - u_{\ell,h} w_{\ell,h} \quad \text{a.e. in } \Omega.$$

From what has been shown to this point, after passing to an appropriate subsequence, the left-hand side converges weakly to  $\bar{z}$ , and the right-hand side converges in  $H$  and thus also in  $L^1(\Omega)$ . We apply [lemma A.1](#) in order to obtain the desired inequality

$$\bar{z} \geq u_\ell \bar{w} + \bar{u} w_\ell - u_\ell w_\ell \quad \text{a.e. in } \Omega.$$

It remains to show that  $(\bar{u}, \bar{w}, \bar{z})$  minimizes [\(McC\)](#). We prove this claim by using a  $\Gamma$ -convergence-type argument. We need to show  $\liminf$ - and  $\limsup$ -inequalities for the objectives when the iterates are restricted to the feasible sets.

For the  $\liminf$ -inequality, let  $u_{hh} \rightarrow u$  in  $U$ ,  $w_h = \sum_{i=1}^{N_h} w_h^i \chi_{Q_h^i} \rightarrow w$  in  $H$ , and  $z_h = \sum_{i=1}^{N_h} z_h^i \chi_{Q_h^i} \rightarrow z$  in  $H$  with  $(u_{hh}, w_h^1, \dots, w_h^{N_h}, z_h^1, \dots, z_h^{N_h})$  being feasible for [\(McC<sub>hh</sub>\)](#). Then the continuity properties of  $j$  and the lower semi-continuity of the TV-seminorm give the  $\liminf$ -inequality

$$j(u, w) + \alpha \text{TV}(w) \leq \liminf_{h \searrow 0} j(u_{hh}, w_h) + \alpha \text{TV}(w_h).$$

For the  $\limsup$ -inequality, let  $(u, w, z)$  be feasible for [\(McC\)](#). We define  $w_h := P_h w$  and  $u_{hh}$  as the unique solution to [\(2\)](#), which exists by [assumption 2.6 2](#). Then the continuity properties of  $j$ , [assumption 2.11 6](#), and [assumption 2.11 3](#) imply

$$j(u, w) + \alpha \text{TV}(w) = \lim_{h \searrow 0} j(u_{hh}, w_h) + \alpha \text{TV}_h(w_h),$$

so that the  $\limsup$ -inequality holds true if we can prove that  $(u_{hh}, w_h, z_h)$  is feasible for [\(McC<sub>hh</sub>\)](#). This follows from the compatibility assumed in [assumption 2.6 3](#).  $\square$

As noted above, such a result requires a modification of the total variation functional dependent on  $h$  for dimensions larger than one. In the one-dimensional case it is possible to prove that  $P_h$  is nonexpansive with respect to the TV-seminorm, which in turn implies the desired  $\Gamma$ -convergence in this case without any modification of TV beforehand. This is shown below.

**Lemma 2.14.** *Let [assumption 2.6](#) hold. Let  $\Omega = (a, b)$  for some  $a < b$ . Let  $f \in \text{BV}(\Omega)$ . Then*

$$\text{TV}(P_h f) \leq \text{TV}(f).$$

*In particular,*

$$\text{TV} \circ P_h \text{ } \Gamma\text{-converges to } \text{TV}.$$

*Proof.* The first claim can be shown by considering the intervals  $Q_i \in \mathcal{Q}_h$  one by one and using the equivalence of TV to the pointwise variation when a *good representative* is chosen; see Definition 3.26, (3.24), and Theorem 3.28 in [\[1\]](#).

Let  $f^n \rightarrow f$  in  $L^1$ . Because  $P_h$  is nonexpansive with respect to the  $L^1$ -norm, we have  $\|P_h f - P_h f^n\|_{L^1} \leq \|f - f^n\|_{L^1}$ . Moreover,  $P_h f \rightarrow f$  in  $L^1(\Omega)$  holds by means of the Lebesgue differentiation theorem. In combination, we obtain  $P_h f^n \rightarrow f$  in  $L^1(\Omega)$ . Consequently, the  $\liminf$ -inequality follows from the lower semi-continuity of TV. The  $\limsup$ -inequality follows from the first claim by choosing the constant sequence that has  $f$  in every element.  $\square$



## 2.4 Optimality-based bound tightening

While [proposition 2.13](#) shows that solutions to the approximate relaxations  $(\text{McC}_{hh})$  of  $(\text{OCP})$  approximate solutions to the true relaxation  $(\text{McC})$ , it is of higher practical importance that the optimal objective values of  $(\text{McC}_h)$  and  $(\text{McC}_{hh})$  are as large as possible in practice. The only way to influence this is the choices of the bounds  $u_\ell^i$  and  $u_u^i$ , which should be as tight as possible in order to have the smallest possible feasible set and in turn the largest possible values for the optimal objectives of  $(\text{McC}_h)$  and  $(\text{McC}_{hh})$  such that (4) and (6) still hold.

Inspecting the arguments in the proof of [theorem 2.10](#) that lead to (6) ((7) in the proof), and similarly in [theorem 2.7](#), we observe that these arguments remain valid when the feasible set of  $(\text{McC}_{hh})$  is shrunk as long as [assumption 2.6 3](#) is preserved. Then the infimum  $m_{(\text{McC}_{hh})}$  increases when the feasible set is shrunk, while the approximation error bound that needs to be subtracted in order to obtain a valid bound on  $m_{(\text{OCP})}$  remains unaffected from this change. Consequently, the lower bound on  $m_{(\text{OCP})}$  is improved by this procedure.

Before proving this, we show the desired property that [assumption 2.6 3](#) is conserved when optimizing the bounds  $u_\ell^i, u_u^i$ .

**Lemma 2.15.** *Let [assumption 2.6](#) hold. Let  $h$  be fixed. Let  $i \in \{1, \dots, N_h\}$  be fixed. Then*

$$\tilde{u}_\ell^i / \tilde{u}_u^i := \min / \max \{ (P_h u)|_{Q_i} \mid (u, w, z_1, \dots, z_{N_h}) \text{ is feasible for } (\text{McC}_h) \}. \quad (8)$$

*are well defined.*

*Moreover,  $\tilde{u}_\ell^i \leq (P_h u)|_{Q_i} \leq \tilde{u}_u^i$  holds for all  $w \in C \cap [w_\ell, w_u]$  and  $u \in U$  that solve (2). In particular, [assumption 2.6](#) holds if  $u_\ell^i$  is replaced by  $\tilde{u}_\ell^i$  and  $u_u^i$  is replaced by  $\tilde{u}_u^i$ .*

*Proof.* We note that  $u \mapsto (P_h u)|_{Q_i}$  is a weakly continuous operation from  $H$  to  $\mathbb{R}$ . Moreover, using the arguments for the existence of solutions to  $(\text{McC}_h)$  in the proof of [theorem 2.7](#), we obtain that the feasible set of  $(\text{McC}_h)$  is nonempty and sequentially compact with respect to weak convergence of  $u$  in  $H$ , weak convergence  $w$  in  $H$ , and convergence of  $z$  in  $H$  (the last one has a finite-dimensional ansatz). Consequently, (8) has a solution, and  $\tilde{u}_\ell^i$  is well defined.

Inspecting [assumption 2.6](#) shows that it remains to prove that [assumption 2.6 3](#) stays valid when replacing  $u_\ell^i$  by  $\tilde{u}_\ell^i$ . For all  $w \in C$  and  $u \in U$  that solve  $Au + (P_h u)(P_h w) = f$ , we define  $z_i := (P_h u)|_{Q_h^i} (P_h w)|_{Q_h^i}$  for  $i \in \{1, \dots, N_h\}$ . Inspecting  $(\text{McC}_h)$ , we obtain that  $(u, w, z_1, \dots, z_{N_h})$  is feasible for  $(\text{McC}_h)$  by means of [assumption 2.6 3](#). Consequently, all  $w \in C$  and  $u \in U$  that solve  $Au + (P_h u)(P_h w) = f$  satisfy

$$\begin{aligned} \tilde{u}_\ell^i &= \min \{ (P_h u)|_{Q_i} \mid (u, w, z_1, \dots, z_{N_h}) \text{ is feasible for } (\text{McC}_h) \} \\ &\leq \inf \{ (P_h u)|_{Q_i} \mid (u, w) \in U \times C, Au + (P_h u)(P_h w) = f \}, \end{aligned}$$

and thus also  $\tilde{u}_\ell^i \leq (P_h u)|_{Q_i}$ , which proves the first claim.

The second well-definedness and claim follow analogously.  $\square$

A similar argument can be made for  $(\text{McC}_{hh})$ .



---

**Algorithm 1** Optimization-based bound tightening (OBBT) for  $(\text{McC}_h)$  (or  $(\text{McC}_{hh})$ )

---

**Input:** Feasible set  $\mathcal{F}^0$  of  $(\text{McC}_h)$  (or  $(\text{McC}_{hh})$ ).

```

1: for  $n = 1, \dots$  do
2:   Choose  $i \in \{1, \dots, N_h\}$ .
3:   Choose  $s \in \{\ell, u\}$ .
4:   if  $s = \ell$  then
5:      $\tilde{u}_\ell^i \leftarrow \min\{(P_h u_h)|_{Q_i} \mid (u_h, w_h, z_h) \in \mathcal{F}^{n-1}\}$ .
6:   else
7:      $\tilde{u}_u^i \leftarrow \max\{(P_h u_h)|_{Q_i} \mid (u_h, w_h, z_h) \in \mathcal{F}^{n-1}\}$ .
8:   end if
9:    $\mathcal{F}^n \leftarrow \mathcal{F}^{n-1}$  with  $u_s^i$  being replaced by  $\tilde{u}_s^i$ .
10: end for

```

---

**Lemma 2.16.** Let *assumption 2.8* hold. Let  $h$  be fixed. Let  $i \in \{1, \dots, N_h\}$  be fixed. Then

$$\tilde{u}_\ell^i / \tilde{u}_u^i := \min / \max\{(P_h u)|_{Q_i} \mid (u, w_1, \dots, w_{N_h}, z_1, \dots, z_{N_h}) \text{ is feasible for } (\text{McC}_{hh})\}.$$

are well defined.

Moreover,  $\tilde{u}_\ell^i \leq (P_h u)|_{Q_i} \leq \tilde{u}_u^i$  holds for all  $w \in C \cap [w_\ell, w_u]$  and  $u \in U$ . In particular, *assumption 2.8* holds if  $u_\ell^i$  is replaced by  $\tilde{u}_\ell^i$  and  $u_u^i$  is replaced by  $\tilde{u}_u^i$ .

*Proof.* The proof parallels the one of *lemma 2.15*. □

We can now state a bound-tightening procedure that successively computes new bounds in order to improve the approximate lower bounds (4) or (6) respectively in *algorithm 1*. Using *lemmas 2.15* and *2.16*, we can prove our main bound-tightening results for  $(\text{McC}_h)$  and  $(\text{McC}_{hh})$ .

**Theorem 2.17.** Let *assumption 2.6* be satisfied. Let  $C$  be a weakly sequentially compact subset of  $H$ . Then *algorithm 1* executed with the feasible set of  $(\text{McC}_h)$  as  $\mathcal{F}^0$  induces a sequence of optimization problems

$$\min_{u, w, z_1, \dots, z_{N_h}} j(u, w) + \alpha \text{TV}(w) \quad \text{s.t.} \quad (u, w, z_1, \dots, z_{N_h}) \in \mathcal{F}^n \quad (\text{McC}_h^n)$$

that satisfy (4) for all  $n \in \mathbb{N}$ . Moreover, the sequence of optimal objective values  $(m_{(\text{McC}_h^n)})_n$  is monotonically nondecreasing.

*Proof.* By assumption, the prerequisites of *lemma 2.15* are satisfied in the first iteration and thus hold inductively for all iterations  $n \in \mathbb{N}$ . This implies that (4) holds for all problems  $(\text{McC}_h^n)$  because *theorem 2.7* asserts (4) under *assumption 2.6*. Because the feasible set is always a subset of the previous one, the infima are monotonically nondecreasing. □

**Theorem 2.18.** Let *assumption 2.8* be satisfied. Then *algorithm 1* executed with the feasible set of  $(\text{McC}_{hh})$  as  $\mathcal{F}^0$  induces a sequence of optimization problems

$$\begin{aligned} \min_{\substack{u, w_1, \dots, w_{N_h}, \\ z_1, \dots, z_{N_h}}} j \left( u, \sum_{i=1}^{N_h} \chi_{Q_h^i} w_i \right) + \alpha \text{TV} \left( \sum_{i=1}^{N_h} \chi_{Q_h^i} w_i \right) \\ \text{s.t.} \quad (u, w_1, \dots, w_{N_h}, z_1, \dots, z_{N_h}) \in \mathcal{F}^n \end{aligned} \quad (\text{McC}_{hh}^n)$$

that satisfy (5) for all  $n \in \mathbb{N}$ . Moreover, the corresponding sequence of optimal objective values  $\{m_{(\text{McC}_{hh}^n)}\}_n$  is monotonically nondecreasing.

*Proof.* The proof parallels the one of theorem 2.18 with the applications of lemma 2.15 and theorem 2.7 being replaced by applications of lemma 2.16 and theorem 2.10.  $\square$

### 3 Application to an elliptic optimal control problem

We now apply our theoretical considerations from section 2 to an instance of (OCP) with an elliptic PDE that is defined on the one-dimensional domain  $\Omega = (0, 1)$ . To this end, let  $C := \{w \in L^\infty(0, 1) \mid w_\ell \leq w(x) \leq w_u \text{ a.e.}\}$  and  $f \in H$  for real constants  $-\pi^2 < w_\ell < w_u < \pi^2$  be fixed.

For these assumptions, we outline the PDE setting in section 3.1 and give ellipticity estimates as well as bounds from above on the norm with constants that are as sharp as we were able to obtain them from the literature. We provide the objective in section 3.2. We then verify Assumptions 2.1, 2.6, 2.8 and 2.11 in section 3.3. We prove an a priori estimate for  $\|u - u_h\|_U$  in the approximate lower bounds (4), (6) in section 3.4. We prove differentiability properties for the control-to-state operator of the PDE in section 3.5. In particular, we verify the assumptions imposed in [21].

#### 3.1 PDE setting

For  $w \in C$ , we are interested in solutions  $u \in U := H_0^1(0, 1)$  to the PDE in weak form

$$\int_0^1 \nabla u \nabla v + \int_0^1 w u v = \int_0^1 f v \quad \text{for all } v \in U. \quad (9)$$

The operator form is

$$-\Delta u + uw = f \text{ on } (0, 1) \text{ with } u(0) = u(1) = 0,$$

that is,  $A = -\Delta$  with homogeneous Dirichlet boundaries and  $N(u, w) = uw$ . We define the bilinear form  $B : U \times U \rightarrow \mathbb{R}$  by

$$B(u, v) := \int_0^1 \nabla u \nabla v + \int_0^1 w u v$$

and obtain

$$B(u, u) \geq \|\nabla u\|_H^2 + w_\ell \|u\|_H^2 \quad \text{for all } u \in U. \quad (10)$$

The existence of (unique) solutions to (9) follows from the Lax–Milgram lemma if

$$\|\nabla u\|_H^2 + w_\ell \|u\|_H^2 \geq c_1(w_\ell) \|\nabla u\|_H^2$$

holds for some  $c_1(w_\ell) > 0$ , which is true if  $w_\ell \geq 0$ . For the case  $w_\ell < 0$ , we recall embedding constants for the Sobolev inequalities for the one-dimensional domain  $(0, 1)$ :

$$\pi^2 \|u\|_H^2 \leq \|\nabla u\|_H^2 \text{ and} \quad (11)$$

$$4 \|u\|_{L^\infty}^2 \leq \|\nabla u\|_H^2. \quad (12)$$

The constant in (11) is optimal and, for example, given in (1.2) in [40] (see [31] for a proof). The constant in (12) can be found in (13) in [35] (choose  $a = \infty$ ,  $b = 2$ ,  $s = 1$  therein). By (11), we obtain that  $0 < c_1(w_\ell) := 1 + w_\ell \pi^{-2}$  because (by our choice above)

$$w_\ell > -\pi^2. \quad (13)$$

The Lax–Milgram lemma also yields that the solution to (9) satisfies

$$\|\nabla u\|_H \leq \frac{1}{c_1(w_\ell)} \|f\|_H \text{ and} \quad (14)$$

$$\|u\|_{L^\infty} \leq \frac{1}{2c_1(w_\ell)} \|f\|_H, \quad (15)$$

where the second inequality follows from (12). We observe that  $u$  is also the weak solution of the PDE  $\int_0^1 \nabla u \nabla v = \int_0^1 g v$  for all  $v \in U$  with the choice  $g = f - wu$ . Because  $g \in H$ , this gives the improved regularity  $u \in H^2(0, 1)$ ; see, for example, Theorem 9.53 in [33].

### 3.2 Objective

We consider a tracking-type objective functional  $j : U \times W \rightarrow \mathbb{R}$  that is defined for  $(u, w) \in U \times W$  as

$$j(u, w) := \frac{1}{2} \|u - u_d\|_H^2. \quad (16)$$

Since our specific choice for  $j$  does not depend on  $w$ , we will abbreviate it as  $j(u)$  in the remainder. If the norm of the input is bounded by a constant  $r_u$ , we obtain that a feasible Lipschitz constant of  $j$  on the ball  $\overline{B_{r_u}(0)}$  is given by  $L_u := r_u + \|u_d\|_H$ . Clearly,  $L_w = 0$ .

### 3.3 Verification of assumptions

We now verify Assumptions 2.1, 2.6, 2.8 and 2.11 one by one.

**Assumption 2.1** Let  $c_2(w_\ell, w_u) := 1 - \max\{|w_\ell|, |w_u|\} \pi^{-2}$ . We choose  $u_u = \frac{1}{2c_2(w_\ell, w_u)} \|f\|_H$  and  $u_\ell = -\frac{1}{2c_2(w_\ell, w_u)} \|f\|_H$ . Then  $c_2(w_\ell, w_u) \leq c_1(w_\ell)$  gives  $|u_u| = |u_\ell| \geq \frac{1}{2c_1(w_\ell)} \|f\|_H$  so that together with the bounds  $w_\ell, w_u$ , we obtain that assumption 2.1 is satisfied because of (15). (We could of course choose the tighter bounds  $u_u = \frac{1}{2c_1(w_\ell)} \|f\|_H$  and  $u_\ell = -\frac{1}{2c_1(w_\ell)} \|f\|_H$  here, but the relaxed bounds using  $c_2(w_\ell, w_u)$  instead of  $c_1(w_\ell)$  will be used to assert the other assumptions below.)

**Assumption 2.6** We consider a sequence of uniform partitions  $\{\mathcal{Q}_{h_n}\}_{n \in \mathbb{N}}$  of the domain  $(0, 1)$  into  $2^{n+1}$  intervals with  $h_n = 2^{-1-n}$  for  $n \in \mathbb{N}$  and  $N_{h_n} = 2^{n+1}$ , which gives assumption 2.6 1.

Let  $n \in \mathbb{N}$ , and abbreviate  $h := h_n$ . For each interval  $I \in \mathcal{Q}_h$ , the projection  $P_I$  is defined as  $P_I g := \chi_I \frac{1}{h} \int_I g$  for  $g \in L^1(0, 1)$ . Clearly,  $P_I$  is nonexpansive in  $L^1(0, 1)$ ,  $H$ , and  $L^\infty(0, 1)$ ; that is,

$$\|P_I g\|_{L^\infty} \leq \|g\|_{L^\infty} \text{ and } \|P_I g\|_H \leq \|g\|_H.$$

The linear projection operator  $P_h$  from (1) satisfies  $P_h(g) = \sum_{I \in \mathcal{Q}_h} P_I g$  and

$$\|g - P_h g\|_H \leq \pi h \|\nabla g\|_H, \quad (17)$$

where the constant  $\pi$  is due to the application of the one-dimensional Poincaré inequality into the interpolation estimate for piecewise constant functions; see, for example, §3.1 in [14]. From (12.24) in Theorem 12.26 in [23], we also obtain

$$\|w - P_h w\|_{L^1} \leq h \text{TV}(w), \quad (18)$$

which we have already used in the proof of [theorem 2.10](#).

For arbitrary, fixed  $w$ , we define the bilinear form  $B_h : U \times U \rightarrow \mathbb{R}$  as

$$B_h(u, v) := \int_0^1 \nabla u \nabla v + \int_0^1 (P_h w)(P_h u) v$$

for  $u, v \in U$  and obtain the coercivity

$$B_h(u, u) \geq \|\nabla u\|_H^2 - \max\{|w_\ell|, |w_u|\} \|u\|_H \|P_h(u)\|_H \geq \|\nabla u\|_H^2 - \max\{|w_\ell|, |w_u|\} \|u\|_H^2.$$

Consequently, the existence of unique solutions to the PDE

$$\int_0^1 \nabla u \nabla v + \int_0^1 (P_h w)(P_h u) v = \int_0^1 f v \quad \text{for all } v \in U. \quad (19)$$

follows from the Lax–Milgram lemma with analogous estimates to (14) and (15), where  $c_1(w_\ell)$  is replaced by  $c_2(w_\ell, w_u) = 1 - \max\{|w_\ell|, |w_u|\} \pi^{-2}$  because  $0 < c_2(w_\ell, w_u)$ . Specifically, we have

$$\|\nabla u\|_H \leq \frac{1}{c_2(w_\ell, w_u)} \|f\|_H \text{ and} \quad (20)$$

$$\|u\|_{L^\infty} \leq \frac{1}{2c_2(w_\ell, w_u)} \|f\|_H. \quad (21)$$

This proves [assumption 2.6 2](#). As for (9), we observe that  $u$  solves  $\int_0^1 \nabla u \nabla v = \int_0^1 g v$  for all  $v \in U$  with the choice  $g = f - (P_h w)(P_h u)$ . Because  $g \in H$ , this gives the improved regularity  $u \in H^2(0, 1)$ ; see, for example, Theorem 9.53 in [33].

Because  $P_h$  is nonexpansive with respect to  $L^\infty(0, 1)$ , we can reuse the bounds  $w_\ell$  and  $w_u$  for  $w_\ell^i$  and  $w_u^i$  for all  $i \in \{1, \dots, N_h\}$ . With the same argument as for [assumption 2.1](#), we choose  $u_u^i = \frac{1}{2c_2(w_\ell, w_u)} \|f\|_H$  and  $u_\ell^i = -\frac{1}{2c_2(w_\ell, w_u)} \|f\|_H$  for all  $i \in \{1, \dots, N_h\}$ . In combination, [assumption 2.6 3](#) is satisfied.

**Assumption 2.8** In our one-dimensional setting, [lemma 2.14](#) implies that [assumption 2.8](#) holds with the choice  $\text{TV}_h := \text{TV}$ .

**Assumptions 2.8 and 2.11** Assumption 2.11 1 is satisfied because of the uniform discretization of the domain into intervals. Assumption 2.11 2 follows from the properties of the total variation seminorm with the choice  $\text{TV}_h := \text{TV}$ ; see Theorem 3.23 in [1]. Assumption 2.11 3 follows from lemma 2.14. Assumption 2.11 4 follows because we have chosen  $u_\ell^i = u_\ell$  and  $u_u^i = u_u$  for all  $i$  and  $h$ . Assumption 2.11 5 follows because we have chosen  $w_\ell^i = w_\ell$  and  $w_u^i = w_u$  for all  $i$  and  $h$ . Assumption 2.11 6 follows from lemma 3.1 that is proven in section 3.4 and a bootstrapping argument / the continuous invertibility of  $A$ .

### 3.4 Estimate on $\|u - u_h\|_H$ and a priori estimates on $m(\text{OCP})$

For our example PDE setting and the tracking-type objective in  $H$ , we are able to obtain lower bounds on (4) and (6) that are quadratic in  $h$  because we can improve over  $\|u - u_h\|_U$  when considering  $\|\cdot\|_H$  instead of  $\|\cdot\|_U$ . We first show the necessary estimates on  $\|u - u_h\|_H$  and subsequently prove the lower bounds on  $m(\text{OCP})$ .

**Lemma 3.1.** *Let assumption 2.6 hold. Let a mesh size  $h$  be fixed. Let  $u$  solve (9) and  $u_h$  solve (19) for the same fixed  $w \in W$ . Then the estimates*

$$\|u - u_h\|_H \leq C_{3/2}^a(w_\ell, w_u, f, w)h^{\frac{3}{2}} + C_{3/2}^b(w_\ell, w_u, f)h^2$$

and

$$\|u - u_h\|_H \leq C_2(w_\ell, w_u, f, w, u_h)h^2 \quad (22)$$

hold for constants  $C_{3/2}^a(w_\ell, w_u, f, w)$ ,  $C_{3/2}^b(w_\ell, w_u, f)$ ,  $C_2(w_\ell, w_u, f, w, u_h) > 0$ .

*Proof.* We observe that the local averaging of  $w$  and  $u$  in the lower-order term in (19) satisfies Galerkin-type orthogonality properties. Specifically, for given  $\phi, \psi, \theta \in H$ , we have

$$\int_0^1 (\phi - P_h \phi)(P_h \psi)(P_h \theta) = \sum_{i=1}^{N_h} \left( \frac{1}{|Q_h^i|} \int_{Q_h^i} \psi \right) \left( \frac{1}{|Q_h^i|} \int_{Q_h^i} \theta \right) \underbrace{\int_{Q_h^i} \phi - P_h \phi}_{=0} = 0. \quad (23)$$

Thus, we use an Aubin–Nitsche duality argument in order to obtain an estimate on the right-hand side of

$$\|u - u_h\|_H = \sup_{g \neq 0} \frac{(u - u_h, g)_H}{\|g\|_H}. \quad (24)$$

To this end, we consider the solution  $p \in U$  to the adjoint PDE

$$\int_0^1 \nabla p \nabla v + \int_0^1 p w v = \int_0^1 g v \quad \text{for all } v \in U \quad (25)$$

for arbitrary  $g \in H$ , which has the same properties as (9) (the only difference is the source term  $g$  instead of  $f$ ). Moreover, we observe with the choice  $v = p$  in (9) and (19) and the

insertion of a suitable zero that

$$\int_0^1 \nabla(u - u_h) \nabla p = - \int_0^1 (u - P_h u_h) w p - \int_0^1 (P_h u_h)(w - P_h w) p = 0. \quad (26)$$

We choose  $v = u - u_h$  in (25) and apply (26) so that we obtain

$$\begin{aligned} (u - u_h, g)_H &= \int_0^1 \nabla p \nabla(u - u_h) + \int_0^1 p w (u - u_h) \\ &= \int_0^1 (u - u_h) w p - \int_0^1 (u - P_h u_h) w p - \int_0^1 (P_h u_h)(w - P_h w) p. \end{aligned}$$

Combining the first two terms on the right-hand side and subtracting  $0 = \int_0^1 (P_h u_h)(w - P_h w)(P_h p)$ , which holds because of (25), we obtain

$$(u - u_h, g)_H = \int_0^1 (P_h u_h - u_h) w p - \int_0^1 (P_h u_h)(w - P_h w)(p - P_h p).$$

Inserting another zero and applying  $0 = \int_0^1 (P_h u_h)(w - P_h w)(P_h p)$  again give

$$\begin{aligned} (u - u_h, g)_H &= \int_0^1 (P_h u_h - u_h)(w - P_h w) p + \int_0^1 (P_h u_h - u_h)(P_h w)(p - P_h p) \\ &\quad - \int_0^1 (P_h u_h)(w - P_h w)(p - P_h p). \end{aligned}$$

Now, we estimate the terms on the right-hand side and obtain with the same estimates that have been used in the preceding subsections the following estimates:

$$\begin{aligned} &\int_0^1 (P_h u_h - u_h)(w - P_h w) p \\ &\leq h\pi \|\nabla u_h\|_H \|w - P_h w\|_H \|p\|_{L^\infty} \quad \text{Hölder, (17)} \\ &\leq \frac{\pi}{c_2(w_\ell, w_u)} \|f\|_H |w_u - w_\ell|^{\frac{1}{2}} \|w - P_h w\|_{L^1}^{\frac{1}{2}} \frac{1}{2c_1(w_\ell)} \|g\|_H \quad \text{Hölder, (15), (20)} \\ &\leq \frac{\pi |w_u - w_\ell|^{\frac{1}{2}}}{2c_1(w_\ell)c_2(w_\ell, w_u)} \text{TV}(w) \|f\|_H h^{\frac{3}{2}} \|g\|_H, \quad (18) \end{aligned}$$

$$\int_0^1 (P_h u_h - u_h)(P_h w)(p - P_h p) \leq \frac{\pi^2 \max\{|w_u|, |w_\ell|\}}{c_1(w_\ell)c_2(w_\ell, w_u)} \|f\|_H h^2 \|g\|_H, \quad \text{Hölder, (15), (17), (20)}$$

and

$$\begin{aligned} &\int_0^1 (P_h u_h)(w - P_h w)(p - P_h p) \\ &\leq \frac{\pi |w_u - w_\ell|^{\frac{1}{2}}}{2c_1(w_\ell)c_2(w_\ell, w_u)} \text{TV}(w) \|f\|_H h^{\frac{3}{2}} \|g\|_H. \quad \text{Hölder, (15), (17), (18), (20)} \end{aligned}$$

Combining these considerations, we obtain from (24)

$$\|u - u_h\|_H \leq \underbrace{\frac{\pi|w_u - w_\ell|^{\frac{1}{2}}}{c_1(w_\ell)c_2(w_\ell, w_u)} \text{TV}(w)\|f\|_H}_{=: C_{3/2}^a(w_\ell, w_u, f, w)} h^{\frac{3}{2}} + \underbrace{\frac{\pi^2 \max\{|w_u|, |w_\ell|\}}{c_1(w_\ell)c_2(w_\ell, w_u)} \|f\|_H}_{=: C_{3/2}^b(w_\ell, w_u, f)} h^2,$$

which is the first of the claimed estimates.

We next prove the second estimate. Because we know that  $u, u_h, p$  are also  $H^2(0, 1)$ -functions (see again Theorem 9.53 in [33]), their derivatives are uniformly bounded, and we can estimate them pointwise. Let  $(\eta_\varepsilon)_\varepsilon$  be a family of standard mollifiers. For  $y \in (0, 1)$ , we define the antiderivative  $N_\varepsilon^y(x) := \int_{-\infty}^x \eta_\varepsilon(z - y) dz$ . We test (19) with  $N_\varepsilon^y$  and deduce

$$\begin{aligned} \int_0^1 \nabla u_h(x) \eta_\varepsilon(x - y) dx &= \int_0^1 f N_\varepsilon^y - \int_0^1 (P_h u_h)(P_h w) N_\varepsilon^y \\ &\leq \int_0^1 |f(x) - (P_h u_h)(x)(P_h w)(x)| \int_{-\infty}^\infty \eta_\varepsilon(x - y) dy dx \\ &= \|f - (P_h u_h)(P_h w)\|_{L^1}. \end{aligned}$$

Driving  $\varepsilon \searrow 0$  and supremizing over the left-hand side, we obtain

$$\|\nabla u_h\|_{L^\infty} \leq \|f - (P_h u_h)(P_h w)\|_{L^1}.$$

An analogous argument and applications of the triangle inequality, Hölder's inequality, (11), and (14) give

$$\|\nabla p\|_{L^\infty} \leq \|g - pw\|_{L^1} \leq \left(1 + \frac{\max\{|w_\ell|, |w_u|\}}{\pi c_1(w_\ell)}\right) \|g\|_H.$$

We also obtain for  $\phi \in W^{1,\infty}(0, 1)$  that

$$\|\phi - P_h \phi\|_{L^\infty} \leq \frac{1}{2} h \|\nabla \phi\|_{L^\infty},$$

where  $\frac{1}{2}$  is the Wirtinger–Sobolev constant that can be found in (13) in [35] (choose  $a = \infty$ ,  $b = \infty$ ,  $s = 1$  therein).

Then, using Hölder's inequality with different exponents, we can estimate the two  $h^{\frac{3}{2}}$ -terms similarly as above but we replace the Hölder conjugates  $(2, 2)$  by  $(1, \infty)$  in order to obtain  $h^2$ -terms, specifically

$$\begin{aligned} \int_0^1 (P_h u_h - u_h)(w - P_h w) p &\leq \frac{1}{2} h \|\nabla u_h\|_{L^\infty} \|w - P_h w\|_{L^1} \|p\|_{L^\infty} \\ &\leq \frac{1}{4c_1(w_\ell)c_2(w_\ell, w_u)} \text{TV}(w) \|f - (P_h u_h)(P_h w)\|_{L^1} h^2 \|g\|_H \end{aligned}$$

and

$$\int_0^1 (P_h u_h)(w - P_h w)(p - P_h p) \leq \frac{c_1(w_\ell) + \max\{|w_\ell|, |w_u|\} \pi^{-1}}{4c_1(w_\ell)c_2(w_\ell, w_u)} \text{TV}(w) \|f\|_H h^2 \|g\|_H.$$

In summary, we obtain

$$\|u - u_h\|_H \leq C_2(w_\ell, w_u, f, w, u_h)h^2$$

with

$$C_2(w_\ell, w_u, f, w, u_h) := \frac{1}{4c_1(w_\ell)c_2(w_\ell, w_u)} \left( 4\pi^2 \max\{|w_u|, |w_\ell|\} \|f\|_H + \right. \\ \left. \text{TV}(w) \|f - (P_h u_h)(P_h w)\|_{L^1} + (c_1(w_\ell) + \max\{|w_\ell|, |w_u|\} \pi^{-1}) \text{TV}(w) \|f\|_H \right), \quad (27)$$

which is the second estimate.  $\square$

We now provide a lower bound on  $m_{(\text{OCP})}$  in terms of  $h^2$  for our guiding example.

**Lemma 3.2.** *Let  $j_0 \geq 0$  be a lower bound on  $J(S(\bar{w}))$  if  $\bar{w}$  minimizes (OCP). Let  $\hat{w} \in W$  be feasible for (OCP). Then*

$$m_{(\text{OCP})} \geq m_{(\text{McC}_{hh})} - c_{quad}(w_\ell, w_u, f, \hat{w}, j_0, u_{\ell,h}, u_{u,h})h^2. \quad (28)$$

*Proof.* For the objective  $j(u) = \frac{1}{2}\|u - u_d\|_H^2$ , we obtain  $\nabla_u j(u) = u - u_d$ . Since we have pointwise lower and upper bounds  $u_\ell$  and  $u_u$  on  $u$ , for example, from (21), we can overestimate the Lipschitz constant of  $j$  on the feasible set with respect to the  $L^2$ -norm by setting  $\tilde{d}_u(x) := \max\{|u_u(x) - u_d(x)|, |u_d(x) - u_\ell(x)|\}$  for a.a.  $x \in (0, 1)$ . Then the Lipschitz constant can be estimated as  $L_u \leq \|\tilde{d}_u\|_H$ .

Using (22) from lemma 3.1, we obtain

$$m_{(\text{OCP})} \geq m_{(\text{McC}_{hh})} - \|\tilde{d}_u\|_H C_2(w_\ell, w_u, f, \bar{w}, \bar{u}_h)h^2.$$

Inspecting  $C_2(w_\ell, w_u, f, \bar{w}, \bar{u}_h)$  in lemma 3.1, we observe that we can overestimate  $\text{TV}(\bar{w}) \leq \frac{j(S(\bar{w})) + \alpha \text{TV}(\bar{w}) - j_0}{\alpha}$ . Moreover, using available bounds on  $P_h \bar{u}_h$ , we can overestimate  $|f(x) - (P_h \bar{u}_h)(x)(P_h \bar{w})(x)|$  pointwise by setting

$$\tilde{d}_f(x) := \max\{|f(x) - u_\ell^i w_\ell|, |f(x) - u_\ell^i w_u|, |f(x) - u_u^i w_\ell|, |f(x) - u_u^i w_u|\}$$

for a.a.  $x \in Q_i$  for all  $Q_i \in \mathcal{Q}_h$ . Thus we obtain the constant  $c_{quad}(w_\ell, w_u, f, \hat{w}, j_0, u_{\ell,h}, u_{u,h})$  by replacing  $\|f - (P_h \bar{u}_h)(P_h \bar{w})\|_{L^1}$  by  $\|\tilde{d}_f\|_{L^1}$  and  $\text{TV}(\bar{w})$  by  $\frac{j(S(\bar{w})) + \alpha \text{TV}(\bar{w}) - j_0}{\alpha}$  in (27) and multiplying the resulting constant by  $\|\tilde{d}_u\|_H$ .  $\square$

*Remark 3.3.* We note that the estimate  $L_u \leq \|\tilde{d}_u\|_H$  can be sharpened by using a posteriori information from the approximate McCormick envelopes in order to bootstrap improved pointwise bounds  $u_\ell$  and  $u_u$  on  $u$ .

### 3.5 Properties of the control-to-state operator for (19)

We denote the nonlinear control-to-state operator that maps  $w \in C$  to the solution of the PDE (19) with the locally averaged state in the bilinearity as  $S : C \rightarrow U$ . It is well defined and uniformly bounded due to the considerations in sections 3.1 and 3.3. We briefly state Lipschitz continuity and differentiability properties of the control-to-state operator so that we verify the assumptions in [21]. Consequently, the algorithmic strategy



from [21, 27] can be used to obtain stationary feasible (primal points) when additional integrality restrictions are imposed on  $w$  in (OCP) while the analysis carried out above allows us to obtain valid (approximate) lower bounds. While the arguments are standard by following, for example, the considerations on control-to-state operators in [39], we provide these proofs here because we were not able to find good references for the specific Lipschitz, embedding, ... constants of the example in this article. We believe, however, that they help make informed assessments of the possible quality and performance of our algorithmic approaches, although some of them may not be tight.

**Lemma 3.4.**  *$S$  is Lipschitz continuous on the set  $C$  as a function  $L^1(0, 1) \rightarrow U$ .*

*Proof.* Let  $w_1, w_2 \in C$ . Let  $u_1 = S(w_1)$ ,  $u_2 = S(w_2)$ . We subtract the weak two formulations, insert a suitable zero, and deduce by means of Hölder's inequality the triangle inequality, (11), (12), and the nonexpansiveness of  $P_h$ :

$$\begin{aligned} \|\nabla(u_1 - u_2)\|_H^2 &= ((P_h w_1)(P_h u_1) - (P_h w_2)(P_h u_1), u_1 - u_2)_H \\ &\quad + ((P_h w_2)(P_h u_1) - (P_h w_2)(P_h u_2), u_1 - u_2)_H \\ &\leq \|u_1 - u_2\|_{L^\infty} \|u_1\|_{L^\infty} \|w_1 - w_2\|_{L^1} + \max\{|w_\ell|, |w_u|\} \|u_1 - u_2\|_H^2 \\ &\leq \left( \frac{\|f\|_H}{4c_2(w_\ell, w_u)} \|\nabla(u_1 - u_2)\|_H \|w_1 - w_2\|_{L^1} + \frac{\max\{|w_\ell|, |w_u|\}}{\pi^2} \|\nabla(u_1 - u_2)\|_H^2 \right). \end{aligned}$$

An equivalent reformulation of this estimate gives

$$\|\nabla(u_1 - u_2)\|_H \leq \underbrace{\frac{\|f\|_H}{4c_2(w_\ell, w_u)^2}}_{=: L_S} \|w_1 - w_2\|_{L^1}. \quad (29)$$

□

Let  $u$  solve (19) for  $w \in C$ . For a given direction  $s \in L^1(0, 1)$  so that  $w + s \in C$  and with  $u = S(w)$ , we denote the (weak) solution to

$$\int_0^1 \nabla q \nabla v + \int_0^1 (P_h q)(P_h w) v = - \int_0^1 (P_h u)(P_h s) v \quad \text{for all } v \in U$$

by  $q$ . We obtain that  $q \in U$  is uniquely defined because  $s \in C - w$  implies  $L^\infty(0, 1)$ -bounds on  $s$ , namely,  $w_\ell - w_u \leq s \leq w_u - w_\ell$ . This gives  $\|(P_h s)(P_h u)\|_H \leq \|w_u - w_\ell\| \|u\|_H$  and a similar analysis as for (19) applies.

**Lemma 3.5.** *The operator  $S : C \rightarrow U$  is Fréchet differentiable with respect to  $L^1(0, 1)$  in  $C$ . Let  $q$  be as above for a given  $s$  such that  $w + s \in C$ . Then  $S'(w)s = q$ . For a fixed  $s \in C$ , the mapping  $w \mapsto S'(w)s$  is Lipschitz continuous in  $C$  with respect to  $L^1(0, 1)$ .*

*Proof.* Clearly, the mapping  $s \mapsto q$  is a bounded linear operator. We need to show

$$\lim_{\|s\|_{L^1} \xrightarrow{s \in C} 0} \frac{\|S(w + s) - S(w) - q\|_U}{\|s\|_{L^1}} \rightarrow 0.$$

Let  $u_s = S(w + s)$ ,  $u = S(w)$ . Let  $d := u_s - u - q$ . Then, we deduce with the considerations

above and [lemma 3.4](#) and the Lipschitz constant  $L_S$  from [\(29\)](#) that

$$\begin{aligned}\|\nabla d\|_H^2 &= (P_h(u - u_s)(P_h s), d)_H - ((P_h d)(P_h w), d)_H \\ &\leq \frac{L_S}{2} \|\nabla d\|_H \|s\|_{L^1} \|s\|_{L^1} + \frac{\max\{|w_\ell|, |w_u|\}}{\pi^2} \|\nabla d\|_H^2,\end{aligned}$$

which yields

$$\|\nabla d\|_H \leq \frac{L_S}{2c_2(w_\ell, w_u)} \|s\|_{L^1}^2.$$

Since  $\|\cdot\|_U$  is equivalent to  $\|\nabla \cdot\|_H$  with the embedding [\(11\)](#), this proves the claim.

For the Lipschitz continuity, let  $w_1, w_2 \in C$ ,  $u_1 = S(w_1)$ ,  $u_2 = S(w_2)$ , and  $q_1 = S'(w_1)s$ ,  $q = S'(w_2)s$ . Then, we obtain with  $r = q_1 - q_2$  that

$$\begin{aligned}\|\nabla r\|_H^2 &= (P_h(u_2 - u_1)(P_h s), r)_H - ((P_h r)(P_h w), r)_H - ((P_h q_2)P_h(w_2 - w_1), r)_H \\ &\leq \frac{L_S}{2} \|w_1 - w_2\|_{L^1} \|s\|_{L^1} \|\nabla r\|_H + \frac{\max\{|w_\ell|, |w_u|\}}{\pi^2} \|\nabla r\|_H^2 + \frac{1}{2} \|\nabla q_2\|_H \|w_1 - w_2\|_{L^1} \|\nabla r\|_H.\end{aligned}$$

With the arguments from the preceding subsections, we obtain

$$\|\nabla q_2\|_H \leq \frac{1}{c_2(w_\ell, w_u)} \|s\|_{L^\infty} \|u_2\|_H \leq \frac{|w_u - w_\ell|}{\pi c_2(w_\ell, w_u)^2} \|f\|_H,$$

so that we obtain the estimate

$$\|\nabla r\|_H \leq \underbrace{\frac{|w_u - w_\ell|}{2c_2(w_\ell, w_u)} \left( L_S + \frac{\|f\|_H}{\pi c_2(w_\ell, w_u)^2} \right)}_{=: L_{S'}} \|w_1 - w_2\|_{L^1}.$$

□

For given directions  $\phi, \psi \in L^1(0, 1)$  so that  $w + \phi, w + \psi \in C$ , we denote the (weak) solution to

$$\begin{aligned}\int_0^1 \nabla \xi \nabla v + \int_0^1 (P_h \xi)(P_h w) v &= \\ - \int_0^1 P_h(S'(w)\phi)(P_h \psi) v - \int_0^1 P_h(S'(w)\psi)(P_h \phi) v &\quad \text{for all } v \in U\end{aligned}\quad (30)$$

by  $\xi$ . As above, we obtain that  $\xi \in U$  is uniquely defined because  $\phi, \psi \in C - w$  imply  $L^\infty(0, 1)$ -bounds on  $\phi, \psi$ , that is,  $w_\ell - w_u \leq \phi \leq w_u - w_\ell$  and  $w_\ell - w_u \leq \psi \leq w_u - w_\ell$ . This implies bounds on the lower-order order terms and a similar analysis as for [\(19\)](#) applies.

**Proposition 3.6.** *Let  $\phi \in C - w$  be given. The operator  $w \mapsto S'(w)\phi$  is continuously Fréchet differentiable with respect to  $L^1(0, 1)$  in  $C$ . Its derivative in direction  $\psi \in C - w$  is given by  $S''(w)[\psi, \phi] = \xi$ , where  $\xi$  is the solution to [\(30\)](#) above. Under these assumptions, it holds that*

$$\|S''(w)[\psi, \phi]\|_H \leq \kappa \|\psi\|_{L^1} \|\phi\|_{L^1}$$

for some  $\kappa > 0$ .

*Proof.* For  $\phi \in C - w$  and  $\psi \in C - w$ , let  $q := S'(w)\phi$ ,  $q_\psi := S'(w + \psi)\phi$ ,  $\tilde{q}_\psi := S'(w)\psi$ ,

$u_\psi = S(w + \psi)$ ,  $u = S(w)$ ,  $\xi$  be the solution to (30). Let  $r := q_\psi - q - \xi$ . Then, we obtain from Hölder's inequality and the embedding constants (11), (12)

$$\begin{aligned}\|\nabla r\|_H^2 &= (P_h(u + \tilde{q}_\psi - u_\psi)(P_h\phi), r)_H + (P_h(q_\psi - q)(P_h\psi), r)_H - ((P_h r)(P_h w), r)_H \\ &\leq \frac{1}{4}\|\nabla(u_\psi - u - \tilde{q}_\psi)\|_H\|\phi\|_{L^1}\|\nabla r\|_H + \frac{1}{4}\|\nabla(q_\psi - q)\|_H\|\psi\|_{L^1}\|\nabla r\|_H \\ &\quad + \frac{\max\{|w_\ell|, |w_u|\}}{\pi^2}\|\nabla r\|_H^2.\end{aligned}$$

Using the constants defined above and the estimates from the proof of lemma 3.5, we obtain

$$4c_2(w_\ell, w_u)\|\nabla r\|_H \leq \frac{L_S}{2c_2(w_\ell, w_u)}\|\psi\|_{L^1}^2\|\phi\|_{L^1} + L_{S'}\|\psi\|_{L^1}^2.$$

Consequently, we obtain

$$\lim_{\|\psi\|_{L^1} \xrightarrow{\psi \in C} 0} \frac{\|S(w + \psi) + S(w) - \xi\|_U}{\|\psi\|_{L^1}} \rightarrow 0,$$

which proves the desired differentiability.

Testing (30) with  $\xi = S''(w)[\psi, \phi]$  and using (11) and similar computations as in the previous arguments, we obtain

$$\|\nabla \xi\|_H^2 \leq \frac{\max\{|w_\ell|, |w_u|\}}{\pi^2}\|\nabla \xi\|_H^2 + \frac{\|f\|_H}{2c_2(w_\ell, w_u)^2}\|\psi\|_{L^1}\|\phi\|_{L^1}\|\nabla \xi\|_H$$

and in turn

$$\|\xi\|_H \leq \underbrace{\frac{\|f\|_H}{2\pi c_2(w_\ell, w_u)^3}}_{=\kappa} \|\psi\|_{L^1}\|\phi\|_{L^1}.$$

The (Lipschitz) continuity of  $w \mapsto S''(w)[\phi, \psi]$  can be shown with arguments similar to but more lengthy than that for the Lipschitz continuity of the mapping  $w \mapsto S'(w)s$  in lemma 3.5.  $\square$

## 4 Computational experiments

We start from the example given in section 3 and provide computational experiments that will serve several purposes. We compare the approximate lower bounds obtained with approximate McCormick envelopes before and after applying the OBBT procedure to a feasible primal point obtained with a gradient-based optimization of (OCP) and a primal point obtained with the SLIP algorithm from [21, 27] in the presence of an additional integrality restriction on  $w$ . We also compare the bounds with lower bounds on (OCP) that can be obtained with much less effort than the OBBT procedure.

We analyze how the bounds we obtain from (McC<sub>hh</sub>) behave when the partitions of the domain assumed in assumption 2.6 1 are uniformly refined, thereby bringing the locally averaged McCormick envelopes closer to a pointwise limit. Moreover, we apply the OBBT procedure algorithm 1 to assess whether and how much the OBBT procedure tightens the bounds for a prescribed termination tolerance as well as to assess its computational effort.

This section is organized as follows. We describe our experiments in section 4.1. In section 4.2 we describe how we approximate (9) and (19) with a very fine discretization

and use the resulting discretized equations as state equations of (OCP), (McC<sub>h</sub>), and (McC<sub>hh</sub>). We then give details on the practical implementation of the OBBT procedure [algorithm 1](#) in [section 4.3](#). The results are provided in [section 4.4](#).

#### 4.1 Experiment description

We consider the instance of (OCP) from [section 3](#) with the following choices of the parameters and fixed inputs. The source term  $f$  of the PDEs (9) and (19) is chosen as  $f(x) := 6$  for all  $x \in [0, 1]$ . Regarding the bounds, we set  $w_\ell := -4$ ,  $w_u := 4$ . For the penalty parameter, we choose  $\alpha := 2.5 \cdot 10^{-4}$ . For the function  $u_d$  in the tracking-type objective (see (16)), we choose

$$\begin{aligned} u_d(x) := & 3 \left( 1.5x(1-x)\chi_{[0,0.25]}(x) + 1.5x(1-x)\chi_{[0.75,1]} \right. \\ & + (0.28125 + 3(x-0.25))\chi_{(0.25,0.4]}(x) \\ & \left. + (0.73125 - 3(x-0.6))\chi_{[0.6,0.75)}(x) + 2\chi_{(0.4,0.6)}(x) \right). \end{aligned}$$

Our main object of interest is to assess the quality of the approximate McCormick envelopes, the effect of the OBBT procedure [algorithm 1](#). To this end, we perform the following computations.

- We execute a local gradient-based NLP solver (L-BFGS-B) using Scipy's implementation [41] in order to obtain a stationary point for (OCP) and thus a low upper bound on  $m_{\text{(OCP)}}$ . In order to be able to apply this method to the nondifferentiable total variation seminorm, the latter is smoothed by using an overestimating Huber regularization for the absolute value in the integrand with smoothing parameter  $10^{-3}$ .
- We add the additional integrality constraint  $w(x) \in \mathbb{Z}$  and execute the SLIP algorithm [21, 27] using the subproblem solver described in [37] in order to obtain a low upper bound for the integrality-constrained version of (OCP).
- We use monotonicity properties of the PDE that we do not exploit in our computations elsewhere in order to compute a pointwise McCormick envelope with bounds  $u_\ell$ ,  $u_u$  that are as tight as possible in order to assess the quality of the approximate McCormick envelopes a posteriori.
- We compute approximate McCormick envelopes, that is, solutions to (McC<sub>hh</sub>), with and without bound tightening (and subsequent improvement of the involved constants) as well as the induced lower bounds on (OCP) for decreasing values of  $h$  using the estimates from [section 3.4](#).

All experiments were executed on a laptop computer with Intel (TM) i7-11850H CPU clocked at 2.5 GHz and 64 GB main memory.

#### 4.2 Baseline PDE discretization with Ritz–Galerkin ansatz

We consider conforming finite elements and thus a finite-dimensional subspace  $U_N \subset U$  with dimension  $N \in \mathbb{N}$ . By means of the Lax–Milgram lemma, we obtain that there exists

a unique solution  $u_N \in U_N$ , the Ritz approximation, of the weak formulation on  $U_N$  of the PDEs (9):

$$\int_0^1 \nabla u_N \nabla v_N + \int_0^1 w u_N v_N = \int_0^1 f v_N \quad \text{for all } v_N \in U_N. \quad (31)$$

and (19)

$$\int_0^1 \nabla u_N \nabla v_N + \int_0^1 (P_h w)(P_h u_N) v_N = \int_0^1 f v_N \quad \text{for all } v_N \in U_N. \quad (32)$$

The solutions to (31) satisfy

$$\|\nabla u_N\|_H \leq \frac{1}{c_1(w_\ell)} \|f\|_H,$$

and the solutions to (32) satisfy

$$\|\nabla u_N\|_H \leq \frac{1}{c_2(w_\ell, w_u)} \|f\|_H.$$

The whole analysis from sections 2 and 3 and, in particular, all of the estimates derived for our example PDE in section 3 still hold when discretizing the variational form of the PDE with a conforming finite-element setting and using a piecewise constant control ansatz for  $w$  on the same grid. The only required change is to insert  $U_N$  as the state space for  $U$ . We use first-order Lagrange elements and a discretization of  $(0, 1)$  into  $N = 2,048$  intervals. Regarding the additional control variable  $z$  in the state equation  $Au + z = f$  of the pointwise envelope (McC), we also choose a piecewise constant ansatz on the 2,048 intervals.

Consequently, in our discretizations of (OCP) and (McC), we have the following discretization that serves as a baseline and substitutes the infinite-dimensional setting in our experiments:

- discretization of  $u$  with first-order Lagrange elements on  $N = 2048$  intervals,
- discretization of  $w$  with piecewise constant functions on  $N = 2,048$  intervals, and
- discretization of  $z$  with piecewise constant functions on  $N = 2,048$  intervals.

*Remark 4.1.* In this *discretize-then-optimize* setting, it is possible to directly employ a branch-and-bound procedure on the discretized version of (OCP) when additional integrality restrictions are imposed on  $w$  such as  $w(x) \in \mathbb{Z}$ . Then (9) is just replaced by (31); the lower bounds are computed by means of McCormick envelopes using (32), and the upper bounds are, for example, computed by means of the SLIP algorithm [21, 26].

### 4.3 Practical implementation of OBBT / algorithm 1

In our implementation of algorithm 1, we first minimize the lower bounds  $u_\ell^i$  one by one along the order of the intervals  $i = 1, \dots, N_h$ . Then we maximize the upper bounds  $u_u^i$  one by one along the order of the intervals  $i = 1, \dots, N_h$ . Then we repeat this process. We terminate when a complete run of minimization of the  $u_\ell^i$  (or maximization of the  $u_u^i$ )

does not produce a tightening of any of the bounds by more than a prescribed tolerance of  $10^{-6}$ .

Moreover, in order to avoid numerical problems and in turn incorrect results, additional safeguarding was necessary. Specifically, when bounds are set to the computed value in [algorithm 1](#) ln. 5 or ln. 7, the value might be slightly too sharp because of the numerical precision of the subproblem solver for the OBBT problems. This, however, can cause the lower bounds to increase and the upper bounds to decrease to incorrect values in further tightenings and the feasible set even contracting to an empty set (when the lower and upper bounds cross each other). In order to avoid this situation, the value  $10^{-7}$  is added to the upper bounds and subtracted from the lower bounds computed by the subproblem solver for the OBBT problems before they are assigned as new bounds and [algorithm 1](#) moves on to tighten the next bound.

Because [algorithm 1](#) requires valid initial bounds, we use the bounds that can be inferred from (15) and our choice of  $f$ . Specifically, we use the initial bounds  $u_\ell^i = -5.0444$  and  $u_u^i = 5.0444$  for all  $i \in \{1, \dots, N_h\}$ .

#### 4.4 Results

We first ran Scipy's implementation [41] of L-BFGS-B on (OCP), where we used the unique solvability of the (discretized) PDE constraint to integrate it into the objective and used adjoint calculus in order to compute the derivative. To use this gradient-based solver, we have smoothed the total variation seminorm using a Huber regularization with smoothing parameter  $10^{-3}$ . We inserted the solution into the nonsmooth objective and obtained an upper bound of  $8.3808 \times 10^{-2}$  on the optimal solution to (OCP).

Then we added the integrality constraint  $w(x) \in \mathbb{Z}$  to (OCP) and executed the SLIP algorithm [21, 27] with the subproblem solver from [37] to compute an upper bound on the optimal solution to (OCP) with the additional integrality constraint  $w(x) \in \mathbb{Z}$ . The resulting upper bound was  $8.5551 \times 10^{-2}$ .

Regarding the lower bounds, we first compute pointwise lower bounds for comparison. For our computational example, we compute the optimal objective value for the uniform bounds  $u_\ell \equiv -5.0444$  and  $u_u \equiv 5.0444$  that are enforced on the nodes of our first-order Lagrange elements. We first compute a lower bound by omitting the total variation term from the objective ( $\alpha = 0$ ), which also gives us an overall lower bound on the tracking-type term in the objective. The resulting linearly constrained convex quadratic program is solved by using Gurobi [18], and the computed lower bound is  $6.7701 \times 10^{-2}$ . Then we consider (McC) with the same bounds but including the total variation term. Again, the resulting linearly constrained convex quadratic program is solved by using Gurobi [18], and the computed lower bound is  $6.8649 \times 10^{-2}$ . Moreover, we can characterize the tightest possible pointwise McCormick envelope exactly. Specifically, the monotonicity properties of the Laplacian yield that the discretization of (9) attains its pointwise minimum  $u_{\min}$  for the choice  $w(x) = w_u = 4$  for all  $x \in \Omega$  and its pointwise maximum  $u_{\max}$  for the choice  $w(x) = w_\ell = -4$  for all  $x \in (0, 1)$ . Consequently, we execute (McC) with the bounds  $u_\ell = u_{\min}$  and  $u_u = u_{\max}$  to obtain the tightest lower bound the McCormick envelope can possibly achieve. We obtain a higher optimal objective value of  $8.3679 \times 10^{-2}$ .

This means that if this perfect information on the bounds  $u_u$  and  $u_\ell$  is available when

Table 1: Exact upper and lower bounds and relative gaps (ratio of the difference between upper and lower bound to the lower bound) for (OCP) and its counterpart with integrality restriction  $w(x) \in \mathbb{Z}$ .

	Upper bounds		Lower bounds		
	SLIP ( $w(x) \in \mathbb{Z}$ )	L-BFGS-B	(McC) (tightest)	(McC)	(McC) ( $\alpha = 0$ )
Value	$8.555 \times 10^{-2}$	$8.381 \times 10^{-2}$	$8.368 \times 10^{-2}$	$6.865 \times 10^{-2}$	$6.770 \times 10^{-2}$
Rel. gap (MINLP)			$2.237 \times 10^{-2}$	$2.462 \times 10^{-1}$	$2.637 \times 10^{-1}$
Rel. gap (NLP)			$1.542 \times 10^{-3}$	$2.208 \times 10^{-1}$	$2.379 \times 10^{-1}$

Table 2: Values for  $c_{quad}$  in (28).

conservative	tight
$5.603 \times 10^4$	$1.132 \times 10^3$

solving (McC), the gap between upper and lower bounds gets reduced from  $1.690 \times 10^{-2}$  to  $1.872 \times 10^{-3}$  in the presence of the constraint  $w(x) \in \mathbb{Z}$  (MINLP case) and from  $1.516 \times 10^{-2}$  to  $1.290 \times 10^{-4}$  in the continuous case. In both cases, the relative gap, which is generally computed as the difference between upper and lower bound divided by the lower bound, gets reduced by an order of magnitude. These results are tabulated in table 1.

In the remaining experiments, we use these results as a baseline to compare them with the (approximate) lower bounds obtained by solving instances of (McC<sub>hh</sub>). We executed our experiments on (McC<sub>hh</sub>) on uniform partitions of the domain  $(0, 1)$  into  $N_h \in \{8, 16, 32, 64, 128, 256, 512, 1024\}$  intervals with the corresponding values  $h = 2^{-N_h}$ .

We now consider the approximate lower bounds that are obtained by solving (McC<sub>hh</sub>) for the aforementioned values of  $h$ . In particular, we assess their approximation quality and the running times that are required to compute them by means of the OBBT procedure for the different values of  $h$ . To obtain a valid lower bound on (OCP), we need to subtract  $c_{quad}h^2$  from the optimal objective value of (McC<sub>hh</sub>); see (28). In practice, the values  $j_0$ ,  $\|\tilde{d}_u\|_H$ , and  $\|\tilde{d}_f\|_H$  are not known precisely because this would imply that tight pointwise McCormick envelopes have already been computed.

We therefore use two constants in our considerations that allow us to present a range where we expect that the a priori estimates lie when all of the other constants are known. The larger and thus more conservative constant is obtained by choosing  $j_0 = 0$  and computing  $\tilde{d}_u$  and  $\tilde{d}_f$  with the conservative bounds  $u_\ell \equiv -5.0444$  and  $u_u \equiv 5.0444$ . The smaller constant is computed by choosing  $j_0 = 6.770 \times 10^{-2}$  as well as the optimal bounds  $u_\ell = u_{\min}$  and  $u_u = u_{\max}$ , which are of course not available in a realistic setting. In our setting, the conservative bound is almost 50 times larger than the tight one. Both values are given in table 2.

For our example, the initial bounds that can be deduced from (15) are  $u_\ell^i = -5.0444$  and  $u_u^i = 5.0444$  for all  $i \in \{1, \dots, N_h\}$ . After executing our implementation of algorithm 1, the lower bounds and upper bounds are much closer to each other and reflect the structure of the possible PDE solutions; that is, they tend to zero toward the boundaries of the computational domain. For the finest computed case with  $h = 2^{-10}$ ,  $N_h = 1024$ , and a termination tolerance of  $10^{-6}$  for our implementation of the OBBT procedure, the lower bounds  $u_\ell^i$  for  $P_h u$  vary between 0 and  $5.280 \times 10^{-1}$  and the upper bounds  $u_u^i$  vary between 0 and 1.276, thereby giving a much tighter envelope in which  $P_h u$  for the solu-






Figure 2: (a) Solutions to (McC) (cyan, solid) and (McC<sub>hh</sub>) for  $h = 2^{-10}$ , (orange, dashed) with  $u_{\min}$ ,  $u_{\max}$  (dark blue, solid), and tracking function  $u_d$  (black, solid). (b) Initial (black) and tightened (blue) lower and upper bounds on  $P_h u$  for solutions  $u$  to (19) for  $h = 2^{-10}$ . The area between the tightened lower and upper bounds, where the solution of the PDE can attain values, is colored with a less intense blue.

Table 3: Running times (in seconds) for the OBBT algorithm for different values of  $h$  on instances of (McC<sub>hh</sub>) and (McC) (mesh size  $2^{-11}$ ).

	$2^{-3}$	$2^{-4}$	$2^{-5}$	(McC <sub>hh</sub> ) with $h = 2^{-6}$	$2^{-7}$	$2^{-8}$	$2^{-9}$	$2^{-10}$	(McC)
Time [s]	8.212	$1.612 \times 10^1$	$3.174 \times 10^1$	$7.026 \times 10^1$	$1.652 \times 10^2$	$4.565 \times 10^2$	$1.594 \times 10^3$	$1.080 \times 10^4$	$2.142 \times 10^4$

tions to (19) can lie. Moreover, the optimal solution computed for (McC<sub>hh</sub>) is close to the optimal solution computed for (McC) for fine values of  $h$ . Both findings are visualized in fig. 2.

Generally, the increase in compute times for the bounds is worse than linear with decreasing values of  $h$ . The runtime for the OBBT procedure is 8.212 seconds for  $h = 2^{-3}$  and increases to  $1.080 \times 10^4$  seconds for  $h = 2^{-10}$  for (McC<sub>hh</sub>) compared with a pointwise bound computation with  $2.142 \times 10^4$  seconds (mesh size  $2^{-11}$ ) for (McC). All running times of the OBBT algorithm are tabulated in table 3.

Next, we assess the approximation quality of the bounds induced by the solutions to (McC<sub>hh</sub>) with and without bound tightening. In both cases we observe that the infima converge numerically to their pointwise counterparts of (McC) that are tabulated in table 1. When subtracting  $c_{quad}h^2$  from each of these values for the conservative and tight choices of  $c_{quad}$  from table 2, we obtain valid lower bounds and observe that only for the smallest grid with  $h = 2^{-10}$  all of the lower bounds are positive and thus beat the trivial lower bound 0 that can be found by inspecting the objective. If the OBBT procedure is applied and a tight choice of  $c_{quad}$  is used, the lower bound is positive for the choices of  $h \leq 2^{-7}$ , and the computed bound beats the pointwise bound without OBBT for  $h = 2^{-9}$  and  $h = 2^{-10}$ , thereby yielding the second and third best lower bounds of all computed lower bounds. All computed values are given in table 4.

We assess the quality of the approximate bounds  $m_{(\text{McC}_{hh})}$  after optimization-based bound tightening with respect to  $m_{(\text{McC})}$ . We observe that the difference as well as the relative difference becomes very small for small values of  $h$ , in particular several magnitudes



Table 4: Optimal objective values achieved for ( $\text{McC}_{hh}$ ) and induced bounds for ( $\text{OCP}$ ) using the estimate (28) with and without OBBT with conservative (c) and tight (t) estimate on  $c_{quad}$ . The bounds that are able to beat the trivial bound 0 that can be directly inferred from the nonnegative objective terms are highlighted in gray.

OBBT	$h$	$2^{-3}$	$2^{-4}$	$2^{-5}$	$2^{-6}$	$2^{-7}$	$2^{-8}$	$2^{-9}$	$2^{-10}$
No	$m(\text{McC}_{hh})$	$+7.153 \times 10^{-2}$	$+7.028 \times 10^{-2}$	$+6.868 \times 10^{-2}$	$+6.867 \times 10^{-2}$	$+6.867 \times 10^{-2}$	$+6.866 \times 10^{-2}$	$+6.865 \times 10^{-2}$	$+6.865 \times 10^{-2}$
Yes	$m(\text{McC}_{hh})$	$+8.373 \times 10^{-2}$	$+8.370 \times 10^{-2}$	$+8.368 \times 10^{-2}$	$+8.368 \times 10^{-2}$	$+8.368 \times 10^{-2}$	$+8.368 \times 10^{-2}$	$+8.368 \times 10^{-2}$	$+8.368 \times 10^{-2}$
No	LB (28) (c)	$-8.753 \times 10^{+2}$	$-2.188 \times 10^{+2}$	$-5.464 \times 10^{+1}$	$-1.361 \times 10^{+1}$	$-3.351$	$-7.862 \times 10^{-1}$	$-1.451 \times 10^{-1}$	$+1.522 \times 10^{-2}$
Yes	LB (28) (c)	$-8.753 \times 10^{+2}$	$-2.188 \times 10^{+2}$	$-5.463 \times 10^{+1}$	$-1.360 \times 10^{+1}$	$-3.336$	$-7.712 \times 10^{-1}$	$-1.300 \times 10^{-1}$	$+3.025 \times 10^{-2}$
No	LB (28) (t)	$-1.761 \times 10^{+1}$	$-4.351$	$-1.037$	$-2.076 \times 10^{-1}$	$-4.106 \times 10^{-4}$	$+5.139 \times 10^{-2}$	$+6.433 \times 10^{-2}$	$+6.757 \times 10^{-2}$
Yes	LB (28) (t)	$-1.760 \times 10^{+1}$	$-4.337$	$-1.022$	$-1.926 \times 10^{-1}$	$+1.460 \times 10^{-2}$	$+6.641 \times 10^{-2}$	$+7.936 \times 10^{-2}$	$+8.260 \times 10^{-2}$

Table 5: Difference and relative difference between  $m(\text{McC})$  for  $u_\ell = u_{\min}$ ,  $u_u = u_{\max}$  and  $m(\text{McC}_{hh})$  for  $u_\ell, u_u$  computed using the OBBT procedure.

$h$	$2^{-3}$	$2^{-4}$	$2^{-5}$	$2^{-6}$	$2^{-7}$	$2^{-8}$	$2^{-9}$	$2^{-10}$
$ m(\text{McC}_{hh}) - m(\text{McC}) $	$+5.106 \times 10^{-5}$	$+2.247 \times 10^{-5}$	$+2.292 \times 10^{-6}$	$+5.740 \times 10^{-7}$	$+1.450 \times 10^{-7}$	$+3.820 \times 10^{-8}$	$+1.176 \times 10^{-8}$	$+3.436 \times 10^{-8}$
$\frac{ m(\text{McC}_{hh}) - m(\text{McC}) }{m(\text{McC})}$	$+6.102 \times 10^{-4}$	$+2.686 \times 10^{-4}$	$+2.738 \times 10^{-5}$	$+6.859 \times 10^{-6}$	$+1.733 \times 10^{-6}$	$+4.565 \times 10^{-7}$	$+1.405 \times 10^{-7}$	$+4.106 \times 10^{-7}$

smaller than the error margin of the a priori estimate. The obtained values are given in table 5.

## 5 Conclusion

We show how to replace nonlinearities that are given in a pointwise fashion in the PDE of an (integer) optimal control problem on the example of bilinear terms by generalizing the idea of McCormick envelopes to the infinite-dimensional setting. To keep the computational effort manageable, in particular with respect to a bound-tightening procedure that tightens the convex relaxations, we introduce a two-level approximation scheme by means of a discretization grid, on which the inequalities that constitute the convex envelope are averaged.

Our computational experiments validate that the computational effort of such a procedure can indeed be significantly reduced for coarser grids. However, the a priori approximation quality of the lower bound improves only over trivial bounds when the mesh size is already quite fine and the involved constants can be estimated well. For example, the OBBT procedure for  $h = 2^{-9}$  yields bounds with (relative) gaps to the upper bounds of +6.338 % and +8.549 % in the NLP and MINLP case compared with theoretically optimal bounds 0.1542 % and +2.237 % at +7.445 % of the computational cost for the pointwise OBBT procedure.

Consequently, in order to use the McCormick envelopes together with a priori estimates in a branch-and-bound procedure as sketched in remark 4.1, it is necessary to obtain estimates on  $\|u - u_h\|_H$  of higher order, and a deliberate analysis of the underlying PDE and its discretization are crucial when applying this method. Similarly, good estimates for the constant  $L_u$  in (6) are important, too.

As mentioned before, we used a priori error estimates in our analysis and computational experiments. Because the actual optimal objective values for ( $\text{McC}_{hh}$ ) were much closer to a true lower bound on ( $\text{OCP}$ ) for larger values of  $h$  (coarser local averaging) than implied by our a priori bounds, we are convinced that the analysis and integration of *a posteriori error estimates* into the procedure is key for future research to be able to use relatively coarse grids for ( $\text{McC}_{hh}$ ) and scale this methodology.

One issue that might arise in practice, in particular for PDEs defined on multidimensional domains, is that  $L^\infty$ -bounds as asserted in (12) for our example might not readily be available. However, we believe that additional interior regularity (see, e.g., Theorem 9.51 in [33]) and nonstandard regularity theory (see, e.g., [17]) can help establish such bounds.

Although this goes significantly beyond the scope of this article, we note that the McCormick envelopes suggest defining a branch-and-bound algorithm in function space in the spirit of the recent article [6], where—depending on current fixations and the overlapping of the approximate upper and lower bounds—one refines both the locally averaged McCormick envelopes and the control discretization adaptively until an acceptable gap is reached.

## Acknowledgments

The authors thank Joachim Stöckler (TU Dortmund) for the pointer to the proof of (12) in [35].

This work was supported by the U.S. Department of Energy, Office of Science, Office of Advanced Scientific Computing Research, Scientific Discovery through the Advanced Computing (SciDAC) Program through the FASTMath Institute under Contract No. DE-AC02-06CH11357.

## A Auxiliary results

**Lemma A.1.** *Let  $f^n \rightarrow f$  in  $L^1(\Omega)$ . Let  $g^n \rightarrow g$  in  $L^1(\Omega)$ . If  $f^n \leq g^n$  (or  $f^n \geq g^n$ ) holds pointwise a.e. for all  $n \in \mathbb{N}$ . Then  $f \leq g$  (or  $f \geq g$ , respectively).*

*Proof.* We prove only the case for the  $\leq$ -inequalities because the other case follows by symmetry of the argument. By way of contradiction, we assume that there exist a measurable set  $A \subset \Omega$ ,  $\varepsilon_1 > 0$ , and  $\varepsilon_2 > 0$  such that  $|A| > \varepsilon_1$  and

$$f(x) > g(x) + \varepsilon_2 \text{ for a.a. } x \in A.$$

Egorov's theorem gives that there exists a measurable set  $B \subset A$  and  $n_0 \in \mathbb{N}$  such that  $|B| > \frac{\varepsilon_1}{2}$ , and for all  $n \geq n_0$ , we obtain

$$f^n(x) > g(x) + \frac{\varepsilon_2}{2} \text{ for a.a. } x \in B.$$

Integrating over  $B$  yields

$$\int_B f^n(x) > \int_B g(x) + |B| \frac{\varepsilon_2}{2}.$$

for all  $n \geq n_0$ . Because  $g^n \rightarrow g$ , there exists  $n_1 \geq n_0$  such that

$$\int_B f^n(x) > \int_B g^n(x) + |B| \frac{\varepsilon_2}{4}$$

holds for all  $n \geq n_1$ . Consequently, there must exist a measurable subset  $C \subset B$  of strictly positive measure that  $f^n(x) > g^n(x)$  holds for  $x \in C$ , which contradicts the assumption that  $f^n \leq g^n$  holds pointwise a.e.  $\square$

## References

- [1] L. Ambrosio, N. Fusco, and D. Pallara. *Functions of bounded variation and free discontinuity problems*, volume 254 of *Oxford Mathematical Monographs*. Clarendon Press Oxford, 2000.
- [2] Martin Philip Bendsoe and Ole Sigmund. *Topology optimization: theory, methods, and applications*. Springer Science & Business Media, 2013.
- [3] Christoph Buchheim. Compact extended formulations for binary optimal control problems. *arXiv preprint arXiv:2401.03942*, 2024.
- [4] Christoph Buchheim, Alexandra Grütering, and Christian Meyer. Parabolic optimal control problems with combinatorial switching constraints—Part I: Convex relaxations. *SIAM Journal on Optimization*, 2024. (to appear).
- [5] Christoph Buchheim, Alexandra Grütering, and Christian Meyer. Parabolic optimal control problems with combinatorial switching constraints—Part II: Outer approximation algorithm. *SIAM Journal on Optimization*, 2024. (to appear).
- [6] Christoph Buchheim, Alexandra Grütering, and Christian Meyer. Parabolic optimal control problems with combinatorial switching constraints—Part III: Branch-and-bound algorithm. *arXiv preprint arXiv:2401.10018*, 2024.
- [7] Christoph Buchheim and Maja Hügging. Bounded variation in binary sequences. In *International Symposium on Combinatorial Optimization*, pages 64–75. Springer, 2022.
- [8] Martin Burger, Yiqiu Dong, and Michael Hintermüller. Exact relaxation for classes of minimization problems with binary constraints. *arXiv preprint arXiv:1210.7507*, 2012.
- [9] Eduardo Casas. Control of an elliptic problem with pointwise state constraints. *SIAM Journal on Control and Optimization*, 24(6):1309–1318, 1986.
- [10] Eduardo Casas. Boundary control of semilinear elliptic equations with pointwise state constraints. *SIAM Journal on Control and Optimization*, 31(4):993–1006, 1993.
- [11] Antonin Chambolle and Jérôme Darbon. A parametric maximum flow approach for discrete total variation regularization. *Image Processing and Analysis with Graphs: Theory and Practice*, 2012.
- [12] Antonin Chambolle and Thomas Pock. Approximating the total variation with finite differences or finite elements. In *Handbook of Numerical Analysis*, volume 22, pages 383–417. Elsevier, 2021.
- [13] Antonin Chambolle, Pauline Tan, and Samuel Vaiter. Accelerated alternating descent methods for Dykstra-like problems. *Journal of Mathematical Imaging and Vision*, 59:481–497, 2017.
- [14] Philippe G Ciarlet. *The finite element method for elliptic problems*. SIAM, 2002.

- [15] Giacomo Cristinelli, José A Iglesias, and Daniel Walter. Conditional gradients for total variation regularization with PDE constraints: a graph cuts approach. *arXiv preprint arXiv:2310.19777*, 2023.
- [16] Claudia D’Ambrosio, Andrea Lodi, Sven Wiese, and Cristiana Bragalli. Mathematical programming techniques in water network optimization. *European Journal of Operational Research*, 243(3):774–788, 2015.
- [17] Konrad Gröger. A  $W^{1,p}$ -estimate for solutions to mixed boundary value problems for second order elliptic differential equations. *Mathematische Annalen*, 283:679–687, 1989.
- [18] Gurobi Optimization, LLC. Gurobi Optimizer Reference Manual, 2020.
- [19] Jaroslav Haslinger and Raino AE Mäkinen. On a topology optimization problem governed by two-dimensional Helmholtz equation. *Computational Optimization and Applications*, 62:517–544, 2015.
- [20] Dorit S Hochbaum. An efficient algorithm for image segmentation, markov random fields and related problems. *Journal of the ACM (JACM)*, 48(4):686–701, 2001.
- [21] Sven Leyffer and Paul Manns. Sequential linear integer programming for integer optimal control with total variation regularization. *ESAIM: Control, Optimisation and Calculus of Variations*, 28(66), 2022.
- [22] Sven Leyffer, Paul Manns, and Malte Winckler. Convergence of sum-up rounding schemes for cloaking problems governed by the helmholtz equation. *Computational Optimization and Applications*, 79:193–221, 2021.
- [23] Francesco Maggi. *Sets of Finite Perimeter and Geometric Variational Problems: An Introduction to Geometric Measure Theory*. Number 135. Cambridge University Press, 2012.
- [24] Paul Manns and Christian Kirches. Multidimensional sum-up rounding for elliptic control systems. *SIAM Journal on Numerical Analysis*, 58(6):3427–3447, 2020.
- [25] Paul Manns and Vanja Nikolić. Homotopy trust-region method for phase-field approximations in perimeter-regularized binary optimal control. *arXiv preprint arXiv:2310.12478*, 2023.
- [26] Paul Manns and Annika Schiemann. On integer optimal control with total variation regularization on multi-dimensional domains. *SIAM Journal on Control and Optimization*, 61(6):3415–3441, 2023.
- [27] Paul Manns and Thomas M Surowiec. On binary optimal control in  $H^s(0, T)$ ,  $s < 1/2$ . *Comptes Rendus. Mathématique*, 361(G9):1531–1540, 2023.
- [28] Garth P McCormick. Computability of global solutions to factorable nonconvex programs: Part I-convex underestimating problems. *Mathematical Programming*, 10(1):147–175, 1976.

- [29] Christian Meyer, Arnd Rösch, and Fredi Tröltzsch. Optimal control of PDEs with regularized pointwise state constraints. *Computational Optimization and Applications*, 33:209–228, 2006.
- [30] Ioannis Papamichail and Claire S Adjiman. Proof of convergence for a global optimization algorithm for problems with ordinary differential equations. *Journal of Global Optimization*, 33:83–107, 2005.
- [31] Lawrence E Payne and Hans F Weinberger. An optimal Poincaré inequality for convex domains. *Archive for Rational Mechanics and Analysis*, 5(1):286–292, 1960.
- [32] Marc E Pfetsch, Armin Fügenschuh, Björn Geißler, Nina Geißler, Ralf Gollmer, Benjamin Hiller, Jesco Humpola, Thorsten Koch, Thomas Lehmann, Alexander Martin, et al. Validation of nominations in gas network optimization: models, methods, and solutions. *Optimization Methods and Software*, 30(1):15–53, 2015.
- [33] Michael Renardy and Robert C Rogers. *An introduction to partial differential equations*, volume 13. Springer Science & Business Media, 2006.
- [34] Annika Schiemann and Paul Manns. Discretization of total variation in optimization with integrality constraints. *arXiv preprint arXiv:2403.08346*, 2024.
- [35] Erhard Schmidt. Über die ungleichung, welche die integrale über eine potenz einer funktion und über eine andere potenz ihrer ableitung verbindet. *Mathematische Annalen*, 117(1):301–326, 1940.
- [36] Joseph K Scott and Paul I Barton. Improved relaxations for the parametric solutions of odes using differential inequalities. *Journal of Global Optimization*, 57(1):143–176, 2013.
- [37] Marvin Severitt and Paul Manns. Efficient solution of discrete subproblems arising in integer optimal control with total variation regularization. *INFORMS Journal on Computing*, 35(4), 2023.
- [38] Adam B Singer and Paul I Barton. Global optimization with nonlinear ordinary differential equations. *Journal of Global Optimization*, 34:159–190, 2006.
- [39] Fredi Tröltzsch. *Optimal control of partial differential equations: theory, methods, and applications*, volume 112. American Mathematical Soc., 2010.
- [40] Andreas Veese and Rüdiger Verfürth. Poincaré constants for finite element stars. *IMA Journal of Numerical Analysis*, 32(1):30–47, 2012.
- [41] Pauli Virtanen, Ralf Gommers, Travis E. Oliphant, Matt Haberland, Tyler Reddy, David Cournapeau, Evgeni Burovski, Pearu Peterson, Warren Weckesser, Jonathan Bright, Stéfan J. van der Walt, Matthew Brett, Joshua Wilson, K. Jarrod Millman, Nikolay Mayorov, Andrew R. J. Nelson, Eric Jones, Robert Kern, Eric Larson, C J Carey, İlhan Polat, Yu Feng, Eric W. Moore, Jake VanderPlas, Denis Laxalde, Josef Perktold, Robert Cimrman, Ian Henriksen, E. A. Quintero, Charles R. Harris, Anne M. Archibald, Antônio H. Ribeiro, Fabian Pedregosa, Paul van Mulbregt, and

SciPy 1.0 Contributors. SciPy 1.0: Fundamental algorithms for scientific computing in Python. *Nature Methods*, 17:261–272, 2020.

The submitted manuscript has been created by UChicago Argonne, LLC, Operator of Argonne National Laboratory (“Argonne”). Argonne, a U.S. Department of Energy Office of Science laboratory, is operated under Contract No. DE-AC02-06CH11357. The U.S. Government retains for itself, and others acting on its behalf, a paid-up nonexclusive, irrevocable worldwide license in said article to reproduce, prepare derivative works, distribute copies to the public, and perform publicly and display publicly, by or on behalf of the Government. The Department of Energy will provide public access to these results of federally sponsored research in accordance with the DOE Public Access Plan <http://energy.gov/downloads/doe-public-access-plan>.