

ON THE DENSITY PATCH PROBLEM FOR THE 2-D INHOMOGENEOUS NAVIER-STOKES EQUATIONS

TIAN TIAN HAO, FENG SHAO, DONGYI WEI, AND ZHIFEI ZHANG

ABSTRACT. In this paper, we first construct a class of global strong solutions for the 2-D inhomogeneous Navier-Stokes equations under very general assumption that the initial density is only bounded and the initial velocity is in $H^1(\mathbb{R}^2)$. With suitable assumptions on the initial density, which includes the case of density patch and vacuum bubbles, we prove that Lions' s weak solution is the same as the strong solution with the same initial data. In particular, this gives a complete resolution of the density patch problem proposed by Lions [25]: *for the density patch data $\rho_0 = 1_D$ with a smooth bounded domain $D \subset \mathbb{R}^2$, the regularity of D is preserved by the time evolution of Lions's weak solution.*

1. INTRODUCTION

We consider the inhomogeneous incompressible Navier-Stokes equations in $\mathbb{R}^+ \times \Omega, \Omega \subseteq \mathbb{R}^d, d = 2, 3$:

$$(1.1) \quad \begin{cases} \partial_t \rho + u \cdot \nabla \rho = 0, \\ \rho(\partial_t u + u \cdot \nabla u) - \Delta u + \nabla P = 0, \\ \operatorname{div} u = 0, \\ (\rho, u)|_{t=0} = (\rho_0, u_0), \end{cases}$$

where ρ, u stand for the density and velocity of the fluid respectively, and P is a scalar pressure function.

Ladyženskaja and Solonnikov [21] first addressed the question of unique solvability of (1.1) in a bounded domain Ω . Under the assumption that $u_0 \in W^{2-\frac{2}{p}, p}(\Omega) (p > d)$ is divergence free and vanishes on $\partial\Omega$ and $\rho_0 \in C^1(\Omega)$ is bounded away from zero, they proved the global well-posedness in dimension $d = 2$, and local well-posedness in dimension $d = 3$. We refer to [1, 2, 3, 4, 9, 11, 18, 27] and references therein for the well-posedness results in the critical functional framework. In all these works, the density has to be continuous, bounded and bounded away from zero.

In the case when $\rho_0 \in L^\infty(\mathbb{R}^d)$ with a positive lower bound and $u_0 \in H^1(\mathbb{R}^d)$, Kazhikov [20] proved the global existence of weak solution to the system (1.1), and Simon [30] removed the lower bound assumption on ρ_0 . While with $u_0 \in H^2(\mathbb{R}^d)$, Danchin and Mucha [12] proved that the system (1.1) has a unique local in time solution. Paicu, Zhang and the fourth author [28] proved the global well-posedness of (1.1) with $\rho_0 \in L^\infty(\mathbb{R}^2)$ bounded away from zero and $u_0 \in H^s(\mathbb{R}^2)$ for any $s > 0$, and $\|u_0\|_{L^2} \|\nabla u_0\|_{L^2}$ small in dimension 3. Furthermore, Zhang [31] established the global existence of strong solution with $\rho_0 \in L^\infty(\mathbb{R}^3)$ bounded away from zero and u_0 small in the critical Besov space $\dot{B}_{2,1}^{\frac{1}{2}}(\mathbb{R}^3)$. See [14, 10] for recent important progress in this direction.

Date: June 13, 2024.

1.1. Lions's density patch problem. In this paper, we only focus on the case of $\Omega = \mathbb{R}^2$, since the global regularity is still a famous open problem even for the 3-D incompressible Navier-Stokes equations.

Given $0 \leq \rho_0 \in L^\infty(\mathbb{R}^2)$, and u_0 satisfying $\operatorname{div} u_0 = 0$, $\sqrt{\rho_0}u_0 \in L^2(\mathbb{R}^2)$, Lions [25] proved that the system (1.1) has a global weak solution satisfying the energy inequality

$$(1.2) \quad \frac{1}{2} \|\sqrt{\rho}u(t)\|_{L^2}^2 + \int_0^t \|\nabla u(s)\|_{L^2}^2 ds \leq \frac{1}{2} \|\sqrt{\rho_0}u_0\|_{L^2}^2.$$

Moreover, for any $0 \leq \alpha \leq \beta < \infty$, the Lebesgue measure $\mu(\{x \in \mathbb{R}^2 : \alpha \leq \rho(t, x) \leq \beta\})$ is independent of t . Furthermore, Lions proposed the following open question(see page 34 in [25]):

We would like to mention another interesting open question: suppose that $\rho_0 = 1_D$ for a smooth domain $D(\subset \Omega)$, i.e., a patch of a homogeneous incompressible fluid "surrounded" by the vacuum(or a bubble of vacuum embedded in the fluid). Then, Theorem 2.1 yields at least one global weak solution and (2.17) implies that, for all $t \geq 0$, $\rho(t) = 1_{D(t)}$ for some set such that $\operatorname{vol}(D(t)) = \operatorname{vol}(D)$. In this case, (2.1)-(2.2) can be reformulated as a somewhat complicated free boundary problem. It is also very natural to ask whether the regularity of D is preserved by the time evolution.

The density patch type problem was first solved by Liao and Zhang [23, 24] for the initial density of the form

$$\rho_0 = \eta_1 1_{\Omega_0} + \eta_2 1_{\Omega_0^c},$$

where (η_1, η_2) is any pair of positive constants and Ω_0 is a bounded, simply connected domain with $W^{3,p}$ boundary, see [15, 16] for related works. When $\eta_2 = 0$, the first breakthrough is due to Danchin and Mucha [13], where they constructed a class of strong solutions for $\rho_0 \in L^\infty(\Omega)$ and $u_0 \in H^1(\Omega)$ without extra compatibility condition in the case when the fluid domain Ω is either a bounded domain or the Torus; in particular, for the density patch data, the regularity of $D \subset \Omega$ is preserved by the time evolution of strong solution they constructed. Prange and Tan [29] extend some results in [13] to the case of $\Omega = \mathbb{R}^2$; moreover, for the density with a bubble of vacuum(i.e., $\eta_1 = 0, \eta_2 > 0$), they can prove the weak-strong uniqueness between Lions's weak solution and strong solution they constructed, thus the regularity of D is preserved by the time evolution of Lions's weak solution in such case. However, for the density patch type data(i.e., $\eta_1 > 0, \eta_2 = 0$), [29] still need to impose the compatibility condition (1.3) on the initial data and a weighted L^∞ bound for the weak solution. Thus, the following original version of Lions' s density patch problem remains open:

For the density patch data $\rho_0 = 1_D$ with a smooth bounded domain $D \subset \mathbb{R}^2$, whether the regularity of D is preserved by the time evolution of Lions's weak solution?

This problem is analogue to the vortex patch problem, which was solved by Chemin [7], see also [6] for a new proof.

1.2. Main results. The goal of this paper is twofolds. First of all, we prove the global existence of strong solution under very general assumption on the initial density.

Theorem 1.1. *Given the initial data (ρ_0, u_0) satisfying $0 \leq \rho_0(x) \leq C_0$ and $\rho_0 \not\equiv 0$, $u_0 \in H^1(\mathbb{R}^2)$, $\operatorname{div} u_0 = 0$, the system (1.1) has a global solution $(\rho, u, \nabla P)$ satisfying the weak energy inequality (1.2), $0 \leq \rho(t, x) \leq C_0$, and the following properties:*

- $\sqrt{\rho}u, \nabla u \in L^\infty(\mathbb{R}^+; L^2(\mathbb{R}^2))$;
- $\sqrt{\rho}\dot{u}, \nabla^2 u, \nabla P, \sqrt{t}\nabla\dot{u} \in L^2(\mathbb{R}^+; L^2(\mathbb{R}^2))$, here $\dot{u} \stackrel{\text{def}}{=} u_t + u \cdot \nabla u$;
- $\nabla u \in L^1([0, T]; L^\infty(\mathbb{R}^2))$, $\rho \in C([0, T]; L^p(B_R))$ for all $1 \leq p < \infty$, $R, T \in (0, \infty)$.

Remark 1.1. (1) Compared with the existence part in [29], here we only require $0 \leq \rho_0(x) \leq C_0$ and $\rho_0 \not\equiv 0^1$. For the case of density patch data (FFV-2 condition in Theorem A, [29]), except for $u_0 \in H^1(\mathbb{R}^2)^2$, [29] also imposed the following key compatibility condition

$$(1.3) \quad -\Delta u_0 + \nabla P_0 = \sqrt{\rho_0} g \quad \text{for } u_0 \in L^2(\mathbb{R}^2), \nabla u_0 \in L^1(\mathbb{R}^2), g \in L^2(\mathbb{R}^2).$$

(2) In [8, 19, 26], the initial density allows the vacuum but requires $\rho_0 \in W^{1,q}$, $q > 2$, which excludes the data of density patch or vacuum bubbles.

(3) In this paper, many new ideas are introduced and the method is robust. We believe that our method could be applied to the well-posed problem for the compressible fluids in the presence of vacuum, see [17, 22] for example.

(4) As $\nabla u \in L^1(0, T; L^\infty(\mathbb{R}^2))$ for any $T > 0$, there exists a unique continuous flow X associated with u globally defined on $\mathbb{R}^+ \times \mathbb{R}^+ \times \mathbb{R}^2$ by

$$(1.4) \quad X(t, s, x) = x + \int_s^t u(\sigma, X(\sigma, s, x)) d\sigma.$$

Then $\rho(t, x) = \rho_0(X(0, t, x))$.

To solve the density patch problem, we need to explore the further regularity of strong solution under one of the following two assumptions:

(H1) ρ_0 has a compact support, thus $(1 + |x|^2)\rho_0 \in L^1(\mathbb{R}^2)$.

(H2) there exist $R_0, c_0 \in (0, \infty)$ such that $\int_{B(x_0, R_0)} \rho_0 dx \geq c_0 > 0$ for all $x_0 \in \mathbb{R}^2$.

Remark 1.2. Indeed, (H1) corresponds to the density patch problem, and (H2) allows vacuum bubbles, especially countable equal size vacuum bubbles. While, the following condition in [29]

$$\max\{\underline{\rho} - \rho_0, 0\} \in L^p(\mathbb{R}^d), \quad \text{for some } \underline{\rho} \in (0, \infty), p \in (d/2, \infty),$$

only allows finitely many vacuum bubbles with an equal non-zero measure.

Theorem 1.2. Under the assumptions of Theorem 1.1, if ρ_0 satisfies (H1) or (H2), then for all $2 \leq m < \infty$, $T > 0$, the solution satisfies $\sqrt{t}\nabla^2 u \in L^2([0, T]; L^m(\mathbb{R}^2))$.

As a corollary, we can prove that the regularity of D is preserved by the time evolution of strong solution with the initial density $\rho_0 = a + b\mathbf{1}_D$.

Corollary 1.1. Under the assumptions of Theorem 1.1, if $\rho_0 = a + b\mathbf{1}_D$ for some bounded domain $D \subset \mathbb{R}^2$ with $C^{1,\alpha}$ regularity ($0 < \alpha < 1$) and $a \geq 0$, $a + b \geq 0$, $(a, b) \in \mathbb{R}^2 \setminus \{(0, 0)\}$, then the solution provided by Theorem 1.1 is such that for all $t \geq 0$, $\rho(t, \cdot) = a + b\mathbf{1}_{D_t}$, with $D_t \stackrel{\text{def}}{=} X(t, 0, D)$, where $X(t, s, \cdot)$ stands for the flow of u that is the unique solution of (1.4). Furthermore, D_t has $C^{1,\alpha}$ regularity.

Remark 1.3. In fact, the open set D in Corollary 1.1 is allowed to be a disjoint union of finitely many open connected sets $D^{(j)}$ ($1 \leq j \leq N$) with $\overline{D^{(i)}} \cap \overline{D^{(j)}} = \emptyset$ for $i \neq j$. In this case, there holds

$$D_t = \bigcup_{j=1}^N D_t^{(j)} \quad \text{with} \quad \overline{D_t^{(i)}} \cap \overline{D_t^{(j)}} = \emptyset \quad \text{for } i \neq j \quad \forall t \geq 0.$$

This follows directly from the fact that

$$(1.5) \quad |X(t, 0, x_1) - X(t, 0, x_2)| \geq |x_1 - x_2| \exp\left(-\int_0^t \|\nabla u(s)\|_{L^\infty} ds\right), \quad \forall x_1, x_2 \in \mathbb{R}^2, \forall t > 0.$$

¹It's natural to assume $\rho_0 \not\equiv 0$, otherwise the problem is trivial and the solution is $\rho \equiv 0$, $u \equiv 0$.

²Indeed, they assume the initial velocity satisfies $\sqrt{\rho_0}u_0 \in L^2(\mathbb{R}^2)$ and $\nabla u_0 \in L^2(\mathbb{R}^2)$.

Indeed, let $\varphi(t) = X(t, 0, x_1) - X(t, 0, x_2)$. By (1.4), we have

$$|\varphi'(t)| \leq \|\nabla u(t)\|_{L^\infty} |\varphi(t)|.$$

Then Grönwall's inequality ensures that

$$|\varphi(t)| \geq |\varphi(0)| \exp\left(-\int_0^t \|\nabla u(s)\|_{L^\infty} ds\right).$$

The second goal of this paper gives a complete resolution of Lions's density patch problem via proving the weak-strong uniqueness and Corollary 1.1.

Theorem 1.3. *Under the assumptions of Theorem 1.1, if ρ_0 satisfies (H1) or (H2), then the weak solution in [25] is unique and coincides with strong solution in Theorem 1.1. In particular, for the density patch data or vacuum bubbles, the regularity of D is preserved by the time evolution of Lions's weak solution.*

Remark 1.4. *At this stage, we only prove the uniqueness of weak solutions for ρ_0 satisfying (H1) or (H2). This gives a partial answer to Lions's open question about the uniqueness of weak solutions in two dimensions (see page 31 in [25]).*

1.3. Sketch of the proof. First of all, it is relatively standard to obtain the following energy estimates

$$\frac{1}{2} \|\sqrt{\rho}u(t)\|_{L^2}^2 + \int_0^t \|\nabla u(s)\|_{L^2}^2 ds \leq \frac{1}{2} \|\sqrt{\rho_0}u_0\|_{L^2}^2,$$

and for $i = 0, 1$,

$$\begin{aligned} \sup_{t \in [0, T]} t^i \|\nabla u\|_{L^2}^2 + \int_0^T t^i \int_{\mathbb{R}^2} \rho |\dot{u}|^2 dx dt &\leq C, \\ \int_0^T t^i (\|\nabla^2 u\|_{L^2}^2 + \|\nabla P\|_{L^2}^2) dt &\leq C. \end{aligned}$$

Here and in what follows, the constant C depends only on $\|\rho_0\|_{L^\infty}$, $\|\sqrt{\rho_0}u_0\|_{L^2}$ and $\|\nabla u_0\|_{L^2}$.

Based on an **important observation** $\partial_i u^k \partial_k u^j \partial_j u^i = 0$ due to an algebraic fact $\text{Tr}(A^3) = 0$ if $\text{Tr}(A) = 0$ for a 2×2 matrix A , we can further show that for $i = 1, 2$ (see Lemma 3.2)

$$\begin{aligned} \sup_{t \in [0, T]} t^i \|\sqrt{\rho} \dot{u}\|_{L^2}^2 + \int_0^T t^i \|\nabla \dot{u}\|_{L^2}^2 dt &\leq C, \\ \sup_{t \in [0, T]} t^i (\|\nabla^2 u\|_{L^2}^2 + \|\nabla P\|_{L^2}^2) &\leq C. \end{aligned}$$

However, due to the presence of vacuum of the density, it is highly nontrivial to obtain $u \in L_{loc}^\infty$ and $\nabla u \in L^1(0, T; L^\infty(\mathbb{R}^2))$ from the above energy estimates, which are crucial to prove the existence of strong solution and weak-strong uniqueness.

To prove $u \in L_{loc}^\infty$, we use the following propagation property of lower bound of the density in the sense that if $\int_{B(x_0, R_0)} \rho_0 dx \geq c_0 > 0$, then for $R \geq R_0 + 2t\|\sqrt{\rho_0}u_0\|_{L^2}/\sqrt{c_0}$,

$$\int_{B(x_0, R)} \rho(t, x) dx \geq c_0/4.$$

Let's emphasize that this property also holds for weak solutions (see Lemma 2.2). This lower bound ensures that there exists a $x_t \in B(x_0, R(t))$ such that $|u(t, x_t)| \leq C_1$. On the other hand, from the energy estimates, we can deduce that

$$|u(t, x) - u(t, y)| \leq C(1+t)^{-\frac{1}{2}} \left[\ln \left(2 + |x - y|/\sqrt{t} \right) \right]^{\frac{1}{2}}.$$

Thus, we can conclude the following bound

$$|u(t, x)| \leq C[\ln(2 + |x - x_0| + t^{-1})]^{\frac{1}{2}} \quad \forall x \in \mathbb{R}^2, t > 0.$$

This bound is completely new and may be of independent interest. See Lemma 3.3 for the detail.

To prove $\nabla u \in L^1(0, T; L^\infty(\mathbb{R}^2))$, **the key ingredient** is to establish the following pointwise inequality of $\nabla u(t, x)$, which is also completely new:

$$|\nabla u(t, x)| \leq Cr^{-1} \|\nabla u(t)\|_{L^2} + C \int_{B(x, r)} \frac{|\rho \dot{u}(t, y)|}{|y - x|} dy.$$

where we further have

$$\int_{B(x, r)} \frac{|\rho \dot{u}(t, y)|}{|y - x|} dy \leq Cr \|\nabla \dot{u}\|_{L^2} + C \|\sqrt{\rho} \dot{u}\|_{L^2}.$$

See Lemma 3.5 for the detail.

To prove the $C^{1, \alpha}$ regularity propagation of the density patch or vacuum bubbles, we need to improve the regularity of u . Indeed, if ρ_0 satisfies (H1) or (H2), then for all $2 \leq m < \infty$, we have $\sqrt{t} \nabla^2 u \in L^2([0, T]; L^m(\mathbb{R}^2))$. For this, the key point is to use **the following important inequality** (see Lemma 2.4)

$$\|\sqrt{\rho}(t, \cdot) f\|_{L^m(\mathbb{R}^2)} \leq C \|(\sqrt{\rho}(t, \cdot) f, \nabla f)\|_{L^2(\mathbb{R}^2)} \quad \forall t \in [0, T],$$

such that

$$\begin{aligned} \|\sqrt{t} \nabla^2 u\|_{L^2([0, T]; L^m(\mathbb{R}^2))} &\leq C \|\sqrt{t} \rho \dot{u}\|_{L^2([0, T]; L^m(\mathbb{R}^2))} \leq C \|\sqrt{t} \sqrt{\rho} \dot{u}\|_{L^2([0, T]; L^m(\mathbb{R}^2))} \\ &\leq C \|(\sqrt{t} \rho \dot{u}, \sqrt{t} \nabla \dot{u})\|_{L^2([0, T]; L^2(\mathbb{R}^2))} \leq C. \end{aligned}$$

For the weak-strong uniqueness, we follow the argument of Prange and Tan [29]. Let $(\bar{\rho}, \bar{u})$ be the strong solution obtained in Theorem 1.1 and (ρ, u) be the weak solution provided by Lions in [25] with same initial data (ρ_0, u_0) . We denote $\delta \rho = \rho - \bar{\rho}$, $\delta u = u - \bar{u}$. First of all, we have

$$D(t) \leq \|\rho_0\|_{L^\infty}^{\frac{1}{2}} \|\sqrt{\rho} \delta u\|_{L^2((0, t) \times \mathbb{R}^2)} \exp\left(\|\nabla \bar{u}\|_{L^1(0, T; L^\infty(\mathbb{R}^2))}\right) \leq C \|\sqrt{\rho} \delta u\|_{L^2((0, t) \times \mathbb{R}^2)},$$

where $D(t) = \sup_{0 < s \leq t} s^{-\frac{1}{2}} \|\delta \rho(s, \cdot)\|_{\dot{H}^{-1}(\mathbb{R}^2)}$. Using a duality argument, we can show that

$$\|\sqrt{\rho} \delta u\|_{L^2((0, T) \times \mathbb{R}^2)}^2 \leq \int_0^T |\langle \delta \rho \dot{u}, v \rangle| dt + \int_0^T |\langle \rho \delta u \cdot \nabla \bar{u}, v \rangle| dt,$$

where v is the solution of the linear backward parabolic system (4.16). By Lemma 2.4, we can prove that

$$\int_0^T |\langle \rho \delta u \cdot \nabla \bar{u}, v \rangle| dt \leq \int_0^T \|\sqrt{\rho} \delta u\|_{L^2} \|\sqrt{\rho} v\|_{L^4} \|\nabla \bar{u}\|_{L^4} dt \leq CT^{\frac{1}{4}} \|\sqrt{\rho} \delta u\|_{L^2((0, T) \times \mathbb{R}^2)}^2.$$

Compared with [29], **the main innovation** is to control the trouble term $\int_0^T |\langle \delta \rho \dot{u}, v \rangle| dt$, which is based on the trilinear estimates in Lemma 4.1 and Lemma 4.2, in a way as

$$\int_0^T |\langle \delta \rho \dot{u}, v \rangle| dt \leq C \|(\sqrt{\rho} v, \nabla v)\|_{L^\infty(0, T; L^2)} \int_0^T D(t) |\ln(\sqrt{t} D(t))|^{\frac{1}{2}} \|(t^{\frac{1}{2}} \sqrt{\rho} \dot{u}, t^{\frac{1}{2}} \nabla \dot{u})\|_{L^2} dt.$$

Thus, we can conclude

$$D(t) \leq C \int_0^t D(s) |\ln(\sqrt{s} D(s))|^{\frac{1}{2}} (\|s^{\frac{1}{2}} \sqrt{\rho} \dot{u}\|_{L^2} + s^{\frac{1}{2}} \|\nabla \dot{u}\|_{L^2}) ds.$$

Then Osgood's lemma ensures that $D(t) = 0$ on $[0, T]$.

Notations.

- (1) $\dot{u} \stackrel{\text{def}}{=} u_t + u \cdot \nabla u$ is the material derivative of the velocity.
- (2) $D_t \stackrel{\text{def}}{=} (\partial_t + u \cdot \nabla)$, $\langle x \rangle \stackrel{\text{def}}{=} \sqrt{1 + |x|^2}$.
- (3) $B(a, R) \stackrel{\text{def}}{=} \{x \in \mathbb{R}^2 : |x - a| < R\}$, $B_R = B(0, R)$, $\forall a \in \mathbb{R}^2$, $R > 0$.
- (4) For matrices $A = (A_{ij})$ and $B = (B_{ij})$, we denote $A : B = \sum_{i,j} A_{ij} B_{ij}$.
- (5) $|D|$ is the Lebesgue measure of D for $D \subset \mathbb{R}^2$.
- (6) We shall always denote C to be an absolute constant which may vary from line to line. The dependence of the constant C will be explicitly indicated if there are any exceptions.

2. SOME IMPORTANT PROPERTIES OF WEAK SOLUTIONS

In this section, we prove some important properties of weak solutions (ρ, u) only using the density equation and the energy inequality (1.2). These properties are also very useful for strong solutions.

Lemma 2.1. *For any $\varphi \in C_c^\infty(\mathbb{R}^2)$, we have*

$$|\|\varphi\sqrt{\rho}(t)\|_{L^2} - \|\varphi\sqrt{\rho_0}\|_{L^2}| \leq t\|\sqrt{\rho_0}u_0\|_{L^2}\|\nabla\varphi\|_{L^\infty}.$$

Proof. Testing the density equation of (1.1) against φ^2 gives

$$\int_{\mathbb{R}^2} \rho(t, x)\varphi^2(x) dx - \int_{\mathbb{R}^2} \rho_0(x)\varphi^2(x) dx = \int_0^t \int_{\mathbb{R}^2} (\rho u)(s, x) \cdot \nabla(\varphi^2)(x) dx ds \quad \forall t \geq 0.$$

Let $I(t) = \|\sqrt{\rho(t)}\varphi\|_{L^2}$. Then by Hölder's inequality, we have

$$\begin{aligned} I(t)^2 &\leq I(0)^2 + 2 \int_0^t \|\sqrt{\rho}u(s)\|_{L^2} \|\sqrt{\rho(s)}\varphi\|_{L^2} \|\nabla\varphi\|_{L^\infty} ds \\ &\leq I(0)^2 + 2\|\sqrt{\rho_0}u_0\|_{L^2} \|\nabla\varphi\|_{L^\infty} \int_0^t I(s) ds, \end{aligned}$$

where we have used the energy inequality (1.2) in the last inequality. Now it follows from Grönwall's lemma that

$$I(t) \leq I(0) + t\|\sqrt{\rho_0}u_0\|_{L^2} \|\nabla\varphi\|_{L^\infty},$$

that is,

$$\|\sqrt{\rho(t)}\varphi\|_{L^2} \leq \|\sqrt{\rho_0}\varphi\|_{L^2} + t\|\sqrt{\rho_0}u_0\|_{L^2} \|\nabla\varphi\|_{L^\infty} \quad \forall t \geq 0.$$

On the other hand, since the density equation of (1.1) is time-reversible, we also have

$$\|\sqrt{\rho_0}\varphi\|_{L^2} \leq \|\sqrt{\rho(t)}\varphi\|_{L^2} + t\|\sqrt{\rho}u(t)\|_{L^2} \|\nabla\varphi\|_{L^\infty} \leq \|\sqrt{\rho(t)}\varphi\|_{L^2} + t\|\sqrt{\rho_0}u_0\|_{L^2} \|\nabla\varphi\|_{L^\infty}$$

for all $t \geq 0$, where we used the energy inequality (1.2) again. \square

As $\rho_0 \not\equiv 0$, there exist $x_0 \in \mathbb{R}^2$ and $R_0 \in (0, \infty)$ such that

$$(2.1) \quad \int_{B(x_0, R_0)} \rho_0 dx \geq c_0 > 0.$$

Lemma 2.2. *Assume (2.1) and $R \geq R_0 + 2t\|\sqrt{\rho_0}u_0\|_{L^2}/\sqrt{c_0}$. Then we have*

$$\int_{B(x_0, R)} \rho(t, x) dx \geq c_0/4.$$

Proof. Fix $t > 0$ and $R \geq R_0 + 2t\|\sqrt{\rho_0}u_0\|_{L^2}/\sqrt{c_0}$. For any $\varepsilon > 0$, we choose a bump function $\varphi \in C_c^\infty(\mathbb{R}^2; [0, 1])$ such that $\varphi|_{B(x_0, R_0)} = 1$, $\text{supp } \varphi \subset B(x_0, R)$ and $\|\nabla\varphi\|_{L^\infty} \leq (1 + \varepsilon)(R - R_0)^{-1} \leq \frac{(1+\varepsilon)\sqrt{c_0}}{2t\|\sqrt{\rho_0}u_0\|_{L^2}}$. By Lemma 2.1, we have

$$\|\sqrt{\rho(t)}\varphi\|_{L^2} \geq \|\sqrt{\rho_0}\varphi\|_{L^2} - t\|\sqrt{\rho_0}u_0\|_{L^2}\|\nabla\varphi\|_{L^\infty} \geq \|\sqrt{\rho_0}\varphi\|_{L^2} - \frac{1+\varepsilon}{2}\sqrt{c_0}.$$

It follows from the support properties of φ that

$$\|\sqrt{\rho(t)}\varphi\|_{L^2} \leq \left(\int_{B(x_0, R)} \rho(t, x) dx \right)^{1/2}, \quad \|\sqrt{\rho_0}\varphi\|_{L^2} \geq \left(\int_{B(x_0, R_0)} \rho_0(x) dx \right)^{1/2} \geq \sqrt{c_0},$$

hence

$$\left(\int_{B(x_0, R)} \rho(t, x) dx \right)^{1/2} \geq \sqrt{c_0} - \frac{1+\varepsilon}{2}\sqrt{c_0}.$$

Letting $\varepsilon \rightarrow 0^+$ completes the proof. \square

Lemma 2.3. *If $\langle x \rangle \rho_0 \in L^2(\mathbb{R}^2)$, then there holds*

$$\|\langle x \rangle \sqrt{\rho}(t)\|_{L^2} \leq \|\langle x \rangle \sqrt{\rho_0}\|_{L^2} + t\|\sqrt{\rho_0}u_0\|_{L^2}.$$

Proof. Let $\eta \in C_c^\infty(\mathbb{R})$ be such that $\eta(r) = r$ for $r \in [-1, 1]$ and $|\eta'| \leq 1$. Then $|\eta(r)| \leq r$ for $r \in \mathbb{R}$. For any $R > 1$, we define $\varphi_R(x) = R\eta(\langle x \rangle/R)$ for $x \in \mathbb{R}^2$, then $\varphi_R \in C_c^\infty(\mathbb{R}^2)$, $|\varphi_R(x)| \leq \langle x \rangle$ and $\nabla\varphi_R(x) = \frac{x}{\langle x \rangle}\eta'(\langle x \rangle/R)$, thus $|\nabla\varphi_R| \leq 1$. By Lemma 2.1, we have

$$\|\varphi_R \sqrt{\rho}(t)\|_{L^2} \leq \|\varphi_R \sqrt{\rho_0}\|_{L^2} + t\|\sqrt{\rho_0}u_0\|_{L^2} \leq \|\langle x \rangle \sqrt{\rho_0}\|_{L^2} + t\|\sqrt{\rho_0}u_0\|_{L^2}.$$

Therefore, letting $R \rightarrow +\infty$ and using $\varphi_R(x) = \langle x \rangle$ for $|x| \leq R - 1$ completes the proof. \square

Lemma 2.4. *Let $p \in [2, \infty)$ and $T \in (0, \infty)$. If ρ_0 satisfies (H1), then there exists a constant $C > 0$ depending only on $T, p, \|\rho_0\|_{L^\infty}, \|\sqrt{\rho_0}\langle x \rangle\|_{L^2}, \|\sqrt{\rho_0}u_0\|_{L^2}$, such that (for $f \in \dot{H}^1(\mathbb{R}^2)$)*

$$(2.2) \quad C^{-1}\|\sqrt{\rho}(t, \cdot)f\|_{L^p(\mathbb{R}^2)} \leq \|f\|_{L^2(B_1)} + \|\nabla f\|_{L^2(\mathbb{R}^2)} \leq C\|(\sqrt{\rho}(t, \cdot)f, \nabla f)\|_{L^2(\mathbb{R}^2)}, \quad \forall t \in [0, T].$$

If ρ_0 satisfies (H2), then there exists a constant $C > 0$ depending only on $T, p, R_0, c_0, \|\rho_0\|_{L^\infty}, \|\sqrt{\rho_0}u_0\|_{L^2}$ such that (for $f \in \dot{H}^1(\mathbb{R}^2)$)

$$(2.3) \quad \|f\|_{H^1(\mathbb{R}^2)} + \|\sqrt{\rho}(t, \cdot)f\|_{L^p(\mathbb{R}^2)} \leq C\|(\sqrt{\rho}(t, \cdot)f, \nabla f)\|_{L^2(\mathbb{R}^2)}, \quad \forall t \in [0, T].$$

Proof. We first show (2.2). By Lemma 2.3 and Lemma A.1, we get

$$\begin{aligned} \|\sqrt{\rho}f\|_{L^p(\mathbb{R}^2)} &= \|\sqrt{\rho}\langle x \rangle^{\frac{1}{p}}\langle x \rangle^{-\frac{1}{p}}f\|_{L^p(\mathbb{R}^2)} \leq \|\sqrt{\rho}\langle x \rangle^{\frac{1}{p}}\|_{L^{\frac{3p}{2}}(\mathbb{R}^2)} \|\langle x \rangle^{-\frac{1}{p}}f\|_{L^{3p}(\mathbb{R}^2)} \\ &= \left(\int_{\mathbb{R}^2} \rho^{\frac{3p}{4}} \langle x \rangle^{\frac{3}{2}} dx \right)^{\frac{2}{3p}} \left(\int_{\mathbb{R}^2} \frac{|f|^{3p}}{\langle x \rangle^3} dx \right)^{\frac{1}{3p}} \\ &\leq C\|\rho\|_{L^\infty}^{\frac{1}{2}-\frac{2}{3p}} \left(\int_{\mathbb{R}^2} \rho \langle x \rangle^2 dx \right)^{\frac{2}{3p}} \left(\|f\|_{L^2(B_1)} + \|\nabla f\|_{L^2(\mathbb{R}^2)} \right) \\ (2.4) \quad &\leq C(\|f\|_{L^2(B_1)} + \|\nabla f\|_{L^2(\mathbb{R}^2)}). \end{aligned}$$

Moreover there exists $R_0 > 1$ and $c_0 > 0$ such that $\int_{B(0, R_0)} \rho_0(x) dx \geq 4c_0$. Let $R = R_0 + 2T\|\sqrt{\rho_0}u_0\|_{L^2}/\sqrt{4c_0}$. Then by Lemma 2.2, we have $\int_{B(0, R)} \rho(t, x) dx \geq c_0 > 0, \forall t \in [0, T]$. Then we get by Lemma A.2 that

$$(2.5) \quad \|f\|_{L^2(B_1)} \leq \|f\|_{L^2(B(0, R))} \leq C(\|\sqrt{\rho}(t, \cdot)f\|_{L^2(B(0, R))} + \|\nabla f\|_{L^2(B(0, R))}) \quad \forall t \in [0, T].$$

Then (2.2) follows from (2.4) and (2.5).

Next we prove (2.3). Firstly, Sobolev embedding implies that

$$(2.6) \quad \|\sqrt{\rho}(t, \cdot)f\|_{L^p(\mathbb{R}^2)} \leq C\|\rho_0\|_{L^\infty}^{\frac{1}{2}}\|f\|_{H^1(\mathbb{R}^2)}.$$

By Lemma 2.2, for $R = R_0 + 2T\|\sqrt{\rho_0}u_0\|_{L^2}/\sqrt{c_0}$, $c_0 \in (0, \infty)$ we have

$$\int_{B(x_0, R)} \rho(t, x) dx \geq \frac{c_0}{4} > 0, \quad \forall x_0 \in \mathbb{R}^2, \forall t \in [0, T].$$

Then by Lemma A.2, for every $t \in [0, T]$, $\rho = \rho(t, \cdot)$ we have

$$\|f\|_{L^2(B(x_0, R))} \leq C(\|\sqrt{\rho}f\|_{L^2(B(x_0, R))} + \|\nabla f\|_{L^2(B(x_0, R))}) \quad \text{for all } x_0 \in \mathbb{R}^2.$$

Integrating it over $x_0 \in \mathbb{R}^2$ gives

$$\int_{\mathbb{R}^2} \|f\|_{L^2(B(x_0, R))}^2 dx_0 \leq C\left(\int_{\mathbb{R}^2} \|\sqrt{\rho}f\|_{L^2(B(x_0, R))}^2 dx_0 + \int_{\mathbb{R}^2} \|\nabla f\|_{L^2(B(x_0, R))}^2 dx_0\right).$$

By the Fubini theorem, we finally obtain

$$(2.7) \quad \|f\|_{L^2(\mathbb{R}^2)} \leq C(\|\sqrt{\rho}f\|_{L^2(\mathbb{R}^2)} + \|\nabla f\|_{L^2(\mathbb{R}^2)}),$$

which along with (2.6) implies (2.3). \square

3. GLOBAL EXISTENCE AND REGULARITY OF STRONG SOLUTION

This section is devoted to the global existence of strong solution and the regularity propagation of density patch and vacuum bubbles.

3.1. A priori energy estimates. Let (ρ, u) be a smooth solution to the system (1.1) on $[0, T] \times \mathbb{R}^2$ satisfying $0 \leq \rho \leq \|\rho_0\|_{L^\infty}$.

First of all, (1.2) and Lemma 2.1–Lemma 2.4 hold for strong solutions. The following result has been proved in [26], Lemma 3.2.

Lemma 3.1. *There exists a positive constant C depending only on $\|\rho_0\|_{L^\infty}$, $\|\sqrt{\rho_0}u_0\|_{L^2}$ and $\|\nabla u_0\|_{L^2}$ such that for $i = 0, 1$, there holds*

$$\begin{aligned} \sup_{t \in [0, T]} t^i \|\nabla u\|_{L^2}^2 + \int_0^T t^i \int_{\mathbb{R}^2} \rho |\dot{u}|^2 dx dt &\leq C, \\ \int_0^T t^i (\|\nabla^2 u\|_{L^2}^2 + \|\nabla P\|_{L^2}^2) dt &\leq C. \end{aligned}$$

Motivated by Lemma 3.3 in [26], we derive the estimates for the material derivative of the velocity, which are crucial for the higher order estimates of the velocity. Here we have an important observation $\partial_i u^k \partial_k u^j \partial_j u^i = 0$ due to an algebraic fact $\text{Tr}(A^3) = 0$ if $\text{Tr}(A) = 0$ for a 2×2 matrix A . Then we can remove the assumption $\rho_0 \in L^1(\mathbb{R}^2)$ in [26].

Lemma 3.2. *There exists a positive constant C depending only on $\|\rho_0\|_{L^\infty}$, $\|\sqrt{\rho_0}u_0\|_{L^2}$ and $\|\nabla u_0\|_{L^2}$ such that for $i = 1, 2$,*

$$\begin{aligned} \sup_{t \in [0, T]} t^i \|\sqrt{\rho}\dot{u}\|_{L^2}^2 + \int_0^T t^i \|\nabla \dot{u}\|_{L^2}^2 dt &\leq C, \\ \sup_{t \in [0, T]} t^i (\|\nabla^2 u\|_{L^2}^2 + \|\nabla P\|_{L^2}^2) &\leq C. \end{aligned}$$

Proof. Applying the material derivative D_t to the momentum equation of (1.1), we get ³

$$\rho(\partial_t \dot{u} + u \cdot \nabla \dot{u}) - \Delta \dot{u} = -\partial_k(\partial_k u \cdot \nabla u) - \partial_k u \cdot \nabla \partial_k u - (\nabla P_t + u \cdot \nabla(\nabla P)).$$

Taking the L^2 inner product with \dot{u} to the above equation, we get by integration by parts and $\operatorname{div} u = 0$ that

$$(3.1) \quad \frac{1}{2} \frac{d}{dt} \int_{\mathbb{R}^2} \rho |\dot{u}|^2 dx + \|\nabla \dot{u}\|_{L^2}^2 = \int_{\mathbb{R}^2} (\partial_k u \cdot \nabla u) \cdot \partial_k \dot{u} dx + \int_{\mathbb{R}^2} \partial_k u \cdot (\partial_k u \cdot \nabla \dot{u}) dx + J,$$

where

$$J \stackrel{\text{def}}{=} - \int_{\mathbb{R}^2} (\nabla P_t + u \cdot \nabla(\nabla P)) \cdot \dot{u} dx.$$

The first two terms on the right hand side can be bounded by

$$(3.2) \quad \int_{\mathbb{R}^2} (\partial_k u \cdot \nabla u) \cdot \partial_k \dot{u} dx + \int_{\mathbb{R}^2} \partial_k u \cdot (\partial_k u \cdot \nabla \dot{u}) dx \leq \frac{1}{4} \|\nabla \dot{u}\|_{L^2}^2 + C \|\nabla u\|_{L^4}^4.$$

As for J , using integration by parts, we get (as $\operatorname{div} u = 0$, $\operatorname{div} \dot{u} = \partial_i u^j \partial_j u^i$)

$$\begin{aligned} J &= \int_{\mathbb{R}^2} (P_t \operatorname{div} \dot{u} + (u \cdot \nabla P) \operatorname{div} \dot{u} + \partial_i u^j \partial_j P \dot{u}^i) dx \\ &= \int_{\mathbb{R}^2} (P_t + u \cdot \nabla P) \partial_i u^j \partial_j \dot{u}^i dx - \int_{\mathbb{R}^2} P \partial_i u^j \partial_j \dot{u}^i dx. \end{aligned}$$

We also have

$$\int_{\mathbb{R}^2} (P_t + u \cdot \nabla P) \partial_i u^j \partial_j \dot{u}^i dx = \frac{d}{dt} \int_{\mathbb{R}^2} P \partial_i u^j \partial_j \dot{u}^i dx - \int_{\mathbb{R}^2} P D_t (\partial_i u^j \partial_j \dot{u}^i) dx,$$

and

$$\begin{aligned} - \int_{\mathbb{R}^2} P D_t (\partial_i u^j \partial_j \dot{u}^i) dx &= - \int_{\mathbb{R}^2} P (\partial_i \dot{u}^j \partial_j \dot{u}^i + \partial_i u^j \partial_j \dot{u}^i) dx \\ &\quad + \int_{\mathbb{R}^2} P (\partial_i u^k \partial_k u^j \partial_j \dot{u}^i + \partial_i u^j \partial_j u^k \partial_k \dot{u}^i) dx. \end{aligned}$$

Notice that $\partial_i u^k \partial_k u^j \partial_j \dot{u}^i = \partial_i u^j \partial_j u^k \partial_k \dot{u}^i = 0$. Then by symmetry, we obtain

$$(3.3) \quad J = \frac{d}{dt} \int_{\mathbb{R}^2} P \partial_i u^j \partial_j \dot{u}^i dx - 3 \int_{\mathbb{R}^2} P \partial_i u^j \partial_j \dot{u}^i dx.$$

Moreover, it follows from $\operatorname{div} u = \operatorname{curl} \nabla \dot{u} = 0$ that

$$(3.4) \quad \left| \int_{\mathbb{R}^2} P \partial_i u^j \partial_j \dot{u}^i dx \right| \leq C \|P\|_{BMO} \|\partial_i u^j \partial_j \dot{u}^i\|_{\mathcal{H}^1} \leq C \|\nabla P\|_{L^2} \|\nabla u\|_{L^2} \|\nabla \dot{u}\|_{L^2}.$$

Here \mathcal{H}^1 denotes the Hardy space.

We denote

$$\Psi(t) \stackrel{\text{def}}{=} \frac{1}{2} \int_{\mathbb{R}^2} \rho |\dot{u}|^2 dx - \int_{\mathbb{R}^2} P \partial_i u^j \partial_j \dot{u}^i dx.$$

Plugging (3.2), (3.3) and (3.4) into (3.1) gives

$$(3.5) \quad \Psi'(t) + \|\nabla \dot{u}\|_{L^2}^2 / 2 \leq C \|\nabla u\|_{L^4}^4 + C \|\nabla P\|_{L^2}^2 \|\nabla u\|_{L^2}^2.$$

Using the fact

$$\left| \int_{\mathbb{R}^2} P \partial_i u^j \partial_j \dot{u}^i dx \right| \leq C \|P\|_{BMO} \|\partial_i u^j \partial_j \dot{u}^i\|_{\mathcal{H}^1} \leq C \|\nabla P\|_{L^2} \|\nabla u\|_{L^2}^2,$$

³Repeated indices represent summation.

and the Stokes estimate

$$(3.6) \quad \|\nabla^2 u\|_{L^2}^2 + \|\nabla P\|_{L^2}^2 \leq C\|\rho\dot{u}\|_{L^2}^2 \leq C\|\sqrt{\rho}\dot{u}\|_{L^2}^2,$$

we infer that

$$(3.7) \quad \frac{1}{4} \int_{\mathbb{R}^2} \rho|\dot{u}|^2 dx - C\|\nabla u\|_{L^2}^4 \leq \Psi(t) \leq \int_{\mathbb{R}^2} \rho|\dot{u}|^2 dx + C\|\nabla u\|_{L^2}^4.$$

By Gagliardo-Nirenberg inequality and (3.6), we have

$$(3.8) \quad \begin{aligned} \|\nabla u\|_{L^4}^4 + \|\nabla P\|_{L^2}^2 \|\nabla u\|_{L^2}^2 &\leq C\|\nabla^2 u\|_{L^2}^2 \|\nabla u\|_{L^2}^2 + \|\nabla P\|_{L^2}^2 \|\nabla u\|_{L^2}^2 \\ &\leq C\|\sqrt{\rho}\dot{u}\|_{L^2}^2 \|\nabla u\|_{L^2}^2. \end{aligned}$$

Multiplying (3.5) by t^i ($i = 1, 2$) and using (3.7)-(3.8), the first inequality can be deduced from Lemma 3.1 and (1.2). The second inequality is a direct consequence of the first inequality and (3.6). \square

3.2. L^∞ bounds of u and ∇u . In order to prove the existence of strong solution, we need to obtain a L^∞ bound of $|u(t, x)|$.

Lemma 3.3. *Assume (2.1). Then there exists a constant $C > 0$ depending only on $\|\rho_0\|_{L^\infty}$, $\|\sqrt{\rho_0}u_0\|_{L^2}$, $\|\nabla u_0\|_{L^2}$, R_0 and c_0 such that*

$$|u(t, x)| \leq C[\ln(2 + |x - x_0| + t^{-1})]^{1/2}, \quad \forall t \in [0, T], \quad \forall x \in \mathbb{R}^2.$$

Proof. Let $R(t) := R_0 + 2t\|\sqrt{\rho_0}u_0\|_{L^2}/\sqrt{c_0}$. Then by Lemma 2.2, we have

$$\int_{B(x_0, R(t))} \rho(t, x) dx \geq c_0/4.$$

So, we obtain

$$\frac{c_0}{4} \inf_{B(x_0, R(t))} |u|^2 \leq \int_{B(x_0, R(t))} \rho|u|^2 dx \leq \|\sqrt{\rho_0}u_0\|_{L^2}^2 \leq C,$$

which implies that there exists a $x_t \in B(x_0, R(t))$ such that $|u(t, x_t)| \leq C_1$.

By Lemma 3.1 and Lemma 3.2, we have

$$\begin{aligned} \|\nabla u(t)\|_{L^2}^2 &= \int_{\mathbb{R}^2} |\widehat{u}(t, \xi)|^2 |\xi|^2 d\xi \leq C(1+t)^{-1}, \\ \|\nabla^2 u(t)\|_{L^2}^2 &= \int_{\mathbb{R}^2} |\widehat{u}(t, \xi)|^2 |\xi|^4 d\xi \leq Ct^{-1}(1+t)^{-1}, \end{aligned}$$

which ensure that

$$(3.9) \quad \int_{\mathbb{R}^2} |\widehat{u}(t, \xi)|^2 |\xi|^2 (1+t|\xi|^2) d\xi \leq C(1+t)^{-1}.$$

Thus, for any $x \neq y \in \mathbb{R}^2$, we have (as $|e^{i\xi \cdot x} - e^{i\xi \cdot y}| \leq \min(|\xi||x-y|, 2)$)

$$(3.10) \quad \begin{aligned} |u(t, x) - u(t, y)| &\leq \int_{\mathbb{R}^2} |\widehat{u}(t, \xi)| |e^{i\xi \cdot x} - e^{i\xi \cdot y}| d\xi \\ &\leq \left(\int_{\mathbb{R}^2} |\widehat{u}(t, \xi)|^2 |\xi|^2 (1+t|\xi|^2) d\xi \right)^{1/2} \left(\int_{\mathbb{R}^2} \frac{\min(|\xi|^2|x-y|^2, 4)}{|\xi|^2(1+t|\xi|^2)} d\xi \right)^{1/2}, \end{aligned}$$

where

$$\int_{\mathbb{R}^2} \frac{\min(|\xi|^2|x-y|^2, 4)}{|\xi|^2(1+t|\xi|^2)} d\xi = \left(\int_{|\xi| \leq |x-y|^{-1}} + \int_{|\xi| > |x-y|^{-1}} \right) \frac{\min(|\xi|^2|x-y|^2, 4)}{|\xi|^2(1+t|\xi|^2)} d\xi$$

$$\leq C \ln \left(2 + \frac{|x-y|}{\sqrt{t}} \right),$$

which along with (3.9) and (3.10) ensures that

$$(3.11) \quad |u(t, x) - u(t, y)| \leq C(1+t)^{-\frac{1}{2}} \left[\ln \left(2 + |x-y|/\sqrt{t} \right) \right]^{\frac{1}{2}}.$$

Therefore, by (3.11), $x_t \in B(x_0, R(t))$, $|u(t, x_t)| \leq C_1$ and $R(t) = R_0 + 2t\|\sqrt{\rho_0}u_0\|_{L^2}/\sqrt{c_0}$, we arrive at

$$\begin{aligned} |u(t, x)| &\leq |u(t, x_t)| + |u(t, x) - u(t, x_t)| \\ &\leq C + C(1+t)^{-\frac{1}{2}} [\ln(2 + t^{-1} + |x - x_0| + R(t))]^{\frac{1}{2}} \\ &\leq C [\ln(2 + |x - x_0| + t^{-1})]^{\frac{1}{2}}. \end{aligned}$$

This completes the proof of the lemma. \square

For the estimate of $\|\nabla u\|_{L^\infty}$, we need the following lemma.

Lemma 3.4. *Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ and $F : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be smooth functions satisfying ⁴*

$$Lf = \operatorname{curl} F, \quad \text{where } L \in \{\Delta, \partial_z^2, \partial_{\bar{z}}^2\},$$

then there exists a constant $C > 0$ independent of f, F such that

$$|f(x)| \leq Cr^{-1}\|f\|_{L^2(B(x,r))} + C \int_{B(x,r)} \frac{|F(y)|}{|y-x|} dy, \quad \forall x \in \mathbb{R}^2, \forall r > 0.$$

Proof. By the translation and dilation, it suffices to show that

$$(3.12) \quad |f(0)| \leq C\|f\|_{L^2(B_1)} + C \int_{B_1} \frac{|F(y)|}{|y|} dy$$

for some constant $C > 0$ independent of f and F . We split the proof into three cases.

Case 1. $L = \Delta$. Let $\eta_0 \in C_c^\infty(\mathbb{R}; [0, 1])$ be such that $\eta_0|_{[-1/2, 1/2]} = 1$ and $\operatorname{supp} \eta_0 \subset [-1, 1]$, and let $\xi_0 \in C_c^\infty(\mathbb{R}^2; [0, 1])$ be given by $\xi_0(x) = \eta_0(|x|^2)$ for all $x \in \mathbb{R}^2$. Thanks to

$$\Delta(\xi_0 f) = 2\operatorname{div}(\nabla \xi_0 f) - \Delta \xi_0 f + \operatorname{curl}(\xi_0 F) + F \cdot \nabla^\perp \xi_0,$$

we have

$$f(0) = \int_{\mathbb{R}^2} K(-y) [2\operatorname{div}(\nabla \xi_0 f) - \Delta \xi_0 f + \operatorname{curl}(\xi_0 F) + F \cdot \nabla^\perp \xi_0](y) dy,$$

where $K(x) = \frac{1}{2\pi} \ln|x|$.

Using the integration by parts and support properties of ξ_0 , we get

$$\begin{aligned} &\left| \int_{\mathbb{R}^2} K(-y) \operatorname{div}(\nabla \xi_0 f)(y) dy \right| + \left| \int_{\mathbb{R}^2} K(-y) (\Delta \xi_0 f)(y) dy \right| \\ &= \left| \int_{B_1 \setminus B_{1/2}} \nabla(K(-y)) \cdot (\nabla \xi_0 f)(y) dy \right| + \left| \int_{B_1 \setminus B_{1/2}} K(-y) (\Delta \xi_0 f)(y) dy \right| \\ &\leq C\|f\|_{L^1(B_1)} \leq C\|f\|_{L^2(B_1)}, \end{aligned}$$

and

$$\left| \int_{\mathbb{R}^2} K(-y) \operatorname{curl}(\xi_0 F)(y) dy \right| + \left| \int_{\mathbb{R}^2} K(-y) (F \cdot \nabla^\perp \xi_0)(y) dy \right|$$

⁴ $\partial_z f = \frac{1}{2}(\partial_1 f - i\partial_2 f)$, $\partial_{\bar{z}} f = \frac{1}{2}(\partial_1 f + i\partial_2 f)$.

$$= \left| \int_{B_1} \nabla^\perp(K(-y)) \cdot (\xi_0 F)(y) dy \right| + \left| \int_{B_1 \setminus B_{1/2}} K(-y)(F \cdot \nabla^\perp \xi_0)(y) dy \right| \leq C \int_{B_1} \frac{|F(y)|}{|y|} dy.$$

Combining above estimates, we obtain (3.12).

Case 2. $L = \partial_z^2$. Notice that $\frac{z}{\bar{z}} \partial_z^2 f = \partial_z(\frac{z}{\bar{z}} \partial_z f - \frac{f}{\bar{z}})$. For $\varepsilon \in (0, 1)$, we let

$$\eta_1(s) := \eta_0(s/\varepsilon^2), \quad \eta(s) := \eta_0(s) - \eta_1(s), \quad \forall s \in \mathbb{R}.$$

Firstly, we have

$$(3.13) \quad \int_{|z| \leq 1} \eta(|z|^2) \frac{z}{\bar{z}} \partial_z^2 f dz = - \int_{|z| \leq 1} \partial_z(\eta(|z|^2)) \left(\frac{z}{\bar{z}} \partial_z f - \frac{f}{\bar{z}} \right) dz = I_1 + I_2 + I_3, \quad \text{where}$$

$$I_1 = \int_{|z| \leq 1} f \partial_z \left(\frac{z}{\bar{z}} \partial_z \eta_0(|z|^2) \right) + \frac{f}{\bar{z}} \partial_z \eta_0(|z|^2) dz,$$

$$I_2 = \int_{|z| \leq \varepsilon} \partial_z \eta_1(|z|^2) \frac{z}{\bar{z}} \partial_z f dz, \quad I_3 = - \int_{|z| \leq \varepsilon} \frac{f}{\bar{z}} \partial_z \eta_1(|z|^2) dz.$$

Since $\partial_z \eta_j(|z|^2) = \eta'_j(|z|^2) \bar{z}$ for $j = 0, 1$, then

$$f \partial_z \left(\frac{z}{\bar{z}} \partial_z \eta_0(|z|^2) \right) + \frac{f}{\bar{z}} \partial_z \eta_0(|z|^2) = (2\eta'_0(|z|^2) + |z|^2 \eta''_0(|z|^2)) f,$$

so we get

$$(3.14) \quad |I_1| \leq C \|f\|_{L^1(B_1)} \leq C \|f\|_{L^2(B_1)},$$

and similarly

$$(3.15) \quad |I_2| = \left| \int_{|z| \leq \varepsilon} \eta'_1(|z|^2) z \partial_z f dz \right| \leq \varepsilon \|\nabla f\|_{L^\infty(B_\varepsilon)} \int_{|z| \leq \varepsilon} |\eta'_1(|z|^2)| dz \leq C \varepsilon \|\nabla f\|_{L^\infty(B_\varepsilon)}.$$

Note that $\int_{|z| \leq \varepsilon} \eta'_1(|z|^2) dz = -\pi$, then

$$(3.16) \quad |I_3 + \pi f(0)| = \left| \int_{|z| \leq \varepsilon} \eta'_1(|z|^2) (f(z) - f(0)) dz \right|$$

$$\leq \varepsilon \|\nabla f\|_{L^\infty(B_\varepsilon)} \int_{|z| \leq \varepsilon} |\eta'_1(|z|^2)| dz \leq C \varepsilon \|\nabla f\|_{L^\infty(B_\varepsilon)}.$$

On the other hand, using integration by parts, we have

$$(3.17) \quad \left| \int_{|z| \leq 1} \eta(|z|^2) \frac{z}{\bar{z}} \partial_z^2 f dz \right| = \left| \int_{|z| \leq 1} \eta(|z|^2) \frac{z}{\bar{z}} \operatorname{curl} F dz \right|$$

$$= \left| \int_{|z| \leq 1} F \cdot \nabla^\perp \left(\eta(|z|^2) \frac{z}{\bar{z}} \right) dz \right| \leq C \int_{|z| \leq 1} \frac{|F(z)|}{|z|} dz,$$

where we have used the fact that

$$\left| \partial_z \left(\eta(|z|^2) \frac{z}{\bar{z}} \right) \right| + \left| \partial_{\bar{z}} \left(\eta(|z|^2) \frac{z}{\bar{z}} \right) \right| = \left| \eta'(|z|^2) z + \eta(|z|^2) \frac{1}{\bar{z}} \right| + \left| \eta'(|z|^2) \frac{z^2}{\bar{z}} - \eta(|z|^2) \frac{z}{\bar{z}^2} \right| \leq \frac{C}{|z|}.$$

Combining (3.13), (3.14), (3.15), (3.16), (3.17) and letting $\varepsilon \rightarrow 0^+$, we obtain (3.12).

Case 3. $L = \partial_z^2$. Notice that $\frac{z}{\bar{z}} \partial_z^2 f = \partial_{\bar{z}}(\frac{z}{\bar{z}} \partial_z f - \frac{f}{\bar{z}})$. The proof is similar to Case 2. \square

Now we are in a position to give a priori estimate of $\|\nabla u\|_{L^1(0,T;L^\infty)}$.

Lemma 3.5. *Assume that $0 \leq \rho \leq C_0$ and u is a smooth solution of the following equation*

$$(3.18) \quad \rho \dot{u} - \Delta u + \nabla P = 0, \quad \operatorname{div} u = 0.$$

Then there is a constant $C > 0$ depending only on C_0 , $\|\sqrt{\rho_0}u_0\|_{L^2}$ and $\|\nabla u_0\|_{L^2}$ such that

$$(3.19) \quad |\nabla u(t, x)| \leq Cr^{-1} \|\nabla u(t)\|_{L^2} + C \int_{B(x, r)} \frac{|\rho \dot{u}(t, y)|}{|y - x|} dy, \quad \forall t \in [0, T], \quad \forall x \in \mathbb{R}^2, \quad \forall r > 0,$$

$$(3.20) \quad \text{and} \quad \int_0^T \|\nabla u(t)\|_{L^\infty} dt \leq C[\ln(1 + T)]^{\frac{1}{2}}.$$

Proof. Let $u = (-\partial_2 \psi, \partial_1 \psi)$, then $\operatorname{curl} u = \Delta \psi$. By (3.18), we have $\operatorname{curl}(\Delta u) = \operatorname{curl}(\rho \dot{u})$, and thus $\Delta^2 \psi = \operatorname{curl}(\rho \dot{u})$. Note that

$$|\nabla u| \sim C(|\psi_1| + |\psi_2| + |\psi_3|),$$

where $\psi_1 := \Delta \psi$, $\psi_2 := \partial_z^2 \psi$ and $\psi_3 := \partial_{\bar{z}}^2 \psi$. Hence, by the fact $\Delta^2 \psi = \operatorname{curl}(\rho \dot{u})$ and $\Delta = 4\partial_z \partial_{\bar{z}}$, we obtain

$$\Delta \psi_1 = \operatorname{curl}(\rho \dot{u}), \quad 16\partial_z^2 \psi_2 = \operatorname{curl}(\rho \dot{u}), \quad 16\partial_{\bar{z}}^2 \psi_3 = \operatorname{curl}(\rho \dot{u}).$$

As $\|\psi_k\|_{L^2} \leq C\|\nabla u\|_{L^2}$ for $k = 1, 2, 3$, then by Lemma 3.4, we get the desired estimate (3.19).

Next we prove (3.20). We claim that there exists a constant $C > 0$ such that

$$(3.21) \quad \|\nabla u\|_{L^\infty} \leq Cr_1^{-1} \|\nabla u\|_{L^2} + Cr_1 C_0 \|\nabla \dot{u}\|_{L^2} + CC_0^{\frac{1}{2}} \|\sqrt{\rho} \dot{u}\|_{L^2}$$

for any $r_1 > 0$. Assume (3.21) for the moment. Let $r_1 = C_0^{-1/2} \|\nabla u\|_{L^2}^{1/2} \|\nabla \dot{u}\|_{L^2}^{-1/2}$. Then we infer that

$$\|\nabla u\|_{L^\infty} \leq C \|\nabla u\|_{L^2}^{1/2} \|\nabla \dot{u}\|_{L^2}^{1/2} + C \|\sqrt{\rho} \dot{u}\|_{L^2}.$$

Now it follows from (1.2), Lemma 3.1, Lemma 3.2 and Hölder's inequality that

$$\begin{aligned} \int_0^T \|\nabla u\|_{L^\infty} dt &\leq \int_0^T \|\nabla u\|_{L^2}^{\frac{1}{2}} \|\nabla \dot{u}\|_{L^2}^{\frac{1}{2}} dt + \int_0^T \|\sqrt{\rho} \dot{u}\|_{L^2} dt \\ &\leq \left(\int_0^T \|\nabla u\|_{L^2}^2 dt \right)^{1/4} \left(\int_0^T t(1+t) \|\nabla \dot{u}\|_{L^2}^2 dt \right)^{1/4} \left(\int_0^T t^{-\frac{1}{2}} (1+t)^{-\frac{1}{2}} dt \right)^{1/2} \\ &\quad + \left(\int_0^T (1+t) \|\sqrt{\rho} \dot{u}\|_{L^2}^2 dt \right)^{1/2} \left(\int_0^T (1+t)^{-1} dt \right)^{1/2} \\ &\leq C[\ln(1 + T)]^{\frac{1}{2}}. \end{aligned}$$

This proves (3.20).

Finally, it remains to prove (3.21). Fix $x \in \mathbb{R}^2$ and $r_1 > 0$. We emphasize that in the proof of (3.21), the constant C is independent of x and r_1 . Let $\operatorname{avg}(\dot{u}) = \frac{1}{|B(x, r_1)|} \int_{B(x, r_1)} \dot{u}(y) dy$. By Poincaré inequality we get that

$$(3.22) \quad \|\dot{u} - \operatorname{avg}(\dot{u})\|_{L^m(B(x, r_1))} \leq C_m r_1^{\frac{2}{m}} \|\nabla \dot{u}\|_{L^2(B(x, r_1))}, \quad \forall m \in [1, +\infty).$$

As a consequence, for all $p > 2$, if we let $p' \in (2, +\infty)$ be such that $\frac{1}{p} + \frac{1}{p'} = 1$, then

$$(3.23) \quad \begin{aligned} \int_{B(x, r_1)} \frac{\rho |\dot{u} - \operatorname{avg}(\dot{u})|}{|y - x|} dy &\leq C_0 \|\dot{u} - \operatorname{avg}(\dot{u})\|_{L^p(B(x, r_1))} \| |\cdot|^{-1} \|_{L^{p'}(B(0, r_1))} \\ &\leq Cr_1 C_0 \|\nabla \dot{u}\|_{L^2}. \end{aligned}$$

On the other hand, as $|\cdot|^{-1} \in L^{2,\infty}(\mathbb{R}^2)$, we get by Hölder inequality and interpolation inequality in the Lorentz space that

$$\int_{B(x,r_1)} \frac{\rho(t,y)}{|y-x|} dy \leq \|\rho\|_{L^{2,1}(B(x,r_1))} \| |\cdot|^{-1} \|_{L^{2,\infty}(\mathbb{R}^2)} \leq C \|\rho\|_{L^\infty}^{1/2} \|\rho\|_{L^1(B(x,r_1))}^{1/2},$$

and by (3.22), we have

$$\begin{aligned} & \|\rho\|_{L^1(B(x,r_1))} |\text{avg}(\dot{u})|^2 = \int_{B(x,r_1)} \rho(y) |\text{avg}(\dot{u})|^2 dy \\ & \leq 2 \int_{B(x,r_1)} \rho |\dot{u} - \text{avg}(\dot{u})|^2 dy + 2 \int_{B(x,r_1)} \rho |\dot{u}|^2 dy \leq Cr_1^2 C_0 \|\nabla \dot{u}\|_{L^2}^2 + C \|\sqrt{\rho} \dot{u}\|_{L^2}^2, \end{aligned}$$

hence

$$(3.24) \quad \begin{aligned} \int_{B(x,r_1)} \frac{\rho(y) |\text{avg}(\dot{u})|}{|y-x|} dy & \leq C \|\rho\|_{L^\infty}^{1/2} \|\rho\|_{L^1(B(x,r_1))}^{1/2} |\text{avg}(\dot{u})| \\ & \leq Cr_1 C_0 \|\nabla \dot{u}\|_{L^2} + CC_0^{\frac{1}{2}} \|\sqrt{\rho} \dot{u}\|_{L^2}. \end{aligned}$$

Thanks to (3.23) and (3.24), we have

$$(3.25) \quad \int_{B(x,r_1)} \frac{|\rho \dot{u}(t,y)|}{|y-x|} dy \leq Cr_1 C_0 \|\nabla \dot{u}\|_{L^2} + CC_0^{\frac{1}{2}} \|\sqrt{\rho} \dot{u}\|_{L^2}.$$

By (3.25), we deduce (3.21) from (3.19). \square

3.3. Proof of global existence. In this subsection, we prove Theorem 1.1. We follow the argument in [13]. The idea is to take advantage of classical result to construct smooth solutions corresponding to smoothed-out approximate data with no vacuum, then to pass to the limit. More precisely, we consider

$$u_0^\varepsilon \in C^\infty(\mathbb{R}^2) \quad \text{with} \quad \text{div} u_0^\varepsilon = 0 \quad \text{and} \quad \rho_0^\varepsilon \in C^\infty(\mathbb{R}^2) \quad \text{with} \quad \varepsilon \leq \rho_0^\varepsilon \leq C_0$$

such that

$$u_0^\varepsilon \rightarrow u_0 \text{ in } H^1, \quad \rho_0^\varepsilon \rightharpoonup \rho_0 \text{ in } L^\infty \text{ weak-}^*, \quad \text{and} \quad \rho_0^\varepsilon \rightarrow \rho_0 \text{ in } L_{\text{loc}}^p \text{ if } p < \infty.$$

As $\rho_0 \not\equiv 0$, there exist $R_0 \in (0, \infty)$ and $c_0 > 0$ such that $\int_{B(0,R_0)} \rho_0 dx > c_0 > 0$. Then $\int_{B(0,R_0)} \rho_0^\varepsilon dx > c_0 > 0$ for ε small enough, thus (2.1) holds for $x_0 = 0$ and ρ_0 replaced by ρ_0^ε . We also have $\|u_0^\varepsilon\|_{L^2} \leq \|u_0\|_{L^2} + 1$ for ε small enough, and $\|\sqrt{\rho_0^\varepsilon} u_0^\varepsilon\|_{L^2} \leq \|\rho_0^\varepsilon\|_{L^\infty}^{1/2} \|u_0^\varepsilon\|_{L^2} \leq C_0^{1/2} (\|u_0\|_{L^2} + 1)$. In light of the classical strong solution theory for the system (1.1), there exists a unique global smooth solution $(\rho^\varepsilon, u^\varepsilon, P^\varepsilon)$ corresponding to data $(\rho_0^\varepsilon, u_0^\varepsilon)$. Thus, the triple $(\rho^\varepsilon, u^\varepsilon, P^\varepsilon)$ satisfies all the a priori estimates of previous subsections uniformly with respect to ε . In particular, by Lemma 3.1, Lemma 3.2 and Lemma 3.3, we have $|u^\varepsilon(t, x)| \leq C[\ln(2+|x|+t^{-1})]^{\frac{1}{2}}$, $\int_0^T \|u^\varepsilon\|_{L^\infty(B_R)}^2 \leq C(R, T)$ and

$$\int_0^T t (\|\nabla \dot{u}^\varepsilon\|_{L^2}^2 + \|\nabla u^\varepsilon\|_{L^2}^2 \|\nabla^2 u^\varepsilon\|_{L^2}^2 + \|u^\varepsilon\|_{L^\infty(B_R)}^2 \|\nabla^2 u^\varepsilon\|_{L^2}^2) dt \leq C(R, T).$$

By the definition, we have

$$\begin{aligned} \|\nabla u_t^\varepsilon\|_{L^2(B_R)} & \leq \|\nabla \dot{u}^\varepsilon\|_{L^2(B_R)} + \|\nabla u^\varepsilon \cdot \nabla u^\varepsilon + u^\varepsilon \cdot \nabla^2 u^\varepsilon\|_{L^2(B_R)} \\ & \leq \|\nabla \dot{u}^\varepsilon\|_{L^2} + C(\|\nabla u^\varepsilon\|_{L^2} + \|u^\varepsilon\|_{L^\infty(B_R)}) \|\nabla^2 u^\varepsilon\|_{L^2} \end{aligned}$$

for any $R > 0$. Thus, we obtain

$$\int_0^T t \|\nabla u_t^\varepsilon\|_{L^2(B_R)}^2 dt \leq C(R, T).$$

Similarly, we have

$$\int_0^T \|\sqrt{\rho^\varepsilon} u_t^\varepsilon\|_{L^2(B_R)}^2 dt \leq C(R, T).$$

Let $R(T) = R_0 + 2TC_0^{1/2}(\|u_0\|_{L^2} + 1)/\sqrt{c_0}$ then $R(T) \geq R_0 + 2T\|\sqrt{\rho_0^\varepsilon}u_0^\varepsilon\|_{L^2}/\sqrt{c_0}$ for ε small enough. By Lemma 2.2 we have $\int_{B(0,R)} \rho^\varepsilon(t, x) dx \geq c_0/4$ for $R \geq R(T)$, $t \in [0, T]$. Then by Lemma A.2 we have $\|u_t^\varepsilon\|_{L^2(B_R)} \leq C_R(\|\nabla u_t^\varepsilon\|_{L^2(B_R)} + \|\sqrt{\rho^\varepsilon}u_t^\varepsilon\|_{L^2(B_R)})$ for $R \geq R(T)$, $t \in [0, T]$. Thus $\int_0^T t \|u_t^\varepsilon\|_{L^2(B_R)}^2 dt \leq C(R, T)$ for $R \geq R(T)$, which also holds for $0 < R < R(T)$ as the left hand side is increasing in R .

By Lemma 3.4 in [13], we know that for all $\alpha \in (0, 1/2)$

$$(3.26) \quad \|u^\varepsilon\|_{H^{1/2-\alpha}(0,T;H^1(B_R))} \leq \|u^\varepsilon\|_{L^2(0,T;H^1(B_R))} + C_{\alpha,T} \|t^{1/2}u_t^\varepsilon\|_{L^2(0,T;H^1(B_R))},$$

where $C_{\alpha,T}$ depending only on α and T . By (1.2) and Lemma 3.1, we also have

$$\|\nabla u^\varepsilon\|_{L^2(0,T;H^1)} \leq C,$$

By the interpolation with (3.26), we can deduce that

$$\|(u^\varepsilon, \nabla u^\varepsilon)\|_{H^{1/2-\alpha}([0,T] \times B_R)} \leq C(R, T) \quad \text{for all } R, T \in (0, +\infty).$$

This implies that up to a subsequence, $u^\varepsilon \rightarrow u$ in $L_{loc}^2([0, +\infty); H_{loc}^1)$. By Theorem 2.5 in [25], one can show that

$$\rho^\varepsilon \rightarrow \rho \text{ in } \mathcal{C}([0, T]; L^p(B_R)) \text{ for all } 1 \leq p < +\infty, T, R \in (0, +\infty).$$

Thus, we can pass to the limit for all the terms in the definition of weak solutions, and the weak energy inequality can be obtained by Fatou property. Moreover Lemma 3.1, Lemma 3.2, (3.20) and Lemma 3.3 also hold for the limit solution (ρ, u) . Furthermore, the moment equation is fulfilled in the strong sense, i.e.,

$$\rho(\partial_t u + u \cdot \nabla u) - \Delta u + \nabla P = 0 \quad \text{in } L_{loc}^2(\mathbb{R}^+; L_{loc}^2(\mathbb{R}^2))$$

for some pressure function P with the estimate in Lemma 3.1.

3.4. Regularity propagation of density patch and vacuum bubbles.

Proof of Theorem 1.2. We assume that ρ_0 satisfies (H1) or (H2). By the Stokes estimate and Lemma 2.4, we have

$$\begin{aligned} \|\sqrt{t}\nabla^2 u\|_{L^2([0,T];L^m(\mathbb{R}^2))} &\leq C\|\sqrt{t}\rho\dot{u}\|_{L^2([0,T];L^m(\mathbb{R}^2))} \leq C\|\sqrt{t}\sqrt{\rho}\dot{u}\|_{L^2([0,T];L^m(\mathbb{R}^2))} \\ &\leq C\|(\sqrt{t}\rho\dot{u}, \sqrt{t}\nabla\dot{u})\|_{L^2([0,T];L^2(\mathbb{R}^2))} \leq C, \end{aligned}$$

where we have used Lemma 3.2 in the last inequality, and $C > 0$ is a constant depending only on $\rho_0, u_0, m \geq 2$ and $T > 0$. \square

Now we prove Corollary 1.1.

Proof. Assume that D corresponds to the level set $\{f_0 > 0\}$ of some function $f_0 : \mathbb{R}^2 \rightarrow \mathbb{R}_{\geq 0}$ with $C^{1,\alpha}$ regularity and $df_0 \neq 0$ on $\{f_0 = 0\}$. Then we have $D_t = f_t^{-1}(0, \infty)$ with $f_t \stackrel{\text{def}}{=} f_0 \circ X(t, 0, \cdot)$, and $df_t \neq 0$ on $\{f_t = 0\}$.

Next we check that ρ_0 satisfies (H1) or (H2). In fact, if $a = 0$, then $\rho_0 = b\mathbf{1}_D$, which implies that ρ_0 satisfies (H1) as D is bounded. If $a > 0$, then $\rho_0 \geq a(1 - \mathbf{1}_D)$, and thus there exist $R_0 > 0, c_0 > 0$ such that for all $x_0 \in \mathbb{R}^2$ we have

$$\int_{B(x_0, R_0)} \rho_0 dx \geq \int_{B(x_0, R_0)} a dx - \int_{\mathbb{R}^2} a\mathbf{1}_D dx = a\pi R_0^2 - a|D| = c_0 > 0,$$

which means that ρ_0 satisfies (H2).

By Theorem 1.1 and Theorem 1.2, there exists a global solution $(\rho, u, \nabla P)$ satisfying $\nabla u \in L^\infty(\mathbb{R}^+; L^2(\mathbb{R}^2))$ and $\sqrt{t}\nabla^2 u \in L^2([0, T]; L^m(\mathbb{R}^2))$ for all $T > 0, m \geq 2$. For $\alpha \in (0, 1)$, we have

$$\begin{aligned} \|\nabla u\|_\alpha &\leq C\|\nabla u\|_{L^2}^{1-s}\|\nabla^2 u\|_{L^m}^s \quad \text{for } m > \frac{2}{1-\alpha}, \quad s = \frac{1+\alpha}{2-2/m} \in (0, 1), \\ \|\nabla u\|_{L^\infty} &\leq C\|\nabla u\|_{L^2}^{1-s'}\|\nabla^2 u\|_{L^m}^{s'} \quad \text{for } s' = \frac{1}{2-2/m} \in (0, s), \end{aligned}$$

which imply that $t^{s/2}\nabla u \in L^2([0, T]; C^{0,\alpha})$ and $\nabla u \in L^1([0, T]; C^{0,\alpha})$. Consequently, the flow $X(t, 0, \cdot)$ is in $C^{1,\alpha}$, which implies that f_t is in $C^{1,\alpha}$. \square

4. WEAK-STRONG UNIQUENESS

Throughout this section, let $(\bar{\rho}, \bar{u})$ be the strong solution obtained in Theorem 1.1 and (ρ, u) be the weak solution provided by Lions in [25] with the same initial data (ρ_0, u_0) . Furthermore, we assume that ρ_0 satisfies (H1) or (H2).

4.1. Trilinear estimates. Let $j \in C_c^\infty(\mathbb{R}^2; [0, 9])$ be fixed such that $\text{supp } j \subset B_{1/4}, \int_{\mathbb{R}^2} j(x) dx = 1$ and we denote $j_\varepsilon(x) := \varepsilon^{-2}j(x/\varepsilon)$ for $\varepsilon > 0$.

Lemma 4.1. *There exists $C > 0$ such that for any $f, f_1, f_2 \in H^1, g \in \dot{H}^{-1} \cap L^\infty$ with $\|g\|_{\dot{H}^{-1}} \leq 1/2$, and any $\varepsilon \in (0, 1/2]$ we have*

$$(4.1) \quad \left| \int_{\mathbb{R}^2} g f^2 dx \right| \leq C(\|g\|_{\dot{H}^{-1}} |\ln \varepsilon|^{1/2} + \varepsilon \|g\|_{L^\infty}) \|f\|_{H^1}^2,$$

$$(4.2) \quad \left| \int_{\mathbb{R}^2} g f_1 f_2 dx \right| \leq C(\|g\|_{\dot{H}^{-1}} |\ln \varepsilon|^{1/2} + \varepsilon \|g\|_{L^\infty}) \|f_1\|_{H^1} \|f_2\|_{H^1}.$$

Proof. For any $\varepsilon \in (0, 1/2]$, we have

$$\begin{aligned} \left| \int_{\mathbb{R}^2} g f^2 dx \right| &= \left| \int_{\mathbb{R}^2} g ((j_\varepsilon * f)^2 + (f - j_\varepsilon * f)^2 + 2(j_\varepsilon * f)(f - j_\varepsilon * f)) dx \right| \\ &\leq \|g\|_{\dot{H}^{-1}} \|(j_\varepsilon * f)^2\|_{\dot{H}^1} + \|g\|_{L^\infty} \|f - j_\varepsilon * f\|_{L^2}^2 + 2\|g\|_{L^\infty} \|j_\varepsilon * f\|_{L^2} \|f - j_\varepsilon * f\|_{L^2} \\ &\leq C\|g\|_{\dot{H}^{-1}} \|j_\varepsilon * f\|_{L^\infty} \|j_\varepsilon * f\|_{\dot{H}^1} + C\varepsilon^2 \|g\|_{L^\infty} \|\nabla f\|_{L^2}^2 + C\varepsilon \|g\|_{L^\infty} \|f\|_{H^1}^2 \\ &\leq C|\ln \varepsilon|^{1/2} \|g\|_{\dot{H}^{-1}} \|f\|_{H^1}^2 + C\varepsilon \|g\|_{L^\infty} \|f\|_{H^1}^2, \end{aligned}$$

where we have used the estimates $\|j_\varepsilon * f\|_{L^\infty} \leq \|\widehat{j_\varepsilon f}\|_{L^1} \leq C|\ln \varepsilon|^{1/2} \|f\|_{H^1}$ and (as $\int_{\mathbb{R}^2} j(x) dx = 1$)

$$(4.3) \quad \|f - j_\varepsilon * f\|_{L^2} \leq C\varepsilon \|f\|_{\dot{H}^1}.$$

This proves (4.1). The proof of (4.2) is similar. \square

Lemma 4.2. *For any $C_0 > 0$, there exists $C > 0$ such that for any $f, f_1, f_2 \in \dot{H}^1$, and $g \in \dot{H}^{-1} \cap L^\infty$ with $\|g\|_{\dot{H}^{-1}} \leq 1/2, \|g\|_{L^\infty} + \int_{\mathbb{R}^2} |g|\langle x \rangle^2 dx \leq C_0$, for any $\varepsilon \in (0, 1/2]$ we have*

$$(4.4) \quad \left| \int_{\mathbb{R}^2} g f^2 dx \right| \leq C(\|g\|_{\dot{H}^{-1}} |\ln \varepsilon|^{1/2} + \varepsilon) (\|f\|_{L^2(B_1)} + \|\nabla f\|_{L^2})^2,$$

$$(4.5) \quad \left| \int_{\mathbb{R}^2} g f_1 f_2 dx \right| \leq C (\|g\|_{\dot{H}^{-1}} |\ln \varepsilon|^{1/2} + \varepsilon) (\|f_1\|_{L^2(B_1)} + \|\nabla f_1\|_{L^2}) (\|f_2\|_{L^2(B_1)} + \|\nabla f_2\|_{L^2}).$$

Proof. Let $\eta(x) \in C_c^\infty(\mathbb{R}^2; [0, 1])$ be fixed such that $\text{supp } \eta \subset B_{1/4}$ and $\eta|_{B_{1/8}} = 1$, and let $\eta_\varepsilon(x) = \eta(\varepsilon x)$ for $\varepsilon \in (0, 1/2]$ and $x \in \mathbb{R}^2$. For $f \in \dot{H}^1$, we denote

$$a_\varepsilon := |B_{1/\varepsilon}|^{-1} \int_{B_{1/\varepsilon}} f(y) dy, \quad \forall \varepsilon \in (0, 1/2].$$

Then Poincaré's inequality implies that

$$(4.6) \quad \|\eta_\varepsilon(j_\varepsilon * f - a_\varepsilon)\|_{L^2(\mathbb{R}^2)} \leq C \varepsilon^{-1} \|\nabla f\|_{L^2(\mathbb{R}^2)}$$

for some absolute constant $C > 0$ independent of f and ε . We claim that

$$(4.7) \quad |a_\varepsilon| \leq C (\|f\|_{L^2(B_1)} + |\ln \varepsilon|^{1/2} \|\nabla f\|_{L^2(\mathbb{R}^2)}) \leq C |\ln \varepsilon|^{1/2} (\|f\|_{L^2(B_1)} + \|\nabla f\|_{L^2(\mathbb{R}^2)}),$$

$$(4.8) \quad g \in L^1(\mathbb{R}^2) \text{ with } \int_{\mathbb{R}^2} g(x) dx = 0,$$

$$(4.9) \quad \int_{\mathbb{R}^2} |g(1 - \eta_\varepsilon^2)(j_\varepsilon * f - a_\varepsilon)^2| dx \leq C \varepsilon \|\nabla f\|_{L^2(\mathbb{R}^2)}^2$$

for some constant $C > 0$ depending only on C_0 .

Now we prove (4.4) by admitting (4.7), (4.8) and (4.9), whose proof will be given as soon as we finish the proof of (4.4). We write $f^2 = -(f - j_\varepsilon * f)^2 + (j_\varepsilon * f)^2 + 2f(f - j_\varepsilon * f)$. First of all, by (4.3) we have

$$(4.10) \quad \int_{\mathbb{R}^2} |g(f - j_\varepsilon * f)^2| dx \leq \|g\|_{L^\infty} \|f - j_\varepsilon * f\|_{L^2}^2 \leq C \varepsilon^2 \|\nabla f\|_{L^2}^2.$$

For the term $(j_\varepsilon * f)^2$, we write

$$\begin{aligned} (j_\varepsilon * f)^2 &= (j_\varepsilon * f - a_\varepsilon)^2 + a_\varepsilon^2 + 2a_\varepsilon(j_\varepsilon * f - a_\varepsilon) \\ &= \eta_\varepsilon^2(j_\varepsilon * f - a_\varepsilon)^2 + (1 - \eta_\varepsilon^2)(j_\varepsilon * f - a_\varepsilon)^2 + a_\varepsilon^2 \\ &\quad + 2a_\varepsilon \eta_\varepsilon^2(j_\varepsilon * f - a_\varepsilon) + 2a_\varepsilon(1 - \eta_\varepsilon^2)(j_\varepsilon * f - a_\varepsilon). \end{aligned}$$

It follows from (A.4) and (4.6) that

$$\begin{aligned} &\left| \int_{\mathbb{R}^2} g \eta_\varepsilon^2(j_\varepsilon * f - a_\varepsilon)^2 dx \right| \leq \|g\|_{\dot{H}^{-1}} \|\eta_\varepsilon^2(j_\varepsilon * f - a_\varepsilon)^2\|_{\dot{H}^1} \\ &\leq C \|g\|_{\dot{H}^{-1}} (\|\nabla \eta_\varepsilon(j_\varepsilon * f - a_\varepsilon)\|_{L^\infty} \|\eta_\varepsilon(j_\varepsilon * f - a_\varepsilon)\|_{L^2} + \|\eta_\varepsilon^2(j_\varepsilon * f - a_\varepsilon)\|_{L^\infty} \|j_\varepsilon * \nabla f\|_{L^2}) \\ &\leq C \|g\|_{\dot{H}^{-1}} (\varepsilon \|\nabla f\|_{L^2} |\ln \varepsilon|^{1/2} \cdot \varepsilon^{-1} \|\nabla f\|_{L^2} + \|\nabla f\|_{L^2} |\ln \varepsilon|^{1/2} \cdot \|\nabla f\|_{L^2}) \\ &\leq C \|g\|_{\dot{H}^{-1}} |\ln \varepsilon|^{1/2} \|\nabla f\|_{L^2}^2. \end{aligned}$$

Also, by (4.8) we have $|\int_{\mathbb{R}^2} g a_\varepsilon^2 dx| = 0$. Moreover, (4.6) and (4.7) imply that

$$\begin{aligned} &\left| \int_{\mathbb{R}^2} g a_\varepsilon \eta_\varepsilon^2(j_\varepsilon * f - a_\varepsilon) dx \right| \leq \|g\|_{\dot{H}^{-1}} |a_\varepsilon| \|\eta_\varepsilon^2(j_\varepsilon * f - a_\varepsilon)\|_{\dot{H}^1} \\ &\leq C \|g\|_{\dot{H}^{-1}} |a_\varepsilon| (\|\nabla \eta_\varepsilon\|_{L^\infty} \|\eta_\varepsilon(j_\varepsilon * f - a_\varepsilon)\|_{L^2} + \|j_\varepsilon * \nabla f\|_{L^2}) \\ &\leq C \|g\|_{\dot{H}^{-1}} |a_\varepsilon| (\varepsilon \cdot \varepsilon^{-1} \|\nabla f\|_{L^2} + \|\nabla f\|_{L^2}) \leq C \|g\|_{\dot{H}^{-1}} |\ln \varepsilon|^{1/2} (\|f\|_{L^2(B_1)} + \|\nabla f\|_{L^2(\mathbb{R}^2)})^2. \end{aligned}$$

And by Hölder's inequality, (4.7) and (4.9), we have

$$\int_{\mathbb{R}^2} |g a_\varepsilon (1 - \eta_\varepsilon^2)(j_\varepsilon * f - a_\varepsilon)| dx$$

$$\begin{aligned}
&\leq |a_\varepsilon| \left(\int_{\mathbb{R}^2} |g|(1 - \eta_\varepsilon^2) dx \right)^{1/2} \left(\int_{\mathbb{R}^2} |g(1 - \eta_\varepsilon^2)(j_\varepsilon * f - a_\varepsilon)^2| dx \right)^{1/2} \\
&\leq |a_\varepsilon| \left(\int_{|x| \geq 1/(8\varepsilon)} |g| dx \right)^{1/2} \sqrt{\varepsilon} \|\nabla f\|_{L^2} \leq C|a_\varepsilon| \left(\varepsilon^2 \int_{\mathbb{R}^2} |g||x|^2 dx \right)^{1/2} \sqrt{\varepsilon} \|\nabla f\|_{L^2} \\
&\leq C|\ln \varepsilon|^{1/2} (\|f\|_{L^2(B_1)} + \|\nabla f\|_{L^2(\mathbb{R}^2)}) \varepsilon \cdot \sqrt{\varepsilon} \|\nabla f\|_{L^2} \leq C\varepsilon (\|f\|_{L^2(B_1)} + \|\nabla f\|_{L^2(\mathbb{R}^2)})^2.
\end{aligned}$$

Therefore, we conclude that (also using (4.9))

$$(4.11) \quad \left| \int_{\mathbb{R}^2} g(j_\varepsilon * f)^2 dx \right| \leq C \left(\|g\|_{\dot{H}^{-1}} |\ln \varepsilon|^{1/2} + \varepsilon \right) \left(\|f\|_{L^2(B_1)} + \|\nabla f\|_{L^2(\mathbb{R}^2)} \right)^2.$$

As for the term $2f(f - j_\varepsilon * f)$, by Hölder's inequality, Lemma A.1 and (4.3) we have

$$(4.12) \quad \left| \int_{\mathbb{R}^2} gf(f - j_\varepsilon * f) dx \right| \leq \left(\int_{\mathbb{R}^2} |g|^3 \langle x \rangle^2 dx \right)^{\frac{1}{3}} \left(\int_{\mathbb{R}^2} \frac{|f(x)|^6}{\langle x \rangle^4} dx \right)^{\frac{1}{6}} \|f - j_\varepsilon * f\|_{L^2} \\ \leq CC_0 \left(\|f\|_{L^2(B_1)} + \|\nabla f\|_{L^2(\mathbb{R}^2)} \right) \varepsilon \|\nabla f\|_{L^2(\mathbb{R}^2)} \leq C\varepsilon \left(\|f\|_{L^2(B_1)} + \|\nabla f\|_{L^2(\mathbb{R}^2)} \right)^2,$$

here we used that $\int_{\mathbb{R}^2} |g|^3 \langle x \rangle^2 dx \leq \|g\|_{L^\infty}^2 \int_{\mathbb{R}^2} |g| \langle x \rangle^2 dx \leq C_0^3$. Hence (4.4) follows from (4.10), (4.11) and (4.12). Similarly, one can prove (4.5). \square

To complete the proof of Lemma 4.2, it suffices to prove (4.7), (4.8) and (4.9).

Proof of (4.7). It follows by taking $R = 1/\varepsilon$ in (A.5) and using $|a_1| = |f(1)| \leq C\|f\|_{L^2(B_1)}$. \square

Proof of (4.8). By $\int_{\mathbb{R}^2} |g| \langle x \rangle^2 dx < +\infty$ we have $g \in L^1(\mathbb{R}^2)$. Thus $\hat{g} \in C(\mathbb{R}^2)$. Since $g \in \dot{H}^{-1}(\mathbb{R}^2)$, we have $|\xi|^{-1} \hat{g}(\xi) \in L^2(\mathbb{R}^2)$. If $\int_{\mathbb{R}^2} g dx \neq 0$ then $\hat{g}(0) \neq 0$ and there exists $r > 0$ such that $|\hat{g}(\xi)| \geq |\hat{g}(0)|/2$ for $|\xi| \leq r$. Thus $\int_{B_r} |\xi|^{-1} \hat{g}(\xi)|^2 d\xi \geq (|\hat{g}(0)|/2)^2 \int_{B_r} |\xi|^{-2} d\xi = +\infty$, which contradicts $|\xi|^{-1} \hat{g}(\xi) \in L^2(\mathbb{R}^2)$. Thus $\hat{g}(0) = 0$ and $\int_{\mathbb{R}^2} g dx = 0$. \square

Proof of (4.9). We denote $\tilde{f} = f - a_\varepsilon$, then $j_\varepsilon * f - a_\varepsilon = j_\varepsilon * \tilde{f}$ (as $\int_{\mathbb{R}^2} j(x) dx = 1$). By Hölder's inequality, we get

$$\begin{aligned}
\int_{\mathbb{R}^2} |g(1 - \eta_\varepsilon^2)(j_\varepsilon * f - a_\varepsilon)^2| dx &= \int_{\mathbb{R}^2} |(1 - \eta_\varepsilon^2)gx^{3/2}| \cdot \frac{|(j_\varepsilon * \tilde{f})(x)|^2}{|x|^{3/2}} dx \\
&\leq \left(\int_{|x| \geq \frac{1}{8\varepsilon}} |g|^{4/3} |x|^2 dx \right)^{\frac{3}{4}} \left(\int_{|x| \geq \frac{1}{8\varepsilon}} \frac{|(j_\varepsilon * \tilde{f})(x)|^8}{|x|^6} dx \right)^{\frac{1}{4}}.
\end{aligned}$$

It follows from Lemma A.1 with a scaling argument and Poincaré's inequality that

$$\begin{aligned}
\left(\int_{|x| \geq \frac{1}{\varepsilon}} \frac{|(j_\varepsilon * \tilde{f})(x)|^8}{|x|^6} dx \right)^{\frac{1}{4}} &\leq C\varepsilon (\varepsilon \|j_\varepsilon * \tilde{f}\|_{L^2(B_{1/\varepsilon})} + \|\nabla(j_\varepsilon * \tilde{f})\|_{L^2(\mathbb{R}^2)})^2 \\
&\leq C\varepsilon (\varepsilon \|\tilde{f}\|_{L^2(B_{1/\varepsilon})} + \varepsilon \|\tilde{f} - j_\varepsilon * \tilde{f}\|_{L^2(\mathbb{R}^2)} + \|\nabla \tilde{f}\|_{L^2(\mathbb{R}^2)})^2 \leq C\varepsilon \|\nabla \tilde{f}\|_{L^2(\mathbb{R}^2)}^2 = C\varepsilon \|\nabla f\|_{L^2(\mathbb{R}^2)}^2.
\end{aligned}$$

Then by $\int_{\mathbb{R}^2} |g|^{4/3} |x|^2 dx \leq \|g\|_{L^\infty}^{1/3} \int_{\mathbb{R}^2} |g||x|^2 dx \leq C_0^{4/3}$, we obtain (4.9). \square

4.2. Proof of weak-strong uniqueness.

Proof of Theorem 1.3. Here we follow the proof from Prange and Tan ([29], Proposition 4.1). We focus on the case when $T \in (0, 1/2)$ is small. We denote $\delta\rho := \rho - \bar{\rho}$ and $\delta u := u - \bar{u}$. Then the system for $(\delta\rho, \delta u)$ reads (here $\dot{\bar{u}} := \partial_t \bar{u} + \bar{u} \cdot \nabla \bar{u}$)

$$(4.13) \quad \begin{cases} \partial_t \delta\rho + \bar{u} \cdot \nabla \delta\rho + \delta u \cdot \nabla \rho = 0, & (t, x) \in \mathbb{R}^+ \times \mathbb{R}^2, \\ \rho(\partial_t \delta u + u \cdot \nabla \delta u) - \Delta \delta u + \nabla \delta P = -\delta\rho \dot{\bar{u}} - \rho \delta u \cdot \nabla \bar{u}, \\ \operatorname{div} \delta u = 0, \\ (\delta\rho, \delta u)|_{t=0} = (0, 0). \end{cases}$$

We set $\phi := (-\Delta)^{-1} \delta\rho$ so that $\|\nabla \phi\|_{L^2(\mathbb{R}^2)} = \|\delta\rho\|_{\dot{H}^{-1}(\mathbb{R}^2)}$. Testing the first equation of (4.13) against ϕ and using the fact $\operatorname{div} \bar{u} = \operatorname{div} \delta u = 0$, following the arguments in [29], for all $t \in [0, T]$ we have

$$(4.14) \quad D(t) \leq \|\rho_0\|_{L^\infty}^{\frac{1}{2}} \|\sqrt{\rho} \delta u\|_{L^2((0,t) \times \mathbb{R}^2)} \exp\left(\|\nabla \bar{u}\|_{L^1(0,T;L^\infty(\mathbb{R}^2))}\right) \leq C \|\sqrt{\rho} \delta u\|_{L^2((0,t) \times \mathbb{R}^2)},$$

where

$$(4.15) \quad D(t) := \sup_{0 < s \leq t} s^{-\frac{1}{2}} \|\delta\rho(s, \cdot)\|_{\dot{H}^{-1}(\mathbb{R}^2)}.$$

Next we use a duality argument to control the difference $\sqrt{\rho} \delta u$ in $L^2((0, t) \times \mathbb{R}^2)$. Let $v(t, x)$ be the solution of the following linear backward parabolic system

$$(4.16) \quad \begin{cases} \rho(\partial_t v + u \cdot \nabla v) + \Delta v + \nabla Q = \rho \delta u, \\ \operatorname{div} v = 0, \\ v|_{t=T} = 0. \end{cases}$$

Then there holds

$$(4.17) \quad \begin{aligned} & \sup_{t \in [0, T]} \|(\sqrt{\rho} v, \nabla v)(t, \cdot)\|_{L^2(\mathbb{R}^2)}^2 + \int_0^T \left(\|\nabla v(t, \cdot)\|_{L^2}^2 + \|(\sqrt{\rho} v, \nabla^2 v, \nabla Q)(t, \cdot)\|_{L^2}^2 \right) dt \\ & \leq C_0 \|\sqrt{\rho} \delta u\|_{L^2((0, T) \times \mathbb{R}^2)}^2. \end{aligned}$$

Testing the equation of δu in the system (4.13) by v yields that

$$(4.18) \quad \|\sqrt{\rho} \delta u\|_{L^2((0, T) \times \mathbb{R}^2)}^2 \leq \int_0^T |\langle \delta\rho \dot{\bar{u}}, v \rangle| dt + \int_0^T |\langle \rho \delta u \cdot \nabla \bar{u}, v \rangle| dt.$$

Firstly, we have (by Lemma 2.4 and (4.17))

$$\begin{aligned} & \int_0^T |\langle \rho \delta u \cdot \nabla \bar{u}, v \rangle| dt \leq \int_0^T \|\sqrt{\rho} \delta u\|_{L^2} \|\sqrt{\rho} v\|_{L^4} \|\nabla \bar{u}\|_{L^4} dt, \\ & \leq \int_0^T \|\sqrt{\rho} \delta u\|_{L^2} \|(\sqrt{\rho} v, \nabla v)\|_{L^2} \|\nabla \bar{u}\|_{L^2}^{\frac{1}{2}} \|\nabla^2 \bar{u}\|_{L^2}^{\frac{1}{2}} dt, \\ & \leq T^{\frac{1}{4}} \|\sqrt{\rho} \delta u\|_{L^2((0, T) \times \mathbb{R}^2)} \|(\sqrt{\rho} v, \nabla v)\|_{L^\infty(0, T; L^2)} \|\nabla \bar{u}\|_{L^\infty(0, T; L^2)}^{\frac{1}{2}} \|\nabla^2 \bar{u}\|_{L^2((0, T) \times \mathbb{R}^2)}^{\frac{1}{2}} \\ & \leq CT^{\frac{1}{4}} \|\sqrt{\rho} \delta u\|_{L^2((0, T) \times \mathbb{R}^2)}^2, \end{aligned}$$

where $C > 0$ depends only on $\|\rho_0\|_{L^\infty}$, $\|\sqrt{\rho_0} u_0\|_{L^2}$ and $\|\nabla u_0\|_{L^2}$. Hence, by adjusting $T \in (0, 1/2)$ to smaller if necessary, it follows from (4.18) that

$$(4.19) \quad \|\sqrt{\rho} \delta u\|_{L^2((0, T) \times \mathbb{R}^2)}^2 \leq 2 \int_0^T |\langle \delta\rho \dot{\bar{u}}, v \rangle| dt.$$

Using (4.14), by adjusting $T \in (0, 1/2)$ to smaller if necessary, we have $D(t) \in (0, 1/2)$ for $t \in (0, T]$. Taking $\varepsilon = \|\delta\rho(t)\|_{\dot{H}^{-1}}$ in Lemma 4.1 if ρ_0 satisfies (H2) (or in Lemma 4.2 if ρ_0 satisfies (H1)), and using Lemma 2.4 and (4.15), we obtain

$$(4.20) \quad \begin{aligned} \int_0^T |\langle \delta\rho\dot{u}, v \rangle| dt &\leq C \int_0^T t^{-\frac{1}{2}} \|\delta\rho(t)\|_{\dot{H}^{-1}} |\ln \varepsilon|^{\frac{1}{2}} \|(t^{\frac{1}{2}}\sqrt{\rho}\dot{u}, t^{\frac{1}{2}}\nabla\dot{u})\|_{L^2} \|(\sqrt{\rho}v, \nabla v)\|_{L^2} dt \\ &\leq C \|(\sqrt{\rho}v, \nabla v)\|_{L^\infty(0, T; L^2)} \int_0^T D(t) |\ln(\sqrt{t}D(t))|^{\frac{1}{2}} \|(t^{\frac{1}{2}}\sqrt{\rho}\dot{u}, t^{\frac{1}{2}}\nabla\dot{u})\|_{L^2} dt. \end{aligned}$$

Thus, it follows from (4.14), (4.17), (4.19) and (4.20) that

$$D(t) \leq C \int_0^t D(s) |\ln(\sqrt{s}D(s))|^{\frac{1}{2}} (\|s^{\frac{1}{2}}\sqrt{\rho}\dot{u}\|_{L^2} + s^{\frac{1}{2}}\|\nabla\dot{u}\|_{L^2}) ds.$$

Let $\gamma(t) := \sqrt{t}(\|\sqrt{\rho}\dot{u}\|_{L^2} + \|\nabla\dot{u}\|_{L^2})$, then by Lemma 3.2 we have

$$\sup_{t \in [0, T]} t \|\sqrt{\rho}\dot{u}\|_{L^2}^2 + \int_0^T t \|\nabla\dot{u}\|_{L^2}^2 dt \leq C \implies \int_0^T t (\|\sqrt{\rho}\dot{u}\|_{L^2}^2 + \|\nabla\dot{u}\|_{L^2}^2) dt \leq C,$$

hence $\gamma \in L^2([0, T])$, $|\ln t|^{\frac{1}{2}}\gamma(t) \in L^1([0, T])$, $T, D(t) \in (0, 1/2)$ and we also have

$$D(t) \leq C \int_0^t D(s) |\ln(\sqrt{s}D(s))|^{\frac{1}{2}} \gamma(s) ds \leq C \int_0^t D(s) |\ln D(s)|^{\frac{1}{2}} |\ln s|^{\frac{1}{2}} \gamma(s) ds, \quad \forall t \in [0, T].$$

Furthermore, the function $r \mapsto r |\ln r|^{\frac{1}{2}}$ is increasing near 0^+ and satisfies $\int_0^{1/2} r^{-1} |\ln r|^{-1/2} dr = +\infty$. Hence we can apply Osgood's lemma ([5], Lemma 3.4) to conclude that $D(t) = 0$ on $[0, T]$. By (4.19), (4.20) and (4.15), we have

$$\sqrt{\rho}\delta u \equiv 0 \quad \text{and} \quad \delta\rho \equiv 0 \quad \text{on} \quad [0, T].$$

Now, a direct L^2 estimate of (4.13) gives $\int_0^T \|\nabla\delta u(s, \cdot)\|_{L^2(\mathbb{R}^2)}^2 ds = 0$, and by $\|\delta u\|_{L^2(B_r)} \leq C(\|\sqrt{\rho}\delta u\|_{L^2} + \|\nabla\delta u\|_{L^2}) = 0$ for all $r > 0$ (see Lemma 2.4), we know that $u = \bar{u}$, $\rho = \bar{\rho}$ a.e. in $[0, T] \times \mathbb{R}^2$.

The uniqueness on the whole time interval $[0, \infty)$ can be obtained by a bootstrap argument. \square

APPENDIX A. TECHNICAL LEMMAS

The following lemma comes from Theorem B.1 in [25].

Lemma A.1. *For $m \in [2, \infty)$ and $\ell \in (1 + \frac{m}{2}, \infty)$, there exists a positive constant C depending only on m and ℓ such that for all $f \in H^1(\mathbb{R}^2)$,*

$$(A.1) \quad \left(\int_{\mathbb{R}^2} \frac{|f|^m}{\langle x \rangle^2} (\log \langle x \rangle)^{-\ell} dx \right)^{1/m} \leq C (\|f\|_{L^2(B_1)} + \|\nabla f\|_{L^2(\mathbb{R}^2)}).$$

The following technical lemma is similar to Proposition A.4 in [29], which will be used constantly.

Lemma A.2. *Assume that $\varrho \in L^\infty(\mathbb{R}^2)$ and $\int_{B(x_0, r_0)} \varrho(x) dx \geq c_0 > 0$. Then for all $p \in [1, \infty)$, there exists a constant $C > 0$ depending only on p , $\|\varrho\|_{L^\infty}$ and c_0 such that for $f \in H_{\text{loc}}^1$,*

$$\|f\|_{L^p(B(x_0, r_0))} \leq Cr_0^{2/p} ((1 + r_0)\|\nabla f\|_{L^2(B(x_0, r_0))} + \|\sqrt{\varrho}f\|_{L^2(B(x_0, r_0))}).$$

Proof. Denote $\bar{f} := \frac{1}{|B(x_0, r_0)|} \int_{B(x_0, r_0)} f(y) dy$. Then Poincaré's inequality implies

$$(A.2) \quad \|f - \bar{f}\|_{L^p(B(x_0, r_0))} \leq Cr_0^{2/p} \|\nabla f\|_{L^2(B(x_0, r_0))}.$$

By (A.2) for $p = 2$, we get

$$\begin{aligned} \sqrt{c_0} |\bar{f}| &\leq \|\sqrt{\varrho} \bar{f}\|_{L^2(B(x_0, r_0))} \leq \|\sqrt{\varrho}(f - \bar{f})\|_{L^2(B(x_0, r_0))} + \|\sqrt{\varrho} f\|_{L^2(B(x_0, r_0))} \\ &\leq C \|\varrho\|_{L^\infty}^{1/2} \|f - \bar{f}\|_{L^2(B(x_0, r_0))} + \|\sqrt{\varrho} f\|_{L^2(B(x_0, r_0))} \\ &\leq Cr_0 \|\varrho\|_{L^\infty}^{1/2} \|\nabla f\|_{L^2(B(x_0, r_0))} + \|\sqrt{\varrho} f\|_{L^2(B(x_0, r_0))}. \end{aligned}$$

Therefore, we have

$$\begin{aligned} \|f\|_{L^p(B(x_0, r_0))} &\leq \|f - \bar{f}\|_{L^p(B(x_0, r_0))} + Cr_0^{2/p} |\bar{f}| \\ &\leq Cr_0^{2/p} \|\nabla f\|_{L^2(B(x_0, r_0))} + Cr_0^{2/p+1} \|\nabla f\|_{L^2(B(x_0, r_0))} + Cr_0^{2/p} \|\sqrt{\varrho} f\|_{L^2(B(x_0, r_0))}. \end{aligned}$$

This completes the proof. \square

Lemma A.3. *Let j_ε be as in Subsection 4.1. There exists a constant $C > 0$ such that for all $R > 0$ and all $f \in \dot{H}^1(\mathbb{R}^2)$, there holds*

$$(A.3) \quad |(j_\varepsilon * f)(x) - f_{(R)}| \leq C \|\nabla f\|_{L^2(\mathbb{R}^2)} \left| \ln \frac{R}{\varepsilon} \right|^{\frac{1}{2}}, \quad \forall |x| \leq \frac{R}{4}, \quad \forall \varepsilon \in \left(0, \frac{R}{2}\right)$$

where $f_{(R)} := \frac{1}{|B_R|} \int_{B_R} f(y) dy$. In particular, for all $\varepsilon \in (0, 1/2]$, all $\varphi \in C_c^\infty(\mathbb{R}^2)$ satisfying $\text{supp } \varphi \subset B_{1/(4\varepsilon)}$, we have

$$(A.4) \quad \|\varphi(j_\varepsilon * f - f_{(1/\varepsilon)})\|_{L^\infty} \leq C |\ln \varepsilon|^{\frac{1}{2}} \|\varphi\|_{L^\infty} \|\nabla f\|_{L^2(\mathbb{R}^2)}.$$

Proof. Let $\zeta \in C_c^\infty(\mathbb{R}^2; [0, 1])$ be fixed such that $\text{supp } \zeta \subset B_1$ and $\zeta|_{B_{3/4}} = 1$. Since $\int_{\mathbb{R}^2} j(x) dx = 1$, we have $(j_\varepsilon * f)(x) - f_{(1)} = \int_{\mathbb{R}^2} j_\varepsilon(x-y)(f(y) - f_{(1)}) dy = j_\varepsilon * (f - f_{(1)})(x)$ for $x \in \mathbb{R}^2$. And by the definition of ζ , for $|x| \leq 1/4$ we have

$$(j_\varepsilon * f)(x) - f_{(1)} = \zeta(x) \left((j_\varepsilon * f)(x) - f_{(1)} \right) = j_\varepsilon * \left(\zeta(f - f_{(1)}) \right)(x) := j_\varepsilon * (\zeta g)(x),$$

where $g = f - f_{(1)} \in \dot{H}^1$ satisfies $\int_{B_1} g(y) dy = 0$. By $\|j_\varepsilon * h\|_{L^\infty} \leq C |\ln \varepsilon|^{1/2} \|h\|_{H^1}$ and Poincaré inequality, we know that

$$\|j_\varepsilon * (\zeta g)\|_{L^\infty} \leq C |\ln \varepsilon|^{\frac{1}{2}} \|\zeta g\|_{H^1} \leq C |\ln \varepsilon|^{\frac{1}{2}} (\|g\|_{L^2(B_1)} + \|\nabla g\|_{L^2(\mathbb{R}^2)}) \leq C |\ln \varepsilon|^{\frac{1}{2}} \|\nabla g\|_{L^2(\mathbb{R}^2)}.$$

This proves (A.3) for $R = 1$, and then a scaling argument gives (A.3) for all $R > 0$. \square

Lemma A.4. *There exists a constant $C > 0$ such that for all $f \in \dot{H}^1(\mathbb{R}^2)$, there holds*

$$(A.5) \quad |f_{(R)} - f_{(1)}| \leq C \|\nabla f\|_{L^2(\mathbb{R}^2)} |\ln R|^{1/2}, \quad \forall R \geq 2.$$

Proof. Fix $\varepsilon_0 = 1/4$. Then Lemma A.3 implies that

$$\begin{aligned} |(j_{\varepsilon_0} * f)(0) - f_{(R)}| &\leq C \|\nabla f\|_{L^2} \left| \ln \frac{R}{\varepsilon_0} \right|^{1/2} \leq C \|\nabla f\|_{L^2} |\ln R|^{1/2}, \\ |(j_{\varepsilon_0} * f)(0) - f_{(1)}| &\leq C \|\nabla f\|_{L^2} |\ln \varepsilon_0|^{1/2} \leq C \|\nabla f\|_{L^2} |\ln R|^{1/2} \end{aligned}$$

for all $R \geq 2$. Then (A.5) follows from the triangle inequality. \square

ACKNOWLEDGMENTS.

D. Wei is partially supported by the National Key R&D Program of China under the grant 2021YFA1001500. Z. Zhang is partially supported by NSF of China under Grant 12288101.

REFERENCES

- [1] H. Abidi, Équation de Navier-Stokes avec densité et viscosité variables dans l'espace critique, *Rev. Mat. Iberoam.*, **23** (2007), 537-586.
- [2] H. Abidi and G. Gui, Global well-posedness for the 2-D inhomogeneous incompressible Navier-Stokes system with large initial data in critical spaces, *Arch. Ration. Mech. Anal.*, **242** (2021), 1533-1570.
- [3] H. Abidi, G. Gui and P. Zhang, On the wellposedness of 3-D inhomogeneous Navier-Stokes equations in the critical spaces, *Arch. Ration. Mech. Anal.*, **204** (2012), 189-230.
- [4] H. Abidi and M. Paicu, Existence globale pour un fluide inhomogène, *Ann. Inst. Fourier (Grenoble)*, **57** (2007), 883-917.
- [5] H. Bahouri, J. Y. Chemin and R. Danchin, *Fourier Analysis and Nonlinear Partial Differential Equations*, Grundlehren der Mathematischen Wissenschaften, vol. **343** (2011), Springer-Verlag, Berlin, Heidelberg.
- [6] A. L. Bertozzi and P. Constantin, Global regularity for vortex patches, *Comm. Math. Phys.*, **152** (1993), 19-28.
- [7] J. Y. Chemin, Persistence de structures géométriques dans les fluides incompressibles bidimensionnels, *Ann. Sci. École Norm. Sup. (4)*, **26** (1993), 517-542.
- [8] H. Choe and H. Kim, Strong solutions of the Navier-Stokes equations for nonhomogeneous incompressible fluids, *Commun. PDE*, **28**(2003), 1183-201
- [9] R. Danchin, Density-dependent incompressible viscous fluids in critical spaces, *Proc. Roy. Soc. Edinburgh Sect. A*, **133** (2003), 1311-1334.
- [10] R. Danchin, Global well-posedness for 2D inhomogeneous viscous flows with rough data via dynamic interpolation, *arXiv:2404.02541*.
- [11] R. Danchin and P. B. Mucha, A Lagrangian approach for the incompressible Navier-Stokes equations with variable density, *Comm. Pure Appl. Math.*, **65** (2012), 1458-1480.
- [12] R. Danchin and P. B. Mucha, Incompressible flows with piecewise constant density, *Arch. Ration. Mech. Anal.*, **207** (2013), 991-1023.
- [13] R. Danchin and P. B. Mucha, The incompressible Navier-Stokes equations in vacuum, *Comm. Pure Appl. Math.*, **72** (2019), 1351-1385.
- [14] R. Danchin and S. Wang, Global unique solutions for the inhomogeneous Navier-Stokes equations with only bounded density in critical regularity spaces, *Comm. Math. Phys.*, **399** (2023), 1647-1688.
- [15] R. Danchin and X. Zhang, On the persistence of Hölder regular patches of density for the inhomogeneous Navier-Stokes equations, *J. Éc. polytech. Math.*, **4** (2017), 781-811
- [16] F. Gancedo and E. García-Juárez, Global regularity of 2D density patches for inhomogeneous Navier-Stokes, *Arch. Ration. Mech. Anal.* **229** (2018), 339-360.
- [17] X. Huang, J. Li and Z. Xin, Global well-posedness of classical solutions with large oscillations and vacuum to the three-dimensional isentropic compressible Navier-Stokes equations, *Comm. Pure Appl. Math.*, **65**(2012), 549-585.
- [18] J. Huang, M. Paicu and P. Zhang, Global wellposedness to incompressible inhomogeneous fluid system with bounded density and non-Lipschitz velocity, *Arch. Ration. Mech. Anal.*, **209** (2013), 631-682.
- [19] X. Huang and Y. Wang, Global strong solution with vacuum to the two-dimensional density-dependent Navier-Stokes system, *SIAM J. Math. Appl.*, **46**(2014), 1771-88
- [20] A. Kazhikhov, Solvability of the initial-boundary value problem for the equations of the motion of an inhomogeneous viscous incompressible fluid, *Dokl. Akad. Nauk SSSR*, **216** (1974), 1008-1010.
- [21] O. A. Ladyženskaja and V.A. Solonnikov, The unique solvability of an initial-boundary value problem for viscous incompressible inhomogeneous fluids. (Russian) Boundary value problems of mathematical physics, and related questions of the theory of functions, 8, *Zap. Naučn. Sem. Leningrad. Otdel. Mat. Inst. Steklov.*, **52** (1975), 52-109, 218-219.
- [22] X. Liao and S. Zodji, Global-in-time well-posedness of the compressible Navier-Stokes equations with striated density, *arXiv:2405.11900*.
- [23] X. Liao and P. Zhang, On the global regularity of the two-dimensional density patch for inhomogeneous incompressible viscous flow, *Arch. Ration. Mech. Anal.*, **220** (2016), 937-981.

- [24] X. Liao and P. Zhang, Global regularity of 2D density patches for viscous inhomogeneous incompressible flow with general density: low regularity case, *Comm. Pure Appl. Math.*, **72**(2019), 835–884.
- [25] P. L. Lions, *Mathematical topics in fluid mechanics. Vol. 1. Incompressible models*, Oxford Lecture Series in Mathematics and its Applications, 3. Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1996.
- [26] B. Lü, X. Shi and X. Zhong, Global existence and large time asymptotic behavior of strong solutions to the Cauchy problem of 2D density-dependent Navier-Stokes equations with vacuum, *Nonlinearity*, **31** (2018), no.6, 2617-2632.
- [27] M. Paicu and P. Zhang, Global solutions to the 3-D incompressible inhomogeneous Navier-Stokes system, *J. Funct. Anal.*, **262** (2012), 3556-3584.
- [28] M. Paicu, P. Zhang and Z. Zhang, Global unique solvability of inhomogeneous Navier-Stokes equations with bounded density, *Comm. Partial Differential Equations*, **38** (2013), 1208-1234.
- [29] C. Prange and J. Tan, Free boundary regularity of vacuum states for incompressible viscous flows in unbounded domains, *arXiv:2310.09288v3*.
- [30] J. Simon, Nonhomogeneous viscous incompressible fluids: existence of velocity, density, and pressure, *SIAM J. Math. Anal.*, **21** (1990), 1093-1117.
- [31] P. Zhang, Global Fujita-Kato solution of the 3-D inhomogeneous incompressible Navier-Stokes system, *Adv. Math.*, **363** (2020), 107007, 43 pp.

(T. Hao) SCHOOL OF MATHEMATICAL SCIENCES, PEKING UNIVERSITY, BEIJING 100871, CHINA
Email address: haotiantian@pku.edu.cn

(F. Shao) SCHOOL OF MATHEMATICAL SCIENCES, PEKING UNIVERSITY, BEIJING 100871, CHINA
Email address: fshao@stu.pku.edu.cn

(D. Wei) SCHOOL OF MATHEMATICAL SCIENCES, PEKING UNIVERSITY, BEIJING 100871, CHINA
Email address: jnwdyi@pku.edu.cn

(Z. Zhang) SCHOOL OF MATHEMATICAL SCIENCES, PEKING UNIVERSITY, BEIJING 100871, CHINA
Email address: zfzhang@math.pku.edu.cn