THE DEFECT OF WEAK APPROXIMATION FOR A REDUCTIVE GROUP OVER A GLOBAL FIELD

MIKHAIL BOROVOI

ABSTRACT. Using ideas of Sansuc and results of Colliot-Thélène and Tate, we compute the defect of weak approximation for a reductive group G over a global field K in terms of the algebraic fundamental group of G.

1. Introduction

1.1. Let K be a global field (a number field or a global function field), and let K^s be a fixed separable closure of K. Let G be a reductive group over K (we follow the convention of SGA3, where reductive groups are assumed to be connected). Let \mathcal{V}_K denote the set of places of K, and let $S \subset \mathcal{V}_K$ be a finite set of places. Consider the group

$$G(K_S) := \prod_{v \in S} G(K_v)$$

where K_v denotes the completion of K at v. The group G(K) embeds diagonally into $G(K_S)$. One says that G has the weak approximation property in S if G(K) is dense in $G(K_S)$. One says that G has the weak approximation property if it has the weak approximation property in S for any finite subset $S \subset \mathcal{V}_K$.

Without assuming that G has the weak approximation property in S, let $\overline{G(K)}_S$ denote the closure of G(K) in $G(K_S)$. Sansuc [San81, §3] showed (in the number field case) that the subgroup $\overline{G(K)}_S$ is normal in $G(K_S)$ and that the quotient group

$$A_S(G) := G(K_S)/\overline{G(K)}_S$$

is a finite abelian group. We say that $A_S(G)$ is the defect of weak approximation for G in S.

1.2. In the number field case, Sansuc computed $A_S(G)$ when G is semisimple, or, more generally, when G admits a *special covering*, that is, G fits into a short exact sequence of special kind

$$(1.3) 1 \to B \to G' \to G \to 1.$$

Here G' is the product of a simply connected semisimple K-group and a quasi-trivial K-torus, and B is a finite abelian K-group. Namely, Sansuc constructed an isomorphism

$$(1.4) A_S(G) \xrightarrow{\sim} \operatorname{H}^1_S(B) := \operatorname{coker} \left[H^1(K, B) \to \prod_{v \in S} H^1(K_v, B) \right].$$

Note that there exist reductive K-groups not admitting a special covering.

Date: July 9, 2024.

²⁰²⁰ Mathematics Subject Classification. 11E72, 20G10, 20G20, 20G25, 20G30.

Key words and phrases. Weak approximation, Galois cohomology, reductive groups, local fields, global fields.

- **1.5.** In the number field case, Colliot-Thélène [CT08, Theorem 9.4(i)] computed the finite group $A(G) := \varprojlim_S A_S(G)$ for any reductive K-group G. He considered a flasque resolution of G, that is, a short exact sequence (1.3) in which G' is a quasi-trivial reductive K-group (see [CT08, §2] or Definition 2.4 below) and B is a flasque K-torus (see [CT08, 0.8]). Colliot-Thélène computed A(G) by a formula similar to (1.4).
- **1.6.** In this paper we compute the defect of weak approximation $A_S(G)$ for a reductive K-group G in terms of the algebraic fundamental group $\pi_1^{\text{alg}}(G)$ introduced in [Bor98, Section 1] (and also by Merkurjev [Mer98] and Colliot-Thélène [CT08]). Let $G^{\text{ss}} = [G, G]$ denote the derived group of G, which is semisimple, and let G^{sc} denote the universal cover of G^{ss} , which is simply connected; see [Brl-T72, Proposition (2:24)(ii)] or [CGP15, Corollary A.4.11]. Consider the composite homomorphism

$$\rho \colon G^{\mathrm{sc}} \twoheadrightarrow G^{\mathrm{ss}} \hookrightarrow G$$
,

which is in general is neither injective nor surjective. For a maximal torus $T \subseteq G$, we denote

$$T^{\mathrm{sc}} = \rho^{-1}(T) \subseteq G^{\mathrm{sc}}.$$

Following [Bor98], we consider the algebraic fundamental group of G defined by

$$\pi_1^{\mathrm{alg}}(G) = \mathsf{X}_*(T)/\rho_*\mathsf{X}_*(T^{\mathrm{sc}})$$

where X_* denotes the cocharacter group. The absolute Galois group $\operatorname{Gal}(K^s/K)$ naturally acts on $\pi_1^{\operatorname{alg}}(G)$, and the Galois module $\pi_1^{\operatorname{alg}}(G)$ is well defined (does not depend on the choice of T up to a transitive system of isomorphisms); see [Bor98, Lemma 1.2]. Note that when G = T is a torus, we have $\pi_1^{\operatorname{alg}}(G) = X_*(T)$.

Write $M = \pi_1^{\mathrm{alg}}(G)$. Let $T \subseteq G$ be a maximal torus. Choose a finite Galois extension L/K in K^s splitting T. Set $\Gamma = \mathrm{Gal}(L/K)$; then the finite group Γ naturally acts on M. Moreover, Γ naturally acts on L and on its set of places \mathcal{V}_L . We compute $A_S(G)$ in terms of the Γ -module M and the Γ -set \mathcal{V}_L .

1.7. Following Tate [Ta66], we consider the group of finite formal linear combinations

$$M[\mathcal{V}_L] = \left\{ \sum_{w \in \mathcal{V}_L} m_w \cdot w \mid m_w \in M \right\}$$

and its subgroup

$$M[\mathcal{V}_L]_0 = \Big\{ \sum m_w \cdot w \in M[\mathcal{V}_L] \mid \sum m_w = 0 \Big\}.$$

Then Γ naturally acts on the groups $M[\mathcal{V}_L]$ and $M[\mathcal{V}_L]_0$. We prove the following theorem.

Theorem 1.8 (Theorem 5.7).

$$A_S(G) \cong \operatorname{coker} \left[H_1(\Gamma, M[\mathcal{V}_L]_0) \to \bigoplus_{v \in S} H_1(\Gamma_w, M) \right]$$

where H_1 denotes group homology. Here for $v \in \mathcal{V}_K$, we choose a place $w \in \mathcal{V}_L$ over v, and we denote by Γ_w the corresponding decomposition group (the stabilizer of w in Γ).

The plan of the rest of this paper is as follows. In Section 2 we prove the Hasse principle and weak approximation for the *quasi-trivial groups* introduced by Colliot-Thélène, Moreover, we introduce L/K-free groups, which are special cases of quasi-trivial groups. In Section 3, when G is the quotient G = G'/B of a quasi-trivial group G' by a smooth central subgroup B, following ideas of Sansuc and Colliot-Thélène, we construct an isomorphism

$$A_S(G) \xrightarrow{\sim} \mathsf{H}^1_S(B).$$

In Section 4, using a result of Tate [Ta66], we compute $\mathsf{H}^1_S(B)$ in the case when B is a K-torus, in terms of the cocharacter group $Y=\mathsf{X}_*(B)$. In Section 5 we consider an L/K-free resolution of G, that is, a short exact sequence

$$1 \to B \to G' \to G \to 1$$

where B is a K-torus and G' is an L/K-free reductive group. We prove our theorems computing the defect of weak approximation $A_S(G)$ in terms of $Y = \mathsf{X}_*(B)$ and in terms of $M = \pi_1^{\mathrm{alg}}(G)$.

2. Quasi-trivial groups and L/K-free groups

2.1. A torus T over a field K is called *quasi-trivial* if its cocharacter group $\mathsf{X}_*(T)$ admits a $\mathsf{Gal}(K^s/K)$ -stable basis. Then by Shapiro's lemma and Hilbert's Theorem 90 we have $H^1(K,T)=1$. If K is a global field, then T has the weak approximation property (because T is a K-rational variety).

Theorem 2.2. Let G be a simply connected semisimple group over a global field K, and let $\mathcal{V}_f(K)$ and $\mathcal{V}_{\infty}(K)$ denote the sets of finite (non-archimedean) and infinite (archimedean) places of K, respectively. Then:

- (i) For all $v \in \mathcal{V}_f(K)$ we have $H^1(K_v, G) = 1$.
- (ii) The localization map

$$\eta \colon H^1(K,G) \to \prod_{v \in \mathcal{V}(K)} H^1(K_v,G) = \prod_{v \in \mathcal{V}_{\infty}(K)} H^1(K_v,G)$$

is bijective.

Proof. Assertion (i) is a theorem of Kneser and of Bruhat and Tits; see Platonov and Rapinchuk [PR94, Theorem 6.4, p. 284] for the number field case, and Bruhat and Tits [BT87, Theorem 4.7(ii)] for the general case. Assertion (ii) is the celebrated Hasse principle of Kneser, Harder, and Chernousov; see Platonov and Rapinchuk [PR94, Theorem 6.6, p. 286] for the number field case, and Harder [Har75] for the function field case. □

Theorem 2.3. Let G be a simply connected semisimple group over a global field K. Then G has the weak approximation property, that is, for any finite subset $S \subset \mathcal{V}(K)$, the group G(K) is dense in $G(K_S) := \prod_{v \in S} G(K_v)$.

Indeed, G has the weak approximation property because it has the strong approximation property. This was proved by Platonov in characteristic 0 (see Platonov and Rapinchuk [PR94, Theorem 7.12, p. 427]), and by G. Prasad [Pra77] in positive characteristic.

Definition 2.4 (Colliot-Thélène [CT08, Definition 2.1]). A reductive group G over a field K is quasi-trivial if its commutator subgroup G^{ss} is simply connected and the quotient torus $G^{tor} := G/G^{tor}$ is a quasi-trivial torus. In other words, G is quasi-trivial if it fits into the exact sequence

$$(2.5) 1 \to G^{\text{sc}} \to G \to G^{\text{tor}} \to 1.$$

where G^{sc} is a simply connected semisimple group, and G^{tor} is a quasi-trivial torus.

Proposition 2.6 (Colliot-Thélène [CT08, Proposition 9.2(i,iii)] in the number field case). Let G be quasi-trivial group over a global field K. Then:

(i) For any $v \in \mathcal{V}_f(K)$ we have $H^1(K_v, G) = 1$.

(ii) The localization map

$$loc_{\infty} \colon H^1(K,G) \to \prod_{v \in \mathcal{V}_{\infty}(K)} H^1(K_v,G)$$

is bijective.

Proof. Concerning (i), from (2.5) we obtain a cohomology exact sequence

$$H^1(K_v, G^{\mathrm{sc}}) \to H^1(K_v, G) \to H^1(K_v, G^{\mathrm{tor}})$$

where $H^1(K_v, G^{\text{tor}})$ is trivial because G^{tor} is a quasi-trivial torus, and $H^1(K_v, G^{\text{sc}})$ is trivial by Theorem 2.2(i). Thus $H^1(K_v, G)$ is trivial.

Concerning (ii), for the case of a number field see [CT08, Proposition 9.2(iii)]. In the case of a function field, from (2.5) we obtain a cohomology exact sequence

$$H^1(K, G^{\mathrm{sc}}) \to H^1(K, G) \to H^1(K, G^{\mathrm{tor}})$$

where $H^1(K, G^{\text{tor}})$ is trivial because G^{tor} is a quasi-trivial torus, and $H^1(K, G^{\text{sc}})$ is trivial in the function field case by Theorem 2.2(ii). Thus $H^1(K, G)$ is trivial, as desired.

Proposition 2.7 (Colliot-Thélène [CT08, Proposition 9.2(ii)] in the number field case). Let G be a quasi-trivial group over a global field K. Then G has the weak approximation property, that is, for any finite subset $S \subset \mathcal{V}(K)$, the group G(K) is dense in $G(K_S)$.

Proof. The proof of [CT08, Proposition 9.2(ii)] for the group A(G) (see Subsection 1.5) in the number field case immediately extends to $A_S(G)$ and to the function field case. However, since Colliot-Theélène's style is concise, we provide a detailed proof here.

If $S' \supseteq S$ are finite subsets of $\mathcal{V}(K)$ and if G has weak approximation property $(WA_{S'})$, then it has (WA_S) . Therefore, we may and shall assume that $S \supseteq \mathcal{V}_{\infty}(K)$.

Let $\mathcal{U} \subseteq G(K_S)$ be an open subset. We show that $G(K) \cap \mathcal{U} \neq \emptyset$. Consider the short exact sequence

$$1 \to G^{\mathrm{sc}} \to G \xrightarrow{\tau} G^{\mathrm{tor}} \to 1$$

and the induced commutative diagram with exact rows

$$G^{\operatorname{sc}}(K) \longrightarrow G(K) \stackrel{\tau}{\longrightarrow} G^{\operatorname{tor}}(K) \longrightarrow H^{1}(K, G^{\operatorname{sc}})$$

$$\downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow \sim$$

$$G^{\operatorname{sc}}(K_{S}) \longrightarrow G(K_{S}) \stackrel{\tau_{S}}{\longrightarrow} G^{\operatorname{tor}}(K_{S}) \longrightarrow H^{1}(K_{S}, G^{\operatorname{sc}})$$

Since the homomorphism $\tau: G \to G^{\text{tor}}$ is surjective, the induced map $\tau_S: G(K_S) \to G^{\text{tor}}(K_S)$ is open. Set $\mathcal{U}^{\text{tor}} = \tau_S(\mathcal{U}) \subseteq G^{\text{tor}}(K_S)$, which is open. By (WA_S) for the quasitrivial torus G^{tor} , there exists an element $g^{\text{tor}} \in G^{\text{tor}}(K) \cap \mathcal{U}^{\text{tor}}$. Since the image of \mathcal{U}^{tor} in $H^1(K_S, G^{\text{sc}})$ is $\{1\}$, and the localization map $H^1(K, G^{\text{sc}}) \to H^1(K_S, G^{\text{sc}})$ is injective, we see that the image of g^{tor} in $H^1(K, G^{\text{sc}})$ is 1, and therefore $g^{\text{tor}} = \tau(g)$ for some $g \in G(K)$. Then $\tau(g) \in \tau_S(\mathcal{U})$, whence

$$\tau(g) = \tau_S(u)$$
 for some $u \in \mathcal{U}$,

and hence u = qc for some $c \in G^{sc}(K_S)$. We have $c = q^{-1}u \in q^{-1}\mathcal{U}$, and therefore

$$\mathcal{U}^{\mathrm{sc}} := G^{\mathrm{sc}}(K_S) \cap g^{-1}\mathcal{U} \neq \varnothing.$$

By (WA_S) for G^{sc} , there exists $g^{sc} \in G^{sc}(K) \cap \mathcal{U}^{sc}$. Then

$$g \cdot g^{\mathrm{sc}} \in G(K) \cap \mathcal{U},$$

as desired. \Box

Definition 2.8. Let T be a torus over a field K, and let L/K be a finite Galois extension. We say that T is L/K-free if T splits over L and the $\mathbb{Z}[\operatorname{Gal}(L/K)]$ -module $\mathsf{X}_*(T)$ is free. Alternatively, T is L/K-free if it is isomorphic to $(R_{L/K}\mathbb{G}_{\mathrm{m},L})^{n_T}$ for some integer $n_T \geq 0$, where $\mathbb{G}_{\mathrm{m},L}$ is the multiplicative group over L and $R_{L/K}$ denotes the Weil restriction of scalars.

Observe that any L/K-free torus is quasi-trivial.

Definition 2.9. Let G be a reductive K-group with derived group $G^{ss} = [G, G]$. Write $G^{tor} = G/G^{ss}$. We say that G is L/K-free if G^{ss} is simply connected and the K-torus G^{tor} is L/K-free.

Observe that any L/K-free reductive K-group is quasi-trivial.

Definition 2.10. Let G be a reductive K-group. An L/K-free resolution of G is a short exact sequence of K-groups

$$(2.11) 1 \to B \to G' \to G \to 1$$

where G' is an L/K-free reductive K-group and B is a K-torus.

Proposition 2.12. Any reductive K-group G that splits over a finite Galois extension L/K, admits an L/K-free resolution.

Proof. We follow the proof of [CT08, Proposition-Definition 3.1] very closely. Since G splits over L, it has a maximal torus T that splits over L. Let Z denote the identity component of the center of G (that is, the radical of G); this is a K-torus splitting over L, because it is a subtorus of the torus T splitting over L. There exists a surjective homomorphism of K-tori $Q \to Z$ with Q being L/K-free. Let $T^{\rm sc} \subset G^{\rm sc}$ denote the preimage of T in $G^{\rm sc}$. Then $T^{\rm sc}$ is isogenous to the K-torus $T^{\rm ss} := T \cap G^{\rm ss}$. We see that both $T^{\rm ss}$ and $T^{\rm sc}$ split over L. We have a natural surjective K-homomorphism $G^{\rm sc} \times Q \to G$. Let Z' denote the kernel of this homomorphism; then Z' is the kernel of the surjective homomorphism $T^{\rm sc} \times Q \to T$, and therefore, Z' is a group of multiplicative type over K, not necessary smooth. It follows that we have a short exact sequence of character groups

$$0 \to \mathsf{X}^*(T) \to \mathsf{X}^*(T^{\mathrm{sc}}) \oplus \mathsf{X}^*(Q) \to \mathsf{X}^*(Z') \to 0;$$


see Milne [Mi17, Theorem 12.9]. We see that the absolute Galois group $Gal(K^s/K)$ acts on the character group $X^*(Z')$ via Gal(L/K).

By Lemma 2.13 below, we can find a short exact sequence

$$1 \to Z' \to B \to F \to 1$$

where B is a K-torus that splits over L, and F is an L/K-free K-torus. The diagonal map defines an embedding of Z' into the product $(G^{\text{sc}} \times Q) \times B$ with central image. Let G' be the quotient of the reductive group $(G^{\text{sc}} \times Q) \times B$ by Z'. Then G' is a reductive K-group, which fits into the following commutative diagram with exact rows and columns

(where we replace the given arrow $Z' \to B$ by its inverse):



The quotient of the group G' by the normal subgroup $G^{\operatorname{sc}} \times 1 \subset G^{\operatorname{sc}} \times Q \subset G'$ is a K-group extension of the K-torus F by the K-torus Q. By [CT08, Section 0.7] such a K-group is a K-torus. Since any extension of an L/K-free torus by an L/K-free torus is a split extension (as one can see on the level of cocharacter groups), we conclude that the quotient of G' by G^{sc} is an L/K-free torus. The derived group G'^{ss} can be identified with G^{sc} , and the group G'^{tor} is an L/K-free torus. Thus the reductive group G' is L/K-free, and the middle row of the diagram is a desired L/K-free resolution of G.

Lemma 2.13. Let Γ be a finite group and M be a finitely generated Γ -module. Then there exists a resolution

$$(2.14) 0 \to M^{-1} \to M^0 \xrightarrow{\phi} M \to 0$$

where both M^{-1} and M^0 are finitely generated torsion-free Γ -modules and such that in addition we can choose any one of M^0 or M^{-1} to be a free Γ -module.

Proof. If we wish M^0 to be such, the construction is very simple. For the finitely generated $\mathbb{Z}[\Gamma]$ -module M, there is a surjection from a finitely generated $free \mathbb{Z}[\Gamma]$ -module $\mathbb{Z}[\Gamma]^n$ for some n:

$$\phi \colon \mathbb{Z}[\Gamma]^n \to M,$$

and clearly ker ϕ is torsion-free and finitely generated.

If we wish M^{-1} to be a finite rank free $\mathbb{Z}[\Gamma]$ -module, then we choose any resolution

$$0 \to \tilde{M}^{-1} \to \tilde{M}^0 \to M \to 0$$

with Γ -modules \tilde{M}^{-1} and \tilde{M}^0 finitely generated and torsion free, for example, the resolution above. By [BK23, Section 5.1] there exists a quasi-isomorphism of complexes

$$(\tilde{M}^{-1} \to \tilde{M}^0) \to (M^{-1} \to M^0)$$

with M^{-1} and M^0 finitely generated and torsion-free, and M^{-1} being a free $\mathbb{Z}[\Gamma]$ -module. Recall that "quasi-isomorphism" means that this morphism of complexes induces isomorphisms of the kernels and of the cokernels. Thus we obtain a commutative diagram

$$0 \longrightarrow \tilde{M}^{-1} \longrightarrow \tilde{M}^{0} \longrightarrow M \longrightarrow 0$$

$$\simeq \downarrow \qquad \qquad \downarrow \simeq \qquad \qquad \downarrow \simeq$$

$$0 \longrightarrow M^{-1} \longrightarrow M^{0} \longrightarrow M^{1} \longrightarrow 0$$

Identifying M^1 with M, we obtain a desired resolution (2.14) of M, with M^{-1} being $\mathbb{Z}[\Gamma]$ -free.

3.
$$A_S(G)$$
 for a quotient of a quasi-trivial group

Theorem 3.1. Let G be a reductive group over a global field K. Consider a short exact sequence

$$(3.2) 1 \to B \to G' \to G \to 1$$

where G' is a quasi-trivial K-group and $B \subset G'$ is a smooth central K-subgroup. Let S be a finite set of places of K. Then the closure $\overline{G(K)}_S$ of G(K) in $G(K_S)$ is a normal subgroup of finite index, and the connecting homomorphism $G(K_S) \to H^1(K_S, B)$ induces an isomorphism

$$(3.3) A_S(G) \xrightarrow{\sim} \mathsf{H}_S^1(B).$$

Proof. From (3.2) we obtain a commutative diagram with exact rows

$$G'(K) \xrightarrow{\varphi} G(K) \xrightarrow{\varphi} H^{1}(K, B) \xrightarrow{\zeta} H^{1}(K, G')$$

$$\downarrow^{j} \qquad \qquad \downarrow^{i} \qquad \qquad \downarrow^{\eta}$$

$$G'(K_{S}) \xrightarrow{\chi} G(K_{S}) \xrightarrow{\chi} H^{1}(K_{S}, B) \xrightarrow{\xi} H^{1}(K_{S}, G')$$

Here we write $H^1(K_S, B) = \prod_{v \in S} H^1(K_v, B)$ and $H^1(K_S, G') = \prod_{v \in S} H^1(K_v, G')$.

Assuming that $S \subset \mathcal{V}(K)$ is a finite subset containing $V_{\infty}(K)$, we see that the groups, sets, and maps in the above diagram have the following properties:

- (1) The map $\eta: H^1(K, G') \to H^1(K_S, G')$ is bijective (Proposition 2.6);
- (2) The homomorphism $j: G'(K) \to G'(K_S)$ has dense image (Proposition 2.7);
- (3) $H^1(K_S, B)$ is a finite abelian group (Milne [Mil06, Corollary I.2.3]).

Setting $X = \chi^{-1}(\operatorname{im} \theta \varphi)$, we obtain a commutative diagram with exact rows

$$G'(K) \xrightarrow{\varrho} G(K) \xrightarrow{\bar{\varphi}} \operatorname{im} \varphi \longrightarrow 1$$

$$\downarrow j \qquad \qquad \downarrow \bar{i} \qquad \qquad \downarrow \bar{\theta}$$

$$G'(K_S) \xrightarrow{\psi} X \xrightarrow{\bar{\chi}} \operatorname{im} \theta \varphi \longrightarrow 1$$

Since φ and θ are group homomorphisms, we see that im $\theta\varphi$ is a subgroup of the finite abelian group $H^1(K_S, B)$, and therefore X is an open normal subgroup of $G(K_S)$. Thus X is a closed subgroup of $G(K_S)$. Clearly, $X \supset i(G(K))$, and so X contains the closure of i(G(K)) in $G(K_S)$. We show that X coincides with this closure, that is, i(G(K)) is dense in X.

Let $\mathcal{U} \subset X$ be a non-empty open subset. Let $u \in \mathcal{U}$. Then by the construction of X, we have $\chi(u) = \theta(\varphi(g))$ for some $g \in G(K)$, whence $\bar{\chi}(u) = \bar{\chi}(\bar{i}(g))$. Then $\bar{\chi}(\bar{i}(g)^{-1} \cdot u) = 1$, whence $\bar{i}(g)^{-1} \cdot u \in \text{im } \psi$, and $\psi^{-1}(\bar{i}(g)^{-1} \cdot u)$ is non-empty. Thus $\psi^{-1}(\bar{i}(g)^{-1} \cdot \mathcal{U})$ is a non-empty open subset of $G'(K_S)$. Since G' has the weak approximation property, there exists $g' \in G'(K)$ with

$$j(g') \in \psi^{-1}(\overline{i}(g)^{-1} \cdot \mathcal{U}).$$

Thus we have found an element $g' \in G'(K)$ such that

$$\bar{i}(g \cdot \varrho(g')) = \bar{i}(g) \cdot \bar{i}(\varrho(g')) = \bar{i}(g) \cdot \psi(j(g')) \in \bar{i}(g) \cdot \left(\bar{i}(g)^{-1} \cdot \mathcal{U}\right) = \mathcal{U}.$$

Set $g_1 = g \cdot \varrho(g') \in G(K)$; then $\bar{i}(g_1) \in \mathcal{U}$. Thus $\bar{i}(G(K))$ is dense in X, and therefore X is the closure of i(G(K)) in $G(K_S)$. Hence this closure is a normal subgroup of $G(K_S)$, which gives a natural group structure on $A_S(G)$, and moreover, this group $A_S(G)$ is a finite abelian group equal to the quotient of im χ by $\theta(\operatorname{im} \varphi)$.

Consider the commutative diagram with exact rows

$$1 \longrightarrow \operatorname{im} \varphi \longrightarrow H^{1}(K, B) \longrightarrow \operatorname{im} \zeta \longrightarrow 1$$

$$\downarrow_{\bar{\theta}} \qquad \qquad \downarrow_{\nu}$$

$$1 \longrightarrow \operatorname{im} \chi \longrightarrow H^{1}(K_{S}, B) \longrightarrow \operatorname{im} \xi \longrightarrow 1$$

We show that the arrow ν is bijective. Since G' is quasi-trivial, by Proposition 2.6(i) we have $H^1(K_v, G') = 1$ for all $v \in \mathcal{V}_f(K)$. Therefore, it suffices to show that ν is surjective when $S = \mathcal{V}_{\infty}(K)$. For this S, the arrow θ is surjective by [San81, Lemma 1.8], and we see from the diagram that ν is surjective as well. Note that the arrow ν is injective by the Hasse principle for G' (Proposition 2.6(ii)). Thus ν is indeed bijective.

The homomorphism $\chi \colon G(K_S) \to H^1(K_S, B)$ induces a homomorphism $\delta \colon A_S(G) \to \mathsf{U}_S^1(B)$. Since the arrow ν is bijective, by the snake lemma the homomorphism coker $\bar{\theta} \to \mathrm{coker} \, \theta$ is bijective. It follows that the homomorphism δ is surjective.

We show that the homomorphism δ is injective. Let $[g] = g \cdot \overline{G(K)}_S \in A_S(G)$ be such that $\delta[g] = 1$, that is, $\chi(g) \in \operatorname{im} \theta$. Write

$$\chi(g) = \theta(b), \quad b \in H^1(K, B).$$

The image of $\chi(g)$ in $\operatorname{im} \xi \subseteq H^1(K_S, G')$ is 1. Since the homomorphism ν is bijective, we see that the image of b in $\operatorname{im} \zeta \subseteq H^1(K, G')$ is 1. It follows that $b \in \operatorname{im} \varphi$ and

$$g \in \chi^{-1}(\operatorname{im}\theta\varphi) = X.$$

Since $X = \overline{G(K)}_S$, we see that [g] = 1 and the homomorphism δ is indeed injective. Thus δ is an isomorphism, as desired.

Remark 3.4. Sansuc [San81, Theorem 3.3] considers a resolution (3.2) in the case when K is a number field and B is a *finite abelian* K-group. Moreover, he assumes that the resolution (3.2) is a *special covering*, that is, that G' is a quasi-trivial K-group of a special kind: $G' = G^{\text{sc}} \times T'$ where T' is a quasi-trivial K-torus. In this case, Sansuc shows that (3.2) induces an isomorphism (3.3). Note that there exist reductive groups admitting no special coverings.

Moreover, Colliot-Thélène [CT08, Theorem 9.4(i)] considers a flasque resolution (3.2) of G when K is a number field. This means that G' is quasi-trivial and B is a flasque torus. By [CT08, Proposition-Definition 3.1], any reductive K-group G admits a flasque resolution. Colliot-Theélène shows that a flasque resolution (3.2) induces an isomorphism (3.3).

Our Theorem 3.1 generalizes [San81, Theorem 3.3] and [CT08, Theorem 9.4(i)]. Sansuc's proof (inspired by Kneser [Kne62]) immediately generalizes to our case. In our proof of Theorem 3.1, we reproduce Sansuc's proof (with details added) using his notations.

4. Computing
$$\mathsf{H}^1_S(B)$$
 for a torus B

Let B be a torus over a global field K with cocharacter group $Y = X_*(B)$. Let L/K be a finite Galois extension splitting T. We write $\Gamma = \text{Gal}(L/K)$. In this section we compute $\mathsf{U}_S^1(B)$ for a finite set S of places of K in terms of the Γ -module Y. We need two lemmas.

Lemma 4.1 (probably well-known). Let Δ be a finite group and A be a Δ -module. If A is torsion-free, then

$$H^{-1}(\Delta, A) = A_{\Delta, Tors}.$$

Proof. By definition,

$$H^{-1}(\Delta, A) = \ker \left[\operatorname{Nm} \colon A_{\Delta} \to A^{\Delta} \right].$$

The group $H^{-1}(\Delta, A)$ is killed by $\#\Delta$ (see [CF86, Section IV.6, Corollary 1 of Proposition 8]), whence $H^{-1}(\Delta, A) \subseteq A_{\Delta,\text{Tors}}$. Since A is torsion-free, so is A^{Δ} , whence

$$H^{-1}(\Delta, A) = \ker \operatorname{Nm} \supseteq A_{\Delta, \operatorname{Tors}}.$$

Thus $H^{-1}(\Delta, A) = A_{\Delta, \text{Tors}}$.

Lemma 4.2. For any place v of K choose a place w of L over v, and denote by $\Gamma_w \subseteq \Gamma$ the corresponding decomposition group (the stabilizer of w in Γ). Then we have canonical isomorphisms

$$H^1(K_v, B) \xrightarrow{\sim} Y_{\Gamma_w, \text{Tors}}, \quad H^1(K, B) \xrightarrow{\sim} (Y[\mathcal{V}_L]_0)_{\Gamma, \text{Tors}}.$$

Proof. We have $H^1(K_v, B) = H^1(L_w/K_v, B)$; see [San81, (1.9.2)]. We have the Tate-Nakayama isomorphism

$$H^{-1}(\Gamma_w, Y) \xrightarrow{\sim} H^1(\Gamma_w, Y \otimes_{\mathbb{Z}} L_w^{\times}) = H^1(L_w/K_v, B);$$

see [Ta66, Theorem on page 717]. By Lemma 4.1, we have $H^{-1}(\Gamma_w, Y) = Y_{\Gamma_w, \text{Tors}}$, whence $H^1(L_w/K_v, B) \cong Y_{\Gamma_w, \text{Tors}}$.

Similarly, we have $H^1(K,B)=H^1(L/K,B);$ see [San81, (1.9.2)]. We have the Tate isomorphism

$$H^{-1}(\Gamma, Y[\mathcal{V}_L]_0) \xrightarrow{\sim} H^1(L/K, B);$$

see [Ta66, Theorem on page 717]. Since $Y[\mathcal{V}_L]_0$ is torsion-free, we conclude by Lemma 4.1 that

$$H^1(L/K, B) \cong H^{-1}(\Gamma, Y[\mathcal{V}_L]_0) = (Y[\mathcal{V}_L]_0)_{\Gamma, \text{Tors}},$$

as desired.

4.3. Consider the groups

$$\Upsilon_S = \bigoplus_{v \in S} Y_{\Gamma_w, \mathrm{Tors}}$$
 and $\Upsilon_{S^{\complement}} = \bigoplus_{v \in S^{\complement}} Y_{\Gamma_w, \mathrm{Tors}}$.

By Lemma 4.2 we have isomorphisms

$$\Upsilon_S = \bigoplus_{v \in S} H^1(K_v, B)$$
 and $\Upsilon_{S^{\complement}} = \bigoplus_{v \in S^{\complement}} H^1(K_v, B)$.

For each $v \in \mathcal{V}_K$, consider the projection homomorphism

$$\pi_v \colon Y_{\Gamma_w, \mathrm{Tors}} \to Y_{\Gamma, \mathrm{Tors}}.$$

We have a natural homomorphism

$$\pi_S \colon \Upsilon_S \to Y_{\Gamma, \text{Tors}}, \ [y_v]_{v \in S} \mapsto \left[\sum_{v \in S} \pi_v(y_v)\right].$$

Similarly, we have a natural homomorphism

$$\pi_{S^{\complement}} \colon \Upsilon_{S^{\complement}} \to Y_{\Gamma, \operatorname{Tors}}.$$

Theorem 4.4. There is a canonical isomorphism

$$\mathsf{Y}^1_S(B) \cong \Upsilon_S / \pi_S^{-1}(\operatorname{im} \pi_{S^{\complement}}).$$

Proof. Consider the following commutative diagram:

$$(4.5) \qquad \stackrel{\sim}{\longrightarrow} H^{-1}(Y[\mathcal{V}_L]_0) \stackrel{\sim}{\longrightarrow} (Y[\mathcal{V}_L]_0)_{\Gamma, \text{Tors}}$$

$$\downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow$$

$$H^1(K_S, B) \stackrel{\sim}{\longrightarrow} \bigoplus_{v \in S} H^{-1}(\Gamma_w, Y) \stackrel{\sim}{\longrightarrow} \Upsilon_S$$

in which the rectangle at left comes from Tate [Ta66, Theorem on page 717]. From (4.5) we obtain a canonical isomorphism

$$\mathrm{H}^1_S(B) \xrightarrow{\sim} \operatorname{coker} \left[\left(Y[\mathcal{V}_L]_0 \right)_{\Gamma, \operatorname{Tors}} \to \Upsilon_S \right].$$

From the short exact sequence

$$0 \to Y[\mathcal{V}_L]_0 \to Y[\mathcal{V}_L] \to Y \to 0$$

we obtain an exact sequence

$$H^{-1}(\Gamma, Y[\mathcal{V}_L]_0) \to \bigoplus_{v \in \mathcal{V}_K} H^{-1}(\Gamma_w, Y) \to H^{-1}(\Gamma, Y),$$

which by Lemma 4.1 gives an exact sequence

$$(Y[\mathcal{V}_L]_0)_{\Gamma,\mathrm{Tors}} \longrightarrow \Upsilon_S \oplus \Upsilon_{S^{\complement}} \xrightarrow{\pi_S + \pi_{S^{\complement}}} Y_{\Gamma,\mathrm{Tors}}.$$

We see that

$$\operatorname{im}\left[\left(Y[\mathcal{V}_L]_0\right)_{\Gamma,\operatorname{Tors}}\to \Upsilon_{\!S}\right]\,=\,\pi_S^{-1}(\operatorname{im}\pi_{S^\complement}),$$

and the theorem follows.

Corollary 4.6 (Sansuc [San81]). Let $S_{nc} \subseteq S$ (resp. $S_c \subseteq S$) denote the subset of places with noncyclic (resp., cyclic) decomposition group in $\Gamma = \operatorname{Gal}(L/K)$. Then the natural epimorphism $\operatorname{H}^1_S(B) \to \operatorname{H}^1_{S_{nc}}(B)$ is an isomorphism.

Proof. By Theorem 4.4, it suffices to show that for any $v \in S_c$, the direct summand $Y_{\Gamma_w, \text{Tors}}$ of Υ_S is contained in $\pi_S^{-1}(\text{im }\pi_{S^\complement})$. Since the decomposition group $\Gamma_w \subset \Gamma$ is cyclic and the set S is finite, by the Chebotarev density theorem, see, for instance, Neukirch [Neu99, Theorem (13.4)], there exist $v' \in S^{\complement}$ and $w' \in \mathcal{V}_L$ over v' with decomposition group $\Gamma_{w'} = \Gamma_w$. Then im $\pi_{w'} = \text{im }\pi_w$, and

$$Y_{\Gamma_w, \text{Tors}} \subseteq \pi_S^{-1}(\text{im } \pi_{S^{\complement}}),$$

as desired. \Box

4.7. Let $CC(\Gamma)$ denote the set of conjugacy classes of subgroups of Γ . We consider the map

$$\varkappa \colon S^{\complement} \to \mathrm{CC}(\Gamma)$$

sending a place $v \in S^{\complement}$ to the conjugacy class of Γ_w for a place w of L over v. We say that the decomposition group Γ_w is maximal if it is not contained is a strictly larger decomposition group $\Gamma_{w'}$ for some $v' \in S^{\complement}$. Let m denote the cardinality of the set of conjugacy classes of maximal decomposition groups Γ_w for $v \in S^{\complement}$. Let $S^{\complement}_{max} \subset S^{\complement}$ be a set of cardinality m such that $\varkappa(S^{\complement}_{max})$ contains (as elements) the conjugacy classes of all maximal decomposition groups. Consider the natural homomorphism

$$\pi_{S_{\max}^{\complement}}: \bigoplus_{v \in S_{\max}^{\complement}} Y_{\Gamma_w, \text{Tors}} \to Y_{\Gamma, \text{Tors}}.$$

Corollary 4.8. There is a canonical isomorphism

$$\mathsf{Y}^1_S(B) \cong \Upsilon_S/\pi_S^{-1}(\operatorname{im} \pi_{S^{\complement}_{\max}}).$$

Proof. Indeed, let $v \in S^{\complement}$; then $\Gamma_w \subseteq \Gamma_{w'}$ for some $v' \in S^{\complement}_{\max}$ and some w' over v'. The natural map

$$\pi_v \colon Y_{\Gamma_w, \mathrm{Tors}} \to Y_{\Gamma, \mathrm{Tors}}$$

factors as follows:

$$\pi_v \colon Y_{\Gamma_w, \text{Tors}} \to Y_{\Gamma_{w', \text{Tors}}} \xrightarrow{\pi_{v'}} Y_{\Gamma, \text{Tors}}.$$

We see that $\operatorname{im} \pi_v \subseteq \operatorname{im} \pi_{v'}$ and therefore

$$\operatorname{im} \pi_{S^{\complement}} = \operatorname{im} \pi_{S^{\complement}_{\max}}.$$

Now the corollary follows from Theorem 4.4.

5.
$$A_S(G)$$
 in terms of $\mathsf{X}_*(B)$ and in terms of $\pi_1^{\mathrm{alg}}(G)$

5.1. Let G be a reductive group over a global field K. Write $M = \pi_1^{\text{alg}}(G)$. Let $T \subseteq G$ be a maximal torus, and let L/K be a finite Galois extension in K^s splitting T; then $\Gamma := \text{Gal}(L/K)$ naturally acts on M.

Choose an L/K-free resolution (2.11) and consider the sequence of Gal(L/K)-modules

$$(5.2) 0 \to Y \to M' \to M \to 0$$

where

$$Y = \pi_1^{\operatorname{alg}}(B) = \mathsf{X}_*(B)$$
 and $M' = \pi_1^{\operatorname{alg}}(G')$.

This sequence is exact; see [CT08, Proposition 6.8] or [BK23, Lemma 6.2.4]. Since Γ acts on Y and on L, it naturally acts on \mathcal{V}_L , on $Y[\mathcal{V}_L]$, and on $Y[\mathcal{V}_L]_0$.

Theorem 5.3. For an L/K-free resolution (2.11) and a finite set of places $S \subset \mathcal{V}_K$, there is a canonical isomorphism

$$A_S(G) \stackrel{\sim}{\longrightarrow} \operatorname{coker} \left[\left(Y[\mathcal{V}_L]_0 \right)_{\Gamma, \operatorname{Tors}} \to \bigoplus_{v \in S} Y_{\Gamma_w, \operatorname{Tors}} \right].$$

Proof. From diagram (4.5) we obtain that

$$\mathsf{H}^1_S(B) \coloneqq \operatorname{coker} \left[H^1(K,B) \to H^1(K_S,B) \right] \cong \operatorname{coker} \left[\left(Y[\mathcal{V}_L]_0 \right)_{\Gamma,\operatorname{Tors}} \to \bigoplus_{v \in S} Y_{\Gamma_w,\operatorname{Tors}} \right],$$

and the theorem follows from Theorem 3.1 giving an isomorphism $A_S(G) \xrightarrow{\sim} \operatorname{H}^1_S(B)$. \square

Theorem 5.4. For an L/K-free resolution (2.11) and a finite set of places $S \subset \mathcal{V}_K$, there is a canonical isomorphism

$$A_S(G) \xrightarrow{\sim} \Upsilon_S/\pi_S^{-1}(\operatorname{im} \pi_{S_{\max}^{\complement}}).$$

Proof. The theorem follows from Theorem 3.1 giving an isomorphism $A_S(G) \xrightarrow{\sim} \mathsf{Y}_S^1(B)$, and Corollary 4.8 giving an isomorphism $\mathsf{Y}_S^1(B) \xrightarrow{\sim} \Upsilon_S/\pi_S^{-1} \big(\operatorname{im} \pi_{S_{\max}^0}\big)$.

Corollary 5.5 (Sansuc [San81]). Assume that G splits over a finite Galois extension L/K, and let $S_{\rm nc} \subseteq S$ denote the subset of places with noncyclic decomposition group in $\Gamma = \operatorname{Gal}(L/K)$. Then the natural epimorphism $A_S(G) \to A_{S_{\rm nc}}(G)$ is an isomorphism.

Proof. Consider the commutative diagram

$$A_{S}(G) \xrightarrow{\sim} \mathsf{Y}_{S}^{1}(B)$$

$$\downarrow \sim$$

$$A_{S_{\mathrm{nc}}}(G) \xrightarrow{\sim} \mathsf{Y}_{S_{\mathrm{nc}}}^{1}(B)$$

in which the horizontal arrows are isomorphisms of Theorem 3.1. By Corollary 4.6 the right-hand vertical arrow is an isomorphism. Thus the left-hand vertical arrow is an isomorphism, as desired.

Corollary 5.6 (Sansuc [San81, Corollary 3.5]).

- (i) If G splits over a finite Galois extension L/K and all places in S have cyclic decomposition group in Gal(L/K), then G has the weak approximation property in
- (ii) G has the weak approximation property in $\mathcal{V}_{\infty}(K)$.

Theorem 5.7. Let G be a reductive group over a global field K, and let M, L, and Γ be as in 5.1. Then there is a canonical isomorphism

$$A_S(G) \xrightarrow{\sim} \operatorname{coker} \left[H_1(\Gamma, M[\mathcal{V}_L]_0) \to \bigoplus_{v \in S} H_1(\Gamma_w, M) \right]$$

where H_1 denotes group homology.

We need a lemma and a corollary.

Lemma 5.8. Let Γ be a finite group, and let N be a finitely generated **free** $\mathbb{Z}[\Gamma]$ -module. Then

- (i) $N_{\Gamma, \text{Tors}} = 0$, and
- (ii) $H_i(\Gamma, N) = 0$ for all i > 1.

Proof. By Shapiro's lemma we can reduce our lemma to the case $\Gamma = 1$ and $M = \mathbb{Z}$, in which assertions (i) and (ii) are obvious.

Corollary 5.9. Let L/K be a finite Galois extension of global fields, and let N be a finitely generated **free** $\mathbb{Z}[\Gamma]$ -module. Then

- (i) $(N[\mathcal{V}_L]_0)_{\Gamma,\text{Tors}} = 0$, and (ii) $H_i(\Gamma, N[\mathcal{V}_L]_0) = 0$ for all $i \ge 1$.

Proof. By [BK23, Theorem A.1.1], the short exact sequence

$$0 \to N[\mathcal{V}_L]_0 \to N[\mathcal{V}_L] \to N \to 0$$

gives rise to a cohomology exact sequence

$$(5.10) \quad \cdots \to H_2(\Gamma, N[\mathcal{V}_L]_0) \to H_2(\Gamma, N[\mathcal{V}_L]) \to H_2(\Gamma, N)$$

$$\to H_1(\Gamma, N[\mathcal{V}_L]_0) \to H_1(\Gamma, N[\mathcal{V}_L]) \to H_1(\Gamma, N)$$

$$\to \left(N[\mathcal{V}_L]_0\right)_{\Gamma \text{ Tors}} \to \left(N[\mathcal{V}_L]\right)_{\Gamma \text{ Tors}} \to N_{\Gamma, \text{Tors}}$$

By Lemma 5.8 we have $H_i(\Gamma, N) = 0$ for all $i \ge 1$. Moreover, we have canonical isomorphisms

$$H_i(\Gamma, N[\mathcal{V}_L] \cong \bigoplus_{v \in \mathcal{V}_K} H_i(\Gamma_w, N) = 0 \text{ for all } i \geq 1,$$

$$(N[\mathcal{V}_L])_{\Gamma, \text{Tors}} \cong \bigoplus_{v \in \mathcal{V}_K} N_{\Gamma_w, \text{Tors}} = 0$$

by Shapiro's lemma and Lemma 5.8 (because the $\mathbb{Z}[\Gamma]$ -free module N is $\mathbb{Z}[\Gamma_w]$ -free). Now the corollary follows from the exactness of (5.10).

Proof of Theorem 5.7. For an L/K-free resolution (2.11), the short exact sequence of Gal(L/K)-modules (5.2) induces a short exact sequence

$$0 \to Y[\mathcal{V}_L]_0 \to M'[\mathcal{V}_L]_0 \to M[\mathcal{V}_L]_0 \to 0$$

and a commutative diagram with exact rows

$$0 = H_{1}(\Gamma, M'[\mathcal{V}_{L}]_{0}) \longrightarrow H_{1}(\Gamma, M[\mathcal{V}_{L}]_{0}) \longrightarrow (Y[V_{L}]_{0})_{\Gamma, \text{Tors}} \longrightarrow (M'[V_{L}]_{0})_{\Gamma, \text{Tors}} = 0$$

$$\downarrow l_{1} \qquad \qquad \downarrow l_{0}$$

$$0 = \bigoplus_{v \in S} H_{1}(\Gamma_{w}, M') \longrightarrow \bigoplus_{v \in S} H_{1}(\Gamma_{w}, M) \longrightarrow \bigoplus_{v \in S} Y_{\Gamma_{w}, \text{Tors}} \longrightarrow \bigoplus_{v \in S} (M')_{\Gamma_{w}, \text{Tors}} = 0$$

In this diagram, the zeros in the bottom row are explained by Lemma 5.8, and the zeros in the top row are explained by Corollary 5.9. The diagram induces an isomorphism coker $l_1 \xrightarrow{\sim}$ coker l_0 . By Theorem 5.3 we have an isomorphism $A_S(G) \cong \operatorname{coker} l_0$, which gives a desired isomorphism $A_S(G) \cong \operatorname{coker} l_1$.

5.11. Consider the groups

$$\Xi_S = \bigoplus_{v \in S} H_1(\Gamma_w, M)$$
 and $\Xi_{S^{\complement}} = \bigoplus_{v \in S^{\complement}} H_1(\Gamma_w, M)$.

For $v \in \mathcal{V}_K$ consider the natural homomorphism

$$\tau_v \colon H_1(\Gamma_w, M) \to H_1(\Gamma, M)$$

We have a natural homomorphism

$$\tau_S \colon \Xi_S \to H_1(\Gamma, M), \ \left[\xi_v\right]_{v \in S} \mapsto \left[\sum_{v \in S} \tau_v(\xi_v)\right].$$

Similarly, we have a natural homomorphism

$$\tau_{S^{\complement}} \colon \Xi_{S^{\complement}} \to H_1(\Gamma, M).$$

Let $S_{\max}^{\complement} \subset S^{\complement}$ be a finite subset as in Subsection 4.7. Consider the natural homomorphism

$$\tau_{S_{\max}^{\complement}} \colon \Xi_{S_{\max}^{\complement}} \to H_1(\Gamma, M).$$

Theorem 5.12. There is a canonical isomorphism

$$A_S(G) \cong \Xi_S/\tau_S^{-1}(\operatorname{im} \tau_{S_{\max}^{\complement}}).$$

Proof. For each $v \in \mathcal{V}_K$, the short exact sequence (5.2) gives rise to a commutative diagram with exact rows

$$0 = H_1(\Gamma_w, M') \longrightarrow H_1(\Gamma_w, M) \longrightarrow Y_{\Gamma_w, \text{Tors}} \longrightarrow (M')_{\Gamma_w, \text{Tors}} = 0$$

$$\tau_v \downarrow \qquad \qquad \downarrow \pi_v$$

$$0 = H_1(\Gamma, M') \longrightarrow H_1(\Gamma, M) \longrightarrow Y_{\Gamma, \text{Tors}} \longrightarrow (M')_{\Gamma, \text{Tors}} = 0$$

In this diagram, the zeros are explained by Lemma 5.8. Now the theorem follows from Theorem 5.4 giving an isomorphism $A_S(G) \xrightarrow{\sim} \Upsilon_S/\pi_S^{-1}(\operatorname{im} \pi_{S_{\operatorname{conv}}^{\mathbb{C}}})$, and the isomorphisms

$$H_1(\Gamma_w, M) \xrightarrow{\sim} Y_{\Gamma_w, \text{Tors}}, \quad H_1(\Gamma, M) \xrightarrow{\sim} Y_{\Gamma, \text{Tors}}$$

coming from the above diagram.

ACKNOWLEDGEMENTS: The author is grateful to Dipendra Prasad for his suggestion to compute the defect of weak approximation for a reductive group. We thank Jean-Louis Colliot-Thélène and Boris Kunyavskiĭ for helpful discussions and email correspondence. This paper was partly written during the author's visit to Max Planck Institute for Mathematics, Bonn, and the author is grateful to the Institute for the hospitality, support, and excellent working conditions.

References

- [Brl-T72] Armand Borel and Jacques Tits, Compléments à l'article: "Groupes réductifs", Inst. Hautes Études Sci. Publ. Math. 41, 253–276 (1972).
- [Bor98] Mikhail Borovoi, Abelian Galois cohomology of reductive groups, Mem. Amer. Math. Soc. 132 (1998), no. 626.
- [BK23] Mikhail Borovoi and Tasho Kaletha, Galois cohomology of reductive groups over global fields, arXiv:2303.04120.
- [BT87] François Bruhat and Jacques Tits, Groupes algébriques sur un corps local. Chapitre III. Compléments et applications à la cohomologie galoisienne, J. Fac. Sci. Univ. Tokyo Sect. IA Math. 34 (1987), no. 3, 671–698.
- [CF86] J. W. S. Cassels and Albrecht Fröhlich (eds.), Algebraic number theory, Academic Press, Inc., London, 1986, Reprint of the 1967 original.
- [CT08] Jean-Louis Colliot-Thélène, Résolutions flasques des groupes linéaires connexes, J. Reine Angew. Math. 618 (2008), 77–133.
- [CGP15] Brian Conrad, Ofer Gabber, and Gopal Prasad, *Pseudo-reductive groups*, 2nd edition. New Mathematical Monographs, 26. Cambridge University Press, Cambridge (2015).
- [Har75] Günter Harder, Über die Galoiskohomologie halbeinfacher algebraischer Gruppen. III, J. Reine Angew. Math. **274/275** (1975), 125–138.
- [Kne62] Martin Kneser, Schwache Approximation in algebraischen Gruppen, in: Colloque sur la théorie des groupes algébriques, Centre Belge rech. Math., Bruxelles, 1962, pp. 41-52.
- [Mer98] Alexander S. Merkurjev, *K-theory and algebraic groups*, European Congress of Mathematics, Vol. II (Budapest, 1996), Progr. Math., vol. 169, Birkhäuser, Basel, 1998, pp. 43–72.
- [Mil06] James S. Milne, Arithmetic duality theorems, Second edition, BookSurge, LLC, Charleston, SC, 2006.
- [Mi17] James S. Milne, Algebraic groups. The theory of group schemes of finite type over a field, Cambridge Stud. Adv. Math., 170 Cambridge University Press, Cambridge, 2017.
- [Neu99] Jürgen Neukirch, Algebraic number theory, Grundlehren Math. Wiss., 322, Springer-Verlag, Berlin, 1999.
- [PR94] Vladimir Platonov and Andrei Rapinchuk, Algebraic groups and number theory, Academic Press, 1994.

- [Pra77] Gopal Prasad, Strong approximation for semi-simple groups over function fields, Ann. of Math. (2) **105** (1977), no. 3, 553–572.
- [San81] Jean-Jacques Sansuc, Groupe de Brauer et arithmétique des groupes algébriques linéaires sur un corps de nombres, J. Reine Angew. Math. **327** (1981), 12–80.
- [Ta66] John Tate, The cohomology groups of tori in finite Galois extensions of number fields, Nagoya Math. J. 27 (1966), 709–719.

Raymond and Beverly Sackler School of Mathematical Sciences, Tel Aviv University, Ramat Aviv, 6997801 Tel Aviv, Israel

 $Email\ address: {\tt borovoi@tauex.tau.ac.il}$