Null hypothesis Bayes factor estimates can be biased in (some) common factorial designs: A simulation study

Daniel J. Schad
1&Shravan Vasishth²

¹ Institute for Mind, Brain and Behavior, HMU Health and Medical University, Potsdam, Germany

² University of Potsdam, Potsdam, Germany

Author Note

Correspondence concerning this article should be addressed to Daniel J. Schad, HMU Health and Medical University, Potsdam, Germany. E-mail: danieljschad@gmail.com. A preprint of this manuscript has been published on arXiv (https://arxiv.org/???). Reproducible code is available from https://osf.io/3g86r/. This study was not preregistered.

Abstract

Bayes factor null hypothesis tests provide a viable alternative to frequentist measures of evidence quantification. Bayes factors for realistic interesting models cannot be calculated exactly, but have to be estimated, which involves approximations to complex integrals. Crucially, the accuracy of these estimates, i.e., whether an estimated Bayes factor corresponds to the true Bayes factor, is unknown, and may depend on data, prior, and likelihood. We have recently developed a novel statistical procedure, namely simulation-based calibration (SBC) for Bayes factors, to test for a given analysis, whether the computed Bayes factors are accurate. Here, we use SBC for Bayes factors to test for some common cognitive designs, whether Bayes factors are estimated accurately. We use the bridgesampling/brms packages as well as the BayesFactor package in R. We find that Bayes factor estimates are accurate and exhibit only little bias in Latin square designs with (a) random effects for subjects only and (b) for crossed random effects for subjects and items, but a single fixed-factor. However, Bayes factor estimates turn out biased and liberal in a 2x2 design with crossed random effects for subjects and items. These results suggest that researchers should test for their individual analysis, whether Bayes factor estimates are accurate. Moreover, future research is needed to determine the boundary conditions under which Bayes factor estimates are accurate or biased, as well as software development to improve estimation accuracy.

Keywords: Bayes factors, Bayesian model comparison, simulation-based calibration, null hypothesis testing, crossed random effects for subjects and items

Null hypothesis Bayes factor estimates can be biased in (some) common factorial designs: A simulation study

Introduction

Bayes factors have emerged as a popular tool for quantifying evidence for hypotheses and comparing models in statistical data analysis (Chow & Hoijtink, 2017; Hoijtink & Chow, 2017; Lee, 2011; Mulder & Wagenmakers, 2016; Vandekerckhove, Rouder, & Kruschke, 2018). Compared to traditional frequentist p-values, Bayes factors offer several advantages, such as their ability to assess the likelihood of the null hypothesis, incorporate prior information, and provide continuous measures of evidence strength. As a result, Bayes factors are increasingly used in many scientific domains, including psychology, neuroscience, and social sciences (Heck et al., 2022).

However, computing exact Bayes factors is often not feasible, as it involves solving complex integrals that cannot be solved analytically. Instead, approximations are necessary, which can lead to noisy or biased estimates of Bayes factors. To address this issue, several software tools have been developed, such as the BayesFactor (Morey & Rouder, 2022) and bridgesampling (Gronau, Singmann, & Wagenmakers, 2020) packages in R, which provide estimates of Bayes factors. Nevertheless, the quality of these estimates are as yet unclear and remain a concern, as biased estimates could lead to erroneous conclusions about the evidence for hypotheses and the relative strengths of models (Schad, Nicenboim, Bürkner, Betancourt, & Vasishth, 2022).

To investigate the accuracy of Bayes factor estimates, Schad et al. (2022) recently proposed a method called Simulation-Based Calibration (SBC) for Bayes factors (for more recent alternatives see Sekulovski, Marsman, & Wagenmakers, 2024). SBC provides a means to assess whether Bayes factors are estimated accurately, or whether there is bias in the Bayes factor estimator, reflecting liberal or conservative bias. In essence, SBC works by defining a model prior (i.e., a priori model probabilities) and a parameter prior for each

BIASED BAYES FACTORS

model (i.e., prior densities for each model parameter), and then doing a simulation many times, where in each simulation, one performs the following steps: (1) sample a model from the model prior, (2) for the sampled model, sample parameters from their prior distributions, (3) use the sampled parameters to simulate data from the model, (4) compute (based on Bayes factor estimates) posterior model probabilities for each of the models given the simulated data. We can then compare the average posterior probabilities (averaged across simulations) with the prior probability; this allows SBC to detect whether the Bayes factor estimates are biased or not. If the average posterior probability is equal to the prior probability, then this is consistent with accurate Bayes factor estimation. If the average posterior deviates from the prior, then this indicates a liberal or conservative bias in the Bayes factor estimation.

An important application domain of Bayes factor estimation is in the analysis of factorial designs using linear mixed-effects models (LMMs; Baayen, Davidson, and Bates (2008)). In particular, nested model comparisons for balanced factorial designs can be analyzed using Bayes factors, which can replace traditional p-values as a means of assessing main effects, interactions, and specific contrasts (Schad, Vasishth, Hohenstein, & Kliegl, 2020). In this context, Bayes factors are often used for null hypothesis tests, where a full model containing all fixed effects of interest is compared to a null model that lacks some effect or a set of effects (van Doorn, Aust, Haaf, Stefan, & Wagenmakers, 2021). This has the potential to replace traditional frequentist p-values that are standardly used to analyze experimental data in many scientific papers.

In this paper, we aim to investigate whether Bayes factor estimates obtained from common factorial designs are biased or accurate using SBC. We build upon our prior work (Schad, Nicenboim, & Vasishth, 2024), which demonstrated that data aggregation can lead to biased inferences in Bayesian linear mixed models and ANOVA. To further assess the accuracy of Bayes factor estimation, we focus on two widely used R packages: bridgesampling (Gronau et al., 2020), in combination with brms (Bürkner, 2017), and BayesFactor (Morey & Rouder, 2022), which is also used in Jeffreys' Amazing Statistics Program (JASP; JASP Team (2024)). Specifically, we apply SBC to a range of experimental repeated measures designs, including a 2x2 design with random effects for subjects, a design with a single fixed factor and crossed random effects for subjects and items, and a 2x2 design with crossed random effects for subjects and items. We summarize simulation studies reported before (Schad et al., 2024) and extend these results with a simulation of the BayesFactor package. By using SBC, we can determine whether the Bayes factor estimates for the fixed effects are unbiased and provide a solid basis for statistical data analysis in these experimental designs. This investigation is important, as biased Bayes factor estimates can lead to erroneous conclusions about the evidence for hypotheses and the relative strengths of models. Overall, our study contributes to the growing literature on the application of Bayesian methods in the cognitive sciences (psychology, linguistics, and related areas).

To foreshadow our results: we found that Bayes factor estimates based on the R-packages bridgesampling/brms as well as for the R-package BayesFactor were accurate or exhibited only little bias for a 2x2 Latin square design with random effects for subjects and for a design with a single fixed factor and crossed random effects for subjects and for items. However, in a 2x2 repeated measures design with crossed random effects for subjects and for items, we found that for both packages (bridgesampling and BayesFactor), the resulting Bayes factor estimates were too large, reflecting a liberal bias. These results suggest that the accuracy of BayesFactor estimates in commonly used software tools cannot be taken for granted. While we provide some initial results about which kinds of designs might be most vulnerable to Bayes factor bias, the accuracy of Bayes factor estimates may vary with design features, statistical models used, and with prior assumptions. Therefore, we recommend (cf. Schad et al., 2022) that SBC for Bayes factors (or another test of its accuracy; Sekulovski et al. (2024)) should be employed to test the accuracy for a given set of design, models, and priors before results from individual analyses are trusted. Moreover, we hope that our work

can motivate improvements in software for estimating Bayes factors to achieve more accurate estimation also in the situations that currently pose problems to the estimation procedures. Until this is achieved, we caution users of respective software to acknowledge potential limitations associated with Bayes factor estimation in their analyses, and to interpret relating claims in the literature with caution.

Next, we provide a brief introduction to Bayesian analyses and Bayes factors, before we report the results of our simulation studies.

Introduction to Bayesian analyses, Bayes factors, and simulation-based calibration

Here, we provide a brief introduction to Bayesian analyses, Bayes factors, and to simulation-based calibration for Bayes factors. This introduction is taken from Schad et al. (2024). Other introductory treatments are available for Bayesian and Bayes factor analyses (Chow & Hoijtink, 2017; Etz, Gronau, Dablander, Edelsbrunner, & Baribault, 2018; Etz & Vandekerckhove, 2018; Hoijtink & Chow, 2017; Lee, 2011; Mulder & Wagenmakers, 2016; Nicenboim & Vasishth, 2016; van Doorn et al., 2021; Vandekerckhove et al., 2018; Vasishth, Nicenboim, Beckman, Li, & Kong, 2018).

A quick review of Bayesian methodology

Statistical analyses in the cognitive sciences often pursue two goals: to estimate parameters and their uncertainties, and to test hypotheses. Both of these goals can be achieved using Bayesian data analysis. Bayesian analyses focus on an "observational" model \mathcal{M} , which specifies the probability density of the data y given the vector of model parameters Θ and the model \mathcal{M} , i.e., $p(y \mid \Theta, \mathcal{M})$, or by dropping the model, $p(y \mid \Theta)$. It is possible to use the observational model to simulate data, by selecting some model parameters Θ and drawing random samples for the data \tilde{y} . When the data is given (fixed, e.g., observed or simulated), then the observational model turns into a likelihood function: $p(y \mid \Theta) = L_y(\Theta)$; this can be used to estimate model parameters or to compute the evidence for the model relative to other models. To estimate parameters, in Bayesian data analyses the likelihood is complemented by the prior, written $p(\Theta)$, that defines the joint probability distribution of all the parameters of the model, which can encode domain expertise. Bayes' rule specifies how to combine the likelihood and the prior to compute the posterior distribution of the model parameters $p(\Theta \mid y, \mathcal{M}_1)$:

$$p(\Theta \mid y, \mathcal{M}_1) = \frac{p(y \mid \Theta, \mathcal{M}_1)p(\Theta \mid \mathcal{M}_1)}{p(y \mid \mathcal{M}_1)}$$
(1)

Here, $p(y \mid \mathcal{M}_1)$ is a normalizing constant termed the "evidence" or "marginal likelihood"; it is the likelihood of the data y based on the model \mathcal{M}_1 averaging over the parameters Θ . The marginal likelihood can only be interpreted relative to other models. It is derived as $p(y \mid \mathcal{M}_1) = \int p(y \mid \Theta, \mathcal{M}_1) p(\Theta \mid \mathcal{M}_1) d\Theta$. There is a key role of priors in this computation. Priors play an important role in Bayesian inference as they can regularize inferences when the data do not inform the likelihood functions strongly. However, they will influence marginal likelihoods even when the data are strongly informative, and are thus even more crucial for Bayes factors than for posterior distributions (Aitkin, 1991; Gelman et al., 2013; Grünwald, 2000; Liu & Aitkin, 2008; Myung & Pitt, 1997; Vanpaemel, 2010).

For very simple models, posterior density functions can be computed analytically. However, for most interesting models this is not possible and we have to rely on approximation methods. One such approach is sampling methods such as Markov Chain Monte Carlo sampling, which is an important method behind popular software implementing Bayesian analysis.

Inference over hypotheses. Bayes factors provide a way to compare any two model hypotheses (i.e., arbitrary hypotheses) against each other by comparing their marginal likelihoods (Betancourt, 2018; Kass & Raftery, 1995; Ly, Verhagen, & Wagenmakers, 2016; Schad et al., 2022). The Bayes factor tells us, given the data and the model priors, how much we need to update our relative belief between the two models.

To derive Bayes factors, we first compute the model posterior, i.e., the posterior probability for a model \mathcal{M}_i given the data:

$$p(\mathcal{M}_i \mid y) = \frac{p(y \mid \mathcal{M}_i) \times p(\mathcal{M}_i)}{p(y \mid \mathcal{M}_1) \times p(\mathcal{M}_1) + p(y \mid \mathcal{M}_2) \times p(\mathcal{M}_2)} = \frac{p(y \mid \mathcal{M}_i) \times p(\mathcal{M}_i)}{p(y)}.$$
 (2)

Here, $p(\mathcal{M}_i)$ is the prior probability for each model *i*. Based on the posterior model probability $p(\mathcal{M}_i \mid y)$, we can compute the model odds for one model over another as:

$$\frac{p(\mathcal{M}_1 \mid y)}{p(\mathcal{M}_2 \mid y)} = \frac{[p(y \mid \mathcal{M}_1) \times p(\mathcal{M}_1)]/p(y)}{[p(y \mid \mathcal{M}_2) \times p(\mathcal{M}_2)]/p(y)} = \frac{p(y \mid \mathcal{M}_1)}{p(y \mid \mathcal{M}_2)} \times \frac{p(\mathcal{M}_1)}{p(\mathcal{M}_2)}$$
(3)

In words:

$$Posterior \ ratio = Bayes \ factor \times prior \ odds \tag{4}$$

The Bayes factor is thus a measure of relative evidence, the comparison of the predictive performance of one model (\mathcal{M}_1) against another one (\mathcal{M}_2) . This comparison (BF_{12}) is a ratio of marginal likelihoods:

$$BF_{12} = \frac{p(y \mid \mathcal{M}_1)}{p(y \mid \mathcal{M}_2)} \tag{5}$$

 BF_{12} indicates the evidence that the data provide for \mathcal{M}_1 over \mathcal{M}_2 ; in other words, which of the two models is more likely to have generated the data, or the relative evidence that we have for \mathcal{M}_1 over \mathcal{M}_2 . Bayes factor values larger than one indicate that \mathcal{M}_1 is more compatible with the data, values smaller than one indicate \mathcal{M}_2 is more compatible with the data, and values close to one indicate that both models are equally compatible with the data.

In the present work, we will consider the case of nested model comparison, where a null model hypothesizes that a model parameter is zero or absent (a point hypothesis), whereas an alternative model hypothesizes that the model parameter is present with some prior distribution and has some value different from exactly zero (a "general" hypothesis).

For most interesting problems and models in cognitive science, Bayes factors cannot be computed analytically; approximations are needed. One major approach is to estimate Bayes factors based on posterior MCMC draws via bridge sampling (Bennett, 1976; Meng & Wong, 1996), implemented in the R package bridgesampling (Gronau et al., 2020), which we use in the present work. We also use Bayes factors computed by the R package BayesFactor (Morey & Rouder, 2018). We chose these two packages for the present study because they are widely used in cognitive science.

Introduction to simulation-based calibration (SBC). We have recently used SBC for Bayes factors (Schad et al., 2022), which is a statistical technique designed to test whether a Bayes factor estimated in a given analysis is accurate, or whether it is biased and deviates from the true Bayes factor. Here, we first provide a short description of SBC (derived from Schad et al., 2022) and then perform SBC to test the accuracy of Bayes factors for analyses of non-/aggregated data in the presence of non-spherical random effects.

We can formulate an SBC-inspired method for Bayes factors (i.e., for model inference), where \mathcal{M} is a true model used to simulate artificial data y, and \mathcal{M}' is a model inferred from the simulated data. SBC for model inference makes use of model priors $p(\mathcal{M})$, which are the prior probabilities for each model before observing any data. SBC can then be formulated as follows (Schad et al., 2022).

$$p(\mathcal{M}') = \sum_{\mathcal{M}} \int p(\mathcal{M}' \mid y) p(y \mid \mathcal{M}) p(\mathcal{M}) \, \mathrm{d}y$$
(6)

We can read this equation sequentially (from right to left): first, we sample a model from the model prior, $p(\mathcal{M})$. Next, we simulate data based on this model, $p(y \mid \mathcal{M})$. This in fact involves two steps: simulating parameters from the parameter prior, $p(\Theta \mid \mathcal{M})$, and then simulating artificial data from the parameters and model, $p(y \mid \Theta, \mathcal{M})$, i.e., $p(y \mid \mathcal{M}) = \int p(y \mid \Theta, \mathcal{M}) \times p(\Theta \mid \mathcal{M}) \, d\Theta$. Next, we estimate the posterior model probabilities from the simulated data, $p(\mathcal{M}' \mid y)$. This again involves several steps: we compute marginal likelihoods, $p(y \mid \mathcal{M}')$, for both models, compute Bayes factors, and compute the posterior probability for each model given the data, $p(\mathcal{M}' \mid y)$, by adding the model prior. That is, we obtain a posterior model probability for each simulated data set. We can now compare, whether the posterior model probability, $p(\mathcal{M}' \mid y)$, – averaged across all simulated data sets – is the same as the prior model probability, $p(\mathcal{M})$.

The key idea is that if the computation of Bayes factors and posterior model probabilities is performed correctly (and of course the data simulation is implemented correctly), then the average posterior probability for a model should be the same as its prior probability. By contrast, if the average posterior probability for a model deviates from its prior probability, then this indicates that the Bayes factor estimate is biased, i.e., that the obtained Bayes factor does not correspond to the true Bayes factor.

For a given run of SBC, to test whether the Bayes factor estimates are biased, we here perform Bayesian t-tests on the posterior model probabilities, i.e., computing null hypothesis Bayes factors to test whether the posterior model probabilities differ from the prior model probability (here either 0.5 or 0.2). For these tests, we use the t-test in the BayesFactor package to test whether SBC indicates bias in the estimation of Bayes factors. Results 1: Cases where Bayes factor estimates are accurate or exhibit slight bias



2x2 design with random effects for subjects: bridgesampling/brms

Figure 1. Results from simulation-based calibration (SBC) for model inference: A simulation example employing a 2 x 2 design based on the two-step decision task. A) The average posterior model probability together with 95 percent confidence intervals is shown for the main effects of reward (last trial was rewarded/not rewarded; random slope variance: SD = 0) and transition frequency (last trial induced a common/rare transition; random slope variance: SD = 0), and their interaction (random slope variance: SD = 0.4), reflecting differing variations of the effects across subjects. The horizontal broken line is the prior probability for the H1, and deviations from this line indicate estimation bias. B) Results from SBC: The question is examined whether the posterior probability of H1 (see panel A) is different from the prior probability of 0.5, using null hypothesis Bayes factor tests, including sensitivity analyses for the prior scale. The vertical dashed line indicates a default Bayes factor with prior scale of sqrt(2)/2. A+B) The results show that the posterior probability for H1 (see panel A) does not diverge from the prior probability (of 0.5), suggesting that Bayes factor estimates are accurate. We first performed SBC for Bayes factors for the bridgesampling package together with brms in a 2×2 design inspired from the 2-step decision-making task (Daw, Gershman, Seymour, Dayan, & Dolan, 2011). In this task, the probability to repeat an action is examined as a function of whether the last trial was rewarded (contrast coding: rewarded = 0.5, not rewarded = -0.5) and whether during the last trial a common (0.5) or a rare (-0.5) transition occurred in the task. Using SBC, we simulated data from a LMM. We simulated data from 10 virtual subjects, and for each subject, we simulated average repetition probabilities for each of five blocks. The model parameters used for simulating the SBC data were inspired by our own data on the task (Schad et al., 2014). We assumed the following prior distributions:

$$\beta_{(Intercept)} \sim Normal(0.7, 0.1)$$
 (7)

$$\beta_{Contrasts} \sim Normal(0, 0.2)$$
 (8)

$$\sigma_{Random\,slopes} \sim Normal_+(0, 0.2) \tag{9}$$

$$\sigma_{Residual} \sim Normal_+(0, 0.5) \tag{10}$$

$$\rho_{Random\,slopes} \sim LKJ(2)$$
(11)

In the simulations, we set random slope variances to SD = 0 for the two main effects, and to SD = 0.4 for the interaction, effectively assuming non-spherical random effects (see Schad et al., 2024). Prior probabilities for the models were set to 50% each. We used 200 runs of SBC.

The results (see Fig. 1) showed that the posterior model probabilities were equal to the prior model probability of 0.5 for both main effects and for the interaction. Indeed, the null hypothesis that posterior model probabilities were equal to 0.5 was supported by Bayesian t-tests (see Fig. 1b) showing Bayes factors of BF01 > 6 for the null for the two main effects

and a Bayes factor of BF01 > 3 for the interaction. These results suggest that Bayes factor estimates were accurate.



2x2 design with random effects for subjects: BayesFactor

Figure 2. Simulation-based calibration for the BayesFactor package, using a 2 x 2 repeated measures design. True random slope variances are simulated to be small for the main effects and large for the interaction. A) Average posterior probabilities for H1 for main effects and the interaction. B) Sensitivity analyses for a Bayesian t-test assessing whether the average posterior probability for H1 (see panel A) differs from the prior value of 0.5. The vertical line indicates the prior scale of a default Bayes factor. A+B) Results show that Bayes factors are accurate and the null hypothesis of no bias is supported by the SBC.

As a next step, we performed SBC to analyze the BayesFactor package. We studied a 2×2 repeated measures design. Given that Bayes factor estimation in the BayesFactor package proceeds considerably more quickly compared to the brms/bridgesampling packages, we assumed 30 simulated subjects with each contributing 16 data points. Model parameters were taken as the default values used in the BayesFactor package.

The results (see Fig. 2) using 1000 SBC simulations showed that the average posterior

probabilities for H1 were close to the prior value of 0.5. Indeed, default Bayesian t-tests showed Bayes factors in support of the null hypothesis of BF01 > 6 for all three tested effects, and BF01 > 10 for two of the three effects. This supports the conclusion that the estimated Bayes factors were accurate in this analysis.

One 2-level factor with crossed random effects for subjects and items: bridgesampling/brms

Next, we simulated data from a Latin square design with one two-level fixed factor and crossed random effects for 42 subjects and 16 items. Across the 125 SBC simulations, the true standard deviation of the random item slopes was increased from 0 to 0.5 in equidistant steps of size 0.004. We set the prior probability of the alternative hypothesis to a smaller value of 0.2 (instead of the previous 0.5) to be able to detect potential liberal biases well without ceiling effects. A Bayesian GLMM with a lognormal likelihood was used in the SBC analyses using the package brms. We included correlated random slopes and intercepts for subjects and for items. For the model parameters, we used the following priors:

$$\beta_{(Intercept)} \sim Normal(6, 0.6) \tag{12}$$

$$\beta_X \sim Normal(0, 0.1) \tag{13}$$

$$\sigma_{Random\,slopes} \sim Normal_+(0, 0.1)$$
 (14)

$$\sigma_{Residual} \sim Normal_+(0, 0.5) \tag{15}$$

$$\rho_{Random\,slopes} \sim LKJ(2)$$
(16)

The results showed (see Fig. 3) that the average posterior model probabilities for the H1 did not differ from the prior value of 0.2. Indeed, default Bayes factor analyses with Bayes factors of BF01 > 3 provided some support for the null hypothesis that the average posterior probability for H1 did not differ from 0.2 and that it did not change with



Figure 3. SBC simulations for a design with crossed random effects for subjects and items and a single two-level fixed factor. A) the posterior probability for the alternative hypothesis (H1) is shown as a function of the true standard deviation of the item random slope. Error bars are confidence intervals, regression lines are from a logistic regression. The prior probability for the null is set to 0.2 (this is chosen - lower than before, where it was 0.5 - for pragmatic reasons to demonstrate a potential increase in the average posterior probabilities compared to the prior of 0.2 and to avoid potential ceiling effects). B) Results from a sensitivity analysis of Bayesian tests of whether the average posterior probability differs from the prior of 0.2 and whether the posterior probability increases with increasing item random slopes. Vertical lines indicate default Bayes factor prior scales. A+B) The results show that the posterior probability for H1 does not differ from the prior value of 0.2, suggesting that Bayes factor estimates are accurate and not biased.

increasing values for the item random slope.



One repeated measures factor with crossed random effects for subjects and

Figure 4. Repeating the previous analyses using the BayesFactor package. A) Results from SBC: the posterior probability for the alternative hypothesis (H1) is shown as a function of the true standard deviation of the item random slope. Error bars are confidence intervals, regression lines are from a logistic regression. The prior probability for the null is set to 0.2. B) Results from sensitivity analyses showing the Bayes factor BF01 in support of the null hypothesis: in the left panel of (B), the null hypothesis is that the average posterior is equal to the prior of 0.2; in the right panel of (B), the null hypothesis is that the influence of the true random slope is zero. The results show that given the high number of simulations, the analysis can detect a numerically small but persistent bias in the Bayes factor estimation: Bayes factors are slightly too small, reflecting a conservative bias.

Next, we used the same experimental design (two-level fixed factor and crossed random effects for subjects and items) to perform SBC for the BayesFactor package, again using default parameter settings for the BayesFactor package. We used n = 500 simulations due to

the relatively high speed of the BayesFactor package in estimating Bayes factors.

As shown in Figure 4, the average posterior probability for H1 was slightly, but persistently smaller than the prior probability of 0.2. This was supported by a default Bayesian t-test with BF10 = 26 for the difference from the prior probability of 0.2. The analysis moreover suggested that this effect did not change with the value of the true random slope variance (BF01 = 3.9). Thus, the relatively high number of simulations revealed a small but persistent conservative bias in the Bayes factor estimate, suggesting that estimated Bayes factors are *slightly* smaller than their true value.

We repeated the above set of simulations with a similar design, but using a four-level fixed factor instead of a two-level fixed factor. The results showed that despite the large number of 500 simulation runs of SBC, the analysis was undecided on whether there was a small bias in the estimated Bayes factors as being too small, or whether the Bayes factor estimates were accurate (see Fig. 5): a default Bayesian t-test yielded a value of BF10 = 1.3 and was thus undecided. Again, the true random slope variance did not seem to influence this result (BF01 = 6.9). The results thus suggest that if there is any bias in the Bayes factor estimates, then this bias is likely of small magnitude. Large biases are unlikely based on the results.

Results 2: Cases where we found (stronger) bias in Bayes factor estimates

Next, we report results in a situation where we found evidence for stronger biases in estimated Bayes factors.

2x2 repeated measures design with crossed random effects for subjects and items: bridgesampling/brms

We extended our previous simulations by performing SBC with 200 simulations with the bridgesampling/brms packages in a 2×2 Latin square design with crossed random



Figure 5. Repeating the previous analyses using a four-level fixed factor instead of a two-level fixed factor. A) Results from SBC: the posterior probability for the alternative hypothesis (H1) is shown as a function of the true standard deviation of the item random slope. Error bars are confidence intervals, regression lines are from a logistic regression. The prior probability for the null is set to 0.2. B) Results from sensitivity analyses showing the Bayes factor BF01 in support of the null hypothesis: in the left panel of (B), the null hypothesis is that the average posterior is equal to the prior of 0.2; in the right panel of (B), the null hypothesis is that the influence of the true random slope is zero. The results show that given the high number of simulations, the analysis is unsure whether there is a small downward bias in the Bayes factor estimation, or whether the analysis is accurate.



Figure 6. Analyses of a 2 x 2 repeated measures design. A) Results from SBC: the posterior probability for the alternative hypothesis (H1) is shown as a function of the true standard deviation of the item random slope for main effects A (upper panel) and B (middle panel) as well as their interaction (lower panel). Error bars are confidence intervals, regression lines are from a logistic regression. The prior probability for the null is set to 0.2. B) Results from sensitivity analyses of Bayes factor tests of whether the average posterior probability diverges from the prior value of 0.2, as well as of the influence of the random slope variance. The results show that in the present design, null hypothesis Bayes factor tests were biased such that estimated Bayes factor values were too large and larger than the true Bayes factor.

The results (see Fig. 6) showed that the average posterior probability for the H1 was larger than the prior probability of 0.2 for both main effects and the interaction. The default Bayes factors for the divergence from 0.2 were BF10 = 58 for main effect A, BF10 = 3342 for main effect B, and BF10 = 50714 for the interaction. These provide strong evidence for a clear divergence of the posterior from the prior, and suggest that Bayes factor estimates are too large, i.e., larger than their true values, and thus provide liberal tests of the alternative hypotheses. These effects were not strongly modulated by the true value of the random slope variance. These results suggest that testing fixed effects (main effects and interactions) in a 2 × 2 design using the bridgesampling/brms packages can run the danger of providing too large Bayes factors that exaggerate the apparent evidence for an effect that is different from zero even if this may not be true.

2x2 repeated measures design with crossed random effects for subjects and items: BayesFactor

We used the same experimental design with a 2×2 fixed effects structure and crossed random effects for 64 subjects and 16 items to perform SBC with 500 simulations for the BayesFactor package. The results (see Fig. 7) showed that the average posterior probability for the alternative hypothesis was much larger than the prior probability of 0.2 for both main effects and the interaction. Bayes factors of the null hypothesis of no divergence were decisive with BF10 > 1e + 10 for all three effects. Moreover, this bias got stronger with increasing random slope variances for main effect B and for the interaction, whereas this effect was not so clear for main effect A.

Discussion

In recent years, very user-friendly software has been developed to estimate Bayes factors in (generalized) linear mixed-effects models in the cognitive sciences, including the bridgesampling/brms packages as well as the BayesFactor package. However, Bayes factors in complex models have to be estimated, with no guarantees for their accuracy. Here, we



Figure 7. Analyses of a 2 x 2 repeated measures design, but now with the BayesFactor package. A) Results from SBC: the posterior probability for the alternative hypothesis (H1) is shown as a function of the true standard deviation of the item random slope for main effects A (upper panel) and B (middle panel) as well as their interaction (lower panel). Error bars are confidence intervals, regression lines are from a logistic regression. The prior probability for the null is set to 0.2. B) Results from sensitivity analyses of Bayes factor tests of whether the average posterior probability diverges from the prior value of 0.2, as well as of the influence of the random slope variance. The results show that in the present design, null hypothesis Bayes factor tests were biased such that estimated Bayes factor values were too large and larger than the true Bayes factor.

BIASED BAYES FACTORS

used simulation-based calibration for Bayes factors (Schad et al., 2022) to test the accuracy of Bayes factor estimation by studying some commonly used factorial experimental designs from the cognitive sciences. The results showed that Bayes factors were estimated accurately or with little bias in a range of well-studied repeated measures Latin square designs, which included (a) a 2×2 design with random effects for subjects and (b) designs with a single fixed factor (with two or four levels) and crossed random effects for subjects and items. This provides important validation of the relevant Bayes factor estimates.

However, when testing main effects and interactions in a 2×2 design with crossed random effects for subjects and items, we found strong biases in Bayes factor estimates indicating that estimated Bayes factors were too large, and larger than the true Bayes factors, leading to liberal statistical null hypothesis tests that may indicate evidence for an effect even when the effect is in fact zero. These biases were present for both studied software packages, including the bridgesampling/brms packages, as well as the BayesFactor package.

These latter results are worrisome for the application of Bayes factor tests in 2×2 designs with crossed random effects for subjects and items - an experimental design that is commonly used in fields such as psycholinguistics and psychology. They suggest that researchers should exercise caution when interpreting results from null hypothesis Bayes factor tests for these types of experimental designs, and they invite a re-evaluation of the published literature based on these approaches.

We think that the results can be highly informative for developers of software tools to estimate Bayes factors in such complex statistical (GLMM) models. They may point to problems in the accurate estimation of Bayes factors in some specific analysis situations, and we hope that the present results will lead to improvements in methodology to overcome the current limitations to provide accurate estimates of Bayes factors also in these complex analysis situations. Our results are quite limited in that we only focussed on a limited set of analysis settings, priors, models, experimental designs, and software packages. There is thus no guarantee that similar results will be obtained with slightly different designs, models, priors, and data sets. The present results therefore highlight that it is good practice to validate the use of Bayes factors when applying them in practice using simulation-based calibration for Bayes factors (Schad et al., 2022) or alternative approaches (Sekulovski et al., 2024). We hope that the present results will lead to a more robust use and application of Bayes factors by highlighting their limitations and encouraging researchers to check for their individual analysis, whether estimated Bayes factors are accurate.

However, performing SBC for every single analysis can be time consuming and difficult to implement for non-expert users. Moreover, software developers need the best possible information to understand the precise conditions under which current methods for Bayes factor estimates are biased versus accurate in order to spot potential problems and improve current technology. Therefore, we think that future research is needed to provide a thorough characterization of the conditions under which Bayes factor estimates are biased versus accurate.

Acknowledgements

We thank Bruno Nicenboim for discussion.

Declarations

• Funding

This work was partly funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation), Sonderforschungsbereich 1287, project number 317633480 (Limits of Variability in Language).

• Conflicts of interest/Competing interests

None

• Ethics approval

Not applicable.

• Consent to participate

Not applicable.

• Consent for publication

Not applicable.

• Availability of data and materials

Not applicable.

• Code availability

The code is available on OSF (https://osf.io/3g86r/).

• Authors' contributions

DJS: Conception, analysis, writing of first draft, review and editing.

SV: Conception, review and editing.

Open Practices Statement

Analysis code is available at https://osf.io/3g86r/. The reported studies were not preregistered.

References

- Aitkin, M. (1991). Posterior Bayes factors. Journal of the Royal Statistical Society: Series B (Methodological), 53(1), 111–128.
- Baayen, R. H., Davidson, D. J., & Bates, D. M. (2008). Mixed-effects modeling with crossed random effects for subjects and items. *Journal of Memory and Language*, 59(4), 390–412.
- Bennett, C. H. (1976). Efficient estimation of free energy differences from Monte Carlo data. Journal of Computational Physics, 22(2), 245–268.
- Betancourt, M. (2018). Calibrating model-based inferences and decisions. arXiv Preprint arXiv:1803.08393.
- Bürkner, P.-C. (2017). brms: An R package for Bayesian multilevel models using Stan. Journal of Statistical Software, 80(1), 1–28. https://doi.org/10.18637/jss.v080.i01
- Chow, S.-M., & Hoijtink, H. (2017). Bayesian estimation and modeling: Editorial to the second special issue on Bayesian data analysis. *Psychological Methods*, 22(4), 609–615.
- Daw, N. D., Gershman, S. J., Seymour, B., Dayan, P., & Dolan, R. J. (2011). Model-based influences on humans' choices and striatal prediction errors. *Neuron*, 69(6), 1204–1215.
- Etz, A., Gronau, Q. F., Dablander, F., Edelsbrunner, P. A., & Baribault, B. (2018).
 How to become a Bayesian in eight easy steps: An annotated reading list.
 Psychonomic Bulletin & Review, 25(1), 219–234.
- Etz, A., & Vandekerckhove, J. (2018). Introduction to Bayesian inference for psychology. *Psychonomic Bulletin & Review*, 25(1), 5–34.
- Gelman, A., Carlin, J. B., Stern, H. S., Dunson, D. B., Vehtari, A., & Rubin, D. B. (2013). Bayesian data analysis. New York: CRC Press.
- Gronau, Q. F., Singmann, H., & Wagenmakers, E.-J. (2020). bridgesampling: An R package for estimating normalizing constants. *Journal of Statistical Software*,

92(10), 1–29. https://doi.org/10.18637/jss.v092.i10

- Grünwald, P. (2000). Model selection based on minimum description length. *Journal* of Mathematical Psychology, 44(1), 133–152.
- Heck, D. W., Boehm, U., Böing-Messing, F., Bürkner, P.-C., Derks, K., Dienes, Z., ... others. (2022). A review of applications of the Bayes factor in psychological research. *Psychological Methods*.
- Hoijtink, H., & Chow, S.-M. (2017). Bayesian hypothesis testing: Editorial to the special issue on Bayesian data analysis. *Psychological Methods*, 22(2), 211–216.
- JASP Team. (2024). JASP (Version 0.18.3)[Computer software]. Retrieved from https://jasp-stats.org/
- Kass, R. E., & Raftery, A. E. (1995). Bayes factors. Journal of the American Statistical Association, 90(430), 773–795.
- Lee, M. D. (2011). How cognitive modeling can benefit from hierarchical Bayesian models. Journal of Mathematical Psychology, 55(1), 1–7.
- Liu, C. C., & Aitkin, M. (2008). Bayes factors: Prior sensitivity and model generalizability. Journal of Mathematical Psychology, 52(6), 362–375.
- Ly, A., Verhagen, J., & Wagenmakers, E.-J. (2016). Harold Jeffreys's default Bayes factor hypothesis tests: Explanation, extension, and application in psychology. *Journal of Mathematical Psychology*, 72, 19–32.
- Meng, X., & Wong, W. H. (1996). Simulating ratios of normalizing constants via a simple identity: A theoretical exploration. *Statistica Sinica*, 6(4), 831–860.
- Morey, R. D., & Rouder, J. N. (2018). BayesFactor: Computation of Bayes factors for common designs. Retrieved from https://CRAN.R-project.org/package=BayesFactor
- Morey, R. D., & Rouder, J. N. (2022). BayesFactor: Computation of bayes factors for common designs. Retrieved from https://CRAN.R-project.org/package=BayesFactor

- Mulder, J., & Wagenmakers, E.-J. (2016). Editors' introduction to the special issue "Bayes factors for testing hypotheses in psychological research: Practical relevance and new developments." *Journal of Mathematical Psychology*, 72, 1–5.
- Myung, I. J., & Pitt, M. A. (1997). Applying Occam's razor in modeling cognition: A Bayesian approach. *Psychonomic Bulletin & Review*, 4(1), 79–95.
- Nicenboim, B., & Vasishth, S. (2016). Statistical methods for linguistic research: Foundational Ideas - Part II. Language and Linguistics Compass, 10(11), 591–613. https://doi.org/10.1111/lnc3.12207
- Schad, D. J., Jünger, E., Sebold, M., Garbusow, M., Bernhardt, N., Javadi, A.-H., ... others. (2014). Processing speed enhances model-based over model-free reinforcement learning in the presence of high working memory functioning. *Frontiers in Psychology*, 1450. https://doi.org/10.3389/fpsyg.2014.01450
- Schad, D. J., Nicenboim, B., Bürkner, P.-C., Betancourt, M., & Vasishth, S. (2022). Workflow techniques for the robust use of Bayes factors. *Psychological Methods*. https://doi.org/10.1037/met0000472
- Schad, D. J., Nicenboim, B., & Vasishth, S. (2024). Data aggregation can lead to biased inferences in Bayesian linear mixed models and Bayesian analysis of variance. *Psychological Methods*. https://doi.org/10.1037/met0000621
- Schad, D. J., Vasishth, S., Hohenstein, S., & Kliegl, R. (2020). How to capitalize on a priori contrasts in linear (mixed) models: A tutorial. *Journal of Memory and Language*, 110, 104038. https://doi.org/10.1016/j.jml.2019.104038
- Sekulovski, N., Marsman, M., & Wagenmakers, E.-J. (2024). A good check on the bayes factor.
- van Doorn, J., Aust, F., Haaf, J. M., Stefan, A. M., & Wagenmakers, E.-J. (2021). Bayes factors for mixed models. *Computational Brain & Behavior*, 1–13.
- Vandekerckhove, J., Rouder, J. N., & Kruschke, J. K. (2018). Bayesian methods for advancing psychological science. *Psychonomic Bulletin & Review*, 25(1), 1–4.

- Vanpaemel, W. (2010). Prior sensitivity in theory testing: An apologia for the Bayes factor. Journal of Mathematical Psychology, 54 (6), 491–498.
- Vasishth, S., Nicenboim, B., Beckman, M. E., Li, F., & Kong, E. (2018). Bayesian data analysis in the phonetic sciences: A tutorial introduction. *Journal of Phonetics*, 71, 147–161. https://doi.org/10.1016/j.wocn.2018.07.008