

INFINITE-DIMENSIONAL FROBENIUS MANIFOLDS UNDERLYING THE GENUS-ZERO UNIVERSAL WHITHAM HIERARCHY

SHILIN MA

ABSTRACT. In this paper, we construct a new class of infinite-dimensional Frobenius manifolds on the spaces of pairs of meromorphic functions that are defined on specific regions of the Riemann sphere. We demonstrate that the principal hierarchy of these Frobenius manifolds serves as an extension of the genus-zero universal Whitham hierarchy.

CONTENTS

1. Introduction	1
1.1. Background	1
1.2. Main results	6
2. Infinite-dimensional Frobenius manifold structure on \mathcal{M}	9
2.1. Flat metric	9
2.2. Hamiltonian structure on $L\mathcal{M}$	11
2.3. Multiplication and Potential	19
3. Principal hierarchy of \mathcal{M}	23
References	27

1. INTRODUCTION

1.1. **Background.** The concept of Frobenius manifold introduced by Dubrovin [1] offers a geometric interpretation of the Witten-Dijkgraaf-Verlinde-Verlinde (WDVV) equation in topological field theory. This manifold has significant applications across various domains of mathematical physics, including Gromov-Witten theory, singularity theory, and integrable systems. For further insights, see [2, 3, 4, 5, 6, 7] and the references cited therein.

A Frobenius manifold of charge d is an n -dimensional manifold M , where each tangent space $T_v M$ is equipped with a Frobenius algebra structure $(A_v = T_v M, \circ, e, \langle \cdot, \cdot \rangle)$ that varies smoothly with $v \in M$. This structure must satisfy the following axioms:

Date: June 13, 2024.

- (1) The bilinear form $\langle \cdot, \cdot \rangle$ provides a flat metric on M , and the unity vector field e satisfies $\nabla e = 0$, where ∇ is the Levi-Civita connection for the flat metric.
- (2) Define a 3-tensor c by $c(X, Y, Z) := \langle X \circ Y, Z \rangle$ with $X, Y, Z \in T_v M$. Then, the 4-tensor $(\nabla_W c)(X, Y, Z)$ is symmetric in $X, Y, Z, W \in T_v M$.
- (3) There exists a vector field E , called the Euler vector field, which satisfies $\nabla^2 E = 0$ and

$$\begin{aligned} [E, X \circ Y] - [E, X] \circ Y - X \circ [E, Y] &= X \circ Y, \\ E(\langle X, Y \rangle) - \langle [E, X], Y \rangle - \langle X, [E, Y] \rangle &= (2 - d)\langle X, Y \rangle \end{aligned}$$

for any vector fields X, Y on M .

On an n -dimensional Frobenius manifold M , we select a set of flat coordinates $t = (t^1, \dots, t^n)$, which simplifies the unity vector field to $e = \frac{\partial}{\partial t^i}$. In this coordinate system, the components of the metric $\langle \cdot, \cdot \rangle$ are given by:

$$\eta_{\alpha\beta} = \left\langle \frac{\partial}{\partial t^\alpha}, \frac{\partial}{\partial t^\beta} \right\rangle, \quad \alpha, \beta = 1, \dots, n,$$

where $\eta_{\alpha\beta}$ defines a constant and non-degenerate $n \times n$ matrix. The inverse of this matrix is denoted by $\eta^{\alpha\beta}$. The metric and its inverse are utilized for index lowering and raising, respectively, with the Einstein summation convention applied to repeated Greek indices.

Furthermore, we denote the components of the 3-tensor c by:

$$c_{\alpha\beta\gamma} = c \left(\frac{\partial}{\partial t^\alpha}, \frac{\partial}{\partial t^\beta}, \frac{\partial}{\partial t^\gamma} \right), \quad \alpha, \beta, \gamma = 1, \dots, n,$$

which allows us to express the product in the Frobenius algebra $T_v M$ in terms of

$$\frac{\partial}{\partial t^\alpha} \circ \frac{\partial}{\partial t^\beta} = c_{\alpha\beta}^\gamma \frac{\partial}{\partial t^\gamma},$$

where the structure coefficients $c_{\alpha\beta}^\gamma$ are obtained by contracting the 3-tensor c with the metric $\eta^{\alpha\beta}$:

$$c_{\alpha\beta}^\gamma = \eta^{\gamma\epsilon} c_{\epsilon\alpha\beta},$$

and satisfy

$$c_{1\alpha}^\beta = \delta_\alpha^\beta, \quad c_{\alpha\beta}^\epsilon c_{\epsilon\gamma}^\sigma = c_{\alpha\gamma}^\epsilon c_{\epsilon\beta}^\sigma.$$

Within the local context of the Frobenius manifold M , there exists a smooth function $F(t)$ that satisfies the following properties:

$$\begin{aligned} c_{\alpha\beta\gamma} &= \frac{\partial^3 F}{\partial t^\alpha \partial t^\beta \partial t^\gamma}, \\ \text{Lie}_E F &= (3 - d)F + \text{quadratic terms in } t. \end{aligned}$$

Hence, $F(t)$ is a solution to the WDVV equation

$$\frac{\partial^3 F}{\partial t^\alpha \partial t^\beta \partial t^\gamma} \eta^{\gamma\epsilon} \frac{\partial^3 F}{\partial t^\epsilon \partial t^\sigma \partial t^\mu} = \frac{\partial^3 F}{\partial t^\alpha \partial t^\sigma \partial t^\gamma} \eta^{\gamma\epsilon} \frac{\partial^3 F}{\partial t^\epsilon \partial t^\beta \partial t^\mu}.$$

The third-order derivatives $c_{\alpha\beta\gamma}$ of $F(t)$ are known as the 3-point correlator functions in the context of topological field theory.

For a Frobenius manifold M , its cotangent space $T_v^* M$ is endowed with a Frobenius algebra structure as well. This structure encompasses an invariant bilinear form and a product, which are defined by:

$$\langle dt^\alpha, dt^\beta \rangle = \eta^{\alpha\beta}, \quad dt^\alpha \circ dt^\beta = \eta^{\alpha\epsilon} c_{\epsilon\gamma}^\beta.$$

Let us define

$$g^{\alpha\beta} = i_E (dt^\alpha \circ dt^\beta),$$

then $(dt^\alpha, dt^\beta) := g^{\alpha\beta}$ establishes a symmetric bilinear form known as the intersection form on $T_v^* M$. The intersection form $g^{\alpha\beta}$ and the invariant bilinear form $\eta^{\alpha\beta}$ together form a pencil of flat metrics

$$g^{\alpha\beta} + \epsilon \eta^{\alpha\beta}$$

parameterized by ϵ . As a result, they give rise to a bi-hamiltonian structure of hydrodynamic type on the loop space $\{S^1 \rightarrow M\}$, expressed as:

$$\{ , \}_2 + \epsilon \{ , \}_1.$$

The deformed flat connection on M , originally introduced by Dubrovin [1], is defined as:

$$\tilde{\nabla}_X Y = \nabla_X Y + z X \circ Y, \quad X, Y \in \text{Vect}(M),$$

This connection can be consistently extended to a flat affine connection on $M \times \mathbb{C}^*$ with the following additional definitions:

$$\begin{aligned} \tilde{\nabla}_X \frac{d}{dz} &= 0, \\ \tilde{\nabla}_{\frac{d}{dz}} \frac{d}{dz} &= 0, \\ \tilde{\nabla}_{\frac{d}{dz}} X &= \partial_z X + E \circ X - \frac{1}{z} \mathcal{V}(X), \end{aligned}$$

where X is a vector field on $M \times \mathbb{C}^*$ that vanishes along the \mathbb{C}^* factor, and $\mathcal{V}(X)$ is defined as

$$\mathcal{V}(X) := \frac{2-d}{2} X - \nabla_X E.$$

There exists a system of deformed flat coordinates $\tilde{v}_1(t, z), \dots, \tilde{v}_n(t, z)$ that can be expressed in terms of

$$(\tilde{v}_1(t, z), \dots, \tilde{v}_n(t, z)) = (\theta_1(t, z), \dots, \theta_n(t, z)) z^\mu z^R.$$

These coordinates are chosen such that the 1-forms

$$\xi_\alpha = \frac{\partial \tilde{v}_\alpha}{\partial t^\beta} dt^\beta, \quad \alpha = 1, \dots, n, \quad \text{and} \quad \xi_{n+1} = dz,$$

constitute a basis of solutions to the system $\tilde{\nabla} \xi = 0$. Here, $\mu = \text{diag}(\mu_1, \dots, \mu_n)$ is a diagonal matrix characterized by

$$\mathcal{V} \left(\frac{\partial}{\partial t^\alpha} \right) = \mu_\alpha \frac{\partial}{\partial t^\alpha}, \quad \alpha = 1, \dots, n,$$

which is called the spectrum of M , and $R = R_1 + \dots + R_m$ is a constant nilpotent matrix satisfying

$$\begin{aligned} (R_s)_\beta^\alpha &= 0 \text{ if } \mu_\alpha - \mu_\beta \neq s, \\ (R_s)_\alpha^\gamma \eta_{\gamma\beta} &= (-1)^{s+1} (R_s)_\beta^\gamma \eta_{\gamma\alpha}. \end{aligned}$$

The functions $\theta_\alpha(t, z)$, being analytic near $z = 0$, can be represented by a power series expansion:

$$\theta_\alpha(t, z) = \sum_{p \geq 0} \theta_{\alpha,p}(t) z^p, \quad \alpha = 1, \dots, n.$$

From this, we derive the following relations:

$$\frac{\partial^2 \theta_{\alpha,p+1}(t)}{\partial t^\beta \partial t^\gamma} = c_{\beta\gamma}^\epsilon(t) \frac{\partial \theta_{\alpha,p}(t)}{\partial t^\epsilon}, \quad (1.1)$$

and

$$\text{Lie}_E \left(\frac{\partial \theta_{\alpha,p}(t)}{\partial t^\beta} \right) = (p + \mu_\alpha + \mu_\beta) \frac{\partial \theta_{\alpha,p}(t)}{\partial t^\beta} + \frac{\partial}{\partial t^\beta} \sum_{s=1}^p \theta_{\epsilon,p-s}(t) (R_s)_\alpha^\epsilon. \quad (1.2)$$

Moreover, the normalization condition¹ is imposed:

$$\theta_{\alpha,0}(t) = \eta_{\alpha\beta} t^\beta. \quad (1.3)$$

Given a system of solutions $\{\theta_{\alpha,p}\}$ to the equations (1.1)-(1.3), the principal hierarchy associated with M is defined by the following system of Hamiltonian equations on the loop space $\{S^1 \rightarrow M\}$:

$$\frac{\partial t^\gamma}{\partial T^{\alpha,p}} = \left\{ t^\gamma(x), \int \theta_{\alpha,p+1}(t) dx \right\}_1 := \eta^{\gamma\beta} \frac{\partial}{\partial x} \left(\frac{\partial \theta_{\alpha,p+1}(t)}{\partial t^\beta} \right), \quad \alpha, \beta = 1, 2, \dots, n, \quad p \geq 0.$$

These commuting flows are tau-symmetric, which means that

$$\frac{\partial \theta_{\alpha,p}(t)}{\partial T^{\beta,q}} = \frac{\partial \theta_{\beta,q}(t)}{\partial T^{\alpha,p}}, \quad \alpha, \beta = 1, 2, \dots, n, \quad p, q \geq 0.$$

Furthermore, these flows can be expressed in a bi-hamiltonian recursion form as

$$\mathcal{R} \frac{\partial}{\partial T^{\alpha,p-1}} = \frac{\partial}{\partial T^{\alpha,p}} \left(p + \mu_\alpha + \frac{1}{2} \right) + \sum_{s=1}^p \frac{\partial}{\partial T^{\epsilon,p-s}} (R_s)_\alpha^\epsilon,$$

¹In the referenced work [1], an additional condition $\langle \nabla \theta_\alpha(t, z), \nabla \theta_\beta(t, -z) \rangle = \eta_{\alpha\beta}$ was considered. However, as it does not significantly alter the properties of the principal hierarchy, we omit it here for computational simplicity

where $\mathcal{R} = \{ , \}_2 \cdot \{ , \}_1^{-1}$ is the recursion operator.

When the associated Frobenius manifold is semisimple, the principal hierarchy can be deformed to a dispersive hierarchy, such that the action of the Virasoro symmetries of the principal hierarchy on the tau function of the result hierarchy is linear. The tau function, determined by the string equation of this hierarchy, gives rise to the partition function of the two-dimensional topological field theory (2D TFT). This significant link between Frobenius manifolds and 2D TFT has been instrumental in advancing the study of both areas. It has also enhanced our comprehension of the geometric and algebraic structures inherent in these domains. For further details, refer to [4, 8, 9].

In recent years, efforts have been made to extend the aforementioned theoretical framework to include integrable systems with two spatial dimensions. Focusing on the dispersionless 2D Toda lattice hierarchy, Carlet, Dubrovin, and Mertens [10] precisely uncovered an infinite-dimensional Frobenius manifold. This manifold exists on a space of pairs of specific meromorphic functions, which are defined on the exterior and interior of the unit circle in the complex plane, respectively.

Building on this approach, several more infinite-dimensional Frobenius manifolds have been constructed. These manifolds underlie the bi-Hamiltonian structures of various integrable systems, including the two-component Kadomtsev-Petviashvili hierarchy of Type B [11], the (N, M)-type Toda lattice hierarchy [12], and the two-component extension of the Kadomtsev-Petviashvili (KP) hierarchy [13]. In a different approach, Raimondo [14] constructed a Frobenius manifold structure related to the dispersionless KP hierarchy on the space of Schwartz function.

The aim of the present paper is to construct infinite-dimensional Frobenius manifolds related to the genus-zero universal Whitham hierarchy. The concept of the universal Whitham hierarchy was introduced by Krichever [15] to describe the deformation of periodic solutions to integrable equations with two spatial dimensions [16]. Subsequent research revealed that this hierarchy has significant applications in various branches of mathematics, such as Seiberg-Witten theory [17, 18, 19, 20, 21], Laplace growth problems [22, 23, 24, 25], and conformal mapping problems [26, 27].

The genus-zero universal Whitham hierarchy serves as an extension of various dispersionless integrable systems, such as the dispersionless KP hierarchy [28] and the dispersionless (N,M)-th KdV hierarchy [29], which has attracted considerable scholarly interest [30, 31, 32, 33, 34, 35, 36, 37, 38]. Consider the following Laurent series of z of the form

$$\lambda_0(z) = z + \sum_{j \geq 1} v_{0,j}(x) z^{-j}, \quad \lambda_i(z) = \sum_{i \geq -1} \hat{v}_{i,j}(x) (z - \varphi_i(x))^i, \quad (1.4)$$

where $i = 1, \dots, m$. The genus-zero universal Whitham hierarchy consists of the following set of evolutionary equations:

$$\frac{\partial \lambda_0(z)}{\partial s^{0,k}} = \{(\lambda_0(z)^k)_+, \lambda_0(z)\}, \quad \frac{\partial \lambda_i(z)}{\partial s^{0,k}} = \{(\lambda_0(z)^k)_+, \lambda_i(z)\}, \quad (1.5)$$

$$\frac{\partial \lambda_0(z)}{\partial s^{i,k}} = \{-(\lambda_i(z)^k)_-, \lambda_0(z)\}, \quad \frac{\partial \lambda_i(z)}{\partial s^{i,k}} = \{-(\lambda_i(z)^k)_-, \lambda_i(z)\}, \quad (1.6)$$

for $k = 1, 2, \dots$, and

$$\frac{\partial \lambda_0(z)}{\partial s^{j,0}} = \{\log(z - \varphi_j(x)), \lambda_0(z)\}, \quad \frac{\partial \lambda_i(z)}{\partial s^{j,0}} = \{\log(z - \varphi_j(x)), \lambda_i(z)\}, \quad (1.7)$$

for $j = 1, \dots, m$, where $(f)_+$ and $(f)_-$ denote the positive and negative parts of the Laurent series expansion of f , respectively, and the Lie bracket $\{ , \}$ is defined as

$$\{f, g\} := \frac{\partial f}{\partial z} \frac{\partial g}{\partial x} - \frac{\partial g}{\partial z} \frac{\partial f}{\partial x}. \quad (1.8)$$

In contrast to the previously discussed integrable hierarchies, the existence of a bi-Hamiltonian structure for the genus-zero Whitham hierarchy remains unclear. Consequently, our investigation involves constructing a new class of infinite-dimensional Frobenius manifolds structure on the phase space of the genus-zero Whitham hierarchy, specifying the form of the associated principal hierarchies, and clarifying their relationship to the Whitham hierarchy. However, the construction of the principal hierarchy for a given infinite-dimensional Frobenius manifold presents a formidable challenge. Although Carlet and Mertens [39] have constructed the principal hierarchy for the Frobenius manifold related to the 2D Toda hierarchy, and Raimondo [14] has done the same for the Frobenius manifold related to the dispersionless KP hierarchy, their methods rely on the special structures of the respective Frobenius manifolds and lack generalizability.

To tackle this challenge, we have noted that the system of partial differential equations (1.1) for certain Frobenius manifolds can be reduced to an algebraic equation involving their superpotential. By finding the homogeneous solutions to this algebraic equation, we can determine the Hamiltonian density of the principal hierarchy. Employing this method, we have successfully constructed the principal hierarchies for a selection of Frobenius manifolds, as detailed in our recent works [40, 41]. Building on this foundation, this paper extends the method used to construct the principal hierarchies for Frobenius manifolds related to the genus-zero universal Whitham hierarchy.

1.2. Main results. Let D_1, \dots, D_m be non-intersecting disks in \mathbb{C} with boundaries $\gamma_1, \dots, \gamma_m$, respectively. Denote $\mathbf{D} = \bigcup_{s=1}^m D_s$, and $\mathring{\mathbf{D}}$ the interior of \mathbf{D} . Let \mathcal{H} be the space of germs of functions that are holomorphic in some neighborhood of

$\bigcup_{s=1}^m \gamma_s$. For any $f(z) \in \mathcal{H}$, we define the positive and negative parts of $f(z)$ as follows:

$$f(z)_+ := \frac{1}{2\pi i} \sum_{s=1}^m \int_{\gamma_s} \frac{f(p)}{p-z} dp, \quad z \in \mathring{\mathbf{D}}, \quad f(z)_- := -\frac{1}{2\pi i} \sum_{s=1}^m \int_{\gamma_s} \frac{f(p)}{p-z} dp, \quad z \in \mathbf{D}^c,$$

where \mathbf{D}^c is the complement of \mathbf{D} in \mathbb{P}^1 . It is clear that both $f(z)_+$ and $f(z)_-$ can be analytically extended beyond the boundary of \mathbf{D} , thereby qualifying them as elements of \mathcal{H} , and $f(z) = f(z)_+ + f(z)_-$. Moreover, this type of decomposition is unique under the condition $f_-(\infty) = 0$. As an example, consider $f(z) \in \mathcal{H}$ that can be analytically continued as a meromorphic function on \mathbf{D} with poles $\{\varphi_1, \dots, \varphi_m\}$ where $\varphi_i \in D_i$ for $i = 1, \dots, m$. Then, we have

$$f(z)_- = \sum_{j=1}^m (f(z))_{\varphi_j, \leq -1},$$

where $(f(z))_{\varphi_j, \leq -1}$ denotes the principal part of $f(z)$ at the pole φ_j .

Set n_0, \dots, n_m to be positive integers, and d_1, \dots, d_m to be non-zero integers. Let \mathcal{M} be a subset of $\mathcal{H} \times \mathcal{H}$, consisting of pairs $(a(z), \hat{a}(z))$, where $a(z)$ can be meromorphically extended to \mathbf{D}^c with a single pole at $\{\infty\}$, and $\hat{a}(z)$ can be meromorphically extended to $\mathring{\mathbf{D}}$ with single poles at $\{\varphi_i\}_{i=1}^m$, where $\varphi_i \in D_i$ for $i = 1, \dots, m$. Moreover, they have the following expansions near their poles:

$$\begin{aligned} a(z) &= z^{n_0} + a_{n_0-2} z^{n_0-2} + a_{n_0-3} z^{n_0-3} + \dots, \quad z \rightarrow \infty, \\ \hat{a}(z) &= \hat{a}_{j, -n_j} (z - \varphi_j)^{-n_j} + \hat{a}_{j, -n_j+1} (z - \varphi_j)^{-n_j+1} + \dots, \quad z \rightarrow \varphi_j, \quad j = 1, \dots, m, \end{aligned}$$

and the following conditions are fulfilled:

- (C1) the coefficient $\hat{a}_{j, -n_j} \neq 0$ for $j = 1, \dots, m$;
- (C2) For any $j = 1, \dots, m$, there exists a holomorphic function $w_j(z)$ defined on some neighborhood of γ_j such that

$$\zeta(z)|_{\gamma_j} = w_j(z)^{d_j}, \quad w'(z)|_{\gamma_j} \neq 0,$$

where $\zeta(z) = a(z) - \hat{a}(z)$, and $w_j(z)$ maps γ_j to a path around the origin with winding number 1.

Such a subset $\mathcal{M} = \mathcal{M}_{n_0, n_1, \dots, n_m}^{d_1, d_2, \dots, d_m}$ can be regarded as an infinite-dimensional manifold. The coordinates of this manifold can be effectively chosen as $\{a_j\}_{j \leq n_0-2} \cup \{\hat{a}_{1,j}\}_{j \geq -n_1} \cup \dots \cup \{\hat{a}_{m,j}\}_{j \geq -n_m} \cup \{\varphi_j\}_{j=1}^m$.

Theorem 1.1. *There exists an infinite-dimensional Frobenius manifold structure $(\mathcal{M}, \circ, e, \eta, E)$ of charge $d = 1 - \frac{2}{n_0}$ on the manifold \mathcal{M} , where the flat metric η , the multiplication structure \circ , the unity vector field e , and the Euler vector field E are given by equalities (2.1), (2.27), (2.28), and (2.37), respectively.*

For the infinite-dimensional Frobenius manifold \mathcal{M} , the flat coordinates of the metric η are given by

$$\{t_{i,s} | 1 \leq i \leq m, s \in \mathbb{Z}\} \cup \{h_{0,j} | 1 \leq j \leq n_0 - 1\} \cup \{h_{k,r} | 1 \leq k \leq m, 0 \leq r \leq n_k\},$$

with detailed definitions provided in Section 2.1.

Theorem 1.2. *The Hamiltonian densities of the principal hierarchy for the infinite-dimensional Frobenius manifold \mathcal{M} are defined as follows:*

$$\theta_{t_{i,s},p} = \frac{1}{2\pi i} \frac{1}{(p+1)!} \frac{d_i}{s+d_i} \int_{\gamma_i} \zeta^{\frac{s}{d_i}} \phi_{p+1} dz, \quad s \neq -d_i;$$

$$\theta_{t_{i,-d_i},p} = \frac{d_i}{2\pi i} \int_{\gamma_i} \frac{a^p}{p!} \log \frac{\zeta^{\frac{1}{d_i}}}{(z-\varphi_i)} dz + d_i \operatorname{Res}_{\infty} \frac{a^p}{p!} \left(\log \frac{a^{\frac{1}{n_0}}}{z-\varphi_i} - \frac{c_p}{n_0} \right) dz - \frac{d_i}{2\pi i} \sum_{i' \neq i} \int_{\gamma_{i'}} \frac{a^p}{p!} \log(z-\varphi_{i'}) dz;$$

$$\theta_{h_{0,j},p} = -\frac{\Gamma\left(1 - \frac{j}{n_0}\right)}{\Gamma\left(2 + p - \frac{j}{n_0}\right)} \operatorname{Res}_{z=\infty} a^{1+p-\frac{j}{n_0}} dz;$$

$$\theta_{h_{k,r},p} = \frac{\Gamma\left(1 - \frac{r}{n_k}\right)}{\Gamma\left(2 + p - \frac{r}{n_k}\right)} \operatorname{Res}_{z=\varphi_k} \hat{a}^{1+p-\frac{r}{n_k}} dz, \quad r \neq n_k;$$

$$\theta_{h_{k,n_k},p} = \frac{n_k}{2\pi i} \int_{\gamma_k} \frac{\hat{a}^p}{p!} \log \frac{\zeta^{\frac{1}{n_k}}}{z-\varphi_k} dz + n_k \operatorname{Res}_{\varphi_k} \frac{\hat{a}^p}{p!} \left(\log \hat{a}^{\frac{1}{n_k}}(z-\varphi_k) - \frac{c_p}{n_k} \right) dz - \frac{n_k}{d_k} \theta_{t_{k,-d_k},p}.$$

Here,

$$\zeta(z) = a(z) - \hat{a}(z), \quad \phi_p(z) = a^p(z) - \hat{a}^p(z), \quad c_0 = 0, \quad c_p = \sum_{s=1}^p \frac{1}{s}.$$

Moreover, the constant matrix μ and $R = R_1$ have the form

$$\mu_u = \begin{cases} \frac{s}{d_i} + \frac{1}{2}, & u = t_{i,s}; \\ \frac{1}{2} - \frac{j}{n_0}, & u = h_{0,j}; \\ \frac{1}{2} - \frac{r}{n_k}, & u = h_{k,r}, \end{cases}$$

and

$$(R_1)_v^u = \begin{cases} \delta_{i,j} - \frac{d_i}{n_0}, & u \in \cup_{j=1}^m \{t_{j,0}, h_{j,0}\}, v = t_{i,-d_i}; \\ \frac{n_i}{n_0} + 1, & u = h_{i,0}, v = h_{i,n_i}; \\ \frac{n_i}{n_0} - \frac{n_i}{d_i}, & u = h_{i,0}, v = h_{i,n_i}; \\ \frac{n_i}{n_0}, & u \in \cup_{j \neq i} \{t_{j,0}, h_{j,0}\}, v = h_{i,n_i}; \\ 0, & \text{for other cases.} \end{cases}$$

Let us now consider the relationship between the principal hierarchy of \mathcal{M} and the genus-zero Whitham hierarchy. The principal hierarchy of \mathcal{M} is given by $\frac{\partial}{\partial T^{\alpha,p-1}} =$

$\mathcal{P}(d\theta_{\alpha,p+1})$ for $\{\theta_{\alpha,p}\}$ defined above and the Hamiltonian operator \mathcal{P} is specified by (2.24). We make the following assumptions regarding the asymptotic behavior of the functions $a(z)$ and $\hat{a}(z)$:

$$a(z) = \lambda_0(z)^{n_0}, \quad z \rightarrow \infty, \quad \hat{a}(z) = \lambda_j(z)^{n_j}, \quad z \rightarrow \varphi_j, \quad j = 1, \dots, m,$$

where the series $\lambda_i(z)$ for $i = 0, \dots, m$ are given by (1.4).

Theorem 1.3. *The principal hierarchy of the infinite-dimensional Frobenius manifold \mathcal{M} can be regarded as an extension of the genus-zero Whitham hierarchy. Specifically, we have the following relations:*

$$\begin{aligned} \frac{\partial}{\partial T^{h_0,j,p}} &= \frac{\Gamma(1 - \frac{j}{n_0})}{\Gamma(2 + p - \frac{j}{n_0})} \frac{\partial}{\partial S^{0,(p+1)n_0-j}}, \quad j = 1, \dots, n_0 - 1; \\ \sum_{k=1}^m \left(\frac{\partial}{\partial T^{t_{k,0,p}}} + \frac{\partial}{\partial T^{h_{k,0,p}}} \right) &= \frac{1}{(p+1)!} \frac{\partial}{\partial S^{0,(p+1)n_0}}, \\ \frac{1}{n_j} \frac{\partial}{\partial T^{h_{i,0,0}}} &= \frac{\partial}{\partial S^{i,0}}, \\ \frac{\partial}{\partial T^{h_{i,j,p}}} &= \frac{\Gamma(1 - \frac{j}{n_i})}{\Gamma(2 + p - \frac{j}{n_i})} \frac{\partial}{\partial S^{i,(p+1)n_i-j}}, \quad i = 1, \dots, m, \quad j = 0, \dots, n_i - 1. \end{aligned}$$

The content of this paper is arranged as follows. In the second chapter, we construct the Frobenius manifold structure on the space \mathcal{M} . We will define the flat metric, the multiplication structure, and the potential on \mathcal{M} , respectively. Chapter three focuses on the principal hierarchy of \mathcal{M} , including the proofs of Theorems 1.2 and 1.3.

2. INFINITE-DIMENSIONAL FROBENIUS MANIFOLD STRUCTURE ON \mathcal{M}

The objective of this section is to construct an infinite-dimensional Frobenius manifold structure on the space \mathcal{M} , encompassing the definition of its flat metric, potential, unity vector field, and Euler vector field.

2.1. Flat metric. At each point $\vec{a} = (a(z), \hat{a}(z)) \in \mathcal{M}$, we identify a vector X in the tangent space $T_{\vec{a}}\mathcal{M}$ with its action $(\partial_X a(z), \partial_X \hat{a}(z))$. Define a bilinear form on $T_{\vec{a}}\mathcal{M}$ as follows:

$$\langle X_1, X_2 \rangle_{\eta} = -\frac{1}{2\pi i} \sum_{j=1}^m \int_{\gamma_j} \frac{\partial_1 \zeta(z) \cdot \partial_2 \zeta(z)}{\zeta'(z)} dz - \left(\text{Res}_{z=\infty} + \sum_{j=1}^m \text{Res}_{z=\varphi_j} \right) \frac{\partial_1 \ell(z) \cdot \partial_2 \ell(z)}{\ell'(z)} dz, \quad (2.1)$$

where

$$\zeta(z) = a(z) - \hat{a}(z), \quad \ell(z) = a(z)_+ + \hat{a}(z)_-,$$

and $\partial_\nu = \partial_{X_\nu}$ denotes the derivative along the vector $X_\nu \in T_{\bar{a}}\mathcal{M}$ for $\nu = 1, 2$. It is clear that the bilinear form is nondegenerate. Let us proceed to construct the flat coordinates for this metric, employing an approach analogous to that presented in [13].

In accordance with condition (C2) of the manifold \mathcal{M} , for each $j = 1, \dots, m$, there exists a holomorphic function $w_j(z)$ defined on some neighborhood of γ_j . This function satisfies the following properties:

$$\zeta(z)|_{\gamma_j} = w_j(z)^{d_j}, \quad w'(z)|_{\gamma_j} \neq 0,$$

where $\zeta(z) = a(z) - \hat{a}(z)$, and $w_j(z)$ maps γ_j to a path encircling the origin with winding number 1. The inverse of these functions is expressible via a Laurent series expansion as follows:

$$z(w_j) = \sum_{s \in \mathbb{Z}} t_{j,s} w_j^s, \quad w_j \in \gamma'_j, \quad j = 1, \dots, m, \quad (2.2)$$

where γ'_j is the image of γ_j under the map w_j . The coefficients $t_{j,s}$ are ascertained by the contour integral

$$t_{j,s} = \frac{1}{2\pi i} \oint_{\gamma'_j} z(w_j) w_j^{-s-1} dw_j, \quad j = 1, \dots, m, \quad s \in \mathbb{Z}.$$

In addition, we introduce a family of functions $\chi_j(z)$, defined in punctured neighborhoods of ∞ for $j = 0$ and φ_j for $j = 1, \dots, m$, respectively. These functions are defined as:

$$\begin{aligned} \chi_0(z) &:= \ell(z)^{\frac{1}{n_0}} = z + b_{0,1}z^{-1} + b_{0,2}z^{-2} + \dots, \quad z \rightarrow \infty, \\ \chi_j(z) &:= \ell(z)^{\frac{1}{n_j}} = \hat{a}_{j,-n_j}^{\frac{1}{n_j}}(z - \varphi_j)^{-1} + b_{j,0} + b_{j,1}(z - \varphi_j) + \dots, \quad z \rightarrow \varphi_j. \end{aligned}$$

The inverse functions of $\chi_j(z)$ can be expressed as Laurent series:

$$z(\chi_0) = \chi_0 - h_{0,1}\chi_0^{-1} - h_{0,2}\chi_0^{-2} - \dots - h_{0,n_0-1}\chi_0^{-n_0+1} + \dots, \quad z \rightarrow \infty, \quad (2.3)$$

$$z(\chi_j) = h_{j,0} + h_{j,1}\chi_j^{-1} + h_{j,2}\chi_j^{-2} + \dots + h_{j,n_j}\chi_j^{-n_j} + \dots, \quad z \rightarrow \varphi_j. \quad (2.4)$$

Then, the variables $\mathbf{t} \cup \mathbf{h}$, where

$$\begin{aligned} \mathbf{t} &:= \{t_{i,s} | 1 \leq i \leq m, s \in \mathbb{Z}\}, \\ \mathbf{h} &:= \{h_{0,j} | 1 \leq j \leq n_0 - 1\} \cup \{h_{k,r} | 1 \leq k \leq m, 0 \leq r \leq n_k\}, \end{aligned}$$

form a set of coordinates for the manifold \mathcal{M} . Moreover, we have

$$\begin{aligned} \frac{\partial \zeta(z)}{\partial t_{i,s}}|_{\gamma_i} &= -\zeta(z)^{\frac{s}{d_i}} \zeta'(z), & \frac{\partial \zeta(z)}{\partial t_{i,s}}|_{\gamma_{i'}} &= 0, \quad i' \neq i, & \frac{\partial \ell(z)}{\partial t_{i,s}} &= 0; \\ \frac{\partial \zeta(z)}{\partial h_{0,j}} &= 0, & \frac{\partial \ell(z)}{\partial h_{0,j}} &= (\ell'(z) \chi_0(z)^{-j})_{\infty, \geq 0}, \\ \frac{\partial \zeta(z)}{\partial h_{k,r}} &= 0, & \frac{\partial \ell(z)}{\partial h_{k,r}} &= -(\ell'(z) \chi_k(z)^{-r})_{\varphi_k, \leq -1}, \end{aligned}$$

which imply the non-vanishing components of the metric are given by

$$\begin{aligned} \left\langle \frac{\partial}{\partial t_{i,s}}, \frac{\partial}{\partial t_{i,s'}} \right\rangle_{\eta} &= -d_i \delta_{-d_i, s+s'}, \quad i = 1, \dots, m, \quad s, s' \in \mathbb{Z}; \\ \left\langle \frac{\partial}{\partial h_{0,j}}, \frac{\partial}{\partial h_{0,j'}} \right\rangle_{\eta} &= n_0 \delta_{n_0, j+j'}, \quad j, j' = 1, \dots, n_0 - 1; \\ \left\langle \frac{\partial}{\partial h_{k,r}}, \frac{\partial}{\partial h_{k,r'}} \right\rangle_{\eta} &= n_k \delta_{n_k, r+r'}, \quad k = 1, \dots, m, \quad r, r' = 0, \dots, n_k. \end{aligned}$$

The proof of these results follows a similar approach to that presented in [13].

Corollary 2.1. *The bilinear form $\langle \cdot, \cdot \rangle_{\eta}$ defines a flat metric on the manifold \mathcal{M} . The associated flat coordinates satisfy*

$$\frac{\partial \vec{a}}{\partial t_{i,s}} = (-\zeta(z)^{\frac{s}{d_i}} \zeta'(z) \mathbf{1}_{\gamma_i})_-, (\zeta(z)^{\frac{s}{d_i}} \zeta'(z) \mathbf{1}_{\gamma_i})_+, \quad s \in \mathbb{Z}; \quad (2.5)$$

$$\frac{\partial \vec{a}}{\partial h_{0,j}} = ((\ell'(z) \chi_0(z)^{-j})_{\infty, \geq 0}, (\ell'(z) \chi_0(z)^{-j})_{\infty, \geq 0}), \quad 1 \leq j \leq n_0 - 1; \quad (2.6)$$

$$\frac{\partial \vec{a}}{\partial h_{k,r}} = (-\ell'(z) \chi_k(z)^{-r})_{\varphi_k, \leq -1}, -\ell'(z) \chi_k(z)^{-r})_{\varphi_k, \leq -1}), \quad 0 \leq r \leq n_k, \quad (2.7)$$

where the functions $\{\mathbf{1}_{\gamma_i}\}_{i=1}^m$ belonging to the space \mathcal{H} are characterized by

$$\mathbf{1}_{\gamma_i}|_{\gamma_j} = \delta_{ij}, \quad i, j = 1, \dots, m.$$

2.2. Hamiltonian structure on LM . In this section, we aim to construct the dispersionless Hamiltonian structure on the loop space of \mathcal{M} corresponding to the flat metric $\langle \cdot, \cdot \rangle_{\eta}$. To achieve this, we must provide a suitable description of the cotangent space $T_{\vec{a}}^* \mathcal{M}$ for any $\vec{a} \in \mathcal{M}$.

For an element $\vec{\omega} = (\omega(z), \hat{\omega}(z))$ in $\mathcal{H} \times \mathcal{H}$, we define the pairing as follows:

$$\langle \vec{\omega}, X \rangle = \frac{1}{2\pi i} \sum_{s=1}^m \int_{\gamma_s} (\omega(z) \partial_X a(z) + \hat{\omega}(z) \partial_X \hat{a}(z)) dz. \quad (2.8)$$

This pairing induces a map from $\mathcal{H} \times \mathcal{H}$ to the cotangent space $T_{\vec{a}}^* \mathcal{M}$. The image of this map, which we refer to as the "canonical cotangent space", is denoted by $\widetilde{T_{\vec{a}}^* \mathcal{M}}$.

Lemma 2.2. Let η be a linear map from $\mathcal{H} \times \mathcal{H}$ to $T_{\vec{a}}\mathcal{M}$ defined as

$$\begin{aligned} \eta \cdot \vec{\omega} = & (a'(z)(\omega(z) + \hat{\omega}(z))_- - (\omega(z)a'(z) + \hat{\omega}(z)\hat{a}'(z))_- , \\ & - \hat{a}'(z)(\omega(z) + \hat{\omega}(z))_+ + (\omega(z)a'(z) + \hat{\omega}(z)\hat{a}'(z))_+). \end{aligned} \quad (2.9)$$

Then, for any $X \in T_{\vec{a}}\mathcal{M}$, we have

$$\langle \vec{\omega}, X \rangle = \langle \eta \cdot \vec{\omega}, X \rangle_{\eta}. \quad (2.10)$$

Proof. Let us denote

$$\eta \cdot \vec{\omega} = (\xi_1(z), \hat{\xi}_1(z)), \quad \partial_X \vec{a} = (\xi_2(z), \hat{\xi}_2(z)).$$

Then, we have

$$\langle \vec{\omega}, X \rangle = \frac{1}{2\pi i} \sum_{j=1}^m \int_{\gamma_j} (\omega(z)\xi_2(z) + \hat{\omega}(z)\hat{\xi}_2(z)) dz = \mathcal{I}_1 + \mathcal{I}_2 + \mathcal{I}_3,$$

where

$$\begin{aligned} \mathcal{I}_1 &= \frac{1}{2\pi i} \sum_{j=1}^m \int_{\gamma_j} (\omega(z)_+ - \hat{\omega}(z)_-) (\xi_2(z) - \hat{\xi}_2(z)) dz, \\ \mathcal{I}_2 &= \frac{1}{2\pi i} \sum_{j=1}^m \int_{\gamma_j} (\omega(z)_- + \hat{\omega}(z)_-) \xi_2(z) dz, \\ \mathcal{I}_3 &= \frac{1}{2\pi i} \sum_{j=1}^m \int_{\gamma_j} (\omega(z)_+ + \hat{\omega}(z)_+) \hat{\xi}_2(z) dz. \end{aligned}$$

From equality (2.9), we derive

$$\xi_1(z) - \hat{\xi}_1(z) = \zeta'_1(z)(\omega(z)_+ - \hat{\omega}(z)_-) \quad (2.11)$$

and

$$\left(\frac{\xi_1(z)}{a'(z)} \right)_{\infty, \geq -n_0+1} = ((\omega(z) + \hat{\omega}(z))_-)_{\infty, \geq -n_0+1}, \quad (2.12)$$

$$\left(\frac{\hat{\xi}_1(z)}{\hat{a}'(z)} \right)_{\varphi_j, \leq n_j} = ((\omega(z) + \hat{\omega}(z))_+)_{\varphi_j, \leq n_j}, \quad j = 1, \dots, m. \quad (2.13)$$

Applying equality (2.11), we have

$$\mathcal{I}_1 = -\frac{1}{2\pi i} \sum_{j=1}^m \int_{\gamma_j} \frac{(\xi_1(z) - \hat{\xi}_1(z))(\xi_2(z) - \hat{\xi}_2(z))}{\zeta'(z)} dz = -\frac{1}{2\pi i} \sum_{j=1}^m \int_{\gamma_j} \frac{\partial_1 \zeta(z) \cdot \partial_2 \zeta(z)}{\zeta'(z)} dz.$$

For \mathcal{I}_2 and \mathcal{I}_3 , following a similar calculation, we obtain

$$\begin{aligned}
\mathcal{I}_2 &= \frac{1}{2\pi i} \sum_{j=1}^m \int_{\gamma_j} ([\omega(z) + \hat{\omega}(z)]_-)_{\infty, \geq -n_0+1} \xi_2(z) dz \\
&= \frac{1}{2\pi i} \sum_{j=1}^m \int_{\gamma_j} \left(\frac{\partial_1 a(z)}{a'(z)} \right)_{\infty, \geq -n_0+1} \partial_2 a(z) dz \\
&= \frac{1}{2\pi i} \sum_{j=1}^m \int_{\gamma_j} \left(\frac{\partial_1 \ell(z)}{a'(z)} \right)_{\infty, \geq -n_0+1} \partial_2 \ell(z) dz \\
&= \frac{1}{2\pi i} \sum_{j=1}^m \int_{\gamma_j} \left(\frac{\partial_1 \ell(z)}{\ell'(z)} \right)_{\infty, \geq -n_0+1} \partial_2 \ell(z) dz \\
&= - \operatorname{Res}_{z=\infty} \frac{\partial_1 \ell(z) \cdot \partial_2 \ell(z)}{\ell'(z)} dz,
\end{aligned}$$

and

$$\begin{aligned}
\mathcal{I}_3 &= \frac{1}{2\pi i} \sum_{j=1}^m \oint_{\gamma_j} ([\omega(z) + \hat{\omega}(z)]_+)_{\varphi_j, \leq n_j} \hat{\xi}_2(z) dz \\
&= -\frac{1}{2\pi i} \sum_{j=1}^m \oint_{\gamma_j} \left(\frac{\partial_1 \hat{a}(z)}{\hat{a}'(z)} \right)_{\varphi_j, \leq n_j} \partial_2 \hat{a}(z) dz \\
&= -\frac{1}{2\pi i} \sum_{j=1}^m \oint_{\gamma_j} \left(\frac{\partial_1 \ell(z)}{\hat{a}'(z)} \right)_{\varphi_j, \leq n_j} \partial_2 \ell(z) dz \\
&= -\frac{1}{2\pi i} \sum_{j=1}^m \oint_{\gamma_j} \left(\frac{\partial_1 \ell(z)}{\ell'(z)} \right)_{\varphi_j, \leq n_j} \partial_2 \ell(z) dz \\
&= - \sum_{j=1}^m \operatorname{Res}_{z=\varphi_j} \frac{\partial_1 \ell(z) \cdot \partial_2 \ell(z)}{\ell'(z)} dz.
\end{aligned}$$

Thus, based on the calculated results of \mathcal{I}_1 , \mathcal{I}_2 , and \mathcal{I}_3 , the lemma is proved. \square

The subsequent lemma establishes the surjectivity of η .

Lemma 2.3. *For any $\vec{\xi} = (\xi(z), \hat{\xi}(z)) \in T_{\vec{a}}\mathcal{M}$, consider $\vec{\omega} = (\omega(z), \hat{\omega}(z)) \in \mathcal{H} \times \mathcal{H}$ such that*

$$\begin{aligned}
\omega(z)_+ &= - \left(\frac{\xi(z) - \hat{\xi}(z)}{\zeta'(z)} \right)_+, \\
\omega(z)_- &= \left(\frac{\xi(z)}{a'(z)} - \left(\frac{\xi(z) - \hat{\xi}(z)}{\zeta'(z)} \right)_- \right)_{\infty, \geq -n_0+1}, \\
\hat{\omega}(z)_- &= \left(\frac{\xi(z) - \hat{\xi}(z)}{\zeta'(z)} \right)_-,
\end{aligned}$$

$$\hat{\omega}(z)_+|_{D_j} = \left(-\frac{\hat{\xi}(z)}{\hat{a}'(z)} + \left(\frac{\xi(z) - \hat{\xi}(z)}{\zeta'(z)} \right)_{+\varphi_j \leq n_j} \right), \quad j = 1, \dots, m,$$

then it follows that $\eta \cdot \vec{\omega} = \vec{\xi}$.

Proof. Let $\vec{\omega} \in \mathcal{H} \times \mathcal{H}$ be as defined in the lemma, and denote $\eta \cdot \vec{\omega} = (I, J)$. Then we have

$$\begin{aligned} I_+ &= (a'(z)[\omega(z) + \hat{\omega}(z)]_-)_+ \\ &= (a'(z)([\omega(z) + \hat{\omega}(z)]_-)_{\infty, \geq -n_0+1})_+ \\ &= (a'(z)\left(\frac{\xi(z)}{a'(z)}\right)_{\infty, \geq -n_0+1})_+ \\ &= \xi(z)_+, \end{aligned}$$

and

$$\begin{aligned} J_- &= -\sum_{j=1}^m (\hat{a}'(z)[\omega(z) + \hat{\omega}(z)]_+)_{\varphi_j, \leq -1} \\ &= -\sum_{j=1}^m (\hat{a}'(z)([\omega(z) + \hat{\omega}(z)]_+)_{\varphi_j, \leq n_j})_{\varphi_j, \leq -1} \\ &= \sum_{j=1}^m (\hat{a}'(z)\left(\frac{\hat{\xi}(z)}{\hat{a}'(z)}\right)_{\varphi_j, \leq n_j})_{\varphi_j, \leq -1} \\ &= \sum_{j=1}^m \hat{\xi}(z)_{\varphi_j, \leq -1} \\ &= \hat{\xi}(z)_-. \end{aligned}$$

Additionally, we observe that

$$\begin{aligned} I - J &= (a'(z) - \hat{a}'(z))(\hat{\omega}(z)_- - \omega(z)_+) \\ &= \zeta'(z)(\hat{\omega}(z)_- - \omega(z)_+) \\ &= \xi(z) - \hat{\xi}(z), \end{aligned}$$

which implies $I_- = \xi(z)_-$ and $J_+ = \hat{\xi}(z)_+$. The lemma is proved. \square

Corollary 2.4. *Let $\vec{\xi} \in T_{\bar{a}}\mathcal{M}$. Then, $\vec{\xi} = 0$ if and only if for any $\vec{\omega} \in \mathcal{H} \times \mathcal{H}$, it holds that $\langle \vec{\omega}, \vec{\xi} \rangle = 0$. Consequently, η induces a surjective map from $\widetilde{T_{\bar{a}}^* \mathcal{M}}$ to $T_{\bar{a}}\mathcal{M}$.*

Proof. Suppose $\vec{\xi} \in T_{\bar{a}}\mathcal{M}$ is such that for any $\vec{\omega} \in \mathcal{H} \times \mathcal{H}$, we have $\langle \vec{\omega}, \vec{\xi} \rangle = 0$. Then, it follows that $\langle \eta \cdot \vec{\omega}, \vec{\xi} \rangle_{\eta} = 0$. Since η is surjective and the metric $\langle \cdot, \cdot \rangle_{\eta}$ is non-degenerate, we deduce that $\vec{\xi} = 0$. This completes the proof. \square

According to the Dubrovin-Novikov theorem, the Hamiltonian operator corresponding to the metric $\langle \cdot, \cdot \rangle_\eta$ is defined by

$$\mathcal{P}(\vec{\omega}) = \eta \cdot \nabla_{\frac{\partial}{\partial x}} \vec{\omega}, \quad (2.14)$$

where ∇ denotes the Levi-Civita connection associated with $\langle \cdot, \cdot \rangle_\eta$, and $\vec{\omega}$ is a covector field on \mathcal{M} that depends on the loop parameter x .

The explicit form of \mathcal{P} can be derived using the following formula:

$$\partial_1 \langle \vec{\omega}, \partial_2 \rangle = \langle \eta \cdot \nabla_{\partial_1} \vec{\omega}, \partial_2 \rangle_\eta + \langle \vec{\omega}, \nabla_{\partial_1} \partial_2 \rangle, \quad \partial_1, \partial_2 \in \mathbf{Vect}(\mathcal{M}).$$

Lemma 2.5. *Let ∂_1 and ∂_2 be vector fields on \mathcal{M} . Then, the covariant derivative of ∂_2 along ∂_1 is given by*

$$\begin{aligned} (\nabla_{\partial_1} \partial_2) \cdot \vec{a} = & \left(\partial_1 \partial_2 a(z) - \left(\frac{\partial_1 \zeta(z) \partial_2 \zeta(z)}{\zeta'(z)} \right)' - \left(\frac{\partial_1 \ell(z) \partial_2 \ell(z)}{\ell'(z)} \right)'_{\infty, \geq 0} - \sum_{j=1}^m \left(\frac{\partial_1 \ell(z) \partial_2 \ell(z)}{\ell'(z)} \right)'_{\varphi_j, \leq -1} \right), \\ & \partial_1 \partial_2 \hat{a}(z) + \left(\frac{\partial_1 \zeta(z) \partial_2 \zeta(z)}{\zeta'(z)} \right)'_+ - \left(\frac{\partial_1 \ell(z) \partial_2 \ell(z)}{\ell'(z)} \right)'_{\infty, \geq 0} - \sum_{j=1}^m \left(\frac{\partial_1 \ell(z) \partial_2 \ell(z)}{\ell'(z)} \right)'_{\varphi_j, \leq -1} \end{aligned} \quad (2.15)$$

Proof. Let us verify that the connection ∇ defined by equality (2.15) is torsion-free and compatible with the metric $\langle \cdot, \cdot \rangle_\eta$. The torsion-free property of the connection ∇ is characterized by the following identity

$$(\nabla_{\partial_1} \partial_2) \cdot \vec{a} - (\nabla_{\partial_2} \partial_1) \cdot \vec{a} = \partial_1 \partial_2 \vec{a} - \partial_2 \partial_1 \vec{a},$$

which can be readily deduced from formula (2.15).

The condition for the connection ∇ to be compatible with the metric $\langle \cdot, \cdot \rangle_\eta$ is given by the following identity:

$$\nabla_{\partial_1} \langle \partial_2, \partial_3 \rangle_\eta = \langle \nabla_{\partial_1} \partial_2, \partial_3 \rangle_\eta + \langle \partial_2, \nabla_{\partial_1} \partial_3 \rangle_\eta. \quad (2.16)$$

To verify this identity, we apply the Leibniz rule and utilize the formula (2.1). This leads to the following computation:

$$\begin{aligned} & \nabla_{\partial_1} \langle \partial_2, \partial_3 \rangle_\eta \\ = & -\frac{1}{2\pi i} \sum_{j=1}^m \int_{\gamma_j} \left(\frac{\partial_1 \partial_2 \zeta(z) \cdot \partial_3 \zeta(z)}{\zeta'(z)} + \frac{\partial_2 \zeta(z) \cdot \partial_1 \partial_3 \zeta(z)}{\zeta'(z)} - \frac{\partial_2 \zeta(z) \cdot \partial_3 \zeta(z) \cdot \partial_1 \zeta'(z)}{(\zeta'(z))^2} \right) dz \\ & - \text{Res}_{z=\infty} \left(\frac{\partial_1 \partial_2 \ell(z) \cdot \partial_3 \ell(z)}{\ell'(z)} + \frac{\partial_2 \ell(z) \cdot \partial_1 \partial_3 \ell(z)}{\ell'(z)} - \frac{\partial_2 \ell(z) \cdot \partial_3 \ell(z) \cdot \partial_1 \ell'(z)}{(\ell'(z))^2} \right) dz \\ & - \sum_{j=1}^m \text{Res}_{z=\varphi_j} \left(\frac{\partial_1 \partial_2 \ell(z) \cdot \partial_3 \ell(z)}{\ell'(z)} + \frac{\partial_2 \ell(z) \cdot \partial_1 \partial_3 \ell(z)}{\ell'(z)} - \frac{\partial_2 \ell(z) \cdot \partial_3 \ell(z) \cdot \partial_1 \ell'(z)}{(\ell'(z))^2} \right) dz. \end{aligned} \quad (2.17)$$

By using the commutativity of ∂_ν and ∂_z for $\nu = 1, 2$, and applying integration by parts, we derive the following results:

$$\begin{aligned}
& \frac{1}{2\pi i} \sum_{j=1}^m \int_{\gamma_j} \frac{\partial_2 \zeta(z) \cdot \partial_3 \zeta(z) \cdot \partial_1 \zeta'(z)}{(\zeta'(z))^2} dz \\
&= -\frac{1}{2\pi i} \sum_{j=1}^m \int_{\gamma_j} \left(\frac{\partial_2 \zeta(z)}{\zeta'(z)} \right)' \frac{\partial_3 \zeta(z) \cdot \partial_1 \zeta(z)}{\zeta'(z)} dz - \frac{1}{2\pi i} \sum_{j=1}^m \int_{\gamma_j} \left(\frac{\partial_3 \zeta(z)}{\zeta'(z)} \right)' \frac{\partial_2 \zeta(z) \cdot \partial_1 \zeta(z)}{\zeta'(z)} dz \\
&= \frac{1}{2\pi i} \sum_{j=1}^m \int_{\gamma_j} \frac{\partial_2 \zeta(z)}{\zeta'(z)} \left(\frac{\partial_3 \zeta(z) \cdot \partial_1 \zeta(z)}{\zeta'(z)} \right)' dz + \frac{1}{2\pi i} \sum_{j=1}^m \int_{\gamma_j} \frac{\partial_3 \zeta(z)}{\zeta'(z)} \left(\frac{\partial_2 \zeta(z) \cdot \partial_1 \zeta(z)}{\zeta'(z)} \right)' dz,
\end{aligned} \tag{2.18}$$

$$\begin{aligned}
& \text{Res}_{z=\infty} \frac{\partial_2 \ell(z) \cdot \partial_3 \ell(z) \cdot \partial_1 \ell'(z)}{(\ell'(z))^2} dz \\
&= \text{Res}_{z=\infty} \frac{\partial_2 \ell(z)}{\ell'(z)} \left(\frac{\partial_3 \ell(z) \cdot \partial_1 \ell(z)}{\ell'(z)} \right)' dz + \text{Res}_{z=\infty} \frac{\partial_3 \ell(z)}{\ell'(z)} \left(\frac{\partial_2 \ell(z) \cdot \partial_1 \ell(z)}{\ell'(z)} \right)' dz,
\end{aligned} \tag{2.19}$$

and

$$\begin{aligned}
& \text{Res}_{z=\varphi_j} \frac{\partial_2 \ell(z) \cdot \partial_3 \ell(z) \cdot \partial_1 \ell'(z)}{(\ell'(z))^2} dz \\
&= \text{Res}_{z=\varphi_j} \frac{\partial_2 \ell(z)}{\ell'(z)} \left(\frac{\partial_3 \ell(z) \cdot \partial_1 \ell(z)}{\ell'(z)} \right)' dz + \text{Res}_{z=\varphi_j} \frac{\partial_3 \ell(z)}{\ell'(z)} \left(\frac{\partial_2 \ell(z) \cdot \partial_1 \ell(z)}{\ell'(z)} \right)' dz.
\end{aligned} \tag{2.20}$$

Substituting equalities (2.18)-(2.20) into (2.17), we obtain

$$\begin{aligned}
& \nabla_{\partial_1} \langle \partial_2, \partial_3 \rangle_\eta \\
&= -\frac{1}{2\pi i} \sum_{j=1}^m \int_{\gamma_j} \frac{\partial_1 \partial_2 \zeta(z) \cdot \partial_3 \zeta(z)}{\zeta'(z)} dz - \frac{1}{2\pi i} \sum_{j=1}^m \int_{\gamma_j} \frac{\partial_2 \zeta(z) \cdot \partial_1 \partial_3 \zeta(z)}{\zeta'(z)} dz \\
&+ \frac{1}{2\pi i} \sum_{j=1}^m \int_{\gamma_j} \frac{\partial_2 \zeta(z)}{\zeta'(z)} \left(\frac{\partial_3 \zeta(z) \cdot \partial_1 \zeta(z)}{\zeta'(z)} \right)' dz + \frac{1}{2\pi i} \sum_{j=1}^m \int_{\gamma_j} \frac{\partial_3 \zeta(z)}{\zeta'(z)} \left(\frac{\partial_2 \zeta(z) \cdot \partial_1 \zeta(z)}{\zeta'(z)} \right)' dz \\
&- \text{Res}_{z=\infty} \frac{\partial_1 \partial_2 \ell(z) \cdot \partial_3 \ell(z)}{\ell'(z)} dz - \text{Res}_{z=\infty} \frac{\partial_2 \ell(z) \cdot \partial_1 \partial_3 \ell(z)}{\ell'(z)} dz \\
&+ \text{Res}_{z=\infty} \frac{\partial_2 \ell(z)}{\ell'(z)} \left(\frac{\partial_3 \ell(z) \cdot \partial_1 \ell(z)}{\ell'(z)} \right)' dz + \text{Res}_{z=\infty} \frac{\partial_3 \ell(z)}{\ell'(z)} \left(\frac{\partial_2 \ell(z) \cdot \partial_1 \ell(z)}{\ell'(z)} \right)' dz \\
&- \sum_{j=1}^m \text{Res}_{z=\varphi_j} \left(\frac{\partial_1 \partial_2 \ell(z) \cdot \partial_3 \ell(z)}{\ell'(z)} + \frac{\partial_2 \ell(z) \cdot \partial_1 \partial_3 \ell(z)}{\ell'(z)} \right) dz \\
&+ \sum_{j=1}^m \text{Res}_{z=\varphi_j} \frac{\partial_2 \ell(z)}{\ell'(z)} \left(\frac{\partial_3 \ell(z) \cdot \partial_1 \ell(z)}{\ell'(z)} \right)' dz + \sum_{j=1}^m \text{Res}_{z=\varphi_j} \frac{\partial_3 \ell(z)}{\ell'(z)} \left(\frac{\partial_2 \ell(z) \cdot \partial_1 \ell(z)}{\ell'(z)} \right)' dz.
\end{aligned} \tag{2.21}$$

On the other hand, formula (2.15) can be equivalently expressed as

$$(\nabla_{\partial_1} \partial_2) \cdot \zeta(z) = \partial_1 \partial_2 \zeta(z) - \left(\frac{\partial_1 \zeta(z) \partial_2 \zeta(z)}{\zeta'(z)} \right)', \quad (2.22)$$

$$(\nabla_{\partial_1} \partial_2) \cdot \ell(z) = \partial_1 \partial_2 \ell(z) - \left(\frac{\partial_1 \ell(z) \partial_2 \ell(z)}{\ell'(z)} \right)'_{\infty, \geq 0} - \sum_{j=1}^m \left(\frac{\partial_1 \ell(z) \partial_2 \ell(z)}{\ell'(z)} \right)'_{\varphi_j, \leq -1}. \quad (2.23)$$

By combining equalities (2.21), (2.22), and (2.23), we can deduce that formula (2.16) holds. This completes the proof. \square

Corollary 2.6. *The Hamiltonian operator (2.14) takes the explicit form:*

$$\begin{aligned} \mathcal{P}(\vec{\omega}) = & (\{\omega(z), a(z)\}_- + \{\hat{\omega}(z), \hat{a}(z)\}_- - \{\omega(z)_- + \hat{\omega}(z)_-, a(z)\}, \\ & - \{\omega(z), a(z)\}_+ - \{\hat{\omega}(z), \hat{a}(z)\}_+ + \{\omega(z)_+ + \hat{\omega}(z)_+, \hat{a}(z)\}), \end{aligned} \quad (2.24)$$

where $\{f, g\} = f' \partial_x g - g' \partial_x f$.

Proof. For any vector fields ∂_1 and ∂_2 , and a covector field $\vec{\omega}$ on \mathcal{M} , we have

$$\partial_1 \langle \vec{\omega}, \partial_2 \rangle = \langle \eta \cdot \nabla_{\partial_1} \vec{\omega}, \partial_2 \rangle_\eta + \langle \vec{\omega}, \nabla_{\partial_1} \partial_2 \rangle. \quad (2.25)$$

By direct computation, we obtain

$$\partial_1 \langle \vec{\omega}, \partial_2 \rangle = \sum_{j=1}^m \int_{\gamma_j} (\partial_1 \omega(z) \partial_2 a(z) + \omega(z) \partial_1 \partial_2 a(z) + \partial_1 \hat{\omega}(z) \partial_2 \hat{a}(z) + \hat{\omega}(z) \partial_1 \partial_2 \hat{a}(z)) dz,$$

and

$$\begin{aligned} & \langle \vec{\omega}, \nabla_{\partial_1} \partial_2 \rangle \\ = & \frac{1}{2\pi i} \sum_{s=1}^m \int_{\gamma_s} \omega(z) \left(\partial_1 \partial_2 a(z) - \left(\frac{\partial_1 \zeta(z) \partial_2 \zeta(z)}{\zeta'(z)} \right)' - \left(\frac{\partial_1 \ell(z) \partial_2 \ell(z)}{\ell'(z)} \right)'_{\infty, \geq 0} - \sum_{j=1}^m \left(\frac{\partial_1 \ell(z) \partial_2 \ell(z)}{\ell'(z)} \right)'_{\varphi_j, \leq -1} \right) dz \\ & + \frac{1}{2\pi i} \sum_{s=1}^m \int_{\gamma_s} \hat{\omega}(z) \left(\partial_1 \partial_2 \hat{a}(z) + \left(\frac{\partial_1 \zeta(z) \partial_2 \zeta(z)}{\zeta'(z)} \right)' - \left(\frac{\partial_1 \ell(z) \partial_2 \ell(z)}{\ell'(z)} \right)'_{\infty, \geq 0} - \sum_{j=1}^m \left(\frac{\partial_1 \ell(z) \partial_2 \ell(z)}{\ell'(z)} \right)'_{\varphi_j, \leq -1} \right) dz \\ = & \frac{1}{2\pi i} \sum_{s=1}^m \int_{\gamma_s} \left(\omega(z) \partial_1 \partial_2 a(z) + \hat{\omega}(z) \partial_1 \partial_2 \hat{a}(z) + \omega'(z)_+ \left(\frac{\partial_1 \zeta(z) \partial_2 \zeta(z)}{\zeta'(z)} \right) - \hat{\omega}'(z)_- \left(\frac{\partial_1 \zeta(z) \partial_2 \zeta(z)}{\zeta'(z)} \right) \right) dz \\ & - \text{Res}_\infty \left(\omega'(z)_- \left(\frac{\partial_1 \ell(z) \partial_2 \ell(z)}{a'(z)} \right) + \hat{\omega}'(z)_- \left(\frac{\partial_1 \ell(z) \partial_2 \ell(z)}{a'(z)} \right) \right) dz \\ & + \sum_{j=1}^m \text{Res}_{\varphi_j} \left(\omega'(z)_+ \left(\frac{\partial_1 \ell(z) \partial_2 \ell(z)}{\hat{a}'(z)} \right) + \hat{\omega}'(z)_+ \left(\frac{\partial_1 \ell(z) \partial_2 \ell(z)}{\hat{a}'(z)} \right) \right) dz. \end{aligned}$$

It follows that

$$\begin{aligned}
& \partial_1 \langle \vec{\omega}, \partial_2 \rangle - \langle \vec{\omega}, \nabla_{\partial_1} \partial_2 \rangle \\
&= \frac{1}{2\pi i} \sum_{s=1}^m \int_{\gamma_s} \left(\partial_1 \omega(z) - \omega'(z)_+ \frac{\partial_1 \zeta(z)}{\zeta'(z)} + \hat{\omega}'(z)_- \frac{\partial_1 \zeta(z)}{\zeta'(z)} \right) \partial_2 a(z) dz \\
&+ \frac{1}{2\pi i} \sum_{s=1}^m \int_{\gamma_s} \left(\partial_1 \hat{\omega}(z) + \omega'(z)_+ \frac{\partial_1 \zeta(z)}{\zeta'(z)} - \hat{\omega}'(z)_- \frac{\partial_1 \zeta(z)}{\zeta'(z)} \right) \partial_2 \hat{a}(z) dz \\
&+ \text{Res}_\infty \left((\omega'(z)_- + \hat{\omega}'(z)_-) \frac{\partial_1 \ell(z)}{a'(z)} \right) \partial_2 a(z) dz \\
&- \sum_{j=1}^m \text{Res}_{\varphi_j} \left((\omega'(z)_+ + \hat{\omega}'(z)_+) \frac{\partial_1 \ell(z)}{\hat{a}'(z)} \right) \partial_2 \hat{a}(z) dz \\
&= \frac{1}{2\pi i} \sum_{s=1}^m \int_{\gamma_s} \frac{((\partial_1 \omega_+ - \partial_1 \hat{\omega}_-) \zeta' - (\omega'_+ - \hat{\omega}'_-) \partial_1 \zeta) \partial_2 \zeta}{\zeta'} dz \\
&- \text{Res}_\infty \frac{((\partial_1 \omega_- + \partial_1 \hat{\omega}_-) \ell' - (\omega'_- + \hat{\omega}'_-) \partial_1 \ell) \partial_2 \ell}{\ell'} dz \\
&- \sum_{j=1}^m \text{Res}_{\varphi_j} \frac{((\omega'_- + \hat{\omega}'_-) \partial_1 \ell - (\partial_1 \omega_- + \partial_1 \hat{\omega}_-) \ell') \partial_2 \ell}{\ell'} dz.
\end{aligned}$$

Let $X = \eta \cdot \nabla_{\partial_1} \vec{\omega}$, then we deduce

$$\partial_X \zeta(z) = \partial_1 \zeta(z) (\omega'(z)_+ - \hat{\omega}'(z)_-) - \zeta'(z) (\partial_1 \omega(z)_+ - \partial_1 \hat{\omega}(z)_-)$$

and

$$\begin{aligned}
\partial_X \ell(z) &= ((\partial_1 \omega(z)_- + \partial_1 \hat{\omega}(z)_-) a'(z) - (\omega'(z)_- + \hat{\omega}'(z)_-) \partial_1 a(z))_{\infty, \geq 0} \\
&+ \sum_{j=1}^m ((\omega'(z)_- + \hat{\omega}'(z)_-) \partial_1 \hat{a}(z) - (\partial_1 \omega(z)_- + \partial_1 \hat{\omega}(z)_-) \hat{a}'(z))_{\varphi_j, \leq -1}.
\end{aligned}$$

Taking $\partial_1 = \partial_x$, we obtain

$$\begin{aligned}
\partial_X \zeta(z) &= \{\omega(z)_+ - \hat{\omega}(z)_-, \zeta(z)\}, \\
\partial_X \ell(z) &= -\{\omega(z)_- + \hat{\omega}(z)_-, a(z)\}_+ + \{\omega(z)_+ + \hat{\omega}(z)_+, \hat{a}(z)\}_-.
\end{aligned}$$

Hence

$$\begin{aligned}
\partial_X a(z) &= \{\omega(z), a(z)\}_- + \{\hat{\omega}(z), \hat{a}(z)\}_- - \{\omega(z)_- + \hat{\omega}(z)_-, a(z)\}, \\
\partial_X \hat{a}(z) &= -\{\omega(z), a(z)\}_+ - \{(z) \hat{\omega}(z), \hat{a}(z)\}_+ + \{\omega(z)_+ + \hat{\omega}(z)_+, \hat{a}(z)\}.
\end{aligned}$$

The corollary is proved. \square

2.3. Multiplication and Potential. For any $\vec{a} \in \mathcal{M}$, in order to construct a multiplication structure on the tangent space $T_{\vec{a}}\mathcal{M}$, we first introduce a multiplication structure on $\mathcal{H} \times \mathcal{H}$.

Lemma 2.7. *Given $\vec{a} \in \mathcal{M}$, we define a multiplication on $\mathcal{H} \times \mathcal{H}$ by*

$$\begin{aligned}\vec{\omega}_1 \circ \vec{\omega}_2 &= (\omega_2(\omega_1 a')_+ - \omega_1(\omega_2 a')_- - \omega_2(\hat{\omega}_1 \hat{a}')_- - \omega_1(\hat{\omega}_2 \hat{a}')_-), \\ &\quad \hat{\omega}_2(\hat{\omega}_1 \hat{a}')_+ - \hat{\omega}_1(\hat{\omega}_2 \hat{a}')_- + \hat{\omega}_1(\omega_2 a')_+ + \hat{\omega}_2(\omega_1 a')_+).\end{aligned}$$

Then, $(\mathcal{H} \times \mathcal{H}, \circ)$ forms a commutative and associative algebra.

Proof. The multiplication \circ obviously satisfies the commutative property. To verify the associative property, first consider the case where $\vec{\omega}_\nu = (\omega_\nu, 0)$, for $\nu = 1, 2, 3$. Using the identity

$$\omega_2(\omega_1 a')_+ = \omega_2 \omega_1 a' - \omega_2(\omega_1 a')_-,$$

we have

$$\begin{aligned}&\vec{\omega}_1 \circ \vec{\omega}_2 \circ \vec{\omega}_3 \\ &= (((\omega_1 a')_+ \omega_2 - (\omega_2 a')_- \omega_1) a')_+ \omega_3 - (\omega_3 a')_- ((\omega_1 a')_+ \omega_2 - (\omega_3 a')_- \omega_1), 0) \\ &\sim (((\omega_1 a')_+ \omega_2 a')_+ \omega_3 - ((\omega_2 a')_- \omega_1 a')_+ \omega_3 - (\omega_3 a')_- (\omega_1 a')_+ \omega_2 + (\omega_3 a')_- (\omega_2 a')_- \omega_1, 0) \\ &= (((\omega_1 a')_+ (\omega_2 a')_+)_+ \omega_3 - (\omega_3 a')_- (\omega_1 a')_+ \omega_2, 0) \\ &\sim (((\omega_1 a')_+ (\omega_2 a')_+)_+ \omega_3 + (\omega_3 a')_+ (\omega_1 a')_+ \omega_2, 0) \\ &= (0, 0),\end{aligned}$$

where we denote $f \sim g$ if $f - g$ is symmetric with respect to ω_2, ω_3 . The above result indicates $\vec{\omega}_1 \circ \vec{\omega}_2 \circ \vec{\omega}_3 = \vec{\omega}_1 \circ \vec{\omega}_3 \circ \vec{\omega}_2$.

Next, consider another case:

$$\begin{aligned}&(\omega_1, 0) \circ (\omega_2, 0) \circ (0, \hat{\omega}_3) \\ &= ((-\omega_1 a')_+ \omega_2 + (\omega_2 a')_- \omega_1)(\hat{\omega}_3 \hat{a}')_-, (((\omega_1 a')_+ \omega_2 - (\omega_2 a')_- \omega_1) a')_+ \hat{\omega}_3) \\ &= ((-\omega_1 a')_+ \omega_2 + (\omega_2 a')_- \omega_1)(\hat{\omega}_3 \hat{a}')_-, (\omega_1 a')_+ (\omega_2 a')_+ \hat{\omega}_3) \\ &= ((-\hat{\omega}_3 \hat{a}')_- \omega_1 a')_+ \omega_2 + (\hat{\omega}_3 \hat{a}')_- (\omega_2 a')_- \omega_1 - ((\omega_1 a')_+ \hat{\omega}_3 \hat{a}')_- \omega_2, (\omega_1 a')_+ (\omega_2 a')_+ \hat{\omega}_3) \\ &= (\omega_1, 0) \circ (0, \hat{\omega}_3) \circ (\omega_2, 0).\end{aligned}$$

The remaining cases can be verified similarly. The lemma is proved. \square

Using the pairing (2.8), the above multiplication induces a linear map from $\mathcal{H} \times \mathcal{H}$ to $T_{\vec{a}}\mathcal{M}$.

Corollary 2.8. *For any $\vec{\xi} = (\xi, \hat{\xi}) \in T_{\vec{a}}\mathcal{M}$, define the linear map from $\mathcal{H} \times \mathcal{H}$ to $T_{\vec{a}}\mathcal{M}$ as*

$$C_{\vec{\xi}} \cdot \vec{\omega} = (a'(\omega \xi + \hat{\omega} \hat{\xi})_- - \xi(\omega a' + \hat{\omega} \hat{a}')_-, -\hat{a}'(\omega \xi + \hat{\omega} \hat{\xi})_+ + \hat{\xi}(\omega a' + \hat{\omega} \hat{a}')_+). \quad (2.26)$$

Then, it holds that

$$\langle \vec{\omega}_1 \circ \vec{\omega}_2, \vec{\xi}_3 \rangle = \langle \vec{\omega}_1, C_{\vec{\xi}_3} \cdot \vec{\omega}_2 \rangle, \quad \vec{\omega}_1, \vec{\omega}_2 \in \mathcal{H}.$$

Define the multiplication on $T_{\vec{a}}\mathcal{M}$ as follows:

$$\vec{\xi}_1 \circ \vec{\xi}_2 = \eta \circ (\eta^{-1}(\vec{\xi}_1) \circ \eta^{-1}(\vec{\xi}_2)) = C_{\vec{\xi}_1} \cdot \eta^{-1}(\vec{\xi}_2). \quad (2.27)$$

It can be verified that the above definition is independent of the choice of $\eta^{-1}(\vec{\xi}) \in \mathcal{H} \times \mathcal{H}$. In fact, if $\vec{\omega} \in \mathcal{H} \times \mathcal{H}$ satisfies $\eta(\vec{\omega}) = 0$, then for any $\vec{\omega}^* \in \mathcal{H} \times \mathcal{H}$ and $\vec{\xi}^* \in T_{\vec{a}}\mathcal{M}$, the following holds:

$$\langle \vec{\omega}, C_{\vec{\xi}^*} \cdot \vec{\omega}^* \rangle = \langle C_{\vec{\xi}^*} \cdot \vec{\omega}, \vec{\omega}^* \rangle = 0.$$

Applying Lemma 2.4, we deduce that $C_{\vec{\xi}^*} \cdot \vec{\omega} = 0$. Hence, the multiplication \circ on $T_{\vec{a}}\mathcal{M}$ is well-defined.

Corollary 2.9. *The unit element of the algebra $(T_{\vec{a}}\mathcal{M}, \circ)$ is given by*

$$e = \begin{cases} \sum_{i=1}^m (\frac{\partial}{\partial t_{i,0}} + \frac{\partial}{\partial h_{i,0}}), & \text{if } n_0 = 1, \\ \frac{1}{n_0} \frac{\partial}{\partial h_{0,n_0-1}}, & \text{if } n_0 \geq 2. \end{cases} \quad (2.28)$$

Proof. From the equalities (2.5) to (2.7) and (2.28), we deduce that

$$\partial_e \vec{a} = \begin{cases} (1 - a'(z), 1 - \hat{a}'(z)), & n_0 = 1, \\ (1, 1), & n_0 \geq 2. \end{cases} \quad (2.29)$$

It is straightforward to verify that

$$C_e \cdot \vec{\omega} = \eta(\vec{\omega}), \quad \vec{\omega} \in \mathcal{H} \times \mathcal{H},$$

which directly leads to

$$e \circ \vec{\xi} = C_e \cdot \eta^{-1}(\vec{\xi}) = \vec{\xi}, \quad \vec{\xi} \in T_{\vec{a}}\mathcal{M}.$$

The corollary is proved. □

Next, we will present the components of the 3-tensor

$$c(\partial', \partial'', \partial''') = \langle \partial' \circ \partial'', \partial''' \rangle_\eta = \langle \eta^{-1}(\partial') \circ \eta^{-1}(\partial''), \partial''' \rangle$$

in the flat coordinates. The proofs of the subsequent conclusions in this subsection closely follow the approach presented in [13] for the corresponding results. To avoid redundancy, we omit the detailed proof process here.

Corollary 2.10. *We have the following expressions for the basis vectors in the flat coordinate system under the map η^{-1} :*

$$\eta^{-1} \frac{\partial}{\partial t_{i,s}} = (\zeta(z)^{\frac{s}{d_i}} \mathbf{1}_{\gamma_i}, -\zeta(z)^{\frac{s}{d_i}} \mathbf{1}_{\gamma_i}), \quad s \in \mathbb{Z}; \quad (2.30)$$

$$\eta^{-1} \frac{\partial}{\partial h_{0,j}} = ((\chi_0(z)^{-j})_{\infty, \geq -n_0+1}, 0), \quad 1 \leq j \leq n_0 - 1; \quad (2.31)$$

$$\eta^{-1} \frac{\partial}{\partial h_{k,r}} = (0, (\chi_k(z)^{-r})_{\varphi_k, \leq n_k} \mathbf{1}_{\gamma_k}), \quad 0 \leq r \leq n_k, \quad (2.32)$$

where η^{-1} is defined by Lemma 2.4. Furthermore, we have

$$\begin{aligned} \eta^{-1} \frac{\partial}{\partial t_{i_1, s_1}} \circ \eta^{-1} \frac{\partial}{\partial t_{i_2, s_2}} &= \left(\delta_{i_1, i_2} \zeta^{\frac{s_1+s_2}{d_{i_1}}} a' \mathbf{1}_{\gamma_{i_1}} - \zeta^{\frac{s_1}{d_{i_1}}} (\zeta^{\frac{s_2}{d_{i_2}}} \zeta' \mathbf{1}_{\gamma_{i_2}})_{-} \mathbf{1}_{\gamma_{i_1}} - \zeta^{\frac{s_2}{d_{i_2}}} (\zeta^{\frac{s_1}{d_{i_1}}} \zeta' \mathbf{1}_{\gamma_{i_1}})_{-} \mathbf{1}_{\gamma_{i_2}}, \right. \\ &\quad \left. - \delta_{i_1, i_2} \zeta^{\frac{s_1+s_2}{d_{i_1}}} \hat{a}' \mathbf{1}_{\gamma_{i_1}} - \zeta^{\frac{s_1}{d_{i_1}}} (\zeta^{\frac{s_2}{d_{i_2}}} \zeta' \mathbf{1}_{\gamma_{i_2}})_{+} \mathbf{1}_{\gamma_{i_1}} - \zeta^{\frac{s_2}{d_{i_2}}} (\zeta^{\frac{s_1}{d_{i_1}}} \zeta' \mathbf{1}_{\gamma_{i_1}})_{+} \mathbf{1}_{\gamma_{i_2}} \right), \\ \eta^{-1} \frac{\partial}{\partial h_{0, j_1}} \circ \eta^{-1} \frac{\partial}{\partial h_{0, j_2}} &= (-\chi_0^{-j_1})_{\infty, \geq -n_0+1} (\chi_0^{-j_2})_{\infty, \geq -n_0+1} a' + (\chi_0^{-j_1})_{\infty, \geq -n_0+1} (\chi_0^{-j_2} \ell')_{\infty, \geq 0} \\ &\quad + (\chi_0^{-j_2})_{\infty, \geq -n_0+1} (\chi_0^{-j_1} \ell')_{\infty, \geq 0}, 0), \\ \eta^{-1} \frac{\partial}{\partial h_{k_1, r_1}} \circ \eta^{-1} \frac{\partial}{\partial h_{k_2, r_2}} &= (0, \delta_{k_1, k_2} (\chi_{k_1}^{-r_1})_{\varphi_{k_1}, \leq n_{k_1}} (\chi_{k_2}^{-r_2})_{\varphi_{k_2}, \leq n_{k_2}} \hat{a}' \mathbf{1}_{\gamma_{k_1}} \\ &\quad - (\chi_{k_1}^{-r_1})_{\varphi_{k_1}, \leq n_{k_1}} (\chi_{k_2}^{-r_2} \ell')_{\varphi_{k_2}, \leq -1} \mathbf{1}_{\gamma_{k_1}} - (\chi_{k_2}^{-r_2})_{\varphi_{k_2}, \leq n_{k_2}} (\chi_{k_1}^{-r_1} \ell')_{\varphi_{k_1}, \leq -1} \mathbf{1}_{\gamma_{k_2}}), \\ \eta^{-1} \frac{\partial}{\partial h_{0, j}} \circ \eta^{-1} \frac{\partial}{\partial h_{k, r}} &= (-\chi_0^{-j})_{\infty, \geq -n_0+1} (\chi_k^{-r} \ell')_{\varphi_k, \leq -1}, (\chi_k^{-r})_{\varphi_k, \leq n_k} (\chi_0^{-j} \ell')_{\infty, \geq 0} \mathbf{1}_{\gamma_k}). \end{aligned}$$

Corollary 2.11. *The components of the symmetric 3-tensor c in the flat coordinates are*

$$\begin{aligned} c(\partial_{t_{i_1, s_1}}, \partial_{t_{i_2, s_2}}, \partial_{t_{i_3, s_3}}) &= \frac{1}{2\pi i} \int_{\gamma_{i_1}} \left(-\delta_{i_1, i_2} \delta_{i_2, i_3} \zeta^{\frac{s_1+s_2+s_3}{d_{i_1}}} \zeta' \mathbf{1}_{\gamma_{i_1}} (\zeta' + \zeta'_- + \ell') dz \right) \\ &\quad + \frac{1}{2\pi i} \sum_{s=1}^m \int_{\gamma_s} \left(\delta_{i_1, i_2} \zeta^{\frac{s_1+s_2}{d_{i_1}}} \zeta' \mathbf{1}_{\gamma_{i_1}} (\zeta^{\frac{s_3}{d_{i_3}}} \zeta' \mathbf{1}_{i_3})_{-} + \text{c.p.}\{(i_1, s_1), (i_2, s_2), (i_3, s_3)\} \right) dz, \\ c(\partial_{h_{0, j_1}}, \partial_{h_{0, j_2}}, \partial_{h_{0, j_3}}) &= -\text{Res}_{z=\infty} \left(-\chi_0^{-j_1-j_2-j_3} \ell' (2\ell' + \zeta'_-) \right) dz \\ &\quad - \text{Res}_{z=\infty} \left(\chi_0^{-j_1-j_2} \ell' (\ell' \chi_0^{-j_3})_{\infty, \geq 0} + \text{c.p.}\{j_1, j_2, j_3\} \right) dz, \\ c(\partial_{h_{k_1, r_1}}, \partial_{h_{k_2, r_2}}, \partial_{h_{k_3, r_3}}) &= \text{Res}_{z=\varphi_{k_1}} \left(-\delta_{k_1, k_2} \delta_{k_2, k_3} \chi_{k_1}^{-r_1-r_2-r_3} \ell' (2\ell' - \zeta'_+) \right) dz \\ &\quad + \sum_{s=1}^m \text{Res}_{\varphi_s} \left(\delta_{k_1, k_2} \chi_{k_1}^{-r_1-r_2} \ell' (\ell' \chi_{k_3}^{-r_3})_{\varphi_{k_3}, \leq -1} + \text{c.p.}\{(k_1, r_1), (k_2, r_2), (k_3, r_3)\} \right) dz, \\ c(\partial_{t_{i_1, s_1}}, \partial_{t_{i_2, s_2}}, \partial_{h_{0, j_3}}) &= -\frac{1}{2\pi i} \int_{\gamma_{i_1}} \delta_{i_1, i_2} \zeta^{\frac{s_1+s_2}{d_{i_1}}} \zeta' (\ell' \chi_0^{-j_3})_{\infty, \geq 0} dz, \end{aligned}$$

$$\begin{aligned}
c(\partial_{t_{i_1, s_1}}, \partial_{t_{i_2, s_2}}, \partial_{h_{k_3, r_3}}) &= \frac{1}{2\pi i} \int_{\gamma_{i_1}} \delta_{i_1, i_2} \zeta^{\frac{s_1+s_2}{d_{i_1}}} \zeta' (\ell' \chi_{k_3}^{-r_3})_{\varphi_{k_3, \leq -1}} dz, \\
c(\partial_{h_{0, j_1}}, \partial_{h_{0, j_2}}, \partial_{t_{i_3, s_3}}) &= \operatorname{Res}_{z=\infty} \chi_0^{-j_1-j_2} \ell' (\zeta^{\frac{s_3}{d_{i_3}}} \zeta' \mathbf{1}_{\gamma_{i_3}})_- dz, \\
c(\partial_{h_{0, j_1}}, \partial_{h_{0, j_2}}, \partial_{h_{k_3, r_3}}) &= \operatorname{Res}_{z=\infty} \chi_0^{-j_1-j_2} \ell' (\ell' \chi_{k_3}^{-r_3})_{\varphi_{k_3, \leq -1}} dz, \\
c(\partial_{h_{k_1, r_1}}, \partial_{h_{k_2, r_2}}, \partial_{t_{i_3, s_3}}) &= - \operatorname{Res}_{z=\varphi_{k_1}} \delta_{k_1, k_2} \chi_{k_1}^{-r_1-r_2} \ell' (\zeta^{\frac{s_3}{d_{i_3}}} \zeta' \mathbf{1}_{\gamma_{i_3}})_+ dz, \\
c(\partial_{h_{k_1, r_1}}, \partial_{h_{k_2, r_2}}, \partial_{h_{0, j_3}}) &= - \operatorname{Res}_{z=\varphi_{k_1}} \delta_{k_1, k_2} \chi_{k_1}^{-r_1-r_2} \ell' (\ell' \chi_0^{-j_3})_{\infty, \geq 0} dz, \\
c(\partial_{t_{i_1, s_1}}, \partial_{h_{0, j_2}}, \partial_{h_{k_3, r_3}}) &= 0.
\end{aligned}$$

To construct a function \mathcal{F} on the manifold \mathcal{M} satisfying

$$\frac{\partial \mathcal{F}}{\partial u \partial v \partial w} = c \left(\frac{\partial}{\partial u}, \frac{\partial}{\partial v}, \frac{\partial}{\partial w} \right), \quad u, v, w \in \mathbf{t} \cup \mathbf{h}, \quad (2.33)$$

we introduce auxiliary functions $V_{1,i}$, $V_{2,j}$, and G on \mathcal{M} . These functions are defined in terms of the coordinates \mathbf{t}_i and \mathbf{h}_0 as follows:

$$\mathbf{t}_i = \{t_{i,s} \mid s \in \mathbb{Z}\}, \quad \mathbf{h}_0 = \{h_{0,j} \mid j = 1, \dots, m\}.$$

Lemma 2.12. *There exist functions $V_{1,i}(\mathbf{t}_i)$, $V_{2,j}(\mathbf{h}_0)$, and $G(\mathbf{h})$ on \mathcal{M} that satisfy*

$$\frac{\partial^2 V_{1,i}}{\partial t_{i,s_1} \partial t_{i,s_2}} = \frac{1}{2\pi i} \int_{\gamma_s} \frac{\zeta'(z) \zeta(z)^{\frac{s_1+s_2}{d_i}}}{z - p_i} dz, \quad i = 1, \dots, m, \quad (2.34)$$

$$\frac{\partial^2 V_{2,j}}{\partial h_{0,j_1} \partial h_{0,j_2}} = - \operatorname{Res}_{z=\infty} \frac{\ell'(z) \chi_0(z)^{-j_1-j_2}}{z - p_j} dz, \quad j = 1, \dots, m, \quad (2.35)$$

$$\frac{\partial^3 G}{\partial u \partial v \partial w} = - \left(\operatorname{Res}_{z=\infty} + \sum_{s=1}^m \operatorname{Res}_{z=\varphi_s} \right) \frac{\partial_u \ell(z) \cdot \partial_v \ell(z) \cdot \partial_w \ell(z)}{\ell'(z)} dz, \quad u, v, w \in \mathbf{h}, \quad (2.36)$$

where p_s is the center of the disk D_s for $s = 1, \dots, m$.

Theorem 2.13. *The function \mathcal{F} on \mathcal{M} , defined by*

$$\begin{aligned}
\mathcal{F} &= \frac{1}{(2\pi i)^2} \sum_{s=1}^m \int_{\gamma_s} \int_{\gamma_s^*} \left(\frac{1}{2} \zeta(z_1) \zeta(z_2) + \zeta(z_1) \ell(z_2) - \ell(z_1) \zeta(z_2) \right) \log \left(\frac{z_1 - z_2}{z_1 - p_s} \right) dz_1 dz_2 \\
&\quad + \frac{1}{(2\pi i)^2} \sum_{s=1}^m \sum_{s' \neq s} \int_{\gamma_s} \int_{\gamma_{s'}} \left(\frac{1}{2} \zeta(z_1) \zeta(z_2) + \zeta(z_1) \ell(z_2) - \ell(z_1) \zeta(z_2) \right) \log(z_1 - z_2) dz_1 dz_2 \\
&\quad - \sum_{s=1}^m \frac{V_{1,s}(\mathbf{t})}{2\pi i} \int_{\gamma_s} \left(\frac{1}{2} \zeta(z) + \ell(z) \right) dz + \sum_{s=1}^m \frac{V_{2,s}(\mathbf{h})}{2\pi i} \int_{\gamma_s} \zeta(z) dz + G(\mathbf{h}),
\end{aligned}$$

satisfies the condition (2.33), where the integration contours γ_s^* are small deformations of γ_s for each $s = 1, \dots, m$, such that the inequality $|z_2 - p_s| < |z_1 - p_s|$ holds for all $z_1 \in \gamma_s$ and $z_2 \in \gamma_s^*$.

Define the Euler vector field E on \mathcal{M} as

$$\partial_E \vec{a} = \left(a(z) - \frac{z}{n_0} a'(z), \hat{a}(z) - \frac{z}{n_0} \hat{a}'(z) \right). \quad (2.37)$$

The formula for this vector field in flat coordinates is given by

$$E = \sum_{i=1}^m \sum_{s \in \mathbb{Z}} \left(\frac{1}{n_0} - \frac{s}{d_i} \right) t_{i,s} \frac{\partial}{\partial t_{i,s}} + \sum_{j=1}^{n_0} \left(\frac{1+j}{n_0} \right) h_{0,j} \frac{\partial}{\partial h_{0,j}} + \sum_{k=1}^m \sum_{r=0}^{n_k} \left(\frac{1}{n_0} + \frac{r}{n_k} \right) h_{k,r} \frac{\partial}{\partial h_{k,r}}.$$

Corollary 2.14. *The function \mathcal{F} defined in Theorem 2.13 satisfies*

$$\text{Lie}_E \mathcal{F} = \left(2 + \frac{2}{n_0} \right) \mathcal{F} + \text{terms quadratic in } \mathbf{t} \cup \mathbf{h}.$$

Combining the results in this section, we have proved the Theorem 1.1.

3. PRINCIPAL HIERARCHY OF \mathcal{M}

In this section, we will give the proof of Theorems 1.2 and 1.3. To substantiate the proof of the Theorem 1.2, the following lemma is indispensable.

Lemma 3.1. *Let $Q_p(a, \hat{a}), p \in \mathbb{N}$, be analytic functions in a and \hat{a} that satisfy*

$$\frac{\partial Q_p(a, \hat{a})}{\partial a} + \frac{\partial Q_p(a, \hat{a})}{\partial \hat{a}} = Q_{p-1}(a, \hat{a}).$$

Define

$$F_{i,p} = \frac{1}{2\pi i} \int_{\gamma_i} Q_{p+1}(a(z), \hat{a}(z)) dz, \quad (3.1)$$

then we have

$$\eta \cdot \nabla_{\partial} dF_{i,p} = C_{\partial}(dF_{i,p-1}), \quad (3.2)$$

where ∂ is any vector field on \mathcal{M} , and the operators $\eta \cdot \nabla_{\partial}$ and C_{∂} are explicitly defined by equalities (2.24) and (2.26), respectively.

Proof. The differential of $F_{i,p}$ at $\vec{a} \in \mathcal{M}$ is given by

$$dF_{i,p}|_{\vec{a}} = \left(\frac{\partial Q_{p+1}}{\partial a} \mathbf{1}_{\gamma_i}, \frac{\partial Q_{p+1}}{\partial \hat{a}} \mathbf{1}_{\gamma_i} \right)|_{\vec{a}} \in \mathcal{H} \times \mathcal{H}.$$

Using the identity

$$\left\{ \frac{\partial Q_p}{\partial a}, a \right\} + \left\{ \frac{\partial Q_p}{\partial \hat{a}}, \hat{a} \right\} = 0,$$

where $\{f, g\} = f'g - g'f$, we obtain

$$\eta \cdot \nabla_{\partial} dF_{i,p} = \left(-\{(Q_p \mathbf{1}_{\gamma_i})_-, a\}, \{(Q_p \mathbf{1}_{\gamma_i})_+, \hat{a}\} \right).$$

For the right-hand side of equality (3.2), we have

$$\begin{aligned}
C_{\partial}(dF_{i,p-1}) &= (a'(\frac{\partial Q_p}{\partial a}\partial a\mathbf{1}_{\gamma_i} + \frac{\partial Q_p}{\partial \hat{a}}\partial \hat{a}\mathbf{1}_{\gamma_i})_- - \partial a(\frac{\partial Q_p}{\partial a}a'\mathbf{1}_{\gamma_i} + \frac{\partial Q_p}{\partial \hat{a}}\hat{a}'\mathbf{1}_{\gamma_i})_- \\
&\quad - \hat{a}'(\frac{\partial Q_p}{\partial a}\partial a\mathbf{1}_{\gamma_i} + \frac{\partial Q_p}{\partial \hat{a}}\partial \hat{a}\mathbf{1}_{\gamma_i})_+ + \partial \hat{a}(\frac{\partial Q_p}{\partial a}a'\mathbf{1}_{\gamma_i} + \frac{\partial Q_p}{\partial \hat{a}}\hat{a}'\mathbf{1}_{\gamma_i})_+) \\
&= (-\{(Q_p\mathbf{1}_{\gamma_i})_-, a\}, \{(Q_p\mathbf{1}_{\gamma_i})_+, \hat{a}\}) \\
&= \eta \cdot \nabla_{\partial} dF_{i,p}.
\end{aligned}$$

Thus, the lemma is proved. \square

Proof of Theorem 1.2. The Hamiltonian density $\theta_{t_{i,s},p}$, for $s \neq -d_i$, takes the form of (3.1), and consequently satisfies equation (3.2), which is equivalent to equation (1.1). For $\theta_{t_{i,-d_i},p}$, consider a subset \mathcal{M}' of \mathcal{M} where $a^{\frac{1}{n_0}}$ can be analytically continued to $\cup_{s=1}^m \gamma_s$, with winding number 1 around γ_i and winding number 0 around γ_s for $s \neq i$. Consequently, $a^{\frac{1}{n_0}} \in \mathcal{H}$, and on \mathcal{M}' we have the following expression for $\theta_{t_{i,-d_i},p}$:

$$\begin{aligned}
\theta_{t_{i,-d_i},p} &= -d_i \operatorname{Res}_{\infty} \frac{c_p}{n_0} \frac{a^p}{p!} dz - \frac{d_i}{2\pi i} \int_{\gamma_i} \frac{a^p}{p!} \log a^{\frac{1}{n_0}} / \zeta^{\frac{1}{d_i}} dz - \frac{d_i}{2\pi i} \sum_{s \neq i} \int_{\gamma_s} \frac{a^p}{p!} \log a^{\frac{1}{n_0}} dz \\
&= -\frac{d_i}{2\pi i} \int_{\gamma_i} \frac{a^p}{p!} (\log a^{\frac{1}{n_0}} / \zeta^{\frac{1}{d_i}} - \frac{c_p}{n_0}) dz - \frac{d_i}{2\pi i} \sum_{s \neq i} \int_{\gamma_s} \frac{a^p}{p!} (\log a^{\frac{1}{n_0}} - \frac{c_p}{n_0}) dz,
\end{aligned}$$

which is a linear combination of functions of the form (3.1), and hence satisfies equation (1.1). Due to the uniqueness of analytic functions, this equation holds on \mathcal{M} .

Using similar methods, the remaining Hamiltonian densities can be shown to satisfy equation (1.1). Specifically, for $\theta_{h_k, n_k, p}$, consider a subset \mathcal{M}' of \mathcal{M} where $a^{\frac{1}{n_0}}$ and $\hat{a}^{\frac{1}{n_k}}$ can be analytically continued to $\cup_{s=1}^m \gamma_s$ and γ_i , respectively, with the appropriate winding numbers. Then, on the subset \mathcal{M}' of \mathcal{M} , the function $\theta_{h_k, n_k, p}$ has the form

$$\theta_{h_k, n_k, p} = \frac{n_k}{2\pi i} \int_{\gamma_k} \frac{\hat{a}^p}{p!} (\log \zeta^{\frac{1}{d_k}} \hat{a}^{\frac{1}{n_k}} - \frac{c_p}{n_k}) dz - \frac{n_k}{d_k} \theta_{t_k, -d_k, p},$$

and hence satisfies equation (1.1). Due to the uniqueness of analytic functions, this property extends to the entire space \mathcal{M} .

We proceed to verify equation (1.2) by introducing the operator $\mathcal{E} = E + \frac{1}{m} z \frac{\partial}{\partial z}$. This yields the following set of relationships:

$$\operatorname{Lie}_{\mathcal{E}} \zeta(z) = \zeta(z), \quad \operatorname{Lie}_{\mathcal{E}} a(z) = a(z), \quad \operatorname{Lie}_{\mathcal{E}} \hat{a}(z) = \hat{a}(z), \quad \operatorname{Lie}_{\mathcal{E}} \phi_p(z) = p\phi_p(z).$$

For any smooth function $f(a, \hat{a})$, the following identity holds:

$$\int_{\gamma_s} \operatorname{Lie}_{\mathcal{E}} f(a, \hat{a}) dz = \operatorname{Lie}_E \int_{\gamma_s} f(a, \hat{a}) dz - \frac{1}{n_0} \int_{\gamma_s} f(a, \hat{a}) dz, \quad s = 1, \dots, m.$$

Employing the aforementioned identity, we can deduce the Lie derivatives of the Hamiltonian densities along E as follows:

$$\begin{aligned} \text{Lie}_E \theta_{t_i, s, p} &= \left(p + 1 + \frac{s}{d_i} + \frac{1}{n_0}\right) \theta_{t_i, s, p}, \quad s \neq -d_i \\ \text{Lie}_E \theta_{t_i, -d_i, p} &= \left(p + \frac{1}{n_0}\right) \theta_{t_i, -d_i, p} + \left(1 - \frac{d_i}{n_0}\right) (\theta_{t_i, 0, p-1} + \theta_{h_{i, 0, p-1}}) - \sum_{i' \neq i} \frac{d_i}{n_0} (\theta_{t_{i'}, 0, p-1} + \theta_{h_{i'}, 0, p-1}), \\ \text{Lie}_E \theta_{h_{0, j, p}} &= \left(p + 1 - \frac{j}{n_0} + \frac{1}{n_0}\right) \theta_{h_{0, j, p}}, \\ \text{Lie}_E \theta_{h_{k, r, p}} &= \left(p + 1 - \frac{r}{n_k} + \frac{1}{n_0}\right) \theta_{h_{k, r, p}}, \quad r \neq n_k, \\ \text{Lie}_E \theta_{h_{k, n_k, p}} &= \left(p + \frac{1}{n_0}\right) \theta_{h_{k, n_k, p}} + \left(\frac{n_k}{n_0} + 1\right) \theta_{h_{k, 0, p-1}} + \left(\frac{n_k}{n_0} - \frac{n_k}{d_k}\right) \theta_{t_{k, 0, p-1}} + \sum_{k' \neq k} \frac{n_k}{n_0} (\theta_{t_{k'}, 0, p-1} + \theta_{h_{k'}, 0, p-1}). \end{aligned}$$

These expressions confirm equation (1.2). The theorem is proved. \square

We now demonstrate the relationship between the principal hierarchy

$$\frac{\partial}{\partial T^{\alpha, p-1}} = \mathcal{P}(d\theta_{\alpha, p+1}),$$

and the Whitham hierarchy given by (1.5)-(1.7). Suppose there exists $(a(z), \hat{a}(z)) \in \mathcal{M}$ such that

$$a(z) = \lambda_0(z)^{n_0}, \quad z \rightarrow \infty, \quad \hat{a}(z) = \lambda_j(z)^{n_j}, \quad z \rightarrow \varphi_j, \quad j = 1, \dots, m,$$

then the Whitham hierarchy is equivalent to the following system:

$$\begin{aligned} \frac{\partial a(z)}{\partial S^{0, p}} &= \{(a(z)^{\frac{p}{n_0}})_{\infty, \geq 0}, a(z)\}, & \frac{\partial \hat{a}(z)}{\partial S^{0, p}} &= \{(a(z)^{\frac{p}{n_0}})_{\infty, \geq 0}, \hat{a}(z)\}, \\ \frac{\partial a(z)}{\partial S^{j, p}} &= \{-(\hat{a}(z)^{\frac{p}{n_j}})_{\varphi_j, \leq -1}, a(z)\}, & \frac{\partial \hat{a}(z)}{\partial S^{j, p}} &= \{-(\hat{a}(z)^{\frac{p}{n_j}})_{\varphi_j, \leq -1}, \hat{a}(z)\}, \\ \frac{\partial a(z)}{\partial S^{j, 0}} &= \{\log(z - \varphi_j), a(z)\}, & \frac{\partial \hat{a}(z)}{\partial S^{j, 0}} &= \{\log(z - \varphi_j), \hat{a}(z)\}. \end{aligned}$$

Proof of Theorem 1.3. Let's first consider the system of equations on the loop space $L\mathcal{M}$ corresponding to the Hamiltonian density $\theta_{h_{0, j, p+1}}$. Since

$$d\theta_{h_{0, j, p+1}}|_{\vec{a}} = (Q_{h_{0, j, p}}, 0)|_{\vec{a}} \in \mathcal{H} \times \mathcal{H},$$

where

$$Q_{h_{0, j, p}} = \frac{\Gamma\left(1 - \frac{j}{n_0}\right)}{\Gamma\left(2 + p - \frac{j}{n_0}\right)} a^{1+p-\frac{j}{n_0}},$$

it follows that

$$\begin{aligned} \mathcal{P}(d\theta_{h_{0, j, p+1}}) &= (\{(Q_{h_{0, j, p}})_+, a\}, \{(Q_{h_{0, j, p}})_+, \hat{a}\}) \\ &= (\{(Q_{h_{0, j, p}})_{\infty, \geq 0}, a\}, \{(Q_{h_{0, j, p}})_{\infty, \geq 0}, \hat{a}\}). \end{aligned}$$

Similarly, let's consider the Hamiltonian density $\theta_{h_k, r, p+1}$, where $r \neq n_k$. Since

$$d\theta_{h_k, r, p+1}|_{\bar{a}} = (0, Q_{h_k, r, p} \mathbf{1}_{\gamma_k})|_{\bar{a}} \in \mathcal{H} \times \mathcal{H},$$

where

$$Q_{h_k, r, p} = \frac{\Gamma\left(1 - \frac{r}{n_k}\right)}{\Gamma\left(2 + p - \frac{r}{n_k}\right)} \hat{a}^{1+p-\frac{r}{n_k}},$$

we can deduce that

$$\begin{aligned} \mathcal{P}(d\theta_{h_k, r, p+1}) &= (\{- (Q_{h_k, r, p} \mathbf{1}_{\gamma_k})_-, a\}, \{- (Q_{h_k, r, p} \mathbf{1}_{\gamma_k})_-, \hat{a}\}) \\ &= (\{- (Q_{h_k, r, p})_{\varphi_k, \leq -1}, a\}, \{- (Q_{h_k, r, p})_{\varphi_k, \leq -1}, \hat{a}\}). \end{aligned}$$

Next, we consider the Hamiltonian density $\theta_{h_k, n_k, p+1}$ that includes logarithmic terms.

Direct computation yields

$$\begin{aligned} \mathcal{P}(d\theta_{h_k, n_k, p+1}) &= (\{-n_k(\log \hat{a}^{\frac{1}{n_k}} a^{\frac{1}{n_0}} \mathbf{1}_{\gamma_k} + \sum_{k' \neq k} \log a^{\frac{1}{n_0}} \mathbf{1}_{\gamma_{k'}})_-, a\}, \\ &\quad \{n_k(\log \hat{a}^{\frac{1}{n_k}} a^{\frac{1}{n_0}} \mathbf{1}_{\gamma_k} + \sum_{k' \neq k} \log a^{\frac{1}{n_0}} \mathbf{1}_{\gamma_{k'}})_+, \hat{a}\}) \\ &= (\{-n_k(\log(\hat{a}^{\frac{1}{n_k}}(z - \varphi_k) + \frac{a^{\frac{1}{n_0}}}{(z - \varphi_k)}) \mathbf{1}_{\gamma_k} + \sum_{k' \neq k} (\log \frac{a^{\frac{1}{n_0}}}{(z - \varphi_k)} + \log(z - \varphi_k)) \mathbf{1}_{\gamma_{k'}})_-, a\}, \\ &\quad \{n_k(\log(\hat{a}^{\frac{1}{n_k}}(z - \varphi_k) + \frac{a^{\frac{1}{n_0}}}{(z - \varphi_k)}) \mathbf{1}_{\gamma_k} + \sum_{k' \neq k} (\log \frac{a^{\frac{1}{n_0}}}{(z - \varphi_k)} + \log(z - \varphi_k)) \mathbf{1}_{\gamma_{k'}})_+, \hat{a}\}) \\ &= (\{-n_k \log \frac{a^{\frac{1}{n_0}}}{(z - \varphi_k)}, a\}, \{n_k(\log \hat{a}^{\frac{1}{n_k}}(z - \varphi_k) \mathbf{1}_{\gamma_k} + \sum_{k' \neq k} \log(z - \varphi_k) \mathbf{1}_{\gamma_{k'}}), \hat{a}\}) \\ &= (\{n_k \log(z - \varphi_k), a\}, \{n_k \log(z - \varphi_k), \hat{a}\}). \end{aligned}$$

In summary, the following relationships are derived:

$$\begin{aligned} \frac{\partial}{\partial S^{0, (p+1)n_0-j}} &= \frac{\Gamma(2 + p - \frac{j}{n_0})}{\Gamma(1 - \frac{j}{n_0})} \frac{\partial}{\partial T^{h_{0,j}, p}}, \quad j = 1, \dots, n_0 - 1; \\ \frac{\partial}{\partial S^{i, (p+1)n_i-j}} &= \frac{\Gamma(2 + p - \frac{j}{n_i})}{\Gamma(1 - \frac{j}{n_i})} \frac{\partial}{\partial T^{h_{i,j}, p}}, \quad i = 1, \dots, m, j = 0, \dots, n_i - 1; \\ \frac{\partial}{\partial S^{0, (p+1)n_0}} &= (p+1)! \sum_{k=1}^m \left(\frac{\partial}{\partial T^{t_{k,0}, p}} + \frac{\partial}{\partial T^{h_{k,0}, p}} \right), \\ \frac{\partial}{\partial S^{i, 0}} &= \frac{1}{n_j} \frac{\partial}{\partial T^{h_{i,0}, 0}}. \end{aligned}$$

The theorem is proved. \square

REFERENCES

- [1] B. Dubrovin. Geometry of 2d topological field theories. In *Integrable Systems and Quantum Groups*, pages 120–348. Springer, 1998.
- [2] I. I. Manin. *Frobenius manifolds, quantum cohomology, and moduli spaces*, volume 47. American Mathematical Soc., 1999.
- [3] C. Hertling. *Frobenius manifolds and moduli spaces for singularities*, volume 151. Cambridge University Press, 2002.
- [4] B. Dubrovin and Y. Zhang. Normal forms of hierarchies of integrable PDEs, Frobenius manifolds and Gromov-Witten invariants. *arXiv preprint math/0108160*, 2001.
- [5] A. Givental and E. Milanov. Simple singularities and integrable hierarchies. In *The Breadth of Symplectic and Poisson Geometry: Festschrift in Honor of Alan Weinstein*, pages 173–201. Springer, 2005.
- [6] S.-Q. Liu, Y. Ruan, and Y. Zhang. BCFG Drinfeld–Sokolov hierarchies and FJRW-theory. *Inventiones mathematicae*, 201:711–772, 2015.
- [7] B. Dubrovin, S.-Q. Liu, D. Yang, and Y. Zhang. Hodge integrals and tau-symmetric integrable hierarchies of Hamiltonian evolutionary PDEs. *Advances in Mathematics*, 293:382–435, 2016.
- [8] S.-Q. Liu, Z. Wang, and Y. Zhang. Variational bihamiltonian cohomologies and integrable hierarchies II: Virasoro symmetries. *Communications in Mathematical Physics*, 395(1):459–519, 2022.
- [9] S.-Q. Liu, Z. Wang, and Y. Zhang. Variational bihamiltonian cohomologies and integrable hierarchies I: foundations. *Communications in Mathematical Physics*, 401(1):985–1031, 2023.
- [10] G. Carlet, B. Dubrovin, and L. P. Mertens. Infinite-dimensional Frobenius manifolds for $2+1$ integrable systems. *Mathematische Annalen*, 349:75–115, 2011.
- [11] C.-Z. Wu and D. Xu. A class of infinite-dimensional Frobenius manifolds and their submanifolds. *International Mathematics Research Notices*, 2012(19):4520–4562, 2012.
- [12] C.-Z. Wu and D. Zuo. Infinite-dimensional Frobenius manifolds underlying the Toda lattice hierarchy. *Advances in Mathematics*, 255:487–524, 2014.
- [13] S. Ma, C.-Z. Wu, and D. Zuo. Infinite-dimensional Frobenius manifolds underlying an extension of the dispersionless Kadomtsev–Petviashvili hierarchy. *Journal of Geometry and Physics*, 161:104006, 2021.
- [14] A. Raimondo. Frobenius manifold for the dispersionless Kadomtsev–Petviashvili equation. *Communications in Mathematical Physics*, 311:557–594, 2012.
- [15] I. M. Krichever. The τ -function of the universal Whitham hierarchy, matrix models and topological field theories. *Communications on Pure and Applied Mathematics*, 47(4):437–475, 1994.
- [16] I. M. Krichever. Method of averaging for two-dimensional” integrable” equations. *Functional analysis and its applications*, 22(3):200–213, 1988.
- [17] A. Gorsky, I. M. Krichever, A. Marshakov, A. Mironov, and A. Morozov. Integrability and Seiberg-Witten exact solution. *Physics Letters B*, 355(3-4):466–474, 1995.
- [18] H. Itoyama and A. Morozov. Integrability and Seiberg-Witten theory curves and periods. *Nuclear Physics B*, 477(3):855–877, 1996.
- [19] H. Itoyama and A. Morozov. Prepotential and the Seiberg-Witten theory. *Nuclear Physics B*, 491(3):529–573, 1997.
- [20] I. M. Krichever and D. H. Phong. On the integrable geometry of soliton equations and $n = 2$ supersymmetric gauge theories. *Journal of Differential Geometry*, 45(2):349–389, 1997.

- [21] A. Gorsky, A. Marshakov, A. Mironov, and A. Morozov. RG equations from Whitham hierarchy. *Nuclear Physics B*, 527(3):690–716, 1998.
- [22] M. Mineev-Weinstein and A. Zabrodin. Whitham–Toda hierarchy in the Laplacian growth problem. *Journal of Nonlinear Mathematical Physics*, 8(Suppl 1):212–218, 2001.
- [23] I. M. Krichever, M. Mineev-Weinstein, P. Wiegmann, and A. Zabrodin. Laplacian growth and Whitham equations of soliton theory. *Physica D: Nonlinear Phenomena*, 198(1-2):1–28, 2004.
- [24] A. Zabrodin. Whitham hierarchy in growth problems. *Theoretical and mathematical physics*, 142(2):166–182, 2005.
- [25] A. Abanov, M. Mineev-Weinstein, and A. Zabrodin. Multi-cut solutions of Laplacian growth. *Physica D: Nonlinear Phenomena*, 238(17):1787–1796, 2009.
- [26] P. B. Wiegmann and A. Zabrodin. Conformal maps and integrable hierarchies. *Communications in Mathematical Physics*, 213:523–538, 2000.
- [27] L. Martínez-Alonso and E. Medina. Genus-zero Whitham hierarchies in conformal-map dynamics. *Physics Letters B*, 641(6):466–473, 2006.
- [28] C.-Z. Wu and X. Zhou. An extension of the Kadomtsev–Petviashvili hierarchy and its Hamiltonian structures. *Journal of Geometry and Physics*, 106:327–341, 2016.
- [29] L. Bonora and C. S. Xiong. The (n, m) -th Korteweg–de Vries hierarchy and the associated W -algebra. *Journal of Mathematical Physics*, 35(11):5781–5819, 1994.
- [30] B. Konopelchenko and L. Martínez-Alonso. $\bar{\partial}$ -equations, integrable deformations of quasiconformal mappings and Whitham hierarchy. *Physics letters A*, 286(2-3):161–166, 2001.
- [31] F. Guil, M. Manas, and L. Martínez-Alonso. The Whitham hierarchies: reductions and hodograph solutions. *Journal of Physics A: Mathematical and General*, 36(14):4047, 2003.
- [32] M. Manas, E. Medina, and L. Martínez-Alonso. On the Whitham hierarchy: dressing scheme, string equations and additional symmetries. *Journal of Physics A: Mathematical and General*, 39(10):2349, 2006.
- [33] L. Martínez-Alonso, E. Medina, and M. Manas. String equations in Whitham hierarchies: τ -functions and Virasoro constraints. *Journal of mathematical physics*, 47(8), 2006.
- [34] K. Takasaki and T. Takebe. Universal Whitham hierarchy, dispersionless Hirota equations and multicomponent KP hierarchy. *Physica D: Nonlinear Phenomena*, 235(1-2):109–125, 2007.
- [35] K. Takasaki and T. Takebe. Löwner equations, Hirota equations and reductions of the universal Whitham hierarchy. *Journal of Physics A: Mathematical and Theoretical*, 41(47):475206, 2008.
- [36] B. M. Szablikowski and M. Błaszak. Dispersionful analog of the Whitham hierarchy. *Journal of mathematical physics*, 49(8), 2008.
- [37] H.-F. Shen, N.-C. Lee, and M.-H. Tu. Kernel formula approach to the universal Whitham hierarchy. *Theoretical and Mathematical Physics*, 165:1456–1469, 2010.
- [38] K. Takasaki, T. Takebe, and L. P. Teo. Non-degenerate solutions of the universal Whitham hierarchy. *Journal of Physics A: Mathematical and Theoretical*, 43(32):325205, 2010.
- [39] G. Carlet and L. P. Mertens. Principal hierarchies of infinite-dimensional Frobenius manifolds: The extended 2d Toda lattice. *Advances in Mathematics*, 278:137–181, 2015.
- [40] S. Ma. Principal hierarchy of infinite-dimensional Frobenius manifold underlying the extended Kadomtsev–Petviashvili hierarchy. *arXiv preprint arXiv:2309.10389*, 2023.
- [41] S. Ma. Principal hierarchy of Frobenius manifolds associated with rational and trigonometric Landau–Ginzburg superpotentials. *arXiv preprint arXiv:2311.17363*, 2023.

SHILIN MA, SCHOOL OF MATHEMATICAL SCIENCE, TSINGHUA UNIVERSITY, BEIJING, 100084,
CHINA

Email address: mashilin@tsinghua.edu.cn