# Regularity of an FI-module via the derivative functor

Cihan Bahran

Department of Mathematics, Bilkent University Ankara 06800, Turkey cihan.bahran@bilkent.edu.tr

#### Abstract

We characterize the regularity of an FI-module using the derivative functors.

## Contents

1	Introduction	1
2	Homological algebra of FI-modules	4
	2.1 Preliminaries	4
	2.2 Proving Theorem A	5

## **1** Introduction

An **FI**-module is a functor  $\mathbf{FI} \to \mathbb{Z}$ -Mod where **FI** is the category of finite sets and injections and  $\mathbb{Z}$ -Mod is the category of abelian groups. We write **FI**-Mod for the functor category [**FI**,  $\mathbb{Z}$ -Mod]. Given an **FI**-module W, we write

 $\deg(W) := \min\{d \ge -1 : W_S = 0 \text{ for } |S| > d\}$  $\in \{-1, 0, 1, 2, 3, \dots\} \cup \{\infty\}.$ 

<sup>2010</sup> Mathematics Subject Classification. Primary 18A25.

Key words and phrases.  ${\bf FI}{\rm -modules},$  regularity.

**FI-homology and regularity.** A systematic way of investigating the "unstable" part of an **FI**-module V is to study its quotient  $H_0^{\mathbf{FI}}(V)$  defined by

$$\mathrm{H}_{0}^{\mathbf{FI}}(V)_{S} := \operatorname{coker}\left(\bigoplus_{A \subsetneq S} V_{A} \to V_{S}\right)$$

The functor  $\mathrm{H}_{0}^{\mathbf{FI}}$ : **FI-Mod**  $\rightarrow$  **FI-Mod** is right exact and has left derived functors  $\mathrm{H}_{i}^{\mathbf{FI}} := \mathrm{L}_{i}\mathrm{H}_{0}^{\mathbf{FI}}$ . Setting  $t_{i}(V) := \mathrm{deg}(\mathrm{H}_{i}^{\mathbf{FI}}(V))$ , we say V is **presented in finite degrees** if  $t_{0}(V)$  and  $t_{1}(V)$  are finite. We also set the **regularity** of V as

$$\operatorname{reg}(V) := \max\{t_i(V) - i : i \ge 1\} \\ \in \{-2, -1, 0, 1, \dots\} \cup \{\infty\}$$

which is finite whenever V is presented in finite degrees [CE17, Theorem A].

The derivative functor, iterated and derived. Given any FI-module V, we write  $\Sigma V$  for the composition

$$\mathbf{FI} \xrightarrow{-\sqcup \{*\}} \mathbf{FI} \xrightarrow{V} \mathbb{Z}$$
-Mod

and call it the **shift** of V. This procedure defines the **shift functor** 

 $\Sigma \colon \mathbf{FI}\text{-}\mathsf{Mod} \to \mathbf{FI}\text{-}\mathsf{Mod}\,.$ 

The inclusions  $S \hookrightarrow S \sqcup \{*\}$  for all finite sets S define a natural transformation

 $\iota\colon \operatorname{id}_{\mathbf{FI}\operatorname{\mathsf{-Mod}}} \to \Sigma \ ,$ 

whose cokernel  $\Delta := \operatorname{coker}(\iota)$  we call the **derivative functor**: it is a right exact functor

 $\Delta \colon \mathbf{FI}\operatorname{\mathsf{-Mod}} o \mathbf{FI}\operatorname{\mathsf{-Mod}}$  .

For every  $i, a \geq 0$ , we write  $H_i^{\Delta^a} := L_i \Delta^a$  for the *i*-th left derived functor of the *a*-fold composite  $\Delta^a$ .

The objective of this paper is to characterize the regularity of an **FI**-module in terms of the derivative functor.

**Definition 1.1.** Given an **FI**-module V and  $i \ge 0$ , we write

width<sup>$$\Delta$$</sup><sub>*i*</sub>(*V*) := sup{deg(H <sup>$\Delta$ a</sup>(*V*)) + *a* : *a* ≥ 0 with H <sup>$\Delta$ a</sup>(*V*) ≠ 0}  
  $\in \{0, 1, 2, ...\} \cup \{-\infty, \infty\}$ 

with the convention  $\sup(\emptyset) = -\infty$ .

**Theorem A.** Let V be an **FI**-module presented in finite degrees which is not  $H_0^{\mathbf{FI}}$ -acyclic. Then for every  $i \ge 1$ , we have

width<sub>*i*</sub><sup>$$\Delta$$</sup>(V) - *i* = reg(V).

**Relationship with previous literature.** The invariant width  ${}_{1}^{\Delta}(V)$  was first considered explicitly in [Ram18, Definition 3.7] (denoted  $\partial$ width there).

The invariants width<sup> $\Delta$ </sup><sub>*i*</sub>(*V*) can be used to conceptually compartmentalize the proof of [CE17, Theorem A], [Ram18, Theorem 3.19] bounding reg(*V*) in terms of  $t_0(V), t_1(V)$  into three parts as follows:

**Theorem 1.2** ([Ram18, Corollary 3.18], based on [CE17, Theorem E]). Every FImodule V presented in finite degrees satisfies

width 
$$_1^{\boldsymbol{\Delta}}(V) \le t_1(V) + \min\{t_0(V), t_1(V)\}$$

**Theorem 1.3** ([CE17, proof of Theorem 4.8]). For every **FI**-module V, there is a weakly decreasing sequence

width<sub>1</sub><sup>$$\Delta$$</sup>(V) - 1 ≥ width<sub>2</sub> <sup>$\Delta$</sup> (V) - 2 ≥ width<sub>3</sub> <sup>$\Delta$</sup> (V) - 3 ≥ · · ·

**Theorem 1.4** ([CE17, proof of Theorem A, pages 2403-2404]). For every FI-module V with  $t_0(V) < t_1(V) < \infty$ , we have

$$\operatorname{reg}(V) \le \max\left(\{t_1(V) - 1\} \cup \{\operatorname{width}_i^{\mathbf{\Delta}}(V) - i : i \ge 2\}\right).$$

These are what partially led me to think something like Theorem A could be true, collapsing the inequalities in the above two results to equalities. It is also perhaps worth mentioning that combining [Ram18, Theorem 4.7, part (2)], [Bah24, Theorem A], [Bah24, Corollary 2.9] with Theorem A, the bound in Theorem 1.2 can be sharpened into the following:

**Theorem 1.5.** Given integers  $a, b \ge -1$  and  $i \ge 1$ , we have

$$\max\left\{ \operatorname{width}_{i}^{\mathbf{\Delta}}(V) \colon \frac{V \text{ is an } \mathbf{FI}\text{-module with}}{t_{0}(V) \leq a, t_{1}(V) \leq b} \right\} = \begin{cases} -\infty & \text{if } a = -1 \text{ or } b \leq 0, \\ i + a + b - 1 & \text{if } 0 \leq a < b, \\ i + 2b - 2 & \text{if } a \geq b \geq 1. \end{cases}$$

Another result which can be interpreted in terms of width  ${}^{\mathbf{\Delta}}_{1}(V)$  is about polynomial conditions on **FI**-modules:

**Theorem 1.6** ([RW17, proof of Proposition 4.18]). Let  $t, w \ge -1$  be integers and V an **FI**-module with  $t_0(V) \le t$  and width  ${}^{\mathbf{\Delta}}_1(V) \le w$ . Then in the sense of [RW17, Definition 4.10], V is of degree t at w.

Due to Theorem 1.6, I had initially thought that establishing Theorem A for i = 1 would be a necessary step to prove the characterization of polynomiality in [Bah23, Theorem B], but later found a more direct argument during writing that paper. The current paper is a result of revisiting the invariants width  $\frac{\Delta}{i}(V)$ .

## 2 Homological algebra of FI-modules

#### 2.1 Preliminaries

In this section we recall relevant homological results from the **FI**-module literature. The first one is a short exact sequence that facilitates inductive arguments about the derived functors  $H_i^{\Delta^a}$ .

**The kernel functor.** Recall from the introduction that the inclusions  $S \hookrightarrow S \sqcup \{*\}$  for all finite sets S define a natural transformation

$$: \operatorname{id}_{\mathbf{FI}\operatorname{-Mod}} \to \Sigma$$
,

whose cokernel is the derivative functor  $\Delta$ . For its kernel, we write  $\mathbf{K} := \ker(\iota)$ : it is a left exact functor

$$\mathbf{K} \colon \mathbf{FI}\operatorname{\mathsf{-Mod}} o \mathbf{FI}\operatorname{\mathsf{-Mod}}$$
 .

**Proposition 2.1** ([CE17, (18)]). For every **FI**-module V and integers  $i, a \ge 1$ , there is a short exact sequence

$$0 \to \mathbf{\Delta} \Big( \mathrm{H}_{i}^{\mathbf{\Delta}^{a-1}}(V) \Big) \to \mathrm{H}_{i}^{\mathbf{\Delta}^{a}}(V) \to \mathbf{K} \Big( \mathrm{H}_{i-1}^{\mathbf{\Delta}^{a-1}}(V) \Big) \to 0$$

of FI-modules.

**Torsion and local cohomology.** We say an **FI**-module V is **torsion** if for every finite set S and  $x \in V_S$ , there is an injection  $\alpha \colon S \hookrightarrow T$  such that  $V_{\alpha}(x) = 0 \in V_T$ . We write

#### $\mathrm{H}^0_\mathfrak{m}\colon \mathbf{FI}\text{-}\mathsf{Mod}\to \mathbf{FI}\text{-}\mathsf{Mod}$

for the functor which assigns an **FI**-module its largest torsion **FI**-submodule; it is left exact. For each  $j \ge 0$ , the *j*-th **local cohomology** functor is the *j*-th derived functor  $\mathrm{H}^{j}_{\mathfrak{m}} := \mathrm{R}^{j}\mathrm{H}^{0}_{\mathfrak{m}}$  of  $\mathrm{H}^{0}_{\mathfrak{m}}$ , for which we set

$$h^{j}(V) := \deg(\mathrm{H}^{j}_{\mathfrak{m}}(V))$$
  
 
$$\in \{-1, 0, 1, \dots\} \cup \{\infty\}$$

for every  $\mathbf{FI}$ -module V.

We recall the characterization of regularity via local cohomology, which then justifies the well-definition of the critical index.

**Theorem 2.2** ([NSS18, Theorem 1.1, Remark 1.3]). Let V be an FI-module presented in finite degrees which is not  $H_0^{FI}$ -acyclic. Then the maximum of the finite set

$$\emptyset \neq \{h^{j}(V) + j : h^{j}(V) \ge 0\} \subseteq \mathbb{Z}_{\ge 0}$$

is equal to reg(V).

**Definition 2.3.** For an **FI**-module presented in finite degrees which is not  $H_0^{FI}$ -acyclic, we define its **critical index** as

 $\operatorname{crit}(V) := \min\{j : h^j(V) \ge 0 \text{ and } h^j(V) + j = \operatorname{reg}(V)\}.$ 

#### 2.2 Proving Theorem A

In this section we prove Theorem A. As ingredients towards that end, we relate the vanishing degrees of the derived functors  $H_i^{\Delta^a}$  with the local cohomology degrees in Proposition 2.4 (the most technical part of the paper) and identify non-vanishing degrees for these functors in Proposition 2.5 using the critical index.

**Proposition 2.4.** For an **FI**-module V presented in finite degrees, the following hold:

- (1) For every  $a \ge 0$  and  $u \ge 0$ , if  $h^u(\Delta^a V) \ge 0$  then we have  $h^u(\Delta^a V) + u + a \le \max\{h^j(V) + j : h^j(V) \ge 0 \text{ and } 0 \le j \le u + a\}$ .
- (2) For every  $a \ge 1$  and  $i \ge 1$ , if  $\operatorname{H}_{i}^{\Delta^{a}}(V) \ne 0$  then we have  $\operatorname{deg} \operatorname{H}_{i}^{\Delta^{a}}(V) + a \le \max\left\{h^{j}(V) + j + i : h^{j}(V) \ge 0 \text{ and } 0 \le j \le a - i\right\}.$

*Proof.* (1) We employ induction on a: the base case a = 0 is immediate so assume  $a \ge 1$ . By [Bah23, Proposition 2.6] applied to  $\Delta^{a-1}V$ , there is an exact sequence

$$\Sigma \mathrm{H}^{u}_{\mathfrak{m}}(\mathbf{\Delta}^{a-1}V) \to \mathrm{H}^{u}_{\mathfrak{m}}(\mathbf{\Delta}^{a}V) \to \mathrm{H}^{u+1}_{\mathfrak{m}}(\mathbf{\Delta}^{a-1}V)$$

of **FI**-modules each of finite degree. If deg  $H^u_m(\Delta^{a-1}V) = h^u(\Delta^{a-1}V) \ge 0$ , then

$$\deg \Sigma \mathrm{H}^{u}_{\mathfrak{m}} \left( \Delta^{a-1} V \right) + u + a = \left( \deg \mathrm{H}^{u}_{\mathfrak{m}} \left( \Delta^{a-1} V \right) - 1 \right) + u + a$$
$$= h^{u} \left( \Delta^{a-1} V \right) + u + a - 1$$
$$\leq \max \left\{ h^{j}(V) + j : h^{j}(V) \ge 0 \text{ and } 0 \le j \le u + a - 1 \right\}$$

by the induction hypothesis. If deg  $H^{u+1}_{\mathfrak{m}}(\Delta^{a-1}V) = h^{u+1}(\Delta^{a-1}V) \ge 0$ , then

$$\deg \mathcal{H}^{u+1}_{\mathfrak{m}}(\boldsymbol{\Delta}^{a-1}V) + u + a = h^{u+1}(\boldsymbol{\Delta}^{a-1}V) + u + a$$
$$= h^{u+1}(\boldsymbol{\Delta}^{a-1}V) + (u+1) + (a-1)$$
$$\leq \max\{h^{j}(V) + j : h^{j}(V) \ge 0 \text{ and } 0 \le j \le u + a\}$$

again by the induction hypothesis. Therefore if deg  $H^u_{\mathfrak{m}}(\Delta^a V) = h^u(\Delta^a V) \ge 0$ , by the exact sequence we have started with, we have that either

$$\deg \mathrm{H}^{u}_{\mathfrak{m}}(\mathbf{\Delta}^{a-1}V) \ge 0 \quad \text{or} \quad \deg \mathrm{H}^{u+1}_{\mathfrak{m}}(\mathbf{\Delta}^{a-1}V) \ge 0$$

Thus

$$h^{u}(\boldsymbol{\Delta}^{a}V) + u + a \leq \max\left\{\deg \boldsymbol{\Sigma} \mathbf{H}^{u}_{\mathfrak{m}}\left(\boldsymbol{\Delta}^{a-1}V\right) + u + a, \ \deg \mathbf{H}^{u+1}_{\mathfrak{m}}\left(\boldsymbol{\Delta}^{a-1}V\right) + u + a\right\}$$
$$\leq \max\left\{h^{j}(V) + j : h^{j}(V) \geq 0 \ \text{and} \ 0 \leq j \leq u + a\right\}$$

as desired.

(2) Let us call the desired statement P(a,i) and employ induction on a + i. For a + i = 2, we have a = i = 1 so that the higher left derived functors of  $\Delta^{a-1} = \mathrm{id}_{\mathbf{FI}-\mathsf{Mod}}$ 

vanish, and Proposition 2.1 reduces to an isomorphism  $H_1^{\Delta}(V) \cong \mathbf{K}(V)$ . Indeed here  $\deg H_1^{\Delta}(V) + 1 = \deg(\mathbf{K}(V)) + 1 = h^0(V) + 1$ ,

verifying P(1, 1). For  $a + i \ge 3$ , there are three cases:

- a = 1 and  $i \ge 2$ : the higher left derived functors of  $\Delta^{a-1} = \mathrm{id}_{\mathbf{FI}-\mathsf{Mod}}$  vanish, and Proposition 2.1 reduces to the fact  $\mathrm{H}_i^{\Delta}(V) = 0$ , therefore P(1, i) is vacuously true.
- $a \ge 2$  and i = 1: By Proposition 2.1 we have a short exact sequence

$$0 \to \mathbf{\Delta} \Big( \mathrm{H}_{1}^{\mathbf{\Delta}^{a-1}}(V) \Big) \to \mathrm{H}_{1}^{\mathbf{\Delta}^{a}}(V) \to \mathbf{K} \big( \mathbf{\Delta}^{a-1}(V) \big) \to 0 \,.$$

If  $\mathrm{H}_{1}^{\mathbf{\Delta}^{a-1}}(V) \neq 0$ , then

$$\begin{split} \operatorname{deg} \mathbf{\Delta} \Big( \mathrm{H}_1^{\mathbf{\Delta}^{a-1}}(V) \Big) + a &\leq \operatorname{deg} \mathrm{H}_1^{\mathbf{\Delta}^{a-1}}(V) + a - 1 \\ &\leq \max \big\{ h^j(V) + j + 1 : h^j(V) \geq 0 \text{ and } 0 \leq j \leq a - 2 \big\} \end{split}$$

by the induction hypothesis P(a-1,1). If  $h^0(\Delta^{a-1}(V)) \ge 0$ , then

$$\deg \mathbf{K} (\Delta^{a-1}(V)) + a = h^0 (\Delta^{a-1}(V)) + (a-1) + 1$$
  
 
$$\leq \max \{ h^j(V) + j : h^j(V) \ge 0 \text{ and } 0 \le j \le a-1 \} + 1$$

by part (1). Therefore if  $H_1^{\Delta^a}(V) \neq 0$ , we have

 $\deg \mathrm{H}_1^{\mathbf{\Delta}^{\!\!\!a}}\!(V) + a \leq \max \bigl\{ h^j(V) + j + 1 : h^j(V) \geq 0 \text{ and } 0 \leq j \leq a-1 \bigr\} \ ,$  establishing P(a,1).

• 
$$a \ge 2$$
 and  $i \ge 2$ : if  $\operatorname{H}_{i}^{\Delta^{a-1}}(V) \ne 0$ , then  
 $\operatorname{deg} \Delta\left(\operatorname{H}_{i}^{\Delta^{a-1}}(V)\right) + a \le \operatorname{deg} \operatorname{H}_{i}^{\Delta^{a-1}}(V) + a - 1$   
 $\le \max\left\{h^{j}(V) + j + i : h^{j}(V) \ge 0 \text{ and } 0 \le j \le (a-1) - i\right\}$ 

by the induction hypothesis P(a-1,i). If  $\operatorname{H}_{i-1}^{\Delta^{a-1}}(V) \neq 0$ , then

$$\deg \mathbf{K} \Big( \mathbf{H}_{i-1}^{\mathbf{\Delta}^{a-1}}(V) \Big) + a \leq \deg \mathbf{H}_{i-1}^{\mathbf{\Delta}^{a-1}}(V) + a = \deg \mathbf{H}_{i-1}^{\mathbf{\Delta}^{a-1}}(V) + (a-1) + 1 \leq \max \Big\{ h^{j}(V) + j + i - 1 : h^{j}(V) \ge 0 \text{ and } 0 \le j \le a - i \Big\} + 1$$

by the induction hypothesis P(a-1, i-1). Therefore if  $H_i^{\Delta^a}(V) \neq 0$ , by Proposition 2.1 either  $H_i^{\Delta^{a-1}}(V) \neq 0$  or  $H_{i-1}^{\Delta^{a-1}}(V) \neq 0$ , and

$$\deg \mathbf{H}_{i}^{\mathbf{\Delta}^{a}}(V) + a = \max\left\{\deg \mathbf{\Delta}\left(\mathbf{H}_{i}^{\mathbf{\Delta}^{a-1}}(V)\right) + a, \deg \mathbf{K}\left(\mathbf{H}_{i-1}^{\mathbf{\Delta}^{a-1}}(V)\right) + a\right\}$$
$$\leq \max\left\{h^{j}(V) + j + i : h^{j}(V) \ge 0 \text{ and } 0 \le j \le a - i\right\},$$

establishing P(a, i).

**Proposition 2.5.** Let V be an **FI**-module presented in finite degrees which is not  $H_0^{\mathbf{FI}}$ -acyclic. Setting  $\rho := \operatorname{reg}(V)$  and  $\gamma := \operatorname{crit}(V)$ , the  $\mathfrak{S}_{\rho-\gamma}$ -modules

(1)  $\operatorname{H}_{\mathfrak{m}}^{\gamma-j}(\Delta^{j}V)_{\rho-\gamma}$  for each  $0 \leq j \leq \gamma$ ,

- (2)  $\mathbf{K}(\mathbf{\Delta}^{\gamma}V)_{\rho-\gamma}$ ,
- (3)  $\operatorname{H}_{i}^{\mathbf{\Delta}^{\gamma+i}}(V)_{\rho-\gamma}$  for each  $i \geq 1$ ,

are all nonzero and pairwise isomorphic.

*Proof.* By Definition 2.3, we have  $h^{\gamma}(V) \ge 0$  and  $h^{\gamma}(V) + \gamma = \rho$ , so that  $0 \le \deg \operatorname{H}^{\gamma}_{\mathfrak{m}}(V) = h^{\gamma}(V) = \rho - \gamma$ 

and hence  $\mathrm{H}^{\gamma}_{\mathfrak{m}}(V)_{\rho-\gamma} \neq 0$ . Next we will show that

$$\mathrm{H}^{\gamma}_{\mathfrak{m}}(V)_{\rho-\gamma} \cong \mathrm{H}^{\gamma-j}_{\mathfrak{m}}(\mathbf{\Delta}^{j}V)_{\rho-\gamma}$$

for every  $0 \leq j \leq \gamma$  by induction on  $\gamma = \operatorname{crit}(V)$ . The base case  $\gamma = 0$  is a tautology, so we assume  $\gamma \geq 1$ . By [Bah23, Proposition 2.10] the **FI**-module  $\Delta V$  is not  $\mathrm{H}_{0}^{\mathbf{FI}}$ -acyclic with

$$\operatorname{reg}(\mathbf{\Delta}V) = \rho - 1$$
 and  $\operatorname{crit}(\mathbf{\Delta}V) = \gamma - 1$ 

Therefore we may apply the induction hypothesis to  $\Delta V$  and get

$$\begin{aligned} \mathrm{H}_{\mathfrak{m}}^{\gamma-1}(\boldsymbol{\Delta}V)_{\rho-\gamma} &= \mathrm{H}_{\mathfrak{m}}^{\gamma-1}(\boldsymbol{\Delta}V)_{(\rho-1)-(\gamma-1)} \\ &\cong \mathrm{H}_{\mathfrak{m}}^{\gamma-1-j'}(\boldsymbol{\Delta}^{j'}\boldsymbol{\Delta}V)_{(\rho-1)-(\gamma-1)} \\ &= \mathrm{H}_{\mathfrak{m}}^{\gamma-(j'+1)}(\boldsymbol{\Delta}^{j'+1}V)_{\rho-\gamma} \end{aligned}$$

as  $\mathfrak{S}_{\rho-\gamma}$ -modules for every  $0 \leq j' \leq \gamma - 1$ . Reindexing as j = j' + 1,

$$\mathrm{H}^{\gamma-1}_{\mathfrak{m}}(\mathbf{\Delta} V)_{\rho-\gamma} \cong \mathrm{H}^{\gamma-j}_{\mathfrak{m}}(\mathbf{\Delta}^{j} V)_{\rho-\gamma}$$

for every  $1 \le j \le \gamma$ . By [Bah23, Proposition 2.6], there is an exact sequence

$$\Sigma \mathrm{H}^{\gamma-1}_{\mathfrak{m}}(V) \to \mathrm{H}^{\gamma-1}_{\mathfrak{m}}(\mathbf{\Delta} V) \to \mathrm{H}^{\gamma}_{\mathfrak{m}}(V) \to \Sigma \mathrm{H}^{\gamma}_{\mathfrak{m}}(V)$$

for which we have

$$\left(\mathbf{\Sigma} \mathbf{H}_{\mathfrak{m}}^{\gamma-1}(V)\right)_{\rho-\gamma} = \mathbf{H}_{\mathfrak{m}}^{\gamma-1}(V)_{\rho-\gamma+1} = 0$$

because  $h^{\gamma-1}(V) + \gamma - 1 < \rho$ , and

$$\Sigma \mathrm{H}^{\gamma}_{\mathfrak{m}}(V))_{\rho-\gamma} = \mathrm{H}^{\gamma}_{\mathfrak{m}}(V)_{\rho-\gamma+1} = 0$$

because  $h^{\gamma}(V) + \gamma = \rho$  by the definition of  $\gamma = \operatorname{crit}(V)$ . Thus evaluating the exact sequence at degree  $\rho - \gamma$  results in an isomorphism

$$\mathrm{H}^{\gamma-1}_{\mathfrak{m}}(\mathbf{\Delta} V)_{\rho-\gamma} \cong \mathrm{H}^{\gamma}_{\mathfrak{m}}(V)_{\rho-\gamma}$$

of  $\mathfrak{S}_{\rho-\gamma}$ -modules, completing the j = 0 case of (1).

For the isomorphism between the modules between (1) and (2), apply [Bah23, Proposition 2.6] to the **FI**-module  $\Delta^{\gamma} V$  and get an exact sequence

$$0 \to \mathbf{K}(\mathbf{\Delta}^{\gamma} V) \to \mathrm{H}^{0}_{\mathfrak{m}}(\mathbf{\Delta}^{\gamma} V) \to \mathbf{\Sigma} \mathrm{H}^{0}_{\mathfrak{m}}(\mathbf{\Delta}^{\gamma} V) \ ,$$

for which we have

$$\left(\Sigma \mathrm{H}^{0}_{\mathfrak{m}}\left(\boldsymbol{\Delta}^{\gamma}V\right)\right)_{\rho-\gamma} = \mathrm{H}^{0}_{\mathfrak{m}}\left(\boldsymbol{\Delta}^{\gamma}V\right)_{\rho-\gamma+1} = 0$$

because by [Bah23, Proposition 2.10] we have  $\operatorname{crit}(\Delta^{\gamma} V) = 0$  and

$$h^0(\mathbf{\Delta}^{\gamma} V) = \operatorname{reg}(\mathbf{\Delta}^{\gamma} V) = \operatorname{reg}(V) - \gamma = \rho - \gamma,$$

resulting in an isomorphism

$$\mathbf{K}(\mathbf{\Delta}^{\gamma}V)_{\rho-\gamma} \cong \mathrm{H}^{0}_{\mathfrak{m}}(\mathbf{\Delta}^{\gamma}V)_{\rho-\gamma}$$

of  $\mathfrak{S}_{\rho-\gamma}$ -modules as desired.

Consider the short exact sequence

$$0 \to \mathbf{\Delta} \Big( \mathbf{H}_{i}^{\mathbf{\Delta}^{\gamma+i-1}}(V) \Big) \to \mathbf{H}_{i}^{\mathbf{\Delta}^{\gamma+i}}(V) \to \mathbf{K} \Big( \mathbf{H}_{i-1}^{\mathbf{\Delta}^{\gamma+i-1}}(V) \Big) \to 0 \tag{\dagger}$$

of **FI**-modules given by Proposition 2.1. For i = 1 it becomes

$$0 \to \mathbf{\Delta} \big( \mathrm{H}_{1}^{\mathbf{\Delta}^{\gamma}}(V) \big) \to \mathrm{H}_{1}^{\mathbf{\Delta}^{\gamma+1}}(V) \to \mathbf{K}(\mathbf{\Delta}^{\gamma}(V)) \to 0 \,. \tag{\dagger\dagger}$$

Here either  $H_1^{\Delta^{\gamma}}(V) = 0$ , or by Proposition 2.4 we have  $H_1^{\Delta^{\gamma}}(V) \neq 0$  and

$$\begin{split} \operatorname{deg} \mathbf{\Delta} \big( \mathrm{H}_{1}^{\mathbf{\Delta}^{\gamma}}(V) \big) + \gamma &\leq \operatorname{deg} \mathrm{H}_{1}^{\mathbf{\Delta}^{\gamma}}(V) + \gamma - 1 \\ &\leq \max \big\{ h^{j}(V) + j + 1 : h^{j}(V) \geq 0 \text{ and } 0 \leq j \leq \gamma - 1 \big\} - 1 \\ &= \max \big\{ h^{j}(V) + j : h^{j}(V) \geq 0 \text{ and } 0 \leq j \leq \gamma - 1 \big\} \\ &< \operatorname{reg}(V) = \rho \end{split}$$

by the definition of  $\gamma = \operatorname{crit}(V)$ . Thus we always have deg  $\Delta(\operatorname{H}_{1}^{\Delta^{\gamma}}(V)) < \rho - \gamma$  and ( $\dagger \dagger$ ) in degree  $\rho - \gamma$  becomes an isomorphism

$$\mathrm{H}_{1}^{\mathbf{\Delta}^{\gamma+1}}(V)_{\rho-\gamma} \cong \mathbf{K}(\mathbf{\Delta}^{\gamma}(V))_{\rho-\gamma}$$

of  $\mathfrak{S}_{\rho-\lambda}$ -modules, bridging the module in (2) with the one in (3) when i = 1. We shall complete the proof by showing

$$\mathrm{H}_{i}^{\mathbf{\Delta}^{\gamma+i}}(V)_{\rho-\gamma} \cong \mathrm{H}_{i-1}^{\mathbf{\Delta}^{\gamma+i-1}}(V)_{\rho-\gamma}$$

for every  $i \ge 2$ . To that end either  $H_i^{\Delta^{\gamma+i-1}}(V) = 0$ , or by part (2) of Proposition 2.4 we have  $H_i^{\Delta^{\gamma+i-1}}(V) \ne 0$  and

$$\begin{split} \operatorname{deg} \mathbf{\Delta} \Big( \mathrm{H}_{i}^{\mathbf{\Delta}^{\gamma+i-1}}(V) \Big) + \gamma + i &\leq \operatorname{deg} \mathrm{H}_{i}^{\mathbf{\Delta}^{\gamma+i-1}}(V) + \gamma + i - 1 \\ &\leq \max \big\{ h^{j}(V) + j + i : h^{j}(V) \geq 0 \text{ and } 0 \leq j \leq \gamma - 1 \big\} \\ &= \max \big\{ h^{j}(V) + j : h^{j}(V) \geq 0 \text{ and } 0 \leq j \leq \gamma - 1 \big\} + i \\ &< \operatorname{reg}(V) + i = \rho + i \end{split}$$

by the definition of critical index. In any case we have

$$\deg \Delta \left( \mathrm{H}_{i}^{\Delta^{\gamma+i-1}}(V) \right) < \rho - \gamma$$

and hence evaluating (†) in degree  $\rho - \gamma$  results in an isomorphism

$$\mathbf{H}_{i}^{\mathbf{\Delta}^{\gamma+i}}(V)_{\rho-\gamma} \cong \mathbf{K} \Big( \mathbf{H}_{i-1}^{\mathbf{\Delta}^{\gamma+i-1}}(V) \Big)_{\rho-\gamma}$$

of  $\mathfrak{S}_{\rho-\gamma}$ -modules. Here by part (2) of Proposition 2.4, the **FI**-module  $W := \mathrm{H}_{i-1}^{\mathbf{\Delta}^{\gamma+i-1}}(V)$ 

satisfies

 $\deg W + \gamma + i - 1 \le \max \left\{ h^j(V) + j + i - 1 : h^j(V) \ge 0 \text{ and } 0 \le j \le \gamma \right\} = \rho + i - 1$  by the definition of  $\gamma = \operatorname{crit}(V)$ . Thus  $\deg W \le \rho - \gamma$  and hence  $(\mathbf{K}W)_{\rho-\gamma} = W_{\rho-\gamma}$ .  $\Box$ 

*Proof of* **Theorem A**. For every  $i \ge 1$ , whenever  $\operatorname{H}_{i}^{\Delta^{a}}(V) \ne 0$  we necessarily have  $a \ge 1$  and part (2) of Proposition 2.4 yields

$$\deg \mathcal{H}_i^{\Delta^a}(V) + a \le \max \left\{ h^j(V) + j + i : h^j(V) \ge 0 \text{ and } 0 \le j \le a - i \right\}$$
$$= \max \left\{ h^j(V) + j : h^j(V) \ge 0 \text{ and } 0 \le j \le a - i \right\} + i$$
$$\le \operatorname{reg}(V) + i$$

where the last line is by Theorem 2.2. As a result, reg(V) + i is an upper bound for the set

$$\mathbf{D}_{i}(V) := \left\{ \operatorname{deg}\left(\operatorname{H}_{i}^{\boldsymbol{\Delta}^{a}}(V)\right) + a : \operatorname{H}_{i}^{\boldsymbol{\Delta}^{a}}(V) \neq 0 \right\},\$$

so that

width<sub>*i*</sub><sup>$$\Delta$$</sup>(*V*) = sup **D**<sub>*i*</sub>(*V*) ≤ reg(*V*) + *i*.

On the other hand, setting  $\gamma := \operatorname{crit}(V)$  and invoking Proposition 2.5, for every  $i \ge 1$ we have  $\operatorname{H}_{i}^{\mathbf{\Delta}^{\gamma+i}}(V) \neq 0$  and

$$\deg\left(\mathbf{H}_{i}^{\mathbf{\Delta}^{\gamma+i}}(V)\right) \geq \operatorname{reg}(V) - \gamma,$$
$$\deg\left(\mathbf{H}_{i}^{\mathbf{\Delta}^{\gamma+i}}(V)\right) + \gamma + i \geq \operatorname{reg}(V) + i.$$

But the left hand side of the last inequality is an element of  $\mathbf{D}_i(V)$ , whence

width<sub>*i*</sub><sup>$$\Delta$$</sup>(V) = sup **D**<sub>*i*</sub>(V) ≥ reg(V) + *i*.

### References

- [Bah23] Cihan Bahran, Polynomial conditions and homology of FI-modules, Pacific Journal of Mathematics 324 (2023), no. 2, 207–226. 3, 5, 7, 8
- [Bah24] \_\_\_\_\_, Regularity and stable ranges of FI-modules, Journal of Algebra 641 (2024), 429–497. 3
- [CE17] Thomas Church and Jordan S. Ellenberg, Homology of FI-modules, Geometry & Topology 21 (2017), no. 4, 2373–2418. 2, 3, 4
- [NSS18] Rohit Nagpal, Steven V. Sam, and Andrew Snowden, Regularity of FI-modules and local cohomology, Proceedings of the American Mathematical Society 146 (2018), no. 10, 4117–4126. 4
- [Ram18] Eric Ramos, Homological invariants of **FI**-modules and **FI**<sub>G</sub>-modules, Journal of Algebra **502** (2018), 163–195. 2, 3

[RW17] Oscar Randal-Williams and Nathalie Wahl, *Homological stability for automorphism groups*, Advances in Mathematics **318** (2017), 534–626. **3**