ON THE EXISTENCE OF SOLUTIONS FOR A CLASS OF SYSTEMS OF INTEGRO-DIFFERENTIAL EQUATIONS WITH THE LOGARITHMIC LAPLACIAN AND DRIFT

Yuming Chen¹, Vitali Vougalter²*

¹ Department of Mathematics, Wilfrid Laurier University Waterloo, Ontario, N2L 3C5, Canada e-mail: ychen@wlu.ca

² * Department of Mathematics, University of Toronto Toronto, Ontario, M5S 2E4, Canada e-mail: vitali@math.toronto.edu

Abstract: In this article, we consider a system of integro-differential equations in $L^2(\mathbb{R}, \mathbb{R}^N)$, which contains the logarithmic Laplacian in the presence of transport terms. The linear operators associated with the system satisfy the Fredholm property. By virtue of a fixed point technique, we demonstrate the existence of solutions. We emphasize that the discussion is more complicated than that of the scalar situation as there are more cumbersome technicalities to overcome.

AMS Subject Classification: 35P05, 45K05, 47G20

Key words: solvability, Fredholm operator, logarithmic Laplacian, integral kernel

1. Introduction

The present work is devoted to the solvability of the following system of integrodifferential equations with $1 \le m \le N$,

$$\left[-\frac{1}{2}\ln\left(-\frac{d^2}{dx^2}\right)\right]u_m +$$

$$b_m \frac{du_m}{dx} + a_m u_m + f_m(x) + \int_{-\infty}^{\infty} K_m(x-y)g_m(u(y))dy = 0, \qquad (1.1)$$

where $x \in \mathbb{R}$ and $a_m, b_m \in \mathbb{R}$, $b_m \neq 0$ are constants. Let us recall that the solvability of a single equation resembling (1.1) with the fractional Laplacian in the diffusion term has been considered by Vougalter and Volpert [33].

Biological processes with nonlocal consumption of resources and intra-specific competition can be described by nonlocal reaction-diffusion equations (see, for example, [1, 4, 21, 29] and references therein). In such time-dependent problems the space variable x corresponds to the cell genotype and $u_m(x,t)$ stand for the cell density distributions for various groups of cells as functions of their genotype and time,

$$u(x,t) = (u_1(x,t), u_2(x,t), ..., u_N(x,t))^T.$$
(1.2)

The evolution of the cell densities occurs via cell proliferation, mutations, transport, and cell influx/efflux. The diffusion terms correspond to the change of genotype due to small random mutations and the integral terms describe large mutations. The functions $g_m(u)$ denote the rates of cell births depending on u (density-dependent proliferation). The kernels $K_m(x-y)$ give the proportions of newly born cells changing their genotypes from y to x. We assume that they depend on the distances between the genotypes. The source terms $f_m(x)$ designate the influxes/effluxes of cells for different genotypes. Since models of this kind describe the distributions of population densities with respect to the genotype, the existence of stationary solutions of the nonlocal reaction-diffusion problems corresponds to the existence of biological species.

We introduce the concept of non-Fredholm operators. Consider the problem

$$-\Delta u + V(x)u - au = f, (1.3)$$

where $u\in E=H^2(\mathbb{R}^d)$ and $f\in F=L^2(\mathbb{R}^d)$, $d\in \mathbb{N}$, a is a constant, and the scalar potential function V(x) either vanishes identically or converges to 0 at infinity. If $a\geq 0$, the origin belongs to the essential spectrum of the operator $A:E\to F$ corresponding to the left side of equation (1.3). Consequently, such an operator fails to satisfy the Fredholm property. Its image is not closed, and, for d>1, the dimension of its kernel and the codimension of its image are not finite. Note that for the Fredholm property we consider bounded but not $H^2(\mathbb{R}^d)$ solutions of the corresponding homogeneous adjoint equation. If V(x) vanishes identically, problem (1.3) has constant coefficients and we can apply the Fourier transform to solve it explicitly. If $f\in L^2(\mathbb{R}^d)$ and $xf\in L^1(\mathbb{R}^d)$, then it has a unique solution in $H^2(\mathbb{R}^d)$ if and only if

$$\left(f(x), \frac{e^{ipx}}{(2\pi)^{\frac{d}{2}}}\right)_{L^2(\mathbb{R}^d)} = 0, \quad p \in S_{\sqrt{a}}^d \quad a.e.$$
(1.4)

(see Lemmas 5 and 6 of [31]). Here $S_{\sqrt{a}}^d$ stands for the sphere in \mathbb{R}^d of radius \sqrt{a} centered at the origin. Hence, though our operator does not satisfy the Fredholm property, the solvability conditions can be formulated similarly. But this similarity is only formal since the range of the operator is not closed. The orthogonality conditions (1.4) are with respect to the standard Fourier harmonics, which solve the

homogeneous adjoint problem for (1.3) when the scalar potential function is trivial. They belong to $L^{\infty}(\mathbb{R}^d)$ but they are not square integrable.

Let us mention that elliptic problems containing non-Fredholm operators have been studied actively in recent years. Approaches in weighted Sobolev and Hölder spaces have been developed in [2, 3, 5, 6, 7]. Particularly, when a=0, the operator A is Fredholm in certain properly chosen weighted spaces (see [2, 3, 5, 6, 7]). However, the situation when $a\neq 0$ is considerably different and the approaches developed in these articles are not applicable. Non-Fredholm Schrödinger type operators were treated with the methods of the spectral and the scattering theory in [16, 28, 30, 32]. Fredholm structures, topological invariants, and applications were discussed in [15]. The articles [22] and [27] deal with the Fredholm and properness properties of quasilinear elliptic systems of the second order and of the operators of this kind on \mathbb{R}^N . The exponential decay and Fredholm properties in second-order quasilinear elliptic systems of equations were covered in [23]. The nonlinear non-Fredholm elliptic equations were discussed in [17, 31, 33]. Considerable applications to the theory of reaction-diffusion equations were developed in [13, 14].

The logarithmic Laplacian $\ln(-\Delta)$ is the operator with Fourier symbol $2 \ln |p|$. It appears as the formal derivative $\partial_s|_{s=0}(-\Delta)^s$ of fractional Laplacians at s=0. The operator $(-\Delta)^s$ has been extensively used, for instance, in the studies of anomalous diffusion problems (to name a few, see [26, 32, 33] and references therein). Spectral properties of the logarithmic Laplacian in an open set of finite measure with Dirichlet boundary conditions were investigated in [25] (see also [11]). The studies of $\ln(-\Delta)$ are crucial to understand the asymptotic spectral properties of the family of fractional Laplacians in the limit $s\to 0^+$. In [24], it was established that this operator enables us to characterize the s-dependence of solutions to fractional Poisson problems for the full range of exponents $s \in (0,1)$.

The operators involved in the left side of system (1.1) are given by

$$L_{a_m,b_m} := \frac{1}{2} \ln \left(-\frac{d^2}{dx^2} \right) - b_m \frac{d}{dx} - a_m, \quad a_m, b_m \in \mathbb{R}, \quad b_m \neq 0$$
 (1.5)

with $1 \le m \le N$ and are considered on $L^2(\mathbb{R})$. By means of the standard Fourier transform, it can be easily derived that the essential spectrum of (1.5) is equal to

$$\lambda_{a_m,b_m}(p) = \ln\left(\frac{|p|}{e^{a_m}}\right) - ib_m p, \quad a_m, b_m \in \mathbb{R}, \quad b_m \neq 0, \quad 1 \le m \le N. \quad (1.6)$$

Evidently, for $p \in \mathbb{R}$, $1 \le m \le N$,

$$|\lambda_{a_m,b_m}(p)| = \sqrt{\ln^2\left(\frac{|p|}{e^{a_m}}\right) + b_m^2 p^2} \ge C > 0,$$
 (1.7)

where C is a constant. Let us note that as distinct from the case without the drift term studied in [18], each operator (1.5) has the Fredholm property. Solvability of

certain nonhomogeneous linear problems involving the logarithmic Schrödinger operators in higher dimensions was considered in [19]. The logarithmic Schrödinger operator and associated Dirichlet problems were discussed in [20]. The logarithmic Laplacian also arises in the geometric context of the 0-fractional perimeter, which has been studied in [12]. The article [34] deals with the symmetry of positive solutions for Lane-Emden systems involving the logarithmic Laplacian.

We set $K_m(x) = \varepsilon_m \mathcal{K}_m(x)$, $1 \le m \le N$, where all $\varepsilon_m \ge 0$ and introduce

$$\varepsilon := \max_{1 < m < N} \varepsilon_m. \tag{1.8}$$

In the biological applications, such small and nonnegative parameters ε_m have the meaning that the integral production terms are small with respect to the others, that is, the frequency of large mutations is sufficiently small. Our first assumption is as follows.

Assumption 1. Let $1 \leq m \leq N$, the constants a_m , $b_m \in \mathbb{R}$ with $b_m \neq 0$. The functions $f_m(x) : \mathbb{R} \to \mathbb{R}$ do not vanish identically on the real line for some $1 \leq m \leq N$ and $f_m(x) \in L^2(\mathbb{R})$. Let us also assume that $\mathcal{K}_m(x) : \mathbb{R} \to \mathbb{R}$ and $\mathcal{K}_m(x) \in L^1(\mathbb{R})$. Moreover,

$$\mathcal{K}^2 := \sum_{m=1}^N \|\mathcal{K}_m\|_{L^1(\mathbb{R})}^2 > 0. \tag{1.9}$$

For a vector function

$$u(x) = (u_1(x), u_2(x), ..., u_N(x))^T,$$
(1.10)

we will use the norm

$$||u||_{L^{2}(\mathbb{R},\mathbb{R}^{N})}^{2} := \sum_{m=1}^{N} ||u_{m}||_{L^{2}(\mathbb{R})}^{2}.$$
 (1.11)

When all the nonnegative parameters ε_m vanish, we obtain the linear system of equations (4.1). By virtue of Lemma 5 further down along with Assumption 1 and Remark 6, system (4.1) has a unique solution

$$u_0(x) \in L^2(\mathbb{R}, \mathbb{R}^N), \tag{1.12}$$

which does not vanish identically on the real line. Let us seek the resulting solution of the nonlinear system of equations (1.1) as

$$u(x) = u_0(x) + u_p(x), (1.13)$$

where

$$u_p(x) := (u_{p,1}(x), u_{p,2}(x), ..., u_{p,N}(x))^T.$$
(1.14)

Obviously, we arrive at the perturbed system

$$\left[\frac{1}{2}\ln\left(-\frac{d^2}{dx^2}\right)\right]u_{p,m} - b_m \frac{du_{p,m}}{dx} - a_m u_{p,m} =$$

$$\varepsilon_m \int_{-\infty}^{\infty} \mathcal{K}_m(x - y) g_m(u_0(y) + u_p(y)) dy \tag{1.15}$$

with $1 \le m \le N$. For technical purpose, we introduce a closed ball in our function space, namely,

$$B_{\rho} := \{ u(x) \in L^{2}(\mathbb{R}, \mathbb{R}^{N}) \mid ||u||_{L^{2}(\mathbb{R}, \mathbb{R}^{N})} \le \rho \}, \quad 0 < \rho \le 1.$$
 (1.16)

Let us look for solutions of the system of equations (1.15) as fixed points of the auxiliary nonlinear problem,

$$\left[\frac{1}{2}\ln\left(-\frac{d^2}{dx^2}\right)\right]u_m - b_m \frac{du_m}{dx} - a_m u_m =$$

$$\varepsilon_m \int_{-\infty}^{\infty} \mathcal{K}_m(x-y)g_m(u_0(y) + v(y))dy, \tag{1.17}$$

where $1 \le m \le N$, in the ball B_{ρ} given by (1.16). The fixed point technique was used in [9] to demonstrate the persistence of pulses for a class of reaction-diffusion equations. For a given vector function v(y), system (1.17) is to be solved for u(x).

The left side of (1.17) involves the Fredholm operators given by (1.5). The solvability of the linear and nonhomogeneous system containing these operators will be considered in the final section of this article. Let us point out that, as distinct from the present case, the equations discussed in [17, 31] involved operators without the Fredholm property and orthogonality relations were used to solve them.

For our nonlinear vector function

$$g(z) := (g_1(z), g_2(z), ..., g_N(z)),$$
 (1.18)

its gradient is given by

$$\nabla g(z) := (\nabla g_1(z), \nabla g_2(z), ..., \nabla g_N(z)),$$
 (1.19)

such that the norm

$$\|\nabla g(z)\|_{L^{\infty}(\mathbb{R}^{N},\mathbb{R}^{N^{2}})} := \sum_{m=1}^{N} \|\nabla g_{m}(z)\|_{L^{\infty}(\mathbb{R}^{N})}.$$
 (1.20)

Let us use the closed ball in the function space,

$$D_M := \{ \nabla g(z) \in L^{\infty}(\mathbb{R}^N, \mathbb{R}^{N^2}) \mid \|\nabla g(z)\|_{L^{\infty}(\mathbb{R}^N, \mathbb{R}^{N^2})} \le M \}, \quad M > 0 \quad (1.21)$$

and impose the following conditions on the nonlinear part of the system of equations (1.1).

Assumption 2. Let $1 \leq m \leq N$, $g_m(z) : \mathbb{R}^N \to \mathbb{R}$, so that $g_m(0) = 0$. Let in addition $\nabla g(z) \in D_M$ and g(z) is nontrivial on \mathbb{R}^N .

Note that the solvability of a local elliptic problem in a bounded domain in \mathbb{R}^N was studied in [8], where the nonlinear function was allowed to have a sublinear growth.

We introduce the operator T_g by $u=T_gv$, where u is a solution of system (1.17). It will be shown that, for $v \in B_\rho$, problem (1.17) has only one solution $u \in B_\rho$. Our first main result is as follows.

Theorem 3. Let Assumptions 1 and 2 hold and

$$0 < \varepsilon \le \frac{\rho C}{M \mathcal{K}(\|u_0\|_{L^2(\mathbb{R},\mathbb{R}^N)} + 1)}. \tag{1.22}$$

Then the map $T_g: B_\rho \to B_\rho$ associated with the system of equations (1.17) is a strict contraction. The unique fixed point $u_p(x)$ of this map T_g is the only solution of system (1.15) in B_ρ .

Evidently, the resulting solution of system (1.1) given by (1.13) will not vanish identically on \mathbb{R} since the source terms $f_m(x)$ are nontrivial for some $1 \leq m \leq N$ and all $g_m(0) = 0$ as we assume.

Our second main statement deals with the continuity of the cumulative solution of the system of equations (1.1) given by (1.13) with respect to the gradient of the nonlinear vector function q. Let us define the following positive technical quantity

$$\sigma := \frac{\varepsilon M \mathcal{K}}{C}.\tag{1.23}$$

Theorem 4. Suppose that the assumptions of Theorem 3 are valid. Let $u_{p,j}(x)$ be the unique fixed point of the map $T_{g_j}: B_\rho \to B_\rho$ for g_j , j=1, 2 and the cumulative solution of system (1.1) with $g(z)=g_j(z)$ be given by

$$u_j(x) := u_0(x) + u_{p,j}(x).$$
 (1.24)

Then

$$||u_{1}(x) - u_{2}(x)||_{L^{2}(\mathbb{R},\mathbb{R}^{N})} \leq \frac{\varepsilon}{1 - \sigma} \frac{\mathcal{K}}{C} (||u_{0}||_{L^{2}(\mathbb{R},\mathbb{R}^{N})} + 1) ||\nabla g_{1}(z) - \nabla g_{2}(z)||_{L^{\infty}(\mathbb{R}^{N},\mathbb{R}^{N^{2}})}.$$
(1.25)

We turn our attention first to the proof of Theorem 3.

2. The existence of the perturbed solution

Proof of Theorem 3. Let us choose an arbitrary $v(x) \in B_{\rho}$ and designate the terms involved in the integral expressions in the right side of the system of equations (1.17) as

$$G_m(x) := g_m(u_0(x) + v(x)), \quad 1 \le m \le N.$$
(2.1)

Throughout the article we will use the standard Fourier transform

$$\widehat{\phi}(p) := \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \phi(x) e^{-ipx} dx. \tag{2.2}$$

Clearly, the upper bound

$$\|\widehat{\phi}(p)\|_{L^{\infty}(\mathbb{R})} \le \frac{1}{\sqrt{2\pi}} \|\phi(x)\|_{L^{1}(\mathbb{R})}$$
 (2.3)

holds. We apply (2.2) to both sides of system (1.17). This yields

$$\widehat{u_m}(p) = \varepsilon_m \sqrt{2\pi} \frac{\widehat{\mathcal{K}}_m(p)\widehat{G}_m(p)}{\ln\left(\frac{|p|}{e^{a_m}}\right) - ib_m p}, \quad 1 \le m \le N.$$
(2.4)

By virtue of estimates (1.7) and (2.3), we easily obtain

$$|\widehat{u_m}(p)| \le \varepsilon_m \frac{\|\mathcal{K}_m\|_{L^1(\mathbb{R})}|\widehat{G_m}(p)|}{C}, \quad 1 \le m \le N,$$
(2.5)

such that

$$||u_m||_{L^2(\mathbb{R})} \le \varepsilon_m \frac{||\mathcal{K}_m||_{L^1(\mathbb{R})}}{C} ||G_m||_{L^2(\mathbb{R})}, \quad 1 \le m \le N.$$
 (2.6)

Obviously,

$$G_m(x) = \int_0^1 \nabla g_m(t(u_0(x) + v(x))).(u_0(x) + v(x))dt, \quad 1 \le m \le N, \quad (2.7)$$

where the dot stands for the scalar product of two vectors in \mathbb{R}^N . Hence

$$|G_m(x)| < M|u_0(x) + v(x)|_{\mathbb{R}^N}, \quad 1 < m < N, \tag{2.8}$$

where $|.|_{\mathbb{R}^N}$ denotes the length of a vector in \mathbb{R}^N . Therefore, for $v(x) \in B_\rho$, we derive

$$||G_m(x)||_{L^2(\mathbb{R})} < M(||u_0||_{L^2(\mathbb{R}\mathbb{R}^N)} + 1), \quad 1 < m < N.$$
 (2.9)

By virtue of bounds (2.6) and (2.9), we arrive at

$$||u||_{L^{2}(\mathbb{R},\mathbb{R}^{N})} \leq \frac{\varepsilon M \mathcal{K}}{C}(||u_{0}||_{L^{2}(\mathbb{R},\mathbb{R}^{N})} + 1) \leq \rho$$
(2.10)

for all the values of the parameter ε , which satisfy (1.22). This implies that $u(x) \in B_{\rho}$ as well.

Let us suppose that for some $v(x) \in B_{\rho}$ there exist two solutions $u_{1,2}(x) \in B_{\rho}$ of the system of equations (1.17). Then the difference vector function $w(x) := u_1(x) - u_2(x) \in L^2(\mathbb{R}, \mathbb{R}^N)$ solves the homogeneous system (4.2). Thus w(x) vanishes a.e. on the real line as discussed with similar arguments for Lemma 5 further down.

This means that the system of equations (1.17) defines a map $T_g: B_\rho \to B_\rho$ for all the values of ε satisfying (1.22). Our goal is to establish that such map is a strict contraction.

We choose arbitrary $v_{1,2}(x) \in B_{\rho}$. By means of the argument above $u_{1,2} := T_g v_{1,2} \in B_{\rho}$ as well when ε satisfies (1.22). By virtue of (1.17), for $1 \le m \le N$,

$$\left[\frac{1}{2}\ln\left(-\frac{d^2}{dx^2}\right)\right]u_{1,m} - b_m \frac{du_{1,m}}{dx} - a_m u_{1,m} =$$

$$\varepsilon_m \int_{-\infty}^{\infty} \mathcal{K}_m(x - y)g_m(u_0(y) + v_1(y))dy, \qquad (2.11)$$

$$\left[\frac{1}{2}\ln\left(-\frac{d^2}{dx^2}\right)\right]u_{2,m} - b_m \frac{du_{2,m}}{dx} - a_m u_{2,m} =$$

$$\varepsilon_m \int_{-\infty}^{\infty} \mathcal{K}_m(x - y)g_m(u_0(y) + v_2(y))dy. \qquad (2.12)$$

Let us define

$$G_{1,m}(x) := g_m(u_0(x) + v_1(x)), \quad G_{2,m}(x) := g_m(u_0(x) + v_2(x))$$
 (2.13)

with $1 \le m \le N$. We apply the standard Fourier transform (2.2) to both sides of systems (2.11) and (2.12) to obtain

$$\widehat{u_{1,m}}(p) = \varepsilon_m \sqrt{2\pi} \frac{\widehat{\mathcal{K}}_m(p)\widehat{G_{1,m}}(p)}{\ln\left(\frac{|p|}{e^{a_m}}\right) - ib_m p}, \quad 1 \le m \le N,$$
(2.14)

$$\widehat{u_{2,m}}(p) = \varepsilon_m \sqrt{2\pi} \frac{\widehat{\mathcal{K}}_m(p)\widehat{G}_{2,m}(p)}{\ln\left(\frac{|p|}{e^{am}}\right) - ib_m p}, \quad 1 \le m \le N.$$
 (2.15)

Let us recall estimates (1.7) and (2.3). This yields

$$|\widehat{u_{1,m}}(p) - \widehat{u_{2,m}}(p)| \le \varepsilon_m \|\mathcal{K}_m\|_{L^1(\mathbb{R})} \frac{|\widehat{G_{1,m}}(p) - \widehat{G_{2,m}}(p)|}{C}, \quad 1 \le m \le N.$$
 (2.16)

Hence

$$||u_{1,m} - u_{2,m}||_{L^2(\mathbb{R})} \le \varepsilon_m \frac{||\mathcal{K}_m||_{L^1(\mathbb{R})}}{C} ||G_{1,m} - G_{2,m}||_{L^2(\mathbb{R})}, \quad 1 \le m \le N. \quad (2.17)$$

Clearly, for $1 \le m \le N$

$$G_{1,m}(x) - G_{2,m}(x) =$$

$$\int_0^1 \nabla g_m(u_0(x) + tv_1(x) + (1-t)v_2(x)).(v_1(x) - v_2(x))dt. \tag{2.18}$$

This allows us to estimate the norm from above as

$$||G_{1,m} - G_{2,m}||_{L^2(\mathbb{R})} \le M||v_1 - v_2||_{L^2(\mathbb{R},\mathbb{R}^N)}, \quad 1 \le m \le N.$$
 (2.19)

By means of bounds (2.17) and (2.19) along with definitions (1.8), (1.9), (1.11) and (1.23), we arrive at

$$||T_q v_1 - T_q v_2||_{L^2(\mathbb{R},\mathbb{R}^N)} \le \sigma ||v_1 - v_2||_{L^2(\mathbb{R},\mathbb{R}^N)}. \tag{2.20}$$

By virtue of inequality (1.22) along with Remark 6 further down, we derive

$$\sigma < 1. \tag{2.21}$$

Therefore, the map $T_g: B_\rho \to B_\rho$ defined by the system of equations (1.17) is a strict contraction for all the values of ε , which satisfy (1.22). Its unique fixed point $u_p(x)$ is the only solution of system (1.15) in the ball B_ρ . The cumulative $u(x) \in L^2(\mathbb{R}, \mathbb{R}^N)$ given by formula (1.13) is a solution of problem (1.1). Obviously, formula (2.10) implies that $||u_p(x)||_{L^2(\mathbb{R}, \mathbb{R}^N)} \to 0$ as $\varepsilon \to 0$.

Let us proceed to demonstrating the validity of the second main statement of the article.

3. The continuity of the cumulative solution

Proof of Theorem 4. Evidently, for all the values of the parameter ε satisfying inequality (1.22), we have

$$u_{p,1} = T_{q_1} u_{p,1}, \quad u_{p,2} = T_{q_2} u_{p,2}.$$
 (3.1)

Since

$$u_{p,1} - u_{p,2} = T_{g_1} u_{p,1} - T_{g_1} u_{p,2} + T_{g_1} u_{p,2} - T_{g_2} u_{p,2},$$
(3.2)

we obtain

$$||u_{p,1} - u_{p,2}||_{L^2(\mathbb{R},\mathbb{R}^N)} \le$$

$$||T_{g_1}u_{p,1} - T_{g_1}u_{p,2}||_{L^2(\mathbb{R},\mathbb{R}^N)} + ||T_{g_1}u_{p,2} - T_{g_2}u_{p,2}||_{L^2(\mathbb{R},\mathbb{R}^N)}.$$
(3.3)

By virtue of estimate (2.20),

$$||T_{g_1}u_{p,1} - T_{g_1}u_{p,2}||_{L^2(\mathbb{R},\mathbb{R}^N)} \le \sigma ||u_{p,1} - u_{p,2}||_{L^2(\mathbb{R},\mathbb{R}^N)}$$
(3.4)

and bound (2.21) is valid. Therefore,

$$(1-\sigma)\|u_{p,1}-u_{p,2}\|_{L^2(\mathbb{R},\mathbb{R}^N)} \le \|T_{g_1}u_{p,2}-T_{g_2}u_{p,2}\|_{L^2(\mathbb{R},\mathbb{R}^N)}. \tag{3.5}$$

Let $\eta(x) := T_{g_1} u_{p,2}$. Obviously, for $1 \le m \le N$,

$$\left[\frac{1}{2}\ln\left(-\frac{d^2}{dx^2}\right)\right]\eta_m - b_m \frac{d\eta_m}{dx} - a_m \eta_m =$$

$$\varepsilon_m \int_{-\infty}^{\infty} \mathcal{K}_m(x-y) g_{1,m}(u_0(y) + u_{p,2}(y)) dy, \tag{3.6}$$

$$\left[\frac{1}{2}\ln\left(-\frac{d^2}{dx^2}\right)\right]u_{p,2,m} - b_m \frac{du_{p,2,m}}{dx} - a_m u_{p,2,m} = \varepsilon_m \int_{-\infty}^{\infty} \mathcal{K}_m(x-y)g_{2,m}(u_0(y) + u_{p,2}(y))dy.$$

$$(3.7)$$

Define

$$G_{1,2,m}(x) := g_{1,m}(u_0(x) + u_{p,2}(x)), \quad G_{2,2,m}(x) := g_{2,m}(u_0(x) + u_{p,2}(x))$$
 (3.8)

with $1 \le m \le N$. Let us apply the standard Fourier transform (2.2) to both sides of systems (3.6) and (3.7). This yields

$$\widehat{\eta_m}(p) = \varepsilon_m \sqrt{2\pi} \frac{\widehat{\mathcal{K}}_m(p) \widehat{G_{1,2,m}}(p)}{\ln\left(\frac{|p|}{e^{a_m}}\right) - ib_m p}, \quad 1 \le m \le N,$$
(3.9)

$$\widehat{u_{p,2,m}}(p) = \varepsilon_m \sqrt{2\pi} \frac{\widehat{\mathcal{K}}_m(p) \widehat{G_{2,2,m}}(p)}{\ln\left(\frac{|p|}{e^{a_m}}\right) - ib_m p}, \quad 1 \le m \le N.$$
 (3.10)

Using (1.7) and (2.3), we derive

$$|\widehat{\eta_m}(p) - \widehat{u_{p,2,m}}(p)| \le \varepsilon_m \frac{\|\mathcal{K}_m\|_{L^1(\mathbb{R})}|\widehat{G_{1,2,m}}(p) - \widehat{G_{2,2,m}}(p)|}{C}, \tag{3.11}$$

such that

$$\|\eta_m - u_{p,2,m}\|_{L^2(\mathbb{R})} \le \varepsilon_m \frac{\|\mathcal{K}_m\|_{L^1(\mathbb{R})} \|G_{1,2,m} - G_{2,2,m}\|_{L^2(\mathbb{R})}}{C},$$
(3.12)

where $1 \le m \le N$. Note that

$$G_{1,2,m}(x) - G_{2,2,m}(x) =$$

$$\int_{0}^{1} \nabla [g_{1,m} - g_{2,m}](t(u_0(x) + u_{p,2}(x))).(u_0(x) + u_{p,2}(x))dt, \quad 1 \le m \le N. \quad (3.13)$$

This enables us to obtain the estimate from above on the norm as

$$||G_{1,2,m} - G_{2,2,m}||_{L^2(\mathbb{R})} \le ||\nabla g_{1,m}(z) - \nabla g_{2,m}(z)||_{L^{\infty}(\mathbb{R}^N)} (||u_0||_{L^2(\mathbb{R},\mathbb{R}^N)} + 1).$$
 (3.14)

Let us use inequalities (3.5), (3.12) and (3.14) along with definitions (1.8) and (1.20). Thus, $||u_{p,1} - u_{p,2}||_{L^2(\mathbb{R},\mathbb{R}^N)}$ can be bounded from above by

$$\frac{\varepsilon}{1-\sigma} \frac{\mathcal{K}}{C} (\|u_0\|_{L^2(\mathbb{R},\mathbb{R}^N)} + 1) \|\nabla g_1(z) - \nabla g_2(z)\|_{L^\infty(\mathbb{R}^N,\mathbb{R}^{N^2})}. \tag{3.15}$$

Therefore, (1.25) is an easy consequence of formulas (1.24) and (3.15).

4. Auxiliary results

Let us establish the solvability of the linear system of equations involving the logarithmic Laplacian, the transport terms, and square-integrable right sides

$$\left[\frac{1}{2}\ln\left(-\frac{d^2}{dx^2}\right)\right]u_m - b_m\frac{du_m}{dx} - a_m u_m = f_m(x), \quad x \in \mathbb{R},\tag{4.1}$$

where $1 \leq m \leq N$ and $a_m, b_m \in \mathbb{R}$, $b_m \neq 0$ are constants. The technical lemma below is the adaptation of the one used in [10] for the studies of the single integrodifferential equation involving the logarithmic Laplacian and the drift term, analogous to system (1.1). We provide it for the convenience of the readers.

Lemma 5. Let $1 \leq m \leq N$, the functions $f_m(x) : \mathbb{R} \to \mathbb{R}$, $f_m(x) \in L^2(\mathbb{R})$ and they are nontrivial for certain $1 \leq m \leq N$. Then the system of equations (4.1) admits a unique solution $u_0(x) \in L^2(\mathbb{R}, \mathbb{R}^N)$.

Proof. First we demonstrate the uniqueness of solutions for system (4.1). Let us suppose that it possesses two solutions $u_1(x)$, $u_2(x) \in L^2(\mathbb{R}, \mathbb{R}^N)$. Evidently, the vector function $w(x) := u_1(x) - u_2(x) \in L^2(\mathbb{R}, \mathbb{R}^N)$ satisfies the homogeneous system of equations

$$\left[\frac{1}{2}\ln\left(-\frac{d^2}{dx^2}\right)\right]w_m - b_m \frac{dw_m}{dx} - a_m w_m = 0, \quad 1 \le m \le N.$$
 (4.2)

Because each operator L_{a_m,b_m} on $L^2(\mathbb{R})$ introduced in (1.5) has only the essential spectrum and no nontrivial zero modes (see (1.6) and (1.7)), the difference of solutions w(x) is trivial on \mathbb{R} .

Let us apply the standard Fourier transform (2.2) to both sides of system (4.1). We arrive at

$$\widehat{u_m}(p) = \frac{\widehat{f_m}(p)}{\ln\left(\frac{|p|}{e^{am}}\right) - ib_m p}, \quad p \in \mathbb{R}, \quad 1 \le m \le N.$$
(4.3)

By means of (1.7), we have

$$|\widehat{u_m}(p)| \le \frac{|\widehat{f_m}(p)|}{C} \in L^2(\mathbb{R}), \quad 1 \le m \le N$$
(4.4)

due to our assumption. Therefore, for the solution of our system of equations (4.1), we have $u(x) \in L^2(\mathbb{R}, \mathbb{R}^N)$. Its Fourier image is given by formula (4.3).

Remark 6. Note that under the assumptions of Lemma 5 above, the unique solution $u_0(x)$ of system (4.1) belonging to $L^2(\mathbb{R}, \mathbb{R}^N)$ is nontrivial because $f_m(x)$ do not vanish identically on the real line for some $1 \le m \le N$.

Acknowledgements

Y.C. was supported by NSERC of Canada (Nos. RGPIN-2019-05892, RGPIN-2024-05593).

Declarations

Ethical approval The results of the work are applicable for both human and animal studies.

Publishing policy The authors have read and understood the publishing policy, and submit this manuscript in accordance with this policy.

Competing interests policy The authors declare that they have no competing interests as defined by Springer, or other interests that might be perceived to influence the results and/or discussion reported in this paper.

Dual publication The results in this manuscript have not been published elsewhere, nor are they under consideration by another publisher.

Authorship The corresponding author has read the Springer journal policies on author responsibilities and submits this manuscript in accordance with those policies.

Author contributions statement Yuming Chen and Vitali Vougalter wrote the manuscript. All authors reviewed the manuscript.

Funding The work was partially supported by the NSERC of Canada (Nos. RGPIN-2019-05892, RGPIN-2024-05593).

Third party material All of the material is owned by the authors and no permissions are required.

Data availability statement All data generated or analyzed during this study are included in this published article.

References

[1] N. Apreutesei, N. Bessonov, V. Volpert, V. Vougalter, *Spatial structures and generalized travelling waves for an integro-differential equation*, Discrete Contin. Dyn. Syst. Ser. B, **13** (2010), no. 3, 537–557.

- [2] C. Amrouche, V. Girault, J. Giroire, *Dirichlet and Neumann exterior problems* for the n-dimensional Laplace operator: an approach in weighted Sobolev spaces, J. Math. Pures Appl. (9), **76** (1997), no. 1, 55–81.
- [3] C. Amrouche, F. Bonzom, *Mixed exterior Laplace's problem*, J. Math. Anal. Appl., **338** (2008), no. 1, 124–140.
- [4] H. Berestycki, G. Nadin, B. Perthame, L. Ryzhik, *The non-local Fisher-KPP equation: travelling waves and steady states*, Nonlinearity, **22** (2009), no. 12, 2813–2844.
- [5] P. Bolley, T.L. Pham, Propriétés d'indice en théorie hölderienne pour des opérateurs différentiels elliptiques dans \mathbb{R}^n , J. Math. Pures Appl. (9), **72** (1993), no. 1, 105–119.
- [6] P. Bolley, T.L. Pham, *Propriété d'indice en théorie Hölderienne pour le problème extérieur de Dirichlet*, Comm. Partial Differential Equations, **26** (2001), no. 1–2, 315–334.
- [7] N. Benkirane, Propriétés d'indice en théorie höldérienne pour des opérateurs elliptiques dans \mathbb{R}^n , C. R. Acad. Sci. Paris Sér. I Math., **307** (1988), no. 11, 577–580.
- [8] H. Brezis, L. Oswald, *Remarks on sublinear elliptic equations*, Nonlinear Anal., **10** (1986), no. 1, 55–64.
- [9] Y. Chen, V. Vougalter, *Persistence of pulses for some reaction-diffusion equations*, Pure Appl. Funct. Anal., **6** (2021), no. 2, 309–315.
- [10] Y. Chen, V. Vougalter, Existence of solutions for a class of integro-differential equations with the logarithmic Laplacian and transport, Preprint (2024), arXiv: 2404.09019.
- [11] H. Chen, T. Weth, *The Dirichlet problem for the logarithmic Laplacian*, Comm. Partial Differential Equations, **44** (2019), no. 11, 1100–1139.
- [12] L. De Luca, M. Novaga, M. Ponsiglione, *The 0-fractional perimeter between fractional perimeters and Riesz potentials*, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5), **22** (2021), no. 4, 1559–1596.
- [13] A. Ducrot, M. Marion, V. Volpert, Systemes de réaction-diffusion sans propriété de Fredholm, C. R. Math. Acad. Sci. Paris, **340** (2005), no. 9, 659–664.
- [14] A. Ducrot, M. Marion, V. Volpert, *Reaction-diffusion problems with non-Fredholm operators*, Adv. Differential Equations, **13** (2008), no. 11–12, 1151–1192.

- [15] M. Efendiev, Fredholm structures, topological invariants and applications. AIMS Ser. Differ. Equ. Dyn. Syst., 3. American Institute of Mathematical Sciences (AIMS), Springfield, MO (2009), 205 pp.
- [16] M. Efendiev, V. Vougalter, Solvability in the sense of sequences for some fourth order non-Fredholm operators, J. Differential Equations, **271** (2021), 280–300.
- [17] M. Efendiev, V. Vougalter, On the solvability of some systems of integrodifferential equations with drift. Nonlinear Syst. Complex., 33, Springer, Cham (2022), 141–167.
- [18] M. Efendiev, V. Vougalter, *Solvability in the sense of sequences for some non-Fredholm operators with the logarithmic Laplacian*, Monatsh. Math., **202** (2023), no. 4, 751–771.
- [19] M. Efendiev, V. Vougalter, Solvability in the sense of sequences for some logarithmic Shrödinger operators in higher dimensions, J. Pseudo-Differ. Oper. Appl., **14** (2023), no. 2, Paper No. 32, 19 pp.
- [20] P.A. Feulefack, *The logarithmic Schrödinger operator and associated Dirichlet problems*, J. Math. Anal. Appl., **517** (2023), no. 2, Paper No. 126656, 33 pp.
- [21] S. Genieys, V. Volpert, P. Auger, *Pattern and waves for a model in population dynamics with nonlocal consumption of resources*, Math. Model. Nat. Phenom., **1** (2006), no. 1, 65–82.
- [22] H.G. Gebran, C.A. Stuart, Fredholm and properness properties of quasilinear elliptic systems of second order, Proc. Edinb. Math. Soc. (2), 48 (2005), no. 1, 91–124.
- [23] H.G. Gebran, C.A. Stuart, Exponential decay and Fredholm properties in second-order quasilinear elliptic systems, J. Differential Equations, **249** (2010), no. 1, 94–117.
- [24] S. Jarohs, A. Saldana, T. Weth, *A new look at the fractional Poisson problem via the logarithmic Laplacian*, J. Funct. Anal., **279** (2020), no. 11, 108732, 50 pp.
- [25] A. Laptev, T. Weth, *Spectral properties of the logarithmic Laplacian*, Anal. Math. Phys., **11** (2021), no. 3, Paper No. 133, 24 pp.
- [26] R. Metzler, J. Klafter, *The random walk's guide to anomalous diffusion: a fractional dynamics approach*, Phys. Rep., **339** (2000), no. 1, 77 pp.

- [27] P.J. Rabier, C.A. Stuart, Fredholm and properness properties of quasilinear elliptic operators on \mathbb{R}^N , Math. Nachr., **231** (2001), 129–168.
- [28] V. Volpert, Elliptic partial differential equations. Volume 1: Fredholm theory of elliptic problems in unbounded domains. Monogr. Math., 101. Birkhäuser/Springer Basel AG, Basel (2011), 639 pp.
- [29] V. Volpert, V. Vougalter, *Emergence and propagation of patterns in nonlocal reaction-diffusion equations arising in the theory of speciation*. Lecture Notes in Math., **2071**, Springer, Heidelberg (2013), 331–353.
- [30] V. Vougalter, V. Volpert, *Solvability conditions for some non-Fredholm operators*, Proc. Edinb. Math. Soc. (2), **54** (2011), no. 1, 249–271.
- [31] V. Vougalter, V. Volpert, *Solvability conditions for some linear and nonlinear non-Fredholm elliptic problems*, Anal. Math. Phys., **2** (2012), no. 4, 473–496.
- [32] V. Vougalter, V. Volpert, On the solvability in the sense of sequences for some non-Fredholm operators related to the anomalous diffusion. Trends Math., Birkhäuser/Springer, Cham (2019), 229–257.
- [33] V. Vougalter, V. Volpert, *Solvability of some integro-differential equations with anomalous diffusion and transport*, Anal. Math. Phys., **11** (2021), no. 3, Paper No. 135, 26 pp.
- [34] R. Zhang, V. Kumar, M. Ruzhansky, Symmetry of positive solutions for Lane-Emden systems involving the logarithmic Laplacian, Acta Appl. Math., 188 (2023), Paper No. 16, 16 pp.