

BRAIDED LOGARITHMIC VERTEX ALGEBRAS

BOJKO BAKALOV AND JUAN J. VILLARREAL

ABSTRACT. We study a family of algebras defined using a locally-finite endomorphism called a braiding map. When the braiding map is semi-simple, the algebra is a generalized vertex algebra, while when the braiding map is locally-nilpotent we have a logarithmic vertex algebra. We describe a method that associates to these algebras non-local Poisson vertex algebras, and we use this relation to build a new example of a generalized vertex algebra motivated by the non-linear Schrödinger non-local Poisson vertex algebra.

1. INTRODUCTION

The following is the main definition of this work (cf. [BV1]).

Definition 1.1. Let V be a vector space. A *braiding map* \mathcal{S} on V is a linear operator

$$(1.1) \quad \mathcal{S} = \sum_{i=1}^L \phi_i \otimes \psi_i, \quad \phi_i, \psi_i \in \text{End}(V),$$

satisfying the following two conditions:

- (1) \mathcal{S} is *symmetric*, i.e., $\mathcal{S} = \sum_{i=1}^L \phi_i \otimes \psi_i = \sum_{i=1}^L \psi_i \otimes \phi_i$;
- (2) $[\phi_i, \psi_j] = [\phi_i, \phi_j] = [\psi_i, \psi_j] = 0$, $1 \leq i, j \leq L$.

Recall that a linear operator φ is called *locally finite* if every vector is contained in some finite-dimensional φ -invariant subspace. A locally finite operator φ has a unique Jordan–Chevalley decomposition $\varphi = \varphi^{(s)} + \varphi^{(n)}$, where $\varphi^{(s)}$ and $\varphi^{(n)}$ commute, $\varphi^{(s)}$ is semi-simple and $\varphi^{(n)}$ is locally nilpotent (for every vector u we have $(\varphi^{(n)})^k u = 0$ for some $k \geq 1$).

Now, we introduce the algebras that we study in this paper, generalizing those of [BV1].

Definition 1.2. A *braided logarithmic vertex algebra* (braided logVA) is a complex vector space V equipped with a vector $\mathbf{1} \in V$, a braiding map $\mathcal{S} = \sum_{i=1}^L \phi_i \otimes \psi_i$ where ϕ_i, ψ_i are locally finite on V , and a family of bilinear products

$$\mu_{(n)}: V \otimes V \rightarrow V, \quad a \otimes b \mapsto a_{(n+\mathcal{S})} b \quad (n \in \mathbb{Z}),$$

satisfying the following identities:

- (i) (*truncation property*) $a_{(n+\mathcal{S})} b = 0$ for $n \gg 0$.
- (ii) (*vacuum axioms*) $\mathcal{S}(\mathbf{1} \otimes a) = 0$, $\mathbf{1}_{(n+\mathcal{S})} a = \delta_{n,-1} a$, $a_{(-1+\mathcal{S})} \mathbf{1} = a$.
- (iii) (*hexagon identity*)

$$(1.2) \quad \mathcal{S}(\mu_{(n)} \otimes I) = (\mu_{(n)} \otimes I)(\mathcal{S}_{13} + \mathcal{S}_{23}), \quad n \in \mathbb{Z}.$$

Date: June 12, 2024.

2010 Mathematics Subject Classification. Primary 17B69; Secondary 17B63, 81R10, 81T40.

(iv) (*Borcherds identity*)

$$\begin{aligned}
(1.3) \quad & \sum_{j \in \mathbb{Z}_+} (-1)^j \mu_{(m+n-j)}(I \otimes \mu_{(k+j)}) \binom{n + \mathcal{S}_{12}}{j} \\
& - \sum_{j \in \mathbb{Z}_+} (-1)^{n+j} \mu_{(n+k-j)}(I \otimes \mu_{(m+j)}) (-1)^{\mathcal{S}_{12}^{(s)}} \binom{n + \mathcal{S}_{12}}{j} P_{12} \\
& = \sum_{j \in \mathbb{Z}_+} \mu_{(m+k-j)}(\mu_{(n+j)} \otimes I) \binom{m + \mathcal{S}_{13}}{j},
\end{aligned}$$

for all $k, m, n \in \mathbb{Z}$, where P is the transposition on $V \otimes V$ and $\mathcal{S}^{(s)}$ is the semi-simple term of the Jordan–Chevalley decomposition of \mathcal{S} (cf. (2.1)).

The name braided vertex algebras appeared in the literature in quantum vertex algebras; see [EK] (see also [BV1, Remark 3.19]). Note that a braided logVA with $\mathcal{S} = 0$ is the same as a vertex algebra [Bo]. More generally, we have:

Theorem 1.3. *Let V be a braided logVA with braiding \mathcal{S} . Then:*

- (i) *If $\mathcal{S} = \mathcal{S}^{(n)}$, then V is a logarithmic vertex algebra as in [BV1].*
- (ii) *If $\mathcal{S} = \mathcal{S}^{(s)}$, then V is a generalized vertex algebra as in [FFR, DL, BK].*

In [FB], Poisson vertex algebras were associated to vertex algebras considering (algebraic) limits. We have the following generalization of this result.

Theorem 1.4. *Let V^ϵ be a logarithmic vertex algebra over $\mathbb{C}[\epsilon]/(\epsilon^2)$ with a state-field map Y^ϵ such that $V^0 = V^\epsilon/\epsilon V_\epsilon$ is a commutative vertex algebra. Then V^0 naturally acquires the structure of a non-local Poisson vertex algebra.*

Recall that a vertex algebra is called *commutative* if all products $\mu_{(n)} = 0$ for $n \geq 0$; then the product $\mu_{(-1)}$ is commutative and associative [Bo]. We review the definition of a non-local Poisson vertex algebra (non-local PVA) in Definition 4.2 below; see [K, DK].

In this work, we describe a new example of a generalized vertex algebra in Theorem 4.6, motivated by the non-linear Schrödinger non-local PVA from [DK]; see Definition 4.4. This generalized vertex algebra is built using an associative algebra with generators $\{w_n, \bar{w}_n\}_{n \in \mathbb{Z}}$ subject to the following relations ($m, k \in \mathbb{Z}$):

$$\begin{aligned}
(1.4) \quad & \sum_{j \geq 0} w_{m-j} w_{k+j} + w_{k-j} w_{m+j} = 0, \\
& \sum_{j \geq 0} \bar{w}_{m-j} \bar{w}_{k+j} + \bar{w}_{k-j} \bar{w}_{m+j} = 0, \\
& w_m \bar{w}_k - w_{m-1} \bar{w}_{k+1} + \bar{w}_k w_m - \bar{w}_{k-1} w_{m+1} = 0.
\end{aligned}$$

In [BV2], we described, in particular, logarithmic vertex algebras quantizing the potential free boson non-local PVA and the potential Virasoro-Magri non-local PVA. For the non-linear Schrödinger non-local PVA the same methods as in [BV2] cannot be used. In this work, the algebraic limits in Theorem 1.4 are used to explain the relation between the generalized vertex algebra defined by (1.4) and the non-linear Schrödinger non-local PVA; see (4.9). We note that algebraic limits have been also used recently to quantize jet schemes of Poisson vertex algebras in terms of asymptotic algebras of chiral differential operators [AKM, AKM2].

This paper is organized as follows. In Section 2, we use the language of fields to describe braided logVAs; see Theorems 2.10 and 1.3. In Section 3, following [BV1], we prove an existence theorem for braided logVAs. In Section 4.1, we prove Theorem 1.4 following ideas of [BV2]. Finally, in Section 4.2, we build the generalized vertex algebra defined by relations (1.4) and we prove that its associated non-local PVA is the non-linear Schrödinger PVA.

2. LOGARITHMIC FIELDS

We denote by $V[[z]][\zeta, z^{-\mathbb{C}}]$ the space of infinite sums $\sum_n f_n(\zeta)z^n$ where $f_n(\zeta) \in V[\zeta]$ and n runs over the union of finitely many sets of the form $\{-d_i + \mathbb{Z}_{\geq 0}\}$ with $d_i \in \mathbb{C}$. The space $V[[z]][\zeta, z^{-\mathbb{C}}]$ is equipped with the usual action of the derivatives ∂_z and ∂_ζ , and the operator

$$D_z := \partial_z + z^{-1}\partial_\zeta,$$

which acts as the total derivative with respect to z if we set $\zeta = \log z$.

Let $\mathcal{S} = \sum_{i=1}^L \phi_i \otimes \psi_i$ be a braiding map where ϕ_i, ψ_i ($1 \leq i \leq L$) are locally finite on V with Jordan–Chevalley decompositions

$$\phi_i = \phi_i^{(s)} + \phi_i^{(n)}, \quad \psi_i = \psi_i^{(s)} + \psi_i^{(n)}.$$

Note that, since all the operators ϕ_i, ψ_j commute with each other, their semisimple and nilpotent part commute with each other too. This implies that \mathcal{S} is locally finite on $V \otimes V$ and

$$(2.1) \quad \mathcal{S}^{(s)} = \sum_{i=1}^L \phi_i^{(s)} \otimes \psi_i^{(s)}, \quad \mathcal{S}^{(n)} = \sum_{i=1}^L \phi_i^{(s)} \otimes \psi_i^{(n)} + \phi_i^{(n)} \otimes \psi_i^{(s)} + \phi_i^{(n)} \otimes \psi_i^{(n)}.$$

We use this decomposition of \mathcal{S} to define the following map on $V \otimes V$:

$$z^{\mathcal{S}} := z^{\mathcal{S}^{(s)}} e^{\zeta \mathcal{S}^{(n)}}.$$

The next lemma can be easily derived from (2.1) and Definition 1.1 (2).

Lemma 2.1. *For any $a, b \in V$, we have*

$$(2.2) \quad z^{\mathcal{S}}(a \otimes b) = z^{\mathcal{S}^{(s)}} e^{\zeta \mathcal{S}^{(n)}}(a \otimes b) = \sum_{j=1}^m \sum_{i=1}^r \frac{1}{j!} z^{d_i} \zeta^j (\mathcal{S}^{(n)})^j (a_i \otimes b_i),$$

where $a \otimes b = \sum_{i=1}^r a_i \otimes b_i$ such that $a_i \in V$ (respectively, $b_i \in V$) are common eigenvectors of all $\phi_j^{(s)}$ (respectively, $\psi_j^{(s)}$), $1 \leq j \leq L$, and

- (i) $\mathcal{S}^{(s)}(a_i \otimes b_i) = d_i(a_i \otimes b_i)$ with $d_i \in \mathbb{C}$;
- (ii) $m \in \mathbb{Z}_{\geq 0}$ is such that $(\mathcal{S}^{(n)})^m(a_i \otimes b_i) = 0$ for all $1 \leq i \leq r$.

2.1. Quantum fields. Now we introduce the notion of a quantum field as in [B].

Definition 2.2. Let V be a vector space. The space of (logarithmic) fields is defined by

$$\text{Fie}(V) = \text{Hom}(V, V[[z]][\zeta, z^{-\mathbb{C}}]).$$

We denote the elements of $\text{Fie}(V)$ as $a(z, \zeta)$ or $a(z)$ for short.

On a braided logVA V , we define for $a, b \in V$ the series (cf. (2.2))

$$(2.3) \quad \begin{aligned} Y(a, z)b &= Y(z)(a \otimes b) := \sum_{n \in \mathbb{Z}} \mu_{(n)}(z^{-n-1-\mathcal{S}}(a \otimes b)) \\ &= \sum_{n \in \mathbb{Z}} \sum_{j=1}^m \sum_{i=1}^r \frac{(-1)^j}{j!} z^{-n-1-d_i} \zeta^j \mu_{(n)}((\mathcal{S}^{(n)})^j(a_i \otimes b_i)). \end{aligned}$$

Note that, by the linearity of $\mu_{(n)}$, we have $a_{(n+\mathcal{S})}b = \sum_{i=1}^r \mu_{(n)}(a_i \otimes b_i) = \sum_{i=1}^r a_{i(n+\mathcal{S})}b_i$. Now, by Definition 1.2 (i) and (2.3), the next proposition is obvious.

Proposition 2.3. *Let V be a braided logVA. Then $Y(a, z)$, defined by (2.3), is a logarithmic field on V for every $a \in V$.*

In a braided logVA V , we define the *translation operator* $T \in \text{End}(V)$ by

$$(2.4) \quad T(a) := a_{(-2+\mathcal{S})}\mathbf{1}, \quad a \in V.$$

Note that $T(\mathbf{1}) = 0$ by definition.

Proposition 2.4. *For all $a \in V$, we have the translation covariance property*

$$(2.5) \quad [T, Y(a, z)] = D_z Y(a, z).$$

Proof. The Borchers identity (1.3) for $m = 0, k = -2$ on $a \otimes b \otimes \mathbf{1}$ gives:

$$\mu_{(n)}(a \otimes Tb) - T\mu_{(n)}(a \otimes b) = -n\mu_{(n-1)}(a \otimes b) - \mu_{(n-1)}(\mathcal{S}(a \otimes b)).$$

Plugging this into (2.3), we obtain the translation covariance. \square

Proposition 2.5. *For any braided logVA V , the map Y defined in (2.3) satisfies $Y(\mathbf{1}, z) = I$, $Y(a, z)\mathbf{1} \in V[[z]]$, and $Y(a, z)\mathbf{1}|_{z=0} = a$.*

Proof. By the hexagon identity (Definition 1.2 (iii)), the components ϕ_i, ψ_i of \mathcal{S} are derivations of all products $\mu_{(n)}$. This implies, in particular, that $\phi_i(\mathbf{1}) = \psi_i(\mathbf{1}) = 0$, because $\mathbf{1}_{(-1+\mathcal{S})}\mathbf{1} = \mathbf{1}$ by the vacuum axiom (Definition 1.2 (ii)). Then the claims of the proposition follow easily from (2.3) and the vacuum axiom. \square

As in [BV1], we will use the notation $z_{ij} = z_i - z_j$, and we introduce the formal variables $\zeta_{ij} = \zeta_{ji}$, which are thought of as $\log|z_{ij}|$. Additionally, we consider the formal expansions for $n \in \mathbb{Z}$

$$(2.6) \quad \begin{aligned} \iota_{z_1, z_2} z_{12}^{n+\mathcal{S}} &= \iota_{z_1, z_2} z_{12}^{n+\mathcal{S}^{(s)}} e^{\zeta_{12}\mathcal{S}^{(n)}} := \sum_{j \in \mathbb{Z}_+} \binom{n+\mathcal{S}}{j} (-1)^j z_1^{n-j+\mathcal{S}^{(s)}} z_2^j, \\ \iota_{z_2, z_1} z_{12}^{n+\mathcal{S}} &= \iota_{z_2, z_1} z_{12}^{n+\mathcal{S}^{(s)}} e^{\zeta_{21}\mathcal{S}^{(n)}} := \sum_{j \in \mathbb{Z}_+} \binom{n+\mathcal{S}}{j} (-1)^{n+j+\mathcal{S}^{(s)}} z_1^j z_2^{n-j+\mathcal{S}}. \end{aligned}$$

In the following proposition, we will utilize the following generalization of delta function introduced in [B]:

$$(2.7) \quad \delta_{\mathcal{S}}(z_1, z_2) := \sum_{n \in \mathbb{Z}} z_1^{-n-1-\mathcal{S}} z_2^{n+\mathcal{S}}.$$

Note that we have

$$(2.8) \quad \frac{1}{j!} D_{z_2}^j \delta_{\mathcal{S}}(z_1, z_2) = \sum_{n \in \mathbb{Z}} \binom{n+\mathcal{S}}{j} z_1^{-n-1-\mathcal{S}} z_2^{n-j+\mathcal{S}}.$$

Proposition 2.6. *Let V be a braided logarithmic vertex algebra. For $n \in \mathbb{Z}$, we have the following Borchers identity*

$$(2.9) \quad \begin{aligned} & \iota_{z_1, z_2} Y(z_1)(I \otimes Y(z_2))z_{12}^{n+\mathcal{S}_{12}} - \iota_{z_2, z_1} Y(z_2)(I \otimes Y(z_1))z_{12}^{n+\mathcal{S}_{12}} \\ &= \sum_{j \in \mathbb{Z}_+} \frac{1}{j!} Y(z_2)(\mu_{(n+j)} \otimes I) D_{z_2}^j \delta_{\mathcal{S}_{13}}(z_1, z_2). \end{aligned}$$

Proof. Using the notation $X(z)(a \otimes b) := \sum_{n \in \mathbb{Z}} \mu_{(n)}(a \otimes b)z^{-n-1}$ we have that (1.3) is equivalent to

$$(2.10) \quad \begin{aligned} & X(z_1)(I \otimes X(z_2)) \sum_{j \in \mathbb{Z}_+} \binom{n+\mathcal{S}_{12}}{j} (-1)^j z_1^{n-j} z_2^j \\ & - X(z_2)(I \otimes X(z_1)) \sum_{j \in \mathbb{Z}_+} \binom{n+\mathcal{S}_{12}}{j} (-1)^{n+j+\mathcal{S}_{12}^{(s)}} z_1^j z_2^{n-j} P_{12} \\ &= \sum_{j \in \mathbb{Z}_+} X(z_2)(\mu_{(n+j)} \otimes I) \sum_{m \in \mathbb{Z}} \binom{m+\mathcal{S}_{13}}{j} z_1^{-m-1} z_2^{m-j}. \end{aligned}$$

We multiply the right-hand side of this identity by $z_1^{-\mathcal{S}_{13}} z_2^{-\mathcal{S}_{23}}$; then from (1.2), (2.6), (2.8), we obtain (2.9). \square

Before we state the locality of braided logVAs, we need the following simple lemma.

Lemma 2.7. *Let \mathcal{S} be as in (2.1). Then for any $a \in V$, there exist $m \in \mathbb{Z}_{\geq 0}$ and an expression $a = \sum_{j=1}^r a_j$ such that*

- (i) $\phi_i^{(s)} a_j = d_{i,j} a_j$ for some $d_{i,j} \in \mathbb{C}$ and all $1 \leq i \leq L$, $1 \leq j \leq r$;
- (ii) $(\phi_i^{(n)})^{m+1} a_j = 0$ for all $1 \leq i \leq L$, $1 \leq j \leq r$.

Furthermore, for any $b \in V$, we have

$$z^{\mathcal{S}}(a \otimes b) = \sum_{j=1}^r \sum_{k=0}^{mL} \sum_{1 \leq i_1, \dots, i_k \leq L} \frac{1}{k!} \zeta^k(\phi_{i_1}^{(n)} \dots \phi_{i_k}^{(n)}) a_j \otimes z^{\sum_{i=1}^L d_{i,j} \psi_i} (\psi_{i_1} \dots \psi_{i_k}) b.$$

Proof. First, note that all ϕ_i commute with each other; hence, all $\phi_i^{(s)}$ commute and can be diagonalized them simultaneously. Then the formula for $z^{\mathcal{S}}$ follows from the identities

$$\begin{aligned} z^{\phi_i^{(s)} \otimes \psi_i}(a \otimes b) &= \sum_{j=1}^r a_j \otimes z^{d_{i,j} \psi_i} b, \\ z^{\phi_i^{(n)} \otimes \psi_i}(a \otimes b) &= \sum_{k=0}^m \frac{\zeta^k}{k!} (\phi_i^{(n)})^k a \otimes (\psi_i)^k b. \end{aligned}$$

\square

Proposition 2.8. *For every $a, b \in V$, there exists $N \in \mathbb{Z}_{\geq 0}$ such that for all $c \in V$*

$$(2.11) \quad \begin{aligned} & Y(z_1)(I \otimes Y(z_2)) \iota_{z_1, z_2} z_{12}^{N+\mathcal{S}}(a \otimes b) \otimes c \\ &= Y(z_2)(I \otimes Y(z_1)) \iota_{z_2, z_1} z_{12}^{N+\mathcal{S}}(b \otimes a) \otimes c. \end{aligned}$$

Proof. First, we rewrite (2.7) as

$$\delta_{\mathcal{S}}(z_1, z_2) = \left(\sum_{n \in \mathbb{Z}} z_1^{-n-1} z_2^n \right) (z_1^{-\mathcal{S}} z_2^{\mathcal{S}}) = \delta(z_1, z_2) \left(\frac{z_2}{z_1} \right)^{\mathcal{S}},$$

where we use the usual delta function from [K]. Then from the Leibniz rule we have

$$D_{z_2}^j \delta_{\mathcal{S}_{13}}(z_1, z_2) = \sum_{k=0}^j \binom{j}{k} D_{z_2}^k (\delta(z_1, z_2)) D_{z_2}^{j-k} \left(\frac{z_2}{z_1} \right)^{\mathcal{S}_{13}}.$$

From Lemma 2.7, we have the finite sum

$$\begin{aligned} z_1^{-\mathcal{S}} z_2^{\mathcal{S}} (a \otimes c) &= \sum_{j=1}^r \sum_{l \geq 0} \sum_{1 \leq i_1, \dots, i_l \leq L} \frac{1}{l!} (\zeta_2 - \zeta_1)^l \\ &\times (\phi_{i_1}^n \cdots \phi_{i_l}^n)(a_j) \otimes z_1^{-\sum_{i=1}^L d'_{i,j} \psi_i} z_2^{\sum_{i=1}^L d'_{i,j} \psi_i} (\psi_{i_1} \cdots \psi_{i_l})(c). \end{aligned}$$

As we mention before, the finiteness condition of the sum is independent of c . Hence, there exists $M \in \mathbb{Z}_{\geq 0}$ such that if $r \geq M$ then for all $l \in \mathbb{Z}_{\geq 0}$ and $c \in V$

$$(\mu_{(r)} \otimes I)(D_{z_2}^l (z_1^{-\mathcal{S}_{13}} z_2^{\mathcal{S}_{13}})(a \otimes b \otimes c)) = 0.$$

Replacing $N = M$ in identity (2.9) gives us the proposition. \square

Remark 2.9. In [BV2, Sect. 4.3], we found an example where \mathcal{S} is locally finite but can not be expressed as a sum $\sum \phi_i \otimes \psi_i$ where each component ϕ_i, ψ_i is locally finite on V . In this case, the integer N necessary for the locality of the fields depends on a, b and c (see also [BV1, Remark 3.27]).

The next theorem gives an equivalent definition of a braided logVA.

Theorem 2.10. *A braided logarithmic vertex algebra is a vector space V , equipped with a vector $\mathbf{1} \in V$, an endomorphism $T \in \text{End}(V)$, a linear map*

$$Y: V \rightarrow \text{Fie}(V), \quad a \mapsto Y(a, z),$$

and a braiding map \mathcal{S} on V , which are subject to the following axioms:

- (i) $Y(\mathbf{1}, z) = I$, $Y(a, z)\mathbf{1} \in V[[z]]$, $Y(a, z)\mathbf{1}|_{z=0} = a$, $T\mathbf{1} = \mathcal{S}(\mathbf{1} \otimes a) = 0$.
- (ii) $[T, Y(a, z)] = \partial_z Y(a, z)$.
- (iii) $\mathcal{S} = \sum_{i=1}^L \phi_i \otimes \psi_i$ where ϕ_i, ψ_i are locally finite on V .
- (iv) $\mathcal{S}(Y(z) \otimes I) = (Y(z) \otimes I)(\mathcal{S}_{13} + \mathcal{S}_{23})$.
- (v) For every $a, b \in V$ there exists $N \in \mathbb{Z}_+$ such that $\forall c \in V$

$$\begin{aligned} (2.12) \quad & Y(z_1)(I \otimes Y(z_2)) \iota_{z_1, z_2}(z_{12}^{N+\mathcal{S}})(a \otimes b) \otimes c \\ &= Y(z_2)(I \otimes Y(z_1)) \iota_{z_2, z_1}(z_{12}^{N+\mathcal{S}})(b \otimes a) \otimes c. \end{aligned}$$

Proof. We have that braided logVAs satisfy the above conditions, by Propositions 2.3, 2.4, 2.5 and 2.8. The other direction follows from Proposition 3.8 in the next section. \square

Now, we prove Theorem 1.3.

Proof of Theorem 1.3. (i) Follows from Theorem 2.10, since this is the definition of a logarithmic vertex algebra in [BV1].

(ii) Following [BK], we define Q, Δ and η . Let $\{a_i\}_{i \in I}$ be a basis of V such that $\mathcal{S}(a_i \otimes a_j) = s_{ij} a_i \otimes a_j$ for some $s_{ij} \in \mathbb{C}$. First, we define the subsets

$$I_i := \{j \in I \mid s_{jl} - s_{il} \in \mathbb{Z} \ \forall l \in I\}.$$

Note that $j \in I_i$ if and only if $I_i = I_j$; hence $I = \bigsqcup I_i$. We define $Q := \{I_i\}$. Now, Q has the structure of an abelian group defined by $(r, t \in \mathbb{Z})$:

$$(2.13) \quad rI_i + tI_j := \{k \in I \mid s_{kl} - rs_{il} - ts_{jl} \in \mathbb{Z} \ \forall l \in I\}.$$

For $\alpha = I_k \in Q$, we define $V_\alpha = \text{span}\{a_j \mid j \in \alpha\}$. Then

$$V = \bigoplus_{\alpha \in Q} V_\alpha.$$

Second, we define $\Delta: Q \times Q \rightarrow \mathbb{C}/\mathbb{Z}$ by

$$(2.14) \quad \Delta(I_i, I_j) := s_{ij} \pmod{\mathbb{Z}}.$$

Note that the map Δ is well defined by the construction of Q . The map Δ is symmetric because \mathcal{S} is symmetric, and Δ is bilinear with respect to the group structure (2.13). Finally, the map $\eta: Q \times Q \rightarrow \mathbb{C}^\times$ is given by $\eta(I_i, I_j) = 1$, the constant map.

It is easy to see that the axioms of a braided logVA give us the axioms of a generalized vertex algebra. \square

3. EXISTENCE OF BRAIDED LOGARITHMIC VERTEX ALGEBRAS

This section follows similar steps as in [BV1, Section 2] where the case of locally nilpotent braiding was considered in detail. Let $\mathcal{S} = \sum_{i=1}^L \phi_i \otimes \psi_i$ be a braiding map on V as in Definition 1.1. Then we have the braiding map on $\text{Fie}(V)$ given by

$$(3.1) \quad \hat{\mathcal{S}} := \sum_{i=1}^L \text{ad}(\phi_i) \otimes \text{ad}(\psi_i).$$

Definition 3.1. We say that a subspace \mathcal{V} of $\text{Fie}(V)$ is $\hat{\mathcal{S}}$ -local if the following two conditions hold:

- (1) $\text{ad}(\phi_i), \text{ad}(\psi_i)$ are locally finite on \mathcal{V} for $1 \leq i \leq L$.
- (2) For any $a, b \in \mathcal{V}$ there is some integer $N \geq 0$ such that

$$(3.2) \quad \mu(\iota_{z_1, z_2}(z_{12}^{N+\hat{\mathcal{S}}})a(z_1) \otimes b(z_2)) = \mu(\iota_{z_2, z_1}(z_{12}^{N+\hat{\mathcal{S}}})b(z_2) \otimes a(z_1)),$$

where μ denotes the composition in $\text{End}(V)$.

Note that on the $\hat{\mathcal{S}}$ -local space \mathcal{V} we can apply Lemma 2.1. Next, we define the $(n + \hat{\mathcal{S}})$ -th products of fields as in [BV1].

Definition 3.2. Let $a, b \in \mathcal{V}$ be local logarithmic fields, and N be from (3.2). For $n \in \mathbb{Z}$, the $(n + \hat{\mathcal{S}})$ -th product $\hat{\mu}_{(n)}(a \otimes b) = a_{(n+\hat{\mathcal{S}})}b$ is defined by:

$$(3.3) \quad \begin{aligned} & (a_{(n+\hat{\mathcal{S}})}b)(z, \zeta) \\ & := \frac{D_{z_1}^{N-1-n}}{(N-1-n)!} \mu\left(\iota_{z_1, z_2}(z_{12}^{N+\hat{\mathcal{S}}})a(z_1, \zeta_1) \otimes b(z_2, \zeta_2)\right) \Big|_{\substack{z_1=z_2=z \\ \zeta_1=\zeta_2=\zeta}} \end{aligned}$$

for $n \leq N - 1$, and $a_{(n+\mathcal{S})}b := 0$ for $n \geq N$. From now on, to simplify the notation, we will suppress the dependence on ζ and understand that setting $z_1 = z$ automatically sets $\zeta_1 = \zeta$.

It is easy to check that $a_{(n+\mathcal{S})}b$ is again a field, and it does not depend on the choice of N satisfying (3.2). Note that, when $b = I$ is the identity operator, we have $N = 0$ and

$$(3.4) \quad (a_{(-1+\mathcal{S})}I)(z) = a(z), \quad (a_{(-2+\mathcal{S})}I)(z) = D_z a(z).$$

Let \mathcal{V} be an \mathcal{S} -local subspace of $\text{Fie}(V)$ containing I . We denote by \mathcal{V}^1 the minimal subspace of $\text{Fie}(V)$ containing $\hat{\mu}_{(n)}(\mathcal{V} \otimes \mathcal{V}) \subset \text{Fie}(V)$ for all $n \in \mathbb{Z}$, i.e.,

$$(3.5) \quad \mathcal{V}^1 = \sum_{n \in \mathbb{Z}} \hat{\mu}_{(n)}(\mathcal{V} \otimes \mathcal{V}).$$

Note that $D_z \mathcal{V} \subset \mathcal{V}^1$, by (3.4). We have the following analog of Dong's Lemma (cf. [BV1, Lemma 2.14]).

Lemma 3.3. *Let $\hat{\mathcal{S}}$ be the braiding map in (3.1) on $\text{Fie}(V)$, and \mathcal{V} be an \mathcal{S} -local subspace of $\text{Fie}(V)$. Then the space \mathcal{V}^1 is $\hat{\mathcal{S}}$ -local.*

The proof is the same as in [BV1, Lemma 2.14] and [BV1, Corollary 2.21], where the case of locally nilpotent braiding was considered.

Applying Lemma 3.3 with \mathcal{V}^1 in place of \mathcal{V} , we construct the local space $\mathcal{V}^2 = (\mathcal{V}^1)^1$. Continuing by induction, we let $\mathcal{V}^{m+1} = (\mathcal{V}^m)^1$, and obtain a sequence of \mathcal{S} -local subspaces

$$(3.6) \quad \mathcal{V} = \mathcal{V}^0 \subset \mathcal{V}^1 \subset \mathcal{V}^2 \subset \dots, \quad \bar{\mathcal{V}} := \bigcup_{m \geq 0} \mathcal{V}^m.$$

The union $\bar{\mathcal{V}}$ is $\hat{\mathcal{S}}$ -local and closed under all $(n + \hat{\mathcal{S}})$ -th products.

Now, we assume that our space V is equipped with a vector $\mathbf{1} \in V$ and a linear operator $T \in \text{End}(V)$ such that $T\mathbf{1} = 0$.

Definition 3.4. A field $a \in \text{Fie}(V)$ is called *translation covariant* if

$$(3.7) \quad [T, a(z)] = D_z a(z).$$

We denote by $\text{Fie}_T(V)$ the space of translation covariant fields.

Additionally, we will assume that the braiding map $\hat{\mathcal{S}}$ in (3.1) satisfies

$$(3.8) \quad \phi_i \mathbf{1} = \psi_i \mathbf{1} = 0, \quad [T, \phi_i] = [T, \psi_i] = 0, \quad i = 1, \dots, L.$$

The following theorem allows us to build examples of braided logVAs.

Theorem 3.5 (Existence Theorem). *Let V be a space equipped with a vector $\mathbf{1} \in V$, a linear operator $T \in \text{End}(V)$ such that $T\mathbf{1} = 0$, and a braiding map $\mathcal{S} = \sum_{i=1}^L \phi_i \otimes \psi_i$ on V satisfying (3.8). Set $\hat{\mathcal{S}} = \sum_{i=1}^L \text{ad}(\phi_i) \otimes \text{ad}(\psi_i)$. Suppose that we have an \mathcal{S} -local subspace of translation covariant fields,*

$$\mathcal{V} = \text{span}\{I, a^i(z) \mid i \in J\} \subset \text{Fie}_T(V),$$

where J is an index set, which is complete in the sense that

$$V = \text{span}\{\mathbf{1}, a_{(n_1+\mathcal{S})}^{i_1} \cdots a_{(n_k+\mathcal{S})}^{i_k} \mathbf{1} \mid k \geq 1, i_1, \dots, i_k \in J, n_1, \dots, n_k \in \mathbb{Z}\}.$$

Then $\bar{\mathcal{V}} \subset \text{Fie}_T(V)$, defined by (3.6), is an $\hat{\mathcal{S}}$ -local space containing \mathcal{V} closed under all $(n + \hat{\mathcal{S}})$ -th products, and for every $v \in V$ there exists a unique field $Y(v, z) \in \bar{\mathcal{V}}$ such that $Y(v, z)\mathbf{1}|_{z=0} = v$. Moreover, $(V, \mathbf{1}, T, Y, \mathcal{S})$ satisfies the conditions of Theorem 2.10, i.e., it is a braided logVA.

Proof. The proof follows the same steps in [BV1, Theorems 2.29, 3.2]. First, note that the $(n + \hat{\mathcal{S}})$ -th product of any two fields $a, b \in \mathcal{V}$ is translation covariant [BV1, Lemma 2.25]:

$$(3.9) \quad a_{(n+\hat{\mathcal{S}})}b \in \text{Fie}_T(V), \quad a, b \in \mathcal{V}, \quad n \in \mathbb{Z}$$

Second, recall that [BV1, Lemma 2.26]:

$$(3.10) \quad a(z)\mathbf{1} \in V[[z]], \quad a \in \text{Fie}_T(V).$$

Hence, the following linear map is well defined:

$$\Theta: \text{Fie}_T(V) \rightarrow V, \quad \Theta(a) := a(z)\mathbf{1}|_{z=0}.$$

We define $U := \text{span}_{\mathbb{C}}\{\Theta(a) \mid a \in \bar{\mathcal{V}}\} \subset V$, and the endomorphisms $a_{(n+\hat{\mathcal{S}})}$ on U by

$$(3.11) \quad \sum_{n \in \mathbb{Z}} z^{-n-1} a_{(n+\hat{\mathcal{S}})}u := \mu\left(z^{\sum_{i=1}^L \text{ad}(\phi_i) \otimes \psi_i} a(z) \otimes u\right), \quad u \in U.$$

Lemma 3.6. *Let $\hat{\mathcal{S}}$ be a braiding map as in (3.1) satisfying (3.8). Then, for any $\hat{\mathcal{S}}$ -local space $\mathcal{V} \subset \text{Fie}_T(V)$, we have*

$$(3.12) \quad \Theta(a_{(n+\hat{\mathcal{S}})}b) = a_{(n+\hat{\mathcal{S}})}\Theta(b), \quad a, b \in \mathcal{V}, \quad n \in \mathbb{Z}.$$

Proof. We will prove that

$$(3.13) \quad \mu\left(z^{\sum_{i=1}^L \text{ad}(\phi_i) \otimes \psi_i} a(z) \otimes \Theta(b)\right) = \sum_{n \in \mathbb{Z}} z^{-n-1} \Theta(a(z)_{(n+\hat{\mathcal{S}})}b(z)).$$

First, note that from (3.10), we have

$$c(z_1, z_2) := \mu\left(\iota_{z_1, z_2}(z_{12}^{N+\hat{\mathcal{S}}})a(z_1) \otimes b(z_2)\right)\mathbf{1} \in V[[z_1, z_2]][[z_1^{-\mathbb{C}}, \zeta_1]].$$

Then, using the locality (3.2), we get $c(z_1, z_2) \in V[[z_1, z_2]]$.

Next, we observe that

$$\Theta(\text{ad}(\psi_i)b(z)) = [\psi_i, b(z)]\mathbf{1}|_{z=0} = \psi_i\Theta(b).$$

Hence,

$$\begin{aligned} \mu\left(z_1^{\text{ad}(\phi_i) \otimes \psi_i} a(z_1) \otimes \Theta(b)\right) &= \mu\left(\iota_{z_1, z_2}(z_{12})^{\hat{\mathcal{S}}} a(z_1) \otimes b(z_2)\right)\mathbf{1}|_{z_2=0} \\ &= \mu\left(\iota_{z_1, z_2}(z_{12}^{-N})(z_{12}^{N+\hat{\mathcal{S}}})a(z_1) \otimes b(z_2)\right)\mathbf{1}|_{z_2=0} \\ &= z_{12}^{-N}c(z_1, z_2)\mathbf{1}|_{z_2=0} \\ &= \sum_{i=1}^r z_1^{-N}c(z_1, 0), \end{aligned}$$

where we used that $\iota_{z_1, z_2} \zeta_{12}|_{z_2=0} = \zeta_1$ and $\iota_{z_1, z_2} z_{12}^{-N}|_{z_2=0} = z_1^{-N}$. Now the coefficient in front of z_1^{N-n-1} in the Taylor expansion of $c(z_1, 0)$ is

$$\begin{aligned} & \frac{1}{(N-n-1)!} D_{z_1}^{N-n-1} c(z_1, 0)|_{z_1=0} \\ &= \frac{1}{(N-n-1)!} (D_{z_1}^{N-n-1} c(z_1, z_2)|_{z_1=z_2=z})|_{z=0} \\ &= \Theta(a_{(n+\hat{\mathcal{S}})} b), \end{aligned}$$

where we used (3.9). This proves (3.13), which together with (3.11) for $u = \Theta(b)$, implies (3.12). \square

The rest of the proof of Theorem 3.5 is the same as in [BV1, Theorems 2.29, 3.2] and is omitted. \square

Finally, we state the following two consequence of Theorem 3.5 and Lemma 3.6; cf. [BV1, Proposition 3.5 and Theorem 3.28].

Proposition 3.7 (*(n + S)-th product identity*). *If $(V, \mathbf{1}, T, Y, \mathcal{S})$ satisfies the assumptions of Theorem 2.10, then for any $a, b \in V$ we have*

$$(3.14) \quad Y(a, z)_{(n+\hat{\mathcal{S}})} Y(b, z) = Y(a_{(n+\mathcal{S})} b, z), \quad n \in \mathbb{Z}.$$

In particular, $Y(Ta, z) = D_z Y(a, z)$.

Proposition 3.8 (*Borcherds identity*). *If $(V, \mathbf{1}, T, Y, \mathcal{S})$ satisfies the assumptions of Theorem 2.10, then the Borcherds identity (2.9) holds.*

4. POISSON LIMITS AND NON-LINEAR SCHRÖDINGER NON-LOCAL PVA

In this section, first we prove Theorem 1.4. Then, we build the generalized vertex algebra defined by (1.4) and we prove that its associated non-local PVA is the non-linear Schrödinger PVA.

4.1. Poisson limits. We start by reviewing some definitions; for more details, we refer to [K, DK, BV2].

Definition 4.1. A *non-local Lie conformal algebra* (abbreviated non-local LCA) is a $\mathbb{C}[\partial]$ -module \mathcal{V} equipped with an admissible λ -bracket

$$\{\cdot, \lambda \cdot\}: \mathcal{V} \otimes \mathcal{V} \rightarrow \mathcal{V}((\lambda^{-1})),$$

satisfying the following axioms:

$$\begin{aligned} (\text{sesqui-linearity}) \quad & \{\partial a, \lambda b\} = -\lambda \{a, \lambda b\}, \quad \{a, \lambda \partial b\} = (\lambda + \partial) \{a, \lambda b\}, \\ (\text{skew-symmetry}) \quad & \{b, \lambda a\} = -\{a, -\lambda - \partial b\}, \\ (\text{Jacobi identity}) \quad & \{a, \lambda \{b, \mu c\}\} = \{\{a, \lambda b\}, \lambda + \mu c\} + \{b, \mu \{a, \lambda c\}\}. \end{aligned}$$

The admissibility of a λ -bracket is a technical condition, which ensures that the compositions in the Jacobi identity are well defined.

Definition 4.2. A *non-local Poisson vertex algebra* (abbreviated non-local PVA) is a non-local LCA \mathcal{V} , which in addition has a commutative associative product

$ab \in \mathcal{V}$ for $a, b \in \mathcal{V}$, with a unit $1 \in \mathcal{V}$, such that ∂ is a derivation of the product and the Leibniz rule holds:

$$\begin{aligned} \text{(derivation)} \quad & \partial(ab) = (\partial a)b + a(\partial b), \\ \text{(Leibniz rule)} \quad & \{a_\lambda bc\} = \{a_\lambda b\}c + b\{a_\lambda c\}. \end{aligned}$$

Now we recall the main result of [BV2]. In [BV2], we consider an increasing *filtration* on a logarithmic vertex algebra V by subspaces $\{0\} \subset F^0 V \subset F^1 V \subset F^2 V \subset \dots$ satisfying the following properties: $\mathbf{1} \in F^0 V$, $T(F^n V) \subset F^n V$ and

$$(4.1) \quad \begin{aligned} \mathcal{S}(F^m V \otimes F^n V) &\subset F^{m+n-1}(V \otimes V) := \sum_{k=0}^{m+n-1} F^k V \otimes F^{m+n-1-k} V, \\ \mu_{(j)}(F^m V \otimes F^n V) &\subset \begin{cases} F^{m+n} V, & j < 0, \\ F^{m+n-1} V, & j \geq 0, \end{cases} \end{aligned}$$

for all $m, n \in \mathbb{Z}_+$, where $F^n V := \{0\}$ for $n < 0$. Then, by [BV2, Theorem 3.3], the *associated graded* $\text{gr } V := \bigoplus_{n \in \mathbb{Z}_+} \text{gr}^n V$, where $\text{gr}^n V := F^n V / F^{n-1} V$, has the structure of a non-local PVA defined as follows. The unit $1 \in \text{gr}^0 V = F^0 V$ is $\mathbf{1}$; the derivation ∂ on $\text{gr } V$ is induced from T ; the commutative associative product on $\text{gr } V$ is induced from $\mu_{(-1)}$; and the λ -bracket on $\text{gr } V$ is given by

$$(4.2) \quad \{a_\lambda b\} := \sum_{n \in \mathbb{Z}_+} \frac{\lambda^n}{n!} \mu_{(n)}(a \otimes b) + \mu_{(-1)} \left(\mathcal{S} \left(\frac{1}{\lambda + \partial} a \otimes b \right) \right).$$

In [FB], Poisson vertex algebras were associated to vertex algebras considering (algebraic) limits. In this section, we follow the same idea. First, we define ideals. An ideal W of a braided logVA V is a subspace invariant under T, ϕ_i, ψ_i ($1 \leq i \leq L$) such that $\mu_{(n)}(a \otimes b) \in W$ for all $a \in V, b \in W, n \in \mathbb{Z}$. As for ordinary vertex algebras, it is not hard to see that V/W has the structure of a braided logVA.

Next, we note that it is straightforward to define the notion of a braided logVA V over a commutative ring R instead of \mathbb{C} , so that V is an R -module and $T, \psi_i, \phi_i, \mu_{(n)}$ ($1 \leq i \leq L, n \in \mathbb{Z}$) commute with the action of R on V . In particular, we will be interested in the case $R = \mathbb{C}[\epsilon]/(\epsilon^2)$.

Example 4.3. If V is a braided logVA, then $V^\epsilon := V \otimes \mathbb{C}[\epsilon]/(\epsilon^2)$ is a braided logVA over $\mathbb{C}[\epsilon]/(\epsilon^2)$, the subspace ϵV^ϵ is an ideal of V^ϵ , and the quotient $V^\epsilon / \epsilon V^\epsilon \cong V$.

Now we prove Theorem 1.4.

Proof of Theorem 1.4. Since $Y^0 = Y^\epsilon \text{ mod } \epsilon$ is commutative, we have

$$(4.3) \quad \begin{aligned} \mathcal{S}(V^\epsilon \otimes V^\epsilon) &\subset \epsilon(V^\epsilon \otimes V^\epsilon), \\ \mu_{(j)}(V^\epsilon \otimes V^\epsilon) &\subset \epsilon V^\epsilon, \quad j \geq 0. \end{aligned}$$

For $a, b \in V^\epsilon$, we define a λ -bracket by

$$\{a_\lambda b\} := \sum_{n \in \mathbb{Z}_+} \frac{1}{\epsilon} \frac{\lambda^n}{n!} \mu_{(n)}(a \otimes b) + \frac{1}{\epsilon} \mu_{(-1)} \left(\mathcal{S} \left(\frac{1}{\lambda + \partial} a \otimes b \right) \right) \quad \text{mod } \epsilon \in V^0((\lambda^{-1})).$$

When $a \in \epsilon V^\epsilon$ or $b \in \epsilon V^\epsilon$, we have $\{a_\lambda b\} = 0$; hence, this descends to a well-defined λ -bracket on the quotient V^0 .

Now we check that the above bracket satisfies each of the conditions of Definitions 4.1 and 4.2. The main point is that the relations (4.3) are similar to (4.1). Hence,

the proof that V^0 is a non-local Poisson vertex algebra is similar to the proof in [BV2, Theorem 3.3].

For instance, let us check sesqui-linearity. From Proposition 3.7, $Y(Ta, z) = D_z Y(a, z)$ on V^ϵ ; hence $\mu_{(n)}(Ta \otimes b) = -\mu_{(n-1)}(n + \mathcal{S})(a \otimes b)$. Then from (4.3) on V^0 , we get that

$$\begin{aligned} \frac{1}{\epsilon} \mu_{(n)}(\partial a \otimes b) &= -\frac{1}{\epsilon} n \mu_{(n-1)}(a \otimes b) \pmod{\epsilon} \text{ for } n \geq 1, \\ \frac{1}{\epsilon} \mu_{(0)}(\partial a \otimes b) &= -\frac{1}{\epsilon} n \mu_{(-1)}(\mathcal{S}(a \otimes b)) \pmod{\epsilon}. \end{aligned}$$

Then as in the proof of [BV2, Proposition 3.5 (i)], we obtain $\{\partial a_\lambda b\} = -\lambda \{a_\lambda b\}$.

Similarly, the skew-symmetry is proved as in the proof of [BV2, Proposition 3.5 (iii)]. The Leibniz rule is proved exactly as in [BV2, Proposition 3.6], and by [BV2, Proposition 3.5] the bracket is admissible. Finally, the Jacobi identity is proved as in [BV2, Proposition 3.8]. Note that identity [BV2, (3.17)] is satisfied mod ϵ multiplying by $\frac{1}{\epsilon^2}$. \square

4.2. The non-linear Schrödinger generalized vertex algebra. We start by reviewing the following example of a non-local PVA from [DK].

Definition 4.4. The *non-linear Schrödinger non-local PVA* is defined as the algebra of differential polynomials $\mathcal{V} := \mathbb{C}[u, v, \partial u, \partial v, \dots]$ with λ -bracket given by:

$$(4.4) \quad \begin{aligned} \{u_\lambda u\} &= v(\lambda + \partial)^{-1}v, & \{u_\lambda v\} &= -v(\lambda + \partial)^{-1}u, \\ \{v_\lambda u\} &= -v(\lambda + \partial)^{-1}u, & \{v_\lambda v\} &= u(\lambda + \partial)^{-1}u. \end{aligned}$$

Note that we can express equivalently the non-linear Schrödinger non-local PVA using the basis $w := u + iv$, $\bar{w} := u - iv$. Then $\mathcal{V} = \mathbb{C}[w, \bar{w}, w', \bar{w}', \dots]$ (where $w' := \partial w$), and the λ -bracket is given by (4.2) for $a, b \in \{w, \bar{w}\}$, where

$$\mathcal{S} = \mathcal{S}^{(s)} := \Phi \otimes \Phi, \quad \Phi(w) = iw, \quad \Phi(\bar{w}) = -i\bar{w},$$

$\mu_{(-1)}$ denotes the product in \mathcal{V} , and

$$(4.5) \quad \mu_{(n)}(a \otimes b) = 0, \quad a, b \in \{w, \bar{w}\}, \quad n \geq 0.$$

We use the above relations to construct a braided logVA V . We will assume that it exists and will derive the properties that V should satisfy.

Lemma 4.5. *Let V be a braided logVA with elements $w, \bar{w} \in V$, whose $(n + \mathcal{S})$ -th products satisfy (4.5). Suppose that there is $\Phi \in \text{Der}(V)$ acting as $\Phi(w) = iw$, $\Phi(\bar{w}) = -i\bar{w}$, such that $\mathcal{S} = \Phi \otimes \Phi$. Then*

$$Y(w, z) = \sum_{n \in \mathbb{Z}} w_{(n+\mathcal{S})} z^{-n-1-i\Phi}, \quad Y(\bar{w}, z) = \sum_{n \in \mathbb{Z}} \bar{w}_{(n+\mathcal{S})} z^{-n-1+i\Phi}$$

where the linear operators $w_n := w_{(n+\mathcal{S})}$, $\bar{w}_n := \bar{w}_{(n+\mathcal{S})}$ satisfy relations (1.4).

Proof. The form of $Y(w, z)$ and $Y(\bar{w}, z)$ follows from (2.3), using that $\mathcal{S}^{(n)} = 0$. From the Borcherds identity (1.3) with $n = 0$ on $w \otimes w \otimes v$, $v \in V$, we have:

$$\begin{aligned} \sum_{j \in \mathbb{Z}_+} (-1)^j \mu_{(m-j)}(I \otimes \mu_{(k+j)}) \binom{-1}{j} (w \otimes w \otimes v) &= \sum_{j \geq 0} w_{(m-j+\mathcal{S})} w_{(k+j+\mathcal{S})} v, \\ \sum_{j \in \mathbb{Z}_+} (-1)^j \mu_{(k-j)}(I \otimes \mu_{(m+j)}) (-1)^{-1} \binom{-1}{j} (w \otimes w \otimes v) &= - \sum_{j \geq 0} w_{(m-j+\mathcal{S})} w_{(k+j+\mathcal{S})} v, \\ \sum_{j \in \mathbb{Z}_+} \mu_{(m+k-j)}(\mu_{(j)} \otimes I) \binom{m+iI \otimes I \otimes \Phi}{j} (w \otimes w \otimes v) &= 0, \end{aligned}$$

where in the last identity we used (4.5). Hence, from (1.3) we obtain the first equation in (1.4). The other two relations in (1.4) are derived analogously from (1.3) for $\bar{w} \otimes \bar{w} \otimes v$ and $w \otimes \bar{w} \otimes v$, respectively. \square

We now use the previous results to construct V . We proceed similarly to the construction in [BV1, Section 4.4], and [BV2, Section 4.3]. We define a unital associative topological algebra \mathcal{A} with generators $\{w_n, \bar{w}_n \mid n \in \mathbb{Z}\}$ subject to the relations (1.4). The algebra \mathcal{A} admits a module V , generated by an element $\mathbf{1} \in V$ such that $w_n \mathbf{1} = \bar{w}_n \mathbf{1} = 0$ for $n \geq 0$, and V is linearly spanned by monomials of the form

$$(4.6) \quad a_{n_1}^1 \cdots a_{n_r}^r \mathbf{1} \quad \text{where} \quad n_i \leq -1, \quad a^i \in \{w, \bar{w}\}, \quad 1 \leq i \leq r.$$

Next, we define linear operators T and Φ on V by $T(\mathbf{1}) = \Phi(\mathbf{1}) = 0$ and

$$(4.7) \quad \begin{aligned} [\Phi, w_n] &= iw_n, & [\Phi, \bar{w}_n] &= -i\bar{w}_n, \\ [T, w_n] &= -nw_{n-1} - iw_{n-1}\Phi, & [T, \bar{w}_n] &= -n\bar{w}_{n-1} + i\bar{w}_{n-1}\Phi, \end{aligned}$$

motivated by the hexagon identity (1.2) and translation covariance (2.5). Additionally, we define the fields

$$(4.8) \quad w(z) := \sum_{n \in \mathbb{Z}} w_n z^{-n-1-i\Phi}, \quad \bar{w}(z) := \sum_{n \in \mathbb{Z}} \bar{w}_n z^{-n-1+i\Phi}.$$

We let the braiding map $\mathcal{S} = \Phi \otimes \Phi$. As before, this gives rise to a braiding map $\hat{\mathcal{S}} := \text{ad}(\Phi) \otimes \text{ad}(\Phi)$ on the space of fields $\text{Fie}(V)$.

Theorem 4.6. *The fields $w(z)$, $\bar{w}(z)$ are translation covariant and $\hat{\mathcal{S}}$ -local, and they generate the structure of a braided logVA on V .*

Proof. We verify each of the hypotheses of Theorem 3.5. By definition we have $\mathbf{1} \in V$ such that $T\mathbf{1} = 0$ and $\mathcal{S}(\mathbf{1} \otimes a) = 0$. By the Jacobi identity, $[[T, \Phi], w_n] = [T, [\Phi, w_n]] - [\Phi, [T, w_n]] = 0$, and similarly, $[[T, \Phi], \bar{w}_n] = 0$. Hence $[T, \Phi] = 0$. Next, we observe that by construction the fields $w(z)$, $\bar{w}(z)$ are translation covariant (cf. (2.5)). We will prove that the space

$$\mathcal{U} = \text{span}\{I, w(z), \bar{w}(z)\} \subset \text{Fie}_T(V)$$

is $\hat{\mathcal{S}}$ -local. We have $z_1^{-i\Phi}w(z_2) = w(z_2)z_1^{1-i\Phi}$ and $z_1^{i\Phi}w(z_2) = w(z_2)z_1^{-1+i\Phi}$. Then using (1.4), we obtain:

$$\begin{aligned}
\mu(\iota_{z_1, z_2}(z_{12}^{\hat{\mathcal{S}}})w(z_1) \otimes w(z_2)) &= \iota_{z_1, z_2}(z_{12}^{-1})w(z_1)w(z_2) \\
&= \sum_{m, k \in \mathbb{Z}} \left(\sum_{j \geq 0} w_{m-j}w_{k+j} \right) z_1^{-m-1-i\Phi} z_2^{-k-1-i\Phi} \\
&= - \sum_{m, k \in \mathbb{Z}} \left(\sum_{j \geq 0} w_{k-j}w_{m+j} \right) z_1^{-m-1-i\Phi} z_2^{-k-1-i\Phi} \\
&= \iota_{z_2, z_1}(z_{12}^{-1})w(z_2)w(z_1) \\
&= \mu(\iota_{z_2, z_1}(z_{12}^{\hat{\mathcal{S}}})w(z_2) \otimes w(z_1)).
\end{aligned}$$

Hence, we have the locality of $w(z)$ with itself as in (3.2) for $N = 0$. The locality of $\bar{w}(z)$, $\bar{w}(z)$ and $w(z)$, $\bar{w}(z)$ is proved in a similar way. \square

Remark 4.7. One can show that the braided logVA V in Theorem 4.6 has a PBW basis consisting of lexicographically ordered monomials of the form (4.6). The proof is similar to the usual Poincaré–Birkhoff–Witt Theorem for Lie algebras (see, e.g., [J]). In the context of logarithmic vertex algebras, a version of the PBW Theorem was proved in [H], while the case of quadratic associative algebras is discussed in [PP].

Finally, following [FB, Section 16.3], we deform the braided logVA in Theorem 4.6 by introducing a formal parameter ϵ . We consider the unital associative topological algebra \mathcal{A}^ϵ with generators $\{w_n, \bar{w}_n \mid n \in \mathbb{Z}\}$ subject to the relations:

$$\begin{aligned}
(4.9) \quad & \sum_{j \geq 0} (-1)^j \binom{-\epsilon}{j} (w_{m-j}w_{k+j} - w_{k-j}w_{m+j}) = 0, \\
& \sum_{j \geq 0} (-1)^j \binom{-\epsilon}{j} (\bar{w}_{m-j}\bar{w}_{k+j} - \bar{w}_{k-j}\bar{w}_{m+j}) = 0, \\
& w_m \bar{w}_k - \epsilon w_{m-1} \bar{w}_{k+1} - \bar{w}_k w_m + \epsilon \bar{w}_{k-1} w_{m+1} = 0.
\end{aligned}$$

As in Theorem 4.6, \mathcal{A}^ϵ has a module V^ϵ spanned by the monomials (4.6). We have operators $T, \Phi \in \text{End}_{\mathbb{C}[e]}(V^\epsilon)$, fields

$$w^\epsilon(z) = \sum w_n z^{-n-1} e^{i\epsilon\Phi\zeta}, \quad \bar{w}^\epsilon(z) = \sum \bar{w}_n z^{-n-1} e^{-i\epsilon\Phi\zeta},$$

a braiding map $\mathcal{S} = \epsilon\Phi \otimes \Phi$, and $\hat{\mathcal{S}} := \epsilon \text{ad}(\Phi) \otimes \text{ad}(\Phi)$ on $\text{Fie}(V^\epsilon)$. The same steps as in the proof of Theorem 4.6 give us that V^ϵ is a braided logVA. Moreover, $V^0 := V^\epsilon/\epsilon V^\epsilon$ is commutative. Then Theorem 1.4 implies that V^0 has the structure of a non-local PVA. It is not hard to see that this non-local PVA is exactly the non-linear Schrödinger non-local PVA from Definition 4.4.

Acknowledgments. The first author was supported in part by a Simons Foundation grant 584741. He is grateful to the University of Bath for the hospitality during August 2023. The second author was supported in part by UKRI grant MR/S032657/1.

REFERENCES

- [AKM] Arakawa, T., Kuwabara, T. and Malikov, F.: *Localization of affine W -algebras* Comm. Math. Phys., **335**:143–182, (2015).
- [AKM2] Arakawa, T., Kuwabara, T. and Möller, S.: *Hilbert Schemes of Points in the Plane and Quasi-Lisse Vertex Algebras with $N=4$ Symmetry* (2023), arXiv:2309.17308.
- [B] Bakalov, B.: *Twisted logarithmic modules of vertex algebras*. Commun. Math. Phys. **345**, 355–383 (2016)
- [BK] Bakalov, B., Kac, V.G.: *Generalized vertex algebras*. In: “Lie theory and its applications in physics VI,” 3–25, ed. V.K. Dobrev et al., Heron Press, Sofia, 2006
- [BV1] Bakalov, B.N., Villarreal, J.: *Logarithmic vertex algebras*. Transf. Groups (2022), <https://doi.org/10.1007/s00031-022-09759-z>
- [BV2] Bakalov, B., Villarreal, J.: *Logarithmic vertex algebras and non-local Poisson vertex algebras*. Commun. Math. Phys. **404**, 185–226 (2023)
- [Bo] Borcherds, R.E.: *Vertex algebras, Kac–Moody algebras, and the Monster*. Proc. Nat. Acad. Sci. USA **83**, 3068–3071 (1986)
- [K] Kac, V.G.: *Introduction to vertex algebras, Poisson vertex algebras, and integrable Hamiltonian PDE*. Adv. Math. **281**, 1025–1099, (2015)
- [DK] De Sole, A., Kac, V.G.: *Non-local Poisson structures and applications to the theory of integrable systems*. Japanese J. Math. **8**, 233–347 (2013)
- [DKV] De Sole A., Kac V.G., Valeri D.: *Dirac reduction for Poisson vertex algebras*. Commun. Math. Phys. **331**, 1155–1190 (2014)
- [DKV2] De Sole A., Kac V.G., Valeri D.: *Integrability of Dirac reduced bi-Hamiltonian equations*. Trends in Contemporary Math., Springer INDAM **8**, 13–32 (2014)
- [DL] Dong, C., Lepowsky, J.: *Generalized vertex algebras and relative vertex operators*. Progress in Math., 112, Birkhäuser Boston, 1993
- [EK] Etingof, P., Kazhdan, D.: *Quantization of Lie bialgebras, Part V: Quantum vertex operator algebras*. Sel. Math., New Ser. **6**, 105–130 (2000)
- [FB] Frenkel, E., Ben-Zvi, D.: *Vertex algebras and algebraic curves*. Math. Surveys and Monographs, 88, Amer. Math. Soc., Providence, RI, 2001; 2nd ed., 2004
- [FFR] Feingold, A.J., Frenkel, I.B., Ries, J.F.X.: *Spinor construction of vertex operator algebras, triality, and $E_8^{(1)}$* . Contemp. Math., 121, Amer. Math. Soc., Providence, RI, 1991
- [H] Hatam, H.: *Advances in logarithmic vertex algebras*. PhD dissertation, North Carolina State University, 2024
- [J] Jacobson, N.: *Lie algebras*. No. 10. Courier Corporation, 1979
- [K] Kac, V.G.: *Vertex algebras for beginners*. University Lecture Series, 10, Amer. Math. Soc., Providence, RI, 1996; 2nd ed., 1998
- [PP] Polishchuk, A., Positselski, L.: *Quadratic algebras*. University Lecture Series, 37, Amer. Math. Soc., Providence, RI, 2005
- [V] Villarreal, J.: *Lambda bracket and intertwiners*. Preprint (2023), arXiv:2310.18872

DEPARTMENT OF MATHEMATICS, NORTH CAROLINA STATE UNIVERSITY, RALEIGH, NC 27695,
UNITED STATES

Email address: `bojko_bakalov@ncsu.edu`

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF BATH, BATH BA2 7AY, UNITED
KINGDOM

Email address: `juanjos3villarreal@gmail.com`