A matricial view of the Collatz conjecture

Pietro Paparella^a

^aDivision of Engineering & Mathematics, University of Washington Bothell, Bothell, 98011, WA, U.S.

Abstract

In this note, it is shown that the nilpotency of submatrices of a certain class of adjacency matrices is equivalent to the Collatz conjecture. Our main result extends the previous work of Alves et al. and clarifies a conjecture made by Cardon and Tuckfield.

Keywords: adjacency matrix, Collatz conjecture, Collatz sequence, directed graph, nilpotent matrix 2020 MSC: 15A18, 15A15, 11B83

1. Introduction

Let $f : \mathbb{N} \longrightarrow \mathbb{N}$ be the function defined by

$$f(n) = \begin{cases} \frac{n}{2}, & n \equiv 0 \mod 2, \\ \frac{3n+1}{2}, & n \equiv 1 \mod 2, \end{cases}$$

and, for a fixed $n \in \mathbb{N}$, consider the sequence $s(n) : \mathbb{N}_0 \longrightarrow \mathbb{N}$ defined by

$$s_k(n) \coloneqq \begin{cases} n, & k = 0, \\ f^k(n), & k > 0 \end{cases}.$$

The function f is called the *(shortcut) Collatz function* and the sequence s(n) is called the *Collatz sequence of* n.

It is well-known that if n is a positive integer, then there are three possibilities for the sequence s(n) [4, Section 2]:

Preprint submitted to Linear Algebra Appl.

Email address: pietrop@uw.edu (Pietro Paparella)

URL: https://faculty.washington.edu/pietrop/ (Pietro Paparella)

- (i) Convergent trajectory. In this case, there is a positive integer p such that $s_p(n) = f^p(n) = 1$. If p is the smallest integer such that $s_p(n) = 1$, then p is called the *total stopping time (of n)*.
- (ii) Nontrivial cycle. In this case, there is a positive integer n > 2 such that $s_k(n) \neq 1$, $\forall k \in \mathbb{N}$ and the sequence s(n) is periodic—i.e., there are positive integers p and q such that $s_p(n) = f^p(n) = f^q(n) = s_q(n)$.
- (iii) Divergent sequence. In this case, the sequence s(n) is unbounded, i.e., $\lim_{k \to \infty} s_k(n) = \infty.$

As is well-known, the *Collatz conjecture* (see, e.g., Lagarias [4] and references therein), asserts that all Collatz sequences are convergent. There are many known reformulations of this problem [5, 6]. The assertion that no nontrivial cyclic trajectories exist will be called the *aperiodic Collatz conjecture*.

Let $A: \mathbb{N} \times \mathbb{N} \longrightarrow \{0, 1\}$ be the infinite matrix defined by

$$a_{ij} = \delta_{f(i),j} = \begin{cases} 1, & j = f(i), \\ 0, & \text{otherwise.} \end{cases}$$

and for $n \in \mathbb{N}$, let A_n be the *n*-by-*n* matrix whose (i, j)-entry is $a_{ij}, 1 \leq i, j \leq n$.

Introduced by Alves et al. [1], the matrix A_n is the adjacency matrix of the directed graph Γ_n with vertices $V_n := \{1, \ldots, n\}$ and (directed) edges $E_n := \{(i, j) \in V_n \times V_n \mid j = f(i)\}$. In addition, they proved the following result.

Theorem 1 ([1, Theorem 1 and Lemma 1]). The aperiodic Collatz conjecture is equivalent to

$$\operatorname{tr}(A_n^p) = \begin{cases} 2, & p \equiv 0 \mod 2, \\ 0, & p \equiv 1 \mod 2. \end{cases}$$
(1)

However, more can be gleaned from carefully examining a submatrix of A_n when n > 2. Since f(2) = 1 and f(1) = 2, it follows that

$$A_2 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix},$$

and

$$A_n = \begin{bmatrix} A_2 & 0\\ B_n & C_n \end{bmatrix},$$

whenever n > 2. More formally, let $\mathbb{N}_3 := \{n \in \mathbb{N} \mid n \geq 3\}$ and let $C : \mathbb{N}_3 \times \mathbb{N}_3 \longrightarrow \{0, 1\}$ be the infinite matrix defined by $c_{ij} = a_{ij}$. For $n \geq 3$, let C_n be the (n-2)-by-(n-2) matrix whose (i, j)-entry is $c_{ij} = a_{ij}, 3 \leq i, j \leq n$. For ease of notation, the rows and columns of C_n are indexed by $\{3, \ldots, n\}$.

Denote by $\mathsf{M}_n(\mathbb{C})$ the algebra of *n*-by-*n* matrices with complex entries. If $A \in \mathsf{M}_n(\mathbb{C})$, then A is called *nilpotent* if there is a positive integer p such that $A^p = 0$. In such a case, the set $\{k \in \mathbb{N} \mid A^k = 0\}$ is nonempty and hence, by the well-ordering principle, contains a smallest element, which is called the *index (of nilpotency) of A*.

Conjecture 2. If $n \ge 3$, then the matrix C_n is nilpotent.

The following result, which is the central aim of this work, reveals the import of Conjecture 2.

Theorem 3. The Collatz conjecture is equivalent to Conjecture 2.

Remark 4. If

$$M = \begin{bmatrix} A & 0 \\ B & C \end{bmatrix} \in \mathsf{M}_n(\mathbb{C}),$$

then a straightforward proof by induction reveals that

$$M^{p} = \begin{bmatrix} A^{p} & 0\\ \sum_{k=0}^{p-1} C^{k} B A^{p-1-k} & C^{p} \end{bmatrix}, \forall p \in \mathbb{N}.$$

Consequently, $\operatorname{tr}(A_n^p) = \operatorname{tr}(A_2^p) + \operatorname{tr}(C_n^p)$ and Conjecture 2 implies (1) since

$$\operatorname{tr}(A_2^p) = \begin{cases} 2, & p \equiv 0 \mod 2, \\ 0, & p \equiv 1 \mod 2. \end{cases}$$

In addition, Conjecture 2 simplifies a similar conjecture posed by Cardon and Tuckfield [2, Conjecture 12].

Because nilpotency has several spectral characterizations, Conjecture 2 places the Collatz conjecture in the arena of spectral graph theory, a serendipitous turn in view of the fact that Collatz is widely considered as a co-founder of the field.

2. Notation & Background

The following standard matrix-theoretic notation is adopted throughout. If $A \in \mathsf{M}_n(\mathbb{C})$, then

- the (i, j)-entry of A is denoted by $a_{ij}, a_{i,j}$, or $(A)_{ij}$;
- |A| denotes the determinant of A;
- p_A denotes the characteristic polynomial of A, i.e., $p_A(t) := |tI A|;$

•
$$\sigma(A) \coloneqq \{\lambda \in \mathbb{C} \mid p_A(\lambda) = 0\}; \text{ and}$$

• $\operatorname{tr}(A) \coloneqq \sum_{k=1}^n a_{kk}.$

The following result lists several characterizations of nilpotency that are relevant to this work (for other characterizations, see, e.g., Horn and Johnson [3]).

Theorem 5. If $A \in M_n(\mathbb{C})$, then the following are equivalent:

- *i)* A is nilpotent;
- *ii)* $A^n = 0$ /3, p. 109/; and
- *iii)* $\operatorname{tr}(A^p) = 0, \ \forall p \in \mathbb{N} \ /3, \ Problem \ 2.4.P10/.$

If $V_n = \{1, \ldots, n\}$ and $E \subseteq V_n \times V_n$, then $\Gamma = (V_n, E)$ is called a *directed* graph or *digraph*. If $A = [a_{ij}]$ is the *n*-by-*n* matrix such that

$$a_{ij} \coloneqq \begin{cases} 1, & (i,j) \in E, \\ 0, & (i,j) \notin E, \end{cases}$$

then A is called the *adjacency matrix of* Γ . If $W = \{(i_0, i_1), \ldots, (i_{\ell-1}, i_\ell)\} \subseteq E, \ell \geq 1$, then W is called a *(directed) walk (of length* ℓ). If $i_0 = i_\ell$, then W is called a *(directed) cycle (of length* ℓ). It is well-known that if $A \in \mathsf{M}_n(\mathbb{C})$ and $p \geq 2$, then

$$(A^p)_{ij} = \sum_{k_1,\dots,k_{p-1}=1}^n \left[\prod_{\ell=1}^p a_{k_{\ell-1},k_\ell}\right],$$

where $k_0 \coloneqq i$ and $k_p \coloneqq j$. In the context of adjacency matrices, the (i, j)entry of A^p yields the number of directed walks of length p of Γ beginning at node i and terminating at node j. In particular, the (i, i)-entry of A^p is the number of directed cycles of length p that originate and terminate at node i.

3. Proof of Main Result

Proof of Theorem 3. Suppose that the Collatz conjecture is true. If n > 2, then the subdigraph of Γ_n with vertices $V_n \setminus \{1, 2\}$ and edges

$$E_n \setminus \{(1,2), (2,1), (4,2)\}$$

does not have any directed cycles. Thus, every diagonal entry of C_n^p is zero. Consequently, $\operatorname{tr}(C_n^p) = 0$, $\forall p \in \mathbb{N}$ and C_n is nilpotent.

Conversely, suppose that C_n is nilpotent $\forall n \in \mathbb{N}_3$ and, for contradiction, that the Collatz conjecture is false.

If there is a nontrivial cycle, then there is a positive integer n > 2 such that $s_k(n) \neq 1$, $\forall k \in \mathbb{N}$ and positive integers p and q such that $s_p(n) = f^p(n) = f^q(n) = s_q(n)$. Assume, without loss of generality, that p < q. If $m \coloneqq s_p(n)$, then $s_{q-p}(m) = m$ and Γ_m contains a cycle of length q - p. Consequently, the (m, m)-entry of $(C_m)^{q-p}$ is positive. This contradicts part iii) of Theorem 5.

Otherwise, if there is a divergent sequence, then there is a positive integer n such that the sequence s(n) is unbounded, i.e., $\lim_{k \to \infty} s_k(n) = \infty$. Thus, if M > 0, then there is a positive integer N such that $m_k := s_k(n) \ge M$, $\forall k \ge N$. If $\ell = \max\{n, N\}$, then there is a walk of length ℓ from n to m_ℓ in Γ_ℓ . But $(C_\ell)^\ell = 0$, a contradiction.

References

- J. F. Alves, M. M. Graça, M. E. Sousa Dias, and J. Sousa Ramos. A linear algebra approach to the conjecture of Collatz. *Linear Algebra Appl.*, 394:277–289, 2005.
- [2] D. A. Cardon and B. Tuckfield. The Jordan canonical form for a class of zero-one matrices. *Linear Algebra Appl.*, 435(11):2942–2954, 2011.
- [3] R. A. Horn and C. R. Johnson. *Matrix analysis*. Cambridge University Press, Cambridge, second edition, 2013.
- [4] J. C. Lagarias. The 3x + 1 problem and its generalizations. Amer. Math. Monthly, 92(1):3–23, 1985.
- [5] J. C. Lagarias. The 3x + 1 problem: An annotated bibliography (1963–1999) (sorted by author), 2011.

[6] J. C. Lagarias. The 3x+1 problem: An annotated bibliography, ii (2000-2009), 2012.