

Formulas of special polynomials involving Bernoulli polynomials derived from matrix equations and Laplace transform

Ezgi Polat and Yilmaz Simsek

Department of Mathematics, Faculty of Science, University of Akdeniz, Antalya TR-07058, Turkey

E-mail. ezgipolat1247@gmail.com and ysimsek@akdeniz.edu.tr

Abstract. The main purpose and motivation of this article is to create a linear transformation on the polynomial ring of rational numbers. A matrix representation of this linear transformation based on standard fundamentals will be given. For some special cases of this matrix, matrix equations including inverse matrices, the Bell polynomials will be given. With the help of these equations, new formulas containing different polynomials, especially the Bernoulli polynomials, will be given. Finally, by applying the Laplace transform to the generating function for the Bernoulli polynomials, we derive some novel formulas involving the Hurwitz zeta function and infinite series.

Keyword. Generating functions, Bernoulli polynomials and numbers, Linear map, Inverse matrix, Laplace transform, Hurwitz zeta function

MSC2020. 05A15, 11B68, 44A10, 47L05

1 Introduction

It is well known that in recent years many different applications of linear transformations have been given not only in the algebraic field of mathematics but also in other applied sciences. To give an example of these different applications, geometric transformations implemented in computer graphics also occur; that is translation, rotation and scaling of $2D$ or $3D$ objects can be done using a transformation matrix. Therefore, linear transformations are also used as a tool to describe change. Many examples can be given for these, some of which are well known to be used in analysis as transformations corresponding to derivatives, transformations corresponding to integrals, or in relativity as a device to keep track of local transformations of reference frames. In addition, other implementation examples can be given, including compiler optimizations of nested loop code and parallelization of compiler techniques.

In recent years, it has also been seen that generating functions for the Bernoulli numbers and polynomials were given by different methods. For example, in complex analysis, generating functions of these numbers and polynomials can be constructed using the Cauchy derivative formula, Cauchy residue theorem and meromorphic functions. These numbers and polynomials are also used in algebraic topology and number theory, as criteria of regular prime numbers

or their appearance in the Todd class which can be seen on complex vector bundles in topological space, the values of zeta functions on even integers, in Milnor's homotopy group related to the characteristic class. They can also appear in K -theory, the Euler-Maclaurin summation formula, Bessel functions, trigonometric functions, cylindrical functions, hypergeometric functions etc. In addition, it is also well-known that these numbers and polynomials are given by the Volkenborn integral on the set of p -adic integers. Perhaps their definitions are also given on other spaces or sets that we have not seen before. Therefore, the main motivation of this paper is to give different computational formulas of polynomials containing the Bernoulli polynomials using not only the linear transformation defined on the polynomials ring of rational numbers and its matrix equations, but also the Laplace transform and the Hurwitz zeta function.

We can use the following notations and definitions:

Let \mathbb{N} , \mathbb{Q} , \mathbb{R} and \mathbb{C} denote a set of positive integers, the ring of rational integers, a set of real numbers, and a set of complex numbers. $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$.

The Bernoulli numbers and polynomials are respectively given by

$$F(t) = \frac{t}{e^t - 1} = \sum_{n=0}^{\infty} B_n \frac{t^n}{n!} \quad (1.1)$$

and

$$F(t, x) = e^{tx} F(t) = \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!} \quad (1.2)$$

(cf. [1, 16, 21]).

The Stirling numbers of the first kind and the second kind, which are denoted by $S_1(n, k)$ and $S_2(m, n)$, respectively, are defined by

$$\frac{(\log(1+t))^k}{k!} = \sum_{n=0}^{\infty} S_1(n, k) \frac{t^n}{n!} \quad (1.3)$$

and

$$\frac{(e^t - 1)^n}{n!} = \sum_{m=0}^{\infty} S_2(m, n) \frac{t^m}{m!} \quad (1.4)$$

(cf. [5, 16, 21]).

The array polynomials are defined by the following generating function:

$$\frac{(e^t - 1)^v e^{tx}}{v!} = \sum_{m=0}^{\infty} S_v^m(x) \frac{t^m}{m!} \quad (1.5)$$

(cf. [3, 4, 17]).

The results of this article, including all sections, are briefly stated as follows:

In Section 2, we defined \mathbb{Q} -linear operator on the polynomials ring $\mathbb{Q}[x]$. By using this operator, we give its matrix representation with respect to the basis of $\mathbb{Q}[x]$: $\{1, x, x^2, x^3, \dots\}$. By using matrix representation, we find some

new classes of special polynomials involving Bernoulli polynomials and Bell polynomials.

In Section 3, we give linear transformation. By applying Cayley–Hamilton theorem, we also give inverse matrix formulas involving the Bell polynomials for this linear transformation. Using this inverse matrix formula, we derive matrix representation for the Bernoulli polynomials.

In Section 4, we give a derivative formulas for linear transformation.

In Section 5, we define a new family of polynomials and its matrix. By using this matrix with its inverse, we also derive the Bernoulli numbers.

In Section 6, applying the Laplace transform to generating function for the Bernoulli polynomials, we give not only infinite series representation for the Bernoulli polynomials, but also we derive some novel formulas including the Stirling numbers and the array polynomials.

Finally, we give conclusion section of this paper.

2 A class of \mathbb{Q} -linear operator on the polynomials ring $\mathbb{Q}[x]$ and their applications

In this section, we define not only the following \mathbb{Q} -linear operator on the polynomials ring $\mathbb{Q}[x]$, but also construct its matrix representation with respect to the basis of $\mathbb{Q}[x]$: $\{1, x, x^2, x^3, \dots\}$. By using this operator and its matrix, we derive some new classes of special polynomials involving the Bernoulli polynomials, the Bell polynomials etc.

Let

$$\mathcal{E} : \mathbb{Q}[x] \rightarrow \mathbb{Q}[x].$$

Let $g(x)$ be integrable function. Then, we define the following linear operator on the polynomials ring $\mathbb{Q}[x]$:

$$\mathcal{E}[g(x)] = \int_{ax+b}^{cx+d} g(u)du, \quad (2.1)$$

where

$$g(x) \in \mathbb{Q}[x],$$

$a, b, m, j \in \mathbb{R}$.

Substituting $g(x) = x^n$ into (2.1), we obtain

$$\mathcal{E}[x^n; a, b, c, d] = \int_{ax+b}^{cx+d} u^n du = \frac{(cx+d)^{n+1} - (ax+b)^{n+1}}{n+1}. \quad (2.2)$$

Remark 1. Substituting $a = c = d = 1$ and $b = 0$ into (2.2), $\mathcal{E}[x^n; 1, 0, 1, 1]$ reduces to \mathbb{Q} -linear map, which is given by Arakawa et al. [1, p. 55].

Using (2.2), we give the following formulas for any polynomials

$$P_n(x) = \sum_{j=0}^n \alpha_j x^j$$

on the ring $\mathbb{Q}[x]$ under the linear operation \mathcal{E} :

$$\begin{aligned}
& \mathcal{E}[P_n(x); a, b, c, d] \tag{2.3} \\
&= \sum_{j=0}^n \binom{j+1}{0} \frac{\alpha_j}{j+1} (d^{j+1} - b^{j+1}) + \sum_{j=0}^n \binom{j+1}{1} \frac{\alpha_j}{j+1} (cd^j - ab^j)x \\
&+ \sum_{j=1}^n \binom{j+1}{2} \frac{\alpha_j}{j+1} (c^2 d^{j-1} - a^2 b^{j-1})x^2 + \sum_{j=2}^n \binom{j+1}{3} \frac{\alpha_j}{j+1} (c^3 d^{j-2} - a^3 b^{j-2})x^3 \\
&+ \sum_{j=3}^n \binom{j+1}{4} \frac{\alpha_j}{j+1} (c^4 d^{j-3} - a^4 b^{j-3})x^4 + \dots \\
&+ \sum_{j=k}^n \binom{j+1}{k-1} \frac{\alpha_j}{j+1} (c^{k+1} d^{j-k} - a^{k+1} b^{j-k})x^{k+1} + \\
&\dots + \alpha_{n-1} \frac{\binom{n}{n}}{n} (c^n d - a^n b)x^n + \alpha_n \frac{\binom{n+1}{n+1}}{n+1} (c^{n+1} - a^{n+1})x^{n+1}.
\end{aligned}$$

With the aid of (2.3), $(n+1) \times (n+2)$ matrix representation of the linear operation $\mathcal{E}[P_n(x); a, b, c, d]$ with respect to basis of $\{1, x, x^2, x^3, \dots\}$ is given by the following theorem:

Theorem 1.

$$M_{\mathcal{E}}(P_n(x); a, b, c, d) = \tag{2.4}$$

$$\begin{bmatrix}
\frac{\binom{1}{0}(d-b)}{1} & \frac{\binom{1}{1}(c-a)}{1} & 0 & 0 & \dots & 0 \\
\frac{\binom{2}{0}(d^2-b^2)}{2} & \frac{\binom{2}{1}(cd-ab)}{2} & \frac{\binom{2}{2}(c^2-a^2)}{2} & 0 & \dots & 0 \\
\frac{\binom{3}{0}(d^3-b^3)}{3} & \frac{\binom{3}{1}(cd^2-ab^2)}{3} & \frac{\binom{3}{2}(c^2d-a^2b)}{3} & \frac{\binom{3}{3}(c^3-a^3)}{3} & \dots & 0 \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\frac{\binom{k+1}{0}(d^{k+1}-b^{k+1})}{k+1} & \frac{\binom{k+1}{1}(cd^k-ab^k)}{k+1} & \frac{\binom{k+1}{2}(c^2d^{k-1}-a^2b^{k-1})}{k+1} & \frac{\binom{k+1}{3}(c^3d^{k-2}-a^3b^{k-2})}{k+1} & \dots & 0 \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
\frac{\binom{n+1}{0}(d^{n+1}-b^{n+1})}{n+1} & \frac{\binom{n+1}{1}(cd^n-ab^n)}{n+1} & \frac{\binom{n+1}{2}(c^2d^{n-1}-a^2b^{n-1})}{n+1} & \frac{\binom{n+1}{3}(c^3d^{n-2}-a^3b^{n-2})}{n+1} & \dots & \frac{\binom{n+1}{n+1}(c^{n+1}-a^{n+1})}{n+1}
\end{bmatrix}$$

If $c = a$, (2.4) is a square matrix and $\det(M_{\mathcal{E}}(P_n(x); a, b, a, d)) := |M_{\mathcal{E}}| \neq 0$. Thus we define the following a new family of polynomials:

Definition 1. We define $Q_n(x)$ polynomial sequences with respect to basis of $\mathbb{Q}[x]: \{1, x, x^2, x^3, \dots\}$:

$$\begin{bmatrix} Q_0(x) \\ Q_1(x) \\ Q_2(x) \\ \vdots \\ Q_{n-1}(x) \end{bmatrix} = \left((M_{\mathcal{E}}(P_n(x); a, b, a, d))^{-1} \right) \begin{bmatrix} 1 \\ x \\ x^2 \\ \vdots \\ x^{n-1} \end{bmatrix}, \tag{2.5}$$

where A^{-1} denotes inverse of the matrix A .

Example 1. Putting $n = 2$ in (2.5), we get

$$M_{\mathcal{E}}(P_2(x); a, b, c, d) = \begin{bmatrix} d-b & c-a & 0 \\ \frac{d^2-b^2}{2} & cd-ab & c^2-a^2 \end{bmatrix},$$

$$\begin{aligned} |M_{\mathcal{E}}(P_2(x); a, b, a, d)| &= a(b-d)^2 \\ &= ab^2 - 2abd + ad^2 \end{aligned}$$

and

$$(M_{\mathcal{E}}(P_2(x); a, b, a, d))^{-1} = \begin{bmatrix} \frac{-1}{b-d} & 0 \\ \frac{(b+d)}{2(ab-ad)} & \frac{-1}{ab-ad} \end{bmatrix}.$$

We have

$$\begin{bmatrix} Q_0(x) \\ Q_1(x) \end{bmatrix} = \begin{bmatrix} \frac{-1}{b-d} & 0 \\ \frac{(b+d)}{2a(b-d)} & \frac{-1}{a(b-d)} \end{bmatrix} \begin{bmatrix} 1 \\ x \end{bmatrix}.$$

From the above equation, we get the following polynomials:

$$\begin{aligned} Q_0(x) &= -\frac{1}{b-d} \\ Q_1(x) &= \frac{(b+d)}{2a(b-d)} - \frac{1}{a(b-d)}x. \end{aligned}$$

Example 2. Putting $n = 3$ in (2.5), we get

$$M_{\mathcal{E}}(P_3(x); a, b, c, d) = \begin{bmatrix} (d-b) & (c-a) & 0 & 0 \\ \frac{(d^2-b^2)}{2} & (cd-ab) & \frac{(c^2-a^2)}{2} & 0 \\ \frac{(d^3-b^3)}{3} & (cd^2-ab^2) & (c^2d-a^2b) & \frac{(c^3-a^3)}{3} \end{bmatrix}.$$

By using inverse matrix method we get

$$(M_{\mathcal{E}}(P_3(x); a, b, a, d))^{-1} = \begin{bmatrix} \frac{-1}{b-d} & 0 & 0 \\ \frac{(b+d)}{2a(b-d)} & \frac{-1}{a(b-d)} & 0 \\ -\frac{(b^2+d^2+\frac{2bd}{3})}{a^2(b-d)} & \frac{(b+d)}{a^2(b-d)} & \frac{-1}{a^2(b-d)} \end{bmatrix}.$$

We have

$$\begin{bmatrix} Q_0(x) \\ Q_1(x) \\ Q_2(x) \end{bmatrix} = \begin{bmatrix} \frac{-1}{b-d} & 0 & 0 \\ \frac{(b+d)}{2a(b-d)} & \frac{-1}{a(b-d)} & 0 \\ -\frac{(b^2+d^2+\frac{2bd}{3})}{a^2(b-d)} & \frac{(b+d)}{a^2(b-d)} & \frac{-1}{a^2(b-d)} \end{bmatrix} \begin{bmatrix} 1 \\ x \\ x^2 \end{bmatrix}.$$

Therefore from the above, we get the following polynomials:

$$\begin{aligned} Q_0(x) &= \frac{-1}{b-d} \\ Q_1(x) &= \frac{(b+d)}{2a(b-d)} - \frac{1}{a(b-d)}x \\ Q_2(x) &= \frac{-\left(\frac{b^2+d^2}{6} + \frac{2bd}{3}\right)}{a^2(b-d)} + \frac{(b+d)}{a^2(b-d)}x + \frac{-1}{a^2(b-d)}x^2. \end{aligned}$$

3 Applications of the linear transformation $\mathcal{E} [P_n(x); a, b, c, d]$

In this section, we give some special values of the linear transformation $\mathcal{E} [P_n(x); a, b, c, d]$. By applying Cayley–Hamilton theorem, we also give inverse matrix formulas involving the Bell polynomials. By the aid of the inverse matrix formula, we give matrix representation for the Bernoulli polynomials.

Substituting $a = c$ into (2.3), we get

$$\begin{aligned} &\mathcal{E} [P_n(x); a, b, a, d] \\ &= \sum_{j=0}^n \binom{j+1}{0} \frac{\alpha_j}{j+1} (d^{j+1} - b^{j+1}) + a \sum_{j=0}^n \binom{j+1}{1} \frac{\alpha_j}{j+1} (d^j - b^j)x + \\ &\quad + a^2 \sum_{j=1}^n \binom{j+1}{2} \frac{\alpha_j}{j+1} (d^{j-1} - b^{j-1})x^2 + a^3 \sum_{j=2}^n \binom{j+1}{3} \frac{\alpha_j}{j+1} (d^{j-2} - b^{j-2})x^3 \\ &\quad + \dots + a^n \alpha_{n-1} \frac{\binom{n+1}{n}}{n+1} (d-b)x^n. \end{aligned}$$

Substituting $a = c$ and $d = b + 1$ into (2.3), we also get

$$\begin{aligned} &\mathcal{E} [P_n(x); a, b, a, b+1] \tag{3.1} \\ &= \sum_{j=0}^n \binom{j+1}{0} \frac{\alpha_j}{j+1} (((b+1)^{j+1} - b^{j+1}) + a(j+1)((b+1)^j - b^j)x) \\ &\quad + a^2 \sum_{j=1}^n \binom{j+1}{2} \frac{\alpha_j}{j+1} ((b+1)^{j-1} - b^{j-1})x^2 + \dots \\ &\quad + a^n \sum_{j=n-1}^n \binom{j+1}{n} \frac{\alpha_j}{j+1} ((b+1)^{j-n+1} - b^{j-n+1})x^n. \tag{3.2} \end{aligned}$$

We assume that $j - n + 1 \geq 0$, otherwise when $j - n + 1 < 0$, these power's numbers omitted in the related sums.

With the aid of (3.1), $n + 1 \times n + 1$ matrix representation of the linear operation $\mathcal{E} [P_n(x); a, b, a, b + 1]$ with respect to basis of $\{1, x, x^2, x^3, \dots\}$ is given by the following corollary:

Corollary 1.

$$M_{\mathcal{E}}(P_n(x); a, b, a, b+1) = \tag{3.3}$$

$$\begin{bmatrix} 1 & 0 & 0 & \dots & 0 \\ \frac{\binom{2}{0}}{2}((b+1)^2 - b^2) & a & 0 & \dots & 0 \\ \frac{\binom{3}{0}}{3}((b+1)^3 - b^3) & \frac{\binom{3}{1}}{3}a((b+1)^2 - b^2) & a^2 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \dots \\ \frac{\binom{n+1}{0}}{n+1}((b+1)^{n+1} - b^{n+1}) & \frac{\binom{n+1}{1}}{n}a((b+1)^n - b^n) & \frac{\binom{n+1}{2}}{n}a^2((b+1)^{n-1} - b^{n-1}) & \dots & a^n \end{bmatrix}.$$

By applying Cayley–Hamilton theorem to (3.3) and using inverse matrix formula, given in formula (*cf.* [12, Eq. (B.10) and Eq. (B.11)]), we show that inverse matrix of the matrix $M_{\mathcal{E}}(P_n(x); a, b, a, d)$ is given by the following theorem.

Lemma 1. *If $c = a$, $M_{\mathcal{E}}(P_n(x); a, b, c, d)$ is a diagonal matrix, then we have*

$$\begin{aligned} & (M_{\mathcal{E}}(P_n(x); a, b, a, d))^{-1} \tag{3.4} \\ &= \frac{1}{|M_{\mathcal{E}}|} \sum_{j=0}^n M_{\mathcal{E}}^j(P_n(x); a, b, a, d) \\ & \times \sum_{a_1, a_2, \dots, a_n} \prod_{v=1}^n \frac{(-1)^{a_v+1}}{v^{a_v} a_v!} (\text{tr}(M_{\mathcal{E}}^v(P_n(x); a, b, a, d)))^{a_v} \end{aligned}$$

$n+1$ is a dimension of matrix $M_{\mathcal{E}}(P_n(x); a, b, a, d)$, $\text{tr}\{M_{\mathcal{E}}(P_n(x); a, b, a, d)\}$ is the trace of main diagonal matrix $M_{\mathcal{E}}(P_n(x); a, b, a, d)$, and also

$$\sum_{a_1, a_2, a_3, \dots, a_n}$$

means that this sum is taken over j and the sets of all $a_v \geq 0$ satisfying the following equation;

$$j + \sum_{m=0}^n m \cdot a_m = n.$$

Lemma 2. *If $c = a$, $M_{\mathcal{E}}(P_n(x); a, b, c, d)$ is a diagonal matrix, then we have*

$$\begin{aligned} & (M_{\mathcal{E}}(P_n(x); a, b, a, d))^{-1} \tag{3.5} \\ &= \frac{1}{|M_{\mathcal{E}}|} \sum_{j=1}^n M_{\mathcal{E}}^{j-1}(P_n(x); a, b, a, d) \frac{(-1)^{n-1}}{(n-j)!} B_{n-j}(w_1, w_2, \dots, w_{n-j}), \end{aligned}$$

where

$$w_k = -(k-1)! \text{tr}\{M_{\mathcal{E}}^k(P_n(x); a, b, a, d)\}$$

and $B_n(w_1, w_2, \dots, w_n)$ denotes the n th complete exponential Bell polynomials, defined by (see [2, 5])

$$B_n(w_1, w_2, \dots, w_n) = n! \sum_{n=1k_1+2k_2+\dots+nk_n} \prod_{j=1}^n \frac{w_j^{k_j}}{(j!)^{k_j} k_j!}.$$

By using (3.4) and $\mathcal{E}^{-1}[x^n; a, b, a, b+1]$, we find another family of polynomials by the following theorem:

Theorem 2.

$$\begin{bmatrix} H_0(x) \\ H_1(x) \\ H_2(x) \\ \vdots \\ H_{n-1}(x) \end{bmatrix} = \left((M_{\mathcal{E}}(P_n(x); a, b, a, b+1))^{-1} \right) \begin{bmatrix} 1 \\ x \\ x^2 \\ \vdots \\ x^{n-1} \end{bmatrix}. \quad (3.6)$$

When $a = c = d = 1$ and $b = 0$, the Bernoulli polynomials, which are also given by the following corollary:

Corollary 2.

$$\begin{bmatrix} B_0(x) \\ B_1(x) \\ B_2(x) \\ \vdots \\ B_{n-1}(x) \end{bmatrix} = \left((M_{\mathcal{E}}(P_n(x); 1, 0, 1, 1))^{-1} \right) \begin{bmatrix} 1 \\ x \\ x^2 \\ \vdots \\ x^{n-1} \end{bmatrix}, \quad (3.7)$$

where

$$\begin{aligned} & (M_{\mathcal{E}}(P_n(x); 1, 0, 1, 1))^{-1} \\ &= \sum_{j=0}^n \frac{M_{\mathcal{E}}^j(P_n(x); 1, 0, 1, 1)}{\det(M_{\mathcal{E}}(P_n(x); 1, 0, 1, 1))} \\ & \quad \times \sum_{a_1, a_2, \dots, a_n} \prod_{v=1}^n \frac{(-1)^{a_v+1}}{v^{a_v} a_v!} (tr(M_{\mathcal{E}}^v(P_n(x); 1, 0, 1, 1)^{a_v})). \end{aligned}$$

Remark 2. When $a = c = d = 1$ and $b = 0$, with the aid of (3.7), we see that

$$B_n(x) = \mathcal{E}^{-1}[x^n; 1, 0, 1, 1]$$

(cf. [1, p. 55]).

4 Derivative formula for linear transformation

$$\mathcal{E}[P_n(x); a, b, c, d]$$

In this section, we give derivative formula of the linear transformation $\mathcal{E}[P_n(x); a, b, c, d]$

Taking derivative of $\mathcal{E}[P_n(x); a, b, c, d]$ with respect to x , we arrive at the following derivative formula:

$$\begin{aligned}
& \frac{d}{dx} \{ \mathcal{E}[P_n(x); a, b, c, d] \} \\
&= \sum_{j=0}^n \binom{j+1}{1} \frac{\alpha_j (cd^j - b^j a)}{j+1} + 2 \sum_{j=1}^n \binom{j+1}{2} \frac{(c^2 d^{j-1} - a^2 b^{j-1}) \alpha_j}{j+1} x \\
&+ 3 \sum_{j=2}^n \binom{j+1}{3} \frac{(c^3 d^{j-2} - a^3 b^{j-2}) \alpha_j}{j+1} x^2 + 4 \sum_{j=3}^n \binom{j+1}{4} \frac{(c^4 d^{j-3} - a^4 b^{j-3}) \alpha_j}{j+1} x^3 \\
&+ \dots + \sum_{j=k}^n \binom{j+1}{k-1} \frac{(k+1)(c^{k+1} d^{j-k} - a^{k+1} b^{j-k}) \alpha_j}{j+1} x^k \\
&+ \dots + (c^n d - a^n b) \alpha_{n-1} x^{n-1} + (c^{n+1} - a^{n+1}) \alpha_n x^n.
\end{aligned}$$

Putting $a = c = d = 1$ and $b = 0$ in the above equation, we have

$$\begin{aligned}
& \frac{d}{dx} \{ \mathcal{E}[P_n(x); 1, 0, 1, 1] \} \\
&= \sum_{j=0}^n \binom{j+1}{1} \frac{\alpha_j}{j+1} + 2 \sum_{j=1}^n \binom{j+1}{2} \frac{\alpha_j}{j+1} x + 3 \sum_{j=2}^n \binom{j+1}{3} \frac{\alpha_j}{j+1} x^2 \\
&+ 4 \sum_{j=3}^n \binom{j+1}{4} \frac{\alpha_j}{j+1} x^3 + \dots + \sum_{j=k}^n \binom{j+1}{k-1} \frac{(k+1) \alpha_j}{j+1} x^k + \dots + \alpha_{n-1} x^{n-1}.
\end{aligned}$$

Combining (3.7) with the above equation and its matrix, we have the following well-known result for the Bernoulli polynomials:

$$\frac{d}{dx} \{ B_n(x) \} = n B_{n-1}(x).$$

5 A new family of polynomials and its matrix

In this section, we define a new family of polynomials. By using these polynomials, we also define a matrix representation of the coefficients of these polynomials. We show that special values of these matrix with its inverse produce the Bernoulli numbers.

By using (2.2), we define the following polynomial:

$$\begin{aligned}
Y_n(x; a, b, c, d) &= (cx + d)^{n+1} - (ax + b)^{n+1} \\
&= \sum_{j=0}^{n+1} \binom{n+1}{j} (c^j d^{n+1-j} - a^j b^{n+1-j}) x^j.
\end{aligned}$$

Some values of $Y_n(x; a, b, c, d)$ are given by

$$\begin{aligned}
Y_0(x; a, b, c, d) &= (c - a)x + d - b \\
Y_1(x; a, b, c, d) &= (c^2 - a^2)x^2 + 2(cd - ab)x + (d^2 - b^2) \\
Y_2(x; a, b, c, d) &= (c^3 - a^3)x^3 + 3(c^2d - a^2b)x^2 + 3(cd^2 - ab^2)x + (d^3 - b^3) \\
&\vdots \\
Y_n(x; a, b, c, d) &= (c^{n+1} - a^{n+1})x^{n+1} + \binom{n+1}{n}(c^nd - a^nb)x^n + \dots \\
&\quad + (d^{n+1} - b^{n+1}).
\end{aligned}$$

We define matrix representation of the aid of coefficients of the above polynomials as follows:

$$M(Y_n(x; a, b, c, d)) = \tag{5.1}$$

$$\begin{bmatrix}
d - b & c - a & 0 & 0 & \dots & 0 \\
d^2 - b^2 & 2(cd - ab) & c^2 - a^2 & 0 & \dots & 0 \\
d^3 - b^3 & 3(cd^2 - ab^2) & 3(c^2d - a^2b) & c^3 - a^3 & \dots & 0 \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
d^{n+1} - b^{n+1} & \binom{n+1}{1}(cd^n - ab^n) & \binom{n+1}{2}(c^2d^{n-1} - a^2b^{n-1}) & \binom{n+1}{3}(c^3d^{n-2} - a^3b^{n-2}) & \dots & c^{n+1} - a^{n+1}
\end{bmatrix}.$$

Thus we get

$$\begin{bmatrix}
Y_0(x; a, b, a, d) \\
Y_1(x; a, b, a, d) \\
Y_2(x; a, b, a, d) \\
\vdots \\
Y_{n-1}(x; a, b, a, d)
\end{bmatrix} = M^{-1}(Y_n(x; a, b, a, d)) \begin{bmatrix}
1 \\
x \\
x^2 \\
\vdots \\
x^{n-1}
\end{bmatrix}.$$

The first row of the inverse of the matrix $M(Y_n(x; a, b, a, d))$ given by equation (5.1) can also be considered as the generating function for a special family of numbers, including Bernoulli numbers maybe other certain family of special numbers. That is, substituting $a = c = d = 1$ and $b = 0$ into inverse matrix $M^{-1}(Y_n(x; 1, 0, 1, 1))$, given by equation (5.1), all entries of the matrix $M^{-1}(Y_n(x; 1, 0, 1, 1))$ are reduced to the Bernoulli numbers, respectively.

Some well-known examples are given as follows:

$$M(Y_6(x; 1, 0, 1, 1)) = \begin{bmatrix}
1 & 0 & 0 & 0 & 0 & 0 & 0 \\
1 & 2 & 0 & 0 & 0 & 0 & 0 \\
1 & 3 & 3 & 0 & 0 & 0 & 0 \\
1 & 4 & 6 & 4 & 0 & 0 & 0 \\
1 & 5 & 10 & 10 & 5 & 0 & 0 \\
1 & 6 & 15 & 20 & 15 & 6 & 0 \\
1 & 7 & 21 & 35 & 35 & 21 & 7
\end{bmatrix}.$$

By using (3.4) inverse matrix method we get

$$M^{-1}(Y_6(x; 1, 0, 1, 1)) = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -\frac{1}{2} & \frac{1}{2} & 0 & 0 & 0 & 0 & 0 \\ \frac{1}{6} & -\frac{1}{2} & \frac{1}{3} & 0 & 0 & 0 & 0 \\ 0 & \frac{1}{4} & -\frac{1}{2} & \frac{1}{4} & 0 & 0 & 0 \\ -\frac{1}{30} & 0 & \frac{1}{3} & -\frac{1}{2} & \frac{1}{5} & 0 & 0 \\ 0 & -\frac{1}{12} & 0 & \frac{5}{12} & -\frac{1}{2} & \frac{1}{6} & 0 \\ \frac{1}{42} & 0 & -\frac{1}{6} & 0 & \frac{1}{2} & -\frac{1}{2} & \frac{1}{7} \end{bmatrix},$$

where we obtained Bernoulli numbers up to $n = 6$ in the first column of the $M(Y_6(x; a, b, a, d))$ matrix (cf. for detail, A027642, see also [9]). On the other hand, other columns or rows of the above matrix also represent different number families. These number sequences may also be in the class of known other families of special number. For this, it is recommended to examine the *oeis* page (<https://oeis.org/>).

6 Applications of the Laplace transform to generating function for the Bernoulli polynomials

In this section, applications of the Laplace transform and the Mellin transformation to (1.2), we give a relation between the Bernoulli polynomials and the Hurwitz zeta function. We also give infinite series representation involving the Bernoulli polynomials.

Let $x \in \mathbb{C}$ with $x = a + ib$ and $\bar{x} = a - ib$ with $a > 0$ and $b > 0$. We set

$$F(-u, \bar{x}) = \sum_{n=0}^{\infty} (-1)^n B_n(a - ib) \frac{u^n}{n!}.$$

Therefore

$$\frac{ue^{-ua}}{1 - e^{-u}} = e^{-uib} \sum_{n=0}^{\infty} (-1)^n B_n(a - ib) \frac{u^n}{n!}. \quad (6.1)$$

Eq. (6.1) can also be written as

$$e^{-ua} \sum_{n=1}^{\infty} \frac{(1 - e^{-u})^{n-1}}{n} = e^{-uib} \sum_{n=0}^{\infty} (-1)^n B_n(a - ib) \frac{u^n}{n!}. \quad (6.2)$$

Combining the above equation with (1.4), we get

$$\begin{aligned} & \sum_{m=0}^{\infty} \sum_{v=0}^m \binom{m}{v} \sum_{n=1}^v \frac{(-1)^{m+n-1} (n-1)! a^{m-v} S_2(v, n-1) u^m}{n m!} \\ &= \sum_{m=0}^{\infty} (-1)^m \sum_{v=0}^m \binom{m}{v} (ib)^{m-v} B_v(a - ib) \frac{u^m}{m!}. \end{aligned}$$

Substituting the following well-known formula into the right-hand side of the above equation

$$\sum_{v=0}^m \binom{m}{v} x^{m-v} B_v(y) = B_m(x+y),$$

we obtain

$$\begin{aligned} & \sum_{m=0}^{\infty} \sum_{v=0}^m \binom{m}{v} \sum_{n=1}^v \frac{(-1)^{m+n-1} (n-1)! a^{m-v} S_2(v, n-1) u^m}{n m!} \\ &= \sum_{m=0}^{\infty} (-1)^m B_m(a) \frac{u^m}{m!}. \end{aligned}$$

By comparing the coefficients $\frac{u^m}{m!}$ on both sides of the above equation, we arrive at the following theorem:

Theorem 3. *Let $m \in \mathbb{N}$. Then we have*

$$B_m(a) = \sum_{v=1}^m \binom{m}{v} \sum_{n=1}^v \frac{(-1)^{n-1} (n-1)! a^{m-v} S_2(v, n-1)}{n}. \quad (6.3)$$

By combining (1.5) with (6.2), by the same method of Cakic and Milovanovic [3], Chang and Ha [4], Simsek[17], we have

$$\sum_{m=0}^{\infty} \sum_{n=1}^{\infty} \frac{(-1)^{n-1} (n-1)!}{n} S_{n-1}^m(a) \frac{(-u)^m}{m!} = \sum_{m=0}^{\infty} \sum_{v=0}^m \binom{m}{v} (ib)^{m-v} B_v(a-ib) \frac{(-u)^m}{m!}.$$

After some calculation, we have the following result, which also proved in [4]:

$$\sum_{n=1}^m \frac{(-1)^{n-1} (n-1)!}{n} S_{n-1}^m(x) = \sum_{v=0}^m \binom{m}{v} (ib)^{m-v} B_v(a-ib). \quad (6.4)$$

Combining the above equation with (6.3) yields

$$\sum_{v=1}^m \sum_{n=1}^v \frac{(-1)^{n-1} \binom{m}{v} (n-1)! a^{m-v} S_2(v, n-1)}{n} = \sum_{n=0}^m \frac{(-1)^n n!}{n+1} S_n^m(x). \quad (6.5)$$

Combining (6.4) and (6.5) with

$$\frac{(-1)^n n!}{n+1} = \sum_{v=0}^n S_1(n, v) B_v, \quad (6.6)$$

(cf. [5, 10, 16, 19]), we arrive at the following theorem:

Theorem 4. *Let $m \in \mathbb{N}_0$. Then we have*

$$B_m(a) = \sum_{n=0}^m \sum_{v=0}^n S_1(n, v) B_v S_n^m(a) \quad (6.7)$$

and

$$\sum_{v=0}^m \binom{m}{v} (ib)^{m-v} B_v(a-ib) = \sum_{n=0}^m \sum_{v=0}^n S_1(n, v) B_v S_n^m(a).$$

Here, we note that different proof of the equation (6.5) was also given by Chang and Ha [4].

By applying the Laplace transform to generating functions involving Bernoulli polynomials, recently many interesting studies have been published (*cf.* [7, 8, 11, 21]). Therefore, by using (6.1), we have

$$\sum_{n=0}^{\infty} u e^{-u(a+n)} = e^{-uib} \sum_{n=0}^{\infty} (-1)^n B_n(a-ib) \frac{u^n}{n!}.$$

Integrate equation the above equation with respect to u from 1 to ∞ , we get

$$\sum_{n=0}^{\infty} \int_0^{\infty} u e^{-u(a+n)} du = \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} B_n(a-ib) \int_0^{\infty} u^n e^{-ibu} du \quad (6.8)$$

with $a > 0$. Applying the Laplace transform of the function $g(u) = u^n$:

$$\mathcal{L}\{g(u)\} = \frac{n!}{y^{n+1}}$$

(where $y > 0$) to the both sides of (6.8), we obtain the following theorem:

Theorem 5. *Let $b > 0$ and $a > 0$. Then we have*

$$\zeta(2, a) = \sum_{n=0}^{\infty} \frac{i^{n-1} B_n(a-ib)}{b^{n+1}}. \quad (6.9)$$

By using (6.1), we have

$$u e^{-au} = \left(e^{-ibu} - e^{-(1+ib)u} \right) \sum_{n=0}^{\infty} (-1)^n B_n(a-ib) \frac{u^n}{n!}.$$

Applying the Laplace transform to the above equation, for $a > 0$ and $b > 0$, we get

$$\begin{aligned} \int_0^{\infty} u e^{-au} du &= \sum_{n=0}^{\infty} (-1)^n B_n(a-ib) \frac{1}{n!} \int_0^{\infty} u^n e^{-ibu} du \\ &\quad - \sum_{n=0}^{\infty} (-1)^n B_n(a-ib) \frac{1}{n!} \int_0^{\infty} u^n e^{-(1+ib)u} du. \end{aligned}$$

After some calculations, for $b > 1$, we arrive at the following theorem:

Theorem 6. *Let $a > 0$ and $b > 1$. Then we have*

$$\sum_{n=0}^{\infty} \sum_{j=0}^n (-1)^n \binom{n}{j} \frac{\left((1+ib)^{n+1} - (ib)^{n+1} \right) (a-ib)^{n-j} B_j}{(ib-b^2)^{n+1}} = \frac{1}{a^2}. \quad (6.10)$$

Remark 3. Putting $n = 1$ into equation (17) in [7], a relationship can be established with equation (6.10).

Substituting $a = 1$ into (6.9), and combining with the following known formula

$$\zeta(2) := \zeta(2, 1) = \sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6},$$

we get the following corollary:

Corollary 3. Let $b > 1$ and $a > 0$. Then we have

$$\sum_{n=0}^{\infty} \frac{i^{n-1} B_n(a - ib)}{b^{n+1}} = \frac{\pi^2}{6}.$$

Replacing t, x by $-t, y - x$ in (1.2) yields

$$F(-t, y - x) = \frac{-te^{-t(y-x)}}{e^{-t} - 1} = \sum_{n=0}^{\infty} (-1)^n B_n(y - x) \frac{t^n}{n!}.$$

Taking the partial derivative k times with respect to y in the above equation, we get

$$\frac{\partial^k}{\partial y^k} \{F(-t, y - x)\} = \frac{(-1)^{k+1} t^{k+1} e^{-t(y-x)}}{e^{-t} - 1}.$$

After some calculations, we obtain

$$\frac{(-1)^{k+1} t^{k+1} e^{-ty}}{e^{-t} - 1} = e^{-tx} \sum_{n=0}^{\infty} \frac{(-1)^n \frac{\partial^k}{\partial y^k} \{B_n(y - x)\}}{n!} t^n.$$

Integrate equation the above equation with respect to u from 1 to ∞ , we get

$$(-1)^{k+1} \int_0^{\infty} \frac{t^{k+1} e^{-ty}}{e^{-t} - 1} dt = \sum_{n=0}^{\infty} \frac{(-1)^n \frac{\partial^k}{\partial y^k} \{B_n(y - x)\}}{n!} \int_0^{\infty} t^n e^{-tx} dt,$$

which yields

$$(-1)^{k+1} (k+1)! \sum_{n=0}^{\infty} \frac{1}{(y+n)^{k+2}} = \sum_{n=0}^{\infty} \frac{(-1)^n}{x^{n+1}} \frac{\partial^k}{\partial y^k} \{B_n(y - x)\}.$$

After some calculations, we arrive at the following theorem:

Theorem 7. Let $k \in \mathbb{N}_0$. Let $y > 0$. Then we have

$$\zeta(k+2, y) = \frac{(-1)^{k+1}}{(k+1)!} \sum_{n=0}^{\infty} \frac{(-1)^n}{x^{n+1}} \frac{\partial^k}{\partial y^k} \{B_n(y - x)\}. \quad (6.11)$$

Substituting $k = 0$ into (6.11), we arrive at (6.9).

Let the ratio of two convergent series be given such that when the first term of the series in the denominator is different from zero, using (1.1), we have the following result:

$$\frac{1 + 0t + 0t^2 + \dots}{1 + \frac{1}{2!}t + \frac{1}{3!}t^2 + \frac{1}{4!}t^3 + \dots} = \sum_{n=0}^{\infty} B_n \frac{t^n}{n!}.$$

From the above equation, we get

$$1 = \left(1 + \frac{1}{2!}t + \frac{1}{3!}t^2 + \frac{1}{4!}t^3 + \dots\right) \left(B_0 + \frac{B_1}{1!}t + \frac{B_2}{2!}t^2 + \frac{B_3}{3!}t^3 + \dots\right).$$

Using the Cauchy product rule on the right-hand side of the above equation yields an infinite system of linear equations with unknown B_k . So, this system is of special form since every $k \in \mathbb{N}_0$ the first $k + 1$ equations contain only the first $k + 1$ unknown B_k . Therefore, the solution of the system of linear equations is given by the following determinant, which is calculated the Bernoulli numbers outside the influence of their generating function:

$$B_n = (-1)^n n! \begin{vmatrix} \frac{1}{2!} & 1 & 0 & 0 & 0 & \dots & 0 \\ \frac{1}{3!} & \frac{1}{2!} & 1 & 0 & 0 & \dots & 0 \\ \frac{1}{4!} & \frac{1}{3!} & \frac{1}{2!} & 1 & 0 & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & 0 \\ \frac{1}{(n+1)!} & \frac{1}{n!} & \frac{1}{(n-1)!} & \frac{1}{(n-2)!} & \frac{1}{(n-3)!} & \dots & \frac{1}{2!} \end{vmatrix}$$

(cf. [14, p. 149], [6]).

7 Conclusion

In this paper, we defined \mathbb{Q} -linear operator on the polynomials ring $\mathbb{Q}[x]$. By using this operator, we gave its matrix representation with respect to the basis of $\mathbb{Q}[x]$: $\{1, x, x^2, x^3, \dots\}$. Using matrix representation, we obtained some new classes of special polynomials involving Bernoulli polynomials and Bell polynomials. We also defined other linear transformations. By applying Cayley–Hamilton theorem, we gave their inverse matrix formulas involving the Bell polynomials. Using this inverse matrix formula, we gave matrix representation for the Bernoulli polynomials. Moreover, applying the Laplace transform to generating function for the Bernoulli polynomials, we obtained both infinite series representation for the Bernoulli polynomials and many new formulas associated with the Stirling numbers and the array polynomials.

The results of this article may potentially be used both in applied sciences and in many branches of mathematics.

References

- [1] T. Arakawa, T. Ibukiyama, M. Kaneko, *Bernoulli Numbers and Zeta Functions*, Springer, 2014.

- [2] E. T. Bell, *Exponential polynomials*, Ann. of Math. 35 (2) (1934) 258–277. Doi:10.2307/1968431.
- [3] N. P. Cakic, G. V. Milovanovic, *On generalized Stirling numbers and polynomials*, Math. Balk. 18 (2004) 241–248.
- [4] C.H. Chang, C.W. Ha, *A multiplication theorem for the Lerch zeta function and explicit representations of the Bernoulli and Euler polynomials*, J. Math. Anal. Appl. 315 (2006) 758–767.
- [5] C.A. Charalambides, *Enumerative Combinatorics*, Chapman and Hall / CRC, New York, 2002.
- [6] H. Chen, *Bernoulli numbers via determinants*, International Journal of Mathematical Education in Science and Technology 34 (2) (2003) 291–297. Doi:10.1080/0020739031000158335.
- [7] D. Gun, Y. Simsek, *Some new identities and inequalities for Bernoulli polynomials and numbers of higher order related to the Stirling and Catalan numbers*, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM 114 (2020). <https://doi.org/10.1007/s13398-020-00899-z>.
- [8] D. Gross, J.F. Shortle, J.M. Thompson, C.M. Harris, *Fundamentals of Queueing Theory*, John Wiley and Sons Inc. Hoboken, New Jersey, 2008.
- [9] G. Helms, *Accessing Bernoulli-numbers by matrix-operations*, http://go.helms-net.de/math/pascal/bernoulli_en.pdf
- [10] D.S. Kim, T Kim, *Daehee numbers and polynomials*, Appl. Math. Sci. (Ruse) 7 (120) (2013) 5969–5976.
- [11] N. Kilar, Y. Simsek, H.M. Srivastava, *Recurrence relations, associated formulas, and combinatorial sums for some parametrically generalized polynomials arising from an analysis of the Laplace transform and generating functions*, Ramanujan J 61 (2023) 731–756. <https://doi.org/10.1007/s11139-022-00679-w>.
- [12] L.A. Kondratyuk, M.I. Krivoruchenko, *Superconducting quark matter in $SU(2)$ colour group*, Z. Phys. A - Hadrons and Nuclei 344 (1992) 99–115.
- [13] I. Kucukoglu, Y. Simsek, *New formulas and numbers arising from analyzing combinatorial numbers and polynomials*, Montes Taurus J. Pure Appl. Math. 3 (3) (2021) 238–259.
- [14] E. Pap, *Complex Analysis through Examples and Exercises*, Springer, Netherlands, 1999. Doi: 10.1007/978-94-017-1106-7.
- [15] E. Polat and Y. Simsek, *Notes on Bernoulli polynomials associated with linear map on the set of rational numbers*, In: Proceedings Book of the 13th Symposium on Generating Functions of Special Numbers and Polynomials and their Applications (GFSNP), ISBN: 978-625-00-1128-7, 2023, pp. 167–169.

- [16] J. Riordan, *Introduction to Combinatorial Analysis*, Princeton University Press, Princeton, 1958.
- [17] Y. Simsek, *Generating functions for generalized Stirling type numbers, Array type polynomials, Eulerian type polynomials and their applications*, Fixed Point Theory Appl. 2013 (87) (2013).
- [18] Y. Simsek, *Explicit formulas for p -adic integrals: Approach to p -adic distribution and some families of special numbers and polynomials*, Montes Taurus J. Pure Appl. Math. 1 (1) (2019) 1–76.
- [19] Y. Simsek, *Formulas for p -adic q -integrals including falling-rising factorials, combinatorial sums and special numbers*, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM 118 (2024). <https://doi.org/10.1007/s13398-024-01592-1>.
- [20] J.L. Steven, *Linear Algebra with Applications*, Pearson, 2015.
- [21] H.M. Srivastava, J. Choi, *Zeta and q -zeta Functions and Associated Series and Integrals*, Elsevier Science Publishers, Amsterdam, London and New York, 2012.
- [22] https://en.wikipedia.org/w/index.php?title=Invertible_matrix.
- [23] <https://math.stackexchange.com/questions/264820/a-relationship-between-matrices-bernoulli-polynomials-and-binomial-coefficient>.