
**NONCOMMUTATIVE
DONOHO-STARK-ELAD-BRUCKSTEIN-RICAUD-TORRÉSANI UNCERTAINTY
PRINCIPLE**

K. MAHESH KRISHNA

School of Mathematics and Natural Sciences
Chanakya University Global Campus
Haraluru Village, Near Kempe Gowda International Airport (BIAL)
Devanahalli Taluk, Bengaluru Rural District
Karnataka 562 110 India
Email: kmaheshak@gmail.com
Date: June 14, 2024

Abstract: Let $\{\tau_n\}_{n=1}^\infty$ and $\{\omega_m\}_{m=1}^\infty$ be two modular Parseval frames for a Hilbert C^* -module \mathcal{E} . Then for every $x \in \mathcal{E} \setminus \{0\}$, we show that

$$(1) \quad \|\theta_\tau x\|_0 \|\theta_\omega x\|_0 \geq \frac{1}{\sup_{n,m \in \mathbb{N}} \|\langle \tau_n, \omega_m \rangle\|^2}.$$

We call Inequality (1) as **Noncommutative Donoho-Stark-Elad-Bruckstein-Ricaud-Torrésani Uncertainty Principle**. Inequality (1) is the noncommutative analogue of breakthrough Ricaud-Torrésani uncertainty principle [*IEEE Trans. Inform. Theory*, 2013]. In particular, Inequality (1) extends Elad-Bruckstein uncertainty principle [*IEEE Trans. Inform. Theory*, 2002] and Donoho-Stark uncertainty principle [*SIAM J. Appl. Math.*, 1989].

Keywords: Uncertainty principle, Parseval frame, Hilbert C^* -module.

Mathematics Subject Classification (2020): 42C15, 46L08.

1. INTRODUCTION

In 1989, Donoho and Stark derived following uncertainty principle which is one of the greatest inequality of all time in both pure and applied Mathematics [3]. For $h \in \mathbb{C}^d$, let $\|h\|_0$ be the number of nonzero entries in h . Let $\widehat{\cdot} : \mathbb{C}^d \rightarrow \mathbb{C}^d$ be the Fourier transform defined by

$$\widehat{(a_j)_{j=0}^{d-1}} := \frac{1}{\sqrt{d}} \left(\sum_{j=0}^{d-1} a_j e^{-\frac{2\pi i j k}{d}} \right)_{k=0}^{d-1}, \quad \forall (a_j)_{j=0}^{d-1} \in \mathbb{C}^d.$$

Theorem 1.1. (Donoho-Stark Uncertainty Principle) [3, 12] For every $d \in \mathbb{N}$,

$$(2) \quad \left(\frac{\|h\|_0 + \|\widehat{h}\|_0}{2} \right)^2 \geq \|h\|_0 \|\widehat{h}\|_0 \geq d, \quad \forall h \in \mathbb{C}^d \setminus \{0\}.$$

By noting that Fourier transform is unitary and unitary operators are in one to one correspondence with orthonormal bases, in 2002, Elad and Bruckstein generalized Inequality (2) to arbitrary orthonormal bases [5]. To state the result we need some notations. Given a collection $\{\tau_j\}_{j=1}^n$ in a finite dimensional Hilbert space \mathcal{H} over \mathbb{K} (\mathbb{R} or \mathbb{C}), we define

$$\theta_\tau : \mathcal{H} \ni h \mapsto \theta_\tau h := (\langle h, \tau_j \rangle)_{j=1}^n \in \mathbb{K}^n.$$

Theorem 1.2. (Elad-Bruckstein Uncertainty Principle) [4, 5] Let $\{\tau_j\}_{j=1}^n, \{\omega_j\}_{j=1}^n$ be two orthonormal bases for a finite dimensional Hilbert space \mathcal{H} . Then

$$\left(\frac{\|\theta_\tau h\|_0 + \|\theta_\omega h\|_0}{2}\right)^2 \geq \|\theta_\tau h\|_0 \|\theta_\omega h\|_0 \geq \frac{1}{\max_{1 \leq j, k \leq n} |\langle \tau_j, \omega_k \rangle|^2}, \quad \forall h \in \mathcal{H} \setminus \{0\}.$$

In 2013, Ricaud and Torr esani showed that orthonormal bases in Theorem 1.2 can be improved to Parseval frames [9]. Recall that a collection $\{\tau_j\}_{j=1}^n$ in a finite dimensional Hilbert space \mathcal{H} is said to be a Parseval frame for \mathcal{H} [1] if

$$\|h\|^2 = \sum_{j=1}^n |\langle h, \tau_j \rangle|^2, \quad \forall h \in \mathcal{H}.$$

Theorem 1.3. (Ricaud-Torr esani Uncertainty Principle) [9] Let $\{\tau_j\}_{j=1}^n, \{\omega_j\}_{j=1}^n$ be two Parseval frames for a finite dimensional Hilbert space \mathcal{H} . Then

$$\left(\frac{\|\theta_\tau h\|_0 + \|\theta_\omega h\|_0}{2}\right)^2 \geq \|\theta_\tau h\|_0 \|\theta_\omega h\|_0 \geq \frac{1}{\max_{1 \leq j, k \leq n} |\langle \tau_j, \omega_k \rangle|^2}, \quad \forall h \in \mathcal{H} \setminus \{0\}.$$

The main purpose of this paper is to generalize and derive a noncommutative version of Theorem 1.3. For this we want generalization of Hilbert spaces known as Hilbert C*-modules. Hilbert C*-modules are first introduced by Kaplansky [7] for modules over commutative C*-algebras and later developed for modules over arbitrary C*-algebras by Paschke [8] and Rieffel [10].

Definition 1.4. [7, 8, 10] Let \mathcal{A} be a unital C*-algebra. A left module \mathcal{E} over \mathcal{A} is said to be a (left) Hilbert C*-module if there exists a map $\langle \cdot, \cdot \rangle : \mathcal{E} \times \mathcal{E} \rightarrow \mathcal{A}$ such that the following hold.

- (i) $\langle x, x \rangle \geq 0, \forall x \in \mathcal{E}$. If $x \in \mathcal{E}$ satisfies $\langle x, x \rangle = 0$, then $x = 0$.
- (ii) $\langle x + y, z \rangle = \langle x, z \rangle + \langle y, z \rangle, \forall x, y, z \in \mathcal{E}$.
- (iii) $\langle ax, y \rangle = a \langle x, y \rangle, \forall x, y \in \mathcal{E}, \forall a \in \mathcal{A}$.
- (iv) $\langle x, y \rangle = \langle y, x \rangle^*, \forall x, y \in \mathcal{E}$.
- (v) \mathcal{E} is complete w.r.t. the norm $\|x\| := \sqrt{\|\langle x, x \rangle\|}, \forall x \in \mathcal{E}$.

We are going to use the following inequality.

Lemma 1.5. [8] (Noncommutative Cauchy-Schwarz inequality) If \mathcal{E} is a Hilbert C*-module over \mathcal{A} , then

$$\langle x, y \rangle \langle y, x \rangle \leq \|\langle y, y \rangle\| \langle x, x \rangle, \quad \forall x, y \in \mathcal{E}.$$

Given a unital C*-algebra \mathcal{A} , define

$$\ell^2(\mathbb{N}, \mathcal{A}) := \left\{ \{a_n\}_{n=1}^\infty : a_n \in \mathcal{A}, \forall n \in \mathbb{N}, \sum_{n=1}^\infty a_n a_n^* \text{ converges in } \mathcal{A} \right\}.$$

Modular \mathcal{A} -inner product on $\ell^2(\mathbb{N}, \mathcal{A})$ is defined as

$$\langle \{a_n\}_{n=1}^\infty, \{b_n\}_{n=1}^\infty \rangle := \sum_{n=1}^\infty a_n b_n^*, \quad \forall \{a_n\}_{n=1}^\infty, \{b_n\}_{n=1}^\infty \in \ell^2(\mathbb{N}, \mathcal{A}).$$

Hence the norm on $\ell^2(\mathbb{N}, \mathcal{A})$ becomes

$$\|\{a_n\}_{n=1}^\infty\| := \left\| \sum_{n=1}^\infty a_n a_n^* \right\|^{\frac{1}{2}}, \quad \forall \{a_n\}_{n=1}^\infty \in \ell^2(\mathbb{N}, \mathcal{A}).$$

2. NONCOMMUTATIVE DONOHO-STARK-ELAD-BRUCKSTEIN-RICAUD-TORRÉSANI UNCERTAINTY PRINCIPLE

We start by recalling the definition of Parseval frames for Hilbert C^* -modules by Frank and Larson [6].

Definition 2.1. [6] *Let \mathcal{E} be a Hilbert C^* -module over a unital C^* -algebra \mathcal{A} . A collection $\{\tau_n\}_{n=1}^\infty$ in \mathcal{E} is said to be a **modular Parseval frame** for \mathcal{E} if*

$$\langle x, x \rangle = \sum_{n=1}^{\infty} \langle x, \tau_n \rangle \langle \tau_n, x \rangle, \quad \forall x \in \mathcal{E}.$$

As shown in [6] a modular Parseval frame $\{\tau_n\}_{n=1}^\infty$ for \mathcal{E} gives an adjointable isometry

$$\theta_\tau : \mathcal{E} \ni x \mapsto \theta_\tau x := \{\langle x, \tau_n \rangle\}_{n=1}^\infty \in \ell^2(\mathbb{N}, \mathcal{A})$$

with adjoint

$$\theta_\tau^* : \ell^2(\mathbb{N}, \mathcal{A}) \ni \{a_n\}_{n=1}^\infty \mapsto \theta_\tau^* \{a_n\}_{n=1}^\infty := \sum_{n=1}^{\infty} a_n \tau_n \in \mathcal{E}.$$

We this preliminaries we can derive noncommutative analogue of Theorem 1.3. In the following theorem, given a subset $\Lambda \subseteq \mathbb{N}$, we set the notation

$$o(\Lambda) := \text{Number of elements in } \Lambda.$$

Theorem 2.2. (*Noncommutative Donoho-Stark-Elad-Bruckstein-Ricaud-Torrésani Uncertainty Principle*) *For any two modular Parseval frames $\{\tau_n\}_{n=1}^\infty$ and $\{\omega_m\}_{m=1}^\infty$ for a Hilbert C^* -module \mathcal{E} , we have*

$$\left(\frac{\|\theta_\tau x\|_0 + \|\theta_\omega x\|_0}{2} \right)^2 \geq \|\theta_\tau x\|_0 \|\theta_\omega x\|_0 \geq \frac{1}{\sup_{n,m \in \mathbb{N}} \|\langle \tau_n, \omega_m \rangle\|^2}, \quad \forall x \in \mathcal{E}, x \neq 0.$$

Proof. Let $x \in \mathcal{E}$ be nonzero. Using Lemma 1.5 and the well-known fact in C^* -algebra that ‘norm respects ordering of positive elements’, we get

$$\begin{aligned}
 \|x\|^2 &= \|\langle x, x \rangle\| = \left\| \sum_{n=1}^{\infty} \langle x, \tau_n \rangle \langle \tau_n, x \rangle \right\| = \left\| \sum_{n \in \text{supp}(\theta_\tau x)} \langle x, \tau_n \rangle \langle \tau_n, x \rangle \right\| \\
 &= \left\| \sum_{n \in \text{supp}(\theta_\tau x)} \left\langle \sum_{m=1}^{\infty} \langle x, \omega_m \rangle \omega_m, \tau_n \right\rangle \left\langle \tau_n, \sum_{k=1}^{\infty} \langle x, \omega_k \rangle \omega_k \right\rangle \right\| \\
 &= \left\| \sum_{n \in \text{supp}(\theta_\tau x)} \left\langle \sum_{m \in \text{supp}(\theta_\omega x)} \langle x, \omega_m \rangle \omega_m, \tau_n \right\rangle \left\langle \tau_n, \sum_{k \in \text{supp}(\theta_\omega x)} \langle x, \omega_k \rangle \omega_k \right\rangle \right\| \\
 &= \left\| \sum_{n \in \text{supp}(\theta_\tau x)} \left(\sum_{m \in \text{supp}(\theta_\omega x)} \langle x, \omega_m \rangle \langle \tau_n, \omega_m \rangle^* \right) \left(\sum_{k \in \text{supp}(\theta_\omega x)} \langle x, \omega_k \rangle \langle \tau_n, \omega_k \rangle \right)^* \right\| \\
 &\leq \left\| \sum_{n \in \text{supp}(\theta_\tau x)} \left\| \sum_{m \in \text{supp}(\theta_\omega x)} \langle \tau_n, \omega_m \rangle \langle \tau_n, \omega_m \rangle^* \right\| \left(\sum_{k \in \text{supp}(\theta_\omega x)} \langle x, \omega_k \rangle \langle \omega_k, x \rangle \right) \right\| \\
 &\leq \left\| \sum_{n \in \text{supp}(\theta_\tau x)} \sum_{m \in \text{supp}(\theta_\omega x)} \|\langle \tau_n, \omega_m \rangle \langle \tau_n, \omega_m \rangle^*\| \left(\sum_{k \in \text{supp}(\theta_\omega x)} \langle x, \omega_k \rangle \langle \omega_k, x \rangle \right) \right\| \\
 &\leq \left(\sup_{n, m \in \mathbb{N}} \|\langle \tau_n, \omega_m \rangle\|^2 \right) \left\| \sum_{n \in \text{supp}(\theta_\tau x)} \sum_{m \in \text{supp}(\theta_\omega x)} 1 \cdot \left(\sum_{k \in \text{supp}(\theta_\omega x)} \langle x, \omega_k \rangle \langle \omega_k, x \rangle \right) \right\| \\
 &\leq \left(\sup_{n, m \in \mathbb{N}} \|\langle \tau_n, \omega_m \rangle\|^2 \right) \left\| o(\text{supp}(\theta_\tau x)) o(\text{supp}(\theta_\omega x)) \left(\sum_{k \in \text{supp}(\theta_\omega x)} \langle x, \omega_k \rangle \langle \omega_k, x \rangle \right) \right\| \\
 &= \left(\sup_{n, m \in \mathbb{N}} \|\langle \tau_n, \omega_m \rangle\|^2 \right) \|\theta_\tau x\|_0 \|\theta_\omega x\|_0 \left\| \sum_{k \in \text{supp}(\theta_\omega x)} \langle x, \omega_k \rangle \langle \omega_k, x \rangle \right\| \\
 &= \left(\sup_{n, m \in \mathbb{N}} \|\langle \tau_n, \omega_m \rangle\|^2 \right) \|\theta_\tau x\|_0 \|\theta_\omega x\|_0 \|\langle x, x \rangle\| \\
 &= \left(\sup_{n, m \in \mathbb{N}} \|\langle \tau_n, \omega_m \rangle\|^2 \right) \|\theta_\tau x\|_0 \|\theta_\omega x\|_0 \|x\|^2.
 \end{aligned}$$

By canceling $\|x\|$ we get the stated inequality. \square

Using Chebotarev theorem, in 2005, Tao [2, 11] improved Theorem 1.1 for prime dimensions d .

Theorem 2.3. (Tao Uncertainty Principle) [11] For every prime p ,

$$\|h\|_0 + \|\widehat{h}\|_0 \geq p + 1, \quad \forall h \in \mathbb{C}^p \setminus \{0\}.$$

In view of Theorem 2.3 we make the following conjecture.

Conjecture 2.4. Let p be a prime and \mathcal{A} be a unital C^* -algebra with invariant basis number property. Let $\widehat{\cdot}: \mathcal{A}^p \rightarrow \mathcal{A}^p$ be the noncommutative Fourier transform defined by

$$\widehat{(a_j)_{j=0}^{p-1}} := \frac{1}{\sqrt{p}} \left(\sum_{j=0}^{p-1} a_j e^{\frac{-2\pi i j k}{p}} \right)_{k=0}^{p-1}, \quad \forall (a_j)_{j=0}^{p-1} \in \mathcal{A}^p.$$

Then

$$\|x\|_0 + \|\widehat{x}\|_0 \geq p + 1, \quad \forall x \in \mathcal{A}^p \setminus \{0\}.$$

REFERENCES

- [1] John J. Benedetto and Matthew Fickus. Finite normalized tight frames. *Adv. Comput. Math.*, 18(2-4):357–385, 2003.
- [2] Tullio Ceccherini-Silberstein, Fabio Scarabotti, and Filippo Tolli. *Discrete harmonic analysis : Representations, number theory, expanders, and the Fourier transform*, volume 172 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2018.
- [3] David L. Donoho and Philip B. Stark. Uncertainty principles and signal recovery. *SIAM J. Appl. Math.*, 49(3):906–931, 1989.
- [4] Michael Elad. *Sparse and redundant representations : From theory to applications in signal and image processing*. Springer, New York, 2010.
- [5] Michael Elad and Alfred M. Bruckstein. A generalized uncertainty principle and sparse representation in pairs of bases. *IEEE Trans. Inform. Theory*, 48(9):2558–2567, 2002.
- [6] Michael Frank and David R. Larson. Frames in Hilbert C^* -modules and C^* -algebras. *J. Operator Theory*, 48(2):273–314, 2002.
- [7] Irving Kaplansky. Modules over operator algebras. *Amer. J. Math.*, 75:839–858, 1953.
- [8] William L. Paschke. Inner product modules over B^* -algebras. *Trans. Amer. Math. Soc.*, 182:443–468, 1973.
- [9] Benjamin Ricaud and Bruno Torr sani. Refined support and entropic uncertainty inequalities. *IEEE Trans. Inform. Theory*, 59(7):4272–4279, 2013.
- [10] Marc A. Rieffel. Induced representations of C^* -algebras. *Advances in Math.*, 13:176–257, 1974.
- [11] Terence Tao. An uncertainty principle for cyclic groups of prime order. *Math. Res. Lett.*, 12(1):121–127, 2005.
- [12] Audrey Terras. *Fourier analysis on finite groups and applications*, volume 43 of *London Mathematical Society Student Texts*. Cambridge University Press, Cambridge, 1999.