### Alternative representation of Magnus series by exact proper operator exponent

Yu. N. Kosovtsov

Lviv, Ukraine email: yunkosovtsov@gmail.com

#### Abstract

In this report the emphasis is on an alternative representation of the Magnus series by proper operator (matrix) exponential solutions to differential equations (systems), both linear and nonlinear ODEs and PDEs. The main idea here is in exact linear representations of the nonlinear DEs. We proceeded from Dyson's time-ordered solutions, and using only generalizations of the well-known Baker-Campbell- Hausdorff (BCH) and Zassenhaus formulae for t-dependent operators directly converted them to simple proper operator exponents. The method being explicit both in terms of the operator and in terms of expressing the formal solution as an ordinary exponential, makes it quite easy to calculate analytical expressions to solutions in the form of a Taylor function series in one variable t. If introduce a mutually invertible change of variable t into the original equations and then find a solution to this new equation in the form with ordinary exponential, one can obtain a completely different Taylor expansion of the desired function. The essence of this method comes down to resuming the series.

### 1 Introduction

Series expansions of solutions to differential equations have been widely investigated in the literature. In 1949, F. J. Dyson [1] published a paper in which he introduced the concept of time-ordered (or chronological) exponent, with the help of which it became possible to concisely represent the solution of a homogeneous linear differential equation with varying coefficients by analogy with a linear differential equation with constant coefficients for which the solutions are represented by an ordinary exponential.

Linear ordinary differential equations with non-constant coefficients (also known as non-autonomous ODE),

$$u' = A(t)u \tag{1}$$

with the initial condition

$$u|_{t=0} = u_0,$$

where A(t) is a linear operator which, generally speaking, depends on t, but does not depend on operator  $\frac{\partial}{\partial t}$  explicitly. Under the assumption that A commutes

with itself at all times, i.e.  $A(t_1)A(t_2)-A(t_2)A(t_1)=0$  for all  $t_1,t_2$ , then solution for u is expressed by the proper exponential of A as  $u(t)=\exp\{\int_0^t d\tau \, A(\tau)\}$ . In general, however, the evolution operator has no known explicit form in terms of A and is rather expressed as

$$u(t) = \mathbf{T} \exp\{\int_0^t d\tau \, A(\tau)\} u_0, \tag{2}$$

were **T** is the so-called time-ordered (or chronological) operator. The action of this operator transforms the series representation of the ordinary operator-exponential function into a operator-valued Peano-Baker series (also known as Dyson series [2]) involving an infinite sum of nested integrals. But this expression cannot give the whole solution as has already been said. This series is frankly unwieldy, hinders theoretical analyses of the solution, is computationally costly to evaluate and purely perturbative from a physical point of view.

Over time, it became clear that there are possibilities for representing solutions in the form of time-ordered exponents to evolution equation for both nonlinear ODEs [3, 14, 15] and nonlinear PDEs [16] and others, as well as their coupled systems. The main idea here is in *exact linear* representations of the *nonlinear* DEs. If for nonlinear ODEs it is well known that they are represented in the form of a linear partial differential equation [14, 15], nonlinear PDEs are *exactly* represented by *linear* differential equations with functional derivatives [16, 17].

Almost immediately after the appearance of Dyson's paper [1] the question arose whether it was possible to find a way to *exactly* represent time-ordered exponents in a simpler form without time-ordering operation.

Magnus proposal [4] with respect to the linear evolution equation

$$Y^{'}(t) = A(t)Y(t)$$

with initial condition Y(0) = I, was to express the solution as the exponential of a certain operator  $\Omega(t)$ ,

$$Y(t) = e^{\Omega(t)}. (3)$$

This is in contrast to the representation (2) in terms of the time-ordering operator introduced by Dyson. At the same time, the time-ordering operator **T** disappears, but the main disadvantage of the Dyson series - bulkiness - only increases significantly.

Over the past 50 years since the appearance of the Magnus article, a significant number of works have appeared devoted to both the development of the Magnus idea itself and its various applications in many fields of science and technology [6, 7, 8, 9, 10, 11, 12, 13].

# 2 Formal solutions to the first order nonlinear ODEs and PDEs by time-ordered operator exponents

Since the representation of solutions for linear ODEs and PDEs in the form (2) is well known, here we present a summary of the results associated with the representation of nonlinear ODEs and PDEs [16], for which linear problems are a special case. For clarity of presentation, we limit ourselves to considering single DEs of the first order in t. Generalizations to DEs of higher orders in t and systems of coupled equations are almost obvious (see [15, 16]).

The starting point for solving differential equations by the operator method is *linear* first-order differential equation for operator  $\mathbf{E} = \mathbf{E}(t)$ :

$$\frac{\partial \mathbf{E}}{\partial t} = \mathbf{L}(t)\mathbf{E} \tag{4}$$

with the initial condition

$$\mathbf{E}|_{t=a} = \mathbf{I},$$

where  $\mathbf{L}(t)$  is a linear operator which, generally speaking, depends on t, but does not depend on operator  $\frac{\partial}{\partial t}$  explicitly,  $\mathbf{I}$  is identity operator. The solution of this equation for t > a was obtained by F. Dyson in the time-ordered (or chronological) exponent form [1]

$$\mathbf{E}(t) = \mathbf{T} \exp\{\int_a^t d\tau \, \mathbf{L}(\tau)\},\,$$

where it is supposed, that exponent represents a power series expansion with following *chronologization* according to the rule

$$\mathbf{T} \left\{ \mathbf{L}(\tau_1) \ \mathbf{L}(\tau_2) \dots \mathbf{L}(\tau_n) \right\} = \mathbf{L}(\tau_{\alpha_1}) \ \mathbf{L}(\tau_{\alpha_2}) \dots \mathbf{L}(\tau_{\alpha_n}).$$

$$\tau_{\alpha_1} \ge \tau_{\alpha_2} \ge \dots \ge \tau_{\alpha_n}$$

Very important role plays the equation adjoint to (4) for operator  $\mathbf{E}^{-1} = \mathbf{E}^{-1}(t)$  (see [3, 16])

$$\frac{\partial \mathbf{E}^{-1}}{\partial t} = -\mathbf{E}^{-1} \mathbf{L}(t)$$

with the initial condition

$$\mathbf{E}^{-1}|_{t=a} = \mathbf{I}.$$

It is easy to see using the Dyson's idea, that its solution is

$$\mathbf{E}^{-1}(t) = \mathbf{T}_0 \exp\{-\int_a^t d\tau \, \mathbf{L}(\tau)\},\,$$

where operator  $\mathbf{T}_0$  is the operator of opposed chronologization with action

$$\mathbf{T}_0 \left\{ \mathbf{L}(\tau_1) \ \mathbf{L}(\tau_2) \dots \mathbf{L}(\tau_n) \right\} = \mathbf{L}(\tau_{\alpha_1}) \ \mathbf{L}(\tau_{\alpha_2}) \dots \mathbf{L}(\tau_{\alpha_n}).$$

$$\tau_{\alpha_1} \le \tau_{\alpha_2} \le \dots \le \tau_{\alpha_n}$$

It appears that the operator  $\mathbf{E}^{-1}(t)$  is inverse to the  $\mathbf{E}(t)$ , i.e.

$$\mathbf{T}_0 \exp\{-\int_a^t d\tau \, \mathbf{L}(\tau)\} \, \mathbf{T} \exp\{\int_a^t d\tau \, \mathbf{L}(\tau)\} = \mathbf{I}.$$

Ordering operators  $\mathbf{T}$  and  $\mathbf{T}_0$  for t < a change places. In some sense in linear problems operators  $\mathbf{T}$  and  $\mathbf{T}_0$  provide description of direct and back scattering accordingly. It be assumed henceforth that t > a.

Time-ordered (chronological) operators have to be introduced because the operators in the integrands with different values of the variable  $\tau$  may not commute, what cause certain difficulties in manipulation with chronological exponents. Nevertheless, examination of algebraical properties of the chronological exponents allows to obtain an extensive family of operator identities [3, 14, 15] which provide possibilities of transformations and analysis of chronological expressions.

The operator  $\Delta$  is derivative if

$$\Delta (\mathbf{A} + \mathbf{B})f = \Delta \mathbf{A}f + \Delta \mathbf{B}f,$$
$$\Delta \mathbf{A}f = (\Delta \mathbf{A})f + \mathbf{A}\Delta f$$

for any differentiable function f and any differentiable linear operators  $\mathbf{A}$  and  $\mathbf{B}$ . In other words, the linear operation is derivative if it satisfies to Leibnitz rule. As we will see later, the chronological operators with derivative operators in exponent, which we will denote for short as

$$\mathbf{E} = \mathbf{T} \exp\{ \int_{a}^{t} d\tau \, \mathbf{\Delta}(\tau) \},$$

play the extremely important role here.

Let us consider particular cases of the derivative operators  $\Delta$  and differential equations, which correspond them.

The (general) solution to the first order nonlinear ODE for u (see [3, 14])

$$\frac{\partial u}{\partial t} = F(t, u), \qquad u|_{t=a} = c,$$
 (5)

where c is an arbitrary constant, in the form with time-ordered exponent is as

$$u(t,c) = \mathbf{T}_0 \exp\{\int_a^t d\tau \, F(\tau,c) \frac{\partial}{\partial c}\} \, c. \tag{6}$$

Here the *linear* derivative operator is

$$\Delta_1(\tau) = F(\tau, c) \frac{\partial}{\partial c}.$$

Note that for nonlinear DEs it differs markedly from the operator A(t) in linear ODE (1) and its solution (2). This feature is characteristic of all nonlinear DEs, since the basis for such a result is the possibility of *exact* representation

of *nonlinear ODE* by *linear PDE*. The equivalent representation of a nonlinear ODE (5) is the following linear PDE

$$\frac{\partial u(t,c)}{\partial t} + F(\tau,c)\frac{\partial u(t,c)}{\partial c} = 0, \qquad u(t,c)|_{t=a} = c.$$

There are other interpretations of the exact linearization [3, 13, 14, 15], leading to the same result.

To nonlinear partial differential equation for u(t,x) of the first order on variable t and arbitrary order on space variables

$$\frac{\partial u}{\partial t} = F(t, x, u, \dots, D_x^{\alpha_j} u), \qquad u|_{t=a} = c(x), \tag{7}$$

where now in ...,  $D_x^{\alpha_j}u$  appears the corresponding set of partial derivatives of u(t,x) on space variables  $x=(x_1,\ldots,x_m)$  and c(x) is an arbitrary function, the (general) time-ordered solution is as follows [16, 17]

$$u(t,x) = \mathbf{T}_0 \exp\{\int_a^t d\tau \int_{\mathbb{R}^m} d^m \zeta \ F(\tau,\zeta,c(\zeta),\dots,D_{\zeta}^{\alpha_j}c(\zeta)) \frac{\delta}{\delta c(\zeta)}\} c(x) ,$$

where  $\frac{\delta}{\delta c(\zeta)}$  is the functional derivative. Here by functional derivative we mean the linear mapping with the following property

$$\frac{\delta}{\delta c(\zeta)}c(x) = Dirac(x - \zeta),$$

the chain rule is also valid in this context.

In this case the *linear* derivative operator is

$$\mathbf{\Delta_2}(\tau) = \int_{\mathbb{R}^m} d\zeta \ F(\tau, \zeta, c(\zeta), \dots D_x^{\alpha_j} c(\zeta)) \frac{\delta}{\delta c(\zeta)}, \qquad \zeta \in \mathbb{R}^m$$

and the equivalent representation of a nonlinear PDE (7) is the following linear DE with functional derivative

$$\frac{\partial u(t,x,c(x))}{\partial t} + \Delta_2(t)u(t,x,c(x)) = 0, \qquad u(t,x,c(x))|_{t=a} = c(x).$$

## 3 Exact representation of time-ordered exponent by proper operator exponent

Here we need generalizations of the well-known Baker-Campbell-Hausdorff (BCH) and Zassenhaus formulaes for t-dependent operators in the sense that the classical BCH formula merges product of two exponential operators into a single one and Zassenhaus formula splits an exponential operator into a product of exponential operators [14, 15].

$$\mathbf{T} \exp\{\int_{a}^{t} d\tau \, \mathbf{B}(\tau)\} \, \mathbf{T} \exp\{\int_{a}^{t} d\tau \, \mathbf{A}(\tau)\} =$$

$$\mathbf{T} \exp\{\int_{a}^{t} d\tau \, [\mathbf{B}(\tau) + \mathbf{T} \exp\{\int_{a}^{\tau} d\xi \, \mathbf{B}(\xi)\} \, \mathbf{A}(\tau) \mathbf{T}_{0} \exp\{-\int_{a}^{\tau} d\xi \, \mathbf{B}(\xi)\}]\} \quad (8)$$

and

$$\mathbf{T} \exp\{\int_{a}^{t} d\tau \left[\mathbf{B}(\tau) + \mathbf{C}(\tau)\right]\} = \mathbf{T} \exp\{\int_{a}^{t} d\tau \, \mathbf{B}(\tau)\} \times \mathbf{T} \exp\{\int_{a}^{t} d\tau \, \mathbf{T}_{0} \exp\{-\int_{a}^{\tau} d\xi \, \mathbf{B}(\xi)\} \, \mathbf{C}(\tau) \, \mathbf{T} \exp\{\int_{a}^{\tau} d\xi \, \mathbf{B}(\xi)\}\}.$$
(9)

By taking inverses of both sides of (8),(9) we can find the similar identities for the inverse ordered operators [15].

$$\begin{split} &\mathbf{T}_0 \exp\{\int_a^t d\tau \, \mathbf{B}(\tau)\} \, \mathbf{T}_0 \exp\{\int_a^t d\tau \, \mathbf{A}(\tau)\} = \\ &\mathbf{T}_0 \exp\{\int_a^t d\tau \, [\mathbf{B}(\tau) + \mathbf{T} \exp\{-\int_a^\tau d\xi \, \mathbf{B}(\xi)\} \mathbf{A}(\tau) \mathbf{T}_0 \exp\{\int_a^\tau d\xi \, \mathbf{B}(\xi)\}]\}. \end{split}$$

The analog of (9) is

$$\mathbf{T}_{0} \exp\{\int_{a}^{t} d\tau \left[\mathbf{B}(\tau) + \mathbf{C}(\tau)\right]\} =$$

$$\mathbf{T}_{0} \exp\{\int_{a}^{t} d\tau \,\mathbf{T}_{0} \exp\{\int_{a}^{\tau} d\xi \,\mathbf{B}(\xi)\} \mathbf{C}(\tau) \mathbf{T} \exp\{-\int_{a}^{\tau} d\xi \,\mathbf{B}(\xi)\}\} \times$$

$$\mathbf{T}_{0} \exp\{\int_{a}^{t} d\tau \,\mathbf{B}(\tau)\}. \tag{10}$$

We will also need the well-known shift operator property

$$\exp\{\alpha \frac{\partial}{\partial s}\}\Phi(s) = \Phi(s+\alpha). \tag{11}$$

Now let us note the following circumstance. Let us consider the following operator chain

$$\exp\{(t-a)\left[\boldsymbol{\Delta}(s) + \frac{\partial}{\partial s}\right]\} \stackrel{(10)}{=}$$

$$\mathbf{T}_{0} \exp\{\int_{a}^{t} d\tau \, e^{\{(\tau-a)\frac{\partial}{\partial s}\}} \boldsymbol{\Delta}(s) e^{\{(a-\tau)\frac{\partial}{\partial s}\}}\} \exp\{(t-a)\frac{\partial}{\partial s}\} \stackrel{(11)}{=}$$

$$\mathbf{T}_{0} \exp\{\int_{a}^{t} d\tau \, \boldsymbol{\Delta}(s+\tau-a)\} \exp\{(t-a)\frac{\partial}{\partial s}\}.$$

So

$$\begin{split} &\mathbf{T}_0 \, \exp\{\int_a^t d\tau \, \boldsymbol{\Delta_1}(s+\tau-a)\} \, \exp\{(t-a)\frac{\partial}{\partial s}\}c = \\ &\mathbf{T}_0 \, \exp\{\int_a^t d\tau \, \boldsymbol{\Delta_1}(s+\tau-a)\}\} \, c = \exp\{(t-a) \, [f(s,c)\, \frac{\partial}{\partial c} + \frac{\partial}{\partial s}]\} \, c \end{split}$$

and since it is obvious that

$$\mathbf{T}_{0} \exp\{\int_{a}^{t} d\tau \, \Delta_{\mathbf{1}}(s+\tau-a)\} c \,|_{s=a} = \mathbf{T}_{0} \exp\{\int_{a}^{t} d\tau \, \Delta_{\mathbf{1}}(\tau)\} c \stackrel{(6)}{=} u(t,c) = \exp\{(t-a) \left[f(s,c) \frac{\partial}{\partial c} + \frac{\partial}{\partial s}\right]\} c \,|_{s=a} .$$

That is, we got the solution to ODE (5) in the form with *proper* operator exponent

$$u(t,c) = \exp\{(t-a)\left[f(s,c)\frac{\partial}{\partial c} + \frac{\partial}{\partial s}\right]\}c|_{s=a}.$$

Denoting

$$U(t, s, c) = \exp\{(t - a) \left[ f(s, c) \frac{\partial}{\partial c} + \frac{\partial}{\partial s} \right] \} c,$$

we obtain from the preceding expressions that the (general) solution of ODE (5) as

$$u(t,c) = U(t,s,c)|_{s=a},$$

where the function U(t, s, c) satisfies the following linear PDE

$$\frac{\partial U(t,s,c)}{\partial t} - f(s,c) \frac{\partial U(t,s,c)}{\partial c} - \frac{\partial U(t,s,c)}{\partial s} = 0, \qquad U(t,c,s) \mid_{t=a} = c.$$

Similarly for nonlinear PDE (7) from

$$\mathbf{T}_{0} \exp\{\int_{a}^{t} d\tau \, \Delta_{2}(s+\tau-a)\} \exp\{(t-a)\frac{\partial}{\partial s}\}c(x) =$$

$$\mathbf{T}_{0} \exp\{\int_{a}^{t} d\tau \, \Delta_{2}(s+\tau-a)\}\} c(x) =$$

$$\exp\{(t-a)\int_{\mathbb{R}^{m}} d\zeta \, [F(s,\zeta,c(\zeta),D_{x}^{\alpha_{j}}c(\zeta))\frac{\delta}{\delta c(\zeta)} + \frac{\partial}{\partial s}]\} c(x)$$

we get (general) solution in the following form with proper operator exponent

$$u(t, c(x)) = \left[\exp\{(t - a) \int_{\mathbb{R}^m} d\zeta \left[F(s, \zeta, c(\zeta), D_x^{\alpha_j} c(\zeta)) \frac{\delta}{\delta c(\zeta)} + \frac{\partial}{\partial s}\right]\} c(x)\right]|_{s=a}.$$
(12)

Denoting now

$$V(t, c(x), s) = \exp\{(t - a) \int_{\mathbb{R}^m} d\zeta \left[ F(s, \zeta, c(\zeta), D_x^{\alpha_j} c(\zeta)) \frac{\delta}{\delta c(\zeta)} + \frac{\partial}{\partial s} \right] \} c(x),$$

we obtain from the preceding expression that the (general) solution of PDE (7) as

$$u(t,c(x)) = V(t,c(x),s)|_{s=a},$$

where the function V(t, c(x), s) satisfies the following linear equation

$$\frac{\partial}{\partial t}V(t,c(x),s) - \Delta_{\mathbf{2}}(s)V(t,c(x),s) - \frac{\partial}{\partial s}V(t,c(x),s) = 0, \qquad V(t,c(x),s)\mid_{t=a} = c(x).$$

The last forms of the operator solutions has some interesting features in the sense that they do not contain ordering operator **T** and even any integration.

So, we solved the problem posed by Magnus, which can be formulated in two ways: how to get rid of time-ordering operators in Dyson's solutions to DEs or how to "correct" the proper operator exponent so that it nevertheless expresses exact solutions and has the simplest *explicit* form.

Magnus, to solve this problem, proceeded from the *nonlinear* equation for  $\Omega(t)$  in (3), which in itself was a difficult task, and obtains an *infinite series* known as Magnus expansion. We directly proceeded from Dyson's time-ordered solutions, and using only *generalizations* of the well-known Baker-Campbell-Hausdorff (BCH) and Zassenhaus formulae for "t-dependent" operators directly converted them to simple proper operator exponents.

Generalizations for differential equations of any order in t and for coupled systems of such equations are quite obvious [15, 16].

In comparison with the cumbersome expressions in Magnus series method, approximate expressions of the solution to u (6), (12) may be expressed immediately in forms of formal Taylor series only on t without solving auxiliary systems of equations. The Maple procedures for analytical calculations of approximate solutions to linear and nonlinear ordinary and partial differential equations (Cauchy problem), which are based on the approach considered here, are presented in [18].

In addition, the method described above, being explicit both in terms of the operator and in terms of expressing the formal solution as an ordinary exponential, makes it quite easy to calculate analytical expressions to solutions in the form of a Taylor function series in one variable t. The resulting Taylor series, generally speaking, does not converge quickly enough and in this version most often there is no small parameter. We do not consider here questions of convergence of the formal series obtained here, we will only limit ourselves to a brief add-on.

# 4 Options for representing solutions in the form of generalized Taylor (Fourier) series

If we introduce a mutually invertible change of variable t into the original equations and then find a solution to this new equation in the form with ordinary exponential, or in the Taylor series form using the method described here and then return to the old variable t, we will obtain a completely different Taylor expansion of the desired function. For example, using the replacement  $\tau = \exp(-pt)$ , where p is an arbitrary parameter (real or complex), we obtain a series for  $\tau = \exp(-pt)$ , and if the real part of p is positive, then this ensures faster convergence of the series, where  $1/(\Re \mathfrak{e}\,p)$  is an artificially introduced necessary small parameter that ensures control of the remainder term. By analogy with the above example, one can use many other mutually invertible substitutions with different rates of convergence of the series. It is easy to see that the essence

of this method comes down to resuming the series.

This approach is illustrated by examples of nontrivial PDE

$$\frac{\partial^2 u(t,x)}{\partial t^2} = u(t,x) \frac{\partial^2 u(t,x)}{\partial t \, \partial x}, \qquad u(0,x) = h(x), \frac{\partial u(t,x)}{\partial t} \mid_{t=0} = g(x).$$
 (13)

Expanding the operator exponent (12) corresponding to the equation (13) into a series and executing the actions of the operators, we obtain a solution to PDE (13) in the form of a Taylor function series (to avoid bulkiness, we limit ourselves to five of its terms)

$$u(t,x) = h + gt + hg' \frac{t^2}{2} + (h^2g'' + g'(hh' + g) \frac{t^3}{6} + (hh'^2g' + 3h^2h'g'' + h^2h''g' + h^3g''' + 2h'gg' + 2hg'^2 + 3hgg'') \frac{t^4}{24} + \dots$$
(14)

Now let's try the above method of series resummation on the same equation (13). Making a mutually invertible change of variable  $t = -\ln(\tau)/p \Leftrightarrow \tau = e^{-pt}$  in equation (13), we get *auxiliary* nonlinear equation

$$W(\tau, x) \frac{\partial^2 W(\tau, x)}{\partial \tau \partial x} + \left(\tau \frac{\partial^2 W(\tau, x)}{\partial \tau^2} + \frac{\partial W(\tau, x)}{\partial \tau}\right) p = 0$$

with new initial conditions for this equation, consistent with the initial conditions of the original PDE.

$$W(1,x) = h(x), \ \frac{\partial W(\tau,x)}{\partial \tau}|_{\tau=1} = -g(x)/p,$$

Using the same method as for the original PDE, we obtain a solution to the auxiliary PDE in the form of a Taylor series in  $\tau$  and, finally, substitute the inverse substitution  $\tau = e^{-pt}$  into the resulting solution. As a result, we obtain a new solution to PDE (13) in the form of a generalized Taylor series

$$u(t,x) = h - \frac{g}{p}(e^{-pt} - 1) + \frac{1}{2p^2}(hg' + gp)(e^{-pt} - 1)^2 + \frac{1}{6p^3}[-h^2g'' + (-g - 3hp - hh')g' - 2gp^2](e^{-pt} - 1)^3 + \frac{1}{24p^4}[h^3g''' + (3hg + 6h^2p + 3h^2h')g'' + h^2h''g' + 2hg'^2 + (hh'^2 + (6hp + 2g)h' + 11hp^2 + 6gp)g' + 6gp^3](e^{-pt} - 1)^4 + \dots$$
(15)

It can be shown that the expansion of the last series (15) into an ordinary Taylor series coincides with the series (14) for any p.

If we put  $p = i\omega$  in (15), where  $\omega \in \mathbb{R}$ , then we get the expansion of the PDE solution (13) in *Fourier series*. Typically, the Fourier expansion is used for linear DEs with coefficients independent of t. The method proposed here without changes allows one to obtain solutions in the form of a Fourier series to linear and nonlinear DEs with parameters depending on t.

### References

- [1] F. J. Dyson, The radiation theories of Tomonaga, Schwinger, and Feynman. Physical Review. 75 486-502, 1949.
- [2] F. J. Dyson. Divergence of Perturbation Theory in Quantum Electrodynamics. Physical Review, 85(4):631-632, 1952.
- [3] A. A. Agrachev and R. V. Gamkrelidze, Exponential representations of flows and chronological calculus, Math. USSR Sbornik, vol. 35, 727-785, 1978.
- [4] W. Magnus. Algebraic aspects in the theory of systems of linear differential equations. Research report, Courant Institute of Mathematical Sciences, New York University, 1953.
- [5] W. Magnus, On the exponential solution of differential equations for a linear operator, Commun. Pure and Appl. Math., vol. 7, 649-673, 1954.
- [6] R. M. Wilcox, Exponential operators and parameter differentiation in quantum physics. J. Math. Phys. 8, 962, 1967.
- [7] B. Mielnik, J. Plebánski, Combinatorial approach to Baker-Campbell-Hausdorff exponents. Ann. Inst. Henri Poincare A XII, 215, 1970.
- [8] R. S. Strichartz, The Campbell-Baker-Hausdorff-Dynkin formula and solutions of differential equations. J. Func. Anal. 72, 320, 1987.
- [9] R.V. Chacon and A.T. Fomenko. Recursion formulas for the Lie integral. Adv. Math., 88 200257, 1991.
- [10] S. Monaco, D.Normand-Cyrot, C. Califano, From chronological calculus to exponential representations of continuous and discrete-time dynamics: A lie-algebraic approach. IEEE Transactions on Automatic Control, 52(12), 2227-2241,2007.
- [11] S. Blanes, F. Casas, J. A. Oteo, and J. Ros, The Magnus expansion and some of its applications. Physics Reports, 470(5):151-238, 2009.
- [12] M. Bauer, R. Chetrite, K. Ebrahimi-Fard, F. Patras, Time-ordering and a generalized Magnus expansion. Letters in Mathematical Physics, 103, 331-350, 2013.
- [13] P-L.Giscard, Algebraic Walk Theory. Université du Littoral Côte d'Opale, 2023. https://hal.science/tel-04323927v1/file/HDR.pdf
- [14] Yu. N. Kosovtsov, The introduction to the operator method for solving differential equations. First-order DE. arXiv: 0202040 [math-ph], 2002.
- [15] Yu. N. Kosovtsov, The Chronological Operator Algebra and formal solutions of differential equations. arXiv: 0409035 [math-ph], 2004.

- [16] Yu. N. Kosovtsov, Formal exact operator solutions to nonlinear differential equations. arXiv: 0910.3923 [math-ph], 2009.
- [17] Yu. N. Kosovtsov, Exact linear representations of the nonlinear Cauchy problems and their smooth solutions. arXiv: 2308.12081 [math-ph], 2023.
- [18] (1) Yu. N. Kosovtsov, Series solutions to non-linear ODEs or systems of them, http://www.maplesoft.com/applications/view.aspx?SID=1426, 2004.
  - (2) Yu. N. Kosovtsov, Formal Series Solutions to Linear PDEs or Systems of Them (Cauchy problem), http://www.maplesoft.com/applications/view.aspx?SID=1687, 2005.
  - (3) Yu. N. Kosovtsov, Formal series solutions to nonlinear DE (ODE or PDE) or systems of them (Cauchy problem), http://www.maplesoft.com/applications/view.aspx?SID=1690, 2005.