LARGE SCALE COMPRESSIBLE TURBULENCE IN THE ISM OF CSWA13, A STAR-FORMING LENSED GALAXY AT Z = 1.87 WITH OUTFLOWING WIND

ITZHAK GOLDMAN^{1,2} ¹Physics Department Afeka College, Tel Aviv 6998812, Israel and ² Astrophysics Department, Tel Aviv University, Tel Aviv 6997801, Israel *Version June 14, 2024*

Abstract

Recently, Keerthi Vasan G. et al. (2024)presented spatially resolved observations of a wind outflow in CSWA13, a gravitationally lensed Star-Forming galaxy at z = 1.87. The gravitational lensing allowed for a substantially improved spatial and kinematic resolution of the wind and of the nebular gas. In this paper we take advantage of the resolved data to test for the existence of turbulence and to study its nature. We derive the spatial structure functions of the residual nebular and wind velocities along the major axis of the galaxy. The structure functions, of both velocity fields, reveal a supersonic compressible large scale turbulence. The turbulent timescale corresponding to the largest scale is about 200 Myr, an order of magnitude larger than the estimated age of the wind and of the young stars. This implies that the turbulence in the ism formed well before the wind and the young stars. Given the large spatial scale of the turbulence, it is plausible that the source of the turbulence is a large scale one e.g. a merger or tidal event that triggered the formation of molecular clouds, in the cores of which the young stars formed. A steepening of the structure functions on the smaller scales provides an estimate of the effective depth along the line of sight of the turbulent layer. The latter turns out to be $\sim 2kpc$.

Subject headings: galaxy outflows, galaxy evolution, interstellar turbulence

1. INTRODUCTION

High redshift galaxies are characterized by high star formation rates as well as outflowing winds, generated by the young stars, e.g. (Bournaud et al. 2009; Hoffmann et al. 2022; Rizzo et al. 2021; Sanders et al. 2023; Shah et al. 2022) The high rate of star formation is attributed to the assembly process of the galaxy, involving gas inflow from the circum galactic medium (CGM) and also by more violent events such as mergers that lead to formation of molecular clouds and eventually to star formation. Such mergers could result in large scale shocks and also in large scale turbulence, created by the shocks and by the instabilities caused by the mergers.

Observations of high redshift galaxies indeed display velocity dispersions that are usually interpreted as manifestation of turbulence. It has been argued that accretion onto disk galaxies can generate large scale turbulence, in particular at the disk outskirts, e. g. (Forbes et al. 2023; Goldman & Fleck 2023). Turbulence can be generated also by mergers and tidal interactions. To establish the existence of turbulence and moreover, to understand its nature, a power spectrum or structure function of the velocity field are needed. This in turn demands observations with high enough spatial resolution which, for galaxies at high redshifts, are challenging.

Gravitational lensing can help with this regard. A recent paper (Keerthi Vasan G. et al. 2024) presented a study of a wind outflow in CSWA13, which is a gravitationally lensed star-forming galaxy at z = 1.87. The gravitational lensing allowed for a substantially improved spatial and kinematic resolution. The authors obtained, among other results, two velocity fields along the major axis of the galaxy: the nebular gas velocity traced by the $C_{||||}$ emission line. that represents also the velocity of the young stars embedded in the nebular gas and the wind velocity traced by the $Si^*_{||}$ florescent emission line. Each of these velocity fields , exhibits a large scale shear.

In the present paper we set to check wether these velocity fields can be used to test for the existence of turbulence, and if so to obtain its characteristics.

The two residual velocity fields are obtained in section 2. The autocorrelation functions are presented in section 3. In section 4 we obtain the structure functions. In the appendix, the theoretical structure function of a quantity that is the result of integration along the line of sight direction, is derived. This provides an estimate of the depth of the turbulent layer. Discussion is presented in section 5.

2. THE RESIDUAL VELOCITY FIELDS

We digitized the velocity curves of Fig. 8 in Keerthi Vasan G. et al. (2024) and obtained the nebular and wind velocity as functions of position along the galactic major axis. The nebular velocity and the wind velocity exhibit each a large scale shear. We subtracted from each velocity field the corresponding large scale shear, and then removed the remaining mean value. Doing so resulted in two residual velocity fields along the major galactic axis. We derived the autocorrelation function and the structure function for each.

Structure functions rather as function of x/D.than

power spectra were employed since the former are more detailed on the smaller and medium spatial scales. They are also more reliable at treating data at the borders of the data domain (Nestingen-Palm et al. 2017).

2.1. The residual nebular velocity along the major galaxy axis

The velocity posses a large scale shear of $96.2km \ s^{-1}/(6.43kpc)$. After subtracting the shear, and the remaining mean value, the residual velocity is obtained and is displayed in Fig. 1.



FIG. 1.— The residual nebular velocity in units of km/s as function of position along the major axis, in units of 207.5pc.

2.2. The residual wind velocity along the major galaxy axis

The velocity posses a large scale shear of $152.2km \ s^{-1}/(6.43kpc)$. After subtracting the shear, and the remaining mean value, the residual velocity is obtained and is displayed in Fig. 2.



FIG. 2.— The residual wind velocity in units of km/s as function of position along the major axis, in units of 207.5pc.

3. AUTOCORRELATION FUNCTIONS

The one-dimensional autocorrelation function, C(x), of a residual velocity v(x) (with a zero mean value) is

$$C(x) = \langle v(x+x')v(x') \rangle,$$
 (1)

The brackets indicate ensemble average which by invoking the ergodic principle can be replaced by an average over x'. Here x denotes the spatial lag between two positions along the major galaxy axis.

3.1. The autocorrelation function of the residual nebular velocity

The computed observational normalized autocorrelation function of the residual nebular velocity is displayed in Fig, 3. It implies that the residual values of the nebular velocity are in fact correlated over large spatial range, comparable to the size of the major axis.



FIG. 3.— The normalized observational autocorrelation function of the residual nebular velocity as function of the spatial lag, in units 207.5pc.

3.2. The autocorrelation function of the residual wind velocity

Fig.4. presents the normalized autocorrelation function of the wind velocity. It exhibits long range correlation similar to that of the nebular velocity.



FIG. 4.— The normalized observational autocorrelation function of the residual wind velocity as function of the spatial lag, in units of 207.5pc.

4. STRUCTURE FUNCTIONS

Long range autocorrelation could be a signature of turbulence. In order to test for the existence of turbulence and understand its nature, we evaluate the structure functions for the two residual velocity fields.

The one-dimensional structure function of a quantity f(x) defined along a straight line is

$$S_f(x) = \langle \left(f(x'+x) - f(x')\right)^2 \rangle = 2C_f(0) - 2C_f(x),$$
(2)

with the lag x being the difference between two positions. $C_f(x)$ is the auto correlation function of f(x) defined as

$$C_f(x) = \langle f(x'+x)f(x) \rangle$$
. (3)

In the following, the computed structure functions of the observational residual nebular velocity, and of the residual wind velocity are presented,

4.1. The observational structure function of the residual nebular velocity

Fig.5 displays the structure function of the observational residual nebular velocity. The blue line has a logarithmic slope of 1. The orange line has a logarithmic slope of 2. In the appendix it is shown that the structure function, evaluated along a lateral line, of a quantity that is an integral over the line of sight direction increases the logarithmic slope by 1 when the lateral lag is smaller than the effective depth of the turbulent layer.

The structure function at the largest lag is $2 < C(0)^2 = 391(km/s)^2$. Thus, the one dimensional rms turbulent velocity is 13.9km/s. Assuming isotropic turbulence, the three-dimensional turbulent velocity is 24km/s



FIG. 5.— The observational structure function of the residual nebular velocity in units of $(km/s)^2$ as function of the spatial lag, in units of 207.5pc. The asymptotes have logarithmic slopes of 1 and 2.

4.2. The observational structure functions of the residual wind velocity

Fig.6 displays the structure function of the residual wind velocity. The blue line has a logarithmic slope of 1. The orange line has a logarithmic slope of 2. This behavior is similar to that of the structure function of the residual nebular velocity.

The structure function at the largest lag is $2 < C(0)^2 = 291(km/s)^2$, implying a one-dimensional rms turbulent velocity of 12.1km/s. Assuming isotropic turbulence, the three-dimensional turbulent velocity is 21km/s



FIG. 6.— The observational structure function of the residual wind velocity, in units of $(km/s)^2$ as function of the spatial lag in units of 207.5*pc*. The asymptotes have logarithmic slopes of 1 and 2.

5. DISCUSSION

5.1. The nature of the turbulence

The observational structure functions of the two residual velocity fields have, each, a logarithmic slope equaling 1 on the large spatial scales and a logarithmic slope of 2 on the small spatial scales. This dependence characterizes compressible turbulence with a one-dimensional power spectrum $\propto k^{-2}$ with k denoting the one -dimensional spatial wave number.

This power spectrum is steeper than the Kolmogorov power spectrum, which corresponds to subsonic incompressible turbulence with a one-dimensional power law with exponent of -5/3 and structure function with logarithmic slope of 2/3 and 5/3 for the large and small scales, respectively.

Such a power spectrum was derived by Burgers (1948) describing a hierarchy of shocks in compressible gas. Compressible turbulence power spectra were observed in HI intensity maps in the Milky Way (MW) galaxy (Green 1993)and in the SMC (Stanimirovic et al. 1999). This power spectrum has been observed also in molecular clouds (Larson 1981; Leung et al. 1982; Dame et al. 1986),in the HII region Sharpless 142 (Roy & Joncas 1985). It has been found in a shocked cloud near the Milky Way galaxy center (Contini & Goldman 2011), and recently in the Gamma ray emission from the large Magellanic Cloud. (Besserglik & Goldman 2021). It has been also obtained in numerical simulations e.g. (Passot et al. 1988; Vázquez-Semadeni et al. 1997; Kritsuk et al. 2007; Federrath et al. 2021).

The steeper slope signals that (unlike in the Kolmogorov spectrum) the rate of energy transfer in the turbulence cascade is not constant but decreases with increasing wavenumber. This is expected in a compressible turbulence since part of the energy at a given wavenumber in the cascade, is diverted to compression of the gas. Indeed, a theoretical derivation of the compressible turbulence power spectrum based on this argument has been obtained (Goldman 2021a).

The three-dimensional rms turbulent velocities estimated in section 4.are supersonic, in line with the shape of the structure functions.

The turbulence timescale of the largest scale eddies is $\sim l_0/v_0 \sim 200$ Myr where l_0 is the largest spatial scale and v_0 is the turbulent velocity on this scale. This timescale represents the eddy correlation time on the largest spatial scale and therefore a lower bound on the time span over which the turbulence was generated. This time span is an order of magnitude larger than the age of the young stars and the outflowing wind. Thus, the turbulence is older than the young stars and the wind that was created by the latter. The timescales of the large scale shears are about 20 Myr, so they formed at the same time as the wind. The emerging picture is that the young stars as well as the wind and the shear were formed on the background of the turbulent interstellar gas. The generating source of this large scale turbulence, must itself be correlated over the largest scale of the turbulence, rather than a collection of smaller scale sources. The probable source is a merger or a close tidal interaction with a smaller galaxy.

5.2. The effective depth of the turbulent region

. The emitted photons that determine the velocities originate from different depths along the line of sight. The issue of power spectra and structure functions of quantities which are the result of integration along the line-of-sight has been addressed by e.g. (Stutzki et al. 1998; Goldman 2000, 2021b; Lazarian & Pogosyan 2000; Miville-Deschênes et al. 2003a). These authors concluded that when the lateral spatial scale is smaller than the depth of the layer, the logarithmic slope of the power spectrum steepens by -1 compared to its value when the lateral scale is large compared to the depth. The logarithmic slope of the structure function increases by 1. This behavior was indeed revealed in observational power spectra of Galactic and extra Galactic turbulence (e.g. Elmegreen et al. (2001),Block et al. (2010), Miville-Deschênes et al. (2003b)) and in solar photospheric turbulence (Abramenko & Yurchyshyn 2020)).

In the appendix, the theoretical structure function of a quantity that is the result of integration along the line of sight, is obtained. For the specific case of compressible turbulence we found that $D = 1.83x_{tr}$, where D is the effective depth of the turbulent layer and x_{tr} is the observational lag marking the slopes transition. The effective depth is the depth of a layer with depth independent of the lateral coordinate, that would yield the observational structure function.

From Fig.5 the observational transition lag for the nebular velocity $(1.31 \pm 0.04)kpc$ yielding an effective depth of $(2.4 \pm 0.07)kpc$. The observational transition lag of the wind velocity structure function, from Fig.6, is $(1.06 \pm 0.04)kpc$ implying an effective depth of $(1.9. \pm 0.07)kpc$.

APPENDIX

THE THEORETICAL STRUCTURE FUNCTION OF A QUANTITY THAT IS THE RESULT OF INTEGRATION ALONG THE LINE OF SIGHT

Consider a function f(x) where x is a straight line in a lateral direction. and is an integral along the line of sight:

$$f(x) = \int_0^D g(x, z) dz.$$
 (A1)

Here, z is the line of sight coordinate and D the depth. A plane parallel geometry is assumed for simplicity.

The autocorrelation function of f(x is:

$$C_{f}(x) = \langle f(x+x')f(x) \rangle =$$
(A2)
$$\int_{0}^{D} \int_{0}^{D} \langle g(x',z)g(x+x',z') \rangle dzdz' =$$
$$\int_{0}^{D} \int_{0}^{D} C_{g}(x,z-z')dzdz'.$$

The autocorrelation function $C_g(x, z - z')$ can be expressed by the two-dimensional power spectrum, $P_2(k_x, k_z)$,

$$C_g(x, z - z') = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{i(k_x x + k_z(z - z'))}$$
(A3)
$$P_2(k_x, k_z) dk_x dk_z.$$

leading to

$$C_f(x) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{ik_x x} P_2(k_x, k_z) \frac{\sin^2(k_z D/2)}{(k_z D/2)^2} dk_x dk_z.$$
(A4)

from equations (2) and (A4) follows the expression for the structure function

$$S_f(x,D) \propto \int_0^\infty \int_0^\infty \sin^2(k_x x/2) \frac{\sin^2(k_z D/2)}{(k_z D/2)^2}$$
(A5)
 $P_2(k_x,k_z) dk_x dk_z.$

In the case of a turbulence with a one-dimensional power spectrum which is a power law with index -m, the two dimensional power spectrum is

$$P_2(k_x, k_z) \propto \left(k_x^2 + k_z^2\right)^{-(m+1)/2.}$$
 (A6)

resulting in

$$S_f(x,D) \propto \int_0^\infty \int_0^\infty \sin^2(k_x x/2) \frac{\sin^2(k_z D/2)}{(k_z D/2)^2}$$
(A7)
$$\left(k_x^2 + k_z^2\right)^{-(m+1)/2} dk_x dk_z.$$

It is convenient to define the dimensionless variables

$$\eta = k_z D/2.$$
; $\mu = k_x D/2.$ (A8)



FIG. 7.— Theoretical structure function, which is an integral along the line of sight, as function of x/D. The straight lines have logarithmic slopes equaling 1 and 2.



FIG. 8.— Theoretical structure function of a quantity, which is an integral along the line of sight, as function of x/D. The straight line has a logarithmic slope of 1.5.

The structure function of equation (A7) can be expressed as

$$S_f(x,D) \propto \int_0^\infty I(\mu) \sin^2\left(\mu x/D\right) d\mu.$$
 (A9)

where $I(\mu)$ is

$$I(\mu) = \int_0^\infty \left(\mu^2 + \eta^2\right)^{-(m+1)/2} \frac{\sin^2 \eta}{\eta^2} d\eta.$$
 (A10)

Equation (A9) implies that the structure function argument is x/D. Also, inspection of equations (A9) and (A10) reveals that for $x \ll D$ the structure function is proportional to x^m while for x >> D it is proportional to x^{m-1} .

A numerical solution for the case of m = 2 is pre-

- Abramenko, V. I. & Yurchyshyn, V. B. 2020, MNRAS, 497, 5405. doi:10.1093/mnras/staa2427.
- Besserglik, D. & Goldman, I. 2021, ApJ, 915, 117.
- doi:10.3847/1538-4357/ac0247.
- Block, D. L., Puerari, I., Elmegreen, B. G., et al. 2010, ApJ, 718, L1. doi:10.1088/2041-8205/718/1/L1.
- Bournaud, F., Elmegreen, B. G., & Martig, M. 2009, ApJ, 707, L1. doi:10.1088/0004-637X/707/1/L1.
- Burgers, j. M. 1948, Advances in Applied Mechanics, 1, 171.
- Contini, M. & Goldman, I. 2011, MNRAS, 411, 792.
- doi:10.1111/j.1365-2966.2010.17719.x.
- Dame, T. M., Elmegreen, B. G., Cohen, R. S., et al. 1986, ApJ, 305, 892. doi:10.1086/164304.
- Elmegreen, B. G., Kim, S., & Staveley-Smith, L. 2001, ApJ, 548, 749.
- Federrath, C., Klessen, R. S., Iapichino, L., et al. 2021, Nature Astronomy, 5, 365. doi:10.1038/s41550-020-01282-z.
- Forbes, J. C., Emami, R., Somerville, R. S., et al. 2023, ApJ, 948, 107. doi:10.3847/1538-4357/acb53e.
- Goldman, I. & Fleck, R. 2023, MNRAS, 521, 2949.
- doi:10.1093/mnras/stad737
- Goldman, I. 2021a, Physics of Fluids, 33, 071706. doi:10.1063/5.0058074.
- Goldman, I. 2021b, MNRAS, 504, 4493.
- doi:10.1093/mnras/stab1227.
- Goldman, I. 2000, ApJ, 541, 701.
- Green, D. A. 1993, MNRAS, 262, 327.
- doi:10.1093/mnras/262.2.327.
- Hoffmann, K., Laigle, C., Chisari, N. E., et al. 2022, MNRAS, 515, 3603. doi:10.1093/mnras/stac1988.
- Kolmogorov, A. 1941, Akademiia Nauk SSSR Doklady, 30, 301.

This paper was built using the Open Journal of Astrophysics IATFX template. The OJA is a journal which sented in Fig.7together power laws with exponents 1 and 2. In order to find the value of x_{tr}/D where x_{tr} denotes the transition lag between the two slopes a power law with exponent of 1.5 is plotted in Fig.8 together with the structure function. The value of x_{tr}/D is taken to be the value for which the logarithmic slope of the structure function equals 1.5, i.e. where the straight line is tangent to the structure function. The result is $x_{tr}/D \sim 0.547$ thus $D \sim 1.83 x_{tr}$.

REFERENCES

- Larson, R. B. 1981, MNRAS, 194, 809.
- doi:10.1093/mnras/194.4.809.
- Keerthi Vasan G., C., Jones, T., Shajib, A. J., et al. 2024, arXiv:2402.00942. doi:10.48550/arXiv.2402.00942.
- Kritsuk, A. G., Norman, M. L., Padoan, P., et al. 2007, ApJ, 665, 416. doi:10.1086/519443.
- Lazarian, A. & Pogosyan, D. 2000, ApJ, 537, 7.
- Leung, C. M., Kutner, M. L., & Mead, K. N. 1982, ApJ, 262, 583. doi:10.1086/160450.
- Miville-Deschênes, M.-A., Levrier, F., & Falgarone, E. 2003a, ApJ, 593, 831.
- Miville-Deschênes, M.-A., Joncas, G., Falgarone, E., et al. 2003b, A&A, 411, 109.
- Shah, E. A., Kartaltepe, J. S., Magagnoli, C. T., et al. 2022, ApJ, 940, 4. doi:10.3847/1538-4357/ac96eb.
- Nestingen-Palm, D., Stanimirović, S., González-Casanova, D. F., et al. 2017, ApJ, 845, 53. doi:10.3847/1538-4357/aa7e78.
- Passot, T., Pouquet, A., & Woodward, P. 1988, A&A, 197, 228. Roy, J.-R. & Joncas, G. 1985, ApJ, 288, 142. doi:10.1086/162772.
- Rizzo, F., Vegetti, S., Fraternali, F., et al. 2021, MNRAS, 507, 3952. doi:10.1093/mnras/stab2295.
- Sanders, R. L., Shapley, A. E., Jones, T., et al. 2023, ApJ, 942,
- 24. doi:10.3847/1538-4357/aca46. Shah, E. A., Kartaltepe, J. S., Magagnoli, C. T., et al. 2022, ApJ, 940, 4. doi:10.3847/1538-4357/ac96eb.
- Stanimirovic, S., Staveley-Smith, L., Dickey, J. M., et al. 1999, MNRAS, 302, 417. doi:10.1046/j.1365-8711.1999.02013.
- Stutzki, J., Bensch, F., Heithausen, A., et al. 1998, A&A, 336, 697.
- Vázquez-Semadeni, E., Ballesteros-Paredes, J., & Rodríguez, L. F. 1997, ApJ, 474, 292. doi:10.1086/303432.

provides fast and easy peer review for new papers in the astro-ph section of the arXiv, making the reviewing process simpler for authors and referees alike. Learn more at http://astro.theoj.org.