INTERPOLATION IN WEIGHTED PROJECTIVE SPACES

SHAHRIYAR ROSHAN-ZAMIR

ABSTRACT. Over an algebraically closed field, the double point interpolation problem asks for the vector space dimension of the projective hypersurfaces of degree d singular at a given set of points. After being open for 90 years, a series of papers by J. Alexander and A. Hirschowitz in 1992–1995 settled this question in what is referred to as the Alexander-Hirschowitz theorem. In this paper we primarily use commutative algebra to lay the groundwork necessary to prove analogous statements in the weighted projective space, a natural generalization of the projective space. We prove the Hilbert function of general simple points in any ndimensional weighted projective space exhibits the expected behavior. We also introduce an inductive procedure for weighted projective space, similar to that originally due to A. Terracini from 1915, to demonstrate an example of a weighted projective plane where the analogue of the Alexander-Hirschowitz theorem holds without exceptions and prove our example is the only such plane. Furthermore, Terracini's lemma regarding secant varieties is adapted to give an interpolation bound for an infinite family of weighted projective planes.

Contents

1. Introduction	2			
2. Weighted projective space and the interpolation problem				
3. Mixed point interpolation in the weighted projective line				
4. Simple point interpolation in the weighted projective space	11			
5. Double point interpolation in the weighted projective plane	13			
6. The Alexander–Hirschowitz theorem in $\mathbb{P}(1,2,3)$	17			
7. Exceptions	24			
8. Open problems	30			
Appendix A. Proofs related to Section 6	30			
Appendix B. Secant varieties to the Veronese variety	34			
References	38			

The author received support from the National Science Foundation grant DMS-210225. Mathematics Subject Classification: 13A02, 13F20, 14N05.

Key Words: Interpolation, Hilbert Function, Weighted Projective Space, Non-standard Grading.

1. INTRODUCTION

Let k be an algebraically closed field. Consider the vector space of homogeneous polynomials of degree d in n+1 variables having singularities at rgeneral points p_1, \ldots, p_r of the projective space \mathbb{P}^n_k . For a polynomial f to be a member of this vector space it must satisfy r(n + 1) linear conditions on its coefficients imposed by the vanishing of all partial derivatives of f at p_1, \ldots, p_r . It is natural to inquire whether these conditions are independent. This problem was considered by A. Hirschowitz and J. E. Alexander in a series of papers [Hir85, Ale88, AH92a, AH92b, AH95] and finally settled in the form stated below, where $H_d(-)$ denotes the dimension of the vector subspace of homogeneous forms of degree d of the input.

Theorem 1.1 (Alexander-Hirschowitz). Let n, d be positive integers. Let X be a general set of r double points in \mathbb{P}^n_k and $I = I_X$ be its defining ideal. Then

$$H_d(R/I) = \min\left\{r(n+1), \binom{n+d}{d}\right\}$$
(1.1)

with exceptions:

- d = 2 and $2 \le r \le n$;
- d = 3, n = 4 and r = 7; and d = 4, $2 \le n \le 4$ and $r = \binom{n+2}{2} 1$.

We say that a set of double points X has expected dimension in degree d if its ideal I satisfies Equation (1.1). Theorem 1.1 has two equivalent formulations in terms of the dimension of higher secant varieties to the Veronese embedding of \mathbb{P}^n and the Big Waring Rank problem ([HM21, Appendix A]) as well as applications in numerical algebraic geometry [DREE+23], deep polynomial neural networks [KTB19], and algebraic statistics [Cer17, Section 6.2]. The original proof of Theorem 1.1 is based on a degeneration technique called the "la méthode d'Horace différentialle", or méthode d'Horace for short, which is based on an earlier degeneration technique by A. Terracini (Theorem 6.2). Méthode d'Horace was later simplified by K. A. Chandler [Cha01, Cha02]. Further expositions and surveys on the Alexander-Hirschowitz theorem and related problems can be found in [BO08, HM21].

Our main goal is to give an analogous statement to Theorem 1.1 in the weighted projective space denoted $\mathbb{P}(a_0,\ldots,a_n)$, where each a_i is a positive integer. This space, surveyed in [Dol82, BR86], arises naturally in contexts such as toric geometry [RT11], algebraic topology [BFNR13], and K-theory [Amr94]. When $a_i = 1$ for all *i*, one can use Definition 2.1 to recover the usual or straight projective space. The author's main approach has been to modify existing techniques from the usual projective space to fit this new setting. Such adaptions require careful, and often non-trivial, considerations of details. This is best exemplified in Appendix B.

In Section 2 we introduce the weighted projective space, its basic properties and the defining ideal of points. Proposition 2.8 shows that unlike the projective space, even a general point in the weighted projective plane need not be generated by a complete intersection ideal; a fact which is critical to computing I in Theorem 1.1. However, having a weight of 1 forces the ideals of general points to be complete intersections, Corollary 2.9, hence allowing for easy calculations of their ideals.

Section 3 concerns an analogue to Theorem 1.1 in the weighted projective line and shows considering this property passes a litmus test, it holds in dimension 1. Its main result is Proposition 3.2 which we now state.

Proposition 3.2. A general set of points with mixed multiplicity in $\mathbb{P}(a, b)$, where gcd(a, b) = 1, has expected dimension in degree d for all $d \in \mathbb{N}$.

Here, mixed multiplicity refers to having different orders of singularity at the points. While for n = 1 Theorem 1.1 does not require the assumption that the points are general, in contrast, our result excludes the point [0:1], hence the hypothesis of Proposition 3.2 requires general points.

Section 4 is dedicated to proving Theorem 4.1 which concerns simple point interpolation in $\mathbb{P}(a_0, a_1, \ldots, a_n)$. The precise result is as follows.

Theorem 4.1. Let $X \subseteq \mathbb{P}(a_0, a_1, \ldots, a_n)$ be a set of r general simple points with defining ideal I_X . Then for all $d \in \mathbb{N}$ we have

$$H_d(S/I_X) = \min\{r, H_d(S)\}.$$

The proof of Theorem 4.1 in the usual projective space is accessible to an advanced undergraduate student; see [Hui18]. But a proof in the setting above is non-trivial and requires tempering with parameter spaces.

In Section 5 we study sets of double points in weighted projective planes of the form $\mathbb{P}(1, b, c)$. The main novelty of this section is taking a different view point towards the problem: instead of studying interpolation in a *fixed space*, we start with a *fixed size* for a set of general double points, say r, and find examples of spaces in which r general double points have expected dimension in every degree d. Proposition 5.4 proves a necessary condition for this to hold.

Proposition 5.4. If r general double points in $\mathbb{P}(1, b, c)$ have expected dimension in every degree, then c < (r+1)b for $b \neq 1$ and $c \leq r+1$ when b = 1.

The main result of Section 5 is a partial converse to Proposition 5.4.

Proposition 5.8. Let $r \in \mathbb{N}$. If rb < c < (r+1)b then a set of r general double points in $\mathbb{P}(1, b, c)$ has expected dimension in every degree d.

The entirety of Section 6 is dedicated to proving an Alexander–Hirschowitz theorem in $\mathbb{P}(1,2,3)$.

Theorem 6.1. Let $X \subseteq \mathbb{P}(1,2,3)$ be a set of r general double points. Then X has expected dimension in every degree d with no exceptions. Moreover $\mathbb{P}(1,2,3)$ is the only weighted projective space where no exceptions occur.

The most striking part of Theorem 6.1 is having *no exceptions* which is not the case even in the usual \mathbb{P}^2 . Indeed, Theorem 7.11 shows $\mathbb{P}(1,2,3)$ is the only such plane. The main inductive procedure of the proof, Theorem 6.2, is the predecessor to méthode d'Horace; a specialization technique due to Terracini from 1915, [Ter15a]. In Theorem 6.2 we observe Terracini's method can be generalized to $\mathbb{P}(1,\ldots,a_n)$. The difference in numerology is key; \mathbb{P}_k^n contains only one hyperplane up to isomorphism. But in $\mathbb{P}(1,\ldots,a_n)$ there are n + 1 hyperplanes with different numerical characteristics.

Theorem 1.1 provides a finite list of exceptions in \mathbb{P}^2 ; 2 points in degree 2 and 5 points in degree 4. In Example 7.2 we show that general double points, even in a "nice" space such as $\mathbb{P}(1, b, c)$ can have erratic behavior. Nonetheless, Theorem 7.7 gives a linear bound, in terms of the weights, on the degree d, after which a set of points of any size expected dimension.

Theorem 7.7. Let $X \subseteq \mathbb{P}(1, b, c)$ be set of r general double points. Then X has expected dimension in degree d if $d \ge 10c$ or if $d \ge 6c$ and $\left|\frac{2c}{b}\right| \ge 5$.

The case of the projective plane of Theorem 1.1 had been previously elucidated by Terracini in [Ter15b] and by Platini in [Pal03]. Our proof of Theorem 7.7 mimics Terracini's approach. However the bound in our result depends on the weights. This is a case where the numerics of weighted projective space work to our disadvantage: as the weights increase, one has more cases to check.

The essential ingredient in proving Theorem 7.7 is the equivalence of double point interpolation and computing the dimension of the higher secant variety to the Veronese embedding of projective space. This is made precise in Theorem 7.5, the proof of which is relegated to Appendix B. The main requisite in proving this equivalence is due to Lasker Proposition B.5.

We end the introduction highlighting three major themes of this paper: the divergence of notions from the usual projective space when they are considered in the weighted projective space, the difference in numerology and the importance of a weight of 1. For example, in Definition 7.4, one observes when $a_i \neq 1$ for all *i*, the *d*-th Veronese map is not defined on all of $\mathbb{P}(a_0,\ldots,a_n)$, hence it is not an embedding; a fact crucial to the analysis of its secant variety. Other examples of this divergence are the defining ideal of a simple point not being complete intersection, addressed earlier, or stabilization of the Hilbert function at the expected number of independent conditions; see Remark 5.2 (2). In all cases, introducing a single weight of 1 controls this divergence to some extent. All specialization techniques used for the interpolation problem rely heavily on numerical conditions [AH95, Cha01, BO08, Pos12]. While for a standard graded polynomial ring the Hilbert function can be computed using a binomial coefficient, no closed formula exists in the non-standard graded setting beyond 2 variables. Therefore applying degeneration techniques becomes very tricky. The remarkable fact that the Hilbert function for the coordinate ring of $\mathbb{P}(1,2,3)$

happens to have a closed formula, given in Remark 6.6, allowed us to follow Terracini's method in this setting.

Acknowledgments: I would like to thank my advisor, Dr. Alexandra Seceleanu, for her support and infinite patience in providing detailed commentary on earlier versions of this paper. Her help was especially instrumental in the final version of Appendix B and the proof of Lemma 6.9. I thank Benjamin Briggs for spending numerous hours helping me with the gory details of abstract tangent spaces, his help was crucial in my initial proof of Theorem 7.5. I thank Ola Sobieska for introducing me to [BR07] and Jack Jeffries for his help fine tuning my lattice point counting skills. I thank Mahrud Sayrafi for his help with some of my Macaulay2 code. I have benefited enormously from my conversations with Jack Huizenga, Nawaj KC, Claudiu Raicu, Andrew Soto Levins, and Ryan Watson. Finally, I thank Paolo Mantero and Tài Huy Hà for writing [HM21] and Giorgio Ottaviani and Maria Brambilla for writing [BO08], their papers are the main inspirations for this work.

2. Weighted projective space and the interpolation problem

2.1. Weighted projective varieties. Fix integers $a_0, \ldots, a_n \in \mathbb{N}$.

Definition 2.1. The weighted projective space $\mathbb{P} = \mathbb{P}(a_0, \ldots, a_n)$ is a quotient of the affine space \mathbb{A}_k^{n+1} under the equivalence relation

$$(p_0, \ldots, p_n) \sim (\lambda^{a_0} p_0, \ldots, \lambda^{a_n} p_n) \text{ for } \lambda \in k \setminus \{0\}.$$
 (2.1)

Denoting $k^{\times} = k \setminus \{0\}$ the multiplicative group of k, and viewing the above as an action of k^{\times} on \mathbb{A}_k^{n+1} allows to write concisely

$$\mathbb{P}(a_0,\ldots,a_n) = (\mathbb{A}_k^{n+1} \setminus \{0\})/k^{\times}.$$

The usual (or straight) projective space \mathbb{P}_k^n can be recovered from this construction as $\mathbb{P}(1, 1, \ldots, 1)$. We reserve the notation \mathbb{P}_k^n for straight projective space and \mathbb{P} or $\mathbb{P}(a_0, \ldots, a_n)$ for weighted projective space.

Let $S = k[x_0, \ldots, x_n]$ be a polynomial ring with (nonstandard) grading given by $\deg(x_i) = a_i$. Observe that endowing S with this grading is equivalent to giving an action on \mathbb{A}_k^{n+1} as in Equation (2.1). Specifically, a polynomial $f \in S$ is homogeneous of degree d if and only if

$$f(\lambda^{a_0} x_0, \dots, \lambda^{a_n} x_n) = \lambda^d f(x_0, \dots, x_n) \text{ for all } \lambda \in k^{\times}.$$
 (2.2)

Therefore the condition f(p) = 0 is well defined on equivalence classes of \sim in Equation (2.1). One can thus alternatively define weighted projective space as the projective variety $\mathbb{P} = \operatorname{Proj}(S)$. More generally, if I is a homogeneous ideal of S, then the projective variety $\operatorname{Proj}(S/I)$ is the quotient $(V(I) \setminus \{0\})/k^{\times}$. Throughout the paper S always represents the (non-standard) graded homogeneous polynomial ring of $\mathbb{P}(a_0, \ldots, a_n)$.

Much like the usual projective space, weighted projective space is covered by the affine charts $U_i = \{p \in \mathbb{P} \mid p_i \neq 0\}$. The coordinate ring of each U_i is the ring of invariant polynomials $k[x_0, \ldots, x_{i-1}, x_{i+1}, \ldots, x_n]^{\mu_{a_i}}$, where the

group μ_{a_i} is identified with the subgroup of k^{\times} of a_i -th roots of 1 and the group acts as follows: an element $\xi \in k$ with $\xi^{a_i} = 1$ maps $x_j \mapsto \xi^{-a_j} x_j$. By Equation (2.2) a polynomial is invariant under this action if and only if its degree is divisible by a_i leading to a ring isomomorphism

$$k[x_0, \dots, x_{i-1}, x_{i+1}, \dots, x_n]^{\mu_{a_i}} \cong S[x_i^{-1}]_0, f \mapsto \frac{f}{x_i^{\deg(f)/a_i}}.$$
 (2.3)

Thus U_i is Spec $(k[x_0,\ldots,x_{i-1},x_{i+1},\ldots,x_n])/\mu_{a_i} = \mathbb{A}^n/\mu_{a_i}$.

Definition 2.2. The *d*-th Veronese subring of S is the graded ring

$$S^{[d]} = \bigoplus_{i \ge 0} S_{id}.$$
 (2.4)

Although $S^{[d]}$ is a proper subring of S, $\operatorname{Proj}(S^{[d]})$ and $\operatorname{Proj}(S)$ are isomorphic projective varieties, because any homogeneous ratio in S can be expressed as a homogeneous ratio in $S^{[d]}$. There are two different conventions in use about degrees in $S^{[d]}$: one can let the elements of the Veronese subring have the same degree they had in S, or one can divide degrees through by d. The advantage of the latter approach is that there exists a Veronese subring $S^{[D]}$ which is generated by its elements of the smallest degree ([Gro60, Lemma 2.1.6]); in this case, after dividing degrees by D, one can view $S^{[D]}$ as being standard graded. This gives rise to an embedding $\mathbb{P} \subset \mathbb{P}^N_k$, where $N = \dim_k(S_D) - 1$; see Remark B.2.

Another application of this ideal is a reduction to sets of weights that satisfy certain a simple condition which we now describe.

Definition 2.3. A weighted projective space $\mathbb{P}(a_0, \ldots, a_n)$ is well formed if no n-1 of a_0, a_1, \ldots, a_n have a common factor greater than 1.

Remark 2.4. Any weighted projective space is isomorphic to a well formed weighted projective space. Two weighted projective spaces with the same weights are isomorphic up to a reordering of the weights. Henceforth the space $\mathbb{P}(a_0, \ldots, a_n)$ is assumed to be well-formed with weights in *increasing* order $a_0 \leq \cdots \leq a_n$ and $a_i \neq 1$ for some $0 \leq i \leq n$.

2.2. Points in weighted projective space. In this section we describe the ideal defining one point in the weighted projective space and its properties.

Definition 2.5. Let $p \in \mathbb{P}$ be a point. The defining ideal of p is the ideal

 $I_p = (f \in S \mid f \text{ homogeneous}, f(p) = 0).$

Lemma 2.6. Fix a point $p \in \mathbb{P}$. Let T = k[t] and consider the graded k-algebra homomorphism

$$\varphi_p: S \to T, x_i \mapsto p_i t^{a_i}.$$

Then $I_p = \ker(\varphi_p)$ and $S/I_p \cong k[t^{a_i} \mid p_i \neq 0]$. Moreover, if $p, q \in \mathbb{P}$ are points such that for all $0 \leq i \leq n$, $p_i = 0$ if and only if $q_i = 0$, then I_p and I_q are isomorphic S-modules.

Proof. Since φ_p is a degree-preserving homomorphism, $\ker(\varphi_p)$ is a homogeneous ideal. If $f \in S$ is homogeneous of degree d then $f \in \ker(\varphi_p)$ if and only if $f(p_i t^{a_i}) = 0$ if and only if $t^d f(p) = 0$ if and only if $f \in I_p$.

The map $\Theta: S \to S, \Theta(x_i) = \frac{p_i}{q_i} x_i$ (where we make the convention $\frac{0}{0} = 1$) is an automorphism on S if and only if for all $0 \le i \le n, p_i = 0 \iff q_i = 0$. It satisfies $\Theta(I_p) = I_q$ thus demonstrating an isomorphism $I_p \cong I_q$.

In the case $p = \mathbf{1} = [1 : 1 : \dots : 1]$ the ideal $I(\mathbf{1}) = \ker(\varphi_{\mathbf{1}})$ is called a monomial curve ideal. This class of ideals has been studied in [Her70], [Bre79] and [GKW91]. An expository survey of affine monomial curves and their computational aspects can be found in [Sen20]. The particular case of affine space curves is best understood. It was studied in [Her70].

Proposition 2.7 ([Her70, Den03]). Consider positive integers (a, b, c) with gcd(a, b, c) = 1. Let r_i for i = 1, 2, 3 be the smallest positive integer such that the following equations admit a solution in non-negative integers.

$$r_1a = k_1b + g_1c$$

$$r_2b = k_2a + g_2c$$

$$r_3c = k_3a + g_3b$$
(2.5)

Then the kernel of the k-algebra map

$$\varphi: k[z, u, v] \to k[t], z \mapsto t^a, u \mapsto t^b, v \mapsto t^c$$

is the ideal

$$I = (z^{r_1} - u^{k_1} v^{g_1}, u^{r_2} - z^{k_2} v^{g_2}, v^{r_2} - z^{k_3} u^{g_3}).$$

Moreover two of the binomials above agree up to sign if and only if $0 \in \{k_1, k_2, k_3, g_1, g_2, g_3\}$.

If in Equation (2.5) $k_i = 0$ or $g_i = 0$ for any *i*, then we say the weights (a, b, c) satisfy the *Herzog Criteria* (HC).

Proposition 2.8. Consider a well formed weighted projective plane $\mathbb{P}(a, b, c)$ with coordinate ring S = k[z, u, v]. Then the following hold:

(1) If $p_i \neq 0$ for $1 \leq i \leq 3$ then $p = [0: p_1: p_2] \Rightarrow I_p = (z, p_2^b u^c - p_1^c v^b)$ $p' = [p_0: 0: p_2] \Rightarrow I_{p'} = (u, p_2^a z^c - p_0^c v^a)$ $p'' = [p_0: p_1: 0] \Rightarrow I_{p''} = (v, p_1^a z^b - p_0^b u^a).$ (2) Let $p = [p_0: p_1: p_2]$ with $p_i \neq 0$ for all i. Then $I_p = (p_1^{k_1} p_2^{g_1} z^{r_1} - p_0^{r_1} u^{k_1} v^{g_1}, p_0^{k_2} p_2^{g_2} u^{r_2} - p_1^{r_2} z^{k_2} v^{g_2}, p_0^{k_3} p_1^{g_3} v^{r_3} - p_2^{r_3} z^{k_3} u^{g_3})$ where r_i, k_i and g_i are described in Equation (2.5)

Proof. Let $p = [0 : p_1 : p_2]$. The polynomial $p_2^b u^c - p_1^c v^b$ is irreducible because the only homogeneous polynomials in k[u, v] of degree less than bc are monomials. Observe that $J = (z, p_2^b u^c - p_1^c v^b) \subseteq I_p$, and J is a prime

ideal of height 2. Since I_p is also a prime ideal of height two, it follows that $J = I_p$. Other cases of (1) are proven similarly.

Statement (2) follows from Lemma 2.6 and Proposition 2.7. Specifically, the proof of Lemma 2.6 yields that $I_p = \Theta(I_1)$ where $\Theta(x_i) = x_i/p_i$ and I_1 is described as I in Proposition 2.7.

Corollary 2.9. All ideals defining points in the weighted projective plane $\mathbb{P}(a, b, c)$ are complete intersections if and only if a, b, c satisfy (HC).

The next proposition shows that ideals defining points in certain higher dimensional weighted projective spaces are also complete intersections.

Proposition 2.10. Consider $\mathbb{P} = \mathbb{P}(1, 1, ..., 1, a_{i+1}, ..., a_n)$ with $a_j \neq 1$ for all $i + 1 \leq j \leq n$ and let $S = k[x_0, ..., x_n]$ be the coordinate ring of \mathbb{P} . Let $p = [p_0 : ... : p_n]$ where $p_t \neq 0$ for some $0 \leq t \leq i$. Then

$$I_p = (p_0 x_t^{a_0} - p_t^{a_0} x_0, \dots, p_n x_t^{a_n} - p_t^{a_n} x_n).$$

Proof. Note that $J = (p_1 x_t^{a_1} - p_t^{a_1} x_1, \dots, p_n x_t^{a_n} - p_t^{a_n} x_n) \subseteq I_p$ and $S/J \cong k[x_t]$ implies J is a prime ideal. Further height(J) = n because J is a complete intersection with n generators. Since I_p is also a prime ideal of height n the equality $J = I_p$ follows.

We end this section by noting that finding explicit generators for the defining ideal of a point in $\mathbb{P}(a_0, \ldots, a_n)$ for $n \geq 3$ is much harder. In fact, outside of the case of Proposition 2.10 we do not know what the minimal generators of ideals of points are.

Problem 2.11. Find explicit generators for the defining ideal of a general point in $\mathbb{P}(a_0, \ldots, a_n)$ for $n \geq 3$.

2.3. Hilbert function. Throughout \mathbb{N} denotes the natural numbers (not including zero), $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$, k is a field and $S = k[x_0, \ldots, x_n]$ is a polynomial ring in variables x_0, \ldots, x_n with coefficients in k. We endow S with a grading given by $\deg(x_i) = a_i \in \mathbb{N}$.

A graded S-module is a module that admits a decomposition $M = \bigoplus_{i \in \mathbb{Z}} M_i$, where M_i denotes the k vector space of homogeneous elements of M of degree *i*. The Hilbert function of M is the function $H_-(M) : \mathbb{N}_0 \to \mathbb{N}_0$ given by $H_i(M) = \dim_k M_i$.

Notation 2.12. Throughout the paper we denote the Hilbert function of the polynomial ring S by $s_i = \dim_k S_i$.

The integer s_d is the number of monomials of degree d in the polynomial ring S. Computing this number is equivalent to finding the number of nonnegative integer solutions to the Diophantine equation

$$\sum_{i=0}^{n} a_i t_i = d, \text{ where } t_i \in \mathbb{N}_0 \text{ for all } 0 \le i \le n.$$

The case of two variables has explicit formulas which are used repeatedly.

Example 2.13. Let S = k[z, u] with $\deg(z) = a$ and $\deg(y) = b$ where $\gcd(a, b) = 1$. Let p and q be positive integers such that aq - bp = 1. By [BR07, Corollary 1.6] we have

$$s_d = \begin{cases} \left\lfloor \frac{qn}{b} \right\rfloor - \left\lfloor \frac{pn}{a} \right\rfloor & \text{if } a \nmid d \\ \left\lfloor \frac{qn}{b} \right\rfloor - \frac{pn}{a} + 1 & \text{if } a \mid d. \end{cases}$$
(2.6)

In particular for a = 1 this yields

$$s_d = \left\lfloor \frac{d}{b} \right\rfloor + 1. \tag{2.7}$$

Remark 2.14. For a graded S-module M, where x_i is a non-zero divisor of degree a_i on M we have the equation

$$H_d(M) = H_{d-a_i}(M) + H_d(M/(x_i \cdot M)).$$
(2.8)

Equation (2.8) is extensively used in the case of $M = S/I_X$ where I_X defines a set of points contained in U_i .

2.4. The interpolation problem. Based on our assumptions on the field, we can define the symbolic powers of a point as follows.

Definition 2.15. Let $p \in \mathbb{P}$ be a point. The ideal defining the multiple point mp (also termed a fat point ideal or a symbolic power ideal) is

$$I_p^{(m)} = \left\{ f \in S \mid \frac{\partial^{m-1} f}{\partial x_0^{u_0} \cdots \partial x_n^{u_n}}(p) = 0 \text{ where } \sum u_t = m - 1 \right\}.$$

Note when the defining ideal of a simple point is a complete intersection the equality $I_p^{(m)} = I_p^m$ holds for all $m \in \mathbb{N}$. Moreover, for a set $X = \{m_i p_i\}_{i=1}^r$ of multiple points the defining ideal is

$$I_X = \bigcap_{i=1}^r I_{p_i}^{(m_i)}.$$

The ideal above contains the equations of hypersurfaces vanishing to order at least m_i at each point p_i . This interpretation leads to the following.

Problem 2.16. (Interpolation Problem) Find the dimension of the vector space of hypersurfaces of a degree d vanishing to order at least m_i at each points p_i in a given finite set. Equivalently, find the Hilbert function the coordinate ring of a set $X = \{m_i p_i\}_{i=1}^r$ of multiple points.

Using Definition 2.15 for a set $X = \{m_i p_i\}_{i=1}^r$ of points with mixed multiplicity we notice the quantity $\sum_{i=1}^r \binom{n+m_i-1}{n}$ is the number of conditions imposed on degree d hypersurfaces by the vanishing of all partial derivatives of appropriate order. It is natural to inquire if these conditions are linearly independent. Clearly, the vanishing conditions can not be linearly independent if their number exceeds s_d , the dimension of the space of all hypersurfaces of degree d. This brings about the next definition which signifies the expected number of linearly independent conditions imposed by X. **Definition 2.17** (AH_n(d) property). Let $X = \{m_i p_i\}_{i=1}^r \subseteq \mathbb{P}(a_0, \ldots, a_n)$ be a set of r points with defining ideal $I_X \subseteq S$. For $d \in \mathbb{N}$, we say X has property $AH_n(d)$ or X is $AH_n(d)$ if

$$H_d(R/I_X) = \min\left\{s_d, \sum_{i=1}^r \binom{n+m_i-1}{n}\right\}.$$

Proposition 2.19 states a well-known equivalent formulations of interpolation in terms of the evaluation map which is now introduced.

Definition 2.18 (Evaluation Map). Let $X = \{m_i p_i\}_{i=1}^r \subseteq \mathbb{P}(a_0, \ldots, a_n)$ be a set of r double points with $m = \sum_{i=1}^r \binom{n+m_i-1}{n}$. For any $d \in \mathbb{N}$, the evaluation map on X in degree d is the map

$$\operatorname{ev}_X: (S)_d \to k^m$$

$$f \to \left(\frac{\partial^{m_i-1}f}{\partial x_0^{u_0}\cdots \partial x_n^{u_n}}(p_i) = 0 \mid \sum u_t = m_i - 1\right)_{i=1}^r.$$

Proposition 2.19. For $X \subseteq \mathbb{P}(a_0, a_1, \ldots, a_n)$ as defined in Definition 2.18 and any $d \in \mathbb{N}_0$ the following are equivalent:

(1) X is $AH_n(d)$, (2) $ev_X : (S)_d \to k^m$ has full rank.

Remark 2.20. Since I_X is the kernel of the evaluation map and S/I_X is its image, it is clear that

$$H_d(S/I_X) \le \min\{s_d, m\}.$$

Therefore if $H_d(S/I_X)$ is equal to s_d or m then that value is automatically the minimum of the two.

3. MIXED POINT INTERPOLATION IN THE WEIGHTED PROJECTIVE LINE

This subsection focuses on interpolation in $\mathbb{P}(a, b)$, where gcd(a, b) = 1and coordinate ring S = k[z, u] where deg(z) = a and deg(u) = b.

Proposition 3.1. Let I be the defining ideal of a point $p = [p_0 : p_1]$. Then

$$I_p = \begin{cases} (p_1^a z^b - p_0^b u^a) & \text{if } p_0, p_1 \neq 0\\ (z) & \text{if } p_0 = 0 \text{ and } p_1 \neq 0\\ (u) & \text{if } p_0 \neq 0 \text{ and } p_1 = 0. \end{cases}$$

Proof. Note I_p is an unmixed height 1 prime ideal, which implies I_p is principal. Therefore it suffices to find an element of smallest degree in I_p . If $p_0 = 0$ or $p_1 = 0$ this element is evidently z and u, respectively. Assuming $p_0, p_1 \neq 0$ implies no pure power of z or u belongs to I_X . Because gcd(a,b) = 1 and $(p_1^a z^b - p_0^b u^a) \subseteq I_X$, the least degree homogenous element in I_X is $p_1^a z^b - p_0^b u^a$.

Proposition 3.2. A set of general points with mixed multiplicity, $X = \{r_i p_i\}_{i=1}^n \subseteq \mathbb{P}(a, b)$, is $AH_1(d)$ for all $d \in \mathbb{N}$, that is

$$H_d(S/I_X) = \min\{s_d, \sum_{i=1}^n r_i\}.$$
(3.1)

Proof. It suffices to demonstrate one such set. (See Remark 5.1.) Let $X = \{r_1p_1, \ldots, r_np_n\}$ where $p_i = [p_i^{(0)} : p_i^{(1)}]$ and $[0 : p_1] \notin X$. Recall that $s_d = H_d(k[z, u])$ is the number of solutions to

$$ax + by = d$$
 where $x, y \in \mathbb{N}_0$. (3.2)

We wish to interpret $H_d(S/I_X)$ in terms of Equation (3.2). For any *i*, Proposition 3.1 implies $I_{p_i} = ((p_i^{(0)})^b z^b - ((p_i^{(1)})^a u^a))$ is a principal prime ideal and thus

$$I_X = \bigcap_{i=1}^n I_{p_i}^{(r_i)} = \bigcap_{i=1}^n \left((p_i^{(0)})^b z^b - ((p_i^{(1)})^a u^a)^{(r_i)} \right)$$

$$\stackrel{*}{=} \left(\prod_{i=1}^n ((p_i^{(0)})^b z^b - ((p_i^{(1)})^a u^a)^{r_i} \right).$$

Identity * follows since each ideal is a complete intersection. Replacing I_X by its initial ideal (u^{ar}) , where $r = \sum_{i=1}^{n} r_i$, we notice that

$$S/I_X = \operatorname{span}_k \{ z^x u^y \mid 0 \le y \le ar - 1, \ 0 \le x \} \underset{k - \operatorname{vector space}}{\cong} k[z] \otimes_k k[u]/(u^{ar}).$$

Hence $H_d(S/I_X)$ is the number of solutions to Equation (3.2) with the restriction that $0 \le y \le ar - 1$. We will show

$$H_d(S/I_X) = \begin{cases} s_d & \text{if } d < b(ar-1) \\ r & \text{if } d \ge b(ar-1). \end{cases}$$
(3.3)

Suppose d < b(ar - 1). Then $0 \le y \le ar - 2$ regardless of whether we are computing s_d or $H_d(S/I_X)$, which implies that $H_d(S/I_X) = s_d$ holds in this case. Let $d \ge b(ar - 1)$. Then y = ar - i for $1 \le i \le ar$. It suffices to count how many values of i yield a solution. From Equation (3.2) we get

$$ax + b(ar - i) = d \Rightarrow bi + d \equiv 0 \pmod{a} \Rightarrow (-b)^{-1}d \equiv i \pmod{a}$$

Hence $[i]_a = [(-b)^{-1}d]_a$ and $1 \le i \le ar$. Therefore $H_d(S/I_X)$ is equal to the number of distinct coset representatives of $[(-b)^{-1}(d)]_a$ in the interval [1, ar]; of which there are r. Hence $H_d(S/I_X) = r$ for $d \ge b(ar - 1)$. Combining Remark 2.20 and Equation (3.3) implies Equation (3.1)

4. SIMPLE POINT INTERPOLATION IN THE WEIGHTED PROJECTIVE SPACE

The objective of this section is to establish simple point interpolation in $\mathbb{P}(a_0, \ldots, a_n)$. Theorem 4.1 is implicitly used in subsequent proofs of Proposition 5.7, Theorem 6.1 and Theorem 7.7. Our proof is inspired by the one in [Hui18].

Theorem 4.1. Let $\mathbb{P} = \mathbb{P}(a_0, a_1, \dots, a_n)$ and $X = \{p_i\}_{i=1}^r$ be a set of general simple points. X is $AH_n(d)$ for all d, that is

$$H_d(S/I_X) = \min\{s_d, r\}$$

$$(4.1)$$

Proof. Let k be an algebraically closed field. We use induction on r, the number of points in X. Let r = 1 and choose $p = [p^{(0)} : \cdots : p^{(n)}]$ such that $p^{(j)} \neq 0$ for all j. By Lemma 2.6, $S/I_p \cong k[t^{a_j} \mid 0 \leq j \leq n]$. The observations below implies Equation (4.1) for $X = \{p\}$ where $A = \{a_j\}_{j=0}^n$ and $\mathcal{N}(A)$ denotes the numerical semigroup generated A.

$$H_d(S/I_X) = \begin{cases} 1 & \text{if } d \in \mathcal{N}(A) \\ 0 & \text{else} \end{cases}, \quad s_d = \begin{cases} \ge 1 & \text{if } d \in \mathcal{N}(A) \\ 0 & \text{else} \end{cases}$$

Let X' be a set of size r-1 for which Equation (4.1) holds for all d. We claim for any fixed D and some choice of a point p, there exists a set $X = X' \cup \{p\}$ such that Equation (4.1) holds for $H_D(S/I_X)$. If $s_D \leq r-1$, choose $p \notin X'$. Then

$$(I_X)_D \subseteq (I_{X'})_D = 0 \Rightarrow H_D(S/I_X) = s_D < r.$$

If $r-1 < s_D$ then there exists $f \in (I_{X'})_D$ such that $f \neq 0$. Choose $p \in \mathbb{P}$ such that $p \notin X' \cup V(f)$. Such choice is possible because $X \cup V(f) \neq \mathbb{P}$ by the Nullstellensatz. In particular, $f(p) \neq 0$. Observe that $(I_X)_D = \{g \in (I_{X'})_D \mid g(p) = 0\}$ is the kernel of the map

$$(I_{X'})_D \to k$$
 where $g \mapsto g(p)$,

which is surjective since $f(p) \neq 0$. Hence $H_D(I_X) = H_D(I_{X'}) - 1$ and $H_D(S/I_X) = H_D(S) - H_D(I_X) = H_D(S) - H_D(I_{X'}) + 1 = H_D(S/I_{X'}) + 1 = r$, where $r = \min\{s_D, r\}$ by choice of D. This proves the claim.

Because the construction of the set X depended on the degree D, the claim above shows for a *fixed* D there exists a set of size r that satisfies Equation (4.1). Equivalently, there exists a non-empty open set in the parameter space $\mathbb{P}(a_0, \ldots, a_n)^r$, denoted C_D , whose points satisfy Equation (4.1). Next we show the existence of a single set Y of size r, independent of D, that satisfies Equation (4.1) for all degrees d.

We argue to obtain an integer t such that $r \leq s_d$ for all $d \geq t - a_n$. The well-formedness of a_i 's, Remark 2.4, implies $gcd(a_0, a_1, \ldots, a_n) = 1$. Since r is fixed and the Hilbert function of S for $d \gg 0$ is given by a quasi-polynomial

$$H_d(S) = P_i(d)$$
 for $d \equiv i \pmod{\operatorname{lcm}(a_0, \dots, a_n)}$

where each P_i is a polynomial of degree $n \ge 1$ such that $P_i(d) > 0$ for $d \gg 0$, it follows that each sequence $\{P_i(d)\}_d$ is eventually increasing. Thus the existence of the desired t is established. Moreover choose the smallest such integer t. Consider

$$C = \bigcap_{0 \le D \le t + \cdot a_n} C_D \cap \left(\mathbb{P}(a_0, \dots, a_n)^r \setminus Z \right),$$

where Z represents the closed set of r-tuples of points in $\mathbb{P}(a_0, \ldots, a_n)$ where each coordinate hyperplane $V(x_j)$ contains at least one of the points or where any two or more of the points coincide.

By construction C is a non-empty open set since it is the intersection of finitely many non-empty open sets and points of C satisfy Equation (4.1) for all $0 \le D \le t + a_n$. We finish the proof by showing this implies points of C satisfy Equation (4.1) for all d. Fix an r-tuple of C which corresponds to a set $Y \subseteq \mathbb{P}$ of r points. Notice some variable x_j is a non-zero divisor on S/I_Y and Equation (2.8) is now applicable. Next we show

$$H_d(S/(I_Y + x_j)) = 0 \text{ for } t \le d \le t + a_n.$$
 (4.2)

Let $d = t + \gamma$ where $0 \le \gamma \le a_n$. Note $t - a_n \le t - a_j$ and $H_{d'}(S/I_Y) = r$ for all $t - a_n \le d' \le t + a_n$ by choice of t. Plugging d in Equation (2.8) gives

$$H_{t+\gamma}(S/I_Y) = H_{t+\gamma-a_j}(S/I_Y) + H_{t+\gamma}(S/(I_Y + x_j)).$$

Which implies Equation (4.2). Finally, $S/(I_Y + x_n)$ is a graded cyclic S-module and since $[S/(I_Y + x_n)]_d \subseteq [S/(I_Y + x_n)]_{d-a_n}S$ for all $d \ge a_n$, Equation (4.2) yields

$$H_d(S/(I_Y + x_j)) = 0 \text{ for } t + a_n \le d.$$

For an arbitrary choice of Y we have shown

$$H_d(S/I_Y) = \begin{cases} r & d \ge t - a_n \\ s_d & d < t - a_n \end{cases}^* \min\{r, s_d\} \text{ for all } d.$$
(4.3)

Where * follows from Remark 2.20. Thus a set of r general simple points satisfies Equation (4.1).

5. Double point interpolation in the weighted projective plane

This section focuses on interpolation in weighted projective planes of the form $\mathbb{P}(1, b, c)$, with coordinate ring k[z, u, v] with $\deg(z) = 1$, $\deg(u) = b$ and $\deg(v) = c$. We work with the evaluation map to show general double points have the $\operatorname{AH}_2(d)$ property for all d, which recall is to show

$$H_d(S/I_X) = \min\{s_d, 3 \cdot |X|\}.$$

Remark 5.1. It is well-known that in order to prove a general set of points has the $AH_n(d)$ property, it suffices to demonstrate one such set. One can further limit the size of this set to at most two values. For a fixed n and d, it suffices to prove the property $AH_n(d)$ for one set of double points X, where |X| = r and $r = \lfloor \frac{1}{n+1} s_d \rfloor$ and/or $r = \lceil \frac{1}{n+1} s_d \rceil$; see [HM21, Corrollary D.4-D.5]. We employ this in the weighted projective plane where n = 2.

Remark 5.2. The existence of a weight of 1 ensures the following:

(1) The defining ideal of a point is a complete intersection. (Corollary 2.9.)

- (2) We can choose $U_0 = \mathbb{P}(1, b, c) \setminus V(z)$ as our locus of interpolation. Note U_0 is a non-empty open dense set in the parameter space, in this case $\mathbb{P}(1, b, c)$. Hence our notion of general refers to sets of points that are contained in U_0 and is made more precise in different results. The advantage is if $X \subseteq U_0$ is a set of points and $H_d(S/I_X) = (n+1)r$ for some d, then Definition 2.18 and Equation (2.8) imply $H_{d'}(S/I_X) = (n+1)r$ for all $d' \geq d$.
- (3) The coordinate ring of $\mathbb{P}(1, b, c)$, S = k[z, u, v], has non-decreasing Hilbert function. Since z is a degree 1 non-zero divisor, Equation (2.8) yields $H_d(S) \ge H_{d-i}(S)$ for all i hence $\{s_i\}$ forms a non-decreasing sequence.

Proposition 5.4 shows to find a space in which r general double points are $AH_2(d)$ for all d, it is necessary for c to not be too large compared to b and r. Lemma 5.3 provides a numerical observation towards this end.

Lemma 5.3. Fix positive integers b, c and r such that rb < c < (r+1)b. For the coordinate ring S of $\mathbb{P}(1,b,c)$, the smallest d such that $s_d = 3r$ is d = (r-1)b + c.

Proof. For d = (r-1)b + c, observe that $\deg(v^2) > d$ and we get

$$k[z, u, v]_d = k[z, u]_d \oplus v \cdot k[z, u]_{d-c}.$$
(5.1)

This implies by Equation (2.7) the claimed identity

$$s_d = \left\lfloor \frac{d}{b} \right\rfloor + 1 + \left\lfloor \frac{d-c}{b} \right\rfloor + 1 = 3r.$$

To see the minimality of d, note for d' < d, an analogous decomposition to Equation (5.1) implies

$$s_{d'} = \left\lfloor \frac{d'}{b} \right\rfloor + 1 + \left\lfloor \frac{d' - c}{b} \right\rfloor + 1 < 3r.$$

Proposition 5.4. If r general double points are $AH_2(d)$ for all d in $\mathbb{P}(1, b, c)$, then

$$\begin{array}{rrl} c & \leq & (r+1) \mbox{ for } b = 1 \\ c & < & (r+1)b \mbox{ for } b \neq 1 \end{array}$$

Proof. Let $X = \{2p_i\}_{i=1}^r$ be a set of points such that $p_i \neq [0:1:0]$ Assume for contradiction that c > (r+1)b. It suffices to show $s_{2rb} \leq 3r$. This claim implies $(I_X)_{2rb} = 0$. But by Proposition 2.8, the square of the ideal of each point, $I_{p_i}^2$, contains a generator, f_i , of degree 2b. Clearly $F = \prod_{i=1}^r f_i \in I_X$ and deg(F) = 2rb. This contradicts X being AH₂(2rb).

To verify $s_{2rb} \leq 3r$, observe deg $(v^2) = 2c > 2rb$ and Equation (5.1) yields

$$s_{2rb} = \left\lfloor \frac{2rb}{b} \right\rfloor + 1 + \left\lfloor \frac{2rb - c}{b} \right\rfloor + 1 \le 2r + 1 + (r - 2) + 1 \le 3r.$$

5.1. Small number of points in $\mathbb{P}(1, b, c)$. Remark 5.5 explains our proof strategy for the rest of Section 5.

Remark 5.5. Let M_d denote the matrix of $ev_X : (S)_d \to k^{3r}$. We wish to demonstrate one set of double points X, of size r, that is $AH_2(d)$ for all d. If there is a degree d such that $s_d = 3r$, then it suffices to check the rank of M_d is maximal for smallest such d. By Remark 5.2 and Proposition 2.19, it follows that X is $AH_2(d')$ for all $d' \ge d$. Moreover, for $d'' \le d$ having a weight of 1 implies

$$(I_X)_d = 0 \Rightarrow (I_X)_{d''} = 0.$$

Combining this with Remark 5.2 – part 3 implies X is $\operatorname{AH}_2(d'')$ for $d'' \leq d$. For certain numbers r the degree where $s_d = 3r$ is identified in Lemma 5.3.

We show M_d has full rank by exhibiting a set of points at which $\det(M_d)$, a polynomial in 2r variables corresponding to the coordinates of r points, is not zero. Thus $\det(M_d)$ is not the zero polynomial. Hence the notion of general will refer to points whose first coordinate is nonzero and last two coordinates are outside of $V(\det(M_d)) \subseteq \mathbb{P}^{2r}$.

In Proposition 5.6 and Proposition 5.7 we give necessary and sufficient condition for 1 point and 2 points to be $AH_2(d)$ for all d.

Proposition 5.6. One general double point in $\mathbb{P}(1, b, c)$ is $AH_2(d)$ for all d if and only if

- (1) c = 2 when b = 1,
- (2) b < c < 2b when $b \neq 1$.

Proof. (⇒) This is a direct consequence of Remark 2.4 and Proposition 5.4. (⇐) Let $X = \{2p\}$ with $p \notin V(z)$. When b = 1 and c = 2 observe that $H_2(S/I_X) = 3$ because $s_2 = 4$ and $H_2(I_X) = 1$. Further $H_1(S/I_X) = 1$. By a similar reasoning to Remark 5.5 X is AH₂(d) for all d. Suppose b < c < 2b. By Lemma 5.3 and Remark 5.5, it suffices to check AH₂(d) for X in degree d = c. By Proposition 2.8, the smallest degree of a generator in I_X is 2b > c. Hence $H_c(S/I_X) = 3 = \min\{s_c, 3\}$.

Our results so far have been characteristic free. In Proposition 5.7 to ensure certain determinants are non-zero we make assumptions on char(k).

Proposition 5.7. Let k be an infinite field with $char(k) \notin \{2,3,b,b+c\}$. Two general double points in $\mathbb{P}(1,b,c)$, where $(1,b,c) \neq (1,2,3)$, are $AH_2(d)$ for all d if and only if

- (1) b = 1 and c = 2 or c = 3,
- (2) 2b < c < 3b, or
- (3) $\frac{3}{2}b < c < 2b$

Proof. (\Leftarrow) If b = 1, c = 3 and X is a set of two double points, one can use the defining equations of the points to show $H_3(S/I_X) = 5 = s_3$ and $H_4(S/I_X) = 6$. By a similar reasoning to Remark 5.5, X is AH₂(d) for all d. Next let 2b < c < 3b or b=1 and c=2. Lemma 5.3 and direct computations imply $s_{b+c} = 6$ and this is the smallest degree in which S has 6 monomials: $z^{b+c}, z^c u, z^{c-b}u^2, z^{c-2b}u^3, z^b v, uv$. The evaluation matrix with respect to this basis for points with coordinates $[1: p_1: p_2]$ and $[1: q_1: q_2]$ is

$$M_{b+c} = \begin{bmatrix} b+c & cp_1 & (c-b)(p_1)^2 & (c-2b)(p_1)^3 & bp_2 & 0\\ 0 & 1 & 2p_1 & 3(p_1)^2 & 0 & p_2\\ b+c & cq_1 & (c-b)(q_1)^2 & (c-2b)(q_1)^3 & bq_2 & 0\\ 0 & 1 & 2q_1 & 3(q_1)^2 & 0 & q_2\\ 0 & 0 & 0 & 0 & 1 & p_1\\ 0 & 0 & 0 & 0 & 1 & q_1 \end{bmatrix}$$

with determinant $\det(M_{b+c}) = (-1)(p_1 - q_1)^5(b+c)^2 \neq 0.$

Finally suppose $\frac{3}{2}b < c < 2b$. Adapting the proof of Lemma 5.3, one can show d = 3b is the first degree where $s_d = 6$. The evaluation matrix with respect to the basis $z^{3b}, z^{2b}u, z^bu^2, u^3, z^{2b-c}uv, z^{3b-c}v$ and the points fixed before is

$$M_{3b} = \begin{bmatrix} 3b & 2b p_1 & b (p_1)^2 & (p_1)^3 & (2b-c) p_1 p_2 & (3b-c) p_2 \\ 0 & 1 & 2p_1 & 3 (p_1)^2 & p_2 & 0 \\ 0 & 0 & 0 & 0 & p_1 & 1 \\ 3b & 2b q_1 & b (q_1)^2 & (q_1)^3 & (2b-c) q_1 q_2 & (3b-c) q_2 \\ 0 & 1 & 2q_1 & 3 (q_1)^2 & q_2 & 0 \\ 0 & 0 & 0 & 0 & q_1 & 1 \end{bmatrix}$$

where det $(M_{3b}) = 3b(p_1 - q_1)^3 \left[3 b (p_1 - q_1)^2 - 2 p_1^2 - 2 p_1 q_1 - 2 q_1^2 \right].$

(⇒) Suppose a set of two general points, X, is AH₂(d). General in this context refers to X not containing any of the points [0:1:0] or [1:0:0]. When b is 1 the result is immediate from Proposition 5.4 so suppose $b \neq 1$ Proposition 5.4 results in c < 3b and if 2b < c < 3b we are done. Suppose b < c < 2b. We wish to show $\frac{3}{2}b < c$. The equality 3b = 2c is impossible by well-formedness and the hypothesis which excludes the possibility b = 2, c = 3. Therefore one may assume for contradiction that 2c < 3b. Direct calculations show $s_{2c} = 6$. By assumption

$$H_{2c}(S/I_X) = \min\{6, s_{2c}\} = 6.$$

This implies $(I_X)_{2c} = 0$. We arrive at a contradiction by constructing a non-zero curve in $(I_X)_{2c}$. Since b < c < 2b it follows $s_c = 3$ and thus there is a degree c curve with defining equation F passing through the two points. Since $F^2 \in [I_X]_{2c}$ we have reached the desired contradiction.

5.2. A sufficient condition for r points. The main goal of this subsection is to give a sufficient condition for the weights so that r general double points in $\mathbb{P}(1, b, c)$ are $\operatorname{AH}_2(d)$ for all d. The proof of Proposition 5.8 is a generalization of case 1 in Proposition 5.7.

Proposition 5.8. Let $r \in \mathbb{N}$. If rb < c < (r+1)b then a set of r general double points in $\mathbb{P}(1, b, c)$ is $AH_2(d)$ for all d.

Proof. Let $X = \{2p_i\}$, where the points $p_i = [1: p_i^{(1)}: p_i^{(2)}]$ for $1 \le i \le r$ satisfy the following conditions:

- (1) $p_i^{(1)} \neq 0$ and $p_i^{(2)} \neq 0$ for all i, (2) $p_i^{(1)} \neq p_{i'}^{(1)}$ for all $i \neq i'$.

Let c = rb + j and d = (r-1)b + c = (2r-1)b + j which, by Lemma 5.3, is the smallest degree d where $s_d = 3r$. By Remark 5.5 it suffices to show $H_d(S/I_X) = 3r.$

We order the monomials of degree d as follows

$$z^{((2r-1)-s)b+j}u^s, z^{((r-1)-t)b}u^tv$$
 where $0 \le s \le 2r-1, \ 0 \le t \le r-1.$

The rows of the evaluation map correspond to $\frac{\partial}{\partial z}|_{p_i}$, $\frac{\partial}{\partial u}|_{p_i}$ and $\frac{\partial}{\partial v}|_{p_i}$ and the columns to monomials as ordered. Based on our ordering of the monomials, the first 2r monomials do not contain v and the last r do.

The matrix M_d of the evaluation map with rows indexed by $\frac{\partial}{\partial z}|_{p_i}, \frac{\partial}{\partial u}|_{p_i}$ in the first block and $\frac{\partial}{\partial v}|_{p_i}$ in the second block becomes

The $2r \times 2r$ minor in the upper left quadrant corresponds to the evaluation map in degree d for the points $[1:p_i^{(1)}] \in \mathbb{P}(1,b)$. Assumption (1) allows us to view the points in $\mathbb{P}(1, b)$ and this minor is non-zero by Proposition 3.2. The bottom right block is the $r \times r$ Vandermonde minor which is non-zero by assumption (2). Hence M_d has full rank which completes the proof. \Box

6. The Alexander-Hirschowitz theorem in $\mathbb{P}(1,2,3)$

This section is dedicated to the proof of the following result.

Theorem 6.1 (Alexander Hirschowitz theorem in $\mathbb{P}(1,2,3)$). Let k be a field where char(k) $\notin \{2,3,5\}$. Let $X \subseteq \mathbb{P}(1,2,3)$ be a set of r general double points with defining ideal I_X . Then X is $AH_2(d)$ for all d with no exceptions. That is

$$H_d(S/I_X) = \min\{s_d, 3r\} \text{ for all } d.$$

Proof. We use strong induction on d. By Remark 5.1, for each d and for each of the values $r = \left|\frac{1}{3}s_d\right|$ and $r = \left|\frac{1}{3}s_d\right|$ it suffices to demonstrate one set of size r that is $AH_2(d)$. Proposition 6.4 proves 6 base cases for the induction, that is any finite set of general double point is $AH_2(d)$ for $0 \le d \le 5$.

For any $d \ge 6$, and either value of r, we satisfy the hypothesis of Theorem 6.2 which constructs a set of size r that is $AH_2(d)$. This entails showing:

- The numerical condition of Theorem 6.2, Equation (6.1), is satisfied by Lemma 6.8. Let q denote the positive integer mentioned in this condition.
- Part (1) of Theorem 6.2 is the content of Proposition 3.2 when $r_i = 2$ for all i.
- Part (2) of Theorem 6.2 follows because by the induction hypothesis a set of general double points of any size in $\mathbb{P}(1,2,3)$ is $\operatorname{AH}_2(t)$ for $1 \leq t \leq d-1$. In particular, a set of r-q general double points in $\mathbb{P}(1,2,3)$ is $\operatorname{AH}_2(d-i)$ and $\operatorname{AH}_2(d-2i)$ for $i \in \{1,2,3\}$. In Lemma 6.9 we show, using Lemma 6.3 and the previous considerations, that adding q general simple points lying on a projective line, to the set of r-q general double point preserves the $\operatorname{AH}_2(d-i)$ property. This constructs the desired set in part (2) of Theorem 6.2.

Theorem 6.2 is the main inductive step used in Theorem 6.1 and in the standard graded case is proven in [HM21] and [BO08]. The assumption on the ambient space in Theorem 6.2 allows us to conclude the defining ideal of a simple point is a complete intersection (Corollary 2.9, Proposition 2.10), a fact which is fundamental to the proof presented in [HM21]. In particular when $n \geq 3$ all general points in \mathbb{P} are considered in $U = \bigcup_{\{j \mid a_j=1\}} U_j$ to ensure simple points are complete intersections.

Theorem 6.2 (Generalized Terracini's Inductive argument). Consider $\mathbb{P} = \mathbb{P}(1, \ldots, a_n)$, with coordinate ring $S = k[x_0, \ldots, x_n]$ where $\deg(x_i) = a_i$. Assume $1 \le q \le r$ and $d \in Z_+$ satisfy one of the following inequalities for at least one *i* in the range $0 \le i \le n$,

$$(n+1)r - s_{d-a_i} \le nq \le \overline{s}_d \qquad or \qquad \overline{s}_d \le nq \le (n+1)r - s_{d-a_i}, \quad (6.1)$$

where $\overline{S} = S/(x_i)$ is the coordinate ring of the hyperplane $L = V(x_i) \cong \mathbb{P}(a_0, \ldots, \hat{a_i}, \ldots, a_n)$ and $\overline{s}_d = \dim_k(\overline{S}_d)$. If

- (1) a set of q general double points in L is $AH_{n-1}(d)$ and
- (2) the union of a set of r q general double points in \mathbb{P} and a set of q general simple points in L is $AH_n(d-a_i)$

then a set of r general double points in \mathbb{P} is $AH_n(d)$.

Proof. See Appendix A.

It is notable that Terracini originally came up with Equation (6.1) to study configurations of points in \mathbb{P}^3_k . For fixed d and r we refer to the existence of q that satisfies one of the inequalities in Equation (6.1) as "the numerical condition" of Theorem 6.2. In the straight projective space there are infinitely many values of r and d where this condition fails. See [BO08, Section 4]. Even in \mathbb{P}^2_k this numerical condition fails for d = 2, 4 and r = 2, 5, respectively, which are exactly the exceptions in Theorem 1.1.

In Lemma 6.8 we show this numerical condition holds for all d in the case of $\mathbb{P}(1,2,3)$ and based on Theorem 7.11 it is the only such space. The main difference with \mathbb{P}^2_k is the different numerics of projective lines in $\mathbb{P}(1,2,3)$.

Part (2) of the hypothesis of Theorem 6.2 is satisfied with the aid of Lemma 6.3 which is proved in Appendix A. Originally due to K. Chandler [Cha01, Lemma 3], this lemma is modified to our setting and shows if a certain numerical criterion is satisfied, then one can add simple points all lying on a hyperplane to a set of points that are $AH_n(d)$ while preserving the property. This is made precise in Lemma 6.3.

Lemma 6.3 (Chandler's Lemma). Let I be a saturated homogeneous ideal in $S = k[x_0, \ldots, x_n]$ with deg $(x_i) = a_i$, the homogeneous coordinate ring of $\mathbb{P}(a_0, \ldots, a_n)$. Assume l is regular on S/I. Let $\overline{S} = S/(l)$ and deg(l) = i. Fix $d \in \mathbb{Z}_+$. Then following are equivalent:

(1) A general set Y_0 of q reduced points in V(l) satisfies

$$H_d(S/(I \cap J_{Y_0})) = H_d(S/I) + q$$

(2) $H_d(S/I) + q \le H_{d-i}(S/I) + H_d(\overline{S}).$

Next we prove 6 base cases of induction used in proof of Theorem 6.1.

Proposition 6.4. Let k be an algebraically closed field such that $\operatorname{char}(k) \notin \{2,3,5\}$. Let X be a set of r double points with I_X as its defining ideal. Assume no point of X lies on V(z). Further assume for distinct points p_i and p_j in X where $p_i = [p_i^{(1)} : p_i^{(2)} : p_i^{(3)}]$ and $p_j = [p_j^{(1)} : p_j^{(2)} : p_j^{(3)}]$, $p_i^{(2)} \neq p_j^{(2)}$ for all $i \neq j$. Then X is $AH_2(d)$ for all $0 \leq d \leq 5$.

Proof. By Remark 5.1 it suffices to consider $r = \lfloor \frac{1}{3}s_d \rfloor$ and/or $r = \lceil \frac{1}{3}s_d \rceil$. Note that for $d \leq 5$ we have $s_d \leq 5$ and thus $r \leq 2$. Anytime r = 1, the result follows from Proposition 5.6, whose proof requires the points not to lie on V(z). For d = 4 or d = 5 and r = 2, the result follows from Proposition 5.7, whose proof requires $p_i^{(2)} \neq p_j^{(2)}$.

We address a subtly in our notion of general and the construction in Theorem 6.2. The condition that no point of X lies on V(z) in Proposition 6.4 does not obstruct the procedure in Theorem 6.2, which requires specializing q points to a hyperplane, L. These q points need to be general in L; but neither condition (1) or (2) requires q general points in $\mathbb{P}(1,2,3)$.

Let $d \ge 6$. Assume any finite set of general double points X is $AH_2(t)$ for $0 \le t \le d-1$. We will use Theorem 6.2, to show for $r = \lfloor \frac{1}{3}s_d \rfloor$ and $r = \lfloor \frac{1}{3}s_d \rfloor$ that there exists a set of r double points that are $AH_2(d)$. This inductive procedure for d = 14 is demonstrated in Example 6.7, after which the rest of this section is dedicated to establishing the numerical condition in Lemma 6.8 and condition (2) in Lemma 6.9. To study Equation (6.1) in the context of the $\mathbb{P}(1,2,3)$, we introduce the following notation.

Notation 6.5. Let S = k[z, u, v] denote the coordinate ring of $\mathbb{P}(1, 2, 3)$ where deg(z) = 1, deg(u) = 2 and deg(v) = 3 and recall that $s_d = \dim_k(S_d)$. Let S' = k[u, v], S'' = k[z, v] and S''' = k[z, u] denote the coordinate rings of the projective lines $L_1 \cong \mathbb{P}(2, 3)$, $L_2 \cong \mathbb{P}(1, 3)$, and $L_3 \cong \mathbb{P}(2, 3)$ respectively, where $L_1 = V(z), L_2 = V(u), L_3 = V(v)$ and denote the Hilbert function for each of these rings in degree d by s'_d, s''_d, s''_d , respectively.

These values can be computed explicitly.

Remark 6.6. For any $d \in \mathbb{N}_0$ we have

(1) $s_d = \lfloor d^2/12 + d/2 + 1 \rfloor$ according to [Sta99, Page 547]. (2) $s_d''' = \lfloor d/2 \rfloor + 1$ cf. Equation (2.7). (3) $s_d'' = \lfloor d/3 \rfloor + 1$ cf. Equation (2.7). (4) $s_d' = \begin{cases} \lfloor d/6 \rfloor & d \equiv 1 \pmod{6} \\ \lfloor d/6 \rfloor + 1 & \text{otherwise.} \end{cases}$ cf. Equation (2.6).

Example 6.7. We exhibit Terracini's technique for d = 14. Since $s_{14} = 24$, by Remark 5.1 it suffices to show 8 double points are AH₂(14). Checking the numerical condition (Equation (6.3) and Equation (6.4)) the inequality $3(8) - s_{11} = 8 \le s_{14}^{\prime\prime\prime} = 8$ is the only one for which q exists; this corresponds to specializing 4 double points to $\mathbb{P}(1,2)$.

The bottom right part, $\mathbb{P}(1, 2)$ with 4 double points, is considered intrinsically in $\mathbb{P}(1, 2)$ and is $AH_1(14)$ by Proposition 3.2, proving (1) in Theorem 6.2. To justify (2) in Theorem 6.2, we need the bottom left figure, regarded in $\mathbb{P}(1, 2, 3)$, to be $AH_2(11)$. First we show 4 general double points are $AH_2(11)$ then by Lemma 6.3 their union with 4 simple points on $\mathbb{P}(1, 2)$ (empty circles) preserves the $AH_2(11)$ property.

Since $s_{11} = 16$, to show that any number of points is AH₂(11) it suffices to demonstrate $\lfloor \frac{1}{3}s_{11} \rfloor = 5$ and $\lfloor \frac{1}{3}s_{11} \rfloor = 6$ points are AH₂(11). We employ Theorem 6.2 again and focus on 5 points which implies 4 points are AH₂(11). Checking Equation (6.3) and Equation (6.4), all three inequalities yield a value q. We choose $15 - s_8 = 5 \leq s_{11}^{\prime\prime\prime} = 6$ which is to specialize 3 double points to $\mathbb{P}(1, 2)$.

By Proposition 6.4, 2 general double points are $AH_2(8)$ in $\mathbb{P}(1,2,3)$. By Lemma 6.9 their union with 3 simple points on $\mathbb{P}(1,2)$ keeps the configuration $AH_2(8)$. Hence any set of general double points in $\mathbb{P}(1,2,3)$ is $AH_2(14)$.

In Lemma 6.8 we establish the numerical condition of Theorem 6.2.

Lemma 6.8. Let $r = \lfloor \frac{1}{3}s_d \rfloor$ or $r = \lceil \frac{1}{3}s_d \rceil$ and $d \ge 6$. Then the following equalities can not hold simultaneously.

$$3r - s_{d-1} = s'_d = positive \ odd \ integer,$$

$$3r - s_{d-2} = s''_d = positive \ odd \ integer,$$

$$3r - s_{d-3} = s'''_d = positive \ odd \ integer$$
(6.2)

Hence there always exists an even number, say 2q, such that at least one of the following inequalities holds.

$$3r - s_{d-1} \le 2q \le s'_d \qquad s'_d \le 2q \le 3r - s_{d-1} 3r - s_{d-2} \le 2q \le s''_d \qquad (6.3) \qquad s''_d \le 2q \le 3r - s_{d-2} 3r - s_{d-3} \le 2q \le s'''_d \qquad s'''_d \le 2q \le 3r - s_{d-3}$$

Furthermore, one can pick q such that $1 \le q \le r$.

Proof. Assume for the sake of contradiction all equalities in Equation (6.2) hold. The identity $s'_d + s_{d-1} = s_d = 3r$ follows from Equation (2.8). This implies s_d is divisible by 3 thus $r = \lfloor 1/3s_d \rfloor = \lceil 1/3s_d \rceil$.

Let d = 6k + i where $k \ge 1$ since $d \ge 6$. Plugging d into the formulas from Remark 6.6 gives

$$\begin{split} \text{I. } s_d &= \lfloor d^2/12 + d/2 + 1 \rfloor = 3k^2 + (3+i)k + 1 + \lfloor i^2/12 + i/2 \rfloor \\ \text{II. } s_d''' &= \lfloor d/2 \rfloor + 1 = 3k + \lfloor i/2 \rfloor + 1 = \text{odd} \\ \text{III. } s_d'' &= \lfloor d/3 \rfloor + 1 = 2k + \lfloor i/3 \rfloor + 1 = \text{odd} \\ \text{IV. } s_d' &= \begin{cases} k + \lfloor i/6 \rfloor & d \equiv 1 \pmod{6} \\ k + \lfloor i/6 \rfloor + 1 & \text{otherwise} \end{cases} = \text{odd} \end{split}$$

Part III implies $i \neq 3, 4, 5$, otherwise the s''_d is not odd. If i = 2 then IV implies k is even which implies part II, 3k + 2, is even. Hence $i \neq 2$. Since 3 divides s_d , I implies $i \neq 0$. Finally suppose i = 1. Part IV implies k must be odd. Hence 3k + 1 is even which implies II can not hold. It follows that

one of the 6 inequalities in Equation (6.3) or Equation (6.4) holds for any $d \ge 6$. This proves the existence of q.

It remains to show one can pick q to satisfy $1 \le q \le r$. For $1 \le q$, it suffices to show either the lower bound in Equation (6.4) or the upper bound in Equation (6.3), namely s'_d , s''_d and s'''_d is at least 2 for $d \ge 6$. For all i and by Equation (2.8)

$$s''_i \le s''_{i+1}, \ s'''_i \le s'''_{i+1},$$

and $s_6'' = 3$ and $s_6''' = 4$. Therefore $1 \le q$ if either of the bottom two inequalities in Equation (6.3) or Equation (6.4) holds. If d = 6 or $d \ge 8$ we also get $s_d' \ge 2$. For d = 7, observe that

$$3r - s_6 = \begin{cases} 0 & \text{if } r = \left\lfloor \frac{1}{3} s_7 \right\rfloor \\ 3 & \text{if } r = \left\lceil \frac{1}{3} s_7 \right\rceil \end{cases}, \ s_7' = 1.$$

For $r = \lfloor \frac{1}{3}s_7 \rfloor$ the inequality $3r - s_6 \leq s'_7$ holds true but we do not have a choice of q. But we can ignore this and use another inequality because

$$3r - s_{d-3} = 3 \le s_d''' = 4$$

also holds when $r = \lfloor \frac{1}{3}s_7 \rfloor$. Therefore $1 \le q$ for either value of r and $d \ge 6$.

To show $q \leq r$, it is sufficient that either the lower bound in Equation (6.3) or the upper bound in Equation (6.4) is less than or equal to 2r; that is $3r - s_{d-i} \leq 2r$ for $i \in \{1, 2, 3\}$. In Equation (6.3) this is sufficient because if 2r is less than or equal to the right hand side, we may choose q to be the smallest number such that 2q is in the given interval. Note that

$$3r - s_{d-i} \le 2r \Leftrightarrow r \le s_{d-i}.$$

Moreover, considering $r = \left\lceil \frac{1}{3} s_d \right\rceil$ and i = 3 is enough because

$$\left\lfloor \frac{1}{3}s_d \right\rfloor \le \left\lceil \frac{1}{3}s_d \right\rceil \le s_{d-3} \le s_{d-2} \le s_{d-1}$$

Replacing $z = x_i$ and M = S in Equation (2.8) for the equivalence (*) yields

$$\left\lceil \frac{1}{3} s_d \right\rceil \le s_{d-3} \Leftrightarrow \frac{1}{3} s_d < s_{d-3} \Leftrightarrow \frac{1}{3} s_d^{\prime\prime\prime} + \frac{1}{3} s_{d-3} < s_{d-3} \Leftrightarrow s_d^{\prime\prime\prime\prime} < 2 \cdot s_{d-3} \quad (6.5)$$

Proposition A.2 finishes the proof by showing Equation (6.5) for $d \ge 6$. \Box

Lemma 6.9 proves when Lemma 6.8 is satisfied a set X of r-q double points and q simple points on L is $AH_2(d-i)$ for $i \in \{1, 2, 3\}$. By Definition 2.17 this means

$$H_{d-i}(S/I_X) = \min\{s_{d-i}, 3(r-q) + q\}.$$

By Lemma 6.8 we know only one of the six inequalities above may be satisfied for any given d and specified values of r. Therefore the proof shows any one of the inequalities implies (2) in Theorem 6.2. The proof makes repeated use of Lemma 6.3 and the induction hypothesis.

Lemma 6.9. Let $d \ge 6$. By Lemma 6.8 at least one of the inequalities

$$\begin{aligned}
3r - s_{d-1} &\leq 2q \leq s'_d & s'_d \leq 2q \leq 3r - s_{d-1} \\
3r - s_{d-2} &\leq 2q \leq s''_d & s''_d \leq 2q \leq 3r - s_{d-2} \\
3r - s_{d-3} &\leq 2q \leq s'''_d & s'''_d \leq 2q \leq 3r - s_{d-3}
\end{aligned}$$
(6.6)

is satisfied and $1 \leq q \leq r$.

Let L denote the projective line with defining variable of degree i where an inequality is satisfied. Assume for any r' < r a general set of r' double points in $\mathbb{P}(1,2,3)$ is both $AH_2(d-i)$ and $AH_2(d-2i)$. Then the union of a set of r-q general double points in $\mathbb{P}(1,2,3)$ and a set of q general simple points of L is $AH_2(d-i)$.

Proof. Denote by \overline{S} the coordinate ring of L where $i \in \{1, 2, 3\}$ is the corresponding index chosen according to the inequality that holds in Eq. (6.6).

Let Y be a set of r-q general double points in $\mathbb{P}(1,2,3)\setminus L$ with defining ideal I_q and $J_q \subseteq S$ be the defining ideal of q general simple points of L perceived in $\mathbb{P}(1,2,3)$. The ideal $I = I_q \cap J_q$ defines the union of Y with q simple points on L.

Case 1: Suppose $3r - s_{d-i} \leq 2q \leq \overline{s}_d$. We want to show

$$H_{d-i}(S/I) = \min\{s_{d-i}, 3(r-q) + q\} = 3r - 2q,$$

where the last equality follows from the assumption $3r - s_{d-i} \leq 2q$. Towards this end we apply Lemma 6.3. Condition (2) in this lemma combined with Y being $AH_2(d-i)$ and $AH_2(d-2i)$ becomes

$$H_{d-i}(S/I_q) + q \le H_{d-2i}(S/I_q) + H_{d-i}(\overline{S})$$

$$\Leftrightarrow 3r - 2q \le \min\{s_{d-2i}, 3(r-q)\} + \overline{s}_{d-i}.$$

If $H_{d-2i}(S/I_q) = s_{d-2i}$, then the right hand side is $s_{d-2i} + \overline{s}_{d-i} = s_{d-i}$ and the inequality $3r - 2q \leq s_{d-i}$ is the assumption of this case. On the other hand, if $H_{d-2i}(S/I_q) = 3(r-q)$, we need to show

$$3r - 2q \le 3(r - q) + \overline{s}_{d-i} \Leftrightarrow q \le \overline{s}_{d-i}$$

Assumptions of this case imply $q \leq \overline{s}_d/2$. Lemma A.3 illustrates

$$\overline{s}_d/2 \le \overline{s}_{d-i},\tag{6.7}$$

which establishes the desired inequality $q \leq \overline{s}_{d-i}$.

Since condition (2) of Lemma 6.3 holds, by the equivalent condition (1) of this result the ideal $I = I_q \cap J_q$ satisfies

$$H_{d-i}(S/I) = H_{d-i}(S/I_q) + q.$$

By hypothesis Y is $\operatorname{AH}_2(d-i)$ and further $3(r-q) < 3r - 2q \leq s_{d-i}$. Hence the displayed identity leads to the desired conclusion

$$H_{d-i}(S/I_q) = 3(r-q) = \min\{s_{d-i}, 3(r-q)\}.$$

Case 2: Suppose $\overline{s}_d \leq 2q \leq 3r - s_{d-i}$. We wish to demonstrate

$$H_{d-i}(S/I) = \min\{s_{d-i}, 3r - 2q\} = s_{d-i}.$$

Since Y is $AH_2(d-i)$ we have

$$H_{d-i}(S/I_q) = \min\{s_{d-i}, 3(r-q)\}.$$

If $H_{d-i}(S/I_q) = s_{d-i}$, we obtain

$$[I_q]_{d-i} = 0 \Rightarrow [I]_{d-i} = [I_q]_{d-i} \cap [J_q]_{d-i} = 0 \Rightarrow H_{d-i}(S/I) = s_{d-i}.$$

Suppose $H_{d-i}(S/I_q) = 3(r-q) < s_{d-i}$. Set $m = s_{d-i} - 3(r-q)$ and note $m \leq q$ by assumption of this case. Let J_m be the defining ideal of m among the chosen simple points on L, J_{q-m} the ideal defining the remaining simple points, and $F_m = I_q \cap J_m$. It is enough to show $(F_m)_{d-i} = 0$. Then

$$[I]_{d-i} = [I_q]_{d-i} \cap [J_m]_{d-i} \cap [J_{q-m}]_{d-i} = [F_m]_{d-i} \cap [J_{q-m}]_{d-i} = 0$$

implies $H_{d-i}(S/I) = s_{d-i}$. Equivalently we want to show

$$H_{d-i}(S/F_m) = H_{d-i}(S/I_q) + m = s_{d-i}.$$
(6.8)

By Lemma 6.3, Equation (6.8) is true if and only if

$$s_{d-i} \le H_{d-2i}(S/I_q) + \overline{s}_{d-i} = \min\{s_{d-2i}, 3(r-q)\} + \overline{s}_{d-i},$$
 (6.9)

where the last equality follows from Y being $AH_2(d-2i)$. It suffices to show $q \leq \overline{s}_{d-i}$. If true, we use the assumption $s_{d-i} \leq 3r - 2q$ to conclude $s_{d-2i} = s_{d-i} - \overline{s}_{d-i} \leq 3(r-q)$. Then the right hand side of Equation (6.9) becomes s_{d-i} and the inequality claimed therein holds as an equality. Therefore we have established Equation (6.8), which settles the claim as explained above. To verify the remaining inequality $q \leq \overline{s}_{d-i}$, observe that

$$q \leq r \leq \left\lceil \frac{1}{3} s_d \right\rceil$$
 and $s_{d-3} \leq s_{d-2} \leq s_{d-1}$.

The inequality $\left[\frac{1}{3}s_d\right] \leq s_{d-3}$ follows from Eq. (6.5) and Proposition A.2.

7. Exceptions

7.1. An interpolation bound for $\mathbb{P}(1, b, c)$. Let k be an algebraically closed field of characteristic 0. The main goal of this section to give an interpolation bound for weighted projective planes of the form $\mathbb{P}(1, b, c)$ in Theorem 7.7. That is to find $t \in \mathbb{N}$ such that for all $d \geq t$ we can guarantee that any set of r general double points X is $\operatorname{AH}_2(d)$, or not exceptional, i.e.

$$H_d(S/I_X) = \min\{s_d, 3r\}.$$

In general double point interpolation in $\mathbb{P}(1, b, c)$ is drastically different than in $\mathbb{P}(1, 2, 3)$ and \mathbb{P}_k^2 as indicated by Example 7.2.

Definition 7.1. For a set of r general double points $X \subseteq \mathbb{P}(a, b, c)$ we define *deficiency of* X *in degree* d as

$$D(X, d) := \min\{s_d, 3r\} - H_d(S/I_X)$$

By Remark 5.2 when at least one weight is 1 we have $D(X, d) \ge 0$. Further using this terminology, Theorem 1.1 states that D(X, d) = 0 in \mathbb{P}_k^2 with exceptions when d = 2, r = 2 and d = 4, r = 5 and deficiency is exactly 1 in each case. Computations with Macaulay2 [GS] illustrate even in $\mathbb{P}(1, b, c)$ this is not the case.

Example 7.2. We demonstrate some examples of exceptional cases in $\mathbb{P}(1, b, c)$.

Weighted \mathbb{P}^2	# of general points	Deficiency
$\mathbb{P}(1,5,9)$	3	$D(X,d) = 1$ for $20 \le d \le 22$
$\mathbb{P}(1,5,26)$	2	D(X,d) = 2 for $d = 25$
		D(X, d) = 1 for $d = 20 - 24, 26 - 30$
$\mathbb{P}(1,4,57)$	4	D(X,d) = 1 for $d = 32 - 35,65 - 68$
		D(X, d) = 2 for $d = 36 - 39, 61 - 64$
		D(X, d) = 3 for $d = 40 - 43,57 - 60$
		D(X, d) = 4 for $d = 44 - 56$

TABLE 1. Exceptional Cases in $\mathbb{P}(1, b, c)$

Example 7.2 shows that even when the ideal of points in a weighted projective plane are defined by complete intersections, their behavior can be different than points in the usual \mathbb{P}_k^2 . Our Macaulay2 experimentation suggests by increasing the difference between the two largest weights one can arbitrarily increase D(X, d) for a suitable number of general double points.

Problem 7.3. If $d \in \mathbb{N}$ is allowed to vary, is D(X,d) bounded in any weighted projective plane?

Proof of Theorem 7.7 mimics Terracini's proof of Theorem 1.1 for n = 2; [Ter11]. It relies on two two prelimenary results, Theorem 7.5 and Lemma 7.6. We introduce the necessary terminology to state these results.

Definition 7.4. Let $S = k[x_0, \ldots, x_n]$ be the coordinate ring of $\mathbb{P}(a_0, \ldots, a_n)$, $N = s_d - 1, m_0, \ldots, m_N$ enumerate the monomials in S_d and I be the kernel of the k-algebra homomorphism

$$\theta: k[y_0, \dots, y_N] \to k[x_0, \dots, x_n]$$
$$y_i \to m_i.$$

The projective variety

$$V^d = \operatorname{Proj}(k[y_0, \dots, y_N]/I) \subset \mathbb{P}_k^N$$

is called the *d*-th Veronese variety of $\mathbb{P}(a_0, \ldots, a_n)$. Geometrically θ induces a map of projective varieties

 $\psi: \mathbb{P}(a_0, \ldots, a_n) \dashrightarrow V^d, \ \psi(p) = [m_0(p): \cdots: m_N(p)]$

defined on $U = \{ p \in \operatorname{Proj}(S) \mid \theta((y_0, \dots, y_N)) \not\subseteq p \}.$

Note that unlike the straight projective space, the Veronese map is only partially defined on $\mathbb{P}(a_0, \ldots, a_n)$. Henceforth the notion of a set of general points X includes $X \subseteq U$.

By $\sigma_r(V^d)$ we denote the *r*-th secant variety to the image of the Veronese map. For a point $p \in V^d$, we let T_pV^d denote the *tangent space of* V^d *at p*. For a precise definition of each see Definition B.7 and Definition B.3.

Theorem 7.5. For a fixed $d \ge a_n$ and a set of r general double points $X \subseteq \mathbb{P}(1, a_1, \ldots, a_n)$, the following are equivalent:

(1) $H_d(S/I_X) = \min\{s_d, r(n+1)\}\)$ (2) $\dim_k(\sigma_r(V^d)) = \min\{s_d - 1, r(n+1) - 1\}.$

Proof. See Appendix B.

Lemma 7.6. (Second Terracini lemma) Let U be as in Definition 7.4. Let $X = \{2p_i\}_{i=1}^r \subseteq U \subseteq \mathbb{P}(a_0, \ldots, a_n)$ be a set of double points with defining ideal I_X and fix $d \ge a_n$. We identify the reduced points p_i with their images on V^d according to the Veronese map.

Assume that $H_d(S/I_X) < \min\{s_d, r(n+1)\}$. Then there is a positive dimensional variety $C \subseteq V^d$ through $p_1, \ldots p_r$ such that if $p \in C$ then $T_pV^d \subseteq \langle T_{p_1}V^d, \ldots, T_{p_r}V^d \rangle$. In particular, by Proposition B.5, every hypersurface of degree d which is singular at p_i is also singular along C.

Proof. See [BO08, Lemma 2.3]

2

Theorem 7.7. A set of r general double points $X \subseteq \mathbb{P}(1, b, c)$ is $AH_2(d)$,

$$H_d(S/I_X) = \min\{s_d, 3r\},$$
 (7.1)

for all $d \ge 10c$. Additionally, Equation (7.1) holds if $d \ge 6c$ and $\left|\frac{2c}{b}\right| \ge 5$.

Proof. Let X be a set of r general double that does not satisfy Equation (7.1) for some degree $d \ge c$. There exists $F \in (I_X)_d$ where $F \ne 0$. By Lemma 7.6, there is a positive dimensional variety $C \subseteq V^d$ through the image of the points in V^d . By abuse of notation we now identify C with its preimage under the Veronese map. By the last part of Lemma 7.6, F is singular along C. Hence F contains a double component through X, namely $F = C^2 H$. Let deg(C) = l. Since one of the weights is 1, Remark 5.2 (3) implies

$$2l \le d \Rightarrow l \le \left\lfloor \frac{d}{2} \right\rfloor \Rightarrow s_l \le s_{\lfloor \frac{d}{2} \rfloor}.$$
 (7.2)

By Lemma 7.6, C vanishes on the reduced set $\{p_i\}_{i=1}^r$. From Theorem 4.1 and Remark 5.1 we get

$$\left\lfloor \frac{1}{3}s_d \right\rfloor \le r \le s_l - 1. \tag{7.3}$$

26

Putting Equation (7.2) and Equation (7.3) together we acquire

$$\left\lfloor \frac{1}{3}s_d \right\rfloor \le s_{\lfloor \frac{d}{2} \rfloor} - 1. \tag{7.4}$$


Hence for values of d where Equation (7.4) does not hold, any set X satisfies Equation (7.1). In Lemma 7.8 we show that the inequalities involving dgiven in the hypothesis force Equation (7.4) to fail. This completes the proof.

Lemma 7.8. If either of the following is true then $\lfloor \frac{1}{3}s_d \rfloor \ge s_{\lfloor \frac{d}{2} \rfloor}$ holds.

(1) $d \ge 10c$ (2) $d \ge 6c$ and $\left\lfloor \frac{2c}{b} \right\rfloor \ge 5$

Proof. We employ an interpretation of the Hilbert function of a nonstandard graded polynomial ring as lattice point counting.

Observe s_d is the number of lattice points on and inside the triangle T with vertices $(0,0), (0,\frac{d}{c}), (\frac{d}{b}, 0)$. Recall s_d is the number solutions to $r_0 + br_1 + cr_2 = d$ where $r_i \in \mathbb{N}_0$. Which are in bijection with lattice points (r_1, r_2) where $br_1 + cr_2 \leq d$, $r_1 \geq 0$ and $r_2 \geq 0$. The latter is the number of lattice points on and inside of T which we now divide into smaller triangles.



Triangle T_1 has vertices at (0,0), $\left(0, \frac{\lfloor \frac{d}{2} \rfloor}{c}\right)$, $\left(\frac{\lfloor \frac{d}{2} \rfloor}{b}, 0\right)$. Our goal is to impose conditions on b and c so that T_1 , T_2 and T_3 have small intersections and T_4 contains enough lattice points to imply the desired inequality. By #T we refer to the number of lattice points on and inside the triangle T. Note that

- #T₁ = s_{⊥d/2},
 T₂ and T₃ are translates of T₁ by integer vectors hence

$$\#T_2 = \#T_3 = \#T_1 = s_{\lfloor \frac{d}{2} \rfloor}.$$

We make the following observations

(1) $T_1 \cap T_2 = \{ \left(\left| \frac{d}{2b} \right|, 0 \right) \}.$

(2)
$$T_2 \cap T_3 = \left\{ \left(\left\lfloor \frac{d}{2b} \right\rfloor, \left\lfloor \frac{d}{2c} \right\rfloor \right) \right\}$$

(3) $T_1 \cap T_3 \subseteq \left\{ \left(t, \left\lfloor \frac{d}{2c} \right\rfloor \right) \mid 0 \le t < \frac{c}{b} \right\}$
(4) $T_1 \cap T_2 \cap T_3 = \emptyset.$

Item (4) follows from (2) once we check the point $\left(\left\lfloor \frac{d}{2b}\right\rfloor, \left\lfloor \frac{d}{2c}\right\rfloor\right)$ is not in T_1 . The slope of the hypotenuse of T_1 is $\frac{-b}{c}$, so the point is in T_1 if and only if $\left\lfloor \frac{d}{2c} \right\rfloor \leq \frac{b}{c} < 1$. But this can't happen since $d \geq 2c$ implies $\left\lfloor \frac{d}{2c} \right\rfloor \geq 1$.

Using (1) – (4) above and denoting by $\#\mathring{T}_4$ the number of lattice points in the interior of T_4 , we obtain

$$s_{d} = \#T$$

$$\geq \#T_{1} + \#T_{2} + \#T_{3} - \#(T_{1} \cap T_{2}) - \#(T_{2} \cap T_{3}) - \#(T_{1} \cap T_{3}) + \#\mathring{T}_{4}$$

$$\geq 3s_{\lfloor \frac{d}{2} \rfloor} - 3 - \lfloor \frac{c}{b} \rfloor + \#\mathring{T}_{4}.$$
(7.5)

We now estimate $\#\mathring{T}_4$. Let (p,q) be the point of intersection between the hypotenuse of T_1 and the line $x = \lfloor \frac{d}{2b} \rfloor$. Using the figure above one can show

$$q < i < \left\lfloor \frac{d}{2c} \right\rfloor$$
 and $\frac{\left\lfloor \frac{d}{2} \right\rfloor}{b} - \frac{ic}{b} < t < \left\lfloor \frac{d}{2b} \right\rfloor \iff (t,i) \in \mathring{T}_4$ (7.6)

For a fixed *i*, we obtain $\lfloor \frac{ic}{b} \rfloor - 1 \ge i - 1$ lattice points in $\#\mathring{T}_4$. Let $d \ge 10c$. From $\lfloor \frac{d}{2c} \rfloor \ge 5$ and Equation (7.6) we get $i \le 4$. Since t_5 is similar to T and $\frac{\lfloor \frac{d}{2} \rfloor}{b} - \lfloor \frac{d}{2b} \rfloor \le 1$ we get q < 1, hence $1 \le i$. Plugging $1 \le i \le 4$ in Equation (7.6) one gets

$$\#\mathring{T}_4 \ge \left\lfloor \frac{c}{b} \right\rfloor + 5.$$

Plugging this into Equation (7.5) yields the following and proves (1).

$$s_d \ge 3s_{\lfloor \frac{d}{2} \rfloor} - \lfloor \frac{c}{b} \rfloor - 3 + \lfloor \frac{c}{b} \rfloor + 5 = 3s_{\lfloor \frac{d}{2} \rfloor} + 2.$$

Suppose $d \ge 6c$ and $\lfloor \frac{2c}{b} \rfloor \ge 5$. By $\lfloor \frac{d}{2c} \rfloor \ge 3$ and Equation (7.6) we get

$$\#\mathring{T}_4 \ge \left\lfloor \frac{c}{b} \right\rfloor - 1 + \left\lfloor \frac{2c}{b} \right\rfloor - 1.$$

By $\lfloor \frac{2c}{b} \rfloor \ge 5$ and Equation (7.5) we arrive at the desired conclusion

$$s_d \ge 3s_{\lfloor \frac{d}{2} \rfloor} - \lfloor \frac{c}{b} \rfloor - 3 + \lfloor \frac{c}{b} \rfloor - 1 + \lfloor \frac{2c}{b} \rfloor - 1 \ge 3s_{\lfloor \frac{d}{2} \rfloor}.$$

7.2. Exceptions in $\mathbb{P}(1, b, c)$. So far we have shown exceptional sets of points in $\mathbb{P}(1, b, c)$ only happen when d < 10c. As a porism of Theorem 7.7 we get Corollary 7.10 which fully classifies exceptional cases of $\mathbb{P}(1, b, c)$ in degrees d where s_d is divisible by 3. Proposition 7.9 gives a sufficient condition for a set of points to fail Equation (7.1).

Proposition 7.9. Fix $d \in \mathbb{N}$. Let $X \subseteq \mathbb{P}(1, a_1, \ldots, a_n)$ be a set of r general simple points. Let J_X and I_X be the defining ideal of X and 2X respectively; with 2X denoting the set of double points supported on X. If $r < s_{\lfloor \frac{d}{2} \rfloor}$ and $(n+1)r \ge s_d$ then X is not $AH_n(d)$.

Proof. Let $e = \lfloor \frac{d}{2} \rfloor$. By Theorem 4.1 and the assumption $r < s_e$, there exists $F \in (J_X)_e$ where $F \neq 0$. The proof follows from the following simple observation.

$$F^2 \in (J_X^2)_{2e} \subseteq (I_X)_{2e}$$

However the expected dimension of $(I_X)_{2e} = 0$ because

$$(n+1) \cdot r \ge s_d \ge s_{2e}$$

where (*) follows since $a_0 = 1$ (Remark 5.2 (3)). Thus X is not $AH_n(d)$.

Corollary 7.10. Let $d \in \mathbb{N}$ be such that $s_d = 3r$. Then a set $X \subseteq \mathbb{P}(1, b, c)$ of r general double points is not $AH_2(d)$ if and only if $r < s_{\lfloor \frac{d}{n} \rfloor}$.

Proof. Let X be as defined in the hypothesis. Using Proposition 7.9 for (\Rightarrow) and Equation (7.4) for (\Leftarrow) , which is a consequence of not being $AH_2(d)$ as shown in the proof of Theorem 7.7, we get

 $\begin{aligned} r &= \frac{1}{3} s_d \leq s_{\lfloor \frac{d}{2} \rfloor} - 1 < s_{\lfloor \frac{d}{2} \rfloor} \\ &\iff X \text{ does does not impose independent conditions in degree } d. \quad \Box \end{aligned}$

Using Corollary 7.10 one can prove $\mathbb{P}(1,2,3)$ is the only space of the form $\mathbb{P}(1,b,c)$ with no exceptional sets of general double points.

Theorem 7.11. Let $X \subseteq \mathbb{P}(1, b, c)$ be a finite set of general double points. Then X is $AH_2(d)$ for all $d \in \mathbb{N}$ if and only if b = 2 and c = 3.

Proof. (\Rightarrow) The proof is broken into the cases b = 1, b = 2 and $b \ge 3$ where r denotes the number of points in X. If b = 1, c = 2 and r = 3 then X is exceptional because $s_2 = 4$ and by Theorem 4.1 there exists a quadratic form F through the 3 points resulting in $F^2 \in [I_X]_4$. But the fact that $s_4 = 9$ yields

$$H_{S/I_X}(4) \le 8 < 9 = \min\{9, 9\}.$$

If $c \geq 3$ by Proposition 5.4 one general double point is exceptional.

For b = 2 and r = 1, Proposition 5.6 implies c < 4 and further 2 < c by Remark 2.4, therefore c = 3 and we are done. Let $b \ge 3$. We contradict the hypothesis by proving it can not hold for r = 1, 2, 3 points and all dsimultaneously. When r = 2, Proposition 5.7, which uses the assumption $b \ne 2$, implies $\frac{3}{2}b < c < 2b$ or 2b < c < 3b. By Proposition 5.6 the latter would contradict 1 point being AH₂(d) for all d therefore $\frac{3}{2}b < c < 2b$. This criteria on c results in $s_{4b} = 9$ and $s_{2b} = 4$ thus Corollary 7.10 implies 3 points are not AH₂(d). Theorem 6.1 yields the backwards direction.

8. Open problems

We end this paper with a list of open problems, some of which have been stated through out the paper.

- (1) With possible restrictions on the weights, find explicit generators for the defining ideal of a general point in $\mathbb{P}(a_0, \ldots, a_n)$ for $n \ge 3$.
- (2) Using the notation introduced in Section 7, is D(X, d) bounded *uni-formly* (independently of d) in any weighted projective plane?
- (3) Let $\mathbb{P}(1, a_1, \ldots, a_n)$ and $g = \text{lcm}(a_1, \ldots, a_n)$. One can show if $g \mid d$ then the Veronese map in Definition 7.4 becomes an embedding. Compute the minimal free resolution of the ideal defining the Veronese variety V^d for such d.
- (4) The well-known SHGH conjecture, see [CHHVT20], states the exceptional degrees for a set $X \subseteq \mathbb{P}^2$ where X contains points with mixed multiplicity. State the analogue of this conjecture for $\mathbb{P}(1, b, c)$.
- (5) We suspect that méthode d'Horace is applicable in $\mathbb{P}(1, a_1, \ldots, a_n)$ for $d \geq \operatorname{lcm}(a_1, \ldots, a_n)$. But we could not adapt the details of the modern proof due to Chandler, in particular the use curvilinear subschemes, in the setting of weighted projective space. We leave it as a conjecture that méthode d'Horace, or a simple modification of it, applies in weighted projective space.

Appendix A. Proofs related to Section 6

This section contains proofs for several foundational results used in Section 6 adapted from [HM21] and [Cha01] as well as proofs for some numerical inequalities we employ in that section. We use Notation 2.12. Our proof of Theorem 6.2 mimics the one presented in [HM21]; with expansion on two details that were not clear to us. The proof of Lemma 6.9 gives an accessible proof of [Cha01, Lemma 3] when all the weights are 1.

Let \mathfrak{m} be the homogeneous maximal ideal of a graded ring and let I be any ideal. The ideal $(I:\mathfrak{m}^{\infty})$ is called *the saturation of* I, denoted I^{sat} .

Lemma A.1 (Generalized Castelnuovo's Inequality). Let S be a not necessarily standard graded polynomial ring, I a homogeneous ideal and l a form of degree k in S. Set $\tilde{I} = I : l$, $\overline{S} = S/(l)$ and $\overline{I} = (I + (l))/(l)$, the ideal of I in \overline{S} . Then

$$H_d(S/I) \ge H_{d-k}(S/\tilde{I}) + H_d(\overline{S}/(\overline{I})^{sat}),$$

with equality if and only if \overline{I} is saturated in \overline{R} .

Proof. See [HM21, Lemma 2.2].

We note the assumption on the ambient space in Theorem 6.2 allows us to conclude the defining ideal of a simple point is a complete intersection (Corollary 2.9, Proposition 2.10) which is implicitly used through out the proof. For a set of reduced points X with defining ideal J_X , let 2X denote the set of double points supported on X where $I_X = J_X^{(2)}$; see Definition 2.15.

Theorem 6.2 (Generalized Terracini's Inductive argument).

Consider $\mathbb{P} = \mathbb{P}(1, a_1, \ldots, a_n)$, with the coordinate ring $S = k[x_0, \ldots, x_n]$ where $\deg(x_i) = a_i$. Assume $1 \leq q \leq r$ and $d \in \mathbb{N}$ satisfy one of the following inequalities for at least one *i* in the range $0 \leq i \leq n$,

$$(n+1)r - s_{d-a_i} \le nq \le \overline{s}_d$$
 or $\overline{s}_d \le nq \le (n+1)r - s_{d-a_i}$, (A.1)

where $\overline{S} = S/(x_i)$ is the coordinate ring of the hyperplane $L = V(x_i) \cong \mathbb{P}(a_0, \ldots, \hat{a_i}, \ldots, a_n)$ and $\overline{s}_d = \dim_k(\overline{S}_d)$. If

- (1) a set of q general double points in L is $AH_{n-1}(d)$ and
- (2) the union of a set of r q general double points in \mathbb{P} and a set of q general simple points in L is $AH_n(d-a_i)$

then a set of r general double points in \mathbb{P} is $AH_n(d)$.

Proof. If $n \geq 3$ all general points in \mathbb{P} are considered in $U = \bigcup_{\{j \mid a_j = 1\}} U_j$. Let Y_1 be a set of q general simple points in L, with $\overline{J_{Y_1}} \subseteq \overline{S}$ as its defining ideal. Note $Y_1 \subseteq \mathbb{P}$ has the defining ideal $J_{Y_1} = (\overline{J_{Y_1}}, x_i)S$. Let Y_2 be a set of r - q general simple points in $\mathbb{P} \setminus L$ with defining ideal $J_{Y_2} \subseteq S$. Let

 $I = J_{Y_1}^{(2)} \cap J_{Y_2}^{(2)}$ be the defining ideal of $2Y = 2Y_1 \cup 2Y_2$ and $\overline{I} = \frac{I + x_i}{x_i}$. We show 2Y has the AH₂(d) property.

For any d, the definition of Hilbert function and Definition 2.18 imply $H_d(S/I) \leq s_d$ and $H_d(S/I) \leq r(n+1)$ hence it suffices to show $H_d(S/I) \geq \min\{s_d, r(n+1)\}$. Since $Y_1 \subseteq L$ and $Y_2 \subseteq \mathbb{P} \setminus L$ we get

$$\tilde{I} := (I : x_i) = (J_{Y_1}^{(2)} \cap J_{Y_2}^{(2)} : x_i) = (J_{Y_1}^{(2)} : x_i) \cap (J_{Y_2}^{(2)} : x_i) = J_{Y_1} \cap J_{Y_2}^{(2)}.$$

To justify the final equality we will show $(J_{Y_1}^{(2)}: x_i) \subseteq J_{Y_1}$. Observe

$$s \in (J_{Y_1}^{(2)} : x_i) = (\bigcap_{j=1}^q J_{p_j}^{(2)} : x_i) = \bigcap_{j=k}^q (J_{p_j}^2 : x_i) \Rightarrow s \cdot x_i \in \bigcap_{i=k}^q J_{p_j}^2.$$

We want to show $s \in J_{p_j}$ for all j.

Fix a j and note $s \cdot x_i = 0$ in $\operatorname{gr}_{J_{p_j}} S$, where $s \in S/J_{p_j}$ and $x_i \in J_{p_j}/J_{p_j}^2$, but clearly $x_i \notin J_{p_j}^2$. Using the generators of J_{p_j} from Corollary 2.9 or Proposition 2.10, we can show $S_{J_{p_j}}$ is a regular ring, hence $\operatorname{gr}_{J_{P_j}} S$ is a domain. Therefore $s \in J_{p_j}$ for all j. Note \tilde{I} defines a set of q general simple points in L and r - q general double points in \mathbb{P} .

Claim: $(\overline{I})^{\text{sat}} = \overline{J_{Y_1}}^{(2)}$. Observe that $\overline{I} \subseteq \overline{J_{Y_1}}$. Since $\overline{S}/\overline{J_{Y_1}}$ is a onedimensional ring, $(\overline{I})^{(\text{sat})}$ is the intersection of minimal components of \overline{I} . These are minimal primes containing $I = J_{Y_1}^{(2)} \cap J_{Y_2}^{(2)}$ and x_i . Since no minimal prime of J_{Y_2} contains x_i , we get $\text{Min}\{(I, x_i)\} = \text{Min}\{J_{Y_1}\}$. For any $\mathfrak{p} \in Min\{(I, x_i)\}$ we get

$$(I, x_i)_{\mathfrak{p}}^{\mathrm{sat}} = (I, x_i)_{\mathfrak{p}} = (J_{Y_1}^{(2)} \cap J_{Y_2}^{(2)}, x_i)_{\mathfrak{p}} = (J_{Y_1}^{(2)}, x_i)_{\mathfrak{p}} \cap (J_{Y_2}^{(2)}, x_i)_{\mathfrak{p}}$$
$$= (J_{Y_1}^{(2)}, x_i)_{\mathfrak{p}} = (J_{Y_1}^{(2)}, x_i)_{\mathfrak{p}}^{\mathrm{sat}}$$

This proves the claim. By Lemma A.1 we get

$$H_d(S/I) \ge H_{d-a_i}(S/\tilde{I}) + H_d(\overline{S}/\overline{I}^{\text{sat}}) = \min\{s_{d-a_i}, (n+1)(r-q) + q\} + \min\{\overline{s}_d, nq\}$$
(A.2)

where the last equality follows from assumptions (1) and (2).

If $(n+1)r - s_{d-a_i} \leq nq \leq \overline{s}_d$ holds, the right hand side of Equation (A.2) becomes (n+1)r. If $\overline{s}_d \leq nq \leq (n+1)r - s_{d-a_i}$ holds, the right hand side of Equation (A.2) is $s_{d-a_i} + \overline{s}_d = s_d$ by Remark 2.14. Note if $H_d(S/I)$ is equal to (n+1)r or s_d then that value is the minimum of the two.

Lemma 6.3 (Chandler's Lemma). Let I be a saturated homogeneous ideal in $S = k[x_0, \ldots, x_n]$ with $\deg(x_i) = a_i$, the homogeneous coordinate ring of $\mathbb{P}(a_0, \ldots, a_n)$. Assume l is regular on S/I. Let $\overline{S} = S/(l)$ and $\deg(l) = i$. Fix $d \in \mathbb{Z}_+$. Then following are equivalent:

(1) A general set Y_0 of q reduced points in V(l) satisfies

$$H_d(S/(I \cap J_{Y_0})) = H_d(S/I) + q.$$
 (A.3)

(2) $H_d(S/I) + q \leq H_{d-i}(S/I) + H_d(\overline{S}).$

Proof. (1) \Rightarrow (2) Since $1 \cdot l \in J_{Y_0}$ and l is regular on S/I we get

$$(I \cap J_{Y_0}: (l)) = (I: (l)) \cap (J_{Y_0}: (l)) = (I: (l)) \cap S = (I: (l)) = I$$

Using this equality and the restriction sequence

$$0 \to S/(I \cap J_{Y_0}: l)(-i) \xrightarrow{\cdot l} S/(I \cap J_{Y_0}) \twoheadrightarrow S/(I \cap J_{Y_0}, l) \to 0$$

we get

$$H_d(S/I \cap J_{Y_0}) = H_{d-i}(S/I) + H_d(S/(I \cap J_{Y_0}, l)) \le H_{d-i}(S/I) + H_d(\overline{S}).$$

 $(2) \Rightarrow (1)$ Suppose there exists a positive integer q satisfying the inequality in (2). We proceed by induction on j where

$$0 \le j \le H_{d-i}(S/(I:l)) + H_d(\overline{S}) - H_d(S/I)$$

to construct a set Y(j) of j reduced points in V(l) that satisfies

$$H_d(S/(I \cap J_{Y(j)})) = H_d(S/I) + j.$$
 (A.4)

If j = 0, let $Y(0) = \emptyset$. For $0 \le j < q$ suppose there is a set $Y(j) \subseteq V(l)$ that satisfies Equation (A.4). Since $j + 1 \le q$ assumption (2) gives

$$j+1 \le q \le H_{d-i}(S/(I:l)) + H_d(\overline{S}) - H_d(S/I)$$

from which it follows

$$H_d(S/(I \cap J_{Y(j)})) - H_{d-i}(S/(I:l)) \le H_d(\overline{S}) - 1.$$

Recall that $(I \cap J_{Y(j)} : l) = (I : l)$ hence

$$H_d(S/(I \cap J_{Y(j)})) - H_{d-i}(S/(I \cap J_{Y(j)}:l)) \le H_d(\overline{S}) - 1$$
 (A.5)

Using Equation (A.5) and the restriction sequence

$$0 \to S/(I \cap J_{Y(j)}:(l))(-i) \xrightarrow{\cdot l} S/(I \cap J_{Y(j)}) \twoheadrightarrow S/(I \cap J_{Y(j)}+l) \to 0$$

We get

$$1 \le H_d(S) - H_d(S/(I \cap J_{Y(j)} + l)) = H_d((I \cap J_{Y(j)} + l)/(l))$$

Therefore there exists $F \in [I \cap J_{Y(j)} + l]_d \setminus [(l)]_d$. Let $p \in V(l) \setminus V(F)$. Such p exists because $F \notin (l)$. Further note

$$F \in ((I \cap J_{Y(j)} + l)/l) \Rightarrow V(F) \supseteq V(I \cap J_{Y(j)}) \cap V(l) \supseteq Y(j).$$

Since $p \notin V(F)$ we get $Y(j+1) = Y(j) \cup \{p\}$ has j+1 distinct points. We obtain

$$H_d(S/(I \cap J_{Y(j+1)})) \ge H_d(S/(I \cap J_{Y(j)})) + 1 = H_d(S/I) + j + 1,$$

where the first inequality follows since $F \in [I \cap J_{Y(j)}]_d \setminus [I \cap J_{Y(j+1)}]_d$ and the last equality follows from the induction hypothesis. Finally, the exact sequence below yields the other inequality.

$$0 \to \frac{S}{I \cap J_{Y(j+1)}} \to S/I \oplus S/J_{Y(j+1)} \to S/I + J_{Y(j+1)} \to 0$$

 $H_d(S/(I \cap J_{Y(j+1)})) \le H_d(S/I) + H_d(S/J_{Y(j+1)}) \le H_d(S/I) + j + 1.$ (A.6)

We have shown the set Y(q) satisfies Equation (A.4). By semicontinuity of Hilbert functions a general set Y_0 of q points then satisfies $H_d(S/(I \cap J_{Y_0})) \ge H_d(S/I) + q$. As shown in Equation (A.6), the opposite inequality is always satisfied, so Equation (A.3) follows.

In the remainder of the section we prove different numerical criteria that are used in Section 6. Notation 6.5 therein is in effect.

Proposition A.2. Let $d \in \mathbb{N}$ satisfy $d \geq 6$. Then $s_d'' < 2 \cdot s_{d-3}$.

Proof. The table below shows the inequality for $6 \le d \le 8$.

	d = 6	d = 7	d = 8
s_d'''	4	4	5
$2 \cdot s_{d-3}$	6	8	10

Next we show $s_{d+3}''' < 2 \cdot s_d$ for all $d \ge 6$. Let d = 6k + i where $1 \le k$ and $0 \le i \le 5$. By Remark 6.6 we get

$$s_{d+3}^{\prime\prime\prime} = 3k + \left\lfloor \frac{i+3}{2} \right\rfloor + 1 \le 3k + 5$$
 since $i \le 5$.

Furthermore, by Remark 6.6 (1) we have $s_{6k} = 3k^2 + 3k + 1 \leq s_d$. Since $1 \leq k$ the strict inequality holds and establishes

$$s_d'' \le 3k + 5 < 2 \cdot s_{6k} = 2 \cdot (3k^2 + 3k + 1) \le 2 \cdot s_d \qquad \Box$$

Lemma A.3. Let $d \in N$ be such that $d \ge 6$. We state and prove all cases of the inequality Equation (6.7) used in the proof of Lemma 6.9.

 $\begin{array}{l} (1) \ \frac{s'_d}{2} \leq s'_{d-1} \\ (2) \ \frac{s''_d}{2} \leq s''_{d-2} \\ (3) \ \frac{s''_d}{2} \leq s'''_{d-3} \end{array}$

Proof. (1) Let d = 6p + i where $p \ge 1$. Using Remark 6.6 we have

$$s'_{d} = \begin{cases} p & i = 1\\ p+1 & \text{otherwise} \end{cases} \quad \text{and} \quad s'_{d-1} = \begin{cases} p & i = 0, 2\\ p+1 & \text{otherwise.} \end{cases}$$

Hence it follows

$$\frac{s'_d}{2} \le \frac{p+1}{2} \le p \le s'_{d-1}.$$

(2) Let d = 3m + j where $m \ge 2$. Using Remark 6.6 we have

$$s''_{d} = \left\lfloor \frac{d}{3} \right\rfloor + 1$$
 and $s''_{d-2} = \begin{cases} m & j = 0, 1 \\ m+1 & j = 2. \end{cases}$

Then

$$\frac{m''_{d}}{2} = \frac{m+1}{2} \le m \le s''_{d-2}.$$

(3) Let d = 2s + t where $s \ge 3$. Recall from Remark 6.6 that

$$s_d^{\prime\prime\prime} = \left\lfloor \frac{d}{2} \right\rfloor + 1$$
 and $s_{d-3}^{\prime\prime} = \begin{cases} s-1 & t=0\\ s & t=1. \end{cases}$

It follows that

$$\frac{s_d''}{2} = \frac{s+1}{2} \le s-1 \le s_{d-3}'''.$$

APPENDIX B. SECANT VARIETIES TO THE VERONESE VARIETY

Definition B.1. Let S be a polynomial ring. Define the *d*-th Veronese variety as $V^d = \operatorname{Proj}(k[S_d])$.

Remark B.2. The Veronese variety can be embedded in (straight) projective space as follows: Let $S = k[x_0, \ldots, x_n]$ and $N = s_d - 1$. Let m_0, \ldots, m_N enumerate the monomials in S_d . Consider the k-algebra homomorphism

$$\begin{array}{c} : k[y_0, \dots, y_N] \to k[x_0, \dots, x_n] \\ y_i \to m_i. \end{array}$$
(B.1)

Set $R = k[y_0, \ldots, y_N]$ to be the domain of θ and I to be the kernel of θ . Then Equation (B.1) gives rise to an isomorphism $R/I \cong k[S_d]$ which in turn induces an isomorphism of projective varieties $V^d \cong \operatorname{Proj}(R/I) \subset \operatorname{Proj}(R) = \mathbb{P}_k^N$.

Geometrically θ induces a map of projective varieties

θ

$$\psi: \mathbb{P} \dashrightarrow V^d, \psi(p) = [m_0(p): \cdots: m_N(p)]$$
(B.2)

defined on $U = \{ p \in \operatorname{Proj}(S) \mid \theta(R_+) \not\subseteq p \}.$

The main goal of this section is to establish the following proposition.

Theorem 7.5. Let $\mathbb{P} = \mathbb{P}(1, a_1, \ldots, a_n)$. For a fixed $d \ge a_n$ the following are equivalent:

(1) if X is a set of r general double points,
$$H_d(S/I_X) = \min\{s_d, r(n+1)\}$$

(2) $\dim_k(\sigma_r(V^d)) = \min\{s_d - 1, r(n+1) - 1\}.$

To show this we make use of tangent spaces which we now define.

Definition B.3. Let p be a point of an affine variety U with defining ideal \mathfrak{m}_p . The *Zariski tangent space* to U at p is the vector space

$$T_p(U) = \operatorname{Hom}_k(\mathfrak{m}_p/\mathfrak{m}_p^2, k).$$

If p is a point of a projective variety X, the projective tangent space to X at p, denoted $\mathbb{T}_p(X)$ is the projectivization of the $T_{\tilde{p}}\tilde{X}$, where \tilde{X} is affine cone over X and \tilde{p} is any point of the affine cone that corresponds to $p \in X$.

Lemma B.4. Consider a map $g : \mathbb{A}_k^n \to \mathbb{A}_k^N$ given by $g(p) = (g_1(p), \dots, g_N(p))$, where $g_i \in k[x_1, \dots, x_n]$. For any $p \in \mathbb{A}^n$ the induced map

$$\mathrm{d}g: T_p\mathbb{A}^n \to T_{g(p)}\mathbb{A}^N \text{ is represented by the Jacobian matrix } \begin{bmatrix} \frac{\partial g_i}{\partial x_j} \big|_p \end{bmatrix}_{\substack{1 \le i \le N \\ 1 \le j \le n}}.$$

We now analyze the projective tangent space to the Veronese variety V^d .

Proposition B.5. Let $\mathbb{P} = \mathbb{P}(1, a_1, \ldots, a_n)$, let $d \ge a_n$ be an integer and denote $N = s_d - 1$. Using the notation introduced in Equation (B.1) and Equation (B.2), consider a point $p \in U_0 = \mathbb{P} \setminus V(x_0)$ and set $U'_0 = V^d \cap (\mathbb{P}^N_k \setminus V(y_0))$. Then

- (1) The inclusion $k[S_d] \hookrightarrow S$ induces an isomorphism between U_0 and U'_0 , both of which are isomorphic to \mathbb{A}^n_k . In particular, the points p and $\psi(p)$ are smooth on \mathbb{P} and V^d , respectively.
- (2) The point $\psi(p)$ is defined and the projective tangent space $\mathbb{T}_{\psi(p)}(V^d)$ is given by

$$\mathbb{T}_{\psi(p)}(V^d) \cong \mathbb{P}\left\langle \left[\frac{\partial m_0}{\partial x_i}(p), \cdots, \frac{\partial m_N}{\partial x_i}(p)\right] \middle| 0 \le i \le n \right\rangle.$$
(B.3)

Proof. (1) Let $S = k[x_0, \ldots, x_n]$ with $\deg(x_i) = a_i$. Recall that $\mathbb{P} = \operatorname{Spec}(S)$. Moreover the coordinate ring of U_0 is $S[x_0^{-1}]_0 \cong k[x_1, \ldots, x_n]$ since $\deg(x_0) = 1$. See the description of the coordinate rings of U_i in Equation (2.3) for a justification of this isomorphism. Since the coordinate ring of U_0 is regular, all points of U_0 are smooth.

Now consider $V^d = \operatorname{Spec}(k[S_d])$. The following are monomials in S_d

$$m_0 = x_0^d, m_1 = x_0^{d-a_1} x_1, \dots, m_n = x_0^{d-a_n} x_n,$$

and every other monomial m in S_d can be written in the form

$$m = \prod_{i=0}^{n} x_{i}^{e_{i}} = \frac{x_{0}^{e_{0}} \prod_{i=1}^{n} m_{i}^{e_{i}}}{x_{0}^{\sum_{i=1}^{n} e_{i}(d-a_{i})}} = \frac{x_{0}^{\sum_{i=0}^{n} e_{i}a_{i}} \prod_{i=1}^{n} m_{i}^{e_{i}}}{x_{0}^{\sum_{i=1}^{n} e_{i}d}} = \frac{m_{0} \prod_{i=1}^{n} m_{i}^{e_{i}}}{m_{0}^{\sum_{i=1}^{n} e_{i}}}.$$

It follows that $k[S_d, x_0^{-d}]_0 \cong k[\frac{m_1}{m_0}, \ldots, \frac{m_n}{m_0}]$. Since $\frac{m_1}{m_0}, \ldots, \frac{m_n}{m_0}$ each contain a distinct variable x_i , they are algebraically independent. Thus the coordinate ring of U'_0 is regular and we deduce that V^d is smooth at q whenever $q \in U'_0$. In particular, the points $\psi(p)$ with $p \in U_0$ are smooth points of V^d .

The inclusion $\iota : k[S_d] \hookrightarrow S$ yields upon inverting powers of x_0 the identity map

$$k\left[\frac{m_1}{m_0},\ldots,\frac{m_n}{m_0}\right] = k[S_d,x_0^{-d}] \to S[x_0^{-1}] = k\left[\frac{x_1}{x_0},\ldots,\frac{x_n}{x_0}\right]$$

since $m_i/m_0 \mapsto x_i/x_0 = m_i/m_0$ for $1 \le i \le n$.


(2) The point $\psi(p)$ is defined since $m_0(p) \neq 0$.

Consider the affine cone $\widetilde{V^d}$ of V^d and the regular map

 $g: \mathbb{A}_k^{n+1} \to \mathbb{A}_k^{N+1}, g(p) = (m_0(p), \dots, m_N(p)).$

The map induced by g on coordinate rings is denoted θ in Equation (B.1).

Let \mathfrak{m} denote the ideal of p in \mathbb{A}_{k}^{n+1} , \mathfrak{m}' denote the ideal of g(p) in \mathbb{A}_{k}^{N+1} , and \mathfrak{m}'' denote the ideal of g(p) in $A' = k[\widetilde{V^{d}}]$. Let $\pi : k[\mathbb{A}_{k}^{N+1}] \to A'$ denote the canonical projection and $\iota : k[S_{d}] \to S = k[A^{n+1}]$ denote the inclusion. These maps satisfy the identity $\theta = \iota \circ \pi$. There are dual commutative diagrams as follows



By part (1), the horizontal map labeled ι and hence also its dual $d\iota$ are isomorphisms. Since π is surjective, $d\pi$ is injective. Hence $T_{g(p)}\widetilde{V^d}$ is isomorphic to the image of $d\pi$ which coincides with the image of dg. By Lemma B.4, the differential $dg: T_p\mathbb{A}_k^{n+1} \to T_{g(p)}\mathbb{A}_k^{N+1}$ is represented by the Jacobian matrix $J = \begin{bmatrix} \frac{\partial m_i}{\partial x_j} \end{bmatrix}_{\substack{0 \le i \le N \\ 0 \le j \le n}}$. Therefore, $T_{\psi(p)}(\widetilde{V^d})$ can be identified with the column space of J and its projectivization yields

$$\mathbb{T}_{\psi(p)}(V^d) \cong \mathbb{P}\left\langle \left[\frac{\partial m_0}{\partial x_j}\Big|_p, \dots, \frac{\partial m_N}{\partial x_j}\Big|_p\right] \mid 0 \le j \le n \right\rangle.$$

In the following, we to denote by $\widetilde{V^d}$ the affine cone over V^d and by \widetilde{q} any affine point that corresponds to the projective point $q \in \mathbb{P}_k^N$. Proposition B.6 in standard graded case was first proven by Lasker in [Las04]. Our proof is an adaption from [BO08].

Proposition B.6 (Lasker's proposition). Let $\mathbb{P} = \mathbb{P}(1, a_1, \ldots, a_n)$, let $d \geq a_n$ be an integer and set $N = s_d - 1$. Let $p \in U_0 = \mathbb{P} \setminus V(x_0)$. Then the orthogonal complement to the tangent space to $\widetilde{V^d}$ at $\widetilde{\psi(p)}$ consists of all hypersurfaces of degree d singular at p. In detail, there is a vector space isomorphism

$$[I_p^{(2)}]_d \cong T_{\widetilde{\psi(p)}}(\widetilde{V^d})^{\perp}.$$

Proof. By the Zariski-Nagata Theorem, $[I_p^{(2)}]_d$ is the vector space of homogeneous polynomials of degree d whose first derivative vanishes on p

$$(I_p^{(2)})_d = \left\{ f \in S_d \mid \frac{\partial f}{\partial x_i}(p) = 0 \right\} = \left\{ \sum_{i=0}^N \lambda_i m_i \mid \sum_{i=0}^N \lambda_i \frac{\partial m_i}{\partial x_i} \big|_p = 0 \right\}.$$

Then for any $\underline{\lambda} \in k^{N+1}$ we have

$$\begin{split} &\sum_{i=0}^{N} \lambda_{i} \frac{\partial m_{i}}{\partial x_{i}} \big|_{p} = 0 \\ \Leftrightarrow (\lambda_{0}, \dots, \lambda_{N}) \cdot \left(\frac{\partial m_{1}}{\partial x_{i}} \big|_{p}, \cdots, \frac{\partial m_{N}}{\partial x_{i}} \big|_{p} \right) = 0 \\ \Leftrightarrow &\underline{\lambda} \in T_{\widetilde{\psi_{0}(p)}}(\widetilde{V^{d}})^{\perp}. \end{split}$$

We now define the k-th secant variety of a projective variety. This is a central tool in establishing Theorem 7.5. For a more thorough study of the secant variety see [Har92, Lecture 8].

Definition B.7 (Secant Variety). Let X be a projective variety. For any nonnegative integer r, the r-secant variety of X, denoted by $\sigma_r(X)$, is defined to be

$$\sigma_r(X) = \overline{\bigcup_{P_1, \dots, P_r \in X} \langle P_1, \dots, P_r \rangle}^{\text{Zariski closure}}$$

Note that $\sigma_r(V_d^n)$ is an irreducible variety for all r. ([Fan90, Remark 1.2])

Lemma B.8 (First Terracini Lemma). Let Y be an a affine variety set, let $p_1, \ldots, p_r \in Y$ be general points and $z \in \langle p_1, \ldots, p_r \rangle$ be a general point. Then

$$T_z(\sigma_r(Y)) = \langle T_{p_1}(Y), \dots, T_{p_r}(Y) \rangle.$$

Proof. See [BO08, Lemma 2.2].

Using arguments identical to [BO08] we prove Theorem 7.5.

Proof of Theorem 7.5. Let $X = \{2p_1, \ldots, 2p_r\}$ be a collection of general double points in \mathbb{P} . Applying Proposition B.6 to get the second equality, we obtain

$$[I_X]_d = \left[\bigcap_{i=1}^r I_{p_i}^{(2)}\right]_d = \bigcap_{i=1}^r \left(T_{\widetilde{\psi(p_i)}}\widetilde{V^d}\right)^{\perp} = \left\langle T_{\widetilde{\psi(p_1)}}\widetilde{V^d}, \dots, T_{\widetilde{\psi(p_r)}}\widetilde{V^d}\right\rangle^{\perp}.$$

Therefore, using Lemma B.8, we obtain

$$\dim([S/I_X]_d) = \dim_k \left\langle T_{\widetilde{\psi(p_1)}} \widetilde{V^d}, \dots, T_{\widetilde{\psi(p_r)}} \widetilde{V^d} \right\rangle = \dim_k T_z(\sigma_k(\widetilde{V^d})).$$

Finally using that $\sigma_r(\widetilde{V^d}) = \widetilde{\sigma_r(V^d)}$ and the irreducibility of $\widetilde{V^d}$, which implies irreducibility of $\sigma_r(\widetilde{V^d})$ for all r [Har92, p. 144, Prop 11.24], we get for z a general point of $\sigma_r(\widetilde{V^d})$

$$\dim T_z(\sigma_k(\widetilde{V^d})) = \dim \sigma_k(\widetilde{V^d}) = \dim(\sigma_k(V^d)) + 1.$$

To summarize, we have obtained the identity

$$\dim([S/I_X]_d) = \dim(\sigma_k(V^d)) + 1,$$

whence the claimed equivalence follows.

References

- [AH92a] J. Alexander and A. Hirschowitz, La méthode d'Horace éclatée: application à l'interpolation en degré quatre, Invent. Math. 107 (1992), no. 3, 585–602. MR 1150603
 [AH92b] ______, Un lemme d'Horace différentiel: application aux singularités hyper-
- [AH92b] _____, Un lemme d'Horace différentiel: application aux singularités hyperquartiques de P⁵, J. Algebraic Geom. 1 (1992), no. 3, 411–426. MR 1158623
 [AH95] ______, Polynomial interpolation in several variables, J. Algebraic Geom. 4 (1995), no. 2, 201–222. MR 1311347
- [Ale88] J. Alexander, Singularités imposables en position générale à une hypersurface projective, Compositio Math. 68 (1988), no. 3, 305–354. MR 971330
- [Amr94] Abdallah Al Amrani, Complex K-theory of weighted projective spaces, Journal of Pure and Applied Algebra **93** (1994), no. 2, 113–127.
- [BFNR13] Anthony Bahri, Matthias Franz, Dietrich Notbohm, and Nigel Ray, The classification of weighted projective spaces, Fundamenta Mathematicae 220 (2013), no. 3, 217–226.
- [BO08] Maria Chiara Brambilla and Giorgio Ottaviani, On the Alexander-Hirschowitz theorem, J. Pure Appl. Algebra 212 (2008), no. 5, 1229–1251. MR 2387598
- [BR86] Mauro Beltrametti and Lorenzo Robbiano, Introduction to the theory of weighted projective spaces, Expositiones Mathematicae 4 (1986).
- [BR07] Matthias Beck and Sinai Robins, *Computing the continuous discretely*, Undergraduate Texts in Mathematics, Springer, New York, 2007, Integer-point enumeration in polyhedra. MR 2271992
- [Bre79] H. Bresinsky, Monomial space curves in A³ as set-theoretic complete intersections, Proc. Amer. Math. Soc. 75 (1979), no. 1, 23–24. MR 529205
- [Cer17] Carlos Enrique Améndola Cerón, Algebraic statistics of gaussian mixtures, Ph.D. thesis, Technische Universität Berlin, 2017.
- [Cha01] Karen A. Chandler, A brief proof of a maximal rank theorem for generic double points in projective space, Trans. Amer. Math. Soc. 353 (2001), no. 5, 1907–1920. MR 1813598
- [Cha02] _____, Linear systems of cubics singular at general points of projective space, Compositio Math. **134** (2002), no. 3, 269–282. MR 1943904
- [CHHVT20] E. Carlini, H.T. Hà, B. Harbourne, and A. Van Tuyl, Ideals of powers and powers of ideals: Intersecting algebra, geometry, and combinatorics, Lecture Notes of the Unione Matematica Italiana, Springer International Publishing, 2020.

- [Den03] Graham Denham, Short generating functions for some semigroup algebras, Electron. J. Combin. 10 (2003), Research Paper 36, 7. MR 2014523 [Dol82] Igor Dolgachev, Weighted projective varieties, Group Actions and Vector Fields (Berlin, Heidelberg) (James B. Carrell, ed.), Springer Berlin Heidelberg, 1982, pp. 34-71. $[DREE^+23]$ Sandra Di Rocco, Parker B Edwards, David Eklund, Oliver Gäfvert, and Jonathan D Hauenstein, Computing geometric feature sizes for algebraic manifolds, SIAM Journal on Applied Algebra and Geometry 7 (2023), no. 4, 716 - 741.[Fan90] Barbara Fantechi, On the superadditivity of secant defects, Bulletin de la Société Mathématique de France **118** (1990), no. 1, 85–100 (en). MR 92c:14049 [GKW91] W. Gastinger, E. Kunz, and R. Waldi, Relation matrices of monomial curves, An. Univ. București Mat. 40 (1991), no. 1-2, 41-53. MR 1220263 A. Grothendieck, Éléments de géométrie algébrique. I. Le langage des [Gro60] schémas., Inst. Hautes Études Sci. Publ. Math. (1960), no. 4, 228. MR 217083 [GS]Daniel R. Grayson and Michael E. Stillman, Macaulay2, a software system for research in algebraic geometry, Available at http://www.math.uiuc.edu/Macaulay2/. [Har92] Joe Harris, Algebraic geometry, Graduate Texts in Mathematics, vol. 133, Springer-Verlag, New York, 1992, A first course. MR 1182558 [Her70] Jürgen Herzog, Generators and relations of abelian semigroups and semigroup rings, Manuscripta Math. 3 (1970), 175–193. MR 269762 [Hir85] André Hirschowitz, La méthode d'Horace pour l'interpolation à plusieurs variables, Manuscripta Math. 50 (1985), 337-388. MR 784148 [HM21] Huy Tài Hà and Paolo Mantero, The Alexander-Hirschowitz theorem and related problems, Commutative algebra, Springer, Cham, [2021] (c)2021, pp. 373-427. MR 4394415 [Hui18] Jack Huizenga, Polynomial Interpolation: An introduction to algebraic geometry, 2018.[KTB19] Joe Kileel, Matthew Trager, and Joan Bruna, On the expressive power of deep polynomial neural networks, Advances in neural information processing systems **32** (2019).
- [Las04] Emanuel Lasker, Zur Theorie der kanonischen Formen, Math. Ann. 58 (1904), no. 3, 434–440. MR 1511244
- [Pal03] F. Palatini, Sulla rappresentazione delle forme ternarie mediante la somma di potenze di forme lineari, Rend. Accad. Lincei V. (1903), no. 12, 378–384 (Italian).
- [Pos12] Elisa Postinghel, A new proof of the Alexander-Hirschowitz interpolation theorem, Annali di Matematica Pura ed Applicata 191 (2012), no. 1, 77–94.
- [RT11] Michele Rossi and Lea Terracini, Weighted projective spaces from the toric point of view with computational applications, arXiv:1112.1677, 2011.
- [Sen20] Indranath Sengupta, Affine monomial curves, 2020.
- [Sta99] Richard P. Stanley, *Enumerative combinatorics*, vol. 2, Cambridge University Press, 1999.
- [Ter11] Alessandro Terracini, Sulle v_k per cui la varietà degli S_h h + 1-secanti ha dimensione minore dell'ordinario, Rend. Circ. Mat. Palermo Selecta Vol. I (1911), no. 31 (Italian).
- [Ter15a] A. Terracini, Sulla rappresentazione delle forme quaternarie mediante somme di potenze di forme lineari, Atti R. Accad. delle Scienze di Torino. Selecta Vol. Selecta I (1915), no. 51 (Italian).

[Ter15b] Alessandro Terracini, Sulla rappresentazione delle coppie di forme ternarie mediante somme di potenze di forme lineari, Annali Mat. Selecta Vol. I (1915), no. 24 (Italian).

Department of Mathematics, University of Nebraska-Lincoln, 1144 T St Lincoln, NE 68588

Email address: sroshanzamir2@huskers.unl.edu