ON FOX'S TRAPEZOIDAL CONJECTURE

SOHEIL AZARPENDAR, ANDRÁS JUHÁSZ, AND TAMÁS KÁLMÁN

ABSTRACT. We investigate Fox's trapezoidal conjecture for alternating links. We show that it holds for diagrammatic Murasugi sums of special alternating links, where all sums involved have length less than three (which includes diagrammatic plumbing). It also holds for links containing a large twist region, which we call twist-concentrated. Furthermore, we show some weaker inequalities between consecutive coefficients of the Alexander polynomial of an alternating 3-braid closure, and extend this to arbitrary alternating links. We then study an extension of the trapezoidal conjecture due to Hirasawa and Murasugi, which states that the stable length of the Alexander polynomial of an alternating link can be bounded from above using the signature. We estabilish this and determine when equality holds for diagrammatic Murasugi sums of special alternating knots where each sum has length less than three, and also for twist-concentrated 3-braids. Finally, we study the behavior of the Hirasawa–Murasugi inequality under concordance.

Contents

1. Overview	2
2. Fox's trapezoidal conjecture	4
2.1. Introduction	4
2.2. Stabilization and highly-twisted links	10
2.3. Plumbings of special alternating knots	17
2.4. A multivariable refinement of the Alexander polynomial for	
alternating 3-braids	34
3. A conjecture of Hirasawa and Murasugi	52
3.1. Introduction	52
3.2. H-M sharp links	58
3.3. Dual inequality and concordance	60
References	62

Date: June 14, 2024.

Key words and phrases. Alexander polynomial, knot, link, log-concave, trapezoidal conjecture, unimodal.

TK was supported by a Japan Society for the Promotion of Science (JSPS) Grant-in-Aid for Scientific Research C (no. 23K03108).

1. Overview

In Section 2, we focus on the following conjecture of Fox [Fox61] about the Alexander polynomial of alternating knots. This is known as Fox's trapezoidal conjecture.

Conjecture 1.1. [Fox61] Let (a_1, \ldots, a_n) be the sequence of absolute values of the coefficients of the Alexander polynomial Δ_K of an alternating link K. With $m = \lfloor \frac{n}{2} \rfloor$, the following properties hold:

- (1) $a_1 \le a_2 \le \dots \le a_m \ge \dots \ge a_{n-1} \ge a_n;$ (2) if $a_i = a_{i+1}$ for i < m, then $a_i = a_{i+1} = \dots = a_m.$

A sequence with such properties is called *trapezoidal*; see Figure 1.

In Subsection 2.1, we go through the main known results about the trapezoidal conjecture in chronological order and briefly mention the methods used to derive each result. The following table summarizes Subsection 2.1.

Results about trapezoidal conjecture		
Reference	Class of alternating links	Method
Parris[Par78]	Alternating pretzel knots	Formula for Δ_K
Hartley[Har79]	Two-bridge links	Relating a formula for Δ_K to extended Schubert diagrams
Murasugi[Mur85]	Alternating arborescent links	Skein formula for plumbing trees and induction
Ozsváth, Szabó[OS03]	$\begin{array}{ll} \text{Alternating} & \text{knots} & \text{with} \\ g(K) \leq 2 \end{array}$	Computation of $HF^+(S_0^3(K))$ using large surgery formula
Jong [Jon10], Stoimenow [Sto16]	Alternating knots with $g(K) \le 4$	Reduction to a finite list of genus generators using $t_2^{\bar{t}}$ move
Alrefai, Chbili[AC21]	Alternating 3-braids of length less than 4	Formula for Δ_K using Burau representation
Hafner, Mészáros, and Vidinas[HMV23]	Special alternating links	Introduction of multivariable refinement of Δ_K

In Subsection 2.2, we use Crowell's formula for the Alexander polynomial and arguments similar to Jong[Jon10] to derive the trapezoidal conjecture for alternating links that contain a (linearly) large twist region. To be more precise, let MT(L) be the size of the biggest coherent twist region in an alternating link; see Definition 2.5 and the paragraph preceding it. We call an alternating link *twist-concentrated* if it satisfies the inequality

$$MT(L) - 3 \ge g(L) + \frac{|L|}{2},$$

 $\mathbf{2}$

where |L| is the number of connected components of L. Such links satisfy Fox's trapezoidal conjecture.

More generally, we show that the first n inequalities of the trapezoidal conjecture follow from the existence of a twist region with n + 3 crossings. The main results of this subsection are Theorems 2.6 and 2.11.

In Subsection 2.3, we use the decomposition of alternating links into diagrammatic Murasugi sums of special alternating links. Murasugi [Mur65] proposed an algorithm for such a decomposition. We use this to extend the result of Hafner, Mészáros, and Vidinas [HMV23] to a certain class of Murasugi sums. In general, it is not possible to derive a formula for the Alexander polynomial of the Murasugi sum of two knots using the Alexander polynomials of the summands. If we restrict the sums to diagrammatic plumbings, then such a formula can be derived as it is done in Proposition 2.21. More generally, this can be done for diagrammatic Murasugi sums of length less than 3; see Subsection 2.3 Using this formula, we prove the trapezoidal conjecture for diagrammatic Murasugi sums of special alternating links such that the lengths of all Murasugi sums are less than 3. The main results of this subsection are Theorems 2.20 and 2.23.

In Subsection 2.4, we focus on alternating 3-braids which form the simplest class of non-special alternating links, and we try to extend the main idea of Hafner, Mészáros, and Vidinas [HMV23]. Their main idea is to construct a multivariable refinement of the Alexander polynomial such that all of its coefficients are equal to 1 and its support is M-convex. This means that this multivariable polynomial will be Lorentzian, and this opens the door to techniques developed by Brändén and Huh [BH20]; see Subsection 2.4 for a detailed explanation. We consider a natural generalization of the multivariable refinement to alternating 3-braids. We show that, unfortunately, this multivariable polynomial cannot be Lorentzian except when the mentioned alternating 3-braid is a connected sum of two torus links. Despite this, one can use combinatorial arguments to derive certain inequalities between its coefficients, which in turn result in inequalities between the coefficients of the Alexander polynomial. These inequalities are generally either weaker than the statement of the trapezoidal conjecture (by a multiplicative factor) or they are independent from it. Finally, since this argument only uses the local combinatorial structure of the alternating 3-braids, one can generalize this to all alternating links. These results are stated in Theorems 2.29 and 2.31.

In Section 3, we discuss a stronger version of the trapezoidal conjecture proposed by Hirasawa and Murasugi [HM14]. The main addition is a conjectured inequality between the stable length of the sequence of coefficients of the Alexander polynomial and the signature of the link, explained in the following. **Conjecture 1.2.** Let *L* be an alternating link and $\Delta_L(t) = \sum_{i=0}^{l-1} (-1)^i a_i t^i$ be its Alexander polynomial. Then there is $i_0 \in \{1, \ldots, l/2 + 1\}$ such that

$$a_0 < \dots < a_{i_0-1} = a_{i_0} = \dots = a_{l-i_0} > \dots > a_{l-1}.$$

Moreover,

$$\left\lfloor \frac{|\sigma(L)|+1}{2} \right\rfloor \ge \left\lfloor \frac{l-2(i_0-1)}{2} \right\rfloor.$$

We call this inequality the *H-M inequality*. We call those alternating links for which the H-M inequality becomes an equality *H-M sharp*. We define the stable length of L to be

$$sl(L) := l - 2(i_0 - 1).$$

In Subsection 3.1, we use the results developed in Section 2 to derive corollaries about the H-M inequality. Using the results of Subsection 2.2, one can show that the H-M inequality for twist-concentrated alternating links L reduces to the problem for the rest of the alternating links. In the case of certain twist-concentrated links for which we can compute the signature, such as twist-concentrated 3-braids, this argument proves the H-M inequality. These results are stated in Corollary 3.6 and Theorem 3.9.

We then use the results of Subsection 2.3, and prove the H-M inequality for the diagrammatic Murasugi sum of special alternating links when the lengths of all Murasugi sums are less than 3. This is the content of Corollary 3.12.

In Subsection 3.2, we investigate the properties of H-M sharp links. Refining the arguments of the previous subsection, one can derive a diagrammatic characterization of H-M sharp knots, which are diagrammatic Murasugi sums of special alternating links such that the lengths of all Murasugi sums are less than 3. This result is stated in Corollary 3.16.

In Subsection 3.3, we use the Fox-Milnor equation to form a relation between the H-M inequality and concordance. Consider the minimum degree of the Alexander polynomial in the algebraic concordance class of an alternating knot K, denoted by $d_c(K)$. It turns out that $d_c(K)+1$ gives an upper bound for the stable length sl(K). The H-M inequality can be viewed as an optimization problem; i.e., finding the maximal stable length in a given algebraic concordance class. From this perspective, computing $d_c(K)$ acts as a dual problem for the H-M inequality. This leads to proof of the H-M inequality for certain concordance classes, as stated in Corollaries 3.21 and 3.24. This perspective relaxes the alternating condition and thus might be useful.

2. Fox's trapezoidal conjecture

2.1. Introduction. A link K in S^3 is called *alternating* if it has a projection D in which the undercrossings and overcrossings alternate as one moves along any component. Let $\Delta_K(t) \in \mathbb{Z}[t]$ be the (single-variable) Alexander polynomial of K. We also use the notation $\tilde{\Delta}_K \in \mathbb{Z}[t, t^{-1}]$ to denote the

symmetrized Alexander polynomial of the link K. Although this is not standard notation, we need it for clarity.

In 1962, Fox conjectured that the sequence of absolute values of the coefficients of Δ_K is trapezoidal for alternating K.

Conjecture 1.1. [Fox61] Let (a_1, \ldots, a_n) be the sequence of absolute values of the coefficients of the Alexander polynomial Δ_K of an alternating link K. With $m = \lfloor \frac{n}{2} \rfloor$, the following properties hold:

- (1) $a_1 \le a_2 \le \dots \le a_m \ge \dots \ge a_{n-1} \ge a_n;$ (2) if $a_i = a_{i+1}$ for i < m, then $a_i = a_{i+1} = \dots = a_m.$

See Figure 1 for an example of a trapezoidal sequence.



FIGURE 1. Graph of a trapezoidal sequence.

Note that the sequence of coefficients of Δ_K is symmetric; i.e., $a_i = a_{n-i+1}$ for $i \in \{1, \ldots, n\}$. Murasugi [Mur85] proved that, for an alternating K, signs of this sequence alternate, which means, using the notation of Conjecture 1.1, that

$$\Delta_K(t) = \pm \sum_{1 \le i \le n} (-1)^{i-1} a_i t^{i-1}.$$

Stoimenow proposed the following strengthening of Conjecture 1.1.

Conjecture 2.1. [Sto05] Let (a_1, \ldots, a_n) be the sequence of absolute values of the coefficients of the Alexander polynomial Δ_K of an alternating link K in S^3 . Then (a_1, \ldots, a_n) is log-concave; i.e., for every $i \in \{2, \ldots, n-1\}$, we have

$$a_{i-1}a_{i+1} \le a_i^2.$$

Parris [Par78] in his PhD thesis used Fox calculus to derive a formula for the Alexander polynomial of pretzel knots which in turn proves the trapezoidal conjecture for this class.

Hartley [Har79] proved the trapezoidal conjecture for two-bridge knots D(p,q). The Alexander polynomial $\Delta_{D(p,q)}$ can be computed by the following formula. Let p, q be coprime integers satisfying 0 < q < p and q odd. Then

(1)
$$\Delta_{D(p,q)}(t) = \sum_{0 \le k \le p-1} (-1)^k t^{s_k} , \ s_k = \sum_{i=0}^k (-1)^{\left\lfloor \frac{iq}{p} \right\rfloor}.$$

Minkus [Min82] proved this formula using Fox calculus on a Schubert normal diagram of D(p,q). Using a similar argument, Hartley provided another formula for $\Delta_{D(p,q)}$ which can be read from an extended diagram obtained from unwinding Schubert's normal diagram; see Figure 2. Note that, in the extended diagram, all of the diagonal segments have slope $\frac{q}{p}$.



FIGURE 2. Hartley's extended diagram [Har79].

Proposition 2.2. [Har79] Let α_i be the number of segments joining the *i*-th and (i + 1)-st vertical lines in the extended diagram of D(p,q). Then

$$\Delta_{D(p,q)} = \sum (-1)^i \alpha_i t^i.$$

Hartley uses other combinatorial properties of the extended diagrams to devise an inductive argument. Chen [Che17] uses the same argument to derive stronger results; see Section 3 for more details.

Murasugi [Mur85] proved the trapezoidal conjecture for arborescent alternating links. These are alternating links that can be constructed from the plumbing of oriented twisted bands over a tree (for details, see Subsection 2.3). Murasugi uses skein relations on plumbing trees to derive an inductive argument. Figure 3 shows an example of such a skein triple.

As depicted in Figure 3, changing the orientation of a crossing reduces the number of twists in a band (weight of a vertex) by 2 and smoothing a crossing, results in a connected sum of the adjacent bands.

Furthermore, Murasugi derived a sufficient criterion for an algebraic link to be alternating. This relies on a trick which relates the plumbing tree Tto the Tait graph of a diagram of the associated algebraic link l(T). More precisely, Murasugi showed that if one augments T by an exterior vertex \hat{v} , and connects \hat{v} to V(T) such that in the augmented graph, the weight of each vertex is equal to the signed count of its adjacent edges, then l(T) is isotopic to the result of applying the median construction to the planar dual of $T \cup \{\hat{v}\}$; see Figure 4.

Ozsváth and Szabó [SO03] used Heegaard Floer homology to prove the trapezoidal conjecture for alternating knots of genus two. They proved that alternating knots are HF-thin; i.e.,

$$\widehat{\mathrm{HFK}}(S^3, K, s) \cong \mathbb{Z}^{|a_s|}$$
 supported in Maslov grading $s + \frac{\sigma}{2}$,



FIGURE 3. Skein relation on plumbing trees.



FIGURE 4. Alternative construction of an algebraic link (both black and green diagrams represent the trefoil).

where $\tilde{\Delta}_K(T) = a_0 + \sum_{s>0} a_s(T^s + T^{-s})$ and σ is the signature of K. The proof comes from construction of a Heegaard diagram for a knot where the Heegaard states are in one-to-one correspondence with Kauffmann states. For alternating knots, this Heegaard diagram has the property that $M(\underline{\mathbf{x}}) - A(\underline{\mathbf{x}}) = \frac{\sigma(K)}{2}$ for any intersection point $\underline{\mathbf{x}}$, where $A(\underline{\mathbf{x}})$ is the Alexander grading and $M(\underline{\mathbf{x}})$ is the Maslov grading of $\underline{\mathbf{x}}$. As a result, all the differentials in the hat version of the knot Floer chain complex vanish. Ozsváth and Szabó go one step further and prove the following.

Proposition 2.3. [SO03] Let K be an alternating knot, oriented so that $\sigma = \sigma(K) \leq 0$. For all s > 0, let

$$\delta(\sigma,s) := max\left(0, \left\lceil \frac{|\sigma| - 2|s|}{4} \right\rceil\right)$$

and $t_s(K) := \sum_{j=1}^{\infty} ja_{|s|+j}$. Then we have a $\mathbb{Z}[U]$ -module isomorphism

$$HF^+(S_0^3(K),s) \cong \mathbb{Z}^{b_s} \oplus \left(\mathbb{Z}[U]/U^{\delta(\sigma,s)}\right),$$

where $(-1)^{s+\frac{\sigma}{2}}b_s = \delta(\sigma, s) - t_s(K)$. Furthermore, for s = 0, we have

$$HF^{+}(S_{0}^{0}(K),0) \cong \mathbb{Z}^{0} \oplus \mathcal{T}_{-1/2}^{+} \oplus \mathcal{T}_{-2\delta(\sigma,0)+\frac{1}{2}}^{+},$$

where $HF^+(S^3) = \mathcal{T}_0^+$ and the subscript denotes a shift in the Q-grading.

Their proof uses the large surgery formula to compute $HF^+(S_n^3(K), s)$ for sufficiently large n, and then applies the surgery exact sequence:

$$\cdot \to \operatorname{HF}^+(S^3) \to \operatorname{HF}^+(S^3_0(K), s) \to \operatorname{HF}^+(S^3_n(K), [s]) \to \cdot$$

Note that a corollary of this computation is that $b_s \ge 0$. Using the inequality $b_0 \ge 0$ for alternating knots of genus two implies the trapezoidal conjecture.

Remark 2.4. One needs to combine the large surgery formula with some homological algebra to calculate $\mathrm{HF}^+(S^3_n(K),s)$. The following (Mayer– Vietoris) exact sequence of subcomplexes of $\mathrm{CFK}^{\infty}(S^3, K)$ is useful:

$$0 \to C\{i \ge 0 \text{ or } j \ge s\} \to C\{i \ge 0\} \oplus C\{j \ge s\} \to C\{i \ge 0 \text{ and } j \ge s\} \to 0$$



FIGURE 5. Exact sequence of subcomplexes of CFK^{∞} .

For an alternating knot, one can use $M(\underline{\mathbf{x}}) - A(\underline{\mathbf{x}}) = \frac{\sigma(K)}{2}$ to conclude that $M([\underline{\mathbf{x}}, i, j]) = M(\underline{\mathbf{x}}) + 2i = A(\underline{\mathbf{x}}) + \frac{\sigma}{2} + 2i = (j - i) + \frac{\sigma}{2} + 2i = i + j + \frac{\sigma}{2}$.

In particular, this means that $H_*(C\{i \ge 0 \text{ and } j \ge s\})$ is supported in degrees higher than $s + \frac{\sigma}{2}$. Combining this fact with the long exact sequence of homologies associated to the above short exact sequence results in:

$$\begin{aligned} H_{\leq s+\frac{\sigma}{2}-2}(C\{i \geq 0 \text{ or } j \geq s\}) &\cong H_{\leq s+\frac{\sigma}{2}-2}(C\{i \geq 0\}) \oplus H_{\leq s+\frac{\sigma}{2}-2}(C\{j \geq s\}) \\ &\cong \mathrm{HF}^{+}_{\leq s+\frac{\sigma}{2}-2}(S^{3}) \oplus \mathrm{HF}^{+}_{\leq -s+\frac{\sigma}{2}-2}(S^{3}). \end{aligned}$$

Recall that, based on the large surgery formula, we have

$$H_*(C\{i \ge 0 \text{ or } j \ge s\}) \cong \mathrm{HF}^+(S^3_n(K), [s])$$

for n sufficiently large.

. .

Ni [Ni21] used these results to prove an extremal case of inequalities in the trapezoidal conjecture. Let K be an alternating knot and $\Delta_K(T) = \sum_{i=-g}^{g} a_i T^i$. Then Ni showed that, if $|a_g| = |a_{g-1}|$, then K or its mirror is the torus knot $T_{2g+1,2}$. This follows from the fact that knot Floer homology detects of fibredness.

Jong[Jon10] gave a combinatorial proof of the trapezoidal conjecture for alternating knots of genus at most two. This proof relies on Stoimenow's characterization of knot diagrams of genus two [Sto08]. Analysis of Seifert graphs proves that all knot diagrams with Seifert genus two come from applying flypes and $\overline{t'_2}$ moves to a finite list of diagrams called generators.



FIGURE 6. The move $\overline{t'_2}$; see Jong [Jon10].

Note that, by a theorem of Gabai [Gab86], the Seifert genus of a reduced alternating diagram is equal to the genus of the knot. As a result, one can conclude that all alternating knots of genus two come from applying flypes and $\overline{t'_2}$ moves to alternating diagrams of Stoimenow's list of generators.

Jong combined this characterization with Crowell's spanning tree formula for the Alexander polynomial of alternating links. We explain this formula in detail in Subsection 2.2. Jong calculated the effect of $\overline{t'_2}$ on Crowell's weighted graph and proved the trapezoidal conjecture by an inductive argument.

Note that Stoimenow's theorem is true for any fixed genus, so one can prove the trapezoidal conjecture by checking it for a finite list of generators. Performing such a task, even with the help of a computer, is not trivial. **Stoimenow** [Sto16] used several computational tricks to prove the trapezoidal conjecture for all alternating knots with genus less than or equal to 4.

Alrefai and Chbili [AC21] used Burau representation to derive closedform formulas for the Alexander polynomial of alternating 3-braids with length less than or equal to 3; i.e., for braid words of the form

$$\sigma_1^{p_1}\sigma_2^{-q_1}\sigma_1^{p_2}\sigma_2^{-q_2}\sigma_1^{p_3}\sigma_2^{-q_3}$$

Similar ideas were used by Alsukaiti and Chbili [AC23] to prove the conjecture for 3-braids of the form $(\sigma_1 \sigma_2^{-1})^n$.

Banfield [Ban22] used Minkus's formula for the Alexander polynomial of two-bridge link (equation 1) to prove that the sequence of coefficients of this polynomial is not only trapezoidal but log-concave. They noticed that Minkus's formula identifies this sequence with the vertical slices of folded Christoffel paths. The Christoffel path with slope $\frac{q}{p}$ is the path in a $p \times q$ grid which is closest to the diagonal and remains under it. The closeness here means that there are no grid points in the region bounded by the path and the diagonal. This path is the best "integer approximation" of the line with slope $\frac{q}{p}$; see Figure 7. Banfield then used facts about the characterization of Christoffel words; i.e., words in $\{x, y\}^*$ generated by following steps of a Christoffel path. Using these, they constructed an inductive argument that completed the proof.

Hafner, Mészáros, and Vidinas [HMV23] proved the trapezoidal conjecture for special alternating knots. Special alternating knots are knots that can be constructed using the median construction on a bipartite plane graph G. Equivalently, these are alternating knots in which all crossings are positive. We talk about their method in detail in Subsection 2.4.



FIGURE 7. Christoffel paths (ordinary and folded); see Ban-field [Ban22].

2.2. Stabilization and highly-twisted links. The aim of this subsection is to prove the trapezoidal conjecture for alternating links with (linearly) large twist regions; see Theorem 2.6. A neighborhood in a link diagram is called an *coherent twist region* if it is locally isomorphic to Figure 8. These twist regions can be constructed by twisting two parallel strands of a link.



FIGURE 8. A coherent twist region.

Definition 2.5. We define the maximal twist number MT(L) of an alternating link L to be the maximum of the number of crossings in an coherent twist region in any reduced alternating diagram of L.

Theorem 2.6. Let L be an alternating link and

$$\Delta_L(t) \doteq \sum_{i=0}^{2g(L)+|L|-1} (-1)^i a_i t^i,$$

where $a_i \ge 0$ for $i \in \{0, \dots, 2g(L) + |L| - 1\}$. If

(1) MT(L) is odd and $MT(L) - 2 \ge g(L) + |L|/2$, or

(2) MT(L) is even and $MT(L) - 3 \ge g(L) + |L|/2$,

then the sequence (a_i) is trapezoidal.

The proof of Theorem 2.6 follows from a stabilization phenomenon for the Alexander polynomial's coefficients after applying a large number of twists. This was shown by Lambert-Cole [LC18] using skein relations. Furthermore, Lambert-Cole generalized this to knot Floer homology. In the rest of this section, we reprove this fact for alternating links using Crowell's formula. This proof gives us some additional information about the stabilized values of the coefficients, which will in turn be used to prove Theorem 2.6. First, we need to describe Crowell's formula for the Alexander polynomial of alternating links.

Let D be an alternating diagram of a link L with m crossings $c_1, ..., c_m$. We can assume that $m \geq 2$. Consider the underlying planar graph of the diagram D, which we denote by G_D and call the *Crowell graph* associated to D. The vertices of G_D are crossings $c_1, ..., c_m$ and the edges are the arcs of the diagram D. We orient the edges of this graph such that the two undercrossing (resp. over-crossing) arcs at each crossing correspond to incoming (resp. outgoing) edges. Furthermore, we label the edges with weights t and 1 such that, as one traverses the over-strand at crossing c_i (following the link's orientation), the weight t (resp. 1) appears on the right (resp. left). We choose an arbitrary vertex, say c_k , of this graph as the root. Let $\mathcal{T}(G_D, c_k)$ be the set of all maximal oriented trees rooted at c_k . For a tree $T \in \mathcal{T}(G_D, c_k)$, we define its weight W(T) to be the product of weights of all of its edges. The *Crowell polynomial* associated to (G_D, c_k) is

$$P_{G_D,c_k}(t) := \sum_{T \in \mathcal{T}(G_D,c_k)} W(T).$$

Note that the Crowell polynomial $P_{G,v}(t)$ can be defined for any weighted, oriented graph G rooted at a vertex v.



FIGURE 9. Crowell's method [Jon10].

Theorem 2.7. [Cro59] Let $\Delta_L(t)$ be the normalized Alexander polynomial of an alternating link L with alternating diagram D. If c_k is a crossing of D, then

$$\Delta_L(-t) \doteq P_{G_D,c_k}(t),$$

where \doteq means equality up to multiplication by units of the Laurent polynomial ring $\mathbb{Z}[t, t^{-1}]$; i.e., by $\pm t^k$.

The absolute values of the coefficients of $\Delta_L(t)$ and $\Delta_L(-t)$ agree. In the rest of this subsection, we are going to focus on the coefficients of the Crowell polynomial.

Assume MT(L) = n, and let R be an coherent twist region with n crossings c_1, \ldots, c_n , labelled from left to right, in some reduced alternating diagram D of L. We define D_k as the link diagram constructed by replacing R with an coherent twist region with k crossings c_1, \ldots, c_k , and let L_k be the corresponding link. Let G_k be the Crowell graph associated with D_k . Let v_1, \ldots, v_k be the vertices of G_k corresponding to the crossings c_1, \ldots, c_k .

Consider the graph $G'_k = G_k \setminus \{v_{k-1}\}$ for $k \geq 3$, where we also remove the edges adjacent to v_{k-1} ; see Figure 10. To define G'_2 , we add an edge to $G_2 \setminus \{v_1\}$, as follows. Consider the two edges to the left of v_1 , and denote the incoming edge by $e_{li} := (v_{li}, v_1)$ and the outgoing edge by $e_{lo} := (v_1, v_{lo})$. Let $G'_2 := G_2 \setminus \{v_1\} \cup \{e_{io}\}$, where $e_{io} := (v_{li}, v_{lo})$. Let the weight of the new edge e_{io} be the product of weights of e_{li} and e_{lo} . We can view the graph G'_k as the non-oriented smoothing of G_k at v_{k-1} . Let $L_k(c_{k-1}, ||)$ denote the 1-smoothing of L_k at c_{k-1} . Ignoring the weights and the edge orientations, G'_k is isomorphic to the Crowell graph corresponding to $L_k(c_{k-1}, ||)$, after deleting two loop edges when $k \geq 3$ and one loop edge when k = 2. Note that the two graphs have different weights and orientations; see Figure 10.

The following lemma is the recursive relation at the heart of the stabilization phenomenon. It is an unoriented skein relation for the Crowell polynomial.

Lemma 2.8. Let $P_k = P_{G_k,v_k}$ and $P'_k = P_{G'_k,v_k}$; i.e., the Crowell polynomials of the graphs G_k and G'_k rooted at v_k . Then

$$P_k = tP_{k-1} + P'_k.$$

Proof. We decompose $\mathcal{T}(G_k, v_k)$ into two sets \mathcal{T}_R and \mathcal{T}_L of oriented spanning trees depending the edge leading to v_{k-1} .

Let \mathcal{T}_R be the set of oriented spanning trees containing the edge (v_k, v_{k-1}) . By contracting this edge to the root vertex v_k and deleting the resulting loop edge, one can build a one-to-one matching between this set and $\mathcal{T}(G_{k-1}, v_{k-1})$. The matching decreases the exponent of t in W(T) by one; see Figure 11.

We write \mathcal{T}_L for the set of oriented spanning trees containing the edge (v_{k-2}, v_{k-1}) . These trees do not contain any other edge adjacent to v_{k-1} . As a result, \mathcal{T}_L is in bijection with $\mathcal{T}(G'_k, v_k)$. This bijection preserves the weights; see Figure 12.



FIGURE 10. Twist region in the Crowell graph.



FIGURE 11. The set \mathcal{T}_R is in bijection with $\mathcal{T}(G_{k-1}, v_{k-1})$. Trees containing the red edge are included in \mathcal{T}_R .



FIGURE 12. The set \mathcal{T}_L is in bijection with $\mathcal{T}(G'_k, v_k)$. Trees containing the red edge are included in \mathcal{T}_L .

By definition of P_k , we have

$$P_k = \sum_{T \in \mathcal{T}(G_k, v_k)} W(T)$$

=
$$\sum_{T \in \mathcal{T}_R} W(T) + \sum_{T \in \mathcal{T}_L} W(T)$$

=
$$\sum_{T \in \mathcal{T}(G_{k-1}, v_{k-1})} tW(T) + \sum_{T \in \mathcal{T}(G'_k, v_k)} W(T)$$

=
$$tP_{k-1} + P'_k,$$

as claimed.

14

Definition 2.9. We say that the sequence of positive integers (a_1, \ldots, a_m) is *half-trapezoidal* if

(1)
$$a_1 \leq a_2 \leq \cdots \leq a_{m-1} \leq a_m;$$

(2) if $a_1 = a_2$, then $a_2 = a_3$.

(2) if $a_i = a_{i+1}$, then $a_i = a_{i+1} = \dots = a_m$;

i.e., if it strictly increases and then possibly stabilizes.

Note that, if (a_1, \ldots, a_n) is a symmetric sequence of positive integers and (a_1, \ldots, a_m) is half-trapezoidal for $m = \lfloor n/2 \rfloor$, then (a_1, \ldots, a_n) is trapezoidal.

Proof of Theorem 2.6. Using the notation of Theorem 2.7 and Lemma 2.8, for $L = L_n$, we have $\Delta_L(-t) \doteq P_n$. Note that when L_n is alternating, L_1 is also alternating and hence $\Delta_{L_1}(-t) \doteq P_1$.

We prove the Theorem 2.6 when n is odd. We are going to use the fact that $|L_n| = |L_1|$ for odd n. The same argument can be repeated for even n, but one needs to use L_2 instead of L_1 .

We show that the sequence of coefficients of P_n is trapezoidal if $n-2 \ge g(L) + |L|/2$. By applying Lemma 2.8 repeatedly, we can deduce that

(2)
$$P_n = t^{n-1}P_1 + (P'_n + tP'_{n-1} + \dots + t^{n-2}P'_2).$$

Note that the only difference between G'_k , for different values of k, is the length of the chain of vertices v_1, \ldots, v_{k-2} . Any tree in $\mathcal{T}(G'_k, v_k)$ must contain the edges $(v_1, v_2), \ldots, (v_{k-3}, v_{k-2})$. Deleting the vertices v_1, \ldots, v_{k-2} and the incident edges from each maximal rooted tree, then replacing e_{li} with e_{io} if both e_{li} and e_{io} are in the original tree results in a one-to-one correspondence between $\mathcal{T}(G'_k, v_k)$ and $\mathcal{T}(G'_2, v_2)$; see Figure 13.

This correspondence preserves weights, so $P'_k = P'_2$. Combining this with equation (2) gives that

(3)
$$P_n = t^{n-1}P_1 + (1 + \dots + t^{n-2})P'_2.$$

Let mindeg(P) and maxdeg(P) be the minimal and maximal power of t that appear in a polynomial P(t), respectively. Let (a'_i) denote the sequence of coefficients of P'_2 and (c'_i) denote the sequence of coefficients of $(1 + \cdots + t^{n-2})P'_2$. The first n-1 terms of (c'_i) are the cumulative sums of the sequence (a'_i) ; i.e.,

$$c'_{\operatorname{mindeg}(P'_2)+j} = a'_{\operatorname{mindeg}(P'_2)} + \dots + a'_{\operatorname{mindeg}(P'_2)+j}$$

for all $j \in \{0, ..., n-2\}$. Note that the coefficients of Crowell polynomials are positive; see Remark 2.10. As a result, their cumulative sums form a half-trapezoidal sequence. This means that the first n-1 coefficients of $(1 + \cdots + t^{n-2})P'_2$ form a half-trapezoidal sequence.

We are going to show that the first n-2 coefficients of P_n agree with the first n-2 terms of (c'_i) and hence they are also half-trapezoidal. We need



FIGURE 13. The bijection between $\mathcal{T}(G'_k, v_k)$ and $\mathcal{T}(G'_2, v_2)$. The oriented spanning trees in $\mathcal{T}(G'_k, v_k)$ contain the red edges. The edges e_{lo} and e_{io} are shown in yellow.

to separate the terms that $(1 + \cdots + t^{n-2})P'_2$ contribute to P_n from the ones coming from $t^{n-1}P_1$. To be more precise, we will prove that

(4)
$$n-1 + \operatorname{mindeg}(P_1) = \operatorname{mindeg}(t^{n-1}P_1) = \operatorname{mindeg}(P_n) + n - 1.$$

In fact, one can prove that, for large enough n, we have mindeg (P_n) = mindeg (P_1) by analysing spanning trees, but to get to our desired lower bound for n, we use an alternative argument. We are going to show that

(5)
$$\max\deg(P_n) = \max\deg(t^{n-1}P_1)$$
, and

(6)
$$\operatorname{span}(P_n) - \operatorname{span}(t^{n-1}P_1) = n - 1$$

Equations (5) and (6) and the fact that $\operatorname{span}(P) = \operatorname{maxdeg}(P) - \operatorname{mindeg}(P)$ give us equation (4).

Note that any oriented spanning tree in $\mathcal{T}(G'_2, v_2)$ is also a spanning tree in $\mathcal{T}(G_1, v_1)$, which means all the terms in P'_2 also contribute to in P_1 . As a result, maxdeg $(P_1) \geq \max \deg(P'_2)$, hence

$$\max\deg(t^{n-1}P_1) > \max\deg((1+\cdots+t^{n-2})P_2').$$

Combining this with equation (3) implies that the term with the highest degree in P_n comes from $t^{n-1}P_1$. This is equation (5).

We can compare the span of P_1 and P_n by comparing the genera of L_1 and L_n . The links are both alternating and the span of the Alexander polynomial of an alternating link J is known to be 2g(J) + |J| - 1. We can write

$$\operatorname{span}(P_n) - \operatorname{span}(P_1) = 2(g(L_n) - g(L_1)) + |L_n| - |L_1|.$$

Since n is odd, L_n and L_1 differ in a number of full twists and $|L_n| = |L_1|$. To compare their genera, we can compare their Seifert graphs (i.e., the result of the Seifert algorithm). The Seifert graph of L_n has n parallel edges (i.e., half-twisted bands) between two vertices (i.e., Seifert cycles) where L_1 has one edge; see Figure 14. Since L_n and L_1 are alternating, their genera is realised by the Seifert algorithm and is equal to half of the rank of the first homology of the Seifert graph. We can deduce that $g(L_n) - g(L_1) = \frac{n-1}{2}$ which gives us equation (6).



FIGURE 14. A twist region is a family of parallel bands between a pair of Seifert cycles (not necessarily different cycles as seen in the right figure).

We have proved that the first n-2 coefficients of P_n come from $(1+\cdots+t^{n-2})P_2'$

and are hence half-trapezoidal. Furthermore, $\operatorname{span}(P_n) = 2g(L_n) + |L_n| - 1$. Now, using the assumption that $n - 2 \ge g(L_n) + |L_n|/2$, we can deduce that all the coefficients in the first half of $\operatorname{span}(P_n)$ are half-trapezoidal. The symmetry of P_n means that it is trapezoidal.

Remark 2.10. It is well know that the coefficient of t^i in the Crowell polynomial P of an alternating knot is positive for $i \in [\operatorname{mindeg}(P), \operatorname{maxdeg}(P)]$. It is non-negative by definition and proven to be non-zero. It turns out that this is true for the Crowell polynomial of any weighted, oriented, rooted graph (G, v). This follows from the fact that $\mathcal{T}(G, v)$ is the basis of a greedoid and hence it has a connected adjacency graph (See Corollary 5.7. of [BKL85]). This means that, for any $T, T' \in \mathcal{T}(G, v)$, there exist two ordered sets of edges $\{e_1, \ldots, e_n\} \subseteq E(T)$ and $\{e'_1, \ldots, e'_n\} \subseteq E(T')$ such that

 $T \setminus \{e_1, \dots, e_t\} \cup \{e'_1, \dots, e'_t\} \in \mathcal{T}(G, v)$

for every $t \in \{1, \ldots, n\}$, and

$$T \setminus \{e_1, \ldots, e_n\} \cup \{e'_1, \ldots, e'_n\} = T'.$$

We end Subsection 2.2 by mentioning some corollaries of Theorem 2.6. First, the proof of Theorem 2.6 immediately implies a more general result:

Theorem 2.11. Let L be an alternating link and

$$\Delta_L(t) \doteq \sum_{i=0}^{2g(L)+|L|-1} (-1)^i a_i t^i.$$

The sequence $\{a_i : 0 \le i \le MT(L) - 3\}$ is half-trapezoidal; i.e., it is strictly increasing until it possibly stabilizes.

It is not straightforward to find knots that satisfy the conditions of Theorem 2.6. If L is a non-trivial link, then $c(L)/2 \ge g(L) + |L|/2$, and it is easy to check whether the stronger condition $MT(L) - 3 \ge c(L)/2$ holds. For instance, if $B = \sigma_{i_1}^{n_1} \cdots \sigma_{i_k}^{n_k}$ is an alternating braid, then $MT(L) = \max\{n_i : 1 \le j \le k\}$. We obtain the following corrollary: **Corollary 2.12.** Let L_B be the closure of the braid $B = \sigma_1^{p_1} \sigma_2^{-q_1} \cdots \sigma_1^{p_n} \sigma_2^{-q_n}$. Assume that

$$p_1 \ge \sum_{i=2}^n p_i + \sum_{j=1}^n q_j + 6.$$

Then the trapezoidal conjecture holds for L_B .

We will see some other corollaries of Theorem 2.6 in the form of bounds on the length of the stable part in Section 3. In light of the results of this subsection, the following definition is useful.

Definition 2.13. We call a link L twist-concentrated if it satisfies the inequality

$$MT(L) - 3 \ge g(L) + |L|/2.$$

2.3. Plumbings of special alternating knots. In this subsection, we prove the trapezoidal conjecture for diagrammatic plumbings of special alternating links, using the fact that the conjecture holds for special alternating links [HMV23]. We explain the proof for a plumbing of a pair of special alternating links, but it can be extended to all diagrammatic plumbings. First, we define the plumbing operation, and, more generally, Murasugi sums. The following is the classical definition of Murasugi sum; see [Kaw96], [Gab85] and [HW05].

Definition 2.14. Consider two links L_1 and L_2 in S^3 with Seifert surfaces S_1 and S_2 . Let $S^3 = B_1 \cup_{S^2} B_2$ be the standard decomposition of the 3-sphere into 3-balls. Assume $S_i \subset B_i$ such that $S_1 \cap S_2$ is a disk $D \subset S^2$. The disk D is the interior of a 2n-gon; i.e., ∂D is decomposed into 2n arcs $a_1, b_1, \ldots, a_n, b_n$, labelled counter-clockwise, such that $a_i \subseteq \partial S_1$ and a_i is properly embedded in S_2 , and $b_i \subseteq \partial S_2$ and b_i is properly embedded in S_1 for $i \in \{1, \ldots, n\}$. The compact oriented surface $S = S_1 \cup_D S_2$ is called the Murasugi sum of S_1 and S_2 along the disk D. The link $L = \partial S$ is called the Murasugi sum of L_1 and L_2 along D; see Figures 15 and 16. Define the gonality of the Murasugi sum to be 2n.



FIGURE 15. A 4-gon Murasugi sum; i.e., plumbing [Kaw96].



FIGURE 16. A 6-gon Murasugi sum [AH21].

Taking 2-gon Murasugi sum is simply the connected sum operation. The 4-gon Mutasugi sum operation is called *plumbing*. As mentioned before, links that can be constructed as boundaries of plumbings of twisted bands over a tree are called *arborescent links*. Figures 3 and 4 show simple examples of arborescent links. General arborescent links look like Figure 17. Another class of such links is that of 2-bridge links; see Figure 18.



FIGURE 17. A general arborescent link [BS10].

Gabai proved a series of important results about Murasugi sums, which we summarize in the following theorem.

Theorem 2.15. [Gab85] The Murasugi sum of two incompressible (resp. minimal genus) Seifert surfaces is incompressible (resp. minimal genus), and, as a result, genus is additive under Murasugi sums along minimal genus Seifert surfaces. Furthermore, S is a fibre surface for L if and only if S_i is a fibre surface for L_i for $i \in \{1, 2\}$.

Definition 2.14 is the most general form of the Murasugi sum operation, but it has some limitations. One of these is that the operation cannot easily be seen in a diagram. Murasugi [Mur65] originally used a more restrictive

18



FIGURE 18. An illustration showing why 2-bridge links are arborescent [HT85].

definition, which we call *diagrammatic Murasugi sum*. We now recall that definition; see also [HW04] and [HW05]. It is more convenient to phrase it in terms of a *Murasugi decomposition algorithm* that decomposes any link diagram as a Murasugi sum, as follows.

Consider a link L with a connected diagram D_L , which consists of crossings c_1, \ldots, c_n and arcs e_1, \ldots, e_{2n} . One can also view the diagram as a 4-valent plane graph with vertices c_i , edges e_j , and regions r_1, \ldots, r_{n+2} . The set $\{e_1, \ldots, e_{2n}\}$ can be decomposed into Seifert cycles T_1, \ldots, T_q using Seifert's algorithm. One can split Seifert cycles into two types. We say that T_i is a type 1 Seifert cycle if it bounds a region r_k , otherwise it is type 2.

If all the Seifert cycles are type 1, then we call the link diagram special. Applying Seifert's algorithm to a special diagram gives a Seifert surface which is obtained by considering the disks bounded by the Seifert cycles of L and attaching half-twisted bands over each crossing of L. Note that this definition is the same as the median construction applied to the adjacency graph of the regions of the same color in a checkerboard coloring of $S^2 \setminus D_L$. We call this graph the *Tait graph*. In the literature, this surface is also known as the *checkerboard* or *Tait surface*. For the Tait surface to be orientable, the Tait graph needs to be bipartite. Figure 19 shows a special diagram and a two-coloring (blue and green) of the vertices of the Tait graph (regions with the same checkerboard color). Any link has a special diagram. See Burde [BZH13], Stoimenow [Sto16], and Hirasawa [Hir95] for algorithms to turn any diagram of a link into a special one, which is shown in Figure 19.

Let C_1, \ldots, C_m be the type 2 Seifert cycles in D. Each C_i is a simple closed curve in S^2 that divides S^2 into disks C_i^+ and C_i^- . Let C_i^+ be the disk containing ∞ . For $\gamma_i \in \{+, -\}$, let $D(\gamma_1, \cdots, \gamma_m)$ be the closure of $C_1^{\gamma_1} \cap \cdots \cap C_m^{\gamma_m}$. Exactly m + 1 of these sets will be nonempty, which we



FIGURE 19. Non-special and special diagrams of a link. The left diagram is non-special as it contains a type 2 Seifert cycle, colored yellow. One can turn the diagram into the special diagram on the right by moving the red segment.

call Seifert domains and denote by D_1, \ldots, D_{m+1} ; see Figure 20. For every $i \in \{1, \ldots, n\}$, at least two of the four regions adjacent to crossing c_i belong to the same Seifert domain D_j for some $j \in \{1, \ldots, m+1\}$. In this case, we say that c_i belongs to D_j . Each vertex belongs to exactly one Seifert domain. For $j \in \{1, \ldots, m+1\}$, let L_j be the link obtained by smoothing all the crossings along ∂D_j that do not belong to D_j . We call $\{L_1, \ldots, L_{m+1}\}$ the Murasugi decomposition of the diagram D_L , and denote it by

$$L = L_1 * L_2 * \cdots * L_{m+1}.$$

Note that one can construct the links L_j by adding vertical segments in the middle of the half-twisted bands which are used in Seifert's algorithm; see Figure 21. In this way, one can see that gluing the canonical Seifert surfaces for $L_1, ..., L_{m+1}$ results in a canonical Seifert surface for L. It is not trivial that this gluing is a Murasugi sum in the sense of Definition 2.14; see Remark 2.16. For an example of a Murasugi decomposition; see Figure 21.

Remark 2.16. Murasugi's *-product is a special case of Murasugi sum in the sense of Definition 2.14. We do not give a proof of this fact, but we show it holds for the example presented in Figure 21. This can be seen in Figure 22. Another example of this fact can be seen in Figure 23.

Remark 2.17. Each of the *-products (i.e., diagrammatic Murasugi sums) in the Murasugi decomposition happens over a Seifert cycle of type 2. To be more precise, consider a Seifert cycle C of type 2 which decomposes S^2 into disks C^+ and C^- . Note that each of the Seifert domains is either in C^+ or in C^- . Assume that $D_1, \ldots, D_i \subseteq C^-$ and $D_{i+1}, \ldots, D_{m+1} \subseteq C^+$. Furthermore, assume C sits between the domains D_i and D_{i+1} . Let L^+ and L^- be the result of the restriction of the link L to C^+ and C^- , respectively. Then

$$L = L^- * L^+,$$



FIGURE 20. Seifert domains and their crossings. Note that none of the crossings belong to D_3 .



FIGURE 21. An example of a Murasugi decomposition. The first row shows two of the Seifert domains and the associated Murasugi summands. Note that the third summand L_3 is trivial and is hence not depicted. The leftmost picture in the second row shows the Murasugi decomposition as a Murasugi sum (related to Figure 22). Other figures in the second row depict Seifert surfaces of the Murasugi summands constructed by cutting the canonical Seifert surface of L.

where the Murasugi sum is taken over the disk associated with the Seifert cycle C in the canonical Seifert surface; see the yellow disks in Figures 22 and 23. This follows from the fact that applying the Murasugi decomposition



FIGURE 22. The *-product is a special case of Murasugi sum (shown in the example of Figure 21).



FIGURE 23. A second example showing that *-product is a special case of Murasugi sum. Note how the gluing disk of the Murasugi sum is slightly different from the one in Figure 22

algorithm to L^- and L^+ gives us

$$L^{-} = L_1 * \cdots * L_i$$
 and $L^{+} = L_{i+1} * \cdots * L_{m+1}$.

Let the gonality of the Murasugi sum over the Seifert cycle C be 2s. We can compute 2s from the diagram as follows. Crossings on C are of two types: ones that belong to D_i and ones that belong to D_{i+1} (shown in red and blue in Figure 24). Moving along C, one alternately encounters x_j crossings belonging to D_i , then y_j crossings belonging to D_{i+1} for $j = 1, \ldots, l$, for some l. Using this notation, we have

(7)
$$s = \sum_{j=1}^{l} x_j + \sum_{j=1}^{l} y_j - l.$$

We call *l* the *length* of the diagrammatic Murasugi sum.



FIGURE 24. Local picture of a type 2 Seifert cycle C.

Murasugi's decomposition has an important property. If D_L is a reduced alternating diagram, then all summands L_i will be alternating as well. This leads to the first part of the following result.

Proposition 2.18. [Mur65] Any alternating link can be written as a *product (i.e., diagrammatic Murasugi sum) of special alternating links. Furthermore, signature and genus are additive under this decomposition.

Note that, applying Seifert's algorithm to an alternating diagram leads to a minimal genus Seifert surface, and the diagrammatic Murasugi sum is taken using such surfaces. Based on Theorem 2.15, genus is additive under Murasugi decomposition of an alternating link. Murasugi proved that signature is also additive under such a decomposition. We will revisit these facts in Subsection 3.3.

Definition 2.19. We say that a link diagram is *decorated* if it has a marked edge, or, equivalently, two marked adjacent regions. Fix a checkerboard coloring of the regions of a decorated link diagram. The black (resp. white) *Tait graph* has vertices the black (resp. white) regions, and we add an edge between two regions through each crossing they share. We use the notation B (resp. W) for this graph. The graph B (resp. W) is rooted at the vertex corresponding to the marked region; see Figure 25. Note that B and W are planar duals.

Let \mathcal{T}_B (resp. \mathcal{T}_W) be the set of rooted spanning trees of B (resp. W), oriented away from the root. There is a natural one-to-one correspondence



FIGURE 25. The white (left) and black (right) Tait graphs in a marked link diagram.

between \mathcal{T}_B and \mathcal{T}_W coming from planar duality. Let $\mathcal{D}_{(B,W)}$ denote the set of all dual pairs in $\mathcal{T}_B \times \mathcal{T}_W$; i.e.,

$$\mathcal{D}_{(B,W)} := \{ (T,T') \in \mathcal{T}_B \times \mathcal{T}_W : T, T' \text{ are planar duals} \}.$$

We call each dual pair a *Kauffman state*, and write $\mathcal{D}_{(B,W)}$ for the set of all Kauffman states.

Kauffman's state-sum formula computes the Alexander polynomial as a weighted sum of Kauffman states. The weights are assigned as follows. A directed edge of either of the Tait graphs passes through a crossing c. It is directed towards one of the four corners around the crossing c. Figure 26 shows the weights assigned to each corner. The weight of a rooted tree is defined as the product of the weights assigned to its edges.



FIGURE 26. Kauffman weights.

Kauffman's state-sum formula for the Alexander polynomial is

(8)
$$\tilde{\Delta}_K(t) = \sum_{(T,T')\in\mathcal{D}_{(B,W)}} w(T) \cdot w(T').$$

We are now ready to prove the main theorem of this subsection.

Theorem 2.20. The trapezoidal conjecture holds for any link $L = L_1 * L_2$ which is a diagrammatic plumbing (4-gon Murasugi sum) of two special alternating links L_1 and L_2 .

Proof. First, we give a diagrammatic characterization of the plumbing. Choose a reduced alternating diagram D of L. Since we only have two summands, there is only one type 2 Seifert cycle C, and the Seifert domains

are the two disks it bounds in S^2 . Using the notation of equation (7), assume first that l = 1. If $y_1 = 1$, the crossing connecting C to its exterior is reducible (also called nugatory), contradicting the assumption that D is reduced. Hence $y_1 \ge 2$. Similarly, $x_1 \ge 2$. By equation (7), the gonality of the Murasugi sum is at least six, which contradicts the fact that we have a 4-gon Murasugi sum. So $l \ge 2$, which, together with equation (7), implies that l = 2 and $x_j = y_j = 1$ for $j \in \{1, 2\}$. So the diagram D looks like Figure 27 in a neighbourhood of C.

Note that Seifert cycles in a special diagram bound regions of the link diagram. As a result, one can describe the diagrammatic plumbing of two special alternating links as follows. Consider the special alternating links L_1 and L_2 , and let R_i be a region of L_i for $i \in \{1,2\}$ such that R_i has degree two as a vertex of the Tait graph of L_i (i.e., it has two crossings along its boundary). Change R_1 to the infinity region (i.e., the region containing $\infty \in S^2 = \mathbb{R}^2 \cup \{\infty\}$) in the diagram of L_1 . Then place the diagram of L_1 inside region R_2 in the diagram of L_2 , identifying the boundaries of R_1 and R_2 according to Figure 27. See Figure 28 for an example.



FIGURE 27. Plumbing in a Murasugi decomposition.

Using this diagrammatic characterization, we can see the effect of the operation on Tait graphs. Let B_{L_i} and W_{L_i} denote the black and white Tait graphs of the special alternating link L_i , respectively, for $i \in \{1, 2\}$. Without loss of generality, assume that R_1 and R_2 are both black regions in the respective colorings of the diagram of L_1 and L_2 . We also use R_1 and R_2 to denote the corresponding vertices of B_{L_1} and B_{L_2} , respectively. Furthermore, we can assume that the checkerboard coloring of L_2 extends to the coloring of L, and let B_L and W_L denote the black and white Tait graphs of L. These graphs can be described as follows. See Figures 29 and 30 for an example and some notation.

The gaph B_L can be constructed from B_{L_2} by replacing the vertex R_2 with the graph W_{L_1} , as follows. Let w and w' be the neighbors of R_1 in W_{L_1} , and let b and b' be the neighbors of R_2 in B_{L_2} . We remove R_1 from W_{L_1} and R_2 from B_{L_2} , then add the edges (b, w) and (b', w'); see Figure 30. We call this operation a *Tait blow-up of the region* R_2 . The graph W_L can be constructed using a similar process, with the roles of the two links reversed.



FIGURE 28. Example of diagrammatic plumbing. The links from left to right are L_2 and L_1 . The red highlighted regions are R_2 and R_1 . The green Seifert cycle is C, whose neighborhood is shown in Figure 27.

The red stars in Figure 30 are the roots (i.e., marked vertices). The yellow vertex y is associated with the unbounded region. It also shows a pair of dual spanning trees $(T, T') \in \mathcal{T}_{B_L} \times \mathcal{T}_{W_L}$, colored red and yellow, respectively. Note that the crossings in Figure 30 both belong to L_2 ; cf. Figure 29.



FIGURE 29. Tait blow-up (Tait graph of a plumbing).

Since T is a spanning tree rooted at b, it contains unique directed paths β from b to b' and ω from w to w'. Depending on β and ω , there are three distinct possibilities for a pair of dual spanning trees $(T, T') \in \mathcal{T}_{B_L} \times \mathcal{T}_{W_L}$; see Figure 31:

Case (i): β passes through w and then w'.

Case (ii): $\beta \subseteq B_{L_2}$ and $\omega \subseteq W_{L_1}$. This case has two subcases:

Case (ii).1: T contains the edge (b, w).

Case (ii).2: T contains the edge (b', w').

Case (iii): $\beta \subseteq B_{L_2}$ and T contains both (b, w) and (b', w').

26



FIGURE 30. A pair of dual trees in $\mathcal{T}_{B_L} \times \mathcal{T}_{W_L}$.

For $* \in \{(i), (ii), (ii), (ii), (ii), (iii), (iii), (iii)\}$, we denote the set of (T, T') in Case * by $\mathcal{D}^*_{(B_L, W_L)}$. By analyzing the possible options for the paths β and ω , we obtain the partition

$$\mathcal{D}_{(B_L,W_L)} = \mathcal{D}_{(B_L,W_L)}^{(i)} \sqcup \mathcal{D}_{(B_L,W_L)}^{(ii)} \sqcup \mathcal{D}_{(B_L,W_L)}^{(iii)}.$$

We now look at these cases one-by-one. In each case, we decompose the dual spanning trees T and T' into dual spanning trees of the Tait graphs of L_1 and L_2 , and relate their weights to the weights of Kauffman states in $\mathcal{D}_{(B_{L_1},W_{L_1})}$ and $\mathcal{D}_{(B_{L_2},W_{L_2})}$.

 $\mathcal{D}_{(B_{L_1},W_{L_1})}$ and $\mathcal{D}_{(B_{L_2},W_{L_2})}$. **Case (i):** The edges of $T \cap W_{L_1}$ form a spanning tree T_1 of W_{L_1} rooted at w. The rest of the edges form a spanning tree T_2 of B_{L_2} after replacing the edges (b,w) and (w',b') with the edges (b,R_2) and (R_2,b') . There is an analogous decomposition of the planar dual T' of T into spanning trees T'_1 of B_{L_1} and T'_2 of W_{L_2} , with one small modification. Consider the vertex x of T'that lies between the summands L_1 and L_2 , and between the two crossings between the two summands. The outgoing edges of x become the outgoing edges of y in T'_1 . See Figure 32 for an illustration of the decomposition.

This decomposition gives us the map

(9)
$$\psi: \mathcal{D}_{(B_L,W_L)}^{(i)} \hookrightarrow \mathcal{D}_{(B_{L_2},W_{L_2})} \times \mathcal{D}_{(W_{L_1},B_{L_1})}.$$
$$(T,T') \mapsto \left((T_2,T'_2), (T_1,T'_1) \right)$$

If T_2 contains the edges (b, R_2) and (R_2, b') , then the construction can be reversed, so ψ is injective. Let

$$\mathcal{D}_{(B_{L_2}, W_{L_2})}^{(i)} := \{ (T_2, T_2') \in \mathcal{D}_{(B_{L_2}, W_{L_2})} : (b, R_2), (R_2, b') \in T_2 \}$$

Then

(...)

(10)
$$\psi \colon \mathcal{D}_{(B_L, W_L)}^{(i)} \xrightarrow{\sim} \mathcal{D}_{(B_{L_2}, W_{L_2})}^{(i)} \times \mathcal{D}_{(W_{L_1}, B_{L_1})}$$



FIGURE 31. The possible cases for the paths β from b to b' and ω from w to w' in a rooted, directed spanning tree T of B_L , together with the dual spanning tree T' of W_L .



FIGURE 32. Decomposition of spanning trees in Case (i).

is a bijection. This map just changes the ends of some of the edges and does not change their position with respect to crossings, hence it preserves the weights of states: If $\psi(T, T') = ((T_2, T'_2), (T_1, T'_1))$, then

(11)
$$w(T) = w(T_2) \cdot w(T_1) \text{ and } w(T') = w(T'_2) \cdot w(T'_1).$$

Case (ii): We decompose dual pairs of spanning trees in $\mathcal{D}_{B_L,W_L}^{(ii).1}$ as in Figures 33 and those in $\mathcal{D}_{B_L,W_L}^{(ii).2}$ as in Figure 34. Similarly to Case (i), we define the following sets of dual pairs of spanning trees:

$$\mathcal{D}_{(B_{L_2}, W_{L_2})}^{(ii).1} := \{ (T_2, T_2') \in \mathcal{D}_{(B_{L_2}, W_{L_2})} : (b, R_2) \in T_2 , (R_2, b') \notin T_2 \},$$

$$\mathcal{D}_{(B_{L_2},W_{L_2})}^{(ii),2} := \{ (T_2,T_2') \in \mathcal{D}_{(B_{L_2},W_{L_2})} : (R_2,b') \in T_2 , (b,R_2) \notin T_2 \}.$$

Decomposing dual pairs of spanning trees as

$$(T, T') \mapsto ((T_2, T'_2), (T_1, T'_1))$$

give bijections

(12)
$$\mathcal{D}_{(B_L,W_L)}^{(ii).1} \xrightarrow{\sim} \mathcal{D}_{(B_{L_2},W_{L_2})}^{(ii).1} \times \mathcal{D}_{(W_{L_1},B_{L_1})}$$

(13)
$$\mathcal{D}_{(B_L,W_L)}^{(ii).2} \xrightarrow{\sim} \mathcal{D}_{(B_{L_2},W_{L_2})}^{(ii).2} \times \mathcal{D}_{(W_{L_1},B_{L_1})}$$

These maps preserve the weights of the Kauffman states in the sense of equation (11).

A spanning tree of B_{L_2} rooted at *b* contains unique directed paths from *b* to *b'* and from *b* to R_2 . There are three possibilities for these paths, giving the partition

$$\mathcal{D}_{(B_{L_2}, W_{L_2})} = \mathcal{D}_{(B_{L_2}, W_{L_2})}^{(i)} \sqcup \mathcal{D}_{(B_{L_2}, W_{L_2})}^{(ii).1} \sqcup \mathcal{D}_{(B_{L_2}, W_{L_2})}^{(ii).2}.$$

We can now compute the weighted sum of all the dual pairs in Cases (i) and (ii). Using the weight-preserving maps in equations (10), (12) and (13), we can write

The last equality comes from the Kauffman state-sum formula. Note that, although the roots of the spanning trees of W_{L_1} is w in Cases (i) and (ii).1 and is w' in Case (ii).2, this does not affect the result of the state-sum.

The product of two symmetric trapezoidal polynomials (i.e., the convolution of their coefficient sequences) with coefficients of alternating signs is also trapezoidal; see [Mur85, Proposition 2.1]. Since the trapezoidal conjecture for special alternating links by Hafner, Mészáros, and Vidinas [HMV23], the sequence of coefficients of $\tilde{\Delta}_{L_1} \cdot \tilde{\Delta}_{L_2}$ is trapezoidal.



FIGURE 33. Decomposition of spanning trees in Case (ii).1.



FIGURE 34. Decomposition of spanning trees in Case (ii).2.

Case (iii): What remains is to show that the weighted sum of all the spanning trees in Case (iii) is also trapezoidal. Let \tilde{L}_2 be the result of deleting R_2 from L_2 and smoothing the two crossings on its boundary. Let $W_{\hat{L}_1}$ be the result of contracting w and w' in W_{L_1} to a single vertex $\{w, w'\}$. The link \hat{L}_1 can be constructed using the median construction from this graph. Note that $B_{\tilde{L}_2}$ sits inside B_L . For a spanning tree T of B_L , let $\tilde{T}_2 := T \cap B_{\tilde{L}_2}$. Due to the definition of Case (iii), the graph \tilde{T}_2 is a rooted spanning tree of $B_{\tilde{L}_2}$. The rest of the edges of T form a spanning forest

of W_{L_1} with two components rooted at w and w'. By contracting w and w', this forest becomes a spanning tree \hat{T}_1 of $W_{\hat{L}_1}$ rooted at $\{w, w'\}$. This decomposition also extends to the planar duals T', \hat{T}'_1 , and \tilde{T}'_2 , with a small modification. The outgoing edges of x turn into the outgoing edges of y in \tilde{T}'_2 . See Figure 35.



FIGURE 35. Decomposition of spanning trees in Case (iii).

Similarly to the previous cases, this construction is reversible, so gives us a bijection

(14)
$$\mathcal{D}_{(B_L,W_L)}^{(iii)} \xrightarrow{\sim} \mathcal{D}_{(B_{\tilde{L}_2},W_{\tilde{L}_2})} \times \mathcal{D}_{(W_{\tilde{L}_1},B_{\tilde{L}_1})}$$
$$(T,T') \mapsto \left((\widetilde{T}_2,\widetilde{T}_2'), (\widehat{T}_1,\widehat{T}_1') \right)$$

This map is also weight-preserving in the sense of equation 11. This part of the argument is a bit different from the previous cases since this decomposition deletes the edges (b, w) and (b', w'). Both of these edges have weight 1, and hence this does not affect the overall weight of the Kauffman states. This decomposition also deletes two of the edges of T'. These are the edges that pass over the other two crossings on the boundary of the type 2 Seifert cycle C. One can see that these two edges also have weight 1.

Using this weight-preservation, we can finally compute the sum of the weights of all the dual pairs of spanning trees in Case (iii) as follows:

$$\sum_{\substack{\mathcal{D}_{(B_{L_{2}},W_{L_{2}})}} w(T) \cdot w(T') =} \left(\sum_{\substack{\mathcal{D}_{(B_{L_{2}},W_{L_{2}})}} w(\widetilde{T}_{2}) \cdot w(\widetilde{T}_{2}')} \right) \left(\sum_{\substack{\mathcal{D}_{(W_{\widehat{L}_{1}},B_{\widehat{L}_{1}})}} w(\widehat{T}_{1}) \cdot w(\widehat{T}_{1}')} \right) = \tilde{\Delta}_{\widetilde{L}_{2}} \cdot \tilde{\Delta}_{\widehat{L}_{1}}$$

This polynomial is also trapezoidal as both L_2 and \tilde{L}_1 have bipartite Tait graphs and hence are special alternating links. This is clear for $B_{\tilde{L}_2}$ = $B_{L_2} \setminus \{R_2\}$. The graph $B_{\widehat{L}_1}$ is also bipartite as its planar dual $W_{\widehat{L}_1}$ comes from contracting two non-adjacent vertices in W_{L_1} . Since B_{L_1} is bipartite, degrees of all vertices in W_{L_1} are even and as a result all of the degrees in $W_{\widehat{L}_1}$ are also even.

Let \tilde{L}_1 be the result of deleting R_1 from L_1 and smoothing the two crossings on its boundary. In $W_{\tilde{L}_1}$, the vertices w and w' (corresponding to neighboring regions of R_1) are contracted to a single vertex $\{w, w'\}$. One can see that $W_{\tilde{L}_1}$ and $W_{\hat{L}_1}$ only differ in some loop edges connected to the vertex $\{w, w'\}$. Loop edges of the Tait graph pass over reducible crossings and do not change the isotopy class of the link. Hence, the links \tilde{L}_1 and \hat{L}_1 are equivalent, so $\tilde{\Delta}_{\tilde{L}_1} = \tilde{\Delta}_{\hat{L}_1}$.

These computations lead to the following final equation for the Alexander polynomial of the diagrammatic plumbing:

(15)
$$\tilde{\Delta}_L = \tilde{\Delta}_{L_1} \cdot \tilde{\Delta}_{L_2} + \tilde{\Delta}_{\tilde{L}_1} \cdot \tilde{\Delta}_{\tilde{L}_2}.$$

We only need to compare the degrees of the two trapezoidal summands $\tilde{\Delta}_{L_1} \cdot \tilde{\Delta}_{L_2}$ and $\tilde{\Delta}_{\tilde{L}_1} \cdot \tilde{\Delta}_{\tilde{L}_2}$ to prove that $\tilde{\Delta}_L$ is trapezoidal. To be precise, we show that

(16)
$$\deg(\tilde{\Delta}_{L_1} \cdot \tilde{\Delta}_{L_2}) = \deg(\tilde{\Delta}_{\tilde{L}_1} \cdot \tilde{\Delta}_{\tilde{L}_2}) + 1;$$

see [Mur85, Proposition 2.1] and note that we are working with symmetrized Alexander polynomials. We determine the degrees of these polynomials by computing the genera of \hat{L}_1 and \tilde{L}_2 . Since the median construction on the black Tait graphs yield minimal genus Seifert surfaces, we only need to compute the rank of the first homology of these graphs. For a graph B, we have

$$\operatorname{rank}(H_1(B)) = \#(\operatorname{edges}) - \#(\operatorname{vertices}) + \#(\operatorname{components}).$$

This formula can be used to deduce that

$$\operatorname{rank}(H_1(B_{\widetilde{L}_2})) = \operatorname{rank}(H_1(B_{L_2})) - 1.$$

Deleting R_2 does not increase the number of components of the black Tait graph since the diagram is reduced and the degree of R_2 is two. Similarly, we have

$$\operatorname{rank}(H_1(B_{\widetilde{L}_1})) = \operatorname{rank}(H_1(B_{L_1})) - 1.$$

This proves equation (16) and concludes the proof of Theorem 2.20. \Box

We end Subsection 2.3 by mentioning some corollaries of Theorem 2.20 and its proof. The formula derived for the Alexander polynomial of diagrammatic plumbings might be of independent interest:

Proposition 2.21. Let the link $L = L_1 * L_2$ be the diagrammatic plumbing of the links L_1 and L_2 . Construct \tilde{L}_1 and \tilde{L}_2 as in the proof of Theorem 2.3; i.e., by deleting the regions involved in the plumbing of L_1 and L_2 . Then

$$\tilde{\Delta}_L = \tilde{\Delta}_{L_1} \cdot \tilde{\Delta}_{L_2} + \tilde{\Delta}_{\tilde{L}_1} \cdot \tilde{\Delta}_{\tilde{L}_2}.$$

As mentioned before, one can use an inductive argument to generalize Theorem 2.20 to the diagrammatic plumbing of several special alternating links.

Theorem 2.22. The trapezoidal conjecture holds for the diagrammatic plumbing $L_1 * \cdots * L_n$ of special alternating links L_1, \ldots, L_n .

Proof. The proof proceeds by considering the innermost Seifert cycle C_m of type 2. The interior disk C_m^+ bounded by C_m is also a Seifert domain and contains a summand of the diagrammatic plumbing. Without loss of generality, assume this summand is L_n and define $L' = L_1 * \cdots * L_{n-1}$.

Now note that a diagrammatic Murasugi decomposition always happens over a tree T. This is due to the inclusion hierarchy of type 2 Seifert cycles, which induces a partial order on the summands. In this Murasugi sum tree, L_n corresponds to a leaf vertex v in T. Let L_i be the summand corresponding to the neighbor of v in T.

We use an induction on the number of summands in the diagrammatic plumbing. Applying Proposition 2.21 to $L = L_n * L'$ gives us

$$\tilde{\Delta}_L = \tilde{\Delta}_{L_n} \cdot \tilde{\Delta}_{L'} + \tilde{\Delta}_{\widetilde{L}_n} \cdot \tilde{\Delta}_{\widetilde{L}'}.$$

The trapezoidal conjecture holds for L_n and \tilde{L}_n which are special alternating. It also holds for L' due to the induction hypothesis. Furthermore, one can show that

$$\widetilde{L}' = L_1 * \cdots * \widetilde{L}_i * \cdots * L_{n-1},$$

and hence the trapezoidal conjecture holds for \widetilde{L}' as well. This concludes the proof.

The proof of Theorem 2.20 does not need the Murasugi sums to be plumbings and a weaker condition also suffices. One can modify the proof to work in the case when all diagrammatic Murasugi sums are of length one or two. This leads to Theorem 2.23.

Theorem 2.23. The trapezoidal conjecture holds for the diagrammatic Murasugi sum $L_1 * \cdots * L_n$ of special alternating links L_1, \ldots, L_n such that the length of all Murasugi sums are less than three.

Remark 2.24. The condition that the length of a Murasugi sum is less than three means that we allow x_1, x_2, y_1, y_2 to be any positive integers, but there are no x_i, y_i for $i \geq 3$, using the notation of Figure 24. The most important change is in the definition of \tilde{L}_1 and \tilde{L}_2 in the formula of Proposition 2.21. In this general setting, one must construct \tilde{L}_1 (resp. \tilde{L}_2) by gluing an unknotted band to L_1 (resp. L_2) inside the region R_1 (resp. R_2) and separating the two groups of twists along its boundary. Gluing these bands changes the Tait graph by a planar contraction of two vertices (or a decomposition of a vertex into two in the dual plane graph), and this is exactly what we need for generalizing the proof of Theorem 2.20. See Figures 36, 37, and 38 for some examples. Note that the gluing construction generalizes the definition in the case of plumbings, as gluing a band to a bigon region results in two kinks that can be removed by Reidmeister 1 moves, hence is equivalent to smoothing the two crossings along the boundary.

Remark 2.25. Suppose one tries to use the same method for Murasugi sums of length at least three. In that case, extra terms appear in the decomposition that cannot be easily grouped together such that they sum up to known trapezoidal polynomials. This can also be seen in Alrefai–Chbili [AC21], where the terms appearing in the formula for the Alexander polynomial of 3-braids of length 3 is much more complicated than 3-braids of length 2.



FIGURE 36. A diagrammatic Murasugi sum of length two.



FIGURE 37. Construction of \widetilde{L}_1 in the example of Figure 36.

2.4. A multivariable refinement of the Alexander polynomial for alternating 3-braids. In this subsection, we prove some inequalities between coefficients of the Alexander polynomial of alternating 3-braids. We start with a combinatorial generalization of the work of Hafner, Mészáros, and Vidinas [HMV23] on special alternating knots. First, we review some of their main ideas.



FIGURE 38. Construction of \widetilde{L}_2 in the example of Figure 36.

Consider a special alternating link L with diagram D. Based on Subsection 2.3 (see Figures 19 and 39), we know that one of the Tait graphs of L is a bipartite plane graph. We denote this graph by B. Furthermore, let G_D be the Crowell graph associated to D.



FIGURE 39. A bipartite Tait graph B, the corresponding special alternating link L, and the Crowell graph G_D ; see [HMV23].

Note that a special alternating link is either positive or negative. Indeed, suppose that there were two consecutive crossings of different signs. See Figure 40, where the boundaries of three different Seifert cycles are highlighted. These Seifert cycles cannot all bound regions of the link diagram, so they cannot all be Seifert cycles of type 1. Hence, all crossings have the same sign in a special alternating diagram.

Up to taking its mirror, one can assume that the special alternating diagram is positive. Since the mirror of a knot has the same Alexander polynomial, we only consider positive diagrams in this subsection.

One of the main ideas of Hafner, Mészáros, and Vidinas is to assign new weights to the edges of G_D . They choose the weights carefully so that



FIGURE 40. Two consecutive crossings of different signs and the Seifert cycles in their neighbourhood.

Crowell's spanning tree weighted sum gives a multi-variable refinement of the Alexander polynomial.

Recall that vertices of B correspond to black regions in G_D . We use the same symbol to refer to a vertex and the corresponding region. The Crowell graph of a special alternating diagram has the following property.

Proposition 2.26. The edges of G_D are oriented such that the boundary of any black region $v \in B$ is oriented clockwise. Furthermore, there is a partition $V(B) = S \sqcup R$ of the vertex set of B such that all the edges on the boundary of a region in S have weight t, and all the edges on the boundary of a region in R have weight 1.

Proof. Using the fact that the link is special alternating, the result follows from a local analysis of the orientations and weights of the edges of the Crowell graph around vertices of the bipartite Tait graph B, as shown in Figure 41; see also Figure 39.



FIGURE 41. A special alternating link (left) and its Crowell graph (right). The blue and red vertices are in S and R, respectively.

We define the multi-variable refinement of the Alexander polynomial as follows. Let $S = \{v_1, \ldots, v_n\}$ and $R = \{v_{n+1}, \ldots, v_{n+m}\}$, and let \widetilde{G}_D be the

same oriented rooted graph as the Crowell graph G_D , but with different weights. Let the weight of the edges along the boundary of the region corresponding to v_i be x_i . Recall that $\mathcal{T}(\tilde{G}_D, c_1)$ is the set of all maximal oriented trees in \tilde{G}_D rooted at c_1 . We define

$$P_{\widetilde{G}_D,c_1}(x_1,\ldots,x_n,x_{n+1},\ldots,x_{n+m}) = \sum_{T \in \mathcal{T}(\widetilde{G}_D,c_1)} W(T),$$

where W(T) is the product of the weights of the edges in T; see Figure 42.



 $W(T) = x_1 x_2 x_3^2 x_5 x_6$



Note that $P_{\widetilde{G}_D,c_1}$ is a refinement of the Crowell polynomial for special alternating links, as

$$P_{\widetilde{G}_D,c_1}(t,\ldots,t,1,\ldots,1) = P_{(G_D,c_1)}(t) = \Delta_L(-t).$$

The advantage of working with this refinement is that all of its non-zero coefficients turn out to be 1. Furthermore, its support has a specific combinatorial convexity property called *M*-convexity; see Definition 2.33. (Here, the support Supp(P) of a multi-variable polynomial P is the set of all $(a_1, \ldots, a_{n+m}) \in \mathbb{Z}^{n+m}$ such that the monomial $x_1^{a_1} \cdots x_{n+m}^{a_{n+m}}$ has non-zero coefficient in P.) These facts follow from a result proved by the third author [Ká13]; see also [HMV23]:

Lemma 2.27. Let $C_1, \ldots, C_n, C_{n+1}, \ldots, C_{n+m}$ be cycles in \widetilde{G}_D such that C_i is the boundary of the region v_i . Let T be any spanning tree in the unoriented graph underlying \widetilde{G}_D . Let $a_i(T)$ be the number of edges of T lying in C_i . Then there exists a unique oriented tree A rooted at c_1 such that $a_i(A) = a_i(T)$ for all $i \in \{1, \ldots, n+m\}$.

Using Lemma 2.27, one can conclude that all non-zero coefficients of $P_{\widetilde{G}_{D,C1}}$ are 1. Furthermore,

$$\operatorname{Supp}(P_{\widetilde{G}_D,c_1}) = \{(a_1(T), a_2(T), \dots, a_{n+m}(T)) : T \in T(\widetilde{G}_D)\},\$$

where $T(\tilde{G}_D)$ is the set of all spanning trees of the unoriented graph underlying \tilde{G}_D . The set $T(\tilde{G}_D)$ is *M*-covex as a subset of $2^{E(\tilde{G}_D)}$. (In fact, it is the basis of the graphical matroid of the unoriented graph underlying \tilde{G}_D .)

These facts were used by Hafner, Mészáros, and Vidinas [HMV23] to prove that $P_{\tilde{G}_D,c_1}$ is *denormalized Lorentzian*, which is a generalization of log-concavity developed by Brändén and Huh [BH20]. This implies that the Alexander polynomial is log-concave.

The main problem with generalizing this idea to non-special diagrams is that Proposition 2.26 does not hold. Type 2 Seifert cycles still correspond to oriented cycles in the Crowell graph, but they contain edges with weights both t and 1. To construct a multi-variable refinement of the Alexander polynomial, one needs to use two variables for the edges of a type 2 Seifert cycle. See Figure 43 for an example. Note that, we also use the term "Seifert cycle" to refer to the corresponding oriented cycle in the Crowell graph.

The simplest class of non-special alternating links is given by closures of alternating 3-braids, which have only one type 2 Seifert cycle. We are going to focus on this family of links in the rest of this subsection.

Let D be the standard diagram of the closure of an alternating 3-braid

$$\sigma_1^{p_1}\sigma_2^{-q_1}\cdots\sigma_1^{p_n}\sigma_2^{-q_n}$$

for p_i , $q_i \in \mathbb{Z}_{>0}$, where σ_1 and σ_2 are the standard generators of the braid group on three strands. This diagram D is depicted on the left of Figure 43 as a long knot, with the second strand of the braid closing at ∞ . This diagram contains two type 1 Seifert cycles, one following the leftmost strand and one following the rightmost. We denote them by L and R, respectively, shown in blue and green in Figure 43. The diagram also has a type 2 Seifert cycle following the middle strand.

In the Crowell graph G_D , the left Seifert cycle is oriented counterclockwise with weight t and the right one is oriented clockwise with weight 1. The middle strand is directed from top to bottom, and it contains two types of edges. The edges ending in L, denoted by EL, with weight 1, and the edges ending in R, denoted by ER, with weight t. The refined Crowell graph \tilde{G}_D is shown on the right of Figure 43. The refined weights are x_1, x_2, x_3 , and x_4 on L, ER, EL, and R, respectively. In order to simplify the notation in the following argument, we use the color-coding of Figure 43. We use colors blue, red, black, green instead of x_1, x_2, x_3, x_4 , respectively.

The refined multi-variable polynomial $P_{\tilde{G}_D,c_1}(x_1, x_2, x_3, x_4)$ is defined analogously to the case of special alternating links; i.e., as the weighted sum of oriented spanning trees rooted at c_1 . Unfortunately, this polynomial has coefficients that are greater than 1, but it has a nice support:

Lemma 2.28. Let P and Q be the number of vertices in L and R, respectively. Note that $P = p_1 + \cdots + p_n$ and $Q = q_1 + \cdots + q_n$. Then

$$Supp(P_{\widetilde{G}_D,c_1}) =$$



FIGURE 43. An alternating 3-braid (left), its Crowell graph (middle), and the refined Crowell graph (right). The root vertex c_1 is denoted by a star.

 $\{ (a_1, a_2, a_3, a_4) \in \mathbb{Z}^4 : a_1, a_3, a_4 \ge 0, \ a_2 > 0, \ a_1 + a_3 = P - 1, \ a_2 + a_4 = Q \}.$

Proof. Consider the braid diagram shown in Figure 43, with the standard height function in the plane. Let c_1 be the topmost vertex of L. Any oriented spanning tree T rooted at c_1 contains a unique edge ending at each vertex other than c_1 . So T must contain at least one edge of type EL. The set of edges in T with weight x_1 or x_3 is precisely the set of edges ending in L. Hence, the number of edges in T with weight x_1 or x_3 is |L| - 1 = P - 1. The same argument shows that the number of edges in T with weight x_2 or x_4 is |R| = Q.

Hence, it suffices to construct a tree $T \in \mathcal{T}(\tilde{G}_D, c_1)$ with weight $W(T) = x_1^{a_1} x_2^{a_2} x_3^{a_3} x_4^{a_4}$ for any $(a_1, a_2, a_3, a_4) \in \mathbb{Z}^4$ with $a_1, a_3, a_4 \geq 0, a_2 > 0, a_1 + a_3 = P - 1$, and $a_2 + a_4 = Q$. Denote the topmost vertex of R by c'_1 . Consider the following subsets of the edge set $E(\tilde{G}_D)$:

 E_1 = the first a_1 edges of L starting at c_1 ,

 $E_2 =$ the topmost a_2 red edges,

 E_3 = the topmost a_3 black edges,

 E_4 = the first a_4 edges of R starting at c'_1 .

Let $E_T = E_1 \cup E_2 \cup E_3 \cup E_4$. We claim that the edges of E_T form the desired oriented rooted tree T.

Since $a_1 + a_3 = P - 1$ and $a_2 + a_4 = Q$, for any vertex c other than the root c_1 , there is exactly one edge in E_T leading to c. Indeed, the edges in E_1

lead to the lowest a_1 vertices of L, and the edges in E_3 lead to the topmost a_3 vertices of L, excluding c_1 . Similarly, edges in E_2 lead to the topmost a_2 vertices of R, and edges in E_4 lead to the lowest a_4 vertices of R. Since $a_2 > 0$, the edges E_T do not contain the cycle R. See the right of Figure 44 for the edges in E_T when $(a_1, a_2, a_3, a_4) = (3, 3, 4, 2)$, for the diagram in Figure 43.

By the above, for any vertex $v \neq c_1$, we can define the parent vertex p(v) of v such that $(p(v), v) \in E_T$; i.e., as the source of the unique edge in E_T leading to v. Note that p(v) is higher than v, unless possibly when $(p(v), v) \in E_1 \cup E_4$; i.e., when (p(v), v) is an edge in one of the cycles L and R. For some $k \in \mathbb{N}$, the sequence

$$(p^k(v),\ldots,p(v),v)$$

is a path from c_1 to v in E_T . So E_T is the edge set of an oriented rooted tree T with the desired weight.



FIGURE 44. An oriented spanning tree T of the refined Crowell graph shown in Figure 43 for $(a_1, a_2, a_3, a_4) = (3, 3, 4, 2)$.

There is a $P \times Q$ matrix $A = [A_{i,j}]$ for $(i, j) \in \{0, \dots, P-1\} \times \{0, \dots, Q-1\}$ such that

(17)
$$P_{\widetilde{G}_D,c_1} = \sum_{\substack{0 \le i \le P-1 \\ 0 \le j \le Q-1}} A_{i,j} \ x_1^{P-1-i} x_2^{Q-j} x_3^i x_4^j.$$

We can recover the Crowell polynomial by setting $x_1 = x_2 = t$ and $x_3 = x_4 = 1$; i.e.,

(18)
$$\Delta_L(-t) \doteq P_{G_D,c_1} = \sum_{0 \le k \le P+Q-2} \left(\sum_{i+j=k} A_{i,j} \right) t^{P+Q-1-k}.$$

The main result of this subsection are inequalities between coefficients $A_{i,j}$, which lead to inequalities between coefficients of the Alexander polynomial.

Theorem 2.29. Let D be the diagram of the closure L of the alternating 3-braid

$$\sigma_1^{p_1}\sigma_2^{-q_1}\cdots\sigma_1^{p_n}\sigma_2^{-q_n}$$

shown in Figure 43. Let $P = p_1 + \cdots + p_n$ and $Q = q_1 + \cdots + q_n$. If $A_{i,j}$ is the $P \times Q$ matrix of coefficients of $P_{\widetilde{G}_D,c_1}$, as above, then

$$nA_{i,j+1} \ge A_{i,j}$$

If

$$\Delta_L(-t) = \sum_{0 \le k \le P + Q - 2} a_k t^k,$$

then $a_{k+1} \ge a_k/n + A_{0,k+1} > a_k/n$ for $k \in \{0, \dots, P-2\}$.

Proof. Let $\mathcal{T}_{i,j}$ be the set of oriented rooted spanning trees of \widetilde{G}_D with weight $x_1^{P-1-i}x_2^{Q-j}x_3^ix_4^j$; i.e., ones contributing to the coefficient $A_{i,j}$. Equivalently, $\mathcal{T}_{i,j}$ is the set of maximal oriented rooted trees in \widetilde{G}_D with *i* black edges and *j* green edges.

By manipulating trees, we build a map $L: \mathcal{T}_{i,j} \to \mathcal{T}_{i,j+1}$ which is at most *n*-to-1. This implies that $nA_{i,j+1} \ge A_{i,j}$. The construction of the map L is as follows.

Recall that c_1 and c'_1 were used to denote the highest vertices of L and R, respectively. Let c_1, \ldots, c_P be the vertices of L, and c'_1, \ldots, c'_Q be the vertices of R, both labelled from top to bottom.

Let $T \in \mathcal{T}_{i,j}$ be an oriented, rooted spanning tree. We call c'_i a maximal vertex if the path in T from c_1 to c'_i only includes vertices in L, except for c'_i . Let the lowermost maximal vertex be c'_m . Consider the longest oriented path of green edges in T originating from c'_m . Note that the path might be of length zero. Assume that this path ends at c'_k and hence does not reach c'_{k-1} . Then there is a red edge in T leading to c'_{k-1} , which we denote by e. The map L is defined by

$$L(T) := T \setminus \{e\} \cup \{(c'_k, c'_{k-1})\}.$$

See Figure 45 for an example, where the maximal vertices are c'_1 and c'_7 , the lowest maximal vertex is $c'_m = c'_7$, and the green path goes from c'_7 to $c'_k = c'_6$.

 $c'_k = c'_6.$ We first show that $\operatorname{Im}(L) \subseteq \mathcal{T}_{i,j+1}$. Based on the construction of L(T), the weight $W(L(T)) = W(T)x_4x_2^{-1}$. Hence, we only need to show that it is an oriented rooted spanning tree. The rooted tree T induces a partial order $>_T$ on the vertices $V(\widetilde{G}_D)$ of \widetilde{G}_D . The relation $v >_T w$ holds if and only if there is a path from v to w in T. For any vertex v, the subset $T_v = \{x \in V(\widetilde{G}_D) : v \ge_T x\}$ forms an oriented subtree of T rooted at v. On the other hand, any partial order on $V(\widetilde{G}_D)$ with c_1 as its maximum also gives an oriented rooted spanning tree of \widetilde{G}_D . Note that the construction of L(T) can be summarized by setting c'_k to be the parent of c'_{k-1} . This operation results in an oriented rooted spanning tree if and only if $c'_k \notin T_{c'_{k-1}}$; see Figure 46. This condition holds for the operation in the construction of L(T). Indeed, if $c'_k \in T_{c'_{k-1}}$, then there is an oriented path from c'_{k-1} to c'_k , which contradicts the assumption that c'_m is a maximal vertex.



FIGURE 45. The map $L: \mathcal{T}_{i,j} \to \mathcal{T}_{i,j+1}$.

We now prove that L is at most *n*-to-1; i.e., we show that, for any $A \in \mathcal{L}_{i,j+1}$, we have $|L^{-1}(A)| \leq n$. To find the elements of $L^{-1}(A)$, first notice that the lowermost maximal vertices of T and L(T) are the same, since the construction of L(T) does not affect the path from c_1 to c'_m , and just turns c'_{k-1} from a possibly maximal vertex to a non-maximal one.

Using this fact, consider the lowermost maximal vertex of A and denote it by c'_m . Denote by J the largest path of green edges in A starting from c'_m . By construction of L, we know that, if $T \in L^{-1}(A)$, then A can be constructed by taking the longest path of green edges in T starting from c'_m and adding an edge at the end. The trees T and A only differ in two edges, one green and one red. Denote the one extra red edge of T by $e_T = (x, y)$. Based on the explanation above, y must be in J.

To construct T from A, we delete the green edge leading to y and replace it with e_T . As a result, x must be outside of J, otherwise the green edge from (y, x) in A is also in T and forms a cycle with e_T . This restricts the options for e_T : it is either one of the edges starting in L and ending in R, or it is an edge leading to the final leaf of J. So there are at most n options for e_T , hence $|L^{-1}(A)| \leq n$. See Figure 47 for an example. This completes our proof of the first inequality. The inequality on the Alexander polynomial coefficients follows from equation (18).



FIGURE 46. Two different examples of the *change of parent* operation in a rooted tree. In both examples, $c'_k \notin T_{c'_{k-1}}$ and the result of the operation is still an oriented rooted tree.



FIGURE 47. Construction of the inverse image $L^{-1}(A)$. The possible options for e_T come from the red edges leading to J from outside. One can see that using a red edge with both ends in J does not work.

We continue this subsection with some corollaries of Theorem 2.29. First, note that, by taking the mirror of the braid diagram and turning it upside

down, one can replace the roles of σ_1 and σ_2 without changing the Alexander polynomial.

Furthermore, one can improve the bound on the number of choices for e_T , and obtain the following strengthening of Theorem 2.29:

Corollary 2.30. Let D be the standard diagram of the closure L of the alternating 3-braid

 $\sigma_1^{p_1} \sigma_2^{-q_1} \cdots \sigma_1^{p_n} \sigma_2^{-q_n}.$ Let $P = p_1 + \cdots + p_n$ and $Q = q_1 + \cdots + q_n$. If $\Delta_L(-t) = \sum_{0 \le k \le P + Q - 2} a_k t^k$

and $[A_{i,j}]$ is the $P \times Q$ matrix of coefficients of $P_{\widetilde{G}_{D,C_1}}$, as above, then

$$a_{k+1} \ge \frac{a_k}{\min(n,k+1)} + A_{0,k+1} > \frac{a_k}{\min(n,k+1)}$$

for $k \in \{0, \dots, \max(P - 2, Q - 2)\}$.

Proof. The majority of the proof is the same as proof of Theorem 2.29. We only improve the upper bound on $|L^{-1}(A)|$. As we discussed, $|L^{-1}(A)|$ is the same as the number of possible choices for the edge e_T , which is a red edge leading to J. An upper bound on the number of these choices is the length of the path J. Each edge in J counts with weight 1 in the computation of W(T). Hence |J| < P + Q - 2 - pw(A), where $W(A) = t^{pw(A)}$.

One might ask if this argument can be generalized to alternating links other than alternating 3-braids. The main ingredient of the proof of Theorem 2.29 is the shape of the Crowell graph around the Seifert cycle R. In fact, this only relies on the Seifert cycle R being of type 1.

The local shape of the diagram around a Seifert cycle of type 1 is depicted in the top of Figure 48. The diagram consists of n twist regions along the boundary of the cycle. The orientations and weights of the Crowell graph are shown on bottom of Figure 48. The edges are colored analogously to a neighborhood of the Seifert cycle R in the Crowell graph of an alternating 3-braid. Using this coloring, the argument of the proof of Theorem 2.29 can be repeated, leading to Theorem 2.31.

Theorem 2.31. Let D be an alternating diagram of a link L. Assume that D contains a Seifert cycle C of type 1 such that there are n twist regions around the C with x_1, \ldots, x_n crossings, respectively. Let $c = x_1 + \cdots + x_n$ and $\Delta_L(-t) = \sum_{0 \le k} a_k t^k$. Then

 $a_{k+1} \ge a_k/n$

for $k \in \{0, \ldots, c-2\}$.

Recall that the construction of the refined Crowell graph for alternating 3-braids (Figure 43) was motivated by the result of Hafner, Mészáros, and

44



FIGURE 48. Top: neighborhood of a type 1 Seifert cycle in an alternating diagram. Bottom: the corresponding Crowell graph, together with a coloring of its edges.

Vidinas [HMV23]. One might try to repeat their argument for alternating 3-braids. Unfortunately, this does not work, as stated in the following Proposition.

Proposition 2.32. Let $P_{\tilde{G}_D,c_1}$ be the refined Crowell polynomial of the closure of an alternating 3-braid $B = \sigma_1^{p_1} \sigma_2^{-q_1} \cdots \sigma_1^{p_n} \sigma_2^{-q_n}$. Then $P_{\tilde{G}_D,c_1}$ is not denormalized Lorentzian unless the 3-braid is a connected sum of two torus links; i.e., $B = \sigma_1^{p_1} \sigma_2^{-q_1}$.

Proposition 2.32 follows from the results we prove in the rest of this section, more specifically Propositions 2.36 and 2.37. To explain these results, we first need to recall some of the basic definitions regarding Lorentzian polynomials.

Definition 2.33. A subset $J \subset \mathbb{N}^n$ is called *M*-convex if, for any $\alpha = (\alpha_1, \ldots, \alpha_n) \in J$ and $\beta = (\beta_1, \ldots, \beta_n) \in J$ such that $\alpha_i > \beta_i$ for some $i \in \{1, \ldots, n\}$, there exist $j \in \{1, \ldots, n\}$ satisfying

$$\alpha_i < \beta_i, \ \alpha - e_i + e_i \in J, \text{ and } \beta - e_i + e_i \in J,$$

where e_i and e_j are the *i*-th and *j*-th standard basis vectors of \mathbb{R}^n , respectively.

Let H_n^d be the space of degree d homogeneous polynomials with real coefficients in the variables x_1, \ldots, x_n . The Hessian of a homogeneous quadratic polynomial $f \in H_n^2$ is the symmetric matrix $H = (H_{ij})_{i,j \in [n]}$, where $H_{ij} = \frac{\partial}{\partial x_i} \frac{\partial}{\partial x_j} f$ and $[n] = \{1, \ldots, n\}$. The set L_n^d of Lorentzian polynomials of degree d in n variables is defined as follows. Let $L_n^1 \subseteq H_n^1$ be the set of all linear polynomials with nonnegative coefficients. Let $L_n^2 \subseteq H_n^2$ be the subset of quadratic polynomials with nonnegative coefficients and M-convex support whose Hessians have at most one positive eigenvalue. For d > 2, define $L_n^d \subseteq H_n^d$ as the subset of polynomials with nonnegative coefficients and M-convex support that satisfy

$$\frac{\partial}{\partial x_{i_1}}\frac{\partial}{\partial x_{i_2}}\cdots\frac{\partial}{\partial x_{i_{d-2}}}f\in L^2_n \text{ for all } i_1,i_2,\ldots,i_{d-2}\in[n].$$

The normalization operator N on $\mathbb{R}[x_1, \ldots, x_n]$ is defined by linear extension of the map

$$N(x^{\alpha}) = \frac{x^{\alpha}}{\alpha!},$$

where $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n, x^{\alpha} := x_1^{\alpha_1} \cdots x_n^{\alpha_n}$, and $\alpha! := \prod_{1 \le i \le n} \alpha_i!$.

Definition 2.34. A polynomial $f \in H_n^d$ is called *denormalized Lorentzian* if $N(f) \in L_n^d$.

Note that

$$\frac{\partial^{\alpha_1}}{\partial x_1^{\alpha_1}} \cdots \frac{\partial^{\alpha_i - 1}}{\partial x_i^{\alpha_i - 1}} \cdots \frac{\partial^{\alpha_j - 1}}{\partial x_j^{\alpha_j - 1}} \cdots \frac{\partial^{\alpha_n}}{\partial x_n^{\alpha_n}} \left(\frac{x^{\alpha}}{\alpha!}\right) = x_i x_j.$$

Hence, the denormalized Lorentzian property can be reformulated as follows. Let $\beta = (\beta_1, \ldots, \beta_n) \in \mathbb{N}^n$ such that $\beta_1 + \cdots + \beta_n = d - 2$. For $f = \sum_{\alpha} a_{\alpha} x^{\alpha} \in$

$$H_n^d$$
, define $f_\beta := \sum_{\alpha \ge \beta} a_\alpha x^\alpha$. Then
$$\frac{\partial^\beta}{\partial x^\beta} N(f) = x^{-\beta} f_\beta = \sum_{\alpha \ge \beta} a_\alpha x^{\alpha - \beta}.$$

So f is denormalized Lorentzian if and only if $x^{-\beta}f_{\beta} \in L_n^2$ for any $\beta = (\beta_1, \ldots, \beta_n) \in \mathbb{N}^n$ such that $\beta_1 + \cdots + \beta_n = d - 2$. This means that f has nonnegative coefficients and M-convex support, and the Hessian of $x^{-\beta}f_{\beta}$ has at most one positive eigenvalue for any β as above.

Now let us consider a polynomial of the form

$$f = \sum_{\substack{0 \le i \le P-1 \\ 0 \le j \le Q-1}} A_{i,j} \ x_1^{P-1-i} x_2^{Q-j} x_3^i x_4^j.$$

The possible choices for β for which $x^{-\beta}f_{\beta}$ is non-zero and quadratic fall into the following three cases.

Case I. We have

$$\beta = (P - 2 - i, Q - 1 - j, i, j)$$

for $i \in \{0, ..., P-2\}$ and $j \in \{0, ..., Q-2\}$. Then

$$x^{-\beta}f_{\beta} = A_{i,j}x_1x_2 + A_{i+1,j}x_2x_3 + A_{i,j+1}x_1x_4 + A_{i+1,j+1}x_3x_4,$$

46

and the Hessian is

(19)
$$\begin{bmatrix} 0 & A_{i,j} & 0 & A_{i,j+1} \\ A_{i,j} & 0 & A_{i+1,j} & 0 \\ 0 & A_{i+1,j} & 0 & A_{i+1,j+1} \\ A_{i,j+1} & 0 & A_{i+1,j+1} & 0 \end{bmatrix}$$

Case II. We have

$$\beta = (0, Q - 2 - j, P - 1, j)$$

for $j \in \{0, \dots, Q-2\}$. Then $x^{-\beta}f_{\beta} = A_{P-1,j}x_2^2 + A_{P-1,j+1}x_2x_4 + A_{P-1,j+2}x_4^2$,

and the Hessian is

(20)
$$\begin{bmatrix} A_{P-1,j} & A_{P-1,j+1} \\ A_{P-1,j+1} & A_{P-1,j+2} \end{bmatrix}$$

Case III. We have

$$\beta = (P - 3 - i, 1, i, Q - 1)$$

for $i \in \{0, \dots, P-3\}$. Then $x^{-\beta} f_{\beta} = A_{i,Q-1} x_1^2 + A_{i+1,Q-1} x_1 x_3 + A_{i+2,Q-1} x_3^2$,

and the Hessian is

(21)
$$\begin{bmatrix} A_{i,Q-1} & A_{i+1,Q-1} \\ A_{i+1,Q-1} & A_{i+2,Q-1} \end{bmatrix}$$

Based on this computation, we can derive a necessary and sufficient condition for f to be denormalized Lorentzian. In the following lemma, we find the signs of the eigenvalues of the matrices occurring as Hessians of $x^{-\beta}f_{\beta}$. The condition is then stated in Proposition 2.36.

Lemma 2.35. For $a, b, c, d \in \mathbb{N}$, let

$$A = \begin{bmatrix} 0 & a & 0 & d \\ a & 0 & b & 0 \\ 0 & b & 0 & c \\ d & 0 & c & 0 \end{bmatrix} \quad and \quad B = \begin{bmatrix} a & b \\ b & c \end{bmatrix}.$$

The matrix B has at most one positive eigenvalue if and only if $b^2 \ge ac$. The matrix A has at most one positive eigenvalue if and only if ac = bd.

Proof. The characteristic polynomial of B is $x^2 - (a + c)x + (ac - b^2)$, so a + c is the sum and $ac - b^2$ is the product of the roots. As $a + c \ge 0$, at least one of the roots is nonnegative. So at most one of the roots is positive if and only if $ac - b^2 \le 0$.

The characteristic polynomial of A is

$$\chi_A = x^4 - (a^2 + b^2 + c^2 + d^2)x^2 + (a^2c^2 + b^2d^2 - 2abcd)$$

The discriminant of this polynomial with respect to x^2 is

$$\Delta_{\chi_A} = (a^2 + b^2 + c^2 + d^2)^2 - 4a^2c^2 - 4b^2d^2 + 8abcd \ge 0.$$

So there exist real numbers r_1 and r_2 such that

$$\chi_A = (x^2 - r_1)(x^2 - r_2),$$

$$r_1 + r_2 = a^2 + b^2 + c^2 + d^2 \ge 0, \text{ and}$$

$$r_1 r_2 = a^2 c^2 + b^2 d^2 - 2abcd \ge 0.$$

Hence $r_1, r_2 \ge 0$. If both of them are positive, then A has exactly two positive eigenvalues. If one of them is zero, then $0 = r_1r_2 = a^2c^2 + b^2d^2 - 2abcd$, so ac = bd. The result follows.

Proposition 2.36. For a $P \times Q$ matrix $A = (A_{ij})$ with positive entries, the polynomial

$$f = \sum_{\substack{0 \le i \le P-1 \\ 0 \le j \le Q-1}} A_{i,j} \ x_1^{P-1-i} x_2^{Q-j} x_3^i x_4^j$$

is denormalized Lorentzian if and only if there exist sequences (z_0, \ldots, z_{P-1}) and (y_0, \ldots, y_{Q-1}) with positive entries such that:

- (1) $A_{i,j} = z_i y_j$,
- (2) $(z_i)_{0 \le i \le P-1}$ is log-concave; i.e., $z_i^2 \ge z_{i-1}z_{i+1}$ for $i \in \{1, \dots, P-2\}$, and
- (3) $(y_j)_{0 \le j \le Q-1}$ is log-concave; i.e., $y_j^2 \ge y_{j-1}y_{j+1}$ for $j \in \{1, \dots, Q-2\}$.

Proof. Note that since the matrix A has positive entries, we have

$$Supp(f) = \{ (P - 1 - i, Q - j, i, j) \in \mathbb{Z}^4 : 0 \le i \le P - 1, \ 0 \le j \le Q - 1 \}.$$

Hence, f has M-convex support. In fact, Supp(f) is exactly the integer points inside a $P \times Q$ rectangle embedded in a 2-dimensional hyperplane in \mathbb{R}^4 .

We now look at the Hessians. Using the criteria of Lemma 2.35 on Hessians of case I, which have form (19), leads to

$$A_{i,j}A_{i+1,j+1} = A_{i+1,j}A_{i,j+1}$$

for $i \in \{0, \ldots, P-2\}$ and $j \in \{0, \ldots, Q-2\}$. This means that all the rows of A are multiples of a vector $\mathbf{y} = (y_0, \ldots, y_{Q-1})$, and all the columns of A are multiples of a vector $\mathbf{z} = (z_0, \ldots, z_{P-1})$. Hence $A_{i,j} = z_i y_j$, possibly after scaling \mathbf{y} or \mathbf{z} .

Using the criteria of Lemma 2.35 on Hessians of case II, which have form (20), and case III, which have form (21), leads to log-concavity of the last row and column of A. This implies that \mathbf{y} and \mathbf{z} are log-concave. \Box

The criteria of Proposition 2.36 are too restrictive. For the refined Crowell polynomial of the closure of an alternating 3-braids $B = \sigma_1^{p_1} \sigma_2^{-q_1} \cdots \sigma_1^{p_n} \sigma_2^{-q_n}$ with coefficient matrix $A = [A_{i,j}]$, a calculation of extremal entries proves that these criteria only hold when n = 1, which means that the closure of the alternating 3-braid is $T_{2,p_1} \# T_{2,-q_1}$. This is an immediate corollary of the following:

Proposition 2.37. The external entries of the matrix $A = [A_{i,j}]$ of coefficients of the refined Crowell polynomial of the closure of an alternating 3-braid $B = \sigma_1^{p_1} \sigma_2^{-q_1} \cdots \sigma_1^{p_n} \sigma_2^{-q_n}$ satisfy the following:

- $A_{i,0} = 1$ for $i \in \{0, \dots, P-1\}$,
- $A_{P-1,j} = 1$ for $j \in \{0, \ldots, Q-1\}$, and
- $A_{0,j} \ge n \text{ for } j \in \{1, \dots, Q-1\}.$

In other words,

$$A = \begin{bmatrix} 1 & \ge n & \cdots & \ge n \\ 1 & . & \cdots & . \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & \cdots & 1 \end{bmatrix}$$

Proof. One can characterize the oriented rooted spanning trees contributing to each term and then count them. We will show that the families $\mathcal{T}_{i,0}$ and $\mathcal{T}_{P-1,i}$ only contain one tree each; see Figure 49.

A tree $T \in \mathcal{T}_{i,0}$ contains all the red edges and no green edge. Let $J = (c_i, c_{i-1}, \ldots, c_{i-t})$ be a maximal path of blue edges in T. Then either $c_i = c_1$, or the parent of vertex c_i in T must be a vertex in R. This leads to a contradiction, as edges of J, red edges, and the black edge leading to c_i form a cycle. Hence, all the blue edges form a path of length P - 1 - i starting at c_1 . The black edges of T are uniquely determined, as they form a disjoint union of paths reaching the vertices in $L \setminus J$.

A tree $T \in \mathcal{T}_{P-1,j}$ contains all the black edges and no blue edge. Let J be the maximal path in T starting at c_1 . This path must go outside L, hence it contains the red edge leading to c'_1 . Deleting this part of the tree, one can think of c'_1 as the root of the remaining subtree, and we repeat the argument of the previous paragraph.

The trees in the family $\mathcal{T}_{0,j}$ contain all the blue edges and no black ones. So the restriction of T to the cycle L is just a path of length P-1. One can then pick one of the n red edges connecting L to R. Assume this edge leads to c'_i . Add an oriented path J consisting of j green edges starting at c'_i . Then one can build an element of $\mathcal{T}_{0,j}$ by adding a disjoint union of paths of red edges leading to vertices in $R \setminus J$. See Figure 50 for an example. \Box

We conclude this subsection with some final remarks. As we discussed, a direct generalization of the multi-variable refinement idea does not work for 3-braids. One can try other weight refinements of the Crowell graph, but even for simple alternating 3-braids the support of the resulting polynomial fails to be M-convex. Another idea is to try to decompose this multi-variable polynomial into denormalized Lorentzian parts; i.e., decompose the $P \times Q$ matrix A into matrices of the form of Proposition 2.36. Finding a general scheme for such a decomposition seems improbable based on small examples. Equations 22 and 23 show a few examples.

In special cases, one can find certain log-concavity properties of the coefficient matrix A. As mentioned before, for alternating knots, being special



FIGURE 49. The trees in $\mathcal{T}_{i,0}$ and $\mathcal{T}_{P-1,j}$.



FIGURE 50. This shows n trees in $\mathcal{T}_{0,j}$.

is equivalent to one of the Tait graphs being bipartite. For alternating links, this is not true. One can change the orientation of some of the components of the link without changing the Tait graph. We call alternating links with a bipartite Tait graph almost special alternating links. The Crowell graph of an almost special alternating link only differs from the Crowell graph of the corresponding special alternating link in some of its weights. Alternating 3-braids $\sigma_1^{p_1}\sigma_2^{-q_1}\cdots\sigma_1^{p_n}\sigma_2^{-q_n}$, where all p_i (or q_i) are even for $i \in \{1, \ldots, n\}$, are examples of such braids. See Figure 51 when $B = \sigma_1^{-2}\sigma_2\sigma_1^{-2}\sigma_2$.

(22)
$$B = \sigma_1 \sigma_2^{-1} \sigma_1 \sigma_2^{-1} \sigma_1 \sigma_2^{-1}, \quad A = \begin{bmatrix} 1 & 3 & 3 \\ 1 & 2 & 3 \\ 1 & 1 & 1 \end{bmatrix}$$

(23)
$$B = \sigma_1^2 \sigma_2^{-2} \sigma_1^2 \sigma_2^{-2} \sigma_1^3 \sigma_2^{-1}, \quad A = \begin{bmatrix} 1 & 3 & 5 & 5 & 3 \\ 1 & 5 & 10 & 11 & 6 \\ 1 & 5 & 11 & 14 & 8 \\ 1 & 4 & 9 & 13 & 8 \\ 1 & 3 & 6 & 9 & 6 \\ 1 & 2 & 3 & 4 & 3 \\ 1 & 1 & 1 & 1 & 1 \end{bmatrix}.$$

Small examples of the refined Crowell polynomial.



FIGURE 51. The diagram D is almost special alternating. The special alternating diagram D' is obtained by changing the orientation of one of the components. The two figures on the right compare their Crowell graphs.

Looking at the example of Figure 51, one notices that, by identifying some of the variables, the refined Crowell polynomials $P_{\tilde{G}_D}$ and $P_{\tilde{G}_{D'}}$ become equal. More precisely, the set of edges with weight x_1, x_2, x_3 in \tilde{G}_D coincide with edges with weight x'_1, \ldots, x'_4 in $\tilde{G}_{D'}$. Hence, we have

$$P_{\widetilde{G}_D}(X, X, X, Y) = P_{\widetilde{G}_{D'}}(X, X, X, X, Y).$$

Based on the work of Hafner, Mészáros, and Vidinas [HMV23], the polynomial $P_{\tilde{G}_{D'}}$ is denormalized Lorentzian. Identifying variables preserves the Lorentzian property by [HMV23, Lemma 2.5], so $P_{\tilde{G}_D}(X, X, X, Y)$ is also Lorentzian. Based on what we know about the general form of $P_{\tilde{G}_D}$ (equation (17)), we can deduce that the sequence of non-zero coefficients of

 $P_{\widetilde{G}_D}(X, X, X, Y)$ is $\sum_j A_{ij}$ for $j \in \{0, \ldots, P-1\}$; i.e., the sum of the rows of the matrix A. We can conclude that this sequence is log-concave:

Theorem 2.38. Let D be the diagram of the closure of an alternating 3-braid word

$$B = \sigma_1^{p_1} \sigma_2^{-q_1} \cdots \sigma_1^{p_n} \sigma_2^{-q_n}.$$

Let $A = (A_{i,j})$ be the $P \times Q$ matrix of coefficients of $P_{\tilde{G}_D,c_1}$. If all p_i are even, then the sum of the rows of A form a log-concave sequence. If all q_i are even, then the sum of the columns of A form a log-concave sequence.

3. A CONJECTURE OF HIRASAWA AND MURASUGI

3.1. Introduction. In 2013, Hirasawa and Murasugi proposed an extension of the trapezoidal conjecture. They conjectured that the signature gives an upper bound on the length of the stable part of the sequence of absolute values of coefficients of the Alexander polynomial of an alternating knot. In 2017, Chen [Che17] generalized their conjecture to alternating links.

Conjecture 1.2. Let *L* be an alternating link and $\Delta_L(t) = \sum_{i=0}^{l-1} (-1)^i a_i t^i$ be its Alexander polynomial. Then there is $i_0 \in \{1, \ldots, l/2 + 1\}$ such that

$$a_0 < \dots < a_{i_0-1} = a_{i_0} = \dots = a_{l-i_0} > \dots > a_{l-1}.$$

Moreover,

52

$$\left\lfloor \frac{|\sigma(L)|+1}{2} \right\rfloor \geq \left\lfloor \frac{l-2(i_0-1)}{2} \right\rfloor.$$

We call this the *H-M inequality*, and the alternating links for which equality holds *H-M sharp*. We define the *stable length of* L to be

$$sl(L) := l - 2(i_0 - 1).$$

This conjecture originates from the study of the roots of the Alexander polynomial of alternating links, which has been the subject of numerous papers, most notably regarding questions related to Hoste's conjecture [LM12]. Hirasawa and Murasugi [HM14] introduced the class of *stable alternating links*, which consists of all alternating links L for which Δ_L has only real roots. Lemma 3.1 implies that Conjecture 1.2 holds for this class of links.

Lemma 3.1 ([HM14]). Let L be a stable alternating link and

$$\Delta_L(t) = \sum_{i=0}^{l-1} (-1)^i a_i t^i.$$

Then $(a_0, a_1, \dots, a_{l-1})$ is a strictly log-concave sequence, and hence it is unimodal; i.e., it is trapezoidal with stable length one. Furthermore, the signature of L is zero.

Chen [Che17] showed that Conjecture 1.2 holds for 2-bridge links, and Alrefai and Chbili [AC21] proved it for closures of alternating 3-braids of length less than four. There also has been some effort to construct families of stable alternating links by plumbing Hopf bands to a disk; see Hirasawa and Murasugi [HM14] and Stoimenow [Sto21].

Note that the H-M inequality is trivial for special alternating links. This comes from the combinatorial formula for the signature of alternating links stated in Lemma 3.2, below. Using Lemma 3.2, we will establish Lemma 3.3 about the signature of special alternating links. Recall that we only consider positive special diagrams, unless otherwise stated.

Lemma 3.2 ([Tra04],[OS03]). Let D be a reduced alternating diagram, c a crossing in D, and D_0 the diagram obtained by taking the oriented smoothing of c. Then

$$\sigma(D_0) = \sigma(D) + \operatorname{sign}(c).$$

Hence

 $\sigma(D) = \#(black \ regions) - \#(positive \ crossings) - 1,$

where the black-white coloring convention for alternating links is as in Figure 52.



FIGURE 52. Black-white coloring convention for regions of an alternating diagram, following Ozsváth and Szabó [OS03].

Lemma 3.3. For a special alternating link L, we have

$$-\sigma(L) = 2g(L) = \deg(\Delta_L).$$

Furthermore, if an alternating link L satisfies this condition, then it is special alternating (up to taking its mirror).

Proof. Suppose that L is a positive special alternating link. Using the second formula of Lemma 3.2, we can compute the signature of L based on its black Tait graph B. The black regions correspond to the vertices of B, and the crossings correspond to the edges. Hence, we have

$$\sigma(L) = |V(B)| - |E(B)| - 1 = -\operatorname{rank}(H_1(B)) = -2g(L).$$

The last equality holds because the black Tait surface, constructed by median construction on B, is a minimal genus Seifert surface.

To prove the converse, we use Proposition 2.18 to decompose L into the diagrammatic Murasugi sum of special alternating components such that genus and signature are additive under this decomposition. The absolute

value of the signature is sub-additive under Murasugi decomposition, so the equality $2g(\Delta_L) = |\sigma(L)|$ holds only when the signatures of all the special alternating components of L have the same sign. Hence, all of these components must be positive (resp. negative) alternating links, which means that L is a positive (resp. negative) alternating link. Consequently, L or its mirror is a special alternating link.

In the rest of this subsection, we prove some new results about the H-M inequality based on our work in Section 2. In subsection 3.2, we focus on the study of H-M sharp knots. In Subsection 3.3, we connect the H-M inequality with concordance. We are going to use the results of Subsection 2.2, so we now summarize these.

Proposition 3.4. Let n be the number of crossings in the largest coherent twist region of L, denoted by R. Let L_i be the alternating link obtained by changing R to a twist region containing i crossings. The polynomials P_1 and P'_2 are Crowell polynomials of the graphs depicted in Figure 53.



FIGURE 53. The Crowell graphs of equations (24) and (25).

If n is odd, then

(24)
$$\Delta_L(-t) = P_n = t^{n-1}P_1 + (1 + \dots + t^{n-2})P'_2.$$

Furthermore, the first n-2 terms of P_n come from only $(1 + \dots + t^{n-2})P'_2$. Similarly, if n is even, then

(25)
$$\Delta_L(-t) = P_n = t^{n-2}P_2 + (1 + \dots + t^{n-3})P_2'$$

and the first n-3 terms of P_n come from only $(1+\cdots+t^{n-3})P'_2$.

Using Proposition 3.4, we now establish the first result of this subsection.

Theorem 3.5. Let L be an alternating link and

$$\Delta_L(T) \doteq \sum_{i=0}^{2g(L)+|L|-1} (-1)^i a_i T^i$$

If $MT(L) - 3 \ge g(L) + |L|/2$, then (a_i) is trapezoidal and $sl(L) = 2g(L) + |L| - 2(\deg(P'_2) - 1),$

where P'_2 is the Crowell polynomial of the graph depicted in Figure 53.

Proof. Let n = MT(L), and assume that n is odd. The trapezoidality of the sequence (a_i) follows from Theorem 2.6. Let $c_0, \ldots, c_{n-2+\deg(P'_2)}$ denote the coefficients of $(1 + \cdots + t^{n-2})P'_2$. Then $(c_i)_{1 \le i \le n-2}$ are cumulative sums of coefficients of P'_2 . Hence, if $n-2 > \deg(P'_2)$, then

$$c_{\deg(P_2')} = \cdots = c_{n-2},$$

and they are all equal to the sum of the coefficients of P'_2 . Using Proposition 3.4, we know that, for $i \in \{0, \ldots, n-2\}$, we have $a_i = c_i$. The symmetry of the sequence (a_i) implies the result. The argument is analogous for n even.

In Subsection 2.2, we used the shape of twist regions in the Seifert graph to prove that $g(L_n) = g(L_1) + \frac{n-1}{2}$; see Figure 14. Using this, one can rewrite the conclusion of Theorem 3.5 as

(26)
$$\operatorname{sl}(L_n) = 2(g(L_1)) + n - 1 + |L_n| - 2(\operatorname{deg}(P_2') - 1).$$

We can use this reformulation and prove Corollary 3.6. Recall that we call an alternating link satisfying MT(L) - 3 > g(L) + |L|/2 a twist-concentrated alternating link.

Corollary 3.6. Using the notation of Proposition 3.4, if L_n is twist-concentrated, then the H-M inequality for L_n follows from the H-M inequality for L_{n-2} .

Proof. Based on equation (26), we have

$$\operatorname{sl}(L_n) = \operatorname{sl}(L_{n-2}) + 2.$$

Without loss of generality, assume that the twisting region consists of negative crossings (as in Figure 8). Using Lemma 3.2, we have

$$\sigma(L_n) = \sigma(L_{n-2}) + 2$$

The combination of these two equalities gives us the conclusion.

Note that, using equation (26), one only needs to compute $\deg(P'_2)$ to obtain an explicit formula for the stable length of alternating twist-concentrated links. One example is twist-concentrated alternating 3-braids, for which the computation is done in Proposition 3.7.

Proposition 3.7. Let $B = \sigma_1^{p_1} \sigma_2^{-q_1} \cdots \sigma_1^{p_n} \sigma_2^{-q_n}$, and let *L* be the corresponding alternating 3-braid. Assume that P'_2 is defined as in Proposition 3.4. Then we have

$$\deg(P_2') = \deg(P_1) - 1 = 2g(L_1) + |L_1| - 2.$$

Proof. Twist regions of L are in one-to-one correspondence with powers of σ_1 and σ_2 in B. Without loss of generality, assume that

$$p_n = \max\{p_1, \cdots, p_n, q_1, \cdots, q_n\}.$$

We use the same notations as in Subsection 2.2. Let $N = |V(G_1)| = |V(G'_2)|$.

We identify rooted spanning trees with the highest weight in G_1 and G'_2 . First, note that edges with weight t, excluding the one that leads to the root v, form a spanning tree $T \in \mathcal{T}(G_1, v)$. As a result, $\deg(P_1) = N - 1$. A similar argument gives that $\operatorname{mindeg}(P_1) = 1$, hence

$$\deg(P_1) = \deg(\Delta_{L_1}) = 2g(L_1) + |L_1| - 2.$$

Note that we showed the existence of these two trees in Lemma 2.28 as well.

Now we compute $\deg(P'_2)$. We are going to show that there is a spanning tree $T' \in \mathcal{T}(G'_2, v)$ with only one edge with weight 1. Let $e_{ou} = (v, v_{ou})$ (resp. $e_{od} = (v, v_{od})$) be the upper (resp. lower) outgoing edge of the root vin G_1 . Also let $e_{iu} = (v_{iu}, v)$ be the upper incoming edge of the root vin G_1 . To construct G'_2 , we deltet the edges e_{ou} and e_{iu} and replace them with an edge $e_{io} = (v_{iu}, v_{ou})$. We are going to construct T' from T using the following modifications. We delete the edge leading to v_{iu} in T and replace it with (v_{od}, v_{iu}) , and then replace e_{ou} with e_{io} . Note that, following this construction, (v_{od}, v_{iu}) will be the only edge with weight 1 in T'. See Figure 54 for an illustration of the proof.

Using Proposition 3.7 and Lemma 3.8, we can deduce the H-M inequality as stated in Theorem 3.9, below.

Lemma 3.8. ([AC21],[Erl99]) Let $B = \sigma_1^{p_1} \sigma_2^{-q_1} \cdots \sigma_1^{p_n} \sigma_2^{-q_n}$ be an alternating 3-braid, and let L_B be its closure. Then

$$\sigma(L_B) = \sum p_i - \sum q_j.$$

Theorem 3.9. Let $B = \sigma_1^{p_1} \sigma_2^{-q_1} \cdots \sigma_1^{p_n} \sigma_2^{-q_n}$ be an alternating 3-braid, and let L_B by its closure. Assume that

$$p_1 \ge \sum_{i\ge 2} p_i + \sum q_j + 6.$$

Then the Hirawasa–Murasugi conjecture holds for L_B .

If Fox's trapezoidal conjecture is true, one can replace the assumption that $MT(L) - 3 \ge g(L) + |L|/2$ in Theorem 3.5 with

$$\mathrm{MT}(L) - 2 > \mathrm{deg}(P_2').$$

We now use the results of Subsection 2.3 to study the stable length and the Hirasawa–Murasugi inequality.

Lemma 3.10. Let K_1 and K_2 be alternating links such that the trapezoidal conjecture holds for both. Assume that $\deg(\Delta_{K_1}) \geq \deg(\Delta_{K_2})$. Then

$$sl(K_1 \# K_2) = \min\{sl(K_1) - \deg(\Delta_{K_2}), 0\}$$

In particular, if $sl(K_1 \# K_2) > 0$, then $sl(K_1) = sl(K_1 \# K_2) + deg(\Delta_{K_2})$.

Proof. The sequence of coefficients of $\Delta_{K_1 \# K_2} = \Delta(K_1) \Delta(K_2)$ is the convolution of the coefficient sequences of $\Delta(K_1)$ and $\Delta(K_2)$. This convolution alongside a sketch of the proof appears in Figure 55.



FIGURE 54. The rooted spanning trees with the highest degree weights in P_1 and P'_2 for an alternating 3-braid.

The trapezoidal inequalities between coefficients become equality only when the shorter sequence falls in the stable part of the longer sequence (blue in Figure 55). As soon as the shorter sequence moves out of this stable part, the associated coefficient drops (red in Figure 55).

One can derive this explicitly by expanding the terms and using the trapezoidal condition, but we do not include the computation here, which is a generalization of the rearrangement inequality. \Box



FIGURE 55. Convolution of two trapezoidal sequences. The top figure is a graph of the sequence of coefficients of Δ_{K_1} . The bottom figure shows different copies of the sequence of coefficients of Δ_{K_2} used for computing the convolution at different points.

A simple corollary of Lemma 3.10 is that one only needs to prove the H-M inequality for prime alternating links.

Corollary 3.11. If the H-M inequality holds for prime alternating links, then it also holds for arbitrary alternating links.

Proof. Assume that the H-M inequality fails for $K_1 \# K_2$. This means that

$$sl(K_1 \# K_2) > |\sigma(K_1 \# K_2)| + 1 = |\sigma(K_1) + \sigma(K_2)| + 1.$$

Using Lemma 3.10, we have

 $sl(K_1) > |\sigma(K_1) + \sigma(K_2)| + deg(\Delta_{K_2}) + 1 \ge |\sigma(K_1)| - |\sigma(K_2)| + deg(\Delta_{K_2}) + 1$, so $sl(K_1) > |\sigma(K_1)| + 1$. Hence, if there exists a non-prime counterexample to the H-M inequality, then there exists a prime counterexample. \Box

One can extend this result to diagrammatic plumbings, and, even more generally, to Murasugi sums of length less than 3:

Corollary 3.12. The H-M inequality holds for any link $L = L_1 * \cdots * L_n$ which is a diagrammatic Murasugi sum of special alternating links L_1, \ldots, L_n , such that the length of all Murasugi sums are less than 3.

Proof. Let $L'_1 = L_1$ and $L'_2 = L_2 * \cdots * L_n$. Then $L = L'_1 * L'_2$, and the length of this Murasugi sum is less than 3. By Theorem 2.23, the trapezoidal conjecture holds for L, L'_1 , and L'_2 .

By Proposition 2.21, for a plumbing, we can write

(27)
$$\tilde{\Delta}_{L_1'*L_2'} = \tilde{\Delta}_{L_1'}\tilde{\Delta}_{L_2'} + \tilde{\Delta}_{\tilde{L}_1'}\tilde{\Delta}_{\tilde{L}_2'}.$$

Since the Alexander polynomials are trapezoidal, one can deduce that

$$\mathrm{sl}(L'_1 * L'_2) = \min\left\{\mathrm{sl}(\Delta_{L'_1} \times \Delta_{L'_2}), \mathrm{sl}\left(\Delta_{\widetilde{L}'_1} \Delta_{\widetilde{L}'_2}\right)\right\}.$$

Recall that signature is additive under diagrammatic Murasugi sum (Proposition 2.18). The result now follows analogously to the proof of Corollary 3.11. The argument for other Murasugi sums of length less than 3 is similar. $\hfill \Box$

3.2. **H-M sharp links.** In this subsection, we study the family of H-M sharp alternating links with non-zero stable lengths. Recall that we call an alternating link L H-M sharp when

$$\left\lfloor \frac{|\sigma(L)|+1}{2} \right\rfloor = \left\lfloor \frac{\mathrm{sl}(L)}{2} \right\rfloor.$$

We start by looking at the results of Subsection 3.1 and analyze when equality holds.

Lemma 3.13. Let K_1 and K_2 be alternating knots satisfying the Hirawasa– Murasugi conjecture, and assume $\deg(\Delta_{K_1}) \geq \deg(\Delta_{K_2})$. The connected sum $K_1 \# K_2$ is H-M sharp if and only if, up to mirroring,

- (1) $sl(K_1) = \sigma(K_1) + 1$,
- (2) $\sigma(K_2) + \deg(\Delta_{K_2}) = 0$, and
- (3) $sl(K_1) \ge \deg(\Delta_{K_2}).$

Furthermore, K_1 is H-M sharp with positive signature and K_2 is special alternating.

Proof. Using Lemma 3.10, we have

$$sl(K_1 \# K_2) = |\sigma(K_1) + \sigma(K_2)| + 1.$$

Hence

 $sl(K_1) \ge |\sigma(K_1) + \sigma(K_2)| + deg(\Delta_{K_2}) + 1 \ge |\sigma(K_1)| - |\sigma(K_2)| + deg(\Delta_{K_2}) + 1,$

which implies that

$$sl(K_1) \ge (|\sigma(K_1)| + 1) + (deg(\Delta_{K_2}) - |\sigma(K_2)|).$$

The assumption that the H-M inequality holds for K_1 gives us the three conditions. The second part of the statement follows from Lemma 3.3.

Repeated application of Lemma 3.13 implies the following:

Corollary 3.14. Assume that the Hirawasa–Murasugi conjecture is true for K_1, \ldots, K_n . Let K be an H-M sharp knot with prime decomposition $K = K_1 \# \cdots \# K_n$, such that $\deg(\Delta_{K_1}) \ge \deg(\Delta_{K_i})$ for $i \ge 2$. Then the following hold up to taking the mirror of K:

- (1) K_1 is H-M sharp with $\sigma(K_1) > 0$,
- (2) K_2, \ldots, K_n are all special alternating knots,

(3)
$$sl(K_1) \ge \sum_{i=2}^n deg(\Delta_{K_i})$$

This characterization is necessary and sufficient.

Similarly to Corollary 3.12, we can generalize the argument of Corollary 3.14 to diagrammatic plumbings $K = K_1 * K_2$, and, even more generally, to diagrammatic Murasugi sums of length less than 3. For this, we need to assume that the trapezoidal conjecture holds for the links \tilde{K}_1 and \tilde{K}_2 , as well as for K_1 and K_2 . Deriving necessary and sufficient criteria in this case is a bit more complicated as we need additional conditions on \tilde{K}_1 and \tilde{K}_2 . Due to Theorem 2.23, the main cases of interest are diagrammatic Murasugi sums of special alternating knots (in particular, Murasugi sums of a pair), when the lengths of all sums are less than 3.

Lemma 3.15. Let K_1 and K_2 be alternating knots satisfying the Hirawasa– Murasugi conjecture, and assume $\deg(\Delta_{K_1}) \geq \deg(\Delta_{K_2})$. Let $K_1 * K_2$ be a diagrammatic Murasugi sum of length less than 3. If the link $K_1 * K_2$ is H-M sharp, then K_1 is H-M sharp with positive signature and K_2 is special alternating.

Repeated application of Lemma 3.15 implies Corollary 3.16:

Corollary 3.16. Let K be an H-M sharp alternating knot with Murasugi decomposition $K = K_1 * \cdots * K_n$, such that the lengths of all Murasugi sums are less than 3. Assume $\deg(\Delta_{K_1}) \ge \deg(\Delta_{K_i})$ for $i \ge 2$. Then the following hold, up to taking the mirror of K:

- (1) K_1 is H-M sharp with $\sigma(K_1) > 0$ (a negative special alternating link),
- (2) K_2, \ldots, K_n are positive special alternating links, and
- (3) $sl(K_1) \ge \sum_{i=2}^{n} \deg(\Delta_{K_i}).$

Note that a special alternating link is H-M sharp if and only if the absolute values of all of the coefficients of the Alexander polynomial are equal. This follows from Lemma 3.3. Alternating torus links $T_{2,q}$ are examples of H-M sharp special alternating links. Another example of such a family comes from taking the boundary of an *n*-twisted, unknotted annulus in S^3 , denoted by $T'_{2,2n}$, which can also be obtained by changing the orientation of one of the components of $T_{2,2n}$. For special alternating knots, H-M sharpness only holds when the knot is $T_{2,2g+1}$; see Ni [Ni21].

Many of the small H-M sharp links come from diagrammatic plumbings of links of the form $T_{2,q}$ and $T'_{2,2n}$. Figure 56 is the list of all H-M sharp prime knots with less than 11 crossings.



FIGURE 56. All H-M sharp prime knots up to 10 crossings.

3.3. **Dual inequality and concordance.** In this subsection, we investigate a connection between the H-M inequality and concordance based on the Fox–Milnor condition, stated in Theorem 3.17.

Theorem 3.17 ([FM66]). For concordant knots K_1 and K_2 , there exists a polynomial $f(t) \in \mathbb{Z}[t]$ and $a \in \mathbb{Z}$ such that

$$\Delta_{K_1}(t)\Delta_{K_2}(t) = t^a f(t) f(t^{-1}).$$

Note that, if one normalizes the Alexander polynomials so that they are both symmetric, then a = 0. All results in this subsection rely on the following:

Lemma 3.18. For a non-zero polynomial $f(t) = \sum_{i=0}^{n} a_i t^i \in \mathbb{Z}[t]$, consider

$$f(t)f(t^{-1}) = \sum_{i=0}^{n} c_i(t^i + t^{-i}).$$

Then $|c_0| > |c_k|$ for all k > 0. In particular, the stable length of $f(t)f(t^{-1})$ is zero.

Proof. We just need to compute c_0 , c_k , and their difference:

$$c_0 = \sum_{i=0}^n a_i^2, \ c_k = \sum_{i=0}^{n-k} a_i a_{i+k},$$

 \mathbf{SO}

$$c_0 - c_k = \sum_{i=0}^{n-k} \frac{1}{2} (a_i - a_{i+k})^2 + \frac{1}{2} \sum_{i=0}^{k-1} a_i^2 + \frac{1}{2} \sum_{i=n-k+1}^n a_i^2 > 0. \qquad \Box$$

We analyzed the stable length of a convolution of two sequences in Figure 55. Together with Lemma 3.18, we obtain Proposition 3.19.

Proposition 3.19. Assume K_1 is an alternating link which is algebraically concordant to K_2 . Then

$$sl(K_1) \le 2\deg(\Delta_{K_2}) + 1.$$

Proof. If $\deg(\tilde{\Delta}_{K_1}) < \deg(\tilde{\Delta}_{K_2})$, then the inequality is obvious. Assume that $\deg(\tilde{\Delta}_{K_1}) \ge \deg(\tilde{\Delta}_{K_2})$. If $\operatorname{sl}(K_1) > 2 \operatorname{deg}(\tilde{\Delta}_{K_2}) + 1$, then at least two copies of the sequence of coefficients of $\tilde{\Delta}_{K_2}$ falls in the stable part of $\tilde{\Delta}_{K_1}$; see Figure 55. Hence, the stable length of $\tilde{\Delta}_{K_1}\tilde{\Delta}_{K_2}$ is non-zero, contradicting Lemma 3.18. Note that the proof does not rely on trapezoidal inequalities, and hence K_2 can be non-alternating. \Box

In the light of Corollary 3.19, we define the following invariant, which we use in Corollary 3.21.

Definition 3.20. Define the *concordance degree* of a knot K as

 $d_c(K) = \min\{\deg(\Delta_{K'}) : K' \text{ algebraically concordant to } K\}.$

Corollary 3.21. The H-M inequality holds for any alternating link K satisfying

$$d_c(K) = |\sigma(K)|.$$

This includes all alternating links which are algebraically concordant to special alternating links.

Proof. Using Proposition 3.19, we have that

$$\operatorname{sl}(K) \le d_c(K) + 1 \Rightarrow \operatorname{sl}(K) \le |\sigma(K)| + 1 \Rightarrow \left\lfloor \frac{\operatorname{sl}(K)}{2} \right\rfloor \le \left\lfloor \frac{|\sigma(K)| + 1}{2} \right\rfloor,$$

where the last inequality is the H-M inequality for K. The second part of Corollary 3.21 follows from Lemma 3.3.

The H-M inequality can be seen as an optimization problem, where we maximize the stable length over all alternating links in an algebraic concordance class. Using inequality (3.3), one can construct a dual of this optimization problem in the form of minimizing the degree of the Alexander polynomial in any algebraic concordance class. In particular, we can search for knots satisfying

$$d_c(K) = \deg(\Delta_K) = |\sigma(K)|.$$

Note that this duality is weak, and the dual problem cannot always prove the H-M inequality:

Example 3.22. Let K be the knot 6_2 . We know that $\sigma(K) = -2$, $\deg(\Delta_K) = 4$, and sl(K) = 3. Note that Δ_K is an irreducible element of $\mathbb{Z}[t]$, which is a UFD. Using the Fox-Milnor condition, one can deduce that, for all links K' algebraically concordant to K, we have $\tilde{\Delta}_K | \tilde{\Delta}_{K'}$. Consequently, $d_c(K) = deg(\Delta_K) = 4$.

We can use the same argument as in Example 3.22 to compute the concordance degree of the figure eight knot.

We now show how to use the knot's symmetry to prove the H-M inequality in its algebraic concordance class. The figure eight knot is negative amphicheiral, and it satisfies the following:

Theorem 3.23 ([Liv05]). If a knot K is concordant to a negative amplicheiral knot, then

$$\tilde{\Delta}_K(t^2) = F(t)F(t^{-1}).$$

Combining this with Lemma 3.18 implies the following:

Corollary 3.24. An alternating knot K which is concordant to a negative amphicheiral knot has stable length zero, and hence satisfies the H-M inequality.

References

- [AC21] Marwa E. Alrefai and Nafaa Chbili. On Fox's trapezoidal conjecture for closed 3-braids. Internat. J. Math., 32(1):Paper No. 2150002, 11, 2021.
- [AC23] Mark E. AlSukaiti and Nafaa Chbili. Alexander and Jones polynomials of weaving 3-braid links and Whitney rank polynomials of Lucas lattice. preprint, arXiv:2303.11398, 2023.
- [AH21] Jared Able and Mikami Hirasawa. Construction and decomposition of knots as murasugi sums of seifert surfaces. *preprint*, *arXiv:2112.00297*, 2021.

62

- [Ban22] Ian M. Banfield. Christoffel words and the strong fox conjecture for two-bridge knots. *preprint*, arXiv:2212.04561, 2022.
- [BH20] Petter Brändén and June Huh. Lorentzian polynomials. Annals of Mathematics, 192(3):821 – 891, 2020.
- [BKL85] Anders Björner, Bernhard Korte, and László Lovász. Homotopy properties of greedoids. Adv. in Appl. Math., 6(4):447–494, 1985.
- [BS10] Francis Bonahon and Laurence Siebenmann. New geometric splittings of classical knots and the classification and symmetries of arborescent knots. *unpublished manuscript*, 2010.
- [BZH13] Gerhard Burde, Heiner Zieschang, and Michael Heusener. Chapter 13: Quadratic forms of a knot, pages 238–265. De Gruyter, Berlin, Boston, 2013.
- [Che17] Wenzhao Chen. On the Alexander polynomial and the signature invariant of two-bridge knots. preprint, arXiv:1712.04993, 2017.
- [Cro59] Richard H. Crowell. Genus of alternating link types. Annals of Mathematics, 69:258, 1959.
- [Erl99] Dieter Erle. Calculation of the signature of a 3-braid link. Kobe J. Math., 16(2):161–175, 1999.
- [FM66] Ralph H. Fox and John W. Milnor. Singularities of 2-spheres in 4-space and cobordism of knots. Osaka Journal of Mathematics, 3(2):257 – 267, 1966.
- [Fox61] R. H. Fox. Some problems in knot theory. In Topology of 3-manifolds and related topics (Proc. The Univ. of Georgia Institute, 1961), pages 168–176. Prentice-Hall, Inc., Englewood Cliffs, NJ, 1961.
- [Gab85] David Gabai. The Murasugi sum is a natural geometric operation. II. In Combinatorial methods in topology and algebraic geometry (Rochester, N.Y., 1982), volume 44 of Contemp. Math., pages 93–100. Amer. Math. Soc., Providence, RI, 1985.
- [Gab86] David Gabai. Genera of the alternating links. *Duke Mathematical Journal*, 53:677–681, 1986.
- [Har79] R. I. Hartley. On two-bridged knot polynomials. Journal of the Australian Mathematical Society, 28(2):241–249, 1979.
- [Hir95] Mikami Hirasawa. The flat genus of links. *Kobe journal of mathematics*, 12(2):155–159, 12 1995.
- [HM14] Mikami Hirasawa and Kunio Murasugi. On stability of Alexander polynomials of knots and links (survey). In *Knots in Poland. III. Part 1*, volume 100 of *Banach Center Publ.*, pages 85–98. Polish Acad. Sci. Inst. Math., Warsaw, 2014.
- [HMV23] Elena S. Hafner, Karola Mészáros, and Alexander Vidinas. Log-concavity of the Alexander polynomial. preprint, arXiv:2303.04733, 2023.
- [HT85] A. Hatcher and W. Thurston. Incompressible surfaces in 2-bridge knot complements. *Inventiones mathematicae*, 79(2):225–246, 1985.
- [HW04] Cam Van Quach Hongler and Claude Weber. On the topological invariance of Murasugi special components of an alternating link. *Mathematical Proceedings* of the Cambridge Philosophical Society, 137:95 – 108, 2004.
- [HW05] Cam Van Quach Hongler and Claude Weber. A Murasugi decomposition for achiral alternating links. *Pacific Journal of Mathematics*, 222:317–336, 2005.
- [Jon10] In Dae Jong. Alexander polynomials of alternating knots of genus two II. J. Knot Theory Ramifications, 19(8):1075–1092, 2010.
- [Kaw96] Akio Kawauchi. A survey of knot theory. Birkhäuser Verlag, Basel, 1996. Translated and revised from the 1990 Japanese original by the author.
- [Ká13] Tamás Kálmán. A version of tutte's polynomial for hypergraphs. Advances in Mathematics, 244:823–873, 2013.
- [LC18] Peter Lambert-Cole. Twisting, mutation and knot Floer homology. Quantum Topol., 9(4):749–774, 2018.

- [Liv05] Charles Livingston. A survey of classical knot concordance. In Handbook of knot theory, pages 319–347. Elsevier B. V., Amsterdam, 2005.
- [LM12] Lilya Lyubich and Kunio Murasugi. On zeros of the Alexander polynomial of an alternating knot. *Topology Appl.*, 159(1):290–303, 2012.
- [Min82] Jerome Minkus. The branched cyclic coverings of 2 bridge knots and links. Mem. Amer. Math. Soc., 35(255):iv+68, 1982.
- [Mur65] Kunio Murasugi. On a certain numerical invariant of link types. *Transactions* of the American Mathematical Society, 117:387–422, 1965.
- [Mur85] Kunio Murasugi. On the Alexander polynomial of alternating algebraic knots. J. Austral. Math. Soc. Ser. A, 39(3):317–333, 1985.
- [Ni21] Yi Ni. A characterization of $T_{2g+1,2}$ among alternating knots. Acta Math. Sin. (Engl. Ser.), 37(12):1841–1846, 2021.
- [OS03] Peter Ozsváth and Zoltán Szabó. Absolutely graded floer homologies and intersection forms for four-manifolds with boundary. Advances in Mathematics, 173(2):179–261, 2003.
- [Par78] Richard L. Parris. Pretzel knots. PhD thesis, Princeton University, 1978. Copyright - Database copyright ProQuest LLC; ProQuest does not claim copyright in the individual underlying works; Last updated - 2023-02-18.
- [SO03] Zoltán Szabó and Peter Ozsváth. Heegaard floer homology and alternating knots. Geometry & Topology, 7:225–254, 2003.
- [Sto05] Alexander Stoimenow. Newton-like polynomials of links. *Enseign. Math. (2)*, 51(3-4):211–230, 2005.
- [Sto08] A. Stoimenow. Knots of (canonical) genus two. Fund. Math., 200(1):1–67, 2008.
- [Sto16] Alexander Stoimenow. *Diagram genus, generators, and applications*. Monographs and Research Notes in Mathematics. CRC Press, Boca Raton, FL, 2016.
- [Sto21] Alexander Stoimenow. Independence polynomials and Alexander-Conway polynomials of plumbing links. Journal of Combinatorial Theory, Series A, 183:105487, 10 2021.
- [Tra04] Paweł Traczyk. A combinatorial formula for the signature of alternating diagrams. Fundamenta Mathematicae, 184:311–316, 2004.

MATHEMATICAL INSTITUTE, UNIVERSITY OF OXFORD, ANDREW WILES BUILDING, RADCLIFFE OBSERVATORY QUARTER, WOODSTOCK ROAD, OXFORD, OX2 6GG, UK Email address: azarpendar@maths.ox.ac.uk

MATHEMATICAL INSTITUTE, UNIVERSITY OF OXFORD, ANDREW WILES BUILDING, RADCLIFFE OBSERVATORY QUARTER, WOODSTOCK ROAD, OXFORD, OX2 6GG, UK Email address: juhasza@maths.ox.ac.uk

DEPARTMENT OF MATHEMATICS, TOKYO INSTITUTE OF TECHNOLOGY, H-214, 2-12-1 Ookayama, Meguro-ku, Tokyo 152-8551, Japan, and International Institute for Sustainability with Knotted Chiral Meta Matter (WPI-SKCM²), Hiroshima University 1-3-1 Kagamiyama, Higashi-Hiroshima, Hiroshima 739-8526, Japan

Email address: kalman@math.titech.ac.jp

64