Expressing turbulent kinetic energy as coarse-grained enstrophy or strain deformations

Damiano Capocci*

Department of Physics, University of Rome Tor Vergata

In turbulent flows, the fluid element gets deformed by chaotic motion due to the formation of sharp velocity gradients. A direct connection between the element of fluid stresses and the energy balance still remains elusive. Here, an exact identity of incompressible turbulence is derived linking the velocity gradient norm across the scales with the total kinetic energy. In the context of three-dimensional (3D) homogeneous turbulence, this relation can be specialised obtaining the expression of total kinetic energy decomposed either in terms of deformations due to strain motion or via the resolved-scale enstrophy of the fluid element. Applied to data from direct numerical simulations (DNS) describing homogeneous and isotropic turbulence, the decomposition reveals that, beyond the scales dominated by the external forcing, extensional and contractile deformations account approximately for 55% and 40% of the kinetic energy of the associated scale while less than the remaining 5% is carried by the indefinite-type stresses. From these two identities one can derive an exact expression for the kinetic energy spectrum which is based solely on real space quantities providing a characterisation of the Kolomogorov constant as well. Numerical evidences show that this formulation of the energy spectrum reproduces the power-law behaviour of the Kolmogorov spectral scaling.

Among the countless scientific contexts that deal with turbulent flows, two characteristics are recurrent: the nonlinearity and the multiscale nature of the velocity field leading to a spatio-temporal chaotic dynamics. Turbulence is a common emerging property in water or air when the value of the viscosity is *small* compared to the related typical sizes and velocities. A turbulent flow is characterised by the formation of small-scale vortical motion from a large scale injection of energy; this phenomenon is usually referred to as the *cascade* of energy across the scales. This energy transfer is responsible of the production of velocity gradient (VG) at every scale which, in turn, determines the deformations that the element of fluid undergoes. Characterising the interplay between turbulent kinetic energy and flow deformation not only pertains to fundamental knowledge but is also relevant for applications ranging from geophysics [1–3] to astrophysics [4–6]. To the author's knowledge, the functional dependence between the total kinetic energy and both the intensity of strain and rotational motions is still unknown.

In this work, by proving a more general identity, two expressions for the total kinetic energy are derived in homogeneous turbulence: one written in terms of the rate of strain deformation and the other based on the squared vorticity, commonly referred as *enstrophy*. As an extension of the latter, a corresponding exact identity for the kinetic energy spectrum is derived through which it is possible to connect the kinetic energy spectrum with filtered enstrophy. The derived identities are verified via data from direct numerical simulation (DNS) and used to reveal the percentages of the purely contractile, the purely extensional and the *indefinite-type* stresses that constitute the kinetic energy across the scales. The validity of the novel representation of the kinetic energy spectrum is tested as well.

The Navier-Stokes equation describing the evolution of an incompressible velocity field $\boldsymbol{u}(\boldsymbol{x},t)$ is the following:

$$\partial_t u_i = -u_j \frac{\partial u_i}{\partial x_j} - \frac{1}{\rho} \frac{\partial p}{\partial x_i} + \nu \nabla^2 u_i + f_i \tag{1}$$

where ν is the kinematic viscosity and ρ the mass density. The linear viscous term $\nu \nabla^2 u_i$ corresponds to the dissipation, being related to the conversion of kinetic energy into heat. The pressure field $p(\boldsymbol{x},t)$ enforces the incompressibility constraint $\partial_i u_i = 0$ and f_i is a generic external force injecting energy into the system. The VG tensor $u_{i,j} = \partial u_i / \partial x_j$ describes the flow geometry in terms of strain rate $S_{ij} = (u_{i,j} + u_{j,i})/2$ and rotation rate $\Omega_{ij} = (u_{i,j} - u_{j,i})/2$, which can also be expressed as the vorticity $\omega_i = \epsilon_{ijk} \Omega_{jk}$. Furthermore, its Frobenius norm determines the rate at which kinetic energy is dissipated into thermal energy in $\varepsilon = \nu u_{i,j} u_{i,j}$. Eq. (1) admits one dimensionless control parameter which is the Reynolds number $Re_L = U L/\nu$, where U and L are characteristic velocity and length scale of the flow, respectively.

By applying a low-pass filter G^{ℓ} , we can separate the large-scale dynamics from the small-scale ones [7]:

$$\overline{u}_i^\ell = G^\ell * u_i, \qquad \tilde{\overline{u}}_i^\ell = \tilde{G}^\ell \cdot \tilde{u}_i \tag{2}$$

where (\cdot) indicates the Fourier transform and * the convolution product. As regards the properties of G^{ℓ} , we require it to be an even function with volume average equal to unity [8].

The application of a low-pass filter to eq. (1) provides:

$$\partial_t \overline{u}_i^\ell = -\overline{u}_j^\ell \frac{\partial \overline{u}_i^\ell}{\partial x_j} - \frac{1}{\rho} \frac{\partial \overline{p}^\ell}{\partial x_i} + \nu \nabla^2 \overline{u}_i^\ell + \overline{f}_i^\ell - \partial_j \tau_{ij}^\ell \qquad (3)$$

^{*} capocci@roma2.infn.it

where $\tau_{ij}^{\ell} = \overline{u_i u_j}^{\ell} - \overline{u}_i^{\ell} \overline{u}_j^{\ell}$ is the so-called subgrid-scale stress (SGS) tensor, representing the effective stress due to the features smaller than ℓ on the resolved-scale velocity. Therefore, we can define the kinetic energy related to scales larger than ℓ i.e. the *resolved-scale energy* as $E_K^{\ell}(\boldsymbol{x},t) = \frac{1}{2} \overline{u}_i^{\ell}(\boldsymbol{x},t) \overline{u}_i^{\ell}(\boldsymbol{x},t)$ and the kinetic energy at scales smaller than ℓ which is the trace $\frac{1}{2} \tau_{ii}^{\ell}(\boldsymbol{x},t)$ as the *unresolved-scale energy*. Note that, by definition, the sum of these two quantities determines the filtered kinetic energy:

$$\frac{1}{2}\overline{u_i u_i}^\ell = E_K^\ell + \frac{1}{2}\tau_{ii}^\ell \tag{4}$$

where we omit the spatio-temporal dependence. If the filtered kernel is non-negative i.e. $G^{\ell}(\mathbf{r}) \geq 0 \ \forall \mathbf{r}$, then $\tau_{ii}(\mathbf{x}) \geq 0 \ \forall \mathbf{x}$, hence τ_{ii} can be correctly interpreted as an energy [9]. We also remind that $\overline{u}^{\ell=0}(\mathbf{x},t) = u(\mathbf{x},t)$ implying that $\tau^{\ell=0} = 0$ and $E_K^{\ell=0} \equiv E_K = \frac{1}{2}u_i(\mathbf{x},t) u_i(\mathbf{x},t)$ which is the kinetic energy.

At this stage, we note that, if a scalar function g admits the Fourier representation [10], it is straightforward to conclude that the relation between the volume averages $\langle \overline{g}^{\ell} \rangle = \langle g \rangle$ is a consequence of the properties of the (generic) filter kernel. This observation becomes relevant for the calculation of the volume average of (4). Indeed, as already noted by [11], we can neglect the filter application on the LHS obtaining:

$$\langle E_K \rangle = \langle E_K^\ell \rangle + \frac{1}{2} \langle \tau_{ii}^\ell \rangle \tag{5}$$

where the filtering parameter, belonging only to the RHS, broadly resembles a *subtraction point* of the renormalization theory [12].

At this point, in order to satisfy the non-negativity of the filter kernel, we center our discussion on the Gaussian kernel:

$$G^{\ell}(\boldsymbol{r}) = \left(2\pi\ell^2\right)^{-3/2} e^{-\boldsymbol{r}^2/2\ell^2}, \qquad \tilde{G}^{\ell}(\boldsymbol{k}) = e^{-\boldsymbol{k}^2\ell^2/2} \quad (6)$$

As a specific feature of the Gaussian filter, there exists an exact decomposition of the SGS stresses [13][14] that reads:

$$\tau_{ik}^{\ell} = \int_{0}^{\ell^{2}} \mathrm{d}\theta \ \overline{\overline{u}_{i,j}^{\sqrt{\theta}}} \ \overline{\overline{u}_{k,j}^{\sqrt{\theta}}}^{\sqrt{\ell^{2}-\theta}} \tag{7}$$

expressing SGS stresses [15] as the sum of the contributions of VG fields from the scales $\leq \ell$ that are in turn projected into the larger scales by the complementary filter $\sqrt{\ell^2 - \theta}$. It follows that the trace of (7) allows us to recast eq. (5) into:

$$\langle E_K \rangle = \langle E_K^\ell \rangle + \frac{1}{2} \int_0^{\ell^2} \mathrm{d}\theta \, \langle \overline{u}_{i,j}^{\sqrt{\theta}} \, \overline{u}_{i,j}^{\sqrt{\theta}} \rangle \tag{8}$$

where the filtering at $\sqrt{\ell^2 - \theta}$ in the averaged trace of (7) can be ignored by virtue of the filter-invariance property of scalar functions mentioned above. In addition, the exchange of the volume averaging and the integral over the

scales is granted by the assumption that the VG norm is bounded. By assuming finite energy, even if we let our system to be characterised by infinite length scales, it would immediately imply that $\lim_{\ell \to \infty} \langle E_K^\ell \rangle = 0$. The prescription imposed by this limit appears as a reasonable assumption for any classical physics system which is well-defined in its range of scales [16]. A detailed and more generalised analysis on both space and scale convergence of (8) will be provided elsewhere [17]. By applying the previous considerations and calculating the limit for infinite filtering-scale on the RHS of (8), we get:

$$\langle E_K \rangle = \frac{1}{2} \int_0^\infty \mathrm{d}\theta \, \langle \overline{u}_{i,j}^{\sqrt{\theta}} \, \overline{u}_{i,j}^{\sqrt{\theta}} \rangle \tag{9}$$

that can be eventually combined with (8) obtaining the more general expression:

$$\langle E_K^\ell \rangle = \frac{1}{2} \int_{\ell^2}^{\infty} \mathrm{d}\theta \, \langle \overline{u}_{i,j}^{\sqrt{\theta}} \, \overline{u}_{i,j}^{\sqrt{\theta}} \rangle. \tag{10}$$

Eqs. (9) and the resolved-scale counterpart (10) can be conceived as an exact gradient-tensor decomposition of kinetic energy. The above equations also go beyond the standard representation of kinetic energy as the squared Euclidean norm of velocity presenting the mean energy budget as the sum over the scales of the squared VG tensor norm.

At this point, in order to quantify eq. (10) we consider data from DNS. Our database [18] describes stationary homogeneous and isotropic turbulence where eq. (1) is evolved in a triply-periodic box with a stochastic forcing active in the wavenumber band $k \in [0.5, 2.4]$ with $k = \pi/\ell$, whose midpoint k_f determines the characteristic forcing scale L_f . This dataset is marked by a $Re_{\lambda} = 327$ where $\lambda = E_K \sqrt{15\nu/\langle \varepsilon \rangle}$ is the Taylor-scale.

In the following quantification, we are relying on one instantaneous configuration of the velocity field. In particular, fig. 1 compares (10) with the *usual* expression of the averaged resolved-scale kinetic energy. Both these two expressions are normalised by the total kinetic energy and displayed as a function of the adimensional parameter L_f/ℓ . In consequence of the validity of (10), we clearly observe that the two displayed profiles are indistinguishable.

As a conclusion of this preliminary analysis, it is worth to underline that in homogeneous turbulence, the first Betchov relation [19]

$$\langle \overline{S}_{ij}^{\ell} \overline{S}_{ij}^{\ell} \rangle = \frac{1}{2} \langle || \overline{\boldsymbol{\omega}}^{\ell} ||^2 \rangle, \qquad (11)$$

holding $\forall \ell$, implies that the mean (filtered) VG norm can be either written in terms of the (filtered) squared strain or via the (filtered) enstrophy.



Figure 1. Resolved scale mean kinetic energy normalised by the unfiltered value $\langle E_K^\ell \rangle / \langle E_K \rangle$ as a function of the dimensionless parameter L_f / ℓ calculated from the standard definition and from eq. (10). The two curves clearly overlap.

I. MEAN KINETIC ENERGY IN TERMS OF FLUID ELEMENT DEFORMATIONS

In this section, as introduced above, we are going to express the mean squared VG appearing on the RHS of (10) only in terms of the resolved-scale strain-rate squared via the application of (11). Additionally, since (11) is an identity involving kinematic invariants, we can choose the reference frame where S_{ij} becomes diagonal, the so-called principal axis frame. We also remark that the strain-rate tensor, being symmetric, has three real eigenvalues λ_i with $i \in \{1, 2, 3\}$ related to a set of orthogonal eigenvectors. Moreover incompressibility imposes the pointwise constraint $\overline{\lambda}_1^{\ell} + \overline{\lambda}_2^{\ell} + \overline{\lambda}_3^{\ell} = 0$. As a consequence, the largest eigenvalue $\overline{\lambda}_1^{\ell} \ge 0$ is always extensional along the direction of its eigenvector and the smallest one $\overline{\lambda}_3^\ell \leq 0$ is always related to a contractile direction while the intermediate $\overline{\lambda}_2^{\epsilon}$ can be either positive or negative from which the definition *indefinite-type*. In light of the preceding discussion, we express (10) via the (sorted) strain-rate tensor eigenvalues obtaining:

$$\langle E_K^\ell \rangle = \int_{\ell^2}^\infty \mathrm{d}\theta \, \langle \, (\overline{\lambda}_1^{\sqrt{\theta}})^2 + (\overline{\lambda}_2^{\sqrt{\theta}})^2 + (\overline{\lambda}_3^{\sqrt{\theta}})^2 \, \rangle \qquad (12)$$

$$=P_1^{\ell} + P_2^{\ell} + P_3^{\ell} \,. \tag{13}$$

where each P_i^{ℓ} represents the contribution of the *i*-th strain-rate eigenvalue to the mean resolved-scale kinetic energy. The above equation describes the energy budget via the intensity of stresses along their principal axes.

By using the abovementioned data, in eq. (12) we can quantify the role of each P_i^{ℓ} across the scales. In fact,



Figure 2. Strain-rate eigenvalues contributions P_i^{ℓ} normalised by the resolved-scale as a function of L_f/ℓ . Both the axes show dimensionless quantities. The values of the curves sum to unity scale by scale. The displayed percentages refer to the smallest resolved-scale. The horizontal thin lines, with their corresponding *y*-axis (rounded) values, are added manually to emphasize the values at which P_i^{ℓ} saturates.

fig. 2 displays the adimensional P_i^{ℓ} normalised by $\langle E_K^{\ell} \rangle$ as a function of the non-dimensional parameter L_f/ℓ describing the percentage of each P_i^{ℓ} in the kinetic energy balance at the scale ℓ . We first notice that, apart from the region $L_f/\ell \lesssim 1$ where the flow geometry is governed by the external forcing, each P_i^{ℓ} is monotonic where $\overline{\lambda}_1^{\ell}$ increases, $\overline{\lambda}_2^{\ell}$ weakly increases, being essentially scale-invariant, while $\overline{\lambda}_3^{\ell}$ decreases. As we progress towards the small scales each contribution saturates. In particular for these scales, the purely extensional motion carries about 54% of the corresponding resolved-scale energy while the purely contractile stresses approximately 41%, as a result the remaining part is due to the indefinite-type deformations.

II. CONNECTING RESOLVED SCALE ENERGY AND ENSTROPHY

In this section, as a complementary procedure, we are going to express the mean squared VG appearing on the RHS of (10) in terms of the resolved-scale enstrophy using (11). Hence, the equivalent expression to (12) yields:

$$\langle E_K^\ell \rangle = \frac{1}{2} \int_{\ell^2}^{\infty} \mathrm{d}\theta \, \langle \, || \overline{\omega}^{\sqrt{\theta}} ||^2 \, \rangle.$$
 (14)

which reveals that, by measuring the resolved-scale enstrophy in real space at each scale, it is possible to retrieve the (resolved-scale) mean kinetic energy. The prior equation becomes definitely more interesting in 2D turbulence where the enstrophy $\zeta = \frac{1}{2} \langle ||\overline{\omega}||^2 \rangle$ is an inviscid conserved quantity; in this scenario the above equation connects two conserved quantities which is non-trivial. It is even less trivial that in 3D flows a conserved (inviscid) quantity can be expressed via the coarse-graining of a non-conserved one. Eq. (14) can be roughly thought of as a sort of *enstrophy-based quantization* of the kinetic energy in analogy with the Hamiltonian operator formulated via ladder operators in quantum field theory [20]. Connected to this interpretation, we underline that approximating the unresolved-scale energy integral of eq. (14) at a given scale ℓ , meaning $\tau_{ii}^{\ell} \approx \frac{1}{2} \ell^2 ||\overline{\omega}^{\ell}||^2$, resembles the expression of the rotational energy of a ring of radius ℓ from rigid body mechanics.

It is remarkable to notice that eq. (14) and in consequence (12), setting $\ell = 0$ are more general. In order to discuss this, we focus on the RHS of eq. (14) disentangling it from its derivation. In this respect we consider the coarse-graining on the vorticity to be implemented by a generic filter kernel G^{ℓ} instead of Gaussian kernel on which the presented methodology is based. In this manner, the RHS of eq. (14) downgrades to:

$$\frac{1}{2} \int_{\ell^2}^{\infty} \mathrm{d}\theta \,\langle \, || \overline{\boldsymbol{\omega}}^{\sqrt{\theta}} ||^2 \,\rangle = \int_{\ell^2}^{\infty} \mathrm{d}\theta \,\int_{\mathbb{R}^3} \mathrm{d}\boldsymbol{k} \,k^2 \,\tilde{u}_i(\boldsymbol{k}) \,\tilde{u}_i^*(\boldsymbol{k}) \,\tilde{G}^{\sqrt{\theta}}(k)^2 \\
= \int_{\mathbb{R}^3} \mathrm{d}\boldsymbol{k} \,\,\tilde{u}_i(\boldsymbol{k}) \,\tilde{u}_i^*(\boldsymbol{k}) \int_{\ell^2 k^2}^{\infty} \mathrm{d}s \,\tilde{G}(\sqrt{s})^2 \\$$
(15)

where $(\cdot)^*$ denotes the complex conjugation. For simplicity, we assume that the velocity field allows the continuous Fourier representation (see [21] for details). Back to the RHS of (15), the kernel dependency on the filtering-scale has been absorbed into its argument i.e. $\tilde{G}^{\sqrt{\theta}}(k) = \tilde{G}(k\sqrt{\theta})$. This effective rescaling holds for the standard filter functions, like those listed in table 13.2 of [8] but in principle this would remove from our discussion *exotic* filter kernel types for which the preceding rescaling does not hold. Like already mentioned above, the inversion of the integrals in the derivation of (15) is a consequence of the assumed convergence of the fields involved.

At this stage, we observe [22, 23] that by setting $\ell = 0$ in (15), the two integrals can be factorised. In this way the RHS of (15) collapses into the product $C_G \langle E_K \rangle$ where C_G is a real number depending on the filter choice and the mean energy is given by the integration of the velocity Fourier modes defined in (2). In such a way, we have generalised eq. (14) to a generic class of filters without introducing the SGS stresses decomposition of (7) and without requiring the associated positive definiteness of its trace. Note that a similar factorization can be derived for the *nth*-derivative of the velocity field [23].

However, depending on the filter kernel, the generalisation originating from the preceding observation does not necessarily hold when (15) is evaluated at $\ell \neq 0$; this condition prevents the calculation of the related kinetic energy spectrum in eq. (18). In order to show this, we consider the *sharp filter* whose Fourier transform is a Heaviside function $G^{\ell}(k) = H(\pi - k \ell)$. For this kernel, eq. (15) becomes $\int_0^{\pi/\ell} dk E(k)(\pi^2 - \ell^2 k^2)$ which does not equal the corresponding mean resolved-scale energy $\langle E_K^{\ell} \rangle = \int_0^{\pi/\ell} dk E(k)$.

III. ADDRESSING THE NON-STATIONARITY

Having obtained an exact expansion of the mean kinetic energy in (14) allows for the dynamical characterization of non-stationary flows $\partial_t \langle E_K(t) \rangle \geq 0$, respectively describing the flow energy increase and the turbulence decay where the time dependence of the energy is purposefully reinstated. For this reason, applying a time derivative to eq. (9) one can obtain an expression involving filtered VG quantities. For simplicity we focus on the homogeneous turbulence case, where the calculation of the time derivative in (9) reads:

$$\partial_t \langle E_K(t) \rangle = \frac{1}{2} \int_0^\infty \mathrm{d}\theta \, \partial_t \, \langle \overline{u}_{i,j}^{\sqrt{\theta}} \overline{u}_{i,j}^{\sqrt{\theta}} \rangle \tag{16}$$

$$= \int_{0}^{\infty} \mathrm{d}\theta \left(-\frac{4}{3} \langle \overline{S}_{ij}^{\sqrt{\theta}} \overline{S}_{jk}^{\sqrt{\theta}} \overline{S}_{ki}^{\sqrt{\theta}} \rangle - \nu \langle \overline{u}_{i,jk}^{\sqrt{\theta}} \overline{u}_{i,jk}^{\sqrt{\theta}} \rangle - \langle \overline{S}_{ij,kk}^{\sqrt{\theta}} \tau_{ij}^{\sqrt{\theta}} \rangle \right)$$

$$+ \langle F^{\sqrt{\theta}} \rangle \right)$$

$$(17)$$

where $F^{\sqrt{\theta}} = \overline{u}_{i,j}^{\sqrt{\theta}} \overline{f}_{i,j}^{\sqrt{\theta}}$ depends on the external forcing and it is set to zero in the case of turbulence decay. As concerns the other terms, the one containing the contraction of three strain-rate tensors is linked to a contribution of *energy transfer* across the scales [13]. Numerical evidence [13][14] shows that it is positive $\forall \ell$ at the stationary state, then it is expected to maintain the same sign in the present non-stationary case as well. As for the third term, it is formed by the contraction between the strain-rate tensor Laplacian and the SGS stresses; being a priori sign indefinite, it may play a relevant role in the above energy evolution equation. Such a term also shows an intrinsic *multi-scale* nature from the presence of the SGS stress tensor which, by virtue of (7), depends on the scales smaller than the integration scale $\sqrt{\theta}$. Finally, it is clear that the term proportional to the viscosity is negative. Applying (17) to data from DNS will assess the importance of such terms in unsteady configurations.

Independently from the quantification of the terms, in case of stationarity i.e. $\partial_t \langle E_K \rangle = 0$, the above expression indicates that the stationarity can, in principle, be reached in two ways: a scale-by-scale stationarity where the contribution of each integrated scale is identically zero or the *scale-time ergodicity* [24] where the RHS of (16) is zero because of a compensation of the nonstationarity of the corresponding scales contributions. It is worth observing that in 2D turbulence eq. (17) simplifies dramatically. Firstly the contraction of the three (resolved scale) strain-rate tensors is identically zero pointwise (see e.g. Appendix C.1 of [25]). Secondly, in the third term, the decomposition of the SGS stresses in (7) comprises of only one term [26][14], as opposed to the general 3D case. It follows that eq. (17) can help to identify the governing physical mechanisms that lead the the turbulence decay or the increase of energy.

IV. AN EXACT IDENTITY FOR THE KINETIC ENERGY SPECTRUM

From the knowledge of the resolved-scale kinetic energy it is possible to obtain the kinetic energy spectrum E(k) through the following formula:

$$E(k) = -\frac{\pi}{k^2} \frac{\partial}{\partial \ell} \langle E_K^\ell \rangle \tag{18}$$

where the time dependence has been omitted. In order to get an expression in terms of velocity gradient, we can plug eq. (14) into the previous equation obtaining:

$$E(k) = \pi^2 k^{-3} \left\langle ||\overline{\boldsymbol{\omega}}^{\pi/k}||^2 \right\rangle \tag{19}$$

which is an exact expression for the kinetic energy spectrum where the wavenumber dependency is both kept by the wavenumber prefactor and by the filter scale that the vorticity is coarse-grained at. The above equation must be compared with the *usual* definition of the kinetic energy spectrum:

$$E^{U}(k) = \frac{1}{2} \int_{\mathbb{R}^{3}} \mathrm{d}\boldsymbol{k} \ \tilde{u}_{i}(\boldsymbol{k}) \, \tilde{u}_{i}^{*}(\boldsymbol{k}) \, \delta(||\boldsymbol{k}|| - k)$$
(20)

where $\delta(\cdot)$ is a delta function. It follows that eq. (19) qualifies as an alternative *representation* of the kinetic energy spectrum [27] which, unlike eq. (20), can be determined directly by real space quantities like the resolvedscale enstrophy. Thus, we can consider the dataset used so far to test eq. (19) by comparing it with the common definition of energy spectrum from eq. (20). Moreover, in turbulence theory, when the study of energy spectra is of interest, it is customary to discuss the corresponding Kolmogorov's scaling [21, 28, 29] which is $E^{KOL}(k) = C \langle \varepsilon \rangle^{2/3} k^{-5/3}$ where C is the Kolmogorov constant, according to the employed dataset its value is nearly 1.6, see [18] for details.

Fig. 3 shows the comparison between eqs. (20), (19) and the Kolmogorov scaling in the evaluation of the kinetic energy spectrum. The first feature we notice is the good agreement of E(k) with Kolmogorov scaling [30] in the wavenumber band $1.5 \leq k/k_f \leq 30.0$. As concerns the comparison between E(k) and $E^U(k)$, we appreciate that the former reproduces the latter in the forcing range i.e. $k/k_f \approx 1$. It is intuitive to recognize both the smoother profile of E(k) and the discrepancy for large k



 10^{1}

 k/k_f

 $E(k)/\langle E_K \rangle L_f$

 10^{0}

Figure 3. Omni-directional kinetic energy spectrum based on one instantaneous configuration and calculated respectively via the two different representations of eqs. (19) and (20). The dotted line refers to the Kolmogorov power-law spectral scaling. All the curves are expressed as functions of the non-dimensional quantity $k/k_f = L_f/\ell$ and normalised by $\langle E_K \rangle L_f$.

with $E^{U}(k)$ as an *inheritance* of the Gaussian filter, according to which filtering at wavenumber k still retains features from < k.

For those scales where $E^{KOL}(k) = E(k)$, we can equate the two corresponding expressions tentatively writing an expression for the Kolmogorov constant:

$$C = \pi^2 \frac{\langle ||\overline{\boldsymbol{\omega}}^{\pi/k}||^2 \rangle}{\langle \varepsilon \rangle^{2/3} k^{4/3}}$$
(21)

that requires further investigations. However a rough estimate would be evaluating (21) at $k/k_f \gg 1$, for instance at the so-called Kolmogorov microscale $\eta = (\nu^3/\varepsilon)^{1/4}$ s.t. $\eta \ll L_f$ [31]. This approximation would provide a $C = \pi^{2/3} \simeq 2.1$ against a $C \approx 1.6$ measured from the compensated energy spectrum [18]. See [32] and references therein for a complete discussion on characterisation of the Kolomogorov constant.

V. CONCLUSIONS AND FUTURE DIRECTIONS

In conclusion, from a more general identity relating the total kinetic energy and the coarse-grained velocity gradient norm based on the Gaussian filter kernel, two (sub)identities are derived for homogeneous turbulence. One expresses the mean energy balance in terms of the strain-rate intensity. Outside the range of scale dominated by the forcing, the purely extensional and contractile deformations provide respectively nearly 55% and

 10^{2}

40% of the resolved-scale energy. The remainder is linked to the intermediate eigenvalues which can be either related to contractions or extensions. Using kinematic constraints, one can obtain a cognate identity showing that in turn the sum of the enstrophy across the scales determines the total kinetic energy. Finally, one can obtain an exact identity for the kinetic energy spectrum in terms of in real space interpretable observables providing a deeper fundamental understanding of the energy spectrum and a guidance for the analytical determination of the Kolmogorov constant. As a consequence of the properties of SGS stresses expansion from (7), this present methodology can be applied to any process characterised by an advective-type non-linearity. On top of that, beyond the strain-vorticity splitting hereby considered, we can also perform different VG decompositions, e.g. [33] [34], to further explore the role of the VG norm in the energy budget across the scales. The analysis of the dynamical equation describing unsteady flows can be deeply expanded to unveil the governing physical mechanisms that

- D. Buaria, A. Pumir, E. Bodenschatz, and P.-K. Yeung, Extreme velocity gradients in turbulent flows, New Journal of Physics 21, 043004 (2019).
- [2] S. A. Thorpe, *The turbulent ocean* (Cambridge university press, 2005).
- [3] J. Pedlosky, *Geophysical fluid dynamics* (Springer Science & Business Media, 2013).
- [4] J. Cho, A. Lazarian, and E. T. Vishniac, *MHD turbulence: scaling laws and astrophysical implications* (Springer, 2003).
- [5] M. L. Goldstein, D. A. Roberts, and W. Matthaeus, Magnetohydrodynamic turbulence in the solar wind, Annual review of astronomy and astrophysics **33**, 283 (1995).
- [6] P. Goldreich and S. Sridhar, Magnetohydrodynamic turbulence revisited, The Astrophysical Journal 485, 680 (1997).
- [7] M. Germano, Turbulence: the filtering approach, Journal of Fluid Mechanics 238, 325 (1992).
- [8] S. B. Pope, *Turbulent Flows* (Cambridge University Press, 2000).
- [9] B. Vreman, B. Geurts, and H. Kuerten, Realizability conditions for the turbulent stress tensor in large-eddy simulation, Journal of Fluid Mechanics 278, 351 (1994).
- [10] Note that, for an unbounded domain, the Fourier representation can be bypassed by requiring that $g(\boldsymbol{x})$ goes to zero as $||\boldsymbol{x}|| \to \infty$. The bypass is formal in periodic boundaries which in turn would imply the expansion in Fourier series. The same considerations applies to the derivation of eq. (7) in case one wants to avoid the Fourier expansion.
- [11] H. Aluie, Coarse-grained incompressible magnetohydrodynamics: analyzing the turbulent cascades, New Journal of Physics 19, 025008 (2017).
- [12] J. Zinn-Justin, *Path integrals in quantum mechanics* (OUP Oxford, 2010).
- [13] P. L. Johnson, Energy transfer from large to small scales in turbulence by multiscale nonlinear strain and vorticity

lead the turbulence decay or the increase of energy.

VI. ACKNOWLEDGMENTS AND FUNDINGS

The author thanks Prof. Luca Biferale and Dr. Massimo Cencini for the financial support provided by both European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 882340) and FieldTurb experiment of the Istituto Nazionale di Fisica Nucleare (INFN). This work used the ARCHER2 UK National Supercomputing Service (www.archer2.ac.uk) with resources provided by the UK Turbulence Consortium (EP-SRC grants EP/R029326/1). The author thanks Dr. Moritz Linkmann for the interesting discussions, encouragement and the support. The author wants to thank Prof. Michael Wilczek and Lukas Bentkamp for their valuable observations and Martin Lellep for editing suggestions.

interactions, Phys. Rev. Lett. 124, 104501 (2020).

- [14] P. L. Johnson, On the role of vorticity stretching and strain self-amplification in the turbulence energy cascade, Journal of Fluid Mechanics 922, A3 (2021).
- [15] Note that the methodology of [13] requires homogeneity or periodic boundaries in order to set to zero the boundary terms and not for the derivation of the decomposition (7) itself.
- [16] The length contraction in general relativity would already pose conceptual difficulties to the present filtering approach.
- [17] However, in the applications, the interest clearly focuses on systems characterised by a maximum allowed scale for which the integration upper bound of (7) does not diverge ensuring a rigorous exchange of integrals to derive (8).
- [18] L. Biferale, F. Bonaccorso, M. Linkmann, and D. Capocci, Turb-hel: an open-access database of helically forced homogeneous and isotropic turbulence, arXiv preprint arXiv:2404.07653 (2024).
- [19] R. Betchov, An inequality concerning the production of vorticity in isotropic turbulence, Journal of Fluid Mechanics 1, 497 (1956).
- [20] M. E. Peskin, An introduction to quantum field theory (CRC press, 2018).
- [21] U. Frisch, Turbulence: the legacy of AN Kolmogorov (Cambridge University Press, 1995).
- [22] L. Bentkamp, Private communication.
- [23] M. Wilczek, Private communication.
- [24] M. Linkmann, Private communication.
- [25] D. Capocci, P. Johnson, S. Oughton, L. Biferale, and M. Linkmann, Energy flux decomposition in magnetohydrodynamic turbulence (2024), arXiv:2402.10125 [physics.flu-dyn].
- [26] Given a symmetric tensor A_{ij} , in 2D one can show that $A_{ij}\tau_{ij}^{\ell} = A_{ij}\int_{0}^{\ell^{2}} dq \,\overline{S_{ij}^{\sqrt{q}} \,\overline{\Omega}_{ij}^{\sqrt{q}}} \,.$ [27] By expressing the filtered in the second sec
- [27] By expressing the filtered enstrophy in Fourier space, one can notice that the rationale of eq. (19) consists in replac-

ing the delta function in (20) by a unit-measure Gaussian.

- [28] A. N. Kolmogorov, Dissipation of Energy in Locally Isotropic Turbulence, Akademiia Nauk SSSR Doklady 32, 16 (1941).
- [29] A. N. Kolmogorov, The Local Structure of Turbulence in Incompressible Viscous Fluid for Very Large Reynolds Numbers, Proceedings of the Royal Society of London Series A 434, 9 (1991).
- [30] Note that by [35], $||\overline{\omega}^{\ell}|| \sim \ell^{-2/3}$ hence eq. (19) scales as $\ell^{5/3} \sim k^{-5/3}$ like the Kolmogorov formula.
- [31] The approximation of this calculation is based on equating the Kolmogorov scaling $E^{KOL}(k)$ with E(k) of eq. (19) at Kolmogorov micro-scale $\eta = (\nu^3/\langle \varepsilon \rangle)^{1/4}$ for which $k_\eta/k_f = 344$. Even though this wavenumber goes beyond the resolution of the analysed dataset, fig. 3

potentially indicates that the two curves do not intersect. From an analytical point of view, *large* values of k would make (21) independent of both the viscosity ν the (resolved-scale) enstrophy where the latter is due to the legitimacy of the approximation $||\overline{\omega}^{\pi/k}|| \approx ||\omega||$.

- [32] D. A. Donzis and K. R. Sreenivasan, The bottleneck effect and the kolmogorov constant in isotropic turbulence, Journal of Fluid Mechanics 657, 171–188 (2010).
- [33] V. Kolář, Vortex identification: New requirements and limitations, International journal of heat and fluid flow 28, 638 (2007).
- [34] R. Das and S. S. Girimaji, Revisiting turbulence smallscale behavior using velocity gradient triple decomposition, New Journal of Physics 22, 063015 (2020).
- [35] G. L. Eyink, Local energy flux and the refined similarity hypothesis, Journal of Statistical Physics 78, 335 (1995).