Angela Zhou University of Southern California Data Sciences and Operations zhoua@usc.edu

6/12/2024

Abstract

Offline reinforcement learning is important in many settings with available observational data but the inability to deploy new policies online due to safety, cost, and other concerns. Many recent advances in causal inference and machine learning target estimation of "causal contrast" functions such as CATE, which is sufficient for optimizing decisions and can adapt to potentially smoother structure. We develop a dynamic generalization of the R-learner (Nie et al., 2021; Lewis and Syrgkanis, 2021) for estimating and optimizing the difference of Q^{π} -functions, $Q^{\pi}(s, 1) - Q^{\pi}(s, 0)$ (which can be used to optimize multiple-valued actions). We leverage orthogonal estimation to improve convergence rates in the presence of slower nuisance estimation rates and prove consistency of policy optimization under a margin condition. The method can leverage black-box nuisance estimators of the Q-function and behavior policy to target estimation of a more structured Q-function contrast.

1 Introduction and Related Work

Learning optimal dynamic treatment rules, or sequential policies for taking actions, is important, although often only observational data is available. Many recent works in offline reinforcement learning develop methodology to evaluate and optimize sequential decision rules, without the ability to conduct online exploration.

An extensive literature on causal inference and machine learning establishes methodologies for learning causal contrasts, such as the conditional average treatment effect (CATE) (Wager and Athey, 2018; Foster and Syrgkanis, 2019; Künzel et al., 2019; Kennedy, 2020), which is sufficient for making optimal decisions. Methods that specifically estimate causal contrasts (such as the CATE), can better adapt to potentially smoother or more structured contrast functions, while methods that instead contrast estimates (by taking the difference of outcome regressions or Q functions) can not. Additionally, estimation of causal contrasts can be improved via orthogonalization or double machine learning (Kennedy, 2022; Chernozhukov et al., 2018). Estimating the causal contrast is both sufficient for optimal decisions and statistically favorable.

In this work, building on recent advances in heterogeneous treatment effect estimation, we focus on estimating analogous causal contrasts for offline reinforcement learning, namely $\tau_t^{\pi}(s) = Q_t^{\pi}(s, 1) - Q_t^{\pi}(s, 0)$, and natural multiple-action generalizations thereof.

The sequential setting offers even more motivation to target estimation of the contrast: additional structure can arise from sparsity patterns induced by the joint (in)dependence of rewards and transition dynamics on the (decompositions of) the state variable. A number of recent works point out this additional structure (Wang, Xiao, Zhu, and Stone, Wang et al.; Wang et al., 2022), for example of a certain transition-reward factorization, first studied by (Dietterich et al., 2018), that admits a sparse Q-function contrast (Pan and Schölkopf, 2023). (Zhou, 2024b) proposes a variant of the underlying blockwise pattern that also admits sparse optimal policies, but proposes a special modification of LASSO. Our method can adapt to such underlying sparsity structure when it is present in the Q-function contrast, in addition to other scenarios where the contrast is smoother than the Q-functions themselves.

The contributions of this work are as follows: We develop a dynamic generalization of the Rlearner (Nie and Wager, 2021; Lewis and Syrgkanis, 2021) for estimating the Q-function contrast. The method wraps around standard estimation procedures in offline reinforcement learning via a sequence of sequential loss minimization problems, which makes it appealingly practical. We show theoretical guarantees of improved convergence rates. Therefore our method leverages behavior policy estimation to improve estimation without suffering from unstable propensity weights. We illustrate benefits of adapting to structure in synthetic examples.

Related work. There is a large body of work on offline policy evaluation and optimization in offline reinforcement learning (Jin et al., 2021; Xie et al., 2021), including approaches that leverage importance sampling or introduce marginalized versions (Jiang and Li, 2016; Thomas et al., 2015; Kallus and Uehara, 2019a; Liu et al., 2018). For Markov decision processes, other papers study statistically semiparametrically efficient estimation of the policy value (Kallus and Uehara, 2019a,b; Xie et al., 2023; Kallus and Zhou, 2022; ??). The literature on dynamic treatment regimes (DTRs) studies a method called advantage learning (Schulte et al., 2014), although DTRs in general lack reward at every timestep, whereas we are particularly motivated on sparsity implications that arise jointly from reward and transition structure. In particular, beyond policy value estimation, we aim to recover the entire contrast function. Prior works that consider policy optimization under a restricted function class can require estimating difficult policy-dependent nuisance functions; we maximize the advantage function without further restricting functional complexity, which requires re-estimating nuisance functions at every timestep (but not every *iteration* of policy optimization, as in (Xu et al., 2021)).

At a high level, our method is similar to the dynamic DR learner studied in (Lewis and Syrgkanis, 2021) in that we extend the R-learner identification approach to a sequential setting, although the estimand is quite different. In particular, they only consider heterogeneity based on a fixed *initial* state, dynamic treatment regimes with terminal rewards, generalize structural nested-mean models (SNMMs) by estimating "blip-to-zero" functions. Consequently, our analysis is similar at a high level. Overall, the most closely related work when it comes to estimating contrast functionals in reinforcement learning is that of (Shi et al., 2022), which derives a pseudo-outcome for estimating the *Q*-function contrast in the infinite horizon setting. We share the same estimand, but in the finite-horizon setting. The estimation strategy is quite different. Crucial differences include: we directly generalize the residualized learning (R-learner) approach, and we work in finite-horizons with the propensity score rather than the hard-to-estimate stationary distribution density ratio (Liu et al., 2018). Note that the (single-stage) R-learner loss function is an overlap-weighted (Li et al., 2019) regression against the doubly-robust score (DR-learner (Kennedy, 2020)). (See (Morzywolek et al., 2023; Chernozhukov et al., 2024) for more discussion).

Advantage functions are a contrast functional widely studied in classical RL (less often in the offline setting) and dynamic treatment regimes (Neumann and Peters, 2008; mur, 2003). However, *which* contrast it evaluates is policy-dependent, and it requires further algorithmic development for

policy learning, unlike our approach that is simply greedy with respect to difference-of-Q functions. (Pan and Schölkopf, 2023) note the analogy with causal contrast estimation and derive a Q-function independent estimator, but in the online setting. (Nie et al., 2021) studies OPE for advantage functions in a special case of optimal stopping. We do make a margin assumption to relate convergence of Q-function contrasts to policy value convergence, analogous to (Shi et al., 2022). (Hu et al., 2024) studies consequences of the margin assumption for fitted-Q-iteration with a tighter analysis. Our approach is better suited for settings with highly structured difference-of-Qs since we introduce auxiliary estimation at every timestep.

2 Method

Problem Setup: We consider a finite-horizon Markov Decision Process on the full-information state space comprised of a tuple $\mathcal{M} = (\mathcal{S}, \mathcal{A}, r, P, \gamma, T)$ of states, actions, reward function r(s, a), transition probability matrix P, discount factor $\gamma < 1$, and time horizon of T steps, where $t = 1, \ldots, T$. We let the state spaces $\mathcal{S} \subseteq \mathbb{R}^d$ be continuous, and assume the action space \mathcal{A} is finite. A policy $\pi : \mathcal{S} \mapsto \Delta(\mathcal{A})$ maps from the state space to a distribution over actions, where $\Delta(\cdot)$ is the set of distributions over (\cdot) , and $\pi(a \mid s)$ is the probability of taking action a in state s. (At times we overload notation a bit so that $\pi(s) \in \mathcal{A}$ indicates the action random variable under π evaluated at state s).

The value function is $V_t^{\pi}(s) = \mathbb{E}_{\pi}[\sum_{t'=t}^T \gamma^{t'-t} R_{t'} \mid s]$, where \mathbb{E}_{π} denotes expectation under the joint distribution induced by the MDP \mathcal{M} running policy π . The state-action value function, or Q function is $Q_t^{\pi}(s) = \mathbb{E}_{\pi}[\sum_{t'=t}^T \gamma R_{t'} \mid s, a]$. These satisfy the Bellman operator, e.g. $Q_t^{\pi}(s, a) = r(s, a) + \gamma \mathbb{E}[V_{t+1}^{\pi}(s_{t+1}) \mid s, a]$. The optimal value and q-functions are denoted V^*, Q^* under the optimal policy. We focus on estimating the difference of Q-functions (each under the same policy), $\tau_t^{\pi}(s) = Q_t^{\pi}(s, 1) - Q_t^{\pi}(s, 0)$. (This differs slightly from the conventional advantage function studied in RL, defined as $Q^{\pi}(s, a) - V^{\pi}(s)$, where the contrast being estimated depends on the policy). We focus on the offline reinforcement learning setting where we have access to a dataset of n offline trajectories, $\mathcal{D} = \{(S_t^{(i)}, A_t^{(i)}, R_t^{(i)} S_{t+1}^{(i)})_{t=1}^T\}_{i=1}^n$, where actions were taken according to some behavior policy π^b . We state some notational conventions. For some generic function f we define the norm $\|f\|_u := \mathbb{E}[\|\psi(X)\|_u]^{1/u}$. In the context of estimation (rather than discussing identification), we denote the true population functions with a \circ subscript, i.e. $\tau_t^{\pi,\circ}$ and so on.

Policy evaluation: Identification. First we overview deriving the estimating moments of our approach. The arguments are broadly a generalization of the so-called residualized R-learner (Nie and Wager, 2021); (Lewis and Syrgkanis, 2021) considers a similar generalization for structural nested mean models without state-dependent heterogeneity. For the purposes of this section, we discuss the true population Q, m, τ functions without notational decoration, which we introduce later on when we discuss estimation.

Denote $\underline{\pi}_{t+1} = \pi_{t+1:T} := \{\pi_{t+1}, \ldots, \pi_T\}$ for brevity. Then $Q_t^{\underline{\pi}_{t+1}}$ indicates the Q_t function under policy π . For brevity, we further abbreviate $Q_t^{\underline{\pi}} := Q_t^{\underline{\pi}_{t+1}}$ when this is unambiguous. We seek to estimate:

$$\tau_t^{\pi}(S_t) = Q_t^{\pi}(S_t, 1) - Q_t^{\pi}(S_t, 0), \tag{1}$$

Note that the Q-function satisfies: $Q_t^{\pi}(S_t, A_t) = \mathbb{E}[R_t + \gamma Q_{t+1}^{\pi}(S_{t+1}, A_{t+1}) \mid S_t, A_t]$. Define

$$\epsilon_t^{(i)}(A_t) = R_t + \gamma Q_{t+1}^{\pi}(S_{t+1}, A_{t+1}) - \{Q_t^{\pi}(S_t, 0) + A_t \tau_t^{\pi}(S_t)\}$$

Under sequential unconfoundedness and Markovian properties, we obtain the conditional moment:

$$\mathbb{E}[\epsilon_t^{(i)}(A_t) \mid S_t, A_t] = 0.$$
⁽²⁾

Define the analogue to the marginal outcome function, which is the state-conditional value function under the behavior policy: $m^{\circ,\pi}(S_t) = V_t^{\pi^b_t, \underline{\pi}_{t+1}} = \mathbb{E}_{\pi^b_t}[R_t + \gamma Q_{t+1}^{\pi}(S_{t+1}, A_{t+1}) \mid S_t]$. Under sequential unconfoundedness,

$$R_t + \gamma Q_{t+1}^{\pi}(S_{t+1}, A_{t+1}) + \epsilon(A_t) = Q_t^{\pi}(S_t, 0) + A_t \tau_t^{\pi}(S_t)$$
$$m_t^{\pi}(S_t) = \mathbb{E}_{\pi^b}[R_t + \gamma Q_{t+1}^{\pi}(S_{t+1}, A_{t+1}) \mid S_t] = Q_t^{\pi}(S_t, 0) + \pi^b(1 \mid S_t)\tau_t^{\pi}(S_t)$$

Hence,

$$R_t + \gamma Q_{t+1}^{\pi}(S_{t+1}, A_{t+1}) - m_t^{\pi}(S_t) = (A - \pi^b(1 \mid S_t))\tau_t^{\pi}(S_t) + \epsilon_t(A_t).$$
(3)

Extension to multiple actions. So far we presented the method with $\mathcal{A} \in \{0, 1\}$ for simplicity, but all methods in this paper will extend to the multi-action case. For multiple actions, fix a choice $a_0 \in \mathcal{A}$, and for $a \in \mathcal{A} \setminus a_0$, define $\tau_t^{\pi}(s, a) \coloneqq \tau_{a,t}^{\pi}(s) = Q_t^{\pi}(s, a) - Q_t^{\pi}(s, a_0)$. For $k \in |\mathcal{A}|$, let $\pi^b(k \mid S_t) = P(A_t = k \mid S_t)$. Redefine $\epsilon_t^{(i)}(A_t) = R_t + \gamma Q_{t+1}^{\pi}(S_{t+1}, A_{t+1}) - \{Q_t^{\pi}(S_t, 0) + \mathbb{I}[A_t = a]\tau_t^{\pi}(S_t)\}$. Then the equivalent of eq. (3) is that $\tau_{a,t}^{\pi}(S_t)$ satisfies:

$$R_t + \gamma Q_{t+1}^{\pi}(S_{t+1}, A_{t+1}) - m_t^{\pi}(S_t) = (\mathbb{I}[A_t = a] - \pi^b(a \mid S_t))\tau_{a,t}^{\pi}(S_t) + \epsilon_t(A_t)$$

The loss function. This motivates the approach based on (penalized) empirical risk minimization:

$$\tau_t(\cdot) \in \operatorname*{argmin}_{\tau} \left\{ \mathbb{E}\left[\left(\{ R_t + \gamma Q_{t+1}^{\pi}(S_{t+1}, A_{t+1}) - m_t^{\pi}(S_t) \} - \gamma \{ A - \pi_t^b(1 \mid S_t) \} \cdot \tau_t^{\pi}(S_t) \right)^2 \right] \right\}$$
(4)

Again, so far we have discussed identification assuming the true Q, m, π^b functions, etc. Next we discuss feasible estimation, and outside of this section we refer to the population-level true nuisance functions as $Q^{\pi,\circ}, m^{\pi,\circ}, \pi^{b,\circ}, \tau^{\pi,\circ}$.

Feasible estimation. In practice, the nuisance functions need to be estimated. We introduce some notation before defining the full estimation algorithm. Let the nuisance vector be denoted $\eta = [\{Q_t^{\pi}\}_{t=1}^T, \{m_t^{\pi}\}_{t=1}^T, \{\pi_t^b\}_{t=1}^T]$. The fitted advantage R-learner for evaluation is a feasible version of the sequential loss minimization approach implied by eq. (4): we describe the algorithm in Algorithm 1. Given an evaluation policy π^e , first fit the nuisance functions: a pilot estimate of the Q function and the behavior policy. Then, evaluate the loss function in eq. (4) and estimate τ_t .

Estimating the nuisance functions. The Q function nuisance can be estimated with a variety of approaches such as fitted-Q-evaluation (Le et al., 2019; Chakraborty and Murphy, 2014; Duan et al., 2021), other approaches in offline reinforcement learning, minimum-distance estimation for conditional moment restrictions/GMM (Kallus and Uehara, 2019a), or the finitehorizon analogous version of DR-learner suggested in (Shi et al., 2022). Estimating the behavior policy is a classic probabilistic classification or multi-class classification problem. Sometimes the offline trajectories might arise from a system with known exploration probabilities, so that the behavior policy might be known.

Cross-fitting. We also introduce cross-fitting, which will differ slightly between policy evaluation and optimization: splitting the dataset \mathcal{D} into K many folds (preserving trajectories, i.e.

Algorithm 1 Dynamic Residualized Difference-of-Q-Evaluation

- 1: Given: π^e , evaluation policy; and for sample splitting, partition of \mathcal{D} into K folds, $\{\mathcal{D}_k\}_{k=1}^K$.
- 2: On \mathcal{D}_k , estimate $\hat{Q}^{\pi^e,k}(s,a)$ and behavior policy $\hat{\pi}_t^{b,k}(s)$. Evaluate the value function via integrating/summing \hat{Q} over the empirical distribution of actions, $a \sim \pi^b$, observed in the data, so that

$$\hat{m}(s) = \mathbb{E}_{\pi_t^b}[R_t + \gamma \hat{Q}_{t+1}^{\pi^e}(S_{t+1}, A_{t+1}) \mid S_t = s].$$

3: for timestep t = T, ..., 1 do 4: $\hat{\tau}_t \in \operatorname{argmin}_{\tau} \left\{ \sum_{k=1}^K \sum_{i \in \mathcal{D}_k} \left(R_t^{(i)} + \gamma \hat{Q}_{t+1}^{\pi,-k}(S_{t+1}^{(i)}, A_{t+1}^{(i)}) - \hat{m}_t^{\pi,-k}(S_t^{(i)}) - \{A_t^{(i)} - \hat{\pi}_t^{b,-k}(1 \mid S_t^{(i)})\}\tau_t(S_t^{(i)}) \right)^2 \right\}$ 5: end for

randomizing over trajectory index i), and learning the nuisance function η^{-k} on $\{\mathcal{D}_{k'}\}_{k'\in\{[K]\setminus k\}}$. (In scenarios with possible confusion we denote the nuisance function $\eta^{(-k)}$ instead. In evaluating the loss-function, we evaluate the nuisance function η^{-k} using data from the held-out kth fold. Given the cross-fitting procedure, we introduce the empirical squared loss function:

$$\hat{\mathcal{L}}_{t}(\tau,\eta) = \sum_{k=1}^{K} \sum_{i \in \mathcal{D}_{k}} \left(R_{t}^{(i)} + \gamma \hat{Q}_{t+1}^{\pi,-k}(S_{t+1}^{(i)}, A_{t+1}^{(i)}) - \hat{m}_{t}^{\pi,-k}(S_{t}^{(i)}) - \{A_{t}^{(i)} - \hat{\pi}_{t}^{b,-k}(1 \mid S_{t}^{(i)})\}\tau_{t}(S_{t}^{(i)}) \right)^{2}$$

and let the population loss function $\mathcal{L}_t(\tau, \eta)$ be the population expectation of the above.

Finally, note that the expectation of the empirical squared loss will not in general be an unbiased estimate of the true squared error, due to the squaring and the expectation over the next transition. Nonetheless, as shown in other works studying fitted Q evaluation/iteration, the resulting Q function (contrast function here) can still be useful for policy optimization.

Lemma 1 (Proxy Sequential R-learner loss: Excess Variance).

$$\mathbb{E}[\hat{\mathcal{L}}_t(\tau^{\pi},\eta)] - \mathcal{L}_t(\tau^{\pi},\eta) = \operatorname{Var}[\max_{a'} Q^{\pi}\left(S_{t+1},a'\right) \mid \pi_t^b]$$

Policy optimization. The sequential loss minimization approach also admits an policy optimization procedure. The policy at every timestep is greedy with respect to the estimated τ . We describe the algorithm in Algorithm 2. We use a slightly different cross-fitting approach for policy optimization. We introduce an additional fold, upon which we alternate estimation of $\hat{\tau}_t$. So, overall we use three folds: one for estimating nuisance functions η , and the other two for estimating $\hat{\tau}_t^{\hat{\pi}_{t+1}}$. On these two other folds, between every timestep, we alternate estimation of $\hat{\tau}_t$ on one of them, in order to break dependence between the estimated optimal forwards policy $\hat{\pi}_{t+1}$ and $\hat{\tau}_t$ (and therefore the greedy policy $\hat{\pi}_t$).

3 Analysis

Our analysis generally proceeds under the following assumptions.

Assumption 1 (Independent and identically distributed trajectories). We assume that the data was collected under a stationary behavior policy, i.e. not adaptively collected from a policy learning over time.

Algorithm 2 Dynamic Residualized Difference-of-Q Optimization

- 1: Given: Partition of \mathcal{D} into 3 folds, $\{\mathcal{D}_k\}_{k=1}^3$.
- 2: Estimate $\hat{\pi}_t^b$ on \mathcal{D}_1 .
- 3: At time T: Set $\hat{Q}_T(s,a) = 0$. Estimate $m_T = \mathbb{E}_{\pi^b}[R_T \mid S_T]$ on \mathcal{D}_1 and $\hat{\tau}_T$ on $\mathcal{D}_{k(T)}$, where k(t) = 2 if t is odd and k(t) = 3 if t is even. Optimize $\hat{\pi}_T(s) \in \arg \max \tau_T(s)$.
- 4: for timestep t = T 1, ..., 1 do
- 5: Estimate $Q^{\hat{\pi}_{t+1}}$ on \mathcal{D}_1 . Evaluate $m_t^{\hat{\pi}_{t+1}}$. Estimate $\hat{\tau}_t^{\hat{\pi}_{t+1}}$ on $\mathcal{D}_{k(t)}$ by minimizing the empirical loss:

$$\hat{\tau}_t(\cdot) \in \operatorname{argmin}_{\tau} \sum_{i \in \mathcal{D}_{k(t)}} \left(R_t^{(i)} + \gamma \hat{Q}_{t+1}^{\pi,(1)}(S_{t+1}^{(i)}, A_{t+1}^{(i)}) - \hat{m}_t^{\pi,(1)}(S_t^{(i)}) - \{A_t^{(i)} - \hat{\pi}_t^{b,(1)}(1 \mid S_t^{(i)})\}\tau_t(S_t^{(i)}) \right)^2$$

6: Optimize $\hat{\pi}_t(s) \in \arg \max \hat{\tau}_t^{\hat{\pi}_{t+1}}(s)$. 7: end for

Assumption 2 (Sequential unconfoundedness). $r(A) \perp A \mid S_t \text{ and } S_{t+1}(a) \perp A_t \mid S_t$

This assumption posits that the state space is sufficient for identification. It is satisfied by design if actions were taken by a previous known behavior policy, i.e. randome xplanation.

Assumption 3 (Boundedness). $V_t \leq B_V, \tau \leq B_\tau$

Assumption 4 (Bounded transition density). Transitions have bounded density: $P(s' | s, a) \leq c$. Let $d_{\pi}(s)$ denote the marginal state distribution under policy π . Assume that $d_{\pi_t^b}(s) < c$, for $t = 1, \ldots, T$.

Next we establish convergence rates for $\hat{\tau}^{\pi}$, depending on convergence rates of the nuisance functions. Broadly we follow the analysis of (Foster and Syrgkanis, 2019; Lewis and Syrgkanis, 2021) for orthogonal statistical learning. The analysis considers some $\hat{\tau}$ with small excess risk relative to the projection onto the function class, i.e. as might arise from an optimization algorithm with some approximation error.

For a fixed evaluation policy π^e , define the projection of the true advantage function onto Ψ^n ,

$$\tau_t^{\pi^e,n} = \arg\inf_{\tau_t \in \Psi_t^n} \|\tau_t - \tau_t^{\circ,\pi^e}\|_2.$$
(5)

For a fixed evaluation policy π^e , define the error of some estimate $\hat{\tau}_t^{\pi^e}$ to projection onto the function class:

$$\nu_t^{\pi^e} = \hat{\tau}_t^{\pi^e} - \tau_t^{n,\pi^e}.$$

Theorem 1 (Policy evaluation). Suppose $\{\sup_{s,t} \mathbb{E}[(A_t - \pi_t^b)(A_t - \pi_t^b) | s]\} \leq C$ and Assumptions 1 to 4. Consider a fixed evaluation policy π^e . Consider any estimation algorithm that produces an estimate $\hat{\tau}^{\pi^e} = (\tau_1^{\pi^e}, \ldots, \tau_T^{\pi^e})$, with small plug-in excess risk at every t, with respect to any generic candidate $\tilde{\tau}^{\pi^e}$, at some nuisance estimate $\hat{\eta}$, i.e.,

$$\mathcal{L}_{D,t}(\hat{\tau}_t^{\pi^e};\hat{\eta}) - \mathcal{L}_{D,t}(\tilde{\tau}_t^{\pi^e};\hat{\eta}) \le \epsilon(\tau_t^n,\hat{\eta}).$$

Let ρ_t denote product error terms:

$$\rho_t^{\pi^e}(\hat{\eta}) = B_\tau^2 \| (\hat{\pi}_t^b - \pi_t^{b,\circ})^2 \|_u + B_\tau \| (\hat{\pi}_t^b - \pi_t^{b,\circ}) (\hat{m}_t^{\pi^e} - m_t^{\pi^e,\circ}) \|_u \\ + \gamma (B_\tau \| (\hat{\pi}_t^b - \pi_t^{b,\circ}) (\hat{Q}_{t+1}^{\pi^e} - Q_{t+1}^{\pi^e,\circ}) \|_u + \| (\hat{m}_t^{\pi^e} - m_t^{\pi^e,\circ}) (\hat{Q}_{t+1}^{\pi^e} - Q_{t+1}^{\pi^e,\circ}) \|_u).$$
(6)

Then, for $\sigma > 0$, and $u^{-1} + \overline{u}^{-1} = 1$,

$$\frac{\lambda}{2} \|\nu_t^{\pi^e}\|_2^2 - \frac{\sigma}{4} \|\nu_t^{\pi^e}\|_{\overline{u}}^2 \le \epsilon(\hat{\tau}_t^{\pi^e}, \hat{\eta}) + \frac{2}{\sigma} \left(\|(\tau^{\pi^e, \circ} - \tau_t^{\pi^e, n})\|_u^2 + \rho_t^{\pi^e}(\hat{\eta})^2 \right).$$

In the above theorem, $\epsilon(\hat{\tau}_t^{\pi^e}, \hat{\eta})$ is the excess risk of the empirically optimal solution. Note that in our setting, this excess risk will be an approximation error incurred from the proxy loss issue described in Lemma 1. The bias term is $\|(\tau^{\pi^e,\circ}-\tau_t^{\pi^e,n})\|_u^2$, which describes the model misspecification bias of the function class parametrizing Q-function contrasts, Ψ . The product error terms $\rho_t^{\pi^e}(\hat{\eta})$ highlight the reduced dependence on individual nuisance error rates. We will instantiate the previous generic theorem for the projection onto Ψ^n , defined in Equation (5), also accounting for the sample splitting. We will state the results with *local Rademacher complexity*, which we now introduce. For generic 1-bounded functions f in a function space $f \in \mathcal{F}, f \in [-1, 1]$, the local Rademacher complexity is defined as follows:

$$\mathscr{R}_{n}(\mathcal{F};\delta) = \mathbb{E}_{\epsilon_{1:n},X_{1:n}} \left[\sup_{f \in \mathcal{F}: \|f\|_{2} \le \delta} \frac{1}{n} \sum_{i=1}^{n} \epsilon_{i} f(X_{i}) \right]$$

The critical radius δ^2 more tightly quantifies the statistical complexity of a function class, and is any solution to the so-called *basic inequality*, $\mathscr{R}_n(\mathcal{F}; \delta) \leq \delta^2$. The star hull of a generic function class \mathcal{F} is defined as $\operatorname{star}(\mathcal{F}) = \{cf : f \in \mathcal{F}, c \in [0, 1]\}$. Bounds on the critical radius of common function classes like linear and polynomial models, deep neural networks, etc. can be found in standard references on statistical learning theory, e.g. (Wainwright, 2019). We can obtain mean-squared error rates for policy evaluation via specializing Theorem 1 to the 2-norm and leveraging results from (Foster and Syrgkanis, 2019)

Theorem 2 (MSE rates for policy evaluation). Suppose $\{\sup_{s,t} \mathbb{E}[(A_t - \pi_t^b)(A_t - \pi_t^b) | s]\} \leq C$ and Assumptions 1 to 4. Consider a fixed policy π^e . Suppose each of $\mathbb{E}[\|(\hat{\pi}_t^b - \pi_t^{b,\circ})\|_2^2]$, $\mathbb{E}[\|(\hat{\pi}_t^b - \pi_t^{b,\circ})(\hat{Q}_{t+1}^{\pi^e} - Q_{t+1}^{\pi^e,\circ})\|_2^2]$, and $\mathbb{E}[\|(\hat{m}_t^{\pi^e} - m_t^{\pi^e,\circ})(\hat{Q}_{t+1}^{\pi^e} - Q_{t+1}^{\pi^e,\circ})\|_2^2]$ are of order $O(\delta_{n/2}^2 + \|\tau_t^{\pi^e,\circ} - \tau_t^{\pi^e,n}\|_2^2)$. Then

$$\mathbb{E}[\|\hat{\tau}_t^{\pi^e} - \tau_t^{\pi^e, \circ}\|_2^2] = O\left(\delta_{n/2}^2 + \|\tau_t^{\pi^e, \circ} - \tau_t^{\pi^e, n}\|_2^2\right)$$

Working with the orthogonalized estimate results in the weaker product-error rate requirements included above. However, our estimating moments do include the Q function nuisances, and quarter-root rates are required for estimating both the Q and π^b functions.

Policy optimization. Convergence of τ_t implies convergence in policy value. We quantify this with the *margin* assumption, which is a low-noise condition that quantifies the gap between regions of different optimal action (Tsybakov, 2004). It is commonly invoked to relate estimation error of plug-in quantities to decision regions, in this case the difference-of-Q functions to convergence of optimal decision values.

Assumption 5 (Margin (Tsybakov, 2004)). Assume there exist some constants $\alpha, \delta_0 > 0$ such that

$$P\left(\max_{a} Q^{*}(s, a) - \max_{a' \in \mathcal{A} \setminus \arg\max_{a} Q^{*}(s, a)} Q^{*}(s, a') \leq \epsilon\right) = O\left(\varepsilon^{\alpha}\right)$$

The probability density in Assumption 5 is evaluated with respect to Lebesgue measure over the state space.

Lemma 2 (Advantage estimation error to policy value via margin.). Suppose Assumptions 2, 4 and 5 (margin assumption holds with α). Suppose that with high probability $\geq 1 - n^{-\kappa}$ for any finite $\kappa > 0$, the following sup-norm convergence holds with some rate $b_* > 0$,

$$\begin{split} \sup_{s \in \mathcal{S}, a \in \mathcal{A}} \left| \hat{\tau}_{t}^{\underline{n}_{t+1}}(s) - \tau_{t}^{\underline{n}_{t+1}, 0}(s) \right| &\leq Kn^{-b*}, \\ then \left| \mathbb{E}[V_{t}^{*}(S_{t}) - V_{t}^{\hat{\pi}_{\tau}}(S_{t})] \right| &\leq \frac{(1 - \gamma^{T-t})}{1 - \gamma} cK^{2} n^{-b*(1+\alpha)} + O(n^{-\kappa}), \\ and \left\{ \int \left(Q_{t}^{*}(s, \pi^{*}(s)) - Q_{t}^{*}(s, \hat{\pi}_{\tau}) \right)^{2} ds \right\}^{1/2} &\leq \frac{(1 - \gamma^{T-t})}{1 - \gamma} cK^{2} n^{-b*(1+\alpha)} + O(n^{-\kappa}). \\ Else, assume that \left(\mathbb{E} \int_{s \in \mathbb{S}} |\hat{\tau}_{t}^{n}(s) - \tau_{t}^{\circ}(s)|^{2} ds \right)^{1/2} &\leq K \left(n^{-b_{*}} \right), \text{ for some rate } b_{*} > 0. \ Then \\ \end{bmatrix}$$

$$\left| \mathbb{E}[V_t^*(S_t) - V_t^{\hat{\pi}_{\hat{\tau}}}(S_t)] \right| = O\left(n^{-b_*\left(\frac{2+2\alpha}{2+\alpha}\right)} \right), \text{ and } \left\{ \int \left(Q_t^*(s, \pi^*(s)) - Q_t^*(s, \hat{\pi}_{\hat{\tau}}) \right)^2 ds \right\}^{1/2} = O\left(n^{-b_*\left(\frac{2+2\alpha}{2+\alpha}\right)} \right)$$

Next we study policy optimization. Studying the convergence of policy optimization requires conditions on convergence of advantage functions from previous steps of the algorithm.

Theorem 3 (Policy optimization bound). Suppose Assumptions 1 to 4. Further, suppose that Q° satisfies Assumption 5 (margin) with $\alpha > 0$. Suppose the product error rate conditions hold for each t for data-optimal policies evaluated along the algorithm, i.e. for each t, for $\hat{\pi}_{t+1}$, each of $\mathbb{E}[\|(\hat{\pi}_t^b - \pi_t^{b,\circ})\|_2^2]$, $\mathbb{E}[\|(\hat{\pi}_t^b - \pi_t^{b,\circ})(\hat{m}_t^{\hat{\pi}_{t+1}} - m_t^{\circ,\hat{\pi}_{t+1}})\|_2^2]$, $\mathbb{E}[\|(\hat{\pi}_t^b - \pi_t^{b,\circ})(\hat{Q}_{t+1}^{\hat{\pi}_{t+2}} - Q_{t+1}^{\circ,\hat{\pi}_{t+2}})\|_2^2]$, and $\mathbb{E}[\|(\hat{m}_t - m_t^{\circ})(\hat{Q}_{t+1}^{\hat{\pi}_{t+2}} - Q_{t+1}^{\circ,\hat{\pi}_{t+2}})\|_2^2]$ are of order $O(\delta_{n/2}^2 + \|\pi_t^{\hat{\pi}_{t+1},\circ} - \pi_t^{\hat{\pi}_{t+1},n}\|_2^2)$. Suppose that for $\hat{\pi}_t$, under the above assumptions, Theorem 2 holds, and the critical radius $\delta_{n/2}$

Suppose that for $\hat{\pi}_t$, under the above assumptions, Theorem 2 holds, and the critical radius $\delta_{n/2}$ and for time t, function class specification error $\|\tau_t^{\hat{\pi}_{t+1},\circ} - \tau_t^{\hat{\pi}_{t+1},n}\|_2$ satisfy the root-mean-squarederror rate conditions: $\rho_t^{(c)}, \rho_t^{(\Psi)}$

$$\delta_{n/2}^2 = K_r^2 n^{-2\rho_t^{(c)}}, \qquad \|\tau_t^{\hat{\pi}_{t+1},\circ} - \tau_t^{\hat{\pi}_{t+1},n}\|_2^2 = K_{\Psi}^2 n^{-2\rho_t^{(\Psi)}},$$

Further define for a generic $t, \ \overline{\rho}_{\geq t}^{(\cdot)} = \min_{t' \geq t} \{\rho_{t'}^{(\cdot)}\}, \ for \ (\cdot) \in \{(c), (\Psi)\}.$ Then, with high probability $\geq n^{-\kappa}, \|\hat{\tau}_t^{\hat{\pi}_{t+1}} - \tau_t^{\circ, \underline{\pi}_{t+1}^*}\| \leq O(\delta_{n/2} + \|\tau_t^{\circ, \underline{\pi}_{t+1}} - \tau_t^{n, \underline{\hat{\pi}}_{t+1}}\|_2) + Kn^{-\mathcal{R}_t}.$ where

$$\mathcal{R}_{k} = \min\left(\rho_{k+1}^{(c)} \cdot \frac{2+2\alpha}{2+\alpha}, \ \rho_{k+1}^{(\Psi)} \cdot \frac{2+2\alpha}{2+\alpha}, \ \left\{\min_{k' \ge k+1}(\rho_{k'}^{(c)}, \rho_{k'}^{(\Psi)})\right\} \cdot \frac{2+2\alpha^{T-k'}}{2+\alpha}\right).$$

Further suppose that $\alpha > 0$ and that for $t' \ge t$, we have that $\rho_t^{(\cdot)} \le \rho_{t'}^{(\cdot)}$, for $(\cdot) \in \{(c), (\Psi)\}$, i.e. the estimation error rate is nonincreasing over time. Then,

$$\|\hat{\tau}_t^{\hat{\underline{\pi}}_{t+1}} - \tau_t^{\circ,\underline{\pi}_{t+1}^*}\| \le O(\delta_{n/2} + \|\tau_t^{\circ,\underline{\hat{\pi}}_{t+1}} - \tau_t^{n,\underline{\hat{\pi}}_{t+1}}\|_2),\tag{7}$$

and

$$\left| \mathbb{E}[V_1^{\pi^*}(S_1) - V_1^{\hat{\pi}_{\hat{\tau}}}(S_1)] \right| = O(n^{-\min\{\overline{\rho}_{\geq 1}^{(c)}, \overline{\rho}_{\geq 1}^{(\Psi)}\}\frac{2+2\alpha}{2+\alpha}})$$

Our method introduces auxiliary estimation at every timestep, so that the exponentiated terms are higher-order relative to the difference-of-Q one-step estimation error at every timestep. Note that (Hu et al., 2024) also establishes margin constants for linear and tabular MDPs.

Table 1: Validation: 1d example, linear FQE vs orthogonal τ estimation. Entries: Mean-squared error of estimated τ function, mean + 1.96 standard errors.

Method (n)	$5.0\cdot 10^1$	$1.3 \cdot 10^{3}$	$2.5 \cdot 10^3$	$3.8\cdot 10^3$	$5.0\cdot 10^3$
FQE	$4 \cdot 10^{-2} \pm 2 \cdot 10^{-2}$	$2 \cdot 10^{-3} \pm 10^{-3}$	$10^{-3} \pm 10^{-4}$	$10^{-3} \pm 10^{-4}$	$4 \cdot 10^{-4} \pm 10^{-4}$
OrthDiff-Q	$6 \cdot 10^{-2} \pm 3 \cdot 10^{-2}$	$3 \cdot 10^{-3} \pm 10^{-3}$	$2 \cdot 10^{-3} \pm 10^{-3}$	$2 \cdot 10^{-3} \pm 10^{-3}$	$2 \cdot 10^{-3} \pm 10^{-4}$
			-	-	



Figure 1: (Left) Reward-relevant/irrelevant factored dynamics of (Zhou, 2024b). (Right) Exogenous-Endogenous MDP model of (Dietterich et al., 2018). The dotted line from a_t to $s_{t+1}^{\rho_c}$ indicates the presence or absence is permitted in the model.

4 Experiments

1d validation. In a very small 1d toy example (Sec 5.1, (Kallus and Uehara, 2019a)) we validate our method. See Appendix C of the appendix for more details.

Adapting to structure in $\tau(s)$. Recent research highlights the joint implications of blockwise conditional independence properties in RL, where some components are considered "exogenous" or irrelevant to rewards and actions (Wang et al., 2022; Wang, Xiao, Zhu, and Stone, Wang et al.; Dietterich et al., 2018). Most papers employ a model-based approach to filter out irrelevant factors with somewhat-intractable mutual information/black-box modified VAEs(Wang et al., 2022). (An exception is (Efroni et al., 2022) which considers sparsity in partially controllable linear models, but without orthogonality). Pretesting approaches such as (Wang, Xiao, Zhu, and Stone, Wang et al.) warrant caution due to poor statistical properties. Additionally, different underlying structures may lead to the same sparsity pattern in the advantage function. (Pan and Schölkopf, 2023) studies whether advantage function estimation can naturally recover the endogenous component under the model of (Dietterich et al., 2018), in an online RL setting. In a similar spirit, we assess the benefits of targeting estimation of the difference-of-Qs in a set of data-generating processes closely related to specific structural models proposed in the literature ((Zhou, 2024b; Dietterich et al., 2018)). We find that orthogonal causal contrast estimation is robust under noisy nuisance functions, as confirmed by our theory, and that it can adapt to a variety of structures.

First we describe the modified Reward-Filtered DGP (left, section 4) of (Zhou, 2024b). In the DGP, $|\mathcal{S}| = 50$ though the first $|\rho| = 10$ dimensions are the reward-relevant sparse component, where ρ is the indicator vector of the sparse support, and $\mathcal{A} = \{0, 1\}$. The reward and states evolve according to $r_t(s, a) = \beta^{\top} \phi_t(s, a) + a * (s_1 + s_2)/2 + \epsilon_r$, $s_{t+1}(s, a) = M_a s + \epsilon_s$, satisfying the graphical restrictions of Section 4. Therefore the transition matrices satisfy the blockwise form



Figure 2: Adapting to structure. Simpler setting where $E[M_1 - M_0] = 0 \cdot I$.



Figure 3: Adapting to structure. Interacted setting where $E[M_1 - M_0] = 0.1 \cdot I$.

 $M_a = \begin{bmatrix} M_a^{\rho \to \rho} & 0\\ M_a^{\rho \to \rho_c} & M_a^{\rho_c \to \rho_c} \end{bmatrix}$, we generate the coefficient matrices M_0, M_1 with independent normal random variables $\sim N(0.2, 1)$. The nonzero mean ensures the beta-min condition. We normalize $M_a^{\rho \to \rho}$ to have spectral radius 1; recovering the sparse component is stable while including distracting dimensions destabilizes. The zero-mean noise terms are normally distributed with standard deviations $\sigma_s = 0.4, \sigma_r = 0.6$. In the estimation, we let $\phi(s, a) = \langle s, s\dot{a}, 1 \rangle$ be the interacted state-action space, i.e. equivalent to fitting a q function separately for every action. The behavior policy is a mixture of logistic (with coefficients generated $\sim N(0, 0.3)$) and 20% probability of uniform random sampling. The evaluation policy is logistic (with coefficients $\sim Unif[-0.5, 0.5]$. (The coefficient vector is fixed within each plot via the random seed).

In Section 4 we compare against a strong set of baselines. In blue is FQE-TL, i.e. naive fitted-Qevaluation with thresholded Lasso (Zhou, 2010, 2009; Belloni and Chernozhukov, 2013). In dotted cyan is FQE-RF, the reward-filtered method of (Zhou, 2024b). Note that with the state-action interacted basis, it is essentially a "T-learner" estimate of reward-thresholded Lasso Q functions of the difference of Q-functions (in the parlance of CATE meta-learners (Künzel et al., 2019)), a very strong baseline when the Q functions are indeed linear. Next we have three variants of our framework: in dot-dashed pink τ -CV, orthogonal difference of Q estimation with loss-function crossvalidated l1 norm regularization on τ , in dotted green τ -TL which uses reward-based thresholding to estimate τ on the recovered support. We also investigate semi-synthetic settings with noisy nuisance functions by adding $N(0, n^{-1/4})$ noise to nuisance function predictions in dotted-red τ -TL- $\hat{\eta}_{\epsilon}$, which also includes sample splitting. For comparison to illustrate a setting with slow nuisance function convergence, we also include in dot-dashed purple FQE-TL- $\hat{\eta}_{\epsilon}$, which adds the $n^{-1/4}$ noise to the first naive FQE baseline. For our methods, we solve the loss function exactly with CVXPY (and l1 norm regularization).

We describe the results left to right. We display the median over 100 replications (fixing the coefficient matrices and vectors, etc. with the same random seed). The y-axis is the normalized MSE (we divide by the square of the range of the true difference of Q_s), and the x axis is the number of episodes from 100 to 1000 on a log-scale. First on the left, we consider the previously mentioned reward-filtered DGP. Obviously the tailored method of (Zhou, 2024a) that was designed for these graphical restrictions does well; it is also well-specified. Nonetheless we see that our methods with thresholded LASSO do well, although are slower because they are not tailored for this graphical restriction. We do see that orthogonality can provide an estimation rate speed up in the case of noisy nuisances, i.e. the red-dotted line with noisy nuisance functions indicates the robustness of orthogonal estimation to slower nuisance function convergence. (The additional sample splitting leads to small-data concerns in the first plot, though as the difference-of-Q signal becomes stronger as in the case with Section 4, these finite sample concerns lessen.)

In all the experiments, we see that naive cross-validation requires a lot more data and converges slowly. This is expected due to 1) a large literature showing how CV Lasso for predictive error doesn't ensure support recovery (while methods like thresholded lasso do ensure support recovery) (Belloni and Chernozhukov, 2013; Hastie et al., 2015) and 2) additional challenges of hyperparameter tuning in offline RL. This illustrates how recovering the exact sparse support is crucial.

Next we instead consider a similar DGP in "Misaligned endo-exo", but change the blockwise conditional independences to follow the exogeneous-endogenous model of (Dietterich et al., 2018) (see Section 4, right). We also introduce some "misalignment" between reward and dynamic sparsity: the entries of β_{dense} are 1 w.p. 0.9 and we add $s^{\top}\beta_{dense}$ to the reward. In this setting, reward sparsity of $R(s, a), a \in \{0, 1\}$ alone is less informative of the underlying sparse component (which is still the first $|\rho| = 10$ dimensions). We see that in small data regimes, this degrades the performance of the reward-filtered thresholded LASSO method of (Zhou, 2024b): it, and vanilla thresholded LASSO FQE (FQE-TL) simply includes too many extraneous dimensions which destabilize estimation. In contrast for small-data regimes, imposing thresholded LASSO on the difference of Q functions, rather than the Q functions themselves remains stable. Again, in the large-sample limit, linear models remain well-specified and performance differences wash out.

The final DGP of "Nonlinear main effects" introduces nonlinear main effects: again we generate a 50% dense vector β_{dense} and we add $s^{\top}\beta_{dense}+3\sin(\pi s_{49}s_{48})+0.5(s_{49}-0.5)^2+0.5(s_{48}-0.5)^2$. (These nonlinear main effects are therefore disjoint from the sparse set and sparse difference-of-Q terms). Our FQE baselines now use kernel ridge regression (KRR), a very strong baseline for nonlinear regression. For small n, FQE wrongly includes extraneous dimensions that destabilize estimation, and our methods directly estimating τ with reward-thresholded-LASSO outperform even KRR for small data sizes. (With large enough data, KRR is well-specified.)

Limitations. To summarize limitations, as with other causal inference approaches, our approach requires certain assumptions, such as causal identification (which could be relaxed with sensitivity analysis). Our approach was also motivated by settings with direct sparsity in contrasted

rewards and dynamics; instead this could be true in some representation of the states/actions.

Conclusions We estimated the contrasts of Q functions with orthogonal estimation which adapts to structure. Important directions for future work include methods that address the proxy loss issue (Antos et al., 2008), model selection, representation-learning, and other techniques from machine learning and causal inference that incorporate inductive bias of learning causal contrasts.

References

- (2003). Optimal dynamic treatment regimes. Journal of the Royal Statistical Society Series B: Statistical Methodology 65(2), 331–355.
- Antos, A., C. Szepesvári, and R. Munos (2008). Learning near-optimal policies with bellman-residual minimization based fitted policy iteration and a single sample path. *Machine Learning* 71, 89–129.
- Belloni, A. and V. Chernozhukov (2013). Least squares after model selection in high-dimensional sparse models. *Bernoulli*.
- Chakraborty, B. and S. A. Murphy (2014). Dynamic treatment regimes. Annual review of statistics and its application 1, 447–464.
- Chernozhukov, V., D. Chetverikov, M. Demirer, E. Duflo, C. Hansen, W. Newey, and J. Robins (2018). Double/debiased machine learning for treatment and structural parameters.
- Chernozhukov, V., C. Hansen, N. Kallus, M. Spindler, and V. Syrgkanis (2024). Applied causal inference powered by ml and ai. rem 12(1), 338.
- Dietterich, T., G. Trimponias, and Z. Chen (2018). Discovering and removing exogenous state variables and rewards for reinforcement learning. In *International Conference on Machine Learning*, pp. 1262–1270. PMLR.
- Duan, Y., C. Jin, and Z. Li (2021). Risk bounds and rademacher complexity in batch reinforcement learning. In International Conference on Machine Learning, pp. 2892–2902. PMLR.
- Efroni, Y., S. Kakade, A. Krishnamurthy, and C. Zhang (2022). Sparsity in partially controllable linear systems. In *International Conference on Machine Learning*, pp. 5851–5860. PMLR.
- Foster, D. J. and V. Syrgkanis (2019). Orthogonal statistical learning. arXiv preprint arXiv:1901.09036.
- Hastie, T., R. Tibshirani, and M. Wainwright (2015). Statistical learning with sparsity. Monographs on statistics and applied probability 143(143), 8.
- Hu, Y., N. Kallus, and M. Uehara (2024). Fast rates for the regret of offline reinforcement learning. Mathematics of Operations Research.
- Jiang, N. and L. Li (2016). Doubly robust off-policy value evaluation for reinforcement learning. Proceedings of the 33rd International Conference on Machine Learning.

- Jin, Y., Z. Yang, and Z. Wang (2021). Is pessimism provably efficient for offline rl? In International Conference on Machine Learning, pp. 5084–5096. PMLR.
- Kallus, N. and M. Uehara (2019a). Double reinforcement learning for efficient off-policy evaluation in markov decision processes. arXiv preprint arXiv:1908.08526.
- Kallus, N. and M. Uehara (2019b). Efficiently breaking the curse of horizon: Double reinforcement learning in infinite-horizon processes. *arXiv preprint arXiv:1909.05850*.
- Kallus, N. and A. Zhou (2022). Stateful offline contextual policy evaluation and learning. In International Conference on Artificial Intelligence and Statistics, pp. 11169–11194. PMLR.
- Kennedy, E. H. (2020). Optimal doubly robust estimation of heterogeneous causal effects. arXiv preprint arXiv:2004.14497.
- Kennedy, E. H. (2022). Semiparametric doubly robust targeted double machine learning: a review. arXiv preprint arXiv:2203.06469.
- Künzel, S. R., J. S. Sekhon, P. J. Bickel, and B. Yu (2019). Metalearners for estimating heterogeneous treatment effects using machine learning. *Proceedings of the national academy of* sciences 116(10), 4156–4165.
- Le, H., C. Voloshin, and Y. Yue (2019). Batch policy learning under constraints. In International Conference on Machine Learning, pp. 3703–3712. PMLR.
- Lewis, G. and V. Syrgkanis (2021). Double/debiased machine learning for dynamic treatment effects. Advances in Neural Information Processing Systems 34.
- Li, F., L. E. Thomas, and F. Li (2019). Addressing extreme propensity scores via the overlap weights. American journal of epidemiology 188(1), 250–257.
- Liu, Q., L. Li, Z. Tang, and D. Zhou (2018). Breaking the curse of horizon: Infinite-horizon off-policy estimation. In Advances in Neural Information Processing Systems, pp. 5356–5366.
- Morzywolek, P., J. Decruyenaere, and S. Vansteelandt (2023). On a general class of orthogonal learners for the estimation of heterogeneous treatment effects. arXiv preprint arXiv:2303.12687.
- Neumann, G. and J. Peters (2008). Fitted q-iteration by advantage weighted regression. Advances in neural information processing systems 21.
- Nie, X., E. Brunskill, and S. Wager (2021). Learning when-to-treat policies. Journal of the American Statistical Association 116(533), 392–409.
- Nie, X. and S. Wager (2021). Quasi-oracle estimation of heterogeneous treatment effects. Biometrika 108(2), 299–319.
- Pan, H.-R. and B. Schölkopf (2023). Learning endogenous representation in reinforcement learning via advantage estimation. In *Causal Representation Learning Workshop at NeurIPS 2023*.
- Schulte, P. J., A. A. Tsiatis, E. B. Laber, and M. Davidian (2014). Q-and a-learning methods for estimating optimal dynamic treatment regimes. *Statistical science: a review journal of the Institute of Mathematical Statistics 29*(4), 640.

- Shi, C., S. Luo, H. Zhu, and R. Song (2022). Statistically efficient advantage learning for offline reinforcement learning in infinite horizons. arXiv preprint arXiv:2202.13163.
- Thomas, P., G. Theocharous, and M. Ghavamzadeh (2015). High confidence policy improvement. In *International Conference on Machine Learning*, pp. 2380–2388.
- Tsybakov, A. B. (2004). Optimal aggregation of classifiers in statistical learning. *The Annals of Statistics* 32(1), 135–166.
- Wager, S. and S. Athey (2018). Estimation and inference of heterogeneous treatment effects using random forests. *Journal of the American Statistical Association* 113(523), 1228–1242.
- Wainwright, M. J. (2019). *High-dimensional statistics: A non-asymptotic viewpoint*, Volume 48. Cambridge university press.
- Wang, T., S. S. Du, A. Torralba, P. Isola, A. Zhang, and Y. Tian (2022). Denoised mdps: Learning world models better than the world itself. arXiv preprint arXiv:2206.15477.
- Wang, Z., X. Xiao, Y. Zhu, and P. Stone. Task-independent causal state abstraction.
- Xie, C., W. Yang, and Z. Zhang (2023). Semiparametrically efficient off-policy evaluation in linear markov decision processes. In *International Conference on Machine Learning*, pp. 38227–38257. PMLR.
- Xie, T., C.-A. Cheng, N. Jiang, P. Mineiro, and A. Agarwal (2021). Bellman-consistent pessimism for offline reinforcement learning. Advances in neural information processing systems 34, 6683– 6694.
- Xu, T., Z. Yang, Z. Wang, and Y. Liang (2021). Doubly robust off-policy actor-critic: Convergence and optimality. In *International Conference on Machine Learning*, pp. 11581–11591. PMLR.
- Zhou, A. (2024a). Optimal and fair encouragement policy evaluation and learning. Advances in Neural Information Processing Systems 36.
- Zhou, A. (2024b). Reward-relevance-filtered linear offline reinforcement learning.
- Zhou, S. (2009). Thresholding procedures for high dimensional variable selection and statistical estimation. Advances in Neural Information Processing Systems 22.
- Zhou, S. (2010). Thresholded lasso for high dimensional variable selection and statistical estimation. arXiv preprint arXiv:1002.1583.

A Proofs

A.1 Preliminaries

Lemma 3 (Excess Variance).

$$\mathbb{E}[\hat{\mathcal{L}}_t(\tau,\eta)] - \mathcal{L}_t(\tau,\eta) = \operatorname{Var}[\max_{a'} Q\left(S_{t+1}, a'\right) \mid \pi_t^b]$$

Proof.

$$\begin{split} & \mathbb{E}[\hat{\mathcal{L}}_{t}(\tau,\eta)] \\ &= \mathbb{E}[\left(\{R_{t} + \gamma Q_{t+1}^{\pi^{e}}(S_{t+1}, A_{t+1}) - m_{t}^{\pi}(S_{t})\} \pm \mathbb{E}[\gamma Q_{t+1}^{\pi^{e}} \mid S_{t}, \pi_{t}^{b}] - \{A - \pi_{t}^{b}(1 \mid S_{t})\} \cdot \tau(S_{t})\right)^{2}] \\ &= \mathbb{E}[\left(\{R_{t} + \gamma \mathbb{E}[Q_{t+1}^{\pi^{e}} \mid S_{t}, \pi_{t}^{b}] - m_{t}^{\pi}(S_{t})\} - \{A - \pi_{t}^{b}(1 \mid S_{t})\} \cdot \tau(S_{t}) + \gamma(Q_{t+1}^{\pi^{e}}(S_{t+1}, A_{t+1}) - \mathbb{E}[Q_{t+1}^{\pi^{e}} \mid S_{t}, \pi_{t}^{b}])\right)^{2}] \\ &= \mathbb{E}[\left(\{R_{t} + \gamma \mathcal{T}Q_{t+1}^{\pi^{e}} - m_{t}^{\pi}\} - \{A - \pi_{t}^{b}(1 \mid S_{t})\} \cdot \tau(S_{t})\right)^{2}] \qquad (\text{squared loss of identifying moment}) \\ &+ \mathbb{E}[\gamma(Q_{t+1}^{\pi^{e}}(S_{t+1}, A_{t+1}) - \mathbb{E}[Q_{t+1}^{\pi^{e}} \mid S_{t}, \pi_{t}^{b}])^{2}] \qquad (\text{residual variance of } Q_{t}(s, a) - R_{t}(s, a)) \\ &+ \mathbb{E}[\left\{R_{t} + \gamma \mathbb{E}[Q_{t+1}^{\pi^{e}} \mid S_{t}, \pi_{t}^{b}] - m_{t}^{\pi}(S_{t}) - \{A - \pi_{t}^{b}(1 \mid S_{t})\} \cdot \tau(S_{t})\right\} \cdot \gamma(Q_{t+1}^{\pi^{e}}(S_{t+1}, A_{t+1}) - \mathbb{E}[Q_{t+1}^{\pi^{e}} \mid S_{t}, \pi_{t}^{b}])] \end{aligned}$$

Note the last term = 0 by iterated expectations and the pull-out property of conditional expectation. \Box

A.2 Orthogonality

Below we will omit the π superscript; the analysis below holds for any valid π . Define $\nu_t = \hat{\tau}_t - \tau_t^n, \nu_t^\circ = \hat{\tau}_t - \tau_t^\circ$. We define for any functional L(f) the Frechet derivative as:

$$D_f L(f)[\nu] = \left. \frac{\partial}{\partial t} L(f+t\nu) \right|_{t=0}$$

Higher order derivatives are denoted as $D_{g,f}L(f,g)[\mu,\nu]$.

Lemma 4 (Universal Orthogonality).

$$D_{\eta,\tau_t}\mathcal{L}_t(\tau_t^n;\tau_{t+1}^n,\eta^*)[\eta-\eta^*,\nu_t]=0$$

Proof of Lemma 4. For brevity, for a generic f, let $\{f\}_{\epsilon}$ denote $f + \epsilon(f - f^{\circ})$. Then the first Frechet derivatives are:

$$\frac{d}{d_{\epsilon_{\tau}}}\mathcal{L}_{t}(\tilde{\tau},\eta^{\circ})[\tau-\tilde{\tau},\eta-\eta^{\circ}] = \mathbb{E}\left[\left\{R_{t}+\gamma\{Q_{t+1}^{\pi^{e},\circ}\}_{\epsilon}-\{m_{t}^{\pi^{e},\circ}\}_{\epsilon}-(A_{t}-\{\pi_{t}^{b,\circ}\}_{\epsilon})\tau\right\}(A_{t}-\{\pi_{t}^{b,\circ}\}_{\epsilon})(\tilde{\tau}-\tau)\right]$$

$$\begin{aligned} \frac{d}{d\epsilon_e} \frac{d}{d\epsilon_\tau} \mathcal{L}_t \left(\tilde{\tau}, \eta^\circ\right) \left[\eta - \eta^\circ, \tau - \tau\right] \Big|_{\epsilon=0} \\ &= \mathbb{E} \left[\left(\pi_t^b - \pi_t^{b,\circ} \right) \tau (\tau - \tilde{\tau}) (A_t - e_t) \right] + \mathbb{E} \left[\left\{ R + \gamma Q_{t+1}^{\pi^e} - m_t^{\pi^e,\circ} - (A_t - e_t) \right\} (\tau - \tilde{\tau}) \cdot - (e_t - e_t^\circ) \right] \\ &= 0 \\ \frac{d}{d\epsilon_{Q_{t+1}}} \frac{d}{d\epsilon_\tau} \mathcal{L}_t \left(\tilde{\tau}, \eta^\circ\right) \left[\eta - \eta^\circ, \tau - \tilde{\tau} \right] \Big|_{\epsilon=0} \\ &= \mathbb{E} [\gamma (Q_{t+1}^{\pi^e} - Q_{t+1}^{\pi^e,\circ}) (A_t - \pi_t^{b,\circ}) (\tau_t - \tilde{\tau}_t)]] \\ &= 0 \\ \frac{d}{d\epsilon_{m_t}} \frac{d}{d\epsilon_\tau} \mathcal{L}_t \left(\tilde{\tau}, \eta^\circ\right) \left[\eta - \eta^\circ, \tau - \tilde{\tau} \right] \Big|_{\epsilon=0} \\ &= \mathbb{E} [-(m_t^{\pi^e} - m_t^{\pi^e,\circ}) (A_t - \pi_t^{b,\circ}) (\tau_t - \tilde{\tau}_t)] \\ &= 0 \end{aligned}$$

Lemma 5 (Second order derivatives). For Q_{t+1}, Q_{t+1}° evaluated at some fixed policy π^e :

$$D_{\eta_{t},\eta_{t}}\mathcal{L}_{t}[\hat{\eta}_{t} - \eta_{t}^{\circ}, \hat{\eta}_{t} - \eta_{t}^{\circ}] \\= \mathbb{E}\left[\tau_{t}^{2}\left(\hat{\pi}_{t}^{b} - \pi_{t}^{b,\circ}\right)^{2}\right] + \mathbb{E}\left[(\hat{\pi}_{t}^{b} - \pi_{t}^{b,\circ})\tau_{t}(\hat{m}_{t} - m_{t}^{\circ})\right] + \mathbb{E}\left[(\hat{\pi}_{t}^{b} - \pi_{t}^{b,\circ})\tau_{t}\gamma(\hat{Q}_{t+1} - Q_{t+1}^{\circ})\right] \\- \mathbb{E}\left[(\hat{m}_{t} - m_{t}^{\circ})\gamma\left(\hat{Q}_{t+1} - Q_{t+1}^{\circ}\right)\right]$$

Proof of Lemma 5. Below, the evaluation policy π^e is fixed and omitted for brevity. Note that

$$D_{e}\mathcal{L}_{D}[\hat{e} - e^{\circ}] = \mathbb{E}[(R_{t} + \gamma Q_{t+1} - \langle \pi^{b}, Q_{t} \rangle + (A - \pi^{b}_{t})\tau_{t})(-\tau_{t})(\hat{e} - e^{\circ})]$$
$$D_{m_{t}}\mathcal{L}_{D}[\hat{m}_{t} - m^{\circ}_{t}] = \mathbb{E}[(R_{t} + \gamma Q_{t+1} - \langle \pi^{b}, Q_{t} \rangle + (A - \pi^{b}_{t})\tau_{t})(-1) * (m_{t} - m^{\circ})]$$

By inspection, note that the nonzero terms of the second-order derivatives are as follows:

$$D_{\pi_{t}^{b},\pi_{t}^{b}}\mathcal{L}_{t}[\hat{\pi}_{t}^{b}-\pi_{t}^{b,\circ},\hat{\pi}_{t}^{b}-\pi_{t}^{b,\circ}] = \mathbb{E}\left[\tau_{t}^{2}\left(\hat{\pi}_{t}^{b}-\pi_{t}^{b,\circ}\right)^{2}\right]$$
$$D_{m_{t},Q_{t+1}}\mathcal{L}_{t}[\hat{Q}_{t+1}-Q_{t+1}^{\circ},\hat{m}_{t}-m_{t}^{\circ}] = \mathbb{E}\left[-\left(\hat{m}_{t}-m_{t}^{\circ}\right)\gamma\left(\hat{Q}_{t+1}-Q_{t+1}^{\circ}\right)\right]\right]$$
$$D_{m_{t},\pi_{t}^{b}}\mathcal{L}_{t}[\hat{\pi}_{t}^{b}-\pi_{t}^{b,\circ},\hat{m}_{t}-m_{t}^{\circ}] = \mathbb{E}\left[\left(\hat{\pi}_{t}^{b}-\pi_{t}^{b,\circ}\right)\tau_{t}\left(\hat{m}_{t}-m_{t}^{\circ}\right)\right]\right]$$
$$D_{Q_{t+1},\pi_{t}^{b}}\mathcal{L}_{t}[\hat{\pi}_{t}^{b}-\pi_{t}^{b,\circ},\hat{Q}_{t+1}-Q_{t+1}^{\circ}] = \mathbb{E}\left[\left(\hat{\pi}_{t}^{b}-\pi_{t}^{b,\circ}\right)\tau_{t}\gamma\left(\hat{Q}_{t+1}-Q_{t+1}^{\circ}\right)\right]$$

By the chain rule for Frechet differentiation, we have that

$$D_{\eta_{t},\eta_{t}}\mathcal{L}_{t}[\hat{\eta}_{t}-\eta_{t}^{\circ},\hat{\eta}_{t}-\eta_{t}^{\circ}] = D_{\pi_{t}^{b},\pi_{t}^{b}}\mathcal{L}_{t}[\hat{\pi}_{t}^{b}-\pi_{t}^{b,\circ},\hat{\pi}_{t}^{b}-\pi_{t}^{b,\circ}] + D_{m_{t},\pi_{t}^{b}}\mathcal{L}_{t}[\hat{\pi}_{t}^{b}-\pi_{t}^{b,\circ},\hat{m}_{t}-m_{t}^{\circ}] + D_{Q_{t+1},\pi_{t}^{b}}\mathcal{L}_{t}[\hat{\pi}_{t}^{b}-\pi_{t}^{b,\circ},\hat{Q}_{t+1}-Q_{t+1}^{\circ}] + D_{m_{t},Q_{t+1}}\mathcal{L}_{t}[\hat{Q}_{t+1}-Q_{t+1}^{\circ},\hat{m}_{t}-m_{t}^{\circ}]$$

A.3 Proof of sample complexity bounds

Proof of Lemma 2.

$$V_t^*(s) - V_t^{\pi_{\hat{\tau}}}(s) = V_t^*(s) - V_t^{\pi_{\hat{\tau}}}(s) \pm Q^{\pi^*}(s, \pi_{\hat{\tau}})$$

= $Q_t^*(s, \pi^*(s)) - Q_t^*(s, \hat{\pi}_{\hat{\tau}}) + Q_t^*(s, \hat{\pi}_{\hat{\tau}}) - V_t^{\hat{\pi}_{\hat{\tau}}}(s)$
 $\leq \gamma \mathbb{E}_{\hat{\pi}_t} \left[V_{t+1}^{\pi^*} - V_{t+1}^{\hat{\pi}_{\hat{\tau}}} \mid s \right] + Q_t^*(s, \pi^*(s)) - Q_t^*(s, \hat{\pi}_{\hat{\tau}})$

Therefore for any t and Markovian policy π inducing a marginal state distribution:

$$\mathbb{E}[V_t^*(s)] - \mathbb{E}[V_t^{\pi_{\hat{\tau}}}(s)] \le \gamma \mathbb{E}\left[\mathbb{E}_{\hat{\pi}_t}[V_{t+1}^{\pi^*} - V_{t+1}^{\hat{\pi}_{\hat{\tau}}} \mid s]\right] + \mathbb{E}[Q_t^*(s, \pi^*) - Q_t^*(s, \hat{\pi}_{\hat{\tau}})]$$
(8)

Assuming bounded rewards implies that $P(s_{t+1} | s, a) \leq c$, which remains true under the stateaction distribution induced by any Markovian policy $\pi(s, a)$, including the optimal policy. Therefore the second term of the above satisfies:

$$\mathbb{E}_{\pi}[Q_t^*(s_t, \pi^*) - Q_t^*(s_t, \hat{\pi}_{\hat{\tau}})] \le c \int \{Q_t^*(s, \pi^*) - Q_t^*(s, \hat{\pi}_{\hat{\tau}})\} ds,$$
(9)

and fixing t = 1, we obtain:

$$\mathbb{E}[Q_1^*(s_1,\pi^*) - Q_1^*(s_1,\hat{\pi}_{\hat{\tau}})] \le c \int \{Q_1^*(s,\pi^*) - Q_1^*(s,\hat{\pi}_{\hat{\tau}})\} \, ds.$$

Next we continue for generic t and bound the right hand side term of eq. (9).

First we suppose we have a high-probability bound on ℓ_{∞} convergence of $\hat{\tau}$. Define the good event

$$\mathcal{E}_g = \left\{ \sup_{s \in \mathcal{S}, a \in \mathcal{A}} \left| \hat{\tau}^{\hat{\pi}_{t+1}}(s) - \tau^{\pi^*_{t+1}, \circ}(s) \right| \le K n^{-b*} \right\}$$

A maximal inequality gives that $P(\mathcal{E}_g) \geq 1 - n^{-\kappa}$. We have that

$$\int \{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, ds = \int \{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}\left[\mathcal{E}_g\right] \, ds + \int \{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}\left[\mathcal{E}_g^c\right] \, ds + \int \{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}\left[\mathcal{E}_g^c\right] \, ds + \int \{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}\left[\mathcal{E}_g^c\right] \, ds + \int \{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}\left[\mathcal{E}_g^c\right] \, ds + \int \{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}\left[\mathcal{E}_g^c\right] \, ds + \int \{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}\left[\mathcal{E}_g^c\right] \, ds + \int \{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}\left[\mathcal{E}_g^c\right] \, ds + \int \{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}\left[\mathcal{E}_g^c\right] \, ds + \int \{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}\left[\mathcal{E}_g^c\right] \, ds + \int \{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}\left[\mathcal{E}_g^c\right] \, ds + \int \{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}\left[\mathcal{E}_g^c\right] \, ds + \int \{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}\left[\mathcal{E}_g^c\right] \, ds + \int \{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}\left[\mathcal{E}_g^c\right] \, ds + \int \{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}\left[\mathcal{E}_g^c\right] \, ds + \int \{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}\left[\mathcal{E}_g^c\right] \, ds + \int \{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}\left[\mathcal{E}_g^c\right] \, ds + \int \{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}\left[\mathcal{E}_g^c\right] \, ds + \int \{Q_t^*(s,\pi^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}\left[\mathcal{E}_g^c\right] \, ds + \int \{Q_t^*(s,\pi^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}\left[\mathcal{E}_g^c\right] \, ds + \int \{Q_t^*(s,\pi^*(s,\pi^*(s,\pi^*(s))) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}\left[\mathcal{E}_g^c\right] \, ds + \int \{Q_t^*(s,\pi^*(s,\pi^*(s,\pi^*(s,\pi^*(s))) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, ds + \int \{Q_t^*(s,\pi^*($$

Assuming boundedness, the bad event occurs with vanishingly small probability $n^{-\kappa}$, which bounds the second term of eq. (14).

For the first term of eq. (14), note that on the good event, if mistakes occur such that $\pi_t^*(s) \neq \hat{\pi}_t(s)$, then the true contrast function is still bounded in magnitude by the good event ensuring closeness of the estimate, so that $|\tau_t^{\pi_{t+1}^*,\circ}(s)| \leq 2Kn^{-b_*}$. And if no mistakes occur, at s the contribution to the integral is 0. Denote the mistake region as

$$\mathcal{S}_m = \{ s \in \mathcal{S} \colon \left| \tau_t^{\pi_{t+1}^*,\circ}(s) \right| \le 2Kn^{-b_*} \}$$

Therefore

$$\int \{Q_t^*(s, \pi^*(s)) - Q_t^*(s, \hat{\pi}_{\hat{\tau}})\} \, ds \leq \int_{s \in \mathcal{S}_m} \{Q_t^*(s, \pi^*(s)) - Q_t^*(s, \hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}\left[s \in \mathcal{S}_m\right] \, \mathbb{I}\left[\mathcal{E}_g\right] \, ds + O(n^{-\kappa}) \tag{11}$$

Note also that (for two actions), if *action* mistakes occur on the good event \mathcal{E}_g , the difference of Q functions must be near the decision boundaries so that we have the following bound on the integrand:

$$|Q^*(s,\pi^*) - Q^*(s,\hat{\pi})| \le |\tau^{\underline{\pi}^*_{t+1},\circ}| \le 2Kn^{-b*}.$$
(12)

Therefore,

$$\int \{Q_t^*(s, \pi^*(s)) - Q_t^*(s, \hat{\pi}_{\hat{\tau}})\} ds \le O(n^{-\kappa}) + Kn^{-b*} \int \mathbb{I}[s \in \mathcal{S}_m] ds$$
$$\le O(n^{-\kappa}) + (Kn^{-b*})(Kn^{-b*\alpha})$$
$$= O(n^{-\kappa}) + (K^2 n^{-b*(1+\alpha)})$$
(13)

where the first inequality follows from the above, and the second from assumption 5 (margin).

Combining eqs. (8) and (13), we obtain:

$$\mathbb{E}[V_t^*(S_t)] - \mathbb{E}[V_t^{\hat{\pi}_{\hat{\tau}}}(S_t)] \le \sum_{t=1}^T \gamma^t c \left\{ \int Q_t^{\hat{\pi}_{\hat{\tau}}}(s, \pi^*(s)) - Q_t^{\hat{\pi}_{\hat{\tau}}}(s, \hat{\pi}_{\hat{\tau}}) ds \right\}$$
$$\le \frac{(1 - \gamma^T)}{1 - \gamma} c T \{ O(n^{-\kappa}) + (K^2 n^{-b*(1+\alpha)}) \}$$

We also obtain analogous results for norm bounds:

$$\begin{split} \left\{ \int \left(Q_t^*(s, \pi^*(s)) - Q_t^*(s, \hat{\pi}_{\hat{\tau}}) \right)^u ds \right\}^{1/u} \\ & \leq \left\{ \int_{s \in \mathcal{S}_m} (Q_t^*(s, \pi^*(s)) - Q_t^*(s, \hat{\pi}_{\hat{\tau}}))^u \mathbb{I}\left[s \in \mathcal{S}_m\right] \mathbb{I}\left[\mathcal{E}_g\right] ds \right\}^{1/u} + O(n^{-\kappa}) \\ & \leq \frac{(1 - \gamma^T)}{1 - \gamma} cT \{ O(n^{-\kappa}) + (K^2 n^{-b*(1+\alpha)}) \} \end{split}$$

The results under an integrated risk bound assumption on convergence of τ follow analogously as (Shi et al., 2022), which we also include for completeness. For a given $\varepsilon > 0$, redefine the mistake region parametrized by ϵ :

$$\mathcal{S}_{\epsilon} = \left\{ \max_{a} Q^*(s, a) - Q^*(s, \hat{\pi}(s)) \le \varepsilon \right\}.$$

Again we obtain the bound by conditioning on the mistake region:

$$\int \{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, ds = \int \{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}[\mathcal{S}_{\epsilon}] \, ds + \int \{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}[\mathcal{S}_{\epsilon}^c] \, ds + \int \{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}[\mathcal{S}_{\epsilon}^c] \, ds + \int \{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}[\mathcal{S}_{\epsilon}^c] \, ds + \int \{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}[\mathcal{S}_{\epsilon}^c] \, ds + \int \{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}[\mathcal{S}_{\epsilon}^c] \, ds + \int \{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}[\mathcal{S}_{\epsilon}^c] \, ds + \int \{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}[\mathcal{S}_{\epsilon}^c] \, ds + \int \{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}[\mathcal{S}_{\epsilon}^c] \, ds + \int \{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}[\mathcal{S}_{\epsilon}^c] \, ds + \int \{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}[\mathcal{S}_{\epsilon}^c] \, ds + \int \{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}[\mathcal{S}_{\epsilon}^c] \, ds + \int \{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}[\mathcal{S}_{\epsilon}^c] \, ds + \int \{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}[\mathcal{S}_{\epsilon}^c] \, ds + \int \{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}[\mathcal{S}_{\epsilon}^c] \, ds + \int \{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}[\mathcal{S}_{\epsilon}^c] \, ds + \int \{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}[\mathcal{S}_{\epsilon}^c] \, ds + \int \{Q_t^*(s,\pi^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}[\mathcal{S}_{\epsilon}^c] \, ds + \int \{Q_t^*(s,\pi^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}[\mathcal{S}_{\epsilon}^c] \, ds + \int \{Q_t^*(s,\pi^*(s,\pi^*(s,\pi^*(s))) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\} \, \mathbb{I}[\mathcal{S}_{\epsilon}^c] \, ds + \int \{Q_t^*(s,\pi^*(s,\pi^*(s,\pi^*(s,\pi^*(s))) - Q_t^*(s,\pi^*$$

Using similar arguments as earlier, we can show by Assumption 5:

$$\int \left\{ Q_t^*\left(s, \pi^*(s)\right) - Q_t^*\left(s, \hat{\pi}_{\hat{\tau}}\right) \right\} \mathbb{I}(s \in \mathcal{S}_*) ds \le \varepsilon \int_x \mathbb{I}(s \in \mathcal{S}_*) ds = O\left(\varepsilon^{1+\alpha}\right).$$

As previously argued, we can show mistakes $\pi_t^*(s) \neq \hat{\pi}_t(s)$ occur only when

$$\max_{a} Q_t^*(s, a) - Q^*(s, \hat{\pi}(s)) \le 2 \left| \hat{\tau}^{\hat{\pi}_{t+1}}(s) - \tau^{\pi_{t+1}^*, \circ}(s) \right|.$$
(15)

It follows that

$$\begin{split} &\int \left\{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\right\} \mathbb{I}\left[s \in \mathcal{S}_{\epsilon}^c\right] ds \\ \leq & \mathbb{E} \int \frac{4 \left|\hat{\tau}^{\hat{\pi}_{t+1}}(s) - \tau^{\pi^*_{t+1},\circ}(s)\right|^2}{\left\{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})\right\}} \mathbb{I}\left[s \in \mathcal{S}_{\epsilon}^c\right] ds \\ \leq & \frac{4}{\varepsilon} \int \left|\hat{\tau}^{\hat{\pi}_{t+1}}(s) - \tau^{\pi^*_{t+1},\circ}(s)\right|^2 ds = O\left(\varepsilon^{-1}|\mathcal{I}|^{-2b_*}\right). \end{split}$$

Combining this together with (E.106) and (E.107) yields that

$$\int \left\{ Q_t^*(s, \pi^*(s)) - Q_t^*(s, \hat{\pi}_{\hat{\tau}}) \right\} ds = O\left(\varepsilon^{1+\alpha}\right) + O\left(\varepsilon^{-1} |\mathcal{I}|^{-2b_*}\right).$$

The result follows by choosing $\varepsilon = n^{-2b_*/(2+\alpha)}$ to balance the two terms. For the norm bound, the first term is analogously bounded as $O(\varepsilon^{1+\alpha})$:

$$\left\{\int (Q_t^*\left(s,\pi^*(s)\right) - Q_t^*\left(s,\hat{\pi}_{\hat{\tau}}\right))^2 \mathbb{I}[s \in \mathcal{S}_*] ds\right\}^{1/2} = O\left(\varepsilon^{1+\alpha}\right).$$

For the second term,

$$\left\{ \int (Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}}))^2 \mathbb{I}\left[s \in \mathcal{S}_{\epsilon}^c\right] ds \right\}^{1/2} \leq \left\{ \int \left(\frac{4 \left| \hat{\tau}^{\hat{\pi}_{t+1}}(s) - \tau^{\pi^*_{t+1},\circ}(s) \right|^2}{Q_t^*(s,\pi^*(s)) - Q_t^*(s,\hat{\pi}_{\hat{\tau}})} \right)^2 \mathbb{I}\left[s \in \mathcal{S}_{\epsilon}^c\right] ds \right\}^{1/2} \\ \leq \frac{4}{\varepsilon} \left\{ \int \left| \hat{\tau}^{\hat{\pi}_{t+1}}(s) - \tau^{\pi^*_{t+1},\circ}(s) \right|^4 ds \right\}^{1/2} = O\left(\varepsilon^{-1} |\mathcal{I}|^{-2b_*}\right).$$

The result follows as previous.

Proof of Theorem 1. In the following, at times we omit the fixed evaluation policy π^e from the notation for brevity. That is, in this proof, $\hat{\tau}_t, \tau_t^n$ are equivalent to $\hat{\tau}_t^{\pi^e}, \tau_t^{n,\pi^e}$. Further define

$$\nu_t = \hat{\tau}_t - \tau_t^n, \nu_t^\circ = \hat{\tau}_t - \tau_t^\circ$$

Strong convexity of the squared loss implies that:

$$D_{\tau_t,\tau_t} \mathcal{L} \left(\tau_t, \hat{\eta} \right) \left[\nu_t, \nu_t \right] \ge \lambda \left\| \nu_t \right\|_2^2$$

therefore

$$\frac{\lambda}{2} \|\nu_t\|_2^2 \leq \mathcal{L}_D(\hat{\tau}_t, \hat{\eta}) - \mathcal{L}_D(\tau_t^n, \hat{\eta}) - D_{\tau_t} \mathcal{L}_D(\tau_t^n, \hat{\eta}) [\nu_t]
\leq \epsilon(\hat{\tau}_t, \hat{\eta}) - D_{\tau_t} \mathcal{L}_D(\tau_t^n, \eta^\circ) [\nu_t]
+ D_{\tau_t} \mathcal{L}_D(\tau_t^n, \eta^\circ) [\nu_t] - D_{\tau_t} \mathcal{L}_D(\tau_t^n, \hat{\eta}) [\nu_t]$$
(16)

We bound each term in turn.

To bound $|D_{\tau_t} \mathcal{L}_D(\tau_t^n, \eta^\circ)[\nu_t]|$, note that

$$D_{\tau_t} \mathcal{L}_D(\tau_t^n, \eta^\circ)[\nu_t] = \mathbb{E}[(R + \gamma Q_{t+1} - V_t^{\pi^b, \pi_{t+1}:T} + \left(A - \pi_t^b\right)\tau_t))\left(A - \pi_t^b\right)\nu_t]$$

and by the properties of the conditional moment at the true τ° ,

$$= \mathbb{E}[(R + \gamma Q_{t+1} - V_t^{\pi^b, \pi_{t+1}:T} + (A - \pi_t^b) \tau_t^\circ)) (A - \pi_t^b) \nu_t] = 0$$

Therefore,

$$D_{\tau_t} \mathcal{L}_D(\tau_t^n, \eta^\circ)[\nu_t] = -\mathbb{E}[(\tau^\circ - \tau_t^n)(A - \pi_t^b)(A - \pi_t^b)(\hat{\tau}_t - \tau_t^n)]$$

Note that in general, for generic p, q, r such that 1/p + 1/q + 1/r = 1 we have that $\mathbb{E}[fgh] \leq ||fg||_{p'} ||h||_r \leq ||f||_p ||g||_q ||h||_r$ where $p' = \frac{pq}{p+q}$ or $\frac{1}{p'} = \frac{1}{p} + \frac{1}{q}$ or $1 = \frac{1}{p/p'} + \frac{1}{q/p'}$.

Therefore,

$$D_{\tau_t} \mathcal{L}_D(\tau_t^n, \eta^{\circ})[\nu_t] \le |D_{\tau_t} \mathcal{L}_D(\tau_t^n, \eta^{\circ})[\nu_t]| \le \mathbb{E}[(\tau^{\circ} - \tau_t^n) \mathbb{E}[(A_t - \pi_t^b)(A_t - \pi_t^b) \mid S_t](\hat{\tau}_t - \tau_t^n)] \le \|(\tau^{\circ} - \tau_t^n)\|_u \|(\hat{\tau}_t - \tau_t^n)\|_{\overline{u}} \cdot \left\{ \sup_s \mathbb{E}[(A_t - \pi_t^b)(A_t - \pi_t^b) \mid s] \right\}$$

where u, \overline{u} satisfy $\frac{1}{u} + \frac{1}{\overline{u}} = 1$. Next we bound $D_{\tau_t} \mathcal{L}_D(\tau_t^n, \eta^\circ)[\nu_t] - D_{\tau_t} \mathcal{L}_D(\tau_t^n, \hat{\eta})[\nu_t]$ by universal orthogonality. By a second order Taylor expansion, we have that, where $\eta_{\epsilon} = \eta^\circ + \epsilon(\hat{\eta} - \eta^\circ)$.

$$D_{\tau_t} \left(\mathcal{L}_D(\tau_t^n, \eta^{\circ}) - \mathcal{L}_D(\tau_t^n, \hat{\eta}) \right) [\nu_t] = \frac{1}{2} \int_0^1 D_{\eta, \eta, \tau_t}(\tau_t^n, \tau_{t+1}^{\circ}, \eta_{\epsilon}) [\hat{\eta} - \eta^{\circ}, \hat{\eta} - \eta^{\circ}, \nu_t]$$

We can deduce from Lemmas 4 and 5 that the integrand is:

$$\mathbb{E}\left[\tau_{t}^{2}\left(\hat{\pi}_{t}^{b}-\pi_{t}^{b,\circ}\right)^{2}\nu_{t}\right] + \mathbb{E}\left[\left(\hat{\pi}_{t}^{b}-\pi_{t}^{b,\circ}\right)\tau_{t}(\hat{m}_{t}-m_{t}^{\circ})\nu_{t}\right] + \mathbb{E}\left[\left(\hat{\pi}_{t}^{b}-\pi_{t}^{b,\circ}\right)\tau_{t}\gamma(\hat{Q}_{t+1}-Q_{t+1}^{\circ})\nu_{t}\right] \\
- \mathbb{E}\left[\left(\hat{m}_{t}-m_{t}^{\circ}\right)\gamma\left(\hat{Q}_{t+1}-Q_{t+1}^{\circ}\right)\nu_{t}\right] \\
\leq B_{\tau}^{2}\|\left(\hat{\pi}_{t}^{b}-\pi_{t}^{b,\circ}\right)^{2}\|_{u}\|\nu_{t}\|_{\overline{u}} + B_{\tau}\|(\hat{\pi}_{t}^{b}-\pi_{t}^{b,\circ})(\hat{m}_{t}-m_{t}^{\circ})\|_{u}\|\nu_{t}\|_{\overline{u}} + \gamma B_{\tau}\|(\hat{\pi}_{t}^{b}-\pi_{t}^{b,\circ})(\hat{Q}_{t+1}-Q_{t+1}^{\circ})\|_{u}\|\nu_{t}\|_{\overline{u}} \\
+ \gamma\|(\hat{m}_{t}-m_{t}^{\circ})(\hat{Q}_{t+1}-Q_{t+1}^{\circ})\|_{u}\|\nu_{t}\|_{\overline{u}}$$

Putting the bounds together, we obtain:

$$\frac{\lambda}{2} \|\nu_{t}\|_{2}^{2} \leq \epsilon(\hat{\tau}_{t},\hat{\eta}) + \|\nu_{t}\|_{\overline{u}} \|(\tau^{\circ} - \tau_{t}^{n})\|_{u}
+ \|\nu_{t}\|_{\overline{u}} \left(B_{\tau}^{2}\|\left(\hat{\pi}_{t}^{b} - \pi_{t}^{b,\circ}\right)^{2}\|_{u} + B_{\tau}\|(\hat{\pi}_{t}^{b} - \pi_{t}^{b,\circ})(\hat{m}_{t} - m_{t}^{\circ})\|_{u} + \gamma B_{\tau}\|(\hat{\pi}_{t}^{b} - \pi_{t}^{b,\circ})(\hat{Q}_{t+1} - Q_{t+1}^{\circ})\|_{u}
+ \gamma\|(\hat{m}_{t} - m_{t}^{\circ})(\hat{Q}_{t+1} - Q_{t+1}^{\circ})\|_{u}\right)$$
(17)

Let $\rho_t^{\pi^e}(\hat{\eta})$ denote the collected product error terms, e.g.

$$\rho_t^{\pi^e}(\hat{\eta}) = B_\tau^2 \| \left(\hat{\pi}_t^b - \pi_t^{b,\circ} \right)^2 \|_u + B_\tau \| (\hat{\pi}_t^b - \pi_t^{b,\circ}) (\hat{m}_t - m_t^\circ) \|_u \\ + \gamma (B_\tau \| (\hat{\pi}_t^b - \pi_t^{b,\circ}) (\hat{Q}_{t+1} - Q_{t+1}^\circ) \|_u + \| (\hat{m}_t - m_t^\circ) (\hat{Q}_{t+1} - Q_{t+1}^\circ) \|_u)$$

Analogously we drop the π^e decoration from ρ_t in this proof. The AM-GM inequality implies that for $x, y \ge 0, \sigma > 0$, we have that $xy \le \frac{1}{2}(\frac{2}{\sigma}x^2 + \frac{\sigma}{2}y^2)$. Therefore

$$\frac{\lambda}{2} \|\nu_t\|_2^2 - \frac{\sigma}{4} \|\nu_t\|_u^2 \le \epsilon(\hat{\tau}_t, \hat{\eta}) + \frac{1}{\sigma} \left(\|(\tau^\circ - \tau_t^n)\|_u + \rho_t(\hat{\eta}) \right)^2 \tag{18}$$

and since $(x+y)^2 \le 2(x^2+y^2)$,

$$\frac{\lambda}{2} \|\nu_t\|_2^2 - \frac{\sigma}{4} \|\nu_t\|_{\overline{u}}^2 \le \epsilon(\hat{\tau}_t, \hat{\eta}) + \frac{2}{\sigma} \left(\|(\tau^\circ - \tau_t^n)\|_u^2 + \rho_t(\hat{\eta})^2 \right)$$

Proof of Theorem 2. Let $\hat{\mathcal{L}}_{S,t}, \hat{\mathcal{L}}_{S',t}$ denote the empirical loss over the samples in S and S'; analogously $\hat{\eta}_S, \hat{\eta}_{S'}$ are the nuisance functions trained on each sample split.

Define the loss function ℓ_t on observation $O = \{(S_t, A_t, R_t, S_{t+1})\}_{t=1}^T$:

$$\ell_t(O;\tau_t;\hat{\eta}) = \left(\{ R_t + \hat{Q}_{t+1}^{\pi_{t+1}^e}(S_{t+1}, A_{t+1}) - \hat{m}_t(S_t) \} - \{ A - \hat{\pi}_t^b(1 \mid S_t) \} \cdot \tau_t(S_t) \right)^2$$

and the centered loss function $\Delta \ell$, centered with respect to $\hat{\tau}_t^n$:

$$\Delta \ell_t(O; \tau_t; \hat{\eta}) = \ell_t(O; \tau_t; \hat{\eta}) - \ell_t(O; \hat{\tau}_t^n; \hat{\eta}).$$

Assuming boundedness, ℓ_t is *L*-Lipschitz constant in τ_t :

$$\left|\Delta\ell_t(O;\tau_t;\hat{\eta}) - \Delta\ell_t(O;\tau_t';\hat{\eta})\right| \le L \|\tau_t - \tau_t\|_2.$$

Note that $\ell(O, \hat{\tau}_t^n, \hat{\eta}) = 0$. Define the centered average losses:

$$\Delta \hat{\mathcal{L}}_{S,t}(\tau_t, \hat{\eta}) = \hat{\mathcal{L}}_{S,t}(\tau_t, \hat{\eta}) - \hat{\mathcal{L}}_{S,t}(\hat{\tau}_t^n, \hat{\eta}) = \hat{\mathbb{E}}_{n/2}^S [\Delta \ell_t(O, \tau_T, \hat{\eta})]$$
$$\Delta \mathcal{L}_{S,t}(\tau_t, \hat{\eta}) = \mathcal{L}_{S,t}(\tau_t, \hat{\eta}) - \mathcal{L}_{S,t}(\hat{\tau}_t^n, \hat{\eta}) = \mathbb{E}[\Delta \ell_t(O, \tau_T, \hat{\eta})]$$

Assume that δ_n is an upper bound on the critical radius of the centered function class $\{\Psi_{t,i}^n - \hat{\tau}_{t,i}^n,$ with $\delta_n = \Omega(\frac{r \log \log n}{n})$, and define $\delta_{n,\xi} = \delta_n + c_0 \sqrt{\frac{\log(c_1 T/\xi)}{n}}$ for some c_0, c_1 . By Lemma 7 (Lemma 14 of (Foster and Syrgkanis, 2019) on local Rademacher complexity

By Lemma 7 (Lemma 14 of (Foster and Syrgkanis, 2019) on local Rademacher complexity decompositions), with high probability 1- ξ , for all $t \in [T]$, and for c_0 a universal constant ≥ 1 .

$$\begin{aligned} |\Delta \mathcal{L}_{S,t}(\hat{\tau}_{t}, \hat{\eta}_{S'}) - \Delta \mathcal{L}_{D,t}(\hat{\tau}_{t}, \hat{\eta}_{S'})| &= |\Delta \mathcal{L}_{S,t}(\hat{\tau}_{t}, \hat{\eta}_{S'}) - \Delta \mathcal{L}_{S,t}(\hat{\tau}_{t}^{n}, \hat{\eta}_{S'}) - (\Delta \mathcal{L}_{D,t}(\hat{\tau}_{t}, \hat{\eta}_{S'}) - \Delta \mathcal{L}_{D,t}(\hat{\tau}_{t}^{n}, \hat{\eta}_{S'}))| \\ &\leq c_{0} \left(rm \delta_{n/2,\xi} \| \hat{\tau}_{t} - \hat{\tau}_{t}^{n} \|_{2}^{2} + rm \delta_{n/2,\xi}^{2} \right) \end{aligned}$$

Assuming realizability of $\hat{\tau}_t$, we have that $\frac{1}{2} \left(\Delta \hat{\mathcal{L}}_{S,t}(\hat{\tau}_t, \hat{\eta}_{S'}) + \Delta \hat{\mathcal{L}}_{S',t}(\hat{\tau}_t, \hat{\eta}_S) \right) \leq 0$. Then with high probability $\geq 1 - 2\xi$:

$$\begin{aligned} &\frac{1}{2} \left(\Delta \mathcal{L}_{D,t}(\hat{\tau}_t, \hat{\eta}_{S'}) + \Delta \mathcal{L}_{D,t}(\hat{\tau}_t, \hat{\eta}_S) \right) \\ \leq &\frac{1}{2} \left| \Delta \mathcal{L}_{D,t}(\hat{\tau}_t, \hat{\eta}_{S'}) - \Delta \mathcal{L}_{S,t}(\hat{\tau}_t, \hat{\eta}_{S'}) + \Delta \mathcal{L}_{D,t}(\hat{\tau}_t, \hat{\eta}_S) - \Delta \mathcal{L}_{S',t}(\hat{\tau}_t, \hat{\eta}_S) \right| \\ \leq &\frac{1}{2} \left| \Delta \mathcal{L}_{D,t}(\hat{\tau}_t, \hat{\eta}_{S'}) - \Delta \mathcal{L}_{S,t}(\hat{\tau}_t, \hat{\eta}_{S'}) \right| + \left| \Delta \mathcal{L}_{D,t}(\hat{\tau}_t, \hat{\eta}_S) - \Delta \mathcal{L}_{S',t}(\hat{\tau}_t, \hat{\eta}_S) \right| \\ \leq &c_0 \left(rm \delta_{n/2,\xi} \| \hat{\tau}_t - \hat{\tau}_t^n \|_2 + rm \delta_{n/2,\xi}^2 \right) \end{aligned}$$

The ϵ excess risk term in Theorem 1 indeed corresponds to one of the loss differences defined here, i.e. $\Delta \mathcal{L}_{D,t}(\hat{\tau}_t, \hat{\eta}_S) := \epsilon(\hat{\tau}_t^n, \hat{\tau}_t, \hat{h}_S)$. Therefore, applying Theorem 1 with $u = \overline{u} = 2$ and $\sigma = \lambda$ with the above bound, and averaging the sample-split estimators, we obtain

$$\frac{\lambda}{4} \|\nu_t\|_2^2 \le \frac{1}{2} \left(\epsilon(\hat{\tau}_t, \hat{\eta}_S) + \epsilon(\hat{\tau}_t, \hat{\eta}_{S'}) \right) + \frac{2}{\lambda} \left(\|\tau_t^{\circ} - \hat{\tau}_t^n\|_2^2 + \sum_{s \in \{S, S'\}} \rho_t(\hat{\eta}_s)^2 \right)$$

We further decompose the excess risk of empirically-optimal $\hat{\tau}_t$ relative to the population minimizer to instead bound by the error of $\hat{\tau}_t$ to the projection onto Ψ , $\hat{\tau}_t^n$, since $\|\hat{\tau}_t - \tau_t^\circ\|_2^2 \leq \|\hat{\tau}_t - \hat{\tau}_t^n\|_2^2 + \|\hat{\tau}_t^n - \tau_t^\circ\|_2^2$, we obtain

$$\frac{\lambda}{4} \|\hat{\tau}_t - \tau_t^{\circ}\|_2^2 \le c_0 \left(rm\delta_{n/2,\xi} \|\hat{\tau}_t - \hat{\tau}_t^n\|_2 + rm\delta_{n/2,\xi}^2 \right) + \frac{8 + \lambda^2}{4\lambda} \|\tau_t^{\circ} - \tau_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S,S'\}} \rho_t(\hat{\eta}_s)^2 \|\hat{\tau}_t - \hat{\tau}_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S,S'\}} \rho_t(\hat{\eta}_s)^2 \|\hat{\tau}_t - \hat{\tau}_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S,S'\}} \rho_t(\hat{\eta}_s)^2 \|\hat{\tau}_t - \hat{\tau}_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S,S'\}} \rho_t(\hat{\eta}_s)^2 \|\hat{\tau}_t - \hat{\tau}_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S,S'\}} \rho_t(\hat{\eta}_s)^2 \|\hat{\tau}_t - \hat{\tau}_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S,S'\}} \rho_t(\hat{\eta}_s)^2 \|\hat{\tau}_t - \hat{\tau}_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S,S'\}} \rho_t(\hat{\eta}_s)^2 \|\hat{\tau}_t - \hat{\tau}_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S,S'\}} \rho_t(\hat{\eta}_s)^2 \|\hat{\tau}_t - \hat{\tau}_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S,S'\}} \rho_t(\hat{\eta}_s)^2 \|\hat{\tau}_t - \hat{\tau}_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S,S'\}} \rho_t(\hat{\eta}_s)^2 \|\hat{\tau}_t - \hat{\tau}_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S,S'\}} \rho_t(\hat{\eta}_s)^2 \|\hat{\tau}_t - \hat{\tau}_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S,S'\}} \rho_t(\hat{\eta}_s)^2 \|\hat{\tau}_t - \hat{\tau}_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S,S'\}} \rho_t(\hat{\eta}_s)^2 \|\hat{\tau}_t - \hat{\tau}_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S,S'\}} \rho_t(\hat{\eta}_s)^2 \|\hat{\tau}_t - \hat{\tau}_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S,S'\}} \rho_t(\hat{\eta}_s)^2 \|\hat{\tau}_t - \hat{\tau}_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S,S'\}} \rho_t(\hat{\eta}_s)^2 \|\hat{\tau}_t - \hat{\tau}_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S,S'\}} \rho_t(\hat{\eta}_s)^2 \|\hat{\tau}_t - \hat{\tau}_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S,S'\}} \rho_t(\hat{\eta}_s)^2 \|\hat{\tau}_t - \hat{\tau}_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S,S'\}} \rho_t(\hat{\eta}_s)^2 \|\hat{\tau}_t - \hat{\tau}_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S,S'\}} \rho_t(\hat{\eta}_s)^2 \|\hat{\tau}_t - \hat{\tau}_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S,S'\}} \rho_t(\hat{\eta}_s)^2 \|\hat{\tau}_t - \hat{\tau}_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S,S'\}} \rho_t(\hat{\eta}_s)^2 \|\hat{\tau}_t - \hat{\tau}_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S,S'\}} \rho_t(\hat{\eta}_s)^2 \|\hat{\tau}_t - \hat{\tau}_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S,S'\}} \rho_t(\hat{\tau}_t - \hat{\tau}_t^n\|$$

Again using the AM-GM inequality $xy \leq \frac{1}{2} \left(\frac{2}{\sigma}x^2 + \frac{\sigma}{2}y^2\right)$, we bound

$$c_{0}\left(rm\delta_{n/2,\xi}\|\hat{\tau}_{t}-\hat{\tau}_{t}^{n}\|_{2}+rm\delta_{n/2,\xi}^{2}\right) \leq \frac{c_{0}}{2}r^{2}m^{2}(1+\frac{2}{\epsilon})\delta_{n/2,\xi}^{2}+\frac{\epsilon}{4}\|\hat{\tau}_{t}-\hat{\tau}_{t}^{n}\|_{2}^{2}$$
$$\leq c_{0}r^{2}m^{2}(1+\frac{1}{\epsilon})\delta_{n/2,\xi}^{2}+\frac{\epsilon}{4}(\|\hat{\tau}_{t}-\tau_{t}^{\circ}\|_{2}^{2}+\|\tau_{t}^{\circ}-\hat{\tau}_{t}^{n}\|_{2}^{2})$$

Therefore,

$$\frac{\lambda - \epsilon}{4} \|\hat{\tau}_t - \tau_t^{\circ}\|_2^2 \le c_0 r^2 m^2 (1 + \frac{1}{\epsilon}) \delta_{n/2,\xi}^2 + \left(\frac{8 + \lambda^2}{4\lambda} + \frac{\epsilon}{4}\right) \|\tau_t^{\circ} - \tau_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S, S'\}} \rho_t(\hat{\eta}_s)^2 \|\tau_s^{\circ} - \tau_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S, S'\}} \rho_t(\hat{\eta}_s)^2 \|\tau_s^{\circ} - \tau_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S, S'\}} \rho_t(\hat{\eta}_s)^2 \|\tau_s^{\circ} - \tau_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S, S'\}} \rho_t(\hat{\eta}_s)^2 \|\tau_s^{\circ} - \tau_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S, S'\}} \rho_t(\hat{\eta}_s)^2 \|\tau_s^{\circ} - \tau_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S, S'\}} \rho_t(\hat{\eta}_s)^2 \|\tau_s^{\circ} - \tau_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S, S'\}} \rho_t(\hat{\eta}_s)^2 \|\tau_s^{\circ} - \tau_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S, S'\}} \rho_t(\hat{\eta}_s)^2 \|\tau_s^{\circ} - \tau_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S, S'\}} \rho_t(\hat{\eta}_s)^2 \|\tau_s^{\circ} - \tau_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S, S'\}} \rho_t(\hat{\eta}_s)^2 \|\tau_s^{\circ} - \tau_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S, S'\}} \rho_t(\hat{\eta}_s)^2 \|\tau_s^{\circ} - \tau_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S, S'\}} \rho_t(\hat{\eta}_s)^2 \|\tau_s^{\circ} - \tau_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S, S'\}} \rho_t(\hat{\eta}_s)^2 \|\tau_s^{\circ} - \tau_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S, S'\}} \rho_t(\hat{\eta}_s)^2 \|\tau_s^{\circ} - \tau_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S, S'\}} \rho_t(\hat{\eta}_s)^2 \|\tau_s^{\circ} - \tau_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S, S'\}} \rho_t(\hat{\eta}_s)^2 \|\tau_s^{\circ} - \tau_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S, S'\}} \rho_t(\hat{\eta}_s)^2 \|\tau_s^{\circ} - \tau_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S, S'\}} \rho_t(\hat{\eta}_s)^2 \|\tau_s^{\circ} - \tau_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S, S'\}} \rho_t(\hat{\eta}_s)^2 \|\tau_s^{\circ} - \tau_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S, S'\}} \rho_t(\hat{\eta}_s)^2 \|\tau_s^{\circ} - \tau_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S, S'\}} \rho_t(\hat{\eta}_s)^2 \|\tau_s^{\circ} - \tau_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S, S'\}} \rho_t(\hat{\eta}_s)^2 \|\tau_s^{\circ} - \tau_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S, S'\}} \rho_t(\hat{\eta}_s)^2 \|\tau_s^{\circ} - \tau_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S, S'\}} \rho_t(\hat{\eta}_s)^2 \|\tau_s^{\circ} - \tau_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S, S'\}} \rho_t(\hat{\eta}_s)^2 \|\tau_s^{\circ} - \tau_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S, S'\}} \rho_t(\hat{\eta}_s)^2 \|\tau_s^{\circ} - \tau_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S, S'\}} \rho_t(\hat{\eta}_s)^2 \|\tau_s^{\circ} - \tau_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S, S'\}} \rho_t(\hat{\eta}_s)^2 \|\tau_s^{\circ} - \tau_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S, S'\}} \rho_t(\hat{\eta}_s)^2 \|\tau_s^{\circ} - \tau_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S, S'\}} \rho_t(\hat{\eta}_s)^2 \|\tau_s^{\circ} - \tau_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S, S'\}} \rho_t(\hat{\eta}_s)^2 \|\tau_s^n\|_2^2 + \frac{$$

Choose $\epsilon \leq \lambda/8$ so that

$$\begin{aligned} \frac{\lambda}{8} \|\hat{\tau}_t - \tau_t^{\circ}\|_2^2 &\leq c_0 r^2 m^2 (1 + \frac{8}{\lambda}) \delta_{n/2,\xi}^2 + \left(\frac{4 + \lambda^2}{2\lambda}\right) \|\tau_t^{\circ} - \tau_t^n\|_2^2 + \frac{2}{\lambda} \sum_{s \in \{S, S'\}} \rho_t(\hat{\eta}_s)^2 \\ &\leq \left(1 + \frac{8}{\lambda} + \frac{\lambda}{2}\right) (c_0 r^2 m^2 \delta_{n/2,\xi}^2 + \|\tau_t^{\circ} - \tau_t^n\|_2^2 + \sum_{s \in \{S, S'\}} \rho_t(\hat{\eta}_s)^2) \end{aligned}$$

and therefore

$$\|\hat{\tau}_t - \tau_t^{\circ}\|_2^2 \le \left(\frac{8}{\lambda}(1+\frac{8}{\lambda})+4\right)\left(c_0 r^2 m^2 \delta_{n/2,\xi}^2 + \|\tau_t^{\circ} - \tau_t^n\|_2^2 + \sum_{s \in \{S,S'\}} \rho_t(\hat{\eta}_s)^2\right)$$

Taking expectations:

$$\mathbb{E}[\|\hat{\tau}_t - \tau_t^{\circ}\|_2^2] \le \left(\frac{8}{\lambda}(1 + \frac{8}{\lambda}) + 4\right) \left(c_0 r^2 m^2 \delta_{n/2}^2 + \|\tau_t^{\circ} - \tau_t^n\|_2^2 + \max_{s \in \{S, S'\}} \mathbb{E}[\rho_t(\hat{\eta}_s)^2]\right)$$

Therefore, if the product error rate terms are all of the same order as the estimation order terms:

$$\begin{split} & \mathbb{E}[\|\hat{\pi}_{t}^{b} - \pi_{t}^{b,\circ}\|_{2}^{2}] = O(\delta_{n/2}^{2} + \|\tau_{t}^{\circ} - \tau_{t}^{n}\|_{2}^{2}) \\ & \mathbb{E}[\|(\hat{\pi}_{t}^{b} - \pi_{t}^{b,\circ})(\hat{m}_{t} - m_{t}^{\circ})\|_{2}^{2}] = O(\delta_{n/2}^{2} + \|\tau_{t}^{\circ} - \tau_{t}^{n}\|_{2}^{2}) \\ & \mathbb{E}[\|(\hat{\pi}_{t}^{b} - \pi_{t}^{b,\circ})(\hat{Q}_{t+1} - Q_{t+1}^{\circ})\|_{2}^{2}] = O(\delta_{n/2}^{2} + \|\tau_{t}^{\circ} - \tau_{t}^{n}\|_{2}^{2}) \\ & \mathbb{E}[\|(\hat{m}_{t} - m_{t}^{\circ})(\hat{Q}_{t+1} - Q_{t+1}^{\circ})\|_{2}^{2}] = O(\delta_{n/2}^{2} + \|\tau_{t}^{\circ} - \tau_{t}^{n}\|_{2}^{2}) \end{split}$$

_	_
_	_

Proof of Theorem 3. Preliminaries We introduce some additional notation. For the analysis of implications of policy optimization, we further introduce notation that parametrizes the time-t loss function with respect to the time-(t + 1) policy. In analyzing the policy optimization, this will be used to decompose the policy error arising from time steps closer to the horizon. Define

$$\mathcal{L}_D(\tau_t^n, \tau_{t+1}', \hat{\eta}) = \mathbb{E}\left[\left(\{R_t + \gamma Q_{t+1}^{\pi_{\tau_{t+1}'}}(S_{t+1}, A_{t+1}) - V_{\pi_t^b, \pi_{\tau_{t+1}}'}(S_t)\} - \{A - \pi_t^b(1 \mid S_t)\} \cdot \tau(S_t)\right)^2\right]$$

where $\pi_{\tau'_{t+1}}(s) \in \operatorname{argmax} \tau'_{t+1}(s)$. That is, the second argument parameterizes the difference-of-Q function that generates the policy that oracle nuisance functions are evaluated at.

Then, for example, the true optimal policy satisfies that $\pi_t^* \in \arg \max \tau_t^\circ(s)$. We define the oracle loss function with nuisance functions evaluated with respect to the optimal policy π^* .

$$\mathcal{L}_D(\tau_t^n, \tau^{\circ}, \hat{\eta}) = \mathbb{E}\left[\left(\{R_t + \gamma Q_{t+1}^{\pi^*_{\tau_{t+1}}}(S_{t+1}, A_{t+1}) - m^{\circ}(S_t)\} - \gamma\{A - \pi_t^b(1 \mid S_t)\} \cdot \tau(S_t)\right)^2\right]$$

In contrast, the empirical policy optimizes with respect to a next-stage *estimate* of the *empirical* best next-stage policy $\hat{\pi}_{\hat{\tau}_{t+1}}$. That is, noting the empirical loss function:

$$\mathcal{L}_D(\tau_t^n, \hat{\tau}_{t+1}, \hat{\eta}) = \mathbb{E}\left[\left(\{R_t + \gamma Q_{t+1}^{\hat{\pi}_{t+1}}(S_{t+1}, A_{t+1}) - m^{\circ}(S_t)\} - \gamma\{A - \pi_t^b(1 \mid S_t)\} \cdot \tau(S_t)\right)^2\right]$$

Step 1: Applying advantage estimation results. At every timestep, the first substep is to estimate the *Q*-function contrast, $\hat{\tau}_t^{\hat{\pi}_{t+1}}$. The assumptions on product error nuisance rates imply that for a fixed $\hat{\pi}_{t+1}$ that we would obtain estimation error

$$\mathbb{E}\left[\|\hat{\tau}_{t}^{\hat{\pi}_{t+1}} - \tau_{t}^{\hat{\pi}_{t+1},\circ}\|_{2}^{2}\right] = O\left(\delta_{n/2}^{2} + \left\|\tau_{t}^{\pi^{e},\circ} - \tau_{t}^{\pi^{e},n}\right\|_{2}^{2}\right)$$

Step 2: Establishing policy consistency. Applying Lemma 2 requires a convergence rate of $\hat{\tau}_t^{\hat{\pi}_{t+1}}$ to $\hat{\tau}_t^{\pi_{t+1}^*}$. The estimation error guarantees on the contrast function, however, are for the policy $\hat{\pi}_{t+1}$. We obtain the required bound via induction. At a high level, the estimation error arising from $\hat{\pi}_{t+1}$ vs π_{t+1}^* too eventually is integrated; so when the margin exponent $\alpha > 0$, these policy error terms are higher-order and vanish at a faster rate.

Importantly, we suppose the product error rate conditions hold for each t for data-optimal policies evaluated along the algorithm, i.e. for each t, for each t, for $\underline{\hat{\pi}}_{t+1}$, each of $\mathbb{E}[\|(\hat{\pi}_t^b - \pi_t^{b,\circ})\|_2^2]$, $\mathbb{E}[\|(\hat{\pi}_t^b - \pi_t^{b,\circ})(\hat{Q}_{t+1}^{\hat{\pi}_{t+1}} - Q_{t+1}^{\circ,\hat{\pi}_{t+2}})\|_2^2]$, and $\mathbb{E}[\|(\hat{m}_t - m_t^{\circ})(\hat{Q}_{t+1}^{\circ,\hat{\pi}_{t+2}} - Q_{t+1}^{\circ,\hat{\pi}_{t+2}})\|_2^2]$, and $\mathbb{E}[\|(\hat{m}_t - m_t^{\circ})(\hat{Q}_{t+1}^{\circ,\hat{\pi}_{t+2}} - Q_{t+1}^{\circ,\hat{\pi}_{t+2}})\|_2^2]$ are of order $O(\delta_{n/2}^2 + \|\tau_t^{\hat{\pi}_{t+1},\circ} - \tau_t^{\hat{\pi}_{t+1},n}\|_2^2)$.

Step 2a: induction hypothesis.

Next we show the induction hypothesis.

First we consider the base case: When t = T, τ_T is independent of the forward policy so that $\|\hat{\tau}_T^{\hat{\pi}} - \tau_T^{\circ,\pi^*}\| = \|\hat{\tau}_T - \tau_T^{\circ}\|$. Then the base case follows by Theorem 2.

Suppose it is true that for timesteps $k \ge t+1$, we have that

$$\|\hat{\tau}_{k}^{\hat{\pi}_{k+1}} - \tau_{k}^{\circ,\underline{\pi}_{k+1}^{*}}\| = O(\delta_{n/2} + \|\tau_{k}^{\circ,\underline{\hat{\pi}}_{k+1}} - \tau_{k}^{n,\underline{\hat{\pi}}_{k+1}}\|_{2}) + Kn^{-\mathcal{R}_{k}},\tag{19}$$

where

$$\mathcal{R}_{k} = \min\left(\rho_{k+1}^{(c)} \cdot \frac{2+2\alpha}{2+\alpha}, \ \rho_{k+1}^{(\Psi)} \cdot \frac{2+2\alpha}{2+\alpha}, \ -\{\min_{k' \ge k+1}(\rho_{k'}^{(c)}, \rho_{k'}^{(\Psi)})\} \cdot \frac{2+2\alpha}{2+\alpha}^{T-k'}\right).$$
(20)

And therefore, applying Lemma 2, that

$$\left| \mathbb{E}[V_k^{\pi^*} - V_k^{\hat{\pi}_{\hat{\tau}}}] \right| = O(n^{-\min\{\rho_k^{(c)}, \rho_k^{(\Psi)}\}\frac{2+2\alpha}{2+\alpha}}) + o(n^{-\min\{\rho_k^{(c)}, \rho_k^{(\Psi)}\}\frac{2+2\alpha}{2+\alpha}}).$$
(21)

We will show that the induction hypothesis implies

$$\|\hat{\tau}_t^{\hat{\underline{\pi}}_{t+1}} - \tau_t^{\circ,\underline{\pi}_{t+1}^*}\| \le O(\delta_{n/2} + \|\tau_t^{\circ,\hat{\underline{\pi}}_{t+1}} - \tau_t^{n,\hat{\underline{\pi}}_{t+1}}\|_2) + Kn^{-\mathcal{R}_t}.$$

and

$$\mathbb{E}[V_k^{\pi^*} - V_k^{\hat{\pi}_{\hat{\tau}}}] = O(n^{-\min\{\rho_k^{(c)}, \rho_k^{(\Psi)}\}\frac{2+2\alpha}{2+\alpha}}) + o(n^{-\min\{\rho_k^{(c)}, \rho_k^{(\Psi)}\}\frac{2+2\alpha}{2+\alpha}})$$

First decompose the desired error $\|\hat{\tau}_t^{\hat{\pi}_{t+1}} - \tau_t^{\circ,\underline{\pi}_{t+1}^*}\|$ as:

$$\|\hat{\tau}_{t}^{\hat{\underline{\pi}}_{t+1}} - \tau_{t}^{\circ,\underline{\pi}_{t+1}^{*}}\| \leq \|\hat{\tau}_{t}^{\hat{\underline{\pi}}_{t+1}} - \tau_{t}^{\circ,\underline{\hat{\pi}}_{t+1}}\| + \|\tau_{t}^{\circ,\underline{\hat{\pi}}_{t+1}} - \tau_{t}^{\circ,\underline{\pi}_{t+1}^{*}}\|$$
(22)

The first term is the policy evaluation estimation error, and under the product error rate assumptions , Theorems 1 and 2 give that $\mathbb{E}[\|\hat{\tau}_t^{\hat{\pi}_{t+1}} - \tau_t^{\circ,\hat{\pi}_{t+1}}\|_2^2] = O(\delta_{n/2}^2 + \|\tau_t^{\circ,\hat{\pi}_{t+1}} - \tau_t^{n,\hat{\pi}_{t+1}}\|_2^2)$. The second term of the above depends on the convergence of the empirically optimal policy $\hat{\pi}$; we use our analysis from Lemma 2 to bound the impact of future estimates of difference-of-Q functions using the induction hypothesis. The following analysis will essentially reveal that the margin assumption of Assumption 5 implies that the error due to the empirically optimal policy is higher-order, and the first term (time-t estimation error of $\hat{\tau}_t$) is the leading term.

As in eq. (8), we have that:

$$V_t^*(s) - V_t^{\pi_{\hat{\tau}}}(s) \le \gamma \mathbb{E}_{\hat{\pi}_t} \left[V_{t+1}^{\pi^*} - V_{t+1}^{\hat{\pi}_{\hat{\tau}}} \mid s_t \right] + Q_t^*(s, \pi^*) - Q_t^*(s, \hat{\pi}_{\hat{\tau}}).$$

Decompose:

$$\|\tau_t^{\circ,\hat{\pi}_{t+1}} - \tau_t^{\circ,\underline{\pi}_{t+1}^*}\| \le \sum_a \|Q_t^{\underline{\pi}_{t+1}^*}(s,a) - Q_t^{\hat{\pi}_{t+1}}(s,a)\|$$

By definition of τ and $\pm V_{t+1}^{\hat{\pi}_{t+1}, \underline{\pi}_{t+2}^*}$, for each a, we have that

$$\begin{aligned} \|Q_{t}^{\underline{\pi}_{t+1}^{*}}(s,a) - Q_{t}^{\underline{\hat{\pi}}_{t+1}}(s,a)\| \\ &= \|\mathbb{E}_{\pi_{t}^{a}}[V_{t+1}^{\underline{\pi}_{t+1}^{*}} - V_{t+1}^{\underline{\hat{\pi}}_{t+1}} \mid S_{t}]\| \\ &\leq \|\mathbb{E}_{\pi_{t}^{a}}[V_{t+1}^{\underline{\pi}_{t+1}^{*}} - V_{t+1}^{\underline{\hat{\pi}}_{t+1},\underline{\pi}_{t+2}^{*}} \mid S_{t}]\| + \|\mathbb{E}_{\pi_{t}^{a}}[V_{t+1}^{\underline{\hat{\pi}}_{t+1},\underline{\pi}_{t+2}^{*}} - V_{t+1}^{\underline{\hat{\pi}}_{t+1}} \mid S_{t}]\| \\ &= \|\mathbb{E}_{\pi_{t}^{a}}[Q_{t+1}^{\underline{\pi}_{t+2}^{*}}(S_{t+1},\pi_{t+1}^{*}) - Q_{t+1}^{\underline{\pi}_{t+2}^{*}}(S_{t+1},\hat{\pi}_{t+1}) \mid S_{t}]\| + \gamma \|\mathbb{E}_{\pi_{t}^{a}}[\mathbb{E}_{\underline{\hat{\pi}}_{t+1}}[V_{t+2}^{\underline{\pi}_{t+2}^{*}} - V_{t+2}^{\underline{\hat{\pi}}_{t+2}} \mid S_{t}]]\| \end{aligned}$$
(23)

$$\leq c \left\{ \int (Q_{t+1}^{\pi_{t+2}^*}(s,\pi_{t+1}^*) - Q_{t+1}^{\pi_{t+2}^*}(s,\hat{\pi}_{t+1}))^2 ds \right\}^{1/2} + \gamma \|\mathbb{E}_{\pi_t^a}[\mathbb{E}_{\hat{\pi}_{t+1}}[V_{t+2}^{\pi_{t+2}^*} - V_{t+2}^{\hat{\pi}_{t+2}} \mid S_t]]\|$$
(24)

where the last inequality follows by Assumption 4 and the policy-convolved transition density.

Next we bound the first term using the margin analysis of Lemma 2 and the inductive hypothesis. Supposing the product error rates are satisfied on the nuisance functions for estimation of $\hat{\tau}_{t+1}$, the induction hypothesis gives that

$$\mathbb{E}[\|\hat{\tau}_{t+1}^{\hat{\underline{\pi}}_{t+2}} - \tau_{t+1}^{\circ,\underline{\pi}_{t+2}^*}\|_2] = O\left(\delta_{n/2} + \|\tau_t^{\pi^e,\circ} - \tau_t^n\|_2 + n^{-\mathcal{R}_{t+1}}\right)$$

The induction hypothesis gives the integrated risk rate assumption on $\hat{\tau}_{t+1}$ to apply Lemma 2,

$$\left\{ \int (Q_{t+1}^{\underline{\pi}_{t+2}^*}(s, \pi_{t+1}^*) - Q_{t+1}^{\underline{\pi}_{t+2}^*}(s, \hat{\pi}_{t+1}))^2 ds \right\}^{1/2} \\ \leq \frac{(1 - \gamma^{T-t-1})}{1 - \gamma} c(T - t - 1) \{ O(n^{-\kappa}) + Kn^{-\min\{r_{t+1}^{(c)}, r_{t+1}^{(\Psi)}, \mathcal{R}_{t+1}\}(1+\alpha)} \}.$$

Combining with the previous analysis, we obtain:

$$\begin{aligned} \|\hat{\tau}_{t}^{\hat{\pi}_{t+1}} - \tau_{t}^{\circ,\underline{\pi}_{t+1}^{*}}\|_{2}^{2} &\leq O(\delta_{t,n/2}^{2} + \|\tau_{t}^{\circ,\underline{\hat{\pi}}_{t+1}} - \tau_{t}^{n,\underline{\hat{\pi}}_{t+1}}\|_{2}^{2}) + O(n^{-\min\{\rho_{t+2}^{(c)},\rho_{t+2}^{(\Psi)},\mathcal{R}_{t+2}\}\frac{2+2\alpha}{2+\alpha}})\} \\ &+ \frac{(1 - \gamma^{T-t-1})}{1 - \gamma}c(T - t - 1)\{O(n^{-\kappa}) + Kn^{-\min\{\rho_{t+1}^{(c)},\rho_{t+1}^{(\Psi)},\mathcal{R}_{t+1}\}\frac{2+2\alpha}{2+\alpha}}\} \end{aligned}$$

$$(25)$$

from eq. (23) and appendix A.3.

Hence we obtain the inductive step and the result follows.

If we further assume that for $t' \geq t$, we have that $\rho_t^{(\cdot)} \leq \rho_{t'}^{(\cdot)}$, for $(\cdot) \in \{(c), (\Psi)\}$, i.e. the estimation error rate is nonincreasing over time, and that $\alpha > 0$ (i.e. Assumption 5, the margin assumption, holds with exponent $\alpha > 0$, then we can see from the result that the integrated risk terms obtain faster rates, hence are higher-order, and the leading term is the auxiliary estimation error of the *Q*-function contrast.

B Results used from other works

Here we collect technical lemmas from other works, stated without proof.

Lemma 6 (Lemma 18 of (Lewis and Syrgkanis, 2021)). Consider any sequence of non-negative numbers a_1, \ldots, a_m satisfying the inequality:

$$a_t \le \mu_t + c_t \max_{j=t+1}^m a_j$$

with $\mu_t, c_t \geq 0$. Let $c := \max_{t \in [m]} c_t$ and $\mu := \max_{t \in [m]} \mu_t$. Then it must also hold that:

$$a_t \le \mu \frac{c^{m-t+1} - 1}{c-1}$$

Lemma 7 (Lemma 14 of (Foster and Syrgkanis, 2019), see also results on local Rademacher complexity (Wainwright, 2019)). Consider a function class \mathcal{F} , with $\sup_{f \in \mathcal{F}} ||f||_{\infty} \leq 1$, and pick any $f^* \in \mathcal{F}$. Let $\delta_n^2 \geq \frac{4d \log(41 \log(2c_2n))}{c_2n}$ be any solution to the inequalities:

$$\forall t \in \{1, \dots, d\} : \mathcal{R} \left(\operatorname{star} \left(\left. \mathcal{F} \right|_t - f_t^{\star} \right), \delta \right) \le \delta^2.$$

Moreover, assume that the loss ℓ is L-Lipschitz in its first argument with respect to the ℓ_2 norm. Then for some universal constants c_5, c_6 , with probability $1 - c_5 \exp(c_6 n \delta_n^2)$,

 $\left|\mathbb{P}_{n}\left(\mathcal{L}_{f}-\mathcal{L}_{f^{\star}}\right)-\mathbb{P}\left(\mathcal{L}_{f}-\mathcal{L}_{f^{\star}}\right)\right|\leq18Ld\delta_{n}\left\{\|f-f^{\star}\|_{2}+\delta_{n}\right\},\quad\forall f\in\mathcal{F}.$

Hence, the outcome \hat{f} of constrained ERM satisfies that with the same probability,

$$\mathbb{P}\left(\mathcal{L}_{\hat{f}} - \mathcal{L}_{f^{\star}}\right) \leq 18Ld\delta_n\left\{\left\|\hat{f} - f^{\star}\right\|_2 + \delta_n\right\}.$$

If the loss \mathcal{L}_f is also linear in f, i.e. $\mathcal{L}_{f+f'} = \mathcal{L}_f + \mathcal{L}_{f'}$ and $\mathcal{L}_{\alpha f} = \alpha \mathcal{L}_f$, then the lower bound on δ_n^2 is not required.

C Experimental details

All experiments were ran either on a Macbook Pro M1 with 16gb RAM and 8 CPU cores or on a computer cluster with 64 CPU cores of 8gb RAM each. Experiments were run in Python using native Python, CVXPY, and scikit-learn. Each figure took approximately 3-10 minutes to generate.

1d validation example (Section 4). Following the specification of (Kallus and Uehara, 2019a, Sec 5.1), we consider a small MDP of T = 30, binary actions, univariate continuous state, initial state distribution $p(s_0) \sim \mathcal{N}(0.5, 0.2)$, transition probabilities $P_t(s_{t+1} | s_t, a_t) \sim \mathcal{N}(s + 0.3a - 0.15, 0.2)$. The target and behavior policies we consider are $\pi^e(a | s) \sim \text{Bernoulli}(p_e)$, $p_e = 0.2/(1 + \exp(-0.1s)) + 0.2U, U \sim \text{Uniform}[0, 1]$ and $\pi^b(a | s) \sim \text{Bernoulli}(p_b)$, $p_b = 0.9/(1 + \exp(-0.1s)) + 0.1U, U \sim \text{Uniform}[0, 1]$. We consider the interacted state-action basis, i.e. fit Qon s + s * a with an intercept. When Q is well-specified, we do nearly exactly recover the right contrast function; although in such a small and well-specified example we do not see benefits of orthogonality.