Linear spectroscopy of collective modes and the gap structure in two-dimensional superconductors

B. A. Levitan,¹ Y. Oreg,¹ E. Berg,¹ M. S. Rudner,² and I. Iorsh^{1,3,4}

¹Department of Condensed Matter Physics, Weizmann Institute of Science, Rehovot 7610001, Israel

²Department of Physics, University of Washington, Seattle, WA 98195-1560, USA

³Faculty of Physics, ITMO University, St. Petersburg 197101, Russia

⁴Department of Physics, Engineering Physics and Astronomy, Queen's University, Kingston, Ontario K7L 3N6, Canada

(Dated: June 14, 2024)

We consider optical response in multi-band, multi-layer two-dimensional superconductors. Within a simple model, we show that collective modes of the condensate, such as Leggett and clapping modes, can be detected in linear response. We show how trigonal warping of the superconducting order parameter can help facilitate detection of clapping modes. Taking rhombohedral trilayer graphene as an example, we consider several possible pairing mechanisms and show that all-electronic mechanisms may produce in-gap clapping modes. These modes, if present, should be detectable in the absorption of microwaves applied via gate electrodes; their detection would constitute strong evidence for unconventional pairing. Lastly, we show that absorption at frequencies above the superconducting gap $2|\Delta|$ also contains a wealth of information about the gap structure. Our results suggest that linear spectroscopy can be a powerful tool for the characterization of unconventional two-dimensional superconductors.

Keywords: Superconductivity, Leggett mode, clapping mode, rhombohedral trilayer graphene, collective mode spectroscopy.

In superconductors, a minimum of two collective modes arise: the Anderson-Bogoliubov-Goldstone (ABG) mode [1–4], corresponding to phase fluctuations of the complex order parameter, and the Higgs mode [5, 6], corresponding to amplitude fluctuations. When the superconducting order parameter has more structure, its enlarged configuration space can produce a richer diversity of collective modes [7–22]. Measurements of the collective mode spectrum can therefore provide a valuable tool for diagnosing the underlying order. However, detection of exotic collective modes is often fraught with difficulties. Means for accessing nontrivial collective modes in linear response are therefore highly desirable. In this paper, we show how linear response to straightforward AC gating can reveal exotic collective modes in superconducting few-layer systems.

As a prototypical example, consider the Leggett mode in multi-band superconductors [8]. One can think of this mode as a momentum-space analogue of the Josephson effect, consisting of relative number/phase fluctuations between the condensates formed from different bands. Assuming that intra-band interactions are attractive, when the Josephson coupling (i.e., inter-band pair swapping interaction) is not too strong, the Leggett mode can arise inside the superconducting gap, and it can therefore be underdamped. However, even when this mode is a well-defined excitation, crystal symmetries often render it invisible to linear electromagnetic response. The Leggett mode is therefore often difficult to access in experiments, requiring nonlinear response techniques such as Raman [23–25] or tunneling spectroscopy [26].

A simple gedankenexperiment illustrates the issue.

Consider two identical superconducting layers (Fig. 1a) coupled by single-particle tunneling J. The tunnel coupling lifts the accidental degeneracy between the identical layers; suppose we are in a regime where electrons occupy both resulting bands. Direct driving of the Leggett mode requires forcing at least one of its conjugate variables, the relative density and relative phase between condensates. We could manipulate the relative phase by modulating the electric potential on each of the two layers with opposite sign, amounting to a displacement field $\mathbf{D} = D\hat{\mathbf{z}}$ out of the plane (Fig. 1a-c). However, before applying the field, the system is invariant under the mirror M_z , which exchanges the two layers (and maps $M_z \mathbf{D} = -\mathbf{D}$). This implies that the spectrum of the system is an even function of D, and for small fields there are thus no energy shifts at linear order; $E(D) = E(0) + O(D^2)$. This picture also hints at what would solve the problem: perturbing around a static background field, $D = D_0 + \delta D(t)$, in which case D_0 breaks M_z and provides energy shifts linear in δD ; $D^2 = D_0^2 + 2D_0\delta D + (\delta D)^2$. We will show that this idea indeed works: the displacement field provides a useful and novel tool for enabling easy access to collective modes in few-layer superconductors. Moreover, the superconducting states in a wide variety of layered two-dimensional materials only arise in specific ranges of values of D_0 away from $D_0 = 0$, naturally giving access to their collective modes as described below.

Besides the Leggett mode, another family of exotic collective modes arise in superconductors with spontaneously-broken time-reversal symmetry. Known as "generalized clapping modes" [9, 11–13], these amount to fluctuations in the uncondensed time-reversed part-




FIG. 1. (a) A two-layer superconductor under a large constant electric displacement field D, driven by a perturbing AC field $\delta D(t)$. (b) Single-electron dispersion in the toy model (black), showing the effect of a static perturbing field (red). (c) Schematic electron-electron scattering between Fermi sheets, indicating the relative phase ($\delta\theta$) and number (δn) fluctuations of the Leggett mode. (d) Absorption spectrum given $\nu_1 = \nu_2$, $g_{11}^{(s)} = g_{22}^{(s)}$, and $\Delta_1 = \Delta_2 = \Delta$, for different values of interband coupling $g_{12}^{(s)}$ and $g^{(p)} = 0$. (e, f) The clapping modes in a chiral *p*-wave superconductor, using the toy model with $g_{\alpha\alpha'}^{(s)} = 0 > g^{(p)}$ and $|V| \gg \mu$. (e) Assuming a $p_x + ip_y$ ground state, the clapping modes at $\mathbf{q} \to 0$ correspond to fluctuations of the form $\Delta_{\mathbf{k}}(t) = \Delta_0 e^{i\varphi_{\mathbf{k}}} + (a(t) + ib(t)) e^{-i\varphi_{\mathbf{k}}}$. Solid lines show a polar plot of the equilibrium order parameter, dashed lines show the distortion associated with each mode. (f) Real (dashed) and imaginary (solid) parts of the clapping mode contribution to the electronic compressibility for small but nonzero \mathbf{q} in the circularly-symmetric model, Eq. (14); see text.

ner of the condensed ground-state channel. For example, in a $p_x + ip_y$ chiral superconductor, clapping modes arise as fluctuations in the $p_x - ip_y$ channel, as shown in Fig. 1e. Direct observation of the clapping modes in linear electromagnetic response is typically not possible at zero wavevector, due to conservation of angular momentum. However, in superconducting rhombohedral trilayer graphene (RTG) [27], which may host a chiral *p*-wave order parameter, the coherence length is likely comparable to the device size. Therefore, finite-size effects may allow even a simple uniform gate to probe the clapping mode, as we will explore.

The rest of this paper is organized as follows. First, using a minimal toy model, we show how Leggett modes can be probed at $\mathbf{q} = 0$ in systems with broken M_z symmetry, and how clapping modes can be probed at small (but nonzero) \mathbf{q} even when a high degree of rotational symmetry is present. We then consider a more realistic model of gated rhombohedral trilaver graphene (RTG). For the RTG model, we show how the observation of an in-gap collective mode can yield insight into the microscopic pairing interactions at play. We consider both conventional (phonon-mediated) and unconventional (electronic fluctuation-mediated) pairing mechanisms; only the unconventional mechanisms produce a chiral order parameter, and its associated clapping modes. We also show how the quasiparticle contribution to the linear response spectrum above $2|\Delta|$ can provide information on the superconducting gap structure.

While previous works have suggested that Leggett modes may aid in identifying the pairing mechanism in high- T_c superconductors (see, e.g., Ref. [28]), our results

show that in the case of 2D superconductors, AC gating can excite Leggett modes at *linear order* in the applied field, greatly simplifying possible experimental realizations. Additionally, other authors [22] have shown that clapping modes can modify the electronic compressibility at $\mathbf{q} \rightarrow 0$, but that this requires breaking of rotational symmetries. Working at finite \mathbf{q} , we focus on the experimentally-relevant case where (discrete) rotational symmetry is preserved, and show that the wavevector required for an appreciable signal corresponds to length scales comparable to device size in the case of RTG. Our results therefore point towards a new experimental tool for the difficult task of characterizing pairing in novel unconventional superconductors.

In order to elucidate the basic ideas at work, we begin with a toy model which captures both the Leggett and clapping modes. The essential physics we are after is captured by the Hamiltonian $H = H_0 + H_{\text{int}}$, with

$$H_{0} = \sum_{\mathbf{k}\sigma} \left(f_{\mathbf{k}t\sigma}^{\dagger} \quad f_{\mathbf{k}b\sigma}^{\dagger} \right) \begin{pmatrix} \xi_{\mathbf{k}0} + V & J \\ J & \xi_{\mathbf{k}0} - V \end{pmatrix} \begin{pmatrix} f_{\mathbf{k}t\sigma} \\ f_{\mathbf{k}b\sigma} \end{pmatrix}$$
$$= \sum_{\mathbf{k}\sigma} \left(c_{\mathbf{k}1\sigma}^{\dagger} \quad c_{\mathbf{k}2\sigma}^{\dagger} \right) \begin{pmatrix} \xi_{\mathbf{k}1} & 0 \\ 0 & \xi_{\mathbf{k}2} \end{pmatrix} \begin{pmatrix} c_{\mathbf{k}1\sigma} \\ c_{\mathbf{k}2\sigma} \end{pmatrix}$$
(1)

describing spin-1/2 fermions ($\sigma \in \{\uparrow,\downarrow\}$) in two layers $(l \in \{t,b\})$, each with bare dispersion $\xi_{\mathbf{k}0} = \epsilon_{\mathbf{k}0} - \mu$ relative to the Fermi level. We assume isotropic $\xi_{\mathbf{k}0}$ for simplicity. Single-particle tunneling (J) couples the layers, and the electric potential drops by 2V from top to bottom (we set the electron charge to 1). Including interlayer tunneling, the band energies are $\xi_{\mathbf{k},\alpha\in\{1,2\}} = \xi_{\mathbf{k}0} \pm \sqrt{J^2 + V^2}$. We assume a strong field, $|V| \gg |J|$,

such that the band splitting is essentially linear in |V|. Our analysis can be applied to any pair of Fermi surfaces with arbitrary dispersions $\xi_{\mathbf{k}\alpha}$, so we leave these general for the most part.

We model interactions using

$$H_{\rm int} = \frac{1}{L^2} \sum_{\substack{\mathbf{kk'q}\\\alpha\alpha'}} \mathcal{V}_{\mathbf{kk'}}^{\alpha\alpha'} c^{\dagger}_{\mathbf{k}\alpha\uparrow} c^{\dagger}_{-\mathbf{k}+\mathbf{q},\alpha\downarrow} c_{-\mathbf{k'}+\mathbf{q},\alpha'\downarrow} c_{\mathbf{k'}\alpha'\uparrow}, \quad (2)$$

focusing on small \mathbf{q} (i.e. the Cooper channel). L is the linear dimension of the sample. For concreteness, we take

$$\mathcal{V}_{\mathbf{k}\mathbf{k}'}^{\alpha\alpha'} = g_{\alpha\alpha'}^{(s)} + 2g^{(p)}\cos(\varphi_{\mathbf{k}} - \varphi_{\mathbf{k}'})\delta_{\alpha\alpha'},\qquad(3)$$

where $g_{\alpha\alpha}^{(s)} < 0$ binds *s*-wave pairs in band α , $g_{12}^{(s)} = \left(g_{21}^{(s)}\right)^*$ Josephson couples the two bands, and $g^{(p)} < 0$ binds *p*-wave pairs. Here $\varphi_{\mathbf{k}}$ is the angle of \mathbf{k} relative to the k_x axis, i.e., $\mathbf{k} = |\mathbf{k}| (\cos \varphi_{\mathbf{k}}, \sin \varphi_{\mathbf{k}})$. Note that with the spin indices as written in Eq. (2), when considering *p*-wave pairing, we implicitly consider the $m_z = 0$ triplet component only; the extension to the more general case is straightforward.

We base our analysis [29] on the imaginary-time path integral. To identify the collective modes, we perform a Hubbard-Stratonovich decoupling in the Cooper channel, and expand the effective action to quadratic order in fluctuations ϕ around the mean-field saddle point [30]. Analytically continuing the Gaussian effective action to real frequency $(i\Omega_m \to \Omega + i0^+)$, the collective modes appear as poles of the bosonic propagator for pair fluctuations. To evaluate the linear response to applied electric fields, we also compute the couplings between the collective modes and such fields.

First we focus on the Leggett mode, by setting $g^{(p)} = 0$ and $g^{(s)}_{\alpha\alpha'} \neq 0$; afterwards we will instead set $g^{(s)}_{\alpha\alpha'} = 0$ and $g^{(p)} < 0$ to focus on the clapping modes. The Leggett mode requires $\mu > \sqrt{J^2 + V^2}$ such that both bands are occupied. Leggett's original paper analyzed essentially this model—at $\mathbf{q} = 0$ and with $g^{(p)} = 0$ —and found its relative-phase mode [8]. We assume $g_{\alpha\alpha} < 0$. Taking $|\Delta_1| \leq |\Delta_2|$ without further loss of generality, we find [29] that an in-gap Leggett mode requires $0 \leq |g_{12}^{(s)}|/(g_{11}^{(s)}g_{22}^{(s)} - |g_{12}^{(s)}|^2) < \nu_2 \arcsin(|\Delta_1/\Delta_2|)/\sqrt{1 - (\Delta_1/\Delta_2)^2}$, where ν_{α} is the density of states per unit area per spin on the Fermi surface of band α in the normal state. (In the symmetric case where $|\Delta_1| = |\Delta_2|$, the second inequality is irrelevant).

We now show how an AC modulation of the displacement field can probe the Leggett mode. The drive field corresponds to a small time-dependent perturbation around the static background, $V \rightarrow V_{\rm DC} + \delta V(t)$. Keeping only terms up to linear order in δV and working in the band basis, driving corresponds to a Hamiltonian term $H_{\delta V} = \sum_{\mathbf{k}\sigma} \delta V(t) (V_{\rm DC}/\sqrt{J^2 + V_{\rm DC}^2}) (c_{\mathbf{k}1\sigma}^{\dagger} c_{\mathbf{k}1\sigma} -$ $c_{\mathbf{k}2\sigma}^{\dagger}c_{\mathbf{k}2\sigma}$). Integrating out the fermions in the presence of the drive yields a quadratic term $\sim \delta V^2$ and source terms $\sim \phi \, \delta V$ in the effective action, on top of the Gaussian action for pair fluctuations $\sim -\phi^{\dagger} \mathcal{D}^{-1} \phi$. Integrating out the fluctuations ϕ then yields the full quadratic free energy functional $\mathcal{F}[\delta V]$, describing a capacitance (dependent on frequency Ω):

$$\frac{1}{L^2}\mathcal{F}[\delta V(\Omega)] = \frac{1}{2}\mathcal{C}(\Omega)\delta V(-\Omega)\delta V(\Omega).$$
(4)

We provide an explicit calculation of the capacitance per unit area $C(\Omega)$ in the Supplemental Material [29]. In the symmetric case where $\Delta_1 = \Delta_2 = \Delta > 0$ and $\nu_1 = \nu_2 = \nu$, the collective-mode contribution is

$$C_{\phi}(\Omega) = \frac{4V_{\rm DC}^2}{V_{\rm DC}^2 + J^2} \frac{(\nu\gamma \sec \gamma)^2}{\nu\gamma \tan \gamma - 2\tilde{g}^{-1}},\tag{5}$$

where

$$\tilde{g}^{-1} = \frac{\left|g_{12}^{(s)}\right|}{g_{11}^{(s)}g_{22}^{(s)} - \left|g_{12}^{(s)}\right|^2} \tag{6}$$

and

$$\gamma = \arcsin\left(\frac{\Omega + i0^+}{2\Delta}\right). \tag{7}$$

The Leggett mode frequency $\Omega_{\rm L}$ solves $\nu \gamma_{\rm L} \tan \gamma_{\rm L} = 2\tilde{g}^{-1}$, with $\sin \gamma_{\rm L} = \Omega_{\rm L}/(2\Delta)$. Near the positive-frequency pole, we can approximate

$$\mathcal{C}_{\phi}(\Omega) \approx \frac{4V_{\rm DC}^2}{V_{\rm DC}^2 + J^2} \frac{\nu\gamma_{\rm L}^2}{\sin\gamma_{\rm L} + \gamma_{\rm L}\sec\gamma_{\rm L}} \frac{2\Delta}{\Omega - \Omega_{\rm L} + i0^+}.$$
(8)

In the general (asymmetric) case, the key results are that at $\mathbf{q} \to 0$: i) The amplitude modes and ABG mode do not contribute to $C(\Omega)$; and ii) The Leggett mode does contribute an in-gap pole to $C(\Omega)$, provided that the background field $V_{\text{DC}} \neq 0$. We make no attempt to explicitly compute the width of the pole, which is controlled by higher-order processes, and is model-dependent. However, we expect the resonance to be narrow, since no quasiparticle excitations are available to facilitate decay through low-order processes.

Fig. 1d shows the total absorption $(-\operatorname{Im} \mathcal{C}, \operatorname{including} quasiparticle and collective-mode contributions) for the symmetric toy model, i.e., for <math>\Delta_1 = \Delta_2, \nu_1 = \nu_2$, and $g_{11}^{(s)} = g_{22}^{(s)}$. In this case, when the interactions become uniform $(g_{12}^{(s)} \to g_{11}^{(s)} = g_{22}^{(s)})$, the absorption peak corresponding to the Leggett mode approaches the gap edge¹.

¹ Since number and phase are canonically-conjugate variables, one expects that Fermi-liquid terms of the form $H_{\rm FL}$ =

We now turn our attention to the clapping modes, setting $g_{\alpha\alpha'}^{(s)} = 0 > g^{(p)}$ and $|V| \gg \mu$. We project into the occupied (lower) band, and hence drop the band index α . The gap equation has two degenerate solutions, corresponding to $p_x \pm ip_y$ pairing. We assume the $p_x + ip_y$ channel condenses, i.e., $\Delta_{\mathbf{k}} = \Delta_0 e^{i\varphi_{\mathbf{k}}}$ with $\Delta_0 > 0$. As detailed in the Supplemental Material [29], fluctuations in the uncondensed $p_x - ip_y$ channel give rise to two real bosons a and b; these are the clapping modes, which in this symmetric minimal model are degenerate at $\Omega = \sqrt{2}\Delta_0$.

To show how an AC electric field can probe the clapping modes, as in the Leggett case, we perturb around the static background, $V \rightarrow V_{\rm DC} + \delta V(t)$. Since we projected out the upper band, the only effect of the modulation is to shift the energy of the lower band; when the displacement field-induced interlayer potential is very large compared with the interlayer hopping matrix element, the electrons only live on one layer, so they only feel the perturbing field through its local potential. Therefore, we calculate the clapping mode contribution to the electronic compressibility Π^{00} . Unlike the Leggett mode, the clapping modes carry angular momentum 2 relative to the ground state; this means that with circular symmetry, their couplings to the scalar potential must vanish at $\mathbf{q} = 0$. Hence, we calculate their compressibility contributions to leading nonvanishing order in $|\mathbf{q}|$.

Our calculation [29] mirrors that of Ref. [22], generalized to nonzero momentum. After Hubbard-Stratonovich decoupling, the fermions encounter the total (fluctuating) pairing field

$$\Delta_{\mathbf{k},q} = \Delta_q^{(+)} e^{i\varphi_{\mathbf{k}}} + \Delta_q^{(-)} e^{-i\varphi_{\mathbf{k}}}.$$
(9)

Here $q = (i\Omega_m, \mathbf{q})$ contains both the bosonic Matsubara frequency $\Omega_m = 2\pi mT$ and momentum \mathbf{q} . In coordinate space, writing $x = (\tau, \mathbf{r})$, the Hubbard-Stratonovich bosons corresponding to the two pairing channels are

$$\Delta^{(+)}(x) = e^{i\theta(x)}(\Delta_0 + h(x))$$
(10)

and

$$\Delta^{(-)}(x) = e^{i\theta(x)}(a(x) + ib(x)).$$
(11)

Here θ is the global phase (ABG) mode, *h* the Higgs mode, and *a* and *b* the clapping modes. After integrating out the fermions, the long-wavelength action at quadratic

order is

$$S = \sum_{q} \left(\sum_{\mu,\nu=0,x,y} \Pi_{q}^{\mu\nu} \mathcal{A}_{-q}^{\mu} \mathcal{A}_{q}^{\nu} - \mathcal{D}_{a,q}^{-1} a_{-q} a_{q} - \mathcal{D}_{b,q}^{-1} b_{-q} b_{q} + \sum_{\mu=0,x,y} \left[\Pi_{q}^{\mu a} \mathcal{A}_{-q}^{\mu} a_{q} + \Pi_{q}^{\mu b} \mathcal{A}_{-q}^{\mu} b_{q} \right] \right), \quad (12)$$

where $\mathcal{A} = (\mathcal{A}^0, \mathcal{A}) = (\mathcal{A}^0 + \partial_\tau \theta, \mathbf{A} - \nabla \theta)$ is the gaugeinvariant combination of the electromagnetic gauge potential A and the spacetime gradient of the global phase θ . Since we are only interested in the impact of the clapping modes on the compressibility, we drop all terms involving the spatial component \mathcal{A} for simplicity. Physically, we expect that this approximation should not drastically change the result, since the ABG mode associated with $\nabla \theta$ is either at a much lower energy (in a neutral superfluid) or much higher energy (after the Anderson-Higgs mechanism in a superconductor) than the clapping mode, for small but nonzero q. We provide explicit expressions for the propagators \mathcal{D}_{Xq} and couplings Π_q^{0X} $(X \in \{a, b\})$ in the Supplemental Material [29]. Note that with circular symmetry, Π_q^{0X} vanishes at $\mathbf{q} = 0$ and first shows up at $O(|\mathbf{q}|^2)$. The product $\Pi_q^{0X} \mathcal{A}_{-q}^0 X_q$ must be scalar, so two factors of momentum are required to balance the angular momentum 2 of the clapping modes.

We obtain the clapping mode contribution to the compressibility $\delta \Pi_q^{00}$ by integrating out those modes, yielding

$$\delta \Pi_q^{00} = \frac{1}{4} \Pi_{-q}^{0a} \mathcal{D}_{a,q} \Pi_q^{0a} + \frac{1}{4} \Pi_{-q}^{0b} \mathcal{D}_{b,q} \Pi_q^{0b}.$$
 (13)

Analytically continuing to real time $(i\Omega_m \to \Omega + i0^+)$, we find to leading order in **q** and for large μ :

 $\delta \Pi_a^{00}$

$$= \frac{\nu}{2} \left(\frac{\pi}{4} \xi_{\rm BCS} |\mathbf{q}|\right)^4 \csc^2 \gamma \sec^4 \gamma \frac{1 - \gamma^2 (\cot \gamma - \tan \gamma)^2}{\gamma (\cot \gamma - \tan \gamma)}.$$
(14)

Here $\xi_{\rm BCS} = v_{\rm F}/(\pi\Delta_0)$ is the BCS coherence length, $v_{\rm F} = \sqrt{2\mu/m}$ is the Fermi velocity, and

$$\gamma = \arcsin\left(\frac{\Omega + i0^+}{2\Delta_0}\right). \tag{15}$$

This result neglects the dispersion of the clapping modes. Note the pole at $\cot \gamma = \tan \gamma$, corresponding to $\Omega = \pm \sqrt{2}\Delta_0$: the clapping modes are visible in the compressibility, and hence in the capacitance. In the vicinity of the positive-frequency pole,

$$\delta \Pi_q^{00} \approx \nu \left(\frac{\pi}{4}\right)^3 \left(\xi_{\rm BCS} |\mathbf{q}|\right)^4 \frac{-\sqrt{2}\Delta_0}{\Omega - \sqrt{2}\Delta_0 + i0^+}.$$
 (16)

 $g_{\rm FL} (n_1 - n_2 - \langle n_1 - n_2 \rangle_0)^2$ might shift the Leggett mode frequency. Here $\langle \dots \rangle_0$ denotes an equilibrium average. As Leggett anticipated [8], attractive interactions of this form $(g_{\rm FL} < 0)$ can pull the mode into the gap even if \hat{g} only has a single attractive eigenvalue (we show this explicitly in the Supplemental Material [29]).




FIG. 2. (a) Electron dispersion of RTG under a static perpendicular electric field corresponding to a potential drop of 20 meV (black) or 30 meV (orange) between the top and bottom layers. Here, k_x is measured relative to the K point. (b) Fermi surfaces of RTG in each of its two valleys, for a potential drop of 20 meV and a Fermi energy of -29.3 meV. The line cut through the dispersion relation (panel (a)) is along the dotted line schematically indicated by the dashed line in panel (b). (c-d) Absorption spectrum and gap structure for (c) phonon-mediated s-wave pairing, and (d) intervalley coherence fluctuation-mediated chiral p-wave pairing. Note the in-gap clapping mode in the p-wave case. The gap curves are colored according to the Fermi surface component on which they reside, matching panel (b).

As explained above, the factor of $|\mathbf{q}|^4$ in Eq. (14) is a consequence of circular symmetry. To obtain a signal at lower order in $|\mathbf{q}|$, something must break that symmetry in order to relax the angular momentum constraint. If the symmetry is broken completely, for example by a $\cos(\zeta)p_x + i\sin(\zeta)p_y$ order parameter with ζ away from high-symmetry values (as in Ref. [22]), then a signal can even arise at $\mathbf{q} \to 0$. A less drastic scenario is that rotation is explicitly broken down to a discrete subgroup dictated by the lattice structure. With C_{3v} symmetry (possessed by RTG, for example), the couplings Π_q^{0X} need only contain a single factor of \mathbf{q} , since $2 = -1 \pmod{3}$. We find [29] that trigonal warping indeed yields a nonvanishing result at $O(|\mathbf{q}|^2)$, of similar qualitative form to Eq. (14), but suppressed by the small factor $1/(\xi_{\text{BCS}}k_{\text{F}})^2 \sim (\Delta_0/\mu)^2$.

To recap, so far we have studied a toy model which captures both the Leggett and clapping modes. We showed that measurements of the AC capacitance (i.e., the response to a perturbing out-of-plane displacement field) can probe both of these classes of mode in 2D few-layer superconductors. Lastly, we pointed out that trigonal warping allows detection of the clapping modes at lower order in $|\mathbf{q}|$ by relaxing the absolute conservation of angular momentum to conservation mod 3, in keeping with C_{3v} symmetry. We now move on to superconducting RTG as an illustrative case study.

We model the single-particle bandstructure of RTG using a 6×6 tight-binding Hamiltonian in its continuum limit, written in the basis of two atoms per unit cell in each of the three layers [29, 31, 32]. We limit ourselves to the range of parameters (static perpendicular displacement field and doping level) where superconductivity was observed [27], focusing on the SC2 phase. In this regime, prior to the onset of superconductivity, RTG contains one annular Fermi sea in each valley (two in total), whose Fermi surfaces show substantial trigonal warping (see Fig. 2b). The specific pairing mechanism responsible for superconductivity in RTG is yet to be determined, so we consider conventional (s wave, mediated by phonons) and unconventional (chiral p wave, mediated by fluctuations of either charge density, as in the Kohn-Luttinger mechanism [33–36], or intervalley coherence [37, 38]) scenarios [29].

Figures 2c,d show our numerical results for the absorption spectrum of superconducting RTG in the SC2 phase, accounting for the Anderson-Higgs mechanism, and working to leading order in $|\Delta|/\mu$ [29] (hence neglecting the weak $O(|\mathbf{q}|^2)$ contribution discussed above, which is permitted by the substantial trigonal warping present in RTG). As in the toy model discussion, the absorption corresponds to the imaginary part of the capacitance. Fig. 2c shows the results for the s-wave case. Fig. 2d shows the results for the IVC-mediated *p*-wave case; the Kohn-Luttinger mechanism gives qualitatively similar results. Most importantly, the all-electronic mechanisms both show an in-gap clapping mode. While the clapping mode is visible only away from $\mathbf{q} = 0$, since superconducting RTG devices are comparable in size to the superconducting coherence length ξ_{BCS} , we expect finite size effects to render the clapping mode visible even without any special patterning of the gate electrodes.

Figs. 2c,d also show that even without the collective modes, the absorption spectrum contains rich information about the gap structure. Extrema of the superconducting gap reveal themselves as sharp features in the absorption.

To conclude, we have shown that linear spectroscopy can be an invaluable tool for studying the pairing mechanisms in 2D superconductors. Using a simple but powerful toy model, we showed how various exotic collective modes should be detectable in linear response, either at $\mathbf{q} = 0$ (in the case of the Leggett mode), or at finite \mathbf{q} (in the case of the clapping modes). By considering the example of superconducting rhombohedral trilayer graphene (RTG) under a background vertical displacement field, we further showed how the observation of an in-gap clapping mode could yield compelling evidence for unconventional, all-electronic pairing mechanisms. We also showed how linear spectroscopy above the quasiparticle excitation gap reveals information on the gap structure, in the form of sharp features at frequencies corresponding to gap extrema over the Fermi surface.

The physics we have described should be accessible in experiments. The superconducting phases in RTG have transition temperatures at the scale of between 50 and 100 mK [27], so one expects the gap and possible in-gap clapping modes to reside at the scale of a few GHz to 10s of GHz, easily within the range of microwave function generators. Further, we expect that similar phenomenology may occur in other quasi-two-dimensional superconductors which are sensitive to a vertical displacement field, such as Bernal bilayer [39, 40] and twisted trilayer [41–44] graphene. Future work should study the collective mode spectroscopy of these systems.

Acknowledgements: B. A. L. was hosted at the Institute for Theoretical Physics at the University of Cologne during a considerable portion of the preparation of this manuscript, and gratefully acknowledges their generous support, as well as the financial support of the Zuckerman STEM Leadership Program. This work was supported by NSF-BSF award DMR-2310312, by the European Union's Horizon 2020 research and innovation programme (grant agreements LEGOTOP No. 788715 and HQMAT No. 817799), and the DFG CRC SFB/TRR183. M. R. gratefully acknowledges support from the Brown Investigator Award, a program of the Brown Science Foundation, the University of Washington College of Arts and Sciences, and the Kenneth K. Young Memorial Professorship.

- P. W. Anderson, Random-phase approximation in the theory of superconductivity, Phys. Rev. 112, 1900 (1958).
- [2] P. W. Anderson, New method in the theory of superconductivity, Phys. Rev. 110, 985 (1958).
- [3] N. Bogoljubov, V. V. Tolmachov, and D. Širkov, A new method in the theory of superconductivity, Fortschritte der physik 6, 605 (1958).
- [4] G. Rickayzen, Collective excitations in the theory of superconductivity, Phys. Rev. 115, 795 (1959).
- [5] P. B. Littlewood and C. M. Varma, Amplitude collective modes in superconductors and their coupling to chargedensity waves, Phys. Rev. B 26, 4883 (1982).
- [6] R. Shimano and N. Tsuji, Higgs mode in superconductors, Annual Review of Condensed Matter Physics 11, 103 (2020).
- [7] A. Bardasis and J. R. Schrieffer, Excitons and plasmons in superconductors, Phys. Rev. 121, 1050 (1961).
- [8] A. Leggett, Number-phase fluctuations in two-band superconductors, Progress of Theoretical Physics 36, 901 (1966).
- [9] P. Wölfle, Order-parameter collective modes in ³He a, Phys. Rev. Lett. 37, 1279 (1976).
- [10] D. S. Hirashima and H. Namaizawa, Collective modes

and a theory of response of d-wave superconductors, Journal of low temperature physics **73**, 137 (1988).

- [11] H.-Y. Kee, Y. B. Kim, and K. Maki, Collective modes and sound propagation in a p-wave superconductor: Sr₂RuO₄, Phys. Rev. B **62**, 5877 (2000).
- [12] S. Higashitani and K. Nagai, Order parameter collective modes in Sr₂RuO₄, Physica B: Condensed Matter 284, 539 (2000).
- [13] A. V. Balatsky, P. Kumar, and J. R. Schrieffer, Collective mode in a superconductor with mixed-symmetry order parameter components, Phys. Rev. Lett. 84, 4445 (2000).
- [14] R. Roy and C. Kallin, Collective modes and electromagnetic response of a chiral superconductor, Phys. Rev. B 77, 174513 (2008).
- [15] Y. Ota, M. Machida, T. Koyama, and H. Aoki, Collective modes in multiband superfluids and superconductors: Multiple dynamical classes, Phys. Rev. B 83, 060507 (2011).
- [16] S.-Z. Lin and X. Hu, Massless Leggett mode in threeband superconductors with time-reversal-symmetry breaking, Phys. Rev. Lett. 108, 177005 (2012).
- [17] V. Stanev, Model of collective modes in three-band superconductors with repulsive interband interactions, Phys. Rev. B 85, 174520 (2012).
- [18] C. M. Puetter and H.-Y. Kee, Identifying spin-triplet pairing in spin-orbit coupled multi-band superconductors, Europhysics Letters 98, 27010 (2012).
- [19] K. Kobayashi, M. Machida, Y. Ota, and F. Nori, Massless collective excitations in frustrated multiband superconductors, Phys. Rev. B 88, 224516 (2013).
- [20] S.-Z. Lin, Ground state, collective mode, phase soliton and vortex in multiband superconductors, Journal of Physics: Condensed Matter 26, 493202 (2014).
- [21] Z. Sun, M. M. Fogler, D. N. Basov, and A. J. Millis, Collective modes and terahertz near-field response of superconductors, Phys. Rev. Res. 2, 023413 (2020).
- [22] N. R. Poniatowski, J. B. Curtis, A. Yacoby, and P. Narang, Spectroscopic signatures of time-reversal symmetry breaking superconductivity, Commun. Phys. 5, 44 (2022).
- [23] G. Blumberg, A. Mialitsin, B. S. Dennis, M. V. Klein, N. D. Zhigadlo, and J. Karpinski, Observation of Leggett's collective mode in a multiband MgB₂ superconductor, Phys. Rev. Lett. **99**, 227002 (2007).
- [24] F. Giorgianni, T. Cea, C. Vicario, C. P. Hauri, W. K. Withanage, X. Xi, and L. Benfatto, Leggett mode controlled by light pulses, Nature Physics 15, 341 (2019).
- [25] S. Z. Zhao, H.-Y. Song, L. L. Hu, T. Xie, C. Liu, H. Q. Luo, C.-Y. Jiang, X. Zhang, X. C. Nie, J.-Q. Meng, Y.-X. Duan, S.-B. Liu, H.-Y. Xie, and H. Y. Liu, Observation of soft Leggett mode in superconducting CaKFe₄As₄, Phys. Rev. B **102**, 144519 (2020).
- [26] Y. G. Ponomarev, S. Kuzmichev, M. Mikheev, M. Sudakova, S. Tchesnokov, N. Timergaleev, A. Yarigin, E. Maksimov, S. Krasnosvobodtsev, A. Varlashkin, <u>et al.</u>, Evidence for a two-band behavior of MgB₂ from pointcontact and tunneling spectroscopy, Solid state communications **129**, 85 (2004).
- [27] H. Zhou, T. Xie, T. Taniguchi, K. Watanabe, and A. F. Young, Superconductivity in rhombohedral trilayer graphene, Nature 598, 434 (2021).
- [28] W. Huang, M. Sigrist, and Z.-Y. Weng, Identifying the dominant pairing interaction in high-T_c FeSe superconductors through Leggett modes, Phys. Rev. B 97, 144507

(2018).

- [29] Please find the details of our analysis and our RTG model in the Supplementary Material at URL_will_be_ inserted_by_publisher.
- [30] S. Sharapov, V. Gusynin, and H. Beck, Effective action approach to the Leggett's mode in two-band superconductors, The European Physical Journal B-Condensed Matter and Complex Systems **30**, 45 (2002).
- [31] F. Zhang, B. Sahu, H. Min, and A. H. MacDonald, Band structure of *ABC*-stacked graphene trilayers, Phys. Rev. B 82, 035409 (2010).
- [32] H. Zhou, T. Xie, A. Ghazaryan, T. Holder, J. R. Ehrets, E. M. Spanton, T. Taniguchi, K. Watanabe, E. Berg, M. Serbyn, and A. F. Young, Half-and quarter-metals in rhombohedral trilayer graphene, Nature **598**, 429 (2021).
- [33] A. Ghazaryan, T. Holder, M. Serbyn, and E. Berg, Unconventional superconductivity in systems with annular fermi surfaces: Application to rhombohedral trilayer graphene, Phys. Rev. Lett. **127**, 247001 (2021).
- [34] T. Cea, P. A. Pantaleón, V. o. T. Phong, and F. Guinea, Superconductivity from repulsive interactions in rhombohedral trilayer graphene: A Kohn-Luttinger-like mechanism, Phys. Rev. B 105, 075432 (2022).
- [35] W. Qin, C. Huang, T. Wolf, N. Wei, I. Blinov, and A. H. MacDonald, Functional renormalization group study of superconductivity in rhombohedral trilayer graphene, Phys. Rev. Lett. **130**, 146001 (2023).
- [36] A. Ghazaryan, T. Holder, E. Berg, and M. Serbyn, Multilayer graphenes as a platform for interaction-driven physics and topological superconductivity, Phys. Rev. B 107, 104502 (2023).
- [37] S. Chatterjee, T. Wang, E. Berg, and M. P. Zaletel, Intervalley coherent order and isospin fluctuation mediated superconductivity in rhombohedral trilayer graphene, Nat. Commun. 13, 6013 (2022).
- [38] Z. Dong, P. A. Lee, and L. S. Levitov, Signatures of cooper pair dynamics and quantum-critical superconductivity in tunable carrier bands, Proceedings of the National Academy of Sciences **120**, e2305943120 (2023).
- [39] H. Zhou, L. Holleis, Y. Saito, L. Cohen, W. Huynh, C. L. Patterson, F. Yang, T. Taniguchi, K. Watanabe, and A. F. Young, Isospin magnetism and spin-polarized superconductivity in bernal bilayer graphene, Science **375**, 774 (2022).
- [40] Y. Zhang, R. Polski, A. Thomson, É. Lantagne-Hurtubise, C. Lewandowski, H. Zhou, K. Watanabe, T. Taniguchi, J. Alicea, and S. Nadj-Perge, Enhanced superconductivity in spin-orbit proximitized bilayer graphene, Nature **613**, 268 (2023).
- [41] G. Chen, A. L. Sharpe, P. Gallagher, I. T. Rosen, E. J. Fox, L. Jiang, B. Lyu, H. Li, K. Watanabe, T. Taniguchi, et al., Signatures of tunable superconductivity in a trilayer graphene moiré superlattice, Nature 572, 215 (2019).
- [42] J. M. Park, Y. Cao, K. Watanabe, T. Taniguchi, and P. Jarillo-Herrero, Tunable strongly coupled superconductivity in magic-angle twisted trilayer graphene, Nature **590**, 249 (2021).
- [43] Z. Hao, A. Zimmerman, P. Ledwith, E. Khalaf, D. H. Najafabadi, K. Watanabe, T. Taniguchi, A. Vishwanath, and P. Kim, Electric field-tunable superconductivity in alternating-twist magic-angle trilayer graphene, Science **371**, 1133 (2021).

[44] H. Kim, Y. Choi, C. Lewandowski, A. Thomson, Y. Zhang, R. Polski, K. Watanabe, T. Taniguchi, J. Alicea, and S. Nadj-Perge, Evidence for unconventional superconductivity in twisted trilayer graphene, Nature 606, 494 (2022).

Supplemental Material: Linear spectroscopy of collective modes and the gap structure in two-dimensional superconductors

(Dated: June 14, 2024)

CONTENTS

I There are a dely I a month and a de	1
I. Toy model: Leggett mode	1
A. Mean-field theory	2
B. Collective modes	3
C. Driving the Leggett mode	5
II. Effect of Fermi-liquid terms on the Leggett mode	7
III. Toy model: Clapping modes	9
A. Mean-field theory	9
B. Collective modes and compressibility	9
C. Trigonal warping	14
IV. Collective modes for a general static interaction	15
A. Setup	15
B. Projection to the Fermi surface	17
1. Evaluating the polarization components	19
2. Evaluating the driving terms	22
V. Single particle Hamiltonian of RTG	28
VI. Interaction potentials	29
VII. Self-consistent electromagnetic response	29
VIII. Clapping mode dispersion for RTG	31
References	31

I. TOY MODEL: LEGGETT MODE

In this section we consider a minimal model containing a Leggett mode,

$$S[\bar{c},c] = S_0 + S_{\rm int} = \sum_{k\alpha\sigma} \bar{c}_{k\alpha\sigma} \left[-i\omega_n + \xi_{\mathbf{k}\alpha} \right] c_{k\alpha\sigma} + \frac{T}{L^2} \sum_{\substack{kk'q\\\alpha\alpha'}} g_{\alpha\alpha'} \bar{c}_{k\alpha\uparrow} \bar{c}_{-k+q,\alpha\downarrow} c_{-k'+q,\alpha'\downarrow} c_{k'\alpha'\uparrow}.$$
 (1)

This action corresponds to the toy model given in the main text, with $g^{(p)} = 0$. L is the linear size of the periodic system. k denotes the three-vector $(i\omega_n, \mathbf{k})$, with $\omega_n = (2n+1)\pi T$ a fermionic Matsubara frequency at temperature T (analogously for k'). q is a similar three-vector, but with a bosonic frequency component $\Omega_m = 2\pi m T$. $\alpha \in \{1, 2\}$ labels the two bands with dispersion relations given by $\xi_{\mathbf{k}\alpha}$ relative to the Fermi level. $\sigma \in \{\uparrow, \downarrow\}$ labels the spin. We assume attractive intra-band interactions $g_{\alpha\alpha} < 0$. Since the single-particle action S_0 conserves charge in each band separately, any phase factor on $g_{12} = g_{21}^*$ may be freely gauged away, so we choose $g_{12} = g_{21} \leq 0$ without loss of generality. This model was studied by Leggett [1]; we present an analysis in modern language to set the stage for the more technically-involved, but conceptually-analogous case with momentum-dependent interactions. The Gaussian pair-fluctuation (GPF) approach we apply is standard [2], so we skip most details and present only the important results.

A. Mean-field theory

The GPF approach is a leading-order expansion around mean-field theory, and begins with a standard Hubbard-Stratonovich decoupling in the Cooper channel. Denoting the corresponding complex auxiliary bosons by Δ_{α} and introducing the Nambu spinors $\Psi_{k\alpha} = (c_{k,\alpha\uparrow} \ \bar{c}_{-k,\alpha\downarrow})^T$, the resulting action is

$$S[\Psi^{\dagger}, \Psi, \overline{\Delta}, \Delta] = -\frac{L^2}{T} \sum_{q} \left(\overline{\Delta}_{q1} \ \overline{\Delta}_{q2} \right) \hat{g}^{-1} \begin{pmatrix} \Delta_{q1} \\ \Delta_{q2} \end{pmatrix} - \sum_{\alpha kq} \Psi^{\dagger}_{\alpha} \mathbb{G}_{\alpha}^{-1} \Psi_{\alpha}.$$
(2)

The inverse fermion propagator is

$$\mathbb{G}_{\alpha;k,k-q}^{-1} = \begin{pmatrix} (i\omega_n - \xi_{\mathbf{k}\alpha})\delta_{q,0} & \Delta_{q,\alpha} \\ \overline{\Delta}_{-q,\alpha} & (i\omega_n + \xi_{\mathbf{k}\alpha})\delta_{q,0} \end{pmatrix}.$$
(3)

The couplings are grouped into the matrix

$$\hat{g} = \begin{pmatrix} g_{11} & g_{12} \\ g_{12} & g_{22} \end{pmatrix}.$$
(4)

We assume that $g_{\alpha\alpha} < 0$ and det $\hat{g} > 0$ such that both eigenvalues are positive, i.e., the interactions are purely attractive. A repulsive eigenvalue would slightly modify the Hubbard-Stratonovich transformation; see section IV. Note that for convenience, we use different Fourier-transform conventions for the fermions vs. the pairing fields. We use the unitary normalization for the fermions,

$$\psi_{\alpha}(\tau, \mathbf{r}) = \sqrt{\frac{T}{L^2}} \sum_{k} e^{i(\mathbf{k} \cdot \mathbf{r} - \omega_n \tau)} c_{k\alpha} \quad \text{and} \quad c_{k\alpha} = \sqrt{\frac{T}{L^2}} \int d\tau \, d^2 \mathbf{r} \, e^{-i(\mathbf{k} \cdot \mathbf{r} - \omega_n \tau)} \psi_{\alpha}(\tau, \mathbf{r}), \tag{5}$$

but the non-unitary normalization for the pairing fields,

$$\Delta(\tau, \mathbf{r}) = \sum_{q} e^{i(\mathbf{q} \cdot \mathbf{r} - \Omega_m \tau)} \Delta_q \quad \text{and} \quad \Delta_q = \frac{T}{L^2} \int d\tau \, d^2 \mathbf{r} \, e^{-i(\mathbf{q} \cdot \mathbf{r} - \Omega_m \tau)} \Delta(\tau, \mathbf{r}). \tag{6}$$

This choice is made so that in the momentum representation, the pairing terms look like $\Delta c^{\dagger}c^{\dagger}$ + H.c. without any normalization factor.

Integrating out the fermion yields the famous "Tr log" action for the HS fields,

$$S[\overline{\Delta}, \Delta] = -\frac{L^2}{T} \sum_{q} \left(\overline{\Delta}_{q1} \ \overline{\Delta}_{q2} \right) \hat{g}^{-1} \begin{pmatrix} \Delta_{q1} \\ \Delta_{q2} \end{pmatrix} - \operatorname{Tr} \log \mathbb{G}^{-1}.$$
(7)

The capital-T trace (Tr) includes three-momentum indices, while the lowercase-t trace (tr) used below is an ordinary matrix trace over Nambu indices only. The mean-field ground state is the saddle point $\delta S/\delta \overline{\Delta}_{\alpha} = 0$, corresponding to the coupled gap equations

$$\Delta_{\alpha} = -\sum_{\alpha'} \int \frac{\mathrm{d}^2 \mathbf{k}'}{(2\pi)^2} g_{\alpha\alpha'} \tanh\left(\frac{E_{\mathbf{k}'\alpha'}}{2T}\right) \frac{\Delta_{\alpha'}}{2E_{\mathbf{k}'\alpha'}},\tag{8}$$

or

$$\frac{g_{22}\Delta_1 - g_{12}\Delta_2}{\det\hat{g}} = -\int \frac{\mathrm{d}^2 \mathbf{k}}{(2\pi)^2} \tanh\left(\frac{E_{\mathbf{k}1}}{2T}\right) \frac{\Delta_1}{2E_{\mathbf{k}1}} \tag{9a}$$

and

$$\frac{-g_{12}\Delta_1 + g_{11}\Delta_2}{\det \hat{g}} = -\int \frac{\mathrm{d}^2 \mathbf{k}}{(2\pi)^2} \tanh\left(\frac{E_{\mathbf{k}2}}{2T}\right) \frac{\Delta_2}{2E_{\mathbf{k}2}}.$$
(9b)

The quasiparticle energies are $E_{\mathbf{k}\alpha} = \sqrt{\xi_{\mathbf{k}\alpha}^2 + |\Delta_{\alpha}|^2}$. Note that the solutions Δ_{α} must be either in phase, or out of phase by π . So, we choose both to be real. Now we trade the momentum integrals for energy integrals via the densities of states ν_{α} , and introduce an energy cutoff Λ :

$$\int \frac{\mathrm{d}^2 \mathbf{k}}{(2\pi)^2} \to \int_{-\Lambda}^{\Lambda} \mathrm{d}\xi \,\nu_\alpha \int_0^{2\pi} \frac{\mathrm{d}\varphi}{2\pi}.$$
(10)

Then we can integrate the gap equations at $T \to 0$ as standard, yielding (for $\Lambda \gg |\Delta_{\alpha}|$)

$$\frac{g_{22}\Delta_1 - g_{12}\Delta_2}{\det \hat{g}} = -\nu_1 \Delta_1 \log\left(\frac{2\Lambda}{|\Delta_1|}\right) \quad \text{and} \quad \frac{-g_{12}\Delta_1 + g_{11}\Delta_2}{\det \hat{g}} = -\nu_2 \Delta_2 \log\left(\frac{2\Lambda}{|\Delta_2|}\right). \tag{11}$$

B. Collective modes

We express both pairing fields in terms of fluctuations ϕ around their mean-field values,

$$\Delta_{q\alpha} = \Delta_{\alpha} \delta_{q,0} + \phi_{q\alpha}. \tag{12}$$

Note that from now on, we are using Δ_{α} to refer to the mean-field gap for each band: all *q*-dependence is in the fluctuations ϕ . We write \mathbb{G}^{-1} in terms of the mean-field inverse propagator \mathcal{G}^{-1} and fluctuation vertex $\mathbb{V}^{(\phi)}$ as $\mathbb{G}_{\alpha;k,k-q}^{-1} = \mathcal{G}_{k\alpha}^{-1} \delta_{q,0} - \mathbb{V}_{\alpha;k,k-q}^{(\phi)}$:

$$\mathcal{G}_{k\alpha}^{-1} = \begin{pmatrix} i\omega_n - \xi_{\mathbf{k}\alpha} & \Delta_\alpha \\ \overline{\Delta}_\alpha & i\omega_n + \xi_{\mathbf{k}\alpha} \end{pmatrix} = \begin{bmatrix} 1\\ \overline{(i\omega_n)^2 - E_{\mathbf{k}\alpha}^2} & \begin{pmatrix} i\omega_n + \xi_{\mathbf{k}\alpha} & -\Delta_\alpha \\ -\overline{\Delta}_\alpha & i\omega_n - \xi_{\mathbf{k}\alpha} \end{pmatrix} \end{bmatrix}^{-1} = \begin{pmatrix} G_{k\alpha} & F_{k\alpha} \\ \overline{F}_{k\alpha} & -G_{-k,\alpha} \end{pmatrix}^{-1}$$
(13)

and

$$\mathbb{V}_{\alpha;k,k-q}^{(\phi)} = \begin{pmatrix} 0 & -\phi_{q\alpha} \\ -\overline{\phi}_{-q,\alpha} & 0 \end{pmatrix}.$$
 (14)

The GPF action is the Gaussian approximation to the $Tr \log$ action in Eq. (7):

$$S[\overline{\phi}, \phi] = -\frac{L^2}{T} \sum_{q} \left(\overline{\Delta}_1 + \overline{\phi}_{q1} \ \overline{\Delta}_2 + \overline{\phi}_{q2}\right) \hat{g}^{-1} \left(\frac{\Delta_1 + \phi_{q1}}{\Delta_2 + \phi_{q2}}\right) - \operatorname{Tr}\log\left[1 - \mathcal{G}\mathbf{V}^{(\phi)}\right]$$
$$\approx -\frac{L^2}{T} \sum_{q} \left(\overline{\phi}_{q1} \ \overline{\phi}_{q2}\right) \hat{g}^{-1} \left(\frac{\phi_{q1}}{\phi_{q2}}\right) + \frac{1}{2} \operatorname{Tr}\left[\mathcal{G}\mathbf{V}^{(\phi)}\mathcal{G}\mathbf{V}^{(\phi)}\right]$$
$$= -\frac{L^2}{T} \sum_{q} \left(\overline{\phi}_{q1} \ \overline{\phi}_{q2}\right) \hat{g}^{-1} \left(\frac{\phi_{q1}}{\phi_{q2}}\right) + \frac{1}{2} \sum_{kq\alpha} \operatorname{tr}\left[\mathcal{G}_{k\alpha}\mathbf{V}^{(\phi)}_{\alpha;k,k-q}\mathcal{G}_{k-q,\alpha}\mathbf{V}^{(\phi)}_{\alpha;k-q,k}\right]$$
$$\equiv \frac{L^2}{T} \sum_{q} \left(\mathcal{R}_{-q,1} \ \mathcal{R}_{-q,2} \ \mathcal{I}_{-q,1} \ \mathcal{I}_{-q,2}\right) \mathcal{D}_{q}^{-1} \left(\frac{\mathcal{R}_{q1}}{\mathcal{I}_{q1}}\right). \tag{15}$$

Note that the linear terms necessarily cancel at the saddle point. Also note that we dropped a constant term $\sim \Delta^{\dagger} g^{-1} \Delta$. In the last line we expressed the complex fluctuations in terms of their real and imaginary parts (describing amplitude and phase fluctuations respectively, since $\Delta_{\alpha} \in \mathbb{R}$),

$$\phi_{q\alpha} = \mathcal{R}_{q\alpha} + i\mathcal{I}_{q\alpha},\tag{16}$$

and introduced the bosonic inverse propagator \mathcal{D}_q^{-1} for Gaussian pair fluctuations:

$$\mathcal{D}_{\mathbf{q}\to0}^{-1}(i\Omega_m) = \begin{pmatrix} \tilde{g}^{-1}\frac{\Delta_2}{\Delta_1} + \nu_1\gamma_1\cot\gamma_1 & -\tilde{g}^{-1} & 0 & 0\\ -\tilde{g}^{-1} & \tilde{g}^{-1}\frac{\Delta_1}{\Delta_2} + \nu_2\gamma_2\cot\gamma_2 & 0 & 0\\ 0 & 0 & \tilde{g}^{-1}\frac{\Delta_2}{\Delta_1} - \nu_1\gamma_1\tan\gamma_1 & -\tilde{g}^{-1}\\ 0 & 0 & -\tilde{g}^{-1} & \tilde{g}^{-1}\frac{\Delta_1}{\Delta_2} - \nu_2\gamma_2\tan\gamma_2 \end{pmatrix}.$$
 (17)

Here

$$\tilde{g}^{-1} = \frac{-g_{12}}{\det \hat{g}}$$
(18)

(recall that $g_{\alpha\alpha'} \leq 0$), and the variables γ_{α} are defined by

$$\sin \gamma_{\alpha} = \frac{i\Omega_m}{2|\Delta_{\alpha}|}.$$
(19)

Along the way we took the $L \to \infty$ and $T \to 0$ limits, and used the gap equations Eq. (11) to simplify some intermediate expressions. We also focus on the $\mathbf{q} \to 0$ case for simplicity. The necessary integrals are then

$$\Pi_{q\alpha}^{\mathcal{R}\mathcal{R}} = T \sum_{i\omega_n} \int \frac{\mathrm{d}^2 \mathbf{k}}{(2\pi)^2} \left(-\frac{1}{2} G_{k\alpha} (G_{-k+q,\alpha} + G_{-k-q,\alpha}) + F_{k\alpha} F_{k-q,\alpha} \right) = \int \frac{\mathrm{d}^2 \mathbf{k}}{(2\pi)^2} \tanh\left(\frac{E_{\mathbf{k}\alpha}}{2T}\right) \frac{1}{E_{\mathbf{k}\alpha}} \frac{2\xi_{\mathbf{k}\alpha}^2}{(i\Omega_m)^2 - 4E_{\mathbf{k}\alpha}^2} \\ \approx \int_{-\Lambda}^{\Lambda} \mathrm{d}\xi \, \nu_\alpha \frac{1}{\sqrt{\xi^2 + \Delta_\alpha^2}} \frac{2\xi^2}{(i\Omega_m)^2 - 4\xi^2 - 4\Delta_\alpha^2} \approx \nu_\alpha \gamma_\alpha \cot\gamma_\alpha - \nu_\alpha \log\left(\frac{2\Lambda}{|\Delta_\alpha|}\right), \quad (20)$$

$$\Pi_{q\alpha}^{\mathcal{II}} = T \sum_{i\omega_n} \int \frac{\mathrm{d}^2 \mathbf{k}}{(2\pi)^2} \left(-\frac{1}{2} G_{k\alpha} (G_{-k+q,\alpha} + G_{-k-q,\alpha}) - F_{k\alpha} F_{k-q,\alpha} \right) = \int \frac{\mathrm{d}^2 \mathbf{k}}{(2\pi)^2} \tanh\left(\frac{E_{\mathbf{k}\alpha}}{2T}\right) \frac{2E_{\mathbf{k}\alpha}}{(i\Omega_m)^2 - 4E_{\mathbf{k}\alpha}^2} \\ \approx \int_{-\Lambda}^{\Lambda} \mathrm{d}\xi \, \nu_\alpha \frac{2\sqrt{\xi^2 + \Delta_\alpha^2}}{(i\Omega_m)^2 - 4\xi^2 - 4\Delta_\alpha^2} \approx -\nu_\alpha \gamma_\alpha \tan\gamma_\alpha - \nu_\alpha \log\left(\frac{2\Lambda}{|\Delta_\alpha|}\right), \quad (21)$$

and

$$\Pi_{q\alpha}^{\mathcal{RI}} = \frac{1}{2i} T \sum_{i\omega_n} \int \frac{\mathrm{d}^2 \mathbf{k}}{(2\pi)^2} \left(G_{k\alpha} G_{-k+q,\alpha} - G_{k\alpha} G_{-k-q,\alpha} \right) = \int \frac{\mathrm{d}^2 \mathbf{k}}{(2\pi)^2} \tanh\left(\frac{E_{\mathbf{k}\alpha}}{2T}\right) \frac{1}{E_{\mathbf{k}\alpha}} \frac{\xi_{\mathbf{k}\alpha} \Omega_m}{4E_{\mathbf{k}\alpha}^2 - (i\Omega_m)^2} \\ \approx \int_{-\Lambda}^{\Lambda} \mathrm{d}\xi \, \nu_\alpha \frac{1}{\sqrt{\xi^2 + \Delta_\alpha^2}} \frac{\xi\Omega_m}{4\xi^2 + 4\Delta_\alpha^2 - (i\Omega_m)^2} = 0.$$
(22)

For Eqns. (20) and (21), the final approximation uses $\Lambda \gg |\Delta_{\alpha}|$. The cutoff can be eliminated from the expressions by using the integrated gap equations (11).

Collective modes correspond to poles of the bosonic propagator \mathcal{D}_q , i.e., zeros of det \mathcal{D}_q^{-1} , after analytically continuing $i\Omega_m \to \Omega + i0^+$. Inside the gap $(|\Omega| < 2|\Delta|)$ the infinitesimal $0^+ > 0$ is immaterial, but outside the gap the branch cuts must be chosen with care to ensure analyticity in the upper half-plane. The amplitude modes solve

$$\frac{\tilde{g}^{-1}}{\Delta_1 \Delta_2} \left(\Delta_1^2 \nu_1 \gamma_1 \cot \gamma_1 + \Delta_2^2 \nu_2 \gamma_2 \cot \gamma_2 \right) + \nu_1 \gamma_1 \cot \gamma_1 \nu_2 \gamma_2 \cot \gamma_2 = 0,$$
(23a)

and the phase modes (ABG and Leggett) solve

$$\frac{\tilde{g}^{-1}}{\Delta_1 \Delta_2} \left(\Delta_1^2 \nu_1 \gamma_1 \tan \gamma_1 + \Delta_2^2 \nu_2 \gamma_2 \tan \gamma_2 \right) - \nu_1 \gamma_1 \tan \gamma_1 \nu_2 \gamma_2 \tan \gamma_2 = 0.$$
(23b)

Eqs. (23) are transcendental, so in general we cannot solve them analytically. Instead we consider some instructive limits, focusing on the phase modes. Recall that we assumed det $\hat{g} > 0$, so $|g_{12}| < \sqrt{g_{11}g_{22}}$. When $g_{12} \to 0$, $\tilde{g}^{-1} \to 0$, and the only solution of Eq. (23b) is $\gamma_{q,\alpha} = 0$, corresponding to $\Omega = 0$. This is to be expected: when the two condensates are completely uncoupled, they each have an independent ABG mode at $\Omega = 0$. Restoring finite g_{12} , now consider the case of completely symmetric bands, where $\nu_1 = \nu_2$, $g_{11} = g_{22} = g_0$, and hence $\Delta_1 = \Delta_2$ and $\gamma_1 = \gamma_2 = \gamma$. In this case, there is an in-gap solution (meaning $\gamma \in \mathbb{R}$) whenever $|g_{12}|^2 < |g_0|$ (recall that $g_0 < 0$ by assumption), i.e., when there are two negative eigenvalues of \hat{g} . This solution (the Leggett mode) gets pushed up to $2|\Delta|$ when $g_{12} \rightarrow g_0$, which is when the second eigenvalue of \hat{g} approaches zero.

The situation changes when the gaps are not equal. Here we assume $\Delta_1 < \Delta_2$ without loss of generality. Again, Eq. (23b) cannot be solved analytically. However, since the left-hand side (LHS) consists only of continuous functions for $|\Omega| < 2|\Delta_1|$, we can determine the *parity* of the number of in-gap solutions by comparing the sign of the LHS just above $\Omega = 0$ to the sign just below $2|\Delta_1|$. If these signs are opposite, then there must be an odd number of zero crossings (i.e., phase modes) between $\Omega = 0$ and $\Omega = 2|\Delta_1|$, not including the global phase mode which always lies at $\Omega = 0$. If the sign just above 0 equals the sign just below $2|\Delta_1|$, then there must be an even number of such modes. Expanding the LHS of Eq. (23b) just above $\Omega = 0$ and just below $\Omega = 2|\Delta_1|$, one finds that the signs are opposite when and only when \tilde{g}^{-1} is smaller than a critical value of $\tilde{g}_{\rm crit}^{-1} = \nu_2 \arcsin(\Delta_1/\Delta_2)/\sqrt{1-(\Delta_1/\Delta_2)^2}$. Therefore, for $\tilde{g}^{-1} > \tilde{g}_{\rm crit}^{-1}$, Eq. (23b) has either no nonzero solutions, or an even number of them. Physically, there is no clear reason for a third phase mode (besides the Leggett and ABG modes) to arise when \tilde{g}^{-1} crosses the critical value, so we expect that in this regime there are no in-gap modes other than the ABG mode. In Fig. 1 we show that there are indeed no nonzero solutions to Eq. (23b) in this regime. We show the dependence of zeros (corresponding to the Leggett mode) as a function of \tilde{g}^{-1} for different values of Δ_2/Δ_1 . We see that for unequal gap widths, there exists a critical $\tilde{g}_{\text{crit}}^{-1}$ such that for $\tilde{g}^{-1} > \tilde{g}_{\text{crit}}^{-1}$ there are no modes in between $\Omega = 0$ and the gap edge. Assuming that $|g_{12}|^2 \ll g_{11}g_{22}$, we can expand Eq. (23b) in Ω , yielding

$$\Omega^2 \frac{\tilde{g}^{-1}(\nu_1 + \nu_2)}{4} - \frac{\Omega^4}{48\Delta_1^2 \Delta_2^2} \left[3\nu_1 \Delta_1 \nu_2 \Delta_2 - 2\tilde{g}^{-1}(\nu_1 \Delta_2^2 + \nu_2 \Delta_1^2) \right] = 0.$$
⁽²⁴⁾

The nontrivial solution is

$$\Omega_{\rm L}^2 = \frac{12\tilde{g}^{-1}(\nu_1 + \nu_2)\Delta_1^2 \Delta_2^2}{3\nu_1 \Delta_1 \nu_2 \Delta_2 - 2\tilde{g}^{-1}(\nu_1 \Delta_2^2 + \nu_2 \Delta_1^2)}.$$
(25)




FIG. 1. Dependence of the Leggett mode frequency on \tilde{g}^{-1} for three values of the gap ratio Δ_2/Δ_1 given $\nu_1 = \nu_2$, obtained by numerically solving Eq. (23b). Note that in each case there is a critical value of \tilde{g}^{-1} beyond which there are no in-gap modes, other than the ABG mode at Omega = 0.

C. Driving the Leggett mode

With the collective modes identified, all that remains is to determine their response to driving fields. The basic idea is that a drive term $\sim \delta V(t)c^{\dagger}c$, upon integrating out the fermions, will yield a source term $\sim \delta V\phi$ for the Hubbard-Stratonovich fluctuations. Integrating out those fluctuations then yields a quadratic free-energy functional $\mathcal{F}[\delta V] = -T \log \mathcal{Z}[\delta V]$, in which the coefficient of δV^2 is the capacitance. \mathcal{Z} is the partition function, $\mathcal{Z}[\delta V] = \int \mathcal{D}[\Psi^{\dagger}, \Psi, \overline{\Delta}, \Delta] e^{-S}$.

As discussed in the main text, the driving Hamiltonian is

$$H_{\delta V} = \sum_{\mathbf{k}} u \,\delta V(t) \begin{pmatrix} c_{1,\mathbf{k}}^{\dagger} & c_{2,\mathbf{k}}^{\dagger} \end{pmatrix} \begin{pmatrix} 1 & 0\\ 0 & -1 \end{pmatrix} \begin{pmatrix} c_{1,\mathbf{k}}\\ c_{2,\mathbf{k}} \end{pmatrix},\tag{26}$$

with $u = V_{\rm DC}/\sqrt{V_{\rm DC}^2 + J^2}$ in the bilayer model. Wick rotating to imaginary time and moving to the Nambu representation, the corresponding action term is

$$S_{\delta V}[\Psi^{\dagger},\Psi] = u \sum_{kq} \delta V_q \left\{ \Psi_{k1}^{\dagger} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \Psi_{k-q,1} - \Psi_{k2}^{\dagger} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \Psi_{k-q,2} \right\}.$$
(27)

Here we assumed that $\delta V(t) = \delta V(-t)$ for simplicity, such that $\delta V_q = \delta V_{-q}$ (since $\delta V(t)$ is real). This contributes an additional vertex to the full fermion inverse propagator $\mathbb{G}^{-1} = \mathcal{G}^{-1} - \mathbb{V}^{(\phi)} - \mathbb{V}^{(\delta V)}$,

$$\mathbb{V}_{\alpha;k,k-q}^{(\delta V)} = u\delta V_q(-1)^{\alpha} \begin{pmatrix} 1 & 0\\ 0 & -1 \end{pmatrix}.$$
(28)

Integrating out the fermion and expanding the Tr log out to second order in all vertices, two new terms result. The δV^2 term,

$$\frac{1}{T}\mathcal{F}_{\rm QP}[\delta V] = \frac{1}{2}\operatorname{Tr}\left[\mathcal{G}\mathbb{V}^{(\delta V)}\mathcal{G}\mathbb{V}^{(\delta V)}\right],\tag{29}$$

corresponds to the contribution of quasiparticles to the free-energy functional for δV , while the $\phi \delta V$ term,

$$S_{\text{drive}}[\overline{\phi}, \phi, \delta V] = \text{Tr}\left[\mathcal{G} \mathbb{V}^{(\phi)} \mathcal{G} \mathbb{V}^{(\delta V)}\right],\tag{30}$$

corresponds to driving of the collective modes. It is useful to immediately go to amplitude and phase variables, which amounts to expanding

$$V_{\alpha;k,k-q}^{(\phi)} = \begin{pmatrix} 0 & -\phi_{q\alpha} \\ -\overline{\phi}_{-q,\alpha} & 0 \end{pmatrix} = -\begin{pmatrix} 0 & \mathcal{R}_{q\alpha} + i\mathcal{I}_{q\alpha} \\ \mathcal{R}_{q\alpha} - i\mathcal{I}_{q\alpha} & 0 \end{pmatrix} \equiv \mathbb{V}_{\alpha;k,k-q}^{(\mathcal{R})} + \mathbb{V}_{\alpha;k,k-q}^{(\mathcal{I})}.$$
 (31)

One finds

$$\operatorname{Tr}\left[\mathcal{G}\mathbb{V}^{(\mathcal{R})}\mathcal{G}\mathbb{V}^{(\delta V)}\right] = \sum_{kq\alpha} 2(-1)^{\alpha+1} u G_{k\alpha} \left\{F_{k+q,\alpha} + F_{k-q,\alpha}\right\} \delta V_{-q} \mathcal{R}_{q\alpha}$$
(32)

and

$$\operatorname{Tr}\left[\mathcal{G}\mathbb{V}^{(\mathcal{I})}\mathcal{G}\mathbb{V}^{(\delta V)}\right] = \sum_{kq\alpha} 2i(-1)^{\alpha} u G_{k\alpha} \left\{F_{k+q,\alpha} - F_{k-q,\alpha}\right\} \delta V_{-q} \mathcal{I}_{q\alpha}.$$
(33)

We again focus on $\mathbf{q} \to 0$ for simplicity (in keeping with the form chosen for $H_{\delta V}$); $q = (i\Omega_m, 0)$ is then just a notationally-convenient way of writing the Matsubara frequency. The amplitude sector decouples from the drive:

$$\frac{T}{L^2} \sum_{k} G_{k\alpha} (F_{k+q,\alpha} + F_{k-q,\alpha}) \xrightarrow{L \to \infty} \int \frac{\mathrm{d}^2 \mathbf{k}}{(2\pi)^2} \tanh\left(\frac{E_{\mathbf{k}\alpha}}{2T}\right) \frac{1}{E_{\mathbf{k}\alpha}} \frac{2\Delta_\alpha \xi_{\mathbf{k}\alpha}}{(i\Omega_m)^2 - 4E_{\mathbf{k}\alpha}^2} \\
\approx \int_{-\Lambda}^{\Lambda} \mathrm{d}\xi \,\nu_\alpha \frac{1}{\sqrt{\xi^2 + \Delta_\alpha^2}} \frac{2\Delta_\alpha \xi}{(i\Omega_m)^2 - 4\xi^2 - 4\Delta_\alpha^2} = 0. \quad (34)$$

The phase sector is affected by the drive:

$$i\frac{T}{L^{2}}\sum_{k}G_{k\alpha}(F_{k+q,\alpha}-F_{k-q,\alpha}) \xrightarrow{L\to\infty} \int \frac{\mathrm{d}^{2}\mathbf{k}}{(2\pi)^{2}} \tanh\left(\frac{E_{\mathbf{k}\alpha}}{2T}\right) \frac{1}{E_{\mathbf{k}\alpha}}\frac{\Delta_{\alpha}\Omega_{m}}{(i\Omega_{m})^{2}-4E_{\mathbf{k}\alpha}^{2}}$$
$$\approx \int_{-\Lambda}^{\Lambda}\mathrm{d}\xi\,\nu_{\alpha}\frac{1}{\sqrt{\xi^{2}+\Delta_{\alpha}^{2}}}\frac{\Delta_{\alpha}\Omega_{m}}{(i\Omega_{m})^{2}-4\xi^{2}-4\Delta_{\alpha}^{2}} = \nu_{\alpha}\frac{i\Omega_{m}}{|\Omega_{m}|}\operatorname{sgn}(\Delta_{\alpha})\gamma_{\alpha}\operatorname{sec}\gamma_{\alpha}, \quad (35)$$

where we took $\Lambda \to \infty$ in the last equality.

Neglecting the amplitude modes (since they aren't affected by δV), including driving, the action for the phase modes at $\mathbf{q} \to 0$ is

$$S_{\text{eff}}[\mathcal{I},\delta V] = \frac{L^2}{T} \sum_{q} \left\{ \left(\mathcal{I}_{-q,1} \ \mathcal{I}_{-q,2} \right) \mathcal{D}_{\mathcal{I},\mathbf{q}\to0}^{-1}(i\Omega_m) \begin{pmatrix} \mathcal{I}_{q1} \\ \mathcal{I}_{q2} \end{pmatrix} + \left(j_{-q,1} \ j_{-q,2} \right) \begin{pmatrix} \mathcal{I}_{q1} \\ \mathcal{I}_{q2} \end{pmatrix} + \left(\mathcal{I}_{-q,1} \ \mathcal{I}_{-q,2} \right) \begin{pmatrix} j_{q1} \\ j_{q2} \end{pmatrix} \right\}, \quad (36)$$

where the sources are

$$j_{-q,1} = u\delta V_{-q}\operatorname{sgn}(\Omega_m)\operatorname{sgn}(\Delta_1)\nu_1\gamma_1\operatorname{sec}\gamma_1, \quad j_{-q,2} = -u\delta V_{-q}\operatorname{sgn}(\Omega_m)\operatorname{sgn}(\Delta_2)\nu_2\gamma_2\operatorname{sec}\gamma_2,$$

$$j_{q1} = u\delta V_q\operatorname{sgn}(\Omega_m)\operatorname{sgn}(\Delta_1)\nu_1\gamma_1\operatorname{sec}\gamma_1, \quad \text{and} \quad j_{q2} = -u\delta V_q\operatorname{sgn}(\Omega_m)\operatorname{sgn}(\Delta_2)\nu_2\gamma_{q,2}\operatorname{sec}\gamma_{q,2}. \quad (37)$$

Performing the Gaussian integration over \mathcal{I}_{α} , we obtain the collective-mode contribution to the free energy functional:

$$\frac{1}{T} \mathcal{F}_{\phi}[\delta V] = -\frac{L^2}{T} \sum_{q} \left(j_{-q,1} \ j_{-q,2} \right) \mathcal{D}_{\mathcal{I},\mathbf{q}\to0}(i\Omega_m) \begin{pmatrix} j_{q1} \\ j_{q2} \end{pmatrix} \\
= -\frac{L^2}{T} u^2 \sum_{q} \delta V_{-q} \delta V_q \left(\operatorname{sgn}(\Delta_1)\nu_1 \gamma_1 \sec \gamma_1 \ -\operatorname{sgn}(\Delta_2)\nu_2 \gamma_2 \sec \gamma_2 \right) \mathcal{D}_{\mathcal{I},\mathbf{q}\to0}(i\Omega_m) \begin{pmatrix} \operatorname{sgn}(\Delta_1)\nu_1 \gamma_1 \sec \gamma_1 \\ -\operatorname{sgn}(\Delta_2)\nu_2 \gamma_2 \sec \gamma_2 \end{pmatrix}. \quad (38)$$

The phase-mode propagator is

$$\mathcal{D}_{\mathcal{I},\mathbf{q}\to0}(i\Omega_m) = \begin{pmatrix} \tilde{g}^{-1}\frac{\Delta_2}{\Delta_1} - \nu_1\gamma_1\tan\gamma_1 & -\tilde{g}^{-1} \\ -\tilde{g}^{-1} & \tilde{g}^{-1}\frac{\Delta_1}{\Delta_2} - \nu_2\gamma_2\tan\gamma_2 \end{pmatrix}^{-1} \\ = \frac{1}{\left(\tilde{g}^{-1}\frac{\Delta_2}{\Delta_1} - \nu_1\gamma_1\tan\gamma_1\right) \left(\tilde{g}^{-1}\frac{\Delta_1}{\Delta_2} - \nu_2\gamma_2\tan\gamma_2\right) - (\tilde{g}^{-1})^2} \begin{pmatrix} \tilde{g}^{-1}\frac{\Delta_1}{\Delta_2} - \nu_2\gamma_2\tan\gamma_2 & \tilde{g}^{-1} \\ \tilde{g}^{-1} & \tilde{g}^{-1}\frac{\Delta_2}{\Delta_1} - \nu_1\gamma_1\tan\gamma_1 \end{pmatrix}.$$
(39)

Analytically continuing $i\Omega_m \to \Omega + i0^+$, the full expression for the collective-mode response is

$$\mathcal{C}_{\phi}(\Omega) = \frac{2V_{\rm DC}^2}{V_{\rm DC}^2 + J^2} \frac{\tilde{g}^{-1} \left(\nu_1 \Delta_1 \gamma_1 \sec \gamma_1 - \nu_2 \Delta_2 \gamma_2 \sec \gamma_2\right)^2 - \nu_1 \Delta_1 \gamma_1 \nu_2 \Delta_2 \gamma_2 \left(\nu_1 \gamma_1 \sec^2 \gamma_1 \tan \gamma_2 + \nu_2 \gamma_2 \sec^2 \gamma_2 \tan \gamma_1\right)}{\tilde{g}^{-1} \left(\nu_1 \Delta_1^2 \gamma_1 \tan \gamma_1 + \nu_2 \Delta_2^2 \gamma_2 \tan \gamma_2\right) - \nu_1 \Delta_1 \gamma_1 \tan \gamma_1 \nu_2 \Delta_2 \gamma_2 \tan \gamma_2}$$
(40)

The result is rather compact in the case of completely-symmetric bands, where we can drop all subscripts $\alpha \in \{1, 2\}$ and find

$$\mathcal{C}_{\phi}[\delta V] = \frac{4V_{\rm DC}^2}{V_{\rm DC}^2 + J^2} \frac{(\nu\gamma\sec\gamma)^2}{\nu\gamma\tan\gamma - 2\tilde{g}^{-1}}.$$
(41)

The Leggett mode satisfies $\nu \gamma_{\rm L} \tan \gamma_{\rm L} = 2\tilde{g}^{-1}$. Around that pole, we can approximate

$$\mathcal{C}_{\phi}(\Omega) \approx \frac{4V_{\rm DC}^2}{V_{\rm DC}^2 + J^2} \frac{\nu(\gamma_{\rm L} \sec \gamma_{\rm L})^2}{\tan \gamma_{\rm L} + \gamma_{\rm L} \sec^2 \gamma_{\rm L}} \frac{1}{\gamma - \gamma_{\rm L}} \approx \frac{4V_{\rm DC}^2}{V_{\rm DC}^2 + J^2} \frac{\nu\gamma_{\rm L}^2}{\sin \gamma_{\rm L} + \gamma_{\rm L} \sec \gamma_{\rm L}} \frac{2\Delta}{\Omega - \Omega_{\rm L} + i0^+}.$$
 (42)

In the asymmetric case, assuming $|g_{12}^{(s)}|^2 \ll g_{11}^{(s)}g_{22}^{(s)}$, we can expand the numerator and denominator for small Ω :

$$\mathcal{C}(\Omega) \approx \frac{V_{\rm DC}^2}{V_{\rm DC}^2 + J^2} \frac{\frac{1}{2}\tilde{g}^{-1}(\nu_1 - \nu_2)^2 - \frac{\Omega^2}{24\Delta_1^2\Delta_2^2} \left[3\nu_1\Delta_1\nu_2\Delta_2(\nu_1 + \nu_2) - 4\tilde{g}^{-1}(\nu_1 - \nu_2)\left(\nu_1\Delta_2^2 - \Delta_1^2\nu_2\right)\right]}{\frac{1}{4}\tilde{g}^{-1}(\nu_1 + \nu_2) - \frac{(\Omega + i0^+)^2}{48\Delta_1^2\Delta_2^2} \left[3\nu_1\Delta_1\nu_2\Delta_2 - 2\tilde{g}^{-1}(\nu_1\Delta_2^2 + \Delta_1^2\nu_2)\right]}.$$
 (43)

Of course, the observable capacitance also has the quasiparticle contribution:

$$\mathcal{F}_{\rm QP}[\delta V] = T \frac{1}{2} \operatorname{Tr} \left[\mathcal{G} \mathbb{V}^{(\delta V)} \mathcal{G} \mathbb{V}^{(\delta V)} \right] = T \sum_{kq\alpha} u^2 \delta V_{-q} \delta V_q \left\{ \frac{1}{2} G_{k\alpha} (G_{k-q,\alpha} + G_{k+q,\alpha}) - F_{k\alpha} F_{k-q,\alpha} \right\}$$
$$\xrightarrow{T \to 0}_{L \to \infty} L^2 \sum_{q\alpha} u^2 \delta V_{-q} \delta V_q \int \frac{\mathrm{d}^2 \mathbf{k}}{(2\pi)^2} \frac{1}{E_{\mathbf{k}\alpha}} \frac{2\Delta_{\alpha}^2}{(i\Omega_m)^2 - 4E_{\mathbf{k}\alpha}^2}$$
$$\approx L^2 \sum_{q\alpha} u^2 \delta V_{-q} \delta V_q \int_{-\infty}^{\infty} \nu_{\alpha} \, \mathrm{d}\xi \, \frac{1}{\sqrt{\xi^2 + \Delta_{\alpha}^2}} \frac{2\Delta_{\alpha}^2}{(i\Omega_m)^2 - 4\xi^2 - 4\Delta_{\alpha}^2}$$
$$= -L^2 \sum_{q\alpha} u^2 \delta V_{-q} \delta V_q \operatorname{sgn}(\Omega_m) \nu_{\alpha} \gamma_{q,\alpha} \operatorname{csc} \gamma_{q,\alpha} \operatorname{sec} \gamma_{q,\alpha}. \quad (44)$$

Anticipating the analytic continuation to the upper half-plane, we can set $sgn(\Omega_m) = 1$. For the case of symmetric bands, we obtain for the total free-energy functional

$$\mathcal{F}[\delta V] = \mathcal{F}_{\phi}[\delta V] + \mathcal{F}_{\rm QP}[\delta V] = L^2 \sum_{q} \delta V_{-q} \delta V_q \frac{V_{\rm DC}^2}{V_{\rm DC}^2 + J^2} 2\nu\gamma \sec^2\gamma \left(\frac{\nu\gamma}{\nu\gamma \tan\gamma - 2\tilde{g}^{-1}} - \cot\gamma\right). \tag{45}$$

We finally analytically continue, and read off the result for the total capacitance per unit area:

$$\mathcal{C}(\Omega) = \frac{V_{\rm DC}^2}{V_{\rm DC}^2 + J^2} 4\nu\gamma \sec^2\gamma \left(\frac{\nu\gamma}{\nu\gamma\tan\gamma - 2\tilde{g}^{-1}} - \cot\gamma\right). \tag{46}$$

The branch cut of the arcsin must be chosen to ensure that the result is analytic on the upper half-plane. Absorption is described by $- \operatorname{Im} \mathcal{C}$, which is plotted in Fig. 1d of the main text.

II. EFFECT OF FERMI-LIQUID TERMS ON THE LEGGETT MODE

In this section we show explicitly that attractive Fermi liquid terms can stabilize the Leggett mode inside the superconducting gap even when the pairing interactions alone would push that mode to the band edge (or, by continuity, beyond). We adopt the toy model considered in Sec. I. The additional term in the action due to the Fermi-liquid (FL) interaction in question takes the simple form

$$\delta S_{\rm FL} = \frac{T}{L^2} \sum_{\substack{kk'q\\\sigma\sigma'}} g_{\rm FL} \left(\bar{c}_{k+q,1\sigma} c_{k,1\sigma} - \bar{c}_{k+q,2\sigma} c_{k,2\sigma} - \langle \delta n_{\mathbf{k}} \rangle_0 \delta_{q,0} \right) \left(\bar{c}_{k'-q,1\sigma'} c_{k',1\sigma'} - \bar{c}_{k'-q,2\sigma'} c_{k',2\sigma'} - \langle \delta n_{\mathbf{k}'} \rangle_0 \delta_{q,0} \right), \quad (47)$$

where $g_{\rm FL} < 0$, $\delta n_{\bf k} = n_{\bf k1} - n_{\bf k2}$, and $\langle \dots \rangle_0$ denotes an equilibrium average; subtracting the averages means that these interactions have no impact on the saddle point. For simplicity we will assume equal densities of states in two

layers $\nu_1 = \nu_2$, and equal intralayer pairing interaction strengths $g_{11} = g_{22}$. Also for simplicity, we focus of $\mathbf{q} = 0$, so $q = (i\Omega_m, \mathbf{q} = 0)$ is again a shorthand for the bosonic Matsubara frequency $i\Omega_m$.

We then perform the Hubbard-Stratonovich transformation simultaneously in the Cooper and Fermi-liquid channels. While the former is introduced in Sec. I, the latter is performed via the boson $\rho_{\alpha,q} \sim \sum_{k\sigma} c^{\dagger}_{k-q,\alpha,\sigma} c_{k,\alpha,\sigma}$. The resulting action is

$$S[\Psi^{\dagger}, \Psi, \overline{\Delta}, \Delta, \rho] = -\frac{L^2}{T} \sum_{q,\alpha,\alpha'} \left(\overline{\Delta}_{q,\alpha} \hat{g}_{\alpha\alpha'}^{-1} \Delta_{q\alpha'} + \rho_{-q,\alpha} \hat{g}_{\mathrm{FL};\alpha\alpha'}^{-1} \rho_{q\alpha'} \right) - \sum_{\alpha kq} \Psi^{\dagger}_{\alpha} \mathbb{G}_{\alpha}^{-1} \Psi_{\alpha}, \tag{48}$$

where

$$\hat{g}_{\rm FL} = g_{\rm FL} \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix}. \tag{49}$$

We note that since \hat{g}_{FL} is singular, the HS transformation is not well defined. However, we however can overcome this difficulty by adding a small negative diagonal part to $\hat{g}_{\text{FL}} : \hat{g}_{\text{FL}} \to \hat{g}_{\text{FL}} + \varepsilon \times \mathbb{I}_{2\times 2}$ with $\varepsilon < 0$, and then take the limit $\varepsilon \to 0$ in the final answer. The inverse fermion propagator is

$$\mathbb{G}_{\alpha;k,k-q}^{-1} = \begin{pmatrix} (i\omega_n - \xi_{\mathbf{k}\alpha})\delta_{q,0} + \rho_{q,\alpha} & \Delta_{q,\alpha} \\ \overline{\Delta}_{-q,\alpha} & (i\omega_n + \xi_{\mathbf{k}\alpha})\delta_{q,0} - \rho_{q,\alpha} \end{pmatrix}.$$
(50)

We then integrate out the fermions to get the effective action $S_{\text{eff}}[\overline{\Delta}, \Delta, \rho]$. As mentioned, the saddle point is the same as for the case without Fermi-liquid interactions. The vertex for density fluctuations is

$$\mathbb{V}_{\alpha;k,k-q}^{(\rho)} = 2 \begin{pmatrix} \rho_{q,\alpha} & 0\\ 0 & -\rho_{q,\alpha} \end{pmatrix}.$$
(51)

We then obtain the effective action in the same manner as in Sec. I, yielding

$$S[\mathcal{R}, \mathcal{I}, \rho] = -\sum_{q} \begin{pmatrix} \mathcal{R}_{-q,1} & \mathcal{R}_{-q,2} & \mathcal{I}_{-q,1} & \mathcal{I}_{-q,2} & \rho_{-q,1} & \rho_{-q,2} \end{pmatrix} \mathcal{D}_{q}^{-1} \begin{pmatrix} \mathcal{R}_{q,1} \\ \mathcal{R}_{q,2} \\ \mathcal{I}_{q,1} \\ \mathcal{I}_{q,2} \\ \rho_{q,1} \\ \rho_{q,2} \end{pmatrix},$$
(52)

where the bosonic inverse propagator \mathcal{D}_q^{-1} can be written in block form as

$$\mathcal{D}_{q}^{-1} = \begin{pmatrix} \hat{g}^{-1} + \Pi^{\mathcal{R}\mathcal{R}} \times \mathbb{I}_{2 \times 2} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \hat{g}^{-1} + \Pi^{\mathcal{I}\mathcal{I}} \times \mathbb{I}_{2 \times 2} & \Pi^{\mathcal{I}\rho} \times \mathbb{I}_{2 \times 2} \\ \mathbf{0} & -\Pi^{\mathcal{I}\rho} \times \mathbb{I}_{2 \times 2} & \hat{g}_{\mathrm{FL}}^{-1} + \Pi^{\rho\rho} \times \mathbb{I}_{2 \times 2} \end{pmatrix},$$
(53)

where $\Pi^{\mathcal{II}}$ and $\Pi^{\mathcal{RR}}$ are given in Sec. I, $\Pi^{\rho\rho} = -8\nu\gamma_q/\sin 2\gamma_q$, and $\Pi^{\mathcal{I}\rho} = -2i\nu\gamma_q/\cos\gamma_q$. The 2 × 2 blocks act in band space. We can see that the amplitude fluctuations \mathcal{R} do not mix with the density fluctuations ρ . The collective mode frequencies are again given by the zeros of the determinant of the inverse propagator, which for the coupled $\{\mathcal{I}, \rho\}$ sector correspond to

$$\gamma_q \tan \gamma_q \left(2\tilde{g}^{-1} - \nu \gamma_q \tan \gamma_q + \frac{16\gamma_q \tilde{g}^{-1} g_{\rm FL} \nu}{\sin \gamma_q \cos \gamma_q} \right) = 0, \tag{54}$$

where $\gamma_q = \arcsin(\Omega/(2\Delta))$. Recall that $\tilde{g}^{-1} = -g_{12}/\det \hat{g}$ (where we chose all $g_{\alpha\alpha'} < 0$), and that for $g_{\rm FL} = 0$, when the determinant of the pairing interaction matrix $\det \hat{g}$ approaches zero, the Leggett mode is pushed to the band edge, $\Omega_{\rm L} \to 2\Delta$. However, for attractive $g_{\rm FL} < 0$ the Leggett mode remains inside the gap even at $\det \hat{g} \to 0$. Specifically, in this limit, the Legett mode frequency solves the equation

$$g_{\rm FL}\nu = -\frac{\sin 2\gamma_q}{16\gamma_q}.$$
(55)

As can be seen, provided that $0 < -g_{FL}\nu < 1/8$, the Fermi-liquid interactions pull the Leggett mode back into the gap even though $g_{12} \rightarrow g_0$ such that the second eigenvalue of \hat{g} approaches zero.

We have thus established that Fermi-liquid interactions can keep the Leggett mode a finite distance inside the gap when the pairing interactions alone would place it ϵ -close to the gap edge. By continuity, we conclude that even when pairing interactions alone would not yield an in-gap Leggett mode at all, Fermi-liquid terms could stabilize such an in-gap mode; nothing drastic should happen when g_{12} crosses its would-be critical value if the Leggett mode retains a finite distance from the gap edge at that value of g_{12} . To establish this rigorously, one would need to modify the Cooper-channel Hubbard-Stratonovich transformation to handle a repulsive eigenvalue (as we do for a more general model in Sec. IV); since the Fermi-liquid effects discussed in this section are not the main purpose of the present paper, we instead invoke the above continuity argument for brevity.

III. TOY MODEL: CLAPPING MODES

The minimal model containing clapping modes is that of a single-band chiral $p_x + ip_y$ superconductor, corresponding to the toy model in the main text with $g_{\alpha\alpha'}^{(s)} = 0$. The interactions are

$$\mathcal{V}_{\mathbf{k}\mathbf{k}'} = 2g\cos(\varphi_{\mathbf{k}} - \varphi_{\mathbf{k}'}) = \sum_{\ell=1,2} g\chi_{\mathbf{k}}^{(\ell)}\chi_{\mathbf{k}'}^{(\ell)},\tag{56}$$

where g < 0, $\varphi_{\mathbf{k}}$ is the angle of \mathbf{k} relative to the k_x axis, i.e., $\mathbf{k} = |\mathbf{k}| (\cos \varphi_{\mathbf{k}}, \sin \varphi_{\mathbf{k}})$, and the form factors are

$$\chi_{\mathbf{k}}^{(1)} = \sqrt{2}\cos(\varphi_{\mathbf{k}}) \quad \text{and} \quad \chi_{\mathbf{k}}^{(2)} = \sqrt{2}\sin(\varphi_{\mathbf{k}}).$$
 (57)

The Cooper-channel Hubbard-Stratonovich decoupling then involves two auxiliary bosons $\Delta_q^{(\ell)}$, yielding the action

$$S = S_{0} + \frac{L^{2}}{T} \sum_{q} \sum_{\ell=1,2} \frac{|\Delta_{q}^{(\ell)}|^{2}}{|g|} - \sum_{kq\ell} \left(\Delta_{q}^{(\ell)} \chi_{\mathbf{k}}^{(\ell)} \bar{c}_{k,\uparrow} \bar{c}_{-k+q,\downarrow} + \overline{\Delta}_{q}^{(\ell)} \chi_{\mathbf{k}}^{(\ell)} c_{-k+q,\downarrow} c_{k,\uparrow} \right)$$

$$= S_{0} + \frac{L^{2}}{T} \sum_{q} \sum_{\ell=1,2} \frac{|\Delta_{q}^{(\ell)}|^{2}}{|g|} - \sum_{kq\ell} \Psi_{k}^{\dagger} \begin{pmatrix} 0 & \Delta_{q}^{(\ell)} \chi_{\mathbf{k}}^{(\ell)} \\ \overline{\Delta}_{-q}^{(\ell)} \chi_{\mathbf{k}-\mathbf{q}}^{(\ell)} & 0 \end{pmatrix} \Psi_{k-q}.$$
(58)

 S_0 is the free action for a single parabolic band.

A. Mean-field theory

It is useful to introduce the chiral combinations

$$\chi_{\mathbf{k}}^{(\pm)} = \frac{\chi_{\mathbf{k}}^{(1)} \pm i\chi_{\mathbf{k}}^{(2)}}{\sqrt{2}} = e^{\pm i\varphi_{\mathbf{k}}}.$$
(59)

The gap equation is

$$\Delta_{\mathbf{k}} = -g \int \frac{\mathrm{d}^2 \mathbf{k}}{(2\pi)^2} \cos(\varphi_{\mathbf{k}} - \varphi_{\mathbf{k}'}) \frac{\Delta_{\mathbf{k}'}}{E_{\mathbf{k}'}}.$$
(60)

With the chiral *p*-wave ansatz $\Delta_{\mathbf{k}} = \Delta_0 \chi_{\mathbf{k}}^{(+)} = \Delta_0 e^{i\varphi_{\mathbf{k}}}$ and introducing an energy cutoff Λ , the mean-field result is standard:

$$\Delta_0 = 2\Lambda e^{1/(g\nu)}.\tag{61}$$

The $\chi^{(+)}$ and $\chi^{(-)}$ channels are degenerate; we assume that the $\chi^{(+)}$ channel condenses. Fluctuations in the $\chi^{(-)}$ channel then give rise to the clapping modes.

B. Collective modes and compressibility

Schematically, the total pairing field is $\Delta = \Delta^{(1)}\chi^{(1)} + \Delta^{(2)}\chi^{(2)} = \Delta^{(+)}\chi^{(+)} + \Delta^{(-)}\chi^{(-)}$. In terms of spacetime coordinates, we parametrize the fluctuations following Ref. [3]:

$$\Delta^{(+)}(x) = e^{i\theta(x)} \left(\Delta_0 + h(x)\right) \tag{62a}$$

and

$$\Delta^{(-)}(x) = e^{i\theta(x)} \left(a(x) + ib(x) \right).$$
(62b)

 $x = (\tau, \mathbf{r})$ contains the spacetime coordinates. θ is the ABG mode, $\Delta_0 \geq 0$ is the mean-field gap, h is the Higgs mode, and a and b are the clapping modes. After minimal coupling to the electromagnetic (U(1)) gauge field $A = (A^0, \mathbf{A})$, we can gauge out slow variations of the pairing phase θ , combining it with \mathcal{A} into the gauge-invariant field $\mathcal{A} = (\mathcal{A}^0, \mathcal{A}) = (\mathcal{A}^0 + \partial_{\tau}\theta, \mathbf{A} - \nabla \theta)$. Before the gauge transformation, expressed in coordinate space, the couplings between the Hubbard-Stratonovich bosons and the fermion is

$$S_{\text{pair}} = -\sum_{\ell} \int_{xx'} \Psi^{\dagger}(x) \begin{pmatrix} 0 & \chi^{(\ell)}(x-x')\Delta^{(\ell)}(x') \\ \overline{\Delta}^{(\ell)}(x)\chi^{(\ell)}(x-x') & 0 \end{pmatrix} \Psi(x').$$
(63)

Assuming that the phase θ varies slowly compared to the scale over which χ vanishes, such that $e^{i\theta(x)}\chi(x-x') \approx e^{i\theta(x')}\chi(x-x')$ —in other words, pairing is essentially local compared to the scale of phase variations—we can remove θ using the unitary $U_{xx'} = \delta(x-x') \exp(-i\theta(x)\tau^z/2)$, which implements the needed gauge transformation as $\mathbb{G}^{-1} \to U\mathbb{G}^{-1}U^{\dagger}$.

After the gauge transformation, one has

$$\Delta^{(1)} = \frac{1}{\sqrt{2}} \left(\Delta^{(+)} + \Delta^{(-)} \right) = \frac{1}{\sqrt{2}} \left(\Delta_0 + h + a + ib \right)$$
(64a)

and

$$\Delta^{(2)} = \frac{i}{\sqrt{2}} \left(\Delta^{(+)} - \Delta^{(-)} \right) = \frac{i}{\sqrt{2}} \left(\Delta_0 + h - a - ib \right)$$
(64b)

The action is then

$$S = \frac{L^2}{T} \sum_{\ell} \frac{|\Delta_q^{(\ell)}|^2}{|g|} - \sum_k \Psi_k^{\dagger} \mathcal{G}_k^{-1} \Psi_k + \sum_{kq} \Psi_k^{\dagger} \left(\mathbb{V}_{k,k-q}^{(\mathcal{A}^0)} + \mathbb{V}_{k,k-q}^{(\mathcal{A})} + \mathbb{V}_{k,k-q}^{(h)} + \mathbb{V}_{k,k-q}^{(a)} + \mathbb{V}_{k,k-q}^{(b)} + \mathbb{V}_{k,k-q}^{(b)} \right) \Psi_{k-q}.$$
(65)

The mean-field inverse propagator is

$$\mathcal{G}_{k}^{-1} = \begin{pmatrix} i\omega_{n} - \xi_{\mathbf{k}} & \Delta_{0}e^{i\varphi_{\mathbf{k}}} \\ \Delta_{0}e^{-i\varphi_{\mathbf{k}}} & i\omega_{n} + \xi_{\mathbf{k}} \end{pmatrix}.$$
(66)

The vertices for the gauge variables are

$$\mathbb{V}_{k,k-q}^{(\mathcal{A}^{0})} = i\mathcal{A}_{q}^{0} \begin{pmatrix} 1 & 0\\ 0 & -1 \end{pmatrix} \quad \text{and} \quad \mathbb{V}_{k,k-q}^{(\mathcal{A})} = -\frac{1}{m}\mathcal{A}_{q} \cdot \left(\mathbf{k} - \frac{1}{2}\mathbf{q}\right) \begin{pmatrix} 1 & 0\\ 0 & 1 \end{pmatrix} + \frac{1}{2m}\sum_{p}\mathcal{A}_{q-p} \cdot \mathcal{A}_{p} \begin{pmatrix} 1 & 0\\ 0 & -1 \end{pmatrix}, \tag{67a}$$

and the vertices for the Higgs and clapping modes are

$$\mathbb{V}_{k,k-q}^{(h)} = h_q \begin{pmatrix} 0 & \chi_{\mathbf{k}}^{(+)} \\ \chi_{\mathbf{k}-\mathbf{q}}^{(-)} & 0 \end{pmatrix}, \quad \mathbb{V}_{k,k-q}^{(a)} = a_q \begin{pmatrix} 0 & \chi_{\mathbf{k}}^{(-)} \\ \chi_{\mathbf{k}-\mathbf{q}}^{(+)} & 0 \end{pmatrix}, \quad \text{and} \quad \mathbb{V}_{k,k-q}^{(b)} = ib_q \begin{pmatrix} 0 & \chi_{\mathbf{k}}^{(-)} \\ -\chi_{\mathbf{k}-\mathbf{q}}^{(+)} & 0 \end{pmatrix}.$$
(67b)

For brevity, we focus only on the part of the effective action which directly relates to the compressibility, and neglect the spatial part of the overall-phase mode, and the Anderson-Higgs mechanism (hence ignoring \mathcal{A} altogether), leaving

$$\frac{T}{L^2}S_{\text{eff}} = \sum_{q} \left(\Pi_q^{00} \mathcal{A}_{-q}^0 \mathcal{A}_q^0 - \mathcal{D}_{a,q}^{-1} a_{-q} a_q - \mathcal{D}_{b,q}^{-1} b_{-q} b_q + \Pi_q^{0a} \mathcal{A}_{-q}^0 a_q + \Pi_q^{0b} \mathcal{A}_{-q}^0 b_q + \Pi_q^{ab} a_{-q} b_q \right),$$
(68)

where

$$\Pi_q^{00} \mathcal{A}_{-q}^0 \mathcal{A}_q^0 = \frac{1}{2} \frac{T}{L^2} \sum_k \operatorname{tr} \Big\{ \mathcal{G}_k \mathbb{V}_{k,k-q}^{(\mathcal{A}^0)} \mathcal{G}_{k-q} \mathbb{V}_{k-q,k}^{(\mathcal{A}^0)} \Big\},$$
(69a)

$$-\mathcal{D}_{a,q}^{-1}a_{-q}a_{q} = \frac{1}{|g|}a_{-q}a_{q} + \frac{1}{2}\frac{T}{L^{2}}\sum_{k} \operatorname{tr}\Big\{\mathcal{G}_{k}\mathbb{V}_{k,k-q}^{(a)}\mathcal{G}_{k-q}\mathbb{V}_{k-q,k}^{(a)}\Big\},\tag{69b}$$

$$-\mathcal{D}_{b,q}^{-1}b_{-q}b_{q} = \frac{1}{|g|}b_{-q}b_{q} + \frac{1}{2}\frac{T}{L^{2}}\sum_{k} \operatorname{tr}\Big\{\mathcal{G}_{k}\mathbb{V}_{k,k-q}^{(b)}\mathcal{G}_{k-q}\mathbb{V}_{k-q,k}^{(b)}\Big\},\tag{69c}$$

$$\Pi_q^{0a} \mathcal{A}_{-q}^0 a_q = \sum_k \frac{T}{L^2} \operatorname{tr} \Big\{ \mathcal{G}_k \mathbb{V}_{k,k-q}^{(a)} \mathcal{G}_{k-q} \mathbb{V}_{k-q,k}^{(\mathcal{A}^0)} \Big\},$$
(69d)

$$\Pi_q^{0b} \mathcal{A}_{-q}^0 b_q = \sum_k \frac{T}{L^2} \operatorname{tr} \Big\{ \mathcal{G}_k \mathbb{V}_{k,k-q}^{(b)} \mathcal{G}_{k-q} \mathbb{V}_{k-q,k}^{(\mathcal{A}^0)} \Big\},$$
(69e)

and

$$\Pi_{q}^{ab} a_{-q} b_{q} = \sum_{k} \frac{T}{L^{2}} \operatorname{tr} \Big\{ \mathcal{G}_{k} \mathbb{V}_{k,k-q}^{(b)} \mathcal{G}_{k-q} \mathbb{V}_{k-q,k}^{(a)} \Big\}.$$
(69f)

Strictly speaking, the collective-mode propagator is a matrix in the full mode space (spanned by a, b, and h). However, since we will evaluate that propagator at $\mathbf{q} = 0$ (as explained below), we can neglect the couplings linking a and b to the Higgs mode; these couplings can easily be shown to vanish at $\mathbf{q} = 0$ in this simple model.

We are interested in the collective-mode contribution to the compressibility which comes from integrating out aand b,

$$\delta \Pi_q^{00} = \frac{1}{4} \Pi_{-q}^{0a} \mathcal{D}_{a,q} \Pi_q^{0a} + \frac{1}{4} \Pi_{-q}^{0b} \mathcal{D}_{b,q} \Pi_q^{0b}, \tag{70}$$

to lowest nonvanishing order in **q**. In the rotationally-symmetric models we consider (either continuous rotations or C_{3v}), the couplings vanish at $O(|\mathbf{q}|^0)$. This means that, neglecting dispersion of the collective modes, we can simply evaluate the propagators at $\mathbf{q} = 0$, and the couplings Π_q^{0X} at leading nonvanishing order. The couplings Π^{0X} are

$$\begin{split} \Pi_{q}^{0a} &= -i\frac{T}{L^{2}}\sum_{k} \left\{ \chi_{\mathbf{k}-\mathbf{q}/2}^{(+)} \left(G_{k-q/2}F_{k+q/2} + F_{k-q/2}G_{-k-q/2} \right) + \chi_{\mathbf{k}+\mathbf{q}/2}^{(-)} \left(\overline{F}_{k-q/2}G_{k+q/2} + G_{-k+q/2}\overline{F}_{k+q/2} \right) \right\} \\ &= i\frac{T}{L^{2}}\sum_{k} \left\{ \chi_{\mathbf{k}-\mathbf{q}/2}^{(+)} \frac{i\omega_{n} - i\Omega_{m}/2 + \xi_{\mathbf{k}-\mathbf{q}/2}}{(i\omega_{n} - i\Omega_{m}/2)^{2} - E_{\mathbf{k}-\mathbf{q}/2}^{2}} \frac{-i\omega_{n} - i\Omega_{m}/2 + \xi_{-\mathbf{k}-\mathbf{q}/2}}{(i\omega_{n} - i\Omega_{m}/2)^{2} - E_{\mathbf{k}-\mathbf{q}/2}^{2}} \\ &+ \chi_{\mathbf{k}-\mathbf{q}/2}^{(+)} \frac{\Delta_{\mathbf{k}-\mathbf{q}/2}}{(i\omega_{n} - i\Omega_{m}/2)^{2} - E_{\mathbf{k}-\mathbf{q}/2}^{2}} \frac{-i\omega_{n} - i\Omega_{m}/2 + \xi_{-\mathbf{k}-\mathbf{q}/2}}{(i\omega_{n} - i\Omega_{m}/2)^{2} - E_{-\mathbf{k}-\mathbf{q}/2}^{2}} \\ &+ \chi_{\mathbf{k}+\mathbf{q}/2}^{(-)} \frac{\overline{\Delta_{\mathbf{k}-\mathbf{q}/2}}}{(i\omega_{n} - i\Omega_{m}/2)^{2} - E_{\mathbf{k}-\mathbf{q}/2}^{2}} \frac{i\omega_{n} + i\Omega_{m}/2 + \xi_{\mathbf{k}+\mathbf{q}/2}}{(i\omega_{n} + i\Omega_{m}/2)^{2} - E_{\mathbf{k}+\mathbf{q}/2}^{2}} \\ &+ \chi_{\mathbf{k}+\mathbf{q}/2}^{(-)} \frac{-i\omega_{n} + i\Omega_{m}/2 + \xi_{-\mathbf{k}+\mathbf{q}/2}}{(-i\omega_{n} + i\Omega_{m}/2)^{2} - E_{\mathbf{k}+\mathbf{q}/2}^{2}} \frac{\overline{\Delta_{\mathbf{k}+\mathbf{q}/2}}}{(i\omega_{n} + i\Omega_{m}/2)^{2} - E_{\mathbf{k}+\mathbf{q}/2}^{2}} \\ &+ \chi_{\mathbf{k}+\mathbf{q}/2}^{(-)} \frac{-i\omega_{n} + i\Omega_{m}/2 + \xi_{-\mathbf{k}+\mathbf{q}/2}}{(-i\omega_{n} + i\Omega_{m}/2)^{2} - E_{\mathbf{k}+\mathbf{q}/2}^{2}} \frac{\overline{\Delta_{\mathbf{k}+\mathbf{q}/2}}}{(i\omega_{n} + i\Omega_{m}/2)^{2} - E_{\mathbf{k}+\mathbf{q}/2}^{2}} \\ &+ \chi_{\mathbf{k}+\mathbf{q}/2}^{(-)} \frac{-i\omega_{n} + i\Omega_{m}/2 + \xi_{-\mathbf{k}+\mathbf{q}/2}}{(i\omega_{n} - i\Omega_{m}/2)^{2} - E_{-\mathbf{k}+\mathbf{q}/2}^{2}} \frac{\overline{\Delta_{\mathbf{k}+\mathbf{q}/2}}}{(i\omega_{n} + i\Omega_{m}/2)^{2} - E_{\mathbf{k}+\mathbf{q}/2}^{2}} \\ &+ \chi_{\mathbf{k}+\mathbf{q}/2}^{(-)} \frac{-i\omega_{n} + i\Omega_{m}/2 + \xi_{-\mathbf{k}+\mathbf{q}/2}}{(i\omega_{n} - i\Omega_{m}/2)^{2} - E_{-\mathbf{k}+\mathbf{q}/2}^{2}} \frac{\overline{\Delta_{\mathbf{k}+\mathbf{q}/2}}}{(i\omega_{n} + i\Omega_{m}/2)^{2} - E_{\mathbf{k}+\mathbf{q}/2}^{2}} \\ &+ \chi_{\mathbf{k}+\mathbf{q}/2}^{(-)} \frac{-i\omega_{n} + i\Omega_{m}/2 + \xi_{-\mathbf{k}+\mathbf{q}/2}}{(i\omega_{n} - i\Omega_{m}/2)^{2} - E_{-\mathbf{k}+\mathbf{q}/2}^{2}} \frac{1}{(i\omega_{n} - i\Omega_{m}/2)^{2} - E_{\mathbf{k}+\mathbf{q}/2}^{2}}} \\ &\times \left\{ \chi_{\mathbf{k}-\mathbf{q}/2}^{(+)} \left[\chi_{\mathbf{k}+\mathbf{q}/2}^{(-)} (i\omega_{n} - i\Omega_{m}/2 + \xi_{\mathbf{k}+\mathbf{q}/2}) + \chi_{\mathbf{k}+\mathbf{q}/2}^{(-)} (-i\omega_{n} - i\Omega_{m}/2 + \xi_{\mathbf{k}+\mathbf{q}/2}) \right] \right\} \\ &\times \left\{ (E_{\mathbf{k}-\mathbf{q}/2} \left\{ \chi_{\mathbf{k}-\mathbf{q}/2}^{(-)} \left\{ \chi_{\mathbf{k}-\mathbf{q}/2}^{(-)} \left\{ \chi_{\mathbf{k}-\mathbf{q}/2}^{(-)} \left\{ \chi_{\mathbf{k}+\mathbf{q}/2}^{(-)} \left\{ \chi_{\mathbf{k}-\mathbf{q}/2}^{(-)} \left\{ \chi_{\mathbf{k}-\mathbf{q}/2}^{(-)} \left\{ \chi_{\mathbf{k}-\mathbf{q}/2}^{(-)}$$

We expand to leading nonvanishing order in ${\bf q}.$ Working at $T\to 0$ so that

$$T\sum_{\omega_n} \to \int_{-\infty}^{\infty} \frac{\mathrm{d}\omega}{2\pi}$$
 (73)

and performing the Fermi surface integrals as standard using

$$\int \frac{\mathrm{d}^2 \mathbf{k}}{(2\pi)^2} \to \nu \int_{-\Lambda}^{\Lambda} \mathrm{d}\xi \int_0^{2\pi} \frac{\mathrm{d}\varphi}{2\pi},\tag{74}$$

one finds

$$\Pi_{q}^{0a} \approx -\frac{i\nu\Delta_{0}\mathbf{q}^{2}}{16m\mu} \Biggl\{ -\frac{8\sin(2\varphi_{\mathbf{q}})(2\mu^{2}-\Omega_{m}^{2}-4\Delta_{0}^{2})}{\Omega_{m}(\Omega_{m}^{2}+4\Delta_{0}^{2})} + \frac{48\mu\cos(2\varphi_{\mathbf{q}})}{\Omega_{m}^{2}} + 2i\arcsin\left(\frac{i\Omega_{m}}{2\Delta_{0}}\right) \frac{1}{|\Omega_{m}|^{3}\left(\Omega_{m}^{2}+4\Delta_{0}^{2}\right)^{3/2}} \Biggl[\Omega_{m}\sin(2\varphi_{\mathbf{q}}) \left(64\Delta_{0}^{4}-16\mu^{2}\Omega_{m}^{2}+\Omega_{m}^{4}+4\Delta_{0}^{2}[5\Omega_{m}^{2}-8\mu^{2}]\right) + 12\mu\cos(2\varphi_{\mathbf{q}}) \left(32\Delta_{0}^{4}+12\Delta_{0}^{2}\Omega_{m}^{2}+\Omega_{m}^{4}\right) \Biggr] \Biggr\}$$
(75)

and

$$\Pi_{q}^{0b} \approx -\frac{i\nu\Delta_{0}\mathbf{q}^{2}}{16m\mu} \left\{ \frac{8\cos(2\varphi_{\mathbf{q}})(2\mu^{2}-\Omega_{m}^{2}-4\Delta_{0}^{2})}{\Omega_{m}(\Omega_{m}^{2}+4\Delta_{0}^{2})} + \frac{48\mu\sin(2\varphi_{\mathbf{q}})}{\Omega_{m}^{2}} + 2i\arcsin\left(\frac{i\Omega_{m}}{2\Delta_{0}}\right)\frac{1}{|\Omega_{m}|^{3}(\Omega_{m}^{2}+4\Delta_{0}^{2})^{3/2}} \left[-\Omega_{m}\cos(2\varphi_{\mathbf{q}})\left(64\Delta_{0}^{4}-16\mu^{2}\Omega_{m}^{2}+\Omega_{m}^{4}+4\Delta_{0}^{2}[5\Omega_{m}^{2}-8\mu^{2}]\right) + 12\mu\sin(2\varphi_{\mathbf{q}})\left(32\Delta_{0}^{4}+12\Delta_{0}^{2}\Omega_{m}^{2}+\Omega_{m}^{4}\right) \right] \right\} \quad (76)$$

where we took the large-cutoff limit $\Lambda \to \infty$. Taking only leading terms in μ , we obtain

$$\Pi_q^{0a} \approx -\sin(2\varphi_{\mathbf{q}})f_q \tag{77a}$$

and

$$\Pi_q^{0b} \approx \cos(2\varphi_{\mathbf{q}}) f_q \tag{77b}$$

with

$$f_q = \frac{\nu \Delta_0 \mathbf{q}^2 \mu}{i\Omega_m m} \frac{1}{4\Delta_0^2 - (i\Omega_m)^2} \left(1 + 2i \arcsin\left[\frac{i\Omega_m}{2\Delta_0}\right] \frac{2\Delta_0^2 - (i\Omega_m)^2}{|\Omega_m|\sqrt{4\Delta_0^2 - (i\Omega_m)^2}} \right). \tag{77c}$$

For the clapping mode propagators $(X \in \{a, b\})$, we have

$$\mathcal{D}_{Xq} = -\frac{1}{|g|} - \Pi_q^{XX} \tag{78}$$

with

$$\Pi_q^{aa} = \frac{1}{2} \frac{T}{L^2} \sum_k \operatorname{tr} \left\{ \mathcal{G}_k \begin{pmatrix} 0 & \chi_{\mathbf{k}}^{(-)} \\ \chi_{\mathbf{k}-\mathbf{q}}^{(+)} & 0 \end{pmatrix} \mathcal{G}_{k-q} \begin{pmatrix} 0 & \chi_{\mathbf{k}-\mathbf{q}}^{(-)} \\ \chi_{\mathbf{k}}^{(+)} & 0 \end{pmatrix} \right\}$$
(79)

and

$$\Pi_{q}^{bb} = -\frac{1}{2} \frac{T}{L^{2}} \sum_{k} \operatorname{tr} \left\{ \mathcal{G}_{k} \begin{pmatrix} 0 & \chi_{\mathbf{k}}^{(-)} \\ -\chi_{\mathbf{k}-\mathbf{q}}^{(+)} & 0 \end{pmatrix} \mathcal{G}_{k-q} \begin{pmatrix} 0 & \chi_{\mathbf{k}-\mathbf{q}}^{(-)} \\ -\chi_{\mathbf{k}}^{(+)} & 0 \end{pmatrix} \right\}.$$
(80)

Explicitly,

$$\begin{aligned} \Pi_{\mathbf{q}=0}^{aa} &= \frac{1}{2} \frac{T}{L^2} \sum_{\mathbf{k}} \operatorname{tr} \left\{ \mathcal{G}_k \left(\chi_{\mathbf{k}}^{(+)} \tau^- + \chi_{\mathbf{k}}^{(-)} \tau^+ \right) \mathcal{G}_{k-q} \left(\chi_{\mathbf{k}}^{(+)} \tau^- + \chi_{\mathbf{k}}^{(-)} \tau^+ \right) \right\} \\ &= \frac{1}{2} \frac{T}{L^2} \sum_{\mathbf{k}} \left\{ \left(\chi_{\mathbf{k}}^{(+)} \right)^2 F_k F_{k-q} + \left(\chi_{\mathbf{k}}^{(-)} \right)^2 \overline{F}_k \overline{F}_{k-q} - \chi_{\mathbf{k}}^{(+)} \chi_{\mathbf{k}}^{(-)} \left(G_k G_{-k+q} + G_{-k} G_{k-q} \right) \right\} \\ &= \frac{T}{L^2} \sum_{\mathbf{k}} \frac{\Delta_0^2 \cos(4\varphi_{\mathbf{k}}) - \frac{1}{2} \left(i\omega_n + \xi_{\mathbf{k}} \right) \left(-i\omega_n + i\Omega_m + \xi_{\mathbf{k}} \right) - \frac{1}{2} \left(-i\omega_n + \xi_{\mathbf{k}} \right) \left(i\omega_n - i\Omega_m + \xi_{\mathbf{k}} \right) \\ &= \left(\frac{T}{L^2} \sum_{\mathbf{k}} \frac{\Delta_0^2 \cos(4\varphi_{\mathbf{k}}) - \frac{1}{2} \left(i\omega_n + \xi_{\mathbf{k}} \right) \left(-i\omega_n + i\Omega_m + \xi_{\mathbf{k}} \right) - \frac{1}{2} \left(-i\omega_n + \xi_{\mathbf{k}} \right) \left(i\omega_n - i\Omega_m + \xi_{\mathbf{k}} \right) \\ &= \left(\frac{T}{L^2} \sum_{\mathbf{k}} \frac{\Delta_0^2 \cos(4\varphi_{\mathbf{k}}) - \frac{1}{2} \left(i\omega_n + \xi_{\mathbf{k}} \right) \left(-i\omega_n + i\Omega_m + \xi_{\mathbf{k}} \right) - \frac{1}{2} \left(-i\omega_n + \xi_{\mathbf{k}} \right) \left(i\omega_n - i\Omega_m \right)^2 - E_{\mathbf{k}}^2 \right] \\ &= \frac{T}{L^2} \sum_{\mathbf{k}} \frac{1}{\left((i\omega_n)^2 - E_{\mathbf{k}}^2 \right] \left[(i\omega_n - i\Omega_m)^2 - E_{\mathbf{k}}^2 \right]} \\ &= \frac{T}{L^2} \sum_{\mathbf{k}} \frac{1}{\left(i\omega_n \right)^2 - E_{\mathbf{k}}^2 \right] \left[(i\omega_n - i\Omega_m)^2 - E_{\mathbf{k}}^2 \right] \\ &= \frac{T}{1} \sum_{\mathbf{k}} \frac{1}{\left(\frac{1}{2} \frac{2\xi_{\mathbf{k}}^2 + \Delta_0^2}{\left(i\omega_m \right)^2 - 4E_{\mathbf{k}}^2 \right)} \\ &= \frac{T}{2} \sum_{\mathbf{k}} \frac{1}{\left(\frac{1}{2} \frac{2\xi_{\mathbf{k}}^2 + \Delta_0^2}{\left(i\Omega_m \right)^2 - 4E_{\mathbf{k}}^2 \right)} \\ &= \nu \gamma \cot \gamma - \frac{\nu}{2} \gamma \sec \gamma \frac{2\Delta_0}{\left(\frac{2\Lambda}{\Delta_0} \right)} \\ &= \nu \gamma \cot \gamma - \frac{\nu}{2} \gamma \sec \gamma \frac{2\Delta_0}{i|\Omega_m|} - \nu \log\left(\frac{2\Lambda}{\Delta_0} \right), \quad (81) \end{aligned}$$

with $\gamma = \arcsin(i\Omega_m/2|\Delta|)$ before analytic continuation. For the *b* mode,

$$\Pi_{\mathbf{q}=0}^{bb} = -\frac{1}{2} \frac{T}{L^2} \sum_{k} \operatorname{tr} \left\{ \mathcal{G}_k \left(\chi_{\mathbf{k}}^{(+)} \tau^- - \chi_{\mathbf{k}}^{(-)} \tau^+ \right) \mathcal{G}_{k-q} \left(\chi_{\mathbf{k}}^{(+)} \tau^- - \chi_{\mathbf{k}}^{(-)} \tau^+ \right) \right\} \\
= -\frac{1}{2} \frac{T}{L^2} \sum_{k} \left\{ \left(\chi_{\mathbf{k}}^{(+)} \right)^2 F_k F_{k-q} + \left(\chi_{\mathbf{k}}^{(-)} \right)^2 \overline{F}_k \overline{F}_{k-q} + \chi_{\mathbf{k}}^{(+)} \chi_{\mathbf{k}}^{(-)} \left(G_k G_{-k+q} + G_{-k} G_{k-q} \right) \right\} \\
= -\frac{T}{L^2} \sum_{k} \frac{\Delta_0^2 \cos(4\varphi_{\mathbf{k}}) + \frac{1}{2} \left(i\omega_n + \xi_{\mathbf{k}} \right) \left(-i\omega_n + i\Omega_m + \xi_{\mathbf{k}} \right) + \frac{1}{2} \left(-i\omega_n + \xi_{\mathbf{k}} \right) \left(i\omega_n - i\Omega_m + \xi_{\mathbf{k}} \right)}{\left[(i\omega_n)^2 - E_{\mathbf{k}}^2 \right]} \\
= \frac{T}{L^2} \sum_{k} \frac{i\omega_n (i\omega_n - i\Omega_m) - \xi_{\mathbf{k}}^2}{\left[(i\omega_n - i\Omega_m)^2 - E_{\mathbf{k}}^2 \right]} = \Pi_{\mathbf{q}=0}^{aa}. \quad (82)$$

Using the mean-field equation, the log terms cancel against the 1/|g| terms from the HS transformation, yielding the clapping mode propagator:

$$\mathcal{D}_{a,\mathbf{q}=0}^{-1}(i\Omega_m) = \mathcal{D}_{b,\mathbf{q}=0}^{-1}(i\Omega_m) = \mathcal{D}_{\mathbf{q}=0}^{-1}(i\Omega_m) = -\nu\gamma\cot\gamma + \frac{\nu}{2}\gamma\sec\gamma\frac{2\Delta_0}{i|\Omega_m|}.$$
(83)

Finally, we obtain the compressibility by integrating out a and b, and then analytically continuing $i\Omega_m \to \Omega + i0^+$ in the upper half-plane:

$$\delta \Pi_{q}^{00} = \frac{1}{4} \Pi_{-q}^{0a} \mathcal{D}_{q} \Pi_{q}^{0a} + \frac{1}{4} \Pi_{-q}^{0b} \mathcal{D}_{q} \Pi_{q}^{0b} = \frac{\nu}{2} \left(\frac{\pi}{4} \xi_{\rm BCS} \mathbf{q}\right)^{4} \csc^{2} \gamma \sec^{4} \gamma \frac{1 - \gamma^{2} (\cot \gamma - \tan \gamma)^{2}}{\gamma (\cot \gamma - \tan \gamma)},\tag{84}$$

where

$$\xi_{\rm BCS} = \frac{v_{\rm F}}{\pi \Delta_0} = \frac{k_{\rm F}}{\pi m \Delta_0} = \sqrt{\frac{2\mu}{m}} \frac{1}{\pi \Delta_0},\tag{85}$$

and after analytic continuation

$$\gamma = \arcsin\left(\frac{\Omega + i0^+}{2\Delta_0}\right). \tag{86}$$

C. Trigonal warping

To consider the effect of trigonal warping, we introduce a second valley. The valleys are labelled by $\lambda \in \{+, -\}$, and **k** labels momentum relative to each valley; we assume symmetry under mapping $\lambda, \mathbf{k} \to \overline{\lambda}, -\mathbf{k}$. Pairing occurs between valleys. The Cooper-channel interactions are

$$S_{\rm int} = \frac{T}{L^2} \sum_{kk'q\lambda} \mathcal{V}_{\mathbf{k}\mathbf{k}'} \bar{c}_{k\lambda\uparrow} \bar{c}_{-k+q,\overline{\lambda}\downarrow} c_{-k'+q,\overline{\lambda}\downarrow} c_{k'\lambda\uparrow}.$$
(87)

As discussed in the main text, the idea is that reducing rotational symmetry from complete U(1) circular symmetry down to discrete C_{3v} symmetry relaxes the conservation of angular momentum down to conservation mod 3, enabling access to the clapping modes at $O(|\mathbf{q}|^2)$. In particular, the couplings Π^{0X} between the clapping modes $X \in \{a, b\}$ and the scalar potential can now show up at $O(|\mathbf{q}|)$ instead of $O(|\mathbf{q}|^2)$.

The first thing to try is the first nontrivial harmonic of a trigonally warped dispersion relation, $\xi_{\mathbf{k}} \to \xi_{\mathbf{k}} + \eta \cos(3\varphi_{\mathbf{k}})$. Repeating the same analysis, we find that at $O(\eta|\mathbf{q}|)$, the couplings Π^{0X} vanish in the large-cutoff limit $\Lambda \to \infty$. However, trigonally-warped band dispersion will lead to a trigonally-warped mean-field gap function $\Delta_{\mathbf{k}}$. Therefore, to capture the leading dependence of the compressibility on the warping, we consider the trigonally-warped gap function $\Delta_{\mathbf{k}} = \left(\Delta_0 + \tilde{\Delta}\cos(3\varphi_{\mathbf{k}})\right) e^{i\varphi_{\mathbf{k}}}$. Of course, even considering only the first nontrivial harmonic, the phase variation would also be modified, but since our goal is simply to point out the qualitative change in the compressibility's momentum dependence $(O(|\mathbf{q}|^2) \text{ vs. } O(|\mathbf{q}|^4))$, we consider only the first nontrivial harmonic of the gap amplitude, which we will show is sufficient. The analysis is completely analogous to the circular case, only particles and holes are taken from opposite valleys (i.e. $\Psi^{\dagger}_{\mathbf{k}\lambda} = \left(c^{\dagger}_{\mathbf{k}\lambda\uparrow} \ c_{-\mathbf{k}\overline{\lambda}\downarrow}\right)$), and we Taylor expand the integrands for Π^{0X} to first order in the warping parameter $\tilde{\Delta}$. Since we only want the leading-order behaviour, we evaluate the collective-mode propagators $\mathcal{D}_a = \mathcal{D}_b$ at zero warping ($\tilde{\Delta} = 0$), since the couplings Π^{0X} vanish at $O(\tilde{\Delta}^0 \mathbf{q}^2)$. To leading order in $|\mathbf{q}|$ and $\tilde{\Delta}$, we obtain

$$\delta\Pi_q^{00} = \frac{\nu}{2} \left(\frac{1}{4} \frac{\tilde{\Delta}}{\Delta_0} \frac{\mathbf{q}}{k_{\rm F}}\right)^2 \frac{\sec^2 \gamma \tan^2 \gamma}{\gamma} \frac{1 - \gamma^2 \tan^2 \gamma}{\cot \gamma - \tan \gamma} \approx \nu \left(\frac{1}{4} \frac{\tilde{\Delta}}{\Delta_0} \frac{\mathbf{q}}{k_{\rm F}}\right)^2 \frac{1 - (\pi/4)^2}{\pi} \frac{-\sqrt{2}\Delta_0}{\Omega - \sqrt{2}\Delta_0 + i0^+} \tag{88}$$

where the approximation holds in the vicinity of the positive-frequency pole. The momentum-dependent factor can be re-expressed as

$$\left(\frac{1}{4}\frac{\tilde{\Delta}}{\Delta_0}\frac{\mathbf{q}}{k_{\rm F}}\right)^2 = \frac{1}{2}\left(\frac{\pi}{4}\frac{\tilde{\Delta}}{\mu}\xi_{\rm BCS}\mathbf{q}\right)^2.$$
(89)

Note that while trigonal warping indeed saves a factor of $(\xi_{BCS} \mathbf{q})^2$ as expected, we pay the price of the factor $(\tilde{\Delta}/\mu)^2$; even in the best-case scenario where $\tilde{\Delta} \approx \Delta_0$, this term is suppressed by a factor $1/(\xi_{BCS} k_F)^2$, which is likely to be quite small.

IV. COLLECTIVE MODES FOR A GENERAL STATIC INTERACTION

A. Setup

In preparation for a our trilayer graphene calculation, we now extend the preceding analyses to the case of arbitrary static interactions. The same procedure could be applied to retarded interactions as well, by promoting the indices of the interaction potential to include timelike components ($\mathbf{k} \rightarrow k = (i\omega_n, \mathbf{k})$). First, we perform a spectral decomposition of the interaction potential in terms of form factors (i.e., eigenfunctions),

$$\mathcal{V}_{\mathbf{k}\mathbf{k}'} = \sum_{\ell} g_{\ell} \chi_{\mathbf{k}}^{(\ell)} \chi_{\mathbf{k}'}^{(\ell)}.$$
(90)

The form factors $\chi_{\mathbf{k}}^{(\ell)}$ may always be taken to be real, and we choose them to be normalized according to

$$\int_{\mathbf{k}\in\mathrm{FS}} \chi_{\mathbf{k}}^{(\ell)} \chi_{\mathbf{k}}^{(\ell')} = \delta_{\ell\ell'}.$$
(91)

Next, we introduce one HS boson $\Delta^{(\ell)}$ for each non-null eigenfunction:

$$S = \frac{L^2}{T} \sum_{q\ell} \frac{|\Delta_q^{(\ell)}|^2}{|g_\ell|} - \sum_{kq} \Psi_k^{\dagger} \left(\frac{(i\omega_n - \xi_k)\delta_{q,0}}{\Phi_{k-q,-q}^{(-)} + i\Phi_{k,q}^{(-)} + i\Phi_{k,q}^{(+)}} \right) \Psi_{k-q}$$
(92)

where the attractive $(g_{\ell} < 0)$ and repulsive $(g_{\ell} > 0)$ channels respectively add up to

$$\Phi_{\mathbf{k},q}^{(\mp)} = \sum_{\ell:g_{\ell} \leq 0} \Delta_q^{(\ell)} \chi_{\mathbf{k}}^{(\ell)}.$$
(93)

Note the factor of *i* in front of the repulsive $(\Phi^{(+)})$ terms. As usual, mean-field theory is the saddle point of *S*. Note that at the saddle point, $\overline{\Phi^{(+)}}$ is *not* the complex conjugate of $\Phi^{(+)}$, but is rather -1 times the conjugate; this may be understood in terms of a deformation of the functional integration contour for the repulsive channels [4]. The saddle-point equation is standard:

$$\Delta_{\mathbf{k}} = \frac{T}{L^2} \sum_{\mathbf{k}'} \mathcal{V}_{\mathbf{k}\mathbf{k}'} \frac{\Delta_{\mathbf{k}'}}{(i\omega_{n'})^2 - E_{\mathbf{k}'}^2} = -\int \frac{\mathrm{d}^2 \mathbf{k}}{(2\pi)^2} \mathcal{V}_{\mathbf{k}\mathbf{k}'} \tanh\left(\frac{E_{\mathbf{k}'}}{2T}\right) \frac{\Delta_{\mathbf{k}'}}{2E_{\mathbf{k}'}},\tag{94}$$

where $\Delta_{\mathbf{k}} = \Phi_{\mathbf{k},q=0}^{(-)} + i\Phi_{\mathbf{k},q=0}^{(+)}$ at the saddle point.

We then expand each HS boson in terms of fluctuations $\mathcal{R}_q^{(\ell)} + i\mathcal{I}_q^{(\ell)}$ around its mean-field value. We have one vertex for each fluctuation channel,

$$\mathbb{V}_{k,k-q}^{(\mathcal{R},\ell)} = \begin{cases}
-\mathcal{R}_{q}^{(\ell)} \begin{pmatrix} 0 & \chi_{\mathbf{k}}^{(\ell)} \\ \chi_{\mathbf{k}-\mathbf{q}}^{(\ell)} & 0 \end{pmatrix} & \text{if } g_{\ell} < 0, \\
-i\mathcal{R}_{q}^{(\ell)} \begin{pmatrix} 0 & \chi_{\mathbf{k}}^{(\ell)} \\ \chi_{\mathbf{k}-\mathbf{q}}^{(\ell)} & 0 \end{pmatrix} & \text{if } g_{\ell} > 0,
\end{cases}$$
(95)

and

$$\mathbf{V}_{k,k-q}^{(\mathcal{I},\ell)} = \begin{cases}
\mathcal{I}_{q}^{(\ell)} \begin{pmatrix} 0 & -i\chi_{\mathbf{k}}^{(\ell)} \\ i\chi_{\mathbf{k}-\mathbf{q}}^{(\ell)} & 0 \\
\mathcal{I}_{q}^{(\ell)} \begin{pmatrix} 0 & \chi_{\mathbf{k}}^{(\ell)} \\ -\chi_{\mathbf{k}-\mathbf{q}}^{(\ell)} & 0 \end{pmatrix} & \text{if } g_{\ell} > 0.
\end{cases}$$
(96)

The collective-mode propagator \mathcal{D} is then a matrix in the space of pairing channels ℓ and fluctuation types \mathcal{R}/\mathcal{I} ; writing $X, Y \in \{\mathcal{R}, \mathcal{I}\}$,

$$\left(\mathcal{D}_{q}^{-1}\right)_{\ell\ell'}^{XY} = -\frac{1}{|g_{\ell}|} \delta_{XY} \delta_{\ell\ell'} - \Pi_{q;\ell\ell'}^{XY}$$

$$\tag{97}$$

where the polarization components correspond to

$$X_{-q}^{(\ell)}\Pi_{q;\ell\ell'}^{XY}Y_{q}^{(\ell')} = \frac{1}{2}\frac{T}{L^{2}}\sum_{k} \operatorname{tr}\Big\{\mathcal{G}_{k}\mathbb{V}_{k,k-q}^{(Y,\ell')}\mathcal{G}_{k-q}\mathbb{V}_{k-q,k}^{(X,\ell)}\Big\}.$$
(98)

Note that *all* terms (not just the diagonal ones) carry a factor of 1/2, since we write the effective action in the symmetric form

$$S_{\text{eff}} = -\sum_{\ell \ell' XY} \sum_{q} X_{-q}^{(\ell)} \left(\mathcal{D}_{q}^{-1} \right)_{\ell \ell'}^{XY} Y_{q}^{(\ell')}$$
(99)

where the sums on X, Y, ℓ and ℓ' are not restricted (e.g. we include both \mathcal{RI} and \mathcal{IR} terms). Introducing

$$\rho_{\ell\ell'} = \begin{cases}
0 & g_{\ell} < 0 \text{ and } g_{\ell'} < 0, \\
1 & g_{\ell} < 0 \text{ and } g_{\ell'} > 0 \text{ or vice-versa}, \\
2 & g_{\ell} > 0 \text{ and } g_{\ell'} > 0,
\end{cases}$$
(100)

one finds

$$\Pi_{q;\ell\ell'}^{\mathcal{RR}} = i^{\rho_{\ell\ell'}} \frac{1}{2} \frac{T}{L^2} \sum_{k} \left\{ -\chi_{\mathbf{k}}^{(\ell)} \chi_{\mathbf{k}}^{(\ell')} G_k (G_{-k+q} + G_{-k-q}) + \chi_{\mathbf{k}}^{(\ell)} \chi_{\mathbf{k}-\mathbf{q}}^{(\ell')} F_k F_{k-q} + \chi_{\mathbf{k}-\mathbf{q}}^{(\ell)} \chi_{\mathbf{k}}^{(\ell')} \overline{F}_k \overline{F}_{k-q} \right\},$$
(101a)

$$\Pi_{q;\ell\ell'}^{\mathcal{II}} = i^{\rho_{\ell\ell'}} \frac{1}{2} \frac{T}{L^2} \sum_{k} \left\{ -\chi_{\mathbf{k}}^{(\ell_{-})} \chi_{\mathbf{k}}^{(\ell_{-})} G_k(G_{-k+q} + G_{-k-q}) - \chi_{\mathbf{k}}^{(\ell_{-})} \chi_{\mathbf{k}-\mathbf{q}}^{(\ell_{-})} F_k F_{k-q} - \chi_{\mathbf{k}-\mathbf{q}}^{(\ell_{-})} \overline{F}_k \overline{F}_{k-q} \right\},$$
(101b)

$$\Pi_{q;\ell\ell'}^{\mathcal{RI}} = i^{\rho_{\ell\ell'}-1} \frac{1}{2} \frac{T}{L^2} \sum_{k} \left\{ \chi_{\mathbf{k}}^{(\ell)} \chi_{\mathbf{k}}^{(\ell')} G_k (G_{-k+q} - G_{-k-q}) + \chi_{\mathbf{k}}^{(\ell)} \chi_{\mathbf{k}-\mathbf{q}}^{(\ell')} F_k F_{k-q} - \chi_{\mathbf{k}-\mathbf{q}}^{(\ell)} \chi_{\mathbf{k}}^{(\ell')} \overline{F}_k \overline{F}_{k-q} \right\},$$
(101c)

and

$$\Pi_{q;\ell\ell'}^{\mathcal{IR}} = i^{\rho_{\ell\ell'}-1} \frac{1}{2} \frac{T}{L^2} \sum_{k} \left\{ -\chi_{\mathbf{k}}^{(\ell)} \chi_{\mathbf{k}}^{(\ell')} G_k (G_{-k+q} - G_{-k-q}) + \chi_{\mathbf{k}}^{(\ell)} \chi_{\mathbf{k}-\mathbf{q}}^{(\ell')} F_k F_{k-q} - \chi_{\mathbf{k}-\mathbf{q}}^{(\ell)} \chi_{\mathbf{k}}^{(\ell')} \overline{F}_k \overline{F}_{k-q} \right\}.$$
(101d)

The Matsubara sums are easy to evaluate. At $T \to 0$, they yield

$$\lim_{T \to 0} T \sum_{i\omega_n} G_k G_{-k+q} = \frac{(E_{\mathbf{k}} - \xi_{\mathbf{k}})(E_{\mathbf{k}} + \xi_{-\mathbf{k}+\mathbf{q}} + i\Omega_m)}{2E_{\mathbf{k}} \left[(i\Omega_m + E_{\mathbf{k}})^2 - E_{-\mathbf{k}+\mathbf{q}}^2 \right]} + \frac{(E_{-\mathbf{k}+\mathbf{q}} + \xi_{-\mathbf{k}+\mathbf{q}})(E_{-\mathbf{k}+\mathbf{q}} - \xi_{\mathbf{k}} - i\Omega_m)}{2E_{-\mathbf{k}+\mathbf{q}} \left[(i\Omega_m - E_{-\mathbf{k}+\mathbf{q}})^2 - E_{\mathbf{k}}^2 \right]} \\
= \frac{(E_{\mathbf{k}} + E_{-\mathbf{k}+\mathbf{q}})(E_{\mathbf{k}}E_{-\mathbf{k}+\mathbf{q}} + \xi_{\mathbf{k}}\xi_{-\mathbf{k}+\mathbf{q}}) + i\Omega_m(\xi_{\mathbf{k}}E_{-\mathbf{k}+\mathbf{q}} + E_{\mathbf{k}}\xi_{-\mathbf{k}+\mathbf{q}})}{2E_{\mathbf{k}}E_{-\mathbf{k}+\mathbf{q}} \left[(E_{\mathbf{k}} + E_{-\mathbf{k}+\mathbf{q}})^2 - (i\Omega_m)^2 \right]} \\
= \frac{(E_{\mathbf{k}} + E_{\mathbf{k}-\mathbf{q}})(E_{\mathbf{k}}E_{\mathbf{k}-\mathbf{q}} + \xi_{\mathbf{k}}\xi_{\mathbf{k}-\mathbf{q}}) + i\Omega_m(\xi_{\mathbf{k}}E_{\mathbf{k}-\mathbf{q}} + E_{\mathbf{k}}\xi_{\mathbf{k}-\mathbf{q}})}{2E_{\mathbf{k}}E_{-\mathbf{k}+\mathbf{q}} \left[(E_{\mathbf{k}} + E_{-\mathbf{k}+\mathbf{q}})^2 - (i\Omega_m)^2 \right]} \quad (102a)$$

and

$$\lim_{T \to 0} T \sum_{i\omega_n} F_k F_{k-q} = -\frac{\Delta_{\mathbf{k}} \Delta_{\mathbf{k}-\mathbf{q}}}{2E_{\mathbf{k}} \left[(i\Omega_m + E_{\mathbf{k}})^2 - E_{\mathbf{k}-\mathbf{q}}^2 \right]} - \frac{\Delta_{\mathbf{k}} \Delta_{\mathbf{k}-\mathbf{q}}}{2E_{\mathbf{k}-\mathbf{q}} \left[(i\Omega_m - E_{\mathbf{k}-\mathbf{q}})^2 - E_{\mathbf{k}}^2 \right]} = \frac{\Delta_{\mathbf{k}} \Delta_{\mathbf{k}-\mathbf{q}} (E_{\mathbf{k}} + E_{\mathbf{k}-\mathbf{q}})}{2E_{\mathbf{k}} E_{\mathbf{k}-\mathbf{q}} \left[(E_{\mathbf{k}} + E_{\mathbf{k}-\mathbf{q}})^2 - (i\Omega_m)^2 \right]}.$$
 (102b)

To compute a response function, we also need the coupling between the pair fluctuations and the perturbing field. Assuming the perturbing drive field d couples to the fermion via a vertex $\mathbb{V}^{(d)}$, integrating out the fermion yields source terms

$$\Pi_{q;\ell}^{dX} d_{-q} X_q^{(\ell)} = \frac{T}{L^2} \sum_k \operatorname{tr} \Big\{ \mathcal{G}_k \mathbb{V}_{k,k-q}^{(X,\ell)} \mathcal{G}_{k-q} \mathbb{V}_{k-q,k}^{(d)} \Big\}.$$
(103)

The desired response function χ_q is then obtained via integration of the pair fluctuations,

$$\chi_q = \Pi_q^{dd} + \delta \Pi_q^{dd} \tag{104}$$

where

$$\delta \Pi_q^{dd} = \frac{1}{4} \sum_{XY\ell\ell'} \Pi_{-q;\ell}^{dX} (\mathcal{D}_q)_{\ell\ell'}^{XY} \Pi_{q;\ell'}^{dY}$$
(105)

and

$$\Pi_{q}^{dd}d_{-q}d_{q} = \frac{1}{2}\frac{T}{L^{2}}\sum_{k} \operatorname{tr}\left\{\mathcal{G}_{k}\mathbb{V}_{k,k-q}^{(d)}\mathcal{G}_{k-q}\mathbb{V}_{k-q,k}^{(d)}\right\}$$
(106)

corresponds to the quasiparticle contribution.

B. Projection to the Fermi surface

We now reduce the problem to a set of Fermi surface integrals. Since the Fermi velocity may vary over the Fermi surface, denoting the location on the Fermi surface as k_{\parallel} , we replace

$$\int \frac{\mathrm{d}^2 \mathbf{k}}{(2\pi)^2} \to \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{(2\pi)^2 v_{k_{\parallel}}} \int_{-\Lambda}^{\Lambda} \mathrm{d}\xi \tag{107}$$

where

$$v_{k_{\parallel}} = \left| \frac{\partial \xi_{\mathbf{k}}}{\partial k_{\perp}} \right| \tag{108}$$

with k_{\perp} the momentum component normal to the Fermi surface. Throughout, we assume that $\mathcal{V}_{\mathbf{k}\mathbf{k}'}$, $\chi_{\mathbf{k}}^{(\ell)}$, and $\Delta_{\mathbf{k}}$ vary at most weakly with k_{\perp} , allowing us to write them as functions of k_{\parallel} only.

The gap equation becomes (at $T \to 0$, so that $tanh(E/2T) \to 1$)

$$\begin{aligned} \Delta_{k_{\parallel}} &= -\int_{\mathrm{FS}} \frac{\mathrm{d}k'_{\parallel}}{(2\pi)^{2} v_{k'_{\parallel}}} \mathcal{V}_{k_{\parallel}k'_{\parallel}} \Delta_{k'_{\parallel}} \int_{-\Lambda}^{\Lambda} \mathrm{d}\xi \, \frac{1}{2\sqrt{\xi^{2} + |\Delta_{k'_{\parallel}}|^{2}}} \\ &= -\int_{\mathrm{FS}} \frac{\mathrm{d}k'_{\parallel}}{(2\pi)^{2} v_{k'_{\parallel}}} \mathcal{V}_{k_{\parallel}k'_{\parallel}} \Delta_{k'_{\parallel}} \frac{1}{2} \log \left(1 + \frac{2\Lambda \left[\Lambda + \sqrt{\Lambda^{2} + |\Delta_{k_{\parallel}}|^{2}} \right]}{|\Delta_{k_{\parallel}}|^{2}} \right) \\ &\approx -\int_{\mathrm{FS}} \frac{\mathrm{d}k'_{\parallel}}{(2\pi)^{2} v_{k'_{\parallel}}} \mathcal{V}_{k_{\parallel}k'_{\parallel}} \Delta_{k'_{\parallel}} \log \left(\frac{2\Lambda}{|\Delta_{k'_{\parallel}}|} \right), \quad (109) \end{aligned}$$

where the approximation assumes the cutoff is much larger than the gap, $\Lambda \gg |\Delta_{k_{\parallel}}|$.

Next we tackle the polarization components. In order to express these as Fermi surface quantities, we must expand in \mathbf{q} . To that end, we express

$$\xi_{\mathbf{k}-\mathbf{q}} \approx \xi_{\mathbf{k}} - \mathbf{v}_{\mathbf{k}} \cdot \mathbf{q}, \qquad \chi_{\mathbf{k}-\mathbf{q}}^{(\ell)} \approx \chi_{\mathbf{k}}^{(\ell)} - \mathbf{u}_{\mathbf{k}}^{(\ell)} \cdot \mathbf{q}, \quad \text{and} \quad \Delta_{\mathbf{k}-\mathbf{q}} \approx \Delta_{\mathbf{k}} - \mathbf{u}_{\mathbf{k}}^{(\Delta)} \cdot \mathbf{q}, \tag{110}$$

where $\mathbf{v} = \nabla_{\mathbf{k}} \xi$, $\mathbf{u}^{(\ell)} = \nabla_{\mathbf{k}} \chi^{(\ell)}$, and $\mathbf{u}^{(\Delta)} = \nabla_{\mathbf{k}} \Delta$. In principle, all three terms contribute to the **q** dependence. However, in practice, the Fermi velocity **v** is likely to be large compared to the "gap velocity" $\mathbf{u}^{(\Delta)}$. If we assume that the gap varies at most slowly in the k_{\perp} direction (normal to the Fermi surface), and varies considerably only parallel to the Fermi surface, then $|\nabla_{\mathbf{k}}\Delta| \approx |\partial_{k_{\parallel}}\Delta| \sim (1/k_{\rm F})|\partial_{\varphi}\Delta|$ assuming that $k_{\rm F}$ does not vary too drastically as a function of the angle φ . Therefore, assuming $|\partial_{\varphi}\Delta| \lesssim |\Delta|$ and denoting $v = |\mathbf{v}|$, we compare

$$|\mathbf{u}^{(\Delta)}| \lesssim \frac{|\Delta|}{k_{\rm F}} \sim \frac{|\Delta|^2 \xi_{\rm BCS}}{\mu} \quad \text{vs.} \quad v \sim |\Delta| \xi_{\rm BCS}, \tag{111}$$

so we expect \mathbf{q} dependence arising from the gap velocity to be smaller by a factor of $|\Delta|/\mu$ relative to the dependence arising from the Fermi velocity. Therefore, since this factor is likely to be very small, in practice we neglect the gap velocity entirely. We make a similar approximation for $\chi^{(\ell)}$. We assumed that $\chi_{\mathbf{k}}^{(\ell)}$ is essentially independent of the normal direction to the Fermi surface. Again using the assumption that $k_{\rm F}$ does not vary too much with φ , we then have $\mathbf{q} \cdot \nabla_{\mathbf{k}} \chi^{(\ell)} \sim (1/k_{\rm F}) \mathbf{q} \cdot \hat{\varphi} \partial_{\varphi} \chi^{(\ell)}$ at the Fermi surface. This contains the small parameter $|\mathbf{q}|/k_{\rm F} \sim |\mathbf{q}|v/\mu \sim$ $\xi_{\rm BCS} |\mathbf{q}| |\Delta|/\mu$. We capture \mathbf{q} dependence to zeroth order in $|\Delta|/\mu$, so we then approximate $\chi_{\mathbf{k}-\mathbf{q}}^{(\ell)} \approx \chi_{\mathbf{k}}^{(\ell)}$, $\Delta_{\mathbf{k}-\mathbf{q}} \approx \Delta_{\mathbf{k}}$, and $\xi_{\mathbf{k}-\mathbf{q}} \approx \xi_{\mathbf{k}} - \mathbf{v}_{\mathbf{k}} \cdot \mathbf{q}$ in all integrals.

Neglecting terms of $O(\mathbf{v} \cdot \mathbf{q}/\Lambda)$, we ultimately find below that the polarization components are

$$\Pi_{q;\ell\ell'}^{\mathcal{RR}} = -i^{\rho_{\ell\ell'}} \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \chi^{(\ell)} \chi^{(\ell')} \left[\tilde{\Lambda} - \left(\cos^2 \theta - \sin^2 z \right) \frac{z}{\sin z \cos z} \right],\tag{112a}$$

$$\Pi_{q;\ell\ell'}^{\mathcal{II}} = -i^{\rho_{\ell\ell'}} \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \chi^{(\ell)} \chi^{(\ell')} \left[\tilde{\Lambda} - \left(\sin^2 \theta - \sin^2 z \right) \frac{z}{\sin z \cos z} \right],\tag{112b}$$

and

$$\Pi_{q;\ell\ell'}^{\mathcal{RI}} = \Pi_{q;\ell\ell'}^{\mathcal{IR}} = i^{\rho_{\ell\ell'}} \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \chi^{(\ell)} \chi^{(\ell')} \sin(2\theta) \frac{z}{2\sin z \cos z},\tag{112c}$$

where

$$\theta = \arg \Delta, \tag{113}$$

$$z = \arcsin\left(\frac{\sqrt{(i\Omega_m)^2 - (\mathbf{v} \cdot \mathbf{q})^2}}{2|\Delta|}\right) \xrightarrow{i\Omega_m \to \Omega + i0^+} \arcsin\left(\frac{\sqrt{(\Omega + i0^+)^2 - (\mathbf{v} \cdot \mathbf{q})^2}}{2|\Delta|}\right),\tag{114}$$

and

$$\tilde{\Lambda} = \operatorname{arcsinh}\left(\frac{\Lambda}{|\Delta|}\right). \tag{115}$$

The drive couplings are

$$\Pi_{q;\ell}^{d\mathcal{R}} = i^{\rho_{\ell}} \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^{2}v} \langle u|\,\hat{o}\,|u\rangle\,\chi^{(\ell)}\sin\theta\frac{i\Omega_{m}}{|\Delta|}\frac{iz}{\sin z\cos z} \frac{i\Omega_{m}\to\Omega+i0^{+}}{i\Omega_{m}\to\Omega+i0^{+}}\,i^{\rho_{\ell}}\int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^{2}v}\,\langle u|\,\hat{o}\,|u\rangle\,\chi^{(\ell)}\sin\theta\frac{\Omega+i0^{+}}{|\Delta|}\frac{iz}{\sin z\cos z}$$
(116a)

and

$$\Pi_{q;\ell}^{d\mathcal{I}} = -i^{\rho_{\ell}} \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^{2}v} \left\langle u \right| \hat{o} \left| u \right\rangle \chi^{(\ell)} \cos \theta \frac{i\Omega_{m}}{|\Delta|} \frac{iz}{\sin z \cos z} \frac{i\Omega_{m} \to \Omega + i0^{+}}{-i^{\rho_{\ell}}} \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^{2}v} \left\langle u \right| \hat{o} \left| u \right\rangle \chi^{(\ell)} \cos \theta \frac{\Omega + i0^{+}}{|\Delta|} \frac{iz}{\sin z \cos z}, \quad (116b)$$

with

$$\rho_{\ell} = \begin{cases} 0 & g_{\ell} < 0, \\ 1 & g_{\ell} > 0. \end{cases}$$
(117)

The quasiparticle response is

$$\Pi_q^{dd} = -\int_{\rm FS} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \left\langle u \right| \hat{o} \left| u \right\rangle^2 \frac{(i\Omega_m)^2}{4|\Delta|^2} \frac{2z}{\sin^3 z \cos z} \tag{118}$$

prior to analytic continuation $(i\Omega_m \to \Omega + i0^+)$. The remaining Fermi surface integration can be performed numerically.

1. Evaluating the polarization components

First we evaluate

$$\lim_{T \to 0} \frac{T}{L^2} \sum_{k} \chi_{\mathbf{k}}^{(\ell)} \chi_{\mathbf{k}}^{(\ell')} G_k G_{-k+q} \\ \approx \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \chi^{(\ell)} \chi^{(\ell')} \int_{-\Lambda}^{\Lambda} \mathrm{d}\xi \left\{ \frac{(E-\xi)(E+\xi-\mathbf{v}\cdot\mathbf{q}+i\Omega_m)}{2E\left[(i\Omega_m+E)^2-(E')^2\right]} + \frac{(E'+\xi-\mathbf{v}\cdot\mathbf{q})(E'-\xi-i\Omega_m)}{2E'\left[(i\Omega_m-E')^2-E^2\right]} \right\}$$
(119)

where $E = \sqrt{\xi^2 + |\Delta|^2}$, and $E' = \sqrt{(\xi - \mathbf{v} \cdot \mathbf{q})^2 + |\Delta|^2}$, and we dropped subscripts on all quantities evaluated at k_{\parallel} for brevity. We can then shift the integration variable $\xi \to \xi + \mathbf{v} \cdot \mathbf{q}$ in the second term, and after dropping terms of $O(\mathbf{v} \cdot \mathbf{q}/\Lambda)$, we obtain

$$\lim_{T \to 0} \frac{T}{L^2} \sum_{k} \chi_{\mathbf{k}}^{(\ell)} \chi_{\mathbf{k}}^{(\ell')} G_k G_{-k+q} \approx \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \chi^{(\ell)} \chi^{(\ell')} \int_{-\Lambda}^{\Lambda} \mathrm{d}\xi \left\{ \frac{(E-\xi)(E+\xi-\mathbf{v}\cdot\mathbf{q}+i\Omega_m)}{2E\left[(i\Omega_m)^2+2i\Omega_m E+2\xi(\mathbf{v}\cdot\mathbf{q})-(\mathbf{v}\cdot\mathbf{q})^2\right]} + \frac{(E+\xi)(E-\xi-\mathbf{v}\cdot\mathbf{q}-i\Omega_m)}{2E\left[(i\Omega_m)^2-2i\Omega_m E-2\xi(\mathbf{v}\cdot\mathbf{q})-(\mathbf{v}\cdot\mathbf{q})^2\right]} \right\} \\
= \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \chi^{(\ell)} \chi^{(\ell')} \int_{-\Lambda}^{\Lambda} \mathrm{d}\xi \operatorname{Re}\left\{ \frac{1}{E} \frac{|\Delta|^2+(E-\xi)(i\Omega_m-\mathbf{v}\cdot\mathbf{q})}{(i\Omega_m)^2+2i\Omega_m E+2\xi(\mathbf{v}\cdot\mathbf{q})-(\mathbf{v}\cdot\mathbf{q})^2} \right\}. \quad (120)$$

Performing the same shift,

$$\lim_{T \to 0} \frac{T}{L^2} \sum_{k} \chi_{\mathbf{k}}^{(\ell)} \chi_{\mathbf{k}-\mathbf{q}}^{(\ell')} F_k F_{k-q} \approx -\int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \chi^{(\ell)} \chi^{(\ell')} \int_{-\Lambda}^{\Lambda} \mathrm{d}\xi \left\{ \frac{\Delta^2}{2E \left[(i\Omega_m)^2 + 2i\Omega_m E + 2\xi(\mathbf{v} \cdot \mathbf{q}) - (\mathbf{v} \cdot \mathbf{q})^2 \right]} + \frac{\Delta^2}{2E \left[(i\Omega_m)^2 - 2i\Omega_m E - 2\xi(\mathbf{v} \cdot \mathbf{q}) - (\mathbf{v} \cdot \mathbf{q})^2 \right]} \right\}$$
$$= -\int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \chi^{(\ell)} \chi^{(\ell')} \int_{-\Lambda}^{\Lambda} \mathrm{d}\xi \Delta^2 \operatorname{Re} \left\{ \frac{1}{E} \frac{1}{(i\Omega_m)^2 + 2i\Omega_m E + 2\xi(\mathbf{v} \cdot \mathbf{q}) - (\mathbf{v} \cdot \mathbf{q})^2} \right\}. \quad (121)$$

We then combine these expressions to find the polarization components, writing $\Delta = |\Delta|e^{i\theta}$:

$$\Pi_{q;\ell\ell'}^{\mathcal{RR}} \approx -i^{\rho_{\ell\ell'}} \frac{1}{2} \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \chi^{(\ell)} \chi^{(\ell')} \int_{-\Lambda}^{\Lambda} \mathrm{d}\xi \, \frac{1}{E} \left[\operatorname{Re} \left\{ \frac{|\Delta|^2 + (E - \xi)(i\Omega_m - \mathbf{v} \cdot \mathbf{q})}{(i\Omega_m)^2 + 2i\Omega_m E + 2\xi(\mathbf{v} \cdot \mathbf{q}) - (\mathbf{v} \cdot \mathbf{q})^2} \right\} + \operatorname{Re} \left\{ \frac{|\Delta|^2 - (E - \xi)(i\Omega_m - \mathbf{v} \cdot \mathbf{q})}{(i\Omega_m)^2 - 2i\Omega_m E - 2\xi(\mathbf{v} \cdot \mathbf{q}) - (\mathbf{v} \cdot \mathbf{q})^2} \right\} + 2\operatorname{Re} \left\{ \Delta^2 \right\} \operatorname{Re} \left\{ \frac{1}{(i\Omega_m)^2 + 2i\Omega_m E + 2\xi(\mathbf{v} \cdot \mathbf{q}) - (\mathbf{v} \cdot \mathbf{q})^2} \right\} \right] = -i^{\rho_{\ell\ell'}} \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \chi^{(\ell)} \chi^{(\ell')} \int_{-\Lambda}^{\Lambda} \mathrm{d}\xi \operatorname{Re} \left\{ \frac{1}{E} \frac{|\Delta|^2 (1 + \cos 2\theta) + \xi(\mathbf{v} \cdot \mathbf{q}) - (\mathbf{v} \cdot \mathbf{q})^2}{(i\Omega_m)^2 + 2i\Omega_m E + 2\xi(\mathbf{v} \cdot \mathbf{q}) - (\mathbf{v} \cdot \mathbf{q})^2} \right\}, \quad (122)$$

$$\Pi_{q;\ell\ell'}^{\mathcal{II}} \approx -i^{\rho_{\ell\ell'}} \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \chi^{(\ell)} \chi^{(\ell')} \int_{-\Lambda}^{\Lambda} \mathrm{d}\xi \operatorname{Re}\left\{\frac{1}{E} \frac{|\Delta|^2 (1 - \cos 2\theta) + \xi(\mathbf{v} \cdot \mathbf{q}) + i\Omega_m E}{(i\Omega_m)^2 + 2i\Omega_m E + 2\xi(\mathbf{v} \cdot \mathbf{q}) - (\mathbf{v} \cdot \mathbf{q})^2}\right\}$$
(123)

$$\begin{aligned} \Pi_{q;\ell\ell'}^{\mathcal{RI}} &\approx i^{\rho_{\ell\ell'}-1} \frac{1}{2} \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \chi^{(\ell)} \chi^{(\ell')} \int_{-\Lambda}^{\Lambda} \mathrm{d}\xi \, \frac{1}{E} \left[\operatorname{Re} \left\{ \frac{|\Delta|^2 + (E-\xi)(i\Omega_m - \mathbf{v} \cdot \mathbf{q})}{(i\Omega_m)^2 + 2i\Omega_m E + 2\xi(\mathbf{v} \cdot \mathbf{q}) - (\mathbf{v} \cdot \mathbf{q})^2} \right\} \\ &- \operatorname{Re} \left\{ \frac{|\Delta|^2 - (E-\xi)(i\Omega_m - \mathbf{v} \cdot \mathbf{q})}{(i\Omega_m)^2 - 2i\Omega_m E - 2\xi(\mathbf{v} \cdot \mathbf{q}) - (\mathbf{v} \cdot \mathbf{q})^2} \right\} \\ &- 2i \operatorname{Im} \left\{ \Delta^2 \right\} \operatorname{Re} \left\{ \frac{1}{(i\Omega_m)^2 + 2i\Omega_m E + 2\xi(\mathbf{v} \cdot \mathbf{q}) - (\mathbf{v} \cdot \mathbf{q})^2} \right\} \right] \\ &= -i^{\rho_{\ell\ell'}-1} \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \chi^{(\ell)} \chi^{(\ell')} \int_{-\Lambda}^{\Lambda} \mathrm{d}\xi \, \frac{1}{E} \left[\operatorname{Re} \left\{ \frac{i\Omega_m \xi + E(\mathbf{v} \cdot \mathbf{q})}{(i\Omega_m)^2 + 2i\Omega_m E + 2\xi(\mathbf{v} \cdot \mathbf{q}) - (\mathbf{v} \cdot \mathbf{q})^2} \right\} \right] \\ &+ i |\Delta|^2 \sin 2\theta \operatorname{Re} \left\{ \frac{1}{(i\Omega_m)^2 + 2i\Omega_m E + 2\xi(\mathbf{v} \cdot \mathbf{q}) - (\mathbf{v} \cdot \mathbf{q})^2} \right\} \right] \end{aligned} \tag{124}$$

and

$$\Pi_{q;\ell\ell'}^{\mathcal{IR}} \approx -i^{\rho_{\ell\ell'}-1} \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \chi^{(\ell)} \chi^{(\ell')} \int_{-\Lambda}^{\Lambda} \mathrm{d}\xi \, \frac{1}{E} \left[-\operatorname{Re} \left\{ \frac{i\Omega_m \xi + E(\mathbf{v} \cdot \mathbf{q})}{(i\Omega_m)^2 + 2i\Omega_m E + 2\xi(\mathbf{v} \cdot \mathbf{q}) - (\mathbf{v} \cdot \mathbf{q})^2} \right\} + i |\Delta|^2 \sin 2\theta \operatorname{Re} \left\{ \frac{1}{(i\Omega_m)^2 + 2i\Omega_m E + 2\xi(\mathbf{v} \cdot \mathbf{q}) - (\mathbf{v} \cdot \mathbf{q})^2} \right\} \right].$$
(125)

To perform the ξ integration, we use the change of variables

$$\frac{\xi}{|\Delta_{k_{\parallel}}|} = \sinh(x_{k_{\parallel}}), \quad \frac{E}{|\Delta|} = \sqrt{\sinh(x)^2 + 1} = \cosh(x), \quad \frac{\mathbf{v} \cdot \mathbf{q}}{|\Delta|} = ia \sinh(ib), \quad \text{and} \quad \frac{i\Omega_m}{|\Delta|} = ia \cosh(ib), \quad (126)$$

where

$$a = \frac{1}{|\Delta|} \sqrt{\Omega_m^2 + (\mathbf{v} \cdot \mathbf{q})^2} \in \mathbb{R}.$$
(127)

Note that since $\sinh(ib) \in i\mathbb{R}$ and $\cosh(ib) \in \mathbb{R}$, it follows that $b \in \mathbb{R}$. (Note that in this section, we never use a and b to refer to clapping modes, and use those letters only to label these variables). Then,

$$\Pi_{q;\ell\ell'}^{\mathcal{R}\mathcal{R}} \approx -i^{\rho_{\ell\ell'}} \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \chi^{(\ell)} \chi^{(\ell')} \int_{-\Lambda}^{\Lambda} \mathrm{d}\xi \operatorname{Re} \left\{ \frac{1}{|\Delta| \cosh(x)} \frac{1 + \cos 2\theta + ia \cosh(x + ib)}{2ia \cosh(x + ib) - a^2} \right\}$$
$$= -i^{\rho_{\ell\ell'}} \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \chi^{(\ell)} \chi^{(\ell')} \int_{-\tilde{\Lambda}}^{\tilde{\Lambda}} \mathrm{d}x \operatorname{Re} \left\{ \frac{1 + \cos 2\theta + ia \cosh(x + ib)}{2ia \cosh(x + y) - a^2} \right\}$$
$$= -i^{\rho_{\ell\ell'}} \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \chi^{(\ell)} \chi^{(\ell')} \int_{-\tilde{\Lambda}}^{\tilde{\Lambda}} \mathrm{d}x \operatorname{Re} \left\{ \frac{1 + \cos 2\theta + ia \cosh(x + ib)}{2ia \cosh(x + y) - a^2} + \frac{1}{2} \right\}$$
$$\approx -i^{\rho_{\ell\ell'}} \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \chi^{(\ell)} \chi^{(\ell')} \left[\tilde{\Lambda} + \left(1 + \cos 2\theta + \frac{1}{2}a^2 \right) \int_{-\infty}^{\infty} \mathrm{d}x \operatorname{Re} \left\{ \frac{1}{2ia \cosh(x + ib) - a^2} \right\} \right], \quad (128)$$

where $\tilde{\Lambda} = \operatorname{arcsinh}(\Lambda/|\Delta|)$, and the final approximation evaluates the convergent part of the integral in the $\Lambda \to \infty$ limit. For the remaining x integral, note that even though $ib \in i\mathbb{R}$ (so a naive variable shift $x \to x - ib$ is not valid without ensuring complex analyticity), one has

$$\int_{-\infty}^{\infty} \mathrm{d}x \, \frac{1}{2ia\cosh(x+ib) - a^2} = -\frac{4\arctan\left(\frac{2i+a}{\sqrt{4+a^2}}\right)}{a\sqrt{4+a^2}} \tag{129}$$

which is independent of b, so we can set b = 0 inside the integral and take the real part explicitly to get a simpler expression:

$$\int_{-\infty}^{\infty} \mathrm{d}x \operatorname{Re}\left\{\frac{1}{2ia\cosh(x+ib)-a^2}\right\} = \frac{1}{2a} \int_{-\infty}^{\infty} \mathrm{d}x \left(\frac{1}{-a+2i\cosh(x)} - \frac{1}{a+2i\cosh(x)}\right) \\ = -\int_{-\infty}^{\infty} \mathrm{d}x \frac{1}{a^2 + 4\cosh^2(x)} = -\frac{\operatorname{arcsinh}(a/2)}{a\sqrt{1+(a/2)^2}}.$$
 (130)

Therefore,

$$\Pi_{q;\ell\ell'}^{\mathcal{RR}} = -i^{\rho_{\ell\ell'}} \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \chi^{(\ell)} \chi^{(\ell')} \left[\tilde{\Lambda} - \left(1 + \cos 2\theta + \frac{1}{2}a^2 \right) \frac{\operatorname{arcsinh}(a/2)}{a\sqrt{1 + (a/2)^2}} \right].$$
(131)

We now define

$$z = i \operatorname{arcsinh}\left(\frac{a}{2}\right) = i \operatorname{arcsinh}\left(\frac{-i\sqrt{(i\Omega_m)^2 - (\mathbf{v} \cdot \mathbf{q})^2}}{2|\Delta|}\right) = \operatorname{arcsin}\left(\frac{\sqrt{(i\Omega_m)^2 - (\mathbf{v} \cdot \mathbf{q})^2}}{2|\Delta|}\right).$$
(132)

We can then express

$$a = 2\sinh(-iz) = -2i\sin z$$
 and $\sqrt{1 + (a/2)^2} = \sqrt{1 - \sin^2 z} = \sqrt{\cos^2 z} = \cos z.$ (133)

The final equality uses

$$\cos z = \cos \left[\arcsin\left(\frac{\sqrt{(i\Omega_m)^2 - (\mathbf{v} \cdot \mathbf{q})^2}}{2|\Delta|}\right) \right] = \sqrt{1 + \frac{\Omega_m^2 + (\mathbf{v} \cdot \mathbf{q})^2}{4|\Delta|^2}} > 0.$$
(134)

Then,

$$\Pi_{q;\ell\ell'}^{\mathcal{RR}} = -i^{\rho_{\ell\ell'}} \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \chi^{(\ell)} \chi^{(\ell')} \left[\tilde{\Lambda} - \left(\cos^2 \theta - \sin^2 z \right) \frac{z}{\sin z \cos z} \right].$$
(135)

Similarly,

$$\Pi_{q;\ell\ell'}^{\mathcal{II}} \approx -i^{\rho_{\ell\ell'}} \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \chi^{(\ell)} \chi^{(\ell')} \left[\tilde{\Lambda} - \left(\sin^2 \theta - \sin^2 z \right) \frac{z}{\sin z \cos z} \right].$$
(136)

For the cross terms,

$$\Pi_{q;\ell\ell'}^{\mathcal{RI}} \approx -i^{\rho_{\ell\ell'}-1} \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \chi^{(\ell)} \chi^{(\ell')} \int_{-\tilde{\Lambda}}^{\tilde{\Lambda}} \mathrm{d}x \left[\operatorname{Re}\left\{ \frac{\sinh(x+ib)}{ia+2\cosh(x+ib)} \right\} + i\sin 2\theta \operatorname{Re}\left\{ \frac{1}{2ia\cosh(x+ib)-a^2} \right\} \right] \\ \approx -i^{\rho_{\ell\ell'}-1} \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \chi^{(\ell)} \chi^{(\ell')} \left[\operatorname{Re}\left\{ \int_{-\tilde{\Lambda}}^{\tilde{\Lambda}} \mathrm{d}x \, \frac{\sinh(x+ib)}{ia+2\cosh(x+ib)} \right\} + i\sin 2\theta \operatorname{Re}\left\{ \int_{-\infty}^{\infty} \mathrm{d}x \, \frac{1}{2ia\cosh(x+ib)-a^2} \right\} \right] \\ = i^{\rho_{\ell\ell'}-1} \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \chi^{(\ell)} \chi^{(\ell')} \left[\sin 2\theta \frac{iz}{2\sin z \cos z} - \operatorname{Re}\left\{ \int_{-\tilde{\Lambda}}^{\tilde{\Lambda}} \mathrm{d}x \, \frac{\sinh(x+ib)}{ia+2\cosh(x+ib)} \right\} \right]$$
(137)

$$\Pi_{q;\ell\ell'}^{\mathcal{IR}} \approx i^{\rho_{\ell\ell'}-1} \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \chi^{(\ell)} \chi^{(\ell')} \left[\sin 2\theta \frac{iz}{2\sin z \cos z} + \mathrm{Re} \left\{ \int_{-\tilde{\Lambda}}^{\tilde{\Lambda}} \mathrm{d}x \, \frac{\sinh(x+ib)}{ia+2\cosh(x+ib)} \right\} \right]. \tag{138}$$

The remaining x integral evaluates to

$$I = \int_{-\tilde{\Lambda}}^{\Lambda} \mathrm{d}x \, \frac{\sinh(x+ib)}{ia+2\cosh(x+ib)} = \frac{1}{2} \left\{ \mathrm{Log} \left[a - 2i\cos\left(b - i\tilde{\Lambda}\right) \right] - \mathrm{Log} \left[a - 2i\cos\left(b + i\tilde{\Lambda}\right) \right] \right\}$$
(139)

where Log is the principal branch of the complex log. We can then express

$$\cos\left(b\mp i\tilde{\Lambda}\right) = \cos(b)\cosh\left(\tilde{\Lambda}\right) \pm i\sin(b)\sinh\left(\tilde{\Lambda}\right).$$
(140)

In the large-cutoff limit $\Lambda \to \infty$, $\tilde{\Lambda} \to \infty$ as well, so $\cosh(\tilde{\Lambda}) \approx \sinh(\tilde{\Lambda}) \approx e^{\tilde{\Lambda}}/2 \equiv \lambda$. Then,

$$I = \frac{1}{2} \left\{ \operatorname{Log} \left[a^2 - 2ia\lambda \left(\cos(b) + i\sin(b) \right) \right] - \operatorname{Log} \left[a^2 - 2ia\lambda \left(\cos(b) - i\sin(b) \right) \right] \right\}.$$
(141)

Noting that $a^2 = (\Omega_m^2 + (\mathbf{v} \cdot \mathbf{q})^2)/|\Delta|^2$, $ia\cos(b) = i\Omega_m/|\Delta|$, and $a\sin(b) = -(\mathbf{v} \cdot \mathbf{q})/|\Delta|$, it follows that

$$I = \frac{1}{2} \left\{ \text{Log} \left[\frac{\Omega_m^2 + (\mathbf{v} \cdot \mathbf{q})^2}{|\Delta|^2} - 2\lambda \frac{i\Omega_m}{|\Delta|} - 2\lambda \frac{\mathbf{v} \cdot \mathbf{q}}{|\Delta|} \right] - \text{Log} \left[\frac{\Omega_m^2 + (\mathbf{v} \cdot \mathbf{q})^2}{|\Delta|^2} - 2\lambda \frac{i\Omega_m}{|\Delta|} + 2\lambda \frac{\mathbf{v} \cdot \mathbf{q}}{|\Delta|} \right] \right\}.$$
 (142)

Since the cutoff is large $(\lambda \gg 1)$, we can approximate

$$I \approx \frac{1}{2} \left\{ \log \left[-2\lambda \frac{i\Omega_m}{|\Delta|} - 2\lambda \frac{\mathbf{v} \cdot \mathbf{q}}{|\Delta|} \right] - \log \left[-2\lambda \frac{i\Omega_m}{|\Delta|} + 2\lambda \frac{\mathbf{v} \cdot \mathbf{q}}{|\Delta|} \right] \right\}.$$
(143)

Notice that the complex arguments of the Logs have the same magnitude, meaning the difference of Logs is purely imaginary. Therefore, taking the real part yields zero, and we conclude that

$$\Pi_{q;\ell\ell'}^{\mathcal{RI}} = \Pi_{q;\ell\ell'}^{\mathcal{IR}} = i^{\rho_{\ell\ell'}} \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \chi^{(\ell)} \chi^{(\ell')} \sin(2\theta) \frac{z}{2\sin z \cos z}.$$
(144)

2. Evaluating the driving terms

We now consider the response to driving. We assume that the drive field d couples locally in real space as

$$H_{\rm drive}(t) = \int d^2 \mathbf{r} \sum_j d(t, \mathbf{r}) o_j \psi_j^{\dagger} \psi_j = \sum_{\mathbf{k} \mathbf{q}j} d_{\mathbf{q}}(t) o_j f_{\mathbf{k}j}^{\dagger} f_{\mathbf{k}-\mathbf{q},j}.$$
 (145)

j indexes the orbitals in the unit cell, spin, and any other internal degree of freedom; o_j encodes how the field d affects the energy of local state j. For our RTG calculation, the relevant index will be the layer.

Suppose the non-interacting part of the problem is diagonalized in terms of the band operators $c_{\mathbf{k}\alpha\sigma}$, where α labels the bands. Assuming no spin-orbit coupling, the band operators correspond to wavefunctions $|u_{\mathbf{k}\alpha}\rangle$, i.e. $u_{\mathbf{k}\alpha j} = \langle j | u_{\mathbf{k}\alpha} \rangle$, with

$$f_{\mathbf{k}j\sigma} = \sum_{\alpha} u_{\mathbf{k}j\alpha} c_{\mathbf{k}\alpha\sigma}.$$
(146)

In the band basis, the driving Hamiltonian is then

$$H_{\text{drive}} = \sum_{\substack{\mathbf{kq}\\j\alpha\alpha'\sigma}} d_{\mathbf{q}}(t) u_{\mathbf{k}\alpha j}^* o_j u_{\mathbf{k}-\mathbf{q},\alpha' j} c_{\mathbf{k}\alpha\sigma}^{\dagger} c_{\mathbf{k}-\mathbf{q},\alpha'\sigma} \equiv \sum_{\mathbf{kq}\alpha\alpha'\sigma} d_{\mathbf{q}}(t) \langle u_{\mathbf{k}\alpha} | \hat{o} | u_{\mathbf{k}-\mathbf{q},\alpha'} \rangle c_{\mathbf{k}\alpha\sigma}^{\dagger} c_{\mathbf{k}-\mathbf{q},\alpha'\sigma}.$$
(147)

 $d(t, \mathbf{r})\hat{o}$ is the one-particle operator corresponding to driving; we assumed \hat{o} is diagonal in the basis of orbitals within the unit cell. The last equation is more general, and would also hold for a drive which induces transitions between orbitals.

For the RTG calculation, we will project to a single active band, in which case the driving terms have one fewer summation:

$$H_{\rm drive} = \sum_{\mathbf{kq}\sigma} d_{\mathbf{q}}(t) \langle u_{\mathbf{k}} | \, \hat{o} \, | u_{\mathbf{k}-\mathbf{q}} \rangle \, c^{\dagger}_{\mathbf{k}\sigma} c_{\mathbf{k}-\mathbf{q},\sigma} = \sum_{\mathbf{kq}} d_{\mathbf{q}}(t) \Psi^{\dagger}_{\mathbf{k}} \begin{pmatrix} \langle u_{\mathbf{k}} | \, \hat{o} \, | u_{\mathbf{k}-\mathbf{q}} \rangle & 0\\ 0 & -\langle u_{-\mathbf{k}+\mathbf{q}} | \, \hat{o} \, | u_{-\mathbf{k}} \rangle \end{pmatrix} \Psi_{\mathbf{k}-\mathbf{q}}.$$
(148)

For simplicity, we assume that the wavefunctions respect $|u_{\bf k}\rangle=|u_{-{\bf k}}\rangle$ such that

$$H_{\rm drive} = d_{\mathbf{q}}(t) \langle u_{\mathbf{k}} | \, \hat{o} \, | u_{\mathbf{k}-\mathbf{q}} \rangle \, \Psi_{\mathbf{k}}^{\dagger} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \Psi_{\mathbf{k}-\mathbf{q}}.$$
(149)

Wick rotating to imaginary time and Fourier transforming to Matsubara frequency, we obtain the vertex in the action,

$$\mathbb{V}_{k,k-q}^{(d)} = -d_q \left\langle u_{\mathbf{k}} \right| \hat{o} \left| u_{\mathbf{k}-\mathbf{q}} \right\rangle \begin{pmatrix} 1 & 0\\ 0 & -1 \end{pmatrix}.$$
(150)

Integrating out the fermion then yields two kinds of terms. The quadratic term corresponds to the quasiparticle response,

$$\frac{1}{L^2} \mathcal{F}_{\text{QP}}[d] = \frac{1}{2} \frac{T}{L^2} \operatorname{Tr} \left\{ \mathcal{G} \mathbb{V}^{(d)} \mathcal{G} \mathbb{V}^{(d)} \right\} = \sum_{q} \Pi_q^{dd} d_{-q} d_q$$

$$= \frac{1}{2} \frac{T}{L^2} \sum_{kq} d_q d_{-q} |\langle u_{\mathbf{k}} | \hat{o} | u_{\mathbf{k}-\mathbf{q}} \rangle |^2 \operatorname{tr} \left\{ \mathcal{G}_k \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \mathcal{G}_{k-q} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \right\}$$

$$= \frac{1}{2} \frac{T}{L^2} \sum_{kq} d_q d_{-q} |\langle u_{\mathbf{k}} | \hat{o} | u_{\mathbf{k}-\mathbf{q}} \rangle |^2 \left(\mathcal{G}_k \mathcal{G}_{k-q} + \mathcal{G}_{-k+q} - \mathcal{F}_k \overline{\mathcal{F}}_{k-q} - \overline{\mathcal{F}}_k \mathcal{F}_{k-q} \right), \quad (151)$$

and the linear terms correspond to driving of the collective modes:

$$\frac{1}{L^{2}}\mathcal{F}_{\phi}[d,\mathcal{R},\mathcal{I}] = \frac{T}{L^{2}}\operatorname{Tr}\left\{\mathcal{G}\left(\mathbb{V}^{(\mathcal{R})} + \mathbb{V}^{(\mathcal{I})}\right)\mathcal{G}\mathbb{V}^{(d)}\right\} = \frac{T}{L^{2}}\sum_{kq\ell}\operatorname{tr}\left\{\mathcal{G}_{k}\left(\mathbb{V}_{k,k-q}^{(\mathcal{R},\ell)} + \mathbb{V}_{k,k-q}^{(\mathcal{I},\ell)}\right)\mathcal{G}_{k}\mathbb{V}_{k-q,k}^{(d)}\right\}$$

$$= \sum_{q\ell}\left(d_{-q}\Pi_{q;\ell}^{d\mathcal{R}}\mathcal{R}_{q}^{(\ell)} + d_{-q}\Pi_{q;\ell}^{d\mathcal{I}}\mathcal{I}_{q}^{(\ell)}\right)$$

$$= \frac{T}{L^{2}}\sum_{kq\ell}i^{\rho_{\ell}}d_{-q}\langle u_{\mathbf{k}-\mathbf{q}}|\,\hat{o}\,|u_{\mathbf{k}}\rangle\left\{\mathcal{R}_{q}^{(\ell)}\left[\chi_{\mathbf{k}-\mathbf{q}}^{(\ell)}\left(F_{k}G_{k-q}+G_{-k}F_{k-q}\right)+\chi_{\mathbf{k}}^{(\ell)}\left(G_{k}\overline{F}_{k-q}+\overline{F}_{k}G_{-k+q}\right)\right]\right\}$$

$$+ i\mathcal{I}_{q}^{(\ell)}\left[-\chi_{\mathbf{k}-\mathbf{q}}^{(\ell)}\left(F_{k}G_{k-q}+G_{-k}F_{k-q}\right)+\chi_{k}^{(\ell)}\left(G_{k}\overline{F}_{k-q}+\overline{F}_{k}G_{-k+q}\right)\right]\right\}. (152)$$

Here we defined $\rho_{\ell} = 0$ if $g_{\ell} < 0$, and $\rho_{\ell} = 1$ if $g_{\ell} > 0$.

The Matsubara sums are again standard, yielding

$$\lim_{T \to 0} T \sum_{\omega_n} G_k G_{k-q} = -\frac{1}{2E_{\mathbf{k}}} \frac{(E_{\mathbf{k}} - \xi_{\mathbf{k}})(i\Omega_m + E_{\mathbf{k}} - \xi_{\mathbf{k}-\mathbf{q}})}{(E_{\mathbf{k}} + i\Omega_m)^2 - E_{\mathbf{k}-\mathbf{q}}^2} - \frac{1}{2E_{\mathbf{k}-\mathbf{q}}} \frac{(E_{\mathbf{k}-\mathbf{q}} - \xi_{\mathbf{k}-\mathbf{q}})(i\Omega_m + \xi_{\mathbf{k}} - E_{\mathbf{k}-\mathbf{q}})}{E_{\mathbf{k}}^2 - (E_{\mathbf{k}-\mathbf{q}} - i\Omega_m)^2}, \quad (153)$$

$$\lim_{T \to 0} T \sum_{\omega_n} G_{-k} G_{-k+q} = -\frac{1}{2E_{\mathbf{k}}} \frac{(E_{\mathbf{k}} + \xi_{\mathbf{k}})(i\Omega_m + E_{\mathbf{k}} + \xi_{\mathbf{k}-\mathbf{q}})}{(E_{\mathbf{k}} + i\Omega_m)^2 - E_{\mathbf{k}-\mathbf{q}}^2} - \frac{1}{2E_{\mathbf{k}-\mathbf{q}}} \frac{(E_{\mathbf{k}-\mathbf{q}} + \xi_{\mathbf{k}-\mathbf{q}})(i\Omega_m - \xi_{\mathbf{k}} - E_{\mathbf{k}-\mathbf{q}})}{E_{\mathbf{k}}^2 - (E_{\mathbf{k}-\mathbf{q}} - i\Omega_m)^2}, \quad (154)$$

$$\lim_{T \to 0} T \sum_{\omega_n} F_k \overline{F}_{k-q} = -\frac{1}{2E_k} \frac{\Delta_k \overline{\Delta}_{k-q}}{(E_k + i\Omega_m)^2 - E_{k-q}^2} + \frac{1}{2E_{k-q}} \frac{\Delta_k \overline{\Delta}_{k-q}}{E_k^2 - (E_{k-q} - i\Omega_m)^2} = \left(\lim_{T \to 0} \sum_{\omega_n} \overline{F}_k F_{k-q}\right)^*, \quad (155)$$

$$\lim_{T \to 0} T \sum_{\omega_n} F_k G_{k-q} = -\frac{\Delta_{\mathbf{k}}}{2E_{\mathbf{k}}} \frac{E_{\mathbf{k}} + i\Omega_m - \xi_{\mathbf{k}-\mathbf{q}}}{(E_{\mathbf{k}} + i\Omega_m)^2 - E_{\mathbf{k}-\mathbf{q}}^2} + \frac{\Delta_{\mathbf{k}}}{2E_{\mathbf{k}-\mathbf{q}}} \frac{E_{\mathbf{k}-\mathbf{q}} - \xi_{\mathbf{k}-\mathbf{q}}}{E_{\mathbf{k}}^2 - (E_{\mathbf{k}-\mathbf{q}} - i\Omega_m)^2},\tag{156}$$

$$\lim_{T \to 0} T \sum_{\omega_n} G_{-k} F_{k-q} = \frac{\Delta_{\mathbf{k}-\mathbf{q}}}{2E_{\mathbf{k}}} \frac{E_{\mathbf{k}} + \xi_{\mathbf{k}}}{(E_{\mathbf{k}} + i\Omega_m)^2 - E_{\mathbf{k}-\mathbf{q}}^2} - \frac{\Delta_{\mathbf{k}-\mathbf{q}}}{2E_{\mathbf{k}-\mathbf{q}}} \frac{\xi_{\mathbf{k}} - i\Omega_m + E_{\mathbf{k}-\mathbf{q}}}{E_{\mathbf{k}}^2 - (E_{\mathbf{k}-\mathbf{q}} - i\Omega_m)^2},\tag{157}$$

$$\lim_{T \to 0} T \sum_{\omega_n} G_k \overline{F}_{k-q} = -\frac{\overline{\Delta}_{\mathbf{k}-\mathbf{q}}}{2E_{\mathbf{k}}} \frac{E_{\mathbf{k}} - \xi_{\mathbf{k}}}{(E_{\mathbf{k}} + i\Omega_m)^2 - E_{\mathbf{k}-\mathbf{q}}^2} - \frac{\overline{\Delta}_{\mathbf{k}-\mathbf{q}}}{2E_{\mathbf{k}-\mathbf{q}}} \frac{\xi_{\mathbf{k}} + i\Omega_m - E_{\mathbf{k}-\mathbf{q}}}{E_{\mathbf{k}}^2 - (E_{\mathbf{k}-\mathbf{q}} - i\Omega_m)^2},\tag{158}$$

and

$$\lim_{T \to 0} T \sum_{\omega_n} \overline{F}_k G_{-k+q} = \frac{\overline{\Delta}_{\mathbf{k}}}{2E_{\mathbf{k}}} \frac{E_{\mathbf{k}} + i\Omega_m + \xi_{\mathbf{k}-\mathbf{q}}}{(E_{\mathbf{k}} + i\Omega_m)^2 - E_{\mathbf{k}-\mathbf{q}}^2} - \frac{\overline{\Delta}_{\mathbf{k}}}{2E_{\mathbf{k}-\mathbf{q}}} \frac{E_{\mathbf{k}-\mathbf{q}} + \xi_{\mathbf{k}-\mathbf{q}}}{E_{\mathbf{k}}^2 - (E_{\mathbf{k}-\mathbf{q}} - i\Omega_m)^2}.$$
(159)

We now project to the Fermi surface using the same approximations as for the polarization components $(\chi_{\mathbf{k}-\mathbf{q}}^{(\ell)} \approx \chi_{\mathbf{k}}^{(\ell)}, \Delta_{\mathbf{k}-\mathbf{q}} \approx \Delta_{\mathbf{k}}, \text{ and } \xi_{\mathbf{k}-\mathbf{q}} \approx \xi_{\mathbf{k}} - \mathbf{v}_{\mathbf{k}} \cdot \mathbf{q})$, with the additional approximation that

$$\langle u_{\mathbf{k}-\mathbf{q}} | \, \hat{o} \, | u_{\mathbf{k}} \rangle \approx \langle u_{\mathbf{k}} | \, \hat{o} \, | u_{\mathbf{k}} \rangle \,,$$

$$\tag{160}$$

which we further assume is essentially constant normal to the Fermi surface (i.e. $|u_{\mathbf{k}}\rangle \approx |u_{k_{\parallel}}\rangle$). We then need the integrals

$$\frac{1}{2} \int \frac{\mathrm{d}^2 \mathbf{k}}{4\pi^2} \left\langle u_{\mathbf{k}} \right| \hat{o} \left| u_{\mathbf{k}} \right\rangle^2 \left(G_k G_{k-q} + G_{-k} G_{-k+q} - F_k \overline{F}_{k-q} - \overline{F}_k F_{k-q} \right)$$
(161)

for the quasiparticle response, and

$$\int \frac{\mathrm{d}^2 \mathbf{k}}{4\pi^2} \left\langle u_{\mathbf{k}} \right| \hat{o} \left| u_{\mathbf{k}} \right\rangle \chi_{\mathbf{k}}^{(\ell)} \left(\pm F_k G_{k-q} \pm G_{-k} F_{k-q} + G_k \overline{F}_{k-q} + \overline{F}_k G_{-k+q} \right)$$
(162)

for the collective mode source terms.

For the quasiparticles,

$$\lim_{T \to 0} \sum_{\omega_n} \frac{1}{2} \int \frac{d^2 \mathbf{k}}{4\pi^2} \langle u_{\mathbf{k}} | \, \hat{o} \, | \, u_{\mathbf{k}} \rangle^2 \left(G_k G_{k-q} + G_{-k+q} - F_k \overline{F}_{k-q} - \overline{F}_k F_{k-q} \right) \\
\approx \int_{FS} \frac{dk_{\parallel}}{4\pi^2 v} \langle u | \, \hat{o} \, | \, u \rangle^2 \int_{-\Lambda}^{\Lambda} d\xi \left\{ -\frac{1}{2E} \frac{E^2 + \xi(\xi - \mathbf{v} \cdot \mathbf{q}) + i\Omega_m E}{(E + i\Omega_m)^2 - (E')^2} \right. \\
\left. + \frac{1}{4E} \frac{|\Delta|^2}{(E + i\Omega_m)^2 - (E')^2} + \frac{1}{4E} \frac{|\Delta|^2}{(E - i\Omega_m)^2 - (E')^2} \right. \\
\left. + \frac{1}{2E'} \frac{(E')^2 + \xi(\xi - \mathbf{v} \cdot \mathbf{q}) - i\Omega_m E'}{E^2 - (E' - i\Omega_m)^2} - \frac{1}{4E'} \frac{|\Delta|^2}{E^2 - (E' - i\Omega_m)^2} - \frac{|\Delta|^2}{4E'} \right\} (163)$$

As before, $E' = \sqrt{(\xi - \mathbf{v} \cdot \mathbf{q})^2 + |\Delta|^2}$. We again shift $\xi \to \xi + \mathbf{v} \cdot \mathbf{q}$ whereever necessary to ensure E' only shows up as $(E')^2$, and drop terms of $O(\mathbf{v} \cdot \mathbf{q}/\Lambda)$, yielding

$$\begin{split} \lim_{T \to 0} \sum_{\omega_n} \frac{1}{2} \int \frac{d^2 \mathbf{k}}{4\pi^2} \left\langle u_{\mathbf{k}} \right| \hat{o} \left| u_{\mathbf{k}} \right\rangle^2 \left(G_k G_{k-q} + G_{-k+q} - F_k \overline{F}_{k-q} - \overline{F}_k F_{k-q} \right) \\ \approx - \int_{FS} \frac{dk_{\parallel}}{4\pi^2 v} \left\langle u \right| \hat{o} \left| u \right\rangle^2 \int_{-\Lambda}^{\Lambda} d\xi \frac{1}{2E} \left\{ \frac{\xi(2\xi - \mathbf{v} \cdot \mathbf{q}) + i\Omega_m E}{(i\Omega_m)^2 + 2i\Omega_m E + 2\xi(\mathbf{v} \cdot \mathbf{q}) - (\mathbf{v} \cdot \mathbf{q})^2} \right. \\ \left. + \frac{\xi(2\xi + \mathbf{v} \cdot \mathbf{q}) - i\Omega_m E}{(i\Omega_m)^2 - 2i\Omega_m E - 2\xi(\mathbf{v} \cdot \mathbf{q}) - (\mathbf{v} \cdot \mathbf{q})^2} \right\} \\ = \int_{FS} \frac{dk_{\parallel}}{4\pi^2 v} \left\langle u \right| \hat{o} \left| u \right\rangle^2 \int_{-\Lambda}^{\Lambda} d\xi \frac{1}{E} \frac{1}{\left[(i\Omega_m)^2 + 2i\Omega_m E + 2\xi(\mathbf{v} \cdot \mathbf{q}) - (\mathbf{v} \cdot \mathbf{q})^2 \right] \left[(i\Omega_m)^2 - 2i\Omega_m E - 2\xi(\mathbf{v} \cdot \mathbf{q}) - (\mathbf{v} \cdot \mathbf{q})^2 \right]}{(164)} \end{split}$$

We handle this integral using the same variable transformation as for the polarization components, $\xi/|\Delta| = \sinh(x)$, $E/|\Delta| = \cosh(x)$, $\mathbf{v} \cdot \mathbf{q}/|\Delta| = ia \sinh(ib)$, and $i\Omega_m/|\Delta| = ia \cosh(ib)$. Taking $\Lambda \to \infty$,

$$\lim_{T \to 0} \sum_{\omega_n} \frac{1}{2} \int \frac{d^2 \mathbf{k}}{4\pi^2} \langle u_{\mathbf{k}} | \, \hat{o} \, | u_{\mathbf{k}} \rangle^2 \left(G_k G_{k-q} + G_{-k} G_{-k+q} - F_k \overline{F}_{k-q} - \overline{F}_k F_{k-q} \right) \\ \approx \int_{\mathrm{FS}} \frac{dk_{\parallel}}{4\pi^2 v} \langle u | \, \hat{o} \, | u \rangle^2 \int_{-\infty}^{\infty} \mathrm{d}x \, \frac{4 \cosh^2(ib)}{(2i \cosh(x+ib)-a) \, (2i \cosh(x+ib)+a)} \\ = -\int_{\mathrm{FS}} \frac{dk_{\parallel}}{4\pi^2 v} \langle u | \, \hat{o} \, | u \rangle^2 \int_{-\infty}^{\infty} \mathrm{d}x \, \frac{\cosh^2(ib)}{\cosh^2(x+ib) + (a/2)^2} \\ = -\int_{\mathrm{FS}} \frac{dk_{\parallel}}{4\pi^2 v} \langle u | \, \hat{o} \, | u \rangle^2 \, \frac{4 \cosh^2(ib)}{a \sqrt{1+(a/2)^2}} \operatorname{arcsinh}\left(\frac{a}{2}\right) \quad (165)$$

Recall that $a = -2i \sin z$ and $\sqrt{1 + (a/2)^2} = \cos z$, so

$$\lim_{T \to 0} \sum_{\omega_n} \frac{1}{2} \int \frac{\mathrm{d}^2 \mathbf{k}}{4\pi^2} \left\langle u_{\mathbf{k}} \right| \hat{o} \left| u_{\mathbf{k}} \right\rangle^2 \left(G_k G_{k-q} + G_{-k} G_{-k+q} - F_k \overline{F}_{k-q} - \overline{F}_k F_{k-q} \right) \\ \approx -\int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \left\langle u \right| \hat{o} \left| u \right\rangle^2 \frac{2z \cosh^2(ib)}{\sin z \cos z}.$$
(166)

Using the definition of $\cosh(ib)$, we further have

$$\cosh^2(ib) = \left(\frac{\Omega_m}{|\Delta|}\frac{1}{a}\right)^2 = \left(\frac{\Omega_m}{|\Delta|}\frac{1}{-2i\sin z}\right)^2 = \frac{(i\Omega_m)^2}{4|\Delta|^2}\frac{1}{\sin^2 z},\tag{167}$$

yielding

$$\lim_{T \to 0} \sum_{\omega_n} \frac{1}{2} \int \frac{\mathrm{d}^2 \mathbf{k}}{4\pi^2} \left\langle u_{\mathbf{k}} \right| \hat{o} \left| u_{\mathbf{k}} \right\rangle^2 \left(G_k G_{k-q} + G_{-k} G_{-k+q} - F_k \overline{F}_{k-q} - \overline{F}_k F_{k-q} \right) \\ \approx - \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \left\langle u \right| \hat{o} \left| u \right\rangle^2 \frac{(i\Omega_m)^2}{4|\Delta|^2} \frac{2z}{\sin^3 z \cos z}.$$
(168)

Now for the collective modes. We handle the + case (for the \mathcal{R} sources) first:

$$\begin{split} \lim_{T \to 0} \sum_{\omega_n} \int \frac{\mathrm{d}^2 \mathbf{k}}{4\pi^2} \left\langle u_{\mathbf{k}} \right| \hat{o} \left| u_{\mathbf{k}} \right\rangle \chi_{\mathbf{k}}^{(\ell)} \left(F_k G_{k-q} + G_{-k} F_{k-q} + G_k \overline{F}_{k-q} + \overline{F}_k G_{-k+q} \right) \\ &= \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \left\langle u \right| \hat{o} \left| u \right\rangle \chi^{(\ell)} \int_{-\Lambda}^{\Lambda} \mathrm{d}\xi \left\{ \frac{1}{2E} \frac{\Delta \left(-i\Omega_m + 2\xi - \mathbf{v} \cdot \mathbf{q} \right) + \overline{\Delta} \left(i\Omega_m + 2\xi - \mathbf{v} \cdot \mathbf{q} \right)}{(E + i\Omega_m)^2 - (E')^2} \\ &- \frac{1}{2E'} \frac{\Delta \left(-i\Omega_m + 2\xi - \mathbf{v} \cdot \mathbf{q} \right) + \overline{\Delta} \left(i\Omega_m + 2\xi - \mathbf{v} \cdot \mathbf{q} \right)}{E^2 - (E' - i\Omega_m)^2} \right\} \\ &= \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \left\langle u \right| \hat{o} \left| u \right\rangle \chi^{(\ell)} \int_{-\Lambda}^{\Lambda} \mathrm{d}\xi \frac{|\Delta|}{2E} \left\{ \frac{e^{i\theta} \left(-i\Omega_m + 2\xi - \mathbf{v} \cdot \mathbf{q} \right) + e^{-i\theta} \left(i\Omega_m + 2\xi - \mathbf{v} \cdot \mathbf{q} \right)}{(i\Omega_m)^2 + 2i\Omega_m E + 2\xi(\mathbf{v} \cdot \mathbf{q}) - (\mathbf{v} \cdot \mathbf{q})^2} \\ &+ \frac{e^{i\theta} \left(-i\Omega_m + 2\xi + \mathbf{v} \cdot \mathbf{q} \right) + e^{-i\theta} \left(i\Omega_m + 2\xi + \mathbf{v} \cdot \mathbf{q} \right)}{(i\Omega_m)^2 - 2i\Omega_m E - 2\xi(\mathbf{v} \cdot \mathbf{q}) - (\mathbf{v} \cdot \mathbf{q})^2} \right\} + O(\mathbf{v} \cdot \mathbf{q}/\Lambda) \\ &\approx - \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \left\langle u \right| \hat{o} \left| u \right\rangle \chi^{(\ell)} \int_{-\Lambda}^{\Lambda} \mathrm{d}\xi \frac{2|\Delta|}{E} \left\{ \cos \theta \left(2\xi + \mathbf{v} \cdot \mathbf{q} \right) i \operatorname{Im} \left[\frac{1}{\Omega_m^2 + 2\xi(\mathbf{v} \cdot \mathbf{q}) + (\mathbf{v} \cdot \mathbf{q})^2 + 2i\Omega_m E} \right] \right\}. \quad (169) \end{aligned}$$

Note that

$$\frac{1}{\Omega_m^2 + 2\xi(\mathbf{v}\cdot\mathbf{q}) + (\mathbf{v}\cdot\mathbf{q})^2 + 2i\Omega_m E} = \frac{\Omega_m^2 + 2\xi(\mathbf{v}\cdot\mathbf{q}) + (\mathbf{v}\cdot\mathbf{q})^2 - 2i\Omega_m E}{\left(\Omega_m^2 + 2\xi(\mathbf{v}\cdot\mathbf{q}) + (\mathbf{v}\cdot\mathbf{q})^2\right)^2 + \left(2\Omega_m E\right)^2},\tag{170}$$

$$\lim_{T \to 0} \sum_{\omega_n} \int \frac{\mathrm{d}^2 \mathbf{k}}{4\pi^2} \langle u_{\mathbf{k}} | \, \hat{o} \, | u_{\mathbf{k}} \rangle \, \chi_{\mathbf{k}}^{(\ell)} \left(F_k G_{k-q} + G_{-k} F_{k-q} + G_k \overline{F}_{k-q} + \overline{F}_k G_{-k+q} \right) \\
= \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \langle u | \, \hat{o} \, | u \rangle \, \chi^{(\ell)} \int_{-\Lambda}^{\Lambda} \mathrm{d}\xi \, \frac{2|\Delta|}{E} \frac{\cos \theta (2\xi + \mathbf{v} \cdot \mathbf{q}) (2i\Omega_m E) + i \sin \theta (i\Omega_m) \left(-(i\Omega_m)^2 + 2\xi (\mathbf{v} \cdot \mathbf{q}) + (\mathbf{v} \cdot \mathbf{q})^2 \right)}{\left(-(i\Omega_m)^2 + 2\xi (\mathbf{v} \cdot \mathbf{q}) + (\mathbf{v} \cdot \mathbf{q})^2 \right)^2 - \left(2i\Omega_m E \right)^2}.$$
(171)

Using again the same variable transformations as before, we obtain

$$\lim_{T \to 0} \sum_{\omega_n} \int \frac{\mathrm{d}^2 \mathbf{k}}{4\pi^2} \langle u_{\mathbf{k}} | \, \hat{o} \, | u_{\mathbf{k}} \rangle \, \chi_{\mathbf{k}}^{(\ell)} \left(F_k G_{k-q} + G_{-k} F_{k-q} + G_k \overline{F}_{k-q} + \overline{F}_k G_{-k+q} \right) \\ = 2 \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \, \langle u | \, \hat{o} \, | u \rangle \, \chi^{(\ell)} \\ \int_{-\tilde{\Lambda}}^{\tilde{\Lambda}} \mathrm{d}x \, \frac{1}{a} \frac{2i \cosh(x) \cosh(ib) \left(2 \sinh(x) + ia \sinh(ib)\right) \cos \theta - \cosh(ib) \left(a^2 + 2ia \sinh(x) \sinh(ib)\right) \sin \theta}{\left[a + 2i \sinh(x) \sinh(ib)\right]^2 + 4 \cosh^2(x) \cosh^2(ib)}.$$
(172)

First the $\cos\theta$ term:

$$\int_{-\tilde{\Lambda}}^{\tilde{\Lambda}} \mathrm{d}x \, \frac{2i\cosh(x)\cosh(ib)\left(2\sinh(x)+ia\sinh(ib)\right)}{\left[a+2i\sinh(x)\sinh(ib)\right]^2+4\cosh^2(x)\cosh^2(ib)} = \frac{i}{2}\cos b \log\left(\frac{a^2+4\cos^2 b-4a(\Lambda/|\Delta|)\sin b+4(\Lambda/|\Delta|)^2}{a^2+4\cos^2 b+4a(\Lambda/|\Delta|)\sin b+4(\Lambda/|\Delta|)^2}\right) \xrightarrow{\Lambda\to\infty} 0.$$
(173)

The sin θ term survives the $\Lambda \to \infty$ limit. We already found the necessary integral when evaluating the polarization components,

$$\operatorname{Re} \int_{-\infty}^{\infty} \mathrm{d}\xi \, \frac{1}{E} \frac{1}{(i\Omega_m)^2 + 2i\Omega_m E + 2\xi(\mathbf{v} \cdot \mathbf{q}) - (\mathbf{v} \cdot \mathbf{q})^2} = \frac{1}{|\Delta|^2} \operatorname{Re} \int_{-\infty}^{\infty} \mathrm{d}x \, \frac{1}{2ia \cosh(x + ib) - a^2} = -\frac{1}{|\Delta|^2} \frac{\operatorname{arcsinh}(a/2)}{a\sqrt{1 + (a/2)^2}} = -\frac{1}{|\Delta|^2} \frac{z}{2\sin z \cos z}.$$
 (174)

The drive coupling for $\mathcal{R}_q^{(\ell)}$ is then

$$\Pi_{q;\ell}^{d\mathcal{R}} = i^{\rho_{\ell}} \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \left\langle u \right| \hat{o} \left| u \right\rangle \chi^{(\ell)} \sin \theta \frac{i\Omega_m}{|\Delta|} \frac{iz}{\sin z \cos z}.$$
(175)

26

For the ${\mathcal I}$ drive terms, we have a similar result:

$$\begin{split} \lim_{T\to 0} \sum_{\omega_n} i \int \frac{\mathrm{d}^2 \mathbf{k}}{4\pi^2} \langle u_{\mathbf{k}} | \hat{o} | u_{\mathbf{k}} \rangle \chi_{\mathbf{k}}^{(\ell)} \left(-F_k G_{k-q} - G_{-k} F_{k-q} + G_k \overline{F}_{k-q} + \overline{F}_k G_{-k+q} \right) \\ &= i \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \langle u | \hat{o} | u \rangle \chi^{(\ell)} \int_{-\Lambda}^{\Lambda} \mathrm{d}\xi \left\{ \frac{1}{2E} \frac{\Delta \left(i\Omega_m - 2\xi + \mathbf{v} \cdot \mathbf{q} \right) + \overline{\Delta} \left(i\Omega_m + 2\xi - \mathbf{v} \cdot \mathbf{q} \right)}{(E + i\Omega_m)^2 - (E')^2} \\ &- \frac{1}{2E'} \frac{\Delta \left(i\Omega_m - 2\xi + \mathbf{v} \cdot \mathbf{q} \right) + \overline{\Delta} \left(i\Omega_m + 2\xi - \mathbf{v} \cdot \mathbf{q} \right)}{E^2 - (E' - i\Omega_m)^2} \right\} \\ &\approx i \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \langle u | \hat{o} | u \rangle \chi^{(\ell)} \int_{-\Lambda}^{\Lambda} \mathrm{d}\xi \frac{|\Delta|}{2E} \left\{ \frac{e^{i\theta} \left(i\Omega_m - 2\xi + \mathbf{v} \cdot \mathbf{q} \right) + e^{-i\theta} \left(i\Omega_m + 2\xi - \mathbf{v} \cdot \mathbf{q} \right)}{(i\Omega_m)^2 + 2i\Omega_m E + 2\xi(\mathbf{v} \cdot \mathbf{q}) - (\mathbf{v} \cdot \mathbf{q})^2} \\ &+ \frac{e^{i\theta} \left(i\Omega_m - 2\xi - \mathbf{v} \cdot \mathbf{q} \right) + e^{-i\theta} \left(i\Omega_m + 2\xi + \mathbf{v} \cdot \mathbf{q} \right)}{(i\Omega_m)^2 - 2i\Omega_m E - 2\xi(\mathbf{v} \cdot \mathbf{q}) - (\mathbf{v} \cdot \mathbf{q})^2} \right\} \\ &= i \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \langle u | \hat{o} | u \rangle \chi^{(\ell)} \int_{-\Lambda}^{\Lambda} \mathrm{d}\xi \frac{|\Delta|}{E} \left\{ \frac{\cos \theta i\Omega_m - i \sin \theta \left(2\xi - \mathbf{v} \cdot \mathbf{q} \right)}{(i\Omega_m)^2 - 2i\Omega_m E - 2\xi(\mathbf{v} \cdot \mathbf{q}) - (\mathbf{v} \cdot \mathbf{q})^2} \right\} \\ &= i \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \langle u | \hat{o} | u \rangle \chi^{(\ell)} \int_{-\Lambda}^{\Lambda} \mathrm{d}\xi \frac{|\Delta|}{E} \left\{ -\cos \theta i\Omega_m - i \sin \theta \left(2\xi - \mathbf{v} \cdot \mathbf{q} \right) \\ &- \sin \theta \left(2\xi + \mathbf{v} \cdot \mathbf{q} \right) \operatorname{Im} \left[\frac{1}{\Omega_m^2 + 2i\Omega_m E - 2\xi(\mathbf{v} \cdot \mathbf{q}) - (\mathbf{v} \cdot \mathbf{q})^2} \right] \right\} \\ &= -i \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \langle u | \hat{o} | u \rangle \chi^{(\ell)} \cos \theta i\Omega_m 2|\Delta| \int_{-\Lambda}^{\Lambda} \mathrm{d}\xi \frac{1}{E} \operatorname{Re} \left[\frac{1}{\Omega_m^2 + 2i\Omega_m E + 2\xi(\mathbf{v} \cdot \mathbf{q}) + (\mathbf{v} \cdot \mathbf{q})^2} \right] \\ &= -i \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \langle u | \hat{o} | u \rangle \chi^{(\ell)} \cos \theta i\Omega_m 2|\Delta| \int_{-\Lambda}^{\Lambda} \mathrm{d}\xi \frac{1}{E} \operatorname{Re} \left[\frac{1}{\Omega_m^2 + 2i\Omega_m E + 2\xi(\mathbf{v} \cdot \mathbf{q}) + (\mathbf{v} \cdot \mathbf{q})^2} \right] \\ &= -i \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \langle u | \hat{o} | u \rangle \chi^{(\ell)} \cos \theta i\Omega_m 2|\Delta| \int_{-\Lambda}^{\Lambda} \mathrm{d}\xi \frac{1}{E} \operatorname{Re} \left[\frac{1}{\Omega_m^2 + 2i\Omega_m E + 2\xi(\mathbf{v} \cdot \mathbf{q}) + (\mathbf{v} \cdot \mathbf{q})^2} \right] \\ &= -i \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \langle u | \hat{o} | u \rangle \chi^{(\ell)} \cos \theta i\Omega_m 2|\Delta| \int_{-\Lambda}^{\Lambda} \mathrm{d}\xi \frac{1}{E} \operatorname{Re} \left[\frac{1}{\Omega_m^2 + 2i\Omega_m E + 2\xi(\mathbf{v} \cdot \mathbf{q}) + (\mathbf{v} \cdot \mathbf{q})^2} \right] \\ &= -i \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \langle u | \hat{o} | u \rangle \chi^{(\ell)} \cos \theta i\Omega_m 2|\Delta| \int_{-\Lambda}^$$

The drive coupling for $\mathcal{I}_q^{(\ell)}$ is then

$$\Pi_{q;\ell}^{d\mathcal{I}} = -i^{\rho_{\ell}} \int \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \left\langle u \right| \hat{o} \left| u \right\rangle \chi^{(\ell)} \cos\theta \frac{i\Omega_m}{|\Delta|} \frac{iz}{\sin z \cos z}.$$
(177)

Finally, we integrate out the fluctuations (\mathcal{R} and \mathcal{I}):

$$\int_{\mathcal{R},\mathcal{I}} \exp\{-S_{\text{eff}}\} = \int_{\mathcal{R},\mathcal{I}} \exp\left\{\sum_{qXY\ell\ell'} X_{-q}^{(\ell)}(\mathcal{D}_{q}^{-1})_{\ell\ell'}^{XY}Y_{q}^{(\ell')} - \sum_{qX\ell} d_{-q}\Pi_{q;\ell}^{dX}X_{q}^{(\ell)}\right\} \\
= \exp\left\{-\sum_{qXY\ell\ell'} \frac{1}{4}d_{-q}d_{q}\Pi_{-q;\ell}^{dX}(\mathcal{D}_{q})_{\ell\ell'}^{XY}\Pi_{q;\ell'}^{dY}\right\}. \quad (178)$$

 \mathcal{D}_q is a matrix in the space of fluctuation types $X, Y \in \{\mathcal{R}, \mathcal{I}\}$ and channels ℓ, ℓ' , obtained by inverting the inverse propagator given in the previous section. To obtain the real-time response function, we analytically continue $i\Omega_m \to \Omega + i0^+$ in all expressions.

V. SINGLE PARTICLE HAMILTONIAN OF RTG

We model the single-particle band structure of rhombohedral trilayer graphene using a 6×6 tight binding model [5]. In the continuum limit, the Bloch Hamiltonian matrix is

$$h_{k} = \begin{pmatrix} \delta + \Delta_{1} + \Delta_{2} & v_{0}\pi^{*} & v_{4}\pi^{*} & v_{3}\pi & 0 & \frac{\gamma_{2}}{2} \\ v_{0}\pi & \Delta_{1} + \Delta_{2} & \gamma_{1} & v_{4}\pi^{*} & 0 & 0 \\ v_{4}\pi & \gamma_{1} & -2\Delta_{2} & v_{0}\pi^{*} & v_{4}\pi^{*} & v_{3}\pi \\ v_{3}\pi^{*} & v_{4}\pi & v_{0}\pi & -2\Delta_{2} & \gamma_{1} & v_{4}\pi^{*} \\ 0 & 0 & v_{4}\pi & \gamma_{1} & -\Delta_{1} + \Delta_{2} & v_{0}\pi^{*} \\ \frac{\gamma_{2}}{2} & 0 & v_{3}\pi^{*} & v_{4}\pi & v_{0}\pi & \delta - \Delta_{1} + \Delta_{2} \end{pmatrix},$$
(179)

where $\pi = \tau k_x + i k_y$ (τ is the valley index) and we use the basis $(A_1, B_1, A_2, B_2, A_3, B_3)$ with A_i and B_i denoting the two sites per unit cell (i.e., two sublattices) on layer *i*. Here $v_i = \sqrt{3}a\gamma_i/2$, a = 2.46Å is the lattice constant of monolayer graphene, Δ_1 is proportional to the perpendicular electric field, and γ_i , δ , Δ_2 are fixed material parameters: $\gamma_0 = 3.1 \text{ eV}$, $\gamma_1 = 0.38 \text{ eV}$, $\gamma_2 = -0.015 \text{ eV}$, $\gamma_3 = -0.29 \text{ eV}$, $\gamma_4 = -0.141 \text{ eV}$, $\delta = -0.0105 \text{ eV}$, $\Delta_2 = -0.0023 \text{ eV}$ [6].




FIG. 2. Dependence of density of states on the electron doping in RTG. Perpendicular electric field corresponds to $\Delta_1 = 20$ meV. Colors correspond to different topology of the Fermi contour




FIG. 3. Fermi surfaces in the K valley for three different electron concentrations. Colors correspond to the colors in Fig. 2.

In Fig. 2 we show the dependence of the density of states at the Fermi level on the electron doping for $\Delta_1 = 20$ meV. There exist three van Hove singularities (vHS) associated with Lifshitz transitions where the topology of the Fermi surface changes. Three different topologies are shown in Fig. 3. In what follows we will consider the hole-doped region in the vicinity of the vHS where the Fermi surface is annular with two concentric Fermi pockets, as shown in the central subfigure of Fig. 3.

VI. INTERACTION POTENTIALS

In the main text we consider two kinds of interaction potentials for RTG. The first is a phonon-mediated interaction, with the potential given by

$$\mathcal{V}_{\mathbf{k},\mathbf{k}'}^{\text{phon}} = -\mathcal{V}_0 |\langle u_{\mathbf{k}} | u_{\mathbf{k}'} \rangle|^2, \tag{180}$$

where $|u_{\mathbf{k}}\rangle$ is the Bloch wavevector of the active (valence) band into which we project. The parameter \mathcal{V}_0 is chosen such that the critical temperature is the same as for the case of IVC fluctuation-mediated superconductity described below.

The second interaction we consider is mediated by the fluctuations of the intervalley coherent order parameter [7]:

$$\mathcal{V}_{\mathbf{k},\mathbf{k}';\mathbf{q}}^{\mathrm{IVC}} = \frac{n_e/(\partial n_e/\partial \mu)}{(\mathbf{k} + \mathbf{k}' - \mathbf{q})^2 + \xi_{IVC}^{-2}} |\tilde{u}_{+,\mathbf{k}}\tilde{u}_{+,\mathbf{k}'}|^2, \tag{181}$$

where \tilde{u}_+ is the component of the Bloch wavefunction near the K valley projected to the layer and sublattice with maximal electron density. Note that this potential depends on $\mathbf{k} + \mathbf{k}'$ as opposed to $\mathbf{k} - \mathbf{k}'$. This is because IVC fluctuations are peaked at the wavevector $\mathbf{K} - \mathbf{K}'$ separating the two valleys (and they hence scatter particles between valleys), and since \mathbf{k} and \mathbf{k}' label the outgoing and incoming momenta near the K valley, measured relative to that valley. In this language, IVC scattering in the Cooper channel takes one electron to $\mathbf{K} + \mathbf{k}$ from $\mathbf{K}' - \mathbf{k}' + \mathbf{q}$ and the other to $\mathbf{K}' - \mathbf{k} + \mathbf{q}$ from $\mathbf{K} + \mathbf{k}'$, i.e. the total momentum transfer is $\mathbf{K} - \mathbf{K}' + \mathbf{k} + \mathbf{k}' - \mathbf{q}$. Assuming that $|\mathbf{q}|\xi_{\text{IVC}} \ll 1$, in practice we will evaluate \mathcal{V}^{IVC} at $\mathbf{q} = 0$. Since we will be interested in $|\mathbf{q}|$ up to the scale of the inverse BCS coherence length, this means that we require $\xi_{\text{IVC}} \ll \xi_{\text{BCS}}$.

VII. SELF-CONSISTENT ELECTROMAGNETIC RESPONSE

In this section, we address the fact that the system responds to the total electric field (including the field induced by polarization), not only the incident field. In the superconducting state, this implements the Anderson-Higgs mechanism, which substantially modifies the dispersion of the global phase (ABG) mode.

For this we will need both the response to the driving electric field (d) calculated previously, and the response to the vector potential. The latter can be obtained by the minimal coupling $\mathbf{k} \to \mathbf{k} - \mathbf{A}$, where j = x, y and we set the electron charge e = 1 as throughout. Neglecting the diamagnetic term (which vanishes for linear dispersion, which we will assume around the Fermi level), the resulting vertices are

$$\mathbb{V}_{k,k-q}^{(j)} \approx A_q^j v_{\mathbf{k}}^j \begin{pmatrix} 1 & 0\\ 0 & 1 \end{pmatrix} \tag{182}$$

where the matrix acts in Nambu space. Here we have neglected the contribution of the quantum metric to the vertex, which becomes substantial in flat band systems such as magic angle twisted bilayer graphene [8, 9], where the contribution from the band dispersion vanishes. However, in the RTG case, the contribution due to band dispersion is substantial and we may thus neglect the geometrical contribution. We then obtain the expression for the driving terms corresponding to the vector potential similarly to the expressions in Eqs. (175) and (177):

$$\Pi_{q;\ell}^{j\mathcal{R}} = -i^{\rho_{\ell}} \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \chi^{(\ell)} \sin \theta \frac{v^j (\mathbf{v} \cdot \mathbf{q})}{|\Delta|} \frac{iz}{\sin z \cos z},\tag{183}$$

$$\Pi_{q;\ell}^{j\mathcal{I}} = i^{\rho_{\ell}} \int_{\mathrm{FS}} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \chi^{(\ell)} \cos\theta \frac{v^j(\mathbf{v}\cdot\mathbf{q})}{|\Delta|} \frac{iz}{\sin z \cos z}.$$
(184)

We now introduce the linear response tensor $\tilde{\Pi}_{q}^{(\mu,\nu)}$, where $\mu, \nu = x, y, d$. The free energy can be written as

$$\mathcal{F} = \sum_{q} \tilde{\Pi}_{q}^{(\mu,\nu)} \tilde{A}_{\mu,q} \tilde{A}_{\nu,-q}, \tag{185}$$

where $A_{\mu} = (A_x, A_y, d)$. We note that the variable *d* corresponds to the drive field, which is the component of the electric field perpendicular to the RTG plane, and not to the scalar potential A^0 ; this is why we put tildes on *A*, since it combines the (gauge-invariant) electric field and (gauge-dependent) vector potential. In what follows we use the Weyl gauge $A^0 = 0$, allowing us to express the vector potential in terms of the electric field: $\mathcal{E}_j = i\Omega A_j$ for $j \in \{x, y\}$,

and $\mathcal{E}_z = d$. Ω is the (real) frequency of the applied field. We may decompose the response tensor $\tilde{\Pi}$ as the sum of collective mode contributions $\tilde{\Pi}_{\rm CM}$ and quasiparticle contributions $\tilde{\Pi}_{\rm QP}$. The former is obtained from an expression analogous to Eq. (178), and the latter is written as

$$\tilde{\Pi}_{\rm QP} = -\int_{\rm FS} \frac{\mathrm{d}k_{\parallel}}{4\pi^2 v} \frac{1}{4|\Delta|^2} \frac{2z}{\sin^3 z \cos z} \begin{pmatrix} v_x^2 (\mathbf{v} \cdot \mathbf{q})^2 & v_x v_y (\mathbf{v} \cdot \mathbf{q})^2 & -\langle u| \, \hat{o} \, |u\rangle \, v_x (\mathbf{v} \cdot \mathbf{q})(i\Omega_m) \\ v_x v_y (\mathbf{v} \cdot \mathbf{q})^2 & v_y^2 (\mathbf{v} \cdot \mathbf{q})^2 & -\langle u| \, \hat{o} \, |u\rangle \, v_y (\mathbf{v} \cdot \mathbf{q})(i\Omega_m) \\ -\langle u| \, \hat{o} \, |u\rangle \, v_x (\mathbf{v} \cdot \mathbf{q})(i\Omega_m) & -\langle u| \, \hat{o} \, |u\rangle \, v_y (\mathbf{v} \cdot \mathbf{q})(i\Omega_m) & \langle u| \, \hat{o} \, |u\rangle^2 \, (i\Omega_m)^2 \end{pmatrix}$$
(186)

before analytically continuing $i\Omega_m \to \Omega + i0^+$. Again, we note that in deriving the response functions to the in-plane components of the vector potential, we neglected the contributions from the quantum geometric tensor, which might become dominant in flat band systems [8, 9]. In the case of RTG however the Fermi velocity is large and we thus assume that the quantum geometric contributions to the current matrix elements can be neglected.

The chosen gauge $A^0 = 0$ allows us to directly relate the response tensor Π to the nonlocal electric polarization tensor $\hat{\alpha} = M^{\dagger} \Pi M$, where $P_i = \alpha_{ij} \mathcal{E}_j$, and $M = \text{diag}(i\Omega, i\Omega, 1)$. We can now solve the classical problem of an electric field \mathcal{E}_0 scattering on a two-dimensional polarizable sheet with polarizability $\hat{\alpha}(\Omega, q_x, q_y)\delta(z)$. The answer is written immediately as

$$\boldsymbol{\mathcal{E}}_{\mathbf{q}}(z) = \hat{G}_{\mathbf{q}} e^{ik_z |z|} \hat{\alpha} (\hat{I} - \hat{G}_{\mathbf{q}} \hat{\alpha})^{-1} \boldsymbol{\mathcal{E}}_{0,\mathbf{q}}$$
(187)

where $\hat{G}_{\mathbf{q}}$ is the dyadic Green's function in a mixed representation (in real space along z and taken at z, z' = 0 and in reciprocal space along x and y). This Green's function is given by

$$\hat{G}_{\mathbf{q}} = \frac{i}{2q_z} \left[\hat{I}\left(\frac{\Omega}{c}\right)^2 - (\mathbf{q}, q_z) \otimes (\mathbf{q}, q_z) \right], \tag{188}$$

where $q_z = \sqrt{(\Omega/c)^2 - \mathbf{q}^2}$ and \hat{I} is a identity matrix.

Eq. (187) can then be used to find the Anderson-Higgs renormalized response function. We rewrite Eq. (187) as

$$\boldsymbol{\mathcal{E}}_{\mathbf{q}}(z) = \hat{G}_{\mathbf{q}} e^{iq_z |z|} \mathbf{P} = \hat{G}_{\mathbf{q}}(z) \tilde{\hat{\alpha}} \boldsymbol{\mathcal{E}}_{0,\mathbf{q}}, \quad \tilde{\hat{\alpha}} = \hat{\alpha} (\hat{I} - \hat{G}_{\mathbf{q}} \hat{\alpha})^{-1}$$
(189)

The zz component of $\tilde{\alpha}$ then yields the renormalized response function to the perpendicular electric field.

VIII. CLAPPING MODE DISPERSION FOR RTG

Fig. 4 shows the absorption spectrum for RTG with IVC-mediated interactions in its *p*-wave superconducting phase, including the Anderson-Higgs mechanism via the self-consistent approach described in Sec. VII, as a function of $\mathbf{q} \parallel \hat{\mathbf{x}}$. Different directions of \mathbf{q} yield only minor differences.




FIG. 4. Absorption spectrum of superconducting RTG with IVC-mediated interactions. Parameters are the same as in Fig. 2 of the main text.

- [1] A. Leggett, Number-phase fluctuations in two-band superconductors, Progress of Theoretical Physics 36, 901 (1966).
- [2] S. Sharapov, V. Gusynin, and H. Beck, Effective action approach to the Leggett's mode in two-band superconductors, The European Physical Journal B-Condensed Matter and Complex Systems **30**, 45 (2002).
- [3] N. R. Poniatowski, J. B. Curtis, A. Yacoby, and P. Narang, Spectroscopic signatures of time-reversal symmetry breaking superconductivity, Commun. Phys. 5, 44 (2022).
- [4] A. Dalal, J. Ruhman, and V. Kozii, Field theory of a superconductor with repulsion, Phys. Rev. B 108, 214521 (2023).
- [5] F. Zhang, B. Sahu, H. Min, and A. H. MacDonald, Band structure of ABC-stacked graphene trilayers, Phys. Rev. B 82, 035409 (2010).
- [6] H. Zhou, T. Xie, A. Ghazaryan, T. Holder, J. R. Ehrets, E. M. Spanton, T. Taniguchi, K. Watanabe, E. Berg, M. Serbyn, and A. F. Young, Half-and quarter-metals in rhombohedral trilayer graphene, Nature 598, 429 (2021).
- [7] S. Chatterjee, T. Wang, E. Berg, and M. P. Zaletel, Inter-valley coherent order and isospin fluctuation mediated superconductivity in rhombohedral trilayer graphene, Nat. Commun. 13, 6013 (2022).
- [8] P. Törmä, S. Peotta, and B. A. Bernevig, Superconductivity, superfluidity and quantum geometry in twisted multilayer systems, Nature Reviews Physics 4, 528 (2022).
- [9] S. A. Chen and K. T. Law, Ginzburg-Landau theory of flat-band superconductors with quantum metric, Phys. Rev. Lett. 132, 026002 (2024).