# THE HIGHER ALGEBRA OF WEIGHTED COLIMITS

HADRIAN HEINE, UNIVERSITY OF OSLO, NORWAY, HADRIAH@MATH.UIO.NO

ABSTRACT. We develop a theory of weighted colimits in the framework of weakly bi-enriched  $\infty$ categories, an extension of Lurie's notion of enriched ∞-categories. We prove an existence result for weighted colimits, study weighted colimits of diagrams of enriched functors, express weighted colimits via enriched coends, characterize the enriched  $\infty$ -category of enriched presheaves as the free cocompletion under weighted colimits and develop a theory of universally adjoining weighted colimits to an enriched  $\infty$ -category. We use the latter technique to construct for every presentably  $\mathbb{E}_{k+1}$ -monoidal  $\infty$ -category  $\mathcal{V}$  for  $1 \leq k \leq \infty$  and class  $\mathcal{H}$  of  $\mathcal{V}$ -weights, with respect to which weighted colimits are defined, a presentably  $\mathbb{E}_k$ -monoidal structure on the  $\infty$ -category of  $\mathcal{V}$ -enriched  $\infty$ -categories that admit  $\mathcal{H}$ -weighted colimits. Varying  $\mathcal{H}$  this  $\mathbb{E}_k$ -monoidal structure interpolates between the tensor product for  $\mathcal{V}$ -enriched  $\infty$ -categories and the relative tensor product for  $\infty$ -categories presentably left tensored over  $\mathcal{V}$ . Studying functoriality in the class of  $\mathcal{V}$ -weights  $\mathcal{H}$  we deduce that forming  $\mathcal{V}$ -enriched presheaves is  $\mathbb{E}_k$ -monoidal with respect to the tensor product on small  $\mathcal{V}$ -enriched  $\infty$ -categories and the tensor product on  $\infty$ -categories presentably left tensored over  $\mathcal{V}$ . As an application we construct a  $\mathcal{V}$ -enriched version of Day-convolution and give a new construction of the tensor product for  $\infty$ categories presentably left tensored over  $\mathcal{V}$  as a  $\mathcal{V}$ -enriched localization of Day-convolution. As further applications we construct a tensor product for Cauchy-complete  $\mathcal{V}$ -enriched  $\infty$ -categories, a tensor product for  $(\infty, 2)$ -categories with (op)lax colimits and a tensor product for stable  $(\infty, n)$ -categories.

#### CONTENTS

1.	Introduction	2
1.1.	Notation and terminology	9
Acknowledgements		
2.	Enriched $\infty$ -categories	10
2.1.	∞-operads	10
2.2.	Weakly enriched $\infty$ -categories	13
2.3.	Enriched functors	16
2.4.	Tensored envelopes	21
2.5.	Enrichment	23
2.6.	Generalized enrichment	28
3.	Weighted colimits	33
3.1.	Weighted colimits for weakly enriched $\infty$ -categories	33
3.2.	Weighted colimits for pseudo-enriched $\infty$ -categories	35
3.3.	Weighted colimits for enriched $\infty$ -categories	37
3.4.	Conical colimits and tensors	38
3.5.	Weighted colimits via enriched coends	40
3.6.	A Bousfield Kan formula for weighted colimits	41
3.7.	Preservation of weighted colimits	42
3.8.	Stability of weighted colimits under restriction of enrichment	44
3.9.	Existence of weighted colimits	45
3.10	. Adjoining weighted colimits	47

HADRIAN HEINE	UNIVERSITY O	F OSLO, NORWAY,	HADRIAH@MATH.UIO.NO
---------------	--------------	-----------------	---------------------

3.11. Weighted colimits of diagrams of enriched functors	56	
4. A monoidal structure for enriched $\infty$ -categories with weighted colimits	57	
4.1. Enriched $\infty$ -categories with weighted colimits	57	
4.2. The tensor product of enriched $\infty$ -categories with weighted colimits	62	
4.3. Closedness of the monoidal structure	69	
5. Applications	72	
5.1. Monoidality of enriched presheaves	72	
5.2. A tensor product for presentable enriched $\infty$ -categories	74	
5.2.1. Presentable enriched $\infty$ -categories	74	
5.2.2. The relative tensor product as an enriched functor $\infty$ -category		
5.3. Enriched Day-convolution	80	
5.4. A tensor product for $(\infty, 2)$ -categories with lax colimits	81	
5.5. A tensor product for Cauchy-complete enriched $\infty$ -categories	84	
5.6. A tensor product for stable $(\infty, n)$ -categories	86	
References		

#### 1. INTRODUCTION

It was long time a problem in algebraic topology to construct a symmetric monoidal structure on the  $\infty$ -category of spectra. Lurie [16] gives a remarkable elegant solution for this problem:

- He axiomatizes the essential properties of the ∞-category of spectra Sp to obtain the notion of stable ∞-category [16, Definition 1.1.1.9.].
- (2) He constructs a tensor product for stable presentable ∞-categories whose tensor unit is the stable ∞-category of spectra Sp [16, 4.8.2.]. Playing the role of the tensor unit the ∞-category of spectra Sp refines to the initial E<sub>∞</sub>-algebra for the tensor product of stable presentable ∞-categories, which is identified with a closed symmetric monoidal structure on Sp.

To achieve (2) Lurie builds a tensor product for presentable  $\infty$ -categories [16, Proposition 4.8.1.17.] that may be thought of as a higher categorical analogue of the tensor product of abelian groups: the higher algebra for this tensor product is a higher categorical analogon of commutative algebra, where the  $\infty$ -category of spaces plays the role of the integers, rings and commutative rings are replaced by monoidal and symmetric monoidal presentable ∞-categories and left, right and bi-modules correspond to left, right and bitensored presentable  $\infty$ -categories. To construct a symmetric monoidal  $\infty$ -category of spectra Lurie extends the notion of spectrum to a notion of spectrum object in any presentable  $\infty$ -category and proves that forming spectrum objects is the universal recipe to turn presentable  $\infty$ -categories to stable presentable  $\infty$ -categories: he proves that forming spectrum objects defines a symmetric monoidal localization on the  $\infty$ -category  $Pr^{L}$  of presentable  $\infty$ -categories and left adjoint functors, whose local objects are the stable presentable  $\infty$ -categories [16, Proposition 4.8.2.18.]. This guarantees that the full subcategory of local objects, the full subcategory of stable presentable  $\infty$ categories, inherits a symmetric monoidal structure from  $Pr^{L}$ , for which Sp is the tensor unit [16, Corollary 4.8.2.19]. This way Lurie constructs a tensor product for spectra giving rise to brave new algebra, a homotopical version of commutative algebra, and a tensor product of stable  $\infty$ -categories giving rise to a higher categorified version of commutative algebra, which is omnipresent in the theory of higher-categorical invariants like higher derived Brauer groups [1] or secondary K-theory [19], and serves as a starting point for tensor triangular geometry [2]. Lurie's approach to a tensor product of stable homotopy types via a tensor product of stable  $\infty$ -categories is very robust and works in various other situations: studying multiplicative infinite loop space machines Gepner-Groth-Nikolaus [5, Theorem 4.6.] adapt Lurie's strategy to preadditive and additive presentable  $\infty$ -categories: the authors produce symmetric monoidal structures on the  $\infty$ -categories  $\operatorname{Mon}_{\mathbb{E}_{\infty}}(S), \operatorname{Grp}_{\mathbb{E}_{\infty}}(S)$  of  $\mathbb{E}_{\infty}$ -spaces and grouplike  $\mathbb{E}_{\infty}$ -spaces via constructing tensor products for preadditive and additive presentable  $\infty$ -categories, respectively, for which the  $\infty$ -categories  $\operatorname{Mon}_{\mathbb{E}_{\infty}}(S)$  and  $\operatorname{Grp}_{\mathbb{E}_{\infty}}(S)$  are the respective tensor units [5, Theorem 5.1.]. In a similar fashion one can produce tensor products for compact spectra, compact grouplike  $\mathbb{E}_{\infty}$ -spaces and compact  $\mathbb{E}_{\infty}$ -spaces by constructing tensor products for small stable  $\infty$ -categories, additive  $\infty$ -categories, preadditive  $\infty$ -category  $\operatorname{Cat}_{\infty}^{\operatorname{rex}}$  of small  $\infty$ -categories having finite colimits [10, Lemma 8.15.] and the  $\infty$ -category  $\operatorname{Cat}_{\infty}^{\operatorname{rex}}$  of small  $\infty$ -categories having finite colimits [10, Lemma 8.15.] and the  $\infty$ -category  $\operatorname{Cat}_{\infty}^{\operatorname{rex}}$  of small  $\infty$ -categories having finite colimits, into which  $\operatorname{Pr}^{\mathrm{L}}$  embeds. Consequently, the tensor products of (presentable) stable, additive and preadditive  $\infty$ -categories are all derived from symmetric monoidal structures on the  $\infty$ -categories lawing fine colomits, into which  $\operatorname{Pr}^{\mathrm{L}}$  embeds. Consequently, the tensor products of (presentable) stable, additive and preadditive  $\infty$ -categories are all derived from symmetric monoidal structures on the  $\infty$ -categories  $\operatorname{Cat}_{\infty}^{\operatorname{cc}}$ ,  $\operatorname{Cat}_{\infty}^{\operatorname{cc}}$  and  $\operatorname{Cat}_{\infty}^{\mathrm{L}}$ , respectively, which are instances of the following powerful construction: for any set  $\mathcal{K}$  of small  $\infty$ -categories Lurie [16, Proposition 4.8.1.3.] constructs a closed symmetric monoidal structure on the subcategory  $\operatorname{Cat}_{\infty}(\mathcal{K}) \subset \operatorname{Cat}_{\infty}$  of small  $\infty$ -categories having  $\mathcal{K}$ -indexed colimits and functors preserving  $\mathcal{K}$ -indexed colimits. This symmetric monoidal structure is characterized by the following key properties:

- (1) The symmetric monoidal structure on  $Cat_{\infty}(\mathcal{K})$  is closed. The internal hom between two  $\infty$ -categories  $\mathcal{C}, \mathcal{D}$  having  $\mathcal{K}$ -indexed colimits is the full subcategory  $\operatorname{Fun}^{\mathcal{K}}(\mathcal{C}, \mathcal{D}) \subset \operatorname{Fun}(\mathcal{C}, \mathcal{D})$  of functors preserving  $\mathcal{K}$ -indexed colimits.
- (2) For  $\mathcal{K} = \emptyset$  the empty set the symmetric monoidal structure on  $Cat_{\infty}(\emptyset) = Cat_{\infty}$  is the cartesian structure.
- (3) For K' ⊂ K an inclusion of subsets of small ∞-categories the subcategory inclusion Cat<sub>∞</sub>(K) ⊂ Cat<sub>∞</sub>(K') admits a symmetric monoidal left adjoint.

Taking  $\mathcal{K}$  to be the set of finite  $\infty$ -categories gives the closed symmetric monoidal structure on  $\mathsf{Cat}_{\infty}^{\mathrm{rex}}$  leading to the tensor product of small stable  $\infty$ -categories, choosing  $\mathcal{K}$  to be the set of finite sets gives the closed symmetric monoidal structure on  $\mathsf{Cat}_{\infty}^{\mathrm{IL}}$  leading to the tensor products of small additive and preadditive  $\infty$ -categories. For  $\mathcal{K}$  the large collection of all small  $\infty$ -categories and the corresponding construction  $\widehat{\mathsf{Cat}}_{\infty}(\mathcal{K})$  for not necessarily small  $\infty$ -categories one obtains the closed symmetric monoidal structure on  $\mathsf{Cat}_{\infty}^{\mathrm{cc}}$  leading to the tensor product of presentable stable  $\infty$ -categories. For general set  $\mathcal{K}$  of small  $\infty$ -categories the tensor product on  $\widehat{\mathsf{Cat}}_{\infty}(\mathcal{K})$  is understood best as interpolating between the tensor product on  $\mathsf{Cat}_{\infty}^{\mathrm{cc}}$  and the cartesian structure on the  $\infty$ -category  $\widehat{\mathsf{Cat}}_{\infty}$  of not necessarily small  $\infty$ -categories a factorization

(1) 
$$\operatorname{Cat}_{\infty}^{\operatorname{cc}} = \widehat{\operatorname{Cat}}_{\infty}(\operatorname{Cat}_{\infty}) \subset \widehat{\operatorname{Cat}}_{\infty}(\mathcal{K}) \subset \widehat{\operatorname{Cat}}_{\infty} = \widehat{\operatorname{Cat}}_{\infty}(\emptyset),$$

where both inclusions admit symmetric monoidal left adjoints. The composite of both left adjoints assigns to any small  $\infty$ -category  $\mathcal{C}$  the presentable  $\infty$ -category  $\mathcal{P}(\mathcal{C})$  of presheaves on  $\mathcal{C}$  [16, Theorem 5.1.5.6.] and so restricts to a symmetric monoidal functor  $\mathsf{Cat}_{\infty} \to \mathrm{Pr}^{\mathrm{L}}$ . This proves in particular that the functor of presheaves is symmetric monoidal.

It is goal of this work to build a theory of colimits and limits in enriched  $\infty$ -category theory, which is an  $\infty$ -categorical extension of the theory of weighted colimits and limits in classical enriched category theory, and to construct an analogoue of Lurie's tensor product on  $\mathsf{Cat}_{\infty}(\mathcal{K})$  in  $\mathcal{V}$ -enriched  $\infty$ -category theory, where  $\mathcal{V}$  is any presentably  $\mathbb{E}_2$ -monoidal  $\infty$ -category. To draw this analogy we follow the subsequent table of analogies, which we explain in the following:

Non-enriched	Enriched
S	ν
∞-categories	$\mathcal{V}$ -enriched $\infty$ -categories
colimits	weighted colimits
the cartesian structure on $Cat_\infty$	the tensor product of $\mathcal{V}$ -enriched $\infty$ -categories
the tensor product on $Pr^{L}$	the relative tensor product on $\mathrm{LMod}_{\mathcal{V}}(\mathrm{Pr}^{\mathrm{L}})$
the $\infty$ -category $\mathcal{P}(\mathcal{C})$ of presheaves	the $\mathcal{V}$ -enriched $\infty$ -category $\mathcal{P}_{\mathcal{V}}(\mathcal{C})$ of $\mathcal{V}$ -enriched presheaves
a set $\mathcal{K}$ of small $\infty$ -categories	a set $\mathcal{H}$ of $\mathcal{V}$ -enriched presheaves (= $\mathcal{V}$ -weights)

To understand this table, we first need to explain the concept of weighted colimit. For that we draw an analogy to colimits in the non-enriched setting. A cocone on a functor  $F: \mathcal{J} \to \mathcal{D}$  between  $\infty$ categories is an object  $X \in \mathcal{D}$  equipped with a natural transformation from F to the constant diagram on X, which is the same datum as a map of presheaves on  $\mathcal{J}$  from the final presheaf to the presheaf  $\mathcal{D}(F(-), X)$ . The colimit of a functor  $F : \mathcal{J} \to \mathcal{D}$ , if it exists, is a cocone colim(F) on F inducing for every object Y of  $\mathcal{D}$  an equivalence  $\mathcal{D}(\operatorname{colim}(F), Y) \to \lim(\mathcal{D}(F(-), Y))$ . The notion of weighted colimit arises by replacing the notion of cocone by the notion of weighted cocone: if  $\mathcal{J}, \mathcal{D}$  are  $\infty$ -categories enriched in a presentably monoidal  $\infty$ -category  $\mathcal{V}$  and F is a  $\mathcal{V}$ -enriched functor  $\mathcal{J} \to \mathcal{D}$ , it is natural to replace presheaves on  $\mathcal{J}$  by  $\mathcal{V}$ -enriched presheaves on  $\mathcal{J}$  and the presheaf  $\mathcal{D}(F(-), X)$  on  $\mathcal{J}$  by the  $\mathcal{V}$ -enriched presheaf  $\operatorname{Mor}_{\mathcal{D}}(F(-),X)$ , where  $\operatorname{Mor}_{\mathcal{D}}(-,-)$  is the morphism object in  $\mathcal{V}$ . Moreover it is natural to replace the final presheaf on  $\mathcal{J}$  by the constant  $\mathcal{V}$ -enriched presheaf on the tensor unit of  $\mathcal{V}$ . But in this situation the following problem arises: there is generally no constant  $\mathcal{V}$ -enriched presheaf on  $\mathcal{J}$  since there is generally no  $\mathcal{V}$ -enriched functor from  $\mathcal{J}$  to the  $\mathcal{V}$ -enriched  $\infty$ -category  $B\mathbb{1}_{\mathcal{V}}$  classifying objects. The solution to this problem is to view any enriched presheaf as a possible analogon for the final presheaf and to consider colimits relative to a V-enriched presheaf on J, which one calls a V-weight on  $\mathcal{J}$  in this context. In analogy to the case of colimits a  $\mathcal{V}$ -enriched cocone on a  $\mathcal{V}$ -enriched functor  $F: \mathcal{J} \to \mathcal{D}$  is an object  $X \in \mathcal{D}$  equipped with a map  $H \to Mor_{\mathcal{D}}(F(-), X)$  of  $\mathcal{V}$ -enriched presheaves on J. To refer to H we call such a V-enriched cocone on F a H-weighted cocone on F. The H-weighted colimit of a  $\mathcal{V}$ -enriched functor  $F : \mathcal{J} \to \mathcal{D}$  if it exists, is a H-weighted cocone colim<sup>H</sup>(F) on F that induces for every object X of  $\mathcal{D}$  an equivalence  $\operatorname{Mor}_{\mathcal{D}}(\operatorname{colim}^{H}(F), Y) \to \operatorname{Mor}_{\mathcal{P}_{\mathcal{V}}(\mathcal{A})}(H, \operatorname{Mor}_{\mathcal{D}}(F(-), X)).$ We call a  $\mathcal{V}$ -weight on a  $\mathcal{V}$ -enriched  $\infty$ -category  $\mathcal{J}$  small if  $\mathcal{J}$  is a small  $\mathcal{V}$ -enriched  $\infty$ -category meaning that the space of objects is small. If any  $\mathcal{V}$ -enriched functor  $F: \mathcal{J} \to \mathcal{D}$  admits a H-weighted colimit, one says that  $\mathcal D$  admits H-weighted colimits. For any collection  $\mathcal H$  of  $\mathcal V$ -weights we say that  $\mathcal D$  admits  $\mathcal{H}$ -weighted colimits if it admits H-weighted colimits for every  $\mathbf{H} \in \mathcal{H}$ .

In nature weighted colimits are ubiquitious. To get a feeling for this concept, it is reasonable to consider two important classes of weighted colimits, which are more elementary to understand, and to think of arbitrary weighted colimits as an interpolation between these two classes. The first sort of weighted colimits are conical colimits, which are the natural enriched analogon of colimits of diagrams in an enriched  $\infty$ -category indexed by a non-enriched  $\infty$ -category. Given a  $\mathcal{V}$ -enriched  $\infty$ -category  $\mathcal{D}$  and a functor F from an  $\infty$ -category K to the underlying  $\infty$ -category of  $\mathcal{D}$  one can consider the colimit colim(F) of F satisfying the universal property that the natural map on mapping spaces

$$\mathcal{D}(\operatorname{colim}(F), Y) \to \lim(\mathcal{D}(F(-), Y))$$

is an equivalence. But one can also use the enrichment and ask for the stronger condition that the natural morphism on morphism objects

## $\operatorname{Mor}_{\mathcal{D}}(\operatorname{colim}(F), Y) \to \lim(\operatorname{Mor}_{\mathcal{D}}(F(-), Y))$

is an equivalence. If this holds, we call F the conical colimit of F. For example any stable  $\infty$ -category C has a canonical enrichment in spectra [10, Theorem 8.1.]. The loops of any object X in C, i.e. the pullback  $0 \times_X 0$  in C, is a conical pullback in the spectral  $\infty$ -category underlying C, which implies that the invertible adjunction  $\Sigma : \mathbb{C} \neq \mathbb{C} : \Omega$  of suspension and loops refines to a spectral adjunction.

A functor F from an  $\infty$ -category K to the underlying  $\infty$ -category of  $\mathcal{D}$  corresponds to a  $\mathcal{V}$ -enriched functor  $F' : K_{\mathcal{V}} \to \mathcal{D}$  from the  $\mathcal{V}$ -enriched  $\infty$ -category  $K_{\mathcal{V}}$  arising from K by tensoring the mapping spaces of K with the tensor unit of  $\mathcal{V}$ . Since  $\mathcal{V}$ -enriched presheaves on  $K_{\mathcal{V}}$  are identified with nonenriched  $\mathcal{V}$ -valued presheaves on K, there is a constant  $\mathcal{V}$ -enriched presheaf on any object of  $\mathcal{V}$  and the conical colimit of F is the colimit of F' weighted with respect to the constant  $\mathcal{V}$ -enriched presheaf on the tensor unit of  $\mathcal{V}$ .

The second sort of weighted colimits are tensors, exclusively defined for  $\mathcal{V}$ -enriched functors  $\mathcal{Bl}_{\mathcal{V}} \to \mathcal{D}$ , which classify objects of  $\mathcal{D}$ . Similarly,  $\mathcal{V}$ -enriched presheaves on  $\mathcal{Bl}_{\mathcal{V}}$  classify objects of  $\mathcal{V}$  and the tensor of an object  $V \in \mathcal{V}$  with an object  $Y \in \mathcal{D}$  is the colimit of the  $\mathcal{V}$ -enriched functor  $\mathcal{Bl}_{\mathcal{V}} \to \mathcal{D}$  classifying Y weighted with respect to the  $\mathcal{V}$ -enriched presheaf on  $\mathcal{Bl}_{\mathcal{V}}$  classifying V. So by definition the tensor of V and Y corepresents the  $\mathcal{V}$ -enriched functor  $\mathcal{D} \to \mathcal{V}, X \mapsto \operatorname{Mor}_{\mathcal{V}}(V, \operatorname{Mor}_{\mathcal{D}}(Y, X))$ . If all tensors exist, the  $\mathcal{V}$ -enrichment endows the underlying  $\infty$ -category of  $\mathcal{D}$  with a closed left  $\mathcal{V}$ -action whose internal hom recovers the  $\mathcal{V}$ -enrichment [10, Theorem 6.7.].

Another example of weighted (co)limits appears in higher category theory under the names lax and oplax (co)limits and lax and oplax pushouts and pullbacks (Definition 5.36, Definition 5.41). Given functors  $\alpha : \mathcal{A} \to \mathbb{C}, \beta : \mathbb{B} \to \mathbb{C}$  of  $\infty$ -categories one can consider the pullback  $\mathcal{A} \times_{\mathbb{C}} \times \mathbb{B}$  whose objects are triples (A  $\in \mathcal{A}, B \in \mathcal{B}, \gamma : \alpha(A) \simeq \beta(B)$ ). But one can also consider the lax pullback  $\mathcal{A} \times_{\mathbb{C}}^{e} \times \mathbb{B}$  whose objects are triples (A  $\in \mathcal{A}, B \in \mathcal{B}, \gamma : \alpha(A) \simeq \beta(B)$ ) as well as the oplax pullback  $\mathcal{A} \times_{\mathbb{C}}^{e} \times \mathcal{B}$  whose objects are triples (A  $\in \mathcal{A}, B \in \mathcal{B}, \gamma : \alpha(A) \to \beta(B)$ ) as well as the oplax pullback  $\mathcal{A} \times_{\mathbb{C}}^{e} \otimes \mathcal{B}$  whose objects are triples (A  $\in \mathcal{A}, B \in \mathcal{B}, \gamma : \alpha(A) \to \beta(B)$ ). In higher category theory the lax and oplax versions of the pullback are much more useful since they make more use of non-invertible morphisms: the pullback of two functors  $* \to \mathbb{C}$  classifying objects A, B of C is the space of equivalences between A and B while the lax pullback is the space of morphisms A  $\to B$  in C and the oplax pullback is the space of morphisms B  $\to A$  in C.

(Op)lax pullbacks can be described as weighted colimits for a simple weight: the (op)lax pullback  $\mathcal{A} \times_{\mathbb{C}}^{\mathrm{lax}} \times \mathcal{B}$  is the limit of the diagram  $\mathcal{A} \to \mathcal{C}, \mathcal{B} \to \mathcal{C}$  of  $\infty$ -categories corresponding to a functor  $F : \{0,1\}^{\triangleright} \to \mathsf{Cat}_{\infty}$  weighted with respect to the functor  $H : \{0,1\}^{\triangleright} \to \mathsf{Cat}_{\infty}$  corresponding to the diagram  $\{0\} \subset [1] \supset \{1\}$ , where we view F and H as functors enriched in  $\mathsf{Cat}_{\infty}$ . Similarly, the oplax (co)limit of a functor  $\mathcal{J} \to \mathcal{C}$  of  $(\infty, 2)$ -categories, studied in [7] for  $\mathcal{J}$  an  $\infty$ -category, is the colimit weighted with respect to a  $\mathsf{Cat}_{\infty}$ -enriched weight on  $\mathcal{J}$  that assigns to any object Z of  $\mathcal{J}$  an (op)lax version of the slice  $\infty$ -category  $\mathcal{J}_{Z/}$ .

In this work we build a theory of weighted colimits and limits that extends the classical theory of weighted (colimits) from enriched category theory to enriched  $\infty$ -category theory. Among many other we prove the following key results:

- (1) We prove an existence result of weighted colimits that splits the existence of weighted colimits in the existence of conical colimits and tensors (Proposition 3.63).
- (2) We prove that enriched left adjoints preserve weighted colimits, enriched right adjoints preserve weighted limits (Proposition 3.55) and the enriched Yoneda-embedding preserves weighted limits (Corollary 3.66).
- (3) We characterize the V-enriched ∞-category of V-enriched presheaves on a small V-enriched ∞-category as the free cocompletion under small weighted colimits (Corollary 3.67).
- (4) We study enriched (co)ends as examples of weighted (co)limits and prove that arbitrary weighted (co)limits can be constructed from tensors and weighted (co)ends (Theorem 3.38).
- (5) We study enriched functoriality of weighted colimits in the weight and the enriched functor (Proposition 3.23, Corollary 3.42).
- (6) We prove a Bousfield-Kan formula for weighted colimits that computes weighted colimits as a geometric realization of colimits indexed by spaces (Proposition 3.43).
- (7) We prove that weighted colimits in enriched ∞-categories of enriched functors are formed object-wise (Theorem 3.94).

- (8) We characterize presentably V-tensored ∞-categories as V-presentable ∞-categories, an enriched analogon of presentability expressed via weighted colimits (Theorem 5.12).
- (9) We develop a framework of universally adjoining weighted colimits to an enriched  $\infty$ -category (Proposition 3.85).
- (10) We study absolute weighted colimits, those which are preserved by any  $\mathcal{V}$ -enriched functor, and construct the Cauchy-completion by universally adjoining absolute weighted colimits to an enriched  $\infty$ -category (Notation 5.48, Proposition 5.54).

To prove these results we use a theory of bi-enriched  $\infty$ -categories of [9] that extends enriched  $\infty$ -category theory and is especially useful to describe enrichment on  $\infty$ -categories of enriched functors, which is for example essential to prove (7). Consequently, we develop our theory of weighted colimits in the more general framework of bi-enriched  $\infty$ -categories. However, our definition of weighted colimits specializes to the definition of weighted colimits for enriched  $\infty$ -categories of [15, § 6] (Remark 3.22).

We use the technique of universally adjoining weighted colimits to an enriched  $\infty$ -category (9) to construct a tensor product for  $\mathcal{V}$ -enriched  $\infty$ -categories that admit weighted colimits for a fixed set of weights:

**Theorem 1.1.** (Theorem 4.42 (1)) Let  $\mathcal{V}$  be a presentably  $\mathbb{E}_{k+1}$ -monoidal  $\infty$ -category for  $0 \le k \le \infty$  and  $\mathcal{H}$  a set of small  $\mathcal{V}$ -weights. There is a presentably  $\mathbb{E}_k$ -monoidal structure on the  $\infty$ -category

#### $\operatorname{Enr}_{\mathcal{V}}(\mathcal{H})$

of small  $\mathcal{V}$ -enriched  $\infty$ -categories having  $\mathcal{H}$ -weighted colimits and  $\mathcal{V}$ -enriched functors preserving  $\mathcal{H}$ -weighted colimits. For every  $\mathcal{C}, \mathcal{D} \in \operatorname{Enr}_{\mathcal{V}}(\mathcal{H})$  the internal hom is the  $\mathcal{V}$ -enriched  $\infty$ -category of  $\mathcal{V}$ -enriched functors  $\mathcal{C} \to \mathcal{D}$  preserving  $\mathcal{H}$ -weighted colimits.

Specializing Theorem 1.1 to (op)lax colimits and (op)lax pushouts we obtain the following corollaries:

**Corollary 1.2.** Let  $\mathcal{K}$  be a set of  $(\infty, 2)$ -categories. There is a presentably symmetric monoidal structure on the  $\infty$ -category

# $\mathsf{Cat}_{(\infty,2)}^{(\mathrm{op})\mathrm{lax}}(\mathcal{K})$

of  $(\infty, 2)$ -categories having K-indexed (op)lax colimits and functors preserving K-indexed (op)lax colimits.

**Corollary 1.3.** There is a presentably symmetric monoidal structure on the  $\infty$ -category

# $\mathsf{Cat}^{\mathrm{push},(\mathrm{op})\mathrm{lax}}_{(\infty,2)}$

of  $(\infty, 2)$ -categories having (op)lax pushouts and functors preserving (op)lax pushouts.

In Theorem 1.1 the condition of smallness on a  $\mathcal{V}$ -enriched  $\infty$ -category is crucial: there is no tensor product of not necessarily small  $\mathcal{V}$ -enriched  $\infty$ -categories having  $\mathcal{H}$ -weighted colimits. The reason is already visible for enrichment in the cartesian monoidal  $\infty$ -category  $\mathcal{S}$  of small spaces: there is no tensor product on the  $\infty$ -category of not necessarily small  $\mathcal{S}$ -enriched, i.e. locally small,  $\infty$ -categories having small colimits and functors preserving small colimits since any such tensor product destroys local smallness. The remedy is to consider presentable  $\infty$ -categories, which are locally small and to which the tensor product restricts, or to consider locally large  $\infty$ -categories having small colimits whose very large  $\infty$ -category carries a closed symmetric monoidal structure. A locally large  $\infty$ -category may be viewed as an  $\infty$ -category enriched in the  $\infty$ -category of large spaces, which arises from the  $\infty$ -category of small spaces  $\mathcal{S}$  by formally adding large filtered colimits. In analogy we may think of a locally large  $\mathcal{V}$ -enriched  $\infty$ -category as an  $\infty$ -category enriched in the formal completion of  $\mathcal{V}$ under large filtered colimits. Lurie [16, Definition 4.2.1.25.] introduces for every monoidal  $\infty$ -category  $\mathcal{V}$  a notion of  $\infty$ -category pseudo-enriched in  $\mathcal{V}$ , which we identify with an  $\infty$ -category enriched in the  $\infty$ -category of presheaves on  $\mathcal{V}$  equipped with Day-convolution [9, Corollary 4.31.]. Similarly, in [9, Definition 3.28.] we introduce for every monoidal  $\infty$ -category  $\mathcal{V}$  compatible with small colimits a notion of  $\infty$ -category quasi-enriched in  $\mathcal{V}$ , which we identify with an  $\infty$ -category enriched in the  $\infty$ -category of presheaves on  $\mathcal{V}$  preserving small limits, a model for the formal completion of  $\mathcal{V}$  under large filtered colimits. Consequently, we may view an  $\infty$ -category quasi-enriched in  $\mathcal{V}$  as a locally large  $\mathcal{V}$ -enriched  $\infty$ -category. We prove analogues of Theorem 1.1 for quasi-enriched and pseudo-enriched  $\infty$ -categories:

#### **Theorem 1.4.** (Theorem 4.42 (2) and (3)) Let $0 \le k \le \infty$ .

(1) Let  $\mathcal{V}$  be an  $\mathbb{E}_{k+1}$ -monoidal  $\infty$ -category compatible with small colimits and  $\mathcal{H}$  a collection of not necessarily small  $\mathcal{V}$ -weights. There is a closed  $\mathbb{E}_k$ -monoidal structure on the  $\infty$ -category

### $\operatorname{QEnr}_{\mathcal{V}}(\mathcal{H})$

of not necessarily small  $\infty$ -categories quasi-enriched in V having  $\mathcal{H}$ -weighted colimits and V-enriched functors preserving  $\mathcal{H}$ -weighted colimits.

(2) Let  $\mathcal{V}$  be a small  $\mathbb{E}_{k+1}$ -monoidal  $\infty$ -category and  $\mathcal{H}$  a set of small  $\mathcal{V}$ -weights. There is a presentably  $\mathbb{E}_k$ -monoidal structure on the  $\infty$ -category

#### $\operatorname{PEnr}_{\mathcal{V}}(\mathcal{H})$

of small  $\infty$ -categories pseudo-enriched in  $\mathcal{V}$  having  $\mathcal{H}$ -weighted colimits and  $\mathcal{V}$ -enriched functors preserving  $\mathcal{H}$ -weighted colimits.

For  $\mathcal{H}$  the collection of all small  $\mathcal{V}$ -weights the  $\infty$ -category  $\operatorname{QEnr}_{\mathcal{V}}(\mathcal{H})$  is the  $\infty$ -category  $\operatorname{QEnr}_{\mathcal{V}}^{cc}$  of large  $\infty$ -categories quasi-enriched in  $\mathcal{V}$  that admit small weighted colimits, which by Theorem 1.4 admits a closed  $\mathbb{E}_k$ -monoidal structure. We identify  $\operatorname{QEnr}_{\mathcal{V}}^{cc}$  with the  $\infty$ -category  $\operatorname{LMod}_{\mathcal{V}}(\operatorname{Cat}_{\infty}^{cc})$  of  $\infty$ -categories left tensored over  $\mathcal{V}$  compatible with small colimits (Remark 5.4). For general set  $\mathcal{H}$  of  $\mathcal{V}$ -weights the tensor product on  $\operatorname{QEnr}_{\mathcal{V}}(\mathcal{H})$  interpolates between the relative tensor product on  $\operatorname{LMod}_{\mathcal{V}}(\operatorname{Cat}_{\infty}^{cc}) \simeq \operatorname{QEnr}_{\mathcal{V}}^{cc}$  and a tensor product of  $\infty$ -categories quasi-enriched in  $\mathcal{V}$ : the inclusion  $\emptyset \subset \mathcal{H} \subset \{ \text{small } \mathcal{V} - \text{weights} \}$  gives lax  $\mathbb{E}_k$ -monoidal inclusions

$$\operatorname{LMod}_{\mathcal{V}}(\operatorname{\mathsf{Cat}}_{\infty}^{\operatorname{cc}}) \subset \operatorname{QEnr}_{\mathcal{V}}(\mathcal{H}) \subset \operatorname{QEnr}_{\mathcal{V}} = \operatorname{QEnr}_{\mathcal{V}}(\emptyset).$$

The  $\mathbb{E}_k$ -monoidal left adjoint of the inclusion  $\operatorname{LMod}_{\mathcal{V}}(\operatorname{Cat}_{cc}^{cc}) \subset \operatorname{QEnr}_{\mathcal{V}}$  sends any small  $\mathcal{V}$ -enriched  $\infty$ -category  $\mathcal{C}$  to the presentably left tensored  $\infty$ -category  $\mathcal{P}_{\mathcal{V}}(\mathcal{C})$  of  $\mathcal{V}$ -enriched presheaves on  $\mathcal{C}$  and so restricts to an  $\mathbb{E}_k$ -monoidal functor  $\operatorname{Enr}_{\mathcal{V}} \to \operatorname{LMod}_{\mathcal{V}}(\operatorname{Pr}^{\mathrm{L}})$  to the  $\infty$ -category of  $\infty$ -categories presentably left tensored over  $\mathcal{V}$ . This way we obtain the following corollary, which connects the higher algebra of  $\mathcal{V}$ -enriched  $\infty$ -categories with the considerably more tractable higher algebra in the  $\infty$ -category  $\operatorname{Pr}^{\mathrm{L}}$  of presentable  $\infty$ -categories:

**Corollary 1.5.** (Corollary 5.5) Let  $\mathcal{V}$  be a presentably  $\mathbb{E}_{k+1}$ -monoidal  $\infty$ -category for  $1 \leq k \leq \infty$ . The functor

$$\mathcal{P}_{\mathcal{V}}: \operatorname{Enr}_{\mathcal{V}} \to \operatorname{LMod}_{\mathcal{V}}(\operatorname{Pr}^{\mathrm{L}})$$

of  $\mathcal{V}$ -enriched presheaves is an  $\mathbb{E}_k$ -monoidal functor, where the left hand side carries the tensor product of  $\mathcal{V}$ -enriched  $\infty$ -categories and the right hand side carries the relative tensor product.

By Corollary 1.5 the functor of  $\mathcal{V}$ -enriched presheaves is  $\mathbb{E}_k$ -monoidal and so preserves  $\mathbb{E}_k$ -algebras. As a consequence the  $\mathcal{V}$ -enriched  $\infty$ -category of  $\mathcal{V}$ -enriched presheaves on any  $\mathbb{E}_k$ -monoidal  $\mathcal{V}$ -enriched  $\infty$ -category is an  $\mathbb{E}_k$ -algebra in  $\mathrm{LMod}_{\mathcal{V}}(\mathrm{Pr}^{\mathrm{L}})$  and the  $\mathcal{V}$ -enriched Yoneda-embedding is  $\mathbb{E}_k$ -monoidal. In particular, for  $k = \infty$  the  $\mathcal{V}$ -enriched  $\infty$ -category of  $\mathcal{V}$ -enriched presheaves on a symmetric monoidal  $\mathcal{V}$ -enriched  $\infty$ -category is a  $\mathcal{V}$ -algebra in  $\mathrm{Pr}^{\mathrm{L}}$ , i.e. a presentably symmetric monoidal  $\infty$ -category under  $\mathcal{V}$ . Hinich [15, Proposition 8.4.3.] constructs for every  $\mathbb{E}_{k+1}$ -monoidal  $\infty$ -category  $\mathcal{V}$  an  $\mathbb{E}_k$ -monoidal  $\mathcal{V}$ -enriched Yoneda-embedding, which satisfies the  $\mathbb{E}_k$ -monoidal version of free cocompletion and therefore is canonically equivalent to our construction of  $\mathbb{E}_k$ -monoidal  $\mathcal{V}$ -enriched Yoneda-embedding. This gives an affirmative answer to a question of [3, Question 1.6.].

We combine Corollary 1.5 with the following theorem to construct a  $\mathcal{V}$ -enriched version of Dayconvolution: Lurie [16, Proposition 4.8.1.17.] gives an explicite formula for the tensor product of presentable  $\infty$ -categories by proving that for every presentable  $\infty$ -categories  $\mathcal{C}, \mathcal{D}$  there is a canonical equivalence

$$\mathcal{C} \otimes \mathcal{D} \simeq \operatorname{Fun}^{\mathrm{R}}(\mathcal{C}^{\operatorname{op}}, \mathcal{D}).$$

where the right hand side is the  $\infty$ -category of right adjoint functors  $\mathcal{C}^{\mathrm{op}}$  to  $\mathcal{D}$ . We prove an enriched version of Lurie's result:

**Theorem 1.6.** (Theorem 5.8, Corollary 5.25) Let  $\mathcal{V}$  be a presentably symmetric monoidal  $\infty$ -category,  $\mathcal{B} \in \operatorname{Enr}_{\mathcal{V}}$  and  $\mathcal{C}, \mathcal{D} \in \operatorname{LMod}_{\mathcal{V}}(\operatorname{Pr}^{L})$ . There are canonical  $\mathcal{V}$ -enriched equivalences

$$\mathcal{C} \otimes_{\mathcal{V}} \mathcal{D} \simeq \operatorname{EnrFun}_{\mathcal{V}}^{\mathrm{R}}(\mathcal{C}^{\mathrm{op}}, \mathcal{D}),$$

$$\mathcal{P}_{\mathcal{V}}(\mathcal{B}) \otimes_{\mathcal{V}} \mathcal{D} \simeq \operatorname{EnrFun}_{\mathcal{V}}(\mathcal{B}^{\operatorname{op}}, \mathcal{D}),$$

where the right hand side is the V-enriched  $\infty$ -category of right adjoint V-enriched functors  $\mathcal{C}^{\mathrm{op}}$  to  $\mathcal{D}$ .

Using that  $\mathbb{E}_{\infty}$ -algebras in  $\mathrm{LMod}_{\mathcal{V}}(\mathrm{Pr}^{\mathrm{L}})$  are  $\mathcal{V}$ -algebras in  $\mathrm{Pr}^{\mathrm{L}}$  we obtain the following corollary:

**Corollary 1.7.** (Corollary 5.31) Let  $\mathcal{V}$  be a presentably symmetric monoidal  $\infty$ -category,  $\mathcal{C}$  a small symmetric monoidal  $\mathcal{V}$ -enriched  $\infty$ -category and  $\mathcal{D}$  a  $\mathcal{V}$ -algebra in  $\operatorname{Pr}^{\mathrm{L}}$ . Then  $\operatorname{EnrFun}_{\mathcal{V}}(\mathcal{C}^{\operatorname{op}}, \mathcal{D})$  carries the structure of a  $\mathcal{V}$ -algebra in  $\operatorname{Pr}^{\mathrm{L}}$ .

We use Corollary 1.7 to characterize the relative tensor product for  $\infty$ -categories presentably left tensored over  $\mathcal{V}$  as a  $\mathcal{V}$ -enriched localization of the  $\mathcal{V}$ -enriched  $\infty$ -category of  $\mathcal{V}$ -enriched presentable  $\infty$ categories: for any small regular cardinal  $\kappa$  a  $\mathcal{V}$ -enriched  $\infty$ -category  $\mathcal{M}$  is  $\mathcal{V}$ -enriched  $\kappa$ -presentable (Definition 5.11) if it admits small weighted colimits and every object of  $\mathcal{M}$  is a small  $\kappa$ -filtered conical colimit of  $\mathcal{V}$ -enriched  $\kappa$ -compact objects of  $\mathcal{M}$ , where an object X of  $\mathcal{M}$  is  $\mathcal{V}$ -enriched  $\kappa$ -compact if the morphism object functor  $\operatorname{Mor}_{\mathcal{M}}(X, -) : \mathcal{M} \to \mathcal{V}$  preserves small  $\kappa$ -filtered conical colimits. We characterize  $\mathcal{V}$ -enriched  $\kappa$ -presentable  $\infty$ -categories:

**Theorem 1.8.** (Theorem 5.12) Let  $\tau \leq \kappa$  be small regular cardinals,  $\mathcal{V}$  a  $\tau$ -compactly generated monoidal  $\infty$ -category and  $\mathcal{M}$  a  $\mathcal{V}$ -enriched  $\infty$ -category. The following conditions are equivalent:

- (1) The  $\mathcal{V}$ -enriched  $\infty$ -category  $\mathcal{M}$  is  $\mathcal{V}$ -enriched  $\kappa$ -presentable.
- (2) The  $\mathcal{V}$ -enriched  $\infty$ -category  $\mathcal{M}$  is a  $\kappa$ -compactly generated left tensored  $\infty$ -category.
- (3) There is a small  $\mathcal{V}$ -enriched  $\infty$ -category  $\mathcal{N}$  and a  $\kappa$ -accessible  $\mathcal{V}$ -enriched localization  $\mathcal{M} \neq \mathcal{P}_{\mathcal{V}}(\mathcal{N})$ .

We use Corollary 1.5 and Theorem 1.8 to deduce the following corollary:

**Corollary 1.9.** (Theorem 5.8) Let  $\mathcal{V}$  be a presentably symmetric monoidal  $\infty$ -category and  $\mathcal{M}, \mathcal{N}$  presentably left tensored  $\infty$ -categories. By Theorem 1.8 there are accessible  $\mathcal{V}$ -enriched localizations  $\mathcal{P}_{\mathcal{V}}(\mathcal{C}) \rightleftharpoons \mathcal{M}, \mathcal{P}_{\mathcal{V}}(\mathcal{D}) \rightleftharpoons \mathcal{N}$ . There is an accessible  $\mathcal{V}$ -enriched localization

$$\mathcal{P}_{\mathcal{V}}(\mathcal{C}\otimes\mathcal{D})\to\mathcal{M}\otimes_{\mathcal{V}}\mathcal{N}.$$

Blumberg-Gepner-Tabuada [4] develop algebraic K-theory of stable  $\infty$ -categories and characterize algebraic K-theory by an universal property. In analogy [19] characterize secondary algebraic K-theory by a universal property based on the notion of 2-stable ( $\infty$ , 2)-category: secondary algebraic K-theory is a categorification of algebraic K-theory that instead of the stable  $\infty$ -category of module spectra Mod<sub>R</sub> over a ring R uses the  $\infty$ -category Mod<sub>R</sub><sup>2</sup> of small (Cauchy-complete) stable  $\infty$ -categories left tensored over Mod<sub>R</sub>, thought of as secondary R-modules. Like Mod<sub>R</sub> is the prime example of a stable  $\infty$ -category,  $\operatorname{Mod}_{R}^{2}$  is the prime example of a 2-stable (∞, 2)-category. An ∞-category is stable if it admits a zero object, pushouts and pullbacks and a commutative square is a pushout square if and only if it is a pullback square. The (∞, 2)-category of small stable ∞-categories is a 2-stable (∞, 2)-category in the following sense: it is enriched in stable ∞-categories (via its canonical Cat<sub>∞</sub>enrichment) and it is a preadditive (∞, 2)-category: it admits a zero object in the enriched sense and conical finite coproducts and products and such agree via the canonical map. More generally, we inductively define (Cauchy-complete) n-stable (∞, n)-categories for every n ≥ 1 as those (Cauchycomplete) preadditive (∞, n)-categories that are enriched in (Cauchy-complete) n-1-stable (∞, n - 1)categories. We prove that the (∞, n + 1)-category Cat<sup>ex</sup><sub>(∞,n)</sub> of small stable (∞, n)-categories is n + 1stable and the (∞, n+1)-category Cat<sup>ex,∧</sup><sub>(∞,n)</sub> of small cauchy-complete stable (∞, n)-categories is Cauchy complete and n + 1-stable (Corollary 5.84). We use Theorem 1.1 to construct a tensor product for n-stable (∞, n)-categories generalizing the tensor product of stable ∞-categories:

**Corollary 1.10.** (Corollary 5.86) Let  $n \ge 0$ . There are canonical presentably symmetric monoidal structures on the  $\infty$ -categories

$$\mathsf{Cat}^{\mathrm{ex}}_{(\infty, n)}, \mathsf{Cat}^{\mathrm{ex}, \wedge}_{(\infty, n)}$$

1.1. Notation and terminology. We fix a hierarchy of Grothendieck universes whose objects we call small, large, very large, etc. We call a space small, large, etc. if its set of path components and its homotopy groups are for any choice of base point. We call an  $\infty$ -category small, large, etc. if its maximal subspace and all its mapping spaces are.

We write

- Set for the category of small sets.
- $\Delta$  for (a skeleton of) the category of finite, non-empty, partially ordered sets and order preserving maps, whose objects we denote by  $[n] = \{0 < ... < n\}$  for  $n \ge 0$ .
- S for the  $\infty$ -category of small spaces.
- $Cat_{\infty}$  for the  $\infty$ -category of small  $\infty$ -categories.
- $Cat_{\infty}^{cc}$  for the  $\infty$ -category of large  $\infty$ -categories with small colimits and small colimits preserving functors.

We often indicate  $\infty$ -categories of large objects by (-), for example we write  $\widehat{S}, \widehat{Cat}_{\infty}$  for the  $\infty$ -categories of large spaces,  $\infty$ -categories.

For any  $\infty$ -category  $\mathcal{C}$  containing objects A, B we write

- $\mathcal{C}(A, B)$  for the space of maps  $A \to B$  in  $\mathcal{C}$ .
- $\mathcal{C}_{/A}$  for the  $\infty$ -category of objects over A,
- Ho(C) for its homotopy category,
- $\mathbb{C}^{\triangleleft}, \mathbb{C}^{\triangleright}$  for the  $\infty$ -category arising from  $\mathbb{C}$  by adding an initial, final object, respectively,
- $\mathcal{C}^{\simeq}$  for the maximal subspace in  $\mathcal{C}$ .

Note that  $Ho(Cat_{\infty})$  is cartesian closed and for small  $\infty$ -categories  $\mathcal{C}, \mathcal{D}$  we write  $Fun(\mathcal{C}, \mathcal{D})$  for the internal hom, the  $\infty$ -category of functors  $\mathcal{C} \to \mathcal{D}$ .

We often call a fully faithful functor  $\mathcal{C} \to \mathcal{D}$  an embedding. We call a functor  $\phi : \mathcal{C} \to \mathcal{D}$  an inclusion (of a subcategory) if one of the following equivalent conditions holds:

- For any  $\infty$ -category  $\mathcal{B}$  the induced map  $\mathsf{Cat}_{\infty}(\mathcal{B}, \mathcal{C}) \to \mathsf{Cat}_{\infty}(\mathcal{B}, \mathcal{D})$  is an embedding.
- The functor  $\phi : \mathcal{C} \to \mathcal{D}$  induces an embedding on maximal subspaces and on all mapping spaces.

Let S be an  $\infty$ -category and  $\mathcal{E} \subset \operatorname{Fun}([1], S)$  a full subcategory. We call a functor  $X \to S$  a cocartesian fibration relative to  $\mathcal{E}$  if for every morphism  $[1] \to S$  that belongs to  $\mathcal{E}$  the pullback  $[1] \times_S X \to [1]$  is a

cocartesian fibration whose cocartesian morphisms are preserved by the projection  $[1] \times_S X \to X$ . We call a functor  $X \to Y$  over S a map of cocartesian fibrations relative to  $\mathcal{E}$  if it preserves cocartesian lifts of morphisms of  $\mathcal{E}$ . We write  $Cat^{\mathcal{E}}_{\infty/S} \subset Cat_{\infty/S}$  for the subcategory of cocartesian fibrations relative to  $\mathcal{E}$  and maps of such.

Acknowledgements. We thank David Gepner and Markus Spitzweck for helpful discussions.

#### 2. Enriched ∞-categories

In this section we recall the theory of enriched  $\infty$ -categories of [10] and further develope this theory for our needs.

2.1.  $\infty$ -operads. To define enriched  $\infty$ -categories we first need to introduce non-symmetric  $\infty$ -operads [6, Definition 2.2.6.], [10, Definition 2.16.].

Notation 2.1. Let Ass :=  $\Delta^{\text{op}}$  be the category of finite non-empty totally ordered sets and order preserving maps. We call a map  $[n] \rightarrow [m]$  in Ass

- inert if it corresponds to a map of  $\Delta$  of the form  $[m] \simeq \{i, i+1, ..., i+m\} \subset [n]$  for some  $i \ge 0$ .
- active if it corresponds to a map of  $\Delta$ , which preserves the minimum and maximum.

For every  $n \ge 0$  there are n inert morphisms  $[n] \rightarrow [1]$ , where the i-th inert morphism  $[n] \rightarrow [1]$  for  $1 \le i \le n$  corresponds to the map  $[1] \simeq \{i - 1, i\} \subset [n]$ .

**Definition 2.2.** A (non-symmetric)  $\infty$ -operad is a cocartesian fibration  $\phi : \mathcal{V}^{\otimes} \to \text{Ass relative to the collection of inert morphisms such that the following conditions hold, where we set <math>\mathcal{V} := \mathcal{V}_{[1]}^{\otimes}$ :

- (1) For every  $n \ge 0$  the family of n inert morphisms  $[n] \rightarrow [1]$  of Ass induces an equivalence  $\mathcal{V}_{[n]}^{\otimes} \rightarrow \mathcal{V}^{\times n}$ .
- (2) For every  $Y, X \in \mathcal{V}^{\otimes}$  lying over  $[m], [n] \in Ass$  the family  $X \to X_j$  for  $1 \le j \le n$  of  $\phi$ -cocartesian lifts of the inert morphisms  $[n] \to [1]$  induces a pullback square

**Definition 2.3.** A monoidal  $\infty$ -category is an  $\infty$ -operad  $\phi : \mathcal{V}^{\otimes} \to Ass$  such that  $\phi$  is a cocartesian fibration.

**Remark 2.4.** A cocartesian fibration  $\phi : \mathcal{V}^{\otimes} \to Ass$  is a monoidal  $\infty$ -category if and only if the first condition of Definition 2.2 holds. The second condition is automatic if  $\phi$  is a cocartesian fibration [10, Remark 2.17.].

**Notation 2.5.** For every  $\infty$ -operad  $\phi : \mathcal{V}^{\otimes} \to Ass$  we set  $\mathcal{V} := \mathcal{V}_{[1]}^{\otimes}$  and call  $\mathcal{V}$  the underlying  $\infty$ -category.

**Notation 2.6.** Let  $\mathcal{V}^{\otimes} \to Ass$  be an  $\infty$ -operad and  $V_1, \dots, V_n, W \in \mathcal{V}$  for  $n \ge 0$ . Let

$$Mul_{\mathcal{V}}(V_1, ..., V_n; W)$$

be the full subspace of  $\mathcal{V}^{\otimes}(V, W)$  spanned by the morphisms  $V \to W$  in  $\mathcal{V}^{\otimes}$  lying over the active morphism  $[1] \to [n]$  in  $\Delta$ , where  $V \in \mathcal{V}^{\otimes}_{[n]} \simeq \mathcal{V}^{\times n}$  corresponds to  $(V_1, ..., V_n)$ .

- **Definition 2.7.** (1) An  $\infty$ -operad  $\phi : \mathcal{V}^{\otimes} \to Ass$  is locally small if the mapping spaces of  $\mathcal{V}^{\otimes}$  are small.
  - (2) An  $\infty$ -operad  $\phi: \mathcal{V}^{\otimes} \to Ass$  is small if it is locally small and  $\mathcal{V}$  is small.

**Remark 2.8.** By the first axiom of Definition 2.2 an  $\infty$ -operad  $\mathcal{V}^{\otimes} \to Ass$  is small if and only if  $\mathcal{V}^{\otimes}$  is small. An  $\infty$ -operad  $\mathcal{V}^{\otimes} \to Ass$  is locally small if and only if the multi-morphism spaces of  $\mathcal{V}^{\otimes} \to Ass$  are small. So an  $\infty$ -operad  $\mathcal{V}^{\otimes} \to Ass$  is small if and only if  $\mathcal{V}$  is small and the multi-morphism spaces of  $\mathcal{V}^{\otimes} \to Ass$  are small.

**Notation 2.9.** For every monoidal  $\infty$ -category  $\mathcal{V}^{\otimes} \to Ass$  we write  $\otimes : \mathcal{V} \times \mathcal{V} \simeq \mathcal{V}_{[2]}^{\otimes} \to \mathcal{V}_{[1]}^{\otimes} = \mathcal{V}$  for the functor induced by the active map  $[1] \to [2]$  in  $\Delta$ .

**Definition 2.10.** Let  $\mathcal{K} \subset \mathsf{Cat}_{\infty}$  be a full subcategory. An  $\infty$ -operad  $\mathcal{V}^{\otimes} \to \mathrm{Ass}$  is compatible with  $\mathcal{K}$ -indexed colimits if  $\mathcal{V}$  admits  $\mathcal{K}$ -indexed colimits and for every  $V_1, ..., V_n, V \in \mathcal{V}$  for  $n \ge 0$  and  $0 \le i \le n$  the presheaf  $\mathrm{Mul}_{\mathcal{V}}(V_1, ..., V_{i-1}, ..., V_n; \mathcal{V})$  on  $\mathcal{V}$  preserves  $\mathcal{K}$ -indexed limits.

**Remark 2.11.** A monoidal  $\infty$ -category  $\mathcal{V}^{\otimes} \to Ass$  is compatible with  $\mathcal{K}$ -indexed colimits if it is compatible with  $\mathcal{K}$ -indexed colimits as an  $\infty$ -operad, i.e.  $\mathcal{V}$  admits  $\mathcal{K}$ -indexed colimits and such are preserved by the tensor product component-wise.

**Definition 2.12.** A monoidal  $\infty$ -category  $\mathcal{V}^{\otimes} \to Ass$  is

- (1) closed if the tensor product  $\otimes : \mathcal{V} \times \mathcal{V} \to \mathcal{V}$  is left adjoint component-wise.
- (2) presentably if  $\mathcal{V}$  is presentable and (1) holds.
- (3)  $\kappa$ -compactly generated for a regular cardinal  $\kappa$  if it is compatible with small colimits,  $\mathcal{V}$  is  $\kappa$ compactly generated and the monoidal structure of  $\mathcal{V}$  restricts to the full subcategory  $\mathcal{V}^{\kappa} \subset \mathcal{V}$ of  $\kappa$ -compact objects.

**Remark 2.13.** By [8, Proposition 7.15.] every presentably monoidal  $\infty$ -category is  $\kappa$ -compactly generated for some regular cardinal  $\kappa$ .

We also consider maps of  $\infty$ -operads:

**Definition 2.14.** Let  $\mathcal{V}^{\otimes} \to Ass, \mathcal{W}^{\otimes} \to Ass$  be  $\infty$ -operads. A map of  $\infty$ -operads  $\mathcal{V}^{\otimes} \to \mathcal{W}^{\otimes}$  is a map of cocartesian fibrations  $\mathcal{V}^{\otimes} \to \mathcal{W}^{\otimes}$  relative to the collection of inert morphisms of Ass.

**Definition 2.15.** A map of  $\infty$ -operade  $\mathcal{V}^{\otimes} \to \mathcal{W}^{\otimes}$  is an embedding if it is fully faithful.

**Definition 2.16.** Let  $\mathcal{V}^{\otimes} \to \mathbb{E}_k, \mathcal{W}^{\otimes} \to Ass$  be monoidal  $\infty$ -categories.

- A lax monoidal functor  $\mathcal{V}^{\otimes} \to \mathcal{W}^{\otimes}$  is a map of  $\infty$ -operads  $\mathcal{V}^{\otimes} \to \mathcal{W}^{\otimes}$ .
- A monoidal functor  $\mathcal{V}^{\otimes} \to \mathcal{W}^{\otimes}$  is a map of cocartesian fibrations  $\mathcal{V}^{\otimes} \to \mathcal{W}^{\otimes}$  over Ass.

Notation 2.17. We fix the following notation:

- Let  $\operatorname{Op}_{\infty} \subset \mathsf{Cat}_{\infty/Ass}$  be the subcategory of  $\infty$ -operads and maps of  $\infty$ -operads.
- Let  $\mathrm{Mon}\subset\mathrm{Op}_\infty$  be the subcategory of monoidal  $\infty\text{-}\mathrm{categories}$  and monoidal functors.
- Let PrMon ⊂ Mon be the subcategory of monoidal ∞-categories compatible with small colimits and monoidal functors preserving small colimits.

Notation 2.18. For every  $\infty$ -operads  $\mathcal{V}^{\otimes} \to Ass, \mathcal{W}^{\otimes} \to Ass$  let

$$\operatorname{Alg}_{\mathcal{V}}(\mathcal{W}) \subset \operatorname{Fun}_{\operatorname{Ass}}(\mathcal{V}^{\otimes}, \mathcal{W}^{\otimes})$$

be the full subcategory of maps of  $\infty$ -operads. If  $\mathcal{V}^{\otimes} \to Ass, \mathcal{W}^{\otimes} \to Ass$  are monoidal  $\infty$ -categories, let

$$\operatorname{Fun}^{\otimes, L}(\mathcal{V}, \mathcal{W}) \subset \operatorname{Fun}^{\otimes}(\mathcal{V}, \mathcal{W}) \subset \operatorname{Alg}_{\mathcal{V}}(\mathcal{W})$$

be the full subcategories of monoidal functors (that induce on underlying  $\infty$ -categories a left adjoint).

**Definition 2.19.** An  $\infty$ -operad  $\mathcal{V}^{\otimes} \to Ass$  admits a tensor unit if the functor  $\operatorname{Mul}_{\mathcal{V}}(\emptyset; -) : \mathcal{V} \to S$  is corepresentable. We call the corepresenting object the tensor unit of  $\mathcal{V}$  and write  $\mathbb{1}_{\mathcal{V}}$  for it.

**Remark 2.20.** Let  $\phi : \mathcal{V}^{\otimes} \to Ass$  be an  $\infty$ -operad that admits a tensor unit and for every  $n \ge 1, 1 \le j \le n$ let  $(V_i^j)_{i=1}^{m_j}$  be a family in  $\mathcal{V}^{\otimes}$ , where  $m_j \ge 0$ . By the axioms of an  $\infty$ -operad there is a morphism

$$(V_{i}^{j})_{i=1}^{m_{1}}, \emptyset, (V_{i}^{j})_{i=1}^{m_{2}}, \emptyset, (V_{i}^{j})_{i=1}^{m_{3}}, ..., \emptyset, (V_{i}^{j})_{i=1}^{m_{n}} \rightarrow (V_{i}^{j})_{i=1}^{m_{1}}, \mathbb{1}_{\mathcal{V}}, (V_{i}^{j})_{i=1}^{m_{2}}, \mathbb{1}_{\mathcal{V}}, (V_{i}^{j})_{i=1}^{m_{3}}, ..., (V_{i}^{j})_{i=1}^{m_{3}}, ..., (V_{i}^{j})_{i=1}^{m_{$$

in  $\mathcal{V}^{\otimes}$ , which is  $\phi$ -cocartesian by [10, Lemma 2.24.] since the morphism  $\emptyset \to \mathbb{1}$  is  $\phi$ -cocartesian.

**Lemma 2.21.** Let  $\phi: \mathcal{V}^{\otimes} \to Ass$  be an  $\infty$ -operad that admits a tensor unit. The  $\infty$ -category  $Alg(\mathcal{V})$ admits an initial object. An associative algebra in  $\mathcal{V}$  is initial if and only if it preserves the tensor unit as a map Ass  $\rightarrow \mathcal{V}^{\otimes}$ .

*Proof.* A map  $A \to B$  in Alg( $\mathcal{V}$ ) is an equivalence if A, B preserve the tensor unit since the induced map  $A([1]) \to B([1])$  in  $\mathcal{V}^{\otimes}$  is a map under  $\emptyset \simeq A([0]) \simeq B([0])$ . Thus it is enough to prove that the functor  $\rho : \operatorname{Alg}(\mathcal{V}) \to \mathcal{V}_{[0]}^{\otimes}$  restricting along the embedding  $[0] \subset \operatorname{Ass}$  admits a fully faithful left adjoint that sends Ø to an algebra preserving the tensor unit. By [17, Lemma 4.3.2.13., Proposition 4.3.2.17.] the functor  $\rho$  has a fully faithful left adjoint if for any  $n \ge 0$  the functor  $\lambda : \{[0]\} \times_{Ass} Ass_{/[n]} \to \mathcal{V}_{[0]}^{\otimes} \subset \mathcal{V}^{\otimes}$ admits a  $\phi$ -colimit and the left adjoint sends  $\emptyset \in \mathcal{V}_{[0]}^{\otimes}$  to the algebra Z in  $\mathcal{V}$  that maps  $[n] \in Ass$  to the  $\phi$ -colimit of  $\lambda$ . Since  $\{[0]\} \times_{Ass} Ass_{/[n]}$  is contractible, a  $\phi$ -colimit of  $\lambda$  is a  $\phi$ -cocartesian lift of the map  $[0] \rightarrow [n]$  in Ass, which exists by Remark 2.20. The algebra Z in  $\mathcal{V}$  sends  $[1] \in Ass$  to the  $\phi$ -cocartesian lift  $\emptyset \to \mathbb{1}_{\mathcal{V}}$  of the map  $[0] \to [1]$  in Ass. So Z preserves the tensor unit.

Via Lemma 2.21 we can make the following definition:

**Notation 2.22.** Let  $n \ge 1$ ,  $1 \le i \le n$  and  $\mathcal{V}_i^{\otimes} \to Ass$  be  $\infty$ -operads such that for every  $1 \le j \le n, j \ne i$  the  $\infty$ -operad  $\mathcal{V}_j^{\otimes} \to Ass$  admits a tensor unit. Let  $\tau_i : \mathcal{V}_i^{\otimes} \to \mathcal{V}_1^{\otimes} \times_{Ass} \dots \times_{Ass} \mathcal{V}_n^{\otimes}$  be the map of  $\infty$ -operads that is the identity on the i-th factor and the constant map  $\mathcal{V}_i^{\otimes} \to Ass \to \mathcal{V}_j^{\otimes}$  preserving the tensor unit on the j-th factor for  $j \neq i$ , where we use Lemma 2.21.

Notation 2.23. Let  $\kappa$  be a small regular cardinal and  $\mathcal{C}$  a small  $\infty$ -category. Let  $\operatorname{Ind}_{\kappa}(\mathcal{C}) \subset \mathcal{P}(\mathcal{C})$  be the full subcategory generated by  $\mathcal{C}$  under small  $\kappa$ -filtered colimits.

**Example 2.24.** For  $\kappa = \emptyset$  we find that  $\operatorname{Ind}_{\emptyset}(\mathcal{C}) = \mathcal{P}(\mathcal{C})$ .

**Remark 2.25.** By [16, Corollary 5.3.5.4.] for every small  $\infty$ -category C that admits  $\kappa$ -small colimits the full subcategory  $\operatorname{Ind}_{\kappa}(\mathcal{C}) \subset \mathcal{P}(\mathcal{C})$  precisely consists of the functors  $\mathcal{C}^{\operatorname{op}} \to \mathcal{S}$  preserving  $\kappa$ -small limits. Thus  $\operatorname{Ind}_{\kappa}(\mathcal{C})$  is a  $\kappa$ -accessible localization with respect to the set of maps  $\operatorname{\{colim}(y \circ H) \rightarrow \operatorname{Ind}_{\kappa}(\mathcal{C})$  $y(\text{colim}(H)) | H : K \to \mathcal{C}, K \kappa\text{-small}\}$ , where  $y : \mathcal{C} \subset \mathcal{P}(\mathcal{C})$  is the Yoneda-embedding. Hence  $\text{Ind}_{\kappa}(\mathcal{C})$  is a presentable  $\infty$ -category.

The next proposition follows from [10, Corollary 8.31.]:

**Proposition 2.26.** Let  $\kappa$  be a small regular cardinal and  $\mathcal{V}^{\otimes} \to Ass$  a small monoidal  $\infty$ -category.

- (1) There is a monoidal  $\infty$ -category  $\operatorname{Ind}_{\kappa}(\mathcal{V})^{\otimes} \to \operatorname{Ass}$  compatible with small  $\kappa$ -filtered colimits and a monoidal embedding  $\mathcal{V}^{\otimes} \to \operatorname{Ind}_{\kappa}(\mathcal{V})^{\otimes}$  inducing the embedding  $\mathcal{V} \to \operatorname{Ind}_{\kappa}(\mathcal{V})$ . (2) For every monoidal  $\infty$ -category  $\mathcal{W}^{\otimes} \to \operatorname{Ass}$  compatible with small  $\kappa$ -filtered colimits the functor

 $\operatorname{Alg}_{\operatorname{Ind}_{\kappa}(\mathcal{V})}(\mathcal{W}) \to \operatorname{Alg}_{\mathcal{V}}(\mathcal{W})$ 

admits a fully faithful left adjoint that lands in the full subcategory  $\operatorname{Alg}_{\operatorname{Ind}_{\kappa}(\mathcal{V})}^{\kappa-\operatorname{fil}}(\mathcal{W})$  of maps of  $\infty$ -operads preserving small  $\kappa$ -filtered colimits. Thus the following functor is an equivalence:

$$\operatorname{Alg}_{\operatorname{Ind}_{\kappa}(\mathcal{V})}^{\kappa-\operatorname{nl}}(\mathcal{W}) \to \operatorname{Alg}_{\mathcal{V}}(\mathcal{W})$$

(3) If  $\mathcal{V}$  admits  $\kappa$ -small colimits, the latter equivalence restricts to an equivalence

$$\operatorname{Alg}_{\operatorname{Ind}_{\kappa}(\mathcal{V})}^{\operatorname{L}}(\mathcal{W}) \to \operatorname{Alg}_{\mathcal{V}}^{\kappa}(\mathcal{W}).$$

In the later sections we will also use symmetric  $\infty$ -operads.

Notation 2.27. We write Comm for the category of finite pointed sets. We write finite pointed sets as  $(n) := \{*, 1, ..., n\}$  for  $n \ge 0$ , where \* is the base point.

Notation 2.28. We call a map  $\theta$  of pointed finite sets  $(n) \rightarrow (m)$ 

- inert if for every  $1 \le i \le m$  the fiber of  $\theta$  over i consists precisely of one element.
- active if it sends only the base point to the base point.

**Remark 2.29.** For every  $n \ge 0$  there are n inert morphisms  $(n) \rightarrow (1)$ , where the i-th inert morphism  $(n) \rightarrow (1)$  for  $1 \le i \le n$  sends i to 1.

**Definition 2.30.** A symmetric  $\infty$ -operad is a cocartesian fibration  $\mathcal{V}^{\boxtimes} \to \text{Comm}$  relative to the collection of inert morphisms that satisfies the analogous conditions like in Definition 2.2, where [n], [m] are replaced by  $\langle n \rangle, \langle m \rangle$ . A symmetric monoidal  $\infty$ -category is a symmetric  $\infty$ -operad  $\mathcal{V}^{\boxtimes} \to \text{Comm}$  that is also a cocartesian fibration.

**Remark 2.31.** A cocartesian fibration  $\mathcal{V}^{\boxtimes} \to \text{Comm}$  is a symmetric monoidal  $\infty$ -category if and only if it satisfies the analogue of condition (1) of Definition 2.2, condition (2) is automatic for cocartesian fibrations like in Remark 2.4.

**Notation 2.32.** By [16, Construction 4.1.2.9.] there is a canonical functor  $\theta$ : Ass  $\rightarrow$  Comm,  $[n] \mapsto \langle n \rangle$  and we write  $\mathcal{V}^{\otimes} \rightarrow$  Ass for the pullback of a symmetric monoidal  $\infty$ -category  $\mathcal{V}^{\otimes} \rightarrow$  Comm along  $\theta$ , which is a monoidal  $\infty$ -category that we call the underlying monoidal  $\infty$ -category. We define maps of symmetric  $\infty$ -operads and (lax) symmetric monoidal functors analogously as in Notation 2.18.

**Example 2.33.** By [16, Proposition 5.1.0.3.] for every  $n \ge 0$  there is a symmetric  $\infty$ -operad  $\mathbb{E}_n \rightarrow \text{Comm}$ , the n-th little disk operad. For n = 1 the  $\mathbb{E}_1$ -operad is the symmetrization of the non-symmetric  $\infty$ -operad Ass. For  $n = \infty$  there is an equivalence between Comm-algebras and Comm-algebras [16, Theorem 5.1.2.2].  $\mathbb{E}_0$ -algebras are precisely objects equipped with a map from the tensor unit.

2.2. Weakly enriched  $\infty$ -categories. To define enriched  $\infty$ -categories we first introduce weakly enriched  $\infty$ -categories following [10], which there were called weakly tensored  $\infty$ -categories.

**Definition 2.34.** A morphism in Ass preserves the minimum (maximum) if it corresponds to a map  $[m] \rightarrow [n]$  in  $\Delta$  sending 0 to 0 (sending m to n).

The next definition is [10, Definition 3.3.]:

**Definition 2.35.** Let  $\mathcal{V}^{\otimes} \to \operatorname{Ass}, \mathcal{W}^{\otimes} \to \operatorname{Ass}$  be  $\infty$ -operads. An  $\infty$ -category weakly bi-enriched in  $\mathcal{V}, \mathcal{W}$  is a map  $\phi = (\phi_1, \phi_2) : \mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  of cocartesian fibrations relative to the collection of inert morphisms of Ass × Ass whose first component preserves the maximum and whose second component preserves the minimum such that the following conditions hold:

(1) for every  $n, m \ge 0$  the map  $[0] \simeq \{n\} \subset [n]$  in the first component and the map  $[0] \simeq \{0\} \subset [m]$  in the second component induce an equivalence

$$\theta: \mathcal{M}^{\circledast}_{[n][m]} \to \mathcal{V}^{\otimes}_{[n]} \times \mathcal{M}^{\circledast}_{[0][0]} \times \mathcal{W}^{\otimes}_{[m]},$$

(2) for every X, Y  $\in \mathcal{M}^{\circledast}$  lying over ([m'], [n']), ([m], [n])  $\in Ass \times Ass$  the cocartesian lift Y  $\rightarrow$  Y' of the map  $[0] \simeq \{m\} \subset [m]$  and  $[0] \simeq \{0\} \subset [n]$  induces a pullback square

**Notation 2.36.** For every weakly bi-enriched  $\infty$ -category  $\phi : \mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  we call  $\mathcal{M} := \mathcal{M}_{[0][0]}^{\circledast}$  the underlying  $\infty$ -category of  $\phi$  and say that  $\phi$  exhibits  $\mathcal{M}$  as weakly bi-enriched in  $\mathcal{V}, \mathcal{W}$ .

**Example 2.37.** Let  $\mathcal{V}^{\otimes} \to Ass$  be an  $\infty$ -operad. We write  $\mathcal{V}^{\otimes} \to Ass \times Ass$  for the pullback of  $\mathcal{V}^{\otimes} \to Ass$  along the functor  $Ass \times Ass \to Ass$ ,  $([n], [m]) \mapsto [n] * [m]$ . The two functors  $Ass \times Ass \times [1] \to Ass$  corresponding to the natural transformations  $(-) * \emptyset \to (-) * (-)$ ,  $\emptyset * (-) \to (-) * (-)$  send the morphism  $\mathrm{id}_{[n], [m]}, 0 \to 1$  to an inert one and so give rise to functors  $\mathcal{V}^{\otimes} \to \mathcal{V}^{\otimes} \times Ass$ ,  $\mathcal{V}^{\otimes} \to Ass \times \mathcal{V}^{\otimes}$  over  $Ass \times Ass$ . The resulting functor  $\mathcal{V}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{V}^{\otimes}$  is an  $\infty$ -category weakly bi-enriched in  $\mathcal{V}, \mathcal{V}$ .

**Example 2.38.** Let  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  be a weakly bi-enriched  $\infty$ -category and  $\mathcal{N} \subset \mathcal{M}$  a full subcategory. Let  $\mathcal{N}^{\circledast} \subset \mathcal{M}^{\circledast}$  be the full subcategory spanned by all objects of  $\mathcal{M}^{\circledast}$  lying over some  $(V, W) \in \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  corresponding some object of  $\mathcal{N} \subset \mathcal{M}$ . The restriction  $\mathcal{N}^{\circledast} \subset \mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  is a weakly bi-enriched  $\infty$ -category, whose underlying  $\infty$ -category is  $\mathcal{N}$ . We call  $\mathcal{N}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  the full weakly bi-enriched subcategory of  $\mathcal{M}^{\circledast}$  spanned by  $\mathcal{N}$ .

**Notation 2.39.** Let  $\emptyset^{\otimes} \subset Ass$  be the full subcategory spanned by  $[0] \in Ass$ . Then  $\emptyset^{\otimes}$  is contractible and  $\emptyset^{\otimes} \subset Ass$  is an  $\infty$ -operad that is the initial  $\infty$ -operad.

**Definition 2.40.** An  $\infty$ -category weakly left enriched in  $\mathcal{V}$  is an  $\infty$ -category weakly bi-enriched in  $\mathcal{V}, \emptyset$ . An  $\infty$ -category weakly right enriched in  $\mathcal{V}$  is an  $\infty$ -category weakly bi-enriched in  $\emptyset, \mathcal{V}$ .

**Notation 2.41.** (Multi-morphism spaces) Let  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  be a weakly bi-enriched  $\infty$ -category and  $V_1, ..., V_n \in \mathcal{V}, X, Y \in \mathcal{M}, W_1, ..., W_m \in \mathcal{W}$  for some  $n, m \ge 0$ . Let

 $Mul_{\mathcal{M}}(V_1, ..., V_n, X, W_1, ..., W_m; Y)$ 

be the full subspace of  $\mathcal{M}^{\circledast}(Z, Y)$  spanned by the morphisms  $Z \to Y$  in  $\mathcal{M}^{\circledast}$  lying over the map  $[0] \simeq \{0\} \subset [n], [0] \simeq \{m\} \subset [m]$  in  $\Delta \times \Delta$ , where  $Z \in \mathcal{M}^{\circledast}_{[n],[m]} \simeq \mathcal{V}^{\times n} \times \mathcal{M} \times \mathcal{W}^{\times m}$  corresponds to  $(V_1, ..., V_n, X, W_1, ..., W_m)$ .

**Remark 2.42.** For every  $\infty$ -operad  $\mathcal{V}^{\otimes} \to Ass$  the projection  $\mathcal{M}^{\otimes} := \mathcal{V}^{\otimes} \to \mathcal{V}^{\otimes}$  gives rise to an embedding  $\mathcal{V}^{\otimes}(Z, Y) \to \mathcal{V}^{\otimes}(Z, Y)$  that restricts to an equivalence

$$Mul_{\mathcal{M}}(V_1, ..., V_n, X, W_1, ..., W_m; Y) \simeq Mul_{\mathcal{V}}(V_1, ..., V_n, X, W_1, ..., W_m; Y).$$

**Definition 2.43.** Let  $\phi : \mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be a weakly bi-enriched  $\infty$ -category.

- (1) We call  $\phi$  locally small if the  $\infty$ -categories  $\mathcal{V}^{\otimes}, \mathcal{W}^{\otimes}, \mathcal{M}^{\otimes}$  are locally small.
- (2) We call  $\phi$  small if  $\phi$  is locally small and  $\mathcal{M}$  is small.
- (3) We call  $\phi$  totally small if  $\phi$  is small and  $\mathcal{V}, \mathcal{W}$  are small.

**Remark 2.44.** Let  $\phi : \mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  be a weakly bi-enriched  $\infty$ -category. The first axiom of Definition 2.35 implies that  $\phi$  is totally small if and only if  $\mathcal{M}^{\circledast}, \mathcal{V}^{\circledast}, \mathcal{W}^{\circledast}$  are small. Remark 2.8 and the axioms of Definition 2.35 imply that  $\phi$  is locally small if and only if the multi-morphism spaces (Notation 2.41) of  $\mathcal{V}^{\circledast} \to \operatorname{Ass}, \mathcal{W}^{\circledast} \to \operatorname{Ass}, \phi$  are small. So  $\phi$  is small if and only if  $\mathcal{M}$  is small and the multi-morphism spaces of  $\mathcal{V}^{\circledast} \to \operatorname{Ass}, \mathcal{W}^{\circledast} \to \operatorname{Ass}, \phi$  are small.

Next we define tensored  $\infty$ -categories.

**Definition 2.45.** Let  $\phi : \mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be a weakly bi-enriched  $\infty$ -category.

- (1) We say that  $\phi : \mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  exhibits  $\mathcal{M}$  as left tensored over  $\mathcal{V}$  if  $\mathcal{V}^{\otimes} \to Ass$  is a monoidal  $\infty$ -category and  $\phi$  is a map of cocartesian fibrations over Ass via projection to the first factor.
- (2) We say that  $\phi : \mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  exhibits  $\mathcal{M}$  as right tensored over  $\mathcal{W}$  if  $\mathcal{W}^{\otimes} \to Ass$  is a monoidal  $\infty$ -category and  $\phi$  is a map of cocartesian fibrations over Ass via projection to the second factor.
- (3) We say that  $\phi : \mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  exhibits  $\mathcal{M}$  as bitensored over  $\mathcal{V}, \mathcal{W}$  if  $\phi$  exhibits  $\mathcal{M}$  as left tensored over  $\mathcal{V}$  and right tensored over  $\mathcal{W}$ .

**Remark 2.46.** Let  $\mathcal{V}^{\otimes} \to Ass, \mathcal{W}^{\otimes} \to Ass$  be monoidal  $\infty$ -categories and  $\phi : \mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a map of cocartesian fibrations over  $Ass \times Ass$ . Then  $\phi : \mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  exhibits  $\mathcal{M}$  as bitensored over  $\mathcal{V}, \mathcal{W}$  if and only if condition (1) of Definition 2.35 holds. Condition (2) is then automatic.

**Example 2.47.** Let  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast}$  be a (weakly) left tensored  $\infty$ -category and  $\mathcal{N}^{\circledast} \to \mathcal{W}^{\circledast}$  a (weakly) right tensored  $\infty$ -category. The functor  $\mathcal{M}^{\circledast} \times \mathcal{N}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  is a (weakly) bitensored  $\infty$ -category.

**Definition 2.48.** Let  $\kappa$  be a small regular cardinal.

- (1) A left tensored  $\infty$ -category  $\phi : \mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\circledast}$  is compatible with  $\kappa$ -small colimits if  $\mathcal{V}^{\otimes} \to Ass$  is compatible with  $\kappa$ -small colimits,  $\mathcal{M}$  admits  $\kappa$ -small colimits, for every  $V \in \mathcal{V}, X \in \mathcal{M}$  the functors  $(-) \otimes X : \mathcal{V} \to \mathcal{M}, V \otimes (-) : \mathcal{M} \to \mathcal{M}$  preserve  $\kappa$ -small colimits and for every  $W_1, ..., W_m \in \mathcal{W}$  for  $m \ge 0$  and  $Y \in \mathcal{M}$  the functor  $Mul_{\mathcal{M}}(-, W_1, ..., W_m; Y) : \mathcal{M}^{op} \to S$  preserves  $\kappa$ -small limits.
- (2) A right tensored  $\infty$ -category  $\phi : \mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  is compatible with  $\kappa$ -small colimits if  $\mathcal{W}^{\circledast} \to Ass$  is compatible with  $\kappa$ -small colimits,  $\mathcal{M}$  admits  $\kappa$ -small colimits, for every  $W \in \mathcal{W}, X \in \mathcal{M}$  the functors  $X \otimes (-) : \mathcal{W} \to \mathcal{M}, (-) \otimes W : \mathcal{M} \to \mathcal{M}$  preserve  $\kappa$ -small colimits and for every  $V_1, ..., V_n \in \mathcal{V}$  for  $n \ge 0$  and  $Y \in \mathcal{M}$  the functor  $Mul_{\mathcal{M}}(V_1, ..., V_n, -; Y) : \mathcal{M}^{op} \to \mathcal{S}$  preserves  $\kappa$ -small limits.
- (3) A bitensored  $\infty$ -category is compatible with  $\kappa$ -small colimits if it is a left and right tensored  $\infty$ -category compatible with  $\kappa$ -small colimits, i.e.  $\mathcal{V}^{\otimes} \to \operatorname{Ass}, \mathcal{W}^{\otimes} \to \operatorname{Ass}$  are compatible with  $\kappa$ -small colimits,  $\mathcal{M}$  admits  $\kappa$ -small colimits and for every  $V \in \mathcal{V}, W \in \mathcal{W}, X \in \mathcal{M}$  the functors  $(-) \otimes X : \mathcal{V} \to \mathcal{M}, X \otimes (-) : \mathcal{W} \to \mathcal{M}, V \otimes (-), (-) \otimes W : \mathcal{M} \to \mathcal{M}$  preserve  $\kappa$ -small colimits.

**Definition 2.49.** Let  $\kappa$  be a small regular cardinal.

- (1) A presentably left tensored  $\infty$ -category is a left tensored  $\infty$ -category  $\phi : \mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  compatible with small colimits such that  $\mathcal{V}, \mathcal{M}$  are presentable.
- (2) A presentably right tensored  $\infty$ -category is a right tensored  $\infty$ -category  $\phi : \mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  compatible with small colimits such that  $\mathcal{M}, \mathcal{W}$  are presentable.
- (3) A presentably bitensored  $\infty$ -category is a presentably left tensored and presentably right tensored  $\infty$ -category.
- (4) A left tensored  $\infty$ -category  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  is  $\kappa$ -compactly generated if  $\mathcal{V}^{\otimes} \to Ass$  and  $\mathcal{M}$  are  $\kappa$ -compactly generated and the left  $\mathcal{V}$ -action on  $\mathcal{M}$  restricts to a left  $\mathcal{V}^{\kappa}$ -action on  $\mathcal{M}^{\kappa}$ .
- (5) A right tensored  $\infty$ -category  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  is  $\kappa$ -compactly generated if  $\mathcal{W}^{\otimes} \to Ass$  and  $\mathcal{M}$  are  $\kappa$ -compactly generated and the right  $\mathcal{W}$ -action on  $\mathcal{M}$  restricts to a right  $\mathcal{W}^{\kappa}$ -action on  $\mathcal{M}^{\kappa}$ .
- (6) A bitensored  $\infty$ -category  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  is  $\kappa$ -compactly generated if the left  $\mathcal{V}$ -action and right  $\mathcal{W}$ -action are  $\kappa$ -compactly generated.

**Remark 2.50.** By [8, Proposition 7.15.] every presentably left tensored, right tensored, bitensored  $\infty$ -category, respectively, is  $\kappa$ -compactly generated for some regular cardinal  $\kappa$ .

In the following we generalize the notions of left, right and bitensored  $\infty$ -categories by introducing the notions of tensors.

**Definition 2.51.** Let  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be a weakly bi-enriched  $\infty$ -category and  $V \in \mathcal{V}, W \in \mathcal{W}, X, Y \in \mathcal{M}$ .

- (1) A multi-morphism  $\psi \in Mul_{\mathcal{M}}(V, X; Y)$  exhibits Y as the left tensor of V, X, denoted by V $\otimes$ X, if for every  $Z \in \mathcal{M}, V_1, ..., V_n \in \mathcal{V}, W_1, ..., W_m \in \mathcal{W}$  for  $n, m \ge 0$  the following map is an equivalence:  $Mul_{\mathcal{M}}(V_1, ..., V_n, Y, W_1, ..., W_m; Z) \rightarrow Mul_{\mathcal{M}}(V_1, ..., V_n, V, X, W_1, ..., W_m; Z).$
- (2) A multi-morphism  $\psi \in Mul_{\mathcal{M}}(X, W; Y)$  exhibits Y as the right tensor of X, W, denoted by  $X \otimes W$ , if for any  $Z \in \mathcal{M}, V_1, ..., V_n \in \mathcal{V}, W_1, ..., W_m \in \mathcal{W}$  for  $n, m \ge 0$  the map is an equivalence:  $Mul_{\mathcal{M}}(V_1, ..., V_n, Y, W_1, ..., W_m; Z) \rightarrow Mul_{\mathcal{M}}(V_1, ..., V_n, X, W, W_1, ..., W_m; Z).$

(3) A multi-morphism  $\tau \in Mul_{\mathcal{M}}(V, Y; X)$  exhibits Y as the left cotensor of V, X, denoted by <sup>V</sup>X, if for any  $Z \in \mathcal{M}, V_1, ..., V_n \in \mathcal{V}, W_1, ..., W_m \in \mathcal{W}$  for  $n, m \ge 0$  the following map is an equivalence:

 $Mul_{\mathcal{M}}(V_1,...,V_n,Z,W_1,...,W_m;Y) \rightarrow Mul_{\mathcal{M}}(V,V_1,...,V_n,Z,W_1,...,W_m;X).$ 

(4) A multi-morphism  $\tau \in Mul_{\mathcal{M}}(Y, W; X)$  exhibits Y as the right cotensor of X, W, denoted by  $X^{W}$ , if for every  $Z \in \mathcal{M}, V_{1}, ..., V_{n} \in \mathcal{V}, W_{1}, ..., W_{m} \in \mathcal{W}$  for  $n, m \ge 0$  the map is an equivalence:

 $\operatorname{Mul}_{\mathcal{M}}(V_1,...,V_n,Z,W_1,...,W_m;Y) \to \operatorname{Mul}_{\mathcal{M}}(V_1,...,V_n,Z,W_1,...,W_m,W;X).$ 

**Remark 2.52.** Let  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  be a weakly bi-enriched  $\infty$ -category,  $V \in \mathcal{V}, W \in \mathcal{W}$  and  $X \in \mathcal{M}$ . If the respective left and right tensors exist, there is a canonical equivalence  $(V \otimes X) \otimes W \simeq V \otimes (X \otimes W)$ . In this case we refer to  $(V \otimes X) \otimes W \simeq V \otimes (X \otimes W)$  as the bitensor of V, X, W. Similarly, if the respective left and right cotensors exist, there is a canonical equivalence  $(X^V)^W \simeq (X^W)^V$  and we refer to the latter object as the bicotensor of V, X, W.

**Definition 2.53.** A weakly bi-enriched  $\infty$ -category  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  admits

- left (co)tensors if for every object  $V \in \mathcal{V}, X \in \mathcal{M}$  there is a left (co)tensor of V and X.
- right (co)tensors if for every object  $W \in W, X \in M$  there is a right (co)tensor of X and W.

**Example 2.54.** Every weakly bi-enriched  $\infty$ -category that is left, right, bitensored admits left tensors, right tensors, left and right tensors, respectively.

**Remark 2.55.** Let  $\phi : \mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be a weakly bi-enriched  $\infty$ -category that admits left and right tensors. Then  $\phi$  is a locally cocartesian fibration.

2.3. Enriched functors. Next we define maps of weakly bi-enriched  $\infty$ -categories. The next definition is [10, Definition 3.18., Definition 3.19.]:

**Definition 2.56.** Let  $\alpha : \mathcal{V}^{\otimes} \to \mathcal{V}'^{\otimes}, \beta : \mathcal{W}^{\otimes} \to \mathcal{W}'^{\otimes}$  be maps of  $\infty$ -operads and  $\phi : \mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}, \phi' : \mathcal{M}'^{\otimes} \to \mathcal{V}'^{\otimes} \times \mathcal{W}'^{\otimes}$  weakly bi-enriched  $\infty$ -categories. An enriched functor  $\phi \to \phi'$  is a commutative square of  $\infty$ -categories over Ass × Ass



such that  $\gamma$  preserves cocartesian lifts of inert morphisms of Ass×Ass whose first component preserves the maximum and whose second component preserves the minimum.

- An enriched functor  $\phi \to \phi'$  is an embedding if  $\alpha, \beta, \gamma$  are fully faithful.
- An enriched functor  $\phi \rightarrow \phi'$  is left (right) linear if it preserves left (right) tensors.
- An enriched functor  $\phi \rightarrow \phi'$  is linear if it is left and right linear.
- An enriched functor  $\phi \to \phi'$  is left  $\mathcal{V}$ -enriched if  $\alpha$  is the identity.
- An enriched functor  $\phi \rightarrow \phi'$  is right W-enriched if  $\beta$  is the identity.
- An enriched functor  $\phi \to \phi'$  is  $\mathcal{V}, \mathcal{W}$ -enriched if it is left  $\mathcal{V}$ -enriched and right  $\mathcal{W}$ -enriched.
- An enriched functor  $\phi \to \phi'$  is left  $\mathcal{V}$ -linear if it is left linear and left  $\mathcal{V}$ -enriched.
- An enriched functor  $\phi \to \phi'$  is right W-linear if it is right linear and right W-enriched.
- An enriched functor  $\phi \rightarrow \phi'$  is  $\mathcal{V}, \mathcal{W}$ -linear if it is left  $\mathcal{V}$ -linear and right  $\mathcal{W}$ -linear (or equivalently  $\mathcal{V}, \mathcal{W}$ -enriched and linear).

Notation 2.57. Let  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}, \mathcal{N}^{\circledast} \to \mathcal{V}'^{\otimes} \times \mathcal{W}'^{\otimes}$  be weakly bi-enriched  $\infty$ -categories.

(1) Let

 $\mathrm{EnrFun}(\mathcal{M},\mathcal{N}) \subset (\mathrm{Fun}(\mathcal{V}^{\otimes},\mathcal{V}'^{\otimes}) \times \mathrm{Fun}(\mathcal{W}^{\otimes},\mathcal{W}'^{\otimes})) \times_{\mathrm{Fun}(\mathcal{M}^{\otimes},\mathcal{V}'^{\otimes} \times \mathcal{W}'^{\otimes})} \mathrm{Fun}(\mathcal{M}^{\otimes},\mathcal{N}^{\otimes})$ 

be the full subcategory of enriched functors.

16

(2) Let

LLinFun( $\mathcal{M}, \mathcal{N}$ ), RLinFun( $\mathcal{M}, \mathcal{N}$ ), LinFun( $\mathcal{M}, \mathcal{N}$ )  $\subset$  EnrFun( $\mathcal{M}, \mathcal{N}$ )

be the full subcategories of left linear, right linear, linear functors, respectively.

Notation 2.58. Let  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}, \mathcal{N}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  be weakly bi-enriched  $\infty$ -categories.

(1) Let

 $\operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}(\mathcal{M},\mathcal{N}) \subset \operatorname{Fun}_{\mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}}(\mathcal{M}^{\otimes},\mathcal{N}^{\otimes})$ 

be the full subcategory of  $\mathcal{V}, \mathcal{W}$ -enriched functors.

(2) Let

 $\mathrm{LLinFun}_{\mathcal{V},\mathcal{W}}(\mathcal{M},\mathcal{N}) \subset \mathrm{EnrFun}_{\mathcal{V},\mathcal{W}}(\mathcal{M},\mathcal{N}),$ 

$$\operatorname{RLinFun}_{\mathcal{V},\mathcal{W}}(\mathcal{M},\mathcal{N}) \subset \operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}(\mathcal{M},\mathcal{N}),$$

$$\operatorname{LinFun}_{\mathcal{V},\mathcal{W}}(\mathcal{M},\mathcal{N}) \subset \operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}(\mathcal{M},\mathcal{N})$$

be the full subcategories of left linear, right linear, linear  $\mathcal{V}, \mathcal{W}$ -enriched functors, respectively.

(3) Let

 $\operatorname{LinFun}_{\mathcal{V},\mathcal{W}}^{\mathrm{L}}(\mathcal{M},\mathcal{N}) \subset \operatorname{LinFun}_{\mathcal{V},\mathcal{W}}(\mathcal{M},\mathcal{N})$ 

be the full subcategory of  $\mathcal{V}$ ,  $\mathcal{W}$ -linear functors whose underlying functor admits a right adjoint.

**Example 2.59.** Let  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be a weakly bi-enriched  $\infty$ -category and K an  $\infty$ -category.

- The functor  $K \times \mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  is a weakly bi-enriched  $\infty$ -category that exhibits  $K \times \mathcal{M}$  as weakly bi-enriched in  $\mathcal{V}, \mathcal{W}$ .
- The pullback  $(\mathcal{M}^{K})^{\circledast} := \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast} \times_{\operatorname{Fun}(K,\mathcal{V}^{\circledast}\times\mathcal{W}^{\circledast})} \operatorname{Fun}(K,\mathcal{M}^{\circledast}) \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  along the diagonal functor  $\mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast} \to \operatorname{Fun}(K,\mathcal{V}^{\circledast}\times\mathcal{W}^{\circledast})$  is a weakly bi-enriched  $\infty$ -category that exhibits  $\operatorname{Fun}(K,\mathcal{M})$  as weakly bi-enriched in  $\mathcal{V},\mathcal{W}$ .

**Remark 2.60.** Let  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}, \mathcal{N}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be weakly bi-enriched  $\infty$ -categories and K an  $\infty$ -category. The canonical equivalences

$$\operatorname{Fun}(\mathrm{K}, \operatorname{Fun}_{\mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}}(\mathcal{M}^{\circledast}, \mathcal{N}^{\circledast})) \simeq \operatorname{Fun}_{\mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}}(\mathrm{K} \times \mathcal{M}^{\circledast}, \mathcal{N}^{\circledast}) \simeq \operatorname{Fun}_{\mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}}(\mathcal{M}^{\circledast}, (\mathcal{N}^{\mathrm{K}})^{\circledast})$$

restrict to equivalences

 $\operatorname{Fun}(\mathrm{K},\operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}(\mathcal{M},\mathcal{N}))\simeq\operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}(\mathrm{K}\times\mathcal{M},\mathcal{N})\simeq\operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}(\mathcal{M},\mathcal{N}^{\mathrm{K}}).$ 

We will often use the following proposition, which is [16, Proposition 4.2.4.2.]:

**Proposition 2.61.** For every bitensored  $\infty$ -category  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  and  $\infty$ -category K the following forgetful functor is an equivalence:

$$\operatorname{LinFun}_{\mathcal{V},\mathcal{W}}(\mathcal{V} \times \mathrm{K} \times \mathcal{W}, \mathcal{M}) \to \operatorname{Fun}(\mathrm{K}, \mathcal{M}).$$

Notation 2.62. Let

$$\omega \operatorname{BEnr} \subset (\operatorname{Op}_{\infty} \times \operatorname{Op}_{\infty}) \times_{\mathsf{Cat}_{\infty/\operatorname{Ass} \times \operatorname{Ass}}} \operatorname{Fun}([1], \mathsf{Cat}_{\infty/\operatorname{Ass} \times \operatorname{Ass}})$$

be the subcategory of weakly bi-enriched  $\infty$ -categories.

**Notation 2.63.** Evaluation at the target restricts to a forgetful functor  $\omega \text{BEnr} \to \text{Op}_{\infty} \times \text{Op}_{\infty}$  whose fibers over  $\infty$ -operads  $\mathcal{V}^{\otimes} \to \text{Ass}$ ,  $\mathcal{W}^{\otimes} \to \text{Ass}$  we denote by  $_{\mathcal{V}}\omega \text{BEnr}_{\mathcal{W}}$ .

Remark 2.64. There is a canonical equivalence

 $\omega \operatorname{BEnr}_{\varnothing, \varnothing} \simeq \operatorname{\mathsf{Cat}}_{\infty}, \ \mathcal{M}^{\circledast} \to \varnothing^{\otimes} \times \varnothing^{\otimes} \mapsto \mathcal{M}^{\circledast}$ 

inv<br/>serse to the functor  $\mathbf{K}\mapsto \varnothing^{\otimes}\times\mathbf{K}\times\varnothing^{\otimes}\simeq\mathbf{K}.$ 

The next proposition is [10, Theorem 3.30.]:

**Proposition 2.65.** The forgetful functor

$$\gamma: \omega \operatorname{BEnr} \to \operatorname{Op}_{\infty} \times \operatorname{Op}_{\infty}$$

is a cartesian fibration. Let  $\psi : \mathcal{M}^{\circledast} \to \mathcal{N}^{\circledast}$  be an enriched functor lying over maps of  $\infty$ -operads  $\alpha : \mathcal{V}^{\otimes} \to \mathcal{V}^{\otimes}, \alpha' : \mathcal{W}^{\otimes} \to \mathcal{W}^{\otimes}$ . The map  $\psi$  is  $\gamma$ -cartesian if and only if the commutative square



is a pullback square, in other words if  $\psi$  induces an equivalence  $\mathcal{M} \simeq \mathcal{N}$  and for every  $V_1, ..., V_n \in \mathcal{V}, X, Y \in \mathcal{M}, W_1, ..., W_m \in \mathcal{W}$  for  $n, m \ge 0$  the following canonical map is an equivalence:

$$\operatorname{Mul}_{\mathcal{M}}(V_{1},...,V_{n},X,W_{1},...,W_{m};Y) \to \operatorname{Mul}_{\mathcal{N}}(\alpha(V_{1}),...,\alpha(V_{n}),\psi(X),\alpha'(W_{1}),...,\alpha'(W_{m});\psi(Y)).$$

Notation 2.66. The category Ass =  $\Delta^{\text{op}}$  carries a canonical involution sending [n] to [n] and a map  $f:[n] \rightarrow [m]$  to the map  $[n] \rightarrow [m], i \mapsto m - f(n-i)$ . The involution on Ass induces an involution on  $\mathsf{Cat}_{\infty/Ass}$  that restricts to an involution  $(-)^{\text{rev}}$  on  $\mathsf{Op}_{\infty} \subset \mathsf{Cat}_{\infty/Ass}$ . Moreover the involution on Ass induces an involution on Ass × Ass by applying the involution on Ass to each factor and switching the factors, which induces an involution on  $(\mathsf{Cat}_{\infty/Ass} \times \mathsf{Cat}_{\infty/Ass}) \times_{\mathsf{Cat}_{\infty/Ass}} \mathsf{Fun}([1], \mathsf{Cat}_{\infty/Ass} \times \mathsf{Ass})$  that restricts to involutions  $(-)^{\text{rev}}$  on  $\omega$ BEnr and BMod, under which LMod corresponds to RMod.

**Remark 2.67.** There is a canonical left action of  $\mathsf{Cat}_{\infty}$  on  $\omega BEnr$  that sends  $K, \mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  to  $K \times \mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$ . The forgetful functor  $\omega BEnr \to \operatorname{Op}_{\infty} \times \operatorname{Op}_{\infty}$  is  $\mathsf{Cat}_{\infty}$ -linear, where the target carries the trivial action. Thus for any small  $\infty$ -operads  $\mathcal{V}^{\circledast} \to \operatorname{Ass}, \mathcal{W}^{\circledast} \to \operatorname{Ass}$  also the fiber  $_{\mathcal{V}}\omega BEnr_{\mathcal{W}}$  carries a left  $\mathsf{Cat}_{\infty}$ -action that acts the same. These actions are closed by Remark 2.60 and so exhibit  $\omega BEnr$  and  $_{\mathcal{V}}\omega BEnr_{\mathcal{W}}$  as left  $\mathsf{Cat}_{\infty}$ -enriched, i.e. as  $(\infty, 2)$ -categories.

## Notation 2.68. Let

#### $\operatorname{LMod},\operatorname{RMod},\operatorname{BMod} \subset \omega\operatorname{BEnr}$

be the subcategories of left tensored, right tensored, bitensored  $\infty$ -categories and enriched functors preserving left tensors, right tensors, left and right tensors, respectively.

#### Notation 2.69. Let

#### $ccLMod \subset LMod$ , $ccRMod \subset RMod$ , $ccBMod \subset BMod$

be the subcategories of left tensored, right tensored, bitensored  $\infty$ -categories, respectively, compatible with small colimits and left linear, right linear, linear functors, respectively, preserving small colimits.

The next proposition follows from [10, Corollary 8.31.]:

**Proposition 2.70.** Let  $\kappa$  be a small regular cardinal and  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  a totally small bitensored  $\infty$ -category.

- (1) There is a bitensored  $\infty$ -category  $\operatorname{Ind}_{\kappa}(\mathcal{M})^{\circledast} \to \operatorname{Ind}_{\kappa}(\mathcal{V})^{\circledast} \times \operatorname{Ind}_{\kappa}(\mathcal{W})^{\circledast}$  compatible with small  $\kappa$ -filtered colimits and a  $\mathcal{V}, \mathcal{W}$ -linear embedding  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times_{\operatorname{Ind}_{\kappa}(\mathcal{V})^{\circledast}} \operatorname{Ind}_{\kappa}(\mathcal{M})^{\circledast} \times_{\operatorname{Ind}_{\kappa}(\mathcal{W})^{\circledast}} \mathcal{W}^{\circledast}$  inducing the embedding  $\mathcal{M} \to \operatorname{Ind}_{\kappa}(\mathcal{M})$  on underlying  $\infty$ -categories.
- (2) For every bitensored  $\infty$ -category  $\mathbb{N}^{\otimes} \to \operatorname{Ind}_{\kappa}(\mathcal{V})^{\otimes} \times \operatorname{Ind}_{\kappa}(\mathcal{W})^{\otimes}$  compatible with small  $\kappa$ -filtered colimits the functor

 $\operatorname{EnrFun}_{\operatorname{Ind}_{\kappa}(\mathcal{V}),\operatorname{Ind}_{\kappa}(\mathcal{W})}(\operatorname{Ind}_{\kappa}(\mathcal{M}),\mathcal{N}) \to \operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}(\mathcal{M},\mathcal{N})$ 

admits a fully faithful left adjoint that lands in the full subcategory

 $\mathrm{EnrFun}_{\mathrm{Ind}_{\kappa}(\mathcal{V}),\mathrm{Ind}_{\kappa}(\mathcal{W})}^{\kappa-\mathrm{fil}}(\mathrm{Ind}_{\kappa}(\mathcal{M}),\mathcal{N})$ 

of enriched functors preserving small  $\kappa$ -filtered colimits. So the following is an equivalence:

 $\operatorname{EnrFun}_{\operatorname{Ind}_{\kappa}(\mathcal{V}),\operatorname{Ind}_{\kappa}(\mathcal{W})}^{\kappa-\operatorname{fil}}(\operatorname{Ind}_{\kappa}(\mathcal{M}),\mathcal{N}) \to \operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}(\mathcal{M},\mathcal{N}).$ 

(3) If M admits  $\kappa$ -small colimits, the latter equivalence restricts to an equivalence

 $\operatorname{EnrFun}_{\operatorname{Ind}_{\kappa}(\mathcal{V}),\operatorname{Ind}_{\kappa}(\mathcal{W})}^{L}(\operatorname{Ind}_{\kappa}(\mathcal{M}),\mathcal{N}) \to \operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}^{\kappa}(\mathcal{M},\mathcal{N}).$ 

Notation 2.71. Let  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be a totally small weakly bi-enriched  $\infty$ -category. We set

 $\mathcal{P}(\mathcal{V})^{\otimes} \coloneqq \mathrm{Ind}_{\varnothing}(\mathcal{V})^{\otimes}, \ \mathcal{P}(\mathcal{M})^{\otimes} \coloneqq \mathrm{Ind}_{\varnothing}(\mathcal{M})^{\otimes} \to \mathcal{P}(\mathcal{V})^{\otimes} \times \mathcal{P}(\mathcal{W})^{\otimes}.$ 

Next we define weakly enriched  $\infty$ -categories of enriched functors:

**Lemma 2.72.** Let  $\mathcal{C} \to T, T \to S$  be functors such that  $\mathcal{C} \to T \to S$  is a cocartesian fibration. The functor  $(-) \times_S \mathcal{C} : \operatorname{Cat}_{\infty/S} \to \operatorname{Cat}_{\infty/C} \to \operatorname{Cat}_{\infty/T}$  admits a right adjoint that we denote by  $\operatorname{Fun}_T^S(\mathcal{C}, -)$ .

**Remark 2.73.** For every functors  $\mathcal{D} \to T$  and  $S' \to S$  there is a canonical equivalence  $S' \times_S \operatorname{Fun}_T^S(\mathcal{C}, \mathcal{D}) \simeq \operatorname{Fun}_{S' \times_S T}^{S'}(S' \times_S \mathcal{C}, S' \times_S \mathcal{D})$  specifying the fibers of the functor  $\operatorname{Fun}_T^S(\mathcal{C}, \mathcal{D}) \to S$ .

The next proposition is [10, Lemma 3.73.]:

**Proposition 2.74.** Let  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast}, \mathcal{N}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  be weakly left, weakly bi-enriched  $\infty$ -categories, respectively. The functor  $\operatorname{Fun}_{\mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}}^{\mathcal{W}^{\otimes}}(\mathcal{M}^{\circledast} \times \mathcal{W}^{\otimes}, \mathcal{N}^{\circledast}) \to \mathcal{W}^{\otimes}$  is a weakly right enriched  $\infty$ -category.

Notation 2.75. Let  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast}, \mathcal{N}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  be weakly left, weakly bi-enriched  $\infty$ -categories, respectively. Let

 $\mathrm{EnrFun}_{\mathcal{V},\varnothing}(\mathcal{M},\mathcal{N})^{\circledast} \subset \mathrm{Fun}_{\mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}}(\mathcal{M}^{\circledast} \times \mathcal{W}^{\otimes},\mathcal{N}^{\circledast}) \to \mathcal{W}^{\otimes}$ 

be the full subcategory weakly right enriched in  $\mathcal{W}$  spanned by

 $\mathrm{EnrFun}_{\mathcal{V},\varnothing}(\mathcal{M},\mathcal{N}) \subset \mathrm{Fun}_{\mathcal{V}^{\otimes}}(\mathcal{M}^{\circledast},\mathcal{N}_{[0]}^{\circledast}) \simeq \mathrm{Fun}_{\mathcal{V}^{\otimes}\times\mathcal{W}^{\otimes}}^{\mathcal{W}^{\otimes}}(\mathcal{M}^{\otimes}\times\mathcal{W}^{\otimes},\mathcal{N}^{\otimes})_{[0]}.$ 

Notation 2.76. Let  $\mathcal{O}^{\circledast} \to \mathcal{W}^{\otimes}, \mathcal{N}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be weakly right, weakly bi-enriched  $\infty$ -categories, respectively. Let

 $\mathrm{EnrFun}_{\varnothing, \mathcal{W}}(\mathcal{O}, \mathcal{N})^{\circledast} \coloneqq (\mathrm{EnrFun}_{\mathcal{W}, \varnothing}(\mathcal{O}^{\mathrm{rev}}, \mathcal{N}^{\mathrm{rev}})^{\mathrm{rev}})^{\circledast} \to \mathcal{V}^{\otimes}$ 

be the weakly left  $\mathcal{V}$ -enriched  $\infty$ -category.

The following proposition is [10, Proposition 3.78.]:

**Proposition 2.77.** Let  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast}, \mathcal{O}^{\circledast} \to \mathcal{W}^{\circledast}, \mathcal{N}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  be weakly left, weakly right, weakly bi-enriched  $\infty$ -categories, respectively. The canonical equivalence

$$\operatorname{Fun}_{\mathcal{W}^{\otimes}}(\mathcal{O}^{\circledast}, \operatorname{Fun}_{\mathcal{V}^{\otimes}\times\mathcal{W}^{\otimes}}(\mathcal{M}^{\circledast}\times\mathcal{W}^{\otimes}, \mathcal{N}^{\circledast})) \simeq \operatorname{Fun}_{\mathcal{V}^{\otimes}\times\mathcal{W}^{\otimes}}(\mathcal{M}^{\circledast}\times\mathcal{O}^{\otimes}, \mathcal{N}^{\circledast})$$

restricts to an equivalence

$$\operatorname{EnrFun}_{\varnothing, W}(\mathcal{O}, \operatorname{EnrFun}_{\mathcal{V}, \varnothing}(\mathcal{M}, \mathcal{N})) \simeq \operatorname{EnrFun}_{\mathcal{V}, W}(\mathcal{M} \times \mathcal{O}, \mathcal{N}).$$

The next remark is [9, Remark 5.13.].

**Remark 2.78.** Let  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes}, \mathcal{N}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}, \mathcal{N}'^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be weakly left, weakly bi-enriched  $\infty$ -categories, respectively. For every  $\mathcal{V}, \mathcal{W}$ -enriched embedding  $\mathcal{N}^{\otimes} \to \mathcal{N}'^{\otimes}$  the induced right  $\mathcal{W}$ -enriched functor  $\operatorname{EnrFun}_{\mathcal{V}, \emptyset}(\mathcal{M}, \mathcal{N})^{\otimes} \to \operatorname{EnrFun}_{\mathcal{V}, \emptyset}(\mathcal{M}, \mathcal{N}')^{\otimes}$  is an embedding.

Next we consider trivial weakly bi-enriched  $\infty$ -categories.

**Definition 2.79.** Let  $\mathcal{V}^{\otimes} \to Ass, \mathcal{W}^{\otimes} \to Ass$  be  $\infty$ -operads.

(1) A weakly bi-enriched  $\infty$ -category  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  is left trivial if for every  $n > 0, m \ge 0$ and  $V_1, ..., V_n \in \mathcal{V}, W_1, ..., W_m \in \mathcal{W}, X, Y \in \mathcal{M}$  the space  $Mul_{\mathcal{M}}(V_1, ..., V_n, X, W_1, ..., W_m; Y)$  is empty.

- (2) A weakly bi-enriched  $\infty$ -category  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  right trivial if for every  $n \ge 0, m > 0$ and  $V_1, ..., V_n \in \mathcal{V}, W_1, ..., W_m \in \mathcal{W}, X, Y \in \mathcal{M}$  the space  $Mul_{\mathcal{M}}(V_1, ..., V_n, X, W_1, ..., W_m; Y)$  is empty.
- (3) A weakly bi-enriched  $\infty$ -category is trivial if it is left trivial and right trivial.

**Remark 2.80.** A weakly bi-enriched  $\infty$ -category  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  is left trivial if and only if the functor  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes}$  factors through the subcategory  $\operatorname{triv}_{\mathcal{V}}^{\otimes} \subset \mathcal{V}^{\otimes}$ . And dually for right triviality.

Notation 2.81. Let  $Ass_{min}, Ass_{max} \subset Ass$  be the subcategories with the same objects and with morphisms the inert morphisms preserving the minimum, maximum, respectively.

Notation 2.82. Let  $\mathcal{V}^{\otimes} \to Ass, \mathcal{W}^{\otimes} \to Ass$  be  $\infty$ -operads and K an  $\infty$ -category.

(1) Let

(2) Let

- $_{\mathcal{V}}\mathrm{triv}^{\otimes} \coloneqq \mathcal{V}^{\otimes} \times_{\mathrm{Ass}} \mathrm{Ass}_{\mathrm{max}} \to \mathcal{V}^{\otimes}.$
- $\operatorname{triv}_{\mathcal{W}}^{\mathfrak{G}} := \mathcal{W}^{\otimes} \times_{\operatorname{Ass}} \operatorname{Ass}_{\min} \to \mathcal{W}^{\otimes}.$
- (3) Let  $K_{\mathcal{V},\mathcal{W}}^{\circledast} := \operatorname{triv}_{\mathcal{V}}^{\circledast} \times \mathbf{K} \times \operatorname{triv}_{\mathcal{W}}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}.$

**Example 2.83.** Let  $\mathcal{V}^{\otimes} \to \operatorname{Ass}, \mathcal{W}^{\otimes} \to \operatorname{Ass}$  be  $\infty$ -operads and K a small  $\infty$ -category. Then  $_{\mathcal{V}}\operatorname{triv}^{\otimes} \to \mathcal{V}^{\otimes}, \operatorname{triv}^{\otimes}_{\mathcal{W}} \to \mathcal{W}^{\otimes}, \operatorname{K}^{\otimes}_{\mathcal{V},\mathcal{W}} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  are trivial weakly left, weakly right enriched, weakly bi-enriched  $\infty$ -categories, respectively. We call  $\operatorname{K}^{\otimes}_{\mathcal{V},\mathcal{W}} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  the trivial weakly bi-enriched  $\infty$ -category on K.

The next Proposition is [9, Proposition 2.83.]:

**Proposition 2.84.** Let  $\mathbb{N}^{\otimes} \to \mathbb{V}^{\otimes} \times \mathbb{W}^{\otimes}$  be a weakly bi-enriched  $\infty$ -category.

- (1) Let  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast}$  be a weakly left enriched  $\infty$ -category. The following functor is an equivalence: EnrFun<sub> $\mathcal{V},\mathcal{W}$ </sub> ( $\mathcal{M} \times \operatorname{triv}_{\mathcal{W}}, \mathcal{N}$ )  $\to$  EnrFun<sub> $\mathcal{V},\mathcal{Q}$ </sub> ( $\mathcal{M}, \mathcal{N}$ ).
  - (2) Let  $\mathcal{M}^{\circledast} \to \mathcal{W}^{\circledast}$  be a weakly right enriched  $\infty$ -category. The following functor is an equivalence:

 $\mathrm{EnrFun}_{\mathcal{V},\mathcal{W}}(_{\mathcal{V}}\mathrm{triv}\times\mathcal{M},\mathcal{N})\to\mathrm{EnrFun}_{\varnothing,\mathcal{W}}(\mathcal{M},\mathcal{N}).$ 

(3) Let  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  be a weakly bi-enriched  $\infty$ -category. The next functor is an equivalence: EnrFun<sub> $\mathcal{V},\mathcal{W}$ </sub>(vtriv×triv<sub> $\mathcal{W},\mathcal{N}$ </sub>)  $\to \mathcal{N}$ .

Next we define enriched adjunctions following  $[9, \S 2.4.]$ . The next definitions are [9, Definitions 2.63., 2.64.]:

**Definition 2.85.** An enriched adjunction is an adjunction in the  $(\infty, 2)$ -category  $\omega$ BEnr of Remark 2.67.

**Definition 2.86.** Let  $\mathcal{V}^{\otimes} \to \operatorname{Ass}, \mathcal{W}^{\otimes} \to \operatorname{Ass}$  be small  $\infty$ -operads. An  $\mathcal{V}, \mathcal{W}$ -enriched adjunction is an adjunction in the  $(\infty, 2)$ -category  $_{\mathcal{V}}\omega \operatorname{BEnr}_{\mathcal{W}}$  of Remark 2.67.

**Remark 2.87.** A  $\mathcal{V}, \mathcal{W}$ -enriched adjunction is an adjunction relative to  $\mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$ .

The next remark is [9, Remark 2.67.]:

**Remark 2.88.** Let  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}, \mathcal{N}^{\circledast} \to \mathcal{V}'^{\otimes} \times \mathcal{W}'^{\otimes}$  be weakly bi-enriched  $\infty$ -categories.

(1) An enriched functor  $G: \mathbb{N}^{\circledast} \to \mathbb{M}^{\circledast}$  lying over maps of  $\infty$ -operads  $\gamma: \mathcal{V}^{(\circledast)} \to \mathcal{V}^{\circledast}, \delta: \mathcal{W}^{(\circledast)} \to \mathcal{W}^{\circledast}$ admits an enriched left adjoint if  $\gamma, \delta$  admit left adjoints  $\alpha, \beta$  relative to Ass, respectively, and for every  $X \in \mathcal{M}$  there is an  $Y \in \mathbb{N}$  and a morphism  $X \to G(Y)$  in  $\mathcal{M}$  such that for any  $V_1, ..., V_n \in \mathcal{V}, W_1, ..., W_m \in \mathcal{W}$  for  $n, m \ge 0$  and  $Z \in \mathbb{N}$  the following map is an equivalence:

$$\operatorname{Mul}_{\mathcal{N}}(\alpha(V_1), ..., \alpha(V_n), Y, \beta(W_1), ..., \beta(W_m); Z) \rightarrow$$

20

$$\begin{split} \operatorname{Mul}_{\mathcal{M}}(\gamma(\alpha(V_{1})),...,\gamma(\alpha(V_{n})),\operatorname{G}(Y),\delta(\beta(W_{1})),...,\delta(\beta(W_{m}));\operatorname{G}(Z)) \rightarrow \\ \operatorname{Mul}_{\mathcal{M}}(V_{1},...,V_{n},X,W_{1},...,W_{m};\operatorname{G}(Z)). \end{split}$$

- (2) An enriched functor  $F: \mathcal{M}^{\otimes} \to \mathcal{N}^{\otimes}$  lying over maps of  $\infty$ -operads  $\alpha: \mathcal{V}^{\otimes} \to \mathcal{V}'^{\otimes}, \beta: \mathcal{W}^{\otimes} \to \mathcal{W}'^{\otimes}$  admits an enriched right adjoint if  $\alpha, \beta$  admit right adjoints  $\gamma, \delta$  relative to Ass, respectively, and for every  $Y \in \mathcal{N}$  there is an  $X \in \mathcal{M}$  and a morphism  $F(X) \to Y$  in  $\mathcal{N}$  such that for any  $V_1, ..., V_n \in \mathcal{V}, W_1, ..., W_m \in \mathcal{W}$  for  $n, m \ge 0$  and  $Z \in \mathcal{M}$  the following map is an equivalence:
  - $\mathrm{Mul}_{\mathcal{M}}(\mathrm{V}_{1},...,\mathrm{V}_{n},\mathrm{Z},\mathrm{W}_{1},...,\mathrm{W}_{m};\mathrm{X}) \rightarrow \mathrm{Mul}_{\mathcal{N}}(\alpha(\mathrm{V}_{1}),...,\alpha(\mathrm{V}_{n}),\mathrm{F}(\mathrm{Z}),\beta(\mathrm{W}_{1}),...,\beta(\mathrm{W}_{m});\mathrm{F}(\mathrm{X}))$

$$\rightarrow \operatorname{Mul}_{\mathcal{N}}(\alpha(V_1), ..., \alpha(V_n), F(Z), \beta(W_1), ..., \beta(W_m); Y).$$

(3) The same holds for  $\mathcal{V}, \mathcal{W}$ -enriched left and right adjoint, where  $\alpha, \beta$  are the identities.

Notation 2.89. Let  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}, \mathcal{N}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  be weakly bi-enriched  $\infty$ -categories. Let

$$\operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}^{\operatorname{L}}(\mathcal{M},\mathcal{N}) \subset \operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}(\mathcal{M},\mathcal{N}), \ \operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}^{\operatorname{R}}(\mathcal{M},\mathcal{N}) \subset \operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}(\mathcal{M},\mathcal{N})$$

be the full subcategories of  $\mathcal{V}, \mathcal{W}$ -enriched functors that admit a  $\mathcal{V}, \mathcal{W}$ -enriched left adjoint, a  $\mathcal{V}, \mathcal{W}$ -enriched right adjoint, respectively.

The next proposition is [9, Proposition 2.111.]:

**Proposition 2.90.** Let  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}, \mathcal{N}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be weakly bi-enriched  $\infty$ -categories. There is a canonical equivalence

$$\operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}^{\operatorname{L}}(\mathcal{M},\mathcal{N})^{\operatorname{op}} \simeq \operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}^{\operatorname{R}}(\mathcal{N},\mathcal{M}).$$

2.4. **Tensored envelopes.** In the following we introduce a tool that reduces questions about weakly enriched  $\infty$ -categories to tensored  $\infty$ -categories. The next propositions follow from [10, Proposition 3.92, Proposition 3.101.]:

**Proposition 2.91.** Let  $\mathcal{V}^{\otimes} \to Ass$  be an  $\infty$ -operad. There is a monoidal  $\infty$ -category  $Env(\mathcal{V})^{\otimes} \to Ass$  and an embedding of  $\infty$ -operads  $\mathcal{V}^{\otimes} \subset Env(\mathcal{V})^{\otimes}$  such that

- (1) Every object of  $\operatorname{Env}(\mathcal{V})$  is equivalent to  $V_1 \otimes \ldots \otimes V_n$  for  $n \ge 0$  and  $V_1, \ldots, V_n \in \mathcal{V} \subset \operatorname{Env}(\mathcal{V})$ .
- (2) For every monoidal  $\infty$ -category  $\mathcal{W}^{\otimes} \to Ass$  the induced functor

$$\operatorname{Alg}_{\operatorname{Env}(\mathcal{V})}(\mathcal{W}) \to \operatorname{Alg}_{\mathcal{V}}(\mathcal{W})$$

admits a fully faithful left adjoint that lands in the full subcategory of monoidal functors. In particular, the following induced functor is an equivalence:

$$\operatorname{Fun}^{\otimes}(\operatorname{Env}(\mathcal{V}),\mathcal{W}) \to \operatorname{Alg}_{\mathcal{V}}(\mathcal{W}).$$

**Definition 2.92.** Let  $\mathcal{V}^{\otimes} \to Ass$  be an  $\infty$ -operad. We call  $Env(\mathcal{V})^{\otimes} \to Ass$  the monoidal envelope.

**Proposition 2.93.** Let  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  be a weakly bi-enriched  $\infty$ -category. There is a bitensored  $\infty$ -category  $\operatorname{BEnv}(\mathcal{M})^{\circledast} \to \operatorname{Env}(\mathcal{V})^{\circledast} \times \operatorname{Env}(\mathcal{W})^{\circledast}$  and an enriched embedding  $\mathcal{M}^{\circledast} \subset \operatorname{BEnv}(\mathcal{M})^{\circledast}$  lying over the embeddings  $\mathcal{V}^{\circledast} \subset \operatorname{Env}(\mathcal{V})^{\circledast}, \mathcal{W}^{\circledast} \subset \operatorname{Env}(\mathcal{W})^{\circledast}$  such that

- (1) Every object of BEnv( $\mathfrak{M}$ ) is equivalent to  $V_1 \otimes ... \otimes V_n \otimes X \otimes W_1 \otimes ... \otimes W_m$  for  $n, m \ge 0$  and  $V_1, ..., V_n \in \mathcal{V} \subset Env(\mathcal{V}), X \in \mathcal{M} \subset BEnv(\mathcal{M}), W_1, ..., W_m \in \mathcal{W} \subset Env(\mathcal{W}).$
- (2) For every bitensored  $\infty$ -category  $\mathbb{N}^{\otimes} \to \operatorname{Env}(\mathbb{V})^{\otimes} \times \operatorname{Env}(\mathbb{W})^{\otimes}$  the functor

 $\operatorname{EnrFun}_{\operatorname{Env}(\mathcal{V}),\operatorname{Env}(\mathcal{W})}(\operatorname{BEnv}(\mathcal{M}),\mathcal{N}) \to \operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}(\mathcal{M},\mathcal{N})$ 

admits a fully faithful left adjoint that lands in the full subcategory of  $\text{Env}(\mathcal{V}), \text{Env}(\mathcal{W})$ -linear functors. In particular, the following induced functor is an equivalence:

 $\operatorname{LinFun}_{\operatorname{Env}(\mathcal{V}),\operatorname{Env}(\mathcal{W})}(\operatorname{BEnv}(\mathcal{M}),\mathcal{N}) \to \operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}(\mathcal{M},\mathcal{N}).$ 

**Definition 2.94.** Let  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  be a weakly bi-enriched  $\infty$ -category. We call  $\operatorname{BEnv}(\mathcal{M})^{\circledast} \to \operatorname{Env}(\mathcal{V})^{\circledast} \times \operatorname{Env}(\mathcal{W})^{\circledast}$  the bitensored envelope.

The next lemma is [10, Lemma 3.93.]:

**Lemma 2.95.** Let  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be a bitensored  $\infty$ -category. The embedding  $\mathcal{V}^{\otimes} \subset \operatorname{Env}(\mathcal{V})^{\otimes}$  admits a left adjoint relative to Ass. The embedding  $\mathcal{M}^{\circledast} \subset \operatorname{BEnv}(\mathcal{M})^{\circledast}$  admits an enriched left adjoint covering the left adjoints of the embeddings  $\mathcal{V}^{\otimes} \subset \operatorname{Env}(\mathcal{V})^{\otimes}, \mathcal{W}^{\otimes} \subset \operatorname{Env}(\mathcal{W})^{\otimes}$ .

For the next definition we use Propositions 2.26 and 2.70:

**Definition 2.96.** Let  $\mathcal{V}^{\otimes} \to Ass, \mathcal{W}^{\otimes} \to Ass$  be  $\infty$ -operads and  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a weakly bi-enriched  $\infty$ -category.

(1) The closed monoidal envelope is

 $\mathcal{P}\mathrm{Env}(\mathcal{V})^{\otimes} \coloneqq \mathcal{P}(\mathrm{Env}(\mathcal{V}))^{\otimes} \to \mathrm{Ass}$ 

(2) The closed bitensored envelope is

 $\mathcal{P}\mathrm{BEnv}(\mathcal{M})^{\otimes} \coloneqq \mathcal{P}(\mathrm{BEnv}(\mathcal{M}))^{\otimes} \to \mathcal{P}\mathrm{Env}(\mathcal{V})^{\otimes} \times \mathcal{P}\mathrm{Env}(\mathcal{W})^{\otimes}.$ 

(3) The closed left tensored envelope of  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  is the full left tensored subcategory

 $\mathcal{P}\mathrm{LEnv}(\mathcal{M})^{\circledast} \subset \mathcal{P}\mathrm{BEnv}(\mathcal{M})^{\circledast} \times_{\mathcal{P}\mathrm{Env}(\mathcal{W})^{\circledast}} \mathcal{W}^{\otimes} \to \mathcal{P}\mathrm{Env}(\mathcal{V})^{\otimes} \times \mathcal{W}^{\otimes}$ 

generated under small colimits by the objects  $V_1 \otimes ... \otimes V_n \otimes X$  for  $n \ge 0$  and  $V_1, ..., V_n \in \mathcal{V}$ and  $X \in \mathcal{M}$ .

(4) The closed right tensored envelope of  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  is the full weakly right enriched subcategory

 $\mathfrak{P}\operatorname{REnv}(\mathfrak{M})^{\circledast} \subset \mathcal{V}^{\otimes} \times_{\mathfrak{P}\operatorname{Env}(\mathcal{V})^{\otimes}} \mathfrak{P}\operatorname{BEnv}(\mathfrak{M})^{\circledast} \to \mathcal{V}^{\otimes} \times \mathfrak{P}\operatorname{Env}(\mathcal{W})^{\otimes}$ 

generated under small colimits by the objects  $X \otimes W_1 \otimes ... \otimes W_m$  for  $m \ge 0$  and  $W_1, ..., W_m \in W, X \in M$ .

The embedding  $\mathcal{M}^{\circledast} \to \mathcal{P}BEnv(\mathcal{M})^{\circledast}$  induces embeddings  $\mathcal{M}^{\circledast} \to \mathcal{P}LEnv(\mathcal{M})^{\circledast}, \mathcal{M}^{\circledast} \to \mathcal{P}REnv(\mathcal{M})^{\circledast}$ . Propositions 2.26, 2.70, 2.91 and 2.93 imply the following corollary:

**Corollary 2.97.** (1) Let  $\mathcal{V}^{\otimes} \to Ass$  be a small  $\infty$ -operad. For every monoidal  $\infty$ -category  $\mathcal{W}^{\otimes} \to Ass$  compatible with small colimits the induced functor

$$\operatorname{Alg}_{\operatorname{\mathcal{P}Env}(\mathcal{V})}(\mathcal{W}) \to \operatorname{Alg}_{\mathcal{V}}(\mathcal{W})$$

admits a fully faithful left adjoint that lands in the full subcategory of monoidal functors that admit a right adjoint. In particular, the following induced functor is an equivalence:

$$\operatorname{Fun}^{\otimes, L}(\operatorname{\mathcal{P}Env}(\mathcal{V}), \mathcal{W}) \to \operatorname{Alg}_{\mathcal{V}}(\mathcal{W})$$

(2) Let  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be a weakly bi-enriched  $\infty$ -category. For every bitensored  $\infty$ -category  $\mathcal{N}^{\otimes} \to \mathcal{P}\mathrm{Env}(\mathcal{V})^{\otimes} \times \mathcal{P}\mathrm{Env}(\mathcal{W})^{\otimes}$  compatible with small colimits the induced functor

 $\mathrm{EnrFun}_{\mathcal{P}\mathrm{Env}(\mathcal{V}),\mathcal{P}\mathrm{Env}(\mathcal{W})}(\mathcal{P}\mathrm{B}\mathrm{Env}(\mathcal{M}),\mathcal{N})\to\mathrm{EnrFun}_{\mathcal{V},\mathcal{W}}(\mathcal{M},\mathcal{N})$ 

admits a fully faithful left adjoint that lands in the full subcategory of  $PEnv(\mathcal{V})$ ,  $PEnv(\mathcal{W})$ -linear functors that admit a right adjoint. In particular, the following functor is an equivalence:

 $\operatorname{LinFun}_{\mathcal{P}\operatorname{Env}(\mathcal{V}),\mathcal{P}\operatorname{Env}(\mathcal{W})}^{L}(\mathcal{P}\operatorname{B}\operatorname{Env}(\mathcal{M}),\mathcal{N}) \to \operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}(\mathcal{M},\mathcal{N}).$ 

**Remark 2.98.** By Corollary 2.97 (2) every enriched functor  $F : \mathcal{M}^{\circledast} \to \mathcal{N}^{\circledast}$  starting at a totally small weakly bi-enriched  $\infty$ -category uniquely extends to a linear functor  $\mathcal{PBEnv}(\mathcal{M})^{\circledast} \to \mathcal{N}^{\circledast}$  that lies over left adjoint monoidal functors and admits an enriched right adjoint, which factors as  $\mathcal{N}^{\circledast} \subset \mathcal{PBEnv}(\mathcal{N})^{\circledast} \xrightarrow{\mathrm{BEnv}(F)^*} \mathcal{PBEnv}(\mathcal{M})^{\circledast}$ .

**Remark 2.99.** By the universal property of Corollary 2.97 (2) there is a canonical equivalence  $\mathcal{PBEnv}(\mathcal{M}^{rev})^{\circledast} \simeq (\mathcal{PBEnv}(\mathcal{M})^{rev})^{\circledast}$  that restricts to an equivalence  $\mathcal{PLEnv}(\mathcal{M}^{rev})^{\circledast} \simeq (\mathcal{PREnv}(\mathcal{M})^{rev})^{\circledast}$ .

**Lemma 2.100.** Let  $F : \mathcal{M}^{\circledast} \to \mathcal{M}'^{\circledast}$  be an enriched functor of totally small weakly bi-enriched  $\infty$ -categories lying over maps of  $\infty$ -operads  $\alpha : \mathcal{V}^{\circledast} \to \mathcal{V}'^{\circledast}, \beta : \mathcal{W}^{\circledast} \to \mathcal{W}'^{\circledast}$ . For every  $V_1, ..., V_n \in \mathcal{V}, W_1, ..., W_m \in \mathcal{W}$  for  $n, m \ge 0$  and  $Z \in \mathcal{M}'$  there is a canonical equivalence of presheaves on  $\mathcal{P}(\mathcal{M})$ :

$$\kappa : \mathcal{P}(\mathcal{M})(-, \operatorname{Mul}_{\mathcal{M}'}(\alpha(V_1), ..., \alpha(V_n), F(-), \beta(W_1), ..., \beta(W_m); Z)) \simeq Mul_{\mathcal{P}BEnv(\mathcal{M})}(V_1, ..., V_n, j_!(-), W_1, ..., W_m; F^*(Z)).$$

*Proof.* The equivalence  $\kappa$  is uniquely determined by its restriction to  $\mathcal{M}$  since source and target of  $\kappa$  preserve small limits. There is a canonical equivalence of presheaves on  $\mathcal{M}$ , where  $j : \mathcal{M} \subset BEnv(\mathcal{M})$  is the canonical embedding:

$$\begin{split} & \mathcal{P}(\mathcal{M})(\mathbf{y}_{\mathcal{M}}(-), \mathrm{Mul}_{\mathcal{M}'}(\alpha(\mathrm{V}_{1}), ..., \alpha(\mathrm{V}_{n}), \mathrm{F}(-), \beta(\mathrm{W}_{1}), ..., \beta(\mathrm{W}_{m}); \mathrm{Z})) \simeq \\ & \mathrm{Mul}_{\mathcal{M}'}(\alpha(\mathrm{V}_{1}), ..., \alpha(\mathrm{V}_{n}), \mathrm{F}(-), \beta(\mathrm{W}_{1}), ..., \beta(\mathrm{W}_{m}); \mathrm{Z}) \simeq \\ & \mathrm{BEnv}(\mathcal{M}')(\alpha(\mathrm{V}_{1}) \otimes ... \otimes \alpha(\mathrm{V}_{n}) \otimes \mathrm{j} \circ \mathrm{F}(-) \otimes \beta(\mathrm{W}_{1}) \otimes ... \otimes \beta(\mathrm{W}_{m}), \mathrm{Z}) \simeq \\ & \mathrm{F}^{*}(\mathrm{Z}) \circ (\mathrm{V}_{1} \otimes ... \otimes \mathrm{V}_{n} \otimes \mathrm{j}(-) \otimes \mathrm{W}_{1} \otimes ... \otimes \mathrm{W}_{m}) \simeq \\ & \mathcal{P}\mathrm{BEnv}(\mathcal{M})(\mathrm{V}_{1} \otimes ... \otimes \mathrm{V}_{n} \otimes (\mathrm{y}_{\mathrm{BEnv}(\mathcal{M})} \circ \mathrm{j}(-)) \otimes \mathrm{W}_{1} \otimes ... \otimes \mathrm{W}_{m}, \mathrm{F}^{*}(\mathrm{Z})) \simeq \\ & \mathrm{Mul}_{\mathcal{P}\mathrm{BEnv}(\mathcal{M})}(\mathrm{V}_{1}, ..., \mathrm{V}_{n}, (\mathrm{j}_{!} \circ \mathrm{y}_{\mathcal{M}}(-)), \mathrm{W}_{1}, ..., \mathrm{W}_{m}; \mathrm{F}^{*}(\mathrm{Z})). \end{split}$$

Proposition 2.93, 2.84 and 2.70 imply the following corollary:

**Corollary 2.101.** Let  $\mathcal{V}^{\otimes} \to Ass, \mathcal{W}^{\otimes} \to Ass$  be small  $\infty$ -operads and K an  $\infty$ -category.

(1) The unique  $Env(\mathcal{V}), Env(\mathcal{W})$ -linear functor

 $\operatorname{BEnv}(\operatorname{K}_{\mathcal{V},\mathcal{W}})^{\circledast} \to \operatorname{Env}(\mathcal{V})^{\circledast} \times \operatorname{K} \times \operatorname{Env}(\mathcal{W})^{\circledast}$ 

extending the functor  $K \to Env(\mathcal{V}) \times K \times Env(\mathcal{W})$  is an equivalence, which is inverse to the unique  $Env(\mathcal{V}), Env(\mathcal{W})$ -linear functor  $Env(\mathcal{V})^{\circledast} \times K \times Env(\mathcal{W})^{\circledast} \to BEnv(K_{\mathcal{V},\mathcal{W}})^{\circledast}$  extending the embedding  $K \subset BEnv(K_{\mathcal{V},\mathcal{W}})$ .

(2) The unique left adjoint  $\operatorname{PEnv}(\mathcal{V})$ ,  $\operatorname{PEnv}(\mathcal{W})$ -linear functor

 $\mathcal{P}BEnv(K_{\mathcal{V},\mathcal{W}})^{\circledast} \rightarrow (\mathcal{P}Env(\mathcal{V}) \otimes \mathcal{P}(K) \otimes \mathcal{P}Env(\mathcal{W}))^{\circledast}$ 

extending the functor  $K \to \mathcal{P}(K) \to \mathcal{P}Env(\mathcal{V}) \otimes \mathcal{P}(K) \otimes \mathcal{P}Env(\mathcal{W})$  is an equivalence, which is inverse to the unique left adjoint  $\mathcal{P}Env(\mathcal{V}), \mathcal{P}Env(\mathcal{W})$ -linear functor

 $(\mathcal{P}Env(\mathcal{V}) \otimes \mathcal{P}(K) \otimes \mathcal{P}Env(\mathcal{W}))^{\circledast} \to \mathcal{P}BEnv(K_{\mathcal{V},\mathcal{W}})^{\circledast}$ 

extending the embedding  $K \subset \mathcal{PBEnv}(K_{\mathcal{V},\mathcal{W}})$ .

2.5. Enrichment. Before defining enriched  $\infty$ -categories we define the more general class of pseudoenriched  $\infty$ -categories that contains all tensored  $\infty$ -categories. We follow [9, § 3]. The next definition is [9, Definition 3.1.]:

**Definition 2.102.** Let  $\phi : \mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be a weakly bi-enriched  $\infty$ -category.

(1) We say that  $\phi$  exhibits  $\mathcal{M}$  as left pseudo-enriched in  $\mathcal{V}$  if  $\mathcal{V}^{\otimes} \to Ass$  is a monoidal  $\infty$ -category and for every  $X, Y \in \mathcal{M}$  and  $V_1, ..., V_n \in \mathcal{V}, W_1, ..., W_m \in \mathcal{W}$  for  $n, m \ge 0$  the map

 $Mul_{\mathcal{M}}(V_1 \otimes ... \otimes V_n, X, W_1, ..., W_m; Y) \rightarrow Mul_{\mathcal{M}}(V_1, ..., V_n, X, W_1, ..., W_m; Y)$ 

induced by the active morphism  $V_1, ..., V_n \to V_1 \otimes ... \otimes V_n$  in  $\mathcal{V}^{\otimes}$  is an equivalence.

(2) We say that  $\phi$  exhibits  $\mathcal{M}$  as right pseudo-enriched in  $\mathcal{W}$  if  $\mathcal{W}^{\otimes} \to Ass$  is a monoidal  $\infty$ -category and for every  $X, Y \in \mathcal{M}$  and  $V_1, ..., V_n \in \mathcal{V}, W_1, ..., W_m \in \mathcal{W}$  for  $n, m \ge 0$  the map

 $Mul_{\mathcal{M}}(V_1, ..., V_n, X, W_1 \otimes ... \otimes W_m; Y) \rightarrow Mul_{\mathcal{M}}(V_1, ..., V_n, X, W_1, ..., W_m; Y)$ 

induced by the active morphism  $W_1, ..., W_m \to W_1 \otimes ... \otimes W_m$  in  $\mathcal{W}^{\otimes}$  is an equivalence.

(3) We say that  $\phi$  exhibits  $\mathcal{M}$  as bi-pseudo-enriched in  $\mathcal{V}, \mathcal{W}$  if  $\mathcal{V}^{\otimes} \to Ass, \mathcal{W}^{\otimes} \to Ass$  are monoidal  $\infty$ -categories and for any X, Y  $\in \mathcal{M}$  and  $V_1, ..., V_n \in \mathcal{V}, W_1, ..., W_m \in \mathcal{W}$  for  $n, m \ge 0$  the following map is an equivalence:

 $Mul_{\mathcal{M}}(V_{1}\otimes ...\otimes V_{n},X,W_{1}\otimes ...\otimes W_{m};Y) \rightarrow Mul_{\mathcal{M}}(V_{1},...,V_{n},X,W_{1},...,W_{m};Y).$ 

**Remark 2.103.** A weakly bi-enriched  $\infty$ -category  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  exhibits  $\mathcal{M}$  as bi-pseudo-enriched in  $\mathcal{V}, \mathcal{W}$  if and only if it exhibits  $\mathcal{M}$  as left pseudo-enriched in  $\mathcal{V}$  and right pseudo-enriched in  $\mathcal{W}$ .

The next lemma is [9, Lemma 3.3.]:

**Lemma 2.104.** (1) A weakly bi-enriched  $\infty$ -category  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  is a left tensored  $\infty$ -category if and only if it is a left pseudo-enriched  $\infty$ -category and admits left tensors.

- (2) A weakly bi-enriched ∞-category M<sup>®</sup> → V<sup>®</sup> × W<sup>®</sup> is a right tensored ∞-category if and only if it is a right pseudo-enriched ∞-category and admits right tensors.
- (3) A weakly bi-enriched ∞-category M<sup>®</sup> → V<sup>®</sup> × W<sup>®</sup> is a bitensored ∞-category if and only if it is a bi-pseudo-enriched ∞-category and admits bitensors.

Notation 2.105. Let

#### LPEnr, RPEnr, BPEnr $\subset \omega$ BEnr

be the full subcategories of left pseudo-enriched, right pseudo-enriched, bi-pseudo-enriched  $\infty$ -categories, respectively.

Notation 2.106. Let  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be a weakly bi-enriched  $\infty$ -category. For every  $X \in \mathcal{M}$  let

$$\Gamma_{\mathcal{M}}(\mathbf{X}, -) : \mathcal{M}^{\circledast} \to (\mathcal{P} \operatorname{Env}(\mathcal{V}) \otimes \mathcal{P} \operatorname{BEnv}(\mathcal{W}))^{\circledast} \simeq \mathcal{P}(\operatorname{Env}(\mathcal{V}) \times \operatorname{Env}(\mathcal{W}))^{\circledast}$$

be the restriction of the enriched right adjoint of the left adjoint  $\operatorname{PEnv}(\mathcal{V}), \operatorname{PEnv}(\mathcal{W})$ -linear functor

$$(\mathcal{P}Env(\mathcal{V}) \otimes \mathcal{P}BEnv(\mathcal{W}))^{\otimes} \to \mathcal{P}BEnv(\mathcal{M})^{\otimes}, (V, W) \mapsto V \otimes X \otimes W.$$

We get a  $\mathcal{V}, \mathcal{W}$ -enriched functor  $\Gamma_{\mathcal{M}} : \mathcal{M}^{\mathrm{op}} \times \mathcal{M}^{\circledast} \to \mathcal{P}(\mathrm{Env}(\mathcal{V}) \times \mathrm{Env}(\mathcal{W}))^{\circledast}$  that we call graph of  $\mathcal{M}$ .

**Remark 2.107.** For every  $X, Y \in M$  and  $V_1, ..., V_n \in V, W_1, ..., W_m \in W$  for  $n, m \ge 0$  there is a canonical equivalence

$$\Gamma_{\mathcal{M}}(X,Y)(V_{1} \otimes ... \otimes V_{n}, W_{1} \otimes ... \otimes W_{m}) \simeq \mathcal{P}BEnv(\mathcal{M})(V_{1} \otimes ... \otimes V_{n} \otimes X \otimes W_{1}, ..., W_{m}, Y)$$
$$\simeq Mul_{\mathcal{M}}(V_{1}, ..., V_{n}, X, W_{1}, ..., W_{m}; Y).$$

The next remark is [9, Remark 3.8.]:

**Remark 2.108.** A totally small weakly bi-enriched  $\infty$ -category  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  exhibits  $\mathcal{M}$  as

(1) left pseudo-enriched in  $\mathcal{V}$  if and only if  $\mathcal{V}^{\otimes} \to Ass$  is a monoidal  $\infty$ -category and for every  $X, Y \in \mathcal{M}$  the object

 $\Gamma_{\mathcal{M}}(\mathbf{X},\mathbf{Y}) \in \mathcal{P}(\operatorname{Env}(\mathcal{V}) \times \operatorname{Env}(\mathcal{W})) \simeq \operatorname{Fun}(\operatorname{Env}(\mathcal{W})^{\operatorname{op}}, \mathcal{P}\operatorname{Env}(\mathcal{V}))$ 

lies in Fun(Env( $\mathcal{W}$ )<sup>op</sup>,  $\mathcal{P}(\mathcal{V})$ ).

(2) right pseudo-enriched in  $\mathcal{W}$  if and only if  $\mathcal{W}^{\otimes} \to Ass$  is a monoidal  $\infty$ -category and for every  $X, Y \in \mathcal{M}$  the object

$$\Gamma_{\mathcal{M}}(\mathbf{X},\mathbf{Y}) \in \mathcal{P}(\operatorname{Env}(\mathcal{V}) \times \operatorname{Env}(\mathcal{W})) \simeq \operatorname{Fun}(\operatorname{Env}(\mathcal{V})^{\operatorname{op}}, \mathcal{P}\operatorname{Env}(\mathcal{W}))$$

lies in Fun(Env( $\mathcal{V}$ )<sup>op</sup>,  $\mathcal{P}(\mathcal{W})$ ).

(3) bi-pseudo-enriched in  $\mathcal{V}, \mathcal{W}$  if and only if  $\mathcal{V}^{\otimes} \to Ass, \mathcal{W}^{\otimes} \to Ass$  are monoidal  $\infty$ -categories and for every X, Y  $\in \mathcal{M}$  the object

$$\Gamma_{\mathcal{M}}(\mathbf{X},\mathbf{Y}) \in \mathcal{P}(\operatorname{Env}(\mathcal{V}) \times \operatorname{Env}(\mathcal{W}))$$

lies in  $\mathcal{P}(\mathcal{V} \times \mathcal{W})$ .

24

Now we are ready to define enriched  $\infty$ -categories. The next definition is [9, Definition 3.9.]:

**Definition 2.109.** Let  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be a weakly bi-enriched  $\infty$ -category.

(1) A left multi-morphism object of  $W_1, ..., W_m, X, Y \in \mathcal{M}$  for  $m \ge 0$  is an object

 $LMulMor_{\mathcal{M}}(X, W_1, ..., W_n; Y) \in \mathcal{V}$ 

such that there is a multi-morphism  $\beta \in Mul_{\mathcal{M}}(LMulMor_{\mathcal{M}}(X, W_1, ..., W_n; Y), X, W_1, ..., W_m; Y)$ that induces for every objects  $V_1, ..., V_n \in \mathcal{V}$  for  $n \ge 0$  an equivalence

 $\operatorname{Mul}_{\mathcal{V}}(V_1, ..., V_n; \operatorname{LMulMor}_{\mathcal{M}}(X, W_1, ..., W_m; Y)) \simeq \operatorname{Mul}_{\mathcal{M}}(V_1, ..., V_n, X, W_1, ..., W_m; Y).$ 

(2) A right multi-morphism object of  $V_1,...,V_n,X,Y\in\mathcal{M}$  for  $n\geq 0$  is an object

 $RMulMor_{\mathcal{M}}(V_1, ..., V_n, X; Y) \in \mathcal{W}$ 

such that there is a multi-morphism  $\alpha \in Mul_{\mathcal{M}}(V_1, ..., V_n, X, RMulMor_{\mathcal{M}}(V_1, ..., V_n, X; Y); Y)$ that induces for every objects  $W_1, ..., W_m \in \mathcal{W}$  for  $m \ge 0$  an equivalence

 $Mul_{\mathcal{W}}(W_1, ..., W_m; RMulMor_{\mathcal{M}}(V_1, ..., V_n, X; Y)) \simeq Mul_{\mathcal{M}}(V_1, ..., V_n, X, W_1, ..., W_m; Y).$ 

The next definition is [9, Definition 3.10.]:

**Definition 2.110.** Let  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be a weakly bi-enriched  $\infty$ -category and X, Y  $\in \mathcal{M}$ .

- (1) A left morphism object of X, Y in  $\mathcal{M}$  is a factorization  $\operatorname{LMor}_{\mathcal{M}}(X, Y) : \operatorname{Env}(\mathcal{W})^{\operatorname{op}} \to \mathcal{V}$  of the functor  $\operatorname{Env}(\mathcal{W})^{\operatorname{op}} \to \mathcal{P}\operatorname{Env}(\mathcal{V})$  corresponding to  $\Gamma_{\mathcal{M}}(X, Y) \in \mathcal{P}(\operatorname{Env}(\mathcal{V}) \times \operatorname{Env}(\mathcal{W}))$ .
- (2) A right morphism object of X, Y in  $\mathcal{M}$  is a factorization  $\operatorname{RMor}_{\mathcal{M}}(X, Y) : \operatorname{Env}(\mathcal{V})^{\operatorname{op}} \to \mathcal{W}$  of the functor  $\operatorname{Env}(\mathcal{V})^{\operatorname{op}} \to \mathcal{P}\operatorname{Env}(\mathcal{W})$  corresponding to  $\Gamma_{\mathcal{M}}(X, Y) \in \mathcal{P}(\operatorname{Env}(\mathcal{V}) \times \operatorname{Env}(\mathcal{W}))$ .

**Remark 2.111.** Let  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be a weakly bi-enriched  $\infty$ -category and  $X, Y \in \mathcal{M}, W_1, ..., W_m \in \mathcal{W}$  for  $m \ge 0$ . A left multi-morphism object  $LMulMor_{\mathcal{M}}(W_1, ..., W_m, X; Y) \in \mathcal{V}$  represents the presheaf  $\Gamma_{\mathcal{M}}(X, Y)(W_1 \otimes ... \otimes W_m, -) \in \mathcal{P}Env(\mathcal{V})$ . Consequently, there is a left morphism object  $LMor_{\mathcal{M}}(X, Y) : Env(\mathcal{W})^{\mathrm{op}} \to \mathcal{V}$  if and only if for every  $W_1, ..., W_m \in \mathcal{W}$  for  $m \ge 0$  there is a left multi-morphism object  $LMulMor_{\mathcal{M}}(W_1, ..., W_m, X; Y) \in \mathcal{V}$ . In this case there is a canonical equivalence

 $\operatorname{LMor}_{\mathcal{M}}(X, Y)(W_1 \otimes ... \otimes W_m) \simeq \operatorname{LMulMor}_{\mathcal{M}}(X, W_1, ..., W_m; Y).$ 

The similar holds for right (multi-) morphism objects.

**Remark 2.112.** Let  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  be a weakly bi-enriched  $\infty$ -category and X, Y  $\in \mathcal{M}$ . If the left and right morphism object of X, Y  $\in \mathcal{M}$  exist, for every  $V_1, ..., V_n \in \mathcal{V}, W_1, ..., W_m \in \mathcal{W}$  for  $n, m \ge 0$  there is a canonical equivalence

 $Mul_{\mathcal{W}}(W_1,...,W_m;RMor_{\mathcal{M}}(X,Y)(V_1\otimes...\otimes V_n))\simeq$ 

$$\begin{split} \operatorname{Mul}_{\mathcal{W}}(W_{1},...,W_{m};\operatorname{RMulMor}_{\mathcal{M}}(V_{1},...,V_{n},X;Y)) &\simeq \operatorname{Mul}_{\mathcal{M}}(V_{1},...,V_{n},X,W_{1},...,W_{m};Y) &\simeq \operatorname{Mul}_{\mathcal{V}}(V_{1},...,V_{n};\operatorname{LMulMor}_{\mathcal{M}}(X,W_{1},...,W_{m};Y)) &\simeq \operatorname{Mul}_{\mathcal{V}}(V_{1},...,V_{n};\operatorname{LMor}_{\mathcal{M}}(X,Y)(W_{1}\otimes...\otimes W_{m})). \end{split}$$
This gives an adjunction

$$\operatorname{LMor}_{\mathcal{M}}(X, Y)^{\operatorname{op}} : \operatorname{Env}(\mathcal{W}) \rightleftharpoons \operatorname{Env}(\mathcal{V})^{\operatorname{op}} : \operatorname{RMor}_{\mathcal{M}}(X, Y),$$

where the left adjoint lands in  $\mathcal{V}^{\text{op}}$  and the right adjoint lands in  $\mathcal{W}$  [9, Remark 3.20.].

The next proposition is [9, Proposition 3.13.]:

**Proposition 2.113.** Let  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be a weakly bi-enriched  $\infty$ -category. The embeddings  $\mathcal{M}^{\otimes} \subset \operatorname{BEnv}(\mathcal{M})^{\otimes}, \operatorname{BEnv}(\mathcal{M})^{\otimes} \subset \mathcal{P}\operatorname{BEnv}(\mathcal{M})^{\otimes}$ 

of weakly bi-enriched  $\infty$ -categories preserve left and right multi-morphism objects.

The next definition is [9, Definition 3.14.]:

**Definition 2.114.** (1) A weakly bi-enriched  $\infty$ -category  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  exhibits  $\mathcal{M}$  as left enriched in  $\mathcal{V}$  if for every X, Y  $\in \mathcal{M}$  there is a left morphism object  $\mathrm{LMor}_{\mathcal{M}}(X, Y) \in \mathcal{V}$ .

- (2) A weakly bi-enriched  $\infty$ -category  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  exhibits  $\mathcal{M}$  as right enriched in  $\mathcal{W}$  if for every X, Y  $\in \mathcal{M}$  there is a right morphism object  $\operatorname{RMor}_{\mathcal{M}}(X, Y) \in \mathcal{W}$ .
- (3) A weakly bi-enriched  $\infty$ -category  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  exhibits  $\mathcal{M}$  as bi-enriched in  $\mathcal{V}, \mathcal{W}$  if it exhibits  $\mathcal{M}$  as left enriched in  $\mathcal{V}$  and right enriched in  $\mathcal{W}$ .

#### Notation 2.115. Let

#### LEnr, REnr, BEnr $\subset \omega$ BEnr

be the full subcategories of left enriched, right enriched, bi-enriched  $\infty$ -categories.

The next lemma is [9, Lemma 3.18.]:

**Lemma 2.116.** A weakly bi-enriched  $\infty$ -category  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  that exhibits  $\mathcal{M}$  as left enriched in a monoidal  $\infty$ -category, right enriched in a monoidal  $\infty$ -category, bi-enriched in monoidal  $\infty$ categories, respectively, exhibits  $\mathcal{M}$  as left pseudo-enriched, right pseudo-enriched, bi-pseudo-enriched, respectively.

**Remark 2.117.** Let  $\mathcal{V}^{\otimes} \to Ass, \mathcal{W}^{\otimes} \to Ass$  be presentably monoidal  $\infty$ -categories,  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a weakly bi-enriched  $\infty$ -category and  $\alpha : \mathcal{V} \times \mathcal{W} \to \mathcal{V} \otimes \mathcal{W}$  the universal functor preserving small colimits component-wise. By the adjoint functor theorem the canonical embedding

 $\alpha^*: \mathcal{V} \otimes \mathcal{W} \subset \widehat{\mathcal{P}}(\mathcal{V} \otimes \mathcal{W}) \to \widehat{\mathcal{P}}(\mathcal{V} \times \mathcal{W}) \simeq \operatorname{Fun}(\mathcal{V}^{\operatorname{op}}, \widehat{\mathcal{P}}(\mathcal{W}))$ 

induces an equivalence  $\mathcal{V} \otimes \mathcal{W} \to \operatorname{Fun}^{R}(\mathcal{V}^{\operatorname{op}}, \mathcal{W})$ . Under this equivalence the left morphism object  $\operatorname{LMor}_{\mathcal{M}}(X, Y)$  of  $X, Y \in \mathcal{M}$  is an object of  $\mathcal{V} \otimes \mathcal{W}$ , whose image under the embedding

$$\alpha^*: \mathcal{V} \otimes \mathcal{W} \subset \widehat{\mathcal{P}}(\mathcal{V} \otimes \mathcal{W}) \to \widehat{\mathcal{P}}(\mathcal{V} \times \mathcal{W})$$

is  $Mul_{\mathcal{M}}(-, X, -; Y)$ . In particular, for every  $V \in \mathcal{V}, W \in \mathcal{W}$  there is a canonical equivalence

(2)  $\mathcal{V} \otimes \mathcal{W}(\alpha(V, W), \operatorname{LMor}_{\mathcal{M}}(X, Y)) \simeq \mathcal{M}(V \otimes X \otimes W, Y) \simeq \operatorname{Mul}_{\mathcal{M}}(V, X, W; Y).$ 

**Example 2.118.** Every presentably bitensored  $\infty$ -category  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  exhibits  $\mathcal{M}$  as bi-enriched in  $\mathcal{V}, \mathcal{W}$ . By Example 2.61 for any  $X \in \mathcal{M}$  there is a unique left adjoint  $\mathcal{V}, \mathcal{W}$ -linear functor  $\mathcal{V} \otimes \mathcal{W} \to \mathcal{M}$ sending the tensor unit to X. The right adjoint sends  $Y \in \mathcal{M}$  to a left morphism object  $\operatorname{LMor}_{\mathcal{M}}(X, Y) \in$  $\mathcal{V} \otimes \mathcal{W}$  by equivalence (2). In particular, for every weakly bi-enriched  $\infty$ -category  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$ and every  $X, Y \in \mathcal{M}$  the object  $\Gamma_{\mathcal{M}}(X, Y) \in \mathcal{P}(\operatorname{Env}(\mathcal{V}) \times \operatorname{Env}(\mathcal{W})) \cong \mathcal{P}\operatorname{Env}(\mathcal{V}) \otimes \mathcal{P}\operatorname{Env}(\mathcal{W})$  is the left morphism object  $\operatorname{LMor}_{\overline{\mathcal{M}}}(X, Y)$  of X and Y, where  $\overline{\mathcal{M}^{\circledast}} \subset \mathcal{P}\operatorname{BEnv}(\mathcal{M})^{\circledast} \to \mathcal{P}\operatorname{Env}(\mathcal{V})^{\otimes} \times \mathcal{P}\operatorname{Env}(\mathcal{W})^{\circledast}$  is the full weakly bi-enriched subcategory spanned by  $\mathcal{M}$ .

Moreover we use the following terminology:

**Definition 2.119.** Let  $\kappa, \tau$  be regular cardinals and  $\phi : \mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a weakly bi-enriched  $\infty$ -category.

- (1) We say that  $\phi$  exhibits  $\mathcal{M}$  as left  $\kappa$ -enriched in  $\mathcal{V}$  if  $\mathcal{V}^{\otimes} \to Ass$  is a monoidal  $\infty$ -category compatible with  $\kappa$ -small colimits,  $\phi$  is a left pseudo-enriched  $\infty$ -category and for every  $X, Y \in \mathcal{M}$  and  $W_1, ..., W_m \in \mathcal{W}$  for  $m \ge 0$  the presheaf  $Mul_{\mathcal{M}}(-, X, W_1, ..., W_m; Y)$  on  $\mathcal{M}$  preserves  $\kappa$ -small limits.
- (2) We say that  $\phi$  exhibits  $\mathcal{M}$  as right  $\tau$ -enriched in  $\mathcal{W}$  if  $\mathcal{W}^{\otimes} \to Ass$  is a monoidal  $\infty$ -category compatible with  $\tau$ -small colimits,  $\phi$  is a right pseudo-enriched  $\infty$ -category and for every X, Y  $\in \mathcal{M}$  and  $V_1, ..., V_n \in \mathcal{V}$  for  $n \ge 0$  the presheaf  $Mul_{\mathcal{M}}(V_1, ..., V_n, X, -; Y)$  on  $\mathcal{M}$  preserves  $\tau$ -small limits.
- (3) We say that  $\phi$  exhibits  $\mathcal{M}$  as  $\kappa, \tau$ -bi-enriched in  $\mathcal{V}, \mathcal{W}$  if it exhibits  $\mathcal{M}$  as left  $\kappa$ -enriched in  $\mathcal{V}$  and right  $\tau$ -enriched in  $\mathcal{W}$ .

Notation 2.120. Let  $\kappa, \tau$  be small regular cardinals. Let

#### <sup> $\kappa$ </sup>LEnr, REnr<sup> $\tau$ </sup>, <sup> $\kappa$ </sup>BEnr<sup> $\tau$ </sup> $\subset \omega$ BEnr

be the full subcategories of left  $\kappa$ -enriched, right  $\tau$ -enriched,  $\kappa, \tau$ -bi-enriched  $\infty$ -categories, respectively.

**Example 2.121.** A weakly bi-enriched  $\infty$ -category  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  exhibits  $\mathcal{M}$  as left (right)  $\emptyset$ -enriched in  $\mathcal{V}$  if and only if it exhibits  $\mathcal{M}$  as left (right) pseudo-enriched.

**Definition 2.122.** Let  $\sigma$  be the strongly inaccessible cardinal corresponding to the small universe and  $\phi : \mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  a weakly bi-enriched  $\infty$ -category.

- (1) We say that  $\phi$  exhibits  $\mathcal{M}$  as left quasi-enriched in  $\mathcal{V}$  if  $\phi$  exhibits  $\mathcal{M}$  as left  $\sigma$ -enriched.
- (2) We say that  $\phi$  exhibits  $\mathcal{M}$  as right quasi-enriched in  $\mathcal{W}$  if  $\phi$  exhibits  $\mathcal{M}$  as right  $\sigma$ -enriched.
- (3) We say that  $\phi$  exhibits  $\mathcal{M}$  as bi-quasi-enriched in  $\mathcal{V}, \mathcal{W}$  if  $\phi$  exhibits  $\mathcal{M}$  as  $\sigma, \sigma$ -bi-enriched.

#### Notation 2.123. Let

## LQEnr, RQEnr, BQEnr $\subset \omega \widehat{BEnr}$

be the full subcategories of left quasi-enriched, right quasi-enriched, bi-quasi-enriched  $\infty$ -categories, respectively, whose underlying  $\infty$ -category is small.

The next theorem is [9, Theorem 5.50.]:

**Theorem 2.124.** Let  $\kappa$  be a small regular cardinal,

- (1) Let  $\mathcal{V}^{\otimes} \to \operatorname{Ass}$  be a small  $\infty$ -operad,  $\mathcal{W}^{\otimes} \to \operatorname{Ass}$  a small monoidal  $\infty$ -category compatible with  $\kappa$ -small colimits,  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes}$  a small weakly left enriched  $\infty$ -category and  $\mathcal{N}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a small right  $\kappa$ -enriched  $\infty$ -category. Then  $\operatorname{EnrFun}_{\mathcal{V}, \emptyset}(\mathcal{M}, \mathcal{N})^{\otimes} \to \mathcal{W}^{\otimes}$  is right  $\kappa$ -enriched.
- (2) Let V<sup>⊗</sup> → Ass be an ∞-operad, M<sup>®</sup> → V<sup>⊗</sup> a small weakly left enriched ∞-category, N<sup>®</sup> → V<sup>⊗</sup> × W<sup>⊗</sup> a right enriched ∞-category and W<sup>⊗</sup> → Ass an ∞-operad that exhibits W as right enriched in W such that W admits small limits and for every W ∈ W the functor RMor<sub>W</sub>(W, -) : W → W preserves small limits. Then EnrFun<sub>V,Ø</sub>(M, N)<sup>®</sup> → W<sup>⊗</sup> is a right enriched ∞-category.

**Remark 2.125.** Let  $\mathcal{V}^{\otimes} \to Ass, \mathcal{W}^{\otimes} \to Ass$  be presentably monoidal  $\infty$ -categories. An  $\infty$ -category left quasi-enriched in  $\mathcal{V}$  is an  $\infty$ -category left enriched in  $\mathcal{V}$  if and only if it is locally small. An  $\infty$ -category right quasi-enriched in  $\mathcal{W}$  is an  $\infty$ -category right enriched in  $\mathcal{W}$  if and only if it is locally small.

Notation 2.126. Let  $\mathcal{M}^{\circledast} \to \mathcal{U}^{\circledast} \times \mathcal{V}^{\circledast}$ ,  $\mathcal{N}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$ ,  $\mathcal{O}^{\circledast} \to \mathcal{W}^{\circledast} \times \mathcal{Q}^{\circledast}$  be presentably bitensored  $\infty$ -categories. We write  $(\mathcal{M} \otimes_{\mathcal{V}} \mathcal{N} \otimes_{\mathcal{W}} \mathcal{O})^{\circledast} \to \mathcal{U}^{\circledast} \times \mathcal{Q}^{\circledast}$  for the relative tensor product of [16, Definition 4.4.2.10.].

The next theorem is [9, Theorem 4.60.]:

**Theorem 2.127.** The forgetful functor  $\gamma : \omega BEnr \to Op_{\infty} \times Op_{\infty}$  is a cocartesian fibration. Let  $\psi : \mathcal{M}^{\otimes} \to \mathcal{N}^{\otimes}$  be an enriched functor lying over maps of  $\infty$ -operads  $\alpha : \mathcal{V}^{\otimes} \to \mathcal{V}'^{\otimes}, \alpha' : \mathcal{W}^{\otimes} \to \mathcal{W}'^{\otimes}$ . The following conditions are equivalent:

- (1) The enriched functor  $\psi : \mathcal{M}^{\otimes} \to \mathcal{N}^{\otimes}$  is  $\gamma$ -cocartesian.
- (2) The underlying functor  $\mathcal{M} \to \mathcal{N}$  is essentially surjective and  $\mathcal{PBEnv}(\mathcal{M})^{\circledast} \to \mathcal{PBEnv}(\mathcal{N})^{\circledast}$  induces an equivalence

 $\mathcal{P}BEnv(\mathcal{N}) \simeq \mathcal{P}Env(\mathcal{V}') \otimes_{\mathcal{P}Env(\mathcal{V})} \mathcal{P}BEnv(\mathcal{M}) \otimes_{\mathcal{P}Env(\mathcal{W})} \mathcal{P}Env(\mathcal{W}').$ 

(3) The functor  $\mathcal{M} \to \mathcal{N}$  is essentially surjective and for every  $X, Y \in \mathcal{M}$  the induced morphism

$$(\alpha,\beta)_!(\Gamma_{\mathcal{M}}(\mathbf{X},\mathbf{Y})) \to \Gamma_{\mathcal{N}}(\psi(\mathbf{X}),\psi(\mathbf{Y}))$$

in  $\mathcal{P}(\operatorname{BEnv}(\mathcal{V}') \times \operatorname{BEnv}(\mathcal{W}'))$  is an equivalence.

The next proposition is [9, Proposition 5.45.]:

**Proposition 2.128.** There is an involution

$$(-)^{\mathrm{op}}: {}_{\mathcal{V}}\omega\mathrm{BEnr}_{\mathcal{W}} \simeq {}_{\mathcal{W}}\omega\mathrm{BEnr}_{\mathcal{V}}$$

fitting into a commutative square



such that for every weakly bi-enriched  $\infty$ -category  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  and  $X, Y \in \mathcal{M}$  there is an equivalence

$$\Gamma_{\mathcal{M}^{\mathrm{op}}}(\mathbf{X},\mathbf{Y}) \simeq \Gamma_{\mathcal{M}}(\mathbf{Y},\mathbf{X}).$$

2.6. Generalized enrichment. The next definition is [9, Definition 4.1.]:

**Definition 2.129.** A small localization pair is a pair ( $\mathcal{V}^{\otimes} \to Ass, S$ ), where  $\mathcal{V}^{\otimes} \to Ass$  is a small  $\infty$ -operad and S is a set of morphisms of  $\mathcal{P}Env(\mathcal{V})$  such that the saturated closure  $\bar{S}$  of S is closed under the tensor product and for every  $V \in \mathcal{V}$  and  $f \in S$  the map  $\mathcal{P}Env(\mathcal{V})(f, V)$  is an equivalence.

**Remark 2.130.** Let  $(\mathcal{V}^{\otimes} \to Ass, S)$  be a small localization pair. Since  $\mathcal{P}Env(\mathcal{V})^{\otimes} \to Ass$  is a presentably monoidal  $\infty$ -category, S is a set and  $\overline{S}$  is closed under the tensor product, the embedding  $S^{-1}\mathcal{P}Env(\mathcal{V})^{\otimes} \subset \mathcal{P}Env(\mathcal{V})^{\otimes}$  of the full suboperad spanned by the S-local objects admits a left adjoint relative to Ass. Moreover by definition the embedding of  $\infty$ -operads  $\mathcal{V}^{\otimes} \subset \mathcal{P}Env(\mathcal{V})^{\otimes}$  induces an embedding of  $\infty$ -operads  $\mathcal{V}^{\otimes} \subset S^{-1}\mathcal{P}Env(\mathcal{V})^{\otimes}$ .

The next definition is [9, Definition 4.3.]:

**Definition 2.131.** Let  $(\mathcal{V}^{\otimes} \to Ass, S)$ ,  $(\mathcal{W}^{\otimes} \to Ass, T)$  be small localization pairs.

- (1) A weakly bi-enriched  $\infty$ -category  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  exhibits  $\mathcal{M}$  as left S-enriched in  $\mathcal{V}$  if for every  $f \in S, X, Y \in \mathcal{M}, W \in \mathcal{P}Env(\mathcal{W})$  the induced map  $\mathcal{P}BEnv(\mathcal{M})(f \otimes X \otimes W, Y)$  is an equivalence.
- (2) A weakly bi-enriched  $\infty$ -category  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  exhibits  $\mathcal{M}$  as right T-enriched in  $\mathcal{W}$  if for every  $g \in T, X, Y \in \mathcal{M}, V \in \mathcal{P}Env(\mathcal{V})$  the map  $\mathcal{P}BEnv(\mathcal{M})(V \otimes X \otimes g, Y)$  is an equivalence.
- (3) A weakly bi-enriched  $\infty$ -category  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  exhibits  $\mathcal{M}$  as S, T-bi-enriched in  $\mathcal{V}, \mathcal{W}$  if it exhibits  $\mathcal{M}$  as left S-enriched in  $\mathcal{V}$  and right T-enriched in  $\mathcal{W}$ .

Notation 2.132. Let  $(\mathcal{V}^{\otimes} \to Ass, S)$ ,  $(\mathcal{W}^{\otimes} \to Ass, T)$  be small localization pairs. Let

# <sup>S</sup><sub>v</sub>LEnr<sub>w</sub>, $_{v}$ REnr<sup>T</sup><sub>w</sub>, <sup>S</sup><sub>v</sub>BEnr<sup>T</sup><sub>w</sub> $\subset _{v}\omega$ BEnr<sub>w</sub>

the full subcategories of left S-enriched, right T-enriched, S,T-bi-enriched ∞-categories, respectively.

**Example 2.133.** Let  $\mathcal{U}^{\otimes} \to Ass$  be a small  $\infty$ -operad. The pair  $(\mathcal{U}^{\otimes} \to Ass, \emptyset)$  is a small localization pair, where  $\emptyset^{-1} \mathcal{P} Env(\mathcal{U})^{\otimes} = \mathcal{P} Env(\mathcal{U})^{\otimes}$ . Let  $(\mathcal{V}^{\otimes} \to Ass, S)$ ,  $(\mathcal{W}^{\otimes} \to Ass, T)$  be small localization pairs. Then

$${}_{\mathcal{V}}^{S}LEnr_{\mathcal{W}} = {}_{\mathcal{V}}^{S}BEnr_{\mathcal{W}}^{\varnothing}, \ {}_{\mathcal{V}}REnr_{\mathcal{W}}^{T} = {}_{\mathcal{V}}^{\varnothing}BEnr_{\mathcal{W}}^{T}.$$

**Notation 2.134.** Let  $\kappa$  be a small regular cardinal and  $\mathcal{V}^{\otimes} \to Ass$  a small monoidal  $\infty$ -category compatible with  $\kappa$ -small colimits. Let  $\operatorname{Enr}_{\mathcal{V}}^{\kappa}$  be the set of morphisms

$$V_1',...,V_\ell',V_1,...,V_n,V_1'',...,V_k'' \to V_1',...,V_\ell',V_1 \otimes ... \otimes V_n,V_1'',...,V_k'',$$

 $V'_1, ..., V'_{\ell}, \operatorname{colim}(\iota \circ F), V''_1, ..., V''_k \to V'_1, ..., V'_{\ell}, \iota(\operatorname{colim}(F)), V''_1, ..., V''_k$ for  $V'_1, ..., V'_{\ell}, V_1, ..., V_n, V''_1, ..., V''_k \in \mathcal{V}$  and  $\ell, n, k \ge 0$ , where  $\iota : \mathcal{V} \subset \mathcal{P}\operatorname{Env}(\mathcal{V})$  is the embedding,  $F: K \to \mathcal{V}$  is a functor and K is  $\kappa$ -small.

Notation 2.135. Let  $\kappa, \tau$  be small regular cardinals.

- (1) If  $\mathcal{V}^{\otimes} \to Ass$  is a small monoidal  $\infty$ -category compatible with  $\kappa$ -small colimits and  $\mathcal{W}^{\otimes} \to Ass$  is a small  $\infty$ -operad, let  $\operatorname{LEnr}_{\kappa} := (\operatorname{Enr}_{\mathcal{V}}^{\kappa}, \emptyset)$ .
- (2) If  $\mathcal{V}^{\otimes} \to Ass$  is a small  $\infty$ -operad and  $\mathcal{W}^{\otimes} \to Ass$  is a small monoidal  $\infty$ -category compatible with  $\kappa$ -small colimits, let  $\operatorname{REnr}_{\kappa} := (\emptyset, \operatorname{Enr}_{\mathcal{W}}^{\kappa})$ .
- (3) If  $\mathcal{V}^{\otimes} \to \operatorname{Ass}, \mathcal{W}^{\otimes} \to \operatorname{Ass}$  are small monoidal  $\infty$ -categories compatible with  $\kappa$ -small colimits,  $\tau$ -small colimits, respectively, let  $\operatorname{BEnr}_{\kappa,\tau} := (\operatorname{Enr}_{\mathcal{V}}^{\kappa}, \operatorname{Enr}_{\mathcal{W}}^{\tau})$ . If  $\kappa, \tau$  are the the strongly inaccessible cardinal corresponding to the small universe, we drop  $\kappa, \tau$ , respectively.

**Example 2.136.** Let  $\kappa$  be a small regular cardinal.

- (1) For every small monoidal  $\infty$ -category  $\mathcal{V}^{\otimes} \to Ass$  compatible with  $\kappa$ -small colimits the pair  $(\mathcal{V}^{\otimes} \to Ass, \operatorname{Enr}_{\mathcal{V}}^{\kappa})$  is a small localization pair, where  $(\operatorname{Enr}_{\mathcal{V}}^{\kappa})^{-1}\mathcal{P}\operatorname{Env}(\mathcal{V})^{\otimes} = \operatorname{Ind}_{\kappa}(\mathcal{V})^{\otimes}$ , which is a monoidal localization of  $\mathcal{P}\operatorname{Env}(\mathcal{V})^{\otimes} \to Ass$ . A weakly bi-enriched  $\infty$ -category  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  exhibits  $\mathcal{M}$  as left  $\operatorname{Enr}_{\mathcal{V}}^{\kappa}$ -enriched in  $\mathcal{V}$  if and only if it exhibits  $\mathcal{M}$  as left  $\kappa$ -enriched in  $\mathcal{V}$ .
- (2) If  $\mathcal{V}^{\otimes} \to Ass$  is a presentably monoidal  $\infty$ -category, a locally small weakly bi-enriched  $\infty$ -category  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  exhibits  $\mathcal{M}$  as left  $Enr_{\mathcal{V}}$ -enriched in  $\mathcal{V}$  if and only if it exhibits  $\mathcal{M}$  as left enriched in  $\mathcal{V}$ .

**Definition 2.137.** Let  $(\mathcal{V}^{\otimes} \to Ass, S)$ ,  $(\mathcal{V}'^{\otimes} \to Ass, S')$  be small localization pairs. A map of localization pairs  $(\mathcal{V}^{\otimes} \to Ass, S) \to (\mathcal{V}'^{\otimes} \to Ass, S')$  is a map of  $\infty$ -operads  $\mathcal{V}^{\otimes} \to \mathcal{V}'^{\otimes}$  such that the induced left adjoint monoidal functor  $\mathcal{P}Env(\mathcal{V}) \to \mathcal{P}Env(\mathcal{V}')$  sends morphisms of S to morphisms of S'.

Notation 2.138. Let  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}, \mathcal{N}^{\otimes} \to \mathcal{V}'^{\otimes} \times \mathcal{W}'^{\otimes}$  be weakly bi-enriched  $\infty$ -categories, S, S', T, T' collections of morphisms of  $\mathcal{P}Env(\mathcal{V}), \mathcal{P}Env(\mathcal{W}), \mathcal{P}Env(\mathcal{W})$ , respectively. Let

 $\operatorname{EnrFun}_{S,S',T,T'}(\mathcal{M},\mathcal{N}) \subset \operatorname{EnrFun}(\mathcal{M},\mathcal{N})$ 

be the full subcategory of enriched functors lying over maps of  $\infty$ -operads whose induced left adjoint monoidal functors on closed monoidal envelopes send S to S' and T to T', respectively.

**Example 2.139.** Let  $\mathcal{V}^{\otimes} \to \operatorname{Ass}, \mathcal{V}'^{\otimes} \to \operatorname{Ass}$  be small monoidal  $\infty$ -categories compatible with  $\kappa$ -small colimits, compatible with  $\kappa'$ -small colimits, respectively, for small regular cardinals  $\kappa < \kappa'$ . A map of  $\infty$ -operade  $\mathcal{V}^{\otimes} \to \mathcal{V}'^{\otimes}$  is a map of localization pairs  $(\mathcal{V}^{\otimes} \to \operatorname{Ass}, \operatorname{Enr}_{\mathcal{V}}^{\kappa}) \to (\mathcal{V}'^{\otimes} \to \operatorname{Ass}, \operatorname{Enr}_{\mathcal{V}'}^{\kappa'})$  if and only if it is a monoidal functor preserving  $\kappa$ -small colimits.

The next proposition is [9, Proposition 4.29., Corollary 4.30.]:

**Proposition 2.140.** Let  $(\mathcal{V}^{\otimes} \to Ass, S), (\mathcal{W}^{\otimes} \to Ass, T)$  be small localization pairs.

(1) The functor

 $_{\mathrm{S}^{-1}\mathcal{P}\mathrm{Env}(\mathcal{V})}\omega\mathrm{B}\mathrm{Enr}_{\mathcal{W}} \to \mathcal{V}\omega\mathrm{B}\mathrm{Enr}_{\mathcal{W}},$ 

which takes pullback along the embedding  $\mathcal{V}^{\otimes} \subset S^{-1}\mathcal{P}Env(\mathcal{V})^{\otimes}$ , restricts to an equivalence

 $_{\mathrm{S}^{-1}\mathcal{P}\mathrm{Env}(\mathcal{V})}\mathrm{LEnr}_{\mathcal{W}} \rightarrow {}_{\mathcal{V}}^{\mathrm{S}}\mathrm{LEnr}_{\mathcal{W}}.$ 

(2) The functor

 $_{\mathcal{V}}\omega \operatorname{BEnr}_{\operatorname{T}^{-1}\mathcal{P}\operatorname{Env}(\mathcal{W})} \rightarrow _{\mathcal{V}}\omega \operatorname{BEnr}_{\mathcal{W}},$ 

which takes pullback along the embedding  $\mathcal{W}^{\otimes} \subset T^{-1}\mathcal{P}Env(\mathcal{W})^{\otimes}$ , restricts to an equivalence

$$\gamma \operatorname{REnr}_{T^{-1} \mathcal{P} \operatorname{Env}(\mathcal{W})} \rightarrow \gamma \operatorname{REnr}_{\mathcal{W}}^{T}$$

(3) The functor

 $\rho:_{\mathrm{S}^{-1}\mathcal{P}\mathrm{Env}(\mathcal{V})}\omega\mathrm{B}\mathrm{Enr}_{\mathrm{T}^{-1}\mathcal{P}\mathrm{Env}(\mathcal{W})} \to \mathcal{V}\omega\mathrm{B}\mathrm{Enr}_{\mathcal{W}},$ 

which takes pullback along the embeddings  $\mathcal{V}^{\otimes} \subset S^{-1}\mathcal{P}Env(\mathcal{V})^{\otimes}, \mathcal{W}^{\otimes} \subset T^{-1}\mathcal{P}Env(\mathcal{W})^{\otimes}, restricts$ to an equivalence

 $_{\mathrm{S}^{-1}\mathcal{P}\mathrm{Env}(\mathcal{V})}\mathrm{B}\mathrm{Enr}_{\mathrm{T}^{-1}\mathcal{P}\mathrm{Env}(\mathcal{W})} \rightarrow {}_{\mathcal{V}}^{\mathrm{S}}\mathrm{B}\mathrm{Enr}_{\mathcal{W}}^{\mathrm{T}}.$ 

Proposition 2.140 gives the following corollary:

**Corollary 2.141.** Let  $\mathcal{V}^{\otimes} \to Ass, \mathcal{W}^{\otimes} \to Ass$  be small  $\infty$ -operads.

(1) The functor

$$\mathcal{P}_{\mathrm{Env}(\mathcal{V})}\mathrm{LEnr}_{\mathcal{W}} \to \mathcal{V}\omega\mathrm{BEnr}_{\mathcal{W}}$$

taking pullback along the embedding  $\mathcal{V}^{\otimes} \subset \operatorname{PEnv}(\mathcal{V})^{\otimes}$  is an equivalence.

(2) The functor

 $_{\mathcal{V}}\operatorname{REnr}_{\mathcal{P}\operatorname{Env}(\mathcal{W})} \rightarrow _{\mathcal{V}}\omega\operatorname{BEnr}_{\mathcal{W}}$ 

taking pullback along the embedding  $\mathcal{W}^{\otimes} \subset \mathcal{P}Env(\mathcal{W})^{\otimes}$  is an equivalence.

(3) The functor

 $\mathcal{P}_{Env}(\mathcal{V})BEnr_{\mathcal{P}Env}(\mathcal{W}) \rightarrow \mathcal{V}\omega BEnr_{\mathcal{W}}$ 

taking pullback along the embeddings  $\mathcal{V}^{\otimes} \subset \mathcal{P}Env(\mathcal{V})^{\otimes}, \mathcal{W}^{\otimes} \subset \mathcal{P}Env(\mathcal{W})^{\otimes}$  is an equivalence.

**Corollary 2.142.** Let  $\kappa, \tau$  be small regular cardinals.

(1) Let  $\mathcal{V}^{\otimes} \to Ass$  be a small monoidal  $\infty$ -category compatible with  $\kappa$ -small colimits and  $\mathcal{W}^{\otimes} \to Ass$ a small  $\infty$ -operad. The functor

$$\operatorname{Ind}_{\kappa}(\mathcal{V})\operatorname{LEnr}_{\mathcal{W}} \to \mathcal{V}^{\kappa}\operatorname{LEnr}_{\mathcal{W}}$$

taking pullback along the monoidal embedding  $\mathcal{V}^{\otimes} \subset \mathrm{Ind}_{\kappa}(\mathcal{V})^{\otimes}$  is an equivalence.

(2) Let  $W^{\otimes} \to Ass$  be a small monoidal  $\infty$ -category compatible with  $\tau$ -small colimits and  $\mathcal{V}^{\otimes} \to Ass$  a small  $\infty$ -operad. The functor

$$v \operatorname{REnr}_{\operatorname{Ind}_{\tau}(W)} \rightarrow v \operatorname{REnr}_{W}^{\tau}$$

taking pullback along the monoidal embedding  $\mathcal{W}^{\otimes} \subset \operatorname{Ind}_{\tau}(\mathcal{W})^{\otimes}$  is an equivalence.

(3) Let  $\mathcal{V}^{\otimes} \to \operatorname{Ass}, \mathcal{W}^{\otimes} \to \operatorname{Ass}$  be small monoidal  $\infty$ -categories compatible with  $\kappa$ -small colimits,  $\tau$ -small colimits, respectively. The functor

$$\operatorname{Ind}_{\kappa}(\mathcal{V})\operatorname{BEnr}_{\operatorname{Ind}_{\kappa}(\mathcal{W})} \to \mathcal{V}\operatorname{BEnr}_{\mathcal{W}}^{\prime}$$

restricting along the monoidal embeddings  $\mathcal{V}^{\otimes} \subset \operatorname{Ind}_{\kappa}(\mathcal{V})^{\otimes}, \mathcal{W}^{\otimes} \subset \operatorname{Ind}_{\tau}(\mathcal{W})^{\otimes}$  is an equivalence.

The next notation is [9, Notation 4.14.]:

Notation 2.143. Let  $(\mathcal{V}^{\otimes} \to Ass, S), (\mathcal{W}^{\otimes} \to Ass, T)$  be small localization pairs (Definition 2.129) and  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a totally small weakly bi-enriched  $\infty$ -category.

(1) Let

$$\mathcal{P}BEnv(\mathcal{M})^{\otimes}_{S T} \subset \mathcal{P}BEnv(\mathcal{M})^{\otimes}$$

be the full weakly bi-enriched subcategory spanned by the presheaves on BEnv( $\mathcal{M}$ ) that are local with respect to the set of morphisms of the form  $f \otimes X \otimes W_1 \dots \otimes W_m$  and  $V_1 \otimes \dots \otimes V_n \otimes X \otimes g$  for  $f \in S, g \in T, X \in \mathcal{M}, V_1, \dots, V_n \in \mathcal{V}, W_1, \dots, W_m \in \mathcal{W}$  for  $n, m \ge 0$ .

(2) Let

$$\mathcal{P}\widetilde{\operatorname{BEnv}}(\mathcal{M})^{\otimes}_{S^{\mathsf{T}}} \coloneqq \mathcal{V}^{\otimes} \times_{S^{-1}\mathcal{P}\operatorname{Env}}(\mathcal{V})^{\otimes} \mathcal{P}\operatorname{BEnv}(\mathcal{M})^{\otimes}_{S^{\mathsf{T}}} \times_{T^{-1}\mathcal{P}\operatorname{Env}}(\mathcal{W})^{\otimes} \mathcal{W}^{\otimes}.$$

If  $S = \emptyset$  or  $T = \emptyset$ , we drop S, T from the notation, respectively.

Remark 2.144. As a consequence of Remark 2.99 there is a canonical equivalence

$$(\mathcal{P}BEnv(\mathcal{M})^{rev}_{S,T})^{\circledast} \simeq \mathcal{P}BEnv(\mathcal{M}^{rev})^{\circledast}_{T,S}$$

**Lemma 2.145.** Let  $(\mathcal{V}^{\otimes} \to \operatorname{Ass}, \operatorname{S}), (\mathcal{W}^{\otimes} \to \operatorname{Ass}, \operatorname{T})$  be small localization pairs and  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a small weakly bi-enriched  $\infty$ -category. The embedding  $\operatorname{PBEnv}(\mathcal{M})^{\otimes}_{\operatorname{S},\operatorname{T}} \subset \operatorname{PBEnv}(\mathcal{M})^{\otimes}$  admits an enriched left adjoint such that the unit lies over the units of the localizations  $\operatorname{S}^{-1}\operatorname{PEnv}(\mathcal{V})^{\otimes} \subset \operatorname{PEnv}(\mathcal{V})^{\otimes}, \operatorname{T}^{-1}\operatorname{PEnv}(\mathcal{W})^{\otimes} \subset \operatorname{PEnv}(\mathcal{W})^{\otimes}$ . In particular,  $\operatorname{PBEnv}(\mathcal{M})^{\otimes}_{\operatorname{S},\operatorname{T}} \to \operatorname{S}^{-1}\operatorname{PEnv}(\mathcal{V})^{\otimes} \times \operatorname{T}^{-1}\operatorname{PEnv}(\mathcal{W})^{\otimes}$  is a presentably bitensored  $\infty$ -category.

**Remark 2.146.** Since  $\mathcal{P}BEnv(\mathcal{M})$  is presentable, the embedding  $\mathcal{P}BEnv(\mathcal{M})_{S,T} \subset \mathcal{P}BEnv(\mathcal{M})$  admits a left adjoint and  $\mathcal{P}BEnv(\mathcal{M})_{S,T} \subset \mathcal{P}BEnv(\mathcal{M})$  precisely consists of the presheaves on  $BEnv(\mathcal{M})$  that are local with respect to the collection  $\Omega$  of morphisms of the form  $f \otimes X \otimes g$  for  $f \in \overline{S}, g \in \overline{T}$  and  $X \in \mathcal{M}$ . The biaction of  $\mathcal{P}Env(\mathcal{V}), \mathcal{P}Env(\mathcal{W})$  on  $\mathcal{P}BEnv(\mathcal{M})$  sends morphisms of  $\overline{S}, \Omega, \overline{T}$  to  $\Omega$ . This implies that the embedding  $\mathcal{P}BEnv(\mathcal{M})^{\mathfrak{B}}_{S,T} \subset \mathcal{P}BEnv(\mathcal{M})^{\mathfrak{B}}$  admits an enriched left adjoint such that the unit lies over the units of the localizations  $S^{-1}\mathcal{P}Env(\mathcal{V})^{\mathfrak{S}} \subset \mathcal{P}Env(\mathcal{V})^{\mathfrak{S}}, T^{-1}\mathcal{P}Env(\mathcal{W})^{\mathfrak{S}} \subset \mathcal{P}Env(\mathcal{W})^{\mathfrak{S}}$ .

**Remark 2.147.** A weakly bi-enriched  $\infty$ -category  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  is a S, T-bi-enriched  $\infty$ -category if and only if  $\mathcal{M}^{\circledast} \subset \mathcal{PBEnv}(\mathcal{M})^{\circledast}$  lands in  $\mathcal{PBEnv}(\mathcal{M})^{\circledast}_{S,T}$ .

**Remark 2.148.** Let  $\mathcal{M}^{\circledast} \to \mathcal{N}^{\circledast}$  be an enriched functor between totally small weakly bi-enriched  $\infty$ -categories lying over maps of small localization pairs  $(\mathcal{V}^{\circledast} \to Ass, S) \to (\mathcal{V}'^{\circledast} \to Ass, S'), (\mathcal{W}^{\circledast} \to Ass, T) \to (\mathcal{W}'^{\circledast} \to Ass, T')$ . The induced left adjoint map of bitensored  $\infty$ -categories  $\mathcal{PBEnv}(\mathcal{M})^{\circledast} \to \mathcal{PBEnv}(\mathcal{N})^{\circledast}$  preserves local equivalences for the corresponding localizations and so descends to a left adjoint map of bitensored  $\infty$ -categories  $\mathcal{PBEnv}(\mathcal{M})^{\circledast}_{S,T} \to \mathcal{PBEnv}(\mathcal{N})^{\circledast}_{S',T'}$ .

The next proposition is [9, Proposition 4.20.]:

**Proposition 2.149.** Let  $(\mathcal{V}^{\otimes} \to \operatorname{Ass}, S), (\mathcal{W}^{\otimes} \to \operatorname{Ass}, T)$  be small localization pairs,  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$ a small S, T-bi-enriched  $\infty$ -category and  $\rho : \mathcal{N}^{\otimes} \to S^{-1} \mathcal{P} \operatorname{Env}(\mathcal{V})^{\otimes} \times T^{-1} \mathcal{P} \operatorname{Env}(\mathcal{W})^{\otimes}$  a bitensored  $\infty$ -category compatible with small colimits.

(1) The enriched functor  $\mathcal{M}^{\otimes} \subset \mathcal{PBEnv}(\mathcal{M})^{\otimes} \to \mathcal{PBEnv}(\mathcal{M})_{S,T}^{\otimes}$  induces a functor

 $\operatorname{EnrFun}_{S^{-1}\mathcal{P}\operatorname{Env}(\mathcal{V}),T^{-1}\mathcal{P}\operatorname{Env}(\mathcal{W})}(\mathcal{P}\operatorname{B}\operatorname{Env}(\mathcal{M})_{S,T},\mathcal{N}) \to \operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}(\mathcal{M},\mathcal{N})$ 

that admits a fully faithful left adjoint. The left adjoint lands in the full subcategory

 $\operatorname{LinFun}_{S^{-1}\mathcal{P}\operatorname{Env}(\mathcal{V}),T^{-1}\mathcal{P}\operatorname{Env}(\mathcal{W})}^{L}(\mathcal{P}\operatorname{B}\operatorname{Env}(\mathcal{M})_{S,T},\mathcal{N}).$ 

(2) The following induced functor is an equivalence:

 $\operatorname{LinFun}_{S^{-1}\mathcal{P}\operatorname{Env}(\mathcal{V}), T^{-1}\mathcal{P}\operatorname{Env}(\mathcal{W})}(\mathcal{P}\operatorname{B}\operatorname{Env}(\mathcal{M})_{S,T}, \mathcal{N}) \to \operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}(\mathcal{M}, \mathcal{N}).$ 

**Notation 2.150.** Let  $\kappa$  be a small regular cardinal and  $\mathbb{C}$  a small  $\infty$ -category. Let  $\mathcal{P}_{\kappa}(\mathbb{C})^{\otimes} \subset \mathcal{P}(\mathbb{C})^{\otimes}$  be the full suboperad spanned by the full subcategory of  $\mathcal{P}(\mathbb{C})$  generated by  $\mathbb{C}$  under  $\kappa$ -small colimits.

The next notation is [9, Notation 4.24.]:

Notation 2.151. Let  $\kappa$  be a small regular cardinal.

(1) Let  $\mathcal{V}^{\otimes} \to Ass$  be a small monoidal  $\infty$ -category compatible with  $\kappa$ -small colimits,  $\mathcal{W}^{\otimes} \to Ass$  a small  $\infty$ -operad and  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a small left  $\kappa$ -enriched  $\infty$ -category. Let

$$\mathcal{B}Env_{\kappa}(\mathcal{M})^{\circledast}_{\operatorname{LEnr}} \subset \mathcal{V}^{\otimes} \times_{\operatorname{Ind}_{\kappa}(\mathcal{V})^{\otimes}} \mathcal{P}BEnv(\mathcal{M})^{\circledast}_{\operatorname{LEnr}} \times_{\mathcal{P}Env(\mathcal{W})^{\otimes}} \mathcal{P}_{\kappa}Env(\mathcal{W})^{\otimes}$$

be the full bitensored subcategory generated by  $\mathcal{M}$  under  $\kappa$ -small colimits.

(2) Let  $\mathcal{V}^{\otimes} \to Ass$  be a small  $\infty$ -operad,  $\mathcal{W}^{\otimes} \to Ass$  a small monoidal  $\infty$ -category compatible with  $\kappa$ -small colimits and  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a small right  $\kappa$ -enriched  $\infty$ -category. Let

 $\mathcal{P}BEnv_{\kappa}(\mathcal{M})^{\circledast}_{REnr} \subset \mathcal{P}_{\kappa}Env(\mathcal{V})^{\otimes} \times_{\mathcal{P}Env(\mathcal{V})^{\otimes}} \mathcal{P}BEnv(\mathcal{M})^{\circledast}_{REnr_{\kappa}} \times_{Ind_{\kappa}(\mathcal{W})^{\otimes}} \mathcal{W}^{\otimes}$ 

be the full bitensored subcategory generated by  $\mathcal{M}$  under  $\kappa$ -small colimits.

(3) Let  $\mathcal{V}^{\otimes} \to Ass, \mathcal{W}^{\otimes} \to Ass$  be small monoidal  $\infty$ -categories compatible with  $\kappa$ -small colimits and  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a small  $\kappa, \kappa$ -bi-enriched  $\infty$ -category. Let

 $\mathcal{P}\mathrm{BEnv}_{\kappa}(\mathcal{M})^{\circledast}_{\mathrm{BEnr}} \subset \mathcal{V}^{\otimes} \times_{\mathrm{Ind}_{\kappa}(\mathcal{V})^{\otimes}} \mathcal{P}\mathrm{BEnv}(\mathcal{M})^{\circledast}_{\mathrm{BEnr}_{\kappa}} \times_{\mathrm{Ind}_{\kappa}(\mathcal{W})^{\otimes}} \mathcal{W}^{\otimes}$ 

be the full bit ensored subcategory generated by  ${\mathcal M}$  under  $\kappa\mbox{-small colimits}.$ 

Next we apply Notation 2.151 in a larger universe:

Notation 2.152. Let  $\sigma$  be the large regular cardinal corresponding to the small universe.

(1) Let  $\mathcal{V}^{\otimes} \to Ass$  be a monoidal  $\infty$ -category compatible with small colimits,  $\mathcal{W}^{\otimes} \to Ass$  an  $\infty$ -operad and  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a left quasi-enriched  $\infty$ -category. Let

$$\mathcal{P}_{\mathcal{V}}(\mathcal{M})^{\otimes} \coloneqq \mathcal{P}BEnv(\mathcal{M})^{\otimes}_{LEnr} \coloneqq \mathcal{P}BEnv_{\sigma}(\mathcal{M})^{\otimes}_{LEnr} \to \mathcal{V}^{\otimes} \times \mathcal{P}Env(\mathcal{W})^{\otimes}$$

(2) Let  $\mathcal{V}^{\otimes} \to Ass$  be an  $\infty$ -operad,  $\mathcal{W}^{\otimes} \to Ass$  a monoidal  $\infty$ -category compatible with small colimits and  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a right quasi-enriched  $\infty$ -category. Let

$$\mathcal{P}BEnv(\mathcal{M})^{\mathfrak{G}}_{BEnr} := \mathcal{P}BEnv_{\sigma}(\mathcal{M})^{\mathfrak{G}}_{BEnr} \to \mathcal{P}Env(\mathcal{V})^{\mathfrak{S}} \times \mathcal{W}^{\mathfrak{S}}.$$

(3) Let  $\mathcal{V}^{\otimes} \to Ass, \mathcal{W}^{\otimes} \to Ass$  be monoidal  $\infty$ -categories compatible with small colimits and  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a bi-quasi-enriched  $\infty$ -category. Let

$$\mathcal{P}_{\mathcal{V},\mathcal{W}}(\mathcal{M})^{\circledast} \coloneqq \mathcal{P}BEnv(\mathcal{M})^{\circledast}_{BEnr} \coloneqq \mathcal{P}BEnv_{\sigma}(\mathcal{M})^{\circledast}_{BEnr} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}.$$

The next proposition is [9, Corollary 4.41.]:

 $\begin{array}{ll} \textbf{Proposition 2.153.} & (1) \ Let \ \mathcal{V}^{\otimes} \rightarrow Ass \ be \ a \ presentably \ monoidal \ \infty-category, \ \mathcal{W}^{\otimes} \rightarrow Ass \ a \ small \\ \infty \ -operad \ and \ \mathcal{M}^{\circledast} \rightarrow \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes} \ a \ small \ left \ enriched \ \infty \ -category. \ Then \end{array}$ 

 $\mathcal{P}_{\mathcal{V}}(\mathcal{M})^{\circledast} = \mathcal{P}BEnv(\mathcal{M})^{\circledast}_{LEnr} \to \mathcal{V}^{\otimes} \times \mathcal{P}Env(\mathcal{W})^{\otimes}$ 

is a presentably left tensored  $\infty$ -category.

(2) Let  $\mathcal{V}^{\otimes} \to Ass$  be a small  $\infty$ -operad,  $\mathcal{W}^{\otimes} \to Ass$  a presentably monoidal  $\infty$ -category and  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a small right enriched  $\infty$ -category. Then

$$\mathcal{P}BEnv(\mathcal{M})^{\otimes}_{BEnr} \to \mathcal{P}Env(\mathcal{V})^{\otimes} \times \mathcal{W}^{\otimes}$$

is a presentably right tensored  $\infty$ -category.

(3) Let  $\mathcal{V}^{\otimes} \to Ass, \mathcal{W}^{\otimes} \to Ass$  be presentably monoidal  $\infty$ -categories and  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a small bi-enriched  $\infty$ -category. Then

$$\mathcal{P}_{\mathcal{V},\mathcal{W}}(\mathcal{M})^{\circledast} \coloneqq \mathcal{P}BEnv(\mathcal{M})^{\circledast}_{BEnr} \coloneqq \mathcal{P}BEnv_{\sigma}(\mathcal{M})^{\circledast}_{BEnr} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$$

is a presentably bitensored  $\infty$ -category.

The next theorem is [9, 5.48.], which gives rise to the enriched Yoneda-embedding.

**Theorem 2.154.** Let  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes}$  be a left enriched  $\infty$ -category. There is a left  $\mathcal{V}$ -linear equivalence

 $|-|: \mathcal{P}_{\mathcal{V}}(\mathcal{M})^{\circledast} \simeq \operatorname{EnrFun}_{\emptyset,\mathcal{V}}(\mathcal{M}^{\operatorname{op}},\mathcal{V})^{\circledast}$ 

sending X to  $\mathrm{LMor}_{\mathcal{P}_{\mathcal{V}}(\mathcal{M})}((-)_{|\mathcal{M}}, X)$ . In particular, there is a left  $\mathcal{V}$ -enriched embedding

 $\mathcal{M}^{\circledast} \to \operatorname{EnrFun}_{\varnothing, \mathcal{V}}(\mathcal{M}^{\operatorname{op}}, \mathcal{V})^{\circledast}$ 

that sends X to  $LMor_{\mathcal{M}}(-, X)$ .

Corollary 2.141 gives the following corollary:

**Corollary 2.155.** Let  $\mathcal{V}^{\otimes} \to Ass$  be a small  $\infty$ -operad and  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes}$  a weakly left enriched  $\infty$ -category. There is a left  $\mathcal{P}Env(\mathcal{V})$ -linear equivalence

$$|-|: \mathcal{P}LEnv(\mathcal{M})^{\circledast} \simeq EnrFun_{\emptyset,\mathcal{V}}(\mathcal{M}^{op},\mathcal{P}Env(\mathcal{V}))^{\circledast}$$

sending X to  $\mathrm{LMor}_{\mathcal{P}\mathrm{LEnv}(\mathcal{M})}((-)|_{\mathcal{M}}, X)$ . In particular, there is a left  $\mathcal{P}\mathrm{Env}(\mathcal{V})$ -enriched embedding

$$\mathcal{M}^{\otimes} \to \operatorname{EnrFun}_{\varnothing,\mathcal{V}}(\mathcal{M}^{\operatorname{op}}, \operatorname{\mathcal{P}Env}(\mathcal{V}))^{\otimes}$$

that sends X to  $LMor_{\overline{M}}(-, X)$ .

#### 3. Weighted colimits

3.1. Weighted colimits for weakly enriched  $\infty$ -categories. In this section we define weighted colimits and study their properties. Weighted colimits are a generalization of conical colimits. For every weakly bi-enriched  $\infty$ -category  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  a functor  $\overline{F} : K^{\triangleleft} \to \mathcal{M}$  is a conical colimit diagram if for every  $V_1, ..., V_n \in \mathcal{V}, W_1, ..., W_m \in \mathcal{W}, Z \in \mathcal{M}$  for  $n, m \ge 0$  the canonical map

(3) 
$$\operatorname{Mul}_{\mathcal{M}}(V_1, ..., V_n, F(\infty), W_1, ..., W_m; Z) \to \lim \operatorname{Mul}_{\mathcal{M}}(V_1, ..., V_n, F(-), W_1, ..., W_m; Z)$$

is an equivalence. By Proposition 2.84 the functor  $F := \overline{F}_{|K} : K \to \mathcal{M}$  underlies a unique  $\mathcal{V}, \mathcal{W}$ -enriched functor  $K^{\circledast}_{\mathcal{V},\mathcal{W}} \to \mathcal{M}^{\circledast}$ , which gives rise to a  $\mathcal{P}Env(\mathcal{V}), \mathcal{P}Env(\mathcal{W})$ -enriched adjunction

$$F_!: \mathcal{P}BEnv(K_{\mathcal{V},\mathcal{W}})^{\circledast} \rightleftharpoons \mathcal{P}BEnv(\mathcal{M})^{\circledast}: F^*.$$

Let \* be the final presheaf on  $K_{\mathcal{V},\mathcal{W}}$ . Lemma 2.100 provides a canonical equivalence

 $\lim \operatorname{Mul}_{\mathcal{M}}(V_{1},...,V_{n},F(-),W_{1},...,W_{m};Z) \simeq \mathcal{P}(K_{\mathcal{V},\mathcal{W}})(*,\operatorname{Mul}_{\mathcal{M}}(V_{1},...,V_{n},F(-),W_{1},...,W_{m};Z) \simeq \mathcal{P}(K_{\mathcal{V},\mathcal{W}})(*,\operatorname{Mul}_{\mathcal{M}}(V_{1},...,V_{n},F(-),W_{1},...,W_{m};Z))$ 

 $Mul_{\mathcal{P}BEnv(K_{\mathcal{V},\mathcal{W}})}(V_1,...,V_n,*,W_1,...,W_m;F^*(Z)) \simeq Mul_{\mathcal{P}BEnv(\mathcal{M})}(V_1,...,V_n,F_!(*),W_1,...,W_m;Z),$ under which the map (3) identifies with the following map:

$$\operatorname{Mul}_{\mathcal{M}}(V_1,...,V_n,F(\infty),W_1,...,W_m;Z) \to \operatorname{Mul}_{\mathcal{P}\operatorname{BEnv}(\mathcal{M})}(V_1,...,V_n,F_!(*),W_1,...,W_m;Z).$$

This motivates the following definition of weighted colimit, where we replace the weakly bi-enriched  $\infty$ -category  $K^{\circledast}_{V,W} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  by any weakly bi-enriched  $\infty$ -category  $\mathcal{J}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  and the final presheaf  $* \in \mathcal{P}(\mathcal{J}) \subset \mathcal{PBEnv}(\mathcal{J})$  by any object of  $\mathcal{PBEnv}(\mathcal{J})$ :

**Definition 3.1.** Let  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}, \mathcal{J}^{\circledast} \to \mathcal{V}'^{\otimes} \times \mathcal{W}'^{\otimes}$  be totally small weakly bi-enriched  $\infty$ -categories,  $F : \mathcal{J}^{\circledast} \to \mathcal{M}^{\circledast}$  an enriched functor,  $Y \in \mathcal{M}$  and  $H \in \mathcal{PBEnv}(\mathcal{J})$ . A morphism  $\psi : F_{!}(H) \to Y$  in  $\mathcal{PBEnv}(\mathcal{M})$  exhibits Y as the H-weighted colimit of F if for every  $Z \in \mathcal{M}, V_{1}, ..., V_{n} \in \mathcal{V}, W_{1}, ..., W_{m} \in \mathcal{W}$  for  $n, m \geq 0$  the following map is an equivalence:

 $Mul_{\mathcal{M}}(V_1,...,V_n,Y,W_1,...,W_m;Z) \rightarrow Mul_{\mathcal{P}BEnv(\mathcal{M})}(V_1,...,V_n,F_!(H),W_1,...,W_m;Z).$ 

Notation 3.2. If a morphism  $\psi$  exists like in Definition 3.1, it is unique, and we say that  $\mathcal{M}$  admits the H-weighted colimit of F and write colim<sup>H</sup>(F) for Y.

Dually, we define weighted limits:

**Definition 3.3.** Let  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}, \mathcal{J}^{\circledast} \to \mathcal{V}'^{\otimes} \times \mathcal{W}'^{\otimes}$  be totally small weakly bi-enriched  $\infty$ -categories,  $F : \mathcal{J}^{\circledast} \to \mathcal{M}^{\circledast}$  an enriched functor,  $Y \in \mathcal{M}$  and  $H \in \mathcal{PBEnv}(\mathcal{J}^{\mathrm{op}})$ . The H-weighted limit of F denoted by  $\lim^{H}(F)$  if it exists, is the H-weighted colimit of  $F^{\mathrm{op}} : (\mathcal{J}^{\mathrm{op}})^{\circledast} \to (\mathcal{M}^{\mathrm{op}})^{\circledast}$ .

We often prove results about weighted colimits that dually give results about weighted limits and viceversa.

**Lemma 3.4.** Let  $\rho : \mathfrak{I}^{\circledast} \to \mathfrak{I}^{\circledast}, F : \mathfrak{I}^{\circledast} \to \mathfrak{M}^{\circledast}$  be enriched functors and  $H \in \mathcal{PBEnv}(\mathfrak{I}), Y \in \mathfrak{M}$ . A morphism  $(F \circ \rho)_!(H) \simeq F_!(\rho_!(H)) \to Y$  in  $\mathcal{PBEnv}(\mathfrak{M})$  exhibits Y as the  $\rho_!(H)$ -weighted colimit of F if and only if it exhibits Y as the H-weighted colimit of  $F \circ \rho$ .

*Proof.* This follows immediately from the definition.

**Remark 3.5.** Let the assumptions like in Lemma 3.4 and let  $F : \mathcal{J}^{\circledast} \to \mathcal{M}^{\circledast}$  lie over maps of  $\infty$ -operads  $\alpha, \beta$ . Then F factors as the canonical enriched functor  $\tau : \mathcal{J}^{\circledast} \to (\alpha, \beta)_!(\mathcal{J})^{\circledast}$  followed by a unique  $\mathcal{V}, \mathcal{W}$ enriched functor  $F' : (\alpha, \beta)_!(\mathcal{J})^{\circledast} \to \mathcal{M}^{\circledast}$ . Thus by Lemma 3.4 a morphism  $F'_!(\tau_!(H)) \simeq F_!(H) \to Y$ exhibits Y as the H-weighted colimit of F if and only if it exhibits Y as the  $\tau_!(H)$ -weighted colimit
of F'. In other words by changing the weight it is enough to define weighted colimits for enriched
functors lying over identity maps of  $\infty$ -operads. On the other hand, the extra freedom in Definition
3.1 is more convenient when studying preservation of weighted colimits by general enriched functors.

**Lemma 3.6.** Let the assumptions like in Definition 3.1 assume that F lies over essentially surjective maps  $\alpha : \mathcal{V}^{\otimes} \to \mathcal{V}^{\otimes}, \beta : \mathcal{W}^{\otimes} \to \mathcal{W}^{\otimes}$  of  $\infty$ -operads. The morphism  $\psi : F_{!}(H) \to Y$  in  $\mathcal{PBEnv}(\mathcal{M})$  exhibits Y as the H-weighted colimit of F if and only if the corresponding morphism  $\psi' : H \to F^{*}(Y)$  in  $\mathcal{PBEnv}(\mathcal{J})$  induces for every  $Z \in \mathcal{M}, V'_{1}, ..., V'_{n} \in \mathcal{V}', W'_{1}, ..., W'_{m} \in \mathcal{W}'$  for  $n, m \geq 0$  an equivalence:

$$\operatorname{Mul}_{\mathcal{M}}(\alpha(V_{1}'),...,\alpha(V_{n}'),Y,\beta(W_{1}'),...,\beta(W_{m}');Z) \to \operatorname{Mul}_{\mathcal{P}\operatorname{BEnv}(\mathcal{J})}(V_{1}',...,V_{n}',H,W_{1}',...,W_{m}';F^{*}(Z)).$$

*Proof.* By adjointness there is an equivalence

$$\begin{aligned} \operatorname{Mul}_{\mathcal{P}\operatorname{BEnv}(\mathcal{M})}(\alpha(V_{1}'),...,\alpha(V_{n}'),\operatorname{F}_{!}(\operatorname{H}),\beta(W_{1}'),...,\beta(W_{m}');\operatorname{Z}) &\simeq \\ \operatorname{Mul}_{\mathcal{P}\operatorname{BEnv}(\mathcal{J})}(V_{1}',...,V_{n}',\operatorname{H},W_{1}',...,W_{m}';\operatorname{F}^{*}(\operatorname{Z})). \end{aligned}$$

**Remark 3.7.** Let the assumptions like in Lemma 3.6 but let  $\alpha, \beta$  not be essentially surjective. Then the condition of Lemma 3.6 is weaker than the one of Definition 3.1 and amounts to say that Y is the H-weighted colimit of the induced  $\mathcal{V}', \mathcal{W}'$ -enriched functor  $\mathcal{J}^{\circledast} \to \mathcal{V}'^{\otimes} \times_{\mathcal{W}^{\otimes}} \mathcal{M}^{\circledast} \times_{\mathcal{W}^{\otimes}} \mathcal{W}'^{\otimes}$ .

We have the following functoriality of weighted colimits:

**Remark 3.8.** Let  $\alpha : F \to G : \mathcal{J}^{\circledast} \to \mathcal{M}^{\circledast}$  be an enriched functor and  $\tau : H \to H'$  a morphism in  $\mathcal{P}Env(\mathcal{J})$ . The morphism  $F_!(H) \xrightarrow{\alpha_!(H)} G_!(H) \xrightarrow{G_!(\tau)} G_!(H') \to \operatorname{colim}^{H'}(G)$  in  $\mathcal{P}Env(\mathcal{M})$  factors as the morphism  $F_!(H) \to \operatorname{colim}^{H}(F)$  followed by a unique morphism  $\operatorname{colim}^{H}(F) \to \operatorname{colim}^{H'}(G)$  in  $\mathcal{M}$ .

**Definition 3.9.** A totally small weight is a quadruple  $(\alpha : \mathcal{V}^{\otimes} \to \mathcal{V}^{\otimes}, \beta : \mathcal{W}^{\otimes} \to \mathcal{W}^{\otimes}, \mathcal{J}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}, \mathcal{H})$ , where  $\alpha, \beta$  are maps of small  $\infty$ -operads,  $\mathcal{J}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  is a totally small weakly bi-enriched  $\infty$ -category and  $\mathcal{H} \in \mathcal{PBEnv}(\mathcal{J})$ . We also call  $(\alpha, \beta, \mathcal{J}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}, \mathcal{H})$  a totally small  $(\alpha, \beta)$ -weight over  $\mathcal{V}, \mathcal{W}$  on  $\mathcal{J}$ .

**Definition 3.10.** Let  $\mathcal{V}^{\otimes} \to Ass, \mathcal{W}^{\otimes} \to Ass \text{ small } \infty\text{-operads.}$ 

- (1) A left weight over  $\mathcal{V}$  is a weight over  $\mathcal{V}, \emptyset$ .
- (2) A right weight over  $\mathcal{W}$  is a weight over  $\emptyset, \mathcal{W}$ .
- (3) A  $\mathcal{V}, \mathcal{W}$ -weight is a weight over  $\mathcal{V}, \mathcal{W}$  such that  $\alpha, \beta$  are the identities.
- (4) A left  $\mathcal{V}$ -weight is a left weight over  $\mathcal{V}, \mathcal{W}$  such that  $\alpha$  is the identity.
- (5) A right  $\mathcal{W}$ -weight is a right weight over  $\mathcal{V}, \mathcal{W}$  such that  $\beta$  is the identity.

**Remark 3.11.** A left weight is a weight  $(\alpha : \mathcal{V}^{\otimes} \to \mathcal{V}^{\otimes}, \beta : \mathcal{W}^{\otimes} \to \mathcal{W}^{\otimes}, \mathcal{J}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}, H \in \mathcal{PBEnv}(\mathcal{J}))$ such that  $\mathcal{W}^{\otimes} \simeq \mathscr{Q}^{\otimes}$ . So also  $\mathcal{W}^{\otimes} \simeq \mathscr{Q}^{\otimes}$ . Consequently, a left weight is equivalently a triple  $(\alpha : \mathcal{V}^{\otimes} \to \mathcal{V}^{\otimes}, \mathcal{J}^{\otimes} \to \mathcal{V}^{\otimes}, \mathcal{H} \in \mathcal{PLEnv}(\mathcal{J}))$ . In particular, we can identify a left weight  $(\alpha : \mathcal{V}^{\otimes} \to \mathcal{V}^{\otimes}, \mathcal{J}^{\otimes} \to \mathcal{V}^{\otimes},$ 

**Definition 3.12.** Let  $\kappa$  be a small regular cardinal. A totally small weight  $(\alpha, \beta, \mathcal{J}^{\circledast} \to \mathcal{V}'^{\otimes} \times \mathcal{W}'^{\otimes}, \mathbf{H})$  is  $\kappa$ -small if  $\mathbf{H}$  belongs to the full subcategory of  $\mathcal{PBEnv}(\mathcal{M})$  generated by  $\mathcal{BEnv}(\mathcal{M})$  under  $\kappa$ -small colimits.

**Definition 3.13.** Let  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be totally small weakly bi-enriched  $\infty$ -category and  $T = (\alpha, \beta, \mathcal{J}^{\circledast} \to \mathcal{V}'^{\otimes} \times \mathcal{W}'^{\otimes}, H)$  a totally small weight over  $\mathcal{V}, \mathcal{W}$  and  $\kappa$  a small regular cardinal.

- (1) We say that  $\mathcal{M}$  admits T-weighted colimits if  $\mathcal{M}$  admits the H-weighted colimit of every enriched functor  $\mathcal{J}^{\circledast} \to \mathcal{M}^{\circledast}$  lying over  $\alpha, \beta$ .
- (2) Let  $\mathcal{H}$  be a collection of totally small weights over  $\mathcal{V}, \mathcal{W}$ . We say that  $\mathcal{M}$  admits  $\mathcal{H}$ -weighted colimits if  $\mathcal{M}$  admits T-weighted colimits for every weight  $T \in \mathcal{H}$ .
- (3) We say that  $\mathcal{M}$  admits  $\kappa$ -small weighted colimits if it admits T-weighted colimits for every  $\kappa$ -small weight T over  $\mathcal{V}, \mathcal{W}$ .

- (4) We say that  $\mathcal{M}$  admits  $\kappa$ -small left weighted colimits if it admits T-weighted colimits for every  $\kappa$ -small left weight T over  $\mathcal{V}$ .
- (5) We say that  $\mathcal{M}$  admits  $\kappa$ -small right weighted colimits if it admits T-weighted colimits for every  $\kappa$ -small right weight T over  $\mathcal{W}$ .
- (6) We say that M admits small weighted colimits if it admits T-weighted colimits for every totally small weight T over V, W.
- (7) We say that  $\mathcal{M}$  admits small left weighted colimits if it admits T-weighted colimits for every small left weight T over  $\mathcal{V}$ .
- (8) We say that M admits small right weighted colimits if it admits T-weighted colimits for every small right weight T over W.

We apply Definition 3.13 in particular to the case that  $\mathcal{H}$  is a collection of left weights over  $\mathcal{V}$  or a collection of right weights over  $\mathcal{W}$  viewed as weights over  $\mathcal{V}, \mathcal{W}$  in the canonical way (Remark 3.11).

3.2. Weighted colimits for pseudo-enriched  $\infty$ -categories. Next we consider weighted colimits for quasi-enriched and pseudo-enriched  $\infty$ -categories. To describe quasi-enriched and pseudo-enriched  $\infty$ -categories on one footage we use the notion of S, T-bi-enriched  $\infty$ -categories for localization pairs  $(\mathcal{V}^{\otimes} \rightarrow \text{Ass}, \text{S}), (\mathcal{W}^{\otimes} \rightarrow \text{Ass}, \text{T})$  (Definition 2.129), which specializes to quasi-enriched and pseudoenriched  $\infty$ -categories and the notion of  $\kappa$ -enriched  $\infty$ -categories for any small regular cardinal  $\kappa$ .

Let  $(\mathcal{W}^{\otimes} \to Ass, S), (\mathcal{W}^{\otimes} \to Ass, T)$  be small localization pairs and  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a small S, Tbi-enriched  $\infty$ -category. By Lemma 2.145 there is an enriched localization  $L : \mathcal{PBEnv}(\mathcal{M})^{\otimes} \neq \mathcal{PBEnv}(\mathcal{M})^{\otimes}_{S,T}$  to the respective  $\infty$ -category of "presheaves" for S, T-enriched  $\infty$ -categories (Proposition 2.149). We have the following description of weighted colimit in this context:

**Lemma 3.14.** Let  $(\mathcal{V}^{\otimes} \to \operatorname{Ass}, S), (\mathcal{W}^{\otimes} \to \operatorname{Ass}, T)$  be small localization pairs and  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$ a small S, T-bi-enriched  $\infty$ -category,  $F : \mathcal{J}^{\otimes} \to \mathcal{M}^{\otimes}$  an enriched functor,  $H \in \operatorname{PBEnv}(\mathcal{J}), Y \in \mathcal{M}$  and  $F_{!}(H) \to Y$  a morphism in  $\operatorname{PBEnv}(\mathcal{M})$ . The following conditions are equivalent:

- (1) The morphism  $F_1(H) \to Y$  in  $\mathcal{PBEnv}(\mathcal{M})$  exhibits Y as the H-weighted colimit of F.
- (2) For every  $Z \in M$ ,  $V_1, ..., V_n \in V$ ,  $W_1, ..., W_m \in W$  for  $n, m \ge 0$  the induced morphism  $L(F_!(H)) \rightarrow Y$  in  $\mathcal{PBEnv}(\mathcal{M})_{S,T}$  induces an equivalence:

 $Mul_{\mathcal{M}}(V_1, ..., V_n, Y, W_1, ..., W_m; Z) \rightarrow Mul_{\mathcal{PBEnv}(\mathcal{M})_{S,T}}(V_1, ..., V_n, L(F_!(H)), W_1, ..., W_m; Z).$ 

If  $H \in \mathcal{PBEnv}(\mathcal{M})_{S,T}$  conditions (1),(2) are equivalent to the following one: the corresponding morphism  $\psi' : H \to F^*(Y)$  in  $\mathcal{PBEnv}(\mathcal{J})_{S,T}$  induces for any  $Z \in \mathcal{M}, V_1, ..., V_n \in \mathcal{V}, W_1, ..., W_m \in \mathcal{W}$  for  $n, m \ge 0$  an equivalence:

 $Mul_{\mathcal{M}}(V_1, ..., V_n, Y, W_1, ..., W_m; Z) \rightarrow Mul_{\mathcal{PBEnv}(\mathcal{J})_{S,T}}(V_1, ..., V_n, H, W_1, ..., W_m; F^*(Z)).$ 

*Proof.* The equivalence between (1) and (2) follows by adjointness via the enriched localization L :  $\mathcal{PBEnv}(\mathcal{M})^{\circledast} \rightleftharpoons \mathcal{PBEnv}(\mathcal{M})^{\circledast}_{S,T}$ . The second part of the lemma follows from the enriched adjunction  $L \circ F_! : \mathcal{PBEnv}(\mathcal{J})^{\circledast}_{S,T} \rightleftharpoons \mathcal{PBEnv}(\mathcal{M})^{\circledast}_{S,T} : F^*$ .

$$\square$$

**Notation 3.15.** Let  $(\mathcal{V}^{\otimes} \to \operatorname{Ass}, S), (\mathcal{W}^{\otimes} \to \operatorname{Ass}, T)$  be small localization pairs and  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a small S, T-bi-enriched  $\infty$ -category. Let  $\mathcal{M}^{\otimes}_{S,T} \subset \mathcal{P}BEnv(\mathcal{M})^{\otimes}_{S,T}$  be the full weakly bi-enriched subcategory spanned by the objects  $T \in \mathcal{P}BEnv(\mathcal{M})_{S,T}$  such that there is a morphism  $T \to T'$  in  $\mathcal{P}BEnv(\mathcal{M})_{S,T}$ , where  $T' \in \mathcal{M}$ , such that for every  $Z \in \mathcal{M}, V_1, ..., V_n \in \mathcal{V}, W_1, ..., W_m \in \mathcal{W}$  for  $n, m \ge 0$  the following map is an equivalence:

$$(4) \quad \operatorname{Mul}_{\mathcal{P}\operatorname{BEnv}(\mathcal{M})_{S,T}}(V_1, ..., V_n, T', W_1, ..., W_m; Z) \to \operatorname{Mul}_{\mathcal{P}\operatorname{BEnv}(\mathcal{M})_{S,T}}(V_1, ..., V_n, T, W_1, ..., W_m; Z).$$

**Remark 3.16.** Let  $\overline{\mathcal{M}}^{\circledast} \subset \mathcal{PBEnv}(\mathcal{M})_{S,T}^{\circledast}$  be the full weakly bi-enriched subcategory spanned by  $\mathcal{M}$ . The embedding  $\overline{\mathcal{M}}^{\circledast} \subset \mathcal{PBEnv}(\mathcal{M})_{S,T}^{\circledast}$  lands in  $\mathcal{M}_{S,T}^{\circledast}$  (taking the identity of T). **Lemma 3.17.** Let  $(\mathcal{V}^{\otimes} \to \operatorname{Ass}, S), (\mathcal{W}^{\otimes} \to \operatorname{Ass}, T)$  be small localization pairs and  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a small S, T-bi-enriched  $\infty$ -category. The embedding  $\overline{\mathcal{M}}^{\circledast} \subset \mathcal{M}^{\circledast}_{S,T}$  admits a  $S^{-1}\mathcal{P}\operatorname{Env}(\mathcal{V}), T^{-1}\mathcal{P}\operatorname{Env}(\mathcal{W})$ -enriched left adjoint  $\phi$ .

*Proof.* Let T ∈ PBEnv( $\mathcal{M}$ )<sub>S,T</sub>. Then there is a morphism T → T' in PBEnv( $\mathcal{M}$ )<sub>S,T</sub>, where T' ∈  $\mathcal{M}$ , such that for every Z ∈  $\mathcal{M}$ , V<sub>1</sub>,..., V<sub>n</sub> ∈  $\mathcal{V}$ , W<sub>1</sub>,..., W<sub>m</sub> ∈  $\mathcal{W}$  for n, m ≥ 0 the map (4) is an equivalence. Thus for every V ∈ S<sup>-1</sup>PEnv( $\mathcal{V}$ ), W ∈ T<sup>-1</sup>PEnv( $\mathcal{W}$ ) the following map is an equivalence since S<sup>-1</sup>PEnv( $\mathcal{V}$ ) is generated under small colimits by tensor products of objects of  $\mathcal{V}$ :

(5) 
$$\mathcal{P}BEnv(\mathcal{M})_{S,T}(V \otimes T' \otimes W, Z) \rightarrow \mathcal{P}BEnv(\mathcal{M})_{S,T}(V \otimes T \otimes W, Z).$$

**Remark 3.18.** Let  $(\mathcal{V}^{\otimes} \to \operatorname{Ass}, S), (\mathcal{W}^{\otimes} \to \operatorname{Ass}, T)$  be small localization pairs and  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a small S, T-bi-enriched  $\infty$ -category. An enriched functor  $F : \mathcal{J}^{\otimes} \to \mathcal{M}^{\otimes}$  admits a H-weighted colimit for a weight  $H \in \mathcal{PBEnv}(\mathcal{J})$  if and only if the functor  $\mathcal{PBEnv}(\mathcal{J}) \xrightarrow{F_1} \mathcal{PBEnv}(\mathcal{M}) \xrightarrow{L} \mathcal{PBEnv}(\mathcal{M})_{S,T}$  lands in  $\mathcal{M}_{S,T}$ . In this case there is a canonical equivalence  $\operatorname{colim}^H(F) \simeq \phi(L(F_1(H)))$ .

**Proposition 3.19.** Let  $(\mathcal{V}^{\otimes} \to \operatorname{Ass}, \operatorname{S}), (\mathcal{W}^{\otimes} \to \operatorname{Ass}, \operatorname{T})$  be small localization pairs and  $\mathcal{J}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}, \mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  small S, T-bi-enriched  $\infty$ -categories.

- (1) Let F: J<sup>®</sup> → M<sup>®</sup> be a V, W-enriched functor such that M admits the H-weighted colimit of F for every H ∈ PBEnv(J)<sub>S,T</sub>. The enriched functor F<sup>\*</sup><sub>|M̄®</sub> : M̄<sup>®</sup> → PBEnv(J)<sup>®</sup><sub>S,T</sub> admits a S<sup>-1</sup>PEnv(V), T<sup>-1</sup>PEnv(W)-enriched left adjoint that sends H to colim<sup>H</sup>(F) and extends F.
- (2) Assume that  $\mathcal{M}$  admits the H-weighted colimit of every  $\mathcal{V}, \mathcal{W}$ -enriched functor  $F : \mathcal{J}^{\circledast} \to \mathcal{M}^{\circledast}$ for every  $H \in \mathcal{PBEnv}(\mathcal{J})_{S,T}$ . The induced functor

 $\mathrm{EnrFun}_{\mathrm{S}^{-1}\mathcal{P}\mathrm{Env}(\mathcal{V}),\mathrm{T}^{-1}\mathcal{P}\mathrm{Env}(\mathcal{W})}(\mathcal{P}\mathrm{B}\mathrm{Env}(\mathcal{J})_{\mathrm{S},\mathrm{T}},\bar{\mathcal{M}}) \to \mathrm{EnrFun}_{\mathcal{V},\mathcal{W}}(\mathcal{J},\mathcal{M})$ 

admits a fully faithful left adjoint that lands in  $\operatorname{LinFun}_{S^{-1}\operatorname{PEnv}(\mathcal{V}),T^{-1}\operatorname{PEnv}(\mathcal{W})}^{L}(\operatorname{PBEnv}(\mathcal{J})_{S,T},\overline{\mathcal{M}})$ . In particular, the following induced functor is an equivalence:

 $\operatorname{LinFun}_{S^{-1}\mathcal{P}\operatorname{Env}(\mathcal{V}), T^{-1}\mathcal{P}\operatorname{Env}(\mathcal{W})}^{L}(\mathcal{P}\operatorname{BEnv}(\mathcal{J})_{S,T}, \overline{\mathcal{M}}) \to \operatorname{EnrFun}_{\mathcal{V}, \mathcal{W}}(\mathcal{J}, \mathcal{M})$ 

Proof. (1): The S<sup>-1</sup>𝔅Env(𝔅), T<sup>-1</sup>𝔅Env(𝔅)-enriched adjunction F<sub>1</sub>: 𝔅BEnv(𝔅)<sup>⊕</sup><sub>S,T</sub> ≥ 𝔅BEnv(𝔅)<sup>⊕</sup><sub>S,T</sub>: F<sup>\*</sup> restricts to a S<sup>-1</sup>𝔅Env(𝔅), T<sup>-1</sup>𝔅Env(𝔅)-enriched adjunction F<sub>1</sub>: 𝔅BEnv(𝔅)<sup>⊕</sup><sub>S,T</sub> ≥ 𝔅<sup>⊕</sup><sub>S,T</sub>: F<sup>\*</sup><sub>|𝔅,T</sub>. Composing the latter with the S<sup>-1</sup>𝔅Env(𝔅), T<sup>-1</sup>𝔅Env(𝔅)-enriched localization L : 𝔅<sup>⊕</sup><sub>S,T</sub> ≥ 𝔅<sup>⊕</sup> 𝔅 𝔅<sup>⊕</sup> ve obtain a S<sup>-1</sup>𝔅Env(𝔅), T<sup>-1</sup>𝔅Env(𝔅)-enriched adjunction L ∘ F<sub>1</sub>: 𝔅BEnv(𝔅)<sup>⊕</sup><sub>S,T</sub> ≥ 𝔅<sup>⊕</sup> × 𝔅<sup>⊕</sup> : F<sup>\*</sup><sub>|𝔅,T</sub>. The restriction of the enriched functor L ∘ F<sub>1</sub>: 𝔅BEnv(𝔅)<sup>⊕</sup><sub>S,T</sub> → 𝔅<sup>⊕</sup> factors as 𝔅<sup>⊕</sup> → 𝔅<sup>⊕</sup> ⊂ 𝔅<sup>⊕</sup><sub>S,T</sub> → 𝔅<sup>⊕</sup> and so is F.

(2): By Proposition 2.149 the induced functor

 $\operatorname{EnrFun}_{S^{-1}\mathcal{P}\operatorname{Env}(\mathcal{V}), T^{-1}\mathcal{P}\operatorname{Env}(\mathcal{W})}(\mathcal{P}\operatorname{B}\operatorname{Env}(\mathcal{J})_{S,T}, \mathcal{P}\operatorname{B}\operatorname{Env}(\mathcal{M})_{S,T}) \to \operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}(\mathcal{J}, \mathcal{P}\operatorname{B}\operatorname{Env}(\mathcal{M})_{S,T})$ 

admits a fully faithful left adjoint, which lands in the full subcategory

 $\operatorname{LinFun}_{\mathrm{S}^{-1}\mathcal{P}\operatorname{Env}(\mathcal{V}),\mathrm{T}^{-1}\mathcal{P}\operatorname{Env}(\mathcal{W})}^{\mathrm{L}}(\mathcal{P}\operatorname{B}\operatorname{Env}(\mathcal{J})_{\mathrm{S},\mathrm{T}},\mathcal{P}\operatorname{B}\operatorname{Env}(\mathcal{M})_{\mathrm{S},\mathrm{T}}).$ 

Let

 $\mathrm{EnrFun}_{\mathrm{S}^{-1}\mathcal{P}\mathrm{Env}(\mathcal{V}),\mathrm{T}^{-1}\mathcal{P}\mathrm{Env}(\mathcal{W})}(\mathcal{P}\mathrm{B}\mathrm{Env}(\mathcal{J})_{\mathrm{S},\mathrm{T}},\mathfrak{M}_{\mathrm{S},\mathrm{T}}) \subset$ 

 $\operatorname{EnrFun}_{S^{-1}\mathcal{P}\operatorname{Env}(\mathcal{V}),T^{-1}\mathcal{P}\operatorname{Env}(\mathcal{W})}(\mathcal{P}\operatorname{B}\operatorname{Env}(\mathcal{J})_{S,T},\mathcal{M}_{S,T})$ 

be the full subcategory of enriched functors sending  $\mathcal J$  to  $\mathcal M$ . In particular, the induced functor

 $\operatorname{EnrFun}_{S^{-1}\mathcal{P}\operatorname{Env}(\mathcal{V}), T^{-1}\mathcal{P}\operatorname{Env}(\mathcal{W})}(\mathcal{P}\operatorname{B}\operatorname{Env}(\mathcal{J})_{S,T}, \mathcal{M}_{S,T}) \to \operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}(\mathcal{J},\mathcal{M})$
admits a fully faithful left adjoint that lands in the full subcategory

 $\operatorname{LinFun}_{S^{-1}\mathcal{P}\operatorname{Env}(\mathcal{V}),T^{-1}\mathcal{P}\operatorname{Env}(\mathcal{W})}^{L}(\mathcal{P}\operatorname{B}\operatorname{Env}(\mathcal{J})_{S,T},\mathcal{M}_{S,T}).$ 

So the result follows from the fact that the equivalence

 $\mathrm{EnrFun}_{\mathrm{S}^{-1}\mathcal{P}\mathrm{Env}(\mathcal{V}),\mathrm{T}^{-1}\mathcal{P}\mathrm{Env}(\mathcal{W})}(\mathcal{P}\mathrm{B}\mathrm{Env}(\mathcal{J})_{\mathrm{S},\mathrm{T}},\mathcal{M}) \simeq \mathrm{EnrFun}_{\mathrm{S}^{-1}\mathcal{P}\mathrm{Env}(\mathcal{V}),\mathrm{T}^{-1}\mathcal{P}\mathrm{Env}(\mathcal{W})}(\mathcal{P}\mathrm{B}\mathrm{Env}(\mathcal{J})_{\mathrm{S},\mathrm{T}},\mathcal{M}_{\mathrm{S},\mathrm{T}})$ 

is inverse to the following functor post-composing with L:

$$\operatorname{EnrFun}_{S^{-1}\mathcal{P}\operatorname{Env}(\mathcal{V}),T^{-1}\mathcal{P}\operatorname{Env}(\mathcal{W})}(\mathcal{P}\operatorname{BEnv}(\mathcal{J})_{S,T},\mathcal{M}_{S,T}) \to \operatorname{EnrFun}_{S^{-1}\mathcal{P}\operatorname{Env}(\mathcal{V}),T^{-1}\mathcal{P}\operatorname{Env}(\mathcal{W})}(\mathcal{P}\operatorname{BEnv}(\mathcal{J})_{S,T},\mathcal{M})$$

**Corollary 3.20.** Let  $(\mathcal{V}^{\otimes} \to \operatorname{Ass}, \operatorname{S}), (\mathcal{W}^{\otimes} \to \operatorname{Ass}, \operatorname{T})$  be small localization pairs and  $\mathcal{J}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a small S, T-bi-enriched  $\infty$ -category,  $\operatorname{H} \in \operatorname{PBEnv}(\mathcal{J})_{\operatorname{S},\operatorname{T}}$  and  $\iota : \mathcal{J}^{\circledast} \subset \operatorname{PBEnv}(\mathcal{J})_{\operatorname{S},\operatorname{T}}^{\circledast}$  the canonical enriched embedding. There is a canonical equivalence  $\operatorname{colim}^{\operatorname{H}} \iota \simeq \operatorname{H}$ .

3.3. Weighted colimits for enriched  $\infty$ -categories. Next we consider weighted colimits for enriched  $\infty$ -categories, which is of special relevance for us. In the following we use Notation 2.151. Let  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  be an  $\infty$ -category left enriched in a presentably monoidal  $\infty$ -category and  $L: \widehat{\mathcal{P}BEnv}(\mathcal{M})^{\circledast} \rightleftharpoons \widehat{\mathcal{P}BEnv}(\mathcal{M})^{\circledast}_{LEnr}$  the enriched localization of Lemma 2.145. For every  $\mathcal{V}, \mathcal{W}$ -enriched functor  $F: \mathcal{J}^{\circledast} \to \mathcal{M}^{\circledast}$  the  $\mathcal{V}, \mathcal{W}$ -enriched functor

$$\widehat{\mathcal{P}}BEnv(\mathcal{J})^{\circledast}_{LEnr} \subset \widehat{\mathcal{P}}BEnv(\mathcal{J})^{\circledast} \xrightarrow{F_1} \widehat{\mathcal{P}}BEnv(\mathcal{M})^{\circledast} \xrightarrow{L} \widehat{\mathcal{P}}BEnv(\mathcal{M})^{\circledast}_{LEnr}$$

restricts to a  $\mathcal{V}, \mathcal{W}$ -enriched functor  $\mathcal{PBEnv}(\mathcal{J})_{\text{LEnr}}^{\circledast} \to \mathcal{PBEnv}(\mathcal{M})_{\text{LEnr}}^{\circledast}$  that admits a  $\mathcal{V}, \mathcal{W}$ -enriched right adjoint F<sup>\*</sup>. Specializing to left enriched  $\infty$ -categories Lemma 3.14 gives the following corollary:

**Corollary 3.21.** Let  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be an  $\infty$ -category left enriched in a presentably monoidal  $\infty$ -category,  $F : \mathcal{J}^{\otimes} \to \mathcal{M}^{\otimes}$  a  $\mathcal{V}, \mathcal{W}$ -enriched functor,  $H \in \mathcal{PBEnv}(\mathcal{J})_{LEnr}, Y \in \mathcal{M}$  and  $F_!(H) \to Y$  a morphism in  $\widehat{\mathcal{P}BEnv}(\mathcal{M})$ . The following conditions are equivalent:

- (1) The morphism  $F_!(H) \to Y$  in  $\widehat{\mathcal{P}}BEnv(\mathcal{M})$  exhibits Y as the H-weighted colimit of F.
- (2) The induced morphism  $L(F_!(H)) \rightarrow Y$  in  $\mathcal{PBEnv}(\mathcal{M})_{LEnr}$  induces for every  $Z \in \mathcal{M}$  an equivalence:

 $\operatorname{LMor}_{\mathcal{M}}(Y, Z) \rightarrow \operatorname{LMor}_{\mathcal{P}\operatorname{BEnv}(\mathcal{M})_{\operatorname{LEnr}}}(L(F_{!}(H)), Z).$ 

(3) The corresponding morphism  $\psi' : H \to F^*(Y)$  in  $\mathcal{PBEnv}(\mathcal{J})_{\text{LEnr}}$  induces for every  $Z \in \mathcal{M}$  an equivalence:

$$\mathrm{LMor}_{\mathcal{M}}(\mathrm{Y},\mathrm{Z}) \to \mathrm{LMor}_{\mathcal{P}\mathrm{BEnv}(\mathcal{J})_{\mathrm{LEnr}}}(\mathrm{H},\mathrm{F}^{*}(\mathrm{Z})).$$

**Remark 3.22.** Hinich [15, § 6] defines a notion of weighted colimit for his model of enriched  $\infty$ -categories [14], which is equivalent to our model by [18] and [10, Theorem 6.7.]. The description of the weigted colimit given by Corollary 3.21 implies that our notion of weighted colimit for left enriched  $\infty$ -categories agrees with Hinich's definition.

One proves the following proposition like Proposition 3.23:

**Proposition 3.23.** Let  $\mathcal{V}^{\otimes} \to Ass$  be a presentably monoidal  $\infty$ -category,  $\mathcal{W}^{\otimes} \to Ass$  a small  $\infty$ -operad and  $\mathcal{J}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}, \mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  small left enriched  $\infty$ -categories.

(1) Let F: J<sup>®</sup> → M<sup>®</sup> be a V, W-enriched functor such that M admits the H-weighted colimit of F for every H ∈ PBEnv(J)<sub>LEnr</sub>. The V, W-enriched functor F<sup>\*</sup><sub>|M<sup>®</sup></sub> : M<sup>®</sup> → PBEnv(J)<sup>®</sup><sub>LEnr</sub> admits a V, W-enriched left adjoint that sends H to colim<sup>H</sup>(F) and extends F.

(2) Assume that  $\mathcal{M}$  admits the H-weighted colimit of every  $\mathcal{V}, \mathcal{W}$ -enriched functor  $F : \mathcal{J}^{\circledast} \to \mathcal{M}^{\circledast}$ for every  $H \in \mathcal{PBEnv}(\mathcal{J})_{LEnr}$ . The induced functor

 $\operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}(\mathcal{P}\operatorname{BEnv}(\mathcal{J})_{\operatorname{LEnr}},\mathcal{M}) \to \operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}(\mathcal{J},\mathcal{M})$ 

admits a fully faithful left adjoint that lands in  $\operatorname{LinFun}_{\mathcal{V},\mathcal{W}}^{L}(\mathcal{P}BEnv(\mathcal{J})_{\operatorname{LEnr}},\mathcal{M})$ . In particular, the following induced functor is an equivalence:

 $\operatorname{LinFun}_{\mathcal{V}}^{L} \mathcal{W}(\mathcal{P}\operatorname{BEnv}(\mathcal{J})_{\operatorname{LEnr}}, \mathcal{M}) \to \operatorname{EnrFun}_{\mathcal{V}, \mathcal{W}}(\mathcal{J}, \mathcal{M})$ 

We use the following enriched variants of weights, which put us in the situation of Corollary 3.21 via Lemma 3.4 and Remark 3.5:

**Definition 3.24.** Let  $\mathcal{V}^{\otimes} \to Ass, \mathcal{W}^{\otimes} \to Ass$  be  $\infty$ -operads.

- (1) A small left enriched weight over  $\mathcal{V}, \mathcal{W}$  is a weight  $(\alpha : \mathcal{V}^{\otimes} \to \mathcal{V}^{\otimes}, \beta : \mathcal{W}^{\otimes} \to \mathcal{W}^{\otimes}, \mathcal{J}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}, \mathcal{H})$  over  $\mathcal{V}, \mathcal{W}$  such that  $\mathcal{V}^{\otimes} \to \operatorname{Ass}, \mathcal{V}^{\otimes} \to \operatorname{Ass}$  are presentably monoidal  $\infty$ -categories,  $\mathcal{W}^{\otimes} \to \operatorname{Ass}, \mathcal{W}^{\otimes} \to \operatorname{Ass}$  are small  $\infty$ -operads,  $\alpha$  is a left adjoint monoidal functor,  $\mathcal{J}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  is a small left enriched  $\infty$ -category and  $\mathcal{H} \in \mathcal{PBEnv}(\mathcal{M})_{\operatorname{LEnr}}$ .
- (2) A small right enriched weight over  $\mathcal{V}, \mathcal{W}$  is a weight  $(\alpha : \mathcal{V}^{\otimes} \to \mathcal{V}^{\otimes}, \beta : \mathcal{W}^{\otimes} \to \mathcal{W}^{\otimes}, \mathcal{J}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}, \mathcal{H})$  over  $\mathcal{V}, \mathcal{W}$  such that  $\mathcal{W}^{\otimes} \to \operatorname{Ass}, \mathcal{W}^{\otimes} \to \operatorname{Ass}$  are presentably monoidal  $\infty$ -categories,  $\mathcal{V}^{\otimes} \to \operatorname{Ass}, \mathcal{V}^{\otimes} \to \operatorname{Ass}$  are small  $\infty$ -operads,  $\beta$  is a left adjoint monoidal functor,  $\mathcal{J}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  is a small right enriched  $\infty$ -category and  $\mathcal{H} \in \mathcal{PBEnv}(\mathcal{M})_{\operatorname{REnr}}$ .
- (3) A small enriched weight over  $\mathcal{V}, \mathcal{W}$  is a weight  $(\alpha : \mathcal{V}^{\prime \otimes} \to \mathcal{V}^{\otimes}, \beta : \mathcal{W}^{\prime \otimes} \to \mathcal{W}^{\otimes}, \mathcal{J}^{\otimes} \to \mathcal{V}^{\prime \otimes} \times \mathcal{W}^{\prime \otimes}, \mathbf{H})$ over  $\mathcal{V}, \mathcal{W}$  such that  $\mathcal{V}^{\otimes} \to \mathrm{Ass}, \mathcal{V}^{\prime \otimes} \to \mathrm{Ass}, \mathcal{W}^{\otimes} \to \mathrm{Ass}, \mathcal{W}^{\prime \otimes} \to \mathrm{Ass}$  are presentably monoidal  $\infty$ -categories,  $\alpha, \beta$  are left adjoint monoidal functors,  $\mathcal{J}^{\otimes} \to \mathcal{V}^{\prime \otimes} \times \mathcal{W}^{\prime \otimes}$  is a small bi-enriched  $\infty$ -category and  $\mathbf{H} \in \mathcal{PBEnv}(\mathcal{M})_{\mathrm{BEnr}}$ .

3.4. Conical colimits and tensors. Next we consider our first example of weighted colimits, which are conical colimits and left and right tensors.

**Definition 3.25.** Let  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be a weakly bi-enriched  $\infty$ -category.

- (1) A functor  $F : K^{\triangleright} \to \mathcal{M}$  is a conical colimit diagram if for every  $V_1, ..., V_n \in \mathcal{V}, W_1, ..., W_m \in \mathcal{W}$ for  $n, m \ge 0$  and  $Y \in \mathcal{M}$  the presheaf  $Mul_{\mathcal{M}}(V_1, ..., V_n, -, W_1, ..., W_m; Y)$  on  $\mathcal{M}$  sends  $F^{op} : (K^{op})^{\triangleleft} \to \mathcal{M}^{op}$  to a limit diagram. An  $\infty$ -category  $\mathcal{M}$  admits K-indexed conical colimits if  $\mathcal{M}$  admits the conical colimit of every functor starting at K.
- (2) A functor  $F: K^{\triangleleft} \to \mathcal{M}$  is a conical limit diagram if for every  $V_1, ..., V_n \in \mathcal{V}, W_1, ..., W_m \in \mathcal{W}$  for  $n, m \ge 0$  and  $X \in \mathcal{M}$  the presheaf  $Mul_{\mathcal{M}}(V_1, ..., V_n, X, W_1, ..., W_m; -)$  on  $\mathcal{M}$  sends F to a limit diagram. An  $\infty$ -category  $\mathcal{M}$  admits K-indexed conical limits if  $\mathcal{M}$  admits the conical limit of every functor starting at K.

**Remark 3.26.** Let  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be a weakly bi-enriched  $\infty$ -category.

- (1) Every conical colimit diagram is a colimit diagram.
- (2) If  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  admits left and right cotensors, every colimit diagram is conical.
- (3) Assume that  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  admits left tensors and let  $G: K \to \mathcal{M}$  be a functor. The colimit of G is conical if and only if forming left tensors preserves the colimit of G and for every  $W_1, ..., W_m \in \mathcal{W}$  for  $m \ge 0$  and  $Y \in \mathcal{M}$  the presheaf  $\operatorname{Mul}_{\mathcal{M}}(-, W_1, ..., W_m; Y)$  on  $\mathcal{M}$  preserves the limit of  $G^{\operatorname{op}}$ .
- (4) Assume that  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  admits left and right tensors and let  $G : K \to \mathcal{M}$  be a functor. The colimit of G is conical if and only if forming left and right tensors preserves the colimit of G.

**Definition 3.27.** Let  $\mathcal{K} \subset \mathsf{Cat}_{\infty}$  be a full subcategory. A weakly bi-enriched  $\infty$ -category  $\phi : \mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  is compatible with  $\mathcal{K}$ -indexed colimits if  $\phi$  admits  $\mathcal{K}$ -indexed conical colimits and for every

 $X, Y \in \mathcal{M}, V_1, ..., V_n \in \mathcal{V}, W_1, ..., W_m \in \mathcal{W}$  for  $n, m \ge 0$  and  $0 \le i \le n, 0 \le j \le m$  the presheaves

$$Mul_{\mathcal{M}}(V_1,...,V_i,-,V_{i+1},...,V_n,X,W_1,...,W_m;Y)$$

$$Mul_{\mathcal{M}}(V_1, ..., V_n, X, W_1, ..., W_j, -, W_{j+1}, ..., W_m; Y)$$

on  $\mathcal{V}, \mathcal{W}$ , respectively, preserve  $\mathcal{K}$ -indexed limits.

**Notation 3.28.** Let K be a small  $\infty$ -category and  $\mathcal{V}^{\otimes} \to Ass, \mathcal{W}^{\otimes} \to Ass$  small  $\infty$ -operads. By Corollary 2.101 the equivalence  $K^{\otimes}_{\mathcal{V},\mathcal{W}} \simeq *^{\otimes}_{\mathcal{V}} \times K \times *^{\otimes}_{\mathcal{W}}$  extends to an equivalence

(6) 
$$\mathcal{PBEnv}(\mathbf{K}_{\mathcal{V},\mathcal{W}}) \simeq \mathcal{PEnv}(\mathcal{V}) \otimes \mathcal{P}(\mathbf{K}) \otimes \mathcal{PEnv}(\mathcal{W}).$$

For every  $V \in \mathcal{P}Env(\mathcal{V}), W \in \mathcal{P}Env(\mathcal{W})$  we define the following weight

 $\langle V, K, W \rangle \in \mathcal{P}BEnv(K_{\mathcal{V},\mathcal{W}})$ 

corresponding to the image of  $(V, *, W) \in \mathcal{P}Env(\mathcal{V}) \times \mathcal{P}(K) \times \mathcal{P}Env(\mathcal{W})$  in  $\mathcal{P}Env(\mathcal{V}) \otimes \mathcal{P}(K) \otimes \mathcal{P}Env(\mathcal{W})$ .

Remark 3.29. Equivalence (6) restricts to equivalences

$$\mathcal{P}\text{LEnv}(K_{\mathcal{V},\mathcal{W}}) \simeq \mathcal{P}\text{Env}(\mathcal{V}) \otimes \mathcal{P}(K) \otimes \mathcal{P}(*_{\mathcal{W}}),$$
$$\mathcal{P}\text{REnv}(K_{\mathcal{V},\mathcal{W}}) \simeq \mathcal{P}(*_{\mathcal{V}}) \otimes \mathcal{P}(K) \otimes \mathcal{P}\text{Env}(\mathcal{W}),$$
$$\mathcal{P}(K_{\mathcal{V},\mathcal{W}}) \simeq \mathcal{P}(*_{\mathcal{V}}) \otimes \mathcal{P}(K) \otimes \mathcal{P}(*_{\mathcal{W}}).$$

Thus  $\langle V, *, \mathbb{1}_{\mathcal{P}Env(\mathcal{W})} \rangle$  is a left weight and  $\langle \mathbb{1}_{\mathcal{P}Env(\mathcal{V})}, *, W \rangle$  is a right weight.

**Definition 3.30.** Let K be a small  $\infty$ -category and  $\mathcal{V}^{\otimes} \to Ass, \mathcal{W}^{\otimes} \to Ass$  small  $\infty$ -operads. The trivial weight on K is the weight  $\langle \mathbb{1}_{\mathcal{P}Env(\mathcal{V})}, K, \mathbb{1}_{\mathcal{P}Env(\mathcal{W})} \rangle$ .

**Lemma 3.31.** Let  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\circledast}$  be a totally small weakly bi-enriched  $\infty$ -category, K an  $\infty$ -category, V  $\in \mathcal{V}, W \in \mathcal{W}$  and F :  $K^{\circledast}_{\mathcal{V}, \mathcal{W}} \to \mathcal{M}^{\circledast}$  a  $\mathcal{V}, \mathcal{W}$ -enriched functor and X  $\in \mathcal{M}$  corresponding to a  $\mathcal{V}, \mathcal{W}$ -enriched functor  $*^{\circledast}_{\mathcal{V}, \mathcal{W}} \to \mathcal{M}^{\circledast}$ .

- (1) The colimit of X weighted at the weight  $\langle V, *, W \rangle$  is the bitensor  $V \otimes X \otimes W$ .
- (2) The colimit of F weighted at the left weight  $\langle V, \star, \mathbb{1}_{PEnv(W)} \rangle$  is the left tensor  $V \otimes X$ .
- (3) The colimit of F weighted at the right weight  $(\mathbb{1}_{PEnv(\mathcal{V})}, *, W)$  is the right tensor  $X \otimes W$ .
- (4) The colimit of F weighted at the trivial weight on K is the conical colimit of  $F: K \to M$ .

*Proof.* We prove (1). The proofs of (2), (3), (4) are similar. The induced left adjoint  $\mathcal{P}Env(\mathcal{V})$ ,  $\mathcal{P}Env(\mathcal{W})$ -linear functor  $F_! : (\mathcal{P}Env(\mathcal{V}) \otimes \mathcal{P}(K) \otimes \mathcal{P}Env(\mathcal{W}))^{\circledast} \simeq \mathcal{P}BEnv(K_{\mathcal{V},\mathcal{W}})^{\circledast} \rightarrow \mathcal{P}BEnv(\mathcal{M})^{\circledast}$  sends  $\langle V, K, W \rangle$  to  $V \otimes F_!(*) \otimes W$ . The final presheaf \* on K is the colimit of the Yoneda-embedding of K so that  $F_!(*)$  is the colimit of F. So for every  $Z \in \mathcal{M}, V_1, ..., V_n \in \mathcal{V}, W_1, ..., W_m$  for  $n, m \ge 0$  there is an equivalence

$$\operatorname{Mul}_{\mathcal{P}BEnv(\mathcal{M})}(V_1, ..., V_n, F_!(\langle V, K, W \rangle), W_1, ..., W_m; Z) \simeq$$

 $Mul_{\mathcal{P}BEnv(\mathcal{M})}(V_1,...,V_n,V,X,W,W_1,...,W_m;Z) \simeq Mul_{\mathcal{M}}(V_1,...,V_n,V,X,W,W_1,...,W_m;Z).$ 

**Lemma 3.32.** Let  $\kappa$  be a small regular cardinal, K a  $\kappa$ -small  $\infty$ -category,  $\mathcal{V}^{\otimes} \to \operatorname{Ass}, \mathcal{W}^{\otimes} \to \operatorname{Ass}$  small  $\infty$ -operads, V in the full subcategory of  $\operatorname{PEnv}(\mathcal{V})$  generated by  $\mathcal{V}$  under  $\kappa$ -small colimits, W in the full subcategory of  $\operatorname{PEnv}(\mathcal{W})$  generated by  $\mathcal{W}$  under  $\kappa$ -small colimits. The weight  $\langle V, K, W \rangle$  is  $\kappa$ -small.

Proof. Every presheaf X on K is the colimit of the functor  $K \times_{\mathcal{P}(K)} \mathcal{P}(K)_{/X} \to K \subset \mathcal{P}(K)$ . Thus the final presheaf on K is the colimit of the Yoneda-embedding  $K \subset \mathcal{P}(K)$  and so a  $\kappa$ -small colimit of representables if K is  $\kappa$ -small. So the result follows since the functor  $\mathcal{P}Env(\mathcal{V}) \times \mathcal{P}(K) \times \mathcal{P}Env(\mathcal{W}) \to \mathcal{P}Env(\mathcal{V}) \otimes \mathcal{P}(K) \otimes \mathcal{P}Env(\mathcal{W})$  preserves small colimits component-wise and the product of two  $\kappa$ -small  $\infty$ -categories is again  $\kappa$ -small.

**Corollary 3.33.** Let  $\kappa$  be a small regular cardinal. If a weakly bi-enriched  $\infty$ -category admits  $\kappa$ -small weighted colimits, it admits  $\kappa$ -small conical colimits and left and right tensors.

3.5. Weighted colimits via enriched coends. Next we discuss a further example of weighted colimits, the enriched coends ([9, Definition 5.70.]) with respect to enrichment in any symmetric monoidal  $\infty$ -category. Moreover we prove that general weighted colimits can be built from tensors and coends if such exist (Proposition 3.38).

In the following we will use symmetric monoidal  $\infty$ -categories. If  $\mathcal{V}^{\otimes} \to Ass$  is the pullback of a symmetric monoidal  $\infty$ -category  $\mathcal{V}^{\boxtimes} \to \operatorname{Comm}$ , there is a canonical monoidal equivalence  $(\mathcal{V}^{\operatorname{rev}})^{\otimes} \simeq \mathcal{V}^{\otimes}$ since  $\theta$  factors as Ass  $\xrightarrow{(-)^{\text{op}}}$  Ass  $\xrightarrow{\theta}$  Comm. The latter induces an equivalence between left and right (weakly) enriched  $\infty$ -categories. In particular, there is no need to distinguish between left and right morphism objects, which we call morphism objects.

Notation 3.34. Let  $\mathcal{V}^{\boxtimes} \to \text{Comm}$  be a symmetric monoidal  $\infty$ -category and  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes}$  a left  $\mathcal{V}$ enriched  $\infty$ -category. By [9, Notation 5.36.] there is a left  $\mathcal{V}$ -enriched functor

$$\operatorname{Mor}_{\mathcal{M}} : (\mathcal{M}^{\operatorname{op}} \otimes \mathcal{M})^{\circledast} \to \mathcal{V}^{\circledast}$$

assigning the morphism object.

**Definition 3.35.** Let  $\mathcal{V}^{\otimes} \to \text{Comm}$  be a symmetric monoidal  $\infty$ -category,  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes}, \mathcal{J}^{\otimes} \to \mathcal{V}^{\otimes}$  left pseudo-enriched  $\infty$ -categories and  $F : (\mathcal{J}^{op} \otimes \mathcal{J})^{\otimes} \to \mathcal{M}^{\otimes}$  a left  $\mathcal{V}$ -enriched functor.

- The V-enriched coend of F, denoted by ∫<sup>J</sup> F, is the Mor<sub>J<sup>op</sup></sub>-weighted colimit of F.
  The V-enriched end of F, denoted by ∫<sub>J</sub> F, is the Mor<sub>J</sub>-weighted limit of F.

**Remark 3.36.** The V-enriched end of F is the Mor<sub>d</sub>-weighted limit of F, which is the Mor<sub>d</sub>-weighted colimit of  $F^{op}: (\mathcal{J} \otimes \mathcal{J}^{op})^{\circledast} \to (\mathcal{M}^{op})^{\circledast}$  and so agrees with the  $\mathcal{V}$ -enriched coend of  $F^{op}$ .

**Construction 3.37.** Let  $\mathcal{V}^{\boxtimes} \to \text{Comm}$  be a presentably symmetric monoidal  $\infty$ -category and  $\mathcal{M}^{\circledast} \to$  $\mathcal{V}^{\otimes}$  a left  $\mathcal{V}$ -enriched  $\infty$ -category that admits left tensors. The composition of left  $\mathcal{V}$ -enriched functors

$$(\mathcal{V}^{\mathrm{op}}\otimes\mathcal{M}^{\mathrm{op}}\otimes\mathcal{M})^{\circledast}\xrightarrow{\mathrm{id}\otimes\mathrm{Mor}_{\mathcal{M}}}(\mathcal{V}^{\mathrm{op}}\otimes\mathcal{V})^{\circledast}\xrightarrow{\mathrm{Mor}_{\mathcal{V}}}\mathcal{V}^{\circledast}$$

corresponds to a left  $\mathcal{V}$ -enriched functor

$$(\mathcal{V} \otimes \mathcal{M})^{\circledast} \to (\operatorname{EnrFun}_{\mathcal{V}}(\mathcal{M}, \mathcal{V})^{\operatorname{op}})^{{\mathfrak{g}}}$$

that induces a functor  $\otimes : (\mathcal{V} \otimes \mathcal{M})^{\circledast} \to \mathcal{M}^{\circledast}$  by the existence of tensors. We obtain an equivalence

(7) 
$$\operatorname{Mor}_{\mathcal{M}}((-) \otimes (-), -) \simeq \operatorname{Mor}_{\mathcal{V}}(-, \operatorname{Mor}_{\mathcal{M}}(-, -))$$

of left  $\mathcal{V}$ -enriched functors  $(\mathcal{M}^{\mathrm{op}} \otimes \mathcal{M}^{\mathrm{op}} \otimes \mathcal{M})^{\circledast} \to \mathcal{V}^{\circledast}$ .

**Theorem 3.38.** Let  $\mathcal{V}^{\boxtimes} \to \text{Comm}$  be a presentably symmetric monoidal  $\infty$ -category,  $\mathcal{J}^{\circledast} \to \mathcal{V}^{\otimes}$  a left  $\mathcal{V}$ -enriched  $\infty$ -category,  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast}$  a left  $\mathcal{V}$ -enriched  $\infty$ -category that admits left tensors,  $F: \mathcal{J}^{\circledast} \to \mathcal{M}^{\circledast}$ a left  $\mathcal{V}$ -enriched functor and  $\mathbf{H} \in \mathcal{P}_{\mathcal{V}}(\mathcal{J})$ . There is a canonical equivalence

$$\operatorname{colim}^{\mathrm{H}}(\mathrm{F}) \simeq \int^{\mathcal{J}} \otimes \circ (|\mathrm{H}| \otimes \mathrm{F}).$$

*Proof.* Equivalence (7) gives for every  $X \in \mathcal{M}$  a canonical equivalence

(8) 
$$\operatorname{Mor}_{\mathcal{M}}(\otimes \circ (|H| \otimes F), X) \simeq \operatorname{Mor}_{\mathcal{V}} \circ (|H|^{\operatorname{op}} \otimes (\operatorname{Mor}_{\mathcal{M}}(-, X) \circ F^{\operatorname{op}}))$$

of  $\mathcal{V}$ -enriched functors  $(\mathcal{J}^{\mathrm{op}} \otimes \mathcal{J})^{\otimes} \to \mathcal{V}^{\otimes}$ . We obtain the following chain of natural equivalences:

$$\operatorname{Mor}_{\mathcal{M}}(\int^{\mathcal{J}} \otimes \circ (|H| \otimes F), X) \simeq \int_{\mathcal{J}} \operatorname{Mor}_{\mathcal{M}}(\otimes \circ (|H| \otimes F), X) \simeq \int_{\mathcal{J}} \operatorname{Mor}_{\mathcal{V}} \circ (|H|^{\operatorname{op}} \otimes (\operatorname{Mor}_{\mathcal{M}}(-, X) \circ F^{\operatorname{op}})) \\ \simeq \operatorname{Mor}_{\operatorname{EnrFun}_{\mathcal{V}, \emptyset}(\mathcal{J}^{\operatorname{op}}, \mathcal{V})}(|H|, \operatorname{Mor}_{\mathcal{M}}(-, X) \circ F^{\operatorname{op}}) \simeq \operatorname{Mor}_{\mathcal{P}_{\mathcal{V}}(\mathcal{J})}(H, F^{*}(X)),$$

where the first equivalence is by Corollary 3.65, the second equivalence is by (8), the third equivalence is by [9, Theorem 5.76.] and the last equivalence is by Theorem 2.154.

# 3.6. A Bousfield Kan formula for weighted colimits.

**Notation 3.39.** Let  $\mathcal{J}^{\otimes} \to \mathcal{V}^{\otimes}$  be a left enriched  $\infty$ -category,  $\mathcal{D}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a weakly bi-enriched  $\infty$ -category that admits right cotensors and  $Y \in \mathcal{J}$ . By [9, Proposition 5.23. (1)] the right  $\mathcal{W}$ -enriched functor  $\operatorname{EnrFun}_{\mathcal{V},\emptyset}(\mathcal{J},\mathcal{D})^{\otimes} \to \mathcal{D}^{\otimes}$  evaluating at Y admits a  $\mathcal{W}$ -enriched right adjoint  $\operatorname{Ran}_{Y}$  that sends  $Z \in \mathcal{D}$  to  $\operatorname{LMor}_{\mathcal{J}}(-,Y)Z$ .

In the following we use the notation of Theorem 2.154.

**Proposition 3.40.** Let  $\mathcal{D}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  be a weakly bi-enriched  $\infty$ -category that admits right cotensors,  $F : \mathcal{J}^{\circledast} \to \mathcal{D}^{\circledast}$  a left  $\mathcal{V}$ -enriched functor and  $H \in \mathcal{P}LEnv(\mathcal{J})$  a left weight on  $\mathcal{J}$ . For every  $X \in \mathcal{D}, W_1, ..., W_m \in \mathcal{W}, m \ge 0$  there is a canonical equivalence

 $Mul_{\mathcal{P}LEnv(\mathcal{D})}(F_{!}(H), W_{1}, ..., W_{m}; X) \simeq Mul_{EnrFun_{\mathcal{V}, \varnothing}(\mathcal{J}, \mathcal{D})}(F, W_{1}, ..., W_{m}; {}^{(-)}X \circ \mid H \mid).$ 

*Proof.* There is a canonical equivalence

$$\operatorname{Mul}_{\operatorname{PLEnv}(\mathcal{D})}(\operatorname{F}_{!}(\operatorname{H}), \operatorname{W}_{1}, ..., \operatorname{W}_{m}; \operatorname{X}) \simeq$$

 $Mul_{\mathrm{EnrFun}_{\mathcal{P}\mathrm{Env}(\mathcal{V}), \varnothing}(\mathcal{P}\mathrm{LEnv}(\mathcal{J}), \mathcal{P}\mathrm{LEnv}(\mathcal{D}))}(F_!, W_1, ..., W_m; {}^{(-)}\mathrm{X} \circ \mathcal{P}\mathrm{LEnv}(\mathcal{J})(-, H)) \simeq$ 

 $\operatorname{Mul}_{\operatorname{EnrFun}_{\mathcal{V},\sigma}(\mathcal{A},\mathcal{D})}(F, W_1, ..., W_m; {}^{(-)}X \circ |H|).$ 

**Corollary 3.41.** Let  $\mathcal{J}^{\circledast} \to \mathcal{V}^{\circledast}$  be a weakly left enriched  $\infty$ -category, H a left weight on  $\mathcal{J}$  and  $\mathcal{D}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  a weakly bi-enriched  $\infty$ -category that admits right cotensors and  $F : \mathcal{J}^{\circledast} \to \mathcal{D}^{\circledast}$  a left  $\mathcal{V}$ -enriched functor that admits a H-weighted colimit. There is a canonical equivalence

$$\operatorname{Mul}_{\mathcal{D}}(\operatorname{colim}^{H}(F), W_{1}, ..., W_{m}; X) \simeq \operatorname{Mul}_{\operatorname{EnrFun}_{\mathcal{V}, \sigma}(\mathcal{J}, \mathcal{D})}(F, W_{1}, ..., W_{m}; {}^{(-)}X \circ | H |).$$

*Proof.* For every  $X \in \mathcal{D}, V_1, ..., V_n \in \mathcal{V}, W_1, ..., W_m \in \mathcal{W}, n, m \ge 0$  there is a canonical equivalence

$$\operatorname{Mul}_{\mathcal{D}}(\operatorname{colim}^{H}(F), W_{1}, ..., W_{m}; X) \simeq \operatorname{Mul}_{\mathcal{P}\operatorname{LEnv}(\mathcal{D})}(F_{!}(H), W_{1}, ..., W_{m}; X) \simeq$$

 $\operatorname{Mul}_{\operatorname{EnrFuny}, \sigma(\mathcal{J}, \mathcal{D})}(F, W_1, ..., W_m; {}^{(-)}X \circ |H|).$ 

**Corollary 3.42.** Let  $\mathcal{J}^{\circledast} \to \mathcal{V}^{\circledast}$  be a weakly left enriched  $\infty$ -category, H a left weight on  $\mathcal{J}$  and  $\mathcal{D}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  a weakly bi-enriched  $\infty$ -category that admits right cotensors and H-weighted colimits. The right  $\mathcal{W}$ -enriched functor

$$\mathcal{D}^{\circledast} \xrightarrow{\operatorname{Ran}_{H}} \operatorname{EnrFun}_{\mathcal{P}\operatorname{Env}(\mathcal{V}),\varnothing}(\mathcal{P}\operatorname{LEnv}(\mathcal{J}), \bar{\mathcal{D}})^{\circledast} \xrightarrow{(-)_{|\mathcal{J}}} \operatorname{EnrFun}_{\mathcal{V},\varnothing}(\mathcal{J}, \mathcal{D})^{\circledast}, X \mapsto {}^{(-)}X \circ |H|$$

admits a W-enriched left adjoint that sends F to  $\operatorname{colim}^{H}(F)$ .

**Proposition 3.43.** Let  $\mathcal{J}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be a small weakly bi-enriched  $\infty$ -category,  $\mathcal{D}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$ a weakly bi-enriched  $\infty$ -category that admits left and right tensors and small conical colimits,  $F : \mathcal{J}^{\circledast} \to \mathcal{D}^{\circledast}$  a  $\mathcal{V}, \mathcal{W}$ -enriched functor and  $H \in \mathcal{PBEnv}(\mathcal{J})$ . Then  $\operatorname{colim}^{H}(F)$  is the colimit of a canonical simplicial object  $\mathcal{Y}$  in  $\mathcal{D}$  such that for every  $n \geq 0$  there is a canonical equivalence in  $\mathcal{N}$ :

$$\mathcal{Y}_{n} \simeq \operatorname{colim}_{Z_{1},...,Z_{n} \in \mathcal{J}^{\simeq}}(|H|(Z_{n})) \otimes \operatorname{LMor}_{\bar{\mathcal{J}}}(Z_{n-1},Z_{n}) \otimes ... \otimes \operatorname{LMor}_{\bar{\mathcal{J}}}(Z_{1},Z_{2}) \otimes F(Z_{1})$$

*Proof.* By [9, Proposition 5.26.] every  $\mathcal{V}, \mathcal{W}$ -enriched functor  $F : \mathcal{J}^{\circledast} \to \mathcal{D}^{\circledast}$  is the colimit of a canonical simplicial object  $\mathcal{X}$  in EnrFun<sub> $\mathcal{V},\mathcal{W}$ </sub> $(\mathcal{J},\mathcal{D})$  such that for every  $n \geq 0$  there is a canonical equivalence in EnrFun<sub> $\mathcal{V},\mathcal{W}$ </sub> $(\mathcal{J},\mathcal{D})$ :

 $\mathfrak{X}_n \simeq \operatorname{colim}_{Z_1, \dots, Z_n \in \mathcal{M}^{\simeq}} LMor_{\bar{\mathcal{A}}}(Z_n, -) \otimes LMor_{\bar{\mathcal{A}}}(Z_{n-1}, Z_n) \otimes \dots \otimes LMor_{\bar{\mathcal{A}}}(Z_2, Z_3) \otimes LMor_{\bar{\mathcal{A}}}(Z_1, Z_2) \otimes F(Z_1).$ 

By Proposition 3.23 there is a left adjoint functor  $\operatorname{colim}^{\mathrm{H}} : \operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}(\mathcal{J},\mathcal{D}) \to \mathcal{D}, F \mapsto \operatorname{colim}^{\mathrm{H}}(F)$ . Let  $\mathcal{Y} := \operatorname{colim}^{\mathrm{H}} \circ \mathcal{X}$ . Consequently, there are canonical equivalences:  $\operatorname{colim}^{\mathrm{H}}(F) \simeq \operatorname{colim}_{\lceil n \rceil \in \Delta^{\circ p}} \mathcal{Y}_n$  and

 $\mathcal{Y}_n \simeq \operatorname{colim}_{Z_1, \dots, Z_n \in \mathcal{J}^\simeq} \operatorname{colim}^H(\operatorname{LMor}_{\bar{\mathcal{J}}}(Z_n, -)) \otimes \operatorname{LMor}_{\bar{\mathcal{J}}}(Z_{n-1}, Z_n) \otimes \dots \otimes \operatorname{LMor}_{\bar{\mathcal{J}}}(Z_1, Z_2) \otimes F(Z_1)$ 

 $\simeq \operatorname{colim}_{Z_1,\ldots,Z_n \in \mathcal{J}^\simeq}(|\operatorname{H}|(Z_n)) \otimes \operatorname{LMor}_{\bar{\mathcal{J}}}(Z_{n-1},Z_n) \otimes \ldots \otimes \operatorname{LMor}_{\bar{\mathcal{J}}}(Z_1,Z_2) \otimes F(Z_1),$ 

where the last equivalence is by Corollary 3.42.

3.7. **Preservation of weighted colimits.** Next we prove that enriched left adjoints preserve weighted colimits. We first define what it means that an enriched functor preserves weighted colimits.

**Definition 3.44.** A small diagram is a triple  $(H, F, \psi)$  consisting of

- an enriched functor  $F: \mathcal{J}^{\otimes} \to \mathcal{M}^{\otimes}$  of totally small weakly bi-enriched  $\infty$ -categories,
- an object  $H \in \mathcal{P}BEnv(\mathcal{J})$ ,
- a morphism  $\psi : F_{!}(H) \to Y$  in  $\mathcal{PBEnv}(\mathcal{M})$ , where  $Y \in \mathcal{M}$ .

More precisely, we call  $(H, F, \psi)$  a small H-weighted diagram on  $\mathcal{M}$ . If  $\alpha, \beta$  are the maps of  $\infty$ -operads underlying  $F : \mathcal{J}^{\circledast} \to \mathcal{M}^{\circledast}$ , we call  $(\alpha, \beta, \mathcal{J}^{\circledast}, H)$  the weight underlying the diagram  $(H, F, \psi)$ . If  $\psi$ exhibits Y as the H-weighted colimit, we call  $(H, F, \psi)$  a small H-weighted colimit diagram on  $\mathcal{M}$ . A small left (right) diagram is a small diagram lying over a left (right) weight.

Definition 3.45. A diagram is a left (right) diagram if the underlying weight is a left (right) weight.

**Definition 3.46.** Let  $\kappa, \tau$  be small regular cardinals. A diagram is a left enriched, right enriched, bi-enriched, left  $\kappa$ -enriched, right  $\tau$ -enriched,  $\kappa, \tau$ -bi-enriched diagram, respectively, if the underlying weight is left enriched, right enriched, bi-enriched, left  $\kappa$ -enriched, right  $\tau$ -enriched.

**Notation 3.47.** An enriched functor  $\theta : \mathcal{M}^{\circledast} \to \mathcal{N}^{\circledast}$  lying over maps of  $\infty$ -operads  $\alpha', \beta'$  sends a diagram (H, F,  $\psi$ ) on  $\mathcal{M}$  lying over the weight ( $\alpha, \beta, \mathcal{J}^{\circledast}, H$ ) to the diagram

$$(\mathrm{H}, \theta \circ \mathrm{F}, \psi' : \theta_!(\mathrm{F}_!(\mathrm{H})) \xrightarrow{\theta_!(\psi)} \theta(\mathrm{Y}))$$

on  $\mathbb{N}$  lying over the weight  $(\alpha' \circ \alpha, \beta' \circ \beta, \mathcal{J}^{\circledast}, \mathbf{H})$ . We call the latter diagram the image of  $(\mathbf{H}, \mathbf{F}, \psi)$ under  $\theta$ . We call the weight  $(\alpha' \circ \alpha, \beta' \circ \beta, \mathcal{J}^{\circledast}, \mathbf{H})$  the image of  $(\alpha, \beta, \mathcal{J}^{\circledast}, \mathbf{H})$  under  $\alpha', \beta'$ .

**Definition 3.48.** Let  $\theta : \mathcal{M}^{\circledast} \to \mathcal{N}^{\circledast}$  be an enriched functor of totally small weakly bi-enriched  $\infty$ categories and  $(H, F : \mathcal{J}^{\circledast} \to \mathcal{M}^{\circledast}, \psi)$  a H-weighted colimit diagram on  $\mathcal{M}$ . We say that  $\theta$  preserves the
H-weighted colimit of F if  $\theta$  sends the H-weighted colimit diagram  $(H, F : \mathcal{J}^{\circledast} \to \mathcal{M}^{\circledast}, \psi)$  to a H-weighted
colimit diagram on  $\mathcal{N}$ .

**Definition 3.49.** Let  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}, \mathcal{N}^{\otimes} \to \mathcal{V}''^{\otimes} \times \mathcal{W}''^{\otimes}$  be totally small weakly bi-enriched  $\infty$ -categories and  $\theta : \mathcal{M}^{\otimes} \to \mathcal{N}^{\otimes}$  an enriched functor.

- For every totally small weight  $(\alpha, \beta, \mathcal{J}^{\circledast}, \mathrm{H})$  over  $\mathcal{V}, \mathcal{W}$  we say that  $\theta$  preserves  $(\alpha, \beta, \mathcal{J}^{\circledast}, \mathrm{H})$ -weighted colimits if  $\theta$  preserves the H-weighted colimit of any enriched functor  $\mathcal{J}^{\circledast} \to \mathcal{M}^{\circledast}$  lying over  $\alpha, \beta$ .
- For every collection ℋ of weights over 𝒱, 𝒱 we say that θ preserves ℋ-weighted colimits if θ preserves the (α, β, 𝔅<sup>®</sup>, H)-weighted colimit for every (α, β, 𝔅<sup>®</sup>, H) ∈ ℋ.
- We say that  $\theta$  preserves  $\kappa$ -small weighted colimits if  $\theta$  preserves the  $(\alpha, \beta, \mathcal{J}^{\circledast}, \mathbf{H})$ -weighted colimit for every  $\kappa$ -small weight  $(\alpha, \beta, \mathcal{J}^{\circledast}, \mathbf{H})$  over  $\mathcal{V}, \mathcal{W}$ .

Notation 3.50. For every weakly bi-enriched  $\infty$ -categories  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}, \mathcal{N}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  and collections  $\Lambda, \Lambda'$  of diagrams in  $\mathcal{M}, \mathcal{N}$ , respectively, let

$$\operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}^{\Lambda,\Lambda}(\mathcal{M},\mathcal{N}) \subset \operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}(\mathcal{M},\mathcal{N})$$

be the full subcategory of  $\mathcal{V}, \mathcal{W}$ -enriched functors sending diagrams of  $\Lambda$  to diagrams of  $\Lambda'$ . Let

$$\operatorname{EnrFun}_{\mathcal{V}}^{\Lambda}{}_{\mathcal{W}}(\mathcal{M},\mathcal{N}) \subset \operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}(\mathcal{M},\mathcal{N})$$

be the full subcategory of  $\mathcal{V}$ ,  $\mathcal{W}$ -enriched functors sending diagrams of  $\Lambda$  to weighted colimit diagrams.

**Notation 3.51.** Let  $\mathcal{V}^{\otimes} \to Ass, \mathcal{W}^{\otimes} \to Ass$  be  $\infty$ -operads,  $\mathcal{N}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a weakly bi-enriched  $\infty$ -category,  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes}$  a weakly left enriched  $\infty$ -category,  $\mathcal{O}^{\otimes} \to \mathcal{W}^{\otimes}$  a weakly right enriched  $\infty$ -category,  $\Lambda$  a collection of diagrams in  $\mathcal{M}$  and  $\Lambda'$  a collection of diagrams in  $\mathcal{O}$ . Let

$$\operatorname{EnrFun}_{\mathcal{V}, \mathcal{A}}^{\Lambda}(\mathcal{M}, \mathcal{N})^{\otimes} \subset \operatorname{EnrFun}_{\mathcal{V}, \mathcal{A}}(\mathcal{M}, \mathcal{N})^{\otimes}$$

be the full weakly right enriched subcategory of left  $\mathcal{V}$ -enriched functors  $\mathcal{M} \to \mathcal{N}$  sending diagrams of  $\Lambda$  to weighted colimit diagrams. Let

$$\operatorname{EnrFun}_{\varnothing,W}^{\Lambda'}(\mathcal{O},\mathcal{N})^{\circledast} \subset \operatorname{EnrFun}_{\varnothing,W}(\mathcal{O},\mathcal{N})^{\circledast}$$

be the full weakly left enriched subcategory of right W-enriched functors  $\mathcal{O} \to \mathcal{N}$  sending diagrams of  $\Lambda'$  to weighted colimit diagrams.

**Definition 3.52.** Let  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be a weakly bi-enriched  $\infty$ -category and  $\mathcal{H}$  a collection of weights over  $\mathcal{V}, \mathcal{W}$ . A collection  $\Lambda$  of diagrams in  $\mathcal{M}$  is  $\mathcal{H}$ -weighted if every diagram of  $\Lambda$  is  $\mathcal{H}$ -weighted for some  $\mathcal{H} \in \mathcal{H}$ .

**Notation 3.53.** Let  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be a weakly bi-enriched  $\infty$ -category. For every collection  $\mathcal{H}$  of weights over  $\mathcal{V}, \mathcal{W}$  let  $\Lambda^{(\mathcal{M}, \mathcal{H})}$  be the collection of H-weighted colimit diagrams in  $\mathcal{M}$  for some  $H \in \mathcal{H}$ .

**Remark 3.54.** Let  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}, \mathcal{N}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be weakly bi-enriched  $\infty$ -categories,  $\mathcal{H}$  a collection of weights over  $\mathcal{V}, \mathcal{W}$  and  $\Lambda$  a collection of  $\mathcal{H}$ -weighted diagrams in  $\mathcal{M}$ . Then  $\operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}^{\Lambda,\Lambda^{(\mathcal{N},\mathcal{H})}}(\mathcal{M},\mathcal{N}) = \operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}^{\Lambda,\Lambda^{(\mathcal{N},\mathcal{H})}}(\mathcal{M},\mathcal{N}).$ 

**Proposition 3.55.** Let  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}, \mathcal{N}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  be weakly bi-enriched  $\infty$ -categories and  $\phi: \mathcal{M}^{\circledast} \to \mathcal{N}^{\circledast}$  a  $\mathcal{V}, \mathcal{W}$ -enriched functor.

- (1) If  $\phi$  admits a  $\mathcal{V}, \mathcal{W}$ -enriched right adjoint,  $\phi$  preserves all weighted colimits.
- (2) If  $\phi$  admits a  $\mathcal{V}, \mathcal{W}$ -enriched left adjoint,  $\phi$  preserves all weighted limits.

*Proof.* We prove (1). (2) is similar. A  $\mathcal{V}, \mathcal{W}$ -enriched adjunction  $\phi : \mathcal{M}^{\circledast} \neq \mathcal{N}^{\circledast} : \gamma$  gives rise to a  $\mathcal{P}Env(\mathcal{V}), \mathcal{P}Env(\mathcal{W})$ -enriched adjunction  $\phi_{!} : \mathcal{P}Env(\mathcal{M})^{\circledast} \neq \mathcal{P}Env(\mathcal{N})^{\circledast} : \gamma_{!}$ . Let  $F : \mathcal{J}^{\circledast} \to \mathcal{M}^{\circledast}$  be an enriched functor,  $Y \in \mathcal{M}, H \in \mathcal{P}BEnv(\mathcal{J})$  and  $\psi : F_{!}(H) \to Y$  a map in  $\mathcal{P}BEnv(\mathcal{M})$  that exhibits Y as the H-weighted colimit of F. We like to see that the map  $\phi_{!}(\psi) : \phi_{!}(F_{!}(H)) \to \phi(Y)$  in  $\mathcal{P}BEnv(\mathcal{N})$  exhibits  $\phi(Y)$  as the H-weighted colimit of  $\phi \circ F$ , i.e. that for any  $Z \in \mathcal{N}, V_{1}, ..., V_{n} \in \mathcal{V}, W_{1}, ..., W_{m} \in \mathcal{W}$  for some  $n, m \geq 0$  the map

$$\operatorname{Mul}_{\mathcal{N}}(V_1, ..., V_n, \phi(Y), W_1, ..., W_m; Z) \rightarrow \operatorname{Mul}_{\mathcal{PBEnv}(\mathcal{N})}(V_1, ..., V_n, \phi_!(F_!(H)), W_1, ..., W_m; Z)$$

is an equivalence. The latter map identifies with the equivalence

 $Mul_{\mathcal{M}}(V_1,...,V_n,Y,W_1,...,W_m;\gamma(Z)) \rightarrow Mul_{\mathcal{P}BEnv(\mathcal{M})}(V_1,...,V_n,F_!(H),W_1,...,W_m;\gamma(Z)).$ 

**Definition 3.56.** Let  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be a weakly bi-enriched  $\infty$ -category,  $\Lambda$  a collection of diagrams in  $\mathcal{M}$  and  $\mathcal{H}$  be a collection of weights over  $\mathcal{V}, \mathcal{W}$ .

- (1) A full weakly bi-enriched subcategory  $\mathcal{N}^{\circledast} \subset \mathcal{M}^{\circledast}$  is closed under diagrams of  $\Lambda$  if for every diagram  $(H, \tau, \psi : \tau_!(H) \to Y)$  in  $\mathcal{M}$  the object  $Y \in \mathcal{M}$  belongs to  $\mathcal{N}$  if  $\tau : \mathcal{J}^{\circledast} \to \mathcal{M}^{\circledast}$  factors through  $\mathcal{N}^{\circledast}$ .
- (2) A full weakly bi-enriched subcategory  $\mathcal{N}^{\otimes} \subset \mathcal{M}^{\otimes}$  is closed under  $\mathcal{H}$ -weighted colimits if it is closed under the collection of  $\mathcal{H}$ -weighted colimit diagrams in  $\mathcal{M}$ .

**Example 3.57.** Let  $\alpha : F \to G : \mathcal{M}^{\otimes} \to \mathcal{N}^{\otimes}$  be a morphism of  $\operatorname{Enr}(\mathcal{M}, \mathcal{N})$  and  $T = (H, \tau : \mathcal{J}^{\otimes} \to \mathcal{M}^{\otimes}, \psi : \tau_{!}(H) \to Y)$  a diagram in  $\mathcal{M}$ . If F, G send T to a weighted colimit diagram and  $\alpha_{\tau(Z)}$  is an equivalence for every  $Z \in \mathcal{J}$ , then using Remark 3.8 the morphism  $\alpha_{Y}$  is an equivalence. In particular, the full subcategory of  $\mathcal{M}$  spanned by all  $X \in \mathcal{M}$  such that  $\alpha_{X}$  is an equivalence, is closed under all weighted colimits that are preserved by F, G.

3.8. Stability of weighted colimits under restriction of enrichment. Next we prove that the existence of weighted colimits is stable under pulling back the enrichment (Corollary 3.60).

**Lemma 3.58.** Let  $\alpha : \mathcal{V}^{\otimes} \to \mathcal{V}^{\otimes}, \beta : \mathcal{W}^{\otimes} \to \mathcal{W}^{\otimes}$  be maps of small  $\infty$ -operads,  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a totally small weakly bi-enriched  $\infty$ -category,  $F : \mathcal{J}^{\otimes} \to (\alpha, \beta)^* (\mathcal{M})^{\otimes} := \mathcal{V}^{\otimes} \times_{\mathcal{V}^{\otimes}} \mathcal{M}^{\otimes} \times_{\mathcal{W}^{\otimes}} \mathcal{W}^{\otimes}$  an enriched functor,  $H \in \mathcal{PBEnv}(\mathcal{J})$  and  $Y \in \mathcal{M}, \mathcal{V}'_1, ..., \mathcal{V}'_n \in \mathcal{V}', \mathcal{W}'_1, ..., \mathcal{W}'_m \in \mathcal{W}'$  for some  $n, m \ge 0$ . The projection  $\rho : (\alpha, \beta)^* (\mathcal{M})^{\otimes} \to \mathcal{M}^{\otimes}$  induces an equivalence

$$\begin{aligned} \operatorname{Mul}_{\mathcal{P}\operatorname{LEnv}((\alpha,\beta)^{*}(\mathcal{M}))}(V'_{1},...V'_{n},F_{!}(H),W'_{1},...,W'_{m};Y) \rightarrow \\ \operatorname{Mul}_{\mathcal{P}\operatorname{LEnv}(\mathcal{M})}(\alpha(V'_{1}),...,\alpha(V'_{n}),\rho_{!}(F_{!}(H)),\beta(W'_{1}),...,\beta(W'_{m});\rho(Y)). \end{aligned}$$

Proof. It is enough to prove that the unit  $Y \to \rho^*(\rho_!(Y)) \simeq \rho^*(\rho(Y))$  in  $\mathcal{PBEnv}((\alpha,\beta)^*(\mathcal{M}))$  is an equivalence. The unit induces at  $V_1 \otimes ... \otimes V_n \otimes Z \otimes W_1 \otimes ... \otimes W_m \in \mathrm{BEnv}((\alpha,\beta)^*(\mathcal{M}))$  the equivalence  $\mathrm{Mul}_{(\alpha,\beta)^*(\mathcal{M})}(V_1,...,V_n,Z,W_1,...,W_m;Y) \to \mathrm{Mul}_{\mathcal{M}}(\alpha(V_1),...,\alpha(V_n),\rho(Z),\beta(W_1),...,\beta(W_m);\rho(Y)).$ 

**Proposition 3.59.** Let  $\alpha : \mathcal{V}^{\otimes} \to \mathcal{V}^{\otimes}, \beta : \mathcal{W}^{\otimes} \to \mathcal{W}^{\otimes}$  be maps of small  $\infty$ -operads,  $\mathcal{J}^{\circledast} \to \mathcal{Q}^{\otimes} \times \mathcal{U}^{\otimes}, \mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  totally small weakly bi-enriched  $\infty$ -categories,  $H \in \mathcal{PBEnv}(\mathcal{J}), F : \mathcal{J}^{\circledast} \to (\alpha, \beta)^{*}(\mathcal{M})^{\circledast} := \mathcal{V}^{\otimes} \times_{\mathcal{V}^{\otimes}} \mathcal{M}^{\otimes} \times_{\mathcal{W}^{\otimes}} \mathcal{W}^{\otimes}$  an enriched functor,  $Y \in \mathcal{M}$  and  $\rho : (\alpha, \beta)^{*}(\mathcal{M})^{\circledast} \to \mathcal{M}^{\circledast}$  the projection.

(1) A morphism

$$\psi: \mathbf{F}_{!}(\mathbf{H}) \to \mathbf{Y}$$

in  $\mathcal{PBEnv}((\alpha,\beta)^*(\mathcal{M}))$  exhibits Y as the H-weighted colimit of F if

 $\rho_!(\psi):\rho_!(F_!(H)) \to \rho(Y)$ 

in  $\mathcal{PBEnv}(\mathcal{M})$  exhibits  $\rho(\mathbf{Y})$  as the H-weighted colimit of  $\rho \circ \mathbf{F}$ .

(2) The converse holds by the uniqueness of weighted colimits if the H-weighted colimit of F exists.

*Proof.* If  $\rho_!(\psi) : \rho_!(F_!(H)) \to \rho(Y)$  exhibits  $\rho(Y)$  as the H-weighted colimit of  $\rho \circ F$ , for every  $Z \in (\alpha, \beta)^*(\mathcal{M})$  and  $V'_1, \dots, V'_n \in \mathcal{V}', W'_1, \dots, \mathcal{W}'_m \in \mathcal{W}'$  for some  $n, m \ge 0$  the canonical map

$$\operatorname{Mul}_{\mathcal{M}}(\alpha(V'_{1}),...,\alpha(V'_{n}),\rho(Y),\beta(W'_{1}),...,\beta(W'_{m});\rho(Z)) \rightarrow$$

 $\operatorname{Mul}_{\operatorname{PLEnv}(\mathcal{M})}(\alpha(V_1'), ..., \alpha(V_n'), \rho_!(F_!(H)), \beta(W_1'), ..., \beta(W_m'); \rho(Z))$ 

is an equivalence. But this map canonically identifies with the map

$$\operatorname{Mul}_{(\alpha,\beta)^*(\mathcal{M})}(V'_1, \dots, V'_n, Y, W'_1, \dots, W'_m; Z) \to \operatorname{Mul}_{\mathcal{P}\operatorname{LEnv}((\alpha,\beta)^*(\mathcal{M}))}(V'_1, \dots, V'_n, F_!(H), W'_1, \dots, W'_m; Z) \xrightarrow{\sigma} \operatorname{Mul}_{\mathcal{P}\operatorname{LEnv}((\alpha,\beta)^*(\mathcal{M}))}(V'_1, \dots, V'_n, F_!(H), W'_1, \dots, W'_m; Z) \xrightarrow{\sigma} \operatorname{Mul}_{\mathcal{P}\operatorname{LEnv}((\alpha,\beta)^*(\mathcal{M}))}(V'_1, \dots, V'_n, F_!(H), W'_1, \dots, W'_m; Z) \xrightarrow{\sigma} \operatorname{Mul}_{\mathcal{P}\operatorname{LEnv}((\alpha,\beta)^*(\mathcal{M}))}(V'_1, \dots, V'_n, F_!(H), W'_1, \dots, W'_m; Z) \xrightarrow{\sigma} \operatorname{Mul}_{\mathcal{P}\operatorname{LEnv}((\alpha,\beta)^*(\mathcal{M}))}(V'_1, \dots, V'_n, F_!(H), W'_1, \dots, W'_m; Z) \xrightarrow{\sigma} \operatorname{Mul}_{\mathcal{P}\operatorname{LEnv}((\alpha,\beta)^*(\mathcal{M}))}(V'_1, \dots, V'_n, F_!(H), W'_1, \dots, W'_m; Z) \xrightarrow{\sigma} \operatorname{Mul}_{\mathcal{P}\operatorname{LEnv}((\alpha,\beta)^*(\mathcal{M}))}(V'_1, \dots, V'_n, F_!(H), W'_1, \dots, W'_m; Z) \xrightarrow{\sigma} \operatorname{Mul}_{\mathcal{P}\operatorname{LEnv}((\alpha,\beta)^*(\mathcal{M}))}(V'_1, \dots, V'_n, F_!(H), W'_1, \dots, W'_m; Z) \xrightarrow{\sigma} \operatorname{Mul}_{\mathcal{P}\operatorname{LEnv}((\alpha,\beta)^*(\mathcal{M}))}(V'_1, \dots, V'_n, F_!(H), W'_1, \dots, W'_m; Z) \xrightarrow{\sigma} \operatorname{Mul}_{\mathcal{P}\operatorname{LEnv}((\alpha,\beta)^*(\mathcal{M}))}(V'_1, \dots, V'_n, F_!(H), W'_1, \dots, W'_m; Z) \xrightarrow{\sigma} \operatorname{Mul}_{\mathcal{P}\operatorname{LEnv}((\alpha,\beta)^*(\mathcal{M}))}(V'_1, \dots, V'_n, F_!(H), W'_1, \dots, W'_m; Z) \xrightarrow{\sigma} \operatorname{Mul}_{\mathcal{P}\operatorname{LEnv}((\alpha,\beta)^*(\mathcal{M}))}(V'_1, \dots, V'_n, F_!(H), W'_1, \dots, W'_m; Z) \xrightarrow{\sigma} \operatorname{Mul}_{\mathcal{P}\operatorname{LEnv}((\alpha,\beta)^*(\mathcal{M}))}(V'_1, \dots, V'_n, F_!(H), W'_1, \dots, W'_m; Z) \xrightarrow{\sigma} \operatorname{Mul}_{\mathcal{P}\operatorname{LEnv}((\alpha,\beta)^*(\mathcal{M}))}(V'_1, \dots, V'_n, F_!(H), W'_1, \dots, W'_m; Z) \xrightarrow{\sigma} \operatorname{Mul}_{\mathcal{P}\operatorname{LEnv}((\alpha,\beta)^*(\mathcal{M}))}(V'_1, \dots, V'_n, F_!(H), W'_1, \dots, W'_m; Z)$$

$$\operatorname{Mul}_{\operatorname{PLEnv}(\mathcal{M})}(\alpha(V_1),...,\alpha(V_n),\rho_!(F_!(H)),\beta(W_1),...,\beta(W_m);\rho(Z)),$$

where  $\sigma$  is induced by  $\rho_!$ :  $\mathcal{PLEnv}((\alpha, \beta)^*(\mathcal{M}))^{\circledast} \to \mathcal{PLEnv}(\mathcal{M})^{\circledast}$ . By Lemma 3.58  $\rho$  is an equivalence. (2) immediately follows from 1. and Lemma 3.58.

**Corollary 3.60.** Let  $\tau : \mathcal{V}^{\otimes} \to \mathcal{V}^{\otimes}, \rho : \mathcal{W}^{\otimes} \to \mathcal{W}^{\otimes}$  be maps of small  $\infty$ -operads,  $\mathcal{J}^{\circledast} \to \Omega^{\otimes} \times \mathfrak{U}^{\otimes}$  a totally small weakly bi-enriched  $\infty$ -category and  $(\alpha, \beta, \mathcal{J}^{\circledast}, \mathbf{H})$  a weight over  $\mathcal{V}', \mathcal{W}'$ . For every totally small weakly bi-enriched  $\infty$ -category  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  that admits  $(\tau \circ \alpha, \rho \circ \beta, \mathcal{J}^{\circledast}, \mathbf{H})$ -weighted colimits the pullback  $\mathcal{V}^{\otimes} \times_{\mathcal{V}^{\otimes}} \mathcal{M}^{\otimes} \times_{\mathcal{W}^{\otimes}} \mathcal{W}^{\otimes}$  admits  $(\alpha, \beta, \mathcal{J}^{\circledast}, \mathbf{H})$ -weighted colimits and the projection  $\mathcal{V}^{\otimes} \times_{\mathcal{V}^{\otimes}} \mathcal{M}^{\otimes} \times_{\mathcal{W}^{\otimes}} \mathcal{M}^{\otimes} \times \mathcal{M}^{\otimes}$  sends  $(\alpha, \beta, \mathcal{J}^{\circledast}, \mathbf{H})$ -weighted colimits to  $(\tau \circ \alpha, \rho \circ \beta, \mathcal{J}^{\circledast}, \mathbf{H})$ -weighted colimits.

3.9. Existence of weighted colimits. In the following we prove an existence result for weighted colimits that splits the existence of weighted colimits in the existence of tensors and conical colimits (Proposition 3.63).

**Proposition 3.61.** Let  $\mathcal{J}^{\otimes} \to \mathcal{Q}^{\otimes} \times \mathcal{U}^{\otimes}$ ,  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be totally small weakly bi-enriched  $\infty$ -categories,  $F: \mathcal{J}^{\otimes} \to \mathcal{M}^{\otimes}$  an enriched functor lying over maps of  $\infty$ -operads  $\alpha: \mathcal{Q}^{\otimes} \to \mathcal{V}^{\otimes}$ ,  $\beta: \mathcal{U}^{\otimes} \to \mathcal{W}^{\otimes}$  and  $\mathcal{K} \subset Cat_{\infty}, \mathcal{T} \subset \mathcal{V}, \mathcal{T}' \subset \mathcal{W}$  full subcategories. Let  $\mathcal{H} \subset \mathcal{PBEnv}(\mathcal{J})$  be the smallest full subcategory containing  $\mathcal{J}$ , closed under K-indexed colimits for any  $K \in \mathcal{K}$  and such that for every  $Y \in \alpha^{-1}(\mathcal{T}) \subset \mathcal{Q}, Z \in \beta^{-1}(\mathcal{T}') \subset \mathcal{U}$  the functors  $Y \otimes (-), (-) \otimes Z : \mathcal{PBEnv}(\mathcal{J}) \to \mathcal{PBEnv}(\mathcal{J})$  preserves  $\mathcal{H}$ .

- (1) If  $\mathfrak{M}$  admits left tensors of objects  $Y \in \mathfrak{T}$  and right tensors of objects  $Z \in \mathfrak{T}'$  and K-indexed conical colimits for any  $K \in \mathfrak{K}$ , then  $\mathfrak{M}$  admits the H-weighted colimit of F for any  $H \in \mathfrak{H}$ .
- (2) An enriched functor φ : M<sup>®</sup> → N<sup>®</sup> preserves the H-weighted colimit of F for any H ∈ H if φ preserves left tensors of objects Y ∈ T and right tensors of objects Z ∈ T' and K-indexed conical colimits.

*Proof.* (1): Let  $\mathcal{H}' \subset \mathcal{PBEnv}(\mathcal{J})$  be the full subcategory spanned by those H ∈  $\mathcal{PBEnv}(\mathcal{J})$  such that  $\mathcal{M}$  admits the H-weighted colimit of F. Then  $\mathcal{H}'$  contains  $\mathcal{J}$ , is closed under K-indexed conical colimits for any K ∈  $\mathcal{K}$  and for every Y ∈  $\alpha^{-1}(\mathcal{T})$ , Z ∈  $\beta^{-1}(\mathcal{T}')$  the functor Y ⊗ (-), (-) ⊗ Z :  $\mathcal{PBEnv}(\mathcal{J}) \to \mathcal{PBEnv}(\mathcal{J})$  preserves  $\mathcal{H}'$ , where we use Remark 3.18. Thus  $\mathcal{H} \subset \mathcal{H}'$ .

(2): Let  $\mathcal{H}' \subset \mathcal{PBEnv}(\mathcal{J})$  be the full subcategory spanned by those  $H \in \mathcal{PBEnv}(\mathcal{J})$  such that  $\mathcal{M}$  admits the H-weighted colimit of F and  $\phi$  preserves the H-weighted colimit of F. Then  $\mathcal{H}'$  contains  $\mathcal{J}$ , is closed under K-indexed conical colimits for any  $K \in \mathcal{K}$ , and for every  $Y \in \alpha^{-1}(\mathcal{T}), Z \in \beta^{-1}(\mathcal{T}')$  the functor  $Y \otimes (-), (-) \otimes Z : \mathcal{PBEnv}(\mathcal{J}) \to \mathcal{PBEnv}(\mathcal{J})$  preserves  $\mathcal{H}'$ . Thus  $\mathcal{H} \subset \mathcal{H}'$ .

Proposition 3.61 implies the following corollary:

**Corollary 3.62.** Let  $\mathcal{J}^{\circledast} \to \mathcal{Q}^{\circledast} \times \mathcal{U}^{\circledast}, \mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  be totally small weakly bi-enriched  $\infty$ -categories,  $F: \mathcal{J}^{\circledast} \to \mathcal{M}^{\circledast}$  an enriched functor and  $\mathcal{K} \subset \mathsf{Cat}_{\infty}$  a full subcategory. Let  $\mathcal{H} \subset \mathsf{PBEnv}(\mathcal{J})$  be the smallest full subcategory containing  $\mathcal{J}$ , closed under K-indexed colimits for any  $K \in \mathcal{K}$  and such that for any  $Y \in \mathcal{Q}, Z \in \mathcal{U}$  the functors  $Y \otimes (-), (-) \otimes Z: \mathsf{PBEnv}(\mathcal{J}) \to \mathsf{PBEnv}(\mathcal{J})$  preserve  $\mathcal{H}$ .

- Assume that M admits left and right tensors and K-indexed colimits for any K ∈ K and that K-indexed colimits are preserved by the functors V ⊗ (-), (-) ⊗ W : M → M for every V ∈ V, W ∈ W. Then M admits the H-weighted colimit of F for any H ∈ H.
- (2) An enriched functor  $\phi : \mathcal{M}^{\circledast} \to \mathcal{N}^{\circledast}$  preserves the H-weighted colimit of F for any  $H \in \mathcal{H}$  if  $\phi$  preserves left and right tensors and K-indexed colimits.

We obtain the following proposition:

**Proposition 3.63.** Let  $\kappa$  be a small regular cardinal and  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a totally small weakly bi-enriched  $\infty$ -category.

- (1) Then  $\mathcal{M}$  admits  $\kappa$ -small weighted colimits if and only if  $\mathcal{M}$  admits left and right tensors and  $\kappa$ -small colimits and forming left and right tensors commutes with  $\kappa$ -small colimits.
- (2) Then  $\mathcal{M}$  admits  $\kappa$ -small left (right) weighted colimits if and only if  $\mathcal{M}$  admits left (right) tensors and  $\kappa$ -small colimits and forming left (right) tensors commutes with  $\kappa$ -small colimits.
- (3) An enriched functor  $\mathcal{M}^{\circledast} \to \mathcal{N}^{\circledast}$  preserves  $\kappa$ -small weighted colimits if and only if it preserves  $\kappa$ -small colimits and left and right tensors.

(4) An enriched functor  $\mathcal{M}^{\circledast} \to \mathcal{N}^{\circledast}$  preserves  $\kappa$ -small left (right) weighted colimits if and only if it preserves  $\kappa$ -small colimits and left (right) tensors.

*Proof.* We prove (1) and (3). The proofs of (2) and (4) are similar. (1): If  $\mathcal{M}$  admits left and right tensors and  $\kappa$ -small colimits and forming left and right tensors commutes with  $\kappa$ -small colimits, then  $\mathcal{M}$  admits  $\kappa$ -small weighted colimits by Corollary 3.62.

The converse is Corollary 3.33 and Remark 3.26. (3) follows like (1) from Corollary 3.62.  $\Box$ 

**Corollary 3.64.** Let  $\kappa$  be a small regular cardinal and  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  a totally small weakly bienriched  $\infty$ -category. Then  $\mathcal{M}$  admits  $\kappa$ -small weighted colimits if and only if  $\mathcal{M}$  admits  $\kappa$ -small weighted colimits of  $\mathcal{V}, \mathcal{W}$ -enriched functors  $K^{\circledast}_{\mathcal{V},\mathcal{W}} \to \mathcal{M}^{\circledast}$  for some  $\infty$ -category K.

**Corollary 3.65.** Let  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be a totally small weakly bi-enriched  $\infty$ -category. The embedding  $\mathcal{M}^{\otimes} \subset \mathcal{P}BEnv(\mathcal{M})^{\otimes}$  preserves weighted limits.

*Proof.* By Proposition 3.63 it is enough to show that the embedding  $\mathcal{M}^{\circledast} \subset \mathcal{P}BEnv(\mathcal{M})^{\circledast}$  preserves left and right cotensors and all conical limits. Let  $W \in \mathcal{W}, Y \in \mathcal{M}$  such that for every  $X \in \mathcal{M}, V_1, ..., V_n \in \mathcal{V}, W_1, ..., W_m \in \mathcal{W}$  the induced map

$$\operatorname{Mul}_{\mathcal{M}}(V_{1},...,V_{n},X,W_{1},...,W_{m};Y^{W}) \rightarrow \operatorname{Mul}_{\mathcal{M}}(V_{1},...,V_{n},X,W_{1},...,W_{m},W;Y)$$

is an equivalence. The latter map identifies with the map

$$\mathcal{P}BEnv(\mathcal{M})(V_1 \otimes ... \otimes V_n \otimes X \otimes W_1 \otimes ... \otimes W_m, Y^W) \to Mul_{\mathcal{P}BEnv(\mathcal{M})}(V_1 \otimes ... \otimes V_n \otimes X \otimes W_1 \otimes ... \otimes W_m, W; Y).$$

Hence the map  $\mathcal{PBEnv}(\mathcal{M})(T, Y^W) \to \mathrm{Mul}_{\mathcal{PBEnv}(\mathcal{M})}(T, W; Y)$  is an equivalence for every  $T \in \mathrm{BEnv}(\mathcal{M})$ . Since the functors  $\mathcal{PBEnv}(\mathcal{M})(-, Y^W)$ ,  $\mathrm{Mul}_{\mathcal{PBEnv}(\mathcal{M})}(-, W; Y) : \mathcal{PBEnv}(\mathcal{M})^{\mathrm{op}} \to S$  preserve small limits, the latter map is also an equivalence for every  $T \in \mathcal{PBEnv}(\mathcal{M})$ . So the right cotensor is preserved by the embedding  $\mathcal{M} \subset \mathcal{PBEnv}(\mathcal{M})$ . The case of the left cotensor is similar. We prove the case of conical limits.

Let  $F: K \to \mathcal{M}$  be a functor such that for every  $X \in \mathcal{M}, V_1, ..., V_n \in \mathcal{V}, W_1, ..., W_m \in \mathcal{W}$  the map

 $\operatorname{Mul}_{\mathcal{M}}(V_{1},...,V_{n},X,W_{1},...,W_{m};\operatorname{lim}(F)) \rightarrow \operatorname{lim}\operatorname{Mul}_{\mathcal{M}}(V_{1},...,V_{n},X,W_{1},...,W_{m};F(-))$ 

is an equivalence. The latter map identifies with the map

 $\mathcal{P}BEnv(\mathcal{M})(V_1 \otimes ... \otimes V_n \otimes X \otimes W_1 \otimes ... \otimes W_m, \lim(F)) \rightarrow$ 

 $\lim \mathbb{P}BEnv(\mathcal{M})(V_1 \otimes ... \otimes V_n \otimes X \otimes W_1 \otimes ... \otimes W_m, F(-)).$ 

Hence the map  $\mathcal{PBEnv}(\mathcal{M})(Y, \lim(F)) \to \lim \mathcal{PBEnv}(\mathcal{M})(Y, F(-))$  is an equivalence for every  $Y \in BEnv(\mathcal{M})$ . Since for every  $Z \in \mathcal{PBEnv}(\mathcal{M})$  the functor  $\mathcal{PBEnv}(\mathcal{M})(-, Z) : \mathcal{PBEnv}(\mathcal{M})^{\mathrm{op}} \to S$  preserves small limits, the latter map is also an equivalence for every  $Y \in \mathcal{PBEnv}(\mathcal{M})$ . So the limit of F is preserved by the embedding  $\mathcal{M} \subset \mathcal{PBEnv}(\mathcal{M})$ .

**Corollary 3.66.** Let  $(\mathcal{V}^{\otimes} \to Ass, S), (\mathcal{W}^{\otimes} \to Ass, T)$  be small localization pairs and  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a small S, T-bi-enriched  $\infty$ -category. The embedding  $\mathcal{M}^{\otimes} \subset \mathcal{PBEnv}(\mathcal{M})^{\otimes}_{S,T}$  preserves weighted limits.

- **Corollary 3.67.** (1) For every left quasi-enriched  $\infty$ -category  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  the embedding  $\mathcal{M}^{\circledast} \subset \mathcal{PBEnv}(\mathcal{M})^{\circledast}_{\mathrm{LEnr}}$  preserves left weighted limits and exhibits  $\mathcal{M}$  as free cocompletion under small left weighted colimits.
  - (2) For every right quasi-enriched ∞-category M<sup>®</sup> → V<sup>®</sup>×W<sup>®</sup> the embedding M<sup>®</sup> ⊂ PBEnv(M)<sup>®</sup><sub>REnr</sub> preserves right weighted limits and exhibits M as free cocompletion under small right weighted colimits.
  - (3) For every bi-quasi-enriched  $\infty$ -category  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  the embedding  $\mathcal{M}^{\circledast} \subset \mathrm{PBEnv}(\mathcal{M})^{\circledast}_{\mathrm{BEnr}}$  preserves weighted limits and exhibits  $\mathcal{M}$  as free cocompletion under small weighted colimits.

3.10. Adjoining weighted colimits. In this section we prove that weighted colimits can be universally adjoined (Proposition 3.85). We start with the following lemmas:

**Lemma 3.68.** Let  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  be a locally small weakly bi-enriched  $\infty$ -category that admits  $\mathcal{H}$ -weighted colimits for some set  $\mathcal{H}$  of small weights over  $\mathcal{V}, \mathcal{W}$  and  $\mathcal{B} \subset \mathcal{M}$  a small full subcategory. The full subcategory of  $\mathcal{M}$  generated by  $\mathcal{B}$  under  $\mathcal{H}$ -weighted colimits is small.

*Proof.* We need to see that the collection of equivalence classes of the full subcategory of  $\mathcal{M}$  generated by  $\mathcal{B}$  under  $\mathcal{H}$ -weighted colimits is a set. Since  $\mathcal{H}$  is a set and for every weight  $(\alpha, \beta, \mathcal{J}^{\circledast} \to \mathcal{V}^{(\otimes)} \times \mathcal{W}^{(\otimes)}, \mathbf{H} \in \mathcal{PBEnv}(\mathcal{J})) \in \mathcal{H}$  the  $\infty$ -category  $\mathcal{J}$  is small, there is a small regular cardinal  $\kappa$  such that for every weight  $(\alpha, \beta, \mathcal{J}^{\circledast} \to \mathcal{V}^{(\otimes)} \times \mathcal{W}^{(\otimes)}, \mathbf{H} \in \mathcal{PBEnv}(\mathcal{J})) \in \mathcal{H}$  the collection of equivalence classes of objects of  $\mathcal{J}$  is  $\kappa$ -small. We will first prove that the full subcategory of  $\mathcal{M}$  generated by  $\mathcal{B}$  under  $\mathcal{H}$ -weighted colimits is the  $\kappa$ -filtered colimit of the sequence  $(\mathcal{B}_{\lambda})_{\lambda < \kappa}$  of full subcategories of  $\mathcal{M}$  inductively defined by:

- (1)  $\mathcal{B}_0 \coloneqq \mathcal{B}$ ,
- (2) B<sub>λ+1</sub> is the union of B<sub>λ</sub> and the full subcategory of M of H-weighted colimits for some weight (α, β, J<sup>®</sup> → V'<sup>®</sup> × W'<sup>®</sup>, H ∈ PBEnv(J)) ∈ H of all enriched functors F : J<sup>®</sup> → M<sup>®</sup>, where the collection of equivalence classes in J is λ-small and F lies over α, β and carries J to B<sub>λ</sub>.
- (3) If  $\lambda < \kappa$  is a limit ordinal,  $\mathcal{B}_{\lambda} \coloneqq \cup_{\lambda' < \lambda} \mathcal{B}_{\lambda'}$ .

Indeed, for any enriched functor  $F : \mathcal{J}^{\circledast} \to \mathcal{M}^{\circledast}$  lying over  $\alpha, \beta$  that carries  $\mathcal{J}$  to  $\operatorname{colim}_{\lambda < \kappa} \mathcal{B}_{\lambda}$  for some weight  $(\alpha, \beta, \mathcal{J}^{\circledast} \to \mathcal{V}'^{\otimes} \times \mathcal{W}'^{\otimes}, H \in \mathcal{PBEnv}(\mathcal{J})) \in \mathcal{H}$  the collection of equivalence classes of objects of  $\mathcal{J}$  is  $\kappa$ -small and so a  $\kappa$ -compact object in the category of small sets. Since taking equivalence classes preserves filtered colimits as a functor from  $\infty$ -categories to sets, there is a  $\lambda < \kappa$  such that F carries  $\mathcal{J}$  to  $\mathcal{B}_{\lambda}$ . So the H-weighted colimit of F belongs to  $\mathcal{B}_{\lambda+1}$ . This shows that  $\operatorname{colim}_{\lambda < \kappa} \mathcal{B}_{\lambda} \subset \mathcal{M}$  is the smallest full subcategory of  $\mathcal{M}$  containing  $\mathcal{B}$  and closed under  $\mathcal{H}$ -weighted colimits.

Therefore we have to see that the colimit  $\operatorname{colim}_{\lambda < \kappa} \mathcal{B}_{\lambda}$  is small. We use induction: by assumption  $\mathcal{B}_0 = \mathcal{B}$  is small. The case of a limit ordinal is clear using that  $\kappa$  is small. For the case of a successor ordinal assume that  $\lambda < \kappa$  and  $\mathcal{B}_{\lambda}$  is small. We want to see that  $\mathcal{B}_{\lambda+1}$  is small. We fix the following notation: for any weight  $(\alpha, \beta, \mathcal{J}^{\circledast} \to \mathcal{V}'^{\otimes} \times \mathcal{W}'^{\otimes}, H \in \mathcal{P}BEnv(\mathcal{J})) \in \mathcal{H}$  let  $\mathcal{B}_{\lambda+1}^{H}$  be the full subcategory of  $\mathcal{M}$  spanned by the objects of  $\mathcal{B}_{\lambda}$  and the H-weighted colimits of all enriched functors  $F : \mathcal{J}^{\circledast} \to \mathcal{M}^{\circledast}$  that lie over  $\alpha, \beta$  and carry  $\mathcal{J}$  to  $\mathcal{B}_{\lambda}$ . Then  $\mathcal{B}_{\lambda+1} \subset \mathcal{M}$  is the union  $\cup_{H \in \mathcal{H}} \mathcal{B}_{\lambda+1}^{H}$ . Since  $\mathcal{H}$  is a set, it is enough to see that  $\mathcal{B}_{\lambda+1}^{H}$  is small. So it is enough to see that the full subcategory of  $\mathcal{M}$  spanned by the objects  $\operatorname{colim}^H(F)$  for  $F \in \operatorname{EnrFun}_{\mathcal{V}',\mathcal{W}'}(\mathcal{J}, (\alpha, \beta)^*(\mathcal{B}_{\lambda}))$  is small, where  $\mathcal{B}_{\lambda}^{\circledast} \subset \mathcal{M}^{\circledast}$  is the full weakly bi-enriched subcategory. This follows from the fact that the  $\infty$ -category  $\operatorname{EnrFun}_{\mathcal{V}',\mathcal{W}'}(\mathcal{J}, (\alpha, \beta)^*(\mathcal{B}_{\lambda}))$  is small because  $(\alpha, \beta)^*(\mathcal{B}_{\lambda})^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  and  $\mathcal{J}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  are small. The first is small since the multi-morphism spaces of  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  are small, and  $\mathcal{V}', \mathcal{W}', \mathcal{B}_{\Lambda}$  are small.

**Lemma 3.69.** Let  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be a small weakly bi-enriched  $\infty$ -category,  $\mathcal{H}$  a set of totally small weights over  $\mathcal{V}, \mathcal{W}$  and  $\Lambda$  a collection of  $\mathcal{H}$ -weighted diagrams in  $\mathcal{M}$ . Then  $\Lambda$  is a set.

Proof. Let  $(\alpha, \beta, \mathcal{J}^{\circledast} \to \mathcal{Q}^{\otimes} \times \mathcal{U}^{\otimes}, \mathrm{H}) \in \mathcal{H}$  be a weight. By assumption  $\mathcal{J}^{\circledast} \to \mathcal{Q}^{\otimes} \times \mathcal{U}^{\otimes}$  is a totally small weakly bi-enriched  $\infty$ -category so that the  $\infty$ -category  $\mathrm{EnrFun}_{\mathcal{Q},\mathcal{U}}(\mathcal{J}, (\alpha, \beta)^*(\mathcal{M}))$  is small. Since  $\mathcal{M}$  is small and for every enriched functor  $\mathrm{F} : \mathcal{J}^{\circledast} \to \mathcal{M}^{\circledast}$  lying over  $\alpha, \beta$  and  $\mathrm{Y} \in \mathcal{M}$  the space  $\mathcal{PB}\mathrm{Env}(\mathcal{J})(\mathrm{H}, \mathrm{F}^{*}(\mathrm{Y}))$  is small.

Notation 3.70. Let  $(\mathcal{V}^{\otimes} \to Ass, S)$ ,  $(\mathcal{W}^{\otimes} \to Ass, T)$  be small localization pairs,  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a small S, T-bi-enriched  $\infty$ -category and  $\Lambda$  a set of diagrams in  $\mathcal{M}$ .

- (1) Let  $\Omega_{\Lambda}$  be the set of morphisms  $\theta$  : colim<sup>H</sup>( $\iota \circ F$ )  $\rightarrow \iota(Y)$  in  $\mathcal{PBEnv}(\mathcal{M})_{S,T}$  adjoint to the morphism  $H \xrightarrow{\lambda} F^*(Y) \rightarrow (\iota \circ F)^*(\iota(Y))$  in  $\mathcal{PBEnv}(\mathcal{J})$  for some diagram ( $F : \mathcal{J}^{\circledast} \rightarrow \mathcal{M}^{\circledast}, Y, \lambda : H \rightarrow F^*(Y)) \in \Lambda$ , where  $\iota : \mathcal{M}^{\circledast} \rightarrow \mathcal{PBEnv}(\mathcal{M})_{S,T}^{\circledast}$  is the canonical embedding.
- (2) Let  $\mathfrak{Q}'_{\Lambda}$  be the set of morphisms in  $\mathcal{PBEnv}(\mathcal{M})^{\textcircled{\otimes}}_{S,T}$  of the form  $V_1 \otimes ... \otimes V_n \otimes \mathfrak{f} \otimes W_1 \otimes ... \otimes W_m$  for  $\mathfrak{f} \in \mathfrak{Q}_{\Lambda}$  and  $V_1, ..., V_n \in \mathcal{V}, \mathcal{W}_1, ..., W_m \in \mathcal{W}$  and  $n, m \ge 0$ .

Notation 3.71. Let  $(\mathcal{V}^{\otimes} \to Ass, S)$ ,  $(\mathcal{W}^{\otimes} \to Ass, T)$  be small localization pairs,  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a small S, T-bi-enriched  $\infty$ -category and  $\Lambda$  a set of diagrams in  $\mathcal{M}$ .

(1) Let

$$\mathcal{P}BEnv_{\Lambda}(\mathcal{M})^{\circledast}_{S,T} \subset \mathcal{P}BEnv(\mathcal{M})^{\circledast}_{S,T}$$

be the full weakly bi-enriched subcategory spanned the  $\mathfrak{Q}'_\Lambda\text{-local objects.}$ 

(2) Let

 $\widetilde{\operatorname{PBEnv}}_\Lambda(\mathcal{M})^{\circledast}_{S,T} \coloneqq \mathcal{V}^{\otimes} \times_{S^{-1}\operatorname{PEnv}(\mathcal{V})^{\otimes}} \operatorname{PBEnv}_\Lambda(\mathcal{M})^{\circledast}_{S,T} \times_{T^{-1}\operatorname{PEnv}(\mathcal{W})^{\otimes}} \mathcal{W}^{\otimes}.$ 

**Remark 3.72.** Note that  $\mathcal{PBEnv}_{\Lambda}(\mathcal{M})^{\circledast}_{S,T} \subset \mathcal{PBEnv}(\mathcal{M})^{\circledast}_{S,T}$  is the full weakly bi-enriched subcategory spanned by all X such that the  $\mathcal{W}, \mathcal{V}$ -enriched functor  $\operatorname{Mor}_{\mathcal{PBEnv}(\mathcal{M})_{S,T}}((-)_{|\mathcal{M}}, X) : (\mathcal{M}^{\operatorname{op}})^{\circledast} \to (S^{-1}\mathcal{PEnv}(\mathcal{V}) \otimes T^{-1}\mathcal{PEnv}(\mathcal{W}))^{\circledast}$  sends every diagram of  $\Lambda$  to a weighted limit diagram.

**Lemma 3.73.** Let  $(\mathcal{V}^{\otimes} \to Ass, S)$ ,  $(\mathcal{W}^{\otimes} \to Ass, T)$  be small localization pairs,  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a small S, T-bi-enriched  $\infty$ -category and  $\Lambda$  a set of diagrams in  $\mathcal{M}$ .

- (1) The full weakly bi-enriched subcategory  $\mathcal{PBEnv}_{\Lambda}(\mathcal{M})^{\circledast}_{S,T} \subset \mathcal{PBEnv}(\mathcal{M})^{\circledast}_{S,T}$  is an accessible  $S^{-1}\mathcal{PEnv}(\mathcal{V}), T^{-1}\mathcal{PEnv}(\mathcal{W})$ -enriched localization.
- (2) The restricted localization functor  $\mathcal{M}^{\circledast} \subset \mathcal{PBEnv}(\mathcal{M})^{\circledast}_{S,T} \xrightarrow{L} \mathcal{PBEnv}_{\Lambda}(\mathcal{M})^{\circledast}_{S,T}$  sends diagrams of  $\Lambda$  to weighted colimit diagrams.

*Proof.* (1): Let  $\iota : \mathcal{M}^{\circledast} \to \mathcal{P}BEnv(\mathcal{M})^{\circledast}_{S,T}$  be the canonical embedding. The saturated class generated by  $\mathcal{Q}'_{\Lambda}$  is closed under small colimits and so closed under the S<sup>-1</sup> $\mathcal{P}Env(\mathcal{V}), T^{-1}\mathcal{P}Env(\mathcal{W})$ -biaction. Consequently, because  $\mathcal{P}BEnv(\mathcal{M})^{\circledast}_{S,T} \to S^{-1}\mathcal{P}Env(\mathcal{V})^{\otimes} \times T^{-1}\mathcal{P}Env(\mathcal{W})^{\otimes}$  is a presentably bitensored ∞-category, the full weakly bi-enriched subcategory  $\mathcal{P}BEnv_{\Lambda}(\mathcal{M})^{\circledast}_{S,T} \subset \mathcal{P}BEnv(\mathcal{M})^{\circledast}_{S,T}$  is an accessible S<sup>-1</sup> $\mathcal{P}Env(\mathcal{V}), T^{-1}\mathcal{P}Env(\mathcal{W})$ -enriched localization.

(2): The induced morphism  $H \xrightarrow{\lambda} F^*(Y) \to (L \circ \iota \circ F)^*(L(\iota(Y)))$  is adjoint to the equivalence

$$\operatorname{colim}^{\mathrm{H}}(\mathrm{L}\circ\iota\circ\mathrm{F})\simeq\mathrm{L}(\operatorname{colim}^{\mathrm{H}}(\iota\circ\mathrm{F}))\xrightarrow{\mathrm{L}(\theta)}\mathrm{L}(\iota(\mathrm{Y}))$$

where  $\theta \in \Omega$ , and so exhibits  $L(\iota(Y))$  as the H-weighted colimit of  $L \circ \iota \circ F$ .

Notation 3.74. Let  $(\mathcal{V}^{\otimes} \to Ass, S)$ ,  $(\mathcal{W}^{\otimes} \to Ass, T)$  be small localization pairs,  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a small S, T-bi-enriched  $\infty$ -category,  $\mathcal{H}$  a set of totally small weights over  $\mathcal{V}, \mathcal{W}$  and  $\Lambda$  a collection of  $\mathcal{H}$ -weighted diagrams in  $\mathcal{M}$ . By Lemma 3.69 (1) the collection  $\Lambda$  is a set. Let

$$\mathcal{P}BEnv^{\mathcal{H}}_{\Lambda}(\mathcal{M})^{\circledast}_{S,T} \subset \mathcal{P}\widetilde{BEnv}_{\Lambda}(\mathcal{M})^{\circledast}_{S,T}$$

be the smallest full weakly bi-enriched subcategory containing the essential image of the restricted  $\mathcal{V}, \mathcal{W}$ -enriched localization functor  $\mathcal{M}^{\circledast} \to \mathcal{P}\widetilde{\operatorname{BEnv}}_{\Lambda}(\mathcal{M})^{\circledast}_{S,T}$  and closed under  $\mathcal{H}$ -weighted colimits.

If  $S = T = \emptyset$ , we drop S, T from the notation. If  $\Lambda = \emptyset$ , we drop  $\Lambda$  from the notation.

**Remark 3.75.** We obtain a  $\mathcal{V}, \mathcal{W}$ -enriched functor  $\mathcal{M}^{\otimes} \to \mathcal{P}BEnv_{\Lambda}^{\mathcal{H}}(\mathcal{M})_{S,T}^{\otimes}$ .

**Remark 3.76.** Lemma 3.73 implies that  $\mathcal{PBEnv}_{\Lambda}(\mathcal{M})^{\circledast}_{S,T} \to S^{-1}\mathcal{PEnv}(\mathcal{V})^{\otimes} \times T^{-1}\mathcal{PEnv}(\mathcal{W})^{\otimes}$  is a presentably bitensored  $\infty$ -category. Hence  $\mathcal{P}\widetilde{BEnv}_{\Lambda}(\mathcal{M})^{\circledast}_{S,T} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  is a S, T-bi-enriched  $\infty$ -category.

48

 $\square$ 

by Proposition 2.140 that admits small weighted colimits. Thus  $\mathcal{PBEnv}^{\mathcal{H}}_{\Lambda}(\mathcal{M})^{\textcircled{B}}_{S,T} \to \mathcal{V}^{\bigotimes} \times \mathcal{W}^{\bigotimes}$  is a small S, T-bi-enriched  $\infty$ -category that admits  $\mathcal{H}$ -weighted colimits by Lemma 3.68.

**Remark 3.77.** By Lemma 3.73 (2) the restricted localization functor  $\mathcal{M}^{\circledast} \to \mathcal{P}BEnv_{\Lambda}(\mathcal{M})^{\circledast}_{S,T}$  and so also  $\mathcal{M}^{\circledast} \to \mathcal{P}BEnv^{\mathcal{H}}_{\Lambda}(\mathcal{M})^{\circledast}_{S,T}$  send diagrams of  $\Lambda$  to  $\mathcal{H}$ -weighted colimit diagrams.

**Remark 3.78.** If  $\Lambda$  consists of weighted colimit diagrams, every object of  $\mathcal{M}$  belongs to  $\mathcal{PBEnv}_{\Lambda}(\mathcal{M})_{S,T}$ . So the restricted localization functor  $\mathcal{M}^{\circledast} \to \mathcal{PBEnv}_{\Lambda}(\mathcal{M})^{\circledast}_{S,T}$  and so  $\mathcal{M}^{\circledast} \to \mathcal{PBEnv}^{\mathcal{H}}_{\Lambda}(\mathcal{M})^{\circledast}_{S,T}$  are embeddings.

For the next notation we use the following terminology:

**Definition 3.79.** A weakly bi-enriched  $\infty$ -category  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  exhibits  $\mathcal{M}$  as

- locally left pseudo-enriched in  $\mathcal{V}$  if  $\mathcal{V}^{\otimes} \to Ass$  admits a tensor unit and the pullback  $Ass \times_{\mathcal{V}^{\otimes}} \mathcal{M}^{\otimes} \times_{\mathcal{W}^{\otimes}} \mathcal{Q}^{\otimes} \to Ass$  along the unique map  $\mathcal{Q}^{\otimes} \to \mathcal{W}^{\otimes}$  and the unique map  $Ass \to \mathcal{V}^{\otimes}$  preserving the tensor unit exhibits  $\mathcal{M}$  as left pseudo-enriched.
- locally right pseudo-enriched in  $\mathcal{W}$  if  $\mathcal{W}^{\otimes} \to Ass$  admits a tensor unit and the pullback  $\emptyset^{\otimes} \times_{\mathcal{V}^{\otimes}} \mathcal{M}^{\otimes} \times_{\mathcal{W}^{\otimes}} Ass \to Ass$  along the unique map  $\emptyset^{\otimes} \to \mathcal{V}^{\otimes}$  and the unique map  $Ass \to \mathcal{W}^{\otimes}$  preserving the tensor unit exhibits  $\mathcal{M}$  as right pseudo-enriched.
- locally bi-pseudo-enriched in  $\mathcal{V}, \mathcal{W}$  if  $\mathcal{V}^{\otimes} \to Ass, \mathcal{W}^{\otimes} \to Ass$  admit a tensor unit and the pullback  $Ass \times_{\mathcal{V}^{\otimes}} \mathcal{M}^{\otimes} \times_{\mathcal{W}^{\otimes}} Ass \to Ass \times Ass$  along the unique maps  $Ass \to \mathcal{V}^{\otimes}, Ass \to \mathcal{W}^{\otimes}$  preserving the tensor unit exhibits  $\mathcal{M}$  as bi-pseudo-enriched.

 $\begin{array}{l} \textbf{Proposition 3.80. Let } n \geq 1 \ and \ (\mathcal{V}_i^{\otimes} \rightarrow Ass, S_i), \ (\mathcal{W}_i^{\otimes} \rightarrow Ass, T_i) \ small \ localization \ pairs \ for \ 1 \leq i \leq n. \\ Let \ \mathcal{M}_i^{\otimes} \rightarrow \mathcal{V}_i^{\otimes} \times \mathcal{W}_i^{\otimes} \ be \ small \ bi-S_i, T_i-enriched \ \infty-categories \ such \ that \ for \ every \ 1 \leq i \leq n \ the \ weakly \\ bi-enriched \ \infty-category \ \mathcal{M}_i^{\otimes} \rightarrow \mathcal{V}_i^{\otimes} \times \mathcal{W}_i^{\otimes} \ is \ locally \ bi-pseudo-enriched \ if \ there \ is \ some \ 1 \leq j \leq n \ with \ j \neq i \\ such \ that \ S_j \ or \ T_j \ do \ not \ only \ contain \ equivalences. \ Let \ \mathcal{N}^{\otimes} \rightarrow \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes} \ be \ a \ weakly \ bi-enriched \ \infty-category, \ where \ \mathcal{V}^{\otimes} := \mathcal{V}_1^{\otimes} \times_{Ass} \ \mathcal{N}_n^{\otimes}, \mathcal{W}^{\otimes} := \mathcal{W}_1^{\otimes} \times_{Ass} \ \mathcal{N}_n^{\otimes}, \mathcal{W}^{\otimes} := \mathcal{W}_1^{\otimes} \times_{Ass} \ \mathcal{N}_n^{\otimes}, \mathcal{W}^{\otimes} := \mathcal{W}_1^{\otimes} \times_{Ass} \ \mathcal{N}_n^{\otimes}, \mathcal{W}^{\otimes} \ same \ that \ for \ every \ 1 \leq i \leq n \ the \\ pullback \ \mathcal{N}_i^{\otimes} := \mathcal{V}_i^{\otimes} \times_{\mathcal{W}^{\otimes}} \ \mathcal{W}_i^{\otimes} \ along \ \tau_i : \mathcal{V}_i^{\otimes} \rightarrow \mathcal{V}^{\otimes}, \tau_i : \mathcal{W}_i^{\otimes} \rightarrow \mathcal{W}^{\otimes} \ is \ a \ S_i, T_i-bi-enriched \ \infty-category \\ that \ admits \ left \ and \ right \ tensors \ and \ small \ colimits \ preserved \ by \ forming \ left \ and \ right \ tensors. \end{array}$ 

(1) The induced functor

$$\mathrm{EnrFun}_{\mathcal{V},\mathcal{W}}(\prod_{1\leq i\leq n}\mathcal{P}\widetilde{\mathrm{BEnv}}(\mathcal{M}_i)_{S_i,T_i},\mathcal{N}) \to \mathrm{EnrFun}_{\mathcal{V},\mathcal{W}}(\prod_{1\leq i\leq n}\mathcal{M}_i,\mathcal{N})$$

admits a fully faithful left adjoint that lands in the full subcategory

$$\mathrm{LinFun}_{\mathcal{V},\mathcal{W}}^{\mathrm{L},\mathrm{L}}(\prod_{1\leq i\leq n} \mathcal{P}\widetilde{\mathrm{BEnv}}(\mathcal{M}_i)_{\mathrm{S}_i,\mathrm{T}_i},\mathcal{N})$$

of  $\mathcal{V}, \mathcal{W}$ -linear functors that admit a right adjoint component-wise. (2) The following induced functor is an equivalence:

$$\mathrm{LinFun}_{\mathcal{V},\mathcal{W}}^{\mathrm{L},\mathrm{L}}(\prod_{1\leq i\leq n}\mathcal{P}\widetilde{\mathrm{BEnv}}(\mathcal{M}_i)_{\mathrm{S}_i,\mathrm{T}_i},\mathcal{N}) \to \mathrm{EnrFun}_{\mathcal{V},\mathcal{W}}(\prod_{1\leq i\leq n}\mathcal{M}_i,\mathcal{N}).$$

*Proof.* (2) follows immediately from (1) because the functor of (2) is conservative. (1): The existence of a fully faithful left adjoint in (1) follows from [9, Proposition 2.62.]. Let  $1 \le i \le n$  and  $X := (X_j \in \mathcal{PBEnv}(\mathcal{M}_j)_{S_i,T_i})_{1 \le j \le n, j \ne i}$  a family of objects. We like to see that the composition

$$\begin{split} \mathrm{EnrFun}_{\mathcal{V},\mathcal{W}}(\prod_{1\leq i\leq n}\mathcal{M}_{i},\mathcal{N}) &\to \mathrm{EnrFun}_{\mathcal{V},\mathcal{W}}(\prod_{1\leq i\leq n}\mathcal{P}\widetilde{\mathrm{BEnv}}(\mathcal{M}_{i})_{\mathrm{S}_{i},\mathrm{T}_{i}},\mathcal{N}) \\ & \xrightarrow{\mathrm{ev}_{\mathrm{X}}} \mathrm{EnrFun}_{\mathcal{V}_{i},\mathcal{W}_{i}}(\mathcal{P}\widetilde{\mathrm{BEnv}}(\mathcal{M}_{i})_{\mathrm{S}_{i},\mathrm{T}_{i}},\mathcal{N}_{i}) \end{split}$$

lands in LinFun<sup>L</sup><sub> $\mathcal{V}_i,\mathcal{W}_i$ </sub> (PBEnv( $\mathcal{M}_i$ )<sub>S<sub>i</sub>,T<sub>i</sub></sub>,  $\mathcal{N}_i$ ). By [10, Proposition 8.29. (5), Proposition 3.42.] the latter functor factors as

$$\mathrm{EnrFun}_{\mathcal{V},\mathcal{W}}(\prod_{1\leq i\leq n}\mathcal{M}_{i},\mathcal{N}) \rightarrow \mathrm{EnrFun}_{\mathcal{V},\mathcal{W}}(\prod_{1\leq j< i}\mathcal{P}\widetilde{\mathrm{BEnv}}(\mathcal{M}_{j})_{S_{j},T_{j}}\times\mathcal{M}_{i}\times\prod_{i< j\leq n}\mathcal{P}\widetilde{\mathrm{BEnv}}(\mathcal{M}_{j})_{S_{j},T_{j}},\mathcal{N}) \xrightarrow{\mathrm{ev}_{X}} \mathcal{P}\widetilde{\mathrm{BEnv}}(\mathcal{M}_{j})_{S_{j},T_{j}}$$

 $\operatorname{EnrFun}_{\mathcal{V}_{i},\mathcal{W}_{i}}(\mathcal{M}_{i},\mathcal{N}_{i}) \to \operatorname{EnrFun}_{\mathcal{V}_{i},\mathcal{W}_{i}}(\mathcal{P}BEnv(\mathcal{M}_{i})_{S_{i},T_{i}},\mathcal{N}_{i}).$ 

So we can reduce to the case n = 1, where we use that  $\mathcal{N}_i^{\otimes} \to \mathcal{V}_i^{\otimes} \times \mathcal{W}_i^{\otimes}$  is a  $S_i, T_i$ -bi-enriched  $\infty$ -category that admits left and right tensors and small colimits preserved by forming left and right tensors.

So let n = 1. By Corollary 2.140 and [9, Lemma 4.36.] the functor  $\mathcal{N}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  is the pullback of a bitensored  $\infty$ -category  $\overline{\mathcal{N}^{\circledast}} \to S^{-1}\mathcal{P}Env(\mathcal{V})^{\otimes} \times T^{-1}\mathcal{P}Env(\mathcal{W})^{\otimes}$  compatible with small colimits. By Corollary 2.140 the functor

 $\beta : \operatorname{EnrFun}_{S^{-1}\mathcal{P}\operatorname{Env}(\mathcal{V}), T^{-1}\mathcal{P}\operatorname{Env}(\mathcal{W})}(\mathcal{P}\operatorname{B}\operatorname{Env}(\mathcal{M})_{S,T}, \overline{\mathcal{N}}) \to \operatorname{EnrFun}_{\mathcal{V}, \mathcal{W}}(\mathcal{P}\operatorname{B}\operatorname{Env}(\mathcal{M})_{S,T}, \mathcal{N})$ 

is an equivalence. Since  $\mathcal{P}\text{Env}(\mathcal{V})$  is generated by  $\mathcal{V}$  under small colimits and the tensor product, also the localization  $S^{-1}\mathcal{P}\text{Env}(\mathcal{V})$  is generated by  $\mathcal{V}$  under small colimits and the tensor product. This implies that the functor

 $\alpha: \mathrm{LinFun}_{\mathrm{S}^{-1}\mathcal{P}\mathrm{Env}(\mathcal{V}), \mathrm{T}^{-1}\mathcal{P}\mathrm{Env}(\mathcal{W})}(\mathcal{P}\mathrm{B}\mathrm{Env}(\mathcal{M})_{\mathrm{S},\mathrm{T}}, \bar{\mathcal{N}}) \to \mathrm{LinFun}_{\mathcal{V},\mathcal{W}}^{\mathrm{L}}(\mathcal{P}\widetilde{\mathrm{B}\mathrm{Env}}(\mathcal{M})_{\mathrm{S},\mathrm{T}}, \mathcal{N})$ 

is the pullback of the functor  $\beta$  and so an equivalence, too. By Proposition 2.149 the composition

$$\operatorname{EnrFun}_{S^{-1}\mathcal{P}\operatorname{Env}(\mathcal{V}), T^{-1}\mathcal{P}\operatorname{Env}(\mathcal{W})}(\mathcal{P}\operatorname{B}\operatorname{Env}(\mathcal{M})_{S,T}, \widetilde{\mathcal{N}}) \xrightarrow{\mathcal{P}} \operatorname{EnrFun}_{\mathcal{V}, \mathcal{W}}(\mathcal{P}\operatorname{B}\operatorname{Env}(\mathcal{M})_{S,T}, \mathcal{N})$$

 $\rightarrow \operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}(\mathcal{M},\mathcal{N})$ 

admits a fully faithful left adjoint that lands in the full subcategory

LinFun<sup>L</sup><sub>S<sup>-1</sup>PEnv( $\mathcal{V}$ ), T<sup>-1</sup>PEnv( $\mathcal{W}$ ) (PBEnv( $\mathcal{M}$ )<sub>S,T</sub>,  $\overline{\mathcal{N}}$ ).</sub>

**Notation 3.81.** For every  $n \ge 1$  and  $1 \le i \le n$  let  $\mathcal{V}_i^{\otimes} \to \operatorname{Ass}, \mathcal{V}_i^{\otimes} \to \operatorname{Ass}$  be small  $\infty$ -operads and  $\mathcal{H}_i$  a set of weights over  $\mathcal{V}_i, \mathcal{W}_i$ . Assume that for every  $1 \le i \le n$  the  $\infty$ -operad  $\mathcal{V}_i^{\otimes} \to \operatorname{Ass}$  admits a tensor unit if for every  $1 \le j \le n, j \ne i$  the set  $\mathcal{H}_j$  does not only consist of right weights. Assume that for every  $1 \le i \le n$  the  $\infty$ -operad  $\mathcal{W}_i^{\otimes} \to \operatorname{Ass}$  admits a tensor unit if for every  $1 \le j \le n, j \ne i$  the set  $\mathcal{H}_j$  does not only consist of right weights. Assume that for every  $1 \le i \le n$  the  $\infty$ -operad  $\mathcal{W}_i^{\otimes} \to \operatorname{Ass}$  admits a tensor unit if for every  $1 \le j \le n, j \ne i$  the set  $\mathcal{H}_j$  does not only consist of left weights. We define a set  $\underline{\mathcal{H}}_i$  of weights over  $\mathcal{V}_1 \times \ldots \times \mathcal{V}_n, \mathcal{W}_1 \times \ldots \times \mathcal{W}_n$  the following way: if  $\mathcal{H}_i$  does not only consist of left weights and does not only consist of right weights,  $\underline{\mathcal{H}}_i$  is the image of  $\mathcal{H}_i$  under the maps of  $\infty$ -operads (using Lemma 2.21):

(9) 
$$\{1\} \times \dots \times \{1\} \times \mathcal{V}_{i} \times \{1\} \times \dots \times \{1\} \to \mathcal{V}_{1} \times \dots \times \mathcal{V}_{n},$$

(10) 
$$\{\mathbb{1}\} \times \dots \times \{\mathbb{1}\} \times \mathcal{W}_i \times \{\mathbb{1}\} \times \dots \times \{\mathbb{1}\} \to \mathcal{W}_1 \times \dots \times \mathcal{W}_n$$

If  $\mathcal{H}_i$  only consists of left weights,  $\underline{\mathcal{H}}_i$  is the image of  $\mathcal{H}_i$  under the map of  $\infty$ -operads (9). If  $\mathcal{H}_i$  only consists of right weights,  $\underline{\mathcal{H}}_i$  is the image of  $\mathcal{H}_i$  under the map of  $\infty$ -operads (10). If  $\mathcal{H}_i$  is empty,  $\underline{\mathcal{H}}_i$  is empty. Let

$$\mathcal{H}_1\boxtimes \ldots \boxtimes \mathcal{H}_n\coloneqq \bigcup_{i=1}^n \underline{\mathcal{H}}_i.$$

If n = 0, we agree that  $\mathcal{H}_1 \boxtimes ... \boxtimes \mathcal{H}_n$  is empty.

**Notation 3.82.** For every  $n \ge 1$  and  $1 \le i \le n$  let  $\mathcal{M}_i^{\circledast} \to \mathcal{V}_i^{\circledast} \times \mathcal{W}_i^{\circledast}$  be a totally small weakly bi-enriched  $\infty$ -category that admits  $\mathcal{H}_i$ -weighted colimits for some set  $\mathcal{H}_i$  of weights over  $\mathcal{V}_i$ ,  $\mathcal{W}_i$  and  $\Lambda_i$  a collection of  $\mathcal{H}_i$ -weighted diagrams in  $\mathcal{M}_i$ . Assume that for every  $1 \le i \le n$  the weakly bi-enriched  $\infty$ -category  $\mathcal{M}_i^{\circledast} \to \mathcal{V}_i^{\circledast} \times \mathcal{W}_i^{\circledast}$  is a weakly locally bi-pseudo-enriched  $\infty$ -category if for every  $1 \le j \le n, j \ne i$  the set  $\mathcal{H}_j$  does not only consist of left weights and does not only consist of right weights. Assume that for every  $1 \le i \le n$  the weakly bi-enriched  $\infty$ -category  $\mathcal{M}_i^{\circledast} \to \mathcal{V}_i^{\circledast} \times \mathcal{W}_i^{\circledast}$  is a locally left (right) pseudo-enriched

 $\infty$ -category if for every  $1 \leq j \leq n, j \neq i$  the set  $\mathcal{H}_j$  only consists of left (right) weights. For every  $1 \leq i \leq n$  we define a set  $\underline{\Lambda}_i$  of H-weighted diagrams in  $\mathcal{M}_1 \times \ldots \times \mathcal{M}_n$  for some  $H \in \underline{\mathcal{H}}_i$  the following way:  $\underline{\Lambda}_i$  is the image of  $\Lambda_i$  under all enriched functors of the form

$$(11) \qquad \qquad \{X_1\}\times\ldots\times\{X_{i-1}\}\times \mathcal{M}_i\times\{X_{i+1}\}\times\ldots\times\{X_n\} \to \mathcal{M}_1\times\ldots\times \mathcal{M}_n$$

for some objects  $X_j \in \mathcal{M}_j$  for  $1 \leq j \leq n, j \neq i$ . If  $\Lambda_i$  is empty,  $\underline{\Lambda}_i$  is empty. Let

$$\Lambda_1 \boxtimes \ldots \boxtimes \Lambda_n \coloneqq \bigcup_{i=1}^n \underline{\Lambda}_i.$$

If n = 0, we agree that  $\Lambda_1 \boxtimes ... \boxtimes \Lambda_n$  is empty.

**Notation 3.83.** Let  $n \ge 1$  and for every  $1 \le i \le n$  let  $\mathcal{V}_i^{\otimes} \to Ass$  be a small  $\infty$ -operad that admits a tensor unit if there is some  $1 \le j \le n$  with  $j \ne i$  such that  $S_j$  does not contain only equivalences. Set  $\mathcal{V}^{\otimes} := \mathcal{V}_1^{\otimes} \times_{Ass} \dots \times_{Ass} \mathcal{V}_n^{\otimes}$ . For every  $1 \le i \le n$  let  $S_i$  be a collection of morphisms of  $\mathcal{P}Env(\mathcal{V}_i)$ . Let

$$S_1 \boxtimes ... \boxtimes S_n$$

be the collection of morphisms g of  $\mathcal{P}Env(\mathcal{V})$  such that there is an  $1 \leq i \leq n$  and a morphism  $f \in S_i$  that is sent to g by the left adjoint monoidal functor  $(\tau_i)_! : \mathcal{P}Env(\mathcal{V}_i)^{\otimes} \to \mathcal{P}Env(\mathcal{V})^{\otimes}$  induced by the map of  $\infty$ -operads  $\tau_i : \mathcal{V}_i^{\otimes} \to \mathcal{V}^{\otimes}$  that is the identity on the i-th factor and the constant map preserving the tensor unit on every other factor.

**Notation 3.84.** Let  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\circledast}, \mathcal{N}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be weakly bi-enriched  $\infty$ -categories,  $\mathcal{H}$  a collection of weights over  $\mathcal{V}, \mathcal{W}, \mathcal{H}'$  a collection of weights over  $\mathcal{V}', \mathcal{W}'$  and  $\Lambda$  a collection of  $\mathcal{H}$ -weighted diagrams in  $\mathcal{M}$ . Let S be a collection of morphisms of  $\mathcal{P}Env(\mathcal{V})$  and T a collection of morphisms of  $\mathcal{P}Env(\mathcal{W})$ . Let

$$\operatorname{EnrFun}_{\mathrm{S},\mathrm{S}',\mathrm{T},\mathrm{T}',\mathcal{H},\mathcal{H}'}^{\Lambda}(\mathcal{M},\mathcal{N}) \subset \operatorname{EnrFun}_{\mathrm{S},\mathrm{S}',\mathrm{T},\mathrm{T}'}(\mathcal{M},\mathcal{N})$$

be the full subcategory of enriched functors sending diagrams of  $\Lambda$  to weighted colimit diagrams and lying over maps of  $\infty$ -operads that send weights of  $\mathcal{H}$  to weights of  $\mathcal{H}'$ .

 $\begin{array}{l} \textbf{Proposition 3.85. Let } n \geq 1. \ \textit{For } 1 \leq i \leq n \ \textit{let } (\mathcal{V}_i^{\otimes} \rightarrow Ass, S_i), (\mathcal{W}_i^{\otimes} \rightarrow Ass, T_i), (\mathcal{V}'^{\otimes} \rightarrow Ass, S'), (\mathcal{W}'^{\otimes} \rightarrow Ass, T') \ \textit{be small localization pairs, } \mathcal{H}_i \ \textit{a set of totally small weights over } \mathcal{V}_i, \mathcal{W}_i \ \textit{and } \mathcal{H}' \ \textit{a collection of weights over } \mathcal{V}', \mathcal{W}'. \ \textit{Let } \mathcal{M}_i^{\circledast} \rightarrow \mathcal{V}_i^{\otimes} \times \mathcal{W}_i^{\otimes} \ \textit{be small } S_i, T_i \text{-}\textit{bi-enriched } \infty \text{-}\textit{categories, } \Lambda_i \ \textit{a collection of } \mathcal{H}_i \text{-}\textit{weighted diagrams in } \mathcal{M}_i \ \textit{and } \mathcal{N}^{\circledast} \rightarrow \mathcal{V}'^{\otimes} \times \mathcal{W}'^{\otimes} \ \textit{a S'}, T' \text{-}\textit{bi-enriched } \infty \text{-}\textit{category that admits } \mathcal{H}' \text{-} weighted \ \textit{colimits. Let } \mathcal{H} \coloneqq \mathcal{H}_1 \boxtimes ... \boxtimes \mathcal{H}_n, S \coloneqq S_1 \boxtimes ... \boxtimes S_n, T \coloneqq T_1 \boxtimes ... \boxtimes T_n. \ \textit{Assume that the following conditions hold for every } 1 \leq i \leq n : \end{array}$ 

- (1)  $\mathcal{M}_{i}^{\circledast} \to \mathcal{V}_{i}^{\circledast} \times \mathcal{W}_{i}^{\circledast}$  is a locally left pseudo-enriched  $\infty$ -category if there is some  $1 \leq j \leq n$  with  $j \neq i$  such that  $S_{j}$  does not contain only equivalences or  $\Lambda_{j}$  does not only consist of right diagrams.
- (2)  $\mathcal{M}_{i}^{\circledast} \to \mathcal{V}_{i}^{\circledast} \times \mathcal{W}_{i}^{\circledast}$  is a locally right pseudo-enriched  $\infty$ -category if there is some  $1 \leq j \leq n$  with  $j \neq i$  such that  $T_{j}$  does not contain only equivalences or  $\Lambda_{j}$  does not only consist of left diagrams.

The induced functor

$$\gamma_{\mathcal{N}}: EnrFun_{S,S',T,T',\mathcal{H},\mathcal{H}'}^{\mathcal{M}_{1}}(\prod_{1 \leq i \leq n} \mathcal{P}BEnv_{\Lambda_{i}}^{\mathcal{H}_{i}}(\mathcal{M}_{i})_{S_{i},T_{i}}, \mathcal{N}) \rightarrow EnrFun_{S,S',T,T',\mathcal{H},\mathcal{H}'}^{\Lambda_{1} \otimes \ldots \otimes \Lambda_{n}}(\prod_{1 \leq i \leq n} \mathcal{M}_{i}, \mathcal{N})$$

is an equivalence. The induced functor

$$\delta_{\mathcal{N}}: EnrFun_{S,S',T,T',\mathcal{H},\mathcal{H}'}(\prod_{1 \leq i \leq n} \mathcal{P}BEnv^{\mathcal{H}_{i}}(\mathcal{M}_{i})_{S_{i},T_{i}}, \mathcal{N}) \rightarrow EnrFun_{S,S',T,T',\mathcal{H},\mathcal{H}'}(\prod_{1 \leq i \leq n} \mathcal{M}_{i}, \mathcal{N})$$

admits a fully faithful left adjoint that lands in  $\operatorname{EnrFun}_{S,S',T,T',\mathcal{H},\mathcal{H}'}^{\mathcal{H}_{1}}(\prod_{1\leq i\leq n} \mathcal{P}\operatorname{BEnv}_{\Lambda_{i}}^{\mathcal{H}_{i}}(\mathcal{M}_{i})_{S_{i},T_{i}}, \mathcal{N}).$ 

Proof. We first reduce to the case that  $\mathcal{N}^{\circledast} \to \mathcal{V}'^{\circledast} \times \mathcal{W}'^{\circledast}$  admits large weighted colimits. By Remarks 3.77, 3.78 there is a  $\mathcal{V}', \mathcal{W}'$ -enriched embedding  $\mathcal{N}^{\circledast} \to \widehat{\mathcal{P}BEnv}_{\Lambda^{\mathcal{H}'}}(\mathcal{N})^{\circledast}_{S',T'}$  that preserves  $\mathcal{H}'$ -weighted colimits (Lemma 3.73 (2)) into a bi-S', T'-enriched  $\infty$ -category that admits large weighted colimits

(Remark 3.76). The functor  $\gamma_{\mathcal{N}}$  is the pullback of  $\gamma_{\widehat{\mathcal{P}BEnv}_{\Lambda^{\mathcal{H}'}}(\mathcal{N})_{S',T'}}$  because  $\mathcal{PBEnv}_{\Lambda_{i}}^{\mathcal{H}_{i}}(\mathcal{M}_{i})_{S_{i},T_{i}}$  is generated by  $L_{i}(\mathcal{M}_{i})$  under  $\mathcal{H}_{i}$ -weighted colimits. Moreover if  $\delta_{\mathcal{P}\overline{BEnv}_{\Lambda^{\mathcal{H}'}}(\mathcal{N})_{S',T'}}$  admits a fully faithful left adjoint that lands in  $\operatorname{EnrFun}_{S,S',T,T',\mathcal{H},\mathcal{H}'}(\prod_{1\leq i\leq n}\mathcal{P}BEnv^{\mathcal{H}_{i}}(\mathcal{M}_{i})_{S_{i},T_{i}},\widehat{\mathcal{P}BEnv}_{\Lambda^{\mathcal{H}'}}(\mathcal{N})_{S',T'})$ , the left adjoint sends  $\operatorname{EnrFun}_{S,S',T,T',\mathcal{H},\mathcal{H}'}(\prod_{1\leq i\leq n}\mathcal{M}_{i},\mathcal{N})$  to  $\operatorname{EnrFun}_{S,S',T,T',\mathcal{H},\mathcal{H}'}(\prod_{1\leq i\leq n}\mathcal{P}BEnv^{\mathcal{H}_{i}}(\mathcal{M}_{i})_{S_{i},T_{i}},\mathcal{N})$  because  $\mathcal{P}BEnv^{\mathcal{H}_{i}}(\mathcal{M}_{i})_{S_{i},T_{i}}$  is generated by  $L_{i}(\mathcal{M}_{i})$  under  $\mathcal{H}_{i}$ -weighted colimits. So in this case the fully faithful left adjoint of  $\delta_{\mathcal{P}\overline{BEnv}_{\Lambda^{\mathcal{H}'}}(\mathcal{N})_{S',T'}}$  restricts to a fully faithful left adjoint of  $\delta_{\mathcal{N}}$  that lands in  $\operatorname{EnrFun}_{S,S',T,T',\mathcal{H},\mathcal{H}'}(\prod_{1\leq i\leq n}\mathcal{P}BEnv^{\mathcal{H}_{i}}(\mathcal{M}_{i})_{S_{i},T_{i}})$ . This way we can assume that  $\mathcal{N}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  admits large weighted colimits.

Let  $\overline{\mathcal{PBEnv}}_{\Lambda_{i}}^{\mathcal{H}_{i}}(\mathcal{M}_{i})_{S_{i},T_{i}}^{\circledast} \subset \mathcal{PBEnv}_{\Lambda_{i}}^{\mathcal{H}_{i}}(\mathcal{M}_{i})_{S_{i},T_{i}}^{\circledast}$  be the preimage of  $\mathcal{PBEnv}_{\Lambda_{i}}^{\mathcal{H}_{i}}(\mathcal{M}_{i})_{S_{i},T_{i}}^{\circledast}$  under the left adjoint  $L_{i}: \mathcal{PBEnv}(\mathcal{M}_{i})_{S_{i},T_{i}}^{\circledast} \to \mathcal{PBEnv}_{\Lambda_{i}}(\mathcal{M}_{i})_{S_{i},T_{i}}^{\circledast}$ . The full subcategory  $\overline{\mathcal{PBEnv}}_{\Lambda_{i}}^{\mathcal{H}_{i}}(\mathcal{M}_{i})_{S_{i},T_{i}}^{\circledast}$  is closed in  $\mathcal{PBEnv}(\mathcal{M}_{i})_{S_{i},T_{i}}^{\circledast}$  under  $\mathcal{H}_{i}$ -weighted colimits because  $\mathcal{PBEnv}_{\Lambda_{i}}^{\mathcal{H}_{i}}(\mathcal{M}_{i})_{S_{i},T_{i}}^{\circledast}$  is closed in  $\mathcal{PBEnv}_{\Lambda_{i}}(\mathcal{M}_{i})_{S_{i},T_{i}}^{\circledast}$  is closed in  $\mathcal{PBEnv}_{\Lambda_{i}}(\mathcal{M}_{i})_{S_{i},T_{i}}^{\circledast}$  restricts to a  $\mathcal{V}_{i}, \mathcal{W}_{i}$ -localization  $L_{i}: \mathcal{PBEnv}(\mathcal{M}_{i})_{S_{i},T_{i}}^{\circledast} \neq \mathcal{PBEnv}_{\Lambda_{i}}(\mathcal{M}_{i})_{S_{i},T_{i}}^{\circledast}$  relative to  $\mathcal{V}_{i}^{\circledast} \times \mathcal{W}_{i}^{\circledast}$  restricts to a  $\mathcal{V}_{i}, \mathcal{W}_{i}$ -localization  $L_{i}': \overline{\mathcal{PBEnv}}_{\Lambda_{i}}(\mathcal{M}_{i})_{S_{i},T_{i}}^{\circledast} \neq \mathcal{PBEnv}_{\Lambda_{i}}(\mathcal{M}_{i})_{S_{i},T_{i}}^{\circledast}$ . Precomposition with  $L_{i}'$  induces an embedding

$$\operatorname{EnrFun}_{S,S',T,T',\mathcal{H},\mathcal{H}'}^{\mathcal{M}_{1}}(\prod_{1\leq i\leq n}\mathcal{P}BEnv_{\Lambda_{i}}^{\mathcal{H}_{i}}(\mathcal{M}_{i})_{S_{i},T_{i}},\mathcal{N}) \hookrightarrow \operatorname{EnrFun}_{S,S',T,T',\mathcal{H},\mathcal{H}'}^{\mathcal{M}_{1}}(\prod_{1\leq i\leq n}\overline{\mathcal{P}BEnv_{\Lambda_{i}}^{\mathcal{H}_{i}}}(\mathcal{M}_{i})_{S_{i},T_{i}},\mathcal{N})$$

whose essential image is the full subcategory  $\operatorname{EnrFun}_{S,S',T,T',\mathcal{H},\mathcal{H}'}^{\mathcal{H}_{1}}(\prod_{1 \leq i \leq n} \overline{\mathcal{P}BEnv}_{\Lambda_{i}}^{\mathcal{H}_{i}}(\mathcal{M}_{i})_{S_{i},T_{i}}, \mathcal{N})'$  of enriched functors inverting all morphisms inverted by  $L \coloneqq \prod_{1 \leq i \leq n} L_{i}$ . The functor  $\gamma_{\mathcal{N}}$  factors as

$$\begin{split} \mathrm{Enr}\mathrm{Fun}_{S,S',\mathrm{T},\mathrm{T}',\mathcal{H},\mathcal{H}'}^{\mathcal{H}_{1}}(\prod_{1\leq i\leq n}\mathcal{P}\mathrm{BEnv}_{\Lambda_{i}}^{\mathcal{H}_{i}}(\mathcal{M}_{i})_{S_{i},\mathrm{T}_{i}},\mathcal{N}) \simeq \mathrm{Enr}\mathrm{Fun}_{S,S',\mathrm{T},\mathrm{T}',\mathcal{H},\mathcal{H}'}^{\mathcal{H}_{1}}(\prod_{1\leq i\leq n}\overline{\mathcal{P}\mathrm{BEnv}}_{\Lambda_{i}}^{\mathcal{H}_{i}}(\mathcal{M}_{i})_{S_{i},\mathrm{T}_{i}},\mathcal{N})' \\ \xrightarrow{\beta} \mathrm{Enr}\mathrm{Fun}_{S,S',\mathrm{T},\mathrm{T}',\mathcal{H},\mathcal{H}'}^{\Lambda_{1}\boxtimes\ldots\boxtimes\Lambda_{n}}(\prod_{1\leq i\leq n}\mathcal{M}_{i},\mathcal{N}). \end{split}$$

The functors  $\gamma_{\mathcal{N}}$  and so  $\beta$  are conservative because  $\mathcal{PBEnv}_{\Lambda_i}^{\mathcal{H}_i}(\mathcal{M}_i)_{S_i,T_i}$  is generated by  $L_i(\mathcal{M}_i)$  under  $\mathcal{H}_i$ -weighted colimits. By [9, Proposition 5.16. (1)] the induced functors

$$\begin{split} &\operatorname{EnrFun}_{S,S',T,T'}(\prod_{1\leq i\leq n}\overline{\mathcal{P}\mathrm{BEnv}}_{\Lambda_{i}}^{\mathcal{H}_{i}}(\mathcal{M}_{i})_{S_{i},T_{i}},\mathcal{N}) \to \operatorname{EnrFun}_{S,S',T,T'}(\prod_{1\leq i\leq n}\mathcal{M}_{i},\mathcal{N}),\\ &\operatorname{EnrFun}_{S,S',T,T'}(\prod_{1\leq i\leq n}\mathcal{P}\overline{\mathrm{BEnv}}(\mathcal{M}_{i})_{S_{i},T_{i}},\mathcal{N}) \to \operatorname{EnrFun}_{S,S',T,T'}(\prod_{1\leq i\leq n}\overline{\mathcal{P}\mathrm{BEnv}}_{\Lambda_{i}}^{\mathcal{H}_{i}}(\mathcal{M}_{i})_{S_{i},T_{i}},\mathcal{N}) \end{split}$$

admit fully faithful left adjoints  $\phi, \psi$ , respectively, since  $\mathcal{N}^{\circledast} \to \mathcal{V}'^{\otimes} \times \mathcal{W}'^{\otimes}$  admits large weighted colimits. Consequently, it will be enough to verify that the functor  $\phi$  takes  $\operatorname{EnrFun}_{S,S',T,T',\mathcal{H},\mathcal{H}'}^{\Lambda_1\boxtimes\ldots\boxtimes\Lambda_n}(\prod_{1\leq i\leq n}\mathcal{M}_i,\mathcal{N})$  to  $\operatorname{EnrFun}_{S,S',T,T',\mathcal{H},\mathcal{H}'}^{\Lambda'_1\boxtimes\ldots\boxtimes\Lambda'^{\mathfrak{K}_n}}(\prod_{1\leq i\leq n}\overline{\mathcal{P}BEnv}_{\Lambda_i}^{\mathcal{H}_i}(\mathcal{M}_i)_{S_i,T_i},\mathcal{N})'$ . Because  $\mathcal{N}^{\circledast} \to \mathcal{V}'^{\otimes} \times \mathcal{W}'^{\otimes}$  is bi-S', T'-enriched, by Propositions 3.80 and 3.61 the induced functor

$$EnrFun_{S,S',T,T'}(\prod_{1\leq i\leq n} \mathcal{P}\widetilde{BEnv}(\mathcal{M}_i)_{S_i,T_i}, \mathcal{N}) \rightarrow EnrFun_{S,S',T,T'}(\prod_{1\leq i\leq n} \mathcal{M}_i, \mathcal{N})$$

admits a fully faithful left adjoint  $\rho$  that lands in the full subcategory of enriched functors preserving small weighted colimits component-wise, which send diagrams of  $\Lambda^{\mathcal{H}_1} \boxtimes ... \boxtimes \Lambda^{\mathcal{H}_n}$  to weighted colimit diagrams. By adjointness the functor  $\rho$  factors as fully faithful left adjoints

$$\begin{split} \mathrm{EnrFun}_{\mathrm{S},\mathrm{S}',\mathrm{T},\mathrm{T}'}(\prod_{1\leq i\leq n}\mathcal{M}_{i},\mathcal{N}) &\xrightarrow{\phi} \mathrm{EnrFun}_{\mathrm{S},\mathrm{S}',\mathrm{T},\mathrm{T}'}(\prod_{1\leq i\leq n}\overline{\mathcal{P}\mathrm{BEnv}}_{\Lambda_{i}}^{\mathcal{H}_{i}}(\mathcal{M}_{i})_{\mathrm{S}_{i},\mathrm{T}_{i}},\mathcal{N}) \\ &\xrightarrow{\psi} \mathrm{EnrFun}_{\mathrm{S},\mathrm{S}',\mathrm{T},\mathrm{T}'}(\prod_{1\leq i\leq n}\mathcal{P}\widetilde{\mathrm{BEnv}}(\mathcal{M}_{i})_{\mathrm{S}_{i},\mathrm{T}_{i}},\mathcal{N}) \end{split}$$

so that  $\phi$  factors as  $\rho$  followed by the induced functor

$$\operatorname{EnrFun}_{S,S',T,T'}(\prod_{1\leq i\leq n} \mathcal{P}\widetilde{\operatorname{BEnv}}(\mathcal{M}_i)_{S_i,T_i}, \mathcal{N}) \to \operatorname{EnrFun}_{S,S',T,T'}(\prod_{1\leq i\leq n} \overline{\mathcal{P}\operatorname{BEnv}}_{\Lambda_i}^{\mathcal{H}_i}(\mathcal{M}_i)_{S_i,T_i}, \mathcal{N}).$$

Since the embedding  $\overline{\mathcal{PBEnv}}_{\Lambda_i}^{\mathcal{H}_i}(\mathcal{M}_i)_{S_i,T_i}^{\mathfrak{B}} \subset \widetilde{\mathcal{PBEnv}}(\mathcal{M}_i)_{S_i,T_i}$  preserves  $\mathcal{H}_i$ -weighted colimits, the embedding  $\prod_{1 \leq i \leq n} \overline{\mathcal{PBEnv}}_{\Lambda_i}^{\mathcal{H}_i}(\mathcal{M}_i)_{S_i,T_i}^{\mathfrak{B}} \subset \prod_{1 \leq i \leq n} \widetilde{\mathcal{PBEnv}}(\mathcal{M}_i)_{S_i,T_i}$  preserves diagrams of  $\Lambda^{\mathcal{H}_1} \boxtimes ... \boxtimes \Lambda^{\mathcal{H}_n}$ . Thus  $\phi$  lands in enriched functors sending diagrams of  $\Lambda^{\mathcal{H}_1} \boxtimes ... \boxtimes \Lambda^{\mathcal{H}_n}$  to weighted colimit diagrams. So we have to prove that an enriched functor

$$\alpha: \overline{\mathcal{P}BEnv}_{\Lambda_1}^{\mathcal{H}_1}(\mathcal{M}_1)_{S_1,T_1}^{\circledast} \times_{(Ass \times Ass)} \dots \times_{(Ass \times Ass)} \overline{\mathcal{P}BEnv}_{\Lambda_n}^{\mathcal{H}_n}(\mathcal{M}_n)_{S_n,T_n}^{\circledast} \to \mathcal{N}^{\circledast}$$

sending diagrams of  $\Lambda^{\mathcal{H}_1} \boxtimes ... \boxtimes \Lambda^{\mathcal{H}_n}$  to weighted colimit diagrams inverts all morphisms inverted by L if its restriction to  $\mathcal{M}_1^{\circledast} \times_{(Ass \times Ass)} ... \times_{(Ass \times Ass)} \mathcal{M}_n^{\circledast}$  sends diagrams of  $\Lambda_1 \boxtimes ... \boxtimes \Lambda_n$  to weighted colimit diagrams. Let  $\mathcal{Q}_i$  be the set of morphisms  $\operatorname{colim}^H(\iota \circ F) \to \iota(Y)$  in  $\mathcal{P}BEnv(\mathcal{M}_i)_{S_i,T_i}$  adjoint to a morphism

$$H \xrightarrow{\lambda} F^*(Y) \to (\iota \circ F)^*(\iota(Y))$$

in  $\mathcal{PBEnv}(\mathcal{J})$  for some diagram  $(F : \mathcal{J}^{\circledast} \to \mathcal{M}_{i}^{\circledast}, Y, \lambda : H \to F^{*}(Y)) \in \Lambda_{i}$ , where  $\iota : \mathcal{M}_{i}^{\circledast} \to \mathcal{PBEnv}(\mathcal{M}_{i})_{S_{i},T_{i}}^{\circledast}$ is the canonical embedding. By construction of the localization the set  $\Omega_{i}$  is the set of generating local equivalences of the localization  $\mathcal{PBEnv}_{\Lambda_{i}}(\mathcal{M}_{i})_{S_{i},T_{i}} \subset \mathcal{PBEnv}(\mathcal{M}_{i})_{S_{i},T_{i}}$ , which are morphisms of the full subcategory  $\overline{\mathcal{PBEnv}}_{\Lambda_{i}}^{\mathcal{H}_{i}}(\mathcal{M}_{i})_{S_{i},T_{i}} \subset \mathcal{PBEnv}(\mathcal{M}_{i})_{S_{i},T_{i}}$  closed under  $\mathcal{H}_{i}$ -weighted colimits. Therefore the enriched functor  $\alpha$  inverts morphisms inverted by L if and only if it inverts any  $(f_{1},...,f_{n}) \in \prod_{1 \leq i \leq n} \Omega_{i}$ . For  $1 \leq i \leq n$  let  $\overline{\mathcal{M}}_{i}^{\circledast} \subset \mathcal{PBEnv}_{\Lambda_{i}}(\mathcal{M}_{i})_{S_{i},T_{i}}^{\circledast}$  be the full weakly bi-enriched subcategory generated by  $\mathcal{M}_{i}$ under  $\mathcal{H}_{i}$ -weighted colimits. Since morphisms of  $\mathcal{PBEnv}(\mathcal{M}_{i})_{S_{i},T_{i}}$  that belong to  $\Omega_{i}$  are morphisms of  $\overline{\mathcal{M}}_{i}$ , the enriched functor  $\alpha$  inverts every  $(f_{1},...,f_{n}) \in \prod_{1 \leq i \leq n} \Omega_{i}$  if for every family  $(X_{j} \in \overline{\mathcal{M}}_{j})_{1 \leq j \leq n, j \neq i}$  the induced enriched functor  $\alpha(X_{1},...,X_{i-1},-,X_{i+1},...,X_{n}) : \overline{\mathcal{PBEnv}}_{\Lambda_{i}}^{\mathcal{H}}(\mathcal{M}_{i})_{S_{i},T_{i}}^{\circledast} \to \mathcal{O}^{\circledast}$  inverts morphisms of  $\Omega_{i}$ . To see the latter, by Remark 3.57 and the fact that  $\alpha$  sends diagrams of  $\Lambda^{\mathcal{H}_{1}} \boxtimes ... \boxtimes \Lambda^{\mathcal{H}_{n}}$  to weighted colimit diagrams, we can assume that  $X_{j} \in \mathcal{M}_{j}$  for  $1 \leq j \leq n, j \neq i$ . In this case the induced enriched functor  $\alpha(X_{1},...,X_{i-1},-,X_{i+1},...,X_{n}) : \overline{\mathcal{PBEnv}}_{\Lambda_{i}}^{\mathcal{H}}(\mathcal{M}_{i})_{S_{i},T_{i}}^{\circledast} \to \mathcal{O}^{\circledast}$  inverts morphisms of  $\Omega_{i}$  because it sends diagrams of  $\Lambda_{i}$  and  $\mathcal{H}_{i}$ -weighted colimit diagrams to weighted colimit diagrams.  $\Box$ 

Specializing Proposition 3.85 gives the following corollaries:

**Corollary 3.86.** Let  $n \ge 1$ . For  $1 \le i \le n$  let  $(\mathcal{V}_i^{\otimes} \to Ass, S_i), (\mathcal{V}'^{\otimes} \to Ass, S')$  be small localization pairs,  $\mathcal{H}_i$  a set of small left weights over  $\mathcal{V}_i$  and  $\mathcal{H}'$  a collection of left weights over  $\mathcal{V}'$ . Let  $\mathcal{M}_i^{\otimes} \to \mathcal{V}_i^{\otimes}$  be small left  $S_i$ -enriched  $\infty$ -categories,  $\Lambda_i$  a collection of  $\mathcal{H}_i$ -weighted diagrams in  $\mathcal{M}_i$  and  $\mathcal{N}^{\otimes} \to \mathcal{V}'^{\otimes} \times \mathcal{W}'^{\otimes}$  a left S'-enriched  $\infty$ -category that admits  $\mathcal{H}'$ -weighted colimits. Let  $\mathcal{H} := \mathcal{H}_1 \boxtimes ... \boxtimes \mathcal{H}_n, S := S_1 \boxtimes ... \boxtimes S_n, T := T_1 \boxtimes ... \boxtimes T_n$ . Assume that for every  $1 \le i \le n$  the weakly left enriched  $\infty$ -category  $\mathcal{M}_i^{\otimes} \to \mathcal{V}_i^{\otimes}$  is locally left pseudo-enriched if there is some  $1 \le j \le n$  with  $j \ne i$  such that  $S_j$  does not contain only equivalences or  $\Lambda_i$  is not empty. The induced functor

$$\gamma_{\mathcal{N}} : EnrFun_{S,S',T,T',\mathcal{H},\mathcal{H}'}^{\Lambda^{\mathcal{H}_{1}} \boxtimes \dots \boxtimes \Lambda^{\mathcal{H}_{n}}} (\prod_{1 \leq i \leq n} \mathcal{P}BEnv_{\Lambda_{i}}^{\mathcal{H}_{i}}(\mathcal{M}_{i})_{S_{i},T_{i}}, \mathcal{N}) \rightarrow EnrFun_{S,S',T,T',\mathcal{H},\mathcal{H}'}^{\Lambda_{1} \boxtimes \dots \boxtimes \Lambda_{n}} (\prod_{1 \leq i \leq n} \mathcal{M}_{i}, \mathcal{N})$$

is an equivalence. The induced functor

$$\delta_{\mathcal{N}}: EnrFun_{S,S',T,T',\mathcal{H},\mathcal{H}'}(\prod_{1 \leq i \leq n} \mathcal{P}BEnv^{\mathcal{H}_{i}}(\mathcal{M}_{i})_{S_{i},T_{i}}, \mathcal{N}) \rightarrow EnrFun_{S,S',T,T',\mathcal{H},\mathcal{H}'}(\prod_{1 \leq i \leq n} \mathcal{M}_{i}, \mathcal{N})$$

admits a fully faithful left adjoint that lands in  $\operatorname{EnrFun}_{S,S',T,T',\mathcal{H},\mathcal{H}'}^{\mathcal{H}_{1}}(\prod_{1\leq i\leq n} \mathcal{P}\operatorname{BEnv}_{\Lambda_{i}}^{\mathcal{H}_{i}}(\mathcal{M}_{i})_{S_{i},T_{i}}, \mathcal{N}).$ 

**Corollary 3.87.** Let  $(\mathcal{V}^{\otimes} \to \operatorname{Ass}, S), (\mathcal{W}^{\otimes} \to \operatorname{Ass}, T)$  be small localization pairs,  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a totally small S, T-bi-enriched  $\infty$ -category,  $\mathcal{H}$  a set of totally small weights over  $\mathcal{V}, \mathcal{W}, \Lambda$  a collection of  $\mathcal{H}$ -weighted diagrams in  $\mathcal{M}$  and  $\mathcal{N}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a S, T-bi-enriched  $\infty$ -category that admits  $\mathcal{H}$ -weighted colimits. The following functor is an equivalence:

$$\operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}^{\Lambda^{\mathcal{H}}}(\mathcal{P}\operatorname{BEnv}_{\Lambda}^{\mathcal{H}}(\mathcal{M})_{S,T},\mathcal{N}) \to \operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}^{\Lambda}(\mathcal{M},\mathcal{N}).$$

**Corollary 3.88.** Let  $(\mathcal{V}^{\otimes} \to \operatorname{Ass}, \operatorname{S})$  be a small localization pair,  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes}$  a totally small left Senriched  $\infty$ -category,  $\mathcal{H}$  a set of small left weights over  $\mathcal{V}$ ,  $\Lambda$  a collection of  $\mathcal{H}$ -weighted left diagrams in  $\mathcal{M}$  and  $\mathcal{N}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a left S-enriched  $\infty$ -category that admits  $\mathcal{H}$ -weighted colimits. The following right  $\mathcal{W}$ -enriched functor is an equivalence:

 $\operatorname{EnrFun}_{\mathcal{V}, \varnothing}^{\Lambda^{\mathcal{H}}}(\mathcal{P}\operatorname{BEnv}_{\Lambda}^{\mathcal{H}}(\mathcal{M})_{\mathrm{S}, \mathrm{T}}, \mathcal{N})^{\circledast} \to \operatorname{EnrFun}_{\mathcal{V}, \varnothing}^{\Lambda}(\mathcal{M}, \mathcal{N})^{\circledast}.$ 

**Notation 3.89.** Let  $\kappa, \tau$  be regular cardinals,  $n \ge 1$  and for every  $1 \le i \le n$  let  $\mathcal{M}_i^{\otimes} \to \mathcal{V}_i^{\otimes} \times \mathcal{W}_i^{\otimes}, \mathcal{N}^{\otimes} \to \mathcal{V}_i^{\otimes} \times \mathcal{W}_i^{\otimes}$  be  $\infty$ -categories weakly bi-enriched in monoidal  $\infty$ -categories compatible with  $\kappa$ -small colimits,  $\tau$ -small colimits, respectively. Let  $\mathcal{H}$  be a collection of weights over  $\prod_{i=1}^n \mathcal{V}_i, \prod_{i=1}^n \mathcal{W}_i$  and  $\mathcal{H}'$  a collection of weights over  $\mathcal{V}', \mathcal{W}'$  and  $\Lambda$  a collection of  $\mathcal{H}$ -weighted diagrams in  $\prod_{i=1}^n \mathcal{M}_i$ . Set  $\mathcal{V}^{\otimes} := \mathcal{V}_1^{\otimes} \times_{\mathrm{Ass}} \dots \times_{\mathrm{Ass}} \mathcal{V}_n^{\otimes}, \mathcal{W}^{\otimes} := \mathcal{W}_1^{\otimes} \times_{\mathrm{Ass}} \dots \times_{\mathrm{Ass}} \mathcal{W}_n^{\otimes}$ . Let

$$\sum_{\substack{\otimes,\kappa \in \mathrm{EnrFun}^{\Lambda}_{\mathcal{H},\mathcal{H}'}(\prod_{1 \leq i \leq n} \mathcal{M}_{i}, \mathcal{N}), \ \mathrm{EnrFun}^{\Lambda}_{\otimes,\tau,\mathcal{H},\mathcal{H}'}(\prod_{1 \leq i \leq n} \mathcal{M}_{i}, \mathcal{N}), \ \underset{\otimes,\kappa \in \mathrm{EnrFun}^{\Lambda}_{\otimes,\tau,\mathcal{H},\mathcal{H}'}(\prod_{1 \leq i \leq n} \mathcal{M}_{i}, \mathcal{N})}{\subset \mathrm{EnrFun}^{\Lambda}_{\mathcal{H},\mathcal{H}'}(\prod_{1 \leq i \leq n} \mathcal{M}_{i}, \mathcal{N})}$$

be the full subcategories of enriched functors lying from the left over a map of  $\infty$ -operads  $\mathcal{V}^{\otimes} \to \mathcal{V}'^{\otimes}$ such that for every  $1 \leq i \leq n$  the composition  $\mathcal{V}_i^{\otimes} \to \mathcal{V}^{\otimes} \to \mathcal{V}'^{\otimes}$  is a monoidal functor preserving  $\kappa$ -small colimits, lying from the right over a map of  $\infty$ -operads  $\mathcal{W}^{\otimes} \to \mathcal{W}'^{\otimes}$  such that for every  $1 \leq i \leq n$  the composition  $\mathcal{W}_i^{\otimes} \to \mathcal{W}'^{\otimes}$  is a monoidal functor preserving  $\tau$ -small colimits, lying from the left and right over a map of  $\infty$ -operads with these properties, respectively. If  $\kappa = \emptyset$ , we drop  $\kappa$ . If  $\kappa$  is the large strongly inaccessible cardinal corresponding to the small universe, we replace  $\kappa$  by cc.

**Corollary 3.90.** Let  $\kappa, \tau$  be small regular cardinals,  $n \ge 1$  and for every  $1 \le i \le n$  let  $\mathcal{M}_i^{\circledast} \to \mathcal{V}_i^{\circledast} \times \mathcal{W}_i^{\circledast}$  be a totally small weakly bi-enriched  $\infty$ -category,  $\mathcal{H}_i$  a set of totally small weights over  $\mathcal{V}_i, \mathcal{W}_i, \Lambda_i$  a collection of  $\mathcal{H}_i$ -weighted diagrams in  $\mathcal{M}_i$  and  $\mathcal{N}^{\circledast} \to \mathcal{V}'^{\circledast} \times \mathcal{W}'^{\circledast}$  a weakly bi-enriched  $\infty$ -category that admits  $\mathcal{H}'$ -weighted colimits for a collection  $\mathcal{H}'$  of weights over  $\mathcal{V}', \mathcal{W}'$ .

(1) If for every  $1 \leq i \leq n$  the weakly bi-enriched  $\infty$ -categories  $\mathcal{M}_i^{\circledast} \to \mathcal{V}_i^{\circledast} \times \mathcal{W}_i^{\circledast}$  and  $\mathcal{N}^{\circledast} \to \mathcal{V}'^{\otimes} \times \mathcal{W}'^{\otimes}$  are left  $\kappa$ -enriched  $\infty$ -categories and  $\mathcal{H}_i, \mathcal{H}'$  consist of left weights, the following functor is an equivalence:

$${}_{\otimes,\kappa} Enr Fun_{\mathcal{H},\mathcal{H}'}^{\Lambda^{\mathcal{H}_1}\boxtimes\ldots\boxtimes\Lambda^{\mathcal{H}_n}}(\prod_{1\leq i\leq n}\mathcal{P}BEnv_{\Lambda_i}^{\mathcal{H}_i}(\mathcal{M}_i)_{LEnr_\kappa},\mathcal{N}) \rightarrow {}_{\otimes,\kappa} Enr Fun_{\mathcal{H},\mathcal{H}'}^{\Lambda_1\boxtimes\ldots\boxtimes\Lambda_n}(\prod_{1\leq i\leq n}\mathcal{M}_i,\mathcal{N}).$$

(2) If for every  $1 \leq i \leq n$  the weakly bi-enriched  $\infty$ -categories  $\mathcal{M}_i^{\circledast} \to \mathcal{V}_i^{\otimes} \times \mathcal{W}_i^{\otimes}$  and  $\mathcal{N}^{\circledast} \to \mathcal{V}'^{\otimes} \times \mathcal{W}'^{\otimes}$  are right  $\tau$ -enriched  $\infty$ -categories and  $\mathcal{H}_i, \mathcal{H}'$  consist of right weights, the following functor is an equivalence:

$$\mathrm{EnrFun}_{\otimes,\tau,\mathcal{H},\mathcal{H}'}^{\Lambda^{\mathscr{H}_1}\boxtimes\ldots\boxtimes\Lambda^{\mathscr{H}_n}}(\prod_{1\leq i\leq n}\mathcal{P}\mathrm{BEnv}_{\Lambda_i}^{\mathscr{H}_i}(\mathcal{M}_i)_{\mathrm{REnr}_\tau},\mathcal{N})\rightarrow\mathrm{EnrFun}_{\otimes,\tau,\mathcal{H},\mathcal{H}'}^{\Lambda_1\boxtimes\ldots\boxtimes\Lambda_n}(\prod_{1\leq i\leq n}\mathcal{M}_i,\mathcal{N}).$$

(3) If for every  $1 \le i \le n$  the weakly bi-enriched  $\infty$ -categories  $\mathcal{M}_i^{\otimes} \to \mathcal{V}_i^{\otimes} \times \mathcal{W}_i^{\otimes}$  and  $\mathcal{N}^{\otimes} \to \mathcal{V}'^{\otimes} \times \mathcal{W}'^{\otimes}$ are  $\kappa, \tau$ -bi-enriched  $\infty$ -categories, the following functor is an equivalence:

$$\underset{\otimes,\kappa}{\otimes,\kappa} EnrFun_{\otimes,\tau,\mathcal{H},\mathcal{H}'}^{\Lambda^{\mathcal{H}_{1}} \boxtimes \ldots \boxtimes \Lambda^{\mathcal{H}_{n}}} \big(\prod_{1 \leq i \leq n} \mathcal{P}BEnv_{\Lambda_{i}}^{\mathcal{H}_{i}}(\mathcal{M}_{i})_{BEnr_{\kappa,\tau}}, \mathcal{N}\big) \rightarrow \underset{\otimes,\kappa}{\otimes,\kappa} EnrFun_{\otimes,\tau,\mathcal{H},\mathcal{H}'}^{\Lambda_{1} \boxtimes \ldots \boxtimes \Lambda_{n}} \big(\prod_{1 \leq i \leq n} \mathcal{M}_{i}, \mathcal{N}\big).$$

For the next lemma we use the following terminology:

**Definition 3.91.** Let  $\mathcal{V}^{\otimes} \to Ass, \mathcal{W}^{\otimes} \to Ass$  be small  $\infty$ -operads and  $\kappa, \tau$  small regular cardinals.

- (1) A totally small left  $\kappa$ -enriched weight over  $\mathcal{V}, \mathcal{W}$  is a weight  $(\alpha : \mathcal{V}^{\otimes} \to \mathcal{V}^{\otimes}, \beta : \mathcal{W}^{\otimes} \to \mathcal{W}^{\otimes}, \mathcal{J}^{\otimes} \to \mathcal{W}^{\otimes}, \mathcal{J}^{\otimes$  $\mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}, \mathrm{H}$ ) over  $\mathcal{V}, \mathcal{W}$  such that  $\mathcal{V}^{\otimes} \to \mathrm{Ass}, \mathcal{V}^{\otimes} \to \mathrm{Ass}$  are small monoidal  $\infty$ -categories compatible with  $\kappa$ -small colimits,  $\mathcal{W}^{\otimes} \to \mathrm{Ass}, \mathcal{W}^{\otimes} \to \mathrm{Ass}$  are small  $\infty$ -operads,  $\alpha$  is a monoidal functor preserving  $\kappa$ -small colimits,  $\mathcal{J}^{\otimes} \to \mathcal{V}'^{\otimes} \times \mathcal{W}'^{\otimes}$  is a small left  $\kappa$ -enriched  $\infty$ -category and  $H \in \mathcal{P}BEnv(\mathcal{M})_{LEnr_{\tau}}$ . Dually, we define totally small right  $\tau$ -enriched weights over  $\mathcal{V}, \mathcal{W}$ .
- (2) A totally small  $\kappa, \tau$ -enriched weight over  $\mathcal{V}, \mathcal{W}$  is a weight  $(\alpha : \mathcal{V}^{\otimes} \to \mathcal{V}^{\otimes}, \beta : \mathcal{W}^{\otimes} \to \mathcal{W}^{\otimes}, \mathcal{J}^{\otimes} \to \mathcal{W}^{\otimes}, \mathcal{J}^{\otimes}$  $\mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}, \mathrm{H}$ ) over  $\mathcal{V}, \mathcal{W}$  such that  $\mathcal{V}^{\otimes} \to \mathrm{Ass}, \mathcal{V}^{\otimes} \to \mathrm{Ass}$  are small monoidal  $\infty$ -categories compatible with  $\kappa$ -small colimits,  $\mathcal{W}^{\otimes} \to Ass, \mathcal{W}^{\otimes} \to Ass$  are monoidal  $\infty$ -categories compatible with  $\tau$ -small colimits,  $\alpha$  is a monoidal functor preserving  $\kappa$ -small colimits,  $\beta$  is a monoidal functor preserving  $\tau$ -small colimits,  $\mathcal{J}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  is a small  $\kappa, \tau$ -bi-enriched  $\infty$ -category and  $H \in \mathcal{P}BEnv(\mathcal{M})_{BEnr_{\kappa,\tau}}.$
- **a 3.92.** (1) Let  $\alpha : \mathcal{V}^{\otimes} \to \mathcal{V}^{\otimes}$  be a left adjoint monoidal functor between presentably monoidal  $\infty$ -categories and  $\beta : \mathcal{W}^{\otimes} \to \mathcal{W}^{\otimes}$  a map of  $\infty$ -operads. By [8, Proposition 7.15.] Lemma 3.92. there are small regular cardinals  $\kappa, \kappa'$  such that  $\mathcal{V}^{\otimes} \to Ass$  is  $\kappa$ -compactly generated,  $\mathcal{V}'^{\otimes} \to Ass$ is  $\kappa'$ -compactly generated and  $\alpha$  sends  $\kappa'$ -compact objects to  $\kappa$ -compact objects. For any left  $enriched \ weight \ H \ on \ a \ small \ left \ enriched \ \infty-category \ \mathcal{J}^{\circledast} \rightarrow \mathcal{V}'^{\otimes} \times \mathcal{W}'^{\otimes} \ there \ is \ a \ canonical \ left \ deft \ deft$  $\kappa'$ -enriched weight  $H' \in \mathcal{PBEnv}(\mathcal{J}_{\kappa'})_{\mathrm{LEnr}_{\kappa'}}$  on  $\mathcal{J}_{\kappa'}^{\circledast} \to (\mathcal{V}'^{\kappa'})^{\otimes} \times \mathcal{W}'^{\otimes}$  such that for any enriched functor  $F: \mathcal{J}^{\circledast} \to \mathcal{M}^{\circledast}$  lying over  $\alpha, \beta$  there is a canonical equivalence

$$\operatorname{colim}^{\mathrm{H}}(\mathrm{F}) \simeq \operatorname{colim}^{\mathrm{H}'}(\mathrm{F}'),$$

where  $F' : \mathcal{J}_{\kappa'}^{\circledast} \to \mathcal{M}_{\kappa}^{\circledast}$  is the restriction of F. (2) Let  $\alpha : \mathcal{V}^{\otimes} \to \mathcal{V}^{\otimes}, \beta : \mathcal{W}^{\otimes} \to \mathcal{W}^{\otimes}$  be left adjoint monoidal functors between presentably monoidal  $\infty$ -categories. By [8, Proposition 7.15.] there are small regular cardinals  $\kappa, \kappa', \tau, \tau'$  such that  $\mathcal{V}^{\otimes} \to Ass \text{ is } \kappa\text{-compactly generated, } \mathcal{V}'^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}^{\otimes} \to Ass \text{ is } \kappa'\text{-compactly generated, } \mathcal{W}$  $\tau$ -compactly generated,  $\mathcal{W}^{\otimes} \to Ass$  is  $\tau'$ -compactly generated,  $\alpha$  sends  $\kappa'$ -compact objects to  $\kappa$ compact objects and  $\beta$  sends  $\tau'$ -compact objects to  $\tau'$ -compact objects. For any enriched weight H on  $\mathcal{J}^{\otimes} \to \mathcal{V}'^{\otimes} \times \mathcal{W}'^{\otimes}$  there is a canonical  $\kappa', \tau'$ -enriched weight  $\mathbf{H}' \in \mathcal{P}BEnv(\mathcal{J}_{\kappa',\tau'})_{BEnr_{\kappa',\tau'}}$ on  $\mathcal{J}^{\circledast}_{\kappa',\tau'} \to (\mathcal{V}'^{\kappa'})^{\circledast} \times (\mathcal{W}'^{\tau'})^{\circledast}$  such that for any enriched functor  $F: \mathcal{J}^{\circledast} \to \mathcal{M}^{\circledast}$  lying over  $\alpha, \beta$ there is a canonical equivalence

$$\operatorname{colim}^{\mathrm{H}}(\mathrm{F}) \simeq \operatorname{colim}^{\mathrm{H}'}(\mathrm{F}'),$$

where  $F' : \mathcal{J}^{\circledast}_{\kappa' \tau'} \to \mathcal{M}^{\circledast}_{\kappa,\tau}$  is the restriction of F.

*Proof.* We prove (1). The proof of (2) is similar. By Corollary 2.142 the projections  $j: \mathcal{J}_{\kappa'}^{\otimes} \to \mathcal{J}^{\otimes}, \iota:$  $\mathcal{M}^{\otimes}_{\kappa} \to \mathcal{M}^{\otimes}$  induce  $\mathcal{V}, \mathcal{W}$ -enriched equivalences

$$L \circ j_{!} : \mathcal{P}BEnv(\mathcal{J}_{\kappa'})^{\mathfrak{B}}_{LEnr, \prime} \simeq \mathcal{P}BEnv(\mathcal{J})^{\mathfrak{B}}_{LEnr}, L \circ \iota_{!} : \mathcal{P}BEnv(\mathcal{M}_{\kappa})^{\mathfrak{B}}_{LEnr, \prime} \simeq \mathcal{P}BEnv(\mathcal{M})^{\mathfrak{B}}_{LEnr}$$

Setting H' :=  $(L \circ \tau_1)^{-1}(H)$  there is an equivalence  $L(F_1(H)) \simeq L \circ F_1 \circ L \circ \tau_1(H') \simeq L \circ \iota_1 \circ L \circ F'_1(H')$ . So  $\psi: L(F_!(H)) \to Y$  is the image under  $L \circ \iota_!$  of a morphism  $\psi': L \circ F'_!(H') \to Y$  in  $\mathcal{PBEnv}(\mathcal{M}_{\kappa})_{LEnr_{\kappa}}$ . The morphism  $\psi$  exhibits Y as the H-weighted colimit of F if and only if  $\psi'$  exhibits Y as the H'-weighted colimit of F' since for every  $Z \in \mathcal{M}, \mathcal{V} \in \mathcal{V}^{\kappa}, W_1, ..., W_m \in \mathcal{W}$  for  $m \ge 0$  there is a commutative square:

To apply Corollary 3.90 we use the following proposition:

- **Proposition 3.93.** (1) Let  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  be a small  $\infty$ -category left enriched in a presentably monoidal  $\infty$ -category,  $\mathcal{H}$  a set of small left enriched weights over  $\mathcal{V}, \mathcal{W}$  and  $\Lambda$  a collection of  $\mathcal{H}$ -weighted diagrams in  $\mathcal{M}$ . Then  $\widehat{\mathcal{P}}Benv_{\Lambda}^{\mathcal{H}}(\mathcal{M})^{\circledast}_{Enr} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  is locally small.
  - (2) Let M<sup>®</sup> → V<sup>®</sup>×W<sup>®</sup> be a small ∞-category right enriched in a presentably monoidal ∞-category, H a set of small right enriched weights over V, W and Λ a collection of H-weighted diagrams in M. Then PBEnv<sup>H</sup><sub>Λ</sub>(M)<sup>®</sup><sub>REnr</sub> → V<sup>®</sup> × W<sup>®</sup> is locally small.
    (3) Let M<sup>®</sup> → V<sup>®</sup> × W<sup>®</sup> be a small ∞-category bi-enriched in presentably monoidal ∞-categories,
  - (3) Let M<sup>®</sup> → V<sup>®</sup> × W<sup>®</sup> be a small ∞-category bi-enriched in presentably monoidal ∞-categories, H a set of small enriched weights over V, W and Λ a collection of H-weighted diagrams in M. Then PBEnv<sup>H</sup><sub>Λ</sub>(M)<sup>®</sup><sub>BEnr</sub> → V<sup>®</sup> × W<sup>®</sup> is locally small.

Proof. We prove (3). The proofs of (1) and (2) are similar. By Notation 2.152 there is an enriched embedding  $j: \mathcal{PBEnv}(\mathcal{M})^{\circledast}_{BEnr} \subset \widehat{\mathcal{P}BEnv}(\mathcal{M})^{\circledast}_{BEnr}$  that preserves small weighted colimits. Let  $\iota: \mathcal{M}^{\circledast} \subset \mathcal{PBEnv}(\mathcal{M})^{\circledast}_{BEnr}$  be the canonical enriched embedding. By Proposition 2.153 the weakly bi-enriched  $\infty$ -category  $\mathcal{PBEnv}(\mathcal{M})^{\circledast}_{BEnr} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  is a presentably bitensored  $\infty$ -category. Let  $\Sigma$  be the collection of the following totally small diagrams: let  $(F: \mathcal{J}^{\circledast} \to \mathcal{M}^{\circledast}, H \in \mathcal{PBEnv}(\mathcal{J})_{BEnr}, Y \in \mathcal{M}) \in \Lambda$ . By [8, Proposition 7.15.] there are small regular cardinals  $\kappa, \kappa', \tau, \tau'$  such that  $\mathcal{V}^{\circledast} \to Ass$  is  $\kappa$ -compactly generated,  $\mathcal{V}^{' \circledast} \to Ass$  is  $\kappa'$ -compactly generated,  $\mathcal{W}^{\otimes} \to Ass$  is  $\tau$ -compactly generated,  $\mathcal{W}^{' \circledast} \to Ass$  is  $\tau'$ -compactly generated,  $\alpha$  sends  $\kappa'$ -compact objects to  $\kappa$ -compact objects and  $\beta$  sends  $\tau'$ -compact objects to  $\tau'$ -compact objects. By Lemma (2) 3.92 there is a canonical  $\kappa', \tau'$ -enriched weight H'  $\in$  $\mathcal{PBEnv}(\mathcal{J}_{\kappa',\tau'})_{BEnr_{\kappa',\tau'}} \circ \mathcal{V}^{'\kappa'})^{\otimes} \times (\mathcal{W}^{\prime\tau'})^{\otimes}$  such that for any enriched functor  $F: \mathcal{J}^{\circledast} \to \mathcal{M}^{\circledast}$ lying over  $\alpha, \beta$  there is a canonical equivalence

(12) 
$$\operatorname{colim}^{\mathrm{H}}(\iota \circ \mathrm{F}) \simeq \operatorname{colim}^{\mathrm{H}'}(\iota \circ \mathrm{F}'),$$

where  $F' : \mathcal{J}^{\circledast}_{\kappa',\tau'} \to \mathcal{M}^{\circledast}_{\kappa,\tau}$  is the restriction of F. Then  $(F' : \mathcal{J}^{\circledast}_{\kappa',\tau'} \to \mathcal{M}^{\circledast}_{\kappa,\tau}, H' \in \mathcal{P}BEnv(\mathcal{J}_{\kappa',\tau'})_{BEnr_{\kappa',\tau'}}, Y \in \mathcal{M})$ . Then  $\Sigma$  is a collection of totally small weights over  $\mathcal{V}, \mathcal{W}$ , which is a set by Lemma 3.69 and by equivalence 12 there is a canonical equivalence  $\widehat{\mathcal{P}}BEnv_{\Lambda}(\mathcal{M})^{\circledast}_{BEnr} \simeq \widehat{\mathcal{P}}BEnv_{\Sigma}(\mathcal{M})^{\circledast}_{BEnr}$  by definition of the latter.

Let  $\mathfrak{Q}'_{\Sigma}$  be the set of Notation 3.70 and  $\mathfrak{PBEnv}_{\Sigma}(\mathfrak{M})^{\circledast}_{\mathrm{BEnr}} \subset \mathfrak{PBEnv}(\mathfrak{M})^{\circledast}_{\mathrm{BEnr}}$  the full weakly bienriched subcategory spanned by the  $\mathfrak{Q}'_{\Sigma}$ -local objects. Since  $\mathfrak{Q}'_{\Sigma}$  is preserved by the  $\mathcal{V}, \mathcal{W}$ -biaction, the full weakly bi-enriched subcategory  $\mathfrak{PBEnv}_{\Sigma}(\mathfrak{M})^{\circledast}_{\mathrm{BEnr}} \subset \mathfrak{PBEnv}(\mathfrak{M})^{\circledast}_{\mathrm{BEnr}}$  is an accessible  $\mathcal{V}, \mathcal{W}$ -enriched localization and so a presentably bitensored  $\infty$ -category. Let  $\mathfrak{PBEnv}_{\Sigma}^{\mathcal{H}}(\mathfrak{M})^{\circledast}_{\mathrm{BEnr}} \subset \mathfrak{PBEnv}_{\Sigma}(\mathfrak{M})^{\circledast}_{\mathrm{BEnr}}$ be the full weakly bi-enriched subcategory generated by  $\mathfrak{M}$  under  $\mathfrak{H}$ -weighted colimits. By definition  $\widehat{\mathcal{P}BEnv}_{\Sigma}(\mathfrak{M})^{\circledast}_{\mathrm{BEnr}}$  is the localization with respect to the set  $j(\mathfrak{Q}'_{\Sigma})$ , the image of  $\mathfrak{Q}'_{\Sigma}$  under j. Hence j preserves and reflects local objects (being an embedding) for the localizations with respect to the sets  $\mathfrak{Q}'_{\Sigma}, j(\mathfrak{Q}'_{\Sigma})$ . So j restricts to an embedding  $\mathfrak{PBEnv}_{\Sigma}(\mathfrak{M})^{\circledast}_{\mathrm{BEnr}} \subset \widehat{\mathcal{P}BEnv}_{\Sigma}(\mathfrak{M})^{\circledast}_{\mathrm{BEnr}}$  preserving small weighted colimits. Consequently, the latter embedding restricts to an equivalence  $\mathfrak{PBEnv}_{\Sigma}^{\mathcal{H}}(\mathfrak{M})^{\circledast}_{\mathrm{BEnr}} \simeq$  $\widehat{\mathcal{P}BEnv}^{\mathcal{H}}_{\Sigma}(\mathfrak{M})^{\circledast}_{\mathrm{BEnr}}.$ 

3.11. Weighted colimits of diagrams of enriched functors. In the following we prove that weighted colimits in enriched  $\infty$ -categories of enriched functors are formed object-wise.

**Theorem 3.94.** Let  $\mathcal{V}^{\otimes} \to \operatorname{Ass}, \mathcal{W}^{\otimes} \to \operatorname{Ass}$  be  $\infty$ -operads,  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes}$  a weakly left enriched  $\infty$ -category,  $\mathcal{N}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a weakly bi-enriched  $\infty$ -category and  $\mathcal{H}$  a set of small right weights over  $\mathcal{W}$ . If  $\mathcal{N}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  admits  $\sigma_{1}(\mathcal{H})$ -weighted colimits, where  $\sigma : \mathcal{Q}^{\otimes} \to \mathcal{V}^{\otimes}$  is the unique map of  $\infty$ -operads, the weakly right enriched  $\infty$ -category  $\operatorname{EnrFun}_{\mathcal{V},\mathcal{Q}}(\mathcal{M}, \mathcal{N})^{\circledast} \to \mathcal{W}^{\otimes}$  admits  $\mathcal{H}$ -weighted colimits and for every  $X \in \mathcal{M}$  the right  $\mathcal{W}$ -enriched functor  $\operatorname{EnrFun}_{\mathcal{V},\mathcal{Q}}(\mathcal{M}, \mathcal{N})^{\circledast} \to \mathcal{N}^{\circledast}$  evaluating at X preserves  $\mathcal{H}$ -weighted colimits. Moreover for every set of diagrams  $\Lambda$  in  $\mathcal{M}$  the full subcategory  $\operatorname{EnrFun}_{\mathcal{V},\mathcal{Q}}(\mathcal{M}, \mathcal{N})^{\circledast} \subset \operatorname{EnrFun}_{\mathcal{V},\mathcal{Q}}(\mathcal{M}, \mathcal{N})^{\circledast}$  is closed under  $\mathcal{H}$ -weighted colimits.

*Proof.* Let  $\mathcal{N}^{\otimes} := \mathcal{P}\widetilde{BEnv}_{\Lambda^{\sigma_1(\mathcal{H})}}(\mathcal{N})^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$ . By Remark 3.78 there is a  $\mathcal{V}, \mathcal{W}$ -enriched embedding  $\mathcal{N}^{\otimes} \subset \mathcal{N}^{\otimes}$  that preserves  $\sigma_1(\mathcal{H})$ -weighted colimits. The latter induces a right  $\mathcal{W}$ -enriched embedding:

$$\operatorname{EnrFun}_{\mathcal{V},\varnothing}(\mathcal{M},\mathcal{N})^{\circledast} \subset \operatorname{EnrFun}_{\mathcal{V},\varnothing}(\mathcal{M},\mathcal{N}')^{\circledast}.$$

By Remark 3.76 the weakly bi-enriched  $\infty$ -category  $\mathcal{N}^{\otimes} = \mathcal{P}\widetilde{\mathrm{BEnv}}_{\Lambda^{\sigma_1}(\mathcal{H})}(\mathcal{N})^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  admits right tensors and small conical colimits. By [9, Proposition 5.15.] the weakly right enriched  $\infty$ category  $\operatorname{EnrFun}_{\mathcal{V},\emptyset}(\mathcal{M},\mathcal{N}')^{\circledast} \to \mathcal{W}^{\circledast}$  admits right tensors and small conical colimits and the forgetful functor  $\operatorname{EnrFun}_{\mathcal{V},\varnothing}(\mathcal{M},\mathcal{N}')^{\circledast} \to (\mathcal{N}'^{\mathcal{M}})^{\circledast}$  preserves right tensors and small conical colimits. Corollary 3.63 guarantees that  $\operatorname{EnrFun}_{\mathcal{V},\emptyset}(\mathcal{M},\mathcal{N}')^{\circledast} \to \mathcal{W}^{\circledast}$  admits small weighted colimits and the forgetful functor  $\operatorname{EnrFun}_{\mathcal{V},\varnothing}(\mathcal{M},\mathcal{N}')^{\circledast} \to (\mathcal{N}'^{\mathcal{M}})^{\circledast}$  preserves small weighted colimits. Since the embedding  $\mathcal{N}^{\circledast} \subset$  $\mathcal{N}^{\otimes}$  preserves  $\sigma_1(\mathcal{H})$ -weighted colimits, by Corollary 3.60 the full weakly right enriched subcategory  $\operatorname{EnrFun}_{\mathcal{V},\emptyset}(\mathcal{M},\mathcal{N})^{\circledast} \subset \operatorname{EnrFun}_{\mathcal{V},\emptyset}(\mathcal{M},\mathcal{N}')^{\circledast}$  is closed under  $\mathcal{H}$ -weighted colimits. Consequently, it is enough to check that  $\operatorname{EnrFun}_{\mathcal{V}, \emptyset}^{\Lambda}(\mathcal{M}, \mathcal{N}')$  is closed in  $\operatorname{EnrFun}_{\mathcal{V}, \emptyset}(\mathcal{M}, \mathcal{N}')$  under small weighted colimits, or equivalently by Corollary 3.63 under small colimits and right tensors. For every left V-enriched functor  $F : \mathcal{M}^{\circledast} \to \mathcal{N}'^{\circledast}$  and  $W \in \mathcal{W}$  the right tensor  $F \otimes W$  factors as  $\mathcal{M}^{\circledast} \xrightarrow{F} \mathcal{N}'^{\circledast} \xrightarrow{(-) \otimes W} \mathcal{N}'^{\circledast}$ . where  $\mathcal{N}^{\otimes} \xrightarrow{(-)\otimes W} \mathcal{N}^{\otimes}$  admits a left  $\mathcal{V}$ -enriched right adjoint and so preserves weighted colimits by Proposition 3.55. The colimit of any functor  $G: K \to \operatorname{EnrFun}_{\mathcal{V}, \emptyset}(\mathcal{M}, \mathcal{N}')$  factors as the left  $\mathcal{V}$ -enriched functor  $\mathcal{M}^{\circledast} \xrightarrow{G'} (\mathcal{N}'^{K})^{\circledast} \xrightarrow{\operatorname{colim}} \mathcal{N}'^{\circledast}$ , where G' corresponds to G under the equivalence Fun(K, EnrFun<sub> $\mathcal{V},\emptyset$ </sub>( $\mathcal{M},\mathcal{N}'$ ))  $\simeq$  EnrFun<sub> $\mathcal{V},\emptyset$ </sub>( $\mathcal{M},\mathcal{N}'^{K}$ ) and colim :  $(\mathcal{N}'^{K})^{\circledast} \to \mathcal{N}'^{\circledast}$  is the left  $\mathcal{V}$ -enriched left adjoint of the left  $\mathcal{V}$ -linear diagonal functor  $\mathcal{N}'^{\circledast} \to (\mathcal{N}'^{K})^{\circledast}$ . Therefore it is enough to observe that the left  $\mathcal{V}$ -enriched functor  $\mathcal{M}^{\circledast} \xrightarrow{G'} (\mathcal{N}'^{K})^{\circledast}$  sends diagrams of  $\Lambda$  to weighted colimit diagrams if the functor  $G: K \to EnrFun_{\mathcal{V}, \emptyset}(\mathcal{M}, \mathcal{N}')$  lands in  $EnrFun_{\mathcal{V}, \emptyset}^{\Lambda}(\mathcal{M}, \mathcal{N}')$ : this follows immediately from the fact that  $(\mathcal{N}^{\prime \mathrm{K}})^{\otimes} \to \mathcal{V}^{\otimes}$  is a left tensored  $\infty$ -category compatible with small colimits and so by Corollary 3.63 admits small weighted colimits, and for every  $Z \in K$  the left  $\mathcal{V}$ -linear functor  $(\mathcal{N}'^K)^{\circledast} \to \mathcal{N}'^{\circledast}$  evaluating at Z preserves small colimits and so preserves small weighted colimits by Proposition 3.63.

### 4. A monoidal structure for enriched ∞-categories with weighted colimits

4.1. Enriched  $\infty$ -categories with weighted colimits. In this section we organize weakly bienriched  $\infty$ -categories equipped with a diagram to a symmetric monoidal  $\infty$ -category.

Notation 4.1. Let  $\mathcal{V}^{\otimes} \to Ass, \mathcal{W}^{\otimes} \to Ass$  be small  $\infty$ -operads and  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a weakly bi-enriched  $\infty$ -category.

- (1) Let  $BD(\mathcal{M})$  be the poset of sets of totally small diagrams on  $\mathcal{M}$  ordered by inclusion.
- (2) Let  $\omega(\mathcal{V}, \mathcal{W})$  be the poset of sets of totally small weights over  $\mathcal{V}, \mathcal{W}$  ordered by inclusion.

Since we can transport diagrams and weights (Notation 3.47), we obtain functors to the category of large posets  $\widehat{\text{Poset}}$ :

$$BD: Ho(\omega BEnr) \to \widehat{Poset}, \ \omega: Ho(Op_{\infty}) \times Ho(Op_{\infty}) \to \widehat{Poset}.$$

Notation 4.2. Let

$$\omega \operatorname{BEnr}_* \to \omega \operatorname{BEnr}$$

be the cocartesian fibrations classifying the composition

$$\omega \operatorname{BEnr} \to \operatorname{Ho}(\omega \operatorname{BEnr}) \xrightarrow{\operatorname{BD}} \widehat{\operatorname{Poset}} \subset \widehat{\operatorname{Cat}}_{\infty}.$$

We call  $\omega BEnr_*$  the  $\infty$ -category of totally small weakly bi-enriched  $\infty$ -categories with diagrams.

Notation 4.3. Let

$$(\operatorname{Op}_{\infty} \times \operatorname{Op}_{\infty})_* \to \operatorname{Op}_{\infty} \times \operatorname{Op}_{\infty}$$

be the cocartesian fibrations classifying the composition

$$\operatorname{Op}_{\infty} \times \operatorname{Op}_{\infty} \to \operatorname{\mathsf{Ho}}(\operatorname{Op}_{\infty} \times \operatorname{Op}_{\infty}) \xrightarrow{\omega} \widehat{\operatorname{Poset}} \subset \widehat{\operatorname{\mathsf{Cat}}}_{\infty}.$$

We call  $(Op_{\infty} \times Op_{\infty})_*$  the  $\infty$ -category of pairs of small  $\infty$ -operads with weights.

**Remark 4.4.** Since  $\omega BEnr, Op_{\infty}$  are locally small and posets are locally small, the  $\infty$ -categories  $\omega BEnr_*, (Op_{\infty} \times Op_{\infty})_*$  are large but locally small: for  $(\mathcal{V}^{\otimes}, \mathcal{W}^{\otimes}, \mathcal{H}), (\mathcal{V}'^{\otimes}, \mathcal{H}'^{\otimes}, \mathcal{H}') \in (Op_{\infty} \times Op_{\infty})_*$  the induced map

$$(\mathrm{Op}_{\infty} \times \mathrm{Op}_{\infty})_{*}((\mathcal{V}^{\otimes}, \mathcal{W}^{\otimes}, \mathcal{H}), (\mathcal{V}^{\otimes}, \mathcal{W}^{\otimes}, \mathcal{H}^{\prime})) \to \mathrm{Op}_{\infty}(\mathcal{V}^{\otimes}, \mathcal{V}^{\prime\otimes}) \times \mathrm{Op}_{\infty}(\mathcal{W}^{\otimes}, \mathcal{W}^{\prime\otimes})$$

has empty or contractible fibers and so is fully faithful. It identifies the source with the full subspace of  $\operatorname{Op}_{\infty}(\mathcal{V}^{\otimes}, \mathcal{V}'^{\otimes}) \times \operatorname{Op}_{\infty}(\mathcal{W}^{\otimes}, \mathcal{W}'^{\otimes})$  spanned by the pairs of maps of  $\infty$ -operads that send weights in  $\mathcal{H}$  to weights in  $\mathcal{H}'$ . Similarly, for every  $(\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}, \Lambda), (\mathcal{M}'^{\circledast} \to \mathcal{V}'^{\otimes} \times \mathcal{W}'^{\otimes}, \Lambda') \in \omega \operatorname{BEnr}_*$  the space  $\omega \operatorname{BEnr}_*((\mathcal{M}^{\circledast}, \Lambda), (\mathcal{M}'^{\circledast}, \Lambda'))$  is the full subspace of  $\omega \operatorname{BEnr}(\mathcal{M}^{\circledast}, \mathcal{M}'^{\circledast})$  spanned by the enriched functors sending diagrams in  $\Lambda$  to diagrams in  $\Lambda'$ .

**Remark 4.5.** The functor  $\omega \text{BEnr}_* \rightarrow \omega \text{BEnr}$  admits a left and right adjoint. The left adjoint equips a totally small weakly bi-enriched  $\infty$ -category with the empty set of diagrams and the right adjoint equips it with the set of all diagrams.

**Construction 4.6.** Let  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  be a weakly bi-enriched  $\infty$ -category. Sending a diagram on  $\mathcal{M}$  to the underlying weight defines an order preserving map

$$BD(\mathcal{M}) \to \omega(\mathcal{V}, \mathcal{W})$$

that determines a transformation from the functor  $BD : Ho(\omega BEnr) \to Poset$  to the functor  $Ho(\omega BEnr) \to Ho(Op_{\infty} \times Op_{\infty}) \xrightarrow{\omega} Poset$ . The latter classifies a map of cocartesian fibrations over  $\omega BEnr$ :

(13) 
$$\omega \operatorname{BEnr}_* \to \omega \operatorname{BEnr} \times_{(\operatorname{Op}_m \times \operatorname{Op}_m)} (\operatorname{Op}_m \times \operatorname{Op}_m)_*.$$

**Lemma 4.7.** The functor  $\omega \operatorname{BEnr}_* \to (\operatorname{Op}_{\infty} \times \operatorname{Op}_{\infty})_*$  is a cocartesian and cartesian fibration and the functor  $\omega \operatorname{BEnr}_* \to \omega \operatorname{BEnr}$  sends (co)cartesian lifts of morphisms in  $(\operatorname{Op}_{\infty} \times \operatorname{Op}_{\infty})_*$  to (co)cartesian lifts in  $\operatorname{Op}_{\infty} \times \operatorname{Op}_{\infty}$ .

*Proof.* We start with proving that the functor is a cocartesian fibration. Let  $(\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}, \Lambda) \in \omega$ BEnr<sub>\*</sub> and  $(\alpha, \beta) : (\mathcal{V}^{\circledast}, \mathcal{W}^{\circledast}, \mathcal{H}) \to (\mathcal{V}'^{\circledast}, \mathcal{W}'^{\otimes}, \mathcal{H}')$  a morphism in  $(Op_{\infty} \times Op_{\infty})_{*}$ , where  $\mathcal{H}$  is the set of underlying weights of  $\Lambda$ . The cocartesian lift  $\mathcal{M}^{\circledast} \to \mathcal{M}'^{\circledast} := (\alpha, \beta)_{!}(\mathcal{M})^{\circledast}$  of  $\alpha, \beta$  sends the set  $\Lambda$  of  $\mathcal{H}$ -weighted diagrams to a set  $\Lambda'$  of  $\mathcal{H}'$ -weighted diagrams. By the choice of  $\Lambda'$  the morphism  $(\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}, \Lambda) \to (\mathcal{M}'^{\circledast} \to \mathcal{V}'^{\otimes} \times \mathcal{W}'^{\otimes}, \Lambda')$  in  $\omega$ BEnr<sub>\*</sub> lies over  $(\alpha, \beta) : (\mathcal{V}^{\otimes}, \mathcal{W}^{\otimes}, \mathcal{H}) \to (\mathcal{V}'^{\otimes}, \mathcal{W}'^{\otimes}, \mathcal{H}')$  in  $(Op_{\infty} \times Op_{\infty})_{*}$  and induces for every  $(\mathcal{N}^{\circledast} \to \mathcal{V}'^{\otimes} \times \mathcal{W}'^{\otimes}, \Lambda'') \in \omega$ BEnr<sub>\*</sub> lying over  $(\mathcal{V}'^{\otimes} \to Ass, \mathcal{W}'^{\otimes} \to Ass, \mathcal{H}'') \in (Op_{\infty} \times Op_{\infty})_{*}$  a pullback square

We continue with proving that the functor is a cartesian fibration. Let  $\Lambda'$  be the set of  $\mathcal{H}'$ -weighted diagrams on  $(\alpha, \beta)^*(\mathcal{M})^{\circledast} \to \mathcal{V}'^{\otimes} \times \mathcal{W}'^{\otimes}$  whose image under the projection  $(\alpha, \beta)^*(\mathcal{M})^{\circledast} \to \mathcal{M}^{\circledast}$  belongs to  $\Lambda$ . By the choice of  $\Lambda'$  the morphism  $((\alpha, \beta)^*(\mathcal{M})^{\circledast} \to \mathcal{V}'^{\otimes} \times \mathcal{W}'^{\otimes}, \Lambda') \to (\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}, \Lambda)$  in  $\omega$ BEnr<sub>\*</sub> lies over the morphism  $(\alpha, \beta) : (\mathcal{V}'^{\otimes}, \mathcal{W}'^{\otimes}, \mathcal{H}') \to (\mathcal{V}^{\otimes}, \mathcal{W}^{\otimes}, \mathcal{H})$  of  $(Op_{\infty} \times Op_{\infty})_{*}$  and induces

for any  $(\mathcal{N}^{\otimes} \to \mathcal{V}''^{\otimes} \times \mathcal{W}''^{\otimes}, \Lambda'') \in \omega \operatorname{BEnr}_*$  lying over  $(\mathcal{V}''^{\otimes} \to \operatorname{Ass}, \mathcal{W}''^{\otimes} \to \operatorname{Ass}, \mathcal{H}'') \in (\operatorname{Op}_{\infty} \times \operatorname{Op}_{\infty})_*$  a pullback square

Notation 4.8. Let  $\mathcal{V}^{\otimes} \to Ass, \mathcal{W}^{\otimes} \to Ass$  be small  $\infty$ -operads. For every set  $\mathcal{H}$  of totally small weights over  $\mathcal{V}, \mathcal{W}$  let  $_{\mathcal{V}} \operatorname{BEnr}_{\mathcal{W}}(\mathcal{H})_*$  be the fiber of the functor  $\omega \operatorname{BEnr}_* \to (\operatorname{Op}_{\infty} \times \operatorname{Op}_{\infty})_*$  over  $\mathcal{H}$ .

Notation 4.9. Let

$$\omega \operatorname{BEnr}_{w} \subset (\operatorname{Op}_{\infty} \times \operatorname{Op}_{\infty})_{*} \times_{(\operatorname{Op}_{\infty} \times \operatorname{Op}_{\infty})} \omega \operatorname{BEnr}$$

be the subcategory whose objects are pairs  $(\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}, \mathcal{H})$  such that  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  admits H-weighted colimits and whose morphisms

$$(\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}, \mathcal{H}) \to (\mathcal{N}^{\circledast} \to \mathcal{V}'^{\otimes} \times \mathcal{W}'^{\otimes}, \mathcal{H}')$$

correspond to an enriched functor  $\mathcal{M}^{\circledast} \to \mathcal{N}^{\circledast}$  that sends  $\mathcal{H}$ -weighted colimits to  $\mathcal{H}'$ -weighted colimits.

**Lemma 4.10.** The inclusion  $\omega \operatorname{BEnr}_{w} \subset (\operatorname{Op}_{\infty} \times \operatorname{Op}_{\infty})_{*} \times_{(\operatorname{Op}_{\infty} \times \operatorname{Op}_{\infty})} \omega \operatorname{BEnr}$  lifts to an embedding  $\omega \operatorname{BEnr}_{w} \rightarrow \omega \operatorname{BEnr}_{*}$ .

*Proof.* We prove (1). The proofs of (2) is similar. Let  $Q \subset \omega BEnr_*$  be the full subcategory spanned by the pairs  $(\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}, \Lambda)$  with the following property: if  $\mathcal{H}$  is the set of weights underlying  $\Lambda$ , then  $\mathcal{M}$  admits  $\mathcal{H}$ -weighted colimits and  $\Lambda$  is the set of all  $\mathcal{H}$ -weighted colimit diagrams on  $\mathcal{M}$ . The functor  $\gamma : \omega \operatorname{BEnr}_* \to (\operatorname{Op}_{\infty} \times \operatorname{Op}_{\infty})_* \times_{(\operatorname{Op}_{\infty} \times \operatorname{Op}_{\infty})} \omega \operatorname{BEnr}$  restricts to a functor  $\gamma' : \mathfrak{Q} \to \omega \operatorname{BEnr}_w$ . The functor  $\gamma'$  is essentially surjective since for any  $(\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}, \mathcal{H})$  we can equip  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$ with the set of all  $\mathcal{H}$ -weighted colimit diagrams on  $\mathcal{M}$ . The functor  $\gamma$  induces embeddings on mapping spaces so that the restriction  $\gamma'$  also does. But by definition of Q the functor  $\gamma'$  induces equivalences on mapping spaces and so is an equivalence. This proves the claim.  $\square$ 

Notation 4.11. Let  $\mathcal{V}^{\otimes} \to Ass, \mathcal{W}^{\otimes} \to Ass$  be small  $\infty$ -operads and  $\mathcal{H}$  a set of totally small weights over  $\mathcal{V}, \mathcal{W}$ . We write

$$_{\mathcal{V}}\omega \operatorname{BEnr}_{\mathcal{W}}(\mathcal{H})$$

for the fiber of the functor  $\omega \operatorname{BEnr}_{w} \to (\operatorname{Op}_{\infty} \times \operatorname{Op}_{\infty})_{*}$  over  $\mathcal{H}$ .

Notation 4.12. Let  $(\mathcal{V}^{\otimes} \to Ass, S)$ ,  $(\mathcal{W}^{\otimes} \to Ass, T)$  be small localization pairs and  $\mathcal{H}$  a set of totally small weights over  $\mathcal{V}, \mathcal{W}$ .

- Let  ${}^{s}_{\mathcal{V}}BEnr_{\mathcal{W}}^{T}(\mathcal{H})$  be the pullback  ${}^{s}_{\mathcal{V}}BEnr_{\mathcal{W}}^{T} \times_{v\omega BEnr_{\mathcal{W}}} v\omega BEnr_{\mathcal{W}}(\mathcal{H}).$  Let  ${}^{s}_{\mathcal{V}}BEnr_{\mathcal{W}}^{T}(\mathcal{H})_{*}$  be the pullback  ${}^{s}_{\mathcal{V}}BEnr_{\mathcal{W}}^{T} \times_{v\omega BEnr_{\mathcal{W}}} v\omega BEnr_{\mathcal{W}}(\mathcal{H})_{*}.$

Let  $(\mathcal{V}^{\otimes} \to Ass, S)$ ,  $(\mathcal{W}^{\otimes} \to Ass, T)$  be small localization pairs and  $\mathcal{H}$  a set of totally small weights over  $\mathcal{V}, \mathcal{W}$ . The embedding  $\omega \text{BEnr}_w \rightarrow \omega \text{BEnr}_*$  over  $(\text{Op}_\infty \times \text{Op}_\infty)_*$  of Lemma 4.10 (1) induces on the fiber over  $(\mathcal{V}, \mathcal{W}, \mathcal{H})$  an embedding  $_{\mathcal{V}}\omega \operatorname{BEnr}_{\mathcal{W}}(\mathcal{H}) \subset _{\mathcal{V}}\omega \operatorname{BEnr}_{\mathcal{W}}(\mathcal{H})_*$  that restricts to an embedding  ${}_{\mathcal{V}}^{S}BEnr_{\mathcal{W}}^{T}(\mathcal{H}) \subset {}_{\mathcal{V}}^{S}BEnr_{\mathcal{W}}^{T}(\mathcal{H})_{*}.$ 

(1) Let  $\mathcal{V}^{\otimes} \to Ass$  be a presentably monoidal  $\infty$ -category,  $\mathcal{W}^{\otimes} \to Ass$  a small Notation 4.13.  $\infty$ -operad and  $\mathcal{H}$  a set of small left enriched diagrams over  $\mathcal{V}, \mathcal{W}$ . Let

$$_{\mathcal{V}} \operatorname{LEnr}_{\mathcal{W}}(\mathcal{H})_* \subset _{\mathcal{V}} \omega \widehat{\operatorname{BEnr}}_{\mathcal{W}}(\mathcal{H})_*$$

be the full subcategory of small left enriched  $\infty$ -categories equipped with a set of small left enriched diagrams.

(2) Let  $\mathcal{W}^{\otimes} \to Ass$  be a presentably monoidal  $\infty$ -category,  $\mathcal{V}^{\otimes} \to Ass$  a small  $\infty$ -operad and  $\mathcal{H}$  a set of small right enriched diagrams over  $\mathcal{V}, \mathcal{W}$ . Let

$$_{\mathcal{V}}\operatorname{REnr}_{\mathcal{W}}(\mathcal{H})_* \subset _{\mathcal{V}}\omega \overline{\operatorname{BEnr}}_{\mathcal{W}}(\mathcal{H})_*$$

be the full subcategory of small right enriched  $\infty$ -categories equipped with a set of small right enriched diagrams.

(3) Let  $\mathcal{V}^{\otimes} \to Ass, \mathcal{W}^{\otimes} \to Ass$  be presentably monoidal  $\infty$ -categories and  $\mathcal{H}$  a set of small bienriched diagrams over  $\mathcal{V}, \mathcal{W}$ . Let

$$_{\mathcal{V}}\operatorname{BEnr}_{\mathcal{W}}(\mathcal{H})_* \subset _{\mathcal{V}}\omega \overline{\operatorname{BEnr}}_{\mathcal{W}}(\mathcal{H})_*$$

be the full subcategory of small bi-enriched  $\infty$ -categories equipped with a set of small bi-enriched diagrams.

**Lemma 4.14.** Let  $\mathcal{C}$  be a presentable  $\infty$ -category and  $\gamma : \mathcal{C} \to \mathsf{Set}$  an accessible functor. Let  $\mathcal{D} \to \mathcal{C}$  be the cocartesian fibration classifying the functor  $\mathcal{C} \xrightarrow{\gamma} \mathcal{S} \xrightarrow{\mathcal{P}} \widehat{\mathsf{Poset}}$ . The  $\infty$ -category  $\mathcal{D}$  is presentable.

*Proof.* For every set X the power set  $\mathcal{P}(X)$  seen as poset is presentable as  $\infty$ -category. The functor

$$\mathfrak{C} \xrightarrow{\gamma} \mathfrak{S} \xrightarrow{\mathfrak{P}} \widehat{\operatorname{Poset}} \subset \widehat{\operatorname{Cat}}_{\infty}$$

is accessible because the power set functor is. Thus the cocartesian fibration  $\mathcal{D} \to \mathcal{C}$  classifies an accessible functor and is a cartesian fibration, whose fibers are presentable. By [7, Theorem 10.3.] this implies the claim.

- **Proposition 4.15.** (1) Let  $(\mathcal{V}^{\otimes} \to \operatorname{Ass}, \operatorname{S}), (\mathcal{W}^{\otimes} \to \operatorname{Ass}, \operatorname{T})$  be small localization pairs and  $\mathcal{H}$  a set of totally small weights over  $\mathcal{V}, \mathcal{W}$ . The  $\infty$ -category  ${}_{\mathcal{V}}^{\mathrm{S}}\operatorname{BEnr}_{W}^{\mathrm{T}}(\mathcal{H})_{*}$  is presentable.
  - (2) Let  $\mathcal{V}^{\otimes} \to Ass$  be a presentably monoidal  $\infty$ -category,  $\mathcal{W}^{\otimes} \to Ass$  a small  $\infty$ -operad and  $\mathcal{H}$  a set of small left enriched diagrams over  $\mathcal{V}, \mathcal{W}$ . The  $\infty$ -category  $_{\mathcal{V}}LEnr_{\mathcal{W}}(\mathcal{H})_*$  is presentable.
  - (3) Let  $\mathcal{W}^{\otimes} \to Ass$  be a presentably monoidal  $\infty$ -category,  $\mathcal{V}^{\otimes} \to Ass$  a small  $\infty$ -operad and  $\mathcal{H}$  a set of small right enriched diagrams over  $\mathcal{V}, \mathcal{W}$ . The  $\infty$ -category  $_{\mathcal{V}} \operatorname{REnr}_{\mathcal{W}}(\mathcal{H})_*$  is presentable.
  - (4) Let V<sup>⊗</sup> → Ass, W<sup>⊗</sup> → Ass be presentably monoidal ∞-categories and H a set of small bienriched diagrams over V, W. The ∞-category <sub>V</sub>BEnr<sub>W</sub>(H)<sub>\*</sub> is presentable.

*Proof.* We apply Lemma 4.14. (1): We apply Lemma 4.14. By [9, Proposition 2.57.] the  $\infty$ -category  ${}_{\mathcal{V}}^{S}\omega \operatorname{BEnr}_{\mathcal{W}}^{T}$  is compactly generated. Let  $\gamma$  be the functor  ${}_{\mathcal{V}}^{S}\omega \operatorname{BEnr}_{\mathcal{W}}^{T} \to \operatorname{Set}$  that sends a S, T-bi-enriched  $\infty$ -category  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  to the set of equivalence classes of H-weighted diagrams on  $\mathcal{M}$  for some  $H \in \mathcal{H}$ . Since  $\mathcal{H}$  is a set of totally small weights, the functor  $\gamma$  is accessible. So we can apply Lemma 4.14 to deduce the claim.

(2): Let  $\kappa$  be a small regular cardinal such that  $\mathcal{V}^{\otimes} \to \operatorname{Ass}$  is a  $\kappa$ -compactly generated monoidal  $\infty$ -category. By Corollary 2.142 (1) taking pullback along the monoidal embedding  $(\mathcal{V}^{\kappa})^{\otimes} \subset \mathcal{V}^{\otimes}$  induces an equivalence  $_{\mathcal{V}}\operatorname{LEnr}_{\mathcal{W}} \simeq \frac{\kappa}{\mathcal{V}^{\kappa}}\operatorname{LEnr}_{\mathcal{W}}$ . Thus  $_{\mathcal{V}}\operatorname{LEnr}_{\mathcal{W}}$  is presentable. Let  $\gamma$  be the functor  $_{\mathcal{V}}\operatorname{LEnr}_{\mathcal{W}} \to \operatorname{Set}$  that sends a left enriched  $\infty$ -category  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  to the set of equivalence classes of H-weighted diagrams on  $\mathcal{M}$  for some  $\mathcal{H} \in \mathcal{H}$ . Since  $\mathcal{H}$  is a set of small left enriched weights, the functor  $\gamma$  is accessible. So we can apply Lemma 4.14 to deduce the claim. (3) is dual to (2). (4): Let  $\kappa, \tau$  be small regular cardinals such that  $\mathcal{V}^{\otimes} \to \operatorname{Ass}$  is a  $\kappa$ -compactly generated monoidal  $\infty$ -category and  $\mathcal{W}^{\otimes} \to \operatorname{Ass}$  is a  $\tau$ -compactly generated monoidal  $\infty$ -category and  $\mathcal{W}^{\otimes} \to \operatorname{Ass}$  is a  $\tau$ -compactly generated monoidal  $\infty$ -category. By Corollary 2.142 (3) taking pullback along the monoidal embeddings  $(\mathcal{V}^{\kappa})^{\otimes} \subset \mathcal{V}^{\otimes}, (\mathcal{W}^{\tau})^{\otimes} \subset \mathcal{W}^{\otimes}$  induces an equivalence  $_{\nu}\operatorname{BEnr}_{\mathcal{W}} \simeq \frac{\kappa}{\mathcal{V}^{\kappa}}\operatorname{BEnr}_{\mathcal{W}^{\tau}}^{\tau}$ . Thus  $_{\nu}\operatorname{BEnr}_{\mathcal{W}}$  is presentable. Let  $\gamma$  be the functor  $_{\nu}\operatorname{BEnr}_{\mathcal{W}} \to \operatorname{Set}$  that sends a bi-enriched  $\infty$ -category  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  to the set of equivalence classes of  $\mathcal{H}$ -weighted diagrams on  $\mathcal{M}$  for some  $\mathcal{H} \in \mathcal{H}$ . Since  $\mathcal{H}$  is a set of small enriched weights, the functor  $\gamma$  is accessible. So we apply Lemma 4.14.

**Remark 4.16.** The  $\infty$ -category  $\omega$ BEnr<sub>\*</sub> admits small colimits. The colimit of a functor  $K \to \omega$ BEnr<sub>\*</sub> is the pair consisting of the colimit of the functor  $F : K \to \omega$ BEnr<sub>\*</sub>  $\to \omega$ BEnr and the set of diagrams that precisely consists of the diagrams that are the image of a diagram on F(k) under the enriched functor F(k)<sup> $\otimes$ </sup>  $\to$  colim(F)<sup> $\otimes$ </sup> for some k  $\in$  K.

Let  $(\mathcal{V}^{\otimes} \to \operatorname{Ass}, S)$ ,  $(\mathcal{W}^{\otimes} \to \operatorname{Ass}, T)$  be small localization pairs and  $\mathcal{H}$  a set of totally small weights over  $\mathcal{V}, \mathcal{W}$ . Similarly, the  $\infty$ -category  $_{\mathcal{V}}^{S}\omega \operatorname{BEnr}_{\mathcal{W}}^{T}(\mathcal{H})_{*}$  admits small colimits. The colimit of a functor  $K \to _{\mathcal{V}}^{S}\omega \operatorname{BEnr}_{\mathcal{W}}^{T}(\mathcal{H})_{*}$  is the pair consisting of the colimit of the functor  $F : K \to _{\mathcal{V}}^{S}\omega \operatorname{BEnr}_{\mathcal{W}}^{T}(\mathcal{H})_{*} \to _{\mathcal{V}}^{S}\omega \operatorname{BEnr}_{\mathcal{W}}^{T}(\mathcal{H})_{*}$  and the set of diagrams that precisely consists of the  $\mathcal{H}$ -weighted diagrams that are the image of a  $\mathcal{H}$ -weighted diagram on F(k) under the  $\mathcal{V}, \mathcal{W}$ -enriched functor  $F(k)^{\otimes} \to \operatorname{colim}(F)^{\otimes}$  for some  $k \in K$ .

**Proposition 4.17.** (1) Let  $(\mathcal{V}^{\otimes} \to Ass, S)$ ,  $(\mathcal{W}^{\otimes} \to Ass, T)$  be small localization pairs and  $\mathcal{H}$  a set of totally small weights over  $\mathcal{V}, \mathcal{W}$ . The embedding

$$^{\mathrm{S}}_{\mathcal{V}}\mathrm{BEnr}^{\mathrm{T}}_{\mathcal{W}}(\mathcal{H}) \subset ^{\mathrm{S}}_{\mathcal{V}}\mathrm{BEnr}^{\mathrm{T}}_{\mathcal{W}}(\mathcal{H})_{*}$$

is accessible and admits a left adjoint.

(2) Let V<sup>⊗</sup> → Ass be a presentably monoidal ∞-category, W<sup>⊗</sup> → Ass a small ∞-operad and ℋ a set of small left enriched diagrams over V, W. The embedding

$$_{\mathcal{V}} \operatorname{LEnr}_{\mathcal{W}}(\mathcal{H}) \subset _{\mathcal{V}} \operatorname{LEnr}_{\mathcal{W}}(\mathcal{H}),$$

is accessible and admits a left adjoint.

(3) Let W<sup>⊗</sup> → Ass be a presentably monoidal ∞-category, V<sup>⊗</sup> → Ass a small ∞-operad and ℋ a set of small right enriched diagrams over V, W. The embedding

$$_{\mathcal{V}}\operatorname{REnr}_{\mathcal{W}}(\mathcal{H}) \subset _{\mathcal{V}}\operatorname{REnr}_{\mathcal{W}}(\mathcal{H})_{*}$$

is accessible and admits a left adjoint.

(4) Let  $\mathcal{V}^{\otimes} \to \operatorname{Ass}, \mathcal{W}^{\otimes} \to \operatorname{Ass}$  be presentably monoidal  $\infty$ -categories and  $\mathfrak{H}$  a set of small bienriched diagrams over  $\mathcal{V}, \mathcal{W}$ . The embedding

$$_{\mathcal{V}}\operatorname{BEnr}_{\mathcal{W}}(\mathcal{H}) \subset _{\mathcal{V}}\operatorname{BEnr}_{\mathcal{W}}(\mathcal{H})_{*}$$

is accessible and admits a left adjoint.

Proof. By Corollary 3.87 the embeddings of the statement admit a left adjoint. So it remains to see that the embeddings of the statement are accessible. We prove this for (1). The proofs of (2)-(4) are similar. Since  $\mathcal{H}$  is small and  ${}_{\mathcal{V}}^{S}BEnr_{\mathcal{W}}^{T}$  is compactly generated by [9, Proposition 2.57.], there is a small regular cardinal  $\kappa$  such that for any weight  $(\mathcal{H}, \mathcal{J}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}) \in \mathcal{H}$  the S, T-bi-enriched  $\infty$ -category  $\mathcal{J}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  is a  $\kappa$ -compact object of  ${}_{\mathcal{V}}^{S}BEnr_{\mathcal{W}}^{T}$ . We will prove that the embedding  ${}_{\mathcal{V}}^{S}BEnr_{\mathcal{W}}^{T}(\mathcal{H}) \subset {}_{\mathcal{V}}^{S}BEnr_{\mathcal{W}}^{T}(\mathcal{H})_{*}$  preserves small  $\kappa$ -filtered colimits.

Let K be a small  $\kappa$ -filtered  $\infty$ -category,  $\Psi : K \to {}^{S}_{\mathcal{V}} \omega BEnr_{\mathcal{W}}^{T}(\mathcal{H})$  a functor and  $(\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}, \Lambda)$  the colimit of  $\Psi$  taken in  ${}^{S}_{\mathcal{V}} \omega BEnr_{\mathcal{W}}^{T}(\mathcal{H})_{*}$ . We will show that  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  admits  $\mathcal{H}$ -weighted colimits and  $\Lambda$  is precisely the set of H-weighted colimit diagrams on  $\mathcal{M}$  for some  $H \in \mathcal{H}$ . Observe that for every  $k \in K$  the canonical map  $\psi : \Psi(k)^{\circledast} \to \mathcal{M}^{\circledast}$  preserves  $\mathcal{H}$ -weighted colimits because for every  $V_1, ..., V_n \in \mathcal{V}, W_1, ..., W_m \in \mathcal{W}$  for  $n, m \ge 0$  and  $X, Y \in \Psi(k)$  the canonical map

$$\operatorname{colim}_{\alpha:k \to \ell} \operatorname{Mul}_{\Psi(\ell)}(V_1, ..., V_n, \Psi(\alpha)(X), W_1, ..., W_m; \Psi(\alpha)(Y)) \to$$

$$\operatorname{Mul}_{\mathcal{M}}(V_1, ..., V_n, \psi(X), W_1, ..., W_m; \psi(Y))$$

is an equivalence, and similarly for  $X,Y\in \mathcal{P}BEnv(\Psi(k))$  the canonical map

 $\operatorname{colim}_{\alpha:k \to \ell} \operatorname{Mul}_{\mathcal{PBEnv}(\Psi(\ell))}(V_1, ..., V_n, \Psi(\alpha)_!(X), W_1, ..., W_m; \Psi(\alpha)_!(Y)) \to$ 

(14) 
$$\operatorname{Mul}_{\mathcal{PBEnv}(\mathcal{M})}(V_1, ..., V_n, \psi_!(X), W_1, ..., W_m; \psi_!(Y))$$

is an equivalence. This implies that  $\Lambda$  consists of H-weighted colimit diagrams on  $\mathcal{M}$  for some  $H \in \mathcal{H}$ . Let  $(H, \mathcal{J}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}) \in \mathcal{H}$  be a weight. Then there is a  $k \in K$  so that for any morphism  $k \to \ell$  in K the S, T-bi-enriched  $\infty$ -category  $\Psi(\ell)^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  admits H-weighted colimits. Let  $F : \mathcal{J}^{\circledast} \to \mathcal{M}^{\circledast}$  be an enriched functor. By assumption  $\mathcal{J}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  is  $\kappa$ -compact so that F factors as  $\mathcal{J}^{\circledast} \xrightarrow{G} \Psi(r)^{\circledast} \to \mathcal{M}^{\circledast}$  for some  $r \in K$ , which we can assume to receive a morphism  $k \to r$ . The map  $G : \mathcal{J}^{\circledast} \to \Psi(r)^{\circledast}$  admits a H-weighted colimit that is sent to the H-weighted colimit of F. So  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  admits  $\mathcal{H}$ -weighted colimits.

It remains to see that every H-weighted colimit diagram  $(H, F : \mathcal{J}^{\circledast} \to \mathcal{M}^{\circledast}, \lambda : F_{!}(H) \to Y)$  on  $\mathcal{M}$  for some  $H \in \mathcal{H}$  belongs to  $\Lambda$ . Since  $F : \mathcal{J}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  is  $\kappa$ -compact, F factors as  $\mathcal{J}^{\circledast} \xrightarrow{G} \Psi(k)^{\circledast} \to \mathcal{M}^{\circledast}$ for some  $k \in K$ . By equivalence (4.1) there is a morphism  $k \to \ell$  in K and a diagram  $\nu$  on  $\Psi(\ell)$ that is transported by the enriched functor  $\Psi(\ell)^{\circledast} \to \mathcal{M}^{\circledast}$  to  $(H, F : \mathcal{J}^{\circledast} \to \mathcal{M}^{\circledast}, \lambda : F_{!}(H) \to Y)$ . As  $\nu$  is transported by  $\Psi(\ell)^{\circledast} \to \mathcal{M}^{\circledast}$  to a  $\mathcal{H}$ -weighted colimit diagram and  $\Psi$  sends morphisms in Kto enriched functors preserving  $\mathcal{H}$ -weighted colimits and for any  $k \in K$  the  $\infty$ -category  $\Psi(k)$  admits  $\mathcal{H}$ -weighted colimits, there is a morphism  $\ell \to r$  such that  $\nu$  is transported by the enriched functor  $\Psi(\ell) \to \Psi(r)$  to a  $\mathcal{H}$ -weighted colimit diagram  $\nu'$ . Since  $\Psi(r) \in {}^{S}_{\mathcal{V}} \omega BEnr_{\mathcal{W}}^{T}(\mathcal{H})$ , we find that  $\nu'$  is a diagram  $\Psi(r)$  is equipped with. Hence  $(H, F : \mathcal{J}^{\circledast} \to \mathcal{M}^{\circledast}, \lambda : F_{!}(H) \to Y)$  belongs to  $\Lambda$ .

**Corollary 4.18.** (1) Let  $(\mathcal{V}^{\otimes} \to \operatorname{Ass}, \operatorname{S}), (\mathcal{W}^{\otimes} \to \operatorname{Ass}, \operatorname{T})$  be small localization pairs and  $\mathcal{H}$  a set of totally small weights over  $\mathcal{V}, \mathcal{W}$ . The  $\infty$ -category  ${}_{\mathcal{V}}^{\mathrm{S}}\operatorname{BEnr}_{\mathcal{W}}^{\mathrm{T}}(\mathcal{H})$  is presentable.

- (2) Let  $\mathcal{V}^{\otimes} \to Ass$  be a presentably monoidal  $\infty$ -category,  $\mathcal{W}^{\otimes} \to Ass$  a small  $\infty$ -operad and  $\mathcal{H}$  a set of small left enriched diagrams over  $\mathcal{V}, \mathcal{W}$ . The  $\infty$ -category  $_{\mathcal{V}}LEnr_{\mathcal{W}}(\mathcal{H})$  is presentable.
- (3) Let  $\mathcal{W}^{\otimes} \to Ass$  be a presentably monoidal  $\infty$ -category,  $\mathcal{V}^{\otimes} \to Ass$  a small  $\infty$ -operad and  $\mathcal{H}$  a set of small right enriched diagrams over  $\mathcal{V}, \mathcal{W}$ . The  $\infty$ -category  $_{\mathcal{V}} \operatorname{Renr}_{\mathcal{W}}(\mathcal{H})$  is presentable.
- (4) Let V<sup>⊗</sup> → Ass, W<sup>⊗</sup> → Ass be presentably monoidal ∞-categories and ℋ a set of small bienriched diagrams over V, W. The ∞-category <sub>V</sub>BEnr<sub>W</sub>(ℋ) is presentable.

**Corollary 4.19.** The functor  $\omega \operatorname{BEnr}_{w} \to (\operatorname{Op}_{\infty} \times \operatorname{Op}_{\infty})_{*}$  is a cocartesian and cartesian fibration and the functor  $\omega \operatorname{BEnr}_{w} \to \omega \operatorname{BEnr}$  sends cartesian lifts of morphisms in  $(\operatorname{Op}_{\infty} \times \operatorname{Op}_{\infty})_{*}$  to cartesian lifts in  $\operatorname{Op}_{\infty} \times \operatorname{Op}_{\infty}$ .

Proof. By Lemma 4.7 the functor  $\omega BEnr_* \rightarrow (Op_{\infty} \times Op_{\infty})_*$  is a cartesian fibration and the functor  $\omega BEnr_* \rightarrow \omega BEnr$  sends cartesian lifts of morphisms in  $(Op_{\infty} \times Op_{\infty})_*$  to cartesian lifts in  $Op_{\infty} \times Op_{\infty}$ . By Corollary 3.60 the latter cartesian fibration restricts to a cartesian fibration  $\omega BEnr_w \rightarrow (Op_{\infty} \times Op_{\infty})_*$  with the same cartesian morphisms. This cartesian fibration is also a cocartesian fibration since the fiber transports admit left adjoints by Proposition 4.17.

4.2. The tensor product of enriched  $\infty$ -categories with weighted colimits. In this subsection we construct a tensor product for left enriched  $\infty$ -categories with weighted diagrams and apply the theory of adjoining weighted colimits (Proposition 3.85) to construct a tensor product for left enriched  $\infty$ -categories with weighted colimits (Theorem 4.42 and Theorem 4.42) from the first one.

**Notation 4.20.** Let  $\mathcal{V}^{\otimes} \to Ass$  be a small  $\infty$ -operad and  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes}$  a weakly left enriched  $\infty$ -category.

- (1) Let  $LD(\mathcal{M})$  be the poset of sets of totally small left diagrams on  $\mathcal{M}$  ordered by inclusion.
- (2) Let  $\omega(\mathcal{V})$  be the poset of sets of totally small left weights over  $\mathcal{V}$  ordered by inclusion.

**Remark 4.21.** (1) The functor

$$\omega : Ho(Mon) \to Poset,$$

is lax symmetric monoidal with respect to cartesian symmetric monoidal structures.

For every  $\mathcal{V}_1^{\otimes}, ..., \mathcal{V}_n^{\otimes} \in Mon$  for  $n \ge 0$  the structure maps

$$\prod_{i=1}^{n} \omega(\mathcal{V}_{i}) \to \omega(\prod_{i=1}^{n} \mathcal{V}_{i})$$

send  $(\mathcal{H}_1, ..., \mathcal{H}_n)$  to  $\mathcal{H}_1 \boxtimes ... \boxtimes \mathcal{H}_n$ .

(2) The functor

$$LD: Ho(LPEnr) \rightarrow Poset$$

is lax symmetric monoidal with respect to cartesian symmetric monoidal structures. For every  $\mathcal{M}_1^{\circledast} \to \mathcal{V}_1^{\circledast}, ..., \mathcal{M}_n^{\circledast} \to \mathcal{V}_n^{\circledast}$  for  $n \ge 0$  the structure maps

$$\prod_{i=1}^n \mathrm{LD}(\mathcal{M}_i) \to \mathrm{LD}\big(\prod_{i=1}^n \mathcal{M}_i\big)$$

send  $(\Lambda_1, ..., \Lambda_n)$  to  $\Lambda_1 \boxtimes ... \boxtimes \Lambda_n$ .

(3) Let  $\nu : \mathsf{Ho}(\text{LPEnr}) \to \mathsf{Ho}(\text{Mon})$  be the forgetful functor. The natural transformation  $\text{LD} \to \omega \circ \nu$  is symmetric monoidal.

Notation 4.22. Let  $\kappa$  be a small regular cardinal. Let  $\operatorname{Mon}_{\kappa}^{\otimes} \subset \operatorname{Mon}^{\times}$  be the symmetric suboperad whose colors are the monoidal  $\infty$ -categories compatible with  $\kappa$ -small colimits and whose multimorphisms  $\mathcal{V}_{1}^{\otimes}, ..., \mathcal{V}_{n}^{\otimes} \to \mathcal{W}^{\otimes}$  correspond to monoidal functors  $\mathcal{V}_{1}^{\otimes} \times_{\operatorname{Ass}} ... \times_{\operatorname{Ass}} \mathcal{V}_{n}^{\otimes} \to \mathcal{W}^{\otimes}$  preserving  $\kappa$ -small colimits component-wise.

Notation 4.23. For  $\sigma$  the strongly inaccessible cardinal corresponding to the small universe we set

$$\operatorname{ccMon}^{\otimes} \coloneqq \widehat{\operatorname{Mon}}_{\sigma}^{\otimes}.$$

Notation 4.24. Let

$$\operatorname{PrMon}^{\otimes} \subset \operatorname{ccMon}^{\otimes}$$

be the full symmetric suboperad spanned by the presentably monoidal  $\infty$ -categories.

By [16, Proposition 4.8.1.15.]  $\operatorname{Mon}_{\kappa}^{\otimes} \to \operatorname{Comm}, \operatorname{PrMon}^{\otimes} \to \operatorname{Comm}$  are symmetric monoidal  $\infty$ -categories.

Notation 4.25. Let  $\kappa$  be a small regular cardinal. Let

 ${}^{\kappa}\mathrm{LEnr}^{\otimes} \subset \mathrm{Mon}_{\kappa}^{\otimes} \times_{\mathrm{Mon}^{\times}} (\mathrm{LPEnr}_{\varnothing})^{\times}$ 

be the full symmetric suboperad spanned by the left  $\kappa$ -enriched  $\infty$ -categories.

**Proposition 4.26.** Let  $\kappa$  be a small regular cardinal. The map of symmetric  $\infty$ -operads

$$^{\kappa} \text{LEnr}^{\otimes} \to \text{Mon}_{\kappa}^{\otimes}$$

is a cocartesian fibration.

*Proof.* We first prove that the symmetric monoidal functor  $\phi$ : (LPEnr<sub>∅</sub>)<sup>×</sup> → Mon<sup>×</sup> is a cocartesian fibration. By Proposition 2.127 the underlying functor is a cocartesian fibration. So by [11, Lemma 2.44.] it is enough to check that the collection of  $\phi$ -cocartesian morphisms is stable under product. This follows from the description of  $\phi$ -cocartesian morphisms of Proposition 2.127. By [9, Corollary 4.54.] for every small monoidal ∞-category  $\mathcal{V}^{\otimes} \to Ass$  compatible with  $\kappa$ -small colimits the full subcategory  $\stackrel{\kappa}{_{V}}LEnr \subset _{\mathcal{V}}LPEnr$  is a localization. Moreover for every monoidal functor  $\mathcal{V}^{\otimes} \to \mathcal{V}'^{\otimes}$  preserving  $\kappa$ -small colimits the induced functor  $_{\mathcal{V}}LPEnr \to _{\mathcal{V}'}LPEnr$  preserves local equivalences. Thus by [16, Lemma 2.2.4.11.] the statement follows if we have shown that for every small monoidal ∞-categories  $\mathcal{V}^{\otimes} \to Ass$ ,  $\mathcal{V}'^{\otimes} \to Ass$  and monoidal functors  $\alpha : \mathcal{V}^{\otimes} \times_{Ass} \mathcal{V}'^{\otimes} \to \mathcal{V}'^{\otimes}$  preserving  $\kappa$ -small colimits component-wise the induced functor  $_{\mathcal{V}}LPEnr \times_{\mathcal{V}'}LPEnr \to _{\mathcal{V}''}LPEnr$  preserves local equivalences. In view of Proposition 2.127 and the description of local equivalences of [9, Corollary

4.54.] this follows from the fact that the functor  $\mathcal{P}(\mathcal{V}) \times \mathcal{P}(\mathcal{V}') \to \mathcal{P}(\mathcal{V} \times \mathcal{V}') \to \mathcal{P}(\mathcal{V}'')$  sends local equivalences for the localization  $\mathrm{Ind}_{\kappa}(\mathcal{V}) \times \mathrm{Ind}_{\kappa}(\mathcal{V}')$  to local equivalences for the localization  $\mathrm{Ind}_{\kappa}(\mathcal{V}')$ . This follows from the fact that  $\alpha$  preserves  $\kappa$ -small colimits component-wise.

Notation 4.27. Let  $\kappa$  be a small regular cardinal. Let

$$^{\kappa} \text{LEnr}^{\otimes}_{\star} \rightarrow ^{\kappa} \text{LEnr}^{\otimes}$$

be the cocartesian fibration of symmetric monoidal  $\infty$ -categories classifying the lax symmetric monoidal functor

$$^{\kappa}\text{LEnr} \subset \text{LPEnr} \rightarrow \text{Ho}(\text{LPEnr}) \xrightarrow{\text{LD}} \widehat{\text{Poset}} \subset \widehat{\text{Cat}}_{\infty}.$$

Let

(15) 
$$(\operatorname{Mon}_{\kappa})^{\otimes}_{*} \to \operatorname{Mon}_{\kappa}^{\otimes}$$

be the cocartesian fibration of symmetric monoidal  $\infty$ -categories classifying the lax symmetric monoidal functor

 $\operatorname{Mon}_{\kappa} \subset \operatorname{Mon} \to \operatorname{Ho}(\operatorname{Mon}) \xrightarrow{\omega} \operatorname{Poset} \subset \operatorname{\widehat{Cat}}_{\infty}.$ 

Notation 4.28. For  $\sigma$  the strongly inaccessible cardinal corresponding to the small universe we set

$$\operatorname{ccMon}_*^{\otimes} := (\operatorname{Mon}_{\sigma})_*^{\otimes}, \operatorname{LQEnr}_*^{\otimes} := {}^{\sigma} \widehat{\operatorname{LEnr}}_*^{\otimes}$$

**Remark 4.29.** By Remark 4.21 (3) the transformation  $LD \rightarrow \omega \circ \nu$  is symmetric monoidal and so classifies a map

(16) 
$${}^{\kappa} \operatorname{LEnr}^{\otimes}_{*} \to (\operatorname{Mon}_{\kappa})^{\otimes}_{*} \times_{\operatorname{Mon}^{\otimes}} {}^{\kappa} \operatorname{LEnr}^{\otimes}$$

of cocartesian fibrations over  ${}^{\kappa}\text{LEnr}^{\otimes}$  and so a symmetric monoidal functor.

**Remark 4.30.** Since the fibers of the cocartesian fibration (15) are posets, for any symmetric  $\infty$ -operad  $\mathcal{O}^{\otimes} \to \text{Comm}$  the canonical functor

$$\operatorname{Alg}_{\mathcal{O}}((\operatorname{Mon}_{\kappa})_{*}) \to \operatorname{Alg}_{\mathcal{O}}(\operatorname{Mon}_{\kappa}) \times_{(\prod_{X \in \mathcal{O}} \operatorname{Mon}_{\kappa})} \prod_{X \in \mathcal{O}} (\operatorname{Mon}_{\kappa})_{*}$$

is fully faithful and the essential image are the triples  $(\mathcal{V}, \mathcal{H})$ , where  $\mathcal{V} \in \operatorname{Alg}_{\mathcal{O}}(\operatorname{Mon}_{\kappa})$  and  $\mathcal{H} := (\mathcal{H}_X)_{X \in \mathcal{O}}$  is a family such that  $\mathcal{H}_X$  is a set of totally small weights over  $\mathcal{V}(X)$  for every  $X \in \mathcal{O}$  subject to the condition that for every multi-morphism  $\alpha : X_1, ..., Y_n \to Y$  in  $\mathcal{O}$  we have  $\alpha_!(\mathcal{H}_{X_1} \boxtimes ... \boxtimes \mathcal{H}_{X_n}) \subset \mathcal{H}_Y$ . Thus we will denote an  $\mathcal{O}$ -algebra in  $(\operatorname{Mon}_{\kappa})_*$  as a pair  $(\mathcal{V}, \mathcal{H})$ .

**Example 4.31.** Let  $\mathbb{O}^{\otimes} \to \operatorname{Comm}$  be an unital monochromatic symmetric  $\infty$ -operad, i.e.  $\mathbb{O}$  is contractible, for example the  $\infty$ -operads  $\mathbb{E}_n$  for  $1 \leq n \leq \infty$ . Then an  $\mathbb{O}$ -algebra in  $(\operatorname{Mon}_{\kappa})^{\otimes}_*$  is a pair  $(\mathcal{V}, \mathcal{H})$ , where  $\mathcal{V} \in \operatorname{Alg}_{\mathbb{O}}(\operatorname{Mon}_{\kappa})$  and  $\mathcal{H}$  is a set of totally small left weights over  $\mathcal{V}$ . The condition that for every multi-morphism  $\alpha$  in  $\mathbb{O}$  corresponding to an n-ary operation we have  $\alpha_!(\mathcal{H} \boxtimes ... \boxtimes \mathcal{H}) \subset \mathcal{H}$  is automatic since the composition  $\{1\} \times ... \times \{1\} \times \mathbb{C} \times \{1\} \times ... \times \{1\} \to \mathbb{C}^{\times n} \xrightarrow{\alpha_!} \mathbb{C}$  is the identity.

**Example 4.32.** Let  $LM^{\otimes} \to Comm$  be the symmetric  $\infty$ -operad governing left modules [16, Notation 4.2.1.6.] and  $\kappa$  a small regular cardinal. A LM-algebra in  $Mon_{\kappa}$  corresponds to a left module in  $Mon_{\kappa}$  [16, Proposition 4.2.2.9.]. So a LM-algebra in  $Mon_{\kappa}$  encodes a left action of an associative algebra in  $Mon_{\kappa}$  - corresponding to two braided monoidal  $\infty$ -categories  $\mathcal{V}^{\boxtimes} \to \mathbb{E}_2$  compatible with  $\kappa$ -small colimits, - on an object  $\mathcal{W}^{\otimes} \to Ass$  of  $Mon_{\kappa}$ . Let  $\mu : \mathcal{V}^{\otimes} \times_{Comm} \mathcal{W}^{\otimes} \to \mathcal{W}^{\otimes}$  be the underlying monoidal left action functor and  $\mathcal{H}$  a set of totally small left weights over  $\mathcal{V}$ . By the axioms of a left action the functor  $\mu(\mathbb{1}_{\mathcal{V}}, -) : \mathcal{W}^{\otimes} \to \mathcal{W}^{\otimes}$  is the identity and so sends weights of  $\mathcal{H}'$  to weights of  $\mathcal{H}'$ . If also  $\mu(-,\mathbb{1}_{\mathcal{W}}) : \mathcal{V}^{\otimes} \to \mathcal{W}^{\otimes}$  sends weights of  $\mathcal{H}$  to weights of  $\mathcal{H}'$ , then by Remark 4.30 the given LM-algebra uniquely refines to a LM-algebra in  $(Mon_{\kappa})_*$ .

#### Notation 4.33. Let

$$\operatorname{PrMon}_{*}^{\otimes} \subset \operatorname{PrMon}^{\otimes} \times_{\operatorname{ccMon}^{\otimes}} \operatorname{ccMon}_{*}^{\otimes}$$

be the full symmetric monoidal subcategory spanned by the presentably monoidal  $\infty$ -categories equipped with a set of small left enriched weights.

## Notation 4.34. Let

$$\operatorname{LEnr}^{\otimes}_{*} \subset \operatorname{PrMon}^{\otimes}_{*} \times_{\operatorname{ccMon}^{\otimes}} \operatorname{LQEnr}^{\otimes}_{*}$$

be the full symmetric monoidal subcategory spanned by the small left enriched  $\infty$ -categories equipped with a set of small left enriched diagrams.

The functor (16) for  $\kappa$  the strongly inaccessible cardinal corresponding to the small universe restricts to a map

(17) 
$$\operatorname{LEnr}^{\otimes}_{*} \to \operatorname{PrMon}^{\otimes}_{*} \times_{\operatorname{PrMon}^{\otimes}} \operatorname{LEnr}^{\otimes}$$

of cocartesian fibrations over  $\mathrm{LEnr}^\otimes$  and so a symmetric monoidal functor.

**Proposition 4.35.** (1) Let  $\kappa$  be a small regular cardinal. The symmetric monoidal functor

$$^{\kappa} \text{LEnr}^{\otimes}_* \to (\text{Mon}_{\kappa})^{\otimes}_*$$

is a cocartesian fibration and the symmetric monoidal functor (16) is a map of cocartesian fibrations over  $(Mon_{\kappa})^{\otimes}_{*}$ .

(2) The symmetric monoidal functor

$$\operatorname{LEnr}^{\otimes}_* \to \operatorname{PrMon}^{\otimes}_*$$

is a cocartesian fibration and the symmetric monoidal functor (17) is a map of cocartesian fibrations over  $\operatorname{PrMon}_*^{\otimes}$ .

*Proof.* We prove (1). The proof of (2) is similar. By Proposition 4.26 the map of symmetric ∞-operads  $\psi$  : <sup>κ</sup>LEnr<sup>⊗</sup> → Mon<sup>⊗</sup><sub>κ</sub> is a cocartesian fibration. Let  $\psi'$  be the underlying functor. We start with proving that the functor <sup>κ</sup>LEnr<sub>\*</sub> → (Mon<sub>κ</sub>)<sub>\*</sub> is a cocartesian fibration. Let  $(\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast}, \Lambda) \in {}^{\kappa}$ LEnr<sub>\*</sub> and  $\alpha : (\mathcal{V}^{\circledast}, \mathcal{H}) \to (\mathcal{V}^{(\ast}, \mathcal{H}')$  a morphism in  $(Mon_{\kappa})_*$ , where  $\mathcal{H}$  is the set of underlying weights of  $\Lambda$ . The  $\psi'$ -cocartesian lift  $\mathcal{M}^{\circledast} \to \mathcal{M}^{(\circledast)} := \alpha_!(\mathcal{M})^{\circledast}$  of  $\alpha$  sends the set  $\Lambda$  of  $\mathcal{H}$ -weighted diagrams to a set  $\Lambda'$  of  $\mathcal{H}'$ -weighted diagrams. By the choice of  $\Lambda'$  the morphism  $(\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes}, \Lambda) \to (\mathcal{M}^{(\circledast} \to \mathcal{V}^{(\circledast}, \Lambda'))$  in <sup>κ</sup>LEnr<sub>\*</sub> lies over  $\alpha : (\mathcal{V}^{\otimes}, \mathcal{H}) \to (\mathcal{V}^{(\otimes}, \mathcal{H}'))$  in  $(Mon_{\kappa})_*$  and induces for every  $(\mathcal{N}^{\circledast} \to \mathcal{V}^{''\otimes}, \Lambda'') \in {}^{\kappa}$ LEnr<sub>\*</sub> lying over  $(\mathcal{V}^{''\otimes} \to \operatorname{Ass}, \mathcal{H}'') \in (\operatorname{Mon}_{\kappa})_*$  a pullback square

Hence the functor  $\phi : {}^{\kappa}\text{LEnr}_{*} \to (\text{Mon}_{\kappa})_{*}$  is a cocartesian fibration. To prove the result, by [11, Lemma 2.44.] it is enough to see that the collection of  $\phi$ -cocartesian morphisms is stable under the tensor product on  ${}^{\kappa}\text{LEnr}_{*}$ , i.e. that for any  $(\mathcal{M}'^{\circledast} \to \mathcal{V}'^{\otimes}, \Lambda'') \in {}^{\kappa}\text{LEnr}_{*}$  and any morphism  $(\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes}, \Lambda) \to (\mathcal{M}'^{\circledast} \to \mathcal{V}'^{\otimes}, \Lambda')$  in  ${}^{\kappa}\text{LEnr}_{*}$  the tensor product is  $\phi$ -cocartesian:

$$((\mathcal{M} \otimes \mathcal{M}'')^{\circledast}, \Lambda \boxtimes \Lambda'') \to ((\mathcal{M}' \otimes \mathcal{M}'')^{\circledast}, \Lambda' \boxtimes \Lambda'').$$

The image of the latter morphism in <sup> $\kappa$ </sup>LEnr is  $\psi$ -cocartesian since  $\psi$  is a cocartesian fibration. Consequently, by the description of  $\phi$ -cocartesian morphisms, it is enough to show that every diagram of  $\Lambda' \boxtimes \Lambda''$  is transported from  $\Lambda \boxtimes \Lambda''$ . This follows immediately from the definition of  $\Lambda' \boxtimes \Lambda''$  and that  $\Lambda'$  is transported from  $\Lambda$  by the  $\psi'$ -cocartesian morphism  $\mathcal{M}^{\circledast} \to \mathcal{M}'^{\circledast}$  that induces an essentially surjective functor  $\mathcal{M} \to \mathcal{M}'$  by Proposition 2.127.

Proposition 4.35 implies the following corollary:

**Corollary 4.36.** Let  $\mathbb{O}^{\otimes} \to \text{Comm}$  be a symmetric  $\infty$ -operad and  $\kappa$  a small regular cardinal.

(1) For every  $\mathfrak{O}$ -algebra  $(\mathcal{V}, \mathcal{H}) : \mathfrak{O}^{\otimes} \to (\mathrm{Mon}_{\kappa})^{\otimes}_{*}$  the  $\mathfrak{O}$ -operad

$${}_{\mathcal{V}}^{\kappa} \mathrm{LEnr}(\mathcal{H})_{*}^{\otimes} \coloneqq \mathcal{O}^{\otimes} \times_{(\mathrm{Mon}_{\kappa})_{*}^{\otimes}} {}^{\kappa} \mathrm{LEnr}_{*}^{\otimes} \to \mathcal{O}^{\otimes}$$

is an O-monoidal  $\infty$ -category.

(2) For every  $\mathfrak{O}$ -algebra  $(\mathfrak{V}, \mathfrak{H}) : \mathfrak{O}^{\otimes} \to \operatorname{PrMon}^{\otimes}_*$  the  $\mathfrak{O}$ -operad

$$_{\mathcal{V}} \operatorname{LEnr}(\mathcal{H})^{\otimes}_{*} \coloneqq \mathcal{O}^{\otimes} \times_{\operatorname{PrMon}^{\otimes}} \operatorname{LEnr}^{\otimes}_{*} \to \mathcal{O}^{\otimes}$$

is an O-monoidal  $\infty$ -category.

Notation 4.37. Let  $\kappa$  be a small regular cardinal. Let

$${}^{\kappa} \operatorname{LEnr}_{\mathrm{w}}^{\otimes} \subset (\operatorname{Mon}_{\kappa})_{*}^{\otimes} \times_{\operatorname{Mon}^{\otimes}} {}^{\kappa} \operatorname{LEnr}^{\otimes}$$

be the symmetric suboperad whose colors are pairs  $(\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes}, \mathcal{H})$  such that  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes}$  admits  $\mathcal{H}$ -weighted colimits and whose multi-morphisms

$$(\mathcal{M}_1^{\circledast} \to \mathcal{V}_1^{\otimes}, \mathcal{H}_1), ..., (\mathcal{M}_n^{\circledast} \to \mathcal{V}_n^{\otimes}, \mathcal{H}_n) \to (\mathcal{N}^{\circledast} \to \mathcal{V}^{\otimes}, \mathcal{H})$$

correspond to an enriched functor  $\mathcal{M}_1^{\otimes} \times_{Ass} \dots \times_{Ass} \mathcal{M}_n^{\otimes} \to \mathcal{N}^{\otimes}$  that sends  $\mathcal{H}_1 \boxtimes \dots \boxtimes \mathcal{H}_n$ -weighted colimits to  $\mathcal{H}$ -weighted colimits.

Notation 4.38. Let  $\sigma$  be the strongly inaccessible cardinal corresponding to the small universe. Let

$$LQEnr_{w}^{\otimes} \coloneqq {}^{\sigma}\widehat{LEnr}_{w}^{\otimes} \to (\widehat{Mon}_{\sigma})_{*}^{\otimes}.$$

Notation 4.39. Let  $\sigma$  be the strongly inaccessible cardinal corresponding to the small universe. Let

$$\operatorname{LEnr}_{w}^{\otimes} \subset \operatorname{LQEnr}_{w}^{\otimes}$$

be the full symmetric suboperad whose colors are the pairs  $(\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast}, \mathcal{H})$  such that  $\mathcal{V}^{\circledast} \to Ass$  is a presentably monoidal  $\infty$ -category,  $\mathcal{H}$  is a set of small left enriched weights over  $\mathcal{V}$  and  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast}$  is a small left enriched  $\infty$ -category that admits  $\mathcal{H}$ -weighted colimits.

Lemma 4.10 implies the following:

**Corollary 4.40.** (1) Let  $\kappa$  be a small regular cardinal. The inclusion

 $^{\kappa} \text{LEnr}_{w}^{\otimes} \subset (\text{Mon}_{\kappa})_{*}^{\otimes} \times_{\text{Mon}_{*}^{\otimes}} ^{\kappa} \text{LEnr}^{\otimes}$ 

lifts to an embedding  ${}^{\kappa}\text{LEnr}_{w}^{\otimes} \to {}^{\kappa}\text{LEnr}_{*}^{\otimes}$ .

(2) The inclusion

$$\operatorname{LEnr}_{\mathrm{w}}^{\otimes} \subset \operatorname{PrMon}_{*}^{\otimes} \times_{\operatorname{PrMon}^{\otimes}} \operatorname{LEnr}^{\otimes}$$

lifts to an embedding  $\operatorname{LEnr}_{w}^{\otimes} \to \operatorname{LEnr}_{*}^{\otimes}$ .

Corollary 4.19 and Corollary 3.60 imply the following:

**Corollary 4.41.** Let  $\kappa$  be a small regular cardinal. The cartesian fibration of Corollary 4.19 restricts to the following cartesian fibrations with the same cartesian morphisms:

 $^{\kappa} \text{LEnr}_{w} \rightarrow (\text{Mon}_{\kappa})_{*}, \text{ LEnr}_{w} \rightarrow \text{PrMon}_{*}.$ 

**Theorem 4.42.** Let  $\mathbb{O}^{\otimes} \to \text{Comm}$  be a symmetric  $\infty$ -operad and  $\kappa$  a small regular cardinal.

(1) Let  $(\mathcal{V}, \mathcal{H}) : \mathbb{O}^{\otimes} \to (\mathrm{Mon}_{\kappa})^{\otimes}_{*}$  be an O-algebra. The full suboperad

$${}_{\mathcal{V}}^{\kappa} \operatorname{LEnr}(\mathcal{H})^{\otimes} \coloneqq \mathcal{O}^{\otimes} \times_{(\operatorname{Mon}_{*})^{\otimes}} {}^{\kappa} \operatorname{LEnr}_{w}^{\otimes} \subset {}_{\mathcal{V}}^{\kappa} \operatorname{LEnr}(\mathcal{H})_{*}^{\otimes}$$

is an accessible localization relative to  $0^{\otimes}$  and so a presentably 0-monoidal  $\infty$ -category.

(2) Let 
$$(\mathcal{V}, \mathcal{H}) : \mathbb{O}^{\otimes} \to (\mathsf{ccMon})^{\otimes}_{*}$$
 be an  $\mathbb{O}$ -algebra. The full suboperad

$$_{\mathcal{V}} LQEnr(\mathcal{H})^{\otimes} := \mathcal{O}^{\otimes} \times_{\mathsf{ccMon}^{\otimes}} LQEnr_{\mathsf{w}}^{\otimes} \subset _{\mathcal{V}} LQEnr(\mathcal{H})_{*}^{\otimes}$$

is an accessible localization relative to  $\mathbb{O}^{\otimes}$  and so a  $\mathbb{O}$ -monoidal  $\infty$ -category. (3) Let  $(\mathcal{V}, \mathcal{H}) : \mathbb{O}^{\otimes} \to \operatorname{PrMon}^{\otimes}_{*}$  be an  $\mathbb{O}$ -algebra. The full suboperad

(18)  $_{\text{vLEnr}}(\mathcal{H})^{\otimes} \coloneqq \mathcal{O}^{\otimes} \times_{\text{PrMon}^{\otimes}} \text{LEnr}_{u}^{\otimes}$ 

$$\operatorname{LEnr}(\mathcal{H})^{\otimes} \coloneqq \mathcal{O}^{\otimes} \times_{\operatorname{PrMon}^{\otimes}_{*}} \operatorname{LEnr}^{\otimes}_{\operatorname{w}} \subset \operatorname{VLEnr}(\mathcal{H})^{\otimes}_{*}$$

is an accessible localization relative to  $0^{\otimes}$  and so a presentably 0-monoidal  $\infty$ -category. The embedding

 $_{\mathcal{V}} \operatorname{LEnr}(\mathcal{H})^{\otimes} \subset _{\mathcal{V}} \operatorname{LQEnr}(\mathcal{H})^{\otimes}$ 

is an O-monoidal functor.

*Proof.* (1): By Corollary 4.17 the embedding of (1) induces fiberwise accessible functors. In view of Corollary 4.18 (1) it remains to see that the embedding of (1) admits a left adjoint relative to  $O^{\otimes}$ . We prove that the full suboperad <sup>κ</sup>LEnr<sup>∞</sup><sub>w</sub> ⊂ <sup>κ</sup>LEnr<sup>∞</sup><sub>\*</sub> is a localization relative to  $(Mon_{\kappa})^{\otimes}_{*}$ . For this it is enough to see that the full suboperad <sup>κ</sup>LEnr<sup>∞</sup><sub>w</sub> ⊂ <sup>κ</sup>LEnr<sup>∞</sup><sub>\*</sub> is a localization relative to Comm and that the local equivalences lie over equivalences in  $(Mon_{\kappa})^{\otimes}_{*}$ . By [16, Lemma 2.2.4.11.] it suffices to prove that the full subcategory <sup>κ</sup>LEnr<sub>w</sub> ⊂ <sup>κ</sup>LEnr<sub>\*</sub> is a localization whose local equivalences lie over equivalences in  $(Mon_{\kappa})^{\otimes}_{*}$  and that the collection of local equivalences is closed under the tensor product provided by the symmetric monoidal ∞-category <sup>κ</sup>LEnr<sup>∞</sup><sub>\*</sub> → Comm. Let  $(\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes}, \Lambda) \in ^{\kappa}LEnr_{*}$  and  $\mathcal{H}$  the set of left diagrams Λ. By Corollary 3.90 (1) the morphism

$$(\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes}, \Lambda) \to (\mathcal{P}BEnv_{\Lambda}^{\mathcal{H}}(\mathcal{M})_{LEnr_{e}}^{\otimes} \to \mathcal{V}^{\otimes}, \mathcal{H})$$

in <sup> $\kappa$ </sup>LEnr<sub>\*</sub> whose target belongs to <sup> $\kappa$ </sup>LEnr<sub>w</sub>, is a local equivalence with respect to the full subcategory <sup> $\kappa$ </sup>BEnr<sup> $\tau$ </sup><sub>w</sub>  $\subset {}^{\kappa}$ BEnr<sup> $\tau$ </sup><sub>\*</sub> and lies over the identity in (Mon<sub> $\kappa$ </sub>)<sup> $\otimes$ </sup><sub>\*</sub>. Moreover by Corollary 3.90 (1) the local equivalences are closed under the tensor product provided by <sup> $\kappa$ </sup>LEnr<sup> $\otimes$ </sup><sub>\*</sub>  $\rightarrow$  Comm.

(2) follows immediately from (1). (3): By Corollary 4.17 the embedding (18) induces fiberwise accessible functors. So it is enough to prove that the accessible localization

$$_{\mathcal{V}} \mathrm{LQEnr}(\mathcal{H})^{\otimes} \subset _{\mathcal{V}} \mathrm{LQEnr}(\mathcal{H})^{\otimes}_{*}$$

relative to  $0^{\otimes}$  of Corollary 4.42 restricts to a localization  ${}_{\mathcal{V}}\text{LEnr}(\mathcal{H})^{\otimes} \subset {}_{\mathcal{V}}\text{LEnr}(\mathcal{H})^{\otimes}_{*}$  relative to  $0^{\otimes}$ . This follows from Lemma 3.93 in view of the construction of the localization functor. Presentability follows from Corollary 4.18 (4).

**Remark 4.43.** Let  $\kappa$  be a small regular cardinal,  $\mathcal{V}^{\otimes} \to \mathbb{E}_2$  a braided monoidal  $\infty$ -category compatible with  $\kappa$ -small colimits, seen as an associative algebra in the monoidal  $\infty$ -category  $\operatorname{Mon}_{\kappa}$ , acting from the left on a monoidal  $\infty$ -category  $\mathcal{W}^{\otimes} \to \operatorname{Ass}$  compatible with  $\kappa$ -small colimits. Let  $\mu : \mathcal{V}^{\otimes} \times_{\operatorname{Ass}} \mathcal{W}^{\otimes} \to \mathcal{W}^{\otimes}$ be the monoidal left action functor. Let  $\mathcal{H}$  be a set of small left weights over  $\mathcal{V}$  and  $\mathcal{H}'$  a set of small left weights over  $\mathcal{W}$  such that  $\mu(-, \mathbb{1}_{\mathcal{W}}) : \mathcal{V}^{\otimes} \to \mathcal{W}^{\otimes}$  sends weights of  $\mathcal{H}$  to weights of  $\mathcal{H}'$ . By Example 4.32 the left action of  $\mathcal{V}^{\otimes} \to \operatorname{Ass}$  on  $\mathcal{W}^{\otimes} \to \operatorname{Ass}$  in  $\operatorname{Mon}_{\kappa}$  corresponds to a LM-algebra in  $*\operatorname{Mon}_{\kappa}$ .

(1) By Theorem 4.42 the  $\infty$ -category  ${}^{\kappa}_{\mathcal{V}} \text{LEnr}(\mathcal{H})$  carries a monoidal structure. For two left  $\kappa$ enriched  $\infty$ -categories  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast}, \mathcal{N}^{\circledast} \to \mathcal{V}^{\circledast}$  that admit  $\mathcal{H}$ -weighted colimits let  $(\mathcal{M} \otimes_{\mathcal{H}}^{\text{LEnr}_{\kappa}} \mathcal{N})^{\circledast}$ the tensor product. By construction there is a canonical equivalence

$$(\mathcal{M} \otimes^{\mathrm{LEnr}_\kappa}_{\mathcal{H}} \mathcal{N})^{\circledast} \simeq \mathcal{P}\mathrm{BEnv}^{\mathcal{H}}_{\otimes_! (\Lambda^{(\mathcal{M}, \mathcal{H})} \boxtimes \Lambda^{(\mathcal{N}, \mathcal{H})})} (\otimes_! (\mathcal{M} \times \mathcal{N}))^{\circledast}_{\mathrm{LEnr}_\kappa}$$

(2) By Theorem 4.42 the  $\infty$ -category  ${}^{\kappa}_{\mathcal{W}} \text{LEnr}(\mathcal{H}')$  carries a left action of  ${}^{\kappa}_{\mathcal{V}} \text{LEnr}(\mathcal{H})$ . For a left  $\kappa$ -enriched  $\infty$ -category  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast}$  that admits  $\mathcal{H}$ -weighted colimits and a left  $\kappa$ -enriched  $\infty$ -category  $\mathcal{N}^{\circledast} \to \mathcal{W}^{\circledast}$  that admits  $\mathcal{H}'$ -weighted colimits let  $(\mathcal{M} \otimes_{\mathcal{H},\mathcal{H}'}^{\text{LEnr}_{\kappa}} \mathcal{N})^{\circledast}$  the left tensor. By

construction there is a canonical equivalence

$$(\mathcal{M} \otimes_{\mathcal{H},\mathcal{H}'}^{\operatorname{LEnr}_{\kappa}} \mathcal{N})^{\circledast} \simeq \mathcal{P}\operatorname{BEnv}_{\mu_{1}(\Lambda^{(\mathcal{M},\mathcal{H})}\boxtimes\Lambda^{(\mathcal{N},\mathcal{H}')})}^{\mathcal{H}'}(\mu_{1}(\mathcal{M} \times \mathcal{N}))_{\operatorname{LEnr}_{\kappa}}^{\circledast}.$$

**Proposition 4.44.** Let  $\mathbb{O}^{\otimes} \to \text{Comm}$  be a symmetric  $\infty$ -operad and  $\kappa$  a small regular cardinal.

(1) Let  $(\mathcal{V}, \mathcal{H}) \to (\mathcal{V}, \mathcal{H}')$  be a map of O-algebras in  $(Mon_{\kappa})^{\otimes}_{*}$  inducing the identity on  $\mathcal{V}$ . The inclusion

$${}_{\mathcal{V}}^{\kappa} \operatorname{LEnr}(\mathcal{H}')^{\otimes} \subset {}_{\mathcal{V}}^{\kappa} \operatorname{LEnr}(\mathcal{H})^{\otimes}$$

admits a left adjoint relative to  $\mathbb{O}^{\otimes}$ .

(2) Let  $(\mathcal{V}, \mathcal{H}) \to (\mathcal{V}, \mathcal{H}')$  be a map of O-algebras in  $(\operatorname{ccMon})^{\otimes}_*$  inducing the identity on  $\mathcal{V}$ . The inclusion

$$_{\mathcal{V}} LQEnr(\mathcal{H}')^{\otimes} \subset _{\mathcal{V}} LQEnr(\mathcal{H})^{\otimes}$$

admits a left adjoint relative to  $\mathcal{O}^{\otimes}$ .

(3) Let  $(\mathcal{V}, \mathcal{H}) \to (\mathcal{V}, \mathcal{H}')$  be a map of O-algebras in  $\operatorname{PrMon}^{\otimes}_*$  inducing the identities on  $\mathcal{V}$ . The inclusion

$$_{\mathcal{V}} \operatorname{LEnr}(\mathcal{H}')^{\otimes} \subset _{\mathcal{V}} \operatorname{LEnr}(\mathcal{H})^{\otimes}$$

admits a left adjoint relative to  $O^{\otimes}$ .

*Proof.* (2) follows from (1) by applying (1) in a larger universe and taking  $\tau = \sigma$  the strongly inaccessible cardinals corresponding to the small universe. (3): The adjunction relative to  $\mathcal{O}^{\otimes}$  of (2) restricts to the adjunction of (3) by construction of the left adjoint and Lemma 3.93.

(1): The map  $(\mathcal{V}, \mathcal{H}) \to (\mathcal{V}, \mathcal{H}')$  of  $\mathcal{O}$ -algebras in  $(Mon_{\kappa})^{\otimes}_{*}$  corresponds to a functor  $\mathcal{O}^{\otimes} \times [1] \to (Mon_{\kappa})^{\otimes}_{*}$  over Comm such that the composition  $\mathcal{O}^{\otimes} \times [1] \to (Mon_{\kappa})^{\otimes}_{*} \to Mon^{\otimes}_{\kappa}$  factors as the projection  $\mathcal{O}^{\otimes} \times [1] \to \mathcal{O}^{\otimes}$  followed by  $\mathcal{V}$ . Let  $\phi : {}^{\kappa}LEnr^{\otimes}_{*} \to (Mon_{\kappa})^{\otimes}_{*}$  be the symmetric monoidal forgetful functor that is a locally cocartesian fibration by Proposition 4.35.

We first prove that every locally  $\phi$ -cocartesian morphism  $\alpha$  of "LEnr<sup>\*</sup><sub> $\alpha$ </sub> lying over an equivalence in Mon<sup> $\otimes$ </sup><sub> $\kappa$ </sub> is  $\phi$ -cocartesian: since  $\alpha$  lies over an equivalence in Mon<sup> $\otimes$ </sup><sub> $\kappa$ </sub>, it belongs to some fiber ("LEnr<sup> $\otimes$ </sup>)<sub>(n)</sub>  $\simeq$  ("LEnr<sub>\*</sub>)<sup>×n</sup> for  $n \ge 0$  and is locally  $\phi_{(n)}$ -cocartesian and thus  $\phi_{(n)}$ -cocartesian because  $\phi_{(n)}$  is a cocartesian fibration by Lemma 4.7. Therefore  $\alpha$  is  $\phi$ -cocartesian if its image  $\otimes(\alpha)$  under the tensor product functor  $\otimes$  : ("LEnr<sub>\*</sub>)<sup>×n</sup>  $\rightarrow$  "LEnr<sub>\*</sub> is  $\phi_{(1)}$ -cocartesian. Since  $\alpha$  is  $\phi_{(n)}$ -cocartesian, by the construction of the tensor product on "LEnr the morphism  $\otimes(\alpha)$  is  $\phi_{(1)}$ -cocartesian if the image of  $\otimes(\alpha)$  under the forgetful functor  $\nu$  : "LEnr<sup> $\otimes$ </sup>  $\Rightarrow$  "LEnr is an equivalence. The functor  $\nu$  is symmetric monoidal and preserves cocartesian morphisms by Lemma 4.7. Thus the image of  $\alpha$  in "LEnr<sup> $\otimes$ </sup> is cocartesian over the equivalence  $\phi(\alpha)$  and so an equivalence. Hence the image of  $\otimes(\alpha)$  in "LEnr is an equivalence.

As every locally  $\phi$ -cocartesian morphism of  $(Mon_{\kappa})^{\otimes}_{*}$  lying over an equivalence in  $Mon^{\otimes}_{\kappa}$  is  $\phi$ cocartesian, the locally cocartesian fibration  $(\mathcal{O}^{\otimes} \times [1]) \times_{(Mon_{\kappa})^{\otimes}_{*}} {}^{\kappa}LEnr_{*}^{\otimes} \to \mathcal{O}^{\otimes} \times [1]$  is a map of
cocartesian fibrations over [1]. By Theorem 4.42 the full subcategory

$$\Xi \coloneqq (\mathbb{O}^{\otimes} \times [1]) \times_{(\mathrm{Mon}_{\kappa})^{\otimes}_{*}} {}^{\kappa} \mathrm{LEnr}^{\otimes}_{\mathrm{w}} \subset (\mathbb{O}^{\otimes} \times [1]) \times_{(\mathrm{Mon}_{\kappa})^{\otimes}_{*}} {}^{\kappa} \mathrm{LEnr_{*}}^{\otimes}$$

is a localization relative to  $\mathbb{O}^{\otimes} \times [1]$ . Therefore also the functor  $\Xi \to \mathbb{O}^{\otimes} \times [1]$  is a locally cocartesian fibration and map of cocartesian fibrations over [1]. Since  $\Xi \to \mathbb{O}^{\otimes} \times [1]$  is a locally cocartesian fibration and [1] has no non-trivial endomorphisms, the functor  $\Xi \to \mathbb{O}^{\otimes}$  is a locally cocartesian fibration and the functor  $\Xi \to \mathbb{O}^{\otimes} \times [1]$  is a map of locally cocartesian fibrations over  $\mathbb{O}^{\otimes}$ .

By Corollary 4.41 the functor  $\Xi \to \mathbb{O}^{\otimes} \times [1]$  induces on the fiber over every  $X \in \mathbb{O}$  a cartesian fibration  $[1] \times_{(Mon_{\kappa})^{\otimes}_{*}} {}^{\kappa} LEnr_{w} \to [1]$  and so  $\Xi \to \mathbb{O}^{\otimes} \times [1]$  is a map of cartesian fibrations over [1] by [11, Lemma 2.44. (2)], where we use that  $\Xi \to \mathbb{O}^{\otimes} \times [1]$  is a map of locally cocartesian fibrations over  $\mathbb{O}^{\otimes}$ . The map  $\Xi \to \mathbb{O}^{\otimes} \times [1]$  of cocartesian and cartesian fibrations over [1] classifies the adjunction of (3).

4.3. Closedness of the monoidal structure. In the following we prove that for every braided monoidal  $\infty$ -category  $\mathcal{V}^{\boxtimes} \to \mathbb{E}_2$ , set  $\mathcal{H}$  of small left weights over  $\mathcal{V}$  and small regular cardinal  $\kappa$  the monoidal structures on  ${}^{\kappa}_{\mathcal{V}} \text{LEnr}(\mathcal{H}), {}_{\mathcal{V}} \text{LEnr}(\mathcal{H})$  of Theorem 4.42 are closed (Corollary 4.61) by giving an explicite construction of the internal hom (Theorem 4.60).

**Notation 4.45.** Let  $\mathcal{V}^{\otimes} \to \operatorname{Ass}, \mathcal{W}^{\otimes} \to \operatorname{Ass}$  be small  $\infty$ -operads,  $\mathcal{H}$  a set of left weights over  $\mathcal{V}$  and  $\mathcal{H}'$  a set of right weights over  $\mathcal{W}$ . Let  $\mathcal{H} \oplus \mathcal{H}'$  be the set of weights over  $\mathcal{V}, \mathcal{W}$  consisting of the images of elements of  $\mathcal{H}$  under the map of  $\infty$ -operads  $\mathcal{V}^{\otimes} \simeq \mathcal{V}^{\otimes} \times \mathcal{Q}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  and the images of elements of  $\mathcal{H}'$  under the map of  $\infty$ -operads  $\mathcal{W}^{\otimes} \simeq \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$ .

For the next notation we use that for every weakly left enriched  $\infty$ -category  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes}$  and  $X \in \mathcal{M}$ there is a unique left enriched functor  $\emptyset^{\otimes} \to \mathcal{M}^{\otimes}$  (starting at the final object of  ${}_{\emptyset}\omega BMod_{\emptyset} \simeq Cat_{\infty}$ ) sending the unique object to X.

Notation 4.46. Let  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast}, \mathcal{O}^{\circledast} \to \mathcal{W}^{\circledast}, \mathcal{N}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  be totally small weakly left enriched, weakly right enriched, weakly bi-enriched  $\infty$ -categories, respectively,  $\Lambda$  a collection of left diagrams in  $\mathcal{M}$  and  $\Lambda'$  a collection of right diagrams in  $\mathcal{O}$ . Let

 $\Lambda \oplus \Lambda'$ 

be the set of diagrams in  $\mathcal{M} \times \mathcal{O}$  consisting of the images of elements of  $\Lambda$  under all enriched functors  $\mathcal{M}^{\circledast} \simeq \mathcal{M}^{\circledast} \times \mathcal{O}^{\circledast} \xrightarrow{\mathcal{M}^{\circledast} \times Y} \mathcal{M}^{\circledast} \times \mathcal{O}^{\circledast}$  for  $Y \in \mathcal{O}$  and the images of elements of  $\Lambda'$  under all enriched functors  $\mathcal{O}^{\circledast} \simeq \mathcal{O}^{\circledast} \times \mathcal{O}^{\circledast} \xrightarrow{X \times \mathcal{O}^{\circledast}} \mathcal{M}^{\circledast} \times \mathcal{O}^{\circledast}$  for  $X \in \mathcal{M}$ .

**Remark 4.47.** Let  $\mathcal{H}$  be a set of left weights over  $\mathcal{V}$  and  $\mathcal{H}'$  a set of right weights over  $\mathcal{W}$  and  $\Lambda$  a collection of  $\mathcal{H}$ -weighted left diagrams in  $\mathcal{M}$  and  $\Lambda'$  a collection of  $\mathcal{H}'$ -weighted right diagrams in  $\mathcal{O}$ . Then  $\Lambda \oplus \Lambda'$  is a collection of  $\mathcal{H} \oplus \mathcal{H}'$ -weighted digrams in  $\mathcal{M} \times \mathcal{O}$ .

**Remark 4.48.** Let  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes}, \mathcal{O}^{\otimes} \to \mathcal{W}^{\otimes}, \mathcal{N}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be totally small weakly left enriched, weakly right enriched, weakly bi-enriched  $\infty$ -categories, respectively,  $\Lambda$  a collection of left diagrams in  $\mathcal{M}$  and  $\Lambda'$  a collection of right diagrams in  $\mathcal{O}$ . The canonical equivalences

 $\operatorname{EnrFun}_{\mathcal{V},\varnothing}(\mathcal{M},\operatorname{EnrFun}_{\varnothing,\mathcal{W}}(\mathcal{O},\mathcal{N})) \simeq \operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}(\mathcal{M}\times\mathcal{O},\mathcal{N}) \simeq \operatorname{EnrFun}_{\varnothing,\mathcal{W}}(\mathcal{O},\operatorname{EnrFun}_{\mathcal{V},\varnothing}(\mathcal{M},\mathcal{N}))$ 

restrict to equivalences

$$\operatorname{EnrFun}_{\mathcal{V},\varnothing}^{\Lambda}(\mathcal{M},\operatorname{EnrFun}_{\varnothing,\mathcal{W}}^{\Lambda'}(\mathcal{O},\mathcal{N})) \simeq \operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}^{\Lambda\oplus\Lambda'}(\mathcal{M}\times\mathcal{O},\mathcal{N}) \simeq \operatorname{EnrFun}_{\varnothing,\mathcal{W}}^{\Lambda'}(\mathcal{O},\operatorname{EnrFun}_{\mathcal{V},\varnothing}^{\Lambda}(\mathcal{M},\mathcal{N})).$$

**Lemma 4.49.** Let  $(\mathcal{V}^{\otimes} \to \operatorname{Ass}, S)$ ,  $(\mathcal{W}^{\otimes} \to \operatorname{Ass}, T)$  be small localization pairs such that S is a set of morphisms of  $\operatorname{Env}(\mathcal{V})$  and T is a set of morphisms of  $\operatorname{Env}(\mathcal{W})$ . Let  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes}$  be a small left S-enriched  $\infty$ -category and  $\mathcal{N}^{\circledast} \to \mathcal{W}^{\otimes}$  a small right T-enriched  $\infty$ -category. Then  $\mathcal{M}^{\circledast} \times \mathcal{N}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  is a S, T-bi-enriched  $\infty$ -category.

*Proof.* We prove that  $\mathcal{M}^{\circledast} \times \mathcal{N}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  is a left S-enriched  $\infty$ -category. The proof of the right T-enrichment is dual. We like to see that for every  $f \in S, X, Y \in \mathcal{M}, X', Y' \in \mathcal{N}, W_1, ..., W_m \in \mathcal{W}$  for  $m \ge 0$  the induced map

 $\mathcal{P}BEnv(\mathcal{M}\times\mathcal{N})(\mathsf{f}\otimes(X,X')\otimes W_1\otimes...\otimes W_m,(Y,Y'))\simeq$ 

 $\mathcal{P}Env(\mathcal{V})(\mathsf{f}, LMulMor_{\mathcal{P}BEnv(\mathcal{M}\times\mathcal{N})}((X, X'), W_1, ..., W_m; (Y, Y')))$ 

is an equivalence. Let  $\delta$  be the diagonal functor  $S \to \mathcal{P}Env(\mathcal{V})$ . There is a canonical equivalence

 $LMulMor_{\mathcal{PBEnv}(\mathcal{M}\times\mathcal{N})}((X,X'),W_1,...,W_m;(Y,Y')) \simeq$ 

$$\operatorname{LMor}_{\mathcal{M}}(\mathbf{X},\mathbf{Y}) \times \delta(\operatorname{Mul}_{\mathcal{N}}(\mathbf{X}',\mathbf{W}_{1},...,\mathbf{W}_{m};\mathbf{Y}'))$$

in  $\mathcal{P}Env(\mathcal{V})$ . Since  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes}$  is a left S-enriched  $\infty$ -category, the following map is an equivalence:

 $\mathcal{P}Env(\mathcal{V})(\mathsf{f}, \mathrm{LMor}_{\mathcal{P}\mathrm{LEnv}(\mathcal{M})}(\mathrm{X}, \mathrm{Y})) \simeq \mathcal{P}\mathrm{LEnv}(\mathcal{M})(\mathsf{f} \otimes \mathrm{X}, \mathrm{Y}).$ 

So it remains to see that the induced map  $\mathcal{P}Env(\mathcal{V})(f, \delta(\operatorname{Mul}_{\mathcal{N}}(X', W_1, ..., W_m; Y'))$  is an equivalence. Because f is representable by assumption, by the Yoneda-lemma the latter map is the equivalence  $\delta(\operatorname{Mul}_{\mathcal{N}}(X', W_1, ..., W_m; Y'))(f)$ .

For the next Notation we use Notation 3.53 and Lemma 4.49:

**Notation 4.50.** Let  $(\mathcal{V}^{\otimes} \to \operatorname{Ass}, \operatorname{S})$ ,  $(\mathcal{W}^{\otimes} \to \operatorname{Ass}, \operatorname{T})$  be small localization pairs such that S is a set of morphisms of  $\operatorname{Env}(\mathcal{V})$  and T is a set of morphisms of  $\operatorname{Env}(\mathcal{W})$ . Let  $\mathcal{H}$  be a set of small left weights over  $\mathcal{V}$  and  $\mathcal{H}'$  a set of small right weights over  $\mathcal{W}$ . Let  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes}$  be a small left S-enriched  $\infty$ -category that admits  $\mathcal{H}$ -weighted colimits and  $\mathcal{O}^{\otimes} \to \mathcal{W}^{\otimes}$  a small right T-enriched  $\infty$ -category that admits  $\mathcal{H}'$ -weighted colimits. Let

$$(\mathfrak{M} \times^{\mathcal{H},\mathcal{H}'}_{S,T} \mathfrak{O})^{\circledast} \coloneqq \mathfrak{PBEnv}^{\Lambda^{(\mathfrak{M} \times \mathfrak{O}, \mathcal{H} \oplus \mathcal{H}')}_{\Lambda^{(\mathfrak{M}, \mathfrak{O})} \oplus \Lambda^{(\mathfrak{O}, \mathcal{H}')}} (\mathfrak{M} \times \mathfrak{O})^{\circledast}_{S,T}.$$

**Theorem 4.51.** Let  $(\mathcal{V}^{\otimes} \to \operatorname{Ass}, \operatorname{S})$ ,  $(\mathcal{W}^{\otimes} \to \operatorname{Ass}, \operatorname{T})$  be small localization pairs such that S is a set of morphisms of  $\operatorname{Env}(\mathcal{V})$  and T is a set of morphisms of  $\operatorname{Env}(\mathcal{W})$ . Let  $\mathcal{H}$  be a set of small left weights over  $\mathcal{V}$  and  $\mathcal{H}'$  a set of small right weights over  $\mathcal{W}$ . Let  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes}$  be a small left S-enriched  $\infty$ -category that admits  $\mathcal{H}$ -weighted colimits and  $\mathcal{O}^{\otimes} \to \mathcal{W}^{\otimes}$  a small right T-enriched  $\infty$ -category that admits  $\mathcal{H}'$ -weighted colimits. There are canonical equivalences

 $\mathrm{EnrFun}_{\mathcal{V},\mathcal{W}}^{\mathcal{H},\mathcal{H}'}(\mathcal{M}\times^{\mathcal{H},\mathcal{H}'}_{\mathrm{S},\mathrm{T}}\mathcal{O},\mathcal{N})\simeq\mathrm{EnrFun}_{\emptyset,\mathcal{W}}^{\mathcal{H}'}(\mathcal{O},\mathrm{EnrFun}_{\mathcal{V},\emptyset}^{\mathcal{H}}(\mathcal{M},\mathcal{N}))\simeq\mathrm{EnrFun}_{\mathcal{V},\emptyset}^{\mathcal{H}}(\mathcal{M},\mathrm{EnrFun}_{\emptyset,\mathcal{W}}^{\mathcal{H}'}(\mathcal{O},\mathcal{N})).$ 

Proof. We prove (1). The proof of (2) is similar. By Remark 4.48 and Notation 3.53 the equivalences

 $\mathrm{EnrFun}_{\mathcal{V},\varnothing}(\mathcal{M},\mathrm{EnrFun}_{\varnothing,\mathcal{W}}(\mathcal{O},\mathcal{N}))\simeq\mathrm{EnrFun}_{\mathcal{V},\mathcal{W}}(\mathcal{M}\times\mathcal{O},\mathcal{N})\simeq\mathrm{EnrFun}_{\varnothing,\mathcal{W}}(\mathcal{O},\mathrm{EnrFun}_{\mathcal{V},\varnothing}(\mathcal{M},\mathcal{N}))$ 

restrict to equivalences

$$\operatorname{EnrFun}_{\mathcal{V},\varnothing}^{\mathcal{H}}(\mathcal{M},\operatorname{EnrFun}_{\varnothing,W}^{\mathcal{H}'}(\mathcal{O},\mathcal{N})) \simeq \operatorname{EnrFun}_{\mathcal{V},W}^{\Lambda^{(\mathcal{M},\mathcal{H})} \oplus \Lambda^{(\mathfrak{O},\mathcal{H}')}}(\mathcal{M} \times \mathcal{O},\mathcal{N})$$
$$\simeq \operatorname{EnrFun}_{\varnothing,W}^{\mathcal{H}'}(\mathcal{O},\operatorname{EnrFun}_{\mathcal{V},\varnothing}^{\mathcal{H}}(\mathcal{M},\mathcal{N})).$$

The  $\mathcal{V}, \mathcal{W}$ -enriched functor  $\mathcal{M}^{\circledast} \times \mathcal{O}^{\circledast} \to (\mathcal{M} \times_{S,T}^{\mathcal{H}, \mathcal{H}'} \mathcal{O})^{\circledast} = \mathcal{P}BEnv_{\Lambda^{(\mathcal{M} \times \mathcal{O}, \mathcal{H} \oplus \mathcal{H}')}}^{\Lambda^{(\mathcal{M} \times \mathcal{O}, \mathcal{H} \oplus \mathcal{H}')}} (\mathcal{M} \times \mathcal{O})_{S,T}^{\circledast}$  induces the following equivalence by Proposition 3.85:

$$\operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}^{\mathcal{H}\oplus\mathcal{H}'}(\mathcal{M}\times_{\mathrm{S},\mathrm{T}}^{\mathcal{H},\mathcal{H}'}\mathcal{O},\mathcal{N})\simeq\operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}^{\Lambda^{(\mathcal{M},\mathcal{H})}\oplus\Lambda^{(\mathcal{O},\mathcal{H}')}}(\mathcal{M}\times\mathcal{O},\mathcal{N}).$$

**Notation 4.52.** Let  $\mathcal{V}^{\otimes} \to Ass, \mathcal{W}^{\otimes} \to Ass$  be small  $\infty$ -operads. Let  $\tau$  be the functor

$$\mathcal{V}^{\otimes} \times_{\mathrm{Ass}} (\mathcal{W}^{\mathrm{rev}})^{\otimes} \simeq \mathcal{V}^{\otimes} \times_{\mathrm{Ass}} \mathcal{W}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$$

induced by projection and the equivalence  $(\mathcal{W}^{rev})^{\otimes} \simeq \mathcal{W}^{\otimes}$ . Taking pullback along  $\tau$  defines a functor

$$\theta: \gamma \omega \operatorname{BEnr}_{\mathcal{W}} \to \gamma_{\times \mathcal{W}^{\operatorname{rev}}} \omega \operatorname{LEnr}_{\emptyset}$$

that restricts to a functor

(19)  $\gamma PBEnr_{\mathcal{W}} \simeq \gamma_{\times \mathcal{W}^{rev}} PLEnr_{\emptyset}.$ 

**Theorem 4.53.** Let  $\mathcal{V}^{\otimes} \to Ass, \mathcal{W}^{\otimes} \to Ass$  be small monoidal  $\infty$ -categories. The functor (19) is an equivalence.

**Notation 4.54.** Let  $\mathcal{V}^{\otimes} \to \operatorname{Ass}, \mathcal{W}^{\otimes} \to \operatorname{Ass}$  be monoidal  $\infty$ -categories,  $\mu : \mathcal{V}^{\otimes} \times_{\operatorname{Ass}} \mathcal{W}^{\otimes} \to \mathcal{W}^{\otimes}$  a monoidal functor and  $\mathcal{N}^{\otimes} \to \mathcal{W}^{\otimes}$  a left pseudo-enriched  $\infty$ -category. By Theorem 4.53 the pullback  $\mu^*(\mathcal{N})^{\otimes} \to \mathcal{V}^{\otimes} \times_{\operatorname{Ass}} \mathcal{W}^{\otimes}$  along  $\mu : \mathcal{V}^{\otimes} \times_{\operatorname{Ass}} \mathcal{W}^{\otimes} \to \mathcal{W}^{\otimes}$  is the pullback of a unique bi-pseudo-enriched  $\infty$ -category  $\mathcal{N}^{\otimes}_{\mu} \to \mathcal{V}^{\otimes} \times (\mathcal{W}^{\operatorname{rev}})^{\otimes}$ . Let  $\widetilde{\mathcal{N}}^{\otimes}_{\mu} \to \mathcal{W}^{\otimes} \times (\mathcal{V}^{\operatorname{rev}})^{\otimes}$  be the corresponding bi-pseudo-enriched  $\infty$ -category via Notation 2.66.

**Notation 4.55.** Let  $\mathcal{V}^{\otimes} \to \operatorname{Ass}, \mathcal{W}^{\otimes} \to \operatorname{Ass}$  be monoidal  $\infty$ -categories and  $\mu : \mathcal{V}^{\otimes} \times_{\operatorname{Ass}} \mathcal{W}^{\otimes} \to \mathcal{W}^{\otimes}$  a monoidal functor. Let  $\mathcal{N}^{\otimes} \to \mathcal{W}^{\otimes}$  be a left pseudo-enriched  $\infty$ -category and  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes}, \mathcal{O}^{\otimes} \to \mathcal{W}^{\otimes}$  weakly left enriched  $\infty$ -categories.

(1) Let

$$\operatorname{EnrFun}_{\mathcal{V}}(\mathcal{M},\mathcal{N})^{\circledast} \to \mathcal{W}^{\otimes}$$

be the weakly left enriched  $\infty$ -category corresponding to the weakly right enriched  $\infty$ -category

 $\operatorname{EnrFun}_{\mathcal{V},\varnothing}(\mathcal{M},\mathcal{N}_{\mu})^{\circledast} \to (\mathcal{W}^{\operatorname{rev}})^{\otimes}.$ 

(2) Let

 $\operatorname{EnrFun}_{\mathcal{W}}(\mathcal{O}, \mathcal{N})^{\otimes} \to \mathcal{V}^{\otimes}$ 

be the weakly left enriched  $\infty$ -category corresponding to the weakly right enriched  $\infty$ -category

EnrFun<sub>$$\mathcal{W}, \emptyset$$</sub>  $(\mathcal{O}, \widetilde{\mathcal{N}}_{\mu})^{\otimes} \to (\mathcal{V}^{\mathrm{rev}})^{\otimes}$ .

Proposition 2.124 has the following consequences:

**Corollary 4.56.** Let  $\kappa$  be a small regular cardinal,  $\mathcal{V}^{\otimes} \to \operatorname{Ass}, \mathcal{W}^{\otimes} \to \operatorname{Ass}$  monoidal  $\infty$ -categories compatible with  $\kappa$ -small colimits and  $\mu : \mathcal{V}^{\otimes} \times_{\operatorname{Ass}} \mathcal{W}^{\otimes} \to \mathcal{W}^{\otimes}$  a monoidal functor preserving  $\kappa$ -small colimits component-wise. Let  $\mathcal{N}^{\otimes} \to \mathcal{W}^{\otimes}$  be a left  $\kappa$ -enriched  $\infty$ -category and  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes}, \mathcal{O}^{\otimes} \to \mathcal{W}^{\otimes}$  weakly left enriched  $\infty$ -categories. The following weakly left enriched  $\infty$ -categories are left  $\kappa$ -enriched:

$$\operatorname{EnrFun}_{\mathcal{V}}(\mathcal{M},\mathcal{N})^{\circledast} \to \mathcal{W}^{\circledast}, \ \operatorname{EnrFun}_{\mathcal{W}}(\mathcal{O},\mathcal{N})^{\circledast} \to \mathcal{V}^{\otimes}.$$

**Corollary 4.57.** Let  $\mathcal{V}^{\otimes} \to \operatorname{Ass}, \mathcal{W}^{\otimes} \to \operatorname{Ass}$  be monoidal  $\infty$ -categories,  $\mu : \mathcal{V}^{\otimes} \times_{\operatorname{Ass}} \mathcal{W}^{\otimes} \to \mathcal{W}^{\otimes}$  a monoidal functor,  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes}, \mathcal{O}^{\otimes} \to \mathcal{W}^{\otimes}$  small weakly left enriched  $\infty$ -categories and  $\mathcal{N}^{\otimes} \to \mathcal{W}^{\otimes}$  a left enriched  $\infty$ -category.

(1) If W admits small limits, the monoidal structure on W is closed and for every  $\mathcal{V} \in \mathcal{V}$  the functor  $\mu(V, -) : W \to W$  admits a right adjoint, the weakly left enriched  $\infty$ -category

 $\operatorname{EnrFun}_{\mathcal{V}}(\mathcal{M},\mathcal{N})^{\otimes} \to \mathcal{W}^{\otimes}$ 

is a left enriched  $\infty$ -category.

(2) If  $\mathcal{V}$  admits small limits, the monoidal structure on  $\mathcal{V}$  is closed and for every  $\mathcal{W} \in \mathcal{W}$  the functor  $\mu(-, \mathcal{W}) : \mathcal{V} \to \mathcal{W}$  admits a right adjoint, the weakly left enriched  $\infty$ -category

$$\operatorname{EnrFun}_{\mathcal{W}}(\mathcal{O},\mathcal{N})^{\circledast} \to \mathcal{V}^{\otimes}$$

is a left enriched  $\infty$ -category.

**Notation 4.58.** Let  $\mathcal{V}^{\otimes} \to \operatorname{Ass}, \mathcal{W}^{\otimes} \to \operatorname{Ass}$  be monoidal  $\infty$ -categories and  $\mu : \mathcal{V}^{\otimes} \times_{\operatorname{Ass}} \mathcal{W}^{\otimes} \to \mathcal{W}^{\otimes}$  a monoidal functor. Let  $\mathcal{N}^{\otimes} \to \mathcal{W}^{\otimes}$  be a left pseudo-enriched  $\infty$ -category and  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes}, \mathcal{O}^{\otimes} \to \mathcal{W}^{\otimes}$  weakly left enriched  $\infty$ -categories,  $\mathcal{H}$  a collection of left weights over  $\mathcal{V}$  and  $\mathcal{H}'$  a collection of left weights over  $\mathcal{W}$ .

(1) Let

$$\operatorname{EnrFun}_{\mathcal{V}}^{\mathcal{H}}(\mathcal{M},\mathcal{N})^{\circledast} \subset \operatorname{EnrFun}_{\mathcal{V}}(\mathcal{M},\mathcal{N})^{\circledast}$$

be the full weakly left enriched subcategory of left  $\mathcal{V}$ -enriched functors  $\mathcal{M} \to \mathcal{N}$  preserving  $\mathcal{H}$ -weighted colimits.

(2) Let

$$\operatorname{EnrFun}_{\mathcal{W}}^{\mathcal{H}'}(\mathcal{O},\mathcal{N})^{\circledast} \subset \operatorname{EnrFun}_{\mathcal{W}}(\mathcal{O},\mathcal{N})^{\circledast}$$

be the full weakly left enriched subcategory of left W-enriched functors  $\mathcal{O} \to \mathcal{N}$  preserving  $\mathcal{H}'$ -weighted colimits.

Proposition 3.94 gives the following corollary:

**Corollary 4.59.** Let  $\mathcal{V}^{\otimes} \to \operatorname{Ass}, \mathcal{W}^{\otimes} \to \operatorname{Ass}$  be monoidal  $\infty$ -categories and  $\mu : \mathcal{V}^{\otimes} \times_{\operatorname{Ass}} \mathcal{W}^{\otimes} \to \mathcal{W}^{\otimes}$  a monoidal functor. Let  $\mathcal{N}^{\otimes} \to \mathcal{W}^{\otimes}$  be a left pseudo-enriched  $\infty$ -category and  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes}, \mathcal{O}^{\otimes} \to \mathcal{W}^{\otimes}$  weakly left enriched  $\infty$ -categories,  $\mathcal{H}$  a collection of left weights over  $\mathcal{V}$  and  $\mathcal{H}'$  a collection of left weights over  $\mathcal{W}$ .

- (1) If N<sup>®</sup> → W<sup>®</sup> admits H'-weighted colimits and µ(1<sub>V</sub>, -): W<sup>®</sup> → W<sup>®</sup> sends weights of H to weights of H', then EnrFun<sub>V,Ø</sub>(M, N)<sup>®</sup> → W<sup>®</sup> admits H'-weighted colimits, EnrFun<sub>V</sub><sup>H</sup>(M, N)<sup>®</sup> ⊂ EnrFun<sub>V</sub>(M, N)<sup>®</sup> is closed under H'-weighted colimits and for every X ∈ M the left W-enriched functor EnrFun<sub>V</sub>(M, N)<sup>®</sup> → N<sup>®</sup> preserves H'-weighted colimits.
- (2) If N<sup>\*</sup> → W<sup>\*</sup> admits H-weighted colimits and µ(-, 1<sub>W</sub>) : V<sup>\*</sup> → W<sup>\*</sup> sends weights of H to weights of H', then EnrFun<sub>W</sub>(O, N)<sup>\*</sup> → V<sup>\*</sup> admits H-weighted colimits, EnrFun<sub>W</sub><sup>H'</sup>(O, N)<sup>\*</sup> ⊂ EnrFun<sub>W</sub>(O, N)<sup>\*</sup> is closed under H-weighted colimits and for every X ∈ O the left V-enriched functor EnrFun<sub>W</sub>(O, N)<sup>\*</sup> → N<sup>\*</sup> preserves H-weighted colimits.

For the next theorem we use Example 4.43:

**Theorem 4.60.** Let  $\kappa$  be a small regular cardinal,  $\mathcal{V}^{\otimes} \to \operatorname{Ass}, \mathcal{W}^{\otimes} \to \operatorname{Ass}$  monoidal  $\infty$ -categories compatible with  $\kappa$ -small colimits and  $\mu : \mathcal{V}^{\otimes} \times_{\operatorname{Ass}} \mathcal{W}^{\otimes} \to \mathcal{W}^{\otimes}$  a monoidal functor preserving  $\kappa$ -small colimits component-wise. Let  $\mathcal{H}, \mathcal{H}'$  be sets of small left weights over  $\mathcal{V}$ , over  $\mathcal{W}$ , respectively, such that  $\mu(-, \mathbb{1}_{\mathcal{W}}) : \mathcal{V}^{\otimes} \to \mathcal{W}^{\otimes}$  sends left weights of  $\mathcal{H}$  to left weights of  $\mathcal{H}'$ . Let  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes}, \mathcal{O}^{\otimes} \to \mathcal{W}^{\otimes}, \mathcal{N}^{\otimes} \to \mathcal{W}^{\otimes}$  be left  $\kappa$ -enriched  $\infty$ -categories that admit  $\mathcal{H}$ -weighted colimits,  $\mathcal{H}'$ -weighted colimits, respectively. There are canonical equivalences

 $\operatorname{EnrFun}_{\mathcal{W}}^{\mathcal{H}'}(\mathcal{M} \otimes_{\mathcal{H} \mathcal{H}'}^{\operatorname{Enr}_{\kappa}} \mathcal{O}, \mathcal{N}) \simeq \operatorname{EnrFun}_{\mathcal{W}}^{\mathcal{H}'}(\mathcal{O}, \operatorname{EnrFun}_{\mathcal{V}}^{\mathcal{H}}(\mathcal{M}, \mathcal{N})) \simeq \operatorname{EnrFun}_{\mathcal{V}}^{\mathcal{H}}(\mathcal{M}, \operatorname{EnrFun}_{\mathcal{W}}^{\mathcal{H}'}(\mathcal{O}, \mathcal{N})).$ 

Proof. By [9, Proposition 5.58.] there are canonical equivalences

 $\operatorname{EnrFun}_{\mathcal{V},\varnothing}(\mathcal{M},\operatorname{EnrFun}_{\mathcal{W}}(\mathcal{O},\mathcal{N})) \simeq \operatorname{EnrFun}_{\mathcal{W},\varnothing}(\mu_{!}(\mathcal{M}\times\mathcal{O}),\mathcal{N}) \simeq \operatorname{EnrFun}_{\mathcal{W},\varnothing}(\mathcal{O},\operatorname{EnrFun}_{\mathcal{V}}(\mathcal{M},\mathcal{N})),$ which restrict to equivalences

$$\operatorname{EnrFun}_{\mathcal{V},\varnothing}^{\mathcal{H}}(\mathcal{M},\operatorname{EnrFun}_{\mathcal{W}}^{\mathcal{H}'}(\mathcal{O},\mathcal{N})) \simeq \operatorname{EnrFun}_{\mathcal{W},\varnothing}^{\mu!(\Lambda^{(\mathcal{M},\mathcal{H})}\boxtimes\Lambda^{(\mathfrak{O},\mathcal{H}')})}(\mu_{!}(\mathcal{M}\times\mathcal{O}),\mathcal{N})$$
$$\simeq \operatorname{EnrFun}_{\mathcal{W},\varnothing}^{\mathcal{H}'}(\mathcal{O},\operatorname{EnrFun}_{\mathcal{V}}^{\mathcal{H}}(\mathcal{M},\mathcal{N})).$$

By construction of the left action as  $(\mathcal{M} \otimes_{\mathcal{H},\mathcal{H}'}^{\mathrm{Enr}_{\kappa}} \mathcal{O})^{\circledast} \simeq \mathcal{P}\mathrm{BEnv}_{\mu_!(\Lambda^{(\mathcal{M},\mathcal{H})} \boxtimes \Lambda^{(\mathcal{O},\mathcal{H}')})}^{\mathcal{H}'}(\mu!(\mathcal{M} \times \mathcal{O}))_{\mathrm{LEnr}_{\kappa}}^{\circledast}$  the left  $\mathcal{W}$ -enriched functor  $\mu_!(\mathcal{M} \times \mathcal{O})^{\circledast} \to (\mathcal{M} \otimes_{\mathcal{H},\mathcal{H}'}^{\mathrm{Enr}_{\kappa}} \mathcal{O})^{\circledast}$  induces an equivalence

$$\operatorname{EnrFun}_{\mathcal{W},\emptyset}^{\mathcal{H}'}(\mathcal{M} \otimes_{\mathcal{H}}^{\operatorname{Enr}_{\kappa}} \mathcal{O}, \mathcal{N}) \simeq \operatorname{EnrFun}_{\mathcal{W},\emptyset}^{\mu_{!}(\Lambda^{(\mathcal{M},\mathcal{H})} \boxtimes \Lambda^{(\mathcal{O},\mathcal{H}')})}(\mu_{!}(\mathcal{M} \times \mathcal{O}), \mathcal{N}).$$

We obtain the following corollary using Corollary 2.124:

**Corollary 4.61.** Let  $1 \leq k \leq \infty$  and  $\kappa$  a small regular cardinal,  $\mathcal{V}^{\boxtimes} \to \mathbb{E}_{k+1}$  an  $\mathbb{E}_{k+1}$ -monoidal  $\infty$ -category compatible with  $\kappa$ -small colimits and  $\mathcal{H}$  a set of small left weights over  $\mathcal{V}$ .

- (1) The  $\mathbb{E}_{k}$ -monoidal structure on  $^{\kappa}_{\mathcal{V}} \text{LEnr}(\mathcal{H})$  of Theorem 4.42 is closed.
- (2) If  $\mathcal{V}^{\boxtimes} \to \mathbb{E}_{k+1}$  is a presentably  $\mathbb{E}_{k+1}$ -monoidal  $\infty$ -category, the  $\mathbb{E}_k$ -monoidal structure on  $_{\mathcal{V}}$ LEnr( $\mathcal{H}$ ) of Theorem 4.42 is closed.

## 5. Applications

5.1. Monoidality of enriched presheaves. In the following we prove that the functor of enriched presheaves is monoidal.

**Notation 5.1.** Let  $\mathcal{O} \to \text{Comm}$  be a symmetric  $\infty$ -operad and  $\mathcal{V} : \mathcal{O}^{\otimes} \to \text{ccMon}^{\otimes}$  an  $\mathcal{O}$ -algebra. Let Lcc be the family that assigns to every  $X \in \mathcal{O}$  the set of totally small left weights over  $\mathcal{V}(X)$ .
**Notation 5.2.** Let  $\operatorname{ccLMod}^{\otimes} \subset (\widehat{\operatorname{LMod}}_{\otimes})^{\times}$  be the symmetric suboperad whose colors are the left tensored  $\infty$ -categories compatible with small colimits and whose multi-morphisms  $\mathcal{M}_{1}^{\otimes}, ..., \mathcal{M}_{n}^{\otimes} \to \mathcal{N}^{\otimes}$  for  $n \geq 0$  correspond to a map  $\mathcal{M}_{1}^{\otimes} \times_{\operatorname{Ass}} ... \times_{\operatorname{Ass}} \mathcal{M}_{n}^{\otimes} \to \mathcal{N}^{\otimes}$  of left tensored  $\infty$ -categories preserving small colimits component-wise.

**Notation 5.3.** Let  $PrLMod^{\otimes} \subset ccLMod^{\otimes}$  be the full symmetric suboperad spanned by the presentably left tensored  $\infty$ -categories.

**Remark 5.4.** There is an identity  $ccLMod^{\otimes} = LQEnr(Lcc)^{\otimes}$ .

**Corollary 5.5.** Let  $\mathcal{V}^{\boxtimes} \to \mathbb{E}_{k+1}$  be an  $\mathbb{E}_{k+1}$ -monoidal  $\infty$ -category compatible with small colimits for  $1 \leq k \leq \infty$ . The functor

$${}_{\mathcal{V}}\mathrm{LQEnr} \to {}_{\mathcal{V}}\mathrm{ccLMod}, (\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes}) \mapsto (\mathcal{P}_{\mathcal{V}}(\mathcal{M})^{\circledast} \to \mathcal{V}^{\otimes})$$

is  $\mathbb{E}_k$ -monoidal and restricts to an  $\mathbb{E}_k$ -monoidal functor  $_{\mathcal{V}} \text{LEnr} \rightarrow _{\mathcal{V}} \text{PrLMod}$ .

Proof. By Corollary 4.44 the inclusion

$$_{\mathcal{V}}\mathrm{ccLMod}^{\otimes} = _{\mathcal{V}}\mathrm{LQEnr}(\mathrm{Lcc})^{\otimes} \subset _{\mathcal{V}}\mathrm{LQEnr}(\varnothing)^{\otimes} =: _{\mathcal{V}}\mathrm{LQEnr}^{\otimes}$$

is a lax  $\mathbb{E}_k$ -monoidal functor that admits a left adjoint relative to  $\mathbb{E}_k$ . The left adjoint sends a small left quasi-enriched  $\infty$ -category  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast}$  to  $\mathcal{P}_{\mathcal{V}}(\mathcal{M})^{\circledast} = \mathcal{P}\widetilde{\mathrm{BEnv}}^{\mathrm{Lcc}}(\mathcal{M})^{\circledast}_{\mathrm{LEnr}}$ .

For the next corollary we fix the following notation:

**Notation 5.6.** Let  $\mathcal{V}^{\otimes} \to \operatorname{Ass}, \mathcal{W}^{\otimes} \to \operatorname{Ass}$  be monoidal  $\infty$ -categories compatible with small colimits. For every weakly left enriched  $\infty$ -categories  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes}, \mathcal{N}^{\otimes} \to \mathcal{W}^{\otimes}$  let

$$(\mathcal{M} \boxtimes \mathcal{N})^{\circledast} \to (\mathcal{V} \otimes \mathcal{W})^{\otimes}$$

be the pushforward of  $\mathcal{M}^{\circledast} \times_{Ass} \mathcal{N}^{\circledast} \to \mathcal{V}^{\otimes} \times_{Ass} \mathcal{W}^{\otimes}$  along the canonical monoidal functor  $\mathcal{V}^{\otimes} \times_{Ass} \mathcal{W}^{\otimes} \to (\mathcal{V} \otimes \mathcal{W})^{\otimes}$  to the tensor product of ccMon.

**Corollary 5.7.** Let  $\mathcal{V}^{\otimes} \to \operatorname{Ass}, \mathcal{W}^{\otimes} \to \operatorname{Ass}$  be monoidal  $\infty$ -categories compatible with small colimits and  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes}, \mathcal{N}^{\otimes} \to \mathcal{W}^{\otimes}$  small left quasi-enriched  $\infty$ -categories. There is a canonical equivalence of  $\infty$ -categories left tensored over  $\mathcal{V} \otimes \mathcal{W}$ :

$$\mathcal{P}_{\mathcal{V}\otimes\mathcal{W}}(\mathcal{M}\boxtimes\mathcal{N})^{\circledast}\simeq(\mathcal{P}_{\mathcal{V}}(\mathcal{M})\otimes\mathcal{P}_{\mathcal{W}}(\mathcal{N}))^{\circledast}.$$

*Proof.* Let  $\mathcal{U}: \mathbb{O}^{\otimes} \to \mathsf{ccMon}^{\otimes}$  be the identity. By Corollary 4.44 and Example 5.1 the lax ccMonmonoidal inclusion

$$\mathsf{ccLMod}^{\otimes} \simeq {}_{\mathcal{U}}\mathsf{ccLMod}^{\otimes} \subset {}_{\mathcal{U}}\mathsf{LQEnr}^{\otimes} \simeq \mathsf{LQEnr}^{\otimes}$$

admits a left adjoint  $\theta$  relative to ccMon<sup> $\otimes$ </sup>. The relative left adjoint  $\theta$  is a map of cocartesian fibrations over ccMon<sup> $\otimes$ </sup>. Hence  $\theta$  induces a commutative square



This square gives the desired equivalence.

5.2. A tensor product for presentable enriched  $\infty$ -categories. Let  $\mathcal{V}^{\otimes} \rightarrow \text{Comm}$  be a presentably symmetric monoidal  $\infty$ -category. By Corollary 4.42 there is a symmetric monoidal structure on  $_{\mathcal{V}}\text{LQEnr}(\text{Lcc}) \simeq _{\mathcal{V}}\text{ccLMod}$ . We will prove the following theorem:

**Theorem 5.8.** Let  $\mathcal{V}^{\boxtimes} \to \operatorname{Comm}$  be a presentably symmetric monoidal  $\infty$ -category and  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast}, \mathcal{N}^{\circledast} \to \mathcal{V}^{\circledast}$  presentably left tensored  $\infty$ -categories. There are small left  $\mathcal{V}$ -enriched  $\infty$ -categories  $\mathbb{C}^{\circledast} \to \mathcal{V}^{\circledast}, \mathcal{D}^{\circledast} \to \mathcal{V}^{\circledast}$ , sets S of morphisms of  $\mathcal{P}_{\mathcal{V}}(\mathbb{C})$  and T of morphisms of  $\mathcal{P}_{\mathcal{V}}(\mathcal{D})$  closed under the left actions and left  $\mathcal{V}$ -enriched equivalences  $\mathcal{M}^{\circledast} \simeq \mathrm{S}^{-1}\mathcal{P}_{\mathcal{V}}(\mathbb{C})^{\circledast}, \mathcal{N}^{\circledast} \simeq \mathrm{T}^{-1}\mathcal{P}_{\mathcal{V}}(\mathcal{D})^{\circledast}$ . The tensor product of  $_{\mathcal{V}}\mathrm{LQEnr}(\mathrm{Lcc}) \simeq _{\mathcal{V}}\mathrm{ccLMod}$  admits the following descriptions:

(1) The relative tensor product:

$$(\mathcal{M} \otimes_{\mathcal{V}} \mathcal{N})^{\circledast} \to \mathcal{V}^{\otimes}.$$

(2) The full subcategory of the internal hom in  $_{\mathcal{V}} \text{LEnr}_{\emptyset}$ :

$$\operatorname{EnrFun}_{\mathcal{V}}^{\mathrm{R}}(\mathcal{M}^{\operatorname{op}},\mathcal{N})^{\circledast} \to \mathcal{V}^{\otimes},$$

(3) The accessible  $\mathcal{V}$ -enriched localization:

$$(\mathbb{S} \boxtimes \mathbb{T})^{-1} \mathcal{P}_{\mathcal{V}}(\mathcal{C} \otimes \mathcal{D})^{\otimes} \to \mathcal{V}^{\otimes}.$$

5.2.1. Presentable enriched  $\infty$ -categories.

**Proposition 5.9.** Let  $\kappa$  be a small regular cardinal,  $\mathcal{V}^{\otimes} \to Ass$  a  $\kappa$ -compactly generated monoidal  $\infty$ -category and  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes}$  a  $\kappa$ -compactly generated  $\infty$ -category left tensored over  $\mathcal{V}$ . The restricted left  $\mathcal{V}$ -enriched Yoneda-embedding admits a  $\mathcal{V}$ -enriched left adjoint and is a left  $\mathcal{V}$ -enriched embedding  $\mathcal{M}^{\otimes} \to \mathcal{P}_{\mathcal{V}}(\mathcal{M}^{\kappa})^{\otimes}$  preserving small  $\kappa$ -filtered conical colimits whose essential image precisely consists of the  $\mathcal{V}$ -enriched presheaves preserving  $\kappa$ -small conical limits and right cotensors with  $\kappa$ -compact objects of  $\mathcal{V}$ .

Proof. The left  $\mathcal{V}$ -enriched embedding  $\mathcal{M}^{\circledast} \to \mathcal{P}_{\mathcal{V}}(\mathcal{M}^{\kappa})^{\circledast}$  is  $\mathcal{V}$ -enriched right adjoint to the left  $\mathcal{V}$ enriched embedding  $(\mathcal{M}^{\kappa})^{\circledast} \subset \mathcal{M}^{\circledast}$  of the full left enriched subcategory of  $\kappa$ -compact objects. Since the embedding  $\mathcal{M}^{\kappa} \subset \mathcal{M}$  preserves  $\kappa$ -compact objects, the right adjoint preserves small  $\kappa$ -filtered colimits.

Let  $\mathcal{N}^{\otimes} \to \mathcal{V}^{\otimes}$  be a presentably left tensored  $\infty$ -category. By Proposition 2.70 the following composition is an equivalence:

$$\operatorname{LinFun}_{\mathcal{V}, \emptyset}^{L}(\mathcal{M}, \mathcal{N}) \to \operatorname{EnrFun}_{\mathcal{V}, \emptyset}^{\prime}(\mathcal{M}^{\kappa}, \mathcal{N}) \to \operatorname{LinFun}_{\mathcal{V}^{\kappa}}^{\kappa}(\mathcal{M}^{\kappa}_{\kappa}, \mathcal{N}_{\kappa}),$$

where  $\operatorname{EnrFun}_{\mathcal{V},\emptyset}(\mathcal{M}^{\kappa},\mathcal{N}) \subset \operatorname{EnrFun}_{\mathcal{V},\emptyset}(\mathcal{M}^{\kappa},\mathcal{N})$  is the full subcategory of left  $\mathcal{V}$ -enriched functors preserving  $\kappa$ -small colimits and left tensors with  $\kappa$ -compact objects of  $\mathcal{V}$ . The second functor in the composition is an equivalence as a consequence of Corollary 2.142 (1). So the first functor in the composition is an equivalence and the statement follows from Proposition 3.85.

**Definition 5.10.** Let  $\kappa$  be a small regular cardinal and  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast}$  a left  $\mathcal{V}$ -enriched  $\infty$ -category. An object X of  $\mathcal{M}$  is  $\mathcal{V}$ -enriched  $\kappa$ -compact if the left  $\mathcal{V}$ -enriched functor  $\operatorname{LMor}_{\mathcal{M}}(X, -) : \mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast}$  preserves small  $\kappa$ -filtered conical colimits.

**Definition 5.11.** Let  $\kappa$  be a small regular cardinal. A left  $\mathcal{V}$ -enriched  $\infty$ -category  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast}$  is  $\mathcal{V}$ -enriched  $\kappa$ -presentable if it admits small weighted colimits and every object of  $\mathcal{M}$  is a small  $\kappa$ -filtered conical colimit of  $\mathcal{V}$ -enriched  $\kappa$ -compact objects of  $\mathcal{M}$ .

**Theorem 5.12.** Let  $\tau \leq \kappa$  be small regular cardinals,  $\mathcal{V}^{\otimes} \to Ass$  a  $\tau$ -compactly generated monoidal  $\infty$ -category and  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes}$  a left  $\mathcal{V}$ -enriched  $\infty$ -category. The following conditions are equivalent:

- (1) The left  $\mathcal{V}$ -enriched  $\infty$ -category  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes}$  is a  $\kappa$ -compactly generated left tensored  $\infty$ -category.
- (2) The left V-enriched ∞-category M<sup>®</sup> → V<sup>®</sup> is a κ-accessible V-enriched localization of P<sub>V</sub>(N)<sup>®</sup> → V<sup>®</sup> for some small left V-enriched ∞-category N<sup>®</sup> → V<sup>®</sup>.

#### (3) The left $\mathcal{V}$ -enriched $\infty$ -category $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes}$ is $\mathcal{V}$ -enriched $\kappa$ -presentable.

*Proof.* Condition (2) implies (1) since  $\mathcal{P}_{\mathcal{V}}(\mathcal{N})^{\circledast} \to \mathcal{V}^{\circledast}$  is a κ-compactly generated left tensored ∞category by Remark 2.153. By Proposition 5.9 condition (1) implies (2). We prove that (1) and (3) are equivalent: in view of Proposition 3.63 and the assumption that the tensor unit of  $\mathcal{V}$  is  $\tau$ -compact and so  $\kappa$ -compact it suffices to see that if  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast}$  is a left tensored ∞-category, the left  $\mathcal{V}$ -action preserves  $\kappa$ -compact objects if and only if for every  $\kappa$ -compact object X of  $\mathcal{M}$  the functor  $\mathrm{LMor}_{\mathcal{M}}(X, -) : \mathcal{M} \to \mathcal{V}$ preserves small  $\kappa$ -filtered colimits. Since  $\mathcal{V}$  is  $\kappa$ -compactly generated, an object X ∈  $\mathcal{M}$  has the property that  $\mathrm{LMor}_{\mathcal{M}}(X, -) : \mathcal{M} \to \mathcal{V}$  preserves small  $\kappa$ -filtered colimits if and only if for every  $\kappa$ -compact object V ∈  $\mathcal{V}$  the left tensor V ⊗ X is  $\kappa$ -compact because  $\mathrm{LMor}_{\mathcal{M}}(V \otimes X, -) \simeq \mathrm{LMor}_{\mathcal{V}}(V, \mathrm{LMor}_{\mathcal{M}}(X, -))$ .

**Lemma 5.13.** Let  $\kappa$  be a small regular cardinal,  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes}, \mathcal{N}^{\circledast} \to \mathcal{W}^{\otimes}$  be  $\tau$ -compactly generated left tensored  $\infty$ -categories such that the tensor product  $(\mathcal{M} \otimes \mathcal{N})^{\circledast} \to \mathcal{V}^{\otimes} \otimes \mathcal{W}^{\otimes}$  in LQEnr(Lcc)  $\simeq$  ccLMod is presentably left tensored. Let S be a set of morphisms of  $\mathcal{M}$  and T a set of morphisms of  $\mathcal{N}$  closed under the left actions with  $\kappa$ -compact objects. Let  $S \boxtimes T$  be the set of morphisms of the form  $f \otimes Y, X \otimes g$  for  $X \in \mathcal{M}^{\kappa}, Y \in \mathcal{N}^{\kappa}, f \in S, g \in T$ . There is a canonical  $\mathcal{V} \otimes \mathcal{W}$ -linear equivalence

$$(S^{-1}\mathcal{M}\otimes T^{-1}\mathcal{N})^{\circledast} \simeq (S\boxtimes T)^{-1}(\mathcal{M}\otimes\mathcal{N})^{\circledast}.$$

*Proof.* Let  $\mathfrak{O} \to \mathcal{V}''^{\otimes}$  be a left tensored  $\infty$ -category compatible with small colimits. The functor

$$\operatorname{EnrFun}(S^{-1}\mathcal{M}\otimes T^{-1}\mathcal{N}, \mathcal{O}) \to \operatorname{EnrFun}(S^{-1}\mathcal{M} \times_{\operatorname{Ass}} T^{-1}\mathcal{N}, \mathcal{O})$$

is fully faithful and the essential image  $\Lambda$  precisely consists of the left enriched functors preserving small weighted colimits component-wise by Proposition 3.85. The functor

(20) 
$$\operatorname{EnrFun}(S^{-1}\mathcal{M} \times_{\operatorname{Ass}} T^{-1}\mathcal{N}, \mathcal{O}) \to \operatorname{EnrFun}(\mathcal{M} \times_{\operatorname{Ass}} \mathcal{N}, \mathcal{O})$$

is fully faithful and the essential image precisely consists of the left enriched functors F such that for every  $X \in \mathcal{M}, Y \in \mathcal{N}$  the left enriched functor  $F(X, -) : \mathcal{N} \to \mathcal{O}$  inverts T and the left enriched functor  $F(-, X) : \mathcal{M} \to \mathcal{O}$  inverts S. The functor (20) restricts to an equivalence between  $\Lambda$  and the full subcategory of EnrFun( $\mathcal{M} \times_{Ass} \mathcal{N}, \mathcal{O}$ ) of left enriched functors F that preserve small weighted colimits component-wise and invert  $S \boxtimes T$ . The functor

$$\operatorname{EnrFun}((\mathbb{S} \boxtimes \mathbb{T})^{-1}(\mathcal{M} \otimes \mathcal{N}), \mathbb{O}) \to \operatorname{EnrFun}(\mathcal{M} \otimes \mathcal{N}, \mathbb{O})$$

is fully faithful and the essential image  $\Theta$  precisely consists of the left enriched functors inverting S $\boxtimes$ T. The functor

(21) 
$$\operatorname{EnrFun}(\mathcal{M} \otimes \mathcal{N}, \mathcal{O}) \to \operatorname{EnrFun}(\mathcal{M} \times_{\operatorname{Ass}} \mathcal{N}, \mathcal{O})$$

is fully faithful and the essential image precisely consists of the left enriched functors preserving small weighted colimits component-wise by Proposition 3.85. The functor (21) restricts to an equivalence between  $\Theta$  and the full subcategory of EnrFun( $\mathcal{M} \times_{Ass} \mathcal{N}, \mathcal{O}$ ) of left enriched functors F preserving small weighted colimits component-wise and inverting  $S \boxtimes T$ .

Corollary 5.7 gives the following corollary:

**Corollary 5.14.** Let  $\kappa$  be a small regular cardinal,  $\mathcal{V}^{\otimes} \to \operatorname{Ass}, \mathcal{W}^{\otimes} \to \operatorname{Ass}$  be  $\kappa$ -compactly generated monoidal  $\infty$ -categories and  $\mathbb{C}^{\otimes} \to \mathcal{V}^{\otimes}, \mathbb{D}^{\otimes} \to \mathcal{W}^{\otimes}$  small left enriched  $\infty$ -categories. Let S be a set of morphisms of  $\mathcal{P}_{\mathcal{V}}(\mathbb{C})$  and T a set of morphisms of  $\mathcal{P}_{\mathcal{W}}(\mathcal{D})$  closed under the left actions with  $\kappa$ -compact objects. Let S  $\boxtimes$  T be the set of morphisms of the form  $f \otimes W \otimes Y, V \otimes X \otimes g$  for  $V \in \mathcal{V}^{\kappa}, W \in \mathcal{W}^{\kappa}, X \in \mathbb{C}, Y \in \mathbb{D}, f \in S, g \in T$ . There is a canonical left  $\mathcal{V} \otimes \mathcal{W}$ -linear equivalence

$$(\mathrm{S}^{-1}\mathcal{P}_{\mathcal{V}}(\mathcal{C})\otimes\mathrm{T}^{-1}\mathcal{P}_{\mathcal{W}}(\mathcal{D}))^{\circledast}\simeq(\mathrm{S}\boxtimes\mathrm{T})^{-1}(\mathcal{P}_{\mathcal{V}\otimes\mathcal{W}}(\mathcal{C}\boxtimes\mathcal{D}))^{\circledast}.$$

5.2.2. The relative tensor product as an enriched functor  $\infty$ -category.

**Lemma 5.15.** Let  $\mathcal{M}^{\otimes} \to \mathcal{U}^{\otimes} \times \mathcal{V}^{\otimes}, \mathcal{N}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be weakly bi-enriched  $\infty$ -categories.

- (1) Then  $\mathcal{M}^{\circledast} \times_{\mathcal{V}^{\otimes}} \mathcal{N}^{\circledast} \to \mathcal{U}^{\otimes} \times \mathcal{W}^{\otimes}$  is a weakly bi-enriched  $\infty$ -category.
- (2) If  $\mathcal{M}^{\circledast} \to \mathcal{U}^{\otimes} \times \mathcal{V}^{\otimes}$  is a left tensored  $\infty$ -category and  $\mathcal{N}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  is a right tensored  $\infty$ -category, then  $\mathcal{M}^{\circledast} \times_{\mathcal{V}^{\otimes}} \mathcal{N}^{\circledast} \to \mathcal{U}^{\otimes} \times \mathcal{W}^{\otimes}$  is a bitensored  $\infty$ -category.
- (3) If M<sup>®</sup> → U<sup>®</sup>×V<sup>®</sup> is a left tensored ∞-category compatible with small colimits and N<sup>®</sup> → V<sup>®</sup>×W<sup>®</sup> is a right tensored ∞-category compatible with small colimits, then M<sup>®</sup>×<sub>V<sup>®</sup></sub> N<sup>®</sup> → U<sup>®</sup>×W<sup>®</sup> is a bitensored ∞-category compatible with small colimits.

*Proof.* Via the enveloping bitensored  $\infty$ -category (1) follows from (2). (2): The functors  $\mathcal{M}^{\circledast} \times \mathcal{W}^{\circledast} \rightarrow \mathcal{U}^{\circledast} \times \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast} \rightarrow \mathcal{U}^{\circledast} \times \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  are maps of cocartesian fibrations relative to the collection of triples, whose first and third component is cocartesian and whose second component is an equivalence. Thus the pullback

$$(\mathcal{M}^{\circledast} \times \mathcal{W}^{\otimes}) \times_{(\mathcal{U}^{\otimes} \times \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes})} (\mathcal{U}^{\otimes} \times \mathcal{N}^{\circledast}) \simeq \mathcal{M}^{\circledast} \times_{\mathcal{V}^{\otimes}} \mathcal{N}^{\circledast} \to \mathcal{U}^{\otimes} \times \mathcal{W}^{\otimes}$$

is a cocartesian fibration relative to the collection of pairs whose first and second component is cocartesian. So it is enough to observe that for every  $U \in \mathcal{U}, W \in \mathcal{W}$  the cocartesian lift  $U \to U'$  in  $\mathcal{U}$  of the morphism  $\{n\} \subset [n]$  in  $\Delta$  and the cocartesian lift  $W \to W'$  in  $\mathcal{W}$  of the morphism  $\{0\} \subset [m]$  in  $\Delta$  induce an equivalence

$$(\mathcal{M}^{\circledast} \times_{\mathcal{V}^{\circledast}} \mathcal{N}^{\circledast})_{U,W} \simeq \mathcal{M}^{\circledast}_{U} \times_{\mathcal{V}^{\circledast}} \mathcal{N}^{\circledast}_{W} \to (\mathcal{M}^{\circledast} \times_{\mathcal{V}^{\circledast}} \mathcal{N}^{\circledast})_{U',W'} \simeq \mathcal{M}^{\circledast}_{U'} \times_{\mathcal{V}^{\circledast}} \mathcal{N}^{\circledast}_{W'}$$

since the functors  $\mathcal{M}_{U}^{\circledast} \to \mathcal{M}_{U'}^{\circledast}$  and  $\mathcal{N}_{W}^{\circledast} \to \mathcal{N}_{W'}^{\circledast}$  are equivalences.

(3): Let  $U \to U'$  be a cocartesian lift in  $\mathcal{U}$  of the morphism  $\{0\} \subset [n]$  in  $\Delta$  and  $W \to W'$  a cocartesian lift in  $\mathcal{W}$  of the morphism  $\{m\} \subset [m]$  in  $\Delta$ . The induced functor

$$(\mathcal{M}^{\circledast} \times_{\mathcal{V}^{\circledast}} \mathcal{N}^{\circledast})_{U,W} \simeq \mathcal{M}_{U}^{\circledast} \times_{\mathcal{V}^{\circledast}} \mathcal{N}_{W}^{\circledast} \rightarrow (\mathcal{M}^{\circledast} \times_{\mathcal{V}^{\circledast}} \mathcal{N}^{\circledast})_{U',W'} \simeq \mathcal{M}_{U'}^{\circledast} \times_{\mathcal{V}^{\circledast}} \mathcal{N}_{W'}^{\circledast}$$

preserves small colimits because the functors  $\mathcal{M}_{U}^{\circledast} \to \mathcal{M}_{U'}^{\circledast}$  and  $\mathcal{N}_{W}^{\circledast} \to \mathcal{N}_{W'}^{\circledast}$  preserve small colimits.

**Notation 5.16.** Let  $\mathcal{M}^{\otimes} \to \mathcal{U}^{\otimes} \times \mathcal{V}^{\otimes}, \mathcal{N}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be bitensored  $\infty$ -categories and  $\sigma : \mathcal{M}^{\otimes} \times_{\mathcal{V}^{\otimes}} \mathcal{N}^{\otimes} \to \mathcal{V}^{\otimes} \to \mathcal{V}^{\otimes} \to \mathcal{N}^{\otimes}$  as the canonical cocartesian fibration. We write

$$(\mathcal{M} \boxtimes_{\mathcal{V}} \mathcal{N})^{\otimes}$$

for the  $\infty$ -category arising from  $\mathcal{M}^{\otimes} \times_{\mathcal{V}^{\otimes}} \mathcal{N}^{\otimes}$  by formally inverting the  $\sigma$ -cocartesian morphisms.

Since the functor  $\mathcal{M}^{\otimes} \times_{\mathcal{V}^{\otimes}} \mathcal{N}^{\otimes} \to \mathcal{U}^{\otimes} \times \mathcal{W}^{\otimes}$  inverts  $\sigma$ -cocartesian morphisms, it induces a functor

$$\tau: (\mathcal{M} \boxtimes_{\mathcal{V}} \mathcal{N})^{\otimes} \to \mathcal{U}^{\otimes} \times \mathcal{W}^{\otimes}$$

**Lemma 5.17.** Let  $\mathcal{M}^{\circledast} \to \mathcal{U}^{\circledast} \times \mathcal{V}^{\circledast}, \mathcal{N}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  be bitensored  $\infty$ -categories. The functor  $\tau : (\mathcal{M} \boxtimes_{\mathcal{V}} \mathcal{N})^{\circledast} \to \mathcal{U}^{\circledast} \times \mathcal{W}^{\circledast}$  is a bitensored  $\infty$ -category and the following canonical functor is  $\mathcal{U}, \mathcal{W}$ -linear:

$$\mathcal{M}^{\circledast} \times_{\mathcal{V}^{\otimes}} \mathcal{N}^{\circledast} \to (\mathcal{M} \boxtimes_{\mathcal{V}} \mathcal{N})^{\circledast}$$

Proof. Since the functor  $\mathcal{M}^{\circledast} \times_{\mathcal{V}^{\otimes}} \mathcal{N}^{\circledast} \to \mathcal{U}^{\circledast} \times \mathcal{W}^{\circledast}$  inverts  $\sigma$ -cocartesian morphisms, every  $\sigma$ -cocartesian morphism belongs to some fiber  $\mathcal{M}^{\circledast}_{[n]} \times_{\mathcal{V}^{\otimes}} \mathcal{N}^{\circledast}_{[m]}$  for some  $[m], [n] \in \Delta$ . Every morphisms  $[n] \to [n'], [m] \to [m']$  in  $\Delta$  induce a left  $\mathcal{V}$ -linear functor  $\mathcal{M}^{\circledast}_{[n']} \times_{\mathcal{V}^{\otimes}} \mathcal{N}^{\circledast}_{[m]} \to \mathcal{M}^{\circledast}_{[m']} \times_{\mathcal{V}^{\otimes}} \mathcal{N}^{\circledast}_{[m]}$  that preserves  $\sigma$ -cocartesian morphisms being left  $\mathcal{V}$ -linear. Let  $\mathcal{T}$  be the collection of  $\sigma$ -cocartesian morphisms. By [13, Proposition 2.1.4.] this implies that  $\tau$  is a map of cocartesian fibrations over Ass  $\times$  Ass, the canonical functor  $\mathcal{M}^{\circledast} \times_{\mathcal{V}^{\otimes}} \mathcal{N}^{\circledast} \to (\mathcal{M} \times_{\mathcal{V}} \mathcal{N})^{\circledast}$  is  $\mathcal{U}, \mathcal{W}$ -linear, and for any  $[n], [m] \in \Delta$  the induced functor

$$(\mathcal{M} \times_{\mathcal{V}} \mathcal{N})^{\circledast}_{[n],[m]} \to (\mathcal{M}^{\circledast}_{[n]} \times_{\mathcal{V}^{\circledast}} \mathcal{N}^{\circledast}_{[m]})[\mathcal{T}] \simeq \mathcal{U}^{\times n} \times (\mathcal{M}^{\circledast}_{[0]} \times_{\mathcal{V}^{\circledast}} \mathcal{N}^{\circledast}_{[0]})[\mathcal{T}] \times \mathcal{W}^{\times n}$$

is an equivalence. Thus  $\tau$  is a bitensored  $\infty$ -category.

 $\Box$ 

**Remark 5.18.** Let  $\mathcal{M}^{\circledast} \to \mathcal{U}^{\otimes} \times \mathcal{V}^{\otimes}, \mathcal{N}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be bitensored  $\infty$ -categories. The cocartesian fibration  $\mathcal{M}^{\circledast}_{[0]} \times_{\mathcal{V}^{\otimes}} \mathcal{N}^{\circledast}_{[0]} \to \mathcal{V}^{\otimes} \to Ass$  classifies the Bar-construction  $\Delta^{\mathrm{op}} \to \mathsf{Cat}_{\infty}, [n] \mapsto \mathcal{M} \times \mathcal{V}^{\times n} \times \mathcal{N}$  of [16, Construction 4.4.2.7.] and  $\mathcal{M} \boxtimes_{\mathcal{V}} \mathcal{N}$  is the colimit of the Bar-construction, the relative tensor product of  $\mathcal{M}^{\circledast}_{[0]} \to \mathcal{V}^{\otimes}$  and  $\mathcal{N}^{\circledast}_{[0]} \to \mathcal{V}^{\otimes}$  [16, Definition 4.4.2.10.].

**Lemma 5.19.** Let  $\mathcal{M}^{\otimes} \to \mathcal{U}^{\otimes} \times \mathcal{V}^{\otimes}, \mathcal{N}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be bitensored  $\infty$ -categories compatible with small colimits. There is a bitensored  $\infty$ -category  $(\mathcal{M} \otimes_{\mathcal{V}} \mathcal{N})^{\otimes} \to \mathcal{U}^{\otimes} \times \mathcal{W}^{\otimes}$  compatible with small colimits and a  $\mathcal{U}, \mathcal{W}$ -linear functor

$$(\mathcal{M} \boxtimes_{\mathcal{V}} \mathcal{N})^{\circledast} \to (\mathcal{M} \otimes_{\mathcal{V}} \mathcal{N})^{\circledast}$$

whose underlying functor preserves small colimits component-wise, that induces for every bitensored  $\infty$ -category  $\mathbb{O}^{\circledast} \to \mathbb{U}^{\otimes} \times \mathbb{W}^{\otimes}$  compatible with small colimits an embedding

$$\operatorname{LinFun}_{\mathcal{U},\mathcal{W}}^{\mathrm{L}}(\mathcal{M} \otimes_{\mathcal{V}} \mathcal{N}, \mathcal{O}) \to \operatorname{LinFun}_{\mathcal{U},\mathcal{W}}(\mathcal{M} \boxtimes_{\mathcal{V}} \mathcal{N}, \mathcal{O}),$$

whose essential image are the U, W-linear functors whose underlying functor preserves small colimits component-wise.

*Proof.* We set  $(\mathcal{M} \otimes_{\mathcal{V}} \mathcal{N})^{\circledast} := \mathcal{P}BEnv_{\Lambda}^{\mathcal{H}} (\mathcal{M} \boxtimes_{\mathcal{V}} \mathcal{N})_{LEnr}^{\circledast} \to \mathcal{U}^{\otimes} \times \mathcal{W}^{\otimes}$ , where  $\mathcal{H}$  is the large collection of small  $(\mathcal{U}, \mathcal{W})$ -weights and  $\Lambda$  is the large collection of all colimit diagrams weighted with respect to weights on  $*_{\mathcal{U},\mathcal{W}}$  together with the following conical diagrams:

$$\begin{split} \mathrm{K}^{\rhd} \to \mathcal{M} \simeq \mathcal{M} \times \{\mathrm{Y}\} \to \mathcal{M} \times \mathcal{N} \subset (\mathcal{M} \times_{\mathcal{V}} \mathcal{N})^{\circledast}, \mathrm{T}^{\rhd} \to \mathcal{N} \simeq \{\mathrm{X}\} \times \mathcal{N} \to \mathcal{M} \times \mathcal{N} \subset \mathcal{M}^{\circledast} \times_{\mathcal{V}^{\otimes}} \mathcal{N}^{\circledast} \to (\mathcal{M} \boxtimes_{\mathcal{V}} \mathcal{N})^{\circledast} \\ \text{for some } \mathrm{X} \in \mathcal{M}, \mathrm{Y} \in \mathcal{N} \text{ and colimit diagrams } \mathrm{K}^{\rhd} \to \mathcal{M}, \mathrm{T}^{\rhd} \to \mathcal{N}. \end{split}$$

**Remark 5.20.** The universal property of Lemma 5.19 implies that the  $\mathcal{U}, \mathcal{W}$ -linear functor  $(\mathcal{M} \boxtimes_{\mathcal{V}} \mathcal{N})^{\circledast} \to (\mathcal{M} \otimes_{\mathcal{V}} \mathcal{N})^{\circledast} \oplus \mathcal{U}^{\circledast} \times \mathcal{W}^{\circledast}$  as the relative tensor product of  $\mathcal{M}^{\circledast} \to \mathcal{U}^{\circledast} \times \mathcal{V}^{\circledast}$  and  $\mathcal{N}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  based on the tensor product on  $\mathsf{Cat}^{cc}_{\infty}$ .

Notation 5.21. Let  $\mathcal{N}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be a weakly bi-enriched  $\infty$ -category,  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes}$  a weakly left enriched  $\infty$ -category and  $\Lambda$  a collection of left diagrams in  $\mathcal{M}^{\mathrm{op}}$ . Let

$$\operatorname{EnrFun}_{\mathcal{V},\emptyset}^{\Lambda-\lim}(\mathcal{M},\mathcal{N})^{\circledast} \subset \operatorname{EnrFun}_{\mathcal{V},\emptyset}(\mathcal{M},\mathcal{N})^{\circledast}$$

be the full weakly right enriched subcategory of left  $\mathcal{V}$ -enriched functors sending diagrams of  $\Lambda$  to  $\mathcal{H}$ -weighted limit diagrams.

**Remark 5.22.** Let  $\kappa$  be a regular cardinal,  $\mathbb{N}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  a  $\kappa$ -compactly generated bitensored  $\infty$ -category,  $\mathbb{M}^{\circledast} \to \mathcal{V}^{\circledast}$  a left enriched  $\infty$ -category and  $\Lambda$  a collection of left diagrams in  $\mathbb{M}^{\mathrm{op}}$  such that  $\mathbb{M}$  is small and  $\mathbb{M}$  admits limits weighted with respect to underlying weights of diagrams of  $\Lambda$ . By [10, Proposition 3.76., Lemma 3.64.] the weakly right enriched  $\infty$ -category  $\mathrm{EnrFun}_{\mathcal{V},\emptyset}(\mathbb{M}^{\mathrm{op}},\mathbb{N})^{\circledast} \to \mathbb{W}^{\circledast}$  is presentably right tensored. The full weakly right enriched subcategory  $\mathrm{EnrFun}_{\mathcal{V},\emptyset}(\mathbb{M}^{\mathrm{op}},\mathbb{N})^{\circledast} \subset \mathrm{EnrFun}_{\mathcal{V},\emptyset}(\mathbb{M}^{\mathrm{op}},\mathbb{N})^{\circledast}$  is the right  $\mathbb{W}$ -linear localization with respect to the set of morphisms

$$\{\theta: \operatorname{colim}^{\mathrm{H}}(\psi_{\mathrm{Z}} \circ \mathrm{F}) \to \psi_{\mathrm{Z}}(\mathrm{Y}) \mid \mathrm{Z} \in \mathcal{N}^{\kappa}, (\mathcal{J}^{\circledast} \to \mathcal{U}^{\otimes} \times \mathcal{Q}^{\otimes}, \mathrm{H} \in \mathcal{P}\mathrm{BEnv}(\mathcal{J})) \in \mathcal{H}, \mathrm{F}: \mathcal{J}^{\circledast} \to \mathcal{M}^{\circledast}\}$$

in EnrFun<sub> $\mathcal{V}, \emptyset$ </sub>( $\mathcal{M}^{\mathrm{op}}, \mathcal{N}$ ), where  $\psi_{\mathbb{Z}}$  is the composition

$$\mathcal{M}^{\circledast} \xrightarrow{\rho} \operatorname{EnrFun}_{\mathcal{V}, \varnothing} (\mathcal{M}^{\operatorname{op}}, \mathcal{V})^{\circledast} \xrightarrow{((-) \otimes \mathbb{Z})_{*}} \operatorname{EnrFun}_{\mathcal{V}, \varnothing} (\mathcal{M}^{\operatorname{op}}, \mathcal{N})^{\circledast},$$

adjoint to the following morphism in  $\mathcal{PBEnv}(\mathcal{J})$ :

$$\mathbf{H} \to \mathbf{F}^*(\mathbf{Y}) \to (\psi_{\mathbf{Z}} \circ \mathbf{F})^*(\psi_{\mathbf{Z}}(\mathbf{Y})) \simeq \mathbf{F}^*(\psi_{\mathbf{Z}}^*(\psi_{\mathbf{Z}}(\mathbf{Y}))).$$

**Proposition 5.23.** Let  $\mathbb{N}^{\otimes} \to \mathbb{V}^{\otimes} \times \mathbb{W}^{\otimes}$  be a presentably bitensored  $\infty$ -category.

Let M<sup>®</sup> → V<sup>®</sup> be a small left enriched ∞-category and Λ a collection of left diagrams in M<sup>op</sup> such that M admits limits weighted with respect to underlying weights of diagrams of Λ. There is a canonical equivalence of ∞-categories presentably right tensored over W:

$$\operatorname{EnrFun}_{\mathcal{V},\mathcal{A}}^{\Lambda-\operatorname{lim}}(\mathcal{M},\mathcal{N})^{\circledast} \simeq (\operatorname{EnrFun}_{\mathcal{V},\mathcal{A}}^{\Lambda-\operatorname{lim}}(\mathcal{M},\mathcal{V}) \otimes_{\mathcal{V}} \mathcal{N})^{\circledast}.$$

(2) Let M<sup>®</sup> → W<sup>®</sup> be a small left enriched ∞-category and Λ a collection of left diagrams in M<sup>op</sup> such that M admits limits weighted with respect to underlying weights of diagrams of Λ. There is a canonical equivalence of ∞-categories presentably left tensored over V:

$$\operatorname{EnrFun}_{\sigma,\mathcal{W}}^{\Lambda-\lim}(\mathcal{M},\mathcal{N})^{\circledast} \simeq (\mathcal{N} \otimes_{\mathcal{W}} \operatorname{EnrFun}_{\sigma,\mathcal{W}}^{\Lambda-\lim}(\mathcal{M},\mathcal{W}))^{\circledast}.$$

*Proof.* We prove (1). Statement (2) is dual. Let  $\alpha : \mathcal{V}^{\circledast} \times_{\mathcal{V}^{\circledast}} \mathcal{N}^{\circledast} \to (\mathcal{V} \otimes_{\mathcal{V}} \mathcal{N})^{\circledast} \simeq \mathcal{N}^{\circledast}$  be the universal  $\mathcal{V}, \mathcal{W}$ -enriched functor. The  $\mathcal{V}, \mathcal{W}$ -enriched evaluation functor  $\mathcal{M}^{\circledast} \times \operatorname{EnrFun}_{\mathcal{V}, \varnothing}(\mathcal{M}, \mathcal{V})^{\circledast} \to \mathcal{V}^{\circledast}$  gives rise to a  $\mathcal{V}, \mathcal{W}$ -enriched functor

$$\mathcal{M}^{\circledast} \times \operatorname{EnrFun}_{\mathcal{V}, \varnothing}(\mathcal{M}, \mathcal{V})^{\circledast} \times_{\mathcal{V}^{\circledast}} \mathcal{N}^{\circledast} \to \mathcal{V}^{\circledast} \times_{\mathcal{V}^{\circledast}} \mathcal{N}^{\circledast} \xrightarrow{\alpha} \mathcal{N}^{\circledast}$$

adjoint to a right W-enriched functor

$$\kappa : \operatorname{EnrFun}_{\mathcal{V}, \emptyset}(\mathcal{M}, \mathcal{V})^{\circledast} \times_{\mathcal{V}^{\circledast}} \mathcal{N}^{\circledast} \to \operatorname{EnrFun}_{\mathcal{V}, \emptyset}(\mathcal{M}, \mathcal{N})^{\circledast}$$

that sends  $F \in EnrFun_{\mathcal{V},\emptyset}(\mathcal{M},\mathcal{V}), Y \in \mathcal{N}$  to  $X \mapsto F(X) \otimes Y$ . The functor  $\kappa$  is right  $\mathcal{W}$ -linear because for every  $F : \mathcal{M} \to \mathcal{V}$  and  $Y \in \mathcal{N}, W \in \mathcal{W}$  the canonical morphism  $\kappa(F, Y) \otimes W \to \kappa(F, Y \otimes W)$  identifies after evaluation at any  $Z \in \mathcal{M}$  with the canonical equivalence

$$(F(Z) \otimes Y) \otimes W \simeq F(Z) \otimes (Y \otimes W).$$

For every  $Y \in \mathbb{N}$  the functor  $\kappa(-, Y) : \operatorname{EnrFun}_{\mathcal{V},\emptyset}(\mathcal{M}, \mathcal{V}) \to \operatorname{EnrFun}_{\mathcal{V},\emptyset}(\mathcal{M}, \mathcal{N}), F \mapsto F(-) \otimes Y$  preserves weighted colimits because the left  $\mathcal{V}$ -linear left adjoint functor  $(-) \otimes Y : \mathcal{V}^{\circledast} \to \mathcal{N}^{\circledast}$  preserves weighted colimits using Proposition 3.94. Since  $\kappa(-, Y) : \operatorname{EnrFun}_{\mathcal{V},\emptyset}(\mathcal{M}, \mathcal{V}) \to \operatorname{EnrFun}_{\mathcal{V},\emptyset}(\mathcal{M}, \mathcal{N})$  preserves weighted colimits, it sends generating local equivalences to local equivalences and so induces a right  $\mathcal{W}$ -linear functor  $\kappa'$  that fits into a commutative square:

Let  $\sigma : \operatorname{EnrFun}_{\mathcal{V},\emptyset}(\mathcal{M},\mathcal{V})^{\circledast} \times_{\mathcal{V}^{\circledast}} \mathcal{N}^{\circledast} \to \mathcal{V}^{\circledast} \to \operatorname{Ass}$  be the canonical functor. We prove next that  $\kappa$  sends  $\sigma$ -cocartesian morphisms to equivalences. This holds if and only if for every  $Z \in \mathcal{M}$  the right  $\mathcal{W}$ -linear functor

$$\operatorname{EnrFun}_{\mathcal{V},\varnothing}(\mathcal{M},\mathcal{V})^{\circledast} \times_{\mathcal{V}^{\circledast}} \mathcal{N}^{\circledast} \xrightarrow{\kappa} \operatorname{EnrFun}_{\mathcal{V},\varnothing}(\mathcal{M},\mathcal{N})^{\circledast} \xrightarrow{\operatorname{ev}_{Z}} \mathcal{N}^{\circledast}$$

inverts  $\sigma$ -cocartesian morphisms. By construction the latter factors as

$$\operatorname{EnrFun}_{\mathcal{V},\varnothing}(\mathcal{M},\mathcal{V})^{\circledast} \times_{\mathcal{V}^{\circledast}} \mathcal{N}^{\circledast} \xrightarrow{\operatorname{evz}} \mathcal{V}^{\circledast} \mathcal{N}^{\circledast} \to \mathcal{V}^{\circledast} \xrightarrow{\alpha} \mathcal{N}^{\circledast}$$

and  $\operatorname{ev}_Z : \operatorname{EnrFun}_{\mathcal{V},\emptyset}(\mathcal{M},\mathcal{V})^{\circledast} \to \mathcal{V}^{\circledast}$  is left  $\mathcal{V}$ -linear and  $\alpha$  universally inverts morphisms cocartesian with respect to  $\mathcal{V}^{\circledast} \times_{\mathcal{V}^{\circledast}} \mathcal{N}^{\circledast} \to \mathcal{V}^{\circledast} \to \operatorname{Ass}$ . Therefore  $\kappa$  inverts  $\sigma$ -cocartesian morphisms. Let  $\sigma'$ :  $\operatorname{EnrFun}_{\mathcal{V},\emptyset}^{\Lambda-\lim}(\mathcal{M},\mathcal{V})^{\circledast} \times_{\mathcal{V}^{\circledast}} \mathcal{N}^{\circledast} \to \mathcal{V}^{\circledast} \to \operatorname{Ass}$  be the canonical functor. Since both vertical functors in square (22) send  $\sigma$ -cocartesian morphisms to  $\sigma'$ -cocartesian morphisms,  $\kappa'$  inverts  $\sigma'$ -cocartesian morphisms. So by Lemma 5.17 the functor  $\kappa'$  induces a right  $\mathcal{W}$ -linear functor

$$\rho: (\mathrm{EnrFun}_{\mathcal{V},\mathcal{O}}^{\Lambda-\mathrm{lim}}(\mathcal{M},\mathcal{V}) \boxtimes_{\mathcal{V}} \mathcal{N})^{\circledast} \to \mathrm{EnrFun}_{\mathcal{V},\mathcal{O}}^{\Lambda-\mathrm{lim}}(\mathcal{M},\mathcal{N})^{\circledast}$$

The right W-linear functor  $\kappa$  induces on underlying  $\infty$ -categories the functor

$$\operatorname{EnrFun}_{\mathcal{V},\varnothing}(\mathcal{M},\mathcal{V})\times\mathcal{N}\to\operatorname{EnrFun}_{\mathcal{V},\varnothing}(\mathcal{M},\mathcal{N}),(F,Y)\mapsto((-)\otimes Y)\circ F$$

that preserves small colimits component-wise. Since both vertical functors in square (22) are left adjoints, the right W-linear functor  $\kappa'$  induces on underlying  $\infty$ -categories a functor preserving small colimits component-wise. So by Lemma 5.19 the functor  $\rho$  induces a right W-linear small colimits preserving functor

$$\lambda: (\mathrm{EnrFun}_{\mathcal{V},\varnothing}^{\Lambda-\mathrm{lim}}(\mathcal{M},\mathcal{V}) \otimes_{\mathcal{V}} \mathcal{N})^{\circledast} \to \mathrm{EnrFun}_{\mathcal{V},\varnothing}^{\Lambda-\mathrm{lim}}(\mathcal{M},\mathcal{N})^{\circledast}$$

that fits into a commutative square of  $\infty$ -categories right tensored over  $\mathcal{W}$ :

The canonical conservative right W-linear functor  $\operatorname{EnrFun}_{\mathcal{V},\emptyset}(\mathcal{M},\mathcal{N})^{\circledast} \to (\mathcal{N}^{\mathcal{M}^{\simeq}})^{\circledast}$  preserves small colimits and admits a W-enriched left adjoint by [9, Proposition 5.23.] and so is monadic by [16, Theorem 4.7.3.5.]. Therefore the right vertical functor in square (23) is monadic, too, where we write F for the left adjoint. By [9, Corollary 4.59.] also the left vertical functor in square (23) is monadic, where the left adjoint is F'  $\otimes_{\mathcal{V}} \mathcal{N}$  when F' is the left adjoint of  $\operatorname{EnrFun}_{\mathcal{V},\emptyset}^{\Lambda-\lim}(\mathcal{M},\mathcal{V}) \to \operatorname{Fun}(\mathcal{M}^{\simeq},\mathcal{V})$ . Therefore by [16, Corollary 4.7.3.16.] it is enough to check that  $\lambda$  preserves the left adjoints, i.e. that the canonical morphism F  $\to \lambda \circ F' \otimes_{\mathcal{V}} \mathcal{N}$  is an equivalence. By Lemma [9, Lemma 5.21.] the  $\infty$ -category Fun( $\mathcal{M}^{\simeq}, \mathcal{N}$ )

is generated under small colimits by the functors of the form  $\mathcal{M}^{\simeq} \xrightarrow{\mathcal{M}^{\simeq}(X,-)} S \xrightarrow{(-)\otimes Y} \mathcal{N}$  for  $X \in \mathcal{M}, Y \in \mathcal{N}$ . Therefore it is enough to check that for every  $X \in \mathcal{M}, Y \in \mathcal{N}$  the canonical morphism

$$\psi: \mathcal{F}(\mathcal{M}^{\simeq}(\mathbf{X}, -) \otimes \mathbf{Y}) \to \kappa'(\mathcal{F}'(\mathcal{M}^{\simeq}(\mathbf{X}, -)), \mathbf{Y})$$

is an equivalence. By [9, Lemma 5.21., Proposition 5.22.] there is a local equivalence  $Mor_{\mathcal{M}}(X, -) \rightarrow F'(\mathcal{M}^{\simeq}(X, -))$  and a commutative square in  $EnrFun_{\mathcal{V}, \emptyset}^{\Lambda-lim}(\mathcal{M}, \mathcal{N})$ :

$$\operatorname{Mor}_{\mathcal{M}}(\mathbf{X},-) \otimes \mathbf{Y} \simeq \kappa(\operatorname{Mor}_{\mathcal{M}}(\mathbf{X},-),\mathbf{Y}) \xrightarrow{} \kappa(\mathbf{F}'(\mathcal{M}^{\simeq}(\mathbf{X},-)),\mathbf{Y}) \xrightarrow{} F(\mathcal{M}^{\simeq}(\mathbf{X},-) \otimes \mathbf{Y}) \xrightarrow{\psi} \kappa'(\mathbf{F}'(\mathcal{M}^{\simeq}(\mathbf{X},-)),\mathbf{Y}),$$

where both vertical morphisms are local equivalences. So  $\psi$  is an equivalence if and only if the top morphism of the square is a local equivalence. The top morphism is a local equivalence since  $\kappa(-, Y)$  preserves local equivalences.

**Proposition 5.24.** Let  $\mathbb{N}^{\circledast} \to \mathbb{V}^{\circledast} \times \mathbb{W}^{\circledast}$  be a presentably bitensored  $\infty$ -category,  $\mathbb{M}^{\circledast} \to \mathbb{W}^{\circledast}$  a small left enriched  $\infty$ -category and  $\Lambda$  a collection of left diagrams in  $\mathbb{M}^{\mathrm{op}}$  such that  $\mathbb{M}$  admits limits weighted with respect to underlying weights of diagrams of  $\Lambda$ . There is a left  $\mathbb{V}$ -linear equivalence:

$$\operatorname{EnrFun}_{\sigma,\mathcal{W}}^{\Lambda-\operatorname{lim}}(\mathcal{M}^{\operatorname{op}},\mathcal{N})^{\circledast} \simeq (\mathcal{N} \otimes_{\mathcal{W}} \mathcal{P}\operatorname{LEnv}_{\Lambda}(\mathcal{M})_{\operatorname{LEnr}})^{\circledast}.$$

*Proof.* By Proposition 5.23 there is a left  $\mathcal{V}$ -linear equivalence:

$$\operatorname{EnrFun}_{\varnothing,W}^{\Lambda-\operatorname{lim}}(\mathcal{M}^{\operatorname{op}},\mathcal{N})^{\circledast} \simeq (\mathcal{N} \otimes_{\mathcal{W}} \operatorname{EnrFun}_{\varnothing,W}^{\Lambda-\operatorname{lim}}(\mathcal{M}^{\operatorname{op}},\mathcal{W}))^{\circledast}.$$

By Corollary 2.142 the left  $\mathcal{W}$ -enriched Yoneda-embedding  $\mathcal{M}^{\circledast} \to \operatorname{EnrFun}_{\varnothing,\mathcal{W}}(\mathcal{M}^{\operatorname{op}},\mathcal{W})^{\circledast}$  uniquely extends to a left  $\mathcal{W}$ -linear equivalence  $\mathcal{P}\operatorname{LEnv}(\mathcal{M})^{\circledast}_{\operatorname{LEnr}} \simeq \operatorname{EnrFun}_{\varnothing,\mathcal{W}}(\mathcal{M}^{\operatorname{op}},\mathcal{W})^{\circledast}$  that restricts to an equivalence  $\mathcal{P}\operatorname{LEnv}_{\Lambda}(\mathcal{M})^{\circledast}_{\operatorname{LEnr}} \simeq \operatorname{EnrFun}_{\varnothing,\mathcal{W}}^{\Lambda-\lim}(\mathcal{M}^{\operatorname{op}},\mathcal{W})^{\circledast}$  by Remark 3.72.

**Corollary 5.25.** Let  $\mathbb{N}^{\otimes} \to \mathbb{V}^{\otimes} \times \mathbb{W}^{\otimes}$  be a presentably bitensored  $\infty$ -category and  $\mathbb{M}^{\otimes} \to \mathbb{W}^{\otimes}$  a small left enriched  $\infty$ -category. There is a left  $\mathbb{V}$ -linear equivalence

$$\operatorname{EnrFun}_{\sigma,\mathcal{W}}(\mathcal{M}^{\operatorname{op}},\mathcal{N})^{\circledast} \simeq (\mathcal{N} \otimes_{\mathcal{W}} \mathcal{P}_{\mathcal{W}}(\mathcal{M}))^{\circledast}$$

**Lemma 5.26.** Let  $\mathbb{N}^{\otimes} \to \mathbb{V}^{\otimes} \times \mathbb{W}^{\otimes}$  be a weakly bi-enriched  $\infty$ -category and  $\mathbb{M}^{\otimes} \to \mathbb{V}^{\otimes}$  a weakly left enriched  $\infty$ -category. There is a canonical left  $\mathbb{W}$ -enriched equivalence:

$$(\operatorname{EnrFun}_{\mathcal{V}, \emptyset}(\mathcal{M}, \mathcal{N})^{\operatorname{op}})^{\mathfrak{S}} \simeq \operatorname{EnrFun}_{\emptyset, \mathcal{V}}(\mathcal{M}^{\operatorname{op}}, \mathcal{N}^{\operatorname{op}})^{\mathfrak{S}}.$$

*Proof.* The desired equivalence is represented by the canonical equivalence

$$\operatorname{EnrFun}_{\mathcal{W},\varnothing}(\mathcal{O}^{\operatorname{op}},\operatorname{EnrFun}_{\mathcal{V},\varnothing}(\mathcal{M},\mathcal{N})^{\operatorname{op}})\simeq\operatorname{EnrFun}_{\varnothing,\mathcal{W}}(\mathcal{O},\operatorname{EnrFun}_{\mathcal{V},\varnothing}(\mathcal{M},\mathcal{N}))$$

 $\simeq \operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}(\mathcal{M} \times \mathcal{O}, \mathcal{N}) \simeq \operatorname{EnrFun}_{\mathcal{W},\mathcal{V}}(\mathcal{O}^{\operatorname{op}} \times \mathcal{M}^{\operatorname{op}}, \mathcal{N}^{\operatorname{op}})$ 

$$\simeq \operatorname{EnrFun}_{\mathcal{W}, \emptyset}(\mathcal{O}^{\operatorname{op}}, \operatorname{EnrFun}_{\emptyset, \mathcal{V}}(\mathcal{M}^{\operatorname{op}}, \mathcal{N}^{\operatorname{op}}))$$

for every weakly right enriched  $\infty$ -category  $\mathcal{O}^{\otimes} \to \mathcal{W}^{\otimes}$ .

**Theorem 5.27.** Let  $\kappa$  be a small regular cardinal,  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast}$  a  $\kappa$ -compactly generated right tensored  $\infty$ -category and  $\mathcal{N}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a presentably bitensored  $\infty$ -category. There is a right  $\mathcal{W}$ -linear equivalence

$$\operatorname{EnrFun}_{\mathcal{V}, \varnothing}^{\mathrm{R}}(\mathcal{M}^{\operatorname{op}}, \mathcal{N})^{\circledast} \simeq (\mathcal{M} \otimes_{\mathcal{V}} \mathcal{N})^{\circledast}.$$

*Proof.* There is a chain of right W-linear equivalences, where  $\kappa$ -colim refers to right  $\mathcal{V}$ -enriched functors preserving  $\kappa$ -small colimits and tensors with  $\kappa$ -compact objects of  $\mathcal{V}$  and  $\kappa$ -lim refers to left  $\mathcal{V}$ -enriched functors preserving  $\kappa$ -small limits and cotensors with  $\kappa$ -compact objects of  $\mathcal{V}$ :

$$\operatorname{EnrFun}_{\mathcal{V},\varnothing}^{R}(\mathcal{M}^{\operatorname{op}},\mathcal{N})^{\circledast} \simeq (\operatorname{LinFun}_{\varnothing,\mathcal{V}}^{L}(\mathcal{M},\mathcal{N}^{\operatorname{op}})^{\operatorname{op}})^{\circledast} \simeq (\operatorname{EnrFun}_{\varnothing,\mathcal{V}}^{\kappa-\operatorname{colim}}(\mathcal{M}^{\kappa},\mathcal{N}^{\operatorname{op}})^{\operatorname{op}})^{\circledast}$$

$$\simeq \operatorname{EnrFun}_{\mathcal{V},\varnothing}^{\kappa-\lim}((\mathcal{M}^{\kappa})^{\operatorname{op}},\mathcal{N})^{\circledast} \simeq (\operatorname{EnrFun}_{\mathcal{V},\varnothing}^{\kappa-\lim}((\mathcal{M}^{\kappa})^{\operatorname{op}},\mathcal{V}) \otimes_{\mathcal{V}} \mathcal{N})^{\circledast} \simeq (\operatorname{Ind}_{\kappa}(\mathcal{M}^{\kappa}) \otimes_{\mathcal{V}} \mathcal{N})^{\circledast}.$$

The first and third equivalence is by Lemma 5.26, the fourth equivalence is by Proposition 5.23, the second and fifth equivalence is by Proposition 5.9.

5.3. Enriched Day-convolution. In the following we equip the  $\infty$ -category of enriched functors between  $\mathbb{E}_k$ -monoidal enriched  $\infty$ -categories with an  $\mathbb{E}_k$ -monoidal structure, which is an enriched version of Day-convolution.

**Definition 5.28.** Let  $1 \leq k \leq \infty$  and  $\mathcal{V}^{\boxtimes} \to \mathbb{E}_{k+1}, \mathcal{W}^{\boxtimes} \to \mathbb{E}_{k+1}$  be presentably  $\mathbb{E}_{k+1}$ -monoidal  $\infty$ -categories.

- (1) Let  $\mathcal{H}$  be a collection of left weights over  $\mathcal{V}$ . An  $\mathbb{E}_k$ -monoidal left  $\mathcal{V}$ -enriched  $\infty$ -category compatible with  $\mathcal{H}$ -weighted colimits is an  $\mathbb{E}_k$ -algebra in  $_{\mathcal{V}}$ LEnr( $\mathcal{H}$ ).
- (2) Let  $\mathcal{H}$  be a collection of right weights over  $\mathcal{W}$ . An  $\mathbb{E}_k$ -monoidal right  $\mathcal{W}$ -enriched  $\infty$ -category compatible with  $\mathcal{H}$ -weighted colimits is an  $\mathbb{E}_k$ -algebra in  $\operatorname{REnr}_{\mathcal{W}}(\mathcal{H})$ .
- (3) Let  $\mathcal{H}$  be a collection of weights over  $\mathcal{W}, \mathcal{W}$ . An  $\mathbb{E}_k$ -monoidal  $\mathcal{V}, \mathcal{W}$ -bi-enriched  $\infty$ -category compatible with  $\mathcal{H}$ -weighted colimits is an  $\mathbb{E}_k$ -algebra in  $_{\mathcal{V}} BEnr_{\mathcal{W}}(\mathcal{H})$ .

We refer to  $\mathbb{E}_1$ -monoidal as monoidal. We refer to Comm-monoidal as symmetric monoidal.

**Example 5.29.** An  $\mathbb{E}_k$ -monoidal  $\mathcal{V}, \mathcal{W}$ -bi-enriched  $\infty$ -category compatible with small weighted colimits is an  $\mathbb{E}_k$ -algebra with respect to the relative tensor product on  $_{\mathcal{V}}$ ccBMod<sub> $\mathcal{W}$ </sub>.

**Corollary 5.30.** Let  $\mathcal{V}^{\boxtimes} \to \mathbb{E}_{k+1}, \mathcal{W}^{\boxtimes} \to \mathbb{E}_{k+1}$  be presentably  $\mathbb{E}_{k+1}$ -monoidal  $\infty$ -categories for  $1 \le k \le \infty$ and  $\mathcal{N}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  an  $\mathbb{E}_k$ -monoidal  $\mathcal{V}, \mathcal{W}$ -bi-enriched  $\infty$ -category compatible with small weighted colimits such that  $\mathcal{N}$  is presentable. Let  $\mathcal{H}$  be a collection of left weights over  $\mathcal{W}$  and  $\mathcal{M}^{\circledast} \to \mathcal{W}^{\otimes}$  an  $\mathbb{E}_k$ -monoidal left  $\mathcal{W}$ -enriched  $\infty$ -category compatible with  $\mathcal{H}$ -weighted colimits. Then

$$\operatorname{EnrFun}_{\emptyset \mathcal{W}}^{\Lambda^{\mathcal{H}}-\lim}(\mathcal{M}^{\operatorname{op}},\mathcal{N})^{\circledast} \to \mathcal{V}^{\circledast}$$

refines to an  $\mathbb{E}_k$ -monoidal left  $\mathcal{V}$ -enriched  $\infty$ -category compatible with small weighted colimits.

Proof. The functor  $\mathcal{P}\text{LEnv}_{\Lambda^{\mathfrak{H}}}(-)_{\text{LEnr}} : {}_{\mathcal{V}}\text{LEnr}_{\varnothing}(\mathcal{H}) \to {}_{\mathcal{V}}\text{ccLMod}_{\varnothing}$  is  $\mathbb{E}_{k}$ -monoidal by Theorem 4.42 and so equips  $\mathcal{P}\text{LEnv}_{\Lambda^{\mathfrak{H}}}(\mathcal{M})^{\mathfrak{B}}_{\text{LEnr}} \to \mathcal{W}^{\mathfrak{B}}$  with an  $\mathbb{E}_{k}$ -algebra structure in  ${}_{\mathcal{V}}\text{ccLMod}_{\varnothing}$ . Since  $\mathcal{V}^{\mathfrak{B}} \to \mathbb{E}_{k+1}, \mathcal{W}^{\mathfrak{B}} \to \mathbb{E}_{k+1}$  are presentably  $\mathbb{E}_{k+1}$ -monoidal  $\infty$ -categories, the functor  ${}_{\mathcal{V}}\text{ccLMod}_{\mathcal{W}} \times {}_{\mathcal{W}}\text{ccLMod}_{\varnothing} \to {}_{\mathcal{V}}\text{ccBMod}_{\varnothing}$  is  $\mathbb{E}_{k}$ -monoidal so that  $\text{EnrFun}_{\mathscr{O},\mathcal{W}}^{\Lambda^{\mathfrak{H}}}(\mathcal{M}^{\mathrm{op}}, \mathcal{N})^{\mathfrak{B}} \simeq (\mathcal{N} \otimes_{\mathcal{W}} \mathcal{P}\text{LEnv}_{\Lambda^{\mathfrak{H}}}(\mathcal{M})_{\text{LEnr}})^{\mathfrak{B}}$  carries the structure of an  $\mathbb{E}_{k}$ -algebra in  ${}_{\mathcal{V}}\text{ccLMod}_{\varnothing}$ , where the latter equivalence is by Theorem 5.24.

**Corollary 5.31.** Let  $\mathcal{V}^{\boxtimes} \to \mathbb{E}_{k+1}, \mathcal{W}^{\boxtimes} \to \mathbb{E}_{k+1}$  be presentably  $\mathbb{E}_{k+1}$ -monoidal  $\infty$ -categories for  $1 \leq k \leq \infty$ and  $\mathcal{N}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  an  $\mathbb{E}_k$ -monoidal  $\mathcal{V}, \mathcal{W}$ -bi-enriched  $\infty$ -category compatible with small weighted colimits such that  $\mathcal{N}$  is presentable. Let  $\mathcal{M}^{\circledast} \to \mathcal{W}^{\otimes}$  be an  $\mathbb{E}_k$ -monoidal left  $\mathcal{W}$ -enriched  $\infty$ -category compatible with  $\mathcal{H}$ -weighted colimits. Then

## $\operatorname{EnrFun}_{\varnothing, \mathcal{W}}(\mathcal{M}^{\operatorname{op}}, \mathcal{N})^{\circledast} \to \mathcal{V}^{\otimes}$

refines to an  $\mathbb{E}_k$ -monoidal left  $\mathcal{V}$ -enriched  $\infty$ -category compatible with small weighted colimits.

Remark 5.32. Let the assumptions like in Corollary 5.31. The tensor product

 $\operatorname{EnrFun}_{\varnothing, W}(\mathcal{M}^{\operatorname{op}}, \mathcal{N}) \times \operatorname{EnrFun}_{\varnothing, W}(\mathcal{M}^{\operatorname{op}}, \mathcal{N}) \to \operatorname{EnrFun}_{\varnothing, W}(\mathcal{M}^{\operatorname{op}}, \mathcal{N})$ 

of the left  $\mathcal{V}$ -enriched Day-convolution factors as

 $\operatorname{EnrFun}_{\varnothing,\mathcal{W}}(\mathcal{M}^{\operatorname{op}},\mathcal{N}) \times \operatorname{EnrFun}_{\varnothing,\mathcal{W}}(\mathcal{M}^{\operatorname{op}},\mathcal{N}) \to \operatorname{EnrFun}_{\varnothing,\mathcal{W}}(\mathcal{M}^{\operatorname{op}},\mathcal{N}) \otimes \operatorname{EnrFun}_{\varnothing,\mathcal{W}}(\mathcal{M}^{\operatorname{op}},\mathcal{N})$ 

 $\simeq \mathcal{N} \otimes_{\mathcal{W}} \mathcal{P}LEnv_{\Lambda^{\mathcal{H}}}(\mathcal{M})_{LEnr} \otimes \mathcal{N} \otimes_{\mathcal{W}} \mathcal{P}LEnv(\mathcal{M})_{LEnr} \\ \simeq (\mathcal{N} \otimes \mathcal{N}) \otimes_{\mathcal{W} \otimes \mathcal{W}} (\mathcal{P}LEnv(\mathcal{M})_{LEnr} \otimes \mathcal{P}LEnv(\mathcal{M})_{LEnr})$ 

 $\simeq (\mathcal{N} \otimes \mathcal{N}) \otimes_{\mathcal{W} \otimes \mathcal{W}} \mathcal{P} LEnv(\mathcal{M} \otimes \mathcal{M})_{LEnr}$ 

 $\rightarrow \mathcal{N} \otimes_{\mathcal{W}} \mathcal{P}LEnv(\mathcal{M})_{LEnr} \simeq EnrFun_{\emptyset,\mathcal{W}}(\mathcal{M}^{\mathrm{op}},\mathcal{N}),$ 

which identifies with the functor

$$\operatorname{EnrFun}_{\varnothing,\mathcal{W}}(\mathcal{M}^{\operatorname{op}},\mathcal{N}) \times \operatorname{EnrFun}_{\varnothing,\mathcal{W}}(\mathcal{M}^{\operatorname{op}},\mathcal{N}) \to \operatorname{EnrFun}_{\varnothing,\mathcal{W}}(\mathcal{M}^{\operatorname{op}}\otimes\mathcal{M}^{\operatorname{op}},\mathcal{N}\otimes\mathcal{N})$$

$$\xrightarrow{(\otimes_{\mathcal{N}})_*} \operatorname{EnrFun}_{\sigma,\mathcal{W}}(\mathcal{M}^{\operatorname{op}} \otimes \mathcal{M}^{\operatorname{op}}, \mathcal{N}) \xrightarrow{(\otimes_{\mathcal{M}})_!} \operatorname{EnrFun}_{\sigma,\mathcal{W}}(\mathcal{M}^{\operatorname{op}}, \mathcal{N}).$$

5.4. A tensor product for  $(\infty, 2)$ -categories with lax colimits. In the following we apply our theory to  $(\infty, 2)$ -categories, which we define as  $Cat_{\infty}$ -enriched  $\infty$ -categories, to define lax and oplax colimits and deduce the existence of a monoidal structure on the  $\infty$ -category of small  $\infty$ -categories that admit  $\mathcal{K}$ -indexed (op)lax colimits for any set  $\mathcal{K}$  of  $(\infty, 2)$ -categories.

**Definition 5.33.** Let  $n \ge 1$ . Let  $Cat_{(\infty,1)} \coloneqq Cat_{(\infty,n+1)} \coloneqq Cat_{(\infty,n+1)} \coloneqq Cat_{(\infty,n)} LEnr.$ 

**Construction 5.34.** By [20, Theorem 6.20.] for every small  $(\infty, 2)$ -category  $\mathcal{C}$  there is an  $(\infty, 2)$ -categorical Grothendieck construction, which can be viewed as a functor of  $(\infty, 2)$ -categories

$$\theta: \operatorname{EnrFun}_{\mathsf{Cat}_{\infty}, \varnothing}(\mathcal{C}^{\operatorname{op}}, \mathsf{Cat}_{\infty})^{\circledast} \to \mathsf{Cat}_{(\infty, 2)}^{\circledast}$$

that sends the final object to  $\mathcal{C}$  and sends a functor of  $(\infty, 2)$ -categories  $F : \mathcal{C}^{op} \to \mathsf{Cat}_{\infty}$  to a functor of  $(\infty, 2)$ -categories  $\mathcal{B} \to \mathcal{C}$  whose fiber over  $X \in \mathcal{C}$  is F(X). We obtain a functor of  $(\infty, 2)$ -categories:

$$\kappa^{\mathbb{C}}: \mathbb{C}^{\circledast} \to \operatorname{EnrFun}_{\mathsf{Cat}_{\infty}}(\mathbb{C}^{\operatorname{op}}, \mathsf{Cat}_{\infty})^{\circledast} \to \mathsf{Cat}_{(\infty, 2)}^{\circledast} \to \mathsf{Cat}_{\infty}^{\circledast}$$

where the first functor is the  $Cat_{\infty}$ -enriched Yoneda-embedding and the second functor is the  $Cat_{\infty}$ enriched right adjoint  $\tau$  of the embedding  $Cat_{\infty} \subset Cat_{(\infty,2)}$  that discards all non-invertible 2-morphisms. The functor  $\kappa^{\mathbb{C}}$  sends  $X \in \mathbb{C}$  to a functor  $\tau(\theta(\mathbb{C})) \to \tau(\mathbb{C})$  whose fiber over any  $Y \in \mathbb{C}$  is  $Mor_{\mathbb{C}}(Y, X)$ .

**Definition 5.35.** Let  $\mathcal{C} \in \mathsf{Cat}_{(\infty,2)}$ .

• The lax weight on  $\mathcal{C}$  is the functor of  $(\infty, 2)$ -categories

$$\kappa^{\mathfrak{C}}: \mathfrak{C} \to \mathsf{Cat}_{\infty}$$

• The oplax weight on  $\mathcal{C}$  is the functor of  $(\infty, 2)$ -categories

$$\lambda^{\mathcal{C}} := (-)^{\mathrm{op}} \circ \kappa^{\mathcal{C}} : \mathcal{C} \to \mathsf{Cat}_{\infty}$$

**Definition 5.36.** Let  $F : \mathcal{C} \to \mathcal{D}$  be a functor of  $(\infty, 2)$ -categories.

- (1) The lax colimit of F, denoted by  $\operatorname{colim}^{\operatorname{lax}}(F)$ , is the colimit of F weighted at the lax weight on  $\mathcal{C}^{\operatorname{op}}$ .
- (2) The oplax colimit of F, denoted by  $\operatorname{colim}^{\operatorname{oplax}}(F)$ , is the colimit of F weighted at the oplax weight on  $\mathcal{C}^{\operatorname{op}}$ .
- (3) The lax limit of F, denoted by lim<sup>lax</sup>(F), is the limit of F weighted at the lax weight on C, i.e. the lax colimit of F<sup>op</sup>.
- (4) The oplax limit of F, denoted by  $\lim^{oplax}(F)$ , is the limit of F weighted at the oplax weight on C, i.e. the oplax colimit of  $F^{op}$ .

**Remark 5.37.** In [7, Definition 2.9.] Gepner-Haugseng define the lax colimit and lax limit of a functor  $\mathcal{C}^{\text{op}} \to \mathsf{Cat}_{\infty}$  under the assumption that  $\mathcal{C}$  is an  $(\infty, 1)$ -category.

**Notation 5.38.** Let  $\mathcal{K} \subset Cat_{(\infty,2)}$  be a full subcategory. Let

$$\operatorname{Cat}_{(\infty,2)}^{\operatorname{laxcolim}}(\mathcal{K}) \coloneqq_{\operatorname{Cat}_{\infty}} \operatorname{LEnr}(\{\kappa^{\mathcal{C}} \mid \mathcal{C} \in \operatorname{Cat}_{(\infty,2)}\}) \subset \operatorname{Cat}_{(\infty,2)} = \operatorname{Cat}_{\infty} \operatorname{LEnr},$$

$$\operatorname{Cat}_{(\infty,2)}^{\operatorname{oplaxcolim}}(\mathcal{K}) \coloneqq \operatorname{Cat}_{\infty} \operatorname{LEnr}(\{\lambda^{\mathcal{C}} \mid \mathcal{C} \in \operatorname{Cat}_{(\infty,2)}\}) \subset \operatorname{Cat}_{(\infty,2)} = \operatorname{Cat}_{\infty} \operatorname{LEnr}$$

be the subcategories of small  $(\infty, 2)$ -categories that admit  $\mathcal{K}$ -indexed (op)lax colimits and functors preserving  $\mathcal{K}$ -indexed (op)lax colimits.

Theorem 4.42 specializes to the following:

**Corollary 5.39.** Let  $\mathcal{K} \subset \mathsf{Cat}_{(\infty,2)}$  be a full subcategory. The  $\infty$ -categories

(24) 
$$\operatorname{Cat}_{(\infty,2)}^{\operatorname{laxcolim}}(\mathcal{K}) \subset \operatorname{Cat}_{(\infty,2)}, \ \operatorname{Cat}_{(\infty,2)}^{\operatorname{oplaxcolim}}(\mathcal{K}) \subset \operatorname{Cat}_{(\infty,2)}$$

carry closed symmetric monoidal structures such that the inclusions (24) are lax symmetric monoidal, where the right hand sides carry the cartesian structure.

**Example 5.40.** The functor  $\kappa^{\{0,1\}^{\triangleright}} : \{0,1\}^{\triangleright} \to \mathsf{Cat}_{\infty}$  corresponds to the diagram  $\{0\} \to \{0,1\}^{\triangleright} \leftarrow \{1\}$ . For every  $(\infty, 2)$ -category  $\mathcal{D}$  and functor  $F : \{0,1\}^{\triangleleft} \to \mathcal{D}$  corresponding to a diagram  $X \leftarrow Z \to Y$  in  $\mathcal{D}$  and every  $T \in \mathcal{D}$  there is a canonical equivalence:

$$\mathcal{D}(\operatorname{colim}^{\operatorname{lax}}(F), T) \simeq \operatorname{Fun}((\{0, 1\}^{\triangleleft})^{\operatorname{op}}, \operatorname{Cat}_{\infty})(\kappa^{\{0, 1\}^{\vee}}, \operatorname{Mor}_{\mathcal{D}}(F(-), T)) \simeq$$

 $\mathcal{D}(X,T) \times_{\mathcal{D}(Z,T)^{\{1\}}} \mathsf{Cat}_{\infty}([1],\mathcal{D}(Z,T)) \times_{\mathcal{D}(Z,T)^{\{0\}}} \mathsf{Cat}_{\infty}([1],\mathcal{D}(Z,T)) \times_{\mathcal{D}(Z,T)^{\{1\}}} \mathcal{D}(Y,T).$ 

So if  $\mathcal{D}$  admits pushouts and tensors with [1], the latter equivalence represents an equivalence

$$\operatorname{colim}^{\operatorname{lax}}(\mathbf{F}) \simeq \mathbf{X} \coprod_{\{1\} \otimes \mathbf{Z}} ([1] \otimes \mathbf{Z}) \coprod_{\{0\} \otimes \mathbf{Z}} ([1] \otimes \mathbf{Z}) \coprod_{\{1\} \otimes \mathbf{Z}} \mathbf{Y}$$

Dually, if  $\mathcal{D}$  admits pullbacks and cotensors with [1], the lax limit of a functor  $G : \{0,1\}^{\triangleright} \to \mathcal{D}$  corresponding to a diagram  $X \to Z \leftarrow Y$  in  $\mathcal{D}$  is  $X \prod_{Z^{\{1\}}} Z^{[1]} \prod_{Z^{\{0\}}} Z^{[1]} \prod_{Z^{\{1\}}} Y$ .

**Definition 5.41.** Let  $\sigma : \{0,1\}^{\triangleright} \to \mathsf{Cat}_{\infty}$  be the functor corresponding to the diagram  $\{0\} \to [1] \leftarrow \{1\}$  and  $\mathcal{D}$  an  $(\infty, 2)$ -category.

- Let  $F : \{0,1\}^{\triangleleft} \to \mathcal{D}$  be a functor corresponding to a diagram  $X \leftarrow Z \to Y$  in  $\mathcal{D}$ . The lax pushout of the diagram  $X \leftarrow Z \to Y$  in  $\mathcal{D}$ , denoted by  $X \coprod_{Z}^{\text{lax}} Y$ , is the  $\sigma$ -weighted colimit of F.
- Let  $F : \{0,1\}^{\triangleleft} \to \mathcal{D}$  be a functor corresponding to a diagram  $X \leftarrow Z \to Y$  in  $\mathcal{D}$ . The oplax pushout of the diagram  $X \leftarrow Z \to Y$  in  $\mathcal{D}$ , denoted by  $X \coprod_{Z}^{\text{oplax}} Y$ , is the  $(-)^{\text{op}} \circ \sigma$ -weighted colimit of F.
- Let  $G : \{0,1\}^{\triangleright} \to \mathcal{D}$  be a functor corresponding to a diagram  $X \to Z \leftarrow Y$  in  $\mathcal{D}$ . The lax pullback of the diagram  $X \to Z \leftarrow Y$  in  $\mathcal{D}$ , denoted by  $X \times_Z^{\text{lax}} Y$ , is the  $\sigma$ -weighted limit of G.
- Let  $G : \{0,1\}^{\triangleright} \to \mathcal{D}$  be a functor corresponding to a diagram  $X \to Z \leftarrow Y$  in  $\mathcal{D}$ . The oplax pullback of the diagram  $X \to Z \leftarrow Y$  in  $\mathcal{D}$ , denoted by  $X \times_Z^{\text{oplax}} Y$ , is the  $(-)^{\text{op}} \circ \sigma$ -weighted limit of G.

**Remark 5.42.** Let  $\mathcal{B}$  be an  $\infty$ -category and  $\gamma, \gamma' : \{0, 1\}^{\triangleright} \to \mathcal{B}$  be functors corresponding to diagrams  $A \to C \leftarrow B, A' \to C' \leftarrow B'$ , respectively. There is a canonical equivalence

$$\operatorname{Fun}(\{0,1\}^{\triangleright}, \mathcal{B})(\gamma, \gamma') \simeq \mathcal{B}(A, A') \times_{\mathcal{B}(A, C')} \mathcal{B}(C, C') \times_{\mathcal{B}(B, C')} \mathcal{B}(B, B').$$

**Example 5.43.** Let  $F : \{0,1\}^{\triangleleft} \rightarrow \mathcal{D}$  be a functor corresponding to a diagram  $X \leftarrow Z \rightarrow Y$  in  $\mathcal{D}$ . By Remark 5.42 there is a canonical equivalence for every  $T \in \mathcal{D}$ :

$$\mathcal{D}(X\coprod_{Z}^{\max}Y,T)\simeq \operatorname{Fun}(\{0,1\}^{\triangleright},\mathsf{Cat}_{\infty})(\sigma,\operatorname{Mor}_{\mathcal{D}}(F(-),T))\simeq$$

 $\mathsf{Cat}_{\infty}([0], \mathrm{Mor}_{\mathcal{D}}(X, T)) \times_{\mathsf{Cat}_{\infty}([0], \mathrm{Mor}_{\mathcal{D}}(Z, T))} \mathsf{Cat}_{\infty}([1], \mathrm{Mor}_{\mathcal{D}}(Z, T))$ 

$$\times_{\mathsf{Cat}_{\infty}([0], \operatorname{Mor}_{\mathcal{D}}(Z, T))} \mathsf{Cat}_{\infty}([0], \operatorname{Mor}_{\mathcal{D}}(Y, T)) \simeq$$

 $\mathcal{D}(X,T) \times_{\mathcal{D}(Z,T)} \mathsf{Cat}_{\infty}([1], \operatorname{Mor}_{\mathcal{D}}(Z,T)) \times_{\mathcal{D}(Z,T)} \mathcal{D}(Y,T).$ 

So if  $\mathcal{D}$  admits pushouts and tensors with [1], we find that

1.....

$$X \coprod_{Z}^{lax} Y \simeq X \coprod_{\{0\} \times Z} ([1] \otimes Z) \coprod_{\{1\} \times Z} Y.$$

Similarly, if  $\mathcal{D}$  admits pushouts and tensors with [1], we find that

$$X \coprod_{Z}^{\text{optax}} Y \simeq X \coprod_{\{1\} \times Z} ([1] \otimes Z) \coprod_{\{0\} \times Z} Y.$$

Dually, if  $\mathcal{D}$  admits pullbacks and cotensors with [1], the lax pullback of the diagram  $X \to Z \leftarrow Y$  in  $\mathcal{D}$  is  $X \times_Z^{\text{lax}} Y \simeq X \prod_{Z^{\{0\}}} Z^{[1]} \prod_{Z^{\{1\}}} Y$  and the oplax pullback is  $X \times_Z^{\text{oplax}} Y \simeq X \prod_{Z^{\{1\}}} Z^{[1]} \prod_{Z^{\{0\}}} Y$ .

Notation 5.44. Let  $Cat_{(\infty,2)}^{lax \text{ push}}, Cat_{(\infty,2)}^{oplax \text{ push}} \subset Cat_{(\infty,2)}$  be the subcategories of  $(\infty, 2)$ -categories that admit (op)lax pushouts and functors preserving (op)lax pushouts.

**Corollary 5.45.** The  $\infty$ -categories  $\operatorname{Cat}_{(\infty,2)}^{\operatorname{lax push}}, \operatorname{Cat}_{(\infty,2)}^{\operatorname{oplax push}}$  carry closed symmetric monoidal structures such that the inclusions  $\operatorname{Cat}_{(\infty,2)}^{\operatorname{lax push}}, \operatorname{Cat}_{(\infty,2)}^{\operatorname{oplax push}} \subset \operatorname{Cat}_{(\infty,2)}$  are lax symmetric monoidal, where the right hand side carries the cartesian structure.

#### 5.5. A tensor product for Cauchy-complete enriched $\infty$ -categories.

**Definition 5.46.** Let  $\mathcal{V}^{\otimes} \to Ass, \mathcal{W}^{\otimes} \to Ass$  be presentably monoidal  $\infty$ -categories and  $\mathcal{J}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a bi-enriched  $\infty$ -category. An enriched weight on  $\mathcal{J}$  is absolute if it is preserved by any  $\mathcal{V}, \mathcal{W}$ -enriched functor between  $\infty$ -categories bi-quasi-enriched in  $\mathcal{V}, \mathcal{W}$ .

**Example 5.47.** For  $\mathcal{V} = \mathcal{S}$  the  $\infty$ -category of spaces the  $\infty$ -category Idem classifying idempotents of [17, Definition 4.4.5.2.] equipped with the trivial left  $\mathcal{S}$ -enriched weight is an absolute left  $\mathcal{S}$ -enriched weight.

**Notation 5.48.** Let  $\mathcal{V}^{\otimes} \to \operatorname{Ass}$ ,  $\mathcal{W}^{\otimes} \to \operatorname{Ass}$  be presentably monoidal  $\infty$ -categories and  $\mathcal{J}^{\oplus} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a bi-enriched  $\infty$ -category. The Cauchy-completion  $\widehat{\mathcal{M}}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  of  $\mathcal{M}^{\oplus} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  is the full bi-enriched subcategory of  $\mathcal{PBEnv}(\mathcal{M})^{\otimes}_{\operatorname{BEnr}} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  spanned by the absolute enriched weights on  $\mathcal{M}$ .

**Remark 5.49.** Let  $F: \mathcal{M}^{\circledast} \to \mathcal{N}^{\circledast}$  be a  $\mathcal{V}, \mathcal{W}$ -enriched functor between bi-enriched  $\infty$ -categories. The functor  $F_{!}: \mathcal{P}BEnv(\mathcal{M})_{BEnr} \to \mathcal{P}BEnv(\mathcal{N})_{BEnr}$  sends  $\widehat{\mathcal{M}}$  to  $\widehat{\mathcal{N}}$  since for every absolute enriched weight H on  $\mathcal{M}$  the  $F_{!}(H)$ -weighted colimit of a  $\mathcal{V}, \mathcal{W}$ -enriched functor  $\phi: \mathcal{N}^{\circledast} \to \mathcal{O}^{\circledast}$  between bi-enriched  $\infty$ -categories is the H-weighted colimit of  $\phi \circ F$  as a consequence of Corollary 3.21.

**Definition 5.50.** Let  $\mathcal{V}^{\otimes} \to \operatorname{Ass}, \mathcal{W}^{\otimes} \to \operatorname{Ass}$  be presentably monoidal  $\infty$ -categories. A bi-enriched  $\infty$ -category  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  is Cauchy-complete if it admits the colimit of any  $\mathcal{V}, \mathcal{W}$ -enriched functor weighted with respect to any absolute enriched weight.

**Proposition 5.51.** Let  $\mathcal{V}^{\otimes} \to Ass, \mathcal{W}^{\otimes} \to Ass$  be presentably monoidal  $\infty$ -categories,  $\mathcal{J}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a bi-enriched  $\infty$ -category and H an enriched weight on  $\mathcal{J}$ . The following conditions are equivalent:

(1) The  $\mathcal{V}, \mathcal{W}$ -enriched functor

84

$$\operatorname{Mor}_{\mathcal{PBEnv}(\mathcal{J})_{\operatorname{BEnr}}}(\operatorname{H}, -) : \mathcal{PBEnv}(\mathcal{J})_{\operatorname{BEnr}}^{\circledast} \to (\mathcal{V} \otimes \mathcal{W})^{\circledast}$$

is linear and preserves small colimits.

- (2) The enriched weight H on  $\mathcal{J}$  is absolute.
- (3) The H-weighted colimit of the  $\mathcal{V}, \mathcal{W}$ -enriched embedding  $\mathcal{J}^{\circledast} \subset \mathcal{M}^{\circledast} := \mathcal{P}BEnv(\mathcal{J})^{\circledast}_{BEnr}$  is preserved by the  $\mathcal{V}, \mathcal{W}$ -enriched embedding  $\mathcal{M}^{\circledast} \subset \mathcal{P}BEnv(\mathcal{M})^{\circledast}_{BEnr}$ .

*Proof.* Assume that (1) holds. Let  $F : \mathcal{J}^{\circledast} \to \mathcal{M}^{\circledast}, G : \mathcal{M}^{\circledast} \to \mathcal{N}^{\circledast}$  be  $\mathcal{V}, \mathcal{W}$ -enriched functors between bi-quasi-enriched  $\infty$ -categories. We like to see that G preserves the H-weighted colimit of F if the latter exists. The universal morphism  $H \to F^*(\operatorname{colim}^H(F))$  gives rise to a morphism

$$\theta: \mathfrak{PBEnv}(\mathcal{M})_{\mathrm{BEnr}}(\mathrm{colim}^{\mathrm{H}}(\mathrm{F}), -) \to \mathfrak{PBEnv}(\mathcal{J})_{\mathrm{BEnr}}(\mathrm{H}, -) \circ \mathrm{F}^{*}$$

in LinFun<sup>L</sup><sub> $\mathcal{V},\mathcal{W}$ </sub> ( $\mathcal{P}BEnv(\mathcal{M})_{BEnr}, \mathcal{V} \otimes \mathcal{W}$ ) since colim<sup>H</sup>(F)  $\in \mathcal{M}$  and by [9, Theorem 4.45.] and assumption (1) and Proposition 2.149. By universal property of the weighted colimit for every X  $\in \mathcal{M}$  the morphism  $\theta_X$  is an equivalence so that  $\theta$  is an equivalence by Corollary 2.97. To see that the canonical morphism colim<sup>H</sup>(G  $\circ$  F)  $\rightarrow$  G(colim<sup>H</sup>(F)) in  $\mathcal{N}$  is an equivalence, we need to see that for every Y  $\in \mathcal{N}$  the canonical map

$$\mathcal{PBEnv}(\mathcal{J})_{\mathrm{BEnr}}(\mathrm{H},\mathrm{F}^{*}(\mathrm{G}^{*}(\mathrm{Y}))) \simeq \mathcal{PBEnv}(\mathcal{M})_{\mathrm{BEnr}}(\mathrm{colim}^{\mathrm{H}}(\mathrm{F}),\mathrm{G}^{*}(\mathrm{Y})) \simeq$$

$$\mathcal{P}BEnv(\mathcal{N})_{BEnr}(G_!(\operatorname{colim}^{H}(F)), Y) \simeq \mathcal{N}(G(\operatorname{colim}^{H}(F)), Y) \rightarrow \mathcal{N}(\operatorname{colim}^{H}(G \circ F), Y) \simeq$$

$$\mathcal{P}BEnv(\mathcal{J})_{BEnr}(H, (G \circ F)^*(Y)) \simeq \mathcal{P}BEnv(\mathcal{J})_{BEnr}(H, F^*(G^*(Y)))$$

is an equivalence. The latter morphism identifies with the identity. This proves (2).

Condition (2) trivially implies (3). Assume that (3) holds. The H-weighted colimit of the  $\mathcal{V}, \mathcal{W}$ enriched embedding  $\mathcal{J}^{\circledast} \subset \mathcal{M}^{\circledast} := \mathcal{P}BEnv(\mathcal{J})^{\circledast}_{BEnr}$  is H. By assumption this H-weighted colimit is preserved by the  $\mathcal{V}, \mathcal{W}$ -enriched embedding  $\mathcal{M}^{\circledast} \subset \mathcal{P}BEnv(\mathcal{M})^{\circledast}_{BEnr}$ . Thus the image of  $H \in \mathcal{P}BEnv(\mathcal{J})_{BEnr} = \mathcal{M}$  in  $\mathcal{P}BEnv(\mathcal{M})_{BEnr}$  is the H-weighted colimit of the composition  $\mathcal{J}^{\circledast} \subset \mathcal{P}BEnv(\mathcal{J})^{\circledast}_{BEnr} = \mathcal{M}^{\circledast} \subset \mathcal{P}BEnv(\mathcal{J})^{\circledast}_{BEnr}$ 

 $\mathcal{PBEnv}(\mathcal{M})^{\circledast}_{\mathrm{BEnr}}$ . The latter composition factors as  $\mathcal{J}^{\circledast} \subset \mathcal{PBEnv}(\mathcal{J})^{\circledast}_{\mathrm{BEnr}} \xrightarrow{\theta_{!}} \mathcal{PBEnv}(\mathcal{M})^{\circledast}_{\mathrm{BEnr}}$ , where the latter  $\mathcal{V}, \mathcal{W}$ -enriched embedding is induced by the  $\mathcal{V}, \mathcal{W}$ -enriched embedding  $\theta : \mathcal{J}^{\circledast} \subset \mathcal{PBEnv}(\mathcal{J})^{\circledast}_{\mathrm{BEnr}} = \mathcal{M}^{\circledast}$ . Since  $\theta_{!}$  preserves weighted colimits, the image H' of H  $\in \mathcal{PBEnv}(\mathcal{J})_{\mathrm{BEnr}} = \mathcal{M}$  in  $\mathcal{PBEnv}(\mathcal{M})_{\mathrm{BEnr}}$  is the image of H  $\in \mathcal{PBEnv}(\mathcal{J})_{\mathrm{BEnr}}$  under  $\theta_{!}$ . Thus there is a canonical equivalence

$$\operatorname{Mor}_{\mathcal{P}\operatorname{BEnv}(\mathcal{J})_{\operatorname{BEnr}}}(\operatorname{H}, -) \simeq \operatorname{Mor}_{\mathcal{P}\operatorname{BEnv}(\mathcal{M})_{\operatorname{BEnr}}}(\theta_{!}(\operatorname{H}), -) \circ \theta_{!} \simeq \operatorname{Mor}_{\mathcal{P}\operatorname{BEnv}(\mathcal{M})_{\operatorname{BEnr}}}(\operatorname{H}', -) \circ \theta_{!}$$
  
of  $\mathcal{V}, \mathcal{W}$ -enriched functors  $\mathcal{P}\operatorname{BEnv}(\mathcal{J})_{\operatorname{BEnr}}^{\mathfrak{G}} \to (\mathcal{V} \otimes \mathcal{W})^{\mathfrak{S}}$ . Thus (1) follows.

**Corollary 5.52.** Let  $\mathcal{V}^{\otimes} \to \operatorname{Ass}$ ,  $\mathcal{W}^{\otimes} \to \operatorname{Ass}$  be presentably monoidal  $\infty$ -categories and  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a small bi-enriched  $\infty$ -category. The Cauchy-completion  $\widehat{\mathcal{M}}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  of  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  is small.

Proof. Since  $\mathcal{V}^{\otimes}\mathcal{W}$  is presentable, the tensor unit of  $\mathcal{V}\otimes\mathcal{W}$  is  $\kappa$ -compact for some small regular cardinal  $\kappa$ . Let  $H \in \widehat{\mathcal{M}}$ . By Proposition 5.51 the  $\mathcal{V}$ ,  $\mathcal{W}$ -enriched functor  $\operatorname{Mor}_{\mathcal{P}\operatorname{BEnv}(\mathcal{J})_{\operatorname{BEnr}}}(H, -) : \mathcal{P}\operatorname{BEnv}(\mathcal{J})_{\operatorname{BEnr}}^{\otimes} \to (\mathcal{V}\otimes\mathcal{W})^{\otimes}$  preserves small colimits and therefore small  $\kappa$ -filtered colimits. Hence the left adjoint of the latter functor, which sends  $V \otimes W$  to  $V \otimes H \otimes W$ , preserves  $\kappa$ -compact objects. Thus H is  $\kappa$ -compact. So  $\widehat{\mathcal{M}}$  is contained in the full subcategory of  $\kappa$ -compact objects of  $\mathcal{P}\operatorname{BEnv}(\mathcal{J})_{\operatorname{BEnr}}$ . Proposition 2.153 guarantees that  $\mathcal{P}\operatorname{BEnv}(\mathcal{J})_{\operatorname{BEnr}}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  is a presentably bitensored  $\infty$ -category and so locally small. Presentability implies that the full subcategory of  $\kappa$ -compact objects of  $\mathcal{P}\operatorname{BEnv}(\mathcal{J})_{\operatorname{BEnr}}$  is small [17, Remark 5.4.2.13.].

**Proposition 5.53.** Let  $\mathcal{V}^{\otimes} \to \operatorname{Ass}, \mathcal{W}^{\otimes} \to \operatorname{Ass}$  be presentably monoidal  $\infty$ -categories. A bi-enriched  $\infty$ -category  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  is Cauchy-complete if and only if every absolute enriched weight on  $\mathcal{M}$  belongs to  $\mathcal{M}$ .

Proof. Assume that every absolute enriched weight on  $\mathcal{M}$  belongs to  $\mathcal{M}$ , let H be an absolute enriched weight on a bi-enriched ∞-category  $\mathcal{J}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  and  $F : \mathcal{J}^{\circledast} \to \mathcal{M}^{\circledast}$  a  $\mathcal{V}, \mathcal{W}$ -enriched functor. The H-weighted colimit of the  $\mathcal{V}, \mathcal{W}$ -enriched functor  $\mathcal{J}^{\circledast} \xrightarrow{F} \mathcal{M}^{\circledast} \subset \mathcal{P}BEnv(\mathcal{M})^{\circledast}_{BEnr}$  is  $F_{!}(H)$  by Proposition 3.23, which is an absolute enriched weight on  $\mathcal{M}$  by Remark 5.49. So by assumption  $F_{!}(H) \in \mathcal{M}$ . Thus the H-weighted colimit of the  $\mathcal{V}, \mathcal{W}$ -enriched functor  $\mathcal{J}^{\circledast} \xrightarrow{F} \mathcal{M}^{\circledast} \subset \mathcal{P}BEnv(\mathcal{M})^{\circledast}_{BEnr}$  lies in  $\mathcal{M}$  and so is the H-weighted colimit of F. Conversely, assume that  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  is Cauchy-complete and let H be an absolute enriched weight on  $\mathcal{M}$ . The H-weighted colimit of the identity of  $\mathcal{M}$ , which exists by Cauchy-completeness, is preserved by any  $\mathcal{V}, \mathcal{W}$ -enriched functor and so is sent by the  $\mathcal{V}, \mathcal{W}$ -enriched embedding  $\mathcal{M}^{\circledast} \subset \mathcal{P}BEnv(\mathcal{M})^{\circledast}_{BEnr}$  to the H-weighted colimit of the  $\mathcal{V}, \mathcal{W}$ -enriched embedding  $\mathcal{M}^{\circledast} \subset \mathcal{P}BEnv(\mathcal{M})^{\circledast}_{BEnr}$  to the H-weighted colimit of the  $\mathcal{V}, \mathcal{W}$ -enriched embedding  $\mathcal{M}^{\circledast} \subset \mathcal{P}BEnv(\mathcal{M})^{\circledast}_{BEnr}$ , which is H by Proposition 3.23. So H belongs to  $\mathcal{M}$ .

**Proposition 5.54.** Let  $\mathcal{V}^{\otimes} \to \operatorname{Ass}$ ,  $\mathcal{W}^{\otimes} \to \operatorname{Ass}$  be presentably monoidal  $\infty$ -categories,  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a bi-enriched  $\infty$ -category and  $\mathcal{N}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a Cauchy-complete bi-enriched  $\infty$ -category. The following induced functor is an equivalence:

$$\rho_{\mathcal{N}}$$
: EnrFun <sub>$\mathcal{V},\mathcal{W}$</sub>  $(\mathcal{M},\mathcal{N}) \rightarrow$  EnrFun <sub>$\mathcal{V},\mathcal{W}$</sub>  $(\mathcal{M},\mathcal{N}).$ 

*Proof.* We prove first that the functor  $\rho_{N}$  is conservative. Let H ∈  $\widehat{\mathcal{M}}$ . Then H ∈  $\mathcal{P}BEnv(\mathcal{M})_{BEnr}$  is the H-weighted colimit of the  $\mathcal{V}$ ,  $\mathcal{W}$ -enriched embedding  $\mathcal{M}^{\circledast} \subset \widehat{\mathcal{M}}^{\circledast}$ . Since H is absolute, every  $\mathcal{V}$ ,  $\mathcal{W}$ -enriched functor F :  $\widehat{\mathcal{M}}^{\circledast} \to \mathcal{N}^{\circledast}$  preserves the H-weighted colimit of the  $\mathcal{V}$ ,  $\mathcal{W}$ -enriched embedding  $\mathcal{M}^{\circledast} \subset \widehat{\mathcal{M}}^{\circledast}$  so that the canonical morphism colim<sup>H</sup>(F<sub>| $\mathcal{M}^{\circledast}$ </sub>) → F(H) is an equivalence. This implies that the functor  $\rho_{\mathcal{N}}$  is conservative and that it is the pullback of the conservative functor  $\rho_{\mathcal{P}BEnv(\mathcal{N})_{BEnr}}$  since  $\mathcal{N}$  is closed in  $\mathcal{P}BEnv(\mathcal{N})_{BEnr}$  under H-weighted colimits. So it is enough to see that  $\rho_{\mathcal{P}BEnv(\mathcal{N})_{BEnr}}$  is an equivalence. This functor admits a fully faithful left adjoint by [9, Proposition 2.62.] and so is an equivalence being conservative.

**Corollary 5.55.** Let  $\mathcal{V}^{\otimes} \to \operatorname{Ass}$ ,  $\mathcal{W}^{\otimes} \to \operatorname{Ass}$  be presentably monoidal  $\infty$ -categories and  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a bi-enriched  $\infty$ -category. The Cauchy-completion of  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  is Cauchy-complete.

*Proof.* Let H ∈ PBEnv( $\widehat{\mathcal{M}}$ )<sub>BEnr</sub> be an absolute enriched weight on  $\widehat{\mathcal{M}}$ . We like to see that H ∈  $\widehat{\mathcal{M}}$ . By Proposition 5.54 the embedding  $\mathcal{M}^{\circledast} \subset \widehat{\mathcal{M}}^{\circledast}$  induces an equivalence  $\mathcal{PBEnv}(\mathcal{M})_{BEnr} \simeq \mathcal{PBEnv}(\widehat{\mathcal{M}})_{BEnr}$ . So H is the image of an enriched weight H' on  $\mathcal{M}$  that is absolute by Proposition 5.51. Thus H' ∈  $\widehat{\mathcal{M}}$ . By Proposition 5.54 the  $\mathcal{V}, \mathcal{W}$ -enriched embedding  $\widehat{\mathcal{M}}^{\circledast} \subset \mathcal{PBEnv}(\mathcal{M})^{\circledast}_{BEnr} \simeq \mathcal{PBEnv}(\widehat{\mathcal{M}})^{\circledast}_{BEnr}$  is the canonical embedding since both are equivalent after restriction to  $\mathcal{M}^{\circledast}$ . So H is the image of H' under the embedding  $\widehat{\mathcal{M}} \subset \mathcal{PBEnv}(\widehat{\mathcal{M}})_{BEnr}$ .

Proposition 3.85 gives the following corollary:

**Corollary 5.56.** Let  $\mathcal{V}^{\otimes} \to \operatorname{Ass}$ ,  $\mathcal{W}^{\otimes} \to \operatorname{Ass}$  be presentably monoidal  $\infty$ -categories and  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a small bi-enriched  $\infty$ -category. The Cauchy-completion  $\widehat{\mathcal{M}}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  is generated by  $\mathcal{M}$  under colimits weighted with respect to small absolute  $\mathcal{V}$ ,  $\mathcal{W}$ -enriched weights.

**Notation 5.57.** Let  $\mathcal{V}^{\otimes} \to Ass$  be a presentably monoidal  $\infty$ -category. Let  $_{\mathcal{V}}LEnr^{\wedge} \subset _{\mathcal{V}}LEnr$  be the full subcategory of Cauchy-complete left  $\mathcal{V}$ -enriched  $\infty$ -categories.

**Corollary 5.58.** Let  $n \ge 1$  and  $\mathcal{V}^{\boxtimes} \to \mathbb{E}_{n+1}$  a presentably  $\mathbb{E}_{n+1}$ -monoidal  $\infty$ -category. The  $\infty$ -category  $_{\mathcal{V}}$ LEnr<sup> $\wedge$ </sup> of Cauchy-complete left  $\mathcal{V}$ -enriched  $\infty$ -categories carries a canonical  $\mathbb{E}_n$ -monoidal structure.

*Proof.* There is an identity  $_{\mathcal{V}}\text{LEnr}^{\wedge} = _{\mathcal{V}}\text{LEnr}^{\wedge}(\mathcal{H})$ , where  $\mathcal{H}$  is the set of absolute left  $\mathcal{V}$ -enriched weights. We apply Theorem 4.42.

### 5.6. A tensor product for stable $(\infty, n)$ -categories.

**Definition 5.59.** A weakly bi-enriched  $\infty$ -category  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  is preadditive if it admits finite conical coproducts, i.e. conical colimits indexed by finite sets, and finite conical products, i.e. conical limits indexed by finite sets, and the morphism  $\emptyset \to \ast$  from the initial to the final object in  $\mathcal{M}$  is an equivalence, and for every X, Y  $\in \mathcal{M}$  the canonical morphism X  $\coprod$  Y  $\to$  X  $\times$  Y in  $\mathcal{M}$  is an equivalence.

**Remark 5.60.** A weakly bi-enriched  $\infty$ -category  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  is preadditive if it admits finite conical coproducts and finite conical products and  $\mathcal{M}$  is preadditive. A weakly bi-enriched  $\infty$ -category is preadditive if and only if the opposite weakly bi-enriched  $\infty$ -category is preadditive.

**Example 5.61.** A weakly bi-enriched  $\infty$ -category  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  is preadditive if it admits left and right cotensors and  $\mathcal{M}$  is preadditive, and so dually a weakly bi-enriched  $\infty$ -category  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  is preadditive if it admits left and right tensors and  $\mathcal{M}$  is preadditive. In particular, for every monoidal  $\infty$ -category  $\mathcal{V}^{\circledast} \to \mathcal{A}$ ss such that  $\mathcal{V}$  is preadditive we have that  $\mathcal{V}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{V}^{\circledast}$  is preadditive.

**Example 5.62.** Every full weakly bi-enriched subcategory of a preadditive weakly bi-enriched  $\infty$ -category closed under finite products is again preadditive.

**Remark 5.63.** Let  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes}$  be a weakly left enriched  $\infty$ -category and  $\mathcal{N}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  a preadditive weakly bi-enriched  $\infty$ -category. The weakly right enriched  $\infty$ -category  $\operatorname{EnrFun}_{\mathcal{V},\emptyset}(\mathcal{M},\mathcal{N})^{\circledast} \to \mathcal{W}^{\otimes}$  is preadditive. This follows from Proposition 3.94. In particular, for every monoidal  $\infty$ -category  $\mathcal{V}^{\otimes} \to \operatorname{Ass}$  such that  $\mathcal{V}$  is preadditive we have that  $\operatorname{EnrFun}_{\mathcal{V},\emptyset}(\mathcal{M},\mathcal{V})^{\circledast} \to \mathcal{V}^{\otimes}$  is preadditive.

**Notation 5.64.** Let  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  be a weakly bi-enriched  $\infty$ -category that admits finite conical products, i.e. conical limits indexed by finite sets. Let  $\operatorname{Cmon}(\mathcal{M})^{\circledast} \subset (\mathcal{M}^{\operatorname{Comm}})^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  be the full weakly bi-enriched subcategory spanned by the commutative monoid objects in  $\mathcal{M}$ , i.e. the functors  $X : \operatorname{Comm} \to \mathcal{M}$  such that for every  $n \ge 0$  the induced morphism  $X(\langle n \rangle) \to X(\langle 1 \rangle)^{\times n}$  is an equivalence.

**Remark 5.65.** Evaluating at  $(1) \in \text{Comm gives a } \mathcal{V}, \mathcal{W}\text{-enriched functor } \text{Cmon}(\mathcal{M})^{\circledast} \to \mathcal{M}^{\circledast}$ .

**Remark 5.66.** Let  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  be a weakly bi-enriched  $\infty$ -category that admits finite conical products and left (right) cotensors. Then  $\operatorname{Cmon}(\mathcal{M})^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  admits left (right) cotensors, which are preserved by the forgetful functor  $\operatorname{Cmon}(\mathcal{M})^{\circledast} \to \mathcal{M}^{\circledast}$ : by Proposition 3.94 we find that  $(\mathcal{N}^{\operatorname{Comm}})^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  admits left (right) cotensors, which are formed object-wise. Thus  $\operatorname{Cmon}(\mathcal{N})^{\circledast}$  is closed in  $(\mathcal{M}^{\operatorname{Comm}})^{\circledast}$  under left (right) cotensors since forming left (right) cotensors are right adjoints.

**Lemma 5.67.** Let  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be a weakly bi-enriched  $\infty$ -category that admits finite conical products. Then  $\operatorname{Cmon}(\mathcal{M})^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  is preadditive.

*Proof.* There is a  $\mathcal{V}, \mathcal{W}$ -enriched embedding  $\mathcal{M}^{\circledast} \subset \mathcal{N}^{\circledast}$  preserving finite conical products into a weakly bi-enriched ∞-category that admits left and right cotensors. For example take  $\mathcal{N}^{\circledast} := \mathcal{P}BEnv(\mathcal{M})^{\circledast}$ . The latter embedding yields an enriched embedding  $Cmon(\mathcal{M})^{\circledast} \subset Cmon(\mathcal{M})^{\circledast}$  that preserves finite conical products. By [5, Proposition 2.3.] the ∞-categories  $Cmon(\mathcal{M}), Cmon(\mathcal{N})$  are preadditive. Thus by Example 5.62 and 5.61 we find that  $Cmon(\mathcal{M})^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  is preadditive if  $Cmon(\mathcal{N})^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  admits left and right cotensors. This holds by Remark 5.66.

**Lemma 5.68.** Let  $\mathcal{V}^{\otimes} \to \operatorname{Ass}$ ,  $\mathcal{W}^{\otimes} \to \operatorname{Ass}$  be  $\infty$ -operads such that  $\mathcal{V}$  and  $\mathcal{W}$  are the empty category or preadditive or  $\mathcal{V}^{\otimes} = \mathcal{W}^{\otimes} = S^{\times}$ . Let  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be a preadditive weakly bi-enriched  $\infty$ -category such that for every  $X \in \mathcal{M}, Y \in \mathcal{PBEnv}(\mathcal{M})$  and  $V_1, ..., V_n \in \mathcal{V}, W_1, ..., W_m \in \mathcal{W}$  for  $n, m \ge 0$  the functors

$$\begin{aligned} \operatorname{Mul}_{\mathcal{P}\operatorname{BEnv}(\mathcal{M})}(V_1,...,(-),...,V_n,X,W_1,...,W_m;Y) &: \mathcal{V}^{\operatorname{op}} \to \mathcal{S}, \\ \operatorname{Mul}_{\mathcal{P}\operatorname{BEnv}(\mathcal{M})}(V_1,...,V_n,X,W_1,...,(-),...,W_m;Y) &: \mathcal{W}^{\operatorname{op}} \to \mathcal{S} \end{aligned}$$

preserve finite products. The  $\mathcal{V}, \mathcal{W}$ -enriched forgetful functor  $\operatorname{Cmon}(\mathcal{M})^{\otimes} \to \mathcal{M}^{\otimes}$  is an equivalence.

*Proof.* We first prove that there is an enriched embedding  $\mathcal{M}^{\circledast} \subset \mathcal{N}^{\circledast}$  preserving finite conical products into a preadditive weakly bi-enriched ∞-category that admits left and right cotensors. Let  $\mathcal{N}^{\circledast} := \mathcal{P}BEnv(\mathcal{M})^{\circledast}$ . The weakly bi-enriched ∞-category  $\mathcal{P}BEnv(\mathcal{M})^{\circledast} \to \mathcal{P}Env(\mathcal{V})^{\otimes} \times \mathcal{P}Env(\mathcal{W})^{\otimes}$  is preadditive by Example 5.61 since  $\mathcal{P}BEnv(\mathcal{M})$  is preadditive what we prove in the following: the jointly conservative functors

$$\operatorname{Mul}_{\operatorname{PBEnv}(\mathcal{M})}(V_1, ..., (-), ..., V_n, X, W_1, ..., W_m; -) : \operatorname{PBEnv}(\mathcal{M}) \to \operatorname{Fun}^{\Pi}(\mathcal{V}^{\operatorname{op}}, S)$$

$$\operatorname{Iul}_{\operatorname{\mathcal{PBEnv}}(\mathcal{M})}(V_1, ..., V_n, X, W_1, ..., (-), ..., W_m; -) : \operatorname{\mathcal{PBEnv}}(\mathcal{M}) \to \operatorname{Fun}^{\Pi}(\mathcal{W}^{\operatorname{op}}, S)$$

M

for  $V_1, ..., V_n \in \mathcal{V}, W_1, ..., W_m \in \mathcal{W}$  for  $n, m \ge 0$  and  $X \in \mathcal{M}$  preserve small limits and small colimits by [9, Theorem 4.45.] and Fun<sup>II</sup>( $\mathcal{V}^{op}, \mathcal{S}$ ), Fun<sup>II</sup>( $\mathcal{W}^{op}, \mathcal{S}$ ) are preadditive if  $\mathcal{V}, \mathcal{W}$  are empty or preadditive by [5, Corollary 2.4.]. The  $\mathcal{V}, \mathcal{W}$ -enriched forgetful functor  $\operatorname{Cmon}(\mathcal{M})^{\circledast} \to \mathcal{M}^{\circledast}$  is the pullback of the  $\mathcal{V}, \mathcal{W}$ -enriched forgetful functor  $\operatorname{Cmon}(\mathcal{N})^{\circledast} \to \mathcal{N}^{\circledast}$  that preserves left and right cotensors by Remark 5.66. The latter is an equivalence since it preserves left and right cotensors and induces an equivalence on underlying  $\infty$ -categories by [5, Proposition 2.3.].

**Proposition 5.69.** Let  $\mathcal{V}^{\otimes} \to \operatorname{Ass}, \mathcal{W}^{\otimes} \to \operatorname{Ass}$  be  $\infty$ -operads such that  $\mathcal{V}, \mathcal{W}$  are empty or preadditive or  $\mathcal{V}^{\otimes} = \mathcal{W}^{\otimes} = \mathcal{S}^{\times}$ . Let  $\mathcal{M}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be a preadditive weakly bi-enriched  $\infty$ -category such that for every  $X \in \mathcal{M}, Y \in \mathcal{PBEnv}(\mathcal{M})$  and  $V_1, ..., V_n \in \mathcal{V}, W_1, ..., W_m \in \mathcal{W}$  for  $n, m \ge 0$  the functors

$$\operatorname{Mul}_{\mathcal{PBEnv}(\mathcal{M})}(V_1, ..., (-), ..., V_n, X, W_1, ..., W_m; Y) : \mathcal{V}^{\operatorname{op}} \to \mathcal{S},$$

 $Mul_{\mathcal{P}BEnv(\mathcal{M})}(V_1,..,V_n,X,W_1,...,(-),...,W_m;Y):\mathcal{W}^{op} \rightarrow \mathcal{S}$ 

preserve finite products and  $\mathbb{N}^{\circledast} \to \mathbb{V}^{\circledast} \times \mathbb{W}^{\circledast}$  a weakly bi-enriched  $\infty$ -category that admits finite conical products. The following induced functor is an equivalence:

$$\theta_{\mathcal{N}} : \operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}^{\Pi}(\mathcal{M},\operatorname{Cmon}(\mathcal{N})) \to \operatorname{EnrFun}_{\mathcal{V},\mathcal{W}}^{\Pi}(\mathcal{M},\mathcal{N}).$$

*Proof.* For every ∞-category K there is a  $\mathcal{V}, \mathcal{W}$ -enriched equivalence  $\operatorname{Cmon}(\mathcal{N}^{K})^{\circledast} \simeq (\operatorname{Cmon}(\mathcal{N})^{K})^{\circledast}$ and the functor  $\theta_{\mathcal{N}}^{K}$  identifies with the functor  $\theta_{\mathcal{N}^{K}}$ . Consequently, it is enough to prove that  $\theta_{\mathcal{N}}$  is a bijection on equivalence classes. Since  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  is preadditive, by Lemma 5.68 the  $\mathcal{V}, \mathcal{W}$ enriched forgetful functor  $\operatorname{Cmon}(\mathcal{M})^{\circledast} \to \mathcal{M}^{\circledast}$  is an equivalence. The functor  $\theta_{\mathcal{N}}$  is essentially surjective since every  $\mathcal{V}, \mathcal{W}$ -enriched functor  $\mathcal{M}^{\circledast} \to \mathcal{N}^{\circledast}$  is the image under  $\theta_{\mathcal{N}}$  of the  $\mathcal{V}, \mathcal{W}$ -enriched functor  $\mathcal{M}^{\circledast} \simeq \operatorname{Cmon}(\mathcal{M})^{\circledast} \to \operatorname{Cmon}(\mathcal{N})^{\circledast}$ . The functor  $\theta_{\mathcal{N}}$  is essentially injective: every  $\mathcal{V}, \mathcal{W}$ -enriched functor  $\phi: \mathcal{M}^{\circledast} \to \operatorname{Cmon}(\mathcal{N})^{\circledast}$  factors as  $\mathcal{M}^{\circledast} \simeq \operatorname{Cmon}(\mathcal{M})^{\circledast} \xrightarrow{\operatorname{Cmon}(\theta_{\mathcal{N}}(\phi))} \operatorname{Cmon}(\operatorname{Cmon}(\mathcal{N}))^{\circledast} \to \operatorname{Cmon}(\mathcal{N})^{\circledast}$ .

Let Fin be the category of finite sets.

Notation 5.70. Let  $\mathcal{V}^{\otimes} \to Ass, \mathcal{W}^{\otimes} \to Ass$  be  $\infty$ -operads. Let

Notation 5.71. Let  $\mathcal{V}^{\otimes} \to Ass, \mathcal{W}^{\otimes} \to Ass$  be  $\infty$ -operads. Let

 $_{\mathcal{V}}$ Preadd $_{\mathcal{W}} \subset _{\mathcal{V}}^{\coprod}$ BEnr $_{\mathcal{W}}$ 

be the full subcategory of preadditive  $\mathcal{V}, \mathcal{W}$ -bi-enriched  $\infty$ -categories. For  $\mathcal{V}^{\otimes} = \mathcal{W}^{\otimes} = \mathcal{S}^{\times}$  we drop  $\mathcal{V}, \mathcal{W}$  from the notation.

By definition  ${}^{\mathrm{II}}_{\mathcal{V}}\omega\operatorname{BEnr}_{\mathcal{W}}, {}^{\mathrm{II}}_{\mathcal{V}}\operatorname{BEnr}_{\mathcal{W}}$  are the subcategories of (weakly)  $\mathcal{V}, \mathcal{W}$ -bi-enriched  $\infty$ -categories that admit finite conical coproducts and  $\mathcal{V}, \mathcal{W}$ -enriched functors preserving finite conical coproducts.

**Proposition 5.72.** Let  $\mathcal{V}^{\otimes} \to \operatorname{Ass}, \mathcal{W}^{\otimes} \to \operatorname{Ass}$  be  $\infty$ -operads. Then  ${}^{\mathrm{II}}_{\mathcal{V}}\omega\operatorname{BEnr}_{\mathcal{W}}, {}^{\mathrm{II}}_{\mathcal{V}}\operatorname{BEnr}_{\mathcal{W}}, {}^{\mathrm{Preadd}}_{\mathcal{W}}$  are preadditive.

*Proof.* The ∞-category  $_{\mathcal{V}}\omega$ BEnr<sub>W</sub> admits finite products: for every small weakly bi-enriched ∞-categories  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}, \mathcal{N}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  the weakly bi-enriched ∞-category  $\mathcal{M}^{\circledast} \times_{(\mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes})} \mathcal{N}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  is the product in  $_{\mathcal{V}}\omega$ BEnr<sub>W</sub>, and id:  $\mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  is the final object. By description of the morphism objects of a product a  $\mathcal{V}, \mathcal{W}$ -enriched functor to a product preserves finite conical coproducts if it components do. Thus the subcategory  $\overset{\Pi}{\mathcal{V}}\omega$ BEnr<sub>W</sub> admits finite products and the inclusion  $\overset{\Pi}{\mathcal{V}}\omega$ BEnr<sub>W</sub> ⊂  $_{\mathcal{V}}\omega$ BEnr<sub>W</sub> preserves finite products. Moreover the final object in  $\overset{\Pi}{\mathcal{V}}\omega$ BEnr<sub>W</sub> is an initial object. Let  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  be a weakly bi-enriched ∞-category that admits finite conical coproducts. Then the  $\mathcal{V}, \mathcal{W}$ -enriched diagonal functor  $\mathcal{M}^{\circledast} \to \mathcal{M}^{\circledast} \times_{(\mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes})} \mathcal{M}^{\circledast}$  admits a  $\mathcal{V}, \mathcal{W}$ -enriched left adjoint  $\mu : \mathcal{M}^{\circledast} \times_{(\mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes})} \mathcal{M}^{\circledast}$  assigning the coproduct. Moreover the unique  $\mathcal{V}, \mathcal{W}$ -enriched functor  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes}$  admits a  $\mathcal{V}, \mathcal{W}$ -enriched left adjoint  $\alpha : \mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes} \to \mathcal{M}^{\circledast}$  assigning the initial  $\mathcal{M}^{\circledast} \times_{(\mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes})} \mathcal{M}^{\circledast}$ 

object. We apply [16, Proposition 2.4.3.19.] and have to verify that the composition  $\mathcal{M}^{\circledast} \xrightarrow{\mathcal{M}^{\circledast} \times_{(\mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes})} \alpha} \mathcal{M}^{\circledast} \times_{(\mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes})} \mathcal{M}^{\circledast} \xrightarrow{\mu} \mathcal{M}^{\circledast}$  is the identity and the composition  $\mathcal{M}^{\circledast} \times_{(\mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes})} \mathcal{N}^{\circledast} \times_{(\mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes})} \mathcal{M}^{\circledast} \times_{(\mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes})} \mathcal{M}^{\circledast} \times_{(\mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes})} \mathcal{M}^{\circledast} \times_{(\mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes})} \mathcal{M}^{\circledast} \times_{(\mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes})} \mathcal{N}^{\circledast} \xrightarrow{\mu^{\times} (\mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes})} \mathcal{M}^{\circledast} \times_{(\mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes})} \mathcal{M}^{\circledast} \times_{(\mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes})} \mathcal{M}^{\circledast} \times_{(\mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes})} \mathcal{N}^{\circledast} \xrightarrow{\mu^{\times} (\mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes})} \mathcal{M}^{\circledast} \times_{(\mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes})} \mathcal{M}^{\varepsilon} \times_{(\mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes})} \mathcal{M}^{\varepsilon} \times_{(\mathcal{V}^{\otimes} \times \mathcal{W}^{\otimes})} \mathcal{M}^{\varepsilon} \times_{(\mathcal{V}^{$ 

**Lemma 5.73.** Let  $\mathcal{D}$  be a presentable  $\infty$ -category and  $\mathcal{C} \subset \mathcal{D}$  a full subcategory such that the embedding  $\iota : \mathcal{C} \subset \mathcal{D}$  admits a right adjoint  $G : \mathcal{D} \to \mathcal{C}$ . The following are equivalent:

- (1) The functor  $\iota : \mathbb{C} \subset \mathbb{D}$  admits a left adjoint  $F : \mathbb{D} \to \mathbb{C}$  such that the composition  $\iota \circ F : \mathbb{D} \to \mathbb{C} \subset \mathbb{D}$  is an accessible functor.
- (2) The full subcategory  $\mathcal{C}$  is closed in  $\mathcal{D}$  under small limits and  $G: \mathcal{D} \to \mathcal{C}$  is an accessible functor.
- (3) The functor  $\iota \circ G : \mathcal{D} \to \mathcal{C} \subset \mathcal{D}$  is accessible and preserves small limits.

In particular, if (1) or (2) holds,  $\mathcal{C}$  is presentable.

*Proof.* The  $\infty$ -category  $\mathcal{C}$  admits small limits and small colimits being a colocalization of an  $\infty$ -category that admits small limits and small colimits. Moreover small limits in  $\mathcal{C}$  are coreflected via G from small limits in  $\mathcal{D}$ . So (2) and (3) are equivalent.

If (1) holds,  $\mathcal{C}$  is closed in  $\mathcal{D}$  under small limits. If (1) holds,  $\mathcal{C} \subset \mathcal{D}$  is an accessible localization so that  $\mathcal{C}$  is presentable. Hence  $G : \mathcal{D} \to \mathcal{C}$  is an accessible functor as a right adjoint functor between presentable  $\infty$ -categories.

Assume that (2) holds. The composition  $R := \iota \circ G : \mathcal{D} \to \mathcal{C} \subset \mathcal{D}$  is an accessible functor that preserves small limits. Since  $\mathcal{D}$  is presentable,  $R : \mathcal{D} \to \mathcal{D}$  admits a left adjoint  $T : \mathcal{D} \to \mathcal{D}$  by the adjoint functor theorem [17, Corollary 5.5.2.9]. Let  $\varepsilon : R = \iota \circ G \to id_{\mathcal{D}}$  be the counit. Because  $\iota : \mathcal{C} \subset \mathcal{D}$  is fully faithful, the unit  $id_{\mathcal{C}} \to G \circ \iota$  is an equivalence so that the composition  $G \circ \epsilon : G \circ \iota \circ G \to G$  and thus  $R \circ \varepsilon : R \circ R \to R$  are equivalences by the triangle identities.

Let X, Y be objects of  $\mathcal{D}$ . The map  $\mathcal{D}(T(X), R(Y)) \to \mathcal{D}(T(X), Y)$  induced by the counit  $\varepsilon(Y) : R(Y) \to Y$  is equivalent to the map

$$\mathcal{D}(\mathrm{T}(\mathrm{X}),\mathrm{R}(\mathrm{Y})) \simeq \mathcal{D}(\mathrm{X},\mathrm{R}\circ\mathrm{R}(\mathrm{Y})) \to \mathcal{D}(\mathrm{X},\mathrm{R}(\mathrm{Y})) \simeq \mathcal{D}(\mathrm{T}(\mathrm{X}),\mathrm{Y})$$

induced by the counit  $R(\varepsilon(Y)) : R \circ R(Y) \to R(Y)$  and thus is an equivalence. Thus T(X) is a colocal object of  $\mathcal{D}$  and so belongs to  $\mathcal{C}$ . So  $T : \mathcal{D} \to \mathcal{D}$  factors as  $F : \mathcal{D} \to \mathcal{C} \subset \mathcal{D}$ . Since  $T : \mathcal{D} \to \mathcal{D}$  is left adjoint to  $R : \mathcal{D} \to \mathcal{D}$ , the functor  $F : \mathcal{D} \to \mathcal{C}$  is left adjoint to  $\iota \simeq R \circ \iota : \mathcal{C} \to \mathcal{D}$ .

**Proposition 5.74.** Let  $\mathcal{V}^{\otimes} \to \operatorname{Ass}, \mathcal{W}^{\otimes} \to \operatorname{Ass}$  be  $\infty$ -operads such that  $\mathcal{V}, \mathcal{W}$  are empty or preadditive or  $\mathcal{V}^{\otimes} = \mathcal{W}^{\otimes} = \mathcal{S}^{\times}$ . The embeddings  $_{\mathcal{V}}\operatorname{Preadd}_{\mathcal{W}} \subset _{\mathcal{V}}^{\mathrm{II}}\operatorname{BEnr}_{\mathcal{W}}, _{\mathcal{V}}\operatorname{Preadd}_{\mathcal{W}}^{\wedge} \subset _{\mathcal{V}}^{\mathrm{II}}\operatorname{BEnr}_{\mathcal{W}}^{\wedge}$  admit a left and right adjoint.

Proof. By Proposition 5.69 the embedding  $_{\mathcal{V}}$ Preadd<sub>W</sub> ⊂  $_{\mathcal{V}}^{\mathrm{H}}$ BEnr<sub>W</sub> admits a right adjoint G such that  $\underset{\mathcal{V}}{\overset{\mathrm{H}}{\mathrm{BEnr}_{\mathcal{W}}} \xrightarrow{\mathrm{G}} _{\mathcal{V}}$ Preadd<sub>W</sub> ⊂  $\underset{\mathcal{V}}{\overset{\mathrm{H}}{\mathrm{BEnr}_{\mathcal{W}}}$  preserves small limits and filtered colimits. By Corollary 4.18 the ∞-category  $\underset{\mathcal{V}}{\overset{\mathrm{H}}{\mathrm{BEnr}_{\mathcal{W}}}$  is presentable. We apply Lemma 5.73 to deduce the existence of a left adjoint of the embedding  $_{\mathcal{V}}$ Preadd<sub>W</sub> ⊂  $\underset{\mathcal{V}}{\overset{\mathrm{H}}{\mathrm{BEnr}_{\mathcal{W}}}$ . By the same proposition the embedding  $_{\mathcal{V}}$ Preadd $_{\mathcal{W}}^{\wedge} \subset \underset{\mathcal{V}}{\overset{\mathrm{H}}{\mathrm{BEnr}_{\mathcal{W}}}$  admits a right adjoint G such that  $\underset{\mathcal{V}}{\overset{\mathrm{H}}{\mathrm{BEnr}_{\mathcal{W}}} \xrightarrow{\mathrm{G}} _{\mathcal{V}}$ Preadd $_{\mathcal{W}}^{\wedge} \subset \underset{\mathcal{V}}{\overset{\mathrm{H}}{\mathrm{BEnr}_{\mathcal{W}}}$  preserves small limits and filtered colimits since for every Cauchy-complete bi-enriched ∞-category  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  the bi-enriched ∞-category ( $\mathcal{M}^{\mathrm{Comm}}$ )<sup> $\circledast</sup> \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  is Cauchy-complete (Proposition 3.94) and the full bi-enriched subcategory Cmon( $\mathcal{M}$ )<sup> $\circledast</sup> \to \mathcal{V}^{\circledast} \times \mathcal{W}^{\circledast}$  of the latter is closed under abosulte colimits and so Cauchy-complete, too. By Corollary 4.18 the ∞-category  $\underset{\mathcal{V}}{\overset{\mathrm{H}}{\mathrm{BEnr}_{\mathcal{W}}^{\wedge}}$  is presentable. We apply Lemma 5.73 to deduce the existence of a left adjoint of the embedding  $_{\mathcal{V}}$ Preadd $_{\mathcal{W}}^{\wedge} \subset \underset{\mathcal{V}}{\overset{\mathrm{H}}{\mathrm{BEnr}_{\mathcal{W}}^{\wedge}}$ .</sup></sup>

Notation 5.75. Let  $\mathcal{V}^{\boxtimes} \to \mathbb{E}_{n+1}$  be a presentably  $\mathbb{E}_{n+1}$ -monoidal  $\infty$ -category corresponding to an  $\mathbb{E}_n$ -algebra in PrMon.

- (1) Let  $\stackrel{\amalg}{_{\mathcal{V}}} \operatorname{LEnr}^{\otimes} := \operatorname{vLEnr}(\operatorname{\mathcal{F}in})^{\otimes}$ .
- (2) Let  $\overset{\text{il}}{\mathcal{V}} \text{LEnr}^{\wedge,\otimes} \coloneqq_{\mathcal{V}} \text{LEnr}(\mathcal{F}\text{in} \cup \{\text{absolute } \mathcal{V}\text{-enriched weights}\})^{\otimes}$ .
- (3) Let  $_{\mathcal{V}}$ Preadd<sup> $\otimes$ </sup>  $\subset \underset{\mathcal{V}}{\overset{\text{II}}{\overset{\text{}}{\overset{}}{\overset{}}}}$  LEnr<sup> $\otimes$ </sup> be the full suboperad spanned by the preadditive left  $\mathcal{V}$ -enriched  $\infty$ -categories.
- (4) Let  $_{\mathcal{V}}\operatorname{Preadd}^{\wedge,\otimes} \subset _{\mathcal{V}}^{\mathrm{II}}\operatorname{Enr}^{\wedge,\otimes}$  be the full suboperad spanned by the preadditive left  $\mathcal{V}$ -enriched  $\infty$ -categories.
- (5) For  $\mathcal{V}^{\otimes} = \mathcal{S}^{\times}$  we drop  $\mathcal{V}$  from the notation.

**Proposition 5.76.** Let  $n \ge 0$  and  $\mathcal{V}^{\boxtimes} \to \mathbb{E}_{n+1}$  a preadditive presentably  $\mathbb{E}_{n+1}$ -monoidal  $\infty$ -category corresponding to an  $\mathbb{E}_n$ -algebra in PrMon or  $\mathcal{V}^{\otimes} = S^{\times}$ .

- (1) The embedding  $_{\mathcal{V}}\operatorname{Preadd}^{\otimes} \subset {}_{\mathcal{V}}^{\mathrm{U}}\operatorname{LEnr}^{\otimes}$  admits a left adjoint relative to  $\mathbb{E}_{\mathrm{n}}$  and the embedding  $_{\mathcal{V}}\operatorname{Preadd} \subset {}_{\mathcal{V}}^{\mathrm{U}}\operatorname{LEnr}$  admits a right adjoint.
- (2) The embedding  $_{\mathcal{V}}\operatorname{Preadd}^{\wedge,\otimes} \subset \overset{\amalg}{_{\mathcal{V}}}\operatorname{LEnr}^{\wedge,\otimes}$  admits a left adjoint relative to  $\mathbb{E}_{n}$  and the embedding  $_{\mathcal{V}}\operatorname{Preadd}^{\wedge} \subset \overset{\amalg}{_{\mathcal{V}}}\operatorname{LEnr}^{\wedge}$  admits a right adjoint.

*Proof.* In view of Proposition 5.74 it is enough to see that for every (Cauchy-complete) left  $\mathcal{V}$ -enriched  $\infty$ -categories  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast}, \mathcal{N}^{\circledast} \to \mathcal{V}^{\circledast}$  having finite conical coproducts the internal hom from  $\mathcal{M}^{\circledast} \to \mathcal{V}^{\circledast}$  to  $\mathcal{N}^{\circledast} \to \mathcal{V}^{\circledast}$  in  $\overset{\mathrm{II}}{_{\mathcal{V}}} \mathrm{LEnr}, \overset{\mathrm{II}}{_{\mathcal{V}}} \mathrm{LEnr}^{\wedge}$ , respectively, is preadditive if  $\mathcal{N}^{\circledast} \to \mathcal{V}^{\circledast}$  is preadditive. This follows from the description of the internal hom (Theorem 4.60) and Remark 5.63.

**Corollary 5.77.** Let  $n \ge 0$  and  $\mathcal{V}^{\boxtimes} \to \mathbb{E}_{n+1}$  a preadditive presentably  $\mathbb{E}_{n+1}$ -monoidal  $\infty$ -category corresponding to an  $\mathbb{E}_n$ -algebra in PrMon or  $\mathcal{V}^{\otimes} = S^{\times}$ . Then  $_{\mathcal{V}}$ Preadd $^{\otimes} \to \mathbb{E}_n, _{\mathcal{V}}$ Preadd $^{\wedge, \otimes} \to \mathbb{E}_n$  are preadditive presentably  $\mathbb{E}_n$ -monoidal  $\infty$ -categories.

*Proof.* This follows from Proposition 5.74 since  ${}^{\text{II}}_{\mathcal{V}} \text{LEnr}^{\otimes} \to \mathbb{E}_n$ ,  ${}^{\text{II}}_{\mathcal{V}} \text{LEnr}^{\wedge,\otimes} \to \mathbb{E}_n$  are presentably  $\mathbb{E}_n$ -monoidal  $\infty$ -categories by Theorem 4.42, which are preadditive by Proposition 5.72 and the fact that  ${}^{\nu}_{\mathcal{V}}\text{Preadd}^{\wedge}$  is closed in  ${}^{\nu}_{\mathcal{V}}\text{Preadd}$  under finite products.

**Definition 5.78.** Let  $\mathcal{V}^{\boxtimes} \to \text{Comm}$  be a preadditive presentably symmetric monoidal  $\infty$ -category or  $\mathcal{V}^{\otimes} = S^{\times}$ . Let  $n \ge 1$ . Let

$$_{\mathcal{V}}\operatorname{Preadd}(1)^{\otimes} \coloneqq _{\mathcal{V}}\operatorname{Preadd}^{\otimes}, \ _{\mathcal{V}}\operatorname{Preadd}(n+1)^{\otimes} \coloneqq _{_{\mathcal{V}}\operatorname{Preadd}(n)}\operatorname{Preadd}^{\otimes},$$

 $_{\mathcal{V}}$ Preadd<sup>(1)</sup>  $\approx _{\mathcal{V}}$ Preadd<sup>(3)</sup>,  $_{\mathcal{V}}$ Preadd<sup>(1)</sup>  $\approx _{\mathcal{V}}$ Preadd<sup>(1)</sup>  $\approx _{\mathcal{V}}$ Preadd<sup>(3)</sup>.

For  $\mathcal{V}^{\otimes}=\mathcal{S}^{\times}$  we drop  $\mathcal{V}$  from the notation and call

$$Preadd(n) \coloneqq {}_{S}Preadd(n)$$

the  $\infty$ -category of n-preadditive  $\infty$ -categories and

$$Preadd^{(n)} := {}_{S}Preadd^{(n)}$$

the  $\infty$ -category of Cauchy-complete n-preadditive  $\infty$ -categories.

**Corollary 5.79.** Let  $\mathcal{V}^{\boxtimes} \to \text{Comm}$  be a preadditive presentably symmetric monoidal  $\infty$ -category or  $\mathcal{V}^{\otimes} = S^{\times}$  and  $n \ge 1$ . Then  $_{\mathcal{V}}\text{Preadd}(n)^{\otimes} \to \text{Comm}, _{\mathcal{V}}\text{Preadd}^{\wedge}(n)^{\otimes} \to \text{Comm}$  are preadditive presentably symmetric monoidal  $\infty$ -categories. In particular,  $_{\mathcal{V}}\text{Preadd}(n)$  is enriched in itself and so belongs to  $_{\mathcal{V}}\text{Preadd}(n+1)$  and  $_{\mathcal{V}}\text{Preadd}^{\wedge}(n)$  is enriched in itself and so belongs to  $_{\mathcal{V}}\text{Preadd}^{\wedge}(n+1)$ .

**Notation 5.80.** Let  $\mathsf{Cat}_{\infty}^{\mathrm{ex},\wedge} \subset \mathsf{Cat}_{\infty}^{\mathrm{ex}} \subset \mathsf{Cat}_{\infty} \simeq {}_{S}\mathsf{LEnr}$  be the subcategories of (idempotent complete) stable  $\infty$ -categories and finite colimits preserving functors.

**Remark 5.81.** The  $\infty$ -categories  $Cat_{\infty}^{ex}, Cat_{\infty}^{ex,\wedge}$  are preadditive because they are full subcategories closed under finite products of the  $\infty$ -category  $_{s}LEnr(\{finite \ \infty-categories\})$  of small  $\infty$ -categories having finite colimits and functors preserving finite colimits, which is a preadditive  $\infty$ -category.

Notation 5.82. Let  $(Cat_{\infty}^{ex,\wedge})^{\otimes} \subset (Cat_{\infty}^{ex})^{\otimes} \subset {}_{s}LEnr(\{\text{finite } \infty-\text{categories}\})^{\otimes}$  be the full symmetric suboperads spanned by the (idempotent complete) stable  $\infty$ -categories.

By [10, Lemma 8.15.] the embeddings  $(Cat_{\infty}^{ex,\wedge})^{\otimes} \subset (Cat_{\infty}^{ex})^{\otimes} \subset {}_{\mathbb{S}}LEnr(\{\text{finite } \infty-\text{categories}\})^{\otimes} \text{ admit a left adjoint relative to Comm so that the symmetric } \infty\text{-operads } (Cat_{\infty}^{ex})^{\otimes} \rightarrow \text{Comm}, (Cat_{\infty}^{ex,\wedge})^{\otimes} \rightarrow \text{Comm are presentably symmetric monoidal } \infty\text{-categories}. The next definition for n = 2 is due to [19]:$ 

**Definition 5.83.** Let  $n \ge 2$ . The  $\infty$ -category of n-stable  $\infty$ -categories is

$$Cat_{(\infty,n)}^{ex} \coloneqq {}_{Cat_{\infty}^{ex}} Preadd(n-1)$$

The  $\infty$ -category of Cauchy-complete n-stable  $\infty$ -categories is

$$\mathsf{Cat}^{\mathrm{ex},\wedge}_{(\infty,n)} \coloneqq {}_{\mathsf{Cat}^{\mathrm{ex},\wedge}_{\infty}} \mathrm{Preadd}^{\wedge}(n-1).$$

For  $\mathcal{V} = \mathsf{Cat}_{\infty}^{\mathrm{ex}}, \mathsf{Cat}_{\infty}^{\mathrm{ex},\wedge}$  Corollary 5.79 gives the following corollary:

**Corollary 5.84.** Let  $n \ge 1$ . Then  $Cat_{(\infty,n)}^{ex}$  is n+1-stable and  $Cat_{(\infty,n)}^{ex,\wedge}$  is Cauchy-complete n+1-stable.

**Remark 5.85.** Let  $n \ge 2$ . By induction the inclusion  $Cat_{\infty}^{ex} \subset Cat_{\infty}$  gives rise to inclusions

(25) 
$$\operatorname{Cat}_{(\infty,n)}^{\operatorname{ex}}, \operatorname{Cat}_{(\infty,n)}^{\wedge,\operatorname{ex}} \subset \operatorname{Cat}_{(\infty,n)}.$$

Theorem 4.42 gives the following corollary:

**Corollary 5.86.** Let  $n \ge 2$ . The  $\infty$ -categories

# $\mathrm{Preadd}(n),\ \mathrm{Preadd}^{\wedge}(n),\ \mathsf{Cat}_{(\infty,n)}^{\mathrm{ex}},\mathsf{Cat}_{(\infty,n)}^{\wedge,\mathrm{ex}}$

carry closed symmetric monoidal structures such that the inclusions (25) are lax symmetric monoidal, where the right hand sides carry the cartesian structure.

#### References

- Benjamin Antieau and David Gepner. Brauer groups and étale cohomology in derived algebraic geometry. Geom. Topol., 18(2):1149–1244, 2014.
- [2] Ko Aoki. Tensor triangular geometry of filtered objects and sheaves. Mathematische Zeitschrift, 303:1–27, 2020.
- [3] Shay Ben-Moshe. Naturality of the ∞-categorical enriched yoneda embedding. Journal of Pure and Applied Algebra, 228(6):107625, 2024.
- [4] Andrew Blumberg, David Gepner, and Goncalo Tabuada. A universal characterization of higher algebraic k-theory. *Geometry and Topology*, 17, 01 2010.
- [5] David Gepner, Moritz Groth, and Thomas Nikolaus. Universality of multiplicative infinite loop space machines. Algebraic & Geometric Topology, 15(6):3107–3153, Dec 2015.
- [6] David Gepner and Rune Haugseng. Enriched ∞-categories via non-symmetric ∞-operads. Advances in Mathematics, 279:575 - 716, 2015.
- [7] David Gepner, Rune Haugseng, and Thomas Nikolaus. Lax colimits and free fibrations in ∞-categories. Documenta Mathematica, 22, 01 2015.
- [8] Rune Haugseng. ∞-operads via symmetric sequences. Mathematische Zeitschrift, 301, 05 2022.
- [9] Hadrian Heine. On bi-enriched  $\infty$ -categories. In Preparation.
- [10] Hadrian Heine. An equivalence between enriched  $\infty$ -categories and  $\infty$ -categories with weak action. Advances in Mathematics, 417:108941, 2023.
- [11] Hadrian Heine. A monadicity theorem for higher algebraic structures. arxiv preprint arXiv: 1712.00555, 2023.
- [12] Hadrian Heine, Alejo Lopez-Avila, and Markus Spitzweck. Infinity categories with duality and hermitian multiplicative infinite loop space machines. arxiv preprint arXiv: 1610.10043, 2020.
- [13] Vladimir Hinich. Dwyer-kan localization revisited. Homology, Homotopy and Applications, 18, 11 2013.
- [14] Vladimir Hinich. Yoneda lemma for enriched  $\infty$ -categories. Advances in Mathematics, 367:107129, 2020.
- [15] Vladimir Hinich. Colimits in enriched ∞-categories and day convolution. arxiv preprint arXiv: 2101.09538, 2021.
- [16] Jacob Lurie. Higher Algebra. available at http://www.math.harvard.edu/ lurie/.
- [17] Jacob Lurie. Higher topos theory, volume 170 of Annals of Mathematics Studies. Princeton University Press, Princeton, NJ, 2009.
- [18] Andrew W. Macpherson. The operad that co-represents enrichment. Homology Homotopy Appl., 23(1):387–401, 2021.
- [19] Aaron Mazel-Gee and Reuben Stern. A universal characterization of noncommutative motives and secondary algebraic k-theory. arxiv preprint arXiv: 2104.04021.
- [20] Joost Nuiten. On straightening for segal spaces. arxiv preprint arXiv:2108.11431, 2023.