

# Graphical proof of Ginibre's inequality

Yuki Tokushige\*

June 17, 2024

## Abstract

In this short note, we will give a new combinatorial proof of Ginibre's inequality for XY models. Our proof is based on multigraph representations introduced by van Engelenburg-Lis (2023) and a new bijection, which can be viewed as a combinatorial interpretation of Ginibre's square trick.

## 1 Introduction

For a finite graph  $\Lambda = (V, E)$  and coupling constants  $J = (J_{xy}) : E \rightarrow (0, \infty)$ , we consider the following Hamiltonian of the XY model: for  $\sigma = (\sigma_x)_{x \in V} \in (S^1)^V$ ,

$$H(\sigma) := - \sum_{xy \in E} J_{xy} \sigma_x \cdot \sigma_y = - \sum_{xy \in E} \frac{J_{xy}}{2} (\sigma_x \bar{\sigma}_y + \bar{\sigma}_x \sigma_y) = - \sum_{\vec{xy} \in \vec{E}} \frac{J_{xy}}{2} \sigma_x \bar{\sigma}_y,$$

where  $S^1 = \{z \in \mathbb{C} : |z| = 1\}$  and  $\vec{E}$  denotes the set of oriented edges of  $\Lambda$ . The corresponding Gibbs measure is defined by

$$d\mu_{J,\Lambda}(\sigma) := \frac{1}{\mathcal{Z}_{J,\Lambda}} e^{-H(\sigma)} d\sigma,$$

where  $d\sigma = \prod_{x \in V} d\sigma_x$  is the uniform probability measure on  $(S^1)^V$ . The normalizing constant  $\mathcal{Z}_{J,\Lambda} := \int_{(S^1)^V} e^{-H(\sigma)} d\sigma$  is called the partition function of the model, and by definition  $\mu_{J,\Lambda}$  is a probability measure on  $(S^1)^V$ . We will use a usual convention to denote the expectation with respect to  $\mu_{J,\Lambda}$  by  $\langle \cdot \rangle_{J,\Lambda}$ .

The XY models on  $\mathbb{Z}^2$  are known to exhibit very exotic phase transitions. This was originally proved by Fröhlich-Spencer [12]. Recently, van Engelenburg-Lis [9] and Aizenman-Harel-Peled-Schapiro [1] gave new proofs for this result using the relation between XY models and height models. See [14, 13, 16, 17] for more recent related works.

In order to state the result of this article, we introduce several notation. For any function  $\varphi : V \rightarrow \mathbb{Z}$ , we will write

$$\sigma^\varphi := \prod_{x \in V} \sigma_x^{\varphi_x}.$$

Furthermore, for  $\sigma = (\sigma_x), \sigma' = (\sigma'_x) \in (S^1)^V$  we define  $\theta = (\theta_x)_{x \in V}, \theta' = (\theta'_x)_{x \in V}$  as elements of  $[0, 2\pi)^V$  satisfying

$$\sigma_x = e^{i\theta_x}, \sigma'_x = e^{i\theta'_x}.$$

---

\*Technical University of Braunschweig, Institut für Mathematische Stochastik, 38106 Universitätsplatz 2 Braunschweig, Germany [yuki.tokushige@tu-braunschweig.de](mailto:yuki.tokushige@tu-braunschweig.de)

2020 *Mathematics Subject Classification.* 82B20.

*Key words and phrases.* XY model, correlation inequality, Ginibre's inequality, random current representation.

We then write

$$\varphi \cdot \boldsymbol{\theta} := \sum_{x \in V} \varphi_x \theta_x, \quad \varphi \cdot \boldsymbol{\theta}' := \sum_{x \in V} \varphi_x \theta'_x.$$

We finally state the claim we will show in this article.

**Theorem 1** ([15]). *For  $\varphi, \psi : V \rightarrow \mathbb{Z}$  with  $\sum_{v \in V} \varphi_v = \sum_{v \in V} \psi_v = 0$ , it holds that*

$$\langle \sigma^{\varphi+\psi} \rangle_{J, \Lambda} + \langle \sigma^{\varphi-\psi} \rangle_{J, \Lambda} \geq 2 \langle \sigma^\varphi \rangle_{J, \Lambda} \langle \sigma^\psi \rangle_{J, \Lambda}.$$

Since  $\sigma^\varphi + \sigma^{-\varphi} = 2 \cos(\varphi \cdot \boldsymbol{\theta})$ , this is equivalent to

$$\langle \cos(\varphi \cdot \boldsymbol{\theta}) \cos(\psi \cdot \boldsymbol{\theta}) \rangle_{J, \Lambda} \geq \langle \cos(\varphi \cdot \boldsymbol{\theta}) \rangle_{J, \Lambda} \langle \cos(\psi \cdot \boldsymbol{\theta}) \rangle_{J, \Lambda}.$$

The result is not new. It is nothing but the celebrated Ginibre's inequality shown in [15]. The aim of this article is to give a new combinatorial proof of Theorem 1, which is based on multigraph representations introduced in [8]. The novelty of this work is a new bijection that will play an important role in our proof. This bijection can be interpreted as an expression of a hidden combinatorial structure of the famous Ginibre's trick [15] from which Theorem 1 was originally derived. It seems to the author an interesting problem to investigate the possibility of applying our combinatorial idea to obtain correlation inequalities of XY models other than Ginibre's inequality.

This paper is organized as follows: in Section 2, we will briefly review random current and multigraph representations for the XY model. In Section 3, we will introduce a new bijection and prove Theorem 1. Finally in Section 4, we will explain how the bijection introduced in Section 3 is related to Ginibre's trick.

## Acknowledgements

The author would like to thank Diederik van Engelenburg for reading an earlier version of the manuscript and giving him suggestions which helped simplify notation and the proof. He would also like to thank Piet Lammers for letting him know multigraph representations introduced in [8]. He is grateful to Takashi Hara and Akira Sakai for their encouragements. He also thanks Tyler Helmuth for his comments.

## 2 Graphical representations and Ginibre's inequality

### 2.1 Random current representations

In this article, a  $\mathbb{Z}_{\geq 0}$ -valued function  $\mathbf{n}$  defined on the set of oriented edges  $\vec{E}$  will be called a *current*. For each current  $\mathbf{n}$ , we define its *amplitude*  $|\mathbf{n}| : E \rightarrow \mathbb{Z}_{\geq 0}$  by

$$|\mathbf{n}|_{xy} := \mathbf{n}_{x\vec{y}} + \mathbf{n}_{y\vec{x}}.$$

We define the *source function*  $\partial \mathbf{n}$  of  $\mathbf{n}$ , which is a  $\mathbb{Z}$ -valued function on  $V$ , by

$$\partial \mathbf{n}_x := \sum_{y \in V : y \sim x} (\mathbf{n}_{x\vec{y}} - \mathbf{n}_{y\vec{x}}).$$

These functions naturally arise in the study of XY models. To see this, we will compute the partition function  $\mathcal{Z}_{J, \Lambda}$  of an XY model with coupling constants  $(J_{xy})$  defined on a domain  $\Lambda$ . Let  $d\sigma = \prod_{x \in V} d\sigma_x$  be the uniform probability measure on  $[0, 2\pi)^V$  and define the weight  $w_J(\mathbf{n})$  of a current  $\mathbf{n}$  by

$$w_J(\mathbf{n}) := \prod_{x\vec{y} \in \vec{E}} \frac{\left(\frac{J_{xy}}{2}\right)^{\mathbf{n}_{x\vec{y}}}}{\mathbf{n}_{x\vec{y}}!}.$$

Then,

$$\begin{aligned}
\mathcal{Z}_{J,\Lambda} &:= \int_{[0,2\pi)^V} \exp \left[ \sum_{xy \in E} \frac{J_{xy}}{2} (\sigma_x \bar{\sigma}_y + \bar{\sigma}_x \sigma_y) \right] d\boldsymbol{\sigma} \\
&= \int_{[0,2\pi)^V} \exp \left[ \sum_{\vec{xy} \in \vec{E}} \frac{J_{xy}}{2} \sigma_x \bar{\sigma}_y \right] d\boldsymbol{\sigma} \\
&= \int_{[0,2\pi)^V} \prod_{\vec{xy} \in \vec{E}} \left( \sum_{\mathbf{n}_{\vec{xy}}=0}^{\infty} \frac{\left[ \frac{J_{xy}}{2} \sigma_x \bar{\sigma}_y \right]^{\mathbf{n}_{\vec{xy}}}}{\mathbf{n}_{\vec{xy}}!} \right) d\boldsymbol{\sigma} \\
&= \sum_{\mathbf{n}=(\mathbf{n}_{\vec{xy}})_{\vec{xy} \in \vec{E}}} w_J(\mathbf{n}) \int_{[0,2\pi)^V} \prod_{\vec{xy} \in \vec{E}} (\sigma_x \bar{\sigma}_y)^{\mathbf{n}_{\vec{xy}}} d\boldsymbol{\sigma} \\
&= \sum_{\mathbf{n}=(\mathbf{n}_{\vec{xy}})_{\vec{xy} \in \vec{E}}} w_J(\mathbf{n}) \prod_{x \in V} \int_{[0,2\pi)} \sigma_x^{\partial \mathbf{n}_x} d\sigma_x \\
&= \sum_{\partial \mathbf{n}=0} w_J(\mathbf{n}).
\end{aligned}$$

A straightforward generalization yields the following well-known equality, which gives random current representations of correlation functions. It is well-known that analogous representations for Ising models are very powerful tools to study critical phenomena [2, 7, 3, 6, 5, 10, 11, 4].

**Proposition 2.1.** *For any function  $\varphi : V \rightarrow \mathbb{Z}$  we have*

$$\langle \sigma^\varphi \rangle_{J,\Lambda} = \frac{1}{\mathcal{Z}_{J,\Lambda}} \sum_{\partial \mathbf{n}=-\varphi} w_J(\mathbf{n}) = \frac{1}{\mathcal{Z}_{J,\Lambda}} \sum_{\partial \mathbf{n}=\varphi} w_J(\mathbf{n}).$$

The second equality follows from the fact that  $w_J$  is invariant with respect to the map  $\mathbf{n}_{\vec{xy}} \mapsto \mathbf{n}_{y\vec{x}}$ .

## 2.2 Multigraph representations and Ginibre's inequality

In this subsection, we will relate a random current representation that appeared in Proposition 2.1 to a counting of directed multigraphs involving two colors, say, red and blue. This idea was used in [8, 9], where the authors analyzed the two dimensional XY model. We then reformulate Ginibre's inequality using this multigraph representation.

Let  $\varphi, \psi : V \mapsto \mathbb{Z}$ . By Proposition 2.1, we have

$$\begin{aligned}
\mathcal{Z}_{J,\Lambda}^2 \langle \sigma^\varphi \rangle_{J,\Lambda} &= \sum_{\substack{\partial \mathbf{n}=\varphi \\ \partial \mathbf{m}=0}} w_J(\mathbf{n}) w_J(\mathbf{m}) \\
&= \sum_{\mathbf{N}: E \rightarrow \mathbb{Z}_{\geq 0}} \sum_{\substack{\partial \mathbf{n}=\varphi \\ \partial \mathbf{m}=0 \\ |\mathbf{n}+\mathbf{m}|=\mathbf{N}}} \left[ \prod_{xy \in E} \frac{\left( \frac{J_{xy}}{2} \right)^{\mathbf{N}_{xy}}}{\mathbf{N}_{xy}!} \right] \left[ \prod_{xy \in E} \frac{\mathbf{N}_{xy}!}{\mathbf{n}_{\vec{xy}}! \mathbf{n}_{y\vec{x}}! \mathbf{m}_{\vec{xy}}! \mathbf{m}_{y\vec{x}}!} \right]. \tag{2.1}
\end{aligned}$$

In the same way, we get

$$\mathcal{Z}_{J,\Lambda}^2 \langle \sigma^\varphi \rangle_{J,\Lambda} \langle \sigma^\psi \rangle_{J,\Lambda} = \sum_{\substack{\partial \mathbf{n}=\varphi \\ \partial \mathbf{m}=\psi}} w_J(\mathbf{n}) w_J(\mathbf{m})$$

$$= \sum_{\mathbf{N}: E \rightarrow \mathbb{Z}_{\geq 0}} \sum_{\substack{\partial \mathbf{n} = \varphi \\ \partial \mathbf{m} = \psi \\ |\mathbf{n} + \mathbf{m}| = \mathbf{N}}} \left[ \prod_{xy \in E} \frac{\left(\frac{J_{xy}}{2}\right)^{\mathbf{N}_{xy}}}{\mathbf{N}_{xy}!} \right] \left[ \prod_{xy \in E} \frac{\mathbf{N}_{xy}!}{\mathbf{n}_{x\vec{y}}! \mathbf{n}_{y\vec{x}}! \mathbf{m}_{x\vec{y}}! \mathbf{m}_{y\vec{x}}!} \right]. \quad (2.2)$$

Let us consider a combinatorial interpretation of

$$\prod_{xy \in E} \frac{\mathbf{N}_{xy}!}{\mathbf{n}_{x\vec{y}}! \mathbf{n}_{y\vec{x}}! \mathbf{m}_{x\vec{y}}! \mathbf{m}_{y\vec{x}}!}. \quad (2.3)$$

To begin with, we define  $\mathbb{G}_{\mathbf{N}}$  to be an unoriented multigraph which, for each  $xy \in E$ , has  $\mathbf{N}_{xy}$  distinguishable edges connecting  $x$  and  $y$ . We then interpret (2.3) as the number of ways to paint edges of  $\mathbb{G}_{\mathbf{N}}$  in red and blue and give them orientations in such a way that:

- for each  $xy \in E$  there are  $|\mathbf{n}|_{xy}$  (resp.  $|\mathbf{m}|_{xy}$ ) red (resp. blue) edges.
- For each  $xy \in E$ , among  $|\mathbf{n}|_{xy}$  red edges connecting  $x$  and  $y$ , there are  $\mathbf{n}_{x\vec{y}}$  edges oriented from  $x$  to  $y$ , and  $\mathbf{n}_{y\vec{x}}$  edges oriented from  $y$  to  $x$ . The same property holds for blue edges when we replace  $\mathbf{n}$  with  $\mathbf{m}$ .

In what follows, for vertices  $x, y \in \mathbb{G}_{\mathbf{N}}$  we write  $\mathbf{r}_{x\vec{y}}$  (resp.  $\mathbf{b}_{x\vec{y}}$ ) for the number of red (resp. blue) edges in  $\mathbb{G}_{\mathbf{N}}$  (with colors and orientation) which are oriented from  $x$  to  $y$ . Then the following equalities follow from the above observations and (2.1), (2.2).

$$\mathcal{Z}_{J,\Lambda}^2 \langle \sigma^\varphi \rangle_{J,\Lambda} = \sum_{\mathbf{N}: E \rightarrow \mathbb{Z}_{\geq 0}} \left[ \prod_{xy \in E} \frac{\left(\frac{J_{xy}}{2}\right)^{\mathbf{N}_{xy}}}{\mathbf{N}_{xy}!} \right] \cdot \#\{\partial \mathbf{r} = \varphi, \partial \mathbf{b} = 0, |\mathbf{r} + \mathbf{b}| = \mathbf{N}\},$$

and

$$\mathcal{Z}_{J,\Lambda}^2 \langle \sigma^\varphi \rangle_{J,\Lambda} \langle \sigma^\psi \rangle_{J,\Lambda} = \sum_{\mathbf{N}: E \rightarrow \mathbb{Z}_{\geq 0}} \left[ \prod_{xy \in E} \frac{\left(\frac{J_{xy}}{2}\right)^{\mathbf{N}_{xy}}}{\mathbf{N}_{xy}!} \right] \cdot \#\{\partial \mathbf{r} = \varphi, \partial \mathbf{b} = \psi, |\mathbf{r} + \mathbf{b}| = \mathbf{N}\}. \quad (2.4)$$

For brevity of notation we will write  $\{\partial \mathbf{r} = \varphi, \partial \mathbf{b} = \psi\}_{\mathbf{N}}$  for  $\{\partial \mathbf{r} = \varphi, \partial \mathbf{b} = \psi, |\mathbf{r} + \mathbf{b}| = \mathbf{N}\}$ , and in the next section we will adopt the same rule for sets of oriented multigraphs involving only one color.

Thus, Theorem 1 will follow from

$$\#\{\partial \mathbf{r} = \varphi \pm \psi, \partial \mathbf{b} = 0\}_{\mathbf{N}} \geq 2\#\{\partial \mathbf{r} = \varphi, \partial \mathbf{b} = \psi\}_{\mathbf{N}} \quad (2.5)$$

for any  $\mathbf{N}: E \rightarrow \mathbb{Z}_{\geq 0}$ , where  $\varphi, \psi: V \rightarrow \mathbb{Z}$  are functions with mean zero and  $\psi \neq 0$ . In Section 3, we will prove (2.5).

### 3 Combinatorial proof of Ginibre's inequality

We will prove (2.5) by applying a certain bijection to

$$\#\{\partial \mathbf{r} = \varphi \pm \psi, \partial \mathbf{b} = 0\}_{\mathbf{N}} \quad \text{and} \quad \#\{\partial \mathbf{r} = \varphi, \partial \mathbf{b} = \psi\}_{\mathbf{N}}.$$

The bijection we will introduce in this section is inspired by the famous trick used in [15], from which Ginibre's inequality was derived. In Section 4, we will expand on this matter by showing computations that underlie combinatorial arguments we will present here.

### 3.1 Proof of (2.5)

For any  $f, g : V \rightarrow \mathbb{Z}$ , we will construct a bijection

$$F : \{\partial \mathbf{r} = f, \partial \mathbf{b} = g\}_{\mathbf{N}} \rightarrow \{\partial \mathbf{r} = f + g\}_{\mathbf{N}} \times \{\partial \mathbf{b} = -f + g\}_{\mathbf{N}}. \quad (3.1)$$

Once we have this bijection, it immediately follows that

$$\#\{\partial \mathbf{r} = f, \partial \mathbf{b} = g\}_{\mathbf{N}} = \#\{\partial \mathbf{r} = f + g\}_{\mathbf{N}} \cdot \#\{\partial \mathbf{b} = -f + g\}_{\mathbf{N}}.$$

We then obtain

$$\begin{aligned} & \#\{\partial \mathbf{r} = \varphi \pm \psi, \partial \mathbf{b} = 0\}_{\mathbf{N}} - 2\#\{\partial \mathbf{r} = \varphi, \partial \mathbf{b} = \psi\}_{\mathbf{N}} \\ &= \#\{\partial \mathbf{r} = \varphi + \psi\}_{\mathbf{N}} \cdot \#\{\partial \mathbf{b} = -\varphi - \psi\}_{\mathbf{N}} + \#\{\partial \mathbf{r} = \varphi - \psi\}_{\mathbf{N}} \cdot \#\{\partial \mathbf{b} = -\varphi + \psi\}_{\mathbf{N}} \\ & \quad - 2\#\{\partial \mathbf{r} = \varphi + \psi\}_{\mathbf{N}} \cdot \#\{\partial \mathbf{b} = -\varphi + \psi\}_{\mathbf{N}} \\ &= (\#\{\partial \mathbf{r} = \varphi + \psi\}_{\mathbf{N}})^2 + (\#\{\partial \mathbf{r} = \varphi - \psi\}_{\mathbf{N}})^2 - 2\#\{\partial \mathbf{r} = \varphi + \psi\}_{\mathbf{N}} \cdot \#\{\partial \mathbf{r} = \varphi - \psi\}_{\mathbf{N}} \\ &= (\#\{\partial \mathbf{r} = \varphi + \psi\}_{\mathbf{N}} - \#\{\partial \mathbf{r} = \varphi - \psi\}_{\mathbf{N}})^2 \geq 0, \end{aligned}$$

which proves (2.5).


### 3.2 Construction of a bijection (3.1)

Suppose that we have an element  $\omega \in \{\partial \mathbf{r} = f, \partial \mathbf{b} = g\}_{\mathbf{N}}$ . We define an element

$$(\omega_{\mathbf{R}}, \omega_{\mathbf{B}}) \in \{\partial \mathbf{r} = f + g\}_{\mathbf{N}} \times \{\partial \mathbf{b} = -f + g\}_{\mathbf{N}}$$

that corresponds to  $\omega$  in the following way: we define  $\omega_{\mathbf{R}} \in \{\partial \mathbf{r} = f + g\}_{\mathbf{N}}$  simply by painting all the edges in  $\omega$  in red. We then define  $\omega_{\mathbf{B}} \in \{\partial \mathbf{b} = -f + g\}_{\mathbf{N}}$  as follows: we retain blue edges in  $\omega$  and reverse orientations of red edges and paint all of them in blue. See Figure 1. It is straightforward to see that this defines a well-defined map

$$\begin{aligned} F : \{\partial \mathbf{r} = f, \partial \mathbf{b} = g\}_{\mathbf{N}} &\rightarrow \{\partial \mathbf{r} = f + g\}_{\mathbf{N}} \times \{\partial \mathbf{b} = -f + g\}_{\mathbf{N}} \\ \omega &\mapsto (\omega_{\mathbf{R}}, \omega_{\mathbf{B}}). \end{aligned}$$



**Figure 1:** A bijection between  $\{\partial \mathbf{r} = \partial \mathbf{b} = 0\}_{\mathbf{N}}$  and  $\{\partial \mathbf{r} = 0\}_{\mathbf{N}} \times \{\partial \mathbf{b} = 0\}_{\mathbf{N}}$

We next describe the map in the opposite direction. Suppose that we have an element  $(\omega_{\mathbf{R}}, \omega_{\mathbf{B}}) \in \{\partial \mathbf{r} = f + g\}_{\mathbf{N}} \times \{\partial \mathbf{b} = -f + g\}_{\mathbf{N}}$ . We then define a new configuration  $\tilde{\omega}$  in the following way: we first obtain an oriented colorless configuration  $\xi$  on  $\mathbb{G}_{\mathbf{N}}$  simply by forgetting the color of  $\omega_{\mathbf{R}}$ . The orientation of  $\tilde{\omega}$  is defined by  $\xi$ . We then paint edges in  $\xi$  according to the following rule: for each oriented edge  $\vec{e}$  in  $\xi$ , let  $e$  be

the corresponding unoriented edge in  $\mathbb{G}_{\mathbf{N}}$ . We then look at two oriented edges in  $\omega_{\mathbf{R}}$  and  $\omega_{\mathbf{B}}$  each of which corresponds to  $e$ . We paint  $\vec{e}$  in blue if they have the same direction, and paint  $\vec{e}$  in red otherwise.

That this operation defines a well-defined map from  $\{\partial \mathbf{r} = f + g\}_{\mathbf{N}} \times \{\partial \mathbf{b} = -f + g\}_{\mathbf{N}}$  to  $\{\partial \mathbf{r} = f, \partial \mathbf{b} = g\}_{\mathbf{N}}$  is verified by the following easy observation: let  $\mathbf{r}(\tilde{\omega})$  (resp.  $\mathbf{b}(\tilde{\omega})$ ) be the red (resp. blue) configuration in  $\tilde{\omega}$ . Then by definition, we have

$$\begin{aligned}\partial(\mathbf{r}(\tilde{\omega})) + \partial(\mathbf{b}(\tilde{\omega})) &= f + g, \\ -\partial(\mathbf{r}(\tilde{\omega})) + \partial(\mathbf{b}(\tilde{\omega})) &= -f + g,\end{aligned}$$

which implies that  $\tilde{\omega}$  is in  $\{\partial \mathbf{r} = f, \partial \mathbf{b} = g\}_{\mathbf{N}}$ . Now it is straightforward to verify  $F(\tilde{\omega}) = (\omega_{\mathbf{R}}, \omega_{\mathbf{B}})$ , which implies that

$$F : \{\partial \mathbf{r} = f, \partial \mathbf{b} = g\}_{\mathbf{N}} \rightarrow \{\partial \mathbf{r} = f + g\}_{\mathbf{N}} \times \{\partial \mathbf{b} = -f + g\}_{\mathbf{N}}$$

is indeed a bijection.

## 4 Relation to Ginibre's trick

In this section, we explain computations underlying what we discussed in Section 3. We begin with proving the following lemma, which will be used in subsequent computations.

**Definition 4.1.** • We introduce a subset  $\tilde{E}$  of the collection of all oriented edges  $\vec{E}$  in  $\Lambda$ , which is defined as follows: we choose a strict total order on  $V$  in an arbitrary way and fix it. For each  $pq \in E$ , we set

$$\tilde{pq} := \overrightarrow{(p \wedge q)(p \vee q)}.$$

By definition, we have  $\tilde{pq} = \tilde{qp}$ . We define  $\tilde{E} := \{\tilde{pq} \in \vec{E} : pq \in E\}$ . Then it holds that  $\#\tilde{E} = 2\#E = 2\#\tilde{E}$ . In what follows, we may identify functions on  $E$  and those on  $\tilde{E}$ .

- For  $\vec{e} = \overrightarrow{pq} \in \vec{E}$ , define  $\text{rev}(\vec{e}) := \overrightarrow{qp}$ .

**Lemma 4.2.** Let  $x, y, z, w \in V$  and  $\mathbf{N} : E \rightarrow \mathbb{Z}_{\geq 0}$ . Then there is a bijection between the following two sets.

1. The set of functions  $\mathbf{k} : \tilde{E} \rightarrow \mathbb{Z}_{\geq 0}$  satisfying  $0 \leq \mathbf{k} \leq \mathbf{N}$  and

$$(\delta_x - \delta_y) + (\delta_z - \delta_w) + \sum_{q \in V : \tilde{pq} \in \tilde{E}} (2\mathbf{k}_{\tilde{pq}} - \mathbf{N}_{\tilde{pq}}) + \sum_{q \in V : \tilde{qp} \in \tilde{E}} (\mathbf{N}_{\tilde{qp}} - 2\mathbf{k}_{\tilde{qp}}) = 0 \quad \text{for any } p \in V. \quad (4.1)$$

2. The set of functions  $\mathbf{r} : \vec{E} \rightarrow \mathbb{Z}_{\geq 0}$  satisfying  $|\mathbf{r}| = \mathbf{N}$  and  $\partial \mathbf{r} = -(\delta_x - \delta_y) - (\delta_z - \delta_w)$ .

*Proof.* Assume that we have a function  $\mathbf{k} : \tilde{E} \rightarrow \mathbb{Z}_{\geq 0}$  satisfying  $0 \leq \mathbf{k} \leq \mathbf{N}$  and (4.1). We define  $\mathbf{r} = (\mathbf{r}_{\vec{e}}) : \vec{E} \rightarrow \mathbb{N}$  by

$$\mathbf{r}_{\vec{e}} := \begin{cases} \mathbf{k}_{\vec{e}} & \text{if } \vec{e} \in \tilde{E}, \\ \mathbf{N}_{\text{rev}(\vec{e})} - \mathbf{k}_{\text{rev}(\vec{e})} & \text{if } \vec{e} \notin \tilde{E}, \end{cases}$$

Then it is obvious that  $|\mathbf{r}| = \mathbf{N}$ . Moreover, the condition (4.1) reads

$$(\delta_x - \delta_y) + (\delta_z - \delta_w) + \sum_{q \in V : p < q} (\mathbf{r}_{\tilde{pq}} - \mathbf{r}_{\tilde{qp}}) + \sum_{q \in V : p > q} (\mathbf{r}_{\tilde{pq}} - \mathbf{r}_{\tilde{qp}}) = 0 \quad \text{for any } p \in V.$$

which is exactly the condition  $\partial \mathbf{r} = -(\delta_x - \delta_y) - (\delta_z - \delta_w)$ .

Suppose that we are given a function  $\mathbf{r} : \vec{E} \rightarrow \mathbb{Z}_{\geq 0}$  with  $|\mathbf{r}| = \mathbf{N}$  and  $\partial \mathbf{r} = -(\delta_x - \delta_y) - (\delta_z - \delta_w)$ . Define  $\mathbf{k} : \tilde{E} \rightarrow \mathbb{Z}_{\geq 0}$  just by restriction. We can show by a similar argument that  $\mathbf{k}$  satisfies  $0 \leq \mathbf{k} \leq \mathbf{N}$  and (4.1).  $\square$

We finally discuss computations that inspired combinatorial arguments in Section 3. For simplicity, we set  $\varphi = \delta_a - \delta_b, \psi = \delta_c - \delta_d$  for  $a, b, c, d \in V$ . Recall that  $\boldsymbol{\theta} = (\theta_s)_{s \in V}, \boldsymbol{\theta}' = (\theta'_s)_{s \in V} \in [0, 2\pi)^V$  are defined by

$$\sigma_s = e^{i\theta_s}, \sigma'_s = e^{i\theta'_s}.$$

We begin with a review of the original proof of Ginibre's inequality given in [15]. We have

$$\begin{aligned} & \mathcal{Z}_{J,\Lambda}^2(\langle \sigma^{\varphi+\psi} \rangle_{J,\Lambda} + \langle \sigma^{\varphi-\psi} \rangle_{J,\Lambda}) \\ &= 2 \int_{([0,2\pi) \times [0,2\pi))^V} \cos(\theta_a - \theta_b) \cos(\theta_c - \theta_d) \exp \left[ \sum_{xy \in E} J_{xy} (\cos(\theta_x - \theta_y) + \cos(\theta'_x - \theta'_y)) \right] d\boldsymbol{\theta} d\boldsymbol{\theta}', \end{aligned}$$

and

$$\begin{aligned} & \mathcal{Z}_{J,\Lambda}^2 \langle \sigma^\varphi \rangle_{J,\Lambda} \langle \sigma^\psi \rangle_{J,\Lambda} \\ &= \int_{([0,2\pi) \times [0,2\pi))^V} \cos(\theta_a - \theta_b) \cos(\theta'_c - \theta'_d) \exp \left[ \sum_{xy \in E} J_{xy} (\cos(\theta_x - \theta_y) + \cos(\theta'_x - \theta'_y)) \right] d\boldsymbol{\theta} d\boldsymbol{\theta}', \quad (4.2) \end{aligned}$$

where  $d\boldsymbol{\theta}, d\boldsymbol{\theta}'$  are uniform probability measures on  $[0, 2\pi)^V$ . Since

$$\begin{aligned} & \exp \left[ \sum_{xy \in E} J_{xy} (\cos(\theta_x - \theta_y) + \cos(\theta'_x - \theta'_y)) \right] \\ &= \sum_{\mathbf{N}: E \rightarrow \mathbb{Z}_{\geq 0}} \prod_{xy \in E} \left[ \frac{(J_{xy})^{\mathbf{N}_{xy}}}{\mathbf{N}_{xy}!} \cdot (\cos(\theta_x - \theta_y) + \cos(\theta'_x - \theta'_y))^{\mathbf{N}_{xy}} \right], \end{aligned}$$

by comparing the coefficients of  $\prod_{xy} (J_{xy})^{\mathbf{N}_{xy}}$  in (2.4) and (4.2) we obtain

$$\begin{aligned} & \#\{\partial \mathbf{r} = \pm(\delta_a - \delta_b) \pm (\delta_c - \delta_d), \partial \mathbf{b} = 0\}_{\mathbf{N}} \\ &= 2^{2+\sum \mathbf{N}_{xy}} \int_{([0,2\pi) \times [0,2\pi))^V} \cos(\theta_a - \theta_b) \cos(\theta_c - \theta_d) \prod_{xy \in E} [\cos(\theta_x - \theta_y) + \cos(\theta'_x - \theta'_y)]^{\mathbf{N}_{xy}} d\boldsymbol{\theta} d\boldsymbol{\theta}', \text{ and} \\ & \#\{\partial \mathbf{r} = \pm(\delta_a - \delta_b), \partial \mathbf{b} = \pm(\delta_c - \delta_d)\}_{\mathbf{N}} \\ &= 2^{2+\sum \mathbf{N}_{xy}} \int_{([0,2\pi) \times [0,2\pi))^V} \cos(\theta_a - \theta_b) \cos(\theta'_c - \theta'_d) \prod_{xy \in E} [\cos(\theta_x - \theta_y) + \cos(\theta'_x - \theta'_y)]^{\mathbf{N}_{xy}} d\boldsymbol{\theta} d\boldsymbol{\theta}'. \quad (4.3) \end{aligned}$$

In the rest of this section, we will compute

$$\begin{aligned} & \int_{([0,2\pi) \times [0,2\pi))^V} (\cos(\theta_a - \theta_b) - \cos(\theta'_a - \theta'_b)) (\cos(\theta_c - \theta_d) - \cos(\theta'_c - \theta'_d)) \\ & \quad \cdot \prod_{xy \in E} [\cos(\theta_x - \theta_y) + \cos(\theta'_x - \theta'_y)]^{\mathbf{N}_{xy}} d\boldsymbol{\theta} d\boldsymbol{\theta}', \quad (4.4) \end{aligned}$$

which, by (4.3), is equal to

$$\begin{aligned} & 2 \cdot 2^{-2-\sum \mathbf{N}_{xy}} (\#\{\partial \mathbf{r} = \pm(\delta_a - \delta_b) \pm (\delta_c - \delta_d), \partial \mathbf{b} = 0\}_{\mathbf{N}} - \#\{\partial \mathbf{r} = \pm(\delta_a - \delta_b), \partial \mathbf{b} = \pm(\delta_c - \delta_d)\}_{\mathbf{N}}) \\ &= 4 \cdot 2^{-2-\sum \mathbf{N}_{xy}} (\#\{\partial \mathbf{r} = \delta_a - \delta_b \pm (\delta_c - \delta_d), \partial \mathbf{b} = 0\}_{\mathbf{N}} - \#\{\partial \mathbf{r} = \delta_a - \delta_b, \partial \mathbf{b} = \pm(\delta_c - \delta_d)\}_{\mathbf{N}}). \quad (4.5) \end{aligned}$$

Since  $\cos \eta + \cos \eta' = 2 \cos \frac{\eta+\eta'}{2} \cos \frac{\eta-\eta'}{2}$ ,  $\cos \eta - \cos \eta' = -2 \sin \frac{\eta+\eta'}{2} \sin \frac{\eta-\eta'}{2}$ , we have

$$(4.4) = 2^{2+\sum \mathbf{N}_{xy}} \int_{((-\pi, \pi]^2)^V} \left( \sin \frac{\theta_{ab} + \theta'_{ab}}{2} \sin \frac{\theta_{ab} - \theta'_{ab}}{2} \right) \cdot \left( \sin \frac{\theta_{cd} + \theta'_{cd}}{2} \sin \frac{\theta_{cd} - \theta'_{cd}}{2} \right) \\ \cdot \prod_{xy \in E} \left[ \cos \frac{\theta_{xy} + \theta'_{xy}}{2} \cos \frac{\theta_{xy} - \theta'_{xy}}{2} \right]^{\mathbf{N}_{xy}} d\boldsymbol{\theta} d\boldsymbol{\theta}',$$

where  $\theta_{zw} := \theta_z - \theta_w$ ,  $\theta'_{zw} := \theta'_z - \theta'_w$  for  $z, w \in V$ .

Define

$$F(\boldsymbol{\eta}) := \sin \frac{\eta_{ab}}{2} \sin \frac{\eta_{cd}}{2} \prod_{xy \in E} \left[ \cos \frac{\eta_{xy}}{2} \right]^{\mathbf{N}_{xy}}, \quad (4.6)$$

then we have

$$(4.4) = 2^{2+\sum \mathbf{N}_{xy}} \int_{((-\pi, \pi]^2)^V} F(\mathbf{u} + \mathbf{v}) F(\mathbf{u} - \mathbf{v}) \frac{d\mathbf{u}}{(2\pi)^{\#V}} \frac{d\mathbf{v}}{(2\pi)^{\#V}},$$

where  $d\mathbf{u}, d\mathbf{v}$  are Lebesgue measures on  $\mathbb{R}^V$ . Let  $\boldsymbol{\alpha} := \mathbf{u} + \mathbf{v}$ ,  $\boldsymbol{\beta} := \mathbf{u} - \mathbf{v}$ , then we have

$$(4.4) = 2^{2+\sum \mathbf{N}_{xy}} \cdot \left( \frac{1}{2} \right)^{\#V} \int_{\mathbb{D}^V} F(\boldsymbol{\alpha}) F(\boldsymbol{\beta}) \frac{d\boldsymbol{\alpha}}{(2\pi)^{\#V}} \frac{d\boldsymbol{\beta}}{(2\pi)^{\#V}},$$

where  $\mathbb{D} \subset \mathbb{R}^2$  is the square  $\{(2\pi, 0), (0, 2\pi), (-2\pi, 0), (0, -2\pi)\}$ . Taking account of the parity of sines and cosines, it is straightforward to check that the integrand  $F(\mathbf{u} + \mathbf{v})F(\mathbf{u} - \mathbf{v})$  is a periodic function with period  $2\pi$  in each coordinate of  $\mathbf{u}, \mathbf{v}$  though each of  $F(\mathbf{u} + \mathbf{v})$  and  $F(\mathbf{u} - \mathbf{v})$  is merely periodic with period  $4\pi$ . Therefore, we get

$$(4.4) = 2^{2+\sum \mathbf{N}_{xy}} \cdot \left( \frac{1}{2} \right)^{\#V} \int_{\mathbb{D}^V} F(\boldsymbol{\alpha}) F(\boldsymbol{\beta}) \frac{d\boldsymbol{\alpha}}{(2\pi)^{\#V}} \frac{d\boldsymbol{\beta}}{(2\pi)^{\#V}} \\ = 2^{2+\sum \mathbf{N}_{xy}} \cdot \left( \frac{1}{4} \right)^{\#V} \left( \int_{(-2\pi, 2\pi]^V} F(\boldsymbol{\eta}) \frac{d\boldsymbol{\eta}}{(2\pi)^{\#V}} \right)^2 \\ = 2^{2+\sum \mathbf{N}_{xy}} \cdot \left( \int_{(-2\pi, 2\pi]^V} F(\boldsymbol{\eta}) \frac{d\boldsymbol{\eta}}{(4\pi)^{\#V}} \right)^2 \geq 0,$$

where  $d\boldsymbol{\eta}$  is a Lebesgue measure on  $(-2\pi, 2\pi]^V$ , which means that  $\frac{d\boldsymbol{\eta}}{(4\pi)^{\#V}}$  is a uniform probability measure on the same space. This is a famous trick introduced in [15] which yields Ginibre's inequality. See [15, page 316-317] for more detailed discussions.

Therefore, in order to compute (4.4) it suffices to compute

$$\int_{((-2\pi, 2\pi]^V} \sin \frac{\eta_{ab}}{2} \sin \frac{\eta_{cd}}{2} \prod_{xy \in E} \left[ \cos \frac{\eta_{xy}}{2} \right]^{\mathbf{N}_{xy}} \frac{d\boldsymbol{\eta}}{(4\pi)^{\#V}}. \quad (4.7)$$

Letting  $\boldsymbol{\phi} = 2\boldsymbol{\eta}$ , we get

$$(4.7) = \int_{((-\pi, \pi]^V} \sin \phi_{ab} \sin \phi_{cd} \prod_{xy \in E} [\cos \phi_{xy}]^{\mathbf{N}_{xy}} d\boldsymbol{\phi},$$



where  $d\phi$  is a uniform probability measure (not a Lebesgue measure) on  $((-\pi, \pi])^V$ . Thus

$$\begin{aligned}
&= \int_{((-\pi, \pi])^V} \left( \frac{e^{\mathbf{i}(\phi_a - \phi_b)} - e^{-\mathbf{i}(\phi_a - \phi_b)}}{2\mathbf{i}} \right) \left( \frac{e^{\mathbf{i}(\phi_c - \phi_d)} - e^{-\mathbf{i}(\phi_c - \phi_d)}}{2\mathbf{i}} \right) \\
&\quad \cdot \prod_{xy \in E} \left[ \frac{e^{\mathbf{i}(\phi_x - \phi_y)} + e^{-\mathbf{i}(\phi_x - \phi_y)}}{2} \right]^{\mathbf{N}_{xy}} d\phi \\
&= -2^{-2-\sum \mathbf{N}_{xy}} \int_{((-\pi, \pi])^V} \left( e^{\mathbf{i}(\phi_a - \phi_b)} - e^{-\mathbf{i}(\phi_a - \phi_b)} \right) \left( e^{\mathbf{i}(\phi_c - \phi_d)} - e^{-\mathbf{i}(\phi_c - \phi_d)} \right) \\
&\quad \cdot \prod_{xy \in E} \left[ e^{\mathbf{i}(\phi_x - \phi_y)} + e^{-\mathbf{i}(\phi_x - \phi_y)} \right]^{\mathbf{N}_{xy}} d\phi.
\end{aligned}$$

Let us compute

$$\int_{((-\pi, \pi])^V} \left( e^{\mathbf{i}(\phi_a - \phi_b)} - e^{-\mathbf{i}(\phi_a - \phi_b)} \right) \left( e^{\mathbf{i}(\phi_c - \phi_d)} - e^{-\mathbf{i}(\phi_c - \phi_d)} \right) \cdot \prod_{pq \in E} \left[ e^{\mathbf{i}(\phi_x - \phi_y)} + e^{-\mathbf{i}(\phi_x - \phi_y)} \right]^{\mathbf{N}_{xy}} d\phi. \quad (4.8)$$

We first compute the integrand of (4.8).

$$\begin{aligned}
&\left( e^{\mathbf{i}(\phi_a - \phi_b)} - e^{-\mathbf{i}(\phi_a - \phi_b)} \right) \left( e^{\mathbf{i}(\phi_c - \phi_d)} - e^{-\mathbf{i}(\phi_c - \phi_d)} \right) \cdot \prod_{xy \in E} \left[ e^{\mathbf{i}(\phi_x - \phi_y)} + e^{-\mathbf{i}(\phi_x - \phi_y)} \right]^{\mathbf{N}_{xy}} \\
&= \sum_{0 \leq \mathbf{k} \leq \mathbf{N}} \binom{\mathbf{N}}{\mathbf{k}} \left( e^{\mathbf{i}(\phi_a - \phi_b)} - e^{-\mathbf{i}(\phi_a - \phi_b)} \right) \left( e^{\mathbf{i}(\phi_c - \phi_d)} - e^{-\mathbf{i}(\phi_c - \phi_d)} \right) \\
&\quad \cdot \prod_{xy \in E} \left( e^{\mathbf{i}(\phi_x - \phi_y)} \right)^{\mathbf{k}_{xy}} \left( e^{-\mathbf{i}(\phi_x - \phi_y)} \right)^{\mathbf{N}_{xy} - \mathbf{k}_{xy}} \\
&= \sum_{0 \leq \mathbf{k} \leq \mathbf{N}} \binom{\mathbf{N}}{\mathbf{k}} \left( e^{\mathbf{i}(\phi_a - \phi_b)} - e^{-\mathbf{i}(\phi_a - \phi_b)} \right) \left( e^{\mathbf{i}(\phi_c - \phi_d)} - e^{-\mathbf{i}(\phi_c - \phi_d)} \right) \\
&\quad \cdot \prod_{p \in V} \left( (e^{\mathbf{i}\varphi_p})^{\sum_{q \in V: \vec{p}\vec{q} \in \vec{E}} (2\mathbf{k}_{\vec{p}\vec{q}} - \mathbf{N}_{\vec{p}\vec{q}}) + \sum_{q \in V: \vec{q}\vec{p} \in \vec{E}} (\mathbf{N}_{\vec{p}\vec{q}} - 2\mathbf{k}_{\vec{p}\vec{q}})} \right) \\
&= \sum_{0 \leq \mathbf{k} \leq \mathbf{N}} \binom{\mathbf{N}}{\mathbf{k}} \left( e^{\pm \mathbf{i}(\phi_a - \phi_b) \pm \mathbf{i}(\phi_c - \phi_d)} - e^{\pm \mathbf{i}(\phi_a - \phi_b) \mp \mathbf{i}(\phi_c - \phi_d)} \right) \\
&\quad \cdot \prod_{p \in V} \left( (e^{\mathbf{i}\phi_p})^{\sum_{q \in V: \vec{p}\vec{q} \in \vec{E}} (2\mathbf{k}_{\vec{p}\vec{q}} - \mathbf{N}_{\vec{p}\vec{q}}) + \sum_{q \in V: \vec{q}\vec{p} \in \vec{E}} (\mathbf{N}_{\vec{p}\vec{q}} - 2\mathbf{k}_{\vec{p}\vec{q}})} \right) \\
&= \sum_{0 \leq \mathbf{k} \leq \mathbf{N}} \binom{\mathbf{N}}{\mathbf{k}} \prod_{p \in V} \left( (e^{\mathbf{i}\phi_p})^{\pm(\delta_a - \delta_b) \pm (\delta_c - \delta_d) + \sum_{q \in V: \vec{p}\vec{q} \in \vec{E}} (2\mathbf{k}_{\vec{p}\vec{q}} - \mathbf{N}_{\vec{p}\vec{q}}) + \sum_{q \in V: \vec{q}\vec{p} \in \vec{E}} (\mathbf{N}_{\vec{p}\vec{q}} - 2\mathbf{k}_{\vec{p}\vec{q}})} \right) \\
&\quad - \sum_{0 \leq \mathbf{k} \leq \mathbf{N}} \binom{\mathbf{N}}{\mathbf{k}} \prod_{p \in V} \left( (e^{\mathbf{i}\phi_p})^{\pm(\delta_a - \delta_b) \mp (\delta_c - \delta_d) + \sum_{q \in V: \vec{p}\vec{q} \in \vec{E}} (2\mathbf{k}_{\vec{p}\vec{q}} - \mathbf{N}_{\vec{p}\vec{q}}) + \sum_{q \in V: \vec{q}\vec{p} \in \vec{E}} (\mathbf{N}_{\vec{p}\vec{q}} - 2\mathbf{k}_{\vec{p}\vec{q}})} \right) \quad (4.9)
\end{aligned}$$

Recall that for any  $\mathbf{m} = (m_p) \in \mathbb{Z}^V$  we have

$$\int_{(-\pi, \pi]^V} \prod_p e^{\mathbf{i}m_p \phi_p} d\phi = \prod_p \left( \int_{(-\pi, \pi]} e^{\mathbf{i}m_p \phi_p} d\phi_p \right) = \begin{cases} 1 & \text{if } \mathbf{m} \equiv \mathbf{0}, \\ 0 & \text{otherwise.} \end{cases}$$

Therefore, by (4.9) and Lemma 4.2 we have

$$(4.8) = \#\{\partial \mathbf{r} = \pm(\delta_a - \delta_b) \pm (\delta_c - \delta_d)\}_{\mathbf{N}} - \#\{\partial \mathbf{r} = \pm(\delta_a - \delta_b) \mp (\delta_c - \delta_d)\}_{\mathbf{N}} \\ = 2\#\{\partial \mathbf{r} = \delta_a - \delta_b + \delta_c - \delta_d\}_{\mathbf{N}} - 2\#\{\partial \mathbf{r} = \delta_a - \delta_b - (\delta_c - \delta_d)\}_{\mathbf{N}},$$

which implies that

$$(4.4) = 4 \cdot 2^{-2-\sum \mathbf{N}_{xy}} \cdot (\{\partial \mathbf{r} = \delta_a - \delta_b + \delta_c - \delta_d\}_{\mathbf{N}} - \{\partial \mathbf{r} = \delta_a - \delta_b - (\delta_c - \delta_d)\}_{\mathbf{N}})^2. \quad (4.10)$$

By (4.5) and (4.10), we get

$$\#\{\partial \mathbf{r} = \delta_a - \delta_b \pm (\delta_c - \delta_d), \partial \mathbf{b} = 0\}_{\mathbf{N}} - \#\{\partial \mathbf{r} = \delta_a - \delta_b, \partial \mathbf{b} = \pm(\delta_c - \delta_d)\}_{\mathbf{N}} \\ = (\#\{\partial \mathbf{r} = \delta_a - \delta_b + \delta_c - \delta_d\}_{\mathbf{N}} - \#\{\partial \mathbf{r} = \delta_a - \delta_b - (\delta_c - \delta_d)\}_{\mathbf{N}})^2.$$

Furthermore, by computing

$$\int_{([0,2\pi) \times [0,2\pi))^V} \exp \left[ \sum_{xy \in E} J_{xy} (\cos(\theta_x - \theta_y) + \cos(\theta'_x - \theta'_y)) \right] d\theta d\theta'$$

in the same way, we can obtain

$$\#\{\partial \mathbf{r} = \partial \mathbf{b} = 0\}_{\mathbf{N}} = (\#\{\partial \mathbf{r} = 0\}_{\mathbf{N}})^2.$$

These equalities are exactly same as what we obtained by a bijective argument in Section 3.

## References

- [1] Harel M Peled R Aizenman, M and J. Schapiro. Depinning in integer-restricted gaussian fields and bkt phases of two-component spin models. *arXiv:2110.09498*.
- [2] M. Aizenman. Geometric analysis of  $\varphi^4$  fields and ising models. i, ii. *Comm. Math. Phys.*, 86(1):1–48, 1982.
- [3] M. Aizenman, D.J. Barsky, and R. Fernández. The phase transition in a general class of ising-type models is sharp. *J. Statist. Phys.*, 47(3-4):343–374, 1987.
- [4] M. Aizenman and H. Duminil-Copin. Marginal triviality of the scaling limits of critical 4d ising and  $\phi_4^4$  models. *Ann. of Math.*, 2(1):163–235, 2021.
- [5] M. Aizenman, H. Duminil-Copin, and V. Sidoravicius. Random currents and continuity of ising model’s spontaneous magnetization. *Comm. Math. Phys.*, 334(no. 2):719–742, 2015.
- [6] M. Aizenman and R. Fernández. On the critical behavior of the magnetization in high-dimensional ising models. *J. Statist. Phys.*, 44(3-4):393–454, 1986.
- [7] M. Aizenman and R. Graham. On the renormalized coupling constant and the susceptibility in  $\varphi_4^4$  field theory and the ising model in four dimensions. *Nuclear Phys. B*, 225(2):261–288, 1983.
- [8] M. Lis D. van Engelenburg. An elementary proof of phase transition in the planar xy model. *arXiv:2110.09465*.
- [9] M. Lis D. van Engelenburg. An elementary proof of phase transition in the planar xy model. *Comm. Math. Phys.*, 399(1):85–104, 2023.

- [10] H. Duminil-Copin and V. Tassion. A new proof of the sharpness of the phase transition for bernoulli percolation and the ising model. *Comm. Math. Phys.*, 343(2):725–745, 2016.
- [11] H. Duminil-Copin and V. Tassion. A new proof of the sharpness of the phase transition for bernoulli percolation on  $\mathbb{Z}^d$ . *Enseign. Math.*, 62(1-2):199–206, 2016.
- [12] J Fröhlich and T Spencer. The kosterlitz-thouless transition in two-dimensional abelian spin systems and the coulomb gas. *Comm. Math. Phys.*, 81(no. 4):527–602, 1981.
- [13] C Garban and A Sepúlveda. Statistical reconstruction of the gaussian free field and kt transition. *Journal of Eur. Math. Soc. (to appear)*.
- [14] C Garban and A Sepúlveda. Quantitative bounds on vortex fluctuations in 2d coulomb gas and maximum of the integer-valued gaussian free field. *Proc. London Math. Soc.*, 127(3):653–708, 2023.
- [15] J. Ginibre. General formulation of griffiths’ inequalities. *Comm. Math. Phys.*, 16:310–328., 1970.
- [16] P Lammers. Bijecting the bkt transition. *arXiv:2301.06905*.
- [17] P Lammers. A dichotomy theory for height functions a dichotomy theory for height functions a dichotomy theory for height functions. *arXiv:2211.14365*.