CTU-SCU/2024005

An alternative to purification in CFT

Xin Jiang,^{*} Peng Wang,[†] Houwen Wu,[‡] and Haitang Yang[§]

College of Physics, Sichuan University, Chengdu, 610065, China

(Dated: June 14, 2024)

In conformal field theories, in contrast to *adding* some auxiliary states into the bipartite mixed state ρ_{AB} as the usual purifications do, we show a pure entangled state ψ_{AB} can be constructed by *subtracting* the undetectable regions. In this pure state ψ_{AB} , the von Neumann entropy $S_{\rm vN}(A)$ naturally captures quantum entanglement between A and B. We verify that $S_{\rm vN}(A)$ is equal to the entanglement wedge cross-section E_W in AdS spacetime, which is conjectured to be the holographic dual of the entanglement of purification. We show such constructed entanglement entropy has a phase transition. The ordinary entanglement entropies of critical and non-critical QFTs are simply limits of the two phases.

INTRODUCTION

Entanglement has fundamental importance for quantum information and quantum gravity. In conformal field theories (CFTs), the entanglement entropy classifies quantum entanglement between complementary parts Aand B in a pure entangled state ψ_{AB} , defined by the von Neumann entropy

$$S_{\rm vN}(A) = -\mathrm{Tr}\rho_A \log \rho_A \tag{1}$$

for a reduced density matrix $\rho_A = \text{Tr}_B |\psi_{AB}\rangle \langle \psi_{AB}|$. However, for a mixed states $\rho_{AB} \neq |\psi_{AB}\rangle \langle \psi_{AB}|$, the entanglement entropy fails to characterize entanglement between A and B, as it yields the same values for entangled and unentangled states.

The typical resolution involves purifying AB by adding auxiliary systems \overline{AB} , known as purification. The entanglement of purification (EoP) [1] is defined by minimizing the entanglement entropy $S_{\rm vN}(AA : BB)$ over all possible purifications. Within the AdS/CFT correspondence [2–4], the EoP is conjectured to be equal to the entanglement wedge cross-section (EWCS) in the AdS bulk [5]. However, the conjecture EoP = EWCS is very hard to verify since in practice, carrying out the optimization over all possible purifications is almost Since the minimization of the EoP is not possible. very difficult, a canonical purification [6] is suggested as $|\sqrt{\rho_{AB}}\rangle \in \mathcal{H}_A \otimes \mathcal{H}_{\bar{A}} \otimes \mathcal{H}_B \otimes \mathcal{H}_{\bar{B}}$, where \bar{A} (\bar{B}) is the reflection of A (B). The reflected entropy $S_R(A:B)$ is then defined as the von Neumann entropy $S(\rho_{A\bar{A}})$. However, the reflected entropy still lacks a clear geometric interpretation, since the bulk dual of $|\sqrt{\rho_{AB}}\rangle$ is considered as the doubled entanglement wedge that remains elusive. Another operationally defined and generally computable candidate is the logarithmic negativity [7] whose explicit form is unknown [8], since it depends on the conformal block that is computationally expensive. Moreover, various measures of mixed states are defined in different ways [9–11], but they are all challenging to compute.

In this letter, for a bipartite mixed state ρ_{AB} , in con-

trast to *adding* extra auxiliary parts into the mixed state as the usual purifications do, we show by *subtracting* the undetectable regions in CFT₂, a pure entangled state ψ_{AB} can be constructed on the doubly connected plane (see Fig. 1). In this resulted pure state ψ_{AB} , it is natural to identify the well defined $S_{vN}(A)$ or $S_{vN}(B)$ as the entanglement measure between A and B. Our proposal is independent of purification or four-point conformal block data, distinguishing it from other bipartite entanglement measures. We verify our proposal holographically and find agreement with the EWCS in the AdS₃ bulk.

ENTANGLEMENT ENTROPY IN DOUBLY CONNECTED REGION

In a CFT₂, consider two disjoint subsystems $A = (a_2, b_1)$ and $B = (-\infty, a_1) \cup (b_2, \infty)$ at time $\tau = 0$, such that $a_1 < a_2 < b_1 < b_2$. The subsystems A and B are non-complementary parts typically in a mixed state ρ_{AB} (see the left panel of Fig. 1). In order to construct a pure state ψ_{AB} from the mixed state ρ_{AB} , we remove two discs in the Euclidean path integral region between A and B, obtaining a two-holed plane, topologically a doubly connected region, as depicted in Fig. 1. We then impose conformal invariant boundary conditions on two edges of the two-holed plane, corresponding to two boundary states $|a, b\rangle$ [12]. In this two-holed plane, two subsystems A and B are now complementary parts in a pure entangled state ψ_{AB} , allowing us to define the entanglement entropy $S_{\rm vN}(A)$ [13].

Given that every doubly connected region can be conformally mapped onto an annulus of the type illustrated in Fig. 2, we first calculate the entanglement entropy $S_{\rm vN}(A)$ between A and B in the annulus. To this end, it is useful to define the width of the annulus,

$$W = \log \frac{r_2}{r_1},\tag{2}$$

where r_1 and r_2 are the inner and outer radii, respectively. The entanglement entropy $S_{vN}(A)$ in the annulus



Figure 1. Left panel: At $\tau = 0$, two subsystems A and B on the x-axis are in a mixed state ρ_{AB} . Right panel: By removing the undetectable parts with two discs having different radii, we obtain a two-holed plane, in which two subsystems A and B are now in a pure entangled state ψ_{AB} . The blue shaded region represents the Euclidean path integral.



Figure 2. The two-holed plane can be conformally mapped onto an annulus with the inner radius r_1 and the outer radius r_2 .

is defined through the Rényi entropy:

$$S^{(n)}(A) = \frac{1}{1-n} \log \operatorname{Tr}_A \rho_A^n,$$
 (3)

$$S_{\rm vN}(A) = \lim_{n \to 1} S^{(n)}(A),$$
 (4)

where $\operatorname{Tr}_A \rho_A^n$ is given by

$$\operatorname{Tr}_A \rho_A^n = \frac{Z_n}{Z_1^n}.$$
(5)

Here, Z_1 is the partition function on a single cover of the annulus, and Z_n is the partition function on the *n*-sheeted cover \mathcal{M}_n obtained by sewing *n* copies of \mathcal{M} along *A*. Using the generator of scale transformations, the annulus partition function can be expressed as [12, 14, 15]:

$$Z_1 = e^{cW/12} \sum_k \langle a|k \rangle \langle k|b \rangle e^{-2\delta_k W}, \tag{6}$$

where c is the central charge, k denotes all allowed scalar operators with dimensions δ_k inside the annulus, and $|a,b\rangle$ are boundary states. Similarly,

$$Z_n = e^{cW/12n} \sum_k \langle a|k\rangle \langle k|b\rangle e^{-2\delta_k W/n}, \qquad (7)$$

since the replicated manifold \mathcal{M}_n is conformally equivalent to an annulus with the width $W_n = W/n$. As usual, only the ground state (k = 0) concerns us in calculating the entanglement entropy,

$$Z_1 = e^{cW/12} \langle a|0\rangle \langle 0|b\rangle, \tag{8}$$

$$Z_n = e^{cW/12n} \langle a|0\rangle \langle 0|b\rangle, \tag{9}$$

and $\operatorname{Tr}_A \rho_A^n$ read:

$$\operatorname{Tr}_{A}\rho_{A}^{n} = e^{\frac{c}{12}(\frac{1}{n}-n)W} \left(\langle a|0\rangle\langle 0|b\rangle\right)^{1-n}.$$
 (10)

Substituting this into the Rényi entropy expression, we obtain:

$$S^{(n)}(A) = \frac{c}{12} \left(1 + \frac{1}{n} \right) W + g_a + g_b,$$

where $g_{a,b} = \log \langle a, b | 0 \rangle$ are the Affleck-Ludwig boundary entropies [16], which encode the information of the undetectable regions and are irrelevant. Thus the entanglement between subsystems A and B is quantified by the universal term as

$$S_{\rm vN}(A:B) = \lim_{n \to 1} S^{(n)}(A) = \frac{c}{6}W.$$
 (11)

Now, we can map the u-plane with two holes onto the annulus in the w-plane via the conformal transformations:

$$v(u) = 2\frac{u-a_1}{a_2-a_1} - 1, \quad w(v) = \frac{v-\gamma}{\gamma v - 1},$$
 (12)

with

$$\gamma = \frac{1 + \alpha\beta + \sqrt{(1 - \alpha^2)(1 - \beta^2)}}{\alpha + \beta},$$

$$\alpha = v(b_1), \quad \beta = v(b_2). \tag{13}$$

The width of the annulus is

$$W = \log \frac{\alpha - \beta}{1 - \alpha\beta + \sqrt{(1 - \alpha^2)(1 - \beta^2)}}.$$
 (14)

In terms of the cross ratio

$$z = \frac{(a_2 - a_1)(b_2 - b_1)}{(b_1 - a_2)(b_2 - a_1)},$$
(15)

plugging (14) into (11), we get

$$S_{\rm vN}(A:B) = \frac{c}{6} \log\left[1 + \frac{2}{z} + 2\sqrt{\frac{1}{z}\left(\frac{1}{z} + 1\right)}\right].$$
 (16)

In the limit $b_1 - a_2 = \ell$ and $a_2 - a_1 = b_2 - b_1 = \epsilon \to 0$ as a UV regulator, this entanglement entropy simplifies to the famous single interval result

$$S_{\rm vN}(A:B) = \frac{c}{3}\log\frac{\ell}{\epsilon}.$$
 (17)



Figure 3. Left panel: Four subsystems ABCD in the *u*-plane. Right panel: by subtracting the undetectable parts A and B with two discs, we obtain an asymmetric annular region in which C and D are in a pure entangled state ψ_{CD} .

On the other hand, for the same system, it is illuminating to switch the roles of A, B and those of the undetectable regions C, D, as depicted in Fig. 3. So, we are now concerned with the entanglement between $C \in (a_1, a_2)$ and $D = (b_1, b_2)$, while A and B are undetectable regions. Following the similar procedure, we subtract A and B with two discs. The resulted manifold is an asymmetric annulus, where C and D become complementary parts in a pure entangled state ψ_{CD} .

We can map the asymmetric annular region in the uplane onto an annulus in the w-plane using the following conformal transformations:

$$\tilde{v}(u) = 2\frac{u-a_1}{b_2-a_1} - 1, \quad w(\tilde{v}) = \frac{\tilde{v}-\tilde{\gamma}}{\tilde{\gamma}\tilde{v}-1},$$
 (18)

with

$$\tilde{\gamma} = \frac{1 + \tilde{\alpha}\tilde{\beta} + \sqrt{(1 - \tilde{\alpha}^2)(1 - \tilde{\beta}^2)}}{\tilde{\alpha} + \tilde{\beta}},$$
(19)

$$\tilde{\alpha} = \tilde{v}(b_1), \quad \tilde{\beta} = \tilde{v}(a_2).$$
 (20)

The width of the annulus is

$$\tilde{W} = \log \frac{1 - \tilde{\alpha}\tilde{\beta} + \sqrt{(1 - \tilde{\alpha}^2)(1 - \tilde{\beta}^2)}}{\tilde{\alpha} - \tilde{\beta}}.$$
 (21)

In terms of the cross ratio (15), the entanglement entropy between C and D is

$$S_{\rm vN}(C:D) = \frac{c}{6}\tilde{W} \\ = \frac{c}{6}\log\left[1+2z+2\sqrt{z(z+1)}\right]. \quad (22)$$

In the limit $b_1 = -a_2 = \epsilon \to 0$ and $b_2 = -a_1 = \xi \to \infty$, we reproduce the half-space entanglement entropy

$$S_{\rm vN}(C:D) = \frac{c}{6}\log\frac{\xi}{\epsilon}.$$
 (23)



Figure 4. In the hyperbolic disk, a quadrilateral is bounded by colored geodesics, and the only two perpendicular geodesics (black and gray lines) satisfy the equation $\sinh \frac{\delta}{2} \sinh \frac{\sigma}{2} = 1$, where σ and δ denote their lengths.

HOLOGRAPHIC INTERPRETATION

Notably, $S_{vN}(C:D)$ is related to $S_{vN}(A:B)$ by

$$z \to \frac{1}{z},$$
 (24)

and satisfy

$$\sinh\left[\frac{3}{c}S_{\rm vN}(A:B)\right]\sinh\left[\frac{3}{c}S_{\rm vN}(C:D)\right] = 1.$$
 (25)

This equation is an exclusive constraint in hyperbolic geometry [17] for the only two perpendicular geodesics inside a quadrilateral, as depicted in Fig. 4. This observation is reminiscent of the AdS/CFT correspondence, where it has been conjectured that the minimal crosssection of the entanglement wedge E_W in the AdS bulk is a holographic dual of the EoP in a given bipartite mixed state [5]. With the same setup, the EWCS could be computed in the AdS₃ bulk, and we immediately find that the EWCS precisely agrees with the entanglement entropy $S_{\rm vN}$ in the doubly connected region:

$$E_W(A:B) = S_{vN}(A:B),$$
 (26)

$$E_W(C:D) = S_{vN}(C:D).$$
 (27)

Thus, the entanglement entropy $S_{\rm vN}$ in the doubly connected region should be regarded as an alternative to EoP.

In addition, our proposal provides a novel interpretation of the entanglement phase transition for mixed states. From Eq. (25), we identify a critical point:

$$S_{\rm vN}(A:B) = S_{\rm vN}(C:D) = S_{\rm vN}^*,$$
 (28)

where the critical value

$$S_{\rm vN}^* = \frac{c}{3} \operatorname{arcsinh} 1 = \frac{c}{6} \log \left(3 + 2\sqrt{2}\right)$$
 (29)

matches the phase transition point of the EWCS [5]. This entanglement phase transition is usually interpreted as the mutual information $I(A : B) = \frac{c}{6} \log z$ between two subsystems, vanishing when $z \leq 1$. We now see from our results this is precisely a transition of the dominant contribution between *s*-channel and *t*-channel [18]. It indicates that the entanglement transition near z = 1signifies a transition from entanglement between A and B to entanglement between C and D, or vice versa.

CONCLUSION AND DISCUSSION

In this letter, we proposed an alternative to purification for bipartite mixed states in CFT_2 . Our proposal is completely determined by the definition of von Neumann entropy and the replica trick, independent of any purification or four-point conformal block data. We confirmed the proposed $S_{\rm vN}$ is the dual of EWCS and equal to EoP.

Our choice of regulators, by removing discs, is *canonical*, although other choices with different shapes may exist. The width W is a *conformal invariant*; if two annuli have different widths, they cannot be conformally mapped onto each other. The set of all doubly connected regions falls into classes of conformally equivalent regions, with each class characterized by the width of that class. Therefore, any other possible regulators with different shapes must be conformally equivalent to our choice.

Our proposal suggests that the von Neumann entropy in the multi-connected regions may correspond to an entanglement measure for multipartite mixed states. Having successfully extracted the EWCS from the annulus CFT, it is of interest to further study the connections between the entanglement wedge and the CFT in multiconnected regions.

In [19], the EWCS for two parallel strips with equal widths l in AdS_{d+1} are given. It turns out our proposal can be used to find the dual entanglement entropy in CFT₄.

Acknowledgments. We are very grateful to Song He, Bo Ning, Ronak M Soni, Tadashi Takayanagi, Qiang Wen, Runqiu Yang and Yang Zhou for reading the first version of this work. Their comments or suggestions help us to improve the manuscript substantially. This work is supported in part by NSFC (Grant No. 12105191, 12275183 and 12275184).

- * domoki@stu.scu.edu.cn
- [†] pengw@scu.edu.cn
- [‡] iverwu@scu.edu.cn
- § hyanga@scu.edu.cn
- Barbara M. Terhal, Michal Horodecki, Debbie W. Leung, and David P. DiVincenzo. The entanglement of purification. J. Math. Phys., 43(9):4286-4298, 2002. arXiv:quant-ph/0202044, doi:10.1063/1.1498001.

- [2] Juan Martin Maldacena. The Large N limit of superconformal field theories and supergravity. Adv. Theor. Math. Phys., 2:231-252, 1998. arXiv:hep-th/9711200, doi:10.1023/A:1026654312961.
- [3] Edward Witten. Anti-de Sitter space and holography. Adv. Theor. Math. Phys., 2:253-291, 1998. arXiv: hep-th/9802150, doi:10.4310/ATMP.1998.v2.n2.a2.
- S. S. Gubser, Igor R. Klebanov, and Alexander M. Polyakov. Gauge theory correlators from noncritical string theory. *Phys. Lett. B*, 428:105–114, 1998.
 arXiv:hep-th/9802109, doi:10.1016/S0370-2693(98) 00377-3.
- [5] Tadashi Takayanagi and Koji Umemoto. Entanglement of purification through holographic duality. *Nature Phys.*, 14(6):573–577, 2018. arXiv:1708.09393, doi:10.1038/ s41567-018-0075-2.
- [6] Souvik Dutta and Thomas Faulkner. A canonical purification for the entanglement wedge cross-section. *JHEP*, 03:178, 2021. arXiv:1905.00577, doi:10.1007/ JHEP03(2021)178.
- [7] Pasquale Calabrese, John Cardy, and Erik Tonni. Entanglement negativity in extended systems: A field theoretical approach. J. Stat. Mech., 1302:P02008, 2013. arXiv: 1210.5359, doi:10.1088/1742-5468/2013/02/P02008.
- [8] Jonah Kudler-Flam and Shinsei Ryu. Entanglement negativity and minimal entanglement wedge cross sections in holographic theories. *Phys. Rev. D*, 99(10):106014, 2019. arXiv:1808.00446, doi:10.1103/PhysRevD.99.106014.
- Kotaro Tamaoka. Entanglement Wedge Cross Section from the Dual Density Matrix. Phys. Rev. Lett., 122(14):141601, 2019. arXiv:1809.09109, doi:10.1103/PhysRevLett.122.141601.
- [10] Ricardo Espíndola, Alberto Guijosa, and Juan F. Pedraza. Entanglement Wedge Reconstruction and Entanglement of Purification. *Eur. Phys. J. C*, 78(8):646, 2018. arXiv:1804.05855, doi:10.1140/epjc/ s10052-018-6140-2.
- Jonathan Harper and Matthew Headrick. Bit threads and holographic entanglement of purification. *JHEP*, 08:101, 2019. arXiv:1906.05970, doi:10.1007/JHEP08(2019) 101.
- [12] John L. Cardy. Boundary Conditions, Fusion Rules and the Verlinde Formula. Nucl. Phys. B, 324:581–596, 1989. doi:10.1016/0550-3213(89)90521-X.
- [13] The two holes can be equivalently understood as finite regulators.
- [14] John L. Cardy. Boundary conformal field theory. 11 2004. arXiv:hep-th/0411189.
- [15] John Cardy and Erik Tonni. Entanglement hamiltonians in two-dimensional conformal field theory. J. Stat. Mech., 1612(12):123103, 2016. arXiv:1608.01283, doi: 10.1088/1742-5468/2016/12/123103.
- [16] Ian Affleck and Andreas W. W. Ludwig. Universal noninteger 'ground state degeneracy' in critical quantum systems. *Phys. Rev. Lett.*, 67:161–164, 1991. doi: 10.1103/PhysRevLett.67.161.
- [17] Peter Buser. Geometry and spectra of compact Riemann surfaces. Birkhäuser Boston, 2010. doi:10.1007/ 978-0-8176-4992-0.
- [18] Thomas Hartman. Entanglement Entropy at Large Central Charge. 3 2013. arXiv:1303.6955.
- [19] Niko Jokela and Arttu Pönni. Notes on entanglement wedge cross sections. JHEP, 07:087, 2019. arXiv:1904. 09582, doi:10.1007/JHEP07(2019)087.