

Dynamics of Spinning Binary at 2PM

Gang Chen^a, Tianheng Wang^{b,c}

^a*Niels Bohr International Academy, Niels Bohr Institute, University of Copenhagen, Blegdamsvej 17, DK-2100 Copenhagen Ø, Denmark*

^b*Center for Theoretical Physics, Seoul National University, 1 Gwana-ro, Gwanak-gu, 08826, Seoul, South Korea*

^c*Institute of Theoretical Physics, Chinese Academy of Sciences, 55 Zhongguancun Road East, Haidian District, 100190, Beijing, China*

E-mail: gang.chen@nbi.ku.dk, tianhengwang@snu.ac.kr

ABSTRACT: We consider the covariant proposal for the gravitational Compton amplitude for a Kerr black hole. Employing the covariant three- and four-point Compton amplitudes, we assemble the classical one-loop integrand on the maximal cut at all orders in spin, utilizing the method of unitarity. Expanding in powers of spin, we evaluate the one-loop amplitude up to $\mathcal{O}(G^2 a^8)$. Supplemented with extra contact contributions derived from the far-zone data of the Teukolsky solutions, the one-loop amplitude is in agreement with results available in the literature. We showcase the classical eikonal in the aligned-spin case at $\mathcal{O}(G^2 a^7)$.

Contents

1	Introduction	1
2	Classical Compton amplitudes	3
2.1	Covariant Compton amplitude to arbitrary spin order	3
2.2	Spin expansion and analysis of contact contributions	5
3	Eikonal with spin at 2PM	8
3.1	Contribution from the covariant Compton amplitude	9
3.2	Contribution from extra contact terms	12
3.3	2PM eikonal in the spin expansion	13
4	Conclusions and outlook	15
A	Extra contact terms	17
B	UV behaviours	18
C	Decomposition of tensor integrals	19

1 Introduction

The successful detection of gravitational waves [1–5] has inspired an explosion of developments in the studies of black hole mergers. Besides numerical relativity [6–8] and the formal approaches focusing on the post-Newtonian expansion [9–28], various amplitude-based approaches [29–37] have proven highly effective in computing higher-order corrections in the Post-Minkowskian (PM) expansion and revealing the underlying structures of gravitational interactions.

These amplitude-based approaches view the dynamics of heavy bodies interacting via weak gravity as that of scattering processes in the classical regime where the gravitons are soft. A multitude of methods [38–55] have been devised to single out the classical order from the full (quantum) amplitudes. Techniques for computing loop amplitudes efficiently facilitate computations at high PM orders. At the point of writing, black hole scattering observables have been computed up to 4PM [36, 56–61] for both non-spinning and spinning binaries and partial results are obtained at 5PM in the non-spinning case [62].

Incorporating spin degrees of freedom is non-trivial, both conceptually and computationally. Challenges appear even at tree level. Although the three-point amplitudes of two massive particles and one graviton are classified in [63], complications enter at four points, namely the gravitational Compton scattering. On the one hand, fundamental questions in quantum field theories with higher-spin particles need to be addressed [64–69], towards a full understanding of these amplitudes. On the other hand, the concept of double copy [70–72] and general considerations on symmetry and locality requirements allow us to bootstrap the Compton amplitudes of arbitrary spins [73, 74], despite incomplete knowledge of the underlying theory. These results can then be tested against approaches developed from other perspectives such as the Teukolsky equation [75–83].

Based on somewhat different assumptions, the Compton amplitude for a Kerr black hole has been computed both as an expansion in powers of spin [84–97] and as a resummed function [73, 74, 98, 99]. They all agree up to the quartic order in the spin expansion, where the spin-shift symmetry is expected to hold. Going to higher orders in spin, they deviate from one another, as the nuances in their respective formalism and assumptions become important. These results provide a pool of data for future investigations to analyze and clarify, which in turn will help to reveal the structures of the underlying theory. The Compton amplitude for a Kerr black hole is also an important ingredient in the calculation of loop amplitudes, which describe the dynamics of Kerr binaries at higher PM orders [86, 87, 100–110].

A complementary line of research [111–118] based on the worldline quantum field theory (WQFT) formalism [42] provides a fast-track to high PM calculations for physical observables and generating functions, up to quadratic orders in spin. Recent development in the worldline formalism focuses on the radial action, which produces scattering observables at 6PM beyond the quadratic order in certain kinematic regions [119]. The gravitational waveform in the spinning case is discussed in [96, 107, 120] up to the quartic order in spin. Besides Einstein gravity, related works on the spinning binary systems in gauge [121] and other gravitational backgrounds [122, 123] have been carried out. Beyond conservative dynamics, tidal and radiation effects have been explored from both amplitude approaches and worldline-based formalisms [124–129]. Recent developments on the self-force expansion [130–134] may also shed light on the dynamics in curved backgrounds.

In this paper, we consider one particular proposal for the Compton amplitude [73, 74], which is obtained using the double-copy and bootstrap. In section 2, we review its structures and make comparisons with other proposals at the level of their spin expansions. In section 3, we compute the one-loop amplitude using this Compton amplitude from the leading singularity and the resulting 2PM eikonal phase.

2 Classical Compton amplitudes

We begin this section with a review of the classical three-point [35, 73, 135, 136] and four-point Compton amplitudes [73, 74] of two massive particles with spins minimally coupled to gravity at tree level. The Compton amplitude follows from the double-copy and factorization requirements and includes contact terms. The contact terms match with physical data at low orders in spin and display certain empirical properties, which allows for an extrapolation. This Compton amplitude is conjectured to hold at all orders in spin. In the second part of this section, we discuss the comparison with the Compton amplitudes computed from other approaches [83, 99]. The differences in the contact terms can be interpreted as being related to the internal structures of the Kerr black hole. Moreover, we demonstrate a simple procedure to obtain expressions in covariant variables for these extra contact terms.

2.1 Covariant Compton amplitude to arbitrary spin order

Here we review the structures of the classical amplitudes of a heavy particle emitting gravitons. Throughout this paper, we restrict our discussions to the kinematics in the heavy-mass limit [41]. The incoming and outgoing massive momenta typically are parameterised as $\bar{p} = mv$ and $\bar{p}' = (mv - q)$ where q^μ denotes the total momenta of the emitted graviton(s). The on-shell conditions $\bar{p}^2 = \bar{p}'^2 = (mv - q)^2 = m^2$ yield $v^2 = 1$ and $v \cdot q = 0$.

The three-point amplitude of two massive particles of arbitrary masses and spins and a graviton is first given in [135] by considering general symmetry constraints. Restricting to the minimal coupling and taking the heavy-mass limit, the amplitude corresponds to the Kerr black hole of an arbitrary classical spin. A compact, covariant form of this three-point amplitude is found in [35, 73, 136]

$$\begin{array}{c} \varepsilon_1, p_1 \\ \text{⋈} \\ \bar{p} \text{---} \bullet \text{---} \bar{p}' \end{array} \quad \mathcal{M}_3(1, \bar{p}', \bar{p}) = -i\kappa(\bar{p} \cdot \varepsilon_1)(\mathbf{w} \cdot \varepsilon_1), \quad (2.1)$$

where $\mathbf{w}^\mu := \cosh(x_1)\bar{p}^\mu - iG_1(x_1)(p_1 \cdot S)^\mu$, with $G_1(x_1) = \frac{\sinh(x_1)}{x_1}$, $x_1 = a \cdot p_1$ and $(p_1 \cdot S)^\mu = p_{1\nu} S^{\nu\mu}$. Here $S^{\mu\nu}$ denotes the spin tensor which describes the classical spin of the black hole and we have introduced the spin-length vector a^μ , with $S^{\mu\nu} = -\epsilon^{\mu\nu\rho\sigma}\bar{p}_\rho a_\sigma$. As $G_1(x_1)$ is an entire function, (2.1) is local in the sense that it is free of unphysical poles involving x_1 in the denominator once expanded in powers of the spin vector.

At four points, we have the classical gravitational Compton scattering of two gravitons and the Kerr black hole, depicted below.

$$(2.2)$$

A covariant form of the Compton amplitude for a Kerr black hole is obtained in [73, 74]. Schematically, this amplitude receives three types of contributions as follows,

$$\mathcal{M}_4(1, 2, \bar{p}', \bar{p}) = -\frac{\mathcal{N}_a(1, 2, \bar{p}', \bar{p}) \mathcal{N}_0(1, 2, \bar{p}, \bar{p}')}{2(p_1 \cdot p_2)} + \frac{\mathcal{N}_r(1, 2, \bar{p}', \bar{p})}{4(\bar{p} \cdot p_1)(\bar{p} \cdot p_2)} + \mathcal{N}_c(1, 2, \bar{p}', \bar{p}). \quad (2.3)$$

Here the first term, as its form suggests, is assembled from the double-copy procedure. The second term is designated to take care of the spin-flip effects. The last denotes the contact contributions which can not be determined from factorizations.

The double-copy term, constructed in [73], is given by \mathcal{N}_a denoting the kinematic numerator of the Yang-Mills Compton amplitude in the heavy-mass limit and its spinless counterpart \mathcal{N}_0 . The numerator \mathcal{N}_a takes the following form,

$$\begin{aligned} \mathcal{N}_a(1, 2, \bar{3}, \bar{4}) &= -\frac{\mathbf{w}_1 \cdot F_1 \cdot F_2 \cdot \mathbf{w}_2}{(p_1 \cdot \bar{p})} + \frac{p_1 \cdot \bar{p} - p_2 \cdot \bar{p}}{2(p_1 \cdot \bar{p})} \times \\ &\left(iG_2(x_1, x_2) (a \cdot F_1 \cdot F_2 \cdot S \cdot p_2) + iG_2(x_1, x_2) (a \cdot F_2 \cdot F_1 \cdot S \cdot p_1) + iG_1(x_{12}) \text{tr}(F_1 \cdot S \cdot F_2) \right. \\ &\left. + G_1(x_1)G_1(x_2) \left((a \cdot F_1 \cdot \bar{p})(a \cdot F_2 \cdot p_1) - (a \cdot F_1 \cdot p_2)(a \cdot F_2 \cdot \bar{p}) - \frac{p_2 \cdot \bar{p} - p_1 \cdot \bar{p}}{2} (a \cdot F_1 \cdot F_2 \cdot a) \right) \right), \end{aligned} \quad (2.4)$$

where $x_{12} = x_1 + x_2$ and

$$G_2(x_1, x_2) \equiv \frac{1}{x_2} \left(\frac{\sinh(x_{12})}{x_{12}} - \cosh(x_2) \frac{\sinh(x_1)}{x_1} \right). \quad (2.5)$$

$G_2(x_1, x_2)$ is also an entire function and hence renders \mathcal{N}_a free of unphysical poles in x_1 , x_2 or x_{12} . The scalar numerator \mathcal{N}_0 follows from the kinematic Hopf algebra [137–142]

$$\mathcal{N}_0(1, 2, \bar{p}, \bar{p}') = -\left(\frac{\bar{p}' \cdot F_1 \cdot F_2 \cdot \bar{p}'}{p_1 \cdot \bar{p}'} \right). \quad (2.6)$$

We have $\mathcal{N}_0 = \mathcal{N}_a|_{a \rightarrow 0}$. These numerators manifestly satisfy the crossing symmetries described by group algebra in actions [143, 144] as demanded by the color-kinematic duality. The double-copy term is consistent with all possible massless factorizations.

The second term in eq. (2.3) complements the double-copy contribution to produce the correct spin-flip effects, starting from the cubic order in spin. It is determined by

the factorization requirement on the physical massive poles. The numerator \mathcal{N}_r reads

$$\begin{aligned}
\mathcal{N}_r(1, 2, \vec{p}', \vec{p}) = & \frac{\left((\partial_{x_1} - \partial_{x_2}) G_1(x_1) G_1(x_2) \right)}{4(\vec{p} \cdot p_1)(\vec{p} \cdot p_2)} \left(\vec{p} \cdot p_2 (\vec{p}^2 (a \cdot F_1 \cdot F_2 \cdot a) (a \cdot F_2 \cdot F_1 \cdot \vec{p}) \right. \\
& + a^2 (\vec{p} \cdot F_1 \cdot F_2 \cdot \vec{p}) (a \cdot F_1 \cdot F_2 \cdot \vec{p}) - (1 \leftrightarrow 2) \Big) \\
& + \left(\frac{i(\partial_{x_1} - \partial_{x_2}) G_2(x_1, x_2)}{4(\vec{p} \cdot p_1)(\vec{p} \cdot p_2)} \right) \left((\vec{p} \cdot p_2) (a \cdot F_2 \cdot F_1 \cdot \vec{p}) ((a \cdot F_2 \cdot \vec{p}) (a \cdot \tilde{F}_1 \cdot \vec{p}) \right. \\
& \left. - (a \cdot F_1 \cdot \vec{p}) (a \cdot \tilde{F}_2 \cdot \vec{p})) + (1 \leftrightarrow 2) \right). \tag{2.7}
\end{aligned}$$

The consistency conditions resulted from all the physical factorizations [87] are satisfied.

It is noted in [73, 74] that the three-point amplitude, the double-copy and the spin-flip contributions to the Compton amplitude share the property that their *degrees* are zero. Here the degree is defined as the maximal power of $1/\chi$, when χ parameterizes the scaling of the spin variable, namely $a^\mu \rightarrow \chi a^\mu$ and $\chi \rightarrow i\infty$.

The contact contribution is not accessible via factorizations and instead is obtained by matching with physical data [81, 145] at $\mathcal{O}(a^4)$ and $\mathcal{O}(a^5)$. Such expressions also exhibit the characteristic above that their degrees are zero at low orders in spin. Imposing this as a general constraint, the following contact contribution is extrapolated to all orders in spin,

$$\begin{aligned}
\mathcal{N}_c(1, 2, \vec{p}', \vec{p}) = & \left(\frac{(\partial_{x_1} - \partial_{x_2})^2}{2!} G_1(x_1) G_1(x_2) \right) \left((a \cdot F_1 \cdot \vec{p}) (a \cdot F_2 \cdot \vec{p}) (a \cdot F_1 \cdot F_2 \cdot a) \right. \\
& \left. - \frac{a^2}{2} ((a \cdot F_1 \cdot F_2 \cdot p) (a \cdot F_2 \cdot F_1 \cdot \vec{p}) + (a \cdot F_1 \cdot F_2 \cdot a) (\vec{p} \cdot F_1 \cdot F_2 \cdot \vec{p})) \right) \tag{2.8} \\
& + \left(\frac{i(\partial_{x_1} - \partial_{x_2})^2}{2!} G_2(x_1, x_2) \right) \left(-\frac{1}{2} ((a \cdot F_1 \cdot F_2 \cdot a) (a \cdot F_2 \cdot \vec{p}) (a \cdot \tilde{F}_1 \cdot \vec{p}) - (1 \leftrightarrow 2)) \right).
\end{aligned}$$

Putting together all three types of contributions, the covariant expression for the Compton amplitude (2.3) is manifestly gauge invariant and free of unphysical poles.

2.2 Spin expansion and analysis of contact contributions

Here we consider the expansion of the Compton amplitude reviewed above in powers of the spin vector and discuss the comparison between this Compton amplitude and known results in the literature computed from two different approaches, in particular, the higher-spin theory [99] and the Teukolsky equation [83]. To avoid confusion, in the remaining part of this section, we refer to eq. (2.3) as \mathcal{M}_{SP} , where the subscript SP indicates that this expression is obtained from bootstrapping with only two parameters

x_1 and x_2 and is speculated to describe the single-particle contributions only in [74]. Likewise, we denote the Compton amplitude computed from the higher-spin theory in [99] as \mathcal{M}_{HS} and the one extracted from the solution to the Teukolsky equation in [83] as $\mathcal{M}_{\text{TS-FZ}}$, where the ‘‘FZ’’ in the subscript indicates that we only consider the contributions obtained from imposing the so-called far-zone asymptotic behaviours.¹

The Compton amplitude derived from the higher-spin theory \mathcal{M}_{HS} contains two entire functions which depend on three parameters [99], among which is the spheroidicity parameter defined as $z = 2\sqrt{-a \cdot a}(\vec{p} \cdot p_1)/m$ [146].² The z -dependence in \mathcal{M}_{HS} follows from the higher-spin framework and shows up as contact terms only in the classical limit. Such terms are speculated to be related to the internal structures of the Kerr black hole [147] or the near-zone physics, which is beyond the scope of \mathcal{M}_{SP} given in eq. (2.3). On top of the discussions on the comparison at the level of the entire functions in [74], we have found further agreement with \mathcal{M}_{HS} when taking $z \rightarrow 0$, namely

$$\mathcal{M}_{\text{SP}}(1^\pm, 2^\mp, \vec{p}', \vec{p}) = \mathcal{M}_{\text{HS}}(1^\pm, 2^\mp, \vec{p}', \vec{p})|_{z \rightarrow 0}. \quad (2.9)$$

This relation is checked order by order up to $\mathcal{O}(a^{20})$.

As for the comparison with [83], it is observed in [74] that the extra contact terms needed for matching with the far-zone data at $\mathcal{O}(a^5)$ can be rewritten as a polynomial in z , although the z -dependence is traded for other parameters in the original expression of $\mathcal{M}_{\text{TS-FZ}}$. These extra contact terms vanish as $z \rightarrow 0$. Here we see that this observation holds up to $\mathcal{O}(a^8)$. In other words, at a given order in spin, we can separate $\mathcal{M}_{\text{TS-FZ}}$ into two parts, the z -independent part which agrees with \mathcal{M}_{SP} in 4 dimensions and the z -dependent part which vanishes as $z \rightarrow 0$,

$$\mathcal{M}_{\text{SP}}(1^\pm, 2^\mp, \vec{p}', \vec{p}) = \mathcal{M}_{\text{TS-FZ}}(1^\pm, 2^\mp, \vec{p}', \vec{p}) - \mathcal{M}_{\text{TS-FZ}}^{(c)}(z), \quad \text{with } \mathcal{M}_{\text{TS-FZ}}^{(c)}(0) = 0. \quad (2.10)$$

$\mathcal{M}_{\text{TS-FZ}}$ is expressed in terms of the spinor-helicity variables. \mathcal{M}_{SP} can be readily rewritten and it is straightforward to find $\mathcal{M}_{\text{TS-FZ}}^{(c)}$ in these variables. But to see that $\mathcal{M}_{\text{TS-FZ}}^{(c)}$ indeed vanishes at a given order in spin as $z \rightarrow 0$, it is more convenient to use an alternative expression in terms of covariant variables. To this end, we adopt a simple procedure as follows.

We begin with a general ansatz in terms of the covariant variables. That is, the ansatz is a linear combination of monomials comprised of powers of the parameter z and

¹We refer to the Teukolsky result with $\alpha \rightarrow 1$ as the ‘‘far-zone data’’ through this manuscript, as indicated in [83]. However, we acknowledge ambiguities in this way of separating the far-zone and near-zone contributions.

²We use a different normalization for z from that in [99].

factors of the forms $\mathbf{v}\cdot\mathbf{v}'$, $\mathbf{v}\cdot F_i\cdot F_j\cdot\mathbf{v}'$ and $\mathbf{v}\cdot F_i\cdot\tilde{F}_j\cdot\mathbf{v}'$ where the vectors \mathbf{v} and \mathbf{v}' can be either a^μ or a momentum. The ansatz must have the correct counting in a^μ and ε_i^μ and satisfy the dimension and parity requirements. Imposing that it should agree with the spinor-helicity expression of $\mathcal{M}_{\text{TS-FZ}}^{(c)}(z)$ for all possible helicity configurations, when restricted to 4 dimensions, we fix the coefficients in the ansatz. Recall that the same-helicity gravitational Compton amplitude is known to all orders in spin, which is captured already in \mathcal{M}_{SP} . Hence $\mathcal{M}_{\text{TS-FZ}}^{(c)}(z)$ must vanish in the same-helicity configurations and receive non-zero contributions only in the opposite-helicity ones.

We note that the monomials satisfying the power counting and parity requirements form an over-complete basis in 4 dimensions, due to the identities involving the Levi-Civita symbols. Such relations only hold in 4 dimensions and may contain coefficients that are rational in inner products of momenta. Hence it is difficult to remove such redundancies in the ansatz completely. Consequently, there is a multitude of covariant expressions for $\mathcal{M}_{\text{TS-FZ}}^{(c)}$ at a given order in spin. They all evaluate to the same in 4 dimensions, but may be different in general dimensions. This leads to certain subtleties when we construct their contribution to one-loop amplitude, which will be discussed in the next section.

We give the explicit expressions for $\mathcal{M}_{\text{TS-FZ}}^{(c)}(z)$ at $\mathcal{O}(a^5)$ and $\mathcal{O}(a^6)$ and higher-order terms are given in Appendix A. These expressions are manifestly gauge invariant and vanishing as $z \rightarrow 0$. We emphasize that they should be viewed as valid in 4 dimensions only, although they are expressed in terms of covariant quantities.

$$\begin{aligned}
\mathcal{M}_{\text{TS-FZ}}^{(c,5)}(z) &= 2ip_1\cdot\bar{p}\cdot a\cdot a \left(a\cdot F_2\cdot\tilde{F}_1\cdot\bar{p} + a\cdot\tilde{F}_1\cdot F_2\cdot\bar{p} \right) \left(\frac{(a\cdot a)(\bar{p}\cdot F_1\cdot F_2\cdot\bar{p})}{12m^2} - \frac{11}{60}(a\cdot F_1\cdot F_2\cdot a) \right) \\
\mathcal{M}_{\text{TS-FZ}}^{(c,6)}(z) &= \frac{(a\cdot a)^3(p_1\cdot\bar{p})^2(\bar{p}\cdot F_1\cdot F_2\cdot\bar{p})^2}{9m^4} + \frac{13(a\cdot a)^3p_1\cdot\bar{p}p_1\cdot F_2\cdot F_1\cdot\bar{p}\bar{p}\cdot F_1\cdot F_2\cdot\bar{p}}{90m^2} \\
&\quad - \frac{13(a\cdot a)^3p_1\cdot\bar{p}p_2\cdot F_1\cdot F_2\cdot\bar{p}\bar{p}\cdot F_1\cdot F_2\cdot\bar{p}}{90m^2} + \frac{2(a\cdot a)^2p_1\cdot\bar{p}a\cdot p_1\bar{p}\cdot F_1\cdot F_2\cdot\bar{p}a\cdot F_2\cdot F_1\cdot\bar{p}}{15m^2} \\
&\quad + \frac{16(a\cdot a)^2p_1\cdot\bar{p}a\cdot p_2\bar{p}\cdot F_1\cdot F_2\cdot\bar{p}a\cdot F_2\cdot F_1\cdot\bar{p}}{45m^2} - \frac{(a\cdot a)^2(p_1\cdot\bar{p})^2a\cdot F_1\cdot F_2\cdot a\bar{p}\cdot F_1\cdot F_2\cdot\bar{p}}{m^2} \\
&\quad - \frac{2(a\cdot a)^2p_1\cdot\bar{p}a\cdot p_2\bar{p}\cdot F_1\cdot F_2\cdot\bar{p}a\cdot F_1\cdot F_2\cdot\bar{p}}{15m^2} - \frac{16(a\cdot a)^2p_1\cdot\bar{p}a\cdot p_1\bar{p}\cdot F_1\cdot F_2\cdot\bar{p}a\cdot F_1\cdot F_2\cdot\bar{p}}{45m^2} \\
&\quad + \frac{14}{45}a\cdot a(p_1\cdot\bar{p})^2(a\cdot F_1\cdot F_2\cdot a)^2 + \frac{22}{45}a\cdot ap_1\cdot\bar{p}a\cdot F_1\cdot F_2\cdot aa\cdot p_1a\cdot F_1\cdot F_2\cdot\bar{p} \\
&\quad - \frac{22}{45}a\cdot ap_1\cdot\bar{p}a\cdot F_1\cdot F_2\cdot aa\cdot p_2a\cdot F_2\cdot F_1\cdot\bar{p}. \tag{2.11}
\end{aligned}$$

Before closing this section, we make several remarks on the various attempts at obtaining the gravitational Compton in the literature. The Teukolsky solution yields

the scattering phase of the graviton scattered off the Kerr black hole, by imposing the asymptotic behaviours of the Teukolsky equations in the near-zone (near the horizon) and the far-zone (at infinity) [81, 83, 145]. The scattering amplitude is then determined by matching at the level of the scattering phase. In principle, not only the leading but also higher-order contributions in the PM expansion can enter the non-perturbative scattering phase. Hence for a meaningful comparison with the tree Compton amplitudes obtained from field-theory approaches, it is crucial to isolate the leading PM contribution in the scattering phase. For a complete comparison between these two types of approaches, loop-level Compton amplitudes are also necessary.

On the other hand, the scattering phase obtained from the Teukolsky equation contains both near-zone and far-zone contributions. The far-zone contribution is rational, whereas the near-zone contains transcendental functions. The Compton amplitudes computed from field-theory approaches are postulated to capture the rational part. At the level of the scattering phase, there naturally is another type of ambiguities arising from rewriting the transcendental functions using various identities, which changes the rational terms. Therefore, it is also important to determine the rational terms in a way that is physically meaningful. In this paper, we consider and match with $\mathcal{M}_{\text{TS-FZ}}$ which is extracted from the far-zone contribution only. In [99], another procedure is discussed which leads to the matching between \mathcal{M}_{HS} and \mathcal{M}_{TS} at a different point. It would be interesting to see if there exists a way of rewriting the transcendental functions such that the remaining rational terms are independent of z .

3 Eikonal with spin at 2PM


In this section, we compute the eikonal phase [102] for the spinning Kerr binary at 2PM. The classical eikonal [35] is an important generating that computes physical observables. In the non-spinning case, several variations, such as the radial action [50–52], the HEFT phase [41] and the WQFT eikonal [42, 45, 148], have also been investigated. At 2PM, they are practically the equivalent, given by Fourier transforming the classical one-loop $2 \rightarrow 2$ scattering amplitude to the impact parameter space. As seen in related works [102, 149], these generating functions generally extend to the spinning case with spin-related subtleties.

The eikonal and the resulting observables receive all types of contributions from the Compton amplitudes, as discussed in section 2. Our main focus is the classical one-loop amplitude is constructed from the covariant Compton amplitude, using unitarity-based methods [41, 150, 151]. We compute the general form of the one-loop integrand and perform the loop integration in the spin expansion explicitly up to $\mathcal{O}(a^8)$. For the

sake of comparing with physical data, we include the contributions from the extra z -dependent contact terms at $\mathcal{O}(a^5)$ and $\mathcal{O}(a^6)$. Altogether, we find perfect agreement with the far-zone data.

3.1 Contribution from the covariant Compton amplitude

The classical one-loop amplitude is computed from the leading singularity as depicted below (with the mirror diagram implied), where the triple cut is given by two massless (graviton) cuts and one massive one,



$$\mathcal{M}_{a_1 a_2}^{(1)} = \frac{1}{2} \sum_{h_i = \pm} \frac{(32\pi G)^2}{(4\pi)^{D/2}} \int \frac{d^D \ell_1}{\pi^{D/2}} \frac{\delta(m_2 v_2 \cdot \ell_1)}{\ell_1^2 \ell_2^2} \quad (3.1)$$

$$\times \left(\mathcal{M}_3^{-h_1}(-\ell_1, v_2) \mathcal{M}_3^{-h_2}(-\ell_2, v_2) \mathcal{M}_4^{h_1 h_2}(\ell_1, \ell_2, v_1) \right).$$

Here the massive cut labelled by a solid red line gives the δ -function and we have formally lifted the massless cuts $\delta(\ell_j^2) \rightarrow i/\ell_j^2$, while keeping in mind that bubble or ultra-local terms, with zero or negative propagator powers for the massless propagators, should vanish. The velocities of the heavy bodies are parameterized such that $v_1^2 = v_2^2 = 1$ and $v_1 \cdot v_2 = \gamma$ as usual. The tree-level amplitudes \mathcal{M}_3 and \mathcal{M}_4 are given by eq. (2.1) and eq. (2.3). We use dimensional regularization in the loop integration, namely $D = 4 - 2\epsilon$. The summation is taken over the helicities of the gravitons and the completeness relation reads

$$\sum_{h=\pm} \varepsilon_k^\mu \varepsilon_k^\nu \varepsilon_{-k}^{*\rho} \varepsilon_{-k}^{*\sigma} = \frac{1}{2} (\mathcal{P}^{\mu\rho} \mathcal{P}^{\nu\sigma} + \mathcal{P}^{\mu\sigma} \mathcal{P}^{\nu\rho}) - \frac{1}{D-2} \mathcal{P}^{\mu\nu} \mathcal{P}^{\rho\sigma}, \quad (3.2)$$

where

$$\mathcal{P}^{\mu\nu} = \eta^{\mu\nu} - \frac{k^\mu n^\nu + n^\mu k^\nu}{k \cdot n} \quad (3.3)$$

with the reference null vector n^μ transverse to the polarization $k \cdot \varepsilon = n \cdot \varepsilon = 0$. The covariant three- and four-point tree amplitudes are expected to hold in general dimensions and in practice we can simply set $\mathcal{P}^{\mu\nu} = \eta^{\mu\nu}$. The reference vector always drops out.

The dependence on the graviton polarization vectors in the tree amplitudes eq. (2.1) and eq. (2.3) always enters through the (dual) field strengths $F_i^{\mu\nu}$ and $\tilde{F}_i^{\mu\nu}$ and is completely manifest. Since the function $G_1(x)$ depends only on momenta and spins,

the above gluing can be performed without expanding in powers of spin. The resulting contribution to the one-loop integrand at all orders in spin is given in the **Ancillary Files**.

The presence of $e^{\ell_1 \cdot a_j}$ factors in eq. (3.1) may seem worrisome. Before expanding in powers of spin, potential divergences due to such factors can be easily remedied by a shift of the spin vector $a_j \rightarrow i\tilde{a}_j$. We illustrate with a simple example that the integral in eq. (3.1) does not behave worse than the non-spinning case does in the ultraviolet under this shift in Appendix B. Having evaluated the integral, we can analytically continue back to real values of a_j . As for the spin expansion of this integral, it can be defined as the Taylor expansion of the well-defined integral at each order. Similar integrals are studied and the resummed expressions are given in special kinematic regions in [121]. Our preliminary analysis of simple examples in such kinematic regions show that the naive evaluation of the spin-expanded integral agrees with the expansion of the resummed one, order by order.

We now proceed to consider the expansion of eq. (3.1) in powers of spin. Expanding out the spin tensor $S^{\mu\nu}$ and using the identity³

$$\epsilon^{\mu_1 \dots \mu_n \alpha_1 \dots \alpha_m} \epsilon_{\mu_1 \dots \mu_n \beta_1 \dots \beta_m} = (-1)^n n! \delta_{\beta_1 \dots \beta_m}^{\alpha_1 \dots \alpha_m}, \quad \delta_{\beta_1 \dots \beta_m}^{\alpha_1 \dots \alpha_m} = m! \delta_{\beta_1}^{[\alpha_1} \dots \delta_{\beta_m}^{\alpha_m]}, \quad (3.4)$$

we can always write the integrand in terms of scalar products and at most one factor of $\epsilon(A, B, C, D) = \epsilon_{\mu\nu\rho\sigma} A^\mu B^\nu C^\rho D^\sigma$. In fact, we find that at even orders in spin $\mathcal{O}(a^{2n})$, the integrand contains only scalar products and at odd orders in spin $\mathcal{O}(a^{2n+1})$, the integrand contains one $\epsilon(A, B, C, D)$ factor. Removing the bubble/ultra-local terms, the spin-expanded expression has two types of structures,

$$\int \frac{d^D \ell_1}{\pi^{D/2}} \frac{\delta(\ell_1 \cdot v_2) (\ell_1 \cdot a_1)^{n_1} (\ell_1 \cdot a_2)^{n_2}}{\ell_1^2 \ell_2^2 (\ell_1 \cdot v_1)^m}, \quad \int \frac{d^D \ell_1}{\pi^{D/2}} \frac{\delta(\ell_1 \cdot v_2) (\ell_1 \cdot a_1)^{n_1} (\ell_1 \cdot a_2)^{n_2} \ell_1^\mu}{\ell_1^2 \ell_2^2 (\ell_1 \cdot v_1)^m}, \quad (3.5)$$

where n, n_1, n_2 and m are integers. The vector ℓ_1^μ in the second term above typically gets contracted with $\epsilon(A, B, C, \cdot)_\mu$ where $A, B, C \in \{q, v_1, v_2, a_1, a_2\}$.

Such one-loop tensor integrals can be readily decomposed into scalar integrals. These scalar integrals are dressed by rank- n tensor structures. As shown in [44], it is convenient to write down such tensor structures in terms of the following variables

$$\theta_1^\mu := \frac{\gamma v_2^\mu - v_1^\mu}{\gamma^2 - 1}, \quad \theta_2^\mu := \frac{\gamma v_1^\mu - v_2^\mu}{\gamma^2 - 1}, \quad \Pi^{\mu\nu} := \eta^{\mu\nu} - \theta_1^\mu v_1^\nu - \theta_2^\mu v_2^\nu - \frac{q^\mu q^\nu}{q^2}, \quad (3.6)$$

where $\Pi^{\mu\nu}$ is the $(D-3)$ -dimensional metric. It is easy to verify that a tensor integral of the form

$$\mathcal{I}_{1,1,m,1}[\ell_1^{\mu_1} \dots \ell_1^{\mu_n}] = \int \frac{d^D \ell_1}{\pi^{D/2}} \frac{\delta(\ell_1 \cdot v_2) \ell_1^{\mu_1} \dots \ell_1^{\mu_n}}{\ell_1^2 \ell_2^2 (\ell_1 \cdot v_1)^m} \quad (3.7)$$

³We use the package **KiHA** for manipulating the expressions [152].

can be decomposed into scalar integrals dressed by tensor structures built from q^μ , θ_1 and $\Pi^{\mu\nu}$ only. Since all but at most one of these indices are to be contracted with a_1 or a_2 , symmetries of the two structures in eq. (3.5) lead to closed-form expressions for the decomposition. We present the explicit decomposition of the first structure in eq. (3.5) here and postpone the derivation to Appendix C,

$$\begin{aligned}
& \mathcal{I}_{1,1,\alpha,1}[(\ell_1 \cdot a_1)^M (\ell_1 \cdot a_2)^{N-M}] \\
= & \sum_{\substack{n_1+n_2+2n_3=N, \\ n_i \geq 0, n_i \in \mathbb{Z}}} \frac{1}{N_{n_3}} \left(\sum_{k=0}^{n_3} \binom{n_3}{k} \frac{1}{(\gamma^2 - 1)^k} \left(\frac{q^2}{2}\right)^{n_2+n_3-k} \mathcal{I}_{1,1,\alpha-n_1-2k,1}[1] \right) \\
& \times \left(\sum_{\text{cond.}} C_{N,M,n_1,n_2,m_1,m_2,m_3} (a_1 \cdot \theta_1)^{m_1} (a_2 \cdot \theta_1)^{n_1-m_1} (a_1 \cdot q)^{m_2} (a_2 \cdot q)^{n_2-m_2} \right. \\
& \left. (a_1 \cdot \Pi \cdot a_1)^{m_3} (a_1 \cdot \Pi \cdot a_2)^{m_4} (a_2 \cdot \Pi \cdot a_2)^{n_3-m_3-m_4} \right), \tag{3.8}
\end{aligned}$$

where $N_{n_3} = (D-3)(D-1)\cdots(D-3+2(n_3-1))$ for $n_3 > 0$ and $N_0 = 1$. The summation in the last bracket is taken over all the solutions to the conditions below

$$m_1 + m_2 + 2m_3 + m_4 = M, \quad 0 \leq m_i \leq n_i, \quad 0 \leq m_4 \leq n_3, \quad m_i \in \mathbb{Z}. \tag{3.9}$$

The coefficients read

$$C_{N,M,n_1,n_2,m_1,m_2,m_3} = \frac{M!}{m_1!m_2!m_3!m_4!} \frac{(N-M)!}{(n_1-m_1)!(n_2-m_2)!(n_3-m_3)!} \frac{1}{2^{n_3-m_4}}. \tag{3.10}$$

The second structure in eq. (3.5) admits a similar expression with one free index, which we also give in Appendix C.

After decomposing the tensor integrals, the resulting scalar integrals are easily cast in a basis of master integrals using the IBP relations.⁴ Up to $\mathcal{O}(a^8)$, we see that only the triangle integral below contributes in 4 dimensions,⁵

$$\mathcal{I}_{1,1,0,1} = \int \frac{d^D \ell_1}{\pi^{D/2}} \frac{\delta(\ell_1 \cdot v_2)}{\ell_1^2 \ell_2^2} = \frac{2^{5-D} \pi^2 (-q^2)^{(D-5)/2} \sec(\pi D/2)}{\Gamma(\frac{D}{2} - 1)}. \tag{3.11}$$

Evaluating the integral, we arrive at the spin-expanded one-loop amplitude computed from the covariant Compton amplitude in eq. (2.3). It is independent of the z -parameter. We find perfect agreements with the literature up to $\mathcal{O}(G^2 a^4)$ [145].

⁴We use the package `LiteRed` for the IBP reduction [153, 154].

⁵The box integral $\mathcal{I}_{1,1,1,1}$ appears after the IBP reduction. But its coefficient vanishes when we restrict to 4 dimensions.

In this case, we have to use the “chiral” version of the completeness relation. Equivalently, we can separate the even and odd terms in spin, as discussed in [122]. At even orders in spin, we have

$$\cosh(\ell_{12} \cdot a_2) (\mathcal{C}_{+-} + \mathcal{C}_{-+}) = \cosh(\ell_{12} \cdot a_2) (\mathcal{C}_{+-} + \mathcal{C}_{++} + \mathcal{C}_{--} + \mathcal{C}_{-+}), \quad (3.15)$$

where $\ell_{12} = \ell_1 - \ell_2$ and $\mathcal{C}_{++} + \mathcal{C}_{--} = 0$. Here we can use the same completeness relation in eq. (3.2), since we have

$$\varepsilon_k^{+\mu} \varepsilon_k^{+\nu} \varepsilon_{-k}^{-\rho} \varepsilon_{-k}^{-\sigma} + \varepsilon_k^{-\mu} \varepsilon_k^{-\nu} \varepsilon_{-k}^{+\rho} \varepsilon_{-k}^{+\sigma} \quad (3.16)$$

for both gravitons. At odd powers in spin, we have instead

$$\sinh(\ell_{12} \cdot a_2) (\mathcal{C}_{+-} - \mathcal{C}_{-+}) = \sinh(\ell_{12} \cdot a_2) (\mathcal{C}_{+-} - \mathcal{C}_{++} + \mathcal{C}_{--} - \mathcal{C}_{-+}). \quad (3.17)$$

Hence we have

$$\varepsilon_k^{+\mu} \varepsilon_k^{+\nu} \varepsilon_{-k}^{-\rho} \varepsilon_{-k}^{-\sigma} - \varepsilon_k^{-\mu} \varepsilon_k^{-\nu} \varepsilon_{-k}^{+\rho} \varepsilon_{-k}^{+\sigma} = \frac{1}{4} \left(\mathcal{P}^{\mu\rho} \tilde{\mathcal{P}}^{\nu\sigma} + \mathcal{P}^{\mu\sigma} \tilde{\mathcal{P}}^{\nu\rho} + \mathcal{P}^{\nu\rho} \tilde{\mathcal{P}}^{\mu\sigma} + \mathcal{P}^{\nu\sigma} \tilde{\mathcal{P}}^{\mu\rho} \right), \quad (3.18)$$

for one of the two gravitons, where

$$\tilde{\mathcal{P}}^{\mu\nu} = \frac{i\epsilon^{\mu\nu\rho\sigma} k_\rho n_\sigma}{k \cdot n}. \quad (3.19)$$

This way we guarantee that the remnant ambiguities introduced in the process of lifting the spinor-helicity expression for $\mathcal{M}_{\text{TS-FZ}}^{(c)}$ to the covariant one have no impact at one loop.

This concludes our discussion on the one-loop contribution from the extra contact terms. Together with the contribution from the covariant Compton amplitude, we find agreement with the far-zone data in [145] up to $\mathcal{O}(G^2 a^6)$, which is the highest-order result to our best knowledge. Comparing with the 2PM calculation from the higher-spin amplitudes [155] may be helpful for finding an all-order-in-spin remedy for the z -dependence.

3.3 2PM eikonal in the spin expansion

Generally speaking, the eikonal exponentiation can be identified with the classical amplitude Fourier transformed to the impact parameter space. The classical eikonal phase is then extracted order by order. Recent works [102, 121] show that the classical eikonal can be viewed as a generator for canonical transformations.⁶

⁶Another work [119] shows that the radial action can be understood in a similar fashion.

Up to 2PM, the relation between the amplitude and the eikonal remains straightforward, namely

$$\chi_\tau^{(\text{nPM})} = \frac{1}{4m_1m_2\sqrt{\gamma^2-1}} \int \frac{d^{D-2}\mathbf{q}}{(2\pi)^{D-2}} e^{-i\mathbf{b}\cdot\mathbf{q}} \mathcal{M}_{a_1a_2}^{(n-1)} U_1 U_2, \quad (3.20)$$

where \mathbf{b} is the impact parameter and τ labels the Thomas-Wigner rotation factors [85, 86]

$$U_i = e^{\frac{i\tau\mathcal{E}_i}{E m_i(E_i+m_i)}}, \quad \mathcal{E}_i := E \mathbf{s}_i \cdot (\mathbf{p} \times \mathbf{q}) \quad (3.21)$$

with E_i denoting the energy of the two black holes and $E = E_1 + E_2$. The bold-faced letters denote the three-vectors. The three-momenta are given in the center-of-mass (COM) frame, namely $p_1 = (E_1, \mathbf{p})$, $p_2 = (E_2, -\mathbf{p})$, $q = (0, \mathbf{q})$, and \mathbf{s}_i is the spatial part of $s_i^\mu = m_i a_i^\mu$. Different values of τ correspond to different choices of the spin supplementary conditions (SSC): $\tau = 0$ corresponds to the covariant SSC $\bar{p}_\mu S^{\mu\nu} = 0$ whereas $\tau = 1$ the canonical SSC. (For more details, see also [84, 145].) Dressing the amplitude with the Thomas-Wigner rotation factors is effectively equivalent to shifting the impact parameter $\mathbf{b} \rightarrow \mathbf{b} + \sum_{i=1,2} \left(\frac{-\tau}{E_i+m_i} \right) \mathbf{p} \times \mathbf{s}_i$. The canonical observables follow from the the canonical SSC.

Here we showcase the explicit expression for the z -independent $\mathcal{O}(G^2 a^7)$ contributions to $\chi_{\tau=0}^{(2\text{PM})}$ in the aligned-spin case, which is given by the covariant Compton amplitude. To compare with the far-zone data, it needs to be supplemented by the contributions from the z -dependent contact terms $\mathcal{M}_{\text{TS-FZ}}^{(c,n=5,6,7)}$, which are given in eq. (2.11) and Appendix A. We parameterize the aligned spins as $a_2^\mu = \xi a_1^\mu$ and hence the powers of a_1 and a_2 are tracked by the power of ξ .

$$\chi_{\tau=0}^{(2\text{PM})} \Big|_{\xi^4, z=0} = - \frac{35\pi G^2 \xi^4 \gamma (24(9\gamma^2-4)m_1 + (114-239\gamma^2)m_2)}{512(\gamma^2-1)^{3/2} |\mathbf{b}|^8} \hat{b} \cdot S_1 \cdot p_2 \left(429(\hat{b} \cdot a_1)^6 - 594a_1^2(\hat{b} \cdot a_1)^4 + 216a_1^4(\hat{b} \cdot a_1)^2 - 16a_1^6 \right), \quad (3.22)$$

$$\chi_{\tau=0}^{(2\text{PM})} \Big|_{\xi^5, z=0} = - \frac{21\pi G^2 \xi^5 \gamma \hat{b} \cdot S_1 \cdot p_2}{512(\gamma^2-1)^{3/2} |\mathbf{b}|^8} \left(2145 \left((38\gamma^2-15)m_1 + 26(1-2\gamma^2)m_2 \right) (\hat{b} \cdot a_1)^6 - 33a_1^2 \left((3421\gamma^2-1351)m_1 + 2340(1-2\gamma^2)m_2 \right) (\hat{b} \cdot a_1)^4 + 36a_1^4 \left((1141\gamma^2-451)m_1 + 780(1-2\gamma^2)m_2 \right) (\hat{b} \cdot a_1)^2 - 8a_1^6 \left((381\gamma^2-151)m_1 + 260(1-2\gamma^2)m_2 \right) \right), \quad (3.23)$$

$$\chi_{\tau=0}^{(2\text{PM})} \Big|_{\xi^6, z=0} = - \frac{7\pi G^2 \xi^6 \gamma \hat{b} \cdot S_1 \cdot p_2}{4096(\gamma^2-1)^{3/2} |\mathbf{b}|^8} \left(-16a_1^6 \left(16(81\gamma^2-26)m_1 + 15(77-149\gamma^2)m_2 \right) \right)$$

$$\begin{aligned}
& -66a_1^2(16(721\gamma^2 - 226)m_1 + 135(77 - 149\gamma^2)m_2)(\hat{b}\cdot a_1)^4 \\
& + 72a_1^4(16(241\gamma^2 - 76)m_1 + 45(77 - 149\gamma^2)m_2)(\hat{b}\cdot a_1)^2 \\
& + 2145(16(16\gamma^2 - 5)m_1 + 3(77 - 149\gamma^2)m_2)(\hat{b}\cdot a_1)^6, \tag{3.24}
\end{aligned}$$

$$\begin{aligned}
\chi_{\tau=0}^{(2\text{PM})} \Big|_{\xi^7, z=0} &= -\frac{\pi G^2 \xi^7 \gamma \hat{b}\cdot S_1 \cdot p_2}{1024(\gamma^2 - 1)^{3/2} |\mathbf{b}|^8} \left(2145(6(8\gamma^2 - 1)m_1 + 7(9 - 17\gamma^2)m_2)(\hat{b}\cdot a_1)^6 \right. \\
& - 198a_1^2((727\gamma^2 - 97)m_1 + 105(9 - 17\gamma^2)m_2)(\hat{b}\cdot a_1)^4 \\
& + 108a_1^4((493\gamma^2 - 73)m_1 + 70(9 - 17\gamma^2)m_2)(\hat{b}\cdot a_1)^2 \\
& \left. - 8a_1^6((513\gamma^2 - 93)m_1 + 70(9 - 17\gamma^2)m_2) \right), \tag{3.25}
\end{aligned}$$

where we define $\hat{b}^\mu = b^\mu/|b|$. The terms that are not listed above can be obtained by swapping the two particles $\chi^{(2\text{PM})}|_{\xi^n} = (\chi^{(2\text{PM})}|_{\xi^{7-n}})_{1\leftrightarrow 2}$.

Following the prescription in [102], the impulse $\Delta\mathbf{p}_\perp$ and the spin kick $\Delta\mathbf{s}_1$ can be generated from the eikonal phase $\chi_{\tau=0} = \chi_{\tau=0}^{(1\text{PM})} + \chi_{\tau=0}^{(2\text{PM})} + \mathcal{O}(G^3)$ via⁷

$$\Delta\mathcal{O} = -\{\mathcal{O}, \chi\} - \frac{1}{2}\{\chi, \{\mathcal{O}, \chi\}\} - \mathcal{D}_{SL}(\chi, \{\mathcal{O}, \chi\}) + \frac{1}{2}\{\mathcal{O}, \mathcal{D}_{SL}(\chi, \chi)\}, \tag{3.26}$$

where $\Delta\mathcal{O} = \{\Delta\mathbf{p}_\perp, \Delta\mathbf{s}_1\}$ and the full impulse is given by $\Delta\mathbf{p} = \Delta p_\parallel \mathbf{p}/|\mathbf{p}| + \Delta\mathbf{p}_\perp$, which satisfies the on-shell condition $(\mathbf{p} + \Delta\mathbf{p})^2 = \mathbf{p}^2$. The Poisson bracket and the spin derivative operators are defined as

$$\{\mathbf{p}_\perp, f\} := -\frac{\partial f}{\partial \mathbf{b}}, \quad \{\mathbf{s}_1, f\} := \frac{\partial f}{\partial \mathbf{s}_1} \times \mathbf{s}_1, \quad \mathcal{D}_{SL}(f, g) := -\mathbf{s}_1 \cdot \left(\frac{\partial f}{\partial \mathbf{s}_1} \frac{\partial g}{\partial \mathbf{L}} \right), \tag{3.27}$$

with $\mathbf{L} := \mathbf{b} \times \mathbf{p}$.

4 Conclusions and outlook

We consider a particular proposal for the classical gravitational Compton amplitude, which follows from bootstrapping techniques and is expressed in covariant variables. This covariant expression is consistent with all possible massless and massive factorization requirements at all orders in spin and contains a contact contribution that is obtained by imposing the same empirical patterns as those observed in other contributions.

⁷It is noted in [121] that the operator \mathcal{D}_{SL} is not necessary once all the constraints are properly imposed.

In this paper, we further the analysis in [74] regarding the contact terms. In particular, we verify the matching between the covariant Compton amplitude and the one derived from the higher-spin theory [99] up to $\mathcal{O}(a^{20})$, when taking the spheroidicity parameter $z \rightarrow 0$. We also devise a simple procedure to find a covariant form which accounts for the missing contact contributions compared to those extracted from the far-zone data. We confirm that such contributions can be written as a polynomial in z and vanish as $z \rightarrow 0$ up to $\mathcal{O}(a^8)$. We believe such patterns extend to higher orders in spin. These observations corroborate somewhat the folklore that the z -dependent contact terms are associated with the internal structures of the Kerr black hole. Further studies on multiple fronts are needed to uncover the precise nature of such contact contributions.

From the covariant Compton amplitude, the calculation of the classical one-loop amplitude on the triple cut using unitarity-based methods is then streamlined. We obtain an all-order-in-spin integrand and evaluate the integral up to $\mathcal{O}(a^8)$ in the spin expansion. We have also included the contributions from z -dependent contact terms, for which the computation involves an extra subtlety. The Fourier transform of the one-loop amplitude to the impact parameter space gives the eikonal phase. We find perfect agreement with the 2PM far-zone data up to $\mathcal{O}(a^6)$. It is certainly interesting to consider the one-loop integral without expanding in spin.

Two directions are natural to look into, the binary dynamics and the waveform at higher PM orders. The binary dynamics at $\mathcal{O}(G^3)$ is given by the two-loop $2 \rightarrow 2$ amplitude and the waveform at $\mathcal{O}(G^{5/2})$ is given by the one-loop amplitude with a graviton radiation. The five-point tree-level gravitational Compton amplitude is needed in both cases. Computing the five-point amplitude to an arbitrary order in spin is by itself a challenge, although it is conceivable that the bootstrapping techniques should carry over. Even without the five-point amplitude at hand, one may consider mass sectors, for instance the “zig-zag” diagrams at two loops in [41], which only involve the three- and four-point amplitudes in the construction of loop amplitudes. The behaviours of the loop amplitudes in these sectors may in turn shed new light on additional constraints for the structures of the gravitational Compton amplitudes.

Acknowledgement

We thank F. Y. Baustista, E. Bjerrum-Bohr, A. Brandhuber, G. R. Brown, J. Gowdy, H. Johansson, J.-W. Kim, S. Lee, P. Pichini, M. Skowronek, G. Travaglini for insightful discussions. TW is grateful to H. Johansson, J.-W. Kim and S. Lee for discussions on their respective works and comments on the manuscript. GC has received funding

from the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement No. 847523 "INTERACTIONS". TW is supported by the NRF grant 2021R1A2C2012350 and the Fellowship of China Postdoctoral Science Foundation (No. 2022M713228).

A Extra contact terms

Here we present the explicit expressions for the extra contact terms that give the z -dependence of the Compton amplitude extracted from the Teukolsky solution at $\mathcal{O}(a^7)$ and $\mathcal{O}(a^8)$. We choose only the contribution obtained from fitting the far-zone asymptotic.

$$\begin{aligned}
\mathcal{M}_{\text{TS-FZ}}^{(c,7)} = & \frac{(284a \cdot p_1 + 211a \cdot p_2) (p_1 \cdot \bar{p})^2 \bar{p} \cdot F_1 \cdot F_2 \cdot \bar{p} \bar{p} \cdot F_1 \cdot \tilde{F}_2 \cdot \bar{p} (a \cdot a)^3}{945m^4} \\
& + \frac{8p_1 \cdot \bar{p} (a \cdot F_1 \cdot p_2 p_1 \cdot F_2 \cdot \bar{p} + a \cdot F_2 \cdot p_1 p_2 \cdot F_1 \cdot \bar{p}) (\bar{p} \cdot F_1 \cdot \tilde{F}_2 \cdot \bar{p} - \bar{p} \cdot F_2 \cdot \tilde{F}_1 \cdot \bar{p}) (a \cdot a)^3}{945m^2} \\
& + \frac{(211a \cdot p_1 + 284a \cdot p_2) (p_1 \cdot \bar{p})^2 \bar{p} \cdot F_1 \cdot F_2 \cdot \bar{p} \bar{p} \cdot F_2 \cdot \tilde{F}_1 \cdot \bar{p} (a \cdot a)^3}{945m^4} \\
& + \frac{p_1 \cdot \bar{p} (99 (a \cdot p_1)^2 m^2 - 101a \cdot p_1 a \cdot p_2 m^2 - 78a \cdot a (p_1 \cdot \bar{p})^2) a \cdot F_1 \cdot F_2 \cdot \bar{p} \bar{p} \cdot F_1 \cdot \tilde{F}_2 \cdot \bar{p} (a \cdot a)^2}{1890m^4} \\
& + \frac{4p_1 \cdot \bar{p} (-33 (a \cdot p_1)^2 m^2 + 23a \cdot p_1 a \cdot p_2 m^2 + 60a \cdot a (p_1 \cdot \bar{p})^2) a \cdot F_2 \cdot F_1 \cdot \bar{p} \bar{p} \cdot F_1 \cdot \tilde{F}_2 \cdot \bar{p} (a \cdot a)^2}{945m^4} \\
& + \frac{p_1 \cdot \bar{p} (-99 (a \cdot p_2)^2 m^2 + 101a \cdot p_1 a \cdot p_2 m^2 + 78a \cdot a (p_1 \cdot \bar{p})^2) a \cdot F_2 \cdot F_1 \cdot \bar{p} \bar{p} \cdot F_2 \cdot \tilde{F}_1 \cdot \bar{p} (a \cdot a)^2}{1890m^4} \\
& - \frac{101a \cdot p_1 (p_1 \cdot \bar{p})^2 a \cdot F_1 \cdot F_2 \cdot a \bar{p} \cdot F_1 \cdot \tilde{F}_2 \cdot \bar{p} (a \cdot a)^2}{135m^2} - \frac{101a \cdot p_2 (p_1 \cdot \bar{p})^2 a \cdot F_1 \cdot F_2 \cdot a \bar{p} \cdot F_2 \cdot \tilde{F}_1 \cdot \bar{p} (a \cdot a)^2}{135m^2} \\
& - \frac{4p_1 \cdot \bar{p} (-33 (a \cdot p_2)^2 m^2 + 23a \cdot p_1 a \cdot p_2 m^2 + 60a \cdot a (p_1 \cdot \bar{p})^2) a \cdot F_1 \cdot F_2 \cdot \bar{p} \bar{p} \cdot F_2 \cdot \tilde{F}_1 \cdot \bar{p} (a \cdot a)^2}{945m^4} \\
& + \frac{212}{945} a \cdot p_1 (p_1 \cdot \bar{p})^2 a \cdot F_1 \cdot F_2 \cdot a a \cdot F_1 \cdot \tilde{F}_2 \cdot a a \cdot a + \frac{212}{945} a \cdot p_2 (p_1 \cdot \bar{p})^2 a \cdot F_1 \cdot F_2 \cdot a a \cdot F_2 \cdot \tilde{F}_1 \cdot a a \cdot a \\
& + \frac{p_1 \cdot \bar{p} (-121 (a \cdot p_1)^2 m^2 - 121 (a \cdot p_2)^2 m^2 + 190a \cdot p_1 a \cdot p_2 m^2 + 372a \cdot a (p_1 \cdot \bar{p})^2)}{630m^2} \\
& \times (a \cdot F_1 \cdot F_2 \cdot a a \cdot \tilde{F}_1 \cdot F_2 \cdot \bar{p} a \cdot a) \tag{A.1}
\end{aligned}$$

$$\begin{aligned}
\mathcal{M}_{\text{TS-FZ}}^{(c,8)} = & \frac{64 (p_1 \cdot \bar{p})^2 p_1 \cdot F_2 \cdot \bar{p} p_2 \cdot F_1 \cdot \bar{p} \bar{p} \cdot F_1 \cdot F_2 \cdot \bar{p} (a \cdot a)^4}{2835m^4} + \frac{32 (p_1 \cdot \bar{p})^2 p_1 \cdot F_2 \cdot F_1 \cdot p_2 \bar{p} \cdot F_1 \cdot F_2 \cdot \bar{p} (a \cdot a)^4}{2835m^2} \\
& + \frac{(p_1 \cdot \bar{p})^2 (m^2 (174 (a \cdot p_1)^2 - 55a \cdot p_2 a \cdot p_1 + 174 (a \cdot p_2)^2) - 278a \cdot a (p_1 \cdot \bar{p})^2) (\bar{p} \cdot F_1 \cdot F_2 \cdot \bar{p})^2 (a \cdot a)^3}{5670m^6}
\end{aligned}$$

$$\begin{aligned}
& + \frac{8(a \cdot p_1 - a \cdot p_2)}{945m^2} \left(p_1 \cdot \bar{p} a \cdot F_1 \cdot p_2 p_1 \cdot F_2 \cdot \bar{p} \bar{p} \cdot F_1 \cdot F_2 \cdot \bar{p} (a \cdot a)^3 + p_1 \cdot \bar{p} a \cdot F_2 \cdot p_1 p_2 \cdot F_1 \cdot \bar{p} \bar{p} \cdot F_1 \cdot F_2 \cdot \bar{p} (a \cdot a)^3 \right) \\
& + \frac{p_1 \cdot \bar{p} (m^2 a \cdot p_1 (936a \cdot p_1 - 883a \cdot p_2) - 483a \cdot a (p_1 \cdot \bar{p})^2) p_1 \cdot F_2 \cdot F_1 \cdot \bar{p} \bar{p} \cdot F_1 \cdot F_2 \cdot \bar{p} (a \cdot a)^3}{11340m^4} \\
& + \frac{p_1 \cdot \bar{p} (a \cdot p_2 (883a \cdot p_1 - 936a \cdot p_2) m^2 + 483a \cdot a (p_1 \cdot \bar{p})^2) p_2 \cdot F_1 \cdot F_2 \cdot \bar{p} \bar{p} \cdot F_1 \cdot F_2 \cdot \bar{p} (a \cdot a)^3}{11340m^4} \\
& + \frac{(p_1 \cdot \bar{p})^2}{5670m^4} \left(-13m^2 (132(a \cdot p_1)^2 - 223a \cdot p_2 a \cdot p_1 + 132(a \cdot p_2)^2) \right. \\
& \left. + 1987a \cdot a (p_1 \cdot \bar{p})^2 \right) a \cdot F_1 \cdot F_2 \cdot a \bar{p} \cdot F_1 \cdot F_2 \cdot \bar{p} (a \cdot a)^2 \\
& + \frac{(p_1 \cdot \bar{p})^2 (2m^2 (255(a \cdot p_1)^2 - 443a \cdot p_2 a \cdot p_1 + 255(a \cdot p_2)^2) - 487a \cdot a (p_1 \cdot \bar{p})^2) (a \cdot F_1 \cdot F_2 \cdot a)^2 a \cdot a}{5670m^2} \\
& + \frac{a \cdot p_1 p_1 \cdot \bar{p} (3m^2 a \cdot p_1 (33a \cdot p_1 - 73a \cdot p_2) - 859a \cdot a (p_1 \cdot \bar{p})^2) a \cdot F_1 \cdot F_2 \cdot a a \cdot F_1 \cdot F_2 \cdot \bar{p} a \cdot a}{1890m^2} \\
& + \frac{a \cdot p_2 p_1 \cdot \bar{p} (3a \cdot p_2 (73a \cdot p_1 - 33a \cdot p_2) m^2 + 859a \cdot a (p_1 \cdot \bar{p})^2) a \cdot F_1 \cdot F_2 \cdot a a \cdot F_2 \cdot F_1 \cdot \bar{p} a \cdot a}{1890m^2} \\
& + \frac{(a \cdot a)^2 p_1 \cdot \bar{p}}{5670m^4} \left(3(-72(a \cdot p_1)^3 + 136a \cdot p_2 (a \cdot p_1)^2 + 83(a \cdot p_2)^2 a \cdot p_1 - 27(a \cdot p_2)^3) m^2 \right. \\
& \left. + a \cdot a (2288a \cdot p_1 + 289a \cdot p_2) (p_1 \cdot \bar{p})^2 \right) a \cdot F_1 \cdot F_2 \cdot \bar{p} \bar{p} \cdot F_1 \cdot F_2 \cdot \bar{p} \\
& + \frac{(a \cdot a)^2 p_1 \cdot \bar{p}}{5670m^4} \left(3m^2 (27(a \cdot p_1)^3 - 83a \cdot p_2 (a \cdot p_1)^2 - 136(a \cdot p_2)^2 a \cdot p_1 + 72(a \cdot p_2)^3) \right. \\
& \left. - a \cdot a (289a \cdot p_1 + 2288a \cdot p_2) (p_1 \cdot \bar{p})^2 \right) a \cdot F_2 \cdot F_1 \cdot \bar{p} \bar{p} \cdot F_1 \cdot F_2 \cdot \bar{p} \tag{A.2}
\end{aligned}$$

B UV behaviours

Here we discuss the UV behaviour of the one-loop integral (3.1). Before expanding in spin, the potential divergences can be avoided, when we shift the spin vector to the imaginary axis $a_j \rightarrow i\tilde{a}_j$. We first illustrate this process with a simple example. Consider the double-copy contribution to the Compton amplitude eq. (2.3), of which the spin-dependence is given in eq. (2.4). Naively, the worst UV behaviour would have come from the term with $G_1(x_1)G_2(x_2)$, yielding the highest UV scaling at one loop as follows,

$$\int \frac{d^D \ell_1}{\pi^{D/2}} \frac{\delta(m_2 v_2 \cdot \ell_1)}{\ell_1^2 \ell_2^2} G_1(\ell_1 \cdot a_1) G_1(\ell_2 \cdot a_1) \cosh(\ell_1 \cdot a_2) \cosh(\ell_2 \cdot a_2) \ell_1^\mu \ell_1^\nu, \tag{B.1}$$

where we have implicitly $\ell_2 = q - \ell_1$ as always. Performing the shift $a_j^\mu = i\tilde{a}_j^\mu$, the exponential of $\ell_1 \cdot a_j$ becomes finite. That is, we have the following scaling as $\ell_1 \rightarrow \Lambda \ell_1$

and $\Lambda \rightarrow \infty$,

$$\cosh(\ell_1 \cdot a_j) \sim \mathcal{O}(\Lambda^0), \quad G_1(\ell_1 \cdot a_j) \sim \mathcal{O}(\Lambda^{-1}), \quad G_2(\ell_1 \cdot a_j, \ell_2 \cdot a_j) \sim \mathcal{O}(\Lambda^{-1}). \quad (\text{B.2})$$

This way, we see that the one-loop integral above is well regulated.

We note that the particular derivatives of the entire functions appearing in eq. (2.3) are more suppressed, namely

$$(\partial_{a_1 \cdot \ell_1} - \partial_{a_1 \cdot \ell_2}) G_2(a_1 \cdot \ell_1, a_1 \cdot \ell_2) \sim \mathcal{O}(\Lambda^{-2}), \quad (\text{B.3})$$

$$(\partial_{a_1 \cdot \ell_1} - \partial_{a_1 \cdot \ell_2}) \left(G_1(a_1 \cdot \ell_1) G_1(a_1 \cdot \ell_2) \right) \sim \mathcal{O}(\Lambda^{-2}). \quad (\text{B.4})$$

Since all the entire functions involved in eq. (2.3) are all finite under the shift of the spin vector, the UV scaling boils down to the remaining factors given in section 2.1, which can be straightforwardly examined. We find that all the terms in eq. (3.1) are indeed well-defined under the shift in a similar fashion, which allows us to evaluate it. In the end, we can analytically continue back to the original real values of the spin vectors.

C Decomposition of tensor integrals

Here we derive the closed-form expression for the decompositions of the tensor integrals (2.11). The variables given in eq. (3.6) satisfy the orthogonality relations below,

$$q \cdot \theta_i = \theta_1 \cdot \theta_2 = 0, \quad v_i \cdot \theta_j = \delta_{ij}, \quad q_\mu \Pi^{\mu\nu} = \theta_{i\mu} \Pi^{\mu\nu} = v_{i\mu} \Pi^{\mu\nu} = 0. \quad (\text{C.1})$$

This yields the orthogonality of the tensor basis constructed from these building blocks.

We denote the (tensor) integrals as

$$\mathcal{I}_{\alpha_1, \alpha_2, \alpha_3, \alpha_4}[f(\ell_1)] \equiv \int \frac{d^D \ell_1}{\pi^{D/2}} \frac{\delta^{(\alpha_4-1)}(\ell_1 \cdot v_2) f(\ell_1)}{\ell_1^{2\alpha_1} (q - \ell_1)^{2\alpha_2} (\ell_1 \cdot v_1)^{\alpha_3}}. \quad (\text{C.2})$$

We consider the special case where we only have non-vanishing integrals with $\alpha_1 = \alpha_2 = \alpha_4 = 1$, since the integrand in eq. (3.1) can not have higher propagator powers in α_1 or α_2 and ultra-local terms (negative values of α_1 or α_2) are excluded. The scalar integral family is simply $\mathcal{I}_{1,1,\alpha_3,1}[1]$. The two types of integrals in (3.5) are $\mathcal{I}_{1,1,\alpha_3,1}[(\ell_1 \cdot a_1)^M (\ell_1 \cdot a_2)^{N-M}]$ and $\mathcal{I}_{1,1,\alpha_3,1}[(\ell_1 \cdot a_1)^M (\ell_1 \cdot a_2)^{N-M} \ell^\mu]$.

We consider the tensor numerator with free indices first. The most general ansatz for the decomposition of $\mathcal{I}_{1,1,\alpha_3,1}[\ell_1^{\mu_1} \cdots \ell_1^{\mu_N}]$ reads

$$\mathcal{I}_{1,1,\alpha_3,1}[\ell_1^{\mu_1} \cdots \ell_1^{\mu_N}] = \sum_{\substack{n_1+n_2+2n_3=N, \\ n_i \geq 0, n_i \in \mathbb{Z}}} \mathbb{C}_{n_1 n_2 n_3} \text{sym} \left[\theta_1^{\mu_1} \cdots \theta_1^{\mu_{n_1}} q^{\mu_{n_1+1}} \cdots q^{\mu_{n_1+n_2}} \right]$$

$$\Pi^{\mu_{n_1+n_2+1}\mu_{n_1+n_2+2}} \dots \Pi^{\mu_{N-1}\mu_N}], \quad (\text{C.3})$$

where $\text{sym}[\dots]$ denotes the sum of distinct tensor structures obtained from index permutations. Generally speaking, the coefficients $\mathbb{C}_{n_1 n_2 n_3}$ can be obtained by contracting both sides of the ansatz with all possible rank- n tensor structures built from $\{v_i^\mu, q_i^\mu, \Pi^{\mu\nu}\}$ and solving the linear equations. Thanks to the orthogonality of the tensor structures, these equations are already diagonalized and we can simply read off the coefficients $\mathbb{C}_{n_1 n_2 n_3}$ as follows,

$$\mathbb{C}_{n_1 n_2 n_3} = \frac{1}{N_{n_3}} \mathcal{I}_{1,1,\alpha_3,1} [(\ell_1 \cdot v_1)^{n_1} (\ell_1 \cdot q)^{n_2} (\ell_1 \cdot \Pi \cdot \ell_1)^{n_3}], \quad (\text{C.4})$$

where we have $N_{n_3} = (D-3)(D-1) \dots (D-3+2(n_3-1))$ for $n_3 > 0$ and $N_0 = 1$. The remaining integral can be rewritten in terms of scalar integrals,

$$\mathcal{I}_{1,1,\alpha_3,1} [(\ell_1 \cdot v_1)^{n_1} (\ell_1 \cdot q)^{n_2} (\ell_1 \cdot \Pi \cdot \ell_1)^{n_3}] = \sum_{k=0}^{n_3} \binom{n_3}{k} \frac{1}{(\gamma^2 - 1)^k} \left(\frac{q^2}{2}\right)^{n_2+n_3-k} \mathcal{I}_{1,1,\alpha_3-n_1-2k,1}, \quad (\text{C.5})$$

where we have repeatedly used $\ell_1 \cdot q = \frac{1}{2}(\ell_1^2 + q^2 - (q - \ell_1)^2)$ and that scaleless integrals are vanishing.

The contraction between the symmetrized tensor structures in (C.3) and the spin vectors can be worked out. We have

$$\begin{aligned} \text{Cont}(N, M, n_1, n_2) &\equiv \text{sym}[\overbrace{\theta_1 \dots \theta_1}^{n_1} \overbrace{q \dots q}^{n_2} \overbrace{\Pi \dots \Pi}^{n_3}] \underbrace{a_1 \dots a_1}_M \underbrace{a_2 \dots a_2}_{N-M} \\ &= \sum_{\text{cond.}} C_{N,M,n_1,n_2,m_1,m_2,m_3} (a_1 \cdot \theta_1)^{m_1} (a_2 \cdot \theta_1)^{n_1-m_1} (a_1 \cdot q)^{m_2} (a_2 \cdot q)^{n_2-m_2} \\ &\quad (a_1 \cdot \Pi \cdot a_1)^{m_3} (a_1 \cdot \Pi \cdot a_2)^{m_4} (a_2 \cdot \Pi \cdot a_2)^{n_3-m_3-m_4}, \end{aligned} \quad (\text{C.6})$$

where the sums are taken over the solutions to the conditions below

$$m_1 + m_2 + 2m_3 + m_4 = M, \quad 0 \leq m_i \leq n_i, \quad 0 \leq m_4 \leq n_3, \quad m_i \in \mathbb{Z}. \quad (\text{C.7})$$

The coefficients read

$$C_{N,M,n_1,n_2,m_1,m_2,m_3} = \frac{M!}{m_1! m_2! m_3! m_4!} \frac{(N-M)!}{(n_1-m_1)! (n_2-m_2)! (n_3-m_3)!} \frac{1}{2^{n_3-m_4}}. \quad (\text{C.8})$$

Hence we have arrived at the decomposition of $\mathcal{I}_{1,1,\alpha_3,1}[(\ell_1 \cdot a_1)^M (\ell_1 \cdot a_2)^{N-M}]$.

The remaining integral $\mathcal{I}_{1,1,\alpha_3,1}[(\ell_1 \cdot a_1)^M (\ell_1 \cdot a_2)^{N-M} \ell_1^\mu]$ is nothing but the above with one free index and can be dealt with similarly. We first rewrite the symmetrized sum of the tensor structures as follows,

$$\begin{aligned}
\text{sym}[\overbrace{\theta_1 \cdots \theta_1}^{n_1} \overbrace{q \cdots q}^{n_2} \overbrace{\Pi \cdots \Pi}^{n_3}] &= \theta_1^{\mu N} \text{sym}[\overbrace{\theta_1 \cdots \theta_1}^{n_1-1} \overbrace{q \cdots q}^{n_2} \overbrace{\Pi \cdots \Pi}^{n_3}] \\
&+ q^{\mu N} \text{sym}[\overbrace{\theta_1 \cdots \theta_1}^{n_1} \overbrace{q \cdots q}^{n_2-1} \overbrace{\Pi \cdots \Pi}^{n_3}] \\
&+ \sum_{i=1}^{N-1} \Pi^{\mu_i \mu_N} \text{sym}[\overbrace{\theta_1 \cdots \theta_1}^{n_1} \overbrace{q \cdots q}^{n_2} \overbrace{\Pi \cdots \Pi}^{n_3-1}]. \quad (\text{C.9})
\end{aligned}$$

Contracting with $a_1^{\mu_1} \cdots a_1^{\mu_M} a_2^{\mu_{M+1}} \cdots a_2^{\mu_{N-M-1}}$, we have

$$\begin{aligned}
&\text{sym}[\overbrace{\theta_1 \cdots \theta_1}^{n_1} \overbrace{q \cdots q}^{n_2} \overbrace{\Pi \cdots \Pi}^{n_3}] \underbrace{a_1 \cdots a_1}_M \underbrace{a_2 \cdots a_2}_{N-1-M} \\
= &\theta_1^{\mu N} \text{Cont}(N-1, M, n_1-1, n_2) + q^{\mu N} \text{Cont}(N-1, M, n_1, n_2-1) \quad (\text{C.10}) \\
&+ M(a_1 \cdot \Pi)^{\mu N} \text{Cont}(N-2, M-1, n_1, n_2) + (N-1-M)(a_2 \cdot \Pi)^{\mu N} \text{Cont}(N-2, M, n_1, n_2).
\end{aligned}$$

References

- [1] LIGO SCIENTIFIC, VIRGO collaboration, *Binary Black Hole Mergers in the first Advanced LIGO Observing Run*, *Phys. Rev. X* **6** (2016) 041015 [[1606.04856](#)].
- [2] LIGO SCIENTIFIC, VIRGO collaboration, *Observation of Gravitational Waves from a Binary Black Hole Merger*, *Phys. Rev. Lett.* **116** (2016) 061102 [[1602.03837](#)].
- [3] LIGO SCIENTIFIC, VIRGO collaboration, *GW151226: Observation of Gravitational Waves from a 22-Solar-Mass Binary Black Hole Coalescence*, *Phys. Rev. Lett.* **116** (2016) 241103 [[1606.04855](#)].
- [4] LIGO SCIENTIFIC, VIRGO collaboration, *GW170104: Observation of a 50-Solar-Mass Binary Black Hole Coalescence at Redshift 0.2*, *Phys. Rev. Lett.* **118** (2017) 221101 [[1706.01812](#)].
- [5] LIGO SCIENTIFIC, VIRGO collaboration, *GW170817: Observation of Gravitational Waves from a Binary Neutron Star Inspiral*, *Phys. Rev. Lett.* **119** (2017) 161101 [[1710.05832](#)].
- [6] F. Pretorius, *Evolution of binary black hole spacetimes*, *Phys. Rev. Lett.* **95** (2005) 121101 [[gr-qc/0507014](#)].

- [7] M. Campanelli, C.O. Lousto, P. Marronetti and Y. Zlochower, *Accurate evolutions of orbiting black-hole binaries without excision*, *Phys. Rev. Lett.* **96** (2006) 111101 [[gr-qc/0511048](#)].
- [8] J.G. Baker, J. Centrella, D.-I. Choi, M. Koppitz and J. van Meter, *Gravitational wave extraction from an inspiraling configuration of merging black holes*, *Phys. Rev. Lett.* **96** (2006) 111102 [[gr-qc/0511103](#)].
- [9] A. Buonanno and T. Damour, *Effective one-body approach to general relativistic two-body dynamics*, *Phys. Rev.* **D59** (1999) 084006 [[gr-qc/9811091](#)].
- [10] W.D. Goldberger and I.Z. Rothstein, *An Effective field theory of gravity for extended objects*, *Phys. Rev. D* **73** (2006) 104029 [[hep-th/0409156](#)].
- [11] B. Kol and M. Smolkin, *Classical Effective Field Theory and Caged Black Holes*, *Phys. Rev. D* **77** (2008) 064033 [[0712.2822](#)].
- [12] J.B. Gilmore and A. Ross, *Effective field theory calculation of second post-Newtonian binary dynamics*, *Phys. Rev. D* **78** (2008) 124021 [[0810.1328](#)].
- [13] S. Foffa and R. Sturani, *Effective field theory calculation of conservative binary dynamics at third post-Newtonian order*, *Phys. Rev. D* **84** (2011) 044031 [[1104.1122](#)].
- [14] S. Foffa, P. Mastrolia, R. Sturani and C. Sturm, *Effective field theory approach to the gravitational two-body dynamics, at fourth post-Newtonian order and quintic in the Newton constant*, *Phys. Rev. D* **95** (2017) 104009 [[1612.00482](#)].
- [15] R.A. Porto and I.Z. Rothstein, *Apparent ambiguities in the post-Newtonian expansion for binary systems*, *Phys. Rev. D* **96** (2017) 024062 [[1703.06433](#)].
- [16] J. Blümlein, A. Maier and P. Marquard, *Five-Loop Static Contribution to the Gravitational Interaction Potential of Two Point Masses*, *Phys. Lett. B* **800** (2020) 135100 [[1902.11180](#)].
- [17] S. Foffa and R. Sturani, *Conservative dynamics of binary systems to fourth Post-Newtonian order in the EFT approach I: Regularized Lagrangian*, *Phys. Rev. D* **100** (2019) 024047 [[1903.05113](#)].
- [18] S. Foffa, R.A. Porto, I. Rothstein and R. Sturani, *Conservative dynamics of binary systems to fourth Post-Newtonian order in the EFT approach II: Renormalized Lagrangian*, *Phys. Rev. D* **100** (2019) 024048 [[1903.05118](#)].
- [19] J. Blümlein, A. Maier, P. Marquard and G. Schäfer, *Fourth post-Newtonian Hamiltonian dynamics of two-body systems from an effective field theory approach*, *Nucl. Phys. B* **955** (2020) 115041 [[2003.01692](#)].
- [20] J. Bluemlein, A. Maier, P. Marquard and G. Schaefer, *Testing binary dynamics in gravity at the sixth post-Newtonian level*, [2003.07145](#).

- [21] D. Bini, T. Damour and A. Geralico, *Sixth post-Newtonian local-in-time dynamics of binary systems*, *Phys. Rev. D* **102** (2020) 024061 [[2004.05407](#)].
- [22] D. Bini, T. Damour and A. Geralico, *Sixth post-Newtonian nonlocal-in-time dynamics of binary systems*, *Phys. Rev. D* **102** (2020) 084047 [[2007.11239](#)].
- [23] J. Blümlein, A. Maier, P. Marquard and G. Schäfer, *The fifth-order post-Newtonian Hamiltonian dynamics of two-body systems from an effective field theory approach*, *Nucl. Phys. B* **983** (2022) 115900 [[2110.13822](#)].
- [24] J. Blümlein, A. Maier, P. Marquard and G. Schäfer, *The fifth-order post-Newtonian Hamiltonian dynamics of two-body systems from an effective field theory approach: potential contributions*, [2010.13672](#).
- [25] S. Foffa, R. Sturani and W.J. Torres Bobadilla, *Efficient resummation of high post-Newtonian contributions to the binding energy*, *JHEP* **02** (2021) 165 [[2010.13730](#)].
- [26] J. Blümlein, A. Maier, P. Marquard and G. Schäfer, *The 6th post-Newtonian potential terms at $O(G_N^4)$* , *Phys. Lett. B* **816** (2021) 136260 [[2101.08630](#)].
- [27] J.-W. Kim, M. Levi and Z. Yin, *Quadratic-in-spin interactions at fifth post-Newtonian order probe new physics*, *Phys. Lett. B* **834** (2022) 137410 [[2112.01509](#)].
- [28] G. Cho, R.A. Porto and Z. Yang, *Gravitational radiation from inspiralling compact objects: Spin effects to fourth Post-Newtonian order*, [2201.05138](#).
- [29] N.E.J. Bjerrum-Bohr, P.H. Damgaard, G. Festuccia, L. Plante and P. Vanhove, *General Relativity from Scattering Amplitudes*, *Phys. Rev. Lett.* **121** (2018) 171601 [[1806.04920](#)].
- [30] C. Cheung, I.Z. Rothstein and M.P. Solon, *From Scattering Amplitudes to Classical Potentials in the Post-Minkowskian Expansion*, *Phys. Rev. Lett.* **121** (2018) 251101 [[1808.02489](#)].
- [31] Z. Bern, C. Cheung, R. Roiban, C.-H. Shen, M.P. Solon and M. Zeng, *Scattering Amplitudes and the Conservative Hamiltonian for Binary Systems at Third Post-Minkowskian Order*, *Phys. Rev. Lett.* **122** (2019) 201603 [[1901.04424](#)].
- [32] Z. Bern, C. Cheung, R. Roiban, C.-H. Shen, M.P. Solon and M. Zeng, *Black Hole Binary Dynamics from the Double Copy and Effective Theory*, *JHEP* **10** (2019) 206 [[1908.01493](#)].
- [33] D. Neill and I.Z. Rothstein, *Classical Space-Times from the S Matrix*, *Nucl. Phys.* **B877** (2013) 177 [[1304.7263](#)].
- [34] A. Cristofoli, R. Gonzo, D.A. Kosower and D. O’Connell, *Waveforms from Amplitudes*, [2107.10193](#).

- [35] Z. Bern, A. Luna, R. Roiban, C.-H. Shen and M. Zeng, *Spinning black hole binary dynamics, scattering amplitudes, and effective field theory*, *Phys. Rev. D* **104** (2021) 065014 [[2005.03071](#)].
- [36] Z. Bern, J. Parra-Martinez, R. Roiban, M.S. Ruf, C.-H. Shen, M.P. Solon et al., *Scattering Amplitudes and Conservative Binary Dynamics at $\mathcal{O}(G^4)$* , *Phys. Rev. Lett.* **126** (2021) 171601 [[2101.07254](#)].
- [37] N.E.J. Bjerrum-Bohr, J.F. Donoghue and B.R. Holstein, *Quantum gravitational corrections to the nonrelativistic scattering potential of two masses*, *Phys. Rev. D* **67** (2003) 084033 [[hep-th/0211072](#)].
- [38] D.A. Kosower, B. Maybee and D. O’Connell, *Amplitudes, Observables, and Classical Scattering*, *JHEP* **02** (2019) 137 [[1811.10950](#)].
- [39] B. Maybee, D. O’Connell and J. Vines, *Observables and amplitudes for spinning particles and black holes*, *JHEP* **12** (2019) 156 [[1906.09260](#)].
- [40] A. Brandhuber, G. Chen, G. Travaglini and C. Wen, *A new gauge-invariant double copy for heavy-mass effective theory*, *JHEP* **07** (2021) 047 [[2104.11206](#)].
- [41] A. Brandhuber, G. Chen, G. Travaglini and C. Wen, *Classical gravitational scattering from a gauge-invariant double copy*, *JHEP* **10** (2021) 118 [[2108.04216](#)].
- [42] G. Mogull, J. Plefka and J. Steinhoff, *Classical black hole scattering from a worldline quantum field theory*, *JHEP* **02** (2021) 048 [[2010.02865](#)].
- [43] G.U. Jakobsen, G. Mogull, J. Plefka and J. Steinhoff, *Classical Gravitational Bremsstrahlung from a Worldline Quantum Field Theory*, *Phys. Rev. Lett.* **126** (2021) 201103 [[2101.12688](#)].
- [44] G.U. Jakobsen and G. Mogull, *Conservative and Radiative Dynamics of Spinning Bodies at Third Post-Minkowskian Order Using Worldline Quantum Field Theory*, *Phys. Rev. Lett.* **128** (2022) 141102 [[2201.07778](#)].
- [45] T. Wang, *Binary dynamics from worldline QFT for scalar QED*, *Phys. Rev. D* **107** (2023) 085011 [[2205.15753](#)].
- [46] J. Parra-Martinez, M.S. Ruf and M. Zeng, *Extremal black hole scattering at $\mathcal{O}(G^3)$: graviton dominance, eikonal exponentiation, and differential equations*, *JHEP* **11** (2020) 023 [[2005.04236](#)].
- [47] P. Di Vecchia, C. Heissenberg, R. Russo and G. Veneziano, *The eikonal approach to gravitational scattering and radiation at $\mathcal{O}(G^3)$* , *JHEP* **07** (2021) 169 [[2104.03256](#)].
- [48] C. Heissenberg, *Infrared divergences and the eikonal exponentiation*, *Phys. Rev. D* **104** (2021) 046016 [[2105.04594](#)].
- [49] P. Di Vecchia, C. Heissenberg, R. Russo and G. Veneziano, *The eikonal operator at*

arbitrary velocities I: the soft-radiation limit, [2204.02378](#).

- [50] P.H. Damgaard, L. Plante and P. Vanhove, *On an exponential representation of the gravitational S-matrix*, *JHEP* **11** (2021) 213 [[2107.12891](#)].
- [51] U. Kol, D. O’connell and O. Telem, *The radial action from probe amplitudes to all orders*, *JHEP* **03** (2022) 141 [[2109.12092](#)].
- [52] N.E.J. Bjerrum-Bohr, L. Planté and P. Vanhove, *Post-Minkowskian radial action from soft limits and velocity cuts*, *JHEP* **03** (2022) 071 [[2111.02976](#)].
- [53] Z. Bern, J.P. Gatica, E. Herrmann, A. Luna and M. Zeng, *Scalar QED as a toy model for higher-order effects in classical gravitational scattering*, [2112.12243](#).
- [54] H. Lee, S. Lee and S. Mazumdar, *Classical observables from partial wave amplitudes*, *JHEP* **06** (2023) 096 [[2303.07638](#)].
- [55] H. Lee, K. Lee and S. Lee, *Poincaré generators at second post-Minkowskian order*, *JHEP* **10** (2023) 044 [[2307.05626](#)].
- [56] C. Dlapa, G. Kälin, Z. Liu and R.A. Porto, *Dynamics of binary systems to fourth Post-Minkowskian order from the effective field theory approach*, *Phys. Lett. B* **831** (2022) 137203 [[2106.08276](#)].
- [57] C. Dlapa, G. Kälin, Z. Liu, J. Neef and R.A. Porto, *Radiation Reaction and Gravitational Waves at Fourth Post-Minkowskian Order*, [2210.05541](#).
- [58] C. Dlapa, G. Kälin, Z. Liu and R.A. Porto, *Bootstrapping the relativistic two-body problem*, *JHEP* **08** (2023) 109 [[2304.01275](#)].
- [59] G.U. Jakobsen, G. Mogull, J. Plefka and B. Sauer, *Tidal effects and renormalization at fourth post-Minkowskian order*, *Phys. Rev. D* **109** (2024) L041504 [[2312.00719](#)].
- [60] Z. Bern, J. Parra-Martinez, R. Roiban, M.S. Ruf, C.-H. Shen, M.P. Solon et al., *Scattering Amplitudes, the Tail Effect, and Conservative Binary Dynamics at $O(G^4)$* , *Phys. Rev. Lett.* **128** (2022) 161103 [[2112.10750](#)].
- [61] P.H. Damgaard, E.R. Hansen, L. Planté and P. Vanhove, *Classical Observables from the Exponential Representation of the Gravitational S-Matrix*, [2307.04746](#).
- [62] M. Driesse, G.U. Jakobsen, G. Mogull, J. Plefka, B. Sauer and J. Usovitsch, *Conservative Black Hole Scattering at Fifth Post-Minkowskian and First Self-Force Order*, [2403.07781](#).
- [63] N. Arkani-Hamed, T.-C. Huang and Y.-t. Huang, *Scattering amplitudes for all masses and spins*, *JHEP* **11** (2021) 070 [[1709.04891](#)].
- [64] Y.M. Zinoviev, *On massive high spin particles in AdS*, [hep-th/0108192](#).
- [65] Y.M. Zinoviev, *On massive spin 2 interactions*, *Nucl. Phys. B* **770** (2007) 83 [[hep-th/0609170](#)].

- [66] Y.M. Zinoviev, *On massive spin 2 electromagnetic interactions*, *Nucl. Phys. B* **821** (2009) 431 [[0901.3462](#)].
- [67] Y.M. Zinoviev, *On spin 3 interacting with gravity*, *Class. Quant. Grav.* **26** (2009) 035022 [[0805.2226](#)].
- [68] A. Ochirov and E. Skvortsov, *Chiral Approach to Massive Higher Spins*, *Phys. Rev. Lett.* **129** (2022) 241601 [[2207.14597](#)].
- [69] L. Cangemi and P. Pichini, *Classical limit of higher-spin string amplitudes*, *JHEP* **06** (2023) 167 [[2207.03947](#)].
- [70] Z. Bern, J.J.M. Carrasco and H. Johansson, *New Relations for Gauge-Theory Amplitudes*, *Phys. Rev.* **D78** (2008) 085011 [[0805.3993](#)].
- [71] Z. Bern, J.J.M. Carrasco and H. Johansson, *Perturbative Quantum Gravity as a Double Copy of Gauge Theory*, *Phys. Rev. Lett.* **105** (2010) 061602 [[1004.0476](#)].
- [72] Z. Bern, J.J. Carrasco, M. Chiodaroli, H. Johansson and R. Roiban, *The Duality Between Color and Kinematics and its Applications*, [1909.01358](#).
- [73] N.E.J. Bjerrum-Bohr, G. Chen and M. Skowronek, *Classical spin gravitational Compton scattering*, *JHEP* **06** (2023) 170 [[2302.00498](#)].
- [74] N.E.J. Bjerrum-Bohr, G. Chen and M. Skowronek, *Covariant Compton Amplitudes in Gravity with Classical Spin*, *Phys. Rev. Lett.* **132** (2024) 191603 [[2309.11249](#)].
- [75] S.A. Teukolsky, *Perturbations of a rotating black hole. 1. Fundamental equations for gravitational electromagnetic and neutrino field perturbations*, *Astrophys. J.* **185** (1973) 635.
- [76] W.H. Press and S.A. Teukolsky, *Perturbations of a Rotating Black Hole. II. Dynamical Stability of the Kerr Metric*, *Astrophys. J.* **185** (1973) 649.
- [77] S.A. Teukolsky and W.H. Press, *Perturbations of a rotating black hole. III - Interaction of the hole with gravitational and electromagnetic radiation*, *Astrophys. J.* **193** (1974) 443.
- [78] H.S. Chia, *Tidal deformation and dissipation of rotating black holes*, *Phys. Rev. D* **104** (2021) 024013 [[2010.07300](#)].
- [79] Y.F. Bautista, A. Guevara, C. Kavanagh and J. Vines, *From Scattering in Black Hole Backgrounds to Higher-Spin Amplitudes: Part I*, [2107.10179](#).
- [80] M.M. Ivanov and Z. Zhou, *Vanishing of Black Hole Tidal Love Numbers from Scattering Amplitudes*, *Phys. Rev. Lett.* **130** (2023) 091403 [[2209.14324](#)].
- [81] Y.F. Bautista, A. Guevara, C. Kavanagh and J. Vines, *Scattering in black hole backgrounds and higher-spin amplitudes. Part II*, *JHEP* **05** (2023) 211 [[2212.07965](#)].
- [82] M.V.S. Saketh, Z. Zhou and M.M. Ivanov, *Dynamical tidal response of Kerr black*

- holes from scattering amplitudes, *Phys. Rev. D* **109** (2024) 064058 [2307.10391].
- [83] Y.F. Bautista, G. Bonelli, C. Iossa, A. Tanzini and Z. Zhou, *Black Hole Perturbation Theory Meets CFT₂: Kerr Compton Amplitudes from Nekrasov-Shatashvili Functions*, [2312.05965](#).
- [84] M.-Z. Chung, Y.-T. Huang, J.-W. Kim and S. Lee, *The simplest massive S-matrix: from minimal coupling to Black Holes*, *JHEP* **04** (2019) 156 [1812.08752].
- [85] M.-Z. Chung, Y.-T. Huang and J.-W. Kim, *Classical potential for general spinning bodies*, *JHEP* **09** (2020) 074 [1908.08463].
- [86] M.-Z. Chung, Y.-t. Huang, J.-W. Kim and S. Lee, *Complete Hamiltonian for spinning binary systems at first post-Minkowskian order*, *JHEP* **05** (2020) 105 [2003.06600].
- [87] W.-M. Chen, M.-Z. Chung, Y.-t. Huang and J.-W. Kim, *The 2PM Hamiltonian for binary Kerr to quartic in spin*, *JHEP* **08** (2022) 148 [2111.13639].
- [88] R. Aoude, K. Haddad and A. Helset, *Searching for Kerr in the 2PM amplitude*, *JHEP* **07** (2022) 072 [2203.06197].
- [89] R. Aoude, K. Haddad and A. Helset, *Classical Gravitational Spinning-Spinless Scattering at $O(G^2S^\infty)$* , *Phys. Rev. Lett.* **129** (2022) 141102 [2205.02809].
- [90] Z. Bern, D. Kosmopoulos, A. Luna, R. Roiban and F. Teng, *Binary Dynamics Through the Fifth Power of Spin at $\mathcal{O}(G^2)$* , [2203.06202](#).
- [91] Z. Bern, D. Kosmopoulos, A. Luna, R. Roiban, T. Scheopner, F. Teng et al., *Quantum Field Theory, Worldline Theory, and Spin Magnitude Change in Orbital Evolution*, [2308.14176](#).
- [92] G. Menezes and M. Sergola, *NLO deflections for spinning particles and Kerr black holes*, *JHEP* **10** (2022) 105 [2205.11701].
- [93] F. Febres Cordero, M. Kraus, G. Lin, M.S. Ruf and M. Zeng, *Conservative Binary Dynamics with a Spinning Black Hole at $O(G^3)$ from Scattering Amplitudes*, *Phys. Rev. Lett.* **130** (2023) 021601 [2205.07357].
- [94] L. Cangemi, M. Chiodaroli, H. Johansson, A. Ochirov, P. Pichini and E. Skvortsov, *Kerr Black Holes Enjoy Massive Higher-Spin Gauge Symmetry*, [2212.06120](#).
- [95] K. Haddad, *Recursion in the classical limit and the neutron-star Compton amplitude*, *JHEP* **05** (2023) 177 [2303.02624].
- [96] S. De Angelis, R. Gonzo and P.P. Novichkov, *Spinning waveforms from KMOC at leading order*, [2309.17429](#).
- [97] Y.-J. Chen, T. Hsieh, Y.-T. Huang and J.-W. Kim, *On-shell approach to (spinning) gravitational absorption processes*, [2312.04513](#).
- [98] L. Cangemi, M. Chiodaroli, H. Johansson, A. Ochirov, P. Pichini and E. Skvortsov,

From higher-spin gauge interactions to Compton amplitudes for root-Kerr,
[2311.14668](#).

- [99] L. Cangemi, M. Chiodaroli, H. Johansson, A. Ochirov, P. Pichini and E. Skvortsov, *Compton Amplitude for Rotating Black Hole from QFT*, [2312.14913](#).
- [100] J. Vines, *Scattering of two spinning black holes in post-Minkowskian gravity, to all orders in spin, and effective-one-body mappings*, *Class. Quant. Grav.* **35** (2018) 084002 [[1709.06016](#)].
- [101] A. Guevara, A. Ochirov and J. Vines, *Scattering of Spinning Black Holes from Exponentiated Soft Factors*, *JHEP* **09** (2019) 056 [[1812.06895](#)].
- [102] D. Kosmopoulos and A. Luna, *Quadratic-in-spin Hamiltonian at $\mathcal{O}(G^2)$ from scattering amplitudes*, *JHEP* **07** (2021) 037 [[2102.10137](#)].
- [103] T. Scheopner and J. Vines, *Dynamical Implications of the Kerr Multipole Moments for Spinning Black Holes*, [2311.18421](#).
- [104] F. Alessio and P. Di Vecchia, *Radiation reaction for spinning black-hole scattering*, *Phys. Lett. B* **832** (2022) 137258 [[2203.13272](#)].
- [105] A. Elkhidir, D. O’Connell, M. Sergola and I.A. Vazquez-Holm, *Radiation and Reaction at One Loop*, [2303.06211](#).
- [106] D. Bini, T. Damour and A. Geralico, *Comparing one-loop gravitational bremsstrahlung amplitudes to the multipolar-post-Minkowskian waveform*, *Phys. Rev. D* **108** (2023) 124052 [[2309.14925](#)].
- [107] A. Brandhuber, G.R. Brown, G. Chen, J. Gowdy and G. Travaglini, *Resummed spinning waveforms from five-point amplitudes*, *JHEP* **02** (2024) 026 [[2310.04405](#)].
- [108] A. Luna, N. Moynihan, D. O’Connell and A. Ross, *Observables from the Spinning Eikonal*, [2312.09960](#).
- [109] H. Lee and S. Lee, *Poincaré invariance of spinning binary dynamics in the post-Minkowskian Hamiltonian approach*, *Class. Quant. Grav.* **40** (2023) 245004 [[2305.10739](#)].
- [110] A. Buonanno, G.U. Jakobsen and G. Mogull, *Post-Minkowskian Theory Meets the Spinning Effective-One-Body Approach for Two-Body Scattering*, [2402.12342](#).
- [111] G.U. Jakobsen, G. Mogull, J. Plefka and J. Steinhoff, *Gravitational Bremsstrahlung and Hidden Supersymmetry of Spinning Bodies*, [2106.10256](#).
- [112] G.U. Jakobsen, G. Mogull, J. Plefka and J. Steinhoff, *SUSY in the sky with gravitons*, *JHEP* **01** (2022) 027 [[2109.04465](#)].
- [113] G.U. Jakobsen and G. Mogull, *Linear response, Hamiltonian, and radiative spinning two-body dynamics*, *Phys. Rev. D* **107** (2023) 044033 [[2210.06451](#)].

- [114] G.U. Jakobsen, G. Mogull, J. Plefka, B. Sauer and Y. Xu, *Conservative scattering of spinning black holes at fourth post-Minkowskian order*, [2306.01714](#).
- [115] G.U. Jakobsen, G. Mogull, J. Plefka and B. Sauer, *Dissipative Scattering of Spinning Black Holes at Fourth Post-Minkowskian Order*, *Phys. Rev. Lett.* **131** (2023) 241402 [[2308.11514](#)].
- [116] C. Shi and J. Plefka, *Classical double copy of worldline quantum field theory*, *Phys. Rev. D* **105** (2022) 026007 [[2109.10345](#)].
- [117] F. Comberiati and C. Shi, *Classical Double Copy of Spinning Worldline Quantum Field Theory*, *JHEP* **04** (2023) 008 [[2212.13855](#)].
- [118] R. Gonzo and C. Shi, *Boundary to bound dictionary for generic Kerr orbits*, *Phys. Rev. D* **108** (2023) 084065 [[2304.06066](#)].
- [119] R. Gonzo and C. Shi, *Scattering and bound observables for spinning particles in Kerr spacetime with generic spin orientations*, [2405.09687](#).
- [120] R. Aoude, K. Haddad, C. Heissenberg and A. Helset, *Leading-order gravitational radiation to all spin orders*, *Phys. Rev. D* **109** (2024) 036007 [[2310.05832](#)].
- [121] J.-H. Kim, J.-W. Kim and S. Lee, *Massive twistor worldline in electromagnetic fields*, [2405.17056](#).
- [122] A. Brandhuber, G.R. Brown, P. Pichini, G. Travaglini and P.V. Matasan, *Spinning binary dynamics in cubic effective field theories of gravity*, [2405.13826](#).
- [123] Z. Bern, E. Herrmann, R. Roiban, M.S. Ruf, A.V. Smirnov, V.A. Smirnov et al., *Amplitudes, Supersymmetric Black Hole Scattering at $\mathcal{O}(G^5)$, and Loop Integration*, [2406.01554](#).
- [124] G. Kälin, Z. Liu and R.A. Porto, *Conservative Dynamics of Binary Systems to Third Post-Minkowskian Order from the Effective Field Theory Approach*, *Phys. Rev. Lett.* **125** (2020) 261103 [[2007.04977](#)].
- [125] G. Kälin, Z. Liu and R.A. Porto, *Conservative Tidal Effects in Compact Binary Systems to Next-to-Leading Post-Minkowskian Order*, *Phys. Rev. D* **102** (2020) 124025 [[2008.06047](#)].
- [126] M.M. Riva and F. Vernizzi, *Radiated momentum in the post-Minkowskian worldline approach via reverse unitarity*, *JHEP* **11** (2021) 228 [[2110.10140](#)].
- [127] C. Dlapa, G. Kälin, Z. Liu and R.A. Porto, *Conservative Dynamics of Binary Systems at Fourth Post-Minkowskian Order in the Large-Eccentricity Expansion*, *Phys. Rev. Lett.* **128** (2022) 161104 [[2112.11296](#)].
- [128] W.D. Goldberger, J. Li and I.Z. Rothstein, *Non-conservative effects on spinning black holes from world-line effective field theory*, *JHEP* **06** (2021) 053 [[2012.14869](#)].

- [129] G. Kälin and R.A. Porto, *Post-Minkowskian Effective Field Theory for Conservative Binary Dynamics*, [2006.01184](#).
- [130] D. Kosmopoulos and M.P. Solon, *Gravitational self force from scattering amplitudes in curved space*, *JHEP* **03** (2024) 125 [[2308.15304](#)].
- [131] C. Cheung, J. Parra-Martinez, I.Z. Rothstein, N. Shah and J. Wilson-Gerow, *Effective Field Theory for Extreme Mass Ratios*, [2308.14832](#).
- [132] J. Wilson-Gerow, *Conservative scattering of Reissner-Nordström black holes at third post-Minkowskian order*, *JHEP* **05** (2024) 265 [[2310.17731](#)].
- [133] G.U. Jakobsen, *Spin and Susceptibility Effects of Electromagnetic Self-Force in Effective Field Theory*, *Phys. Rev. Lett.* **132** (2024) 151601 [[2311.04151](#)].
- [134] A. Klemm, C. Nega, B. Sauer and J. Plefka, *CY in the Sky*, [2401.07899](#).
- [135] N. Arkani-Hamed, Y.-t. Huang and D. O’Connell, *Kerr black holes as elementary particles*, *JHEP* **01** (2020) 046 [[1906.10100](#)].
- [136] H. Johansson and A. Ochirov, *Double copy for massive quantum particles with spin*, *JHEP* **09** (2019) 040 [[1906.12292](#)].
- [137] A. Brandhuber, G. Chen, H. Johansson, G. Travaglini and C. Wen, *Kinematic Hopf Algebra for Bern-Carrasco-Johansson Numerators in Heavy-Mass Effective Field Theory and Yang-Mills Theory*, *Phys. Rev. Lett.* **128** (2022) 121601 [[2111.15649](#)].
- [138] A. Brandhuber, G.R. Brown, G. Chen, J. Gowdy, G. Travaglini and C. Wen, *Amplitudes, Hopf algebras and the colour-kinematics duality*, *JHEP* **12** (2022) 101 [[2208.05886](#)].
- [139] G. Chen, G. Lin and C. Wen, *Kinematic Hopf algebra for amplitudes and form factors*, *Phys. Rev. D* **107** (2023) L081701 [[2208.05519](#)].
- [140] G. Chen, L. Rodina and C. Wen, *Kinematic Hopf algebra for amplitudes from higher-derivative operators*, *JHEP* **02** (2024) 096 [[2310.11943](#)].
- [141] G. Chen, L. Rodina and C. Wen, *Kinematic Hopf algebra and BCJ numerators at finite α'* , [2403.04614](#).
- [142] N.E.J. Bjerrum-Bohr, G. Chen, Y. Miao and M. Skowronek, *Color-Kinematic Numerators for Fermion Compton Amplitudes*, [2404.15265](#).
- [143] G. Chen, H. Johansson, F. Teng and T. Wang, *On the kinematic algebra for BCJ numerators beyond the MHV sector*, *JHEP* **11** (2019) 055 [[1906.10683](#)].
- [144] G. Chen, H. Johansson, F. Teng and T. Wang, *Next-to-MHV Yang-Mills kinematic algebra*, *JHEP* **10** (2021) 042 [[2104.12726](#)].
- [145] Y.F. Bautista, *Dynamics for Super-Extremal Kerr Binary Systems at $\mathcal{O}(G^2)$* , [2304.04287](#).

- [146] S.R. Dolan, *Scattering and Absorption of Gravitational Plane Waves by Rotating Black Holes*, *Class. Quant. Grav.* **25** (2008) 235002 [[0801.3805](#)].
- [147] J.-W. Kim and J. Steinhoff, *Spin supplementary condition in quantum field theory: covariant SSC and physical state projection*, *JHEP* **07** (2023) 042 [[2302.01944](#)].
- [148] G.U. Jakobsen, G. Mogull, J. Plefka and B. Sauer, *All things retarded: radiation-reaction in worldline quantum field theory*, *JHEP* **10** (2022) 128 [[2207.00569](#)].
- [149] V. Witzany, *Hamilton-Jacobi equation for spinning particles near black holes*, *Phys. Rev. D* **100** (2019) 104030 [[1903.03651](#)].
- [150] N.E.J. Bjerrum-Bohr, P.H. Damgaard, L. Planté and P. Vanhove, *Classical gravity from loop amplitudes*, *Phys. Rev. D* **104** (2021) 026009 [[2104.04510](#)].
- [151] N.E.J. Bjerrum-Bohr, P.H. Damgaard, L. Planté and P. Vanhove, *The Amplitude for Classical Gravitational Scattering at Third Post-Minkowskian Order*, [2105.05218](#).
- [152] “Kinematic hopf algebra.”
- [153] R.N. Lee, *Presenting LiteRed: a tool for the Loop InTEgrals REDuction*, [1212.2685](#).
- [154] R.N. Lee, *LiteRed 1.4: a powerful tool for reduction of multiloop integrals*, *J. Phys. Conf. Ser.* **523** (2014) 012059 [[1310.1145](#)].
- [155] L. Bohnenblust, L. Cangemi, H. Johansson and P. Pichini, *to appear*, .