

When Pearson χ^2 and other divisible statistics are not goodness-of-fit tests

Sara Algeri¹ and Estate V. Khmaladze²

¹ *School of Statistics, University of Minnesota, Minneapolis, MN, USA.*

Email: salgeri@umn.edu

² *School of Mathematics and Statistics, Victoria University of Wellington, Wellington, New Zealand.*

Email: estate.khmaladze@vuw.ac.nz

Summary. Thousands of experiments are analyzed and papers are published each year involving the statistical analysis of grouped data. While this area of statistics is often perceived – somewhat naively – as saturated, several misconceptions still affect everyday practice, and new frontiers have so far remained unexplored. Researchers must be aware of the limitations affecting their analyses and what are the new possibilities in their hands.

Motivated by this need, the article introduces a unifying approach to the analysis of grouped data which allows us to study the class of divisible statistics – that includes Pearson’s χ^2 , the likelihood ratio as special cases – with a fresh perspective. The contributions collected in this manuscript span from modeling and estimation to distribution-free goodness-of-fit tests.

Perhaps the most surprising result presented here is that, in a sparse regime, all tests proposed in the literature are dominated by a class of weighted linear statistics.

Keywords: Goodness-of-fit, Divisible statistics, Grouped data, Counting experiments, Poisson statistics, Empirical processes, Distribution-free tests

1. Introduction

The rise of goodness-of-fit in statistics can be traced back to a correspondence between Edgeworth and Pearson at the end of the 19th century [Cfr. [Stigler, 1990](#)] which led to Pearson’s seminal paper on the χ^2 -test [[Pearson, 1900](#)]. The latter can be considered one of the pillars of data analysis and, similarly to other foundational tools such as regression via least squares and the analysis of variance, it is still dominant in everyday practice.

Given K mutually exclusive groups or cells, labeled by x_k , let $\nu(x_k)$ be the frequencies observed in each group. Pearson’s χ^2 aims to test the hypothesis

$$H_0 : E[\nu(x_k)] = m_\theta(x_k) \quad \text{for all } k \in \{k = 1, \dots, K\} \quad (1)$$

where the expected frequencies, $m_\theta(x_k)$, reflect our understanding of the phenomenon under study and may depend on a – possibly unknown – parameter $\theta \in \mathbb{R}^p$. The prescribed test statistic is

$$\sum_{k=1}^K \frac{(\nu(x_k) - m_\theta(x_k))^2}{m_\theta(x_k)},$$

and it is known to be χ^2 distributed under H_0 if $\min_{x_k} \{m_\theta(x_k)\} \rightarrow \infty$.

Following Pearson’s result, many different statistics have been introduced in the literature as substitutes to the χ^2 -test or supplementary to it [Cfr., [Cochran, 1952](#)]. They can be classified into two main kinds: those requiring the data to be in grouped form and those that apply to continuous data.

Theoretical unification in the analysis of continuous data. An important feature of goodness-of-fit statistics for continuous data is that they can all be specified as functionals of the *empirical process*. Letting y_1, \dots, y_n be i.i.d. realizations of a real-valued random variable Y , distributed according to the distribution F , the empirical process $w_n(y)$ is such that

$$w_n(y) = \sqrt{n}[F_n(y) - F(y)],$$

with $F_n(y) = \frac{1}{n} \sum_{i=1}^n \mathbb{1}_{\{y_i \leq y\}}$ being the empirical distribution function. A prominent example of a goodness-of-fit test based on $w_n(y)$ is the Kolmogorov-Smirnov statistic [Kolmogorov, 1933, Smirnov, 1939]

$$\sup_y |w_n(y)|.$$

Also very prominent are the Cramer-von Mises [Smirnov, 1937] and Anderson-Darling [Anderson and Darling, 1954] statistics given by

$$\int w_n^2(y) dF(y) \quad \text{and} \quad \int \frac{w_n^2(y)}{\sqrt{F(y)(1-F(y))}} dF(y),$$

respectively. Several other constructs used in the analysis of continuous data can be expressed as linear functionals of $w_n(y)$ [e.g., Wellner, 1992]. Such unification has led to notable advancements in different areas of statistics and has enabled developments in the theory and practice of goodness-of-fit which continue at present.

An attempt of unification in the analysis of grouped data. Alternatives to Pearson in the analysis of grouped data include the likelihood ratio

$$\sum_{k=1}^K \nu(x_k) \ln \frac{m_\theta(x_k)}{\nu(x_k)},$$

the linear statistic

$$\sum_{k=1}^K [\nu(x_k) - m_\theta(x_k)],$$

and the (cumulative) spectrum

$$\sum_{k=1}^K \mathbb{1}_{\{\nu(x_k) \leq r\}} \quad \text{with } r \in \mathbb{N},$$

which plays a central role in occupancy and diversity problems [e.g., Magurran, 1988, Mirakhmedov, 2007, Barbour and Gnedin, 2009, Khmaladze, 2011]. The commonality of these and many other tests is that they can all be expressed as the sum of a function of the observed and expected frequencies. For this reason, they have been referred to in the literature as “divisible statistics” (or sometimes “separable statistics”) [Cfr. Medvedev, 1970].

Given a function $g(z, t)$, usually differentiable in t , a divisible statistic based on such function has the form

$$\frac{1}{\sqrt{K}} \sum_{k=1}^K g(\nu(x_k), m_\theta(x_k)). \quad (2)$$

The study of the class of divisible statistics as such began in the second half of the 20th century [e.g., Ivchenko and Medvedev, 1979, 1981, Khmaladze, 1984, Mnatsakanov, 1986, 1988]. Of particular interest is their behavior in a sparse regime in which, letting $T = \sum_{k=1}^K m_\theta(x_k)$ be the expected sample size, K and T are assumed to increase simultaneously so that

$$\frac{T}{K} \rightarrow c \in (0, \infty), \quad \text{as } T \rightarrow \infty \text{ and } K \rightarrow \infty, \quad (3)$$

and c can be interpreted as the average number of expected events per cell. One of the most notable results on divisible statistics under (3) is that, if θ is known, no divisible statistic can detect all local departure from the null, somewhat invalidating their adequacy for goodness-of-fit. However, power can be restored by considering partial sums [Cfr. Khmaladze, 1984].

The existing literature on the class of divisible statistics focuses on goodness-of-fit for simple hypotheses. The effect of parameter estimation, on the other hand, has only been investigated on a

case-by-case basis [Cfr., [Cressie and Read, 1989](#)] or for sub-classes such as that of power-divergence statistics [[Cressie and Read, 1984](#), [Müller and Osius, 2003](#)]. No parametric result on the class of divisible statistics as a whole has yet been introduced nor an analog to the empirical process has been proposed for the analysis of grouped data.

Divisible statistics anew: statistical motivation and scientific relevance. The primary motivation of this manuscript is the need to unify the theory of statistical inference for grouped data to an extent comparable to what is enabled by the theory of empirical processes in the continuous regime.

To serve this purpose, in [Section 3](#) we show that it is possible to express different statistics involved in the analysis of grouped data as linear functionals based on the same random measure. This construction not only brings the desired unification but also unveils new phenomena, not visible in the analysis of i.i.d. observations. For example, a somewhat unanticipated result is presented in [Section 6.2](#): in a sparse regime, all tests based on divisible statistics are dominated by a class of weighted linear statistics.

In what follows, we assume the observed frequencies are independent Poisson random variables and [\(3\)](#) holds – that is, asymptotically, the frequencies remain Poisson and do not reach the Gaussian limit. We also assume that, given a compact subset of \mathcal{X} of \mathbb{R}^d with Lebesgue measure $\mu(\mathcal{X})$, the grouping leads to an exhaustive partition of \mathcal{X} into bins.

The scientific motivation behind these assumptions is rooted in the physical sciences. Specifically, in physics and astronomy, the detector mechanism often leads to a discretization of the area under study. As a result, the data consists of event counts observed in the bins – be it channels, pixels, voxels, or other segments – of the discretized space. Moreover, even when the detector resolution is so high that leads to a nearly continuous data stream, the statistical analysis is often conducted by partitioning the data into bins. This choice may be dictated by the need for fast computations and numerical stability when fitting the models under study, to improve the signal-to-noise ratio, and/or to simplify the model assessment stage. In X -ray astronomy, for example, the images and the spectra collected by the Chandra X -ray Observatory [[Swartz et al., 2010](#)] are used to study the properties of celestial objects. In high-energy physics, the energy spectra produced by the Large Hadron Collider are at the core of searches for new physics beyond the Standard Model; of particular interest are thinly populated high-energy bins in which the new phenomena are often expected to appear [e.g., [Belfonte et al., 2021](#)]. In astroparticle physics, the images and the spectra generated by the Fermi Large Area Telescope [[Atwood et al., 2009](#)] enable the study of subatomic particles. In these and many other instances, the data consists of Poisson-distributed photon counts observed over several bins. Hence, the goal is to estimate and validate the hypothesized mean function of a Poisson process defined over a discretized space.

In addition to their scientific relevance, the Poisson and binned data assumptions allow us to convey our findings simply and intuitively. Yet, extensions to the multinomial scheme or situations in which the cells are defined by categorical variables are also possible in the proposed setup.

2. Modeling framework

To formally define the quantities introduced in [Section 1](#), consider a Poisson process $N_r(A)$, $A \subseteq \mathcal{X}$, with mean

$$E[N_r(A)] = \Lambda_r(A, \beta) = T \int_A \lambda(x, \beta) \mu(dx), \quad (4)$$

where $\lambda(x, \beta)$ is the function describing the spatial spread of events over \mathcal{X} which depends on $\beta \in \mathbb{R}^{p-1}$. The parameter T determines the size of the sample – that is, in our context “large samples” means $T \rightarrow \infty$. It is reasonable to assume that $\lambda(x, \beta)$ is independent of T and

$$\int_{\mathcal{X}} \lambda(x, \beta) \mu(dx) = 1 \quad \text{for all } \beta \in \mathbb{R}^{p-1}.$$

Without loss of generality, let \mathcal{X} be a d -dimensional hyperrectangle or a finite union of d -dimensional hyperrectangles. When the region of interest is not rectangular, but merely a bounded set in \mathbb{R}^d , we can cover it with a finite union of rectangles \mathcal{R} and assume $\lambda(x, \beta) = 0$ for all $x \in \mathcal{R} \setminus \mathcal{X}$. Subsequently,

we can construct a grid on \mathcal{X} composed of K disjoint squares, or bins, denoted by $\{\Delta[x_k]\}_{k=1}^K$ and centered at points x_k . Assume the bins to be equally spaced with volume

$$\delta = \frac{\mu(\mathcal{X})}{K}.$$

The observed frequencies correspond to the increments of the process $N_T(A)$ over each bin, i.e.,

$$\nu(x_k) = N_T(\Delta[x_k]), \quad \text{for } k = 1, \dots, K,$$

where the dependence of $\nu(x_k)$ on T is dropped to simplify the notation. Their expectation is

$$m_\theta(x_k) = \Lambda_T(\Delta[x_k], \beta) = T \int_{\Delta[x_k]} \lambda(x, \beta) \mu(dx) \sim c\mu(\mathcal{X})\lambda(x_k, \beta), \quad (5)$$

in which the asymptotic equality follows from (3). The collection of frequencies $\{\nu(x_k)\}_{k=1}^K$, which, from some point of view, one would like to call a Poisson brush, is an unusual object. Indeed, $\nu(x_k)$ counts the number of jump points of N_T in the bin $\Delta[x_k]$; yet, as the bin shrinks, these points do not vanish because T increases at the same time; but where do the limiting jump points live? The answer is, we believe, associated with the notion of fold-up derivatives of shrinking sets and the local Poisson processes (Cfr. Khmaladze [2007], Khmaladze and Weil [2008]). However, we will not digress in this direction here.

In practice, β is often unknown or unspecified. Moreover, from a technical standpoint, it is convenient to treat T as an unknown parameter to be estimated, along with β . Since under (3) T may grow without limit, we estimate it in relative terms by considering $c \sim \frac{T}{K}$. Therefore, $\theta = (c, \beta)^\top \in \Theta \subseteq \mathbb{R}^p$ is the focus of estimation.

3. Divisible statistics as linear functionals based on the same random measure

As described in Section 1, classical goodness-of-fit statistics for testing (1) fall under the class of divisible statistics defined as in (2). This definition, however, does not include other important statistics, especially in the context of estimation.

A broader class of divisible statistics. Let $m_\theta(x)$ denote a step-wise constant function defined as

$$m_\theta(x) = m_\theta(x_k) \quad \text{for all } x \in \Delta[x_k].$$

To broaden the class of divisible statistics, we propose a revision of its traditional definition in which we allow the structure of $g(z, t)$ to vary over \mathcal{X} . This could be done by expressing g as a function of three arguments, i.e., $g(x, z, t)$. However, since the third argument depends on x through the deterministic function $m_\theta(x)$, we write $g_\theta(x, z)$ to denote the ‘‘inhomogenous’’ version of $g(z, t)$. We can then rely on the definition:

$$\frac{1}{\sqrt{K}} \sum_{k=1}^K g_\theta(x_k, \nu(x_k)), \quad \text{with } g_\theta(x_k, \nu(x_k)) = g(x_k, \nu(x_k), m_\theta(x_k)), \quad (6)$$

to include not only goodness-of-fit statistics but also estimating equations, projections, partial sums, and many other forms of statistics that can be employed in the analysis of grouped data. For example, when different weights are assigned to different bins, we may choose

$$g_\theta(x, z) = \omega(x)g(z, t), \quad (7)$$

leading back to (2) when $\omega(x) = 1$. As shown in Section 4, the estimating equations of a particular class of estimators for θ are themselves divisible statistics defined by (7) with $\omega(x)$ depending on θ . Partial sums can be constructed by choosing $\omega(x) = \mathbb{1}_{\{x \in A\}}$. Examples of inhomogenous divisible statistics in which $g_\theta(x, z)$ is structurally different from (7) are those defined by projections that play a central role when θ is estimated.

Divisible statistics as stochastic integrals. The concept of divisible statistics can be extended by expressing them as linear functionals based on the random measure

$$v_{\theta,K}(A, z) = \frac{1}{\sqrt{K}} \sum_{k=1}^K \mathbb{1}_{\{x_k \in A\}} \left[\mathbb{1}_{\{\nu(x_k) \leq z\}} - P(z|m_{\theta}(x_k)) \right], \quad A \subseteq \mathcal{X}, z \in \mathbb{N} \quad (8)$$

in which $P(z|t)$ is the Poisson distribution function with mean t . Notice that, for fixed A and z , $v_{\theta,K}(A, z)$ is itself a divisible statistic generalizing the (centered) cumulative spectrum. It counts the number of bins in A with frequencies $\nu(x_k)$ less than or equal to z while centering the random terms, $\mathbb{1}_{\{\nu(x_k) \leq z\}}$, by their expectation.

To show the connection between $v_{\theta,K}$ and the classical empirical process for i.i.d. observations, rewrite (8) as

$$v_{\theta,K}(A, z) = \sqrt{K} \left[\frac{1}{K} \sum_{k=1}^K \mathbb{1}_{\{x_k \in A\}} \mathbb{1}_{\{\nu(x_k) \leq z\}} - \frac{1}{K} \sum_{k=1}^K \mathbb{1}_{\{x_k \in A\}} P(z|m_{\theta}(x_k)) \right].$$

The first average in k serves a purpose similar to that of the empirical distribution function. The second average represents the expected value of such empirical distribution given the collection of x_k . When rescaling the latter by a factor of $\mu(\mathcal{X})$ we obtain the measure

$$\mu_{\theta,K}(A, z) = \delta \sum_{k=1}^K \mathbb{1}_{\{x_k \in A\}} P(z|m_{\theta}(x_k)),$$

which assigns to each bin in A the Poisson probability $P(z|m_{\theta}(x_k))$. As $K \rightarrow \infty$, $\mu_{\theta,K}(A, z)$ converges weakly to

$$\mu_{\theta}(A, z) = \int_A P(z|m_{\theta}(x)) \mu(dx).$$

Similarly, the marginal $\mu_K(A) = \mu_K(A, \infty) = \delta \sum_{k=1}^K \mathbb{1}_{\{x_k \in A\}}$ converges to $\mu(A)$. For example, letting $E_{\theta}[\cdot]$ be the expectation taken with respect to a Poisson random variable, $\nu(x)$, with mean $m_{\theta}(x)$, as $K \rightarrow \infty$,

$$\iint g_{\theta}(x, z) \mu_{\theta,K}(dx, dz) = \int E_{\theta}[g_{\theta}(x, \nu(x))] \mu_K(dx) \rightarrow \iint g_{\theta}(x, z) \mu_{\theta}(dx, dz) = \int E_{\theta}[g_{\theta}(x, \nu(x))] \mu(dx).$$

Now note that for centered divisible statistics we can write

$$\frac{1}{\sqrt{K}} \sum_{k=1}^K \left[g_{\theta}(x_k, \nu(x_k)) - E_{\theta}[g_{\theta}(x_k, \nu(x_k))] \right] = \iint g_{\theta}(x, z) v_{\theta,K}(dx, dz) = v_{\theta,K}(g_{\theta}), \quad (9)$$

so that the class of divisible statistics becomes the function-parametric process based on $v_{\theta,K}$ in (8). The natural range for the functions g_{θ} indexing such a process is the Hilbert space

$$\mathcal{L}_2(\mu_{\theta,K}) = \{g_{\theta}(x, z) : \langle g_{\theta}, 1 \rangle = 0, \langle g_{\theta}, g_{\theta}' \rangle < \infty\}$$

with inner product $\langle g_{\theta}, g_{\theta}' \rangle = \iint g_{\theta}(x, z) g_{\theta}'(x, z) \mu_{\theta,K}(dx, dz)$. Here, the orthogonality condition $\langle g_{\theta}, 1 \rangle = 0$ indicates that we can restrict our attention to functions g_{θ} that are centered.

The representation in (9) allows us to express seemingly different statistics as the values of the function-parametric process $v_{\theta,K}(g_{\theta})$. For example, such different statistics as Pearson's χ^2 statistic and the spectrum become not so different and subject to the same treatment.

4. Parameter estimation

When testing the validity of a given model $m_{\theta}(x)$ without a prescribed value of θ , one would need to replace the latter with an estimator. Therefore, it is useful to study how estimation affects the structure of divisible statistics.

Let b_θ be a p -dimensional vector function with linearly independent components in $\mathcal{L}_2(\mu_{\theta,K})$ and consider the system of estimating equations

$$v_{\hat{\theta},K}(b_{\hat{\theta}}) = 0. \quad (10)$$

For example, when using Maximum Likelihood Estimation (MLE), b_θ is equal to the score function

$$\psi_\theta(x, z) = \frac{\partial}{\partial \theta} \ln p(z|m_\theta(x)) = \frac{\dot{m}_\theta(x)}{m_\theta(x)}(z - m_\theta(x)),$$

with $p(z|t)$ denoting the Poisson probability mass function with rate t and $\dot{m}_\theta(x) = \frac{\partial}{\partial \theta} m_\theta(x)$. When estimating θ via least squares, the minimization

$$\hat{\theta} = \arg \min_{\theta} \sum_{k=1}^K (\nu(x_k) - m_\theta(x_k))^2,$$

leads to $b_\theta(x, z) = -2\dot{m}_\theta(x)(z - m_\theta(x))$.

Under mild regularity conditions [e.g., [van der Vaart, 2000](#), Chapter 5], there exists a consistent root $\hat{\theta}$ of (10) and for this estimator the Taylor expansion

$$v_{\hat{\theta},K}(g_{\hat{\theta}}) = v_{\theta,K}(g_\theta) + \sqrt{K}(\hat{\theta} - \theta)^\top \frac{1}{\sqrt{K}} \frac{\partial}{\partial \theta} \iint g_\theta(x, z) v_{\theta,K}(dx, dz) + o_P(1)$$

is valid for any $g_\theta \in \mathcal{L}_2(\mu_{\theta,K})$. Also, assume that if the differentiation in θ can be moved under the integral sign and the derivative $\dot{g}_\theta(x, z) = \frac{\partial}{\partial \theta} g_\theta(x, z)$ exists for all $\theta \in \Theta$ then

$$\frac{1}{\sqrt{K}} \frac{\partial}{\partial \theta} \iint g_\theta(x, z) v_{\theta,K}(dx, dz) = \frac{1}{\sqrt{K}} v_{\theta,K}(\dot{g}_\theta) - \frac{1}{\mu(\mathcal{X})} \iint g_\theta(x, z) \frac{\partial}{\partial \theta} \mu_{\theta,K}(dx, dz)$$

in which the first term on the right-hand side is $o_P(1)$ and the second term is proportional to

$$\iint g_\theta(x, z) \frac{\partial}{\partial \theta} P(dz|m_\theta(x)) \mu_K(dx) = \int E_\theta [g_\theta(x, \nu(x)) \psi_\theta(x, \nu(x))] \mu_K(dx) = \langle g_\theta, \psi_\theta \rangle.$$

Therefore, the Taylor expansion above can be rewritten as

$$v_{\hat{\theta},K}(g_{\hat{\theta}}) = v_{\theta,K}(g_\theta) - \langle g_\theta, \psi_\theta^\top \rangle \frac{\sqrt{K}(\hat{\theta} - \theta)}{\mu(\mathcal{X})} + o_P(1). \quad (11)$$

The components of b_θ live in $\mathcal{L}_2(\mu_{\theta,K})$, thus, the asymptotic representation in (11) can also be applied to the left-hand side of the estimating equations in (10). By equating the resulting expansion to zero, we obtain,

$$\frac{\sqrt{K}(\hat{\theta} - \theta)}{\mu(\mathcal{X})} = \langle b_\theta, \psi_\theta^\top \rangle^{-1} v_{\theta,K}(b_\theta) + o_P(1) \quad (12)$$

and substituting the right-hand side of this equality into (11) gives

$$\begin{aligned} v_{\hat{\theta},K}(g_{\hat{\theta}}) &= v_{\theta,K}(\Pi g_\theta) + o_P(1) \\ \text{with } v_{\theta,K}(\Pi g_\theta) &= v_{\theta,K}(g_\theta) - \langle g_\theta, \psi_\theta^\top \rangle \langle b_\theta, \psi_\theta^\top \rangle^{-1} v_{\theta,K}(b_\theta), \end{aligned} \quad (13)$$

in which the notation Πg_θ is used to emphasize that the linear transformation involving g_θ is a projection. The properties of the latter are formalized in [Proposition 1](#).

PROPOSITION 1. *The linear transformation*

$$\Pi g_\theta(x, z) = g_\theta(x, z) - \langle g_\theta, \psi_\theta^\top \rangle \langle b_\theta, \psi_\theta^\top \rangle^{-1} b_\theta(x, z)$$

is a projection of g_θ orthogonal to ψ_θ and parallel to b_θ , i.e.,

$$\langle \Pi g_\theta, \psi_\theta \rangle = 0 \quad \text{and} \quad \Pi b_\theta(x, z) = 0. \quad (14)$$

One can show that Π is a projection operator by verifying that

$$\Pi\Pi g_\theta(x, z) = \Pi g_\theta(x, z).$$

The second equality in (14) tells us that b_θ belongs to the kernel of the operator Π . When $b_\theta \neq \psi_\theta$, however, the kernel of Π may not coincide with the orthogonal complement of its image. Hence, in general, the projection is not orthogonal, but it is, for example, when the parameters are estimated via MLE. For the latter, the leading term in (13) can be expressed as

$$v_{\theta, \mathcal{K}}(\Pi g_\theta) = v_{\theta, \mathcal{K}}(g_\theta) - \sum_{j=1}^p \langle g_\theta, s_j \rangle v_{\theta, \mathcal{K}}(s_j),$$

in which s_j denotes the j -th coordinate of the orthonormalized score function

$$s(x, z) = \langle \psi_\theta, \psi_\theta^\top \rangle^{-1/2} \psi_\theta(x, z)$$

and $\langle \psi_\theta, \psi_\theta^\top \rangle^{-1/2}$ is the inverse square root of the Fisher information matrix.

The projection $v_{\theta, \mathcal{K}}(\Pi g_\theta)$ belongs to the class of divisible statistics defined by (9). This fact has several implications of both practical and theoretical nature. For instance, it allows us to describe the asymptotic behavior of divisible statistics under local alternatives when testing parametric hypotheses (Cfr. Section 6.1). It also offers notable computational advantages when deriving the null distribution of divisible statistics through numerical simulations (Cfr. Section 7.2). Moreover, as demonstrated in Section 7.3, it enables the construction of asymptotically distribution-free goodness-of-fit tests.

Let us now choose b_θ equal to the right-hand side of (7). That is, for a given choice of g , we consider the class of estimators defined by

$$v_{\hat{\theta}, \mathcal{K}}(\omega g) = 0. \quad (15)$$

For instance, when g gives the linear statistic, two members of such a class are the MLE and the least squares estimator with $\omega(x) = \frac{\dot{m}_\theta(x)}{m_\theta(x)}$ and $\omega(x) = -2\dot{m}_\theta(x)$, respectively.

The estimator with the minimum variance among those given by (15) can be constructed by choosing $\omega(x)$ equal to

$$\gamma_\theta(x) = \frac{E_\theta[g(\nu(x), m_\theta(x))\psi_\theta(x, \nu(x))]}{E_\theta[g^2(\nu(x), m_\theta(x))]}.$$
 (16)

This result is formalized in Proposition 2.

PROPOSITION 2. *Let $\hat{\theta}$ be a consistent root of the estimating equations in (15) and let u be a vector in \mathbb{R}^p . For a given choice of ωg , the asymptotic variance of $u^\top \sqrt{K}(\hat{\theta} - \theta)$ is the smallest when $\omega = \gamma_\theta$.*

PROOF. By the definition of γ_θ we have that

$$\langle \omega g, \psi_\theta^\top \rangle = \langle \omega g, g\gamma_\theta^\top \rangle.$$

Therefore, the asymptotic representation of $\hat{\theta}$ is

$$\frac{\sqrt{K}(\hat{\theta} - \theta)}{\mu(\mathcal{X})} \sim \langle \omega g, g\gamma_\theta^\top \rangle^{-1} v_{\theta, \mathcal{K}}(\omega g). \quad (17)$$

Let us now consider the random variables

$$V = u^\top \langle \omega g, g\gamma_\theta^\top \rangle^{-1} v_{\theta, \mathcal{K}}(\omega g) \quad \text{and} \quad W = u^\top \langle \gamma_\theta g, g\gamma_\theta^\top \rangle^{-1} v_{\theta, \mathcal{K}}(\gamma_\theta g).$$

Their covariance is

$$E_\theta[VW] = u^\top \langle \omega g, g\gamma_\theta^\top \rangle^{-1} \langle \omega g, g\gamma_\theta^\top \rangle \langle \gamma_\theta g, g\gamma_\theta^\top \rangle^{-1} u = u^\top \langle \gamma_\theta g, g\gamma_\theta^\top \rangle^{-1} u = E_\theta[W^2].$$

It follows that

$$0 \leq E_\theta[(V - W)^2] = E_\theta[V^2] - E_\theta[W^2];$$

therefore, the variance of W is always smaller than or equal to that of V . ■

When g defines the linear statistic, we have that $\gamma_\theta g = \psi_\theta$. When g is not linear, for each x fixed, $\gamma_\theta g$ gives the best mean-square prediction of ψ_θ given g . For example, when g defines the centred cumulative spectrum, $\gamma_\theta g$ is

$$\frac{r - m_\theta(x)}{1 - p(r|m_\theta(x))} \frac{\dot{m}_\theta(x)}{m_\theta(x)} [\mathbb{1}_{\{z \leq r\}} - P(r|m_\theta(x))],$$

and for the centred Pearson's χ^2 statistic, $\gamma_\theta g$ specifies as

$$\frac{\dot{m}_\theta(x)}{2m_\theta(x) + 1} \left[\frac{(z - m_\theta(x))^2}{m_\theta(x)} - 1 \right].$$

The class of estimators based on $\gamma_\theta g$ offers an alternative to the MLE based on a function g of choice. Nonetheless, as shown in Section 6.1, the MLE is often preferable for goodness-of-fit.

5. Contiguous alternatives and adequacy for goodness-of-fit

A statistical test can be considered adequate for a given testing problem if its power is larger than the significance level. Therefore, to determine its adequacy, we must first identify which alternatives should be examined when studying its power.

In the context of goodness-of-fit, the usual class of alternatives considered for such a study is that of *contiguous alternatives*. The notion of contiguity, introduced by Le Cam [1960], allows us to define sequences of alternatives that converge to the null hypothesis as the sample size increases, thereby describing local deviations from $m_\theta(x)$ due to fainter and fainter signals. Yet, these alternatives remain distinguishable from the null hypothesis.

Consider the distribution function $F_\beta(A) = \int_A \lambda(x, \beta) \mu(dx)$. When θ is known, we say that

$$\tilde{m}_{\theta, T}(x) = m_\theta(x) \left[1 + \frac{h_T(x)}{\sqrt{T}} \right] \quad (18)$$

defines a sequence of *contiguous alternatives* to $m_\theta(x)$ if, for a given sequence of functions, $h_T \in L_2(F_\beta)$, we have

$$\|h_T\|_{F_\beta}^2 \rightarrow \alpha \in (0, \infty), \quad \text{as } T \rightarrow \infty. \quad (19)$$

Moreover, if

$$\|h_T - h\|_{F_\beta}^2 \rightarrow 0, \quad \text{as } T \rightarrow \infty, \quad (20)$$

for some $h \in L_2(F_\beta)$, we say that $\tilde{m}_{\theta, T}(x)$ defines a sequence of *converging contiguous alternatives* to $m_\theta(x)$.

The constant α in (19) can be taken equal to one without loss of essential content. Sequences of $\tilde{m}_{\theta, T}(x)$ with the same limiting function can be considered equivalent and can be labeled by functions h in the unit ball in $L_2(F)$.

The function h defining a given sequence of alternatives can be interpreted as the functional direction from which $\tilde{m}_{\theta, T}(x)$ approaches $m_\theta(x)$. For example, a common problem arising in physics and astronomy is the detection of a faint signal on top of a background. Assume the background density, $\lambda_b(x)$, to be fully specified and let $\lambda_s(x, x_0)$ be the density of the signal, with x_0 being its known location. We may think of $\lambda_s(x, x_0)$ as a Gaussian bump with mean x_0 . The intensity function for the alternative model, inclusive of both background and signal, can be specified as

$$\lambda_b(x) + \eta_T \lambda_s(x, x_0),$$

where $\eta_T \in \mathbb{R}$ is the signal strength relative to T – that is, the total number of expected events when the signal is present is $T + T\eta_T$. Assume η_T to be appropriately small, i.e., $\eta_T = T^{-\frac{1}{2}}$. Hence, the background+signal model above can be rewritten as in (18), with $h_T(x)$ converging to

$$h(x) = \frac{\lambda_s(x, x_0)}{\lambda_b(x)}.$$

The latter defines the direction from which the model inclusive of the signal approaches the background-only model, whereas η_T quantifies the closeness of these two models as a function of the sample size.

When testing parametric hypotheses and θ is allowed to vary, the class of converging contiguous alternatives is much wider compared to the case of simple hypotheses. Such a class includes all sequences of alternatives indexed by functions h in the unit ball in any of the spaces $L_2(F_\beta)$, $\beta \in \mathbb{R}^{p-1}$. Within this class, of particular interest are alternatives satisfying:

$$\int h(x)F_\beta(dx) = 0 \quad \text{and} \quad \int h(x)\frac{\dot{\lambda}(x, \beta)}{\lambda(x, \beta)}F_\beta(dx) = 0, \quad (21)$$

with $\dot{\lambda}(x, \beta) = \frac{\partial}{\partial \beta}\lambda(x, \beta)$. The first condition excludes sequences of alternatives that differ from $\lambda(x, \beta)$ only by a scaling factor and are accounted for through the estimation of c . In other words, we focus on alternatives that only affect the spatial spread of the data over \mathcal{X} , and thus, the function

$$\lambda(x, \beta) \left[1 + \frac{h(x)}{\sqrt{T}} \right]$$

integrates to one. The second condition in (21) excludes alternatives indexed by functions $h \in L_2(F_\beta)$ which are linear combinations of the coordinates of $\dot{\lambda}(x, \beta)/\lambda(x, \beta)$. The latter describes how $\lambda(x, \beta)$ varies for different values of β [Cfr. Amari, 1985, Kass and Vos, 2011]. Such deviations do not describe departures from the parametric family as such and are incorporated into the null through the estimation of β . Alternatives for which both conditions in (21) hold are of the form

$$\hat{h}(x) = h(x) - \frac{\dot{m}_\theta(x)}{m_\theta(x)} \Gamma^{-1} \int \frac{\dot{m}_\theta(y)}{m_\theta(y)} h(y) F_\beta(dy), \quad (22)$$

with $\Gamma = \int \frac{\dot{m}_\theta(x)}{m_\theta(x)} \frac{\dot{m}_\theta(x)}{m_\theta(x)}^\top F_\beta(dx) = \frac{1}{c} \langle \psi_\theta, \psi_\theta^\top \rangle$.

In the literature, the use of contiguous alternatives typically requires the validity of (20). This condition is sufficient for contiguity but not necessary. On the contrary, (19) is both sufficient and necessary [Cfr. Oosterhoff and Van Zwet, 1979]. This implies that there exists a sub-class of alternatives that are contiguous even without (20). In this case, it is not possible to identify a direction from which $\tilde{m}_{\theta_T}(x)$ approaches $m_\theta(x)$, and thus, the sequence of alternatives diverges or oscillates without limit around the null. These alternatives, interesting in themselves, cannot be detected via goodness-of-fit, neither in the continuous nor in the grouped data regime, but they remain discernible from the null through specifically adjusted empirical processes. For this reason, they have been called *chimeric alternatives* [Cfr. Khmaladze, 1998].

Therefore, in the attempt to formally define the adequacy of a statistical test for goodness-of-fit, we rely on the class of converging contiguous alternatives.

DEFINITION 1. (Adequacy for goodness-of-fit) *We say that a statistical test is adequate for goodness-of-fit if its power exceeds the significance level for all converging contiguous alternatives.*

When testing simple hypotheses in a sparse regime, tests based on a single divisible statistic, including Pearson's χ^2 , are known not to satisfy Definition 1 [Cfr. Khmaladze, 1984]. As shown in the sections that follow, the problem persists when testing parametric hypotheses.

In the remainder of the manuscript, we will only consider alternatives within the class of converging contiguous alternatives; thus, we will simply refer to them as “alternatives”.

6. On the behavior of divisible statistics under contiguous alternatives

6.1. Distribution of divisible statistics under converging contiguous alternatives

The stochastic representation of divisible statistics introduced in Section 3 allows us to determine their power on the basis of the asymptotic behavior of $v_{\theta,K}(A, z)$ under the alternatives.

For fixed A and z , $v_{\theta_K}(A, z)$ is the sum of independent random variables, and by the central limit theorem, it is asymptotically Gaussian. Under the null, it has zero mean, and its variance is asymptotically equal to

$$\frac{1}{\mu(\mathcal{X})} \int_A P(z|m_\theta(x)) [1 - P(z|m_\theta(x))] \mu(dx). \quad (23)$$

The first two moments of $v_{\theta_K}(A, z)$ under the alternatives are given in Proposition 3.

PROPOSITION 3. Denote with $\tilde{E}_{\theta,T}[\cdot]$ and $\tilde{V}_{\theta,T}[\cdot]$, respectively, the expectation and the variance taken with respect to the distribution of a Poisson random variable with rate $\tilde{m}_{\theta,T}(x)$. If (3) holds,

$$\tilde{E}_{\theta,T}[v_{\theta_K}(A, z)] \sim \frac{1}{\mu(\mathcal{X})\sqrt{c}} \int_A h(x) \sum_{i=0}^z (i - m_\theta(x)) p(i|m_\theta(x)) \mu(dx) \quad (24)$$

and $\tilde{V}_{\theta,T}[v_{\theta_K}(A, z)]$ is asymptotically equal to (23).

PROOF. Consider

$$\tilde{E}_{\theta,T}[v_{\theta_K}(A, z)] = \frac{\sqrt{K}}{\mu(\mathcal{X})} \int_A [P(z|\tilde{m}_{\theta,T}(x)) - P(z|m_\theta(x))] \mu_K(dx) \quad (25)$$

in which the difference in the square brackets is asymptotically equal to

$$\begin{aligned} \frac{m_\theta(x)h(x)}{\sqrt{T}} \frac{\partial}{\partial \theta} P(z|m_\theta(x_k)) &= \frac{m_\theta(x)h(x)}{\sqrt{T}} \sum_{i=1}^z \frac{\partial \ln p(i|m_\theta(x))}{\partial m_\theta(x)} p(i|m_\theta(x)) \\ &= \frac{h(x)}{\sqrt{T}} \sum_{i=1}^z (i - m_\theta(x)) p(i|m_\theta(x)). \end{aligned}$$

By plugging the last expression into (25) we obtain (24).

The variance under the alternatives is

$$\tilde{V}_{\theta,T}[v_{\theta_K}(A, z)] = \frac{1}{\mu(\mathcal{X})} \int_A P(z|\tilde{m}_{\theta,T}(x)) [1 - P(z|\tilde{m}_{\theta,T}(x))] \mu_K(dx)$$

and converges to (23), as $T \rightarrow \infty$. This also follows from Le Cam's third lemma [Cfr. Hajek and Sidak, 1967, Chapter 6]. ■

Proposition 3 tells us that the asymptotic distribution of $v_{\theta_K}(A, z)$ under the alternatives differs from the null only in the mean. Therefore, to assess the adequacy of divisible statistics for goodness-of-fit, we focus on their shift.

Consider the covariance function

$$C(x; g_\theta) = E_\theta[g_\theta(x, \nu(x))(\nu(x) - m_\theta(x))]$$

taken at each $x \in \mathcal{X}$. When θ is given,

$$\tilde{E}_{\theta,T}[v_{\theta_K}(g_\theta)] = \iint g_\theta(x, z) \tilde{E}_{\theta,T}[v_{\theta_K}(dx, dz)] \sim \frac{\int C(x; g_\theta) h(x) \mu(dx)}{\mu(\mathcal{X})\sqrt{c}} \quad (26)$$

in which the asymptotic equality follows from (24).

Assume that the representation in (13) is valid. When $\hat{\theta}$ solves the estimating equations in (10), the limiting shift is

$$\tilde{E}_{\theta,T}[v_{\theta_K}(\Pi g_\theta)] \sim \frac{\int C(x; g_\theta) h(x) \mu(dx)}{\mu(\mathcal{X})\sqrt{c}} - \langle g_\theta, \psi_\theta^\top \rangle \langle b_\theta, \psi_\theta^\top \rangle^{-1} \frac{\int C(x; b_\theta) h(x) \mu(dx)}{\mu(\mathcal{X})\sqrt{c}}. \quad (27)$$

Moreover, when θ is estimated via MLE,

$$\int C(x; \psi_\theta) h(x) \mu(dx) = c \int \frac{\dot{m}_\theta(x)^\top}{m_\theta(x)} h(x) F_\beta(dx).$$

Thus, the second term in (27) can be expressed as

$$\frac{1}{\mu(\mathcal{X})\sqrt{c}} \int C(x; g_\theta) \left[\frac{\dot{m}_\theta(x)^\top}{m_\theta(x)} \Gamma^{-1} \int \frac{\dot{m}_\theta(x)^\top}{m_\theta(x)} h(x) F_\beta(dx) \right] \mu(dx)$$

and the shift reduces to

$$\frac{\int C(x; g_\theta) \hat{h}(x) \mu(dx)}{\mu(\mathcal{X})\sqrt{c}},$$

with \hat{h} as in (22). Therefore, for any choice of g_θ , only the part of h orthogonal to $\frac{\dot{m}_\theta}{m_\theta}$ with respect to F_β can be detected. Moreover, for sequences of alternatives satisfying the conditions in (21), the limiting shift is the same when testing simple or parametric hypotheses. When the parameters are estimated, however, the variance of the resulting projection is smaller than the variance of $v_{\theta, \kappa}(g_\theta)$. As a result, against this class of alternatives, the MLE leads to higher power compared to the case in which θ is given.

The same does not hold, in general, for other estimators. Specifically, when substituting \hat{h} into (27), the second term on the right-hand side may not be zero if $b_\theta \neq \psi_\theta$. Hence, compared to the simple hypotheses case, the power can still increase, but only when the projection induced by parameter estimation leads to a larger reduction in the variance than in the shift.

6.2. Every divisible statistic is dominated by a weighted linear statistic.

The previous section suggests that regardless of the choice of g_θ , estimating θ via MLE leads to higher power against alternatives for which $h = \hat{h}$. Nevertheless, once a certain function g_θ has been selected, it may be possible to ask if, besides estimation, something else can be done to increase the power of the divisible statistic given by this function. As we shall see, the study of geometric aspects of $v_{\hat{\theta}, \kappa}(g_{\hat{\theta}})$ leads to a somewhat unusual answer to this question.

Consider the asymptotic representation of $v_{\hat{\theta}, \kappa}(g_{\hat{\theta}})$ in (13), with particular focus on the structure of $\langle g_\theta, \psi_\theta^\top \rangle$, i.e.,

$$\langle g_\theta, \psi_\theta^\top \rangle = \int_A \frac{\dot{m}_\theta(x)^\top}{m_\theta(x)} C(x; g_\theta) \mu_\kappa(dx).$$

The inner product above is always zero if $C(x; g_\theta) = 0$ for all $x \in \mathcal{X}$. In other words, if we choose

$$g_\theta^\perp(x, z) = g_\theta(x, z) - \frac{C(x; g_\theta)}{m_\theta(x)} (z - m_\theta(x))$$

then $C(x; g_\theta^\perp) = 0$ and for all $x \in \mathcal{X}$ thus, $\langle g_\theta^\perp, \psi_\theta \rangle = 0$ and $v_{\hat{\theta}, \kappa}(g_{\hat{\theta}}^\perp) = v_{\theta, \kappa}(g_\theta^\perp) + o_P(1)$. Therefore, the asymptotic behavior of divisible statistics based on g_θ^\perp is the same when evaluated at $\hat{\theta}$ or at θ . This is true for any estimator solving (10), not just for the MLE.

At first, this may appear to be a useful property, but there is a substantial pitfall. Specifically, since $C(x; g_\theta^\perp) = 0$ for all $x \in \mathcal{X}$, the shift is zero for all $h \in L_2(F_\beta)$, $\beta \in \mathbb{R}^p$. Therefore, the asymptotic distribution of $v_{\theta, \kappa}(g_\theta^\perp)$ is the same under the null and the alternatives, which makes it not useful. Denote with g_θ^{\parallel} the function

$$g_\theta^{\parallel}(x, z) = \frac{C(x; g_\theta)}{m_\theta(x)} (z - m_\theta(x)) \quad (28)$$

and express $v_{\hat{\theta}, \kappa}(g_{\hat{\theta}})$ as

$$v_{\hat{\theta}, \kappa}(g_{\hat{\theta}}) = v_{\hat{\theta}, \kappa}(g_{\hat{\theta}}^\perp) + v_{\hat{\theta}, \kappa}(g_{\hat{\theta}}^{\parallel}).$$

Since under the alternatives the limiting shift of $v_{\hat{\theta}_{\mathcal{K}}}(g_{\hat{\theta}}^{\perp})$ is zero, tests based on $v_{\hat{\theta}_{\mathcal{K}}}(g_{\hat{\theta}}^{\parallel})$ have the same asymptotic shift as those based on $v_{\hat{\theta}_{\mathcal{K}}}(g_{\hat{\theta}})$, but they are not affected by the additional randomness due to $v_{\hat{\theta}_{\mathcal{K}}}(g_{\hat{\theta}}^{\perp})$. It follows that under (3), tests based on $v_{\hat{\theta}_{\mathcal{K}}}(g_{\hat{\theta}}^{\parallel})$ have higher power than those based on $v_{\hat{\theta}_{\mathcal{K}}}(g_{\hat{\theta}})$ for all alternatives.

6.3. No single divisible statistic is adequate for goodness-of-fit

Despite the advantages given by parameter estimation against alternatives of the form in (22) and the use of $g_{\hat{\theta}}^{\parallel}$ in increasing the power of divisible statistics, neither of these phenomena guarantees their adequacy for goodness-of-fit.

Consider the limiting shifts in (26) and (27). They are linear functionals on $L_2(F_{\beta})$; hence, for any h lying in the subspaces of $L_2(F_{\beta})$ annihilated by these functionals, the limiting power will be equal to the significance level. It follows that no single divisible statistic satisfies Definition 1.

For example, for the (centered) Pearson's χ^2 statistic, $C(x; g_{\theta})$ is identically equal to one. Hence, the shift in (26) is zero whenever

$$\int h(x)\mu(dx) = 0; \quad (29)$$

that is, regardless of the model being tested, Pearson's χ^2 has no power against alternatives approaching the null from a direction h orthogonal to one with respect to the Lebesgue measure.

Furthermore, let F_{β} be the uniform distribution. For any choice of g defining homogeneous divisible statistics, $C(x; g)$ is constant in x . Thus, the same condition in (29) states that the shift in (26) is zero, and no homogeneous divisible statistic can detect departures from uniformity. Unfortunately, the first condition in (21) implies that these are also the only alternatives of interest when testing parametric hypotheses.

In addition to identifying the alternatives towards which the power is lost, the shift also allows us to determine the direction where the power concentrates. Specifically, let us decompose h as

$$h(x) = h^{\perp}(x) + h^{\parallel}(x), \quad (30)$$

in which h^{\perp} denotes the part of h orthogonal to $C(x; g_{\theta})$ with respect to the Lebesgue measure, and h^{\parallel} is the part parallel to it. The asymptotic shift in (26) depends solely on h^{\parallel} , i.e.,

$$\frac{\int C(x; g_{\theta})h(x)\mu(dx)}{\mu(\mathcal{X})\sqrt{c}} = \frac{\int C(x; g_{\theta})h^{\parallel}(x)\mu(dx)}{\mu(\mathcal{X})\sqrt{c}}. \quad (31)$$

This implies that alternatives for which h is proportional to $C(x; g_{\theta})$ have the largest shift.

When estimating the parameters via MLE, the same decomposition holds for \hat{h} . For example, when choosing Pearson's χ^2 statistic, $C(x; g_{\theta}) = 1$ for all $x \in \mathcal{X}$; thus, the power concentrates towards constant deviations from the null. Therefore, when testing parametric hypotheses, only the constant part of \hat{h} , i.e.,

$$\hat{h}^{\parallel}(x) = \int \hat{h}(y)\mu(dy)$$

could be detected. This phenomenon is illustrated through the following three examples.

Example I. Let F_{β} be the distribution of an exponential random variable with rate $\beta = 1.5$ and truncated over the range $\mathcal{X} = [0, 1]$, assuming $c = 5$ and $T = 500$. We are interested in deviations from the direction

$$h_1(x) = a \ln x + b,$$

it corresponds to the normalized score function for the shape parameter of a (truncated) gamma distribution and approaches our exponential model at a rate of \sqrt{T} . Here, we choose $a = 0.87$ and $b = 1.21$ for normalization purposes. Hence, h_1 satisfies the first condition in (21) – that is, it affects the spatial spread of the data but has no impact on the expected sample size described by c . When c and β are estimated via MLE, the power of Pearson's χ^2 obtained by means of 100,000 Monte Carlo replicates is approximately equal to the significance level, here chosen to be 5%.




Fig. 1. Mean functions under the null (black solid lines), alternatives (purple dashed lines), and the corresponding decomposition obtained by replacing h in (18) with \hat{h} (red chained lines) and \hat{h}' (blue dotted lines) for Example I (left panel), Example II (central panel), and Example III (right panel). In all three examples, the alternative defined by \hat{h}' is too weak to be discerned from the null.

To better understand this result, the left panel of Figure 1 shows the graphs of the models for the expected counts under the null (black solid line) and under the alternatives defined by h_1 (purple dashed line), \hat{h}_1 (red chained line), and \hat{h}'_1 (blue dotted line). For this example,

$$\frac{\dot{\lambda}(x, \beta)}{\lambda(x, \beta)} = \varphi(\beta) - x \quad (32)$$

where $\varphi(\beta)$ is a constant depending on β . This tells us that the linear part of h_1 describes deviations within the parametric family. Thus, when the parameters are estimated, only the alternative defined by \hat{h}_1 , which is orthogonal to x with respect to F_β , is detectable by divisible statistics. When using Pearson's χ^2 , however, the power is limited to the constant deviation due to \hat{h}'_1 , which is too small to be distinguished from the null.

Example II. We consider the same truncated exponential null model described in Example I, but we now focus on the direction

$$h_2(x) = ae^{-\frac{1}{2}\left(\frac{x-x_0}{\sigma}\right)^2 + \beta x}.$$

Similarly to the background+signal example in Section 5, h_2 is the direction from which the model inclusive of a Gaussian bump centered at $x_0 = 0.5$ and with standard deviation $\sigma = 0.05$ approaches our exponential background. The coefficient a is chosen equal to 1.66 to ensure that h_2 has a unit norm. In this case, h_2 describes deviations from the null affecting both the spatial spread and the expected sample size. The power of Pearson's statistic estimated via MLE and obtained using 100,000 Monte Carlo replicates is approximately 5% – the same as the significance level.

The central panel of Figure 1 shows that the alternatives defined by h_2 (purple dashed lines) and \widehat{h}_2 (red chained line) differ by a small scaling factor. That is because, in this case, the part of h_2 collinear with (32), and responsible for deviations within the parametric family, is negligible. It follows that the Gaussian bump could be detected by divisible statistics. Pearson χ^2 , however, concentrates its power towards the constant shift due \widehat{h}_2'' (blue dotted line) which is too weak to be discerned from the null.

Example III. Luminosity functions can often be described using power-laws and broken power-laws. Discerning between the two is a common problem arising in X -ray astronomy [e.g., Gladstone et al., 2009, Humphrey and Buote, 2004, Marlowe et al., 2014]. In statistical terms, a power-law is equal to a Pareto type I distribution, while a broken power-law can be constructed by introducing a cutpoint to induce a change in the slope. Let our null model correspond to a power-law with slope $\beta = 2$ and truncated over the interval $\mathcal{X} = [1, 2]$. Set $c = 5$ and $T = 500$. Our interest is in the direction,

$$h_3(x) = a(\ln \xi - \ln x)\mathbb{1}_{\{x \geq \xi\}} + b$$

from which a broken power-law with a cutpoint at $\xi = 1.4$ approaches the simple power-law model. Set $a = 5.58$ and $b = -0.40$ to ensure that h_3 has norm one. The deviations induced by h_3 affect both the sample size and the spatial spread of the data. The power of Pearson's χ^2 with estimated parameters is once again 5%.

The right panel of Figure 1 displays the graphs of the mean models under the null (black solid line) and the alternatives defined by h_3 (purple dashed line), \widehat{h}_3 (red chained line), and \widehat{h}_3'' (blue dotted line). In this case,

$$\frac{\dot{\lambda}(x, \beta)}{\lambda(x, \beta)} = \phi(\beta) - \ln x$$

where $\phi(\beta)$ is a constant depending on β . Comparing the above expression with that of h_3 it is apparent that most of the departure from the null entailed h_3 consists of deviations within the parametric family. Hence, the only part that remains discernable from the null is the faint effect of \widehat{h}_3 . Moreover, since the shift of Pearson's χ^2 is given by the – even weaker – constant part \widehat{h}_3'' , the power is the same as the significance level.

7. Goodness-of-fit via divisible statistics

7.1. On the construction of goodness-of-fit tests based on partial sums

The results of Section 6.3 demonstrate that tests based on a single divisible statistic are inadequate for goodness-of-fit. Nevertheless, power can be restored by relying on a family of divisible statistics.

As shown by Khmaladze [1984] for the case of simple hypotheses, one possible choice is the process of partial sums

$$v_{\theta, K}(g_\theta \mathbb{1}_A) = \int_A \int g_\theta(x, z) v_{\theta, K}(dx, dz) = \frac{1}{\sqrt{K}} \sum_{x_k \in A} g_\theta(x_k, \nu(x_k)), \quad (33)$$

in which $g_\theta \mathbb{1}_A$ denotes functions of the form $g_\theta(x, z)\mathbb{1}_{\{x \in A\}}$. When testing parametric hypotheses, the process of interest is

$$v_{\theta, K}(g_\theta \mathbb{1}_A) = v_{\theta, K}(\Pi g_\theta \mathbb{1}_A) + o_P(1) \quad (34)$$

in which the functions $\Pi g_\theta \mathbb{1}_A$ can be written explicitly as

$$\Pi g_\theta(x, z)\mathbb{1}_{\{x \in A\}} = g_\theta(x, z)\mathbb{1}_{\{x \in A\}} - \langle g_\theta \mathbb{1}_A, \psi_\theta^\top \rangle \langle b_\theta, \psi_\theta^\top \rangle^{-1} b_\theta(x, z).$$

In this paper, we choose the sets A to be members of the *scanning family* [Cfr. Khmaladze, 1993], i.e., an increasing sequence of subsets, $\mathcal{A} = \{A_t\}_{0 \leq t \leq 1}$, of \mathcal{X} such that

- (a) $A_s \subseteq A_t$ if $s \leq t$;
- (b) $\mu(A_0) = 0, \mu(A_1) = \mu(\mathcal{X})$;
- (c) $\mu(A_t)$ is absolutely continuous in t .

It follows that one can label the functions in $\mathcal{L}_2(\mu_{\theta,K})$ indexing the processes in (33)-(34) using the one-dimensional time parameter $t \in [0, 1]$. As shown in Section 7.3, this also simplifies the construction of asymptotically distribution-free goodness-of-fit tests.

The process $v_{\theta,K}(g_\theta \mathbb{1}_A)$ consists of partial sums of independent random variables; hence, it converges weakly to a Brownian motion [Cfr. Pyke, 1983, Kenneth and Pyke, 1986]. The projection $v_{\theta,K}(\Pi g_\theta \mathbb{1}_A)$, on the other hand, converges weakly to a projected Brownian motion. Under the null, the mean function of such processes is identically equal to zero for all $A \in \mathcal{A}$; whereas their shift under the alternatives can be derived by replacing g_θ with $g_\theta \mathbb{1}_A$ in (26) and (27). For instance, (26) becomes

$$\frac{\int_A C(x; g_\theta) h(x) \mu(dx)}{\mu(\mathcal{X}) \sqrt{c}}.$$

Since the integral is taken over all $A \in \mathcal{A}$, it is identically equal to zero only if $C(x; g_\theta) = 0$ for all $x \in \mathcal{X}$ or if the function h satisfies rather exotic conditions. Therefore, in practice, any choice of g_θ that differs from g_θ^\perp (Cfr. Section 6.2) would give a non-zero shift over at least one set $A \in \mathcal{A}$. Consequently, goodness-of-fit tests satisfying Definition 1 can be constructed by considering omnibus functionals of the partial sum processes in (33)-(34) as test statistics.

For example, let us start with a collection of hyperrectangles

$$(-\infty, y_t] = (-\infty, y_{1t}] \times \cdots \times (-\infty, y_{dt}]$$

with $y_t = (y_{1t}, \dots, y_{dt}) \in \mathcal{X}$ such that, if $s \leq t$, $y_{is} \leq y_{it}$, for all $i = 1, \dots, d$. Then choose A_t to be $(-\infty, y_t] \cap \mathcal{X}$. A possible test statistic for parametric goodness-of-fit is

$$\widehat{\text{KS}} = \max_t |v_{\hat{\theta},K}(g_{\hat{\theta}}^{\parallel} \mathbb{1}_t)| \quad (35)$$

in which $\mathbb{1}_t$ denotes the indicator function $\mathbb{1}_{\{x \in (-\infty, y_t] \cap \mathcal{X}\}}$ and $g_{\hat{\theta}}^{\parallel}$ is constructed as in (28). The statistic in (35) is the Kolmogorov-Smirnov statistic for testing parametric hypotheses in the binned data regime. We investigate its power and compare it with that of divisible statistics using a numerical example.

Numerical example. Let $K = 100$, $c = 5$, and $\lambda(x, \beta)$ is chosen to be a normal density with mean β , variance $\sigma^2 = 0.04$, truncated over the interval $\mathcal{X} = [0, 1]$. Assuming the true value of β to be 0.5, the alternative considered is defined by the functional direction

$$h_4(x) = \frac{a}{2\sigma^2} \left[\left(\frac{x - \beta}{\sigma} \right)^2 - 1 \right]$$

and, for $a = 0.07$, it corresponds to the normalized derivative of the log-likelihood of our (truncated) normal model taken with respect to σ^2 – hence, the conditions in (21) are satisfied. The resulting null and alternative models are shown on the left panel of Figure 2.

We assess numerically the power of Pearson's χ^2 and that of the corresponding weighted linear statistic based on

$$g_\theta^{\parallel}(x, z) = \frac{(z - m_\theta(x))}{m_\theta(x)}. \quad (36)$$

Begin with the case in which θ is given. For Pearson's χ^2 , the power estimated using 100,000 Monte Carlo replicates and choosing the significance level to be 5% is approximately 7%; whereas, for $v_{\theta,K}(g_\theta^{\parallel})$ the power is 11%. As expected, if we free ourselves from the extra randomness due to $v_{\theta,K}(g_\theta^\perp)$, the power increases. When θ is estimated via MLE, the power of $v_{\hat{\theta},K}(g_{\hat{\theta}}^{\parallel})$ amounts to 14% – that is, compared to Pearson's statistic with θ known, the use of g_θ^{\parallel} and estimation via MLE together lead to twice as much power. Finally, when relying on the Kolmogorov-Smirnov statistic in (35), the power is 17%.

In this example, the Kolmogorov-Smirnov statistic based on $v_{\hat{\theta},K}(g_{\hat{\theta}}^{\parallel} \mathbb{1}_t)$ exhibits higher power than $v_{\hat{\theta},K}(g_{\hat{\theta}}^{\parallel})$. However, we should not expect this result to be true in general. Specifically, a test based on functionals of the partial sums process should be preferred to a test based on a single divisible statistic – such as $v_{\hat{\theta},K}(g_{\hat{\theta}}^{\parallel})$ – because the latter does not satisfy Definition 1. Nevertheless, it is possible




Fig. 2. Left panel: Null model (black solid line) and contiguous alternative model (red dashed line) considered in our numerical example. Right panel: Comparing the bootstrapped distribution of the Kolmogorov-Smirnov statistics \hat{KS} and \bar{KS} , as defined in (35) and (38), using 100,000 replicates.

that, for some alternatives, a single divisible statistic may exhibit higher power than functionals of the partial sums process. What holds, in general, is that, under (3), statistics based on $g_{\hat{\theta}}^{\prime\prime}$ always exhibit higher asymptotic power than those based on $g_{\hat{\theta}}$. Moreover, parameter estimation via MLE leads to an increase of power in detecting alternatives of the form in (22) compared to the case in which θ is given.

7.2. The projected parametric bootstrap

While the construction of asymptotically distribution-free tests for parametric hypotheses is possible (Cfr. Section 7.3), one can alternatively derive the null distribution of functionals of $v_{\hat{\theta},K}(g_{\hat{\theta}}\mathbf{1}_A)$ via the parametric bootstrap. As discussed below, the computational effort required by the latter can be substantially reduced when relying on the projected process of partial sums.

Let us focus on the situation in which the process of partial sums is indexed by functions $g_{\hat{\theta}}^{\prime\prime}\mathbf{1}_A$, and the parameters are estimated via MLE. Similar considerations can be made for any other choice of g_{θ} and other estimators solving the estimating equations in (10).

Denote with $\hat{\theta}_{\text{obs}}$ the MLE of θ obtained on the set of data collected by the experiment. We are interested in simulating the null distribution of the process

$$v_{\hat{\theta},K}(g_{\hat{\theta}}^{\prime\prime}\mathbf{1}_A) = \frac{1}{\sqrt{K}} \sum_{x_k \in A} \frac{C(x_k; g_{\hat{\theta}})}{m_{\hat{\theta}}(x_k)} (\nu(x_k) - m_{\hat{\theta}}(x_k)) \quad (37)$$

and its functionals. To account for the randomness associated with $\hat{\theta}$, the classical parametric bootstrap entails maximizing the log-likelihood on each bootstrap sample and re-evaluating $v_{\hat{\theta},K}(g_{\hat{\theta}}^{\prime\prime}\mathbf{1}_A)$ at each bootstrap estimate of θ . Conversely, to generate bootstrap replicates of the projected process of partial sums (Cfr. Section 4),

$$v_{\theta,K}(\Pi g_{\theta}^{\prime\prime}\mathbf{1}_A) = v_{\theta,K}(g_{\theta}^{\prime\prime}\mathbf{1}_A) - \sum_{j=1}^p \langle g_{\theta}^{\prime\prime}\mathbf{1}_A, s_j \rangle v_{\theta,K}(s_j),$$

the estimation of θ is replaced by the evaluation of the second term on the right-hand side of each bootstrap sample. Nonetheless, the inner products involved need only to be computed once at $\theta = \hat{\theta}_{\text{obs}}$. Furthermore, since neither $v_{\theta,K}(g_{\theta}^{\prime\prime}\mathbf{1}_A)$ nor the random variables $v_{\theta,K}(s_j)$ depend on $\hat{\theta}$, one needs not to re-evaluate $m_{\theta}(x)$ on each bootstrap estimate of θ , often leading to a substantial computational gain.

The computational benefits of the projected empirical process, particularly in the context of testing multivariate distributions in the i.i.d. setting, have been discussed in Algeri [2022]. To illustrate this aspect in the binned data regime, we rely on the setup of the numerical example introduced in Section

7.3. We compare the bootstrapped null distribution of the Kolmogorov-Smirnov statistics in (35) with that of

$$\overline{\text{KS}} = \max_t |v_{\hat{\theta}_K}(\Pi g_{\theta} // \mathbf{1}_t)|, \quad (38)$$

where t labels the sets $(-\infty, y_t] \cap \mathcal{X}$ (Cfr. Section 7.1) and $g_{\theta} //$ specifies as in (36). The results obtained are shown on the right panel of Figure 2. Not surprisingly, the two bootstrapped distributions are effectively overlapping. The projected process $v_{\hat{\theta}_K}(\Pi g_{\theta} // \mathbf{1}_t)$, however, provides substantial computational gain compared to $v_{\hat{\theta}_K}(g_{\theta} // \mathbf{1}_t)$. Specifically, simulating the null distribution of $\overline{\text{KS}}$ using 100,000 bootstrap replicates required only 5.51 seconds of (system+user) CPU time; approximately 3.3 times faster than simulating that of $\widehat{\text{KS}}$ and which required 18.38 seconds of CPU time.

7.3. Asymptotically distribution-free goodness-of-fit tests via unitary operators

The projected bootstrap can be used to reduce the CPU time needed to simulate the null distribution of goodness-of-fit statistics. At the same time, it is also possible to construct goodness-of-fit tests that are asymptotically distribution-free by adapting the unitary transformation for empirical processes introduced in Khmaladze [2016] to divisible statistics.

Given the prominence of the linear statistics, we consider the functions

$$\ell_t(x, z) = \frac{z - m_{\theta}(x)}{\sqrt{m_{\theta}(x)}} \mathbf{1}_{\{x \in A_t\}}$$

in which the sets A_t are members of the scanning family \mathcal{A} and are chosen so that

$$\frac{\mu(A_t)}{\mu(\mathcal{X})} = t. \quad (39)$$

Under (3) and if θ is known, the null distribution of the process $v_{\hat{\theta}_K}(\ell_t)$ converges weakly to that of a standard Brownian motion. When θ is estimated via MLE, $v_{\hat{\theta}_K}(\ell_t)$ is asymptotically equivalent to $v_{\hat{\theta}_K}(\Pi \ell_t)$ and its limiting null distribution is that of a projected Brownian motion orthogonal to the normalized score function s of the model we wish to test. Its covariance will depend on such s . Thus, it cannot be employed for distribution-free testing.

However, it is possible to construct a linear operator in $\mathcal{L}_2(\mu_{\theta_K})$, denoted by U_p , which maps the process $v_{\hat{\theta}_K}(\Pi \ell_t)$ into a process in t with standard distribution, unconnected with the model we are testing. The operator U_p will depend on the model, but the distribution of the resulting process will be free from it. In its spirit, the situation is not unlike the one encountered in the i.i.d. data regime in which the classical time-transformation $t = F(x)$ depends on the hypothetical F but maps F -empirical processes into the uniform empirical process which is free from F .

The geometric idea behind the choice of the operator U is clear: many different parametric models with p -dimensional estimated parameters will asymptotically lead to the projected Brownian motions with the kernel of the projector, although different for different models of the same dimension p . But then one can choose one such projection to be orthogonal to a fixed collection of p orthonormal functions and map particular projections depending on particular models into this chosen standard projection. This can be done with the help of a unitary operator.

Let $r = \{r_j\}_{j=1}^p$ form a fixed orthonormal system in $\mathcal{L}_2(\mu_{\theta_K})$. Unlike the coordinates of orthonormalized score function $s = \{s_j\}_{j=1}^p$, this new system will stay the same for many different models. Consider the projection of ℓ_t orthogonal to r :

$$\Pi_r \ell_t(x, z) = \ell_t(x, z) - \sum_{j=1}^p \langle \ell_t, r_j \rangle r_j(x, z).$$

Let U_p be the unitary operator that maps r into s and, therefore, functions orthogonal to r into functions orthogonal to s . Then

$$\Pi U_p \Pi_r \ell_t(x, z) = U_p \Pi_r \ell_t(x, z) = U_p \ell_t(x, z) - \sum_{j=1}^p \langle U_p \ell_t, s_j \rangle s_j(x, z),$$




Fig. 3. A realization of the limiting process of $v_{\theta, \kappa}(U_p \Pi_r \ell_t)$ given by a succession of rescaled independent standard Brownian bridges, defined as in (42), when $p = 4$.

and if we consider the process $v_{K, \hat{\theta}}(U \Pi_r \ell_t)$ we find that it is the process that serves our purpose since

$$v_{\hat{\theta}, K}(U_p \Pi_r \ell_t) \sim v_{\theta, K}(\Pi U_p \Pi_r \ell_t) = v_{\theta, K}(U_p \Pi_r \ell_t).$$

The variance of the process on the right-hand side, and therefore its covariance, depends on r and does not involve s :

$$E_{\theta}[v_{\theta, K}^2(U_p \Pi_r \ell_t)] = \frac{\|U_p \Pi_r \ell_t\|^2}{\mu(\mathcal{X})} = \frac{\|\Pi_r \ell_t\|^2}{\mu(\mathcal{X})} = \frac{1}{\mu(\mathcal{X})} \left[\|\ell_t\|^2 - \sum_{j=1}^p \langle \ell_t, r_j \rangle^2 \right]. \quad (40)$$

In this manuscript, we focus on one particular choice of r , which, as we shall see, allows us to derive the asymptotic null distribution of goodness-of-fit statistics in closed form.

Specifically, let $\{A_{t_j}\}_{j=1}^p$ be the sequence of p sets within the scanning family \mathcal{A} and satisfying (39) for $t_j = j/p$. Denote with $\{B_j\}_{j=1}^p$ the collection of disjoint sets

$$B_j = A_{t_j} \setminus A_{t_{j-1}},$$

with $B_1 = \mathcal{X}$ if $p = 1$; for any p , $\{B_j\}_{j=1}^p$ provides an exhaustive partition of \mathcal{X} . Choose

$$r_j(x, z) = \frac{\mathbb{1}_{\{x \in B_j\}}}{\sqrt{\mu_{\kappa}(B_j)}} \frac{z - m_{\theta}(x)}{\sqrt{m_{\theta}(x)}}. \quad (41)$$

For this choice of $\{r_j\}_{j=1}^p$ the asymptotic null distribution of $v_{\theta, \kappa}(U_p \Pi_r \ell_t)$ has a remarkably convenient structure and the expression in (40) become simple. Specifically, we will see that, asymptotically, the process $v_{\theta, \kappa}(U_p \Pi_r \ell_t)$ is a sequence of independent standard Brownian bridges, each living on the corresponding set B_j .

One can split $\|\ell_t\|^2$ in the form of the sum

$$\|\ell_t\|^2 = \sum_{j=1}^p \|\mathbb{1}_{B_j} \ell_t\|^2,$$

and rewrite (40) as

$$\frac{1}{\mu(\mathcal{X})} \sum_{j=1}^p [\|\mathbb{1}_{B_j} \ell_t\|^2 - \langle \ell_t, r_j \rangle^2].$$

It can be shown that if either $B_j \subseteq A_t$ or $B_j \cap A_t = \emptyset$, the corresponding summands will be equal zero, and therefore, for any given t , only the summand for which $t_{j-1} < t < t_j$ will remain. For this summand

$$\frac{\|\mathbb{1}_{B_j} \ell_t\|^2}{\mu(\mathcal{X})} \rightarrow t - t_{j-1} \quad \text{while} \quad \frac{\langle \ell_t, r_j \rangle^2}{\mu(\mathcal{X})} \rightarrow p(t - t_{j-1})^2,$$

and eventually, we obtain

$$E_\theta[v_{\theta,K}^2(U_p \Pi_r \ell_t)] \rightarrow t - t_{i-1} - p(t - t_{i-1})^2, \quad \text{with} \quad t_{i-1} = \frac{[pt]}{p},$$

and $i - 1 = [pt]$ denotes the integer part of pt .

When comparing the last expression with the variance of the standard Brownian bridge on $[0, 1]$, $u(t)$, rescaled to set B_i , i.e.,

$$u_i(t) = \frac{1}{\sqrt{p}} u(p(t - t_{i-1})), \quad t_{i-1} \leq t \leq t_i, \quad (42)$$

we find that they are the same. Therefore, the process $v_{\theta,K}(U_p \Pi_r \ell_t)$ is indeed asymptotically a succession of rescaled independent standard Brownian bridges. For illustration purposes, one realization of such a process when $p = 4$ is provided in Figure 3.

The distribution of many goodness-of-fit statistics from such a process is analytically known. For example, consider

$$\text{KS}^* = \max_t |v_{\theta,K}(U_p \Pi_r \ell_t)|. \quad (43)$$

Since the maximum of a standard Brownian bridge is known to follow the Kolmogorov distribution [Kolmogorov, 1933], here denoted with $\mathcal{K}(\cdot)$, we have that, as $K \rightarrow \infty$,

$$P(\text{KS}^* \leq y | H_0) \rightarrow P\left(\max_i \left\{ \max_t |\sqrt{p} u_i(t)| \right\} \leq \sqrt{p} y\right) = \left[\mathcal{K}(\sqrt{p} y)\right]^p. \quad (44)$$

The partial sums process, however, is observed over discrete times; hence, its maximum is always smaller than or equal to the maximum of the Brownian bridge, leading to a shift in the distribution. A numerical study has revealed that when the number of sets A_t considered is the same as the number of bins, such a shift amounts to $\frac{0.6}{\sqrt{K}}$. As demonstrated through the example below, when adding such a quantity to KS^* , its null distribution coincides with

$$\left[\mathcal{K}\left(\sqrt{p}\left(y + \frac{0.6}{\sqrt{K}}\right)\right)\right]^p.$$

even for $K = O(10)$. A similar result can be obtained for situations in which the partial sums are taken over less than K sets.

It remains to address how to construct the unitary operator U_p . Consider the unitary operator $U_{a,b}$ defined as

$$U_{a,b} = I - \frac{\langle \cdot, a - b \rangle}{1 - \langle a, b \rangle} [a(x, z) - b(x, z)]$$

where $a, b \in \mathcal{L}_2(\mu_{\theta,K})$ and $\|a\| = \|b\| = 1$. $U_{a,b}$ maps a into b , b into a , and leaves functions orthogonal to both a and b unchanged. Proceed by constructing a set of auxiliary functions \tilde{r}_j such that $\tilde{r}_1 = r_1$ and, for $j > 1$, set

$$\tilde{r}_j(x, z) = U_{j-1} r_j(x, z) \quad \text{with} \quad U_j = U_{\tilde{r}_j, s_j} \dots U_{\tilde{r}_1, s_1}. \quad (45)$$

For example,

$$\tilde{r}_2(x, z) = U_{\tilde{r}_1, s_1} r_2(x, z), \quad \tilde{r}_3(x, z) = U_{\tilde{r}_2, s_2} U_{\tilde{r}_1, s_1} r_3(x, z), \quad \text{etc.}$$

Each function \tilde{r}_j is orthogonal to all s_i with $i < j$.

The operator U_p , constructed as in (45), is a product of unitary operators; hence, it is unitary, and it can be shown that $U_p r_j = s_j$ for all $j = 1, \dots, p$. Furthermore, given that Π_r and U_p are linear




Fig. 4. Simulated null distributions of the test statistic \overline{KS} in (38) (left panels) and its transformed counterpart KS^* in (43) shifted by $\frac{0.6}{\sqrt{K}}$ (right panels), for the exponential (orange dashed lines) and the normal (blue chained lines) models for $K = 50, 100, 1000$. For comparison, the limiting distribution in (44) is also plotted in both panels (black solid lines). Each simulation has been conducted using 100,000 replicates.

operators, $v_{\theta,K}(U_p\Pi_r\ell_t)$ is a function-parametric process with values given by weighted linear statistics. Thus, the corresponding $v_{\theta,K}(U_p\Pi_r\ell_t^\perp)$ component (Cfr. Section 6.2) is always zero.

Numerical example. To validate the performance of the approach described in this section in retrieving distribution-freeness, consider the exponential and normal models introduced, respectively, in Example I of Section 6.3 and in the numerical example of Section 7.1. In both instances, assume $\theta = (c, \beta)^\top$ to be unknown. Furthermore, to assess how large K needs to be for our asymptotic result to hold, we consider $K = 50, 100, 1000$. The left panels of Figure 4 show the simulated null distributions of the statistic \overline{KS} in (38) with $g_\theta/\mathbf{1}_t = \ell_t$ and $t \in \{0, \frac{1}{K}, \frac{2}{K}, \dots, 1\}$. Due to the absence of distribution-freeness, regardless of how large K is, these null distributions differ substantially when testing the exponential model (orange dashed line) and the normal model (blue chained line). As expected, they also deviate from the limiting distribution in (44) (black solid lines). Conversely, the right panels of Figure 4 show that, for all the values of K considered, the simulated null distributions of the KS^* in (43), adjusted by a factor of $\frac{0.6}{\sqrt{K}}$, is the same when testing either the exponential or the normal model. Moreover, both distributions coincide with the limit in (44).

8. Summary of the main results

This article proposes a revision of the concept of divisible statistics that includes several forms of statistics routinely used in the analyses of grouped data and enables their representation as linear functionals of the random measure $v_{\theta,K}(A, z)$. By studying the latter, researchers can acquire a comprehensive perspective on the behavior of the class of divisible statistics as a whole.

In the context of goodness-of-fit in a sparse regime, such a study has revealed that no single divisible statistic is adequate for goodness-of-fit; however, tests based on functionals of the process of partial sums are. It has also shown that, for any given divisible statistic, it is possible to determine where the power is lost and where it concentrates. Moreover, there exist alternatives for which parameter estimation leads to higher power compared to test for simple hypotheses. The most unexpected finding (for us) is that a class of weighted linear statistics dominates all divisible statistics.

These facts imply that it is beneficial to rely on omnibus functionals of the process in (37) to conduct goodness-of-fit. Their null distribution can be efficiently derived using the projected parametric bootstrap. Test statistics with an asymptotic null distribution free from the model and known in closed form can also be constructed.

The theoretical framework introduced in this manuscript is illustrated through the lenses of a Poisson-binned data analysis. Nevertheless, it naturally lends itself to a broader study of statistics used in the analysis of non-identically distributed data.

Code availability

The R code needed to replicate the results presented in this manuscript is available in the Supplementary Material.

Acknowledgments

S.A. is grateful to the School of Mathematics and Statistics at the Victoria University of Wellington for providing the resources and fostering a welcoming environment at the beginning of this project. She also thanks Bob Cousins, Vinay Kashyap, Knut Morå, and Lawrence Rudnick for the useful discussions on goodness-of-fit problems arising in physics and astronomy.

Funding

This research is supported by the Office of the Vice President for Research at the University of Minnesota and by the Marsden grant VUW1616.

References

- S. Algeri. K-2 rotated goodness-of-fit for multivariate data. *Physical Review D*, 105:035030, 2022.
- S.-i. Amari. *Differential-Geometrical Methods in Statistics*. Lecture Notes in Statistics. Springer, 1985.
- T. Anderson and D. Darling. A test of goodness of fit. *Journal of the American Statistical Association*, 49(268):765–769, 1954.
- W. B. Atwood et al. The Large Area Telescope on the Fermi Gamma-Ray Space Telescope Mission. *The Astrophysical Journal*, 697(2):1071, 2009.
- A. Barbour and A. Gnedin. Small counts in the infinite occupancy scheme. *Electronic Journal of Probability*, 14:365–384, 2009.
- S. Belfonte et al. Search for resonant and nonresonant new phenomena in high-mass dilepton final states at $\sqrt{s} = 13$ TeV. *Journal of High Energy Physics*, 2021(07):1–62, 2021.
- W. Cochran. The χ^2 test of goodness of fit. *The Annals of Mathematical Statistics*, 23(3):315–345, 1952.
- N. Cressie and T. Read. Multinomial goodness-of-fit tests. *Journal of the Royal Statistical Society: Series B (Methodological)*, 46(3):440–464, 1984.
- N. Cressie and T. Read. Pearson’s X^2 and the loglikelihood ratio statistic G^2 : A comparative review. *International Statistical Review*, 57(1):19–43, 1989.
- J. C. Gladstone, T. Roberts, and C. Done. The ultraluminous state. *Monthly Notices of the Royal Astronomical Society*, 397(4):1836–1851, 2009.
- J. Hajek and Z. Sidak. *Theory of Rank Tests*. First Edition. Academia, Prague & Academic Press, New York, 1967.
- P. Humphrey and D. Buote. A Chandra view of the normal S0 galaxy NGC 1332. I. an unbroken, steep power-law luminosity function for the low-mass X-ray binary population. *The Astrophysical Journal*, 612(2):848, 2004.
- G. Ivchenko and Y. Medvedev. Separable statistics and hypothesis testing. the case of small samples. *Theory of Probability and Its Applications*, 23(4):764–775, 1979.
- G. Ivchenko and Y. Medvedev. Decomposable statistics and hypothesis testing for grouped data. *Theory of Probability and Its Applications*, 25(3):540–551, 1981.
- R. Kass and P. Vos. *Geometrical Foundations of Asymptotic Inference*. John Wiley & Sons, 2011.
- A. Kenneth and R. Pyke. A uniform central limit theorem for set-indexed partial-sum processes with finite variance. *The Annals of Probability*, 14(2):582–597, 1986.
- E. Khmaladze. Martingale limit theorems for divisible statistics. *Theory of Probability and Its Applications*, 28(3):530–548, 1984.
- E. Khmaladze. Goodness of fit problem and scanning innovation martingales. *The Annals of Statistics*, 21(2):798–829, 1993.
- E. Khmaladze. Goodness of fit tests for “chimeric” alternatives. *Statistica Neerlandica*, 52(1):90–111, 1998.
- E. Khmaladze. Differentiation of sets in measure. *Journal of Mathematical Analysis and Applications*, 334(2):1055–1072, 2007.

- E. Khmaladze. Convergence properties in certain occupancy problems including the Karlin-Rouault law. *Journal of Applied Probability*, 48(4):1095–1113, 2011.
- E. Khmaladze. Unitary transformations, empirical processes and distribution free testing. *Bernoulli*, 22(1):563–588, 2016.
- E. Khmaladze and W. Weil. Local empirical processes near boundaries of convex bodies. *Annals of the Institute of Statistical Mathematics*, 60(4):813–842, 2008.
- A. Kolmogorov. Sulla determinazione empirica di una legge di distribuzione. *Giornale dell’Istituto Italiano degli Attuari*, 4:83–91, 1933.
- L. Le Cam. Locally asymptotically normal families of distributions. *University of California Publications in Statistics*, 3:37–98, 1960.
- A. Magurran. *Ecological Diversity and its Measurement*. Princeton University Press, 1988.
- H. Marlowe, P. Kaaret, C. Lang, H. Feng, F. Grise, N. Miller, D. Cseh, S. Corbel, and R. F. Mushotzky. Spectral state transitions of the ultraluminous X-ray source IC 342 X-1. *Monthly Notices of the Royal Astronomical Society*, 444(1):642–650, 2014.
- Y. Medvedev. Some theorems on the asymptotic distribution of the χ^2 statistic. *Doklady Akademii Nauk*, 192(5):987–989, 1970.
- S. Mirakhmedov. Asymptotic normality associated with generalized occupancy problems. *Statistics and Probability Letters*, 77(15):1549–1558, 2007.
- R. Mnatsakanov. A functional limit theorem for additively separable statistics in the case of very rare events. *Theory of Probability and Its Applications*, 30(3):622–626, 1986.
- R. Mnatsakanov. On the convergence of separable statistics to a Wiener process. *Theory of Probability and Its Applications*, 32(1):152–157, 1988.
- U. Müller and G. Osius. Asymptotic normality of goodness-of-fit statistics for sparse Poisson data. *Statistics: A Journal of Theoretical and Applied Statistics*, 37(2):119–143, 2003.
- J. Oosterhoff and W. Van Zwet. A note on contiguity and Hellinger distance. *Reidel, Dordrecht*, 157:166, 1979.
- K. Pearson. On the criterion that a given system of deviations from the probable in the case of a correlated system of variables is such that it can be reasonably supposed to have arisen from random sampling. *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science*, 50(302):157–175, 1900.
- R. Pyke. A uniform central limit theorem for partial-sum processes indexed by sets. *London Mathematical Society Lecture Notes Series*, 79:219–240, 1983.
- N. Smirnov. On the distribution of the von Mises ω^2 -criterion. *Matematicheskii Sbornik*, 5:973–993, 1937.
- N. Smirnov. On the estimation of the discrepancy between empirical curves of distribution for two independent samples. *Moscow University Mathematics Bulletin*, 2(2):3–14, 1939.
- S. Stigler. *The History of Statistics*. Harvard University Press, 1990.
- D. Swartz, S. Wolk, and A. Fruscione. Chandra’s first decade of discovery. *Proceedings of the National Academy of Sciences*, 107(16):7127–7134, 2010.
- A. van der Vaart. *Asymptotic Statistics*, volume 3. Cambridge University Press, 2000.
- J. Wellner. Empirical processes in action: a review. *International Statistical Review*, 60(3):247–269, 1992.