

TOPOLOGICAL ISOTOPY AND FINITE TYPE INVARIANTS

SERGEY A. MELIKHOV

ABSTRACT. In 1974, D. Rolfsen asked: If two PL links in S^3 are isotopic (=homotopic through embeddings), then are they PL isotopic? We prove that they are PL isotopic to another pair of links which are indistinguishable from each other by finite type invariants. Thus if finite type invariants separate PL links in S^3 , then Rolfsen’s problem has an affirmative solution. In fact, we show that finite type invariants separate PL links in S^3 if and only if certain 6 rather diverse conjectures hold simultaneously.

We also show that if v is a finite type invariant (or more generally a colored finite type invariant) of PL links, and v is invariant under PL isotopy, then v assumes the same value on all sufficiently close C^0 -approximations of any given topological link; moreover, the extension of v by continuity to topological links is an invariant of isotopy. Some specific invariants of this kind are discussed.

1. INTRODUCTION

1.1. **PL isotopy versus topological isotopy.** Fifty years ago D. Rolfsen [75] posed the following problem. (It can be argued that this question is also implicit in J. Milnor’s 1957 paper [67; Remark 2 and Theorem 10].)

Rolfsen’s Problem. “If L_0 and L_1 are PL links¹ connected by a topological isotopy, are they PL isotopic?”

It has been long known that some PL knots are topologically slice but not PL slice (see [39; comments on problems 1.36–1.38]) and some PL links are connected by a topological I-equivalence (=“non-locally-flat concordance”), but not by a PL I-equivalence [77; Example 7], [25] (see also [16; §12], [37; Theorem 1.9]). In fact, there exists a PL link which is topologically slice, but does not bound PL disks disjointly embedded in the 4-ball (see [21], [39; comments on problems 1.38–1.39] and [77; Theorem 3]).

More recently, it was shown by the author that not every topological link in S^3 is isotopic to a PL link [57]. In the present paper we prove

Theorem A. *If L_0 and L_1 are PL links in S^3 connected by a topological isotopy, then they are PL isotopic to links L'_0 and L'_1 which are not separated from each other by finite type invariants.*

The definition of finite type invariants is recalled in §4.

¹It is not clear whether these links are meant to be in S^3 (as is implicitly assumed in the assertion “all PL knots are isotopic to one another” mentioned in the previous problem in the same list of problems) or in an arbitrary 3-manifold. For the sake of definiteness we will assume the former interpretation.

Thus Rolfsen’s Problem is solved affirmatively provided that the following conjecture holds. (By “links” we will mean “PL links” in the present paper.)

Conjecture (L). *Finite type invariants separate links in S^3 .*

This is a natural extension of a well-known conjecture of V. Vassiliev:

Conjecture (K) (Vassiliev [84], [85]). *Finite type invariants separate knots in S^3 .*

It should be noted that finite type invariants fail to separate knots in the Whitehead manifold, and more generally in any contractible open 3-manifold which is embeddable in \mathbb{R}^3 but not homeomorphic to \mathbb{R}^3 [18], [19].

In the case of knots and links in S^3 , some weight to Conjectures (K) and (L) is added by the close relation between finite type invariants and configuration space integrals (see [41]). Some numeric evidence exists (see e.g. [44]) for the AJ Conjecture, which implies [17] that the colored Jones polynomial, and hence finite type invariants [64], detect the unknot. Some numeric evidence exists (see e.g. [92], [89]) for the Volume Conjecture, which also implies (see [80]) that the colored Jones polynomial detects the unknot.

On the other hand, it remains unknown, according to [13; Preface and §3.2.4], whether finite type invariants of knots detect the orientation of non-invertible knots (i.e. those knots whose equivalence class depends on their orientation), and whether the signature of a knot is dominated by finite type invariants (see, however, [24] for some numeric evidence for a conjectural domination). Closely related to Conjectures (K) and (L) is the following problem: Can one extract at least one finite type invariant from the celebrated categorified invariants, such as Khovanov homology (which is known to detect the unknot) and the link Floer homology of Ozsváth–Szabó, other than the ones already extractable from what they categorify (i.e. the Jones polynomial and the Conway potential function, respectively, in the two said cases)? While there is some work in this direction [6], [38], [91], it appears that the extraction problem turned out to be unexpectedly difficult.

1.2. Extension of invariants. Our approach to topological isotopy is based on the notion of *n-quasi-isotopy*, which was introduced by the author (see [60], [61]) but can be traced back to a number of older constructions, from the Penrose–Whitehead–Zeeman–Irwin trick to Casson handles to the Homma–Bryant proof of the Chernavsky–Miller codimension three PL approximation theorem (see [60; Remarks (i)–(iii) in §1.5]). We recall the definition in §3.

Theorem 1.1. [60; Theorem 1.3] *Let $h_t: \Theta \rightarrow M$ be a topological isotopy of a closed 1-manifold in a 3-manifold. Then for each n there exists an $\epsilon > 0$ such that every PL homotopy $f_t: \Theta \rightarrow M$ which is ϵ -close to h_t and has only finitely many double points occurring at distinct time instances is an n -quasi-isotopy.*

Since the proof is only sketched in [60], we include a more detailed proof in §3.

Our next result involves *colored finite type invariants* of links, introduced by Kirk and Livingston [40] (they did not use the word “colored”). An invariant v of links is said to

be of *colored type n* if its standard extension to singular links (see §4) vanishes on those singular links with $n + 1$ double points whose all double points are self-intersections of components (rather than intersections of distinct components). Colored type n invariants agree on any two links related by self C_n -equivalence (see Remark 5.7). One reason for our interest in colored finite type invariants is the following

Kirk–Livingston Conjecture. [40; p. 1333] *There exists an infinite family of linearly independent \mathbb{Z} -valued colored type 2 invariants of two-component links with $\text{lk} = 0$.*

The following is the core result of the present paper.

Theorem B. *Let v be a type n invariant, or more generally a colored type n invariant of links in an oriented 3-manifold. If v is invariant under PL isotopy, then it is invariant under n -quasi-isotopy.*

Theorems 1.1 and B imply

Corollary 1.2. *Let v be a finite type invariant, or more generally a colored finite type invariant of links in an oriented 3-manifold. Suppose that v is invariant under PL isotopy.*

Then v assumes the same value on all sufficiently close C^0 -approximations of any given topological link. Moreover,² the extension of v by continuity to topological links is an invariant of isotopy.

Example 1.3. Let L be a link in S^3 with components K_1, \dots, K_m . The hypothesis of Corollary 1.2 is satisfied for each coefficient of each of the following 3 formal power series:

- $\bar{\nabla}_L(z) := \frac{\nabla_L(z)}{\nabla_{K_1}(z) \cdots \nabla_{K_m}(z)}$, where ∇_L is the Conway polynomial (see §10 for the details).
- $\bar{\nabla}_L(z_1, \dots, z_m) := \frac{\nabla_L(z_1, \dots, z_m)}{\nabla_{K_1}(z_1) \cdots \nabla_{K_m}(z_m)}$, where ∇_L is the multi-variable Conway polynomial of [58], which contains the same information as the Conway potential function (see [58] for the details). The coefficients of ∇_L and $\bar{\nabla}_L$ are not finite type invariants, but they are colored finite type invariants.
- $\bar{Z}(L) := \frac{Z(L)}{Z(K_1) \cdots Z(K_m)}$, where Z is the Kontsevich integral (see Remark 9.3) and the meaning of the fraction is explained in the proof of Theorem 9.4.

Corollary 1.2 implies that $\bar{\nabla}_L$, $\bar{\nabla}_L$ and \bar{Z} extend by continuity to invariants of isotopy of topological links. In the case of $\bar{\nabla}_L$ and $\bar{\nabla}_L$ more can be said: since ∇_L and ∇_L are genuine polynomials, $\bar{\nabla}_L$ and $\bar{\nabla}_L$ are rational power series for every PL link L . Their extensions to topological links need not be rational (because exactly how close is “sufficiently close” C^0 -approximation is determined individually for each coefficient),

²This “Moreover” can be replaced by “Consequently”, that is, the second assertion of Corollary 1.2 can in fact be deduced directly from the first one (using that $[0, 1]$ is compact).

but whenever one of them is not rational for some topological link \mathcal{L} , we immediately know that \mathcal{L} is not isotopic to any PL link.

The latter observation is applied in a subsequent paper by the author [59] to obtain some progress on another 50-year old problem of Rolfsen [75]: *Is every topological knot isotopic to a PL knot? In particular, is the Bing sling isotopic to a PL knot?* For example, we show in [59], using $\bar{\nabla}_L(z)$ (extended to topological links by means of Corollary 1.2), that the Bing sling is not isotopic to any PL knot through topological knots that are intersection of solid tori. Also we show in [59], using $\bar{\nabla}_L(z_1, z_2)$ (extended to topological links by means of Corollary 1.2), that the Bing sling is not isotopic to any PL knot by an isotopy which extends to an isotopy, or just a link homotopy of two-component links with linking number 1.

Although the results of [59] go much further than this, Rolfsen’s Bing sling problem remains open. In fact the results of [59] seem to suggest that the invariant $\bar{\nabla}_L(z_1, z_2)$ is simply not powerful enough to suffice for its solution.

Remark 1.4. It was noted already by Traldi [83; Theorem 1 and §5] and Rolfsen [78] that for any link polynomial f_L which is multiplicative under addition of a local knot, the rational function $f_L/(f_{K_1} \cdots f_{K_m})$ is invariant under PL isotopy. Thus similarly to Example 1.3 one can deal with appropriate power series expansions of the two-variable HOMFLY and Kauffman polynomials whose coefficients are finite type invariants (see [7; Theorem 3], [48], [36]). But it seems that the currently most promising source of invariants which one could try to feed into Corollary 1.2 is the Akutsu–Deguchi–Ohtsuki polynomials, which generalize the multivariable Alexander polynomial and in the case of knots contain the same information as the colored Jones polynomial [87; p. 24], [88].

Remark 1.5. Another interesting question which one could try to solve by applying Corollary 1.2 is the Isotopic Realization Problem in S^3 [60; Problem 1.1] (see also [53; Questions I and III]): *Is it true that for every continuous map $f: \Theta \rightarrow S^3$, where Θ is a compact 1-manifold, there exists a homotopy $h_t: \Theta \rightarrow S^3$ such that $h_1 = f$ and h_t is injective for $t < 1$?* Potential counterexamples include the infinite connected sum of Whitehead string links [60; Figure 1] and a pair of “linked” wild arcs [53; Figure 1].³ As pointed out in [60], to solve the problem negatively it suffices to find an invariant of 1-quasi-isotopy, or of k -quasi-isotopy for some fixed k , which “detects accumulation of complexity” in the sense explained in [60; Problem 1.5]. In terms of invariants with values in an abelian group, “detecting accumulation of complexity” translates roughly into an infinite series of linearly independent invariants (for instance, the infinite family of coefficients of Conway polynomials of knots yields a lower bound on knot genus), so in view of Theorem B, the desired invariant almost certainly exists if the Kirk–Livingston Conjecture is true. But there might be other approaches as well, as the gap between k -quasi-isotopy and colored type k invariants, well-defined up to PL isotopy, appears to be very wide (see Examples 4.12 and 4.13).

³Beware that the argument in [53; Example 1.3] is erroneous, as explained in [60; end of §1.2].

Example 1.6. Let us discuss in more detail the formal power series $\bar{\nabla}_L$ of Example 1.3. It is well-known (see §10) that for an m -component PL link L its Conway polynomial is of the form

$$\nabla_L(z) = z^{m-1}(c_0 + c_1z^2 + c_2z^4 + \cdots + c_nz^{2n}).$$

Hence its $\bar{\nabla}_L$ is of the form

$$\bar{\nabla}_L(z) = z^{m-1}(\alpha_0 + \alpha_1z^2 + \alpha_2z^4 + \dots).$$

It is not hard to see that $\alpha_k(L)$ is of type $m - 1 + 2k$ and of colored type $2k$. With more work one can show that $\alpha_k(L)$ is of colored type $2k - 1$ for $k > 0$ (Proposition 10.7) but $\alpha_2(L)$ is not of colored type 2 for two-component links of linking number 0 (Proposition 10.13). On the other hand, we show that $\alpha_k(L)$ is invariant under k -quasi-isotopy (Theorem 10.14). Thus the converse to Theorem B is not true.

If L is a 2-component link, then $\alpha_0(L) = c_0(L)$ is its linking number and $\alpha_1(L) = c_1(L) - c_0(L)(c_1(K_1) + c_1(K_2))$ is its generalized Sato–Levine invariant, which generates the group of colored type 1 invariants of two-component links with a fixed linking number modulo type 0 invariants [40]. The generalized Sato–Levine invariant emerged independently in the work of Traldi [83; §10], Polyak–Viro (see [4]), Kirk–Livingston [40] (see also [49]), Akhmetiev (see [3]) and Nakanishi–Ohyama [70]. It is proved in [70] (see also [56] for an alternative proof) that α_0 and α_1 constitute a complete set of invariants of self C_2 -equivalence; and it is shown in [56; Corollary 5.2] that self C_2 -equivalence is the same thing as $\frac{1}{2}$ -quasi-isotopy.

Let us note that the invariance $\alpha_1(L)$ under 1-quasi-isotopy along with Theorem 1.1 already suffice to prove that there exists a PL knot in the solid torus $S^1 \times D^2$ which is homotopic, but not topologically isotopic to the core circle [60; proof of Theorem 2.2]. By contrast, every PL knot in the solid torus which is homotopic to the core circle is topologically I -equivalent to the core circle, i.e. they cobound an embedded annulus in $S^1 \times D^2 \times I$ [25] (see also [16; §12], [37; Theorem 1.9]); this is also true of topological knots in the solid torus [26] (see also [59; Appendix A]).

When L is a 3-component link, $\alpha_1(L)$ is also of some interest. Upon adding to $\alpha_1(L)$ a correction term, which is a function of $\alpha_0(\Lambda)$ and $\alpha_1(\Lambda)$ for 2-component sublinks Λ of L , we obtain an invariant $\gamma(L)$ which has a rather neat crossing change formula for self-intersections of a component (Proposition 10.10). This crossing change formula originally appeared as a side remark in an early preprint version of the present paper [54], but it has since found some applications in magnetohydrodynamics [1], [2].

1.3. Reduction of conjectures. In order to state some further consequences of Theorem B we introduce the following

Conjecture (L/K). *PL isotopy classes of links in S^3 are separated by those finite type invariants that are well-defined up to PL isotopy.*

Since two links are PL isotopic if and only if they are equivalent under the equivalence relation generated by ambient isotopy and insertion of local knots (see [76; Theorem

4.2]), this can be seen as the “links modulo knots” version of the Vassiliev Conjecture (Conjecture (K)).

Is (L) equivalent to the conjunction of (K) and (L/K)? This is not clear. But we prove

Theorem C. *(L) is equivalent to the conjunction of (K) and (L/K):*

(a) *if only rational finite type invariants are considered;*

(b) *if string links are considered instead of links in S^3 .*


For a more accurate formulation see Theorems 9.2 and 9.4. Part (a) is an easy consequence of the (not so easy) theory of the Kontsevich integral; part (b) is proved by using the clasper theory of Gusarov and Habiro.

Now we resume the discussion of consequences of Theorem B.

Corollary 1.7. *If W is any contractible open 3-manifold other than \mathbb{R}^3 , then the analogue of Conjecture (L/K) for links in W fails.*

Proof. W is known to contain a knot K which is not PL isotopic to the unknot U , but n -quasi-isotopic to U for all finite n (and in fact even for $n = \omega$) [61; Proposition 2.3]. Then by Theorem B, K is not separated from U by finite type invariants, well-defined up to PL isotopy. \square

Corollary 1.8. *Conjecture (L/K) is equivalent to the conjunction of the 4 conjectures asserting that the following 4 implications can be reversed for any links L, L' in S^3 :*



Proof. The 4 implications do hold: the first one and the last one are trivial, the second one follows from Theorem 1.1 and the third one from Theorem B. Conjecture (L/K) is the composite of the converses of the 4 implications. Hence it is implied by these 4 converses, and at the same time implies that all 5 assertions in the frame are equivalent. \square

Corollary 1.9. *If Conjecture (L/K) holds, then topologically isotopic links in S^3 are PL isotopic.*

While Corollary 1.9 and Theorem C do not quite suffice to get Theorem A, a minor modification of their proofs does. On this way we also get a modification of Corollary 1.8, which we now prepare to state.

If two string links are not separated by type n invariants, then it is easy to see that their closures are also not separated by type n invariants (cf. Lemma 4.1). The converse


is false, and for a good reason: a nontrivial string link may have trivial closure.⁴ But if we fix the link and allow the string link to vary, the converse is still false. For instance, the Borromean rings are not separated from the unlink by type 2 invariants [32; Proposition 7.4(2)], but whenever we represent both as closures of string links, these string links are always separated by type 2 invariants, since it is known that string links that are not separated by type 2 invariants are C_3 -equivalent (see [52]), but the Borromean rings are not C_3 -equivalent to the unlink [32; Proposition 7.4(1)].

Moreover, given a sequence L_1, \dots, L_k of links such that L_1 is the Borromean rings and L_k is the unlink, and a representation for each pair L_i, L_{i+1} as closures of string links $\Lambda_{2i-1}, \Lambda_{2i}$, then at least one pair $\Lambda_{2i-1}, \Lambda_{2i}$ will be separated by type 2 invariants, as the same argument with C_3 -equivalence shows. In this event we say that the Borromean rings and the unlink “are separated by type 2 invariants of string links”. More formally, given a class C of invariants of string links, and given links L and L' in S^3 , we say that L and L' are *not separated by invariants of string links of class C* if L and L' are equivalent with respect to the equivalence relation generated by ambient isotopy and the relation “to be closures of string links that are not separated by invariants of class C ”. If two links in S^3 are not separated by type n invariants of string links, then by the previous arguments (cf. Lemma 4.1) they are not separated by type n invariants (of links in S^3).

Conjecture (HM). *If two links in S^3 are not separated by finite type invariants, then for each n they are not separated by type n invariants of string links.*

The question of validity of Conjecture (HM) is presumably included in a problem of N. Habegger and J.-B. Meilhan [30; Problem 5.4], which however can be made precise in multiple ways as it is stated in informal language.

Theorem D. *(L) is equivalent to the conjunction of (K), (HM) and the 4 conjectures asserting that the following 4 implications can be reversed for any links L, L' in S^3 :*



⁴For instance, the connected sum $W \#_{\rho} W$ of the Whitehead string link and its reflection is a non-trivial string link, as detected by a type 3 invariant (see [60; proof of Theorem 2.2] or [56; Example 3.3]). On the other hand, the closure of $W \#_{\rho} W$ (which is a connected sum of the Whitehead link with its reflection along bands, orthogonal to the mirror) is easily seen to be the unlink.

1.4. **Organization of material.** The paper is organized as follows.

- A detailed proof of Theorem 1.1 is included in §3.
- Theorem B is proved in §4.
- The main result of §5 says roughly that finite type invariants are dominated by those finite type invariants that are additive under insertion of local knots (Theorem 5.2). The proof is based on the clasper calculus.
- Theorem A is proved in §6, based on the results of §§3–5.
- Theorem D is proved in §7, based on the results of §§3–6.
- The main result of §8 says roughly that every string link admits a unique decomposition into a string link which has no local knots and a string link which has nothing but local knots (Theorem 8.3(b)).
- §8 also includes an example of a string link which has no local knots, while its closure has a local knot (Example 8.1).
- Theorem C is proved in §9, based on the results of §5 and §8.
- §10 is devoted to examples, illustrating colored finite type invariants and n -quasi-isotopy in a more practical light in the context of the Conway polynomial.

As a byproduct of the proofs of the main results, the paper also contains alternative proofs of some known results:

- §8 contains an alternative proof of a classic 1958 theorem of Y. Hashizume, which says that every non-split link admits a unique decomposition into prime links (Theorem 8.9). Our proof is considerably shorter than the original proof and avoids most of its case exhaustion.
- §5 includes a simple visual proof that type n invariants are invariant under C_n -equivalence (Proposition 5.6). While the proofs of this fact in the literature are not very difficult, the new proof appears to be still easier.
- §5 also contains an alternative proof of a 2009 theorem by G. Massuyeau, which provides a partial converse to the previous item (Theorem 5.8). Our argument proceeds in the original language of string links (as opposed to Massuyeau’s language of homology cylinders) and is a correction of K. Habiro’s attempted proof of a slightly weaker result.
- §4 contains a particularly simple (but asymmetric) form of the Leibniz rule for finite type invariants, which I did not see in the literature.

2. BASIC NOTIONS

By a *link* we mean a PL embedding of a closed 1-manifold in 3-manifold. Thus every link is of the form $mS^1 \rightarrow M$, where $mS^1 = \{1, \dots, m\} \times S^1$. More generally, a *tangle* is a proper PL embedding $L: \Theta \rightarrow M$ of compact 1-manifold in a 3-manifold, where *proper* means that $L^{-1}(\partial M) = \partial\Theta$. Two tangles are called *equivalent* if they are ambient isotopic keeping ∂M fixed. An embedding $\Xi: \partial\Theta \rightarrow \partial M$ is called a *boundary pattern*, and a tangle or more generally a map $L: \Theta \rightarrow M$ is said to be *of boundary pattern* Ξ if $L|_{\partial\Theta} = \Xi$. A *string link* is a tangle $mI \rightarrow I^3$, where $I = [0, 1]$ and $mI = \{1, \dots, m\} \times I$,

of the *string link boundary pattern* $\Xi_m: \partial(mI) \rightarrow \partial I^3$, given by $\Xi_m(k, i) = (\frac{k}{m+1}, \frac{1}{2}, i)$. The *string unlink* $U: mI \rightarrow I^3$ is given by $U(k, t) = (\frac{k}{m+1}, \frac{1}{2}, t)$. The *closure* of a string link $L: mI \rightarrow I^3$ is the link $mS^1 \cong mI \cup_{\partial} mI \xrightarrow{L \cup U} I^3 \cup_{\partial} I^3 \cong S^3$. The *unlink* is the closure of the string unlink. A *knot* is a 1-component link, and the *unknot* is the 1-component unlink.

3. n -QUASI-ISOTOPY AND STRONG n -QUASI-ISOTOPY

We recall the definitions of n -quasi-isotopy and strong n -quasi-isotopy [60], [61]. Let $f: \Theta \rightarrow M$ be a PL map of a 1-manifold in a 3-manifold with precisely one double point $x = f(p) = f(q)$. It is called a *strong n -quasi-embedding* if in addition to the singleton $B_0 := \{x\}$ there exist PL 3-balls $B_1, \dots, B_n \subset M$ and arcs $J_0, \dots, J_n \subset \Theta$ such that $f^{-1}(B_i) \subset J_i$ for each $i \leq n$ and $B_i \cup f(J_i) \subset B_{i+1}$ for each $i < n$. Let us note that since J_n is an arc, B_n can intersect the image of only one component of Θ (namely, the one which contains J_n). Next, f is called an *n -quasi-embedding* if in addition to the singleton $P_0 := \{x\}$ there exist compact polyhedra $P_1, \dots, P_n \subset M$ and arcs $J_0, \dots, J_n \subset \Theta$ such that $f^{-1}(P_i) \subset J_i$ for each $i \leq n$ and $P_i \cup f(J_i) \subset P_{i+1}$ for each $i < n$, and moreover the inclusion $P_i \cup f(J_i) \rightarrow P_{i+1}$ is null-homotopic for each $i < n$. It should be noted that like before, P_n can intersect the image of only one component of Θ . A PL homotopy $f_t: \Theta \rightarrow M$ is called a (*strong*) *n -quasi-isotopy* if it contains only finitely many double points, all occurring at distinct time instances $t_1, \dots, t_k \in I$, and each f_{t_i} is a (*strong*) n -quasi-embedding. Two tangles $L_0, L_1: \Theta \rightarrow M$ are (*strongly*) *n -quasi-isotopic* if they are of the same boundary pattern Ξ and are connected by a (*strong*) quasi-isotopy $L_t: \Theta \rightarrow M$ such that each L_t is a proper map of boundary pattern Ξ .




FIGURE 1. The n th Milnor link for $n = 4$.

Example 3.1. The n th Milnor link M_n (see Figure 1) is easily seen to be strongly $(n - 1)$ -quasi-isotopic to the unlink. It is not n -quasi-isotopic to the unlink, which can be shown either “geometrically”, by using $(n + 1)$ -cobordism and Cochran’s derived invariant β^n [61; Corollary 3.6] or “algebraically”, by using the fundamental group and Milnor’s invariant $\bar{\mu}(\underbrace{11 \dots 11}_{2n} 22)$ [62; Theorem 2.12].

Example 3.2. The n th Whitehead link W_n (see Figure 2) is the n th (untwisted left-handed) Whitehead double of either component of the Hopf link (due to the symmetry of




FIGURE 2. The n th Whitehead link for $n = 3$.

the Whitehead link, it does not matter which component is being doubled). It is easily seen to be $(n - 1)$ -quasi-isotopic to the unlink. In fact, there are two $(n - 1)$ -quasi-isotopies from W_n to the unlink (each with one double point), and is not hard to show [60; §1.2] that each of them is not an n -quasi-isotopy and not a strong 1-quasi-isotopy.

Conjecture 3.3. [61] (a) W_n is not n -quasi-isotopic to the unlink.
 (b) W_n is not strongly 1-quasi-isotopic to the unlink.

Let us note that either (a) or (b) would imply (see Theorem 3.5) that each W_n is not topologically isotopic to the unlink. It is known that W_n is topologically slice, and hence topologically I-equivalent to the unlink for $n \geq 3$ [21].

Remark 3.4. Let us discuss some approaches to Conjecture 3.3. The Conway polynomial vanishes on W_n for $n \geq 2$ (this follows from Lemma 10.15 and the Conway skein relation (C)). For similar reasons the multi-variable Conway polynomial also vanishes on W_n for $n \geq 2$. To compute the Jones polynomial of the Whitehead double a link L along some component K is the same task as to compute the Jones polynomial of the link obtained from L by adding a parallel copy of K (this follows from the Jones skein relation and the formula [47; p. 26] for reversing the orientation of a component). So this is closely related to the task of computation of the colored Jones polynomial. A formula for the colored Jones polynomial of a Whitehead double of a knot is given in [80] and [92; §2], and the colored Jones polynomial of a twisted version of W_n is calculated in [89; §4]. See also [22] for a related computation.

Many further examples and results on n -quasi-isotopy can be found in [60] and [61]. See also Theorem 10.14 below.

Theorem 1.1 is a consequence of the following

Theorem 3.5. [60; Theorem 1.3] *Let $h_t: \Theta \rightarrow M$ be a topological isotopy of a compact 1-manifold in a 3-manifold. Then for each n there exists an $\epsilon > 0$ such that every PL*

homotopy $f_t: \Theta \rightarrow M$ which is ϵ -close to h_t and has only finitely many double points occurring at distinct time instances is a strong n -quasi-isotopy.

Since the proof sketch given in [60] covers explicitly only the case $n = 1$, we include the details here. (The case $n = 1$ might be the most important one for the purposes of [60; Problem 1.1], but it certainly does not suffice for the purposes of the present paper.)

Proof. To simplify the argument we use a metric on M whose closed ϵ -balls are PL balls. For instance, for $M = \mathbb{R}^3$ the usual l_∞ metric will do. In general, one such metric on M is given by the following construction (see [55] for its further discussion). Let K be a triangulation of M and V the set of its vertices. The set $\mathbb{R}[V]$ of all finite formal linear combinations $\lambda_1 v_1 + \dots + \lambda_k v_k$, where $v_i \in V$ and $\lambda_i \in \mathbb{R}$, is a vector space over \mathbb{R} , which can be endowed with the l_∞ metric. Let $f: M \rightarrow \mathbb{R}[V]$ be defined by sending the barycenter of every simplex σ of K to the sum of the vertices of σ and extending linearly to every simplex of the barycentric subdivision of K . It is easy to see that f is an embedding and that the f -image of each k -simplex σ of K is a union of $k + 1$ of k -dimensional faces of the cube $[0, 1]^{k+1} \subset \mathbb{R}^{k+1} = \mathbb{R}[\text{vertices of } \sigma]$. It follows that the induced metric on M is as desired.

Lemma 3.6. *For every $\delta > 0$ there exists a $\gamma = \gamma(\delta) > 0$ such that for any triple $(p, q, t) \in \Theta \times \Theta \times I$ such that $h_t(p)$ and $h_t(q)$ are γ -close, p and q belong to the same component C of Θ and are not antipodal in it (in the event that $C \cong S^1$), and the h_t -image of the shortest arc $[p, q] \subset C$ between p and q is of diameter $< \delta$.*

Proof. Given a $\delta > 0$, let $V_\delta \subset \Theta \times \Theta \times I$ denote the set of all triples (p, q, t) such that p and q belong to the same component of Θ and are not antipodal in it, and $h_t([p, q])$ is of diameter $< \delta$. It is easy to see that V_δ is open and contains $\Delta_\Theta \times I$, where $\Delta_X = \{(x, x) \in X \times X\}$. Then $\Theta \times \Theta \times I \setminus V_\delta$ is compact, and hence so is its image K_δ under the map $H: \Theta \times \Theta \times I \rightarrow M \times M$, defined by $H(x, y, t) = (h_t(x), h_t(y))$. Since h_t is an isotopy, K_δ is disjoint from Δ_M . Then there exists a $\gamma > 0$ such that $M \times M \setminus K_\delta$ contains every pair $(x, y) \in M \times M$ such that x and y are at a distance $\leq \gamma$. \square

Let ϵ_{n+1} be any positive real number. Assuming that ϵ_i is defined, let $\epsilon_{i-1} = \frac{1}{2}\gamma(\frac{\epsilon_i}{2})$ be given by Lemma 3.6. Let us note that each $\epsilon_{i-1} \leq \frac{\epsilon_i}{4}$. Finally, set $\epsilon = \epsilon_0$.

Let $f_t: (\Theta, \partial\Theta) \rightarrow (M, \partial M)$ be a PL homotopy, ϵ -close to h_t and possessing only finitely many double points that occur at distinct time instances $t_1, \dots, t_k \in I$. Let us fix some $j \in \{1, \dots, k\}$, let us write $f = f_{t_j}$ and $h = h_{t_j}$, and let $x := f(p) = f(q)$, $p \neq q$, be the corresponding double point. Let $B_0 = \{x\}$ and for $i = 1, \dots, n$ let B_i be the closed ball of radius ϵ_i about x . Thus $B_0 \subset \dots \subset B_n$.

Since h is ϵ -close to f , the distance between $h(p)$ and $h(q)$ is at most $2\epsilon = \gamma(\frac{\epsilon_1}{2})$. Hence by Lemma 3.6 p and q belong to the same component of Θ and are not antipodal in it, and $h([p, q])$ is of diameter $< \frac{\epsilon_1}{2}$. Set $J_0 = [p, q]$. Then $f(J_0)$ is of diameter $< \frac{\epsilon_1}{2} + 2\epsilon \leq \epsilon_1$. Hence it lies in B_1 .

Let $i \in \{1, \dots, n\}$. Given any $r \in \Theta$ such that $f(r) \in B_i$, the distance from $f(r)$ to $x = f(p)$ is at most ϵ_i . Hence the distance from $h(r)$ to $h(p)$ is at most $\epsilon_i + 2\epsilon \leq 2\epsilon_i =$

$\gamma(\frac{\epsilon_{i+1}}{2})$. Then by Lemma 3.6 r belongs to the same component of Θ as p and is not antipodal to p in it, and $h([p, r])$ is of diameter $< \frac{\epsilon_{i+1}}{2}$. Hence $f([p, r])$ is of diameter $< \frac{\epsilon_{i+1}}{2} + 2\epsilon \leq \epsilon_{i+1}$. Hence it lies in B_{i+1} , unless $i = n$ (in which case B_{i+1} is undefined, because it is not needed). Let J_i be the union of all arcs of the form $[p, r]$, where $r \in \Theta$ is such that $f(r) \in B_i$. Clearly J_i is closed, connected and does not contain the antipode of p , so it is a closed arc. By construction $f^{-1}(B_i) \subset J_i$ and if $i < n$ also $f(J_i) \subset B_{i+1}$. \square

4. FINITE TYPE INVARIANTS AND COLORED FINITE TYPE INVARIANTS


By a *singular tangle* we mean a proper PL map $L: \Theta \rightarrow M$ of a compact 1-manifold in a 3-manifold, which has no triple points and only finitely many double points, and every its double point $L(p) = L(q)$ is *rigid* in the sense that the tangent vectors $dL_p(1)$ and $dL_q(1)$ exist (i.e. L has no kinks at p and q) and are linearly independent. Two singular tangles $L, L': \Theta \rightarrow M$ are called *equivalent* if they are ambient isotopic keeping ∂M fixed and so that all the double points remain rigid at every time instant.

For the rest of this section we fix a compact 1-manifold Θ , an oriented 3-manifold M and a boundary pattern $\Xi: \partial\Theta \rightarrow \partial M$, and by “tangles” or “singular tangles” mean only those of this boundary pattern. (For example, these can be links in S^3 or string links.)

Given an invariant v of tangles, which assumes values in an abelian group G , we claim that it uniquely extends to a G -valued invariant v^\times of singular tangles satisfying the relation

$$v^\times(L) = v^\times(L_+) - v^\times(L_-), \tag{V}$$

where L, L_+ and L_- agree outside a small ball, and inside this ball they are as follows:⁵



Indeed, the value of v^\times on a given singular tangle can be computed by inductively resolving its double points in some order, and applying the relation (V) at each stage. This results in the formula

$$v^\times(L) = \sum_{\epsilon_1, \dots, \epsilon_n \in \{1, -1\}} \epsilon_1 \cdots \epsilon_n v(L_{\epsilon_1 \dots \epsilon_n}) \tag{V'}$$

for every singular tangle L with n double points, where $L_{\epsilon_1 \dots \epsilon_n}$ is the tangle obtained by resolving L at all its double points, with the resolution at the i th double point being of the sign ϵ_i . But it is easy to see that this formula is not affected by reordering the double points of L . (And this is where we need the group G to be abelian.)

⁵Let us note that L_+ and L_- can be distinguished from each other without using the plane diagram. Namely, the frame $(d(L_\epsilon)_p(1), d(L_\epsilon)_q(1), L_\epsilon(p) - L_\epsilon(q))$ in $T_{L(p)}M$ preserves its sign when p and q are interchanged (and also when the orientation of Θ is reversed), but reverses the sign when ϵ is reversed.

A G -valued invariant v of tangles is said to be of *type n* if v^\times vanishes on all singular tangles with $n + 1$ double points (and therefore also on all singular tangles with $> n$ double points). It is said to be of *finite type* if it is of type n for some n . Finite type invariants were introduced by M. Gusarov (cf. the editor's comment in [29]) and independently by V. Vasiliev [84]; the case of links appears in [27].

The following lemmas follow easily from the definition.

Lemma 4.1. *Let v be a type n invariant of m -component links in S^3 . Let v_c be an invariant of m -component string links defined by evaluating v on the the closure. Then v_c is a type n invariant.*

Lemma 4.2. (a) *Let v be a type n invariant of knots in S^3 . Let v_i^λ be an invariant of m -component links in S^3 defined by evaluating v on the i th component. Then v_i^λ is a type n invariant.*

(b) *Let v be a type n invariant of m -component links in S^3 . Let v_i^k be an invariant of knots in S^3 defined by evaluating v on the totally split link whose i th component is the given knot and the other components are unknotted. Then v_i^k is a type n invariant.*

It is easy to see that G -valued type n invariants of tangles $\Theta \rightarrow M$ of boundary pattern Ξ form an abelian group. In particular, G -valued type n invariants of m -component links in S^3 form an abelian group $G_{m,n}$. Lemma 4.2 identifies the direct product of m copies of $G_{1,n}$ with a direct summand of $G_{m,n}$.

An alternative notion of finite type invariants was introduced by Kirk and Livingston [40]. Let us recall that a *link map* is a map which sends distinct connected components to disjoint sets. A G -valued invariant v of tangles is said to be of *colored type n* if v^\times vanishes on those singular tangles with $n + 1$ double points that are link maps (and therefore also on those singular tangles with $> n$ double points that are link maps). The invariant v is said to be of *colored finite type* if it is of colored type n for some n .

Example 4.3. The colored type 0 invariant $v(L) = (-1)^{\text{lk}(L)}$ is not of finite type.

Indeed, arguing by induction, it is easy to see that $v^\times(L_s) = \pm 2^k$ for any singular link L_s with k intersections between distinct components and no self-intersections.

Example 4.4. The coefficients of the Conway polynomial are of finite type, but their colored type is usually less than their type (see Lemma 10.3). The coefficients of the multi-variable Conway polynomial are “naturally occurring” invariants of colored finite type that are not of finite type [58].

It is sometimes more natural to consider a different filtration on colored finite type invariants. Assuming that Θ consists of m connected components $\Theta_1, \dots, \Theta_m$, the invariant v is said to be of *type (k_1, \dots, k_m)* if v^\times vanishes on all singular links with $k_1 + 1$ self-intersections of Θ_1 , on all singular links with $k_2 + 1$ self-intersections of Θ_2 , and so on. It is easy to see that every type (k_1, \dots, k_m) invariant is a colored type $k_1 + \dots + k_m$ invariant; and every colored type n invariant is a type (n, \dots, n) invariant. Thus v is of colored finite type if and only if it is of type (k_1, \dots, k_m) for some k_1, \dots, k_m .

Lemma 4.5. *Let R be a ring and u, v be R -valued invariants of tangles $\Theta \rightarrow M$ with a fixed boundary pattern. Then*

$$(uv)^\times(L) = \sum_{\substack{S \cup T = \Delta_L \\ S \cap T = \emptyset}} u^\times(L_{S+})v^\times(L_{T-}),$$

where for a singular tangle L with a finite set Δ_L of double points and for an $S \subset \Delta_L$, by L_{S+} (resp. L_{S-}) we denote the partial resolution of L where the double points in S are resolved positively (resp. negatively) and the double points in $\Delta_L \setminus S$ remain unresolved.

When L has only one double point, the Leibniz rule of Lemma 4.5 specializes to

$$(uv)^\times(L) = u(L_+)v^\times(L) + u^\times(L)v(L_-).$$

A more symmetric, but more complicated Leibniz rule is proved in [86; Theorem 3.2]. We will use the specific form of our Leibniz rule in the proof of Proposition 10.13.

Proof. Let us note that the lemma holds trivially for non-singular tangles. Let us assume that it holds for singular tangles with fewer double points than L . Pick some $x \in \Delta_L$. Then by (V) $(uv)^\times(L) = (uv)^\times(L_{\{x\}+}) - (uv)^\times(L_{\{x\}-})$. Applying the induction hypothesis and abbreviating $(L_{S\epsilon})_{T\delta}$ by $L_{S\epsilon, T\delta}$, the latter expression takes the form

$$\sum_{\substack{S \cup T = \Delta_L \setminus \{x\} \\ S \cap T = \emptyset}} u^\times(L_{\{x\}+, S+})v^\times(L_{\{x\}+, T-}) - u^\times(L_{\{x\}-, S+})v^\times(L_{\{x\}-, T-}).$$

If we subtract $u^\times(L_{\{x\}+, S+})v^\times(L_{\{x\}-, T-})$ from the left hand term and add it to the right hand term, we obtain

$$\sum_{\substack{S \cup T = \Delta_L \setminus \{x\} \\ S \cap T = \emptyset}} u^\times(L_{\{x\}+, S+})v^\times(L_{T-}) + u^\times(L_{S+})v^\times(L_{\{x\}-, T-}).$$

But this clearly equals the right hand side of (4.5). \square

Corollary 4.6. [28], [86] (a) *If u, v are of (colored) types k, l respectively, then uv is of (colored) type $k + l$.*

(b) *If u, v are of types (k_1, \dots, k_m) and (l_1, \dots, l_m) respectively, then uv is of type $(k_1 + l_1, \dots, k_m + l_m)$.*

The non-colored case is proved in [28; Lemma 5.8], [86; Corollary 3.3] and [13; 3.2.3] (see also [7; Exercise 3.10]), and the proofs in [86], [13] work in both colored cases.

Remark 4.7. Given a *coloring* of Θ , that is, a continuous map $c: \Theta \rightarrow \{1, \dots, \chi\}$, one may define a c -link map as a map which sends distinct point-inverses of c to disjoint sets. (One may think of c as dipping the components of Θ into paintpots numbered $1, \dots, \chi$.) This leads to invariants of c -type n . The case of a constant coloring corresponds to type n invariants, and the case of a coloring which is itself a link map corresponds to colored type n invariants. These are the only two cases that are needed in the present paper.

Theorem B is a consequence of the following

Theorem 4.8. *Let v be a type n invariant, or more generally a type (n, \dots, n) invariant of tangles. If v is invariant under PL isotopy, then it is invariant under n -quasi-isotopy.*

Proof. *Case $n = 0$.* The relation of 0-quasi-isotopy is the same as that of link homotopy. On the other hand, every type $(0, \dots, 0)$ invariant assumes the same values on link homotopic tangles. This completes the proof of the theorem in the case $n = 0$. \square

Step I (includes the case $n = 1$). Assume that $n \geq 1$ and that v is invariant under PL isotopy. It suffices to show that v^\times vanishes on every proper n -quasi-embedding $f: \Theta \rightarrow M$ of boundary pattern Ξ which is a singular tangle (i.e. its double point is rigid). Let $x = f(p) = f(q)$ be the double point of f . Let P_0, \dots, P_n and J_0, \dots, J_n be as in the definition of an n -quasi-embedding. We may assume that $\partial J_0 = \{p, q\}$. By passing to small regular neighborhoods we may further assume that P_1, \dots, P_n are compact 3-manifolds (with boundary) and that each J_i , $i < n$, lies in the interior of an arc J_i^+ such that $P_i \cup f(J_i^+)$ lies in P_{i+1} and is null-homotopic in it (and also $f^{-1}(P_i) \subset J_i$, as before). Moreover, it is easy to see that $\overline{P_0} = \{x\}$ lies in the interior of a PL ball P_0^+ which meets $f(\Theta \setminus J_0)$ in the arc $f(J_0^+ \setminus J_0)$, lies in P_1 and is such that the inclusion $P_0^+ \cup f(J_0^+) \rightarrow P_1$ is null-homotopic.

Since $P_0^+ \cup f(J_0)$ is null-homotopic in P_1 , there exists a homotopy $f_t: \Theta \rightarrow M$ keeping $\overline{\Theta \setminus J_0}$ fixed and keeping J_0 within P_1 , from $f_0 = f$ to a map f_1 sending J_0 into the ball P_0^+ and such that $f_1(J_0)$ is a small circle bounding a small embedded disk in P_0^+ , whose interior is disjoint from $f(J_0^+ \setminus J_0)$. Moreover, since P_0^+ and P_1 are 3-manifolds, we may assume that x is a rigid double point of f_t for each $t \in I$, and apart from this permanent double point, the homotopy f_t has only finitely many additional double points that occur at distinct time instants $t_1, \dots, t_l \in (0, 1)$ and are rigid. Thus each f_{t_i} is a singular tangle with two double points, and f_t for each $t \notin \{t_1, \dots, t_l\}$ is a singular tangle with one double point.

Since the two resolutions L_+ , L_- of the singular tangle $L := f_1$ are ambient isotopic, every colored finite type invariant, when extended to singular tangles, vanishes on f_1 . (This relation is known as “the one-term relation” or “the framing independence relation” in the theory of finite type invariants, cf. [13; §4].)

On the other hand, since $f^{-1}(P_1)$, and hence also each $f_{t_i}^{-1}(P_1)$, lies in the arc J_1 , the two double points of each f_{t_i} are self-intersections of the same component. Hence every type $(1, \dots, 1)$ invariant u , when extended to singular tangles, vanishes on each f_{t_i} . Therefore $u^\times(f) = u^\times(f_1)$, but from the one-term relation we also know that $u^\times(f_1) = 0$. This completes the proof of the theorem in the case $n = 1$. \square

The following lemma can be regarded as a generalization of the one-term relation.

Lemma 4.9. *Let L be a singular tangle with n double points, where $n \geq 1$, and suppose that all its double points lie in the interior of a PL ball B such that $L^{-1}(B)$ is an arc. Then for every invariant u of PL isotopy, $u^\times(L) = 0$.*

Proof. Pick any double point of L and consider the two resolutions L_+ and L_- of L at this double point.

If $n = 1$, then L_+ and L_- are non-singular tangles which meet the ball B in possibly distinct local knots and agree outside B . Hence they are PL isotopic, and therefore $u^\times(L) = u(L_+) - u(L_-) = 0$.

If $n > 1$, then, arguing by induction, we may assume that the lemma is established for all smaller values of n . Then $u^\times(L_+) = u^\times(L_-) = 0$ by the induction hypothesis, and hence $u^\times(L) = 0$. \square

Step II (includes the case $n = 2$). To continue the proof of the theorem for $n \geq 2$, we fix some $i \in \{1, \dots, l\}$ and study the jump of v^\times on the singular tangle f_{t_i} .

Since $f^{-1}(P_1)$ lies in J_1 , so does $f_{t_i}^{-1}(P_1)$, and consequently both double points of f_{t_i} lie in the graph $G := f_{t_i}(J_1)$. Let g be an embedding of the graph $G^+ := f_{t_i}(J_1^+)$ into $P_0^+ \cup f(J_1^+ \setminus J_0)$ which restricts to an embedding of G into the ball P_0^+ , to the identity on the two points $f_{t_i}(\partial J_1^+) = f(\partial J_1^+)$ and to a homeomorphism between the two arcs $f_{t_i}(\overline{J_1^+ \setminus J_1})$ and the two arcs $f(\overline{J_1^+ \setminus J_0^+})$. Since G^+ and $g(G^+)$ both lie in $P_1 \cup f(J_1^+)$, which is in turn null-homotopic in P_2 , the inclusion $G^+ \rightarrow P_2$ is homotopic to g by a homotopy $g_t: G^+ \rightarrow P_2$ keeping $f(\partial J_1^+)$ fixed.⁶ The composition $J_1^+ \xrightarrow{f_{t_i}} f_{t_i}(J_1^+) = G^+ \xrightarrow{g_t} P_2$ extends by the identity on $\Theta \setminus J_1^+$ to a homotopy $f_{t_i;t}: \Theta \rightarrow M$ from $f_{t_i;0} = f_{t_i}$ to a map $f_{t_i;1}$ which has precisely two double points, both contained in P_0^+ , and is such that $f_{t_i;1}^{-1}(P_0^+) = J_1$. Moreover, since P_0^+ and P_2 are 3-manifolds, we may assume that the two double points of f_{t_i} continue as rigid double points of $f_{t_i;t}$ for each $t \in I$, and apart from these two permanent double points, the homotopy $f_{t_i;t}$ has only finitely many additional double points that occur at distinct time instants $t_{i1}, \dots, t_{ik_i} \in (0, 1)$ and are rigid. Thus each $f_{t_i;t_{ij}}$ is a singular tangle with three double points and $f_{t_i;t}$ for each $t \notin \{t_{i1}, \dots, t_{ik_i}\}$ is a singular tangle with two double points.

Since both double points of $f_{t_i;1}$ are contained in P_0^+ and $f_{t_i;1}^{-1}(P_0^+)$ is the arc J_1 , by Lemma 4.9 every colored finite type invariant, when extended to singular tangles, vanishes on $f_{t_i;1}$.

On the other hand, since $f^{-1}(P_2)$, and hence also each $f_{t_i;t_{ij}}^{-1}(P_2)$, lies in the arc J_2 , the three double points of each $f_{t_i;t_{ij}}$ are self-intersections of the same component. Hence every type $(2, \dots, 2)$ invariant u , when extended to singular tangles, vanishes on each $f_{t_i;t_{ij}}$. Therefore $u^\times(f_{t_i}) = u^\times(f_{t_i;1})$, but from Lemma 4.9 we also know that $u^\times(f_{t_i;1}) = 0$. Thus u^\times vanishes on each f_{t_i} . Consequently $u^\times(f) = u^\times(f_1)$, but from the one-term relation we also know that $u^\times(f_1) = 0$. This completes the proof of the theorem in the case $n = 2$. \square

The proof of the general case is similar to the case $n = 2$ (and if the reader feels that reading the proof of the case $n = 2$ was enough for her to understand the proof of the general case, she is probably right), but for completeness we include the details.

Step III (Case $n \geq 2$). Suppose that for each $k = 1, \dots, n - 1$ we have constructed a collection C_k of homotopies $f_{s_1, \dots, s_{k-1}; t}$ which go through singular tangles with k double

⁶Indeed, if $X \supset A \subset Y$, the inclusion $A \rightarrow Y$ is null-homotopic and the inclusion $A \rightarrow X$ is a cofibration, then it is easy to see that the inclusion $A \rightarrow Y$ extends to a map $X \rightarrow Y$.

points, apart from finitely many critical levels, which are singular tangles with $k + 1$ double points, so that the following conditions hold.

- (1) $f_0 = f$, and for every $k \in \{1, \dots, n - 2\}$ a singular tangle is the initial level $f_{s_1, \dots, s_k; 0}$ of a homotopy of the form $f_{s_1, \dots, s_k; t} \in C_{k+1}$ if and only if it is a critical level, corresponding to $t = s_k$, of a homotopy of the form $f_{s_1, \dots, s_{k-1}; t} \in C_k$.
- (2) The terminal level $f_{s_1, \dots, s_k; 1}$ of every homotopy $f_{s_1, \dots, s_k; t} \in C_{k+1}$ has all its double points in the ball P_0^+ and satisfies $f_{s_1, \dots, s_k; 1}^{-1}(P_0^+) = J_k$.
- (3) Every level of every homotopy $f_{s_1, \dots, s_{k-1}; t} \in C_k$ agrees with f outside J_{k-1}^+ , and sends J_{k-1}^+ into P_k .

Let us show that the given collections C_1, \dots, C_{n-1} can be extended by a collection C_n so that the collections C_1, \dots, C_n satisfy the same conditions with n replaced by $n + 1$.

Suppose that s_{n-1} is a critical time instant of a homotopy $f_{s_1, \dots, s_{n-2}; t} \in C_{n-1}$. Let $S = (s_1, \dots, s_{n-1})$. Thus f_S is a singular tangle with n double points, which agrees with f outside J_{n-2}^+ and sends J_{n-2}^+ into P_{n-1} . Since $f^{-1}(P_{n-1})$ lies in J_{n-1} , so does $f_S^{-1}(P_{n-1})$, and consequently all double points of f_S lie in the graph $G := f_S(J_{n-1})$. Let g be an embedding of the graph $G^+ := f_S(J_{n-1}^+)$ into $P_0^+ \cup f(\overline{J_{n-1}^+ \setminus J_0})$ which restricts to an embedding of G into the ball P_0^+ , to the identity on the two points $f_S(\partial J_{n-1}^+) = f(\partial J_{n-1}^+)$ and to a homeomorphism between the two arcs $f_S(\overline{J_{n-1}^+ \setminus J_{n-1}})$ and the two arcs $f(\overline{J_{n-1}^+ \setminus J_0^+})$. Since G^+ and $g(G^+)$ both lie in $P_{n-1} \cup f(J_{n-1}^+)$, which is in turn null-homotopic in P_n , the inclusion $G^+ \rightarrow P_n$ is homotopic to g by a homotopy $g_t: G^+ \rightarrow P_n$ keeping $f(\partial J_{n-1}^+)$ fixed. The composition $J_{n-1}^+ \xrightarrow{f_S} f_S(J_{n-1}^+) = G^+ \xrightarrow{g_t} P_n$ extends by the identity on $\overline{\Theta \setminus J_{n-1}^+}$ to a homotopy $f_{S;t}: \Theta \rightarrow M$ from $f_{S;0} = f_S$ to a map $f_{S;1}$ which has precisely n double points, all contained in P_0^+ , and is such that $f_{S;1}^{-1}(P_0^+) = J_{n-1}$. Moreover, since P_0^+ and P_n are 3-manifolds, we may assume that the n permanent double points, the homotopy $f_{S;t}$ has only finitely many additional double points that occur at distinct time instances $u_1, \dots, u_k \in (0, 1)$ and are rigid. Thus each $f_{S;u_j}$ is a singular tangle with $n + 1$ double points and $f_{S;t}$ for each $t \notin \{u_1, \dots, u_k\}$ is a singular tangle with n double points. This completes the construction of the desired collection C_n .

Now that the collections C_1, \dots, C_n have been constructed, we are ready to evaluate the given invariant v of type (n, \dots, n) , which is well-defined up to PL isotopy, on the n -quasi-embedding f . For each $k = 1, \dots, n$ and each homotopy $f_{S;t} \in C_k$ we have $v^\times(f_{S;1}) = 0$ by condition (2) and Lemma 4.9. Since v is a type (n, \dots, n) invariant, and each critical level $f_{S;u_j}$ of each homotopy $f_{S;t} \in C_n$ has $n + 1$ double points, all of them being self-intersections of the same component of Θ (indeed, they lie in $f_{S;u_j}(J_n)$ by condition (3)), we also get that $v^\times(f_{S;0}) = v^\times(f_{S;1})$. Thus $v^\times(f_{S;0}) = 0$ for each $f_{S;t} \in C_n$. Hence by condition (1) we obtain that v^\times vanishes on each critical level of each homotopy in C_{n-1} . Proceeding in the same fashion, we eventually obtain that $v(f) = 0$. \square

Remark 4.10. The proof of Theorem 4.8 can be modified so as to work under a weaker hypothesis, with n -quasi-isotopy replaced by *virtual n -quasi-isotopy*, which is defined in [60; end of §3]. The modified construction is geometrically more intuitive, but its accurate description would be longer and more complicated. We give only a rough sketch of the modified construction. The homotopy f_t is the same as before. Using the notation from [60; proof of Lemma 3.1], each homotopy $f_{t_j,t}$ can be visualized as shifting the arc $f(I'_j)$ onto the arc $F(I_j)$ and then taking the image of J_1 into a ball along the track of the null-homotopy F . Thus $f_{t_j,1}$ still satisfies the hypothesis of Lemma 4.9, but, unless $I'_j \subset J_0$, the image of $f_{t_j,1}$ is now different: instead of being the same as the image of f outside the ball P_0^+ , it now coincides with the image of f only outside a small neighborhood of the arc $f(I'_j)$.

Remark 4.11. The proof of Theorem 4.8, as well as its modification sketched in the previous remark, can be seen to yield a stronger conclusion: if tangles L and L' are n -quasi-isotopic (or just virtually n -quasi-isotopic), then they are not just indistinguishable by type (n, \dots, n) invariants that are well-defined up to PL isotopy, but also *geometrically (n, \dots, n) -equivalent* in the following sense. Let \mathcal{LM} denote the space of all singular links which are link maps. (The topology is the restriction of the C^0 topology on the space of maps, modified so as to ensure that double points remain rigid under homotopies given by paths in this space.) Let \mathcal{LM}_n denote the subset of \mathcal{LM} consisting of singular links with precisely n double points, and let $\mathcal{LM}_{(n,\dots,n)}$ denote the subset of \mathcal{LM}_n consisting of singular links whose all double points occur on the same component. On the other hand, let \mathcal{LM}° denote the subset of \mathcal{LM} consisting of those singular links L whose all double points lie in a ball B such that $L^{-1}(B)$ is an arc. We call two elements of \mathcal{LM}_n *geometrically k -equivalent* if they are homotopic in the space $\mathcal{LM}_n \cup \mathcal{LM}_{n+1,k}$, where $\mathcal{LM}_{i,0} = \mathcal{LM}_i$ and $\mathcal{LM}_{i,k}$ for $k > 0$ consists of those elements of \mathcal{LM}_i that are geometrically $(k-1)$ -equivalent to an element of $\mathcal{LM}_i \cap \mathcal{LM}^\circ$. Two elements of $\mathcal{LM}_{(n,\dots,n)}$ are *geometrically (k, \dots, k) -equivalent* if the same holds with $\mathcal{LM}_{(i,\dots,i)}$ in place of \mathcal{LM}_i . It is shown in the proof of Theorem 4.8 that if links L and L' are geometrically (k, \dots, k) -equivalent, then they are not separated by type (k, \dots, k) invariants. The same argument works to show that if L and L' are geometrically k -equivalent, then they are not separated by colored type k invariants.

Example 4.12. (a) It is easy to see that geometric 1-equivalence implies weak 1-quasi-isotopy.⁷ For 2-component links the converse also holds (see [66; Figure 2]).

(b) It is shown in [40] that 2-component links are not separated by colored type 1 invariants if and only if they have the same linking number and the same generalized Sato–Levine invariant (see Example 1.6 concerning the latter). However, it seems highly unlikely that these two invariants constitute a complete set of invariants of weak 1-quasi-isotopy (see, in particular, [60; Problem 1.5], which also makes sense for weak

⁷Weak n -quasi-isotopy is defined similarly to n -quasi-isotopy but with “null-homotopic” replaced by “induces zero homomorphisms on reduced integral homology” (cf. [60], [61], see also [60; Figure 2(d)]).

1-quasi-isotopy in place of 1-quasi-isotopy). In fact they constitute a complete set of invariants of $\frac{1}{2}$ -quasi-isotopy (see Example 1.6).

(c) Geometric $(1, \dots, 1)$ -equivalence is clearly the same thing as 1-quasi-isotopy.

Example 4.13. Let $M_k = (K, Q)$ be the k th Milnor link (see Figure 1), where Q is the “long” component, and let M_k^W , $k \geq 2$, be a Whitehead double of M_k along Q . Thus $M_k^W = (K, Q')$, where Q' is the image of the Whitehead curve in $S^1 \times D^2$ under some homeomorphism of $S^1 \times D^2$ with a regular neighborhood of Q . Unclasping the clasp of Q' yields a 1-quasi-isotopy h_t from M_k^W to the unlink. It is not a 2-quasi-isotopy (not even a weak 2-quasi-isotopy), since M_k is not a boundary link, as detected by Cochran’s derived invariants (see [61; §3 and §1.2]). It is natural to conjecture that M_k^W is not 2-quasi-isotopic, nor even weakly 2-quasi-isotopic to the unlink. However, it is not hard to see that h_t yields a geometric (k, \dots, k) -equivalence between M_k^W and the unlink.

5. LOCALLY ADDITIVE INVARIANTS AND C_n -EQUIVALENCE

A tangle is called *totally split* if its components are contained in pairwise disjoint 3-balls. If K and L are m -component links and K is totally split, then the (componentwise) connected sum $L\#K$ is easily seen to be well-defined. Also, for any string links L and L' their connected sum $L\#L'$ is well defined.

We call an invariant v of m -component (string) links *locally additive* if $v(L\#K) = v(L) + v(K)$ whenever K is totally split. This implies in particular that v vanishes on the trivial (string) link.

For every link L in S^3 (respectively, for every string link) there is a unique, up to ambient isotopy, totally split link in S^3 (respectively, totally split string link) K_L whose components are ambient isotopic to the respective components of L .

Lemma 5.1. *Let A and B be m -component links in S^3 (respectively, string links). Then A and B are not separated by type n invariants that are well-defined up to PL isotopy if and only if A and B are PL isotopic to (string) links that are not separated by locally additive type n invariants.*

Proof. “If”. It suffices to observe that if an invariant is well-defined up to PL isotopy, then it is locally additive.

“Only if”. Clearly, $A' := A\#K_B$ and $B' := B\#K_A$ are PL isotopic to A and B (respectively). Let v be a locally additive type n invariant of m -component (string) links. Then \bar{v} , defined by $\bar{v}(L) = v(L) - v(K_L)$, is a type n invariant which is well-defined up to PL isotopy. (Let us note that $v(K_L) = (v_1^\kappa)_1^\lambda(L) + \dots + (v_m^\kappa)_m^\lambda(L)$ in the notation of Lemma 4.2.) We have $\bar{v}(A') = \bar{v}(A) = \bar{v}(B) = \bar{v}(B')$. Since $K_{A'} = K_{B'}$, we obtain that $v(A') = \bar{v}(A') + v(K_{A'}) = \bar{v}(B') + v(K_{B'}) = v(B')$. \square

Theorem 5.2. *If two m -component string links are separated by invariants of type n , then they are separated by locally additive invariants of type r for some $r = r(m, n)$.*

Theorem 5.2 will be proved in the present section by building on Habiro’s proof [32; Theorem 6.18] of T. Stanford’s theorem [79; Theorem 2.43] (cf. [32; Remark 3.19])

that two knots are separated by type n invariants if and only if they are separated by additive type n invariants. (The rational version of this theorem was originally proved by Gusarov [28; Theorem 5.2]; see also [81] for an alternative proof.) Habiro’s proof is in turn based on the study of C_n -equivalence.




FIGURE 3. The C_n -move for $n = 4$.

Two tangles $L, L': \Theta \rightarrow M$ are said to be related by a C_n -move if they agree outside a 3-ball B , and intersect B along the $(n + 1)$ -strand tangles $T, T': (n + 1)I \rightarrow I^3$, which are shown in Figure 3 in the case $n = 4$ and are similarly drawn for arbitrary $n \geq 1$. If all the $n + 1$ strands of T belong to the same component of L , then the C_n -move is said to be a *self C_n -move*. L and L' are said to be (*self*) C_n -equivalent if they are related by a sequence of (*self*) C_n -moves and ambient isotopies.

Remark 5.3. It is easy to see that the C_n -move is “Brunnian” in the sense that it becomes trivial (i.e. is realized by an ambient isotopy) if for any $i \in \{1, \dots, n + 1\}$, the i th strand is erased from the picture (both on the left and on the right).

Remark 5.4. The literature contains at least four different definitions of a C_n -move, all leading to the same notion of C_n -equivalence:

- (1) Habiro’s second definition [32; §3.2 (see also §7)]: a “ C_n -move” on L is the operation of taking a band connected sum of L with an arbitrary n -tuply iterated untwisted Bing doubling Λ of the Hopf link,⁸ along bands which intersect a ball containing Λ in a specified way.
- (2) The operation of taking a band connected sum of L with Milnor’s $(n + 1)$ -component Brunnian link M_{n+1} , along bands which intersect a ball containing M_n in a specified way. This is a special case of Habiro’s second definition and is called “band sum with one-branched C_n -chord” in [68]. This definition is used for instance in [13].
- (3) Habiro’s original definition (from his unpublished master’s thesis, written in Japanese): a “ C_n -move” is an arbitrary n -tuply iterated Bing doubling of a crossing change⁹ (for the details see the introduction of [68]).

⁸An n -tuply iterated untwisted Bing doubling of a 2-component link is defined in [15; §7.4]. It is determined by a rooted unitrivalent tree with n trivalent vertices. The two (non-iterated) Bing doublings of the Hopf link are easily seen to be ambient isotopic to the Borromean rings and in particular to each other. This implies that an iterated untwisted Bing doubling of the Hopf link depends only on the tree and not on its root.

⁹The two (non-iterated) Bing doublings of the crossing change are easily seen to be equivalent to the Δ -move and in particular to each other. This implies that an iterated Bing doubling of the crossing change is fully determined by a unitrivalent tree and does not depend on its root.

- (4) C_n -move in the sense of Figure 3. This is a special case of Habiro’s original definition and is called a “one-branched C_n -move” in [68] and [81]. (But in later papers it is often called just a “ C_n -move”.)

A proof that (3) and (4) lead to the same notion of C_n -equivalence can be found in [81; Lemma 2.2]. For a proof that (1) and (3) lead to the same notion of C_n -equivalence see [81; Lemma 3.6]. As noted in [68; Lemma 2.1], the latter argument also shows that (2) and (4) lead to the same notion of C_n -equivalence. Moreover, as noted in [68; Lemma 2.1] and in [81; third line of the proof of Lemma 2.2], the arguments in [81; proofs of Lemmas 2.2 and 3.6] preserve the set of components of L that are involved in the moves, so in particular they apply to self C_n -moves in place of C_n -moves.

It should also be noted that Gusarov’s definition of an n -variation [29] is close to the definition (1) of a C_n -move. Gusarov’s previous definition of an n -equivalence [28], which developed Ohyaama’s notion of n -triviality, is not directly related to either of the definitions (1)–(4).

Remark 5.5. It is well-known that a C_{n+1} -move is realized by a sequence of C_n -moves (see [13; §14.2.1]).

Proposition 5.6 (Gusarov [29; 10.3], Habiro [32; 6.8]). *If two tangles are C_{n+1} -equivalent, then they are not separated by invariants of type n .*

Although the literature also contains other proofs of Proposition 5.6 (see in particular [13; §14.2.3] for a comparatively short one), the following very simple proof does not seem to appear anywhere.

Proof. Let v be an invariant of tangles with a fixed boundary pattern and v^\times be its standard extension to singular tangles with the same boundary pattern. Figure 4 shows a generic homotopy with two double points between two tangles L and L' related by a C_n -move, $n \geq 2$. The total change of v under this homotopy equals the difference of the values of v^\times on the two singular tangles L_\times and L'_\times which occur in it. Clearly L_\times and L'_\times are homotopic, so if v is a type 1 invariant, then $v^\times(L_\times) = v^\times(L'_\times)$ and hence $v(L) = v(L')$.

For $n \geq 3$ Figure 5 shows a generic homotopy with four new double points between the singular tangles L_\times and L'_\times . The total change of v^\times under this homotopy is the difference of its values on the doubly singular tangles $L_{\times\times}$ and $L'_{\times\times}$ plus the difference of its values on the other two doubly singular tangles $L_{\times\tilde{\times}}$ and $L'_{\times\tilde{\times}}$. Clearly the former two are homotopic to each other and the latter two are homotopic to each other; so if v is a type 2 invariant, then $v^\times(L_{\times\times}) = v^\times(L'_{\times\times})$ and $v^\times(L_{\times\tilde{\times}}) = v^\times(L'_{\times\tilde{\times}})$. Then $v^\times(L_\times) = v^\times(L'_\times)$ and hence $v(L) = v(L')$.

Proceeding in the same fashion, we will eventually obtain that if v is a type $n - 1$ invariant, then $v(L) = v(L')$. □

Remark 5.7. The proof of Proposition 5.6 also shows that if two links or string links are self C_{n+1} -equivalent, then they are not separated by invariants of type (n, \dots, n) .




FIGURE 4. Type 1 invariants are invariant under C_n -move for $n \geq 2$.




FIGURE 5. Type 2 invariants are invariant under C_n -move for $n \geq 3$.

Gusarov and Habiro also proved that the converse to Proposition 5.6 holds for knots [32; Theorem 6.18], [29; Remark 10.5] (the rational version was originally proved by Gusarov [28; Theorem 5.2]; alternative proofs appear in [79], [23], [90]) but does not hold for links of two components [29; Remark 10.8] and of more than two components [32; Proposition 7.4]. However, they conjectured that it holds for string links [32; Conjecture 6.13], [29; Remark 10.7], and this conjecture has been verified for $n \leq 4$ [52]. Gusarov proved a version of this conjecture for partially defined invariants [29; Theorem 10.4] and G. Massuyeau proved the following rational version of this conjecture: If two string links L, L' are not separated by rational invariants of type n , then there exists a k such that $\underbrace{L \# \dots \# L}_k$ and $\underbrace{L' \# \dots \# L'}_k$ are C_{n+1} -equivalent [51; Theorem 1.1] (see also [13; 12.6.3]).

He also proved the following:

Theorem 5.8 (Massuyeau [51; Corollary 1.3]). *There exists an $r = r(m, n)$ such that if two m -component string links are not separated by invariants of type r , then they are C_n -equivalent.*

It appears to be unknown whether Theorem 5.8 holds for (closed) links (compare [30] and in particular Problem 5.4 in there).

In fact, Massuyeau's results in [51] are stated in a more general setting of homology cylinders, and his proofs are also in that setting. For our purposes, it is preferable to understand the proof of Theorem 5.8 in the original language of string links. To this end we look more closely at Habiro's paper [32].

A slightly weaker statement, asserting that if two string links are not separated by finite type invariants then they are C_k -equivalent for all k , is also found in Habiro's paper [32; Corollary 6.12] (this result was also announced by Gusarov [29; Remark 10.7] but apparently not written up because of his premature death). However, it is deduced from [32; Theorem 6.11], whose proof in [32] contains an error. Namely, the author claims: "Since J is the augmentation ideal of the group ring of the nilpotent group $\mathcal{L}_1(\Sigma, n)/C_k$, we have $\bigcap_{l=1}^{\infty} J^l = \{0\}$." But it is not true that if J is the augmentation ideal of the (integral) group ring of a nilpotent group G , then $\bigcap_{l=1}^{\infty} J^l = \{0\}$. For instance, this statement is false for $G = \mathbb{Z}/pq$, where p and q are distinct primes (see [10; proof of Theorem 1]). Also, already for $G = \mathbb{Z}$ none of the powers J^l is zero itself, which is why Habiro's assertion [32; Corollary 6.12] is weaker than Theorem 5.8. In the case where Σ is a disk, Habiro's group $\mathcal{L}_1(\Sigma, n)/C_k$ is finitely generated [32; Theorem 5.4(3)], and in this case we will show that the error can be corrected, and moreover the result can be improved to Theorem 5.8, by using the following lemma.

Lemma 5.9 (Hartley [see below]; Massuyeau [51; Corollary 4.8]). *Let G be a finitely generated nilpotent group. Let $d_n G = G \cap (1 + \Delta^n)$, where Δ is the augmentation ideal of $\mathbb{Z}G$. Then $d_n G = 1$ for some finite n .*

It is easy to see that $\gamma_n G \subset d_n G$ for all n and it is known that $d_n G = \gamma_n G$ for $n \leq 3$ (see [71; V.5.3 and V.5.10]). It had been an open problem in group theory for about 30

years whether this holds for all n , until E. Rips solved it in 1972; it is now known that for each $n \geq 4$ there exists a finitely presented group G with $\gamma_n G = 1$ and $d_n G \neq 1$ (see [65; 2.11]); also there exists a finitely presented G with $\gamma_5 G = 1$ and $d_6 G \neq 1$ [65; 2.15]. However, $\gamma_n G$ always has a finite index in $d_n G$ and this index is bounded above by a certain explicit function of n (see [65; 2.17]). According to [65; Problem 2.51], it is unknown whether Lemma 5.9 holds when G is not finitely generated.

Remark 5.10. (a) A short proof of Lemma 5.9 for finite G appears in [10; Corollary 1].

(b) It is also easy to prove that $\bigcap_n d_n G = 1$ under the hypothesis of Lemma 5.9. Indeed, suppose that $\bigcap_{n=1}^{\infty} d_n G$ contains an element $g \neq 1$. Since G is finitely generated and nilpotent, it is polycyclic and hence residually finite (see [73; 5.2.18 and 5.4.17]). Thus G admits a homomorphism f onto a finite group Q such that $f(g) \neq 1$. Clearly, Q is nilpotent (see [33; 10.3.1]) and $f(g) \in \bigcap_{n=1}^{\infty} d_n Q$, which contradicts the case of Lemma 5.9 where G is finite [10; Corollary 1].

(c) Lemma 5.9 can be seen to follow from a result of Hartley [34]. In more detail, let T be the torsion subgroup of G . Since G is finitely generated and nilpotent, it contains no infinitely ascending chains of subgroups (see [73; 5.2.18]), and it follows that T is finitely generated. Since T is a finitely generated nilpotent torsion group, it is finite (see [73; 5.2.18 or 5.4.11]). Since T is a finite nilpotent group, by the case of Lemma 5.9 where G is finite [10; Corollary 1] $d_m T = 1$ for some finite m . Since G is nilpotent, by Hartley's theorem [34; Corollary A1] this implies that $d_n G = 1$ for some finite n .

Alternative proof of Theorem 5.8. This is a correction of [32; proof of Theorem 6.11].

Let us fix an $m > 1$, and let \mathcal{L} be the monoid of m -component string links. Since the relation of C_k -equivalence, viewed as a subset of $\mathcal{L} \times \mathcal{L}$, is a submonoid of $\mathcal{L} \times \mathcal{L}$, the quotient G_k of \mathcal{L} by the relation of C_k -equivalence is a monoid. Moreover, the following was shown by Habiro (that G_k is a group, and that it is nilpotent was also shown by Gusarov [29; 9.2, 9.4(3)]):

Lemma 5.11 (Habiro [32; 5.4(1,3)]). *G_k is a finitely generated nilpotent group.*

It is easy to see that the group ring $\mathbb{Z}G_k$ is isomorphic to the quotient $\mathbb{Z}\mathcal{L}/W_k$, where W_k is the two-sided ideal of the monoid ring $\mathbb{Z}\mathcal{L}$ generated by all elements of the form $L - L'$, where L is C_k -equivalent to L' .

On the other hand, let V_k be the two-sided ideal of $\mathbb{Z}\mathcal{L}$ generated by all singular links with k double points, where a singular link L with $k+1$ double points is identified with an alternating sum of links by the formula (V') .¹⁰ Thus a type k invariant on m -component string links with values in an abelian group A can be thought of as a homomorphism $\mathbb{Z}\mathcal{L} \rightarrow A$ which vanishes on V_{k+1} .

Proposition 5.6 implies that $W_k \subset V_k$ (actually it is obtained in [32] as a consequence of this inclusion). It is easy to see that $W_1 = V_1$. Moreover, W_1 coincides also with the augmentation ideal Δ of $\mathbb{Z}\mathcal{L}$, that is, the kernel of the augmentation homomorphism $\epsilon: \mathbb{Z}\mathcal{L} \rightarrow \mathbb{Z}$, which is given by $\epsilon(\mathcal{L}) = 1$. (Indeed, Δ is easily seen to be generated by

¹⁰Thus V_k is $J_k(D^2, m)$ in Habiro's notation; W_k is his $J_{k,1}(D^2, m)$ and \mathcal{L} is his $\mathcal{L}(D^2, m) = \mathcal{L}_1(D^2, m)$.

elements of the form $L_0 - L_1$, where $L_0, L_1 \in \mathcal{L}$, and if L_t is a generic homotopy from L_0 to L_1 , and L_{t_1}, \dots, L_{t_k} are its critical levels, which are singular links with one point each, then $L_0 - L_1 = \epsilon_1 L_{t_1} + \dots + \epsilon_k L_{t_k}$ for some signs $\epsilon_i \in \{1, -1\}$.) In the direction of the reverse inclusion “ $V_k \subset W_k$ ”, Habiro shows, in particular:

Lemma 5.12 (Habiro [32]). $V_{q(n-1)} \subset \Delta^q + W_n$.

Proof. This is a special case of [32; Proposition 6.10], which is seen by using that if $k_1 + \dots + k_l = q(n-1)$, then either $l \geq q$ or some $k_i \geq n$. \square

Let Δ_n be the augmentation ideal of $\mathbb{Z}G_n$. Since the projection $p_n: \mathbb{Z}\mathcal{L} \rightarrow \mathbb{Z}G_n$ commutes with the two augmentation homomorphisms, $p_n(\Delta) \subset \Delta_n$. Since $\ker p_n = W_n$, we get that $p_n(\Delta^l + W_n) \subset \Delta_n^l$ for all l . On the other hand, by Lemma 5.9 there exists a $q = q(m, n)$ such that $(1 - G_n) \cap \Delta_n^q = 0$. Therefore $(1 - \mathcal{L}) \cap (\Delta^q + W_n) \subset \ker p_n = W_n$. Since $V_{q(n-1)} \subset \Delta^q + W_n$, we get that $(1 - \mathcal{L}) \cap V_{q(n-1)} \subset (1 - \mathcal{L}) \cap (\Delta^q + W_n) \subset W_n$.

Let $r = q(n-1) - 1$. We may assume that $q, n \geq 2$, whence $r \geq n-1$. Suppose that $L, L' \in \mathcal{L}$ are not separated by invariants of type r . Thus $L - L' \in V_{r+1}$. Let $\bar{L} \in \mathcal{L}$ be a string link which is inverse to L modulo C_{r+1} -equivalence. Since V_{r+1} is a right ideal, $L\bar{L} - L'\bar{L} \in V_{r+1}$. On the other hand $L\bar{L}$ is C_{r+1} -equivalent to $1 \in \mathcal{L}$, and therefore $1 - L\bar{L} \in W_{r+1} \subset V_{r+1}$. Then $1 - L'\bar{L} \in V_{r+1}$. Consequently $1 - L'\bar{L} \in (1 - \mathcal{L}) \cap V_{r+1} \subset W_n$. Since W_n is a right ideal, $L - L'\bar{L}L \in W_n$. On the other hand $\bar{L}L$ is C_{r+1} -equivalent to $1 \in \mathcal{L}$, and therefore $\bar{L}L - 1 \in W_{r+1} \subset W_n$. Since W_n is a left ideal, $L'\bar{L}L - L' \in W_n$. Thus $L - L' \in W_n$, and so L is C_n -equivalent to L' . \square

Theorem 5.13. *There exists an $r = r(m, n)$ such that if two m -component string links are not separated by locally additive invariants of type r , then they are C_n -equivalent.*

Let us note that $L\#K = K\#L$ if K and L are m -component string links and K is totally split. We will call this property the *local commutativity* of string links. It is proved similarly to the usual proof that the monoid of knots is commutative (see [11; Figure 7.3] or [20; p. 140]).

Proof. This is an elaboration on the previous proof (the alternative proof of Theorem 5.8), additionally employing some ideas from [32; proof of Theorem 6.18]. We will use the notation of the previous proof, except that q and r will be chosen differently.

Let \mathcal{K} be the submonoid of \mathcal{L} consisting of all totally split m -component string links. Clearly, an invariant $v: \mathcal{L} \rightarrow A$ is locally additive if and only if its additive extension $\bar{v}: \mathbb{Z}\mathcal{L} \rightarrow A$ vanishes on all elements of the form $L\#K - L - K$, where $L \in \mathcal{L}$ and $K \in \mathcal{K}$. This is equivalent to saying that \bar{v} vanishes on $1 \in \mathcal{L}$ and on all elements of the form $(1 - L)(1 - K)$, where $L \in \mathcal{L}$ and $K \in \mathcal{K}$. Since elements of the form $1 - L$, where $L \in \mathcal{L}$, additively generate the augmentation ideal Δ of $\mathbb{Z}\mathcal{L}$, and elements of the form $1 - K$, where $K \in \mathcal{K}$, additively generate the augmentation ideal Λ of $\mathbb{Z}\mathcal{K}$, elements of the form $(1 - L)(1 - K)$ additively generate $\Delta\Lambda$. Thus v is locally additive if and only if \bar{v} vanishes on $1 \in \Lambda$ and on $\Delta\Lambda$. Let us note that Λ is not an ideal of $\mathbb{Z}\mathcal{L}$, but $\Delta\Lambda$ is a two-sided ideal of $\mathbb{Z}\mathcal{L}$. (Indeed, since Δ a left ideal, so is $\Delta\Lambda$; similarly, $\Lambda\Delta$ is a right ideal, but $\Lambda\Delta = \Delta\Lambda$ due to the local commutativity of string links.)

Let H_n be the image of \mathcal{K} in G_n , and let Λ_n be the augmentation ideal of $\mathbb{Z}H_n$. It equals $p_n(\Lambda)$ since both are additively generated by elements of the form $1 - h$, where $h \in H_n$. It follows that $p_n(\Delta\Lambda) = \Delta_n\Lambda_n$, and consequently $\mathbb{Z}\mathcal{L}/(W_n + \Delta\Lambda) = \mathbb{Z}G_n/(\Delta_n\Lambda_n)$. Our next goal is to get an explicit description of this quotient ring.

By the local commutativity H_n lies in the center of G_n , and in particular it is a normal subgroup. On the other hand, the totally split string link K_L corresponding to a given string link L may be described so as to be well-defined not just up to ambient isotopy, but as a specific string link. Consequently a self- C_n -move on L induces a self- C_n -move on K_L ; whereas a C_n -move on L which is not a self- C_n -move induces an ambient isotopy on K_L (see Remark 5.3). Thus the C_n -equivalence class of K_L is determined by the C_n -equivalence class of L . This yields an epimorphism $G_n \rightarrow H_n$, which restricts to the identity on H_n . Its kernel Q_n is a normal subgroup of G_n such that $Q_n H_n = G_n$ and $Q_n \cap H_n = 1$. Hence $G_n = Q_n \times H_n$.

Let R_n be the ring whose additive group is $(\mathbb{Z}Q_n) \oplus H_n$, with multiplication given by $(r_1, h_1) \cdot (r_2, h_2) = (r_1 r_2, h_1^{\epsilon(r_2)} h_2^{\epsilon(r_1)})$. (Here H_n is written multiplicatively, even though it is abelian. It is straightforward to verify that the multiplication is associative and distributive with respect to the addition. Clearly, the multiplicative identity is $(1, 1)$.) A homomorphism of the additive groups $\varphi_n: \mathbb{Z}G_n \rightarrow R_n$ is given on the additive generators by $\varphi_n(qh) = (q, h)$ for any $q \in Q_n$ and $h \in H_n$ and is easily seen to be a ring homomorphism (it suffices to check its multiplicativity on the additive generators). It can be described in general by $\varphi_n(m_1 q_1 h_1 + \cdots + m_k q_k h_k) = (m_1 q_1 + \cdots + m_k q_k, h_1^{m_1} \cdots h_k^{m_k})$, where $h_i \in H_n$, $q_i \in Q_n$ and $m_i \in \mathbb{Z}$.

It is easy to see that $\ker \varphi_n$ contains all elements of the form $qh'h - qh' - h + 1$ where $q \in Q_n$ and $h', h \in H_n$. These are the same as elements of the form $(g - 1)(h - 1)$, where $g \in G_n$ and $h \in H_n$. Hence $\ker \varphi_n$ contains $\Delta_n\Lambda_n$. To prove the reverse inclusion it suffices to show that the quotient map $\mathbb{Z}G_n \rightarrow \mathbb{Z}G_n/(\Delta_n\Lambda_n)$ factors through φ_n . The relations $(h - 1)(h' - 1) = 0$ for $h, h' \in H_n$ can be rewritten as $h + h' = hh' + 1$ and imply $m_1 h_1 + \cdots + m_k h_k = h_1^{m_1} \cdots h_k^{m_k} + (m_1 + \cdots + m_k - 1)$ for $h_i \in H_n$ and $m_i \in \mathbb{Z}$. The relations $(q - 1)(h - 1) = 0$ for $q \in Q_n$ and $h \in H_n$ can be rewritten as $qh = q + h - 1$ and imply $m_1 q_1 h_1 + \cdots + m_k q_k h_k = m_1 q_1 + \cdots + m_k q_k + m_1 h_1 + \cdots + m_k h_k - (m_1 + \cdots + m_k) = m_1 q_1 + \cdots + m_k q_k - 1 + h_1^{m_1} \cdots h_k^{m_k}$ for $q_i \in Q_n$, $h_i \in H_n$ and $m_i \in \mathbb{Z}$. In particular, we get $(m_1 q_1 + \cdots + m_k q_k)h = m_1 q_1 + \cdots + m_k q_k - 1 + h^{m_1 + \cdots + m_k}$, or $rh = r - 1 + h^{\epsilon(r)}$ for $r = m_1 q_1 + \cdots + m_k q_k \in \mathbb{Z}Q_n$ and $h \in H_n$. Hence given $r_1, r_2 \in \mathbb{Z}Q_n$ and $h_1, h_2 \in H_n$, we obtain $(r_1 - 1 + h_1)(r_2 - 1 + h_2) = (r_1 - 1)(r_2 - 1) + r_1 + r_2 - 2 + h_1^{\epsilon(r_2)} + h_2^{\epsilon(r_1)} - h_1 - h_2 + h_1 h_2 = r_1 r_2 - 1 + h_1^{\epsilon(r_2)} h_2^{\epsilon(r_1)}$. If we introduce the notation $(r, h) = r - 1 + h$ for $r \in \mathbb{Z}Q_n$ and $h \in H_n$, this becomes $(r_1, h_1) \cdot (r_2, h_2) = (r_1 r_2, h_1^{\epsilon(r_2)} h_2^{\epsilon(r_1)})$. This shows that the ring R_n is a quotient of $\mathbb{Z}G_n$ by an ideal contained in $\Delta_n\Lambda_n$, with $\varphi_n: \mathbb{Z}G_n \rightarrow R_n$ being the quotient map.

Thus we have proved that $\ker \varphi_n = \Delta_n\Lambda_n$, and hence the kernel of the composition $\mathbb{Z}\mathcal{L} \xrightarrow{p_n} \mathbb{Z}G_n \xrightarrow{\varphi_n} R_n$ equals $W_n + \Delta\Lambda$. Let Γ_n be the augmentation ideal of $\mathbb{Z}Q_n$. Using the identity $qh - 1 = (q - 1) + (h - 1) + (q - 1)(h - 1)$ it is easy to see that $\Delta_n^l \subset \Gamma_n^l + \Delta_n\Lambda_n$

for $l \geq 2$. Therefore $\varphi_n(\Delta_n^l) \subset \Gamma_n^l$ for $l \geq 2$. Hence $\varphi_n p_n(\Delta^l + W_n + \Delta\Lambda) = \varphi_n p_n(\Delta^l) \subset \Gamma_n^l$ for $l \geq 2$. By Lemma 5.11 Q_n is nilpotent (as a subgroup of G_n) and finitely generated (as a quotient of G_n); hence by Lemma 5.9 there exists a $q = q(m, n) \geq 2$ such that $(1 - Q_n) \cap \Gamma_n^q = 0$. This implies that $\varphi_n(1 - G_n) \cap \Gamma_n^q = 0$. (Indeed, given any $q \in Q_n$ and $h \in H_n$, if $\varphi_n(1 - qh) = (1 - q, 1/h)$ lies in $\Gamma_n^q \subset \mathbb{Z}Q_n$, then it equals $1 - q$ and hence lies in $1 - Q_n$.) Therefore $(1 - \mathcal{L}) \cap (\Delta^q + W_n + \Delta\Lambda) \subset \ker(\varphi_n p_n) = W_n + \Delta\Lambda$. On the other hand, by Lemma 5.12 $V_{q(n-1)} \subset \Delta^q + W_n$. Hence $(1 - \mathcal{L}) \cap (V_{q(n-1)} + \Delta\Lambda) \subset (1 - \mathcal{L}) \cap (\Delta^q + W_n + \Delta\Lambda) \subset W_n + \Delta\Lambda$.

Let $r = q(n - 1) - 1$. Suppose that all locally additive type r invariants take the same values on $L, L' \in \mathcal{L}$. Then $L - L'$ lies in the subgroup of the additive group of $\mathbb{Z}\mathcal{L}$ generated by 1 and by the ideal $V_{r+1} + \Delta\Lambda$. Since ϵ vanishes on $L - L'$ and on $V_{r+1} + \Delta\Lambda$, but not on 1, it follows that $L - L' \in V_{r+1} + \Delta\Lambda$. Arguing like in the previous proof, but now using that $(1 - \mathcal{L}) \cap (V_{r+1} + \Delta\Lambda) \subset W_n + \Delta\Lambda$, we obtain that $L - L' \in W_n + \Delta\Lambda$. Then $\varphi_n p_n(L) = \varphi_n p_n(L')$. But φ_n is injective on G_n (by the definition). Hence $p_n(L) = p_n(L')$. Therefore L is C_n -equivalent to L' . \square

Proof of Theorem 5.2. By Proposition 5.6 and Theorem 5.13. \square

6. PROOF OF THEOREM A

It does not seem to be easy to prove (or disprove) that for each n there exists an $r = r(n)$ such that if two links in S^3 are (strongly) r -quasi-isotopic, then they are ambient isotopic to the closures of some (strongly) n -quasi-isotopic string links¹¹ (compare Example 8.1). But it is easy to prove a slightly weaker assertion.

We say two links in S^3 are *(strongly) n -quasi-isotopic via string links* if they are equivalent with respect to the equivalence relation generated by ambient isotopy and the relation “to be closures of (strongly) n -quasi-isotopic string links”.

Proposition 6.1. (a) *If two links in S^3 are n -quasi-isotopic, $n \geq 1$, then they are $(n - 1)$ -quasi-isotopic via string links.*

(b) *If two links in S^3 are strongly n -quasi-isotopic, then they are strongly n -quasi-isotopic via string links.*

Proof. (a). Let $L, L': mS^1 \rightarrow S^3$ be the given links, related by an n -quasi-isotopy. We may assume that the n -quasi-isotopy has only one double point. Let $f: mS^1 \rightarrow S^3$ be the corresponding n -quasi-embedding. It suffices to show that f is ambient isotopic to the closure of an $(n - 1)$ -quasi-embedding $mI \rightarrow I^3$ of the string link boundary pattern. Let $P_0 \subset \cdots \subset P_n \subset S^3$ and $J_0 \subset \cdots \subset J_n \subset mS^1$ be given by the definition an n -quasi-embedding. Since the inclusion $P_{n-1} \rightarrow P_n$ is null-homotopic, it induces zero homomorphisms on the reduced cohomology groups. Hence by the Alexander duality the inclusion $S^3 \setminus P_n \rightarrow S^3 \setminus P_{n-1}$ induces zero homomorphisms on the reduced homology groups. This implies in particular that any two points in $S^3 \setminus P_n$ are connected by an arc in $S^3 \setminus P_{n-1}$. Let us recall that $f^{-1}(P_n)$ lies in the arc J_n . Hence each component of

¹¹I am indebted to M. Il'insky for pointing out a gap in what I thought was a proof of this assertion.

$f(mS^1)$ contains some point which lies in $S^3 \setminus P_n$. Let $f(x_i)$ be such a point in the i th component of $f(mS^1)$. Since $S^3 \setminus P_{n-1}$ is a 3-manifold, it contains a tree T with leaves at the points $f(x_i)$. Since $f(mS^1)$ is a 1-manifold, T may be assumed to meet it only in the leaves. Let B be a regular neighborhood of T , disjoint from $P_{n-1} \cup f(J_{n-1})$ and meeting each component of $f(mS^1)$ in an arc. Then $Q := \overline{S^3 \setminus B}$ is a ball containing $P_{n-1} \cup f(J_{n-1})$ and such that $f^{-1}(Q)$ meets each component of mS^1 in an arc. Hence f is ambient isotopic to the closure of an $(n-1)$ -quasi-embedding $mI \rightarrow I^3$ of the string link boundary pattern. \square

(b). This is similar to the proof of (a), if the following observation is used. Given a strong n -quasi-embedding $f: mS^1 \rightarrow S^3$ and the balls $B_1 \subset \cdots \subset B_n \subset S^3$ given by the definition a strong n -quasi-embedding, $S^3 \setminus B_n$ is connected (since it is an open ball). \square

Remark 6.2. As observed by J. Levine [45; Proposition 6 (stated slightly differently)], given a link L in S^3 , the choice of a string link whose closure is ambient isotopic to L is equivalent to the choice of a PL 2-disk D in S^3 which meets every component K_i of L transversely in a single point p_i . Let us note that D deformation retracts onto a tree T with leaves p_1, \dots, p_n and with one vertex of valency m .

Lemma 6.3. *Let A and B be m -component links in S^3 or string links. Suppose that $K_A = K_B$ or more generally K_A and K_B are not separated by locally additive type n invariants. Then A and B are not separated by type n invariants that are well-defined up to PL isotopy if and only if A and B are not separated by locally additive type n invariants.*

Proof. Similarly to the proof of Lemma 5.1. \square

Lemma 6.4. *Let A and B be m -component links in S^3 , and let $A' = A \# K_B$ and $B' = B \# K_A$ (which are PL isotopic to A and B respectively). Then A and B are not separated by type n invariants of string links that are well-defined up to PL isotopy if and only if A' and B' are not separated by locally additive type n invariants of string links.*

Proof. “If”. Since every invariant of PL isotopy is locally additive, the hypothesis implies that A' and B' are not separated by type n invariants of string links that are well-defined up to PL isotopy. On the other hand, let α and β be some string links whose closures are A and B , respectively. Then α is PL isotopic to $\alpha' := \alpha \# K_\beta$, whose closure is A' . Hence A and A' are not separated by type n invariants of string links that are well-defined up to PL isotopy. Similarly for B' and B .

“Only if”. We are given string links L_1, \dots, L_{2k} such that each L_{2i-1} is not separated from L_{2i} by type n invariants, well-defined up to PL isotopy; the closures of L_{2i} and L_{2i+1} are ambient isotopic whenever both are defined; and the closures of L_1, L_{2k} are ambient isotopic to A and B , respectively. Let us note that $K_{L_{2i}} = K_{L_{2i+1}}$ since these two string links are totally split and their closures are ambient isotopic. Let $\Lambda_1 = K_{L_{2k}}$ and $\Lambda_{2k} = K_{L_1}$. Then the closures of $L_1 \# \Lambda_1$ and $L_{2k} \# \Lambda_{2k}$ are ambient isotopic

to A' and B' , respectively. For $i = 1, \dots, k-1$ let Λ_{2i} be a totally split string link representing $[K_{L_{2i}}]^{-1}[K_{L_1} \# K_{L_{2k}}]$ in the direct product of m copies of Gusarov's group of C_{n+1} -equivalence classes of knots [28] (see also [32]), and let $\Lambda_{2i+1} = \Lambda_{2i}$. Then the closures of $L_{2i} \# \Lambda_{2i}$ and $L_{2i+1} \# \Lambda_{2i+1}$ are ambient isotopic whenever both are defined, and $K_{L_{2i}} \# \Lambda_{2i}$ and $K_{L_{2i+1}} \# \Lambda_{2i+1}$ are ambient isotopic themselves. Also $K_{L_i} \# \Lambda_i$ is C_{n+1} -equivalent to $K_{L_1} \# K_{L_{2k}}$, and hence (see Proposition 5.6) is not separated from it by type n invariants. Then by Lemma 6.3 each $L_{2i-1} \# \Lambda_{2i-1}$ is not separated from $L_{2i} \# \Lambda_{2i}$ by locally additive type n invariants. Writing $L'_i = L_i \# \Lambda_i$, the sequence of string links L'_1, \dots, L'_{2k} shows that A' and B' are not separated by locally additive type n invariants of string links. \square

Proof of Theorem A. The previous results can be seen to yield the implications (1)–(9) shown in Figure 6 for every links A, B in S^3 , where $A' = A \# K_B$ and $B' = B \# K_A$.

In more detail, the implication (1) holds by Theorem 3.5; (2) is obvious; (3) holds by Proposition 6.1; (4) holds by Theorem 4.8 (the case of string links); and (5) is obvious. The equivalence (6) holds by Lemma 6.4; and (7) follows from Theorem 5.2. The implication (8) holds by Lemma 4.1; and (9) is obvious. \square




FIGURE 6. Some implications, where $A' = A \# K_B$ and $B' = B \# K_A$

7. PROOF OF THEOREM D

Theorem 7.1 (Rolfsen [74]). *If L and L' are links in S^3 which are PL isotopic, and $K_L = K_{L'}$, then L and L' are ambient isotopic.*

An alternative proof of Theorem 7.1 will be discussed in §8. Let us note that the assertion of Theorem 7.1 fails for links in $S^1 \times S^2$ [75; Example 2].

Lemma 7.2. *Let A and B be m -component links in S^3 and K be a totally split m -component link in S^3 . Then A and B are not separated by locally additive type n invariants of string links if and only if $A\#K$ and $B\#K$ are not separated by locally additive type n invariants of string links.*

Proof. “Only if”. We are given string links L_1, \dots, L_{2k} such that the closures of L_1, L_{2k} are ambient isotopic to A and B , respectively; the closures of L_{2i} and L_{2i+1} are ambient isotopic whenever both are defined; and each L_{2i-1} is not separated from L_{2i} by locally additive type n invariants. Let Λ be a totally split string link whose closure is ambient isotopic to K , and let $L'_i = L_i\#\Lambda$. Then the sequence of string links L'_1, \dots, L'_{2k} shows that $A\#K$ and $B\#K$ are not separated by locally additive type n invariants of string links (it is here that we need the local additivity).

“If”. Let K' be a totally split string link representing $[K]^{-1}$ in the direct product of m copies of Gusarov’s group of C_{n+1} -equivalence classes of string knots [28] (see also [32]). Then by the “only if” part $A\#K\#K'$ and $B\#K\#K'$ are not separated by locally additive type n invariants of string links. Let L be any string link whose closure is ambient isotopic to A . Let Λ and Λ' be totally split string links whose closures are ambient isotopic to K and K' , respectively. Then the closure of $L\#\Lambda\#\Lambda'$ is ambient isotopic to $A\#K\#K'$. On the other hand, $\Lambda\#\Lambda'$ is C_{n+1} -equivalent to the string unlink. Hence $L\#\Lambda\#\Lambda'$ is C_{n+1} -equivalent to L , and consequently (see Proposition 5.6) they are not separated by type n invariants. Thus A and $A\#K\#K'$ are not separated by type n invariants of string links. Similarly, $B\#K\#K'$ and B are not separated by type n invariants of string links. \square

Proof of Theorem D. The implications (1)–(9) in Figure 6 were established in the proof of Theorem A, whereas the implication (0) is obvious.

Let (X) denote the converse to the composite of the implications (0)–(9) (i.e. “if A and B are PL isotopic to links which are not separated by finite type invariants, then A and B are PL isotopic”). Then (X) clearly follows from (L), and by Theorem 7.1, (X) \wedge (K) implies (L).

Let (Y) denote the converse to the composite of the implications (8) and (9) (i.e. “if A and B are PL isotopic to links that are not separated by finite type invariants, then for each n , A' and B' are not separated by type n invariants of string links”).

Let us show that (Y) \wedge (K) implies (HM). Suppose that A and B are not separated by finite type invariants. Then by (K) $K_A = K_B$, and by (Y) for each n , A' and B' are not separated by type n invariants of string links. We have $A' = A\#K$ and $B' = B\#K$, where $K = K_A = K_B$, so by Lemma 7.2 for each n , A and B are not separated by locally

additive type n invariants of string links. Then by Theorem 5.2 for each n , A and B are not separated by type n invariants of string links.

Let us show that $(\text{HM}) \wedge (\text{K})$ implies (Y). Suppose that A, B are PL isotopic to links \bar{A}, \bar{B} (respectively), which are not separated by finite type invariants. Then by (K) $K_{\bar{A}} = K_{\bar{B}}$, and by (HM) for each n , \bar{A} and \bar{B} are not separated by type n invariants of string links. The link A' is PL isotopic to A and hence to \bar{A} , and similarly B' is PL isotopic to \bar{B} . Therefore $\bar{A} \# Q$ is ambient isotopic to $A' \# Q'$ and $\bar{B} \# R$ is ambient isotopic to $B' \# R'$ for some totally split links Q, Q', R and R' . Then $\bar{A} \# Q \# R$ is ambient isotopic to $A' \# Q' \# R$ and $\bar{B} \# Q \# R$ is ambient isotopic to $B' \# Q' \# R'$. Since $K_{\bar{A}} = K_{\bar{B}}$, we have $K_{\bar{A}} \# Q \# R = K_{\bar{B}} \# Q \# R$, and therefore also $K_{A'} \# Q' \# R = K_{B'} \# Q' \# R'$. But on the other hand, we have $K_{A'} = K_A \# K_B = K_{B'}$. Hence by the uniqueness of factorization into prime knots, $Q' \# R = Q \# R'$. Thus, writing $S = Q \# R$ and $S' = Q' \# R' = Q \# R'$, we get that $\bar{A} \# S = A' \# S'$ and $\bar{B} \# S = B' \# S'$. Since for each n , \bar{A} and \bar{B} are not separated by type n invariants of string links, it follows by a double application of Lemma 7.2 that for each n , A' and B' are not separated by additive type n invariants of string links. Then by Theorem 5.2 for each n , A' and B' are not separated by type n invariants of string links.

Let (Z) denote the converse to the composite of the implications (0)–(5) (i.e. “if A and B can be represented for each n as closures of string links that are not separated by type n invariants, well-defined up to PL isotopy, then A and B are PL isotopic”). Then we get that $(\text{L}) \Leftrightarrow (\text{K}) \wedge (\text{X}) \Leftrightarrow (\text{K}) \wedge (\text{Y}) \wedge (\text{Z}) \Leftrightarrow (\text{K}) \wedge (\text{HM}) \wedge (\text{Z})$. \square

8. GEOMETRIC FACTORIZATIONS

In order to prove part (a) of Theorem C, we will need the string link version of Rolfsen’s Theorem 7.1. The following example indicates that passing from links in S^3 to string links is far from being automatic for such purposes.

A tangle $L: \Theta \rightarrow M$ is said to *have a local knot* if there exists a 3-ball B in M which meets $L(\Theta)$ in an arc such that the union of this arc with an arc in ∂B which has the same endpoints is a nontrivial knot.




FIGURE 7. A string link with no local knots whose closure has a local knot.

Example 8.1. There exists a 3-component string link L such that L has no local knots, but its closure has a local knot. See Figure 7, where L is shown in the middle and its closure on the right.

Indeed, it is easy to see that the closure of L has a local knot. To show that L itself has no local knots, we consider a non-standard two-component closure \bar{L} of L , shown on the left in Figure 7. If L has a local knot, then so does \bar{L} . Moreover, if \bar{L} has a local knot, then this local knot must be one of the prime factors of its components. But the only prime factor of the components of \bar{L} is the trefoil knot. Thus it suffices to show that \bar{L} has no local knot which is the trefoil.

The Jones polynomial of \bar{L} is $V_L(t) = t^{-7/2} - t^{-5/2} - t^{-1/2} - t^{1/2} - t^{5/2} + t^{7/2}$. The Jones polynomial of the trefoil knot T is $V_T(t) = t + t^3 - t^4$. If T is a local knot in \bar{L} , then $V_L(t)$ must be divisible by $V_T(t)$ in $\mathbb{Z}[t^{\pm 1/2}]$ (see [47; p. 29]). But it is easy to see that $V_L(t) = t^{-7/2}(t^4(t^3 - t^2 - 1) - (t^3 + t - 1))$ is not divisible by $V_T(t) = (-t)(t^3 - t^2 - 1)$.

Remark 8.2. Let us note that the non-locality of the trefoil in \bar{L} is not detected by the Conway polynomial and the Conway potential function. In fact their values on a possibly twisted Whitehead double of a two-component link (K, K') along K depend only on $\text{lk}(K, K')$ and on the knot K' (see [59; Corollary 7.10]).

Theorem 8.3. (a) *Every link in S^3 admits a unique decomposition into a connected sum of a link which has no local knots and a totally split link.*

(b) *Every string link admits a unique decomposition into a connected sum of a string link which has no local knots and a totally split string link.*

Example 8.1 suggests that (b) is not a consequence of (a), but has to be proved independently. In fact, (a) is an easy consequence of Hashizume's 1958 theorem on uniqueness of factorization of links into prime links (see Theorem 8.9). Our proof of (b) is a variation on a new, comparatively short proof of Hashizume's theorem, which we include below.

Corollary 8.4. *If L and L' are links in S^3 or string links which are PL isotopic, and $K_L = K_{L'}$, then L and L' are ambient isotopic.*

The case of links in S^3 is the same as Rolfsen's Theorem 7.1, but now it comes with an alternative proof.

Proof. Theorem 8.3 yields a decomposition $L = \Gamma(L) \# \Lambda(L)$ up to ambient isotopy, where $\Gamma(L)$ has no local knots and $\Lambda(L)$ is totally split. It is easy to see that $\Gamma(L)$ is invariant under adding local knots to L . Therefore it is invariant under PL isotopy.

Since L and L' are PL isotopic, $\Gamma(L) = \Gamma(L')$. In particular, $K_{\Gamma(L)} = K_{\Gamma(L')}$. But clearly $K_L = K_{\Gamma(L)} \# \Lambda(L)$, and similarly for L' . Since $K_L = K_{L'}$, we obtain (using the uniqueness of the decomposition of a knot into prime factors) that $\Lambda(L) = \Lambda(L')$. Since L represents the ambient isotopy type of $\Gamma(L) \# \Lambda(L)$ and similarly for L' , we conclude that L is ambient isotopic to L' . \square

A link L in S^3 is called *split* if there exists a PL 3-ball $B \subset S^3$ such that L is disjoint from ∂B but meets both B and $S^3 \setminus B$. A string link $L: mI \rightarrow I^3$ is called *split* if there exists a properly embedded PL 2-disk $D \subset I^3$ such that L is disjoint from D but meets both components of $I^3 \setminus D$.

Proposition 8.5. (a) *For every link L in S^3 there exists a family of pairwise disjoint PL 3-balls in S^3 such that L lies in their union and meets each of the balls in a non-split sublink.*

(b) *For every string link L there exists a family of pairwise disjoint properly embedded PL 2-disks in I^3 such that L is disjoint from their union Δ and meets each component of $I^3 \setminus \Delta$ in a non-split sublink.*

Proof. (a). Let B_0 be some PL ball containing L . If L is non-split, this completes the proof.

Lemma 8.6. *If L is split and lies in a PL ball B_0 , then there exist disjoint PL balls B_+ and B_- in B_0 such that L lies in their union and meets both of them.*

Proof. Since L is split, exists a PL ball $B \subset S^3$ such that L is disjoint from ∂B but meets both B and $S^3 \setminus B$. Let B_1 be a PL ball containing both B_0 and B in its interior. By the PL annulus theorem there exists a PL homeomorphism $B_1 \rightarrow B_0$ keeping L fixed. It sends B onto a PL ball B_+ lying in the interior of B_0 . Since L is a 1-manifold, there exists an arc J in $\overline{B_0} \setminus \overline{B_+} \setminus L$ joining ∂B_+ to ∂B_0 . Let N be a regular neighborhood of $B_+ \cup J$ in B_0 , disjoint from L , and let $B_- = \overline{B_0} \setminus N$. \square

Let us note that each of $L \cap B_+$ and $L \cap B_-$ has fewer components than L . Hence by repeatedly applying Lemma 8.6 we eventually obtain a desired family of balls. \square

(b). The assertion follows similarly to (a) from the following lemma.

Lemma 8.7. *If L is split and lies in one component of $I^3 \setminus D_0$ for some properly embedded PL 2-disk $D_0 \subset I^3$, then there exists a properly embedded PL 2-disk $D \subset I^3$, disjoint from D_0 and such that L is disjoint from D but meets both components of $I^3 \setminus D$.*

Proof. Since L is split, there exists a properly embedded PL 2-disk $D' \subset I^3$ such that L is disjoint from D' but meets both components of $I^3 \setminus D'$. On the other hand, let B_0 be the closure of the component of $I^3 \setminus D_0$ containing L . By Alexander's Schoenflies theorem the 2-sphere ∂B_0 is unknotted in S^3 , and hence B_0 is a PL 3-ball. Then there is a PL homeomorphism $h: B_0 \rightarrow I^3$ which is the identity outside a small neighborhood of D_0 in B_0 . In particular, this neighborhood can be chosen to be disjoint from L . We may further assume that $h(D_0)$ is a small disk, which is disjoint from D' . Then $D := h^{-1}(D')$ is disjoint from D_0 and retains the properties of D . \square

\square

By a *punctured 3-sphere* we shall mean the 3-manifold $P_n := \overline{S^3 \setminus (B_1 \cup \dots \cup B_n)}$ for some $n \geq 0$, where B_1, \dots, B_n are pairwise disjoint PL balls. A *beaded link*, or more

specifically an n -beaded link, is a tangle $L: \Theta \rightarrow P_n$ of a boundary pattern $\Xi: \partial(nI) \rightarrow \partial P_n$ that sends each $\{k\} \times \partial I$ into ∂B_k . The *closure* of a beaded link L is the link in S^3 obtained by gluing L with any embedding $\bar{\Xi}: nI \rightarrow \partial P_n$ that sends each $\{k\} \times I$ into ∂B_k . It is easy to see that the choice of a particular embedding $\bar{\Xi}$ is irrelevant in the sense that the closure is well-defined up to ambient isotopy. We call a beaded link a *beaded (un)knot* if its closure is a knot (respectively, an unknot). The *genus* $g(L)$ of a beaded link L is the genus of its closure \bar{L} , that is, the minimal genus of a *connected* Seifert surface spanned by the link \bar{L} .

Given an n -beaded link $L: \Theta \rightarrow P_n$ and a PL 2-sphere Σ in the interior of P_n which meets $L(\Theta)$ transversely in two points, by Alexander's Schoenflies theorem it bounds two complementary 3-balls in S^3 and hence partitions L into an $(i+1)$ -beaded link L' and an $(n-i+1)$ -beaded link L'' for some $i \leq n$. When $n=0$, the link L can be easily reconstructed from the (oriented) 1-beaded links L' and L'' , and in this case we write $L = L' \# L''$. On the other hand, given a link $L: mS^1 \rightarrow S^3$ and an index $i \in \{1, \dots, m\}$, we can represent L as the closure of a 1-beaded link $L[i]: (mS^1 \setminus \{i\} \times S^1) \cup I \rightarrow B^3$, which is easily seen to be well-defined up to ambient isotopy. Given another link $L': \tilde{m}S^1 \rightarrow S^3$ and another index $j \in \{1, \dots, \tilde{m}\}$, we get the link $L[i] \# L'[j]: (m+m'-1)S^1 \rightarrow S^3$, which is known as the Hashizume connected sum¹² $L \#_{i,j} L'$ of L and L' along the selected components [35]. This yields a well-defined operation on pairs $([L], i)$, where $[L]$ is the ambient isotopy type of the link L in S^3 , and i is a selected component of L . It is easy to see that this operation is associative and that the unknot is its unit.

If K and K' are knots, then $K \#_{1,1} K'$ is their usual connected sum $K \# K'$. It is well-known that connected sum of knots is commutative, and the same proof (see [11; Figure 7.3] or [20; p. 140]) works for the Hashizume connected sum. It is also well-known that $g(K \# K') = g(K) + g(K')$, and the 13-line proof of this fact in [20; p. 141] works verbatim to show that $g(L \#_{i,j} L') = g(L) + g(L')$ for any links L, L' in S^3 and any i, j . This in turn implies that whenever a PL 2-sphere Σ in the interior of P_n meets an n -beaded link L transversely in two points and hence partitions it into beaded links L' and L'' , we have $g(L) = g(L') + g(L'')$. Using this, it is easy to prove the non-cancellation property:

Lemma 8.8. *Let L be an n -beaded link, and suppose that the interior of P_n contains a PL 2-sphere Σ which meets L transversely in two points and partitions it into beaded links L_1 and L_2 so that L_2 is not a beaded unknot. Then L_1 has either fewer components than L or genus smaller than that of L . In particular, L is not a beaded unknot.*

Proof. If L_1 has the same number of components as L , then L_2 is a beaded knot. But L_2 is not a beaded unknot, so $g(L_2) \geq 1$. Then $g(L) = g(L_1) + g(L_2)$ implies $g(L_1) < g(L)$. \square

¹²The operation of Hashizume connected sum should not be confused with the componentwise connected sum, which was used in the previous sections and will be used in the following sections. In this connection see also [9], where a theorem of uniqueness of decomposition into prime factors is proved for the componentwise connected sum of 2-component string links.

A link L in S^3 is called *prime* if it is non-split and not an unknot, but for every representation of L as $A\#_{i,j}B$, either A or B is the unknot. A link in S^3 is *composite* if it is non-split, not an unknot and not a prime link. Thus a non-split link L is composite if and only if S^3 contains a PL 2-sphere which meets L in two points and partitions it into two 1-beaded links L_1, L_2 neither of which is a beaded unknot. (The “only if” assertion uses the case $n = 0$ of Lemma 8.8.)

Theorem 8.9 (Hashizume [35]). *Every non-split link L in S^3 admits a unique decomposition into prime factors L_0, \dots, L_r (well-defined up to ambient isotopy and up to a permutation of the factors) with respect to the operation of Hashizume connected sum.*

Moreover, S^3 contains r pairwise disjoint 2-spheres, each meeting L transversely in two points, which partition L into $r + 1$ beaded links whose closures are L_0, \dots, L_r .

Hashizume’s theorem plays a fundamental role in knot theory as it is needed to make sense out of the knot tables, which traditionally list only prime links. Nevertheless I was unable to find its proof in any textbook, even though a number of textbooks include the proof of uniqueness of factorization of knots into prime knots. One potential reason is that textbook authors might be somewhat terrified by Hashizume’s original proof [35], which takes about 16 pages. (About a half of these 16 pages consist of pictures, which could be related to the fact that much of the proof proceeds by exhaustion of cases.) Our proof of Hashizume’s theorem aims at being a bit more conceptual (the idea to work with punctured spheres comes from [61; proof of Lemma 2.2]).

Alternative proof of Theorem 8.9. If L is prime, there is nothing to prove. Else we get the desired collection of r spheres by repeatedly applying the following lemma along with Lemma 8.8.

Lemma 8.10. *Let L be an n -beaded link whose closure is a composite link. Then the interior of P_n contains a PL 2-sphere Σ which meets L transversely in two points and partitions it into beaded links L_1, L_2 , neither of which is a beaded unknot.*

Proof. The case $n = 0$ holds by the definition of a composite link. Suppose that the lemma holds for $(n - 1)$ -beaded links. By adjoining to L an arc J in ∂B_n we obtain an $(n - 1)$ -beaded link \bar{L} in $P_{n-1} = P_n \cup B_n$. We may identify B_n with the cone over its boundary, $c * \partial B_n$. Then there is an ambient isotopy of P_{n-1} keeping L fixed, moving J along $c * J$ and taking it onto $c * \partial J$. Let \hat{L} be the $(n - 1)$ -beaded link in P_{n-1} obtained by adjoining $c * \partial J$ to L . Now by the induction hypothesis the interior of P_{n-1} contains a PL 2-sphere Σ' which meets \hat{L} transversely in two points and partitions it into two beaded links neither of which is a beaded unknot. We may assume that Σ' is disjoint from the cone point c . Let h be a self-homeomorphism of the pair (P_{n-1}, \hat{L}) with support in a small neighborhood of B_n , which squeezes B_n radially onto a small ball ϵB_n about c , disjoint from Σ' . Then $\Sigma := h^{-1}(\Sigma')$ lies in the interior of P_n , meets \hat{L} transversely in two points and partitions it into two beaded links neither of which is a beaded unknot. The same holds if we replace \hat{L} with \bar{L} , and consequently also if we replace it with L . \square

The uniqueness of the decomposition now follows from the following lemma.

Lemma 8.11. *Let L be a non-split link in S^3 and let Σ be the union of a collection of r pairwise disjoint PL 2-spheres Σ_i , each meeting L transversely in two points, such that the $r + 1$ beaded links cut out of L by Σ have prime closures. Let L/Σ denote the unordered collection of the ambient isotopy types of these closures.*

(a) *Let Σ' be another union of the same kind. Then there exists yet another union Σ'' of the same kind such that $\Sigma'' \cap \Sigma = \emptyset$ and $L/\Sigma'' = L/\Sigma'$.*

(b) $L/\Sigma'' = L/\Sigma$.

Proof. (a). We may assume that Σ intersects Σ' transversely in a closed 1-manifold M . The proof is by induction on the number of components of M . If M is empty, then we let $\Sigma'' = \Sigma'$. Suppose that M is non-empty. If we fix some orientations on mS^1 , on S^3 and on Σ , then each component of Σ meets L in two points of the opposite signs, and $\Sigma \cap L = \Delta_+ \cup \Delta_-$, where Δ_+ consists of the positive intersections and Δ_- of the negative ones. Now M contains a component C that is innermost in $\Sigma \setminus \Delta_+$ in the sense that C bounds a disk D in $\Sigma \setminus \Delta_+$ whose interior is disjoint from M . Thus the interior of D is also disjoint from Σ' . Let D_+ and D_- be the two disks bounded by C in Σ' . The 2-spheres $D \cup D_+$ and $D \cup D_-$ are unknotted in S^3 by Alexander's Schoenflies theorem, and hence bound in S^3 balls B_+ and B_- (respectively) such that $B_+ \cap B_- = D$ and $B_+ \cup B_-$ is a 3-ball bounded by the 2-sphere $D_+ \cup D_-$. Let B be the other 3-ball bounded in S^3 by the latter 2-sphere, and let H be a 2-handle attached to B along D . We may assume that H is thin enough to be disjoint from L . Then $\partial(B \cup H)$ consists of two disjoint 2-spheres S_+ and S_- , which lie in B_+ and B_- respectively.

If D is disjoint from Δ_- , then C has zero linking number with every component of L , and consequently both points of intersection between L and the 2-sphere $D_+ \cup D_-$ must lie in one of the disks D_+ and D_- , say in D_+ . Then the 2-sphere $D \cup D_-$ is disjoint from L . Since L is non-split, at least one of the 3-balls B_- and $B \cup B_+$ bounded by this 2-sphere must also be disjoint from L . But L definitely meets $D_+ = B \cap B_+$, so it must be disjoint from B_- . Then we define Σ'' as the modification of Σ' obtained by replacing the 2-sphere $D_+ \cup D_-$ with the 2-sphere S_+ . Then the intersection of Σ'' with Σ is a subset of $M \setminus C$, and hence has fewer components than M . Since L is disjoint from B_- , it is easy to see that $L/\Sigma'' = L/\Sigma'$.

If D meets Δ_- , then it meets L in one point, and consequently both D_+ and D_- must also meet the same component of L , each in one point (by considering the linking numbers). The closure of the component of $S^3 \setminus \Sigma'$ that contains the interior of D is a punctured 3-sphere P_n (for some $n \geq 1$) such that ∂P_n contains the 2-sphere $D_+ \cup D_-$. Now P_n intersects L in an n -beaded link Λ , which is in turn partitioned by D into an i -beaded link Λ_+ in the punctured 3-sphere $P_n \cap B_+$, and an $(n - i)$ -beaded link Λ_- in the punctured 3-sphere $P_n \cap B_-$, for some $i \leq n$. It is easy to see that the closure of Λ is a Hashizume connected sum of the closures of Λ_+ and Λ_- . Since the closure of Λ is prime, some Λ_ϵ must be a beaded unknot, say, Λ_- . Then we define Σ'' as the modification of Σ' obtained by replacing the 2-sphere $D_+ \cup D_-$ with the 2-sphere S_+ .

(It is easy to see that S_+ lies in $P_n \cap B_+$). The intersection of Σ'' with Σ is a subset of $M \setminus C$, and hence has fewer components than M . Since L meets $P_n \cap B_-$ in a beaded unknot, it is easy to see that $L/\Sigma'' = L/\Sigma'$. \square

(b). Let Q_0, \dots, Q_r be the closures of the components of $S^3 \setminus \Sigma$ and let R_0, \dots, R_s be those of $S^3 \setminus \Sigma''$. Each $P_{ij} := Q_i \cap R_j$ is either empty or a punctured 3-sphere (indeed, $S^3 \setminus (Q_i \cap R_j)$ equals $(S^3 \setminus Q_i) \cup (S^3 \setminus R_j)$, which is a union of balls whose boundaries are pairwise disjoint). Since each $R_j \cap R_k$ is either empty or a common boundary component, the same is true of each $P_{ij} \cap P_{ik}$; and similarly for each $P_{ij} \cap P_{kj}$. Let L_i and L'_j be the closures of the beaded links cut out of L by Q_i and R_j , respectively. Each P_{ij} intersects L either in \emptyset or in a beaded link, whose closure will be denoted Λ_{ij} . (The closure of \emptyset is formally set to be \emptyset .) Then each L_i can be described as a certain Hashizume connected sum of those links Λ_{ij} , $j = 1, \dots, s$, that are nonempty. Since L_i is prime, it follows by induction that there exists a j_i such that Λ_{ij} is either empty or the unknot for all $j \neq j_i$, whereas $\Lambda_{ij_i} = L_i$ (up to ambient isotopy). Similarly, for each j there exists an i_j such that Λ_{ij} is either empty or the unknot for all $i \neq i_j$, whereas $\Lambda_{i_j j} = L'_j$. Since L_i is not an unknot, $i_{j_i} = i$, and since L'_j is not an unknot, $j_{i_j} = j$. Thus $i \mapsto j_i$ is a bijection such that $L_i = L'_{j_i}$. \square

\square

Proof of Theorem 8.3. (a). For an m -component link L in S^3 let us consider its unique partition into non-split sublinks (Proposition 8.5(a)) and the unique decomposition of each non-split sublink into prime factors (Theorem 8.9). By discarding those prime factors that are knots we obtain an m -component link $\Gamma(L)$ (the ‘‘global part’’ of L); the discarded factors can also be put together to form a totally split m -component link $\Lambda(L)$ (the ‘‘local part’’ of L). Clearly, $\Gamma(L)$ has no local knots, and $L = \Gamma(L) \# \Lambda(L)$ (up to ambient isotopy). \square

(b). It suffices to show that every string link (not necessarily non-split) admits a unique decomposition into a connected sum of a string link which has no local knots and prime local knots (the latter being well-defined up to a permutation). This can be done along the lines of the proof of Theorem 8.9.

In more detail, we are given a string link $L: mI \rightarrow I^3$. The most important amendment in the proof is that all the punctured spheres will now be contained in the interior of I^3 , apart from the ‘‘outer’’ one, which lies in I^3 and contains ∂I^3 ; this outer punctured sphere will intersect L in a tangle with no local knots, and the ‘‘inner’’ ones in beaded knots whose closures are prime. If the genus of a string link is defined to be the genus of its closure, then the proof of the existence of a decomposition goes through without essential changes. The amended proof of Lemma 8.11(a) still has two cases: D is disjoint from Δ_- and D meets Δ_- . In the first case we use that, even though the given string L may be non-split, it contains no closed components. In the second case we now have to consider two subcases: P_n is an inner punctured sphere and P_n is the outer punctured sphere. The first subcase is treated like before. In the second subcase

one of the two balls B_+ , B_- lies in the interior of I^3 . If B_ϵ lies in the interior of I^3 , then since P_n intersects L in a tangle with no local knots, Λ_ϵ must be a beaded unknot. The remainder of the proof of the second subcase proceeds like before. The amended proof of Lemma 8.11(b) goes through with aid of the following observation: if Q_i is the outer punctured sphere and R_j is an inner punctured sphere, then Λ_{ij} must be either empty or the unknot, using that L_i is a tangle with no local knots. \square

9. PROOF OF THEOREM C

The following more precise version of Lemma 5.1 is proved by the same argument.

Lemma 9.1. *Let A and B be m -component links in S^3 or string links. Then A and B are not separated by type n invariants that are well-defined up to PL isotopy if and only if $A' := A\#K_B$ and $B' := B\#K_A$ are not separated by locally additive type n invariants.*

Let us reformulate part (b) of Theorem C in more detail:

Theorem 9.2. *Finite type invariants separate string links if and only if finite type invariants separate string knots (or equivalently knots in S^3) and finite type invariants, well-defined up to PL isotopy, separate PL isotopy classes of string links.*

Proof. By Lemma 9.1 and Theorem 5.2, the string link version of (L) implies the string link version of (L/K).

By Corollary 8.4 the conjunction of the string link versions of (K) and (L/K) implies the string link version of (L). \square

Remark 9.3. In order to prepare for the proof of part (a) of Theorem C let us review some basics of the Kontsevich integral. Since much of the literature on the Kontsevich integral focuses on the case of knots, we focus on references which cover the case of links (in the present remark, all knots and links are in S^3 , and all tangles are in I^3).

- A detailed exposition of M. Kontsevich’s original (analytical) definition for knots appears in [13]. A generalization of the Kontsevich integral to tangles, along with a proof of its invariance for tangles, is sketched in [42; §1]; when guided by this sketch, some details of the proof can be found in [7] and, when guided by [7], some further details can be found in [13] and [12].
- There is also a combinatorial version of the Kontsevich integral, first described by P. Cartier, Le–Murakami and S. Piunikhin (independently); its construction for “non-associative tangles”, which include links, along with a proof of invariance, can be found in [8] (concerning the definition see also [13]).
- That the two versions of the Kontsevich integral are equal for links is proved in [42] (see also [13]) and by a different method in [5].
- It is shown in [43] (see also [13]) that the combinatorial Kontsevich integral Z of links assumes only rational values. In more detail, let V_m be the vector space of all rational formal linear combinations of chord diagrams on m circles, let A_m be its quotient by the 4-term and the 1-term relations,¹³ which is graded by the

¹³See [13] concerning chord diagrams and the 4-term and the 1-term relations.

number of chords: $A_m = \bigoplus_{n=0}^{\infty} A_{mn}$, and let \hat{A}_m be the graded completion of A_m , that is, $\prod_{n=0}^{\infty} A_{mn}$. Then, as explained in [42], [43] and [31], $Z(L)$ takes values in \hat{A}_m , where m is the number of components of L .

- A proof sketch that the combinatorial Kontsevich integral is a universal rational¹⁴ finite type invariant of links appears in [8]. As noted in [43; §5] and [31; Remark 3.3], the proof in [7] that the analytic Kontsevich integral is a universal rational finite type invariant of knots (see also [13] concerning this proof) carries over to the case of links.
- As noted in [43; Theorem 5], the combinatorial Kontsevich integral, when suitably normalized (see [13; §8.7.2] concerning the normalization), is multiplicative with respect to the Hashizume connected sum of links and the similarly defined connected sum of chord diagrams along selected components: $Z(L\#_{i,j}L') = Z(L)\#_{i,j}Z(L')$.¹⁵ The same formula for the analytical Kontsevich integral can be seen to follow from the multiplicativity of the preliminary Kontsevich integral under composition of tangles (see [13; 8.4.3]).
- Since connected sum is single-valued for chord diagrams on one circle, A_1 is an algebra, and \hat{A}_1 is an algebra where every element of the form $1+(\text{terms of positive degrees})$ is invertible (cf. [13]). In particular, $Z(K)$ is invertible for every knot K .

Let us reformulate part (a) of Theorem C in more detail:

Theorem 9.4. *Rational finite type invariants separate links in S^3 if and only if rational finite type invariants separate knots in S^3 and rational finite type invariants, well-defined up to PL isotopy, separate PL isotopy classes of links in S^3 .*

Proof. Let $\bar{Z}(L) = \frac{Z(L)}{Z(K_1) \cdots Z(K_m)}$, where K_1, \dots, K_m are the components of L , and the fraction is understood more precisely as $(\cdots (Z(L)\#_{1,1}Z(K_1)^{-1}) \cdots)\#_{n,1}Z(K_1)^{-1}$. Then $\bar{Z}(L)$ is invariant under PL isotopy; also its coefficients are easily seen to be finite type invariants.

Let us show that the rational version of (L) implies the rational version of (L/K). Suppose that A and B are links in S^3 which are not separated by rational finite type invariants, well-defined up to PL isotopy. Then in particular $\bar{Z}(A) = \bar{Z}(B)$. Let $A' = A\#K_B$ and $B' = B\#K_A$. Then A' is PL isotopic to A , so $\bar{Z}(A') = \bar{Z}(A)$; and similarly $\bar{Z}(B') = \bar{Z}(B)$. On the other hand, $K_{A'}$ is ambient isotopic to $K_{B'}$, so the denominators of the equal fractions $\bar{Z}(A')$ and $\bar{Z}(B')$ are equal. Hence so are their numerators $Z(A')$

¹⁴By a rational invariant we mean an invariant with values in a vector space over \mathbb{Q} .

¹⁵In more detail, if d is a chord diagram on m circles and $i \in \{1, \dots, m\}$, then $d[i]$ denotes the chord diagram on $m-1$ circles and one arc, obtained by removing from the i th circle a small open arc disjoint from all chords. This $d[i]$ is well-defined (i.e. does not depend on the choice of the small open arc) due to the 4-term relation (see [13; proof of Lemma in §4.4.3]). Also, if d and d' are chord diagrams on oriented 1-manifolds Θ and Θ' with $\partial\Theta = \partial I = \partial\Theta'$, then $d\#d'$ denotes the chord diagram on the closed 1-manifold $\Theta \cup_{\partial\Theta=\partial\Theta'} \Theta'$ consisting of all chords of d and all chords of d' . Both operators extend linearly over the graded completions of spaces of chord diagrams. Finally, $d\#_{i,j}d'$ denotes $d[i]\#d'[j]$.

and $Z(B')$. Now by the universality of Z , the rational version of (L) implies that A' is ambient isotopic to B' . Hence A is PL isotopic to B .


By Corollary 8.4 the conjunction of the rational versions of (K) and (L/K) implies the rational version of (L). \square

10. REDUCED CONWAY POLYNOMIAL

The *Conway polynomial* ∇_L of a link L in S^3 is a normalized sign-refined version of the Alexander polynomial; it can be defined for instance in terms of Seifert matrices (see [47]). The Conway polynomial is characterized (see [47; Theorem 8.6]) by two axioms:

$$\begin{aligned} \nabla_{\text{unknot}} &= 1, \\ \nabla_{L_+}(z) - \nabla_{L_-}(z) &= z\nabla_{L_0}(z), \end{aligned} \tag{C}$$

where L_0 , L_+ and L_- agree outside a small ball, and inside this ball they are as follows:¹⁶



Implicit in this figure is the singular link L (see the figure related to formula (V)). Let us note that L_0 has one more component than L_+ and L_- if the intersection in L is a self-intersection of some component, and one less component if the intersection involves distinct components. We will refer to L_+ and L_- as the positive and negative *resolutions* of L , and L_0 as the *smoothing* of L .

Suppose that we are given a class \mathcal{C} of links whose Conway polynomials we know, and a link L whose Conway polynomial we want to compute. Let us define a *computation tree* for L with respect to \mathcal{C} . A computation tree T_0 of order 0 is a homotopy h_t from L to some $L' \in \mathcal{C}$ through links and singular links with one double point. The singular links of h_t are called the *buds* of T_0 . A computation tree T_n of order $n > 1$ consists of a computation tree T_{n-1} of order $n - 1$, and, for each bud Λ of T_{n-1} , of a homotopy h_t^Λ from the smoothing Λ_0 of Λ to some $\Lambda' \in \mathcal{C}$ through links and singular links with one double point. The singular links of each h_t^Λ are called the *buds* of T_0 . We also call L the *root*, the homotopies added when forming T_k the *order k branches*, and their final links, which are in \mathcal{C} , the *order k leaves* of T_n .

Since every link is homotopic to a trivial link, for every link L there exists a computation tree of an arbitrarily high order, whose leaves are trivial links. If we want the computation to be guaranteed to terminate in finitely many steps, a slightly more elaborate construction is needed (see Remark 10.4 below).

Lemma 10.1. (cf. [47; Proposition 8.7]) (a) $\nabla_L(z) = 0$ for any split link L .

¹⁶As explained in §4, L_+ and L_- can be distinguished from each other without using the plane diagram.

(b) The Conway polynomial of an m -component link L is of the form

$$\nabla_L(z) = z^{m-1}(c_0 + c_1z^2 + c_2z^4 + \cdots + c_rz^{2r}).$$

(c) $c_0(K) = 1$ for a knot K , and $c_0(L) = \text{lk}(L)$ for a 2-component link L .

Proof (sketch). (a). Represent L as Λ_0 , where Λ_+ and Λ_- are ambient isotopic. \square

(b). We need to show that the coefficient of ∇_L at z^n is zero if $n < m - 1$ or n has the same parity as m . If $n < m - 1$, there is a computation tree for L of order n whose all leaves are split links. In general there is a computation tree for L of order n whose leaves are split links and unknots; each leaf of an order $\equiv m \pmod{2}$ has an even number of components, and so cannot be an unknot. \square

(c). Use any computation tree for L of order 0, respectively 1, whose leaves are split links and unknots. \square

Remark 10.2. It is easy to see that for 3-component links $c_0(L) = ab + bc + ca$, where a, b, c are the linking numbers of the 2-component sublinks (cf. [49]). For any number of components, $c_0(L)$ is a symmetric polynomial in the pairwise linking numbers l_{ij} . Namely, $c_0(L) = \det(\Lambda^{(p)})$ for any $p \in \{1, \dots, m\}$, where $\Lambda = (\lambda_{ij})$ is the $m \times m$ matrix with entries

$$\lambda_{ij} = \begin{cases} -l_{ij}, & \text{if } i \neq j \\ \sum_{k \neq i} l_{ik} & \text{if } i = j \end{cases}$$

and $\Lambda^{(p)}$ denotes the $(m-1) \times (m-1)$ matrix obtained by removing from Λ the p th row and the p th column (Hosokawa, Hartley and Hoste; see [46; Proposition 3.2] for a proof and [50] for the references). Hartley and Hoste also reformulated the same expression as a sum over all spanning trees T of the complete graph K_m :

$$c_0(L) = \sum_T \prod_{\{i,j\} \in E(T)} l_{ij}.$$

Lemma 10.3. For an m -component link L each $c_n(L)$ is of type $m - 1 + 2n$ and of colored type $2n$.

The first assertion is well-known [7; Theorem 2].

Proof. The skein relation (C) is reminiscent of the formula (V) in §4 describing the extension of a link invariant to singular links. The two formulas together imply the relation $\nabla_{\times}^{\times} = z\nabla_{\zeta}^{\times}$, where ∇^{\times} is the standard extension of ∇ to singular links. It follows that $\nabla_{\Lambda}^{\times}$ is divisible by z^{n+1} for any singular link Λ with $n + 1$ double points. Therefore the coefficient of ∇_L at z^n is a type n invariant.

Now suppose that Λ is a singular link with m components and $n + 1$ double points, whose all double points are self-intersections of components. Let us color the components of Λ in distinct colors. Then after smoothing each double point we obtain a non-singular link $\Lambda_{0\dots 0}$ which is still colored in m colors (each component being colored only in one color, and each of the m colors being used). This means that $\Lambda_{0\dots 0}$ has at least m

components. Then by Lemma 10.1(b) $\nabla_{\Lambda_0 \dots 0}$ is divisible by z^{m-1} . Hence $\nabla_{\Lambda}^{\times}$ is divisible by $z^{(m-1)+(n+1)}$. Since Λ was an arbitrary singular link with $n+1$ double points that are all self-intersections of components, the coefficient of ∇_L at z^{m-1+n} is a colored type n invariant. \square

Remark 10.4. For every link L there exists a finite computation tree for L which has no buds, and whose leaves are trivial links. Namely, given a plane diagram D of L , this D is also a plane diagram of a trivial link L' (see [72; Theorem 3.8]).¹⁷ In order to get from L to L' , we only need to switch some of the crossings in D . For each singular link Λ in this homotopy, its smoothing Λ_0 has a plane diagram which has one less crossing than D . Then we repeat the same construction for this plane diagram, and so on. This process terminates after finitely many steps, since the number of crossings in the plane diagrams being considered decreases in each step. One consequence of this computation is that the degree of ∇_L is bounded from above by the number of crossings in D .

Lemma 10.5. (cf. [47; Proposition 16.2(ii)]) $\nabla_{L\#_{i,j}L'} = \nabla_L \nabla_{L'}$.

Here $L\#_{i,j}L'$ is the connected sum of L and L' along selected components (see §8).

Proof (sketch). The computation tree for L described in Remark 10.4 yields $\nabla_L = \sum_{k=0}^n z^{2k} \sum_{i=1}^{r_k} \epsilon_{ki} \nabla_U$, where U is the unknot and $\epsilon_{ki} = \pm 1$. If the same computation tree is applied to a plane diagram of $L\#_{i,j}L'$ obtained from the diagram D of L by adjoining a diagram of L' lying in a small disk, so that the contents of the small disk is kept intact, then it yields $\nabla_{L\#_{i,j}L'} = \sum_{k=0}^n z^{2k} \sum_{i=1}^{r_k} \epsilon_{ki} \nabla_{L'}$, where r_k and ϵ_{ki} are same as before. Since $\nabla_U = 1$, the two expressions combine to yield the desired result. \square

Since $c_0(K) = 1$ for every knot K , we have the formal power series

$$\bar{\nabla}_L(z) := \frac{\nabla_L(z)}{\nabla_{K_1}(z) \cdots \nabla_{K_m}(z)},$$

where K_1, \dots, K_m are the components of L .

From Lemma 10.5 we obtain

Corollary 10.6. [83], [78] $\bar{\nabla}_L$ is invariant under PL isotopy.

Proposition 10.7. If L is an m -component link, then $\bar{\nabla}_L$ is of the form

$$\bar{\nabla}_L(z) = z^{m-1}(\alpha_0 + \alpha_1 z^2 + \alpha_2 z^4 + \dots),$$

with each $\alpha_i(L) = c_i(L) - (\alpha_{i-1}(L)c_1(K) + \dots + \alpha_0(L)c_i(K))$, where $K = K_1\# \dots \# K_m$.

Proof. By Lemma 10.5 $\nabla_{K_1} \cdots \nabla_{K_m} = \nabla_K$. Then the definition of $\bar{\nabla}_L$ implies $\bar{\nabla}_L = \nabla_L - \bar{\nabla}_L(\nabla_K - 1)$. The assertion follows from this formula and Lemma 10.1(b). \square

¹⁷Thinking of D as lying in the xy plane, each component of L' consists of arc which projects homeomorphically onto the z axis and a straight line segment whose projection onto the xy -plane is disjoint from the crossings of D (cf. [72; Figure 3.10]). The projections of the components on the z axis are pairwise disjoint.

Corollary 10.8. *If L is a 2-component link, then $\alpha_0(L) = c_0(L)$ is its linking number and $\alpha_1(L) = c_1(L) - c_0(L)(c_1(K_1) + c_1(K_2))$ is its generalized Sato–Levine invariant.*

See Example 1.6 concerning the generalized Sato–Levine invariant. The following crossing change formula for the generalized Sato–Levine invariant is well-known.

Proposition 10.9. [49] (see also [83; Theorem 10.2], [40], [3]) *Let $L = (K, K')$ be a 2-component link. Then the jump of $\alpha_1(L)$ under a self-intersection of K is*

$$\alpha_1(L_+) - \alpha_1(L_-) = \text{lk}(P, K') \text{lk}(Q, K'),$$

where P and Q are the components of the smoothing K_0 of the singular knot.

Proof. We have

$$\alpha_1(L_+) - \alpha_1(L_-) = (c_1(L_+) - c_1(L_-)) - c_0(L_+)c_0(K_0) = c_0(L_0) - c_0(L_+)c_0(K_0),$$

using that $c_0(L_+) = c_0(L_-)$. Now

$$c_0(L_0) = \text{lk}(P, K') \text{lk}(Q, K') + \text{lk}(P, K') \text{lk}(P, Q) + \text{lk}(Q, K') \text{lk}(P, Q).$$

The latter two summands are canceled by $c_0(L_+)c_0(K_0) = \text{lk}(K_+, K') \text{lk}(P, Q)$. The remaining summand is the one in the statement. \square

In order to similarly understand $\alpha_1(L)$ for a 3-component link L , it is convenient to introduce a correction term. Given a 3-component link L , let us consider

$$\gamma(L) := \alpha_1(L) - \sum_{(\Lambda, \Lambda')} \alpha_1(\Lambda) \alpha_0(\Lambda'),$$

where (Λ, Λ') runs over all ordered pairs of distinct 2-component sublinks of L .

Proposition 10.10. *Let $L = (K, K', K'')$ be a 3-component link. Then the jump of $\gamma(L)$ under a self-intersection of K is*

$$\gamma(L_+) - \gamma(L_-) = \text{lk}(K', K'') (\text{lk}(P, K') \text{lk}(Q, K'') + \text{lk}(Q, K') \text{lk}(P, K'')),$$

where P and Q are the components of the smoothing K_0 of the singular knot.

Proof. We have $\alpha_1(L) = c_1(L) - c_0(L)(c_1(K) + c_1(K') + c_1(K''))$. Then

$$\alpha_1(L_+) - \alpha_1(L_-) = (c_1(L_+) - c_1(L_-)) - c_0(L_+)c_0(K_0) = c_0(L_0) - c_0(L_+)c_0(K_0),$$

using that $c_0(L_+) = c_0(L_-)$. The formula of Remark 10.2 expresses $c_0(L_0)$ as a sum of 16 summands (corresponding to the 15 spanning trees of K_4). It can be checked that 8 of these summands cancel with $c_0(L_+)c_0(K_0)$ and another 6 cancel with the jump of the correction term. The remaining 2 summands are the ones in the statement. \square

Remark 10.11. P. M. Akhmetiev obtained some applications of Proposition 10.10 in magnetohydrodynamics [1], [2], building on his previous applications of the generalized Sato–Levine invariant.

Proposition 10.12. *For an m -component link L each $\alpha_n(L)$ is*

(a) *of type $m - 1 + 2n$ and of colored type $2n$;*

(b) of colored type $2n - 1$ if $n > 0$.

Proof. (a). By Lemma 10.5 $\nabla_K = \nabla_{K_1}(z) \cdots \nabla_{K_m}(z)$, where K_1, \dots, K_m are the components of L and $K = K_1 \# \dots \# K_m$. Hence $c_i(K) = \sum_{i_1 + \dots + i_m = i} c_{i_1}(K_1) \cdots c_{i_m}(K_m)$. By Lemma 10.3 each $c_k(K_j)$ is a type $2k$ invariant of K_j , and hence a type $2k$ invariant of L (cf. Lemma 4.2(a)). Then by Corollary 4.6(a) $c_i^*(L) := c_i(K)$ is a type $2i$ invariant of L .

By Proposition 10.7 each $\alpha_n(L) = c_n(L) - (\alpha_{n-1}(L)c_1^*(L) + \cdots + \alpha_0(L)c_n^*(L))$. By Lemma 10.3 $\alpha_0(L) = c_0(L)$ is of type $m - 1$ and of colored type 0. Now it follows by induction, using Corollary 4.6(a), that $\alpha_n(L)$ is a type $m - 1 + 2n$ invariant and a colored type $2n$ invariant of L . \square

(b). We will use the notation from the proof of (a). We need to show that $\alpha_k(L)$ is of colored type $2k - 1$ for each $k > 0$. Suppose that this is known for $i = 1, \dots, n - 1$. Then it follows from Corollary 4.6(a) that $(\alpha_{n-1}(L)c_1^*(L) + \cdots + \alpha_1(L)c_{n-1}^*(L))$ is of colored type $2n - 1$. Since $\alpha_n(L) = c_n(L) - (\alpha_{n-1}(L)c_1^*(L) + \cdots + \alpha_0(L)c_n^*(L))$, it remains to show that $c_n(L) - c_0(L)c_n^*(L)$ is a colored type $2n - 1$ invariant of L .

Let $\nabla_L^* = \nabla_K$. Thus we have $\nabla_L^* = c_0^*(L) + c_1^*(L)z^2 + \cdots + c_n^*(L)z^{2n}$ for some $n = n(L)$, where $c_0^*(L) = 1$. Then $c_n(L) - c_0(L)c_n^*(L)$ is the coefficient at z^{m-1+2n} of the polynomial $\hat{\nabla}_L := \nabla_L - z^{m-1}c_0(L)\nabla_L^*$. More generally, for a link L colored in m colors so that no color is omitted, let $\nabla_L^* = \nabla_{L^1}(z) \cdots \nabla_{L^m}(z)$, where L^i is the sublink of L of the i th color, and let $\hat{\nabla}_L = \nabla_L - z^{m-1}\lambda(L)\nabla_L^*$, where $\lambda(L) = \sum_T \prod_{\{i,j\} \in E(T)} \text{lk}(L^i, L^j)$, the sum being over all spanning trees T of the complete graph K_m (cf. Remark 10.2).

Let Λ be a singular link with m components and $2n$ double points, whose all double points are self-intersections of components. The standard extension $\hat{\nabla}^\times$ of $\hat{\nabla}$ to singular links is easily seen to satisfy $\hat{\nabla}_\Lambda^\times = \nabla_\Lambda^\times - z^{m-1}\lambda(\Lambda)\nabla_\Lambda^{*\times}$, where λ is extended by continuity to singular links which are link maps. This implies the relation $\nabla_{\times}^\times = z\nabla_{\wr}^\times$. Let $\Lambda_{0\dots 0}$ be the non-singular link obtained by smoothing all the double points of Λ . As explained in the proof of Lemma 10.3 $\Lambda_{0\dots 0}$ is colored in m colors so that no color is omitted.

If $\Lambda_{0\dots 0}$ has precisely m components, then it follows from Remark 10.2 that $\hat{\nabla}_{\Lambda_{0\dots 0}}$ is divisible by z^m . If $\Lambda_{0\dots 0}$ has more than m components, then by Lemma 10.1(b) $\nabla_{\Lambda_{0\dots 0}}$ is divisible by z^m and also $\nabla_{\Lambda_{0\dots 0}^i}$ is divisible by z for at least one index i . Hence again $\hat{\nabla}_{\Lambda_{0\dots 0}}$ is divisible by z^m .

Since $\hat{\nabla}_{\Lambda_{0\dots 0}}$ is divisible by z^m , $\hat{\nabla}_\Lambda^\times$ is divisible by z^{m+2n} . Since Λ was an arbitrary singular link with $2n$ double points whose all double points are self-intersections of components, the coefficient of $\hat{\nabla}_L$ at z^{m-1+2n} is a colored type $2n - 1$ invariant. \square

Proposition 10.13. $\alpha_2(L)$ is not of colored type 2, nor even of type $(2, 2)$ for two-component links of linking number 0.

Proof. For a two-component link L of linking number 0 we have $c_0(L) = 0$ and hence $\alpha_2(L) = c_2(L) - c_1(L)(c_1(K_1) + c_1(K_2))$. Let $L_{\times \times \times}$ be a two-component singular link with 3 double points and linking number 0 such that all its double points are on the first component, and after smoothing any two of them we get a 2-component (and not

4-component) link. If we regard $v_i(L) := c_1(K_i)$ as an invariant of L , then we have $\alpha_2^\times(L_{\times\times\times}) = c_2^\times(L_{\times\times\times}) - (c_1v_1)^\times(L_{\times\times\times}) - (c_2v_2)^\times(L_{\times\times\times})$. Clearly $c_2^\times(L_{\times\times\times}) = c_0(L_{000})$, where L_{000} is the 3-component link obtained by smoothing all double points of $L_{\times\times\times}$. Lemma 4.5 expresses $(c_1v_i)^\times(L_{\times\times\times})$ as the sum of all products of the form $c_1^\times(L_\alpha)v_i^\times(L_\beta)$, where L_α is obtained by resolving positively some number i of the double points of $L_{\times\times\times}$ and L_β by resolving negatively the remaining $3 - i$ double points of $L_{\times\times\times}$. By Lemma 10.3 $c_1(L)$ is of colored type 2, so $c_1^\times(L_{\times\times\times}) = 0$. But since $v_2(L)$ depends only on the second component, v_2^\times vanishes on any singular link that has at least one double point on the first component, so from Lemma 4.5 we get $(c_1v_2)^\times(L_{\times\times\times}) = 0$. Since v_1 is of colored type 2 (cf. Lemma 4.2(a)), we have $v_1^\times(L_{\times\times\times}) = 0$. If the first double point of $L_{\times\times\times}$ is resolved positively or negatively, then the resulting singular link $L_{\pm\times\times}$ satisfies $c_1^\times(L_{\pm\times\times}) = c_0(L_{\pm00})$, where $L_{\pm00}$ is the two-component link obtained by smoothing both double points of $L_{\pm\times\times}$, but $c_0(L_{\pm00}) = \text{lk}(L_{\pm00}) = \text{lk}(L_{\pm\times\times}) = 0$. Also, $v_1^\times(L_{\pm\times\times}) = c_0(K_{\pm00})$, where $K_{\pm00}$ is the first component of $L_{\pm00}$, but we know that $c_0(K_{\pm00}) = 1$ (see Lemma 10.1(c)). Similar arguments apply if the second or the third double point of $L_{\times\times\times}$ is resolved positively or negatively. Thus from Lemma 4.5 we get

$$\begin{aligned} (c_1v_1)^\times(L_{\times\times\times}) &= c_1^\times(L_{++\times})v_1^\times(L_{\times\times-}) + c_1^\times(L_{+\times+})v_1^\times(L_{\times-\times}) + c_1^\times(L_{\times++})v_1^\times(L_{-\times\times}) \\ &= c_1^\times(L_{++\times}) + c_1^\times(L_{+\times+}) + c_1^\times(L_{\times++}) = c_0(L_{++0}) + c_0(L_{+0+}) + c_0(L_{0++}), \end{aligned}$$

where $+$ and $-$ indicate the positive and the negative resolution of the corresponding double point, and 0 its smoothing. Thus

$$\alpha_2^\times(L_{\times\times\times}) = c_0(L_{000}) - c_0(L_{++0}) - c_0(L_{+0+}) - c_0(L_{0++}).$$

Let $\varphi: S^1 \rightarrow \mathbb{R}^2$ be a self-transverse C^1 -approximation with 3 double points of the double cover $S^1 \rightarrow S^1$ of the clockwise oriented unit circle $S^1 \subset \mathbb{R}^2$, and let A, B, C, D denote the 4 bounded components of $\mathbb{R}^2 \setminus \varphi(S^1)$ such that D contains the origin. The composition $K_{\times\times\times}: S^1 \xrightarrow{\varphi} \mathbb{R}^2 \subset \mathbb{R}^3$ is a singular knot. Let $K': S^1 \hookrightarrow S^3 \setminus K_{\times\times\times}(S^1)$ be a knot in the complement of $K_{\times\times\times}$ linking the clockwise oriented boundaries of A, B, C, D with linking numbers a, b, c, d , respectively, so that $a + b + c + 2d = 0$. Finally, define $L_{\times\times\times}$ to be the union of $K_{\times\times\times}$ and K' . Then $\text{lk}(L_{\times\times\times}) = a + b + c + 2d = 0$ and by smoothing any two of the 3 double points of $L_{\times\times\times}$ we get a 2-component link, so that our previous formula for $\alpha_2^\times(L_{\times\times\times})$ applies. It is easy to see that by smoothing any one of the double points of $K_{\times\times\times}$, we get a link of linking number 1, and by smoothing all three we get a link of linking number 0. Then using the formula for c_0 of a 3-component link (see Remark 10.2), we find that $c_0(L_{000}) = (a + b + c + d)d$ and one of $c_0(L_{++0})$, $c_0(L_{+0+})$ and $c_0(L_{0++})$ equals $(a + b + d)(c + d) + (a + b + d) + (c + d)$ and the other two are obtained from it by cyclically permuting a, b and c . Since $(a + b + d) + (c + d) = 0$ and $(a + b + d)(c + d) = -(c + d)^2$, we get

$$\alpha_2^\times(L_{\times\times\times}) = (a + b + c + d)d + (a + d)^2 + (b + d)^2 + (c + d)^2.$$

This expression is nonzero e.g. for $a = 1, b = 2, c = -3, d = 0$. \square

It was shown in [60; proof of Theorem 2.2] that α_1 is invariant under 1-quasi-isotopy. By Proposition 10.12 and Theorem 4.8 $\alpha_n(L)$ is invariant under $(2n - 1)$ -quasi-isotopy. However, by Proposition 10.13 the following cannot be deduced from Theorem 4.8.

Theorem 10.14. $\alpha_n(L)$ is invariant under n -quasi-isotopy.

Proof. Let L be a singular link which is an n -quasi-embedding, and let m be the number of its components. Let P_0, \dots, P_n and J_0, \dots, J_n be as in the definition of an n -quasi-embedding. We may assume that the P_i are connected. By passing to small regular neighborhoods we may assume that P_1, \dots, P_n are connected 3-manifolds with boundary, and P_0 lies in a 3-ball P_0^+ such that $P_0^+ \cup L(J_0)$ is null-homotopic in P_1 . We may assume that the images of L_+ , L_- and L_0 lie in $P_0^+ \cup L$. For the purposes of the present proof let us redefine P_0 as P_0^+ . Let K be the component of L containing the double point. Then K_+ , K_- and K_0 lie in $P_0 \cup K = P_0 \cup K_-$.

Lemma 10.15. [59; Corollary 7.9] *Let L be a link in S^3 and let A be an embedded annulus in $S^3 \setminus L$ whose core has zero linking number with L . Then $\nabla_{L \cup \partial A}(z) = 0$.*

Lemma 10.16. *For each $k \leq n$ there exists a computation tree T_{2k} of order $2k$ for K_+ , whose branches are homotopies with values in $P_k \cup K_-$ and whose leaves are split links, copies of K_- and 3-component links of the form $K_- \cup \partial A$, where A is an embedded annulus in $P_k \setminus K_-$ whose core has zero linking number with K_- .*

Proof. Let T_0 consist of the straight line homotopy between K_+ and K_- , which has one singular link, namely, K . Arguing by induction, we may assume that T_{2k} has been constructed. Let X be a bud of T_{2k} . Let Q_1, \dots, Q_μ be the components of its smoothing X_0 . Thus Q_1, \dots, Q_μ lie in $P_k \cup K_-$, and one of them, say Q_1 , contains $K_- \setminus P_k$. Then Q_1 is homotopic to K_- in $P_{k+1} \cup K_-$ by some homotopy $Q_{1,t}$, and the other ones are null-homotopic in P_{k+1} by some homotopies $Q_{2,t}, \dots, Q_{\mu,t}$. Since K_+ is a knot, X has an odd number of components, and then X_0 has an even number of components. Thus $\mu \geq 2$. Using the null-homotopy $Q_{2,t}$ and that P_{k+1} is a 3-manifold, it is easy to construct a homotopy h_t^X with values in $P_{k+1} \cup K_-$ from X_0 to a split link. Using again that P_{k+1} is a 3-manifold, we may assume that this homotopy goes only through links and singular links with one double point. Let T_{2k+1} consist of T_{2k} and the homotopies h_t^X for all buds X of T_{2k} . Let Y be a bud of T_{2k+1} , that is, a singular link of some h_t^X .

Let us first consider the case where the double point of Y is an intersection of $Q_{2,s}$ for some $s \in (0, 1)$ with some Q_i , $i \neq 2$. If $i \geq 3$, then the null-homotopies $Q_{2,t|t \in [s, 1]}$, and $Q_{i,t}$ can be combined to get a null-homotopy of $Q_{2,s} \cup Q_i$ in P_{k+1} . This in turn yields a homotopy with values in $P_{k+1} \cup K_-$ from Y_0 to a split link. If $i = 1$ and $\mu = 2$, then using the same homotopies $Q_{2,t|t \in [s, 1]}$ and $Q_{i,t}$ one can similarly construct a homotopy with values in $P_{k+1} \cup K_-$ from Y_0 to K_- . If $i = 1$ and $\mu \geq 3$, then using the null-homotopy $Q_{3,t}$ it is easy to construct a homotopy with values in $P_{k+1} \cup K_-$ from Y_0 to a split link.

It remains to consider the case where the double point of Y is a self-intersection of $Q_{2,s}$ for some $s \in (0, 1)$. If $\mu \geq 3$, then using the null-homotopy $Q_{3,t}$ it is easy to construct a homotopy with values in $P_{k+1} \cup K_-$ from Y_0 to a split link. If $\mu = 2$, then Y_0 is a

3-component link consisting of Q_0 and two other knots C, D , and the null-homotopy $Q_{2,t}|_{t \in [s,1]}$ can be used to construct a homotopy with values in $P_{k+1} \cup K_-$ from C to $-D$, i.e. to D with orientation reversed. This in turn yields a homotopy with values in $P_{k+1} \cup K_-$ from Y_0 to a link of the form $Q_1 \cup \partial A$, where A is an embedded annulus in P_{k+1} , disjoint from Q_1 . Since P_{k+1} is connected, by a further homotopy with values in $P_{k+1} \cup K_-$ we may assume that the core of A has zero linking number with K_- . \square

Let T_{2n} be a computation tree given by Lemma 10.16. Since K_0 is a 2-component link, each leaf of T_{2n} of an odd order must have an even number of components, so it must be a split link. Let Z be a leaf of T_{2n} of order $2k$ which is of the form $K_- \cup \partial A$. Then the annulus A lies in P_k , so in the case $k < n$ it is null-homotopic in P_n . Then its core α has zero linking number with $L_- \setminus K_-$. Since α also has zero linking number with K_- , we have $\text{lk}(\alpha, L_-) = 0$. This does not work if Z is a leaf of order $2n$, but in this case, since P_n is connected, we can amend Z by a homotopy with values in $P_n \cup K_-$ so that $\text{lk}(\alpha, K_-)$ is any given number. If we choose this number to be $-\text{lk}(\alpha, L_- \setminus K_-)$, then we get $\text{lk}(\alpha, L_-) = 0$. This amendment, when carried out for all leaves of T_{2n} of order $2n$ which are of the form $K_- \cup \partial A$, results in a new computation tree T'_{2n} , whose branches of orders $< 2n$ are the same as those of T_{2n} . By adding the identical homotopies of $L_- \setminus K_-$, we may regard T'_{2n} as a computation tree of order $2n$ for L_+ , whose leaves are split links, copies of L_- , and links of the form $L_- \cup \partial A$, where A is an embedded annulus in $S^3 \setminus L_-$ whose core has zero linking number with L_- .

Taking into account Lemma 10.15, from T_{2n} we get

$$\nabla_{K_+} = \sum_{k=0}^n z^{2k} \left(\sum_{i=1}^{r_k} \epsilon_{ki} \nabla_{K_-} \right) + z^{2n+1} P$$

for some $P \in \mathbb{Z}[z]$, where $r_0 = 1$ and $\epsilon_{01} = +1$, and for $k > 0$ the signs ϵ_{ki} are the signs of some of the singular links in the homotopies that are branches of T_{2n} of order $2k - 1$. The polynomial P is a sum with signs of the Conway polynomials of the buds of T_{2n} . Similarly, from T'_{2n} we get

$$\nabla_{L_+} = \sum_{k=0}^n z^{2k} \left(\sum_{i=1}^{r_k} \epsilon_{ki} \nabla_{L_-} \right) + z^{2n+1} Q$$

for some $Q \in \mathbb{Z}[z]$, where the r_k and the ϵ_{ki} are the same as in the previous sum, since they are determined by the branches of orders $< 2n$, which are shared by T_{2n} and T'_{2n} . The polynomial Q is a sum with signs of the Conway polynomials of the buds of T'_{2n} . These buds all have at least m components, so by Proposition 10.1(b) $Q = z^{m-1} Q_0$ for some $Q_0 \in \mathbb{Z}[z]$. Let $R = \sum_{k=0}^n z^{2k} \sum_{i=1}^{r_k} \epsilon_{ki}$. Then $\nabla_{K_+} = R \nabla_{K_-} + z^{2n+1} P$ and $\nabla_{L_+} = R \nabla_{L_-} + z^{(m-1)+(2n+1)} Q_0$. Hence

$$\frac{z^{1-m} \nabla_{L_+}}{\nabla_{K_+}} = \frac{z^{1-m} \nabla_{L_-} R + z^{2n+1} Q_0}{\nabla_{K_-} R + z^{2n+1} P} = \frac{z^{1-m} \nabla_{L_-}}{\nabla_{K_-}} + z^{2n+1} S$$

for some $S \in \mathbb{Z}[[z]]$. Therefore $z^{1-m} \bar{\nabla}_{L_+} = z^{1-m} \bar{\nabla}_{L_-} + z^{2n+1} S_0$ for some $S_0 \in \mathbb{Z}[[z]]$, and the assertion follows. \square

Remark 10.17. The most difficult case in the proof of Lemma 10.16 is a self-intersection of the null-homotopy of Q_2 in the absence of Q_3 . However, if we start from a *strong* k -quasi-isotopy, then this difficulty disappears (because a knot lying in a ball is null-homotopic within this ball), and one might even wonder if anything at all prevents building a computation tree of any order under the hypothesis of strong 1-quasi-isotopy. In fact, the only thing which may go wrong with it is the following: in the homotopy from Q_1 to K_- we may get a self-intersection with Q_1 . If one lobe of the resulting singular link lies entirely within the ball B_1 (from the definition of a strong 1-quasi-isotopy), this is still not a problem, but it may happen that none of the two lobes lies in B_1 .

Since $c_k \equiv \alpha_k \pmod{\gcd(c_0, \dots, c_{k-1})}$, Theorem 10.14 implies the case $l = 1$, and also the 2-component case of the following

Theorem 10.18. [61; Corollary 3.5] *Set $\lambda = \lceil \frac{(l-1)(m-1)}{2} \rceil$. The residue class of $c_{\lambda+k}$ modulo \gcd of $c_\lambda, \dots, c_{\lambda+k-1}$ and all $\bar{\mu}$ -invariants of length $\leq l$ is invariant under $(\lfloor \frac{\lambda}{m-1} \rfloor + k)$ -quasi-isotopy of m -component links.*

(Here $\lceil x \rceil = n$ if $x \in [n - \frac{1}{2}, n + \frac{1}{2})$, and $\lfloor x \rfloor = n$ if $x \in (n - \frac{1}{2}, n + \frac{1}{2}]$ for $n \in \mathbb{Z}$.)

One special case of Theorem 10.18 not covered by Theorem 10.14 asserts that for 3-component links the residue class of c_k modulo the greatest common divisor Δ_k of all $\bar{\mu}$ -invariants of length $\leq k + 1$ is invariant under $\lfloor \frac{k}{2} \rfloor$ -quasi-isotopy. When $k = 1$, this is saying that the residue class of c_1 modulo the \gcd of the pairwise linking numbers is invariant under 0-quasi-isotopy, that is, link homotopy. This residue class is also known as $\bar{\mu}(123)^2$ (Murasugi–Traldi [69], [82] and Cochran [14]; see also [46], [50]).

Can one find an integer lift of $\bar{\mu}(123)^2$, invariant under link homotopy, among coefficients of rational functions in $\bar{\nabla}_\Lambda$ for sublinks Λ of L ? The answer is negative, since the desired integer invariant would be a finite type (specifically, type 4) invariant of link homotopy which is not a function of the pairwise linking numbers, but this is impossible for 3-component links by [63]. Alternatively, using Proposition 10.10 one can see it directly, by checking that the jump of $\gamma(L)$ cannot be canceled by the jump of any polynomial expression in the coefficients of ∇_Λ , homogeneous of degree 4.

ACKNOWLEDGEMENTS

I would like to thank P. M. Akhmetiev, I. Dynnikov and M. Il'insky for useful discussions and thoughtful remarks.

REFERENCES

- [1] P. M. Akhmet'ev, *On a higher integral invariant for closed magnetic lines, revisited*, J. Geom. Phys. **170** (2021), Paper No. 104379, 15pp. [↑1.6](#), [10.11](#)
- [2] ———, *Topological meaning of the slope of the Kolmogorov spectrum of magnetic turbulence: M_5 -invariant of magnetic lines and its combinatorial formula*, J. Geom. Phys. **178** (2022), Paper No. 104583, 14pp. [↑1.6](#), [10.11](#)
- [3] P. M. Akhmet'ev and D. Repovš, *A generalization of the Sato–Levine invariant*, Tr. Mat. Inst. Steklova **221** (1998), 69–80; [MathNet](#); English transl., Proc. Steklov Inst. Math. **221** (1998), 60–70. [↑1.6](#), [10.9](#)

- [4] P. M. Akhmet'ev, J. Malešič, and D. Repovš, *A formula for the generalized Sato–Levine invariant*, Mat. Sb. **192** (2001), no. 1, 3–12; [MathNet](#); English transl., Sb. Math. **192** (2001), 1–10. [↑1.6](#)
- [5] D. Altschüler and L. Freidel, *On universal Vassiliev invariants*, Comm. Math. Phys. **170** (1995), 41–62. [arXiv:hep-th/9403053](#). [↑9.3](#)
- [6] B. Audoux, *Singular link Floer homology*, Algebr. Geom. Topol. **9** (2009), 495–535. [arXiv:0705.2377v6](#) (2017). [↑1.1](#)
- [7] D. Bar-Natan, *On the Vassiliev knot invariants*, Topology **34** (1995), 423–472. [Author's webpage](#). [↑1.4](#), [4](#), [9.3](#), [10](#)
- [8] ———, *Non-associative tangles*, Geometric Topology (Athens, GA, 1993), AMS/IP Stud. Adv. Math., vol. 2.1, Amer. Math. Soc., Providence, RI, 1997, pp. 139–183. [Author's webpage](#). [↑9.3](#)
- [9] R. Blair, J. Burke, and R. Koytcheff, *A prime decomposition theorem for the 2-string link monoid*, J. Knot Theory Ram. **24** (2015), 1550005. [arXiv:1308.1594](#). [↑12](#)
- [10] J. Buckley, *On the D-series of a finite group*, Proc. Amer. Math. Soc. **18** (1967), 185–186. [Journal](#). [↑5](#), [5.10](#)
- [11] G. Burde and H. Zieschang, *Knots*, 2nd ed., De Gruyter Studies in Math., vol. 5, de Gruyter, Berlin, 2003. [↑5](#), [8](#)
- [12] S. Chmutov and S. Duzhin, *The Kontsevich integral*, Acta Appl. Math. **66** (2001), 155–190. [CiteSeerX](#). [↑9.3](#)
- [13] S. Chmutov, S. Duzhin, and J. Mostovoy, *Introduction to Vassiliev Knot Invariants*, Cambridge Univ. Press, Cambridge, 2012. [↑1.1](#), [4](#), [4](#), [2](#), [5.5](#), [5](#), [5](#), [9.3](#), [13](#), [15](#)
- [14] T. D. Cochran, *Concordance invariance of coefficients of Conway's link polynomial*, Invent. Math. **82** (1985), 527–541. [EuDML](#). [↑10](#)
- [15] ———, *Derivatives of links: Milnor's concordance invariants and Massey's products*, Mem. Amer. Math. Soc. **84** (1990), no. 427. [↑8](#)
- [16] R. J. Daverman, *Decompositions of Manifolds*, Pure Appl. Math., vol. 124, Academic Press, Orlando, FL, 1986. [↑1.1](#), [1.6](#)
- [17] N. M. Dunfield and S. Garoufalidis, *Non-triviality of the A-polynomial for knots in S^3* , Algebr. Geom. Topol. **4** (2004), 1145–1153. [arXiv:math/0405353](#). [↑1.1](#)
- [18] M. Eisermann, *Invariants de Vassiliev et conjecture de Poincaré*, C. R. Math. Acad. Sci. Paris **334** (2002), 1005–1010 (French, contains abridged English version). [Journal](#). [↑1.1](#)
- [19] ———, *Vassiliev invariants and the Poincaré conjecture*, Topology **43** (2004), 1211–1229. [Journal](#). [↑1.1](#)
- [20] R. H. Fox, *A quick trip through knot theory*, Topology of 3-Manifolds and Related Topics (Proc. The Univ. of Georgia Institute, 1961), Prentice-Hall, Inc., Englewood Cliffs, NJ, 1961, pp. 120–167. [A. Ranicki's web archive](#). [↑5](#), [8](#)
- [21] M. H. Freedman, *Whitehead₃ is a "slice" link*, Invent. Math. **94** (1988), 175–182. [EuDML](#). [↑1.1](#), [3](#)
- [22] L. Funar, *TQFT and Whitehead's manifold*, J. Knot Theory Ram. **6** (1997), 13–30. [Author's homepage](#). [↑3.4](#)
- [23] ———, *On knots having the same invariants up to a certain degree* (1999). [Author's webpage](#). [↑5](#)
- [24] S. Garoufalidis, *Does the Jones polynomial determine the signature of a knot?*. [arXiv:math/0310203](#). [↑1.1](#)
- [25] C. H. Giffen, *Disciplining dunce hats in 4-manifolds* (1976). Unpublished typeset manuscript. [↑1.1](#), [1.6](#)
- [26] ———, *F-isotopy and I-equivalence of links* (1976). Unpublished handwritten manuscript. [↑1.6](#)
- [27] M. N. Gusarov, *A new form of the Conway–Jones polynomial of oriented links*, Zap. Nauchn. Sem. LOMI **193** (1991), 4–9, 161; [MathNet](#). English transl. in *Topology of Manifolds and Varieties*, Adv. Soviet Math., vol. 18, Amer. Math. Soc., Providence, RI, 1994, pp. 167–172. [↑4](#)

- [28] ———, *On the n -equivalence of knots and invariants of finite degree*, Zap. Nauchn. Sem. POMI **208** (1993), 152–173, 221–222; [MathNet](#); English transl., J. Math. Sci. **81** (1996), 2549–2561. Another English transl. in *Topology of Manifolds and Varieties*, Adv. Soviet Math., vol. 18, Amer. Math. Soc., Providence, RI, 1994, pp. 173–192. [↑4.6](#), [4](#), [5](#), [5.4](#), [5](#), [6](#), [7](#)
- [29] ———, *Variations of knotted graphs. The geometric technique of n -equivalence*, Algebra i Analiz **12** (2000), no. 4, 79–125; [MathNet](#); English transl., St. Petersburg Math. J. **12** (2001), 569–604. [Goussarov memorial page maintained by D. Bar-Natan](#). [↑4](#), [5.4](#), [5.6](#), [5](#), [5](#), [5](#)
- [30] N. Habegger and J.-B. Meilhan, *On the classification of links up to finite type*, Topology and Physics, Nankai Tracts Math., vol. 12, World Sci., Hackensack, NJ. [arXiv:0804.3548](#). [↑1.3](#), [5](#)
- [31] N. Habegger and G. Masbaum, *The Kontsevich integral and Milnor’s invariants*, Topology **39** (2000), 1253–1289. [Journal](#). [↑9.3](#)
- [32] K. Habiro, *Claspers and finite type invariants of links*, Geom. Topol. **4** (2000), 1–83. [↑1.3](#), [5](#), [1](#), [5.6](#), [5](#), [5](#), [5.11](#), [5](#), [5.12](#), [5](#), [5](#), [6](#), [7](#)
- [33] M. Hall, *The Theory of Groups*, The Macmillan Co., New York, 1959. [↑5.10](#)
- [34] B. Hartley, *Powers of the augmentation ideal in group rings of infinite nilpotent groups*, J. London Math. Soc. **25** (1982), 43–61. [↑5.10](#)
- [35] Y. Hashizume, *On the uniqueness of the decomposition of a link*, Osaka Math. J. **10** (1958), 283–300. [ProjectEuclid](#). [↑8](#), [8.9](#), [8](#)
- [36] L. Helme-Guizon, *The coefficients of the HOMFLYPT and the Kauffman polynomials are pointwise limits of Vassiliev invariants*, J. Knot Theory Ramifications **16** (2007), 203–215. [arXiv:math/0503185](#). [↑1.4](#)
- [37] J. Hillman, *Algebraic Invariants of Links*, Series on Knots and Everything, vol. 32; 2nd ed., vol. 52, World Sci., River Edge, NJ, 2002/2012. Chapter 1 available at [author’s webpage](#). [↑1.1](#), [1.6](#)
- [38] N. Ito and J. Yoshida, *On the four-term relation on Khovanov homology*. [arXiv:2202.08527](#). [↑1.1](#)
- [39] R. Kirby, *Problems in low-dimensional topology*, Geometric Topology (Athens, GA, 1993), AMS/IP Stud. Adv. Math., vol. 2.2, Amer. Math. Soc., Providence, RI, 1997, pp. 35–473. [C. Rourke’s webpage](#). [↑1.1](#)
- [40] P. Kirk and C. Livingston, *Vassiliev invariants of two component links and the Casson–Walker invariant*, Topology **36** (1997), 1333–1353. [Journal](#). [↑1.2](#), [2](#), [1.6](#), [4](#), [4.12](#), [10.9](#)
- [41] D. Kosanović, *Embedding calculus and grope cobordism of knots*. [arXiv:2010.05120](#). [↑1.1](#)
- [42] Thang Q. T. Le and J. Murakami, *Representation of the category of tangles by Kontsevich’s iterated integral*, Comm. Math. Phys. **168** (1995), 535–562. [ProjectEuclid](#). [↑9.3](#)
- [43] ———, *The universal Vassiliev–Kontsevich invariant for framed oriented links*, Compositio Math. **102** (1996), 41–64. [arXiv:hep-th/9401016](#). [↑9.3](#)
- [44] Thang T. Q. Le and Anh T. Tran, *On the AJ conjecture for knots*, Indiana Univ. Math. J. **64** (2015), 1103–1151. [arXiv:1111.5258](#). [↑1.1](#)
- [45] J. Levine, *The $\bar{\mu}$ -invariants of based links*, Differential Topology (Siegen, 1987), Lecture Notes in Math., vol. 1350, Springer, Berlin, 1988, pp. 87–103. [↑6.2](#)
- [46] ———, *The Conway polynomial of an algebraically split link*, KNOTS ’96 (Tokyo), World Sci. Publ., River Edge, NJ, 1997, pp. 23–29. [↑10.2](#), [10](#)
- [47] W. B. R. Lickorish, *An Introduction to Knot Theory*, Grad. Texts in Math., vol. 175, Springer-Verlag, New York, 1997. [↑3.4](#), [8.1](#), [10](#), [10.1](#), [10.5](#)
- [48] J. Lieberum, *The number of independent Vassiliev invariants in the Homfly and Kauffman polynomials*, Doc. Math. **5** (2000), 275–299. [arXiv:math/9806064](#). [↑1.4](#)
- [49] C. Livingston, *Enhanced linking numbers*, Amer. Math. Monthly **110** (2003), no. 5, 361–385. [MAA](#). [↑1.6](#), [10.2](#), [10.9](#)

- [50] G. Masbaum, *Matrix-tree theorems and the Alexander–Conway polynomial*, Invariants of Knots and 3-Manifolds (Kyoto, 2001), Geom. Topol. Monogr., vol. 4, Geom. Topol. Publ., Coventry, 2002, pp. 201–214. [arXiv:math/0211063](#). [↑10.2](#), [10](#)
- [51] G. Massuyeau, *Finite-type invariants of 3-manifolds and the dimension subgroup problem*, J. Lond. Math. Soc. **75** (2007), 791–811. [arXiv:math/0605497](#). [↑5](#), [5.8](#), [5](#), [5.9](#)
- [52] J.-B. Meilhan and A. Yasuhara, *Characterization of finite type string link invariants of degree < 5* , Math. Proc. Cambridge Philos. Soc. **148** (2010), 439–472. [arXiv:0904.1527](#). [↑1.3](#), [5](#)
- [53] S. A. Melikhov, *On maps with unstable singularities*, Topology Appl. **120** (2002), 105–156. [arXiv:math/0101047](#). [↑1.5](#), [3](#)
- [54] ———, *Colored finite type invariants and a multi-variable analogue of the Conway polynomial* (2003). [arXiv:math.GT/0312007v2](#) (an early preprint version of [58] and of the present paper). [↑1.6](#)
- [55] S. A. Melikhov, *Infinite-dimensional uniform polyhedra*. [arXiv:1109.0346](#). [↑3](#)
- [56] S. A. Melikhov, *Self C_2 -equivalence of two-component links and invariants of link maps*, J. Knot Theory Ram. (2018), 1842012. [arXiv:1711.03514](#). [↑1.6](#), [4](#)
- [57] ———, *Topological isotopy and Cochran’s derived invariants*, Topology, Geometry, and Dynamics — V. A. Rokhlin Memorial, Contemp. Math., vol. 772, Amer. Math. Soc., Providence, RI, 2021, pp. 249–266. [arXiv:2011.01409](#). [↑1.1](#)
- [58] S. A. Melikhov, *The two-variable Conway polynomial and Cochran’s derived invariants*. [arXiv:?](#). [↑1.3](#), [4.4](#), [54](#)
- [59] ———, *Is every knot isotopic to the unknot?*. [arXiv:?](#). [↑1.3](#), [1.6](#), [8.2](#), [10.15](#)
- [60] S. A. Melikhov and D. Repovš, *n -Quasi-isotopy. I. Questions of nilpotence*, J. Knot Theory Ram. **14** (2005), 571–602. [arXiv:math/0103113](#). [↑1.2](#), [1.1](#), [1.2](#), [1.5](#), [3](#), [1.6](#), [4](#), [3](#), [3.2](#), [3](#), [3.5](#), [3](#), [4.10](#), [4.12](#), [7](#), [10](#)
- [61] ———, *n -Quasi-isotopy. II. Comparison*, J. Knot Theory Ram. **14** (2005), 603–626. [arXiv:math/0103114](#). [↑1.2](#), [1.3](#), [3](#), [3.1](#), [3.3](#), [3](#), [7](#), [4.13](#), [8](#), [10.18](#)
- [62] S. A. Melikhov and R. V. Mikhailov, *n -Quasi-isotopy: III. Engel conditions*. [arXiv:math.GT/0201022](#). [↑3.1](#)
- [63] B. Mellor and D. Thurston, *On the existence of finite type link homotopy invariants*, J. Knot Theory Ramifications **10** (2001), 1025–1039. [arXiv:math/0010206](#). [↑10](#)
- [64] P. M. Melvin and H. R. Morton, *The coloured Jones function*, Comm. Math. Phys. **169** (1995), 501–520. [ProjectEuclid](#). [↑1.1](#)
- [65] R. Mikhailov and I. B. S. Passi, *Lower Central and Dimension Derives of Groups*, Lecture Notes in Math., vol. 1952, Springer, Berlin, 2009. [↑5](#)
- [66] J. Milnor, *Link groups*, Ann. Math. **59** (1954), 177–195. [↑4.12](#)
- [67] ———, *Isotopy of links*, Algebraic Geometry and Topology: A Symposium in Honor of S. Lefschetz, Princeton Univ. Press, Princeton, NJ, 1957, pp. 280–306. [↑1.1](#)
- [68] H. A. Miyazawa and A. Yasuhara, *Classification of n -component Brunnian links up to C_n -move*, Topology Appl. **153** (2006), 1643–1650. [arXiv:math/0412490](#). [↑2](#), [3](#), [4](#), [5.4](#)
- [69] K. Murasugi, *On Milnor’s invariant for links*, Trans. Amer. Math. Soc. **124** (1966), 94–110. [Journal](#). [↑10](#)
- [70] Y. Nakanishi and Y. Ohyama, *Delta link homotopy for two component links. III*, J. Math. Soc. Japan **55** (2003), 641–654. [J-STAGE](#). [↑1.6](#)
- [71] I. B. S. Passi, *Group Rings and their Augmentation Ideals*, Lecture Notes in Math., vol. 715, Springer, Berlin, 1979. [↑5](#)
- [72] V. V. Prasolov and A. B. Sossinsky, *Knots, Links, Braids and 3-Manifolds*, MCCME, Moscow, 1997; English transl. in Transl. Math. Mono., vol. 154, Amer. Math. Soc., Providence, RI, 1997. [↑10.4](#), [17](#)

- [73] D. J. S. Robinson, *A Course in the Theory of Groups*, 2nd ed., Grad. Texts in Math., vol. 80, Springer, New York, 1996. [↑5.10](#)
- [74] D. Rolfsen, *Isotopy of links in codimension two*, J. Indian Math. Soc. **36** (1972), 263–278. [↑7.1](#)
- [75] ———, *Some counterexamples in link theory*, Canadian J. Math. **26** (1974), 978–984. [Journal](#). [↑1.1](#), [1.3](#), [7](#)
- [76] ———, *Localized Alexander invariants and isotopy of links*, Ann. Math. **101** (1975), 1–19. [↑1.3](#)
- [77] ———, *Piecewise-linear I-equivalence of links*, Low-Dimensional Topology (Chelwood Gate, 1982), London Math. Soc. Lecture Note Ser., vol. 95, Cambridge Univ. Press, Cambridge, 1985, pp. 161–178. [↑1.1](#)
- [78] ———, *PL link isotopy, essential knotting and quotients of polynomials*, Canad. Math. Bull. **34** (1991), 536–541. [Journal](#). [↑1.4](#), [10.6](#)
- [79] T. Stanford, *Vassiliev invariants and knots modulo pure braid subgroups* (1998). [arXiv:math/9805092](#). [↑5](#), [5](#)
- [80] T. Tanaka, *The colored Jones polynomials of doubles of knots*, J. Knot Theory Ram. **17** (2008), 925–937. [↑1.1](#), [3.4](#)
- [81] K. Taniyama and A. Yasuhara, *Band description of knots and Vassiliev invariants*, Math. Proc. Cambridge Philos. Soc. **133** (2002), 325–343. [arXiv:math/0003021](#). [↑5](#), [4](#), [5.4](#)
- [82] L. Traldi, *Milnor’s invariants and the completions of link modules*, Trans. Amer. Math. Soc. **284** (1984), 401–424. [Journal](#). [↑10](#)
- [83] ———, *Conway’s potential function and its Taylor series*, Kobe J. Math. **5** (1988), 233–263. [Lafayette Repository](#). [↑1.4](#), [1.6](#), [10.6](#), [10.9](#)
- [84] V. A. Vassiliev, *Cohomology of knot spaces*, Theory of Singularities and its Applications, Adv. Soviet Math., vol. 1, Amer. Math. Soc., Providence, RI, 1990, pp. 23–69. [ResearchGate](#). [↑\(K\)](#), [4](#)
- [85] ———, *Complements of Discriminants of Smooth Maps: Topology and Applications*, Transl. Math. Mono., vol. 98, Amer. Math. Soc., Providence, RI, 1992/1994. Extended Russian ed., Fазis, Moscow, 1997. [↑\(K\)](#)
- [86] S. Willerton, *Vassiliev invariants and the Hopf algebra of chord diagrams*, Math. Proc. Cambridge Philos. Soc. **119** (1996), 55–65. [↑4](#), [4.6](#), [4](#)
- [87] S. Willetts, *A unification of the ADO and colored Jones polynomials of a knot*, Quantum Topol. **13** (2022), 137–181. [arXiv:2003.09854](#). [↑1.4](#)
- [88] ———, *Vassiliev invariants for knots from the ADO polynomials*. [arXiv:2105.09786](#). [↑1.4](#)
- [89] Ka Ho Wong, *Volume conjecture, geometric decomposition and deformation of hyperbolic structures*. [arXiv:1912.11779](#). [↑1.1](#), [3.4](#)
- [90] A. Yasuhara, *Brunnian local moves of knots and Vassiliev invariants*, Fund. Math. **190** (2006), 289–297. [arXiv:math/0412489](#). [↑5](#)
- [91] J. Yoshida, *On Khovanov homology and Vassiliev theory*. Kyoto University Research Information Repository, [Preprint No. 2191-03](#). [↑1.1](#)
- [92] Hao Zheng, *Proof of the volume conjecture for Whitehead doubles of a family of torus knots*, Chinese Ann. Math. Ser. B **28** (2007), no. 4, 375–388. [arXiv:math/0508138](#). [↑1.1](#), [3.4](#)

STEKLOV MATHEMATICAL INSTITUTE OF RUSSIAN ACADEMY OF SCIENCES, MOSCOW, RUSSIA
 Email address: melikhov@mi-ras.ru