

IS EVERY KNOT ISOTOPIC TO THE UNKNOT?

SERGEY A. MELIKHOV

ABSTRACT. In 1974, D. Rolfsen asked: Is every knot in S^3 isotopic (=homotopic through embeddings) to a PL knot or, equivalently, to the unknot? In particular, is the Bing sling isotopic to a PL knot? We show that the Bing sling is not isotopic to any PL knot:

- (1) by an isotopy which extends to an isotopy of 2-component links with $\text{lk} = 1$;
- (2) through knots that are intersections of nested sequences of solid tori.

There are also stronger versions of these results. In (1), the additional component may be allowed to self-intersect, and even to get replaced by a new one as long as it represents the same conjugacy class in $G/[G', G'']$, where G is the fundamental group of the complement to the original component. In (2), the “solid tori” can be replaced by “boundary-link-like handlebodies”, where a handlebody $V \subset S^3$ of genus g is called boundary-link-like if $\pi_1(S^3 \setminus \bar{V})$ admits a homomorphism to the free group F_g such that the composition $\pi_1(\partial V) \rightarrow \pi_1(S^3 \setminus \bar{V}) \rightarrow F_g$ is surjective.

1. INTRODUCTION

Fifty years ago¹ D. Rolfsen [78] posed the following problem.

Rolfsen’s Problem. “All PL knots are isotopic to one another.² Is this true of wild knots? A knot in the 3-space which is not isotopic to a tame knot would have to be so wild as to fail to pierce a disk at each of its points.³ The ‘Bing sling’ [[12]] is such a candidate.”

The “Bing sling” is R. H. Bing’s example of a wild knot that “pierces no disk” [12]. It is the intersection of a chain of solid tori $T_1 \supset T_2 \supset \dots$, where T_1 is unknotted in S^3 and each T_{i+1} is a suitable regular neighborhood of a certain PL knot K_i in T_i (see Figure 1). We discuss this construction and its outcome in more detail in §5.

Until recently, Rolfsen’s Problem has been mostly approached with geometric methods. D. Gillman constructed a knot that pierces no disk but lies on a disk (and hence is isotopic to the unknot); but he also proved that no subarc of the Bing sling lies on a disk [35]. M. Brin constructed a knot that at each point is locally equivalent to the

¹In his AMS Math Review of [18], published in 1985, R. Daverman speaks of this problem as “being around for almost 20 years”. Although it is clear from [70] (1957), [32; p. 171] (1962), [35] (1964), [41] (1966) that Rolfsen’s problem was probably known to a number of people long before it was stated explicitly in Rolfsen’s 1974 paper, I’m not aware of its earlier explicit appearances in the literature.

²See for instance Figure 1.1 in the Burde–Zieschang textbook [20].

³This assertion of Rolfsen is easy to prove in the case where the disk is tame (see Lemma 2.1). A proof of the general case was found recently by R. Ancel (see [4] for a sketch).




FIGURE 1. The Bing sling.

Bing sling, but is isotopic to the unknot [18]. The most striking result was obtained by C. Giffen:

Giffen's Theorem. (see Appendix A) *Every knot in the solid torus T which represents a generator of $H_1(T)$ is I -equivalent to the core circle of T .*

Two m -component (possibly wild) links in a 3-manifold M are called *I -equivalent* if they cobound a topological embedding of m disjoint annuli in $M \times I$. (The embedding is not assumed to be locally flat, so this is not the same as topological concordance.)

An easy, though not entirely obvious (cf. [81; Question 1]) consequence of Giffen's Theorem is

Corollary. (see Appendix A) *Every knot in S^3 is I -equivalent to the unknot.*

The following was shown in a recent paper by the author.

Theorem 1.1 ([65]). *Not every link in S^3 is I -equivalent to a PL link.*

Corollary. *Not every link in S^3 is isotopic to a PL link.*

The proof of Theorem 1.1 in [65] can be called geometric. It is based on Cochran's derived invariants β_i , which are extended to I -equivalence invariants of topological links

by using infinite *homological* Seifert surfaces. By using these, it is shown in [65] that there exist uncountably many 2-component links of linking number 0, with unknotted first component, which are not I -equivalent to each other.

By contrast, the proofs of the main results of the present paper can be called algebraic. They are based on the following result (and its version for colored links):

Theorem 1.2 ([67; Corollary 1.2]). *Let v be a finite type invariant of PL links which is invariant under PL isotopy. Then*

(a) *v assumes the same value on all sufficiently close PL C^0 -approximations of any given topological link;*

(b) *the extension of v by continuity to topological links is an invariant of isotopy.*

The proof of Theorem 1.2 in [67] is in turn based on the notion of n -quasi-isotopy, which was introduced by the author [63] but can be traced back to a number of older constructions (from the Penrose–Whitehead–Zeeman–Irwin trick to Casson handles), most notably to the Homma–Bryant proof of the Chernavsky–Miller codimension three PL approximation theorem (see [63; Remarks (i)–(iii) in §1.5]).

Theorem A. *Let \mathcal{B} be the Bing sling, and let Q be a PL knot in $S^3 \setminus \mathcal{B}$ with $\text{lk}(Q, \mathcal{B}) = 1$. Then (Q, \mathcal{B}) is not isotopic to any PL link.*

While this is a special case of the results to follow, we single it out since its proof is considerably easier. Apart from Theorem 1.2 it uses only the Conway polynomial ∇_L (and some of its properties). Or rather its reduced version $\nabla_{(K_1, K_2)}(z) / \nabla_{K_1}(z) \nabla_{K_2}(z)$, which is easily seen to be an invariant of PL isotopy.

A minor elaboration on the proof of Theorem A yields

Theorem B. *The Bing sling is not isotopic to any PL knot through knots that are intersections of nested sequences of solid tori.*

Not every wild knot is the intersection of a nested sequence of solid tori. There are at least four proofs of this fact in the literature.

- (1) J. Milnor [72; Theorem 10 and assertions (1) and (2) on p. 303] constructed a 2-component wild link which is isotopic to the unlink, but not through links satisfying property (S) , where a 2-component link (Q, \mathcal{K}) is said to have *property (S)* if \mathcal{K} lies in a solid torus T disjoint from Q and such that \mathcal{K} represents a generator of $H_1(T)$. His proof is based on the fundamental group. But it is easy to see that (Q, \mathcal{K}) satisfies property (S) if \mathcal{K} is the intersection of a nested sequence of solid tori (see Lemma 11.2).
- (2) J. McPherson [61] proved that Fox’s “remarkable simple closed curve” is not the intersection of any nested sequence of solid tori. The proof is based on the observation (found already in [19]) that the Alexander module of a wild knot can have rank > 1 .
- (3) S. Kojima and M. Yamasaki [53; assertions (1), (3) of Theorem 4 and the first line of §7] proved that a certain 2-component wild link whose first component is

PL (a variation of Milnor’s wild link) is isotopic to the unlink but not F-isotopic to any PL link. In particular, it does not satisfy Milnor’s property (S). The proof is based on the observation that the knot module of a wild knot need not be a torsion module (which is equivalent to the observation mentioned in the previous item, see [48; Proposition 7.3.4(1)]).

- (4) Since the first component of the link in Theorem 1.1 is PL, it follows from Giffen’s Theorem and Theorem 1.1 that it does not satisfy Milnor’s property (S).

Theorem A’. *The Bing sling is not isotopic to any PL knot by any isotopy which extends to an isotopy of 2-component links with linking number 1.*

This is like Theorem A, except that the knot Q and the terminal knot of the isotopy which Q undergoes are now allowed to be wild. This may seem like a minor improvement, but here is one application. It is well-known that the Whitehead link is symmetric (i.e. its components may be interchanged by an ambient isotopy); it is easy to check that its counterparts among links of linking number 1, the Mazur link and the fake Mazur link (see Figure 7 below and comments in footnote 22 to Figure 6 below) are also symmetric. (For a symmetric diagram of the Mazur link see [1; Figure 4].) Even though the Bing sling is basically made up of copies of the fake Mazur link (see §5), one consequence of Theorem A’ is

Corollary. *Let \mathcal{B} be the Bing sling, and let Q be a PL knot in $S^3 \setminus \mathcal{B}$ with $\text{lk}(Q, \mathcal{B}) = 1$. Then (Q, \mathcal{B}) is not isotopic to (\mathcal{B}, Q) .⁴*

But the real difference between Theorems A and A’ is that a minor extension of the proof of the latter yields

Theorem C. *The Bing sling is not isotopic to any PL knot by any isotopy which extends to a link homotopy of 2-component links with linking number 1.*

We recall that a *link homotopy* is a homotopy of a link in which the link components remain disjoint.

The proofs of Theorems A’ and C are based on a version of Theorem 1.2 for *colored* finite type invariants and on the 2-variable Conway polynomial $\nabla_L(u, v)$ introduced in [68]. More precisely we make use of a 1-parameter subfamily β_1, β_2, \dots of the coefficients of its reduced version $\nabla_{(K_1, K_2)}(u, v) / \nabla_{K_1}(u) \nabla_{K_2}(v)$, which in the case $\text{lk} = 0$ coincides with the sequence of Cochran’s derived invariants. However, in our case $\text{lk} = 1$ these integers β_i exhibit a much subtler geometric behavior than the original Cochran’s invariants, as we will see shortly.

Although the proofs of Theorems A, B, C were written up only in 2023, these results were announced in my 2005 talk at the conference “Manifolds and their Mappings” in Siegen, Germany. A question which I asked in the same talk was recently answered by A. Zastrow [97], [98]:

⁴Such an isotopy, if it existed, would provide an elegant solution to Rolfsen’s Problem.

Zastrow’s Example. *There exists an isotopy from the unknot to itself which does not extend to any link homotopy of 2-component links with linking number 1.*

We include a short self-contained review of Zastrow’s Example (see Example 2.2), as it will be used in the proof of Theorem E.

To counter the difficulty highlighted by Zastrow’s Example, we consider relations weaker than link homotopy.

The finite derived series of a group G is usually defined by $G^{(0)} = G$ and $G^{(n)} = [G^{(n-1)}, G^{(n-1)}]$ for positive integer n . But sometimes it is expanded to include half-integer terms, defined by $G^{(n+0.5)} = [G^{(n)}, G^{(n-1)}]$ for positive integer n . Speaking geometrically, it is well-known and easy to see that a loop $\varphi: S^1 \rightarrow X$ represents an element of $\pi_1(X)^{(n)}$ if and only if φ bounds a map $\Gamma \rightarrow X$ of a grope of height⁵ n (see [25; Definition 7.9 and Lemma 7.10], where gropes of half-integer height are included).

We call 2-component links n -bordant if they are equivalent under the equivalence relation generated by the following two relations:

- (1) link homotopy which restricts to an isotopy on the second component;
- (2) $(Q, \mathcal{K}) \sim (Q', \mathcal{K})$ if Q and Q' represent the same conjugacy class in $G/G^{(n)}$, where $G = \pi_1(S^3 \setminus \mathcal{K})$.

Obviously, links (Q_0, \mathcal{K}_0) and (Q_1, \mathcal{K}_1) are n -bordant if and only if they are related by a sequence of link homotopies which extend consecutive time intervals of a single isotopy \mathcal{K}_t between \mathcal{K}_0 and \mathcal{K}_1 , with “jumps” between consecutive link homotopies related by the relation (2). Such a sequence of link homotopies will be called an n -bordism between (Q_0, \mathcal{K}_0) and (Q_1, \mathcal{K}_1) .

By using 2-bordism in place of link homotopy we can avoid Zastrow’s Example:

Theorem E’. *If L and L' are links of linking number 1, then every isotopy between their second components extends to a 2-bordism between L and L' .*

As discussed in §3, from the geometric viewpoint this result was initially very surprising and hard to believe. Eventually, there are 3 independent proofs, none of them particularly hard (one based on the commutator identities, another on the knot module, and a geometric one, based on Seifert surfaces).

On the other hand, Corollary to Theorem 1.1 holds with 2-bordism in place of isotopy:

Theorem D’. *There exists a 2-component link in S^3 with linking number 0 which is not 2-bordant to any link whose second component is PL .*

Like the proof of Theorem 1.1, the proof of Theorem D’ is based on Cochran’s derived invariants β_i . So one could hope to obtain a similar result for links of linking number 1 by using their β_i . One issue with this approach (which was one reason why it took me 18 years to find time for writing up the results announced in 2005) was that while the geometry of Cochran’s original invariants β_i was well-understood since the 1980s, virtually nothing was known about the geometry of the β_i for links of linking number

⁵Not to be confused with gropes of class n , which are similarly related to the lower central series of G .

1. But finally some basic understanding has been gained [68; Theorem D]), and as its application we obtain the following result.

Theorem D. *The Bing sling is not isotopic to a PL knot by any isotopy which extends to a 2.5-bordism of 2-component links with linking number 1.*

A more precise version of Theorem D says that the power series $C_L(z) = \sum_{i=1}^{\infty} \beta_i z^i$ is rational if the second component of L is PL and not rational if it is the Bing sling, but $C_L - C_{L'}$ is rational if L and L' are 2.5-bordant links of linking number 1. But in fact, $C_L - C_{L'}$ is also rational if L and L' are 2-bordant links of linking number 0 — and this is where Theorem D' comes from. However, Theorem E' and Theorem 14.2 imply that $C_L - C_{L'}$ need not be rational for 2-bordant links L and L' of linking number 1.⁶ A further blow is provided by the following example.

Theorem E. *There exists an isotopy from the unknot to itself which does not extend to any 2.5-bordism of 2-component links with linking number 1.*

Remark 1.3. The proof of Theorem D shows that it can be improved by replacing $G/[G', G'']$, which occurs in the definition of 2.5-bordism, with $G/[G', T]$, where T is preimage in G' of the torsion submodule of the link module G'/G'' . Unfortunately, Theorem E can also be improved in the same way (see Proposition 4.6).

Let us now turn to our final result, which applies the techniques of Theorem D to the setting where we refrain from inserting a new component, like in Theorem B. As noted above, a wild knot need not be the intersection of a nested sequence of solid tori. But it is well-known that not only every wild knot, but every 1-dimensional compact ANR subset X of S^3 is the intersection of a nested sequence of handlebodies [60; Lemmas 2, 3] (see also [52; Theorem]).⁷ Before we discuss these nested sequences, let us recall one classical fragment of the theory of knotted handlebodies.

Theorem 1.4. *Let V be a genus g PL handlebody in S^3 and let $W = \overline{S^3 \setminus V}$. The following are equivalent:*

- (1) W admits a retraction onto a PL embedded wedge of g circles;
- (2) $\pi_1(W)$ admits an epimorphism onto the free group F_g ;
- (3) $\pi_1(W)/\gamma_\omega \pi_1(W)$ is a free group,⁸

⁶This observation refutes a premature announcement from my 2005 talk in Siegen. At that time I wrongly believed that Cochran's theorem on additivity of the β_i under band connected sum [24; Theorem 5.4] extends to links of any linking number.

⁷Indeed, X is certainly the intersection of a nested sequence of closed polyhedral neighborhoods $M_1 \supset M_2 \supset \dots$. Each M_i has a handle decomposition with 0-, 1- and 2-handles. To eliminate the 2-handles it suffices to "puncture" them, i.e. to make their cocore 1-disks disjoint from \mathcal{K} . This can always be done since X is tame in the sense of Shtan'ko (or in other words has embedding dimension 1, see [86]) by [60; Lemma 3] (see also [16; Satz 4], [87; Theorem 4]). In fact, a 1-dimensional compact subset X of S^3 is the intersection of a nested sequence of *thin* handlebodies (i.e. such that for each $\epsilon > 0$ almost all of the handlebodies admit ϵ -retractions onto graphs) if and only if X is tame in the sense of Shtan'ko (see [17; p. 176]).

⁸ $\gamma_\omega G$ denotes the intersection of the finite lower central series $\gamma_n G$ of the group G , where $\gamma_1 G = G$ and $\gamma_{n+1} G = [\gamma_n G, G]$.

- (4) W contains pairwise disjoint properly⁹ PL embedded surfaces $\Sigma_1, \dots, \Sigma_g$ whose union is non-separating in W .

Proof. This is similar to Smythe's characterizations of homology boundary links [89]. Formally, (1) \Leftrightarrow (2) is a special case of [46; Corollary to Theorem 2] and (2) \Leftrightarrow (4) is a special case of [45; Theorem 2.1]. Finally, (2) \Rightarrow (3) follows from [46; Corollary to Theorem 1] and the converse is obvious (since the rank of a free group is determined by its abelianization). \square

Theorem 1.5. *Let V be a genus g PL handlebody in S^3 and let $W = \overline{S^3 \setminus V}$. The following are equivalent:*

- (1) W admits a retraction onto a PL embedded wedge of g circles contained in ∂W ;
- (2) $\pi_1(W)$ admits a homomorphism to the free group F_g such that the composition $\pi_1(\partial W) \rightarrow \pi_1(W) \rightarrow F_g$ is surjective;
- (3) $\pi_1(W)/\gamma_\omega \pi_1(W)$ is a free group and $\pi_1(\partial W)$ maps surjectively onto it;
- (4) W contains pairwise disjoint properly PL embedded surfaces $\Sigma_1, \dots, \Sigma_g$ such that $\partial \Sigma_1 \cup \dots \cup \partial \Sigma_g$ is non-separating in ∂W ;
- (5) W admits a map onto a handlebody U extending a homeomorphism $\partial W \rightarrow \partial U$.

As noted in [8], using Perelman's solution of the Poincaré Conjecture one can reformulate some of the equivalent conditions of Theorem 1.5 in a more rigid way (see Theorem 17.12). A further condition equivalent to those of Theorem 1.5, in terms of Dehn surgery, is given in [76].

Proof. This is similar to Smythe's characterizations of boundary links [89]. More formally, a proof of (4) \Leftrightarrow (5) is discussed in [55; Theorem 2], and (5) \Leftrightarrow (1) is established in [46; Theorem 3], whose proof proceeds by showing that (5) \Rightarrow (1) \Rightarrow (4) \Rightarrow (5). The equivalence of (1) and (3) follows from that in Theorem 1.4 along with [44; Theorems 2, 3]. The equivalence of (2) and (3) follows from the proof of that in Theorem 1.4. \square

Knotted handlebodies satisfying any of the equivalent conditions of Theorem 1.5 will be called *boundary-link-like* handlebodies; while those satisfying the conditions of Theorem 1.4 can be called *HBL-like*.

Examples. It is easy to see that every genus 1 handlebody is boundary-link-like. Moreover, given a boundary-link-like handlebody, it will remain boundary-link-like if we add any local knot to it. A regular neighborhood in \mathbb{R}^3 of a connected graph in $\mathbb{R}^2 \times \{0\}$ is always a boundary-link-like handlebody. More generally, if $G \subset \mathbb{R}^3$ is an embedded graph which contains a boundary link L whose complement in G is a tree which is disjoint from some collection of disjoint Seifert surfaces for L , then a regular neighborhood of G is a boundary-link-like handlebody. (See [90; Example 4.4] for a specific example.)

Examples of handlebodies that are not boundary-link-like but HBL-like; or not HBL-like can be found in [55], [46], [44], [90], [8; §8], [38].

⁹An embedding $g: P \rightarrow Q$ between manifolds with boundary is called *proper* if $g^{-1}(\partial Q) = \partial P$.

Theorem F. *The Bing sling is not isotopic to any PL knot through knots that are intersections of nested sequences of boundary-link-like handlebodies.*

The proof consists of two steps.

- (1) If a knot \mathcal{K} is the intersection of a nested sequence of boundary-link-like handlebodies, then its knot module¹⁰ is a torsion module (Corollary 18.2).
- (2) The Bing sling is not isotopic to any PL knot through knots whose knot module is a torsion module (Corollary 16.3).

The proof of (1) in fact works under a weaker hypothesis. Instead of assuming that each handlebody V in the given nested sequence is boundary-link-like, it suffices to assume that $[\mathcal{K}] \in H_1(V)$ can be represented by a PL knot $K \subset \partial V$ which bounds a Seifert surface Σ in $W := \overline{S^3 \setminus V_i}$ such that each knot in Σ is null-homologous in W . However, we show that in the case where V has genus 2, this apparently weaker hypothesis is in fact equivalent to the hypothesis that V is boundary-link-like. This provides yet another characterization of boundary-link-like handlebodies in the case of genus 2 (Theorem 17.10).

Remark 1.6. J. McPherson [61] constructed for any $g \geq 1$ a wild knot which is the intersection of a nested sequence of handlebodies of genus $g + 1$, but not of those of genus g . In fact, if \mathcal{K} is a wild knot whose knot module has rank ≥ 1 and which contains a tame arc, then, as noted in [19], a connected sum \mathcal{K}_∞ of a null-sequence of copies of \mathcal{K} has knot module of infinite rank; then by [61; Theorem 2] \mathcal{K}_∞ is not the intersection of any nested sequence of handlebodies of bounded genus.

2. SOME BASIC EXAMPLES

Lemma 2.1. *Let K be a knot in S^3 which meets an embedded PL disk D^2 in a single point p of D^2 so that D^2 locally separates K at p .¹¹ Then K is isotopic to the unknot.*

Proof. We may identify (S^3, D^2) with the quotient of $(B^3, \partial B^3)$ by $\rho|_{\partial B^3}$, where B^3 is the unit 3-ball in \mathbb{R}^3 and $\rho: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is the reflection $(x, y, z) \mapsto (-x, y, z)$. The preimage of K in B^3 is an arc J_0 with endpoints in ∂B^3 , which may be assumed to be $(\pm 1, 0, 0)$. Let $J_1 = [-1, 1] \times \{0\} \times \{0\}$ and for $0 < t < 1$ let J_t be the union of $([-1, -1 + t] \cup [1 - t, 1]) \times \{0\} \times \{0\}$ with the image of J_0 under the homothety $H_t: \mathbb{R}^3 \rightarrow \mathbb{R}^3, v \mapsto (1 - t)v$. It is easy to see that the family of arcs J_t is parameterized by an isotopy and the image of this isotopy in S^3 is an isotopy from K to an unknot. \square

Example 2.2 (Zastrow's Example). Let $K_{1/2}$ be Rolfsen's wild knot [78; Figure 1], which can be described as the visible outline of a standard drawing of Alexander's horned sphere (cf. [30; Figure 2.11]) — in other words, an equatorial circle of the horned sphere which passes through all its wild points. Thus $K_{1/2}$ has a Cantor set C of wild

¹⁰That is, $H_1(\widetilde{S^3 \setminus \mathcal{K}})$. It is closely related to the Alexander module, see [48; Proposition 7.1.2(1)].

¹¹This means that for every sufficiently small neighborhood U of p , not all connected components of $(K \cap U) \setminus \{p\}$ lie in the same component of $U \setminus D^2$. Let us note that this condition implies that $p \notin \partial D^2$.

points, and its complement $K_{1/2} \setminus C$ consists of countably many tame arcs. Let A_0 and A_1 be the two “longest” ones of all these arcs, and let D_i be a small embedded PL disk which meets $K_{1/2}$ in a single point $x \in A_i$ so that D_i locally separates K at x . Let us orient each D_i so that $\text{lk}(K_{1/2}, \partial D_i) = +1$. By applying the construction in the proof of Lemma 2.1 to the disk D_1 we obtain an isotopy from $K_{1/2}$ to an unknot, which will be the second half $K_{[1/2, 1]}$ of the desired isotopy K_t . Its first half $K_{[0, 1/2]}$ will be the time-reversed version of the isotopy from $K_{1/2}$ to an unknot yielded by the same construction in the proof of Lemma 2.1, this time applied to the disk D_0 . Thus K_i for $i = 0, 1$ is an unknot disjoint from ∂D_i , and moreover $(K_i, \partial D_i)$ is the Hopf link.

Lemma 2.3. $X := S^3 \times I \setminus \bigcup_{t \in I} K_t \times \{t\}$ deformation retracts onto $(S^3 \setminus K_{1/2}) \times \{\frac{1}{2}\}$.

Proof. In the notation of the proof of Lemma 2.1, it suffices to construct a deformation retraction r_t of $B^3 \times I \setminus \bigcup_{t \in I} J_t \times \{t\}$ onto $(B^3 \setminus J_0) \times \{0\}$ which restricts to a $(\rho \times \text{id}_I)$ -equivariant deformation retraction of $(\partial B^3 \setminus \partial J_0) \times I$ onto $(\partial B^3 \setminus \partial J_0) \times \{0\}$. (Note that $\partial J_t = \partial J_0$ for all $t \in I$.) It can be defined by $r_t(v, 1-s) = (\min(\frac{st}{s}, \frac{1}{\|v\|})v, 1-s_t)$, where s_t denotes the linear motion from $s_0 = s$ to $s_1 = 1$, that is, $s_t = (1-t)s + t$. Let us note that r_t is the restriction of a deformation retraction R_t of $B^3 \times I \setminus \{0\} \times \{1\}$ onto $B^3 \times \{0\}$ which is defined by the same formula. This R_t also restricts to a deformation retraction of the cone $C := \bigcup_{t \in [0, 1]} (1-t)\mathring{B}^3 \times \{t\}$ onto the open unit ball $\mathring{B}^3 \times \{0\}$, and to a deformation retraction of $B^3 \times I \setminus (C \cup \{0\} \times \{1\})$ onto $\partial B^3 \times \{0\}$. \square

Suppose that there exists a homotopy J_t of a knot J_0 such that $\text{lk}(J_0, K_0) = 1$ and $J_t \cap K_t = \emptyset$ for each $t \in I$. Since K_0 and K_1 are unknots, we may assume that $J_0 = \partial D_0$ and $J_1 = \partial D_1$. The homotopy $J_t \times \{t\}$ between $J_0 \times \{0\}$ and $J_1 \times \{1\}$ has its values in X and extends by using the cylinders $J_0 \times [0, \frac{1}{2}]$ and $J_1 \times [\frac{1}{2}, 1]$ (which also lie in X) to a homotopy J_t^+ with values in X between $J_0 \times \{\frac{1}{2}\}$ and $J_1 \times \{\frac{1}{2}\}$. The image of J_t^+ under the deformation retraction then yields a homotopy J'_t in $S^3 \setminus K_{1/2}$ between J_0 and J_1 . The image of J'_t , being disjoint from $K_{1/2}$, is in particular disjoint from its subset C , so it must also be disjoint from $K_{1/2} \cup N_n$ for some neighborhood N_n of C , which can be chosen to be a union of 2^n disjoint 3-balls.

Now let us recall that $K_{1/2}$ was an equatorial circle of the horned sphere, which in turn is the boundary of Alexander’s horned ball B . If H is an unknotted 1-handle in B connecting its polar caps and disjoint from the ball N_0 , then it is easy to see that $(B \cup N_n) \setminus H$ deformation retracts onto $K_{1/2} \cup N_n$. Moreover, the deformation retraction extends to a homotopy of S^3 which restricts to an isotopy on the complement of $(B \cup N_n) \setminus H$. Hence $\pi_1(S^3 \setminus (K_{1/2} \cup N_n)) \simeq \pi_1(S^3 \setminus ((B \cup N_n) \setminus H)) \simeq \mathbb{Z} * \pi_1(S^3 \setminus (B \cup N_n))$, where the free factor \mathbb{Z} is generated by the class x of a loop passing through the hole H drilled in the horned ball B . Let also $m \in \pi_1(S^3 \setminus (B \cup N_n))$ be the image of the class of a meridian of the solid torus $B \cup N_0$. Then J_0 and J_1 are in the free homotopy classes of x and xm respectively. If they are freely homotopic, then $xm = x^g$ for some $g \in \pi_1(S^3 \setminus (B \cup N_n))$, or equivalently $m = x^{-1}x^g$. Thus m belongs to the normal closure of the free factor \mathbb{Z} of $\mathbb{Z} * \pi_1(S^3 \setminus (B \cup N_n))$. On the other hand, m in fact belongs to

the other free factor $\pi_1(S^3 \setminus (B \cup N_n))$. It follows from the normal form theorem for free products (see e.g. [39; §1.2] or [57; Theorem 4.1]) that this can happen only if $m = 1$ [57; Exercise 4.1.4]. But it is well-known that the image of m in $\pi_1(S^3 \setminus B)$ is nontrivial [14].

Example 2.4. It is not hard to show that every PL isotopy of the first component of the Hopf link extends to a PL isotopy of the entire link (see [43; Theorem 2]; compare [97; Example 2.2]). On the other hand:

Proposition 2.5. *For $n = 0, 1$ there exists a PL link of linking number n and a PL isotopy of its first component which does not extend to a topological isotopy of the link.*

Proof for $n = 0$. The desired link consists of any nontrivial PL knot K_0 (for example, the trefoil) and any PL knot J_0 in $S^3 \setminus K_0$ which is null-homologous but not null-homotopic in $S^3 \setminus K_0$. Let K_t be the usual PL isotopy from K_0 to an unknot K_1 (given, for instance, by a PL version of the proof of Lemma 2.1, where the round ball B^3 is replaced by the cube $[-1, 1]^3$). Suppose that there exists a homotopy J_t such that $J_t \cap K_t = \emptyset$ for each $t \in I$. Since K_1 is the unknot and J_1 is null-homologous in $S^3 \setminus K_1$, it is null-homotopic in $S^3 \setminus K_1$. Hence $J_0 \times \{0\}$ is null-homotopic in $X := S^3 \times I \setminus \bigcup_{t \in [0, 1]} K_t \times \{t\}$. But it is easy to see that X deformation retracts onto $(S^3 \setminus K_0) \times \{0\}$ (by a version of the construction in the proof of Lemma 2.3 where B^3 is replaced by $[-1, 1]^3$ and hence the Euclidean norm by the l_∞ norm). Hence J_0 is null-homotopic in $S^3 \setminus K_0$, which is a contradiction. \square

Proof for $n = 1$. This is similar to the proof for $n = 0$, if we additionally use the following well-known lemma.

Lemma 2.6. *There exists a PL knot K and a PL knot J in $S^3 \setminus K$ which has linking number 1 with K but is not freely homotopic to a meridian¹² of K .*

There are several proofs of this Lemma in the literature: a geometric proof in [92; Theorem 3.9] and several algebraic proofs in [88]. (See also Example 4.4 below, which provides a not so easy proof of a stronger assertion.) For completeness, here is a simplified exposition of a proof found in [88; p. 5 of the arXiv version].

Proof. Let K be the trefoil knot. A Wirtinger presentation of $\pi_1(S^3 \setminus K)$ is

$$\langle x, y, z \mid z = x^{-1}yx, y = z^{-1}xz, x = y^{-1}zy \rangle.$$

As usual, one of the relations is redundant. By expressing z from the first relation and substituting for z in the second relation, we simplify the presentation as $\langle x, y \mid xyx = yxy \rangle$. From the latter relation we get $(xy)(xy)(xy) = (xyx)(yxy) = (xyx)(xyx)$, or $p^3 = q^2$, where $p = xy$ and $q = xyx$. Conversely, we can recover x and y as $x = p^{-1}q$ and $y = q^{-1}p^2$, so the presentation further simplifies as $\langle p, q \mid p^3 = q^2 \rangle$.

¹²A meridian of K is the boundary of a small embedded PL 2-disk which intersects K in precisely one point and locally separates K at this point.

This group surjects onto free product $(\mathbb{Z}/p) * (\mathbb{Z}/q) = \langle p, q \mid p^3 = q^2 = 1 \rangle$. We will use the well-known theorem about normal form of cyclically reduced elements of a free product (see e.g. [57; Theorem 4.2]). The element $x^2y^{-1} = p^{-1}qp^{-1}qp^{-2}q$ of the free product is cyclically reduced and alternates 6 times between the free factors \mathbb{Z}/p and \mathbb{Z}/q . Hence it is not conjugate to the meridian $x = p^{-1}q$ (which alternates only 2 times between the free factors), and therefore it is not conjugate to any meridian (because all meridians are conjugate to each other). Hence any knot J in the complement of the trefoil K representing the conjugacy class of x^2y^{-1} is not freely homotopic to a meridian of K . \square

\square

Example 2.7. The following theorem was proved in [63; Theorem 1.3 + the proof of Theorem 2.2]: there exists a PL knot in the solid torus $S^1 \times D^2$ which is homotopic, but not topologically isotopic to the core circle (by an isotopy within the solid torus).

3. PROOF OF THEOREM E'

Theorem E' follows from Lemma 3.1 and Theorem 3.2.

Lemma 3.1. *Let X_t be an isotopy of a knot, $t \in [0, 1]$. Then there exist numbers $0 = t_0 < \dots < t_n = 1$ and PL knots K_0, \dots, K_{n-1} such that each K_i is disjoint from X_t for all $t \in [t_i, t_{i+1}]$, and also $\text{lk}(K_i, X_{t_i}) = 1$.*

Moreover, the t_i may be chosen to belong to a given dense subset S of $[0, 1]$.

Proof. If S is not given, let $S = \mathbb{Q} \cap [0, 1]$. For each $s \in S$ let Q_s be some PL knot disjoint from X_s and such that $\text{lk}(Q_s, X_s) = 1$. Then there exists an $\varepsilon_s > 0$ such that Q_s is disjoint from X_t for each $t \in (s - \varepsilon_s, s + \varepsilon_s) \cap [0, 1]$. Since S is dense in $[0, 1]$, the intervals $J_s := (s - \varepsilon_s, s + \varepsilon_s) \cap [0, 1]$ for $s \in S$ form an open cover of $[0, 1]$. Then it has a finite subcover J_{s_1}, \dots, J_{s_n} . By discarding those of the J_{s_i} that can be discarded we may assume that none of the J_{s_i} can be discarded. Then each $J_{s_i} \cap J_{s_{i+1}}$ is nonempty; let t_i be some point in it. It remains to set $t_0 = 0, t_n = 1$ and $K_i = Q_{s_i}$. \square

Theorem 3.2. *Let \mathcal{K} be a knot in S^3 and let Q_1, Q_2 be PL knots in $S^3 \setminus \mathcal{K}$ such that $\text{lk}(Q_i, \mathcal{K}) = 1$. Then Q_1 and Q_2 represent the same conjugacy class in the quotient of $\pi_1(S^3 \setminus \mathcal{K})$ modulo the second commutator subgroup.*

Geometrically, this theorem is very surprising.¹³ In the geometric language it says that the connected sum $Q_1 \#_b (-Q_2)$ along some band disjoint from \mathcal{K} bounds a singular grope of height 2 in $S^3 \setminus \mathcal{K}$. It is not hard to construct a wild knot \mathcal{K} , using Fox–Artin wild arcs or Fox’s “remarkable simple closed curve”, which has two wild points, and its two tame arcs A, B are “linked” with each other so that (in contrast to Zastrow’s example, see Example 2.2) any meridian of A visually appears to be highly unlikely to

¹³The author has tried to disprove it for a week for a certain wild knot K by finding an appropriate epimorphism of $\pi_1(S^3 \setminus K)$ onto a finite group, and exhausted all finite groups of order ≤ 100 in this search.

cobound a grope of height 2 with any meridian of B . But Theorem 3.3 says that such a grope does exist!¹⁴

Lemma 3.3. *If G is a metabelian group with $G/G' \simeq \mathbb{Z}$, and $a, b \in G$ are such that the abelianization map $G \rightarrow G/G'$ sends a and b to the same generator of \mathbb{Z} , then a is conjugate to b .*

Proof. The idea of this proof comes from [7; proof of Theorem 2.3].

Let us consider the subgroup $[a, G']$ of G generated by all commutators of the form $[a, c]$ where $c \in G'$.¹⁵ Using the identity $[x, yz] = [x, z][x, y][[x, y], z]$ and the hypothesis that G' is abelian it is easy to see that $[a, c][a, d] = [a, cd]$ and $[a, c]^{-1} = [a, c^{-1}]$ for $c, d \in G'$. Hence every element of $[a, G']$ (and not just its generators) can be represented in the form $[a, c]$ for some $c \in G'$. On the other hand, using the identities $x^y = x[x, y]$ and $[x, y] = [y, x]^{-1}$ and the metabelian Hall–Witt identity $[[x, y], z][[y, z], x][[z, x], y] = 1$,¹⁶ we get $[a, c]^g = [a, c][[a, c], g] = [a, c][[g, a], c]^{-1}[[c, g], a]^{-1} = [a, c][a, [c, g]]$. Hence $[a, G']$ is normal in G .

Given $g, h \in G$, we have $g = a^m c$ and $h = a^n d$ for some $m, n \in \mathbb{Z}$ and $c, d \in G'$. Using the identities $[x, yz] = [x, z][x, y][[x, y], z]$ and $[xy, z] = [x, z][[x, z], y][y, z]$ we get $[g, h] = [a^m, d][c, a^n] = [a^m, d][a^n, c]^{-1}$. Then using the identity $[xy, z] = [x, z]^y[y, z]$ and the normality of $[a, G']$ we get that $[g, h] \in [a, G']$. Hence $G' = [a, G']$. Since $a^{-1}b \in G' = [a, G']$, we have $a^{-1}b = [a, c] = a^{-1}a^c$ for some $c \in G'$. Hence $b = a^c$. \square

Theorem 3.2 is a special case of Lemma 3.3. But in the geometric context there is also a more illuminating proof of Theorem 3.2, based on the knot module.

Let K be a knot in S^3 . Let $X = S^3 \setminus K$ and let $p: \tilde{X} \rightarrow X$ be the infinite cyclic covering. The group $H_1(X) \simeq \mathbb{Z} \simeq \langle t \mid \rangle$ acts on \tilde{X} by covering transformations.

Lemma 3.4. *The action by $t - 1$ on $H_1(\tilde{X})$ and on $H_2(\tilde{X})$ is by automorphisms.*

Proof. The proof for PL knots found in [68; Lemma 4.1(a)] also works for topological knots, since so does the Alexander duality (using Čech cohomology; see for instance [64; Theorem 4.3] or [66; Theorem 27.1(a)]). \square

Proof of Theorem 3.2. Let $G = \pi/\pi''$, where $\pi = \pi_1(S^3 \setminus K)$. Let $a, b \in G$ be some members of the conjugacy classes represented by $[Q_1]$ and $[Q_2]$. Let $A = \pi'/\pi''$. Since $\pi/\pi' \simeq \mathbb{Z}$ is a free group, the short exact sequence $1 \rightarrow A \rightarrow G \rightarrow \mathbb{Z} \rightarrow 1$ splits, and so G is a semidirect product $G \simeq A \rtimes_{\varphi} \mathbb{Z}$ with respect to some automorphism φ of A ; namely, $\varphi(g) = g^a$. Clearly, A with this action of \mathbb{Z} is the knot module $H_1(\tilde{X})$; more

¹⁴A. Zastrow was able to explicitly construct such a grope for a certain wild knot satisfying the above specifications (private communication, August 2023). It is different from the wild knot of the previous footnote. This had been before I found the geometric proof of Theorem 3.2 (see Theorem 3.6).

¹⁵Hereafter $[x, y] = x^{-1}y^{-1}xy$ and $x^y = y^{-1}xy$.

¹⁶The usual Hall–Witt identity $[[x, y], z^x][[y, z], x^y][[z, x], y^z] = 1$ holds in all groups; the first factor $[[x, y], z^x] = [[x, y], z[z, x]] = [[x, y], [z, x]][[x, y], z][[[x, y], z], [z, x]]$ simplifies as $[[x, y], z]$ in metabelian groups, and similarly for the second and third ones.

precisely, $\varphi(g) = t \cdot g$ in the additive notation, where t is the image of a in $G/G' \simeq \mathbb{Z}$. Moreover, the action by $1 - t$ is an automorphism of A (see Lemma 3.4(a)).

We want to show that $a^g = b$ for some $g \in A$. This can be rewritten as $a^{-1}g^{-1}ag = a^{-1}b$, or $(g^{-1})^a g = c$, where $c = a^{-1}b \in A$. In the additive notation, this becomes $g - t \cdot g = c$, or $(1 - t)g = c$. But since the action by $1 - t$ is invertible, such a g does exist. \square

The second proof of Theorem 3.2 can actually be turned into a geometric proof (see Theorem 3.6 below) by using the techniques of the following lemma.

Lemma 3.5. [6] *Let $L = (K_1, K_2)$ be a PL link with $\text{lk}(L) = 1$. Then there exist Seifert surfaces F_i for K_i , intersecting transversely and such that $F_1 \cap F_2$ is a single arc with one endpoint in K_1 and another in K_2 .*

The proof is quite simple, so we include it for the reader's convenience.

Proof. Let F_i be an arbitrary Seifert surface for K_i . We may assume that F_1 and F_2 intersect transversely. Since $\text{lk}(L) = 1$, there are n negative and $n+1$ positive intersection points in $F_1 \cap K_2$ for some non-negative integer n . If $n > 0$, at least one of the arcs into which these intersection points split K_2 has endpoints of opposite signs. By attaching a thin tube to F_1 going along this arc we eliminate these two intersection points. By continuing in this fashion we eliminate all of the intersection points in $F_1 \cap K_2$ but one. By similarly attaching tubes to F_2 going along arcs of K_1 we eliminate all intersection points in $F_2 \cap K_1$ except one. The two remaining points in $\partial(F_1 \cap F_2)$ must cobound an arc J in $F_1 \cap F_2$. If $F_1 \cap F_2$ also contains closed components, then there is one, call it C , such that some point of C can be joined to an interior point of J by an arc A in F_1 which is otherwise disjoint from $F_1 \cap F_2$ and from K_1 . Since J is non-separating in F_1 , one can choose A so that both its ends approach F_2 from the same side. Then by attaching a thin tube to F_2 going along A we will replace J and C by their connected sum. By continuing in this fashion we will eliminate all closed components of $F_1 \cap F_2$. \square

Theorem 3.6. *Let \mathcal{K} be a knot in S^3 and let Q_1, Q_2 be smooth knots in $S^3 \setminus \mathcal{K}$ such that $\text{lk}(Q_i, \mathcal{K}) = 1$. Then the connected sum $Q_1 \#_b (-Q_2)$ along some band disjoint from \mathcal{K} bounds a Seifert surface F in $S^3 \setminus \mathcal{K}$ such that each knot in F is null-homologous in $S^3 \setminus \mathcal{K}$.*

Proof. Since the link $L := Q_1 \cup (-Q_2)$ is null-homologous in $S^3 \setminus \mathcal{K}$, it bounds a smooth Seifert surface S in $S^3 \setminus \mathcal{K}$.¹⁷ Let Σ be a smooth h -Seifert surface bounded by \mathcal{K} in S^3 (see [65; Lemma 2.1]). We may assume that S and Σ intersect transversely. Since $\text{lk}(Q_i, \mathcal{K}) = 1$, by attaching tubes to Σ as in the proof of Lemma 3.5 we can eliminate

¹⁷This follows similarly to [65; Lemma 2.1] and [83; §1]. In more detail, let N be a tubular neighborhood of L in $S^3 \setminus \mathcal{K}$ and let $L_+ \subset \partial N$ be a zero pushoff of L , which is null-homologous in $S^3 \setminus L$. Since L_+ is also null-homologous in $S^3 \setminus \mathcal{K}$, it is null-homologous in $S^3 \setminus (\mathcal{K} \cup L)$ (by the Mayer-Vietoris exact sequence), and hence also in $S^3 \setminus N \setminus \mathcal{K}$. Then by Lemma 17.8 L_+ bounds an orientable smooth surface in $S^3 \setminus N \setminus \mathcal{K}$. Using the obvious pair of annuli cobounded by L and L_+ in N , this extends to a smooth Seifert surface of L in $S^3 \setminus \mathcal{K}$.

all its intersections with Q_i except one, call it p_i . The two remaining intersection points p_1 and p_2 must cobound an arc J in $\Sigma \cap S$. If $\Sigma \cap S$ also contains closed components, then by attaching tubes to Σ along arcs in S as in the proof of Lemma 3.5, using that J is non-separating in S , we can make them absorbed into J . Now $S \cap \Sigma$ consists of a single arc J . Let b be its regular neighborhood in S . Then $Q := Q_1 \#_b (-Q_2)$ bounds the Seifert surface $F := \overline{S \setminus b}$. By construction F is disjoint from Σ . Hence every knot in F has zero linking number with \mathcal{K} . \square

4. PROOF OF THEOREM E

We recall that a group G is called n -solvable if $G^{(n)} = 1$. Let us note that 1-solvable, 1.5-solvable and 2-solvable groups are also known as abelian, nilpotent of class 2 and metabelian, respectively.

Example 4.1. There exist a finitely presented 2.5-solvable group G such that $G/G' \simeq \mathbb{Z}$ and $a, b \in G$ such that the abelianization map $G \rightarrow G/G'$ sends a and b to the same generator of \mathbb{Z} , but a is not conjugate to b .

First suppose that G is the desired group. Since $G/G' \simeq \mathbb{Z}$, we have a short exact sequence $1 \rightarrow G' \rightarrow G \rightarrow \mathbb{Z} \rightarrow 1$, which splits since \mathbb{Z} is free. Hence $G \simeq G' \rtimes_{\varphi} \mathbb{Z}$ for some automorphism φ of G' . On the other hand, saying that G is 2.5-solvable, that is, $[G', G''] = 1$, amounts to saying that $H := G'$ is 1.5-solvable, that is, $[H, H'] = 1$ (in other words, H nilpotent of class 2).

Thus it makes sense to seek G in the form $G = H \rtimes_{\varphi} \mathbb{Z}$ for some automorphism φ of H , where H is nilpotent of class 2. If we show that $G' = H$, then we will get that $[G', G''] = [H, H'] = 1$, that is, G is 2.5-solvable. Let $\bar{\varphi}$ be the automorphism of the abelianization H/H' determined by φ .

Lemma 4.2. *If $\bar{\varphi} - \text{id}: H/H' \rightarrow H/H'$ is surjective, then $G' = H$.*

Proof. Geometrically, $G = \pi_1(MT, pt)$, where MT is the mapping torus of a self-map $\Phi: (BH, pt) \rightarrow (BH, pt)$ of the classifying space $BH = K(H, 1)$ representing the automorphism φ (see Lemma 4.3(b)). Using the isomorphism $H_i(MT, BH) \simeq H_i(BH \times I, BH \times \partial I) \simeq H_{i-1}(BH)$, the exact sequence of the pair (MT, BH) can be written as

$$H_1(BH) \xrightarrow{\bar{\varphi} - \text{id}} H_1(BH) \xrightarrow{\subset_*} H_1(MT) \rightarrow H_0(BH) \xrightarrow{\text{id} - \text{id}} H_0(BH).$$

Since $\bar{\varphi} - \text{id}$ is surjective, the map $H_1(BH) \rightarrow H_1(MT)$ is trivial, and it follows that $H_1(MT) \simeq \mathbb{Z}$. In other words, the map $H/H' \rightarrow G/G'$ is trivial, and also $G/G' \simeq \mathbb{Z}$.¹⁸

¹⁸Algebraically, the same conclusions can be similarly derived from Bieri's long exact sequence [11]: $H_1(H) \xrightarrow{\bar{\varphi} - \text{id}} H_1(H) \rightarrow H_1(G) \rightarrow H_0(H) \xrightarrow{\text{id} - \text{id}} H_0(H)$. In fact, the two long exact sequences are the same. Indeed, since Φ is a homotopy equivalence, MT is the total space of an approximate fibration over S^1 [28]. Consequently, since S^1 is aspherical, so is MT . Alternatively, with the of choice H that will interest us below, BH can be chosen to be a closed 3-manifold (the quotient of the continuous Heisenberg group by the discrete Heisenberg group), and Φ can be chosen to be a homeomorphism [54; Theorem 0.7]. In this case MT will be the total space of a (genuine) fibration over S^1 .

It follows from the first observation that H lies in G' . Then G/G' is a quotient of G/H ; but both are isomorphic to \mathbb{Z} , which is a Hopfian group, so we get that $H = G'$. \square

Lemma 4.3. (a) *If $\langle x_1, \dots, x_n \mid r_1, \dots, r_m \rangle$ is a presentation of a group H and φ is an automorphism of H , then a presentation of $H \rtimes_{\varphi} \mathbb{Z}$ is given by*

$$\langle x_1, \dots, x_n, t \mid r_1 = 1, \dots, r_m = 1, x_1^t = \varphi(x_1), \dots, x_n^t = \varphi(x_n) \rangle.$$

(b) *Let X be a CW complex, $b \in X$ a point and $f: (X, b) \rightarrow (X, b)$ a map such that $f_*: \pi_1(X, b) \rightarrow \pi_1(X, b)$ is an isomorphism. Let T_f be the mapping torus of f , that is, $T_f = X \times [0, 1] / \sim$, where $(x, 0) \sim (f(x), 1)$. Then $\pi_1(T_f, b) \simeq \pi_1(X, b) \rtimes_{f_*} \mathbb{Z}$.*

This is well-known. Part (a) says that “a semidirect product with \mathbb{Z} is special kind of an HNN-extension” and part (b) is related to the assertion that “an HNN-extension is the fundamental group of a certain graph of groups”.

Proof. (a). It follows from the “inner” and “outer” definitions of a semidirect product that H and $\mathbb{Z} = \langle t \mid \rangle$ can be identified with subgroups of $H \rtimes_{\varphi} \mathbb{Z}$, that every element of $H \rtimes_{\varphi} \mathbb{Z}$ can be expressed in the form $t^k h$ for some $h \in H$ and $k \in \mathbb{Z}$, and that $t^k h \cdot t^l h' = t^{k+l} \varphi^l(h) h'$ for $h, h' \in H$ and $k, l \in \mathbb{Z}$. In particular, $h^t = \varphi(h)$ for all $h \in H$, and it follows that the relations $x_i^t = \varphi(x_i)$ and $r_j = 1$ do hold in $H \rtimes_{\varphi} \mathbb{Z}$. Conversely, if $h^t = \varphi(h)$ holds for $h = x_i$, then it clearly holds for all $h \in H$, which in turn implies the multiplication law, $t^k h \cdot t^l h' = t^{k+l} \varphi^l(h) h'$. \square

(b). We may assume that b is a 0-cell of X and there are no other 0-cells, and that f is a cellular map. Then T_f can be identified with the CW complex obtained from X by attaching one $(n+1)$ -cell corresponding to each n -cell of X in the obvious way; namely, $D^n \xrightarrow{e} X$ yields $D^n \times I \xrightarrow{e \times \text{id}_I} X \times I \rightarrow T_f$. It is a well-known consequence of the Seifert–van Kampen theorem that a presentation of $\pi_1(X, b)$ is determined by the 2-skeleton of X (with generators given by the 1-cells and relators by the 2-cells). A presentation of $\pi_1(T_f, b)$ is similarly determined by the 2-skeleton of T_f , and now the assertion follows from (a). \square

Now let, in fact, H be the free nilpotent group of class 2 on two generators, which has the presentation $\langle x, y \mid [[x, y], x] = [[x, y], y] = 1 \rangle$ and is also known as the discrete Heisenberg group. By using the commutator collecting process (see [36; §11]) it is easy to see that every element of H can be uniquely written as $x^l y^m [y, x]^n$ for some $l, m, n \in \mathbb{Z}$, and $x^l y^m [y, x]^n \cdot x^{l'} y^{m'} [y, x]^{n'} = x^{l+l'} y^{m+m'} [y, x]^{n+n'+ml'}$. It follows that $(x^l y^m [y, x]^n)^{-1} = x^{-l} y^{-m} [y, x]^{-n+lm}$.

Let $\varphi: H \rightarrow H$ be given¹⁹ by $\varphi(x) = y^{-1}$ and $\varphi(y) = yx$. Let us note that φ is an isomorphism, with φ^{-1} given by $\varphi^{-1}(y) = x^{-1}$ and $\varphi^{-1}(x) = xy$. The abelianization $\bar{\varphi}: H/H' \rightarrow H/H'$ of φ is given by the matrix $\begin{pmatrix} 0 & 1 \\ -1 & 1 \end{pmatrix}$. Therefore $\bar{\varphi} - \text{id}: H/H' \rightarrow H/H'$ is given by the matrix $\begin{pmatrix} -1 & 1 \\ -1 & 0 \end{pmatrix}$, and so is also an isomorphism.

¹⁹Since H is a free nilpotent group of class 2, every map of the set of its generators to a class 2 nilpotent group K extends to a homomorphism $H \rightarrow K$.

It is easy to see that $\varphi([y, x]) = [yx, y^{-1}] = [x, y^{-1}] = [y, x]^{y^{-1}}$, which equals $[y, x]$ in H . Hence $\varphi(x^l y^m [y, x]^n) = y^{-l} (yx)^m [y, x]^n = x^m y^{m-l} [y, x]^{n-lm+m(m+1)/2}$, where $(yx)^k = x^k y^k [y, x]^{k(k+1)/2}$ by using the commutator collecting process.

The semidirect product $G = H \rtimes_{\varphi} \mathbb{Z}$ consists of all products of the form $t^k h$, where $h \in H$ and $k \in \mathbb{Z}$, and satisfies the relation $h^t = \varphi(h)$, or equivalently $ht = t\varphi(h)$ for all $h \in H$ (see the proof of Lemma 4.3(a)).

Finally, let us show that $a := t$ and $b := txy$ are not conjugate by any $g \in G$. We have $g = t^k x^l y^m [y, x]^n$ for some $k, l, m, n \in \mathbb{Z}$. Then

$$\begin{aligned} a^g &= t^{t^k x^l y^m [y, x]^n} = x^{-l} y^{-m} [y, x]^{-n+lm} t x^l y^m [y, x]^n = t\varphi(x^{-l} y^{-m} [y, x]^{-n+lm}) x^l y^m [y, x]^n \\ &= t x^{-m} y^{-m+l} [y, x]^{-n+lm-(-l)(-m)+(-m)(-m+1)/2} x^l y^m [y, x]^n = t x^{l-m} y^l [y, x]^{l(l-m)+m(m-1)/2}. \end{aligned}$$

If $a^g = b$, then $l - m = 1$, $l = 1$ and $l(l - m) + m(m - 1)/2 = 0$. The first two equations imply $m = 0$, and then the third equation implies $1 = 0$.




FIGURE 2.

Example 4.4. The fundamental group $\pi := \pi_1(S^3 \setminus Q)$ of the trefoil knot Q contains elements a, b which map to the same generator of $H_1(S^3 \setminus Q) \simeq \mathbb{Z}$ but their images in $\pi/\pi^{(2.5)}$ are not conjugate.

Indeed, the trefoil is well-known to be a fibered knot, more precisely $S^3 \setminus Q$ fibers over S^1 with fiber being the punctured torus (see [80; pp. 327–332], see also [20; 5.12]). Hence $\pi \simeq \pi' \rtimes_{\psi} \mathbb{Z}$, where the commutator subgroup π' is isomorphic to the free group $F_2 = \langle x, y \mid \rangle$, and ψ is its automorphism induced by the monodromy self-homeomorphism of the fiber (see [20; 5.4], see also [80; p. 332]). In order to prepare for the next example, we recall a computation of the monodromy ψ (this computation is a slight variation of [20; 5.15]). Figure 2 shows a Seifert surface S of the minimal genus for the trefoil K ; loops x, y in S whose homotopy classes freely generate $\pi_1(S)$; and the generators of the Wirtinger presentation $\langle p, q, r \mid pq^{-1} = r^{-1}p, qr^{-1} = p^{-1}q, rp^{-1} = q^{-1}r \rangle$ for π . (The first relation corresponds to the upper crossing, the second relation to the middle crossing and the third relation to the lower crossing in the figure. As is well-known, any one of the relations is redundant.) It is easy to see from the figure that $x = r^{-1}p$ and $y = p^{-1}q$. By using the relations we also get $x = pq^{-1}$ and $y = qr^{-1}$. Let us note that ψ depends on an

identification of \mathbb{Z} with a subgroup of π ; if, for instance, the generator t of \mathbb{Z} is identified with p , then $\psi(g) = g^p$. We have $x^p = p^{-1}pqp^{-1}p = q^{-1}p = y^{-1}$ and $y^p = p^{-1}qr^{-1}p = yx$. Thus $\psi(x) = y^{-1}$ and $\psi(y) = yx$.

Since the automorphism φ of the previous example is given by the same formula, we obtain a homomorphism $f: \pi \rightarrow G$, where G is the group of the previous example. Clearly f is surjective. Since G is 2.5-solvable, f factors through the quotient $\pi/\pi^{(2.5)}$.²⁰ Now the assertion follows from the previous example.




FIGURE 3.

Example 4.5. There exists a wild knot $\mathcal{K} \subset S^3$ and its meridians $\partial D_1, \partial D_2$ ²¹ such that the conjugacy classes in $\pi := \pi_1(S^3 \setminus \mathcal{K})$ corresponding to ∂D_1 and ∂D_2 have different images in $\pi/\pi^{(2.5)}$.

The wild knot \mathcal{K} is shown in Figure 3. It is the union of two closed arcs with common endpoints: a tame arc A (which intersects the disk D_1) and a wild arc A' (which intersects the disk D_2). Since A is tame, $S^3 \setminus A$ is homeomorphic to \mathbb{R}^3 , which is in turn homeomorphic to the infinite cyclic cover of $S^3 \setminus B$, where B is an unknot. Moreover, it is easy to see that the composite homeomorphism can be chosen so that $\mathcal{K} \setminus A$ gets identified with the preimage under the covering map $S^3 \setminus A \rightarrow S^3 \setminus B$ of the knot $Q \subset S^3 \setminus B$ which is shown on the left hand side of Figure 4. Let $L = B \cup Q$. The images of ∂D_i under the covering map are the knots J_i in $S^3 \setminus L$, which are also shown on the left hand side of Figure 4. Hence it suffices to show that the conjugacy classes in $\bar{\pi} := \pi_1(S^3 \setminus L)$ corresponding to J_1 and J_2 have different images in $\bar{\pi}/\bar{\pi}^{(2.5)}$. For this it in turn suffices to show that the conjugacy classes in $\hat{\pi} := \pi_1(S^3 \setminus Q)$ corresponding to J_1 and J_2 have different images in $\hat{\pi}/\hat{\pi}^{(2.5)}$. The knot Q is easily seen to be equivalent to the trefoil, and the right hand side of Figure 4 shows the images of J_1 and J_2 under the equivalence.

²⁰In fact, the epimorphism $g: \pi/\pi^{(2.5)} \rightarrow G$ is an isomorphism. Indeed, since $\pi^{(2.5)} = [\pi'', \pi'] = [F'_2, F_2]$, the image of F_2 in $\pi/\pi^{(2.5)}$ is the free nilpotent group H of class 2, and it follows that $\ker g = 1$.

²¹That is, each ∂D_i bounds an embedded PL disk $D_i \subset S^3$ which meets \mathcal{K} in a single point and locally separates it at that point.

Then in the notation of the previous example (Example 4.4), J_2 can be represented by p and J_1 by p^2r^{-1} in $\pi_1(S^3 \setminus Q)$. Since $p^2r^{-1} = pxy$, in the notation of Example 4.1 these correspond to t and txy . Thus the assertion follows from the previous two examples.




FIGURE 4.

Proof of Theorem E. Theorem E is proved by the construction of Zastrow's Example (see Example 2.2) if instead of Rolfsen's wild knot and its meridional disks D_0 and D_1 we use the wild knot of Example 4.5 and its meridional disks D_1 and D_2 . \square

The strengthening of Theorem E stated in Remark 1.3 follows from the proof of Theorem E, taking into account the following observation.

Proposition 4.6. *The knot module $H_1(\widetilde{S^3 \setminus \mathcal{K}})$ of the wild knot \mathcal{K} shown in Figure 3 is $\mathbb{Z}[t^{\pm 1}]$ -torsion free.*

Proof. As explained in Example 4.5, \mathcal{K} contains a closed tame arc A such that the pair $(S^3 \setminus A, \mathcal{K} \setminus A)$ is homeomorphic to $(\widetilde{S^3 \setminus B}, p^{-1}(Q))$, where $L = B \cup Q$ is the 2-component link in S^3 shown in Figure 4 (on the left) and $p: S^3 \setminus B \rightarrow S^3 \setminus B$ is the infinite cyclic covering. It follows that the infinite cyclic covering $q: \widetilde{S^3 \setminus \mathcal{K}} \rightarrow S^3 \setminus \mathcal{K}$ is fiberwise equivalent to the projection $\widetilde{S^3 \setminus L} \rightarrow (\widetilde{S^3 \setminus B}) \setminus p^{-1}(Q)$ from the universal abelian covering of $S^3 \setminus L$ to its infinite cyclic covering which extends to the infinite cyclic covering of $S^3 \setminus B$. Thus $H_1(\widetilde{S^3 \setminus \mathcal{K}})$ is a $\mathbb{Z}[s^{\pm 1}, t^{\pm 1}]$ -module, where s and t are the positive generators of the deck transformation groups of the coverings p and q , respectively.

The infinite cyclic covering $\widetilde{S^3 \setminus B}$ can be regarded as a copy of \mathbb{R}^3 , which contains the periodic open knot $p^{-1}(Q)$. Figure 5 shows a periodic Seifert surface F of this open knot in this \mathbb{R}^3 , such that $F = p^{-1}(\Sigma \setminus B)$, where Σ is a Seifert surface for Q which meets B transversely in one point. The identifying homeomorphism between $S^3 \setminus A$ and $\widetilde{S^3 \setminus B}$ can be chosen so that $\mathcal{F} := F \cup A$ is a Seifert surface for \mathcal{K} in S^3 . Let us note that $S^3 \setminus \mathcal{F}$ is identified with $\mathbb{R}^3 \setminus F$.




FIGURE 5.

The upper half of Figure 5 shows representative cycles for classes $a, b \in H_1(F)$ and for classes $\alpha, \beta, \gamma, \delta \in H_1(\mathbb{R}^3 \setminus F)$. Clearly, a basis in $H_1(F)$ over $\mathbb{Z}[s^{\pm 1}]$ is given by a and b . It follows from the Alexander duality (see [68; §5]) that a basis of $H_1(\mathbb{R}^3 \setminus F)$ over $\mathbb{Z}[s^{\pm 1}]$ is given by α and β . It is easy to see that

$$\begin{aligned} \alpha + \gamma &= \beta \\ \beta + \delta &= s\beta. \end{aligned} \tag{*}$$

Now let $a^{\pm}, b^{\pm} \in H_1(\mathbb{R}^3 \setminus F)$ be the classes of the parallel \pm -pushoffs of representative cycles of a and b . Their representative cycles are shown in the lower half of Figure 5. It is easy to see that

$$\begin{aligned} a^+ &= \alpha & b^+ &= \beta + \gamma - s^{-1}\beta \\ a^- &= -\gamma & b^- &= \beta - \delta. \end{aligned}$$

Using (*), we can eliminate γ and δ :

$$\begin{aligned} a^+ &= \alpha & b^+ &= (2 - s^{-1})\beta - \alpha \\ a^- &= \alpha - \beta & b^- &= (2 - s)\beta. \end{aligned} \tag{**}$$

Let $\tilde{\mathcal{F}}$ be some fixed lift of $\mathcal{F} \setminus \mathcal{K}$ in the infinite cyclic cover $\widetilde{S^3 \setminus \mathcal{K}}$ and let $\tilde{a}^\pm, \tilde{b}^\pm, \tilde{\alpha}$ and $\tilde{\beta}$ be the classes in $H_1(\widetilde{S^3 \setminus \mathcal{K}})$ of the lifts of some representative cycles of a^\pm, b^\pm, α and β respectively lying between $\tilde{\mathcal{F}}$ and $t\tilde{\mathcal{F}}$, where t is the positive generator of the group of deck transformations of $\widetilde{S^3 \setminus \mathcal{K}}$. Then \tilde{a}^\pm and \tilde{b}^\pm can be expressed in terms of $\tilde{\alpha}$ and $\tilde{\beta}$ similarly to (**), and a presentation of $H_1(\widetilde{S^3 \setminus \mathcal{K}})$ as a $\mathbb{Z}[s^{\pm 1}, t^{\pm 1}]$ -module is given by $\langle \tilde{\alpha}, \tilde{\beta} \mid \tilde{a}^- = t\tilde{a}^+, \tilde{b}^- = t\tilde{b}^+ \rangle$; that is,

$$\langle \tilde{\alpha}, \tilde{\beta} \mid (\tilde{\alpha} - \tilde{\beta}) = t\tilde{\alpha}, (2 - s)\tilde{\beta} = t(2 - s^{-1})\tilde{\beta} - t\tilde{\alpha} \rangle.$$

From the first relation we have $\tilde{\beta} = (1 - t)\tilde{\alpha}$. Substituting this into the second relation and dividing by t , we get $(1 - t^{-1})(s - 2)\tilde{\alpha} + (1 - t)(s^{-1} - 2)\tilde{\alpha} + \tilde{\alpha} = 0$. Expanding out, we arrive at the following presentation of $H_1(\widetilde{S^3 \setminus \mathcal{K}})$ as a $\mathbb{Z}[s^{\pm 1}, t^{\pm 1}]$ -module:

$$\langle \tilde{\alpha} \mid (1 - t^{-1})s\tilde{\alpha} + (1 - t)s^{-1}\tilde{\alpha} + (2t + 2t^{-1} - 3)\tilde{\alpha} = 0 \rangle.$$

Writing $x_i = s^i\tilde{\alpha}$, we get a presentation of $H_1(\widetilde{S^3 \setminus \mathcal{K}})$ as a $\mathbb{Z}[t^{\pm 1}]$ -module:

$$\langle x_i, i \in \mathbb{Z} \mid (1 - t^{-1})x_{i+1} + (1 - t)x_{i-1} + (2t + 2t^{-1} - 3)x_i = 0, i \in \mathbb{Z} \rangle. \quad (***)$$

Since the multiplication by $1 - t$ in $H_1(\widetilde{S^3 \setminus \mathcal{K}})$ is an isomorphism (see Lemma 3.4(a)), this abelian group is also a module over the ring $\mathbb{Z}[t^{\pm 1}]_{[\frac{1}{1-t}]}$ (that is, the localization of $\mathbb{Z}[t^{\pm 1}]$ with respect to the multiplicatively closed set $\{(1 - t)^i \mid i = 0, 1, 2, \dots\}$). The same presentation (***) works for $H_1(\widetilde{S^3 \setminus \mathcal{K}})$ as a module over $\mathbb{Z}[t^{\pm 1}]_{[\frac{1}{1-t}]}$. But over this ring the relation can be rewritten as

$$x_{i+1} = -(1 - t^{-1})((1 - t)x_{i-1} + (2t + 2t^{-1} - 3)x_i)$$

and also as

$$x_{i-1} = -(1 - t)((1 - t^{-1})x_{i+1} + (2t + 2t^{-1} - 3)x_i),$$

and it follows that $H_1(\widetilde{S^3 \setminus \mathcal{K}})$ is a free module of rank two over $\mathbb{Z}[t^{\pm 1}]_{[\frac{1}{1-t}]}$, freely generated by x_0 and x_1 (for instance). In particular, this implies that $H_1(\widetilde{S^3 \setminus \mathcal{K}})$ contains no $\mathbb{Z}[t^{\pm 1}]$ -torsion. \square

5. EXACTLY WHAT IS THE BING SLING?

Let K_0 be an unknot in S^3 , and let T_1 be its regular neighborhood in S^3 . Let J be a copy of the fake²² Mazur curve in the unknotted solid torus T_1 (see Figure 6). Given an $i \in \{1, 2, \dots\}$ and assuming that a solid torus T_i in S^3 has been defined, let K_i be the preimage of J under an r_i -fold cover $p_i: T_i \rightarrow T_1$ and let T_{i+1} be a regular neighborhood K_i in T_i . (In Figure 1, $r_1 = 6$.) We will always assume the following hypothesis:

²²It differs from the original Mazur curve [59] in one crossing, the one which is marked by an asterisk in Figure 6. See [63; §2.2] for a discussion of one curve versus another. See also [1] for a discussion of the Mazur curve versus Zeeman's curve, which is a mirror image of the fake Mazur curve.

- (0) each p_i is untwisted, in the sense of being regular homotopic (with values in T_1) to the pullback $T_1 \rightarrow T_1$ of the orientation-preserving r_i -fold cover $S^1 \rightarrow S^1$ along the projection $T_1 \simeq S^1 \times D^2 \rightarrow S^1$.




FIGURE 6. The fake Mazur curve.

While Bing himself did not explicitly discuss in his paper [12] any of the choices involved, it is not hard to see, as explained in the book of R. Daverman and G. Venema [30; §2.8.5], that the numbers r_i , the covers p_i (as specific maps, not just up to a fiberwise homeomorphism) and the regular neighborhoods T_i can be chosen so that $\bigcap_{i=1}^{\infty} T_i$ is homeomorphic to S^1 and thus we get an embedding \mathcal{B} of S^1 . However, with their approach (which is based on a general lemma [30; Proposition 2.2.2] that the uniform limit of a sequence of embeddings will be an embedding if they converge rapidly enough) it seems to be challenging to describe any one resulting sequence r_1, r_2, \dots in closed terms, and it appears that the equivalence class of \mathcal{B} (up to ambient isotopy) will depend not only on the integers r_i , but also on metric properties of the maps p_i . Which could lead some readers to ask: “OK, but exactly what do you mean by the Bing sling? Is it a well-defined topological object? Can it be described explicitly enough that one can evaluate at least one algebraic invariant on it and obtain an answer in closed form?”

We will take a different, more tailor-made approach, which presumably settles these questions in a satisfactory way. Let us note that the meridional disk D of T_1 shown in Figure 6 has r_i distinct lifts D_{ij} into T_i , which partition T_i into r_i (closed) fundamental domains Φ_{ij} of the group of deck transformations of the cover p_i . (Figure 1 hints at the six disks D_{1j} and the six cylinders Φ_{1j} , as well as five of the disks D_{2j} and four of the cylinders Φ_{2j} .) Suppose that

- (1) each Φ_{ij} is of diameter $\leq \epsilon_i$, where $\epsilon_n \rightarrow 0$ as $n \rightarrow \infty$;
- (2) each $\Phi_{ij} \cap T_{i+1}$ has only 3 connected components and each of them is a union of q_i of the $\Phi_{i+1,k}$'s, where $q_i := \frac{r_{i+1}}{3r_i}$ is required to be an integer.

(In Figure 1, $q_1 = 4$.)

Theorem 5.1. *Suppose that conditions (0), (1) and (2) hold. Then*

- (a) *the knots K_i , when suitably parameterized, uniformly converge to an embedding \mathcal{B} of S^1 whose image is $\bigcap_{i=1}^{\infty} T_i$;*

(b) the equivalence class of \mathcal{B} (up to ambient isotopy) depends only on the integers r_i .
Moreover,

(c) given a sequence of positive integers r_1, r_2, \dots such that each $\frac{r_{i+1}}{3r_i}$ is an integer greater than 1, there exists a sequence of K_i and T_i as above, satisfying (0), (1) and (2).

The approach in the proof of (a) will still work, albeit with more tedious details, if instead of (2) we assume the following weaker hypothesis:

(2') each $\Phi_{ij} \cap T_{i+1}$ has only 3 connected components and each of them is a union of some of the $\Phi_{i+1,k}$'s.

However, it is expected that (b) will be false if we replace (2) with (2').²³

Although (c) tells us that for Bing's construction to work, the sequence r_1, r_2, \dots need not be particularly rapidly growing after all (for instance, $r_i = 6^{i-1}$ will do), we will nevertheless need to assume in the present paper (apart from the present section) that it is quite rapidly growing, because of our use of Theorem 9.2. To be specific, the proofs of Theorems A and B work under the hypothesis that each $r_{i+1} \geq (r_i^2 + 3r_i)^2$ (see Remark 10.2), and the proofs of Theorems C, D and F work under the slightly weaker hypothesis $r_{i+1} \geq (r_i^2 - r_i + 1)^2 + 1$ (see Remark 13.3). For instance, $r_i = 3^{5^{i-1}}$ satisfies all our hypotheses.

Despite the fact that the wild knot \mathcal{B} produced by Bing's construction depends on the choice of the integers r_i , it is customary in the literature to speak of *the* Bing sling²⁴ (rather than many different Bing slings) and we will keep with this tradition, in order to avoid peripheral details in statements of theorems. But this forces us to include the condition of rapid growth of the r_i in the definition of "the Bing sling". Thus *the Bing sling* is defined for the purposes of this paper by conditions (0), (1), (2) and the condition that each $r_{i+1} \geq (r_i^2 + 3r_i)^2$.

Proof of Theorem 5.1. (a). Let us make the notation D_{ij}, Φ_{ij} more precise (so far they are defined only up to a permutation of the second indices). First we choose an arbitrary lift D_1 of D in T_1 . To continue, we need the basepoint b shown in Figure 6 (one of the three intersection points of D with the fake Mazur curve J). Assuming that a lift D_i of D in T_i has been defined, it contains a unique lift b_i of b in T_i . This b_i lies in K_i and hence also in $D_i \cap T_{i+1}$. Then by (2) there is a unique lift D_{i+1} of D in T_{i+1} which contains b_i . Let us note that the sequence of disks defined in this way is nested: $D_1 \supset D_2 \supset \dots$. Moreover, since by (1) their diameters tend to zero, $\bigcap_{i=1}^{\infty} D_i$ consists of a single point b_{∞} , which is also the limit of the sequence b_1, b_2, \dots . Now let $D_{i0} = D_i$. Assuming that a lift D_{ij} of D in T_i has been defined, we define $D_{i,j+1}$ to be the next one in the positive direction (determined by the orientation of the fake Mazur curve J). Thus $D_{ij} = D_{i,j+r_i}$,

²³Let us note a potential connection with Rolfsen's Problem: if (b) were true when (2) is replaced with (2'), then this would provide, in particular, an isotopy from the Bing sling (as defined more precisely below) to a somewhat differently looking wild knot, which might be easier to isotope to the unknot by some geometric construction.

²⁴The first appearance of this terminology that I'm aware of is in R. H. Fox's list of problems in knot theory [32; p. 171].

and using this formula we also define D_{ij} for all $j < 0$. Also let Φ_{ij} be the fundamental domain bounded by D_{ij} and $D_{i,j+1}$, and let b_{ij} be the unique lift of b in T_i that lies in D_{ij} . (Thus $b_{i0} = b_i$.)

The fake Mazur curve can be parameterized by an embedding $k: S^1 \rightarrow T_1$, which sends $0 \in S^1 = \mathbb{R}/\mathbb{Z}$ onto b . Let \tilde{k} be the composition $\mathbb{R} \xrightarrow{q} \mathbb{R}/\mathbb{Z} = S^1 \xrightarrow{k} T_1$. We may assume that \tilde{k} sends $\frac{1}{3}$ and $\frac{2}{3}$ onto the other two points of intersection of J and D . The embedding k , when precomposed with the r_i -fold cover $S^1 \rightarrow S^1$, $s \mapsto r_i s$, lifts to an embedding $k_i: S^1 \rightarrow T_i$ which parameterizes K_i . We may assume that $k_i(0) = b_i$. Let \tilde{k}_i be the composition $\mathbb{R} \xrightarrow{q} S^1 \xrightarrow{k_i} T_i$. Then $\tilde{k}_i(\frac{j}{r_i}) = b_{ij}$. Moreover, it is clear from Figure 6 that $\tilde{k}_i(\frac{j+1/3}{r_i})$ lies in the same disk D_{ij} which contains b_{ij} , whereas $\tilde{k}_i(\frac{j+2/3}{r_i})$ lies in the disk $D_{i,j+1}$ which contains $b_{i,j+1} = \tilde{k}_i(\frac{j+1}{r_i})$. Hence \tilde{k}_i sends $[\frac{j+1/3}{r_i}, \frac{j+2/3}{r_i}]$ onto an arc in Φ_{ij} with one endpoint in D_{ij} and another in $D_{i,j+1}$. Also, $[\frac{j}{r_i}, \frac{j+1/3}{r_i}]$ must go onto an arc in $\Phi_{i,j-1}$ with both endpoints in D_{ij} ; and $[\frac{j+2/3}{r_i}, \frac{j+1}{r_i}]$ onto an arc in $\Phi_{i,j+1}$ with both endpoints in $D_{i,j+1}$. Thus \tilde{k}_i sends each arc $J_{ij} := [\frac{j}{r_i}, \frac{j+1}{r_i}]$ onto an arc in the interior of $\Psi_{ij} := \Phi_{i,j-1} \cup \Phi_{ij} \cup \Phi_{i,j+1}$.

Next, we have $\frac{j}{r_i} = \frac{3jq_i}{r_{i+1}}$, and consequently $J_{ij} = [\frac{j}{r_i}, \frac{j+1}{r_i}] = [\frac{3jq_i}{r_{i+1}}, \frac{3(j+1)q_i}{r_{i+1}}] = J_{i+1,3jq_i} \cup \dots \cup J_{i+1,3(j+1)q_i-1}$. Hence $\tilde{k}_{i+1}(J_{ij})$ lies in $\Psi_{i+1,3jq_i} \cup \dots \cup \Psi_{i+1,3(j+1)q_i-1}$, which is the same as $\Phi_{i+1,3jq_i-1} \cup \dots \cup \Phi_{i+1,3(j+1)q_i}$. But it follows from (2) and from our choice of the lift $D_{0,i+1}$ that the latter union lies in Ψ_{ij} .²⁵ Arguing by induction, it is easy to get from this that each \tilde{k}_l for $l > i$ also sends J_{ij} into Ψ_{ij} . Hence $d(k_i, k_l)$ for $l > i$ is bounded above by the maximum of the diameters of $\Psi_{i1}, \dots, \Psi_{ir_i}$, which by (1) does not exceed $3\epsilon_i$. Since $C^0(S^1, T_1)$ is complete, k_1, k_2, \dots uniformly converge to a continuous map $\mathcal{B}: S^1 \rightarrow T_1$.

Let us show that \mathcal{B} is injective (since S^1 is compact, this will imply that \mathcal{B} is a homeomorphism onto its image). First let us note that $r_i \geq 3^{i-1}$ by (2). Therefore, given distinct points $x, y \in [0, 1)$, there exists an i such that $x \in J_{ij}$ and $y \in J_{ik}$ where $j - k \not\equiv 0, \pm 1, \pm 2, \pm 3 \pmod{r_i}$. Then Ψ_{ij} and Ψ_{ik} are disjoint, but $\tilde{k}_l(x) \in \tilde{k}_l(J_{ij}) \subset \Psi_{ij}$ and $\tilde{k}_l(y) \in \tilde{k}_l(J_{ik}) \subset \Psi_{ik}$ for each $l \geq i$. Writing $\bar{x} = q(x) \in S^1$, we get $\mathcal{B}(\bar{x}) \in \Psi_{ij}$ and $\mathcal{B}(\bar{y}) \in \Psi_{ik}$. Thus $\mathcal{B}(\bar{x}) \neq \mathcal{B}(\bar{y})$.

Let us show that the image of \mathcal{B} equals $\bigcap_{i=1}^{\infty} T_i$. The images of k_i, k_{i+1}, \dots lie in T_i , and hence so does that of \mathcal{B} . Therefore the image of \mathcal{B} lies in $\bigcap_{i=1}^{\infty} T_i$. Let us prove the converse inclusion. Since each T_i is a union of the fundamental domains Φ_{ij} , every point $y \in \bigcap_{i=1}^{\infty} T_i$ is the intersection of some sequence of the form $\Phi_{1j_1} \supset \Phi_{2j_2} \supset \dots$. Let us pick some points $x_i \in J_{ij_i}$. Then $\tilde{k}_l(x_i) \in \Psi_{ij_i}$ for each $l \geq i$ and hence $\mathcal{B}(\bar{x}_i) \in \Psi_{ij_i}$ for each i . Also $y \in \bigcap_{i=1}^{\infty} \Psi_{ij_i}$, and it follows from (1) that $\mathcal{B}(\bar{x}_i) \rightarrow y$. But the image of \mathcal{B} is compact (it is homeomorphic to S^1) and hence closed. Thus it contains y . \square

(b). We assume familiarity with the proof of (a). Let \mathcal{B} and \mathcal{B}' be two embeddings as in (a) corresponding to the same sequence r_1, r_2, \dots . Let T_i, p_i, Φ_{ij} and Ψ_{ij} be as above, and

²⁵In more detail, $\Phi_{i+1,k}$ lies in $\Phi_{i,j-1}$ if $3jq_i \leq k < (3j+1)q_i$; in $\Phi_{i,j+1}$ if $(3j+2)q_i \leq k < (3j+3)q_i$; and in Φ_{ij} if $(3j+1)q_i \leq k < (3j+2)q_i$ as well as for $k = 3jq_i - 1$ and $k = (3j+3)q_i$.

T'_i, p'_i, Φ'_{ij} and Ψ'_{ij} be defined similarly. Let us note that $T'_1 = T_1$ by construction. Since both p_1 and p'_1 are untwisted r_1 -fold covers, there is a homeomorphism $h_1: T_1 \rightarrow T_1$ which sends each Φ'_{1j} onto Φ_{1j} and extends to an orientation preserving PL homeomorphism $S^3 \rightarrow S^3$. Next, it follows from (2), using that both p_2 and p'_2 are untwisted r_2 -fold covers, that there is a homeomorphism $h_2: T_1 \rightarrow T_1$ which keeps ∂T_1 fixed, preserves each Φ_{1j} and sends each $h_1(\Phi'_{2j})$ onto Φ_{2j} . Similarly, there is a homeomorphism $h_3: T_1 \rightarrow T_1$ which keeps $\overline{T_1 \setminus T_2}$ fixed, preserves each Φ_{2j} and sends each $h_2 \circ h_1(\Phi'_{3j})$ onto Φ_{3j} . Continuing in the same fashion, we obtain homeomorphisms $h_4, h_5, \dots: T_1 \rightarrow T_1$. Let $H_k = h_k \circ \dots \circ h_1$. Then $H_k(\Phi'_{ij}) = \Phi_{ij}$ for all $i \leq k$ and H_{k+1} disagrees with H_k only on the interior of T'_k . Also $h_l \circ \dots \circ h_{k+1}$ preserves each Φ_{kj} for $l > k$, and hence by (1) H_l is ε_k -close to H_k for all $l > k$. Since $C^0(T_1, T_1)$ is complete, H_1, H_2, \dots uniformly converge to a continuous map $H: T_1 \rightarrow T_1$.

For each k the restriction of H to $\overline{T_1 \setminus T'_k}$ is a finite composition of homeomorphisms, hence a homeomorphism onto $\overline{T_1 \setminus T_k}$. Therefore H restricts to a homeomorphism between $T_1 \setminus \mathcal{B}'(S^1)$ and $T_1 \setminus \mathcal{B}(S^1)$. Also we have $H(\Phi'_{ij}) = \Phi_{ij}$ for all i and j . It follows that $H(\Psi'_{ij}) = \Psi_{ij}$ for all i and j . On the other hand, since each $r_i \geq 3^{i-1}$, every point $x \in [0, 1)$ is the intersection of some sequence of the form $J_{1,j_1} \supset J_{2,j_2} \supset \dots$. Then $\mathcal{B}(\bar{x})$ lies in the intersection of the sequence $\Psi_{1,j_1} \supset \Psi_{1,j_2} \supset \dots$ and in fact equals this intersection by (1). Similarly $\mathcal{B}'(\bar{x}) = \bigcap_{i=1}^{\infty} \Psi'_{ij_i}$. Hence H sends $\mathcal{B}'(\bar{x})$ onto $\mathcal{B}(\bar{x})$. Thus $H \circ \mathcal{B}' = \mathcal{B}$. In particular, this implies that H restricts to an injective map $\mathcal{B}'(S^1) \rightarrow \mathcal{B}(S^1)$. Since it also restricts to an injective map $T_1 \setminus \mathcal{B}'(S^1) \rightarrow T_1 \setminus \mathcal{B}(S^1)$, we get that H is injective. Since T_1 is compact, H is a homeomorphism. Since h_1 extends to an orientation preserving PL homeomorphism of S^3 and each h_i for $i > 1$ keeps ∂T_1 fixed, H also extends to an orientation preserving homeomorphism $\bar{H}: S^3 \rightarrow S^3$, which is PL outside T_1 . Then \bar{H} is isotopic to the identity by the Alexander trick. \square

(c). Let us fix some $\varepsilon > 0$. By scaling we may assume that the length of the unknot K_0 equals 1. Then the untwisted r_1 -fold cover $p_1: T_1 \rightarrow T_1$ may be chosen so that each $\Phi_{1j} \cap K_0$ is an arc of length $\frac{1}{r_1}$. Hence, as long as the regular neighborhood T_1 of K_0 is chosen to be thin enough, the diameter of each Φ_{1j} will be bounded above by $\frac{1+\varepsilon}{r_1}$.

On the other hand, if the regular neighborhood T_1 of K_0 is chosen to be thin enough and the fake Mazur curve J is drawn appropriately in T_1 (with a sufficiently small clasp), then we may assume that its parameterization $k: S^1 \rightarrow J \subset T_1$ (see the proof of (a)) is ε -close to the map $S^1 \xrightarrow{f} S^1 = K_0 \subset T_1$, where f is defined as follows. Let $\tilde{f}: [0, 1] \rightarrow \mathbb{R}$ be defined by $f(0) = 0$, $f(\frac{1}{6}) = -\frac{1}{2}$, $f(\frac{5}{6}) = \frac{3}{2}$, $f(1) = 1$ and extending linearly. (Thus $f(\frac{1}{3}) = 0$, $f(\frac{1}{2}) = \frac{1}{2}$ and $f(\frac{2}{3}) = 1$.) Then the composition $[0, 1] \xrightarrow{\tilde{f}} \mathbb{R} \xrightarrow{q} \mathbb{R}/\mathbb{Z} = S^1$ factors as $[0, 1] \subset \mathbb{R} \xrightarrow{q} S^1 \xrightarrow{f} S^1$ for a unique map f . The identification $S^1 = K_0$ is chosen so that $f(0) = b$ (the point b is shown in Figure 6). Clearly, the length of the loop $S^1 \xrightarrow{f} S^1 = K_0 \subset T_1$ equals 3. While this does not formally imply any bound on the length of J , it follows that J can be drawn so that its length does not exceed $3 + \varepsilon$.

The map $S^1 \xrightarrow{f} S^1 = K_0 \subset T_1$ precomposed with the r_1 -fold cover $S^1 \rightarrow S^1$, $s \mapsto r_1 s$, lifts with respect to the cover p_1 to a map $f_1: S^1 \rightarrow K_0 \subset T_1$. We may assume that the parameterization $k_1: S^1 \rightarrow K_1 \subset T_1$ of K_1 (see the proof of (a)) is ϵ -close to this f_1 . The loop f_1 is of length 3, and the three paths that each Φ_{1j} cuts out of this loop are of total length $\frac{3}{r_1}$. (These three paths have equal lengths.) It follows that p_1 can be chosen so that the three arcs in $\Phi_{1j} \cap K_1$ are of total length $\leq \frac{3+\epsilon}{r_1}$. Now the untwisted r_2 -fold cover $p_2: T_2 \rightarrow T_1$ may certainly be chosen to satisfy the case $i = 1$ of condition (2), i.e. so that each $\Phi_{1j} \cap T_2$ has only three components and each of them is a union of q_1 of $\Phi_{2,k}$'s. Then we may as well choose p_2 so that each $\Phi_{2,k} \cap K_1$ is an arc of length $\leq \frac{3+\epsilon}{3q_1 r_1} = \frac{3+\epsilon}{r_2}$. Hence, as long as the regular neighborhood T_2 of K_1 is chosen to be thin enough, the diameter of each $\Phi_{2,k}$ will be bounded above by $\frac{(3+\epsilon)+\epsilon}{r_2}$.

Now suppose inductively that for some i each $\Phi_{ij} \cap K_{i-1}$ is an arc of length $\leq \frac{(3+\epsilon)^{i-1}}{r_i}$. The map $S^1 \xrightarrow{f} S^1 = K_0 \subset T_1$ precomposed with the r_i -fold cover $S^1 \rightarrow S^1$, $s \mapsto r_i s$, lifts with respect to the untwisted r_i -fold cover $p_i: T_i \rightarrow T_1$ to a map $f_i: S^1 \rightarrow K_{i-1} \subset T_i$. Since the arc $\Phi_{ij} \cap K_{i-1}$ is of length $\leq \frac{(3+\epsilon)^{i-1}}{r_i}$, the three paths that Φ_{ij} cuts out of f_i are of total length $\leq \frac{3(3+\epsilon)^{i-1}}{r_i}$. It follows that p_i can be chosen so that the three arcs in $\Phi_{ij} \cap K_i$ are of total length $\leq \frac{3(3+\epsilon)^{i-1}+\epsilon}{r_i} \leq \frac{(3+\epsilon)^i}{r_i}$. Arguing as before, we get that T_{i+1} and p_{i+1} can be chosen so that each $\Phi_{i+1,j} \cap K_i$ is an arc of length $\leq \frac{(3+\epsilon)^i}{3q_i r_i} = \frac{(3+\epsilon)^i}{3r_{i+1}}$ and the diameter of each $\Phi_{i+1,j}$ is bounded above by $\frac{(3+\epsilon)^i+\epsilon}{r_{i+1}}$.

Now setting $\epsilon = 0.5$, we have proved that there exists a sequence of K_i and T_i with each Φ_{ij} of diameter $\leq \frac{3.5^{i-1}+0.5}{r_i} \leq \frac{4^{i-1}}{r_i}$. Since $r_1 \geq 1$ and by the hypothesis each $\frac{r_{i+1}}{3r_i} \geq 2$, each $r_i \geq 6^{i-1}$. Then condition (1) holds with $\epsilon_i = (4/6)^{i-1}$. \square

6. CONWAY POTENTIAL FUNCTION

For an m -component PL link L in S^3 let $\Omega_L(x_1, \dots, x_m)$ denote its Conway potential function. Several explicit constructions of Ω_L are known (see references in the introduction of [68]). We have $\Omega_L(x_1, \dots, x_m) \in \mathbb{Z}[x_1^{\pm 1}, \dots, x_m^{\pm 1}]$ for $m > 1$. Also $\Omega_L(x, \dots, x) = \frac{\nabla_K(x - x^{-1})}{x - x^{-1}}$, where $\nabla_K(z) \in \mathbb{Z}[z]$ is called the Conway polynomial. If the components of L are colored in n colors according to a coloring function $c: \{1, \dots, m\} \rightarrow \{1, \dots, n\}$, then $\Omega_L(x_1, \dots, x_n)$ denotes $\Omega_L(x_{c(1)}, \dots, x_{c(m)})$.

Let us recall Conway's identities I (involving strands of the same color) and II (involving strands of possibly distinct colors):²⁶

$$\Omega_{i \times i} - \Omega_{i \times i} = (x_i - x_i^{-1}) \Omega_i \downarrow_i \quad (\text{I})$$

$$\Omega_{i \times j} + \Omega_{i \times j} = (x_i x_j + x_i^{-1} x_j^{-1}) \Omega_i \downarrow_j \quad (\text{II})$$

²⁶To be precise, (I) appears in the papers by Kauffman [49] and Hartley [37] (and in a preliminary form also in the paper by Alexander [2]) and differs in a sign from the identity announced by Conway [26].

We will also frequently use the formula²⁷ for a connected sum with the Hopf link:

$$\Omega \left(\text{Hopf link with strand } j \text{ and } i \right) = (x_i - x_i^{-1}) \cdot \Omega \Big|_i \quad (\Phi)$$

A basis for computation is given by²⁸

$$\Omega_L = 0 \text{ for every split link } L \quad (\text{S})$$

and by the value of Ω on the Hopf link:

$$\Omega \left(\text{Hopf link} \right) = 1. \quad (\text{H})$$

We refer to [21] or [37] for the proofs of the relations (I), (II), (Φ), (S) and (H).²⁹ In the case of two colors, it is easy to calculate Ω by using these five relations as axioms [75; pp. 166–168]. The case of more than two colors is more complicated.³⁰

Let us note that in presence of (Φ), an equivalent form of (H) is given by the value of Ω on the unknot:

$$\Omega_{\bigcirc_i} = \frac{1}{x_i - x_i^{-1}}.$$

Also (H), (II) and (S) easily imply

$$\Omega \left(\text{Hopf link with crossing} \right) = -1 \quad (\text{H}')$$

and similarly (Φ), (II) and (S) imply

$$\Omega \left(\text{Hopf link with crossing} \right) = -(x_i - x_i^{-1}) \cdot \Omega \Big|_i \quad (\Phi')$$

Example 6.1. Let us compute Ω_M , where $M = (J, B)$ is the fake Mazur link (see Figure 7).


M can be made into the Hopf link H by one crossing change (see Figure 7). Then by (I) we have $\Omega_M - \Omega_H = -(x - x^{-1})\Omega_L$, where L is the 3-component link shown in Figure 7. Hence $\Omega_M = 1 - (x - x^{-1})\Omega_L$. By (Φ') we have $\Omega_L = -(y - y^{-1})\Omega_{H_2}$, where

²⁷Here the component that is shown entirely need not be the only component of color j . In fact, the case of the formula where it is the only component of color j easily implies the general case.

²⁸A link L in S^3 is called *split* if there exists a PL 3-ball in S^3 containing a nonempty collection of components of L but not all of them, and disjoint from the remaining components of L .


²⁹Proofs of (I) and (II) from the definition of [21] can also be found in [27; §7.9]. The relation (S) is proved only in a special case in [21], but the general case is proved similarly. The proof of (S) from the definition of [37], which is omitted in [37], can be based on the fact that one of the relations in the Wirtinger presentation of a link group is redundant (see [37; (2.2)]), and consequently for a suitable plane diagram of a split link there are two redundant relations. The relation (Φ) is not proved in [37] but can be easily verified by the methods of [37].

³⁰In the case of three colors Ω can be calculated by using these five axioms, Conway's third identity (which involves four terms, each with three strands) and the value of Ω on the Borromean rings [75]. In the general case several axiomatic characterizations of Ω are known, but they are either non-local [94] or contain a local identity that involves six or more terms [47], [73].

FIGURE 7. The fake Mazur link M and a computation of Ω_M (beginning).

H_2 is shown in Figure 8. By (II) we have $\Omega_{H_2} + \Omega_U = (xy + x^{-1}y^{-1})\Omega_H$, where U is the 2-component unlink (see Figure 8), whence $\Omega_{H_2} = xy + x^{-1}y^{-1}$. Thus we obtain

$$\Omega_M(x, y) = 1 + (xy + x^{-1}y^{-1})(x - x^{-1})(y - y^{-1}).$$

FIGURE 8. The computation of Ω_M (conclusion).

Remark 6.2. Up to a sign, Ω_L is determined by the multi-variable Alexander polynomial Δ_L .³¹ The sign can be recovered for knots (using Theorem 7.2); for 2-component links it is determined by the sign of the linking number when it is nonzero (using Corollary 7.4), and in general by the sign of any shortest non-vanishing $\bar{\mu}$ -invariant of L of the type $\bar{\mu}(1 \dots 12 \dots 2)$ [91; 8.3]. However no such description of the sign of Ω_L is known for links of 3 or more components (compare [56; 1.2, 1.6], [91; 8.2], [93; 1.8], [51]).

This sign might seem to be a minor issue, but it is indispensable for a recursive computation of Ω_L (see Example 6.1), and at the same time it is, in a sense, a much subtler invariant than the multi-variable Alexander polynomial. Indeed, Δ_L is an invariant of the $\mathbb{Z}[\pi_L/\pi'_L]$ -module π'_L/π''_L , where $\pi_L = \pi_1(S^3 \setminus L)$ and $G' = [G, G]$.³² By contrast, the sign of Ω_L is not an invariant of π_L , and not even of $S^3 \setminus L$, because it generally depends on the orientations of S^3 and of the components of L (see [37; 5.6, 5.7]). Nevertheless, the definition of Ω_L in terms of Seifert surfaces [21] deals (as explained in [23]) with a certain presentation of the same module G'/G'' ; and the two definitions of Ω_L in terms of

³¹Namely, $\Omega_L(x_1, \dots, x_m) = \pm x_1^{k_1} \dots x_m^{k_m} \Delta_L(x_1^2, \dots, x_m^2)$ for $m > 1$ and $(x - x^{-1})\Omega_K(x) = \pm x^k \Delta_K(x^2)$ for a knot K , where the integers k, k_1, \dots, k_m are uniquely determined by the symmetry relation $\Omega_L(x_1, \dots, x_m) = (-1)^m \Omega_L(x_1^{-1}, \dots, x_m^{-1})$.

³²Namely, Δ_L is the gcd of all $n \times n$ minors of a presentation matrix of this module, where n is the number of generators. Note that $\mathbb{Z}[\pi_L/\pi'_L] \simeq \mathbb{Z}[H_1(S^3 \setminus L)] \simeq \mathbb{Z}[t_1^{\pm 1}, \dots, t_m^{\pm 1}]$ and $\pi'_L/\pi''_L \simeq H_1(\widetilde{S^3 \setminus L})$, where \tilde{X} denotes the universal abelian cover of X .

the Fox calculus [37], [9; §2] deal with certain presentations of a closely related module.³³ All three presentations have to be chosen very carefully, because interchanging any two relators would have the effect of changing the sign of Ω_L ; and if adding a redundant relator 1 is allowed, then any two relators can be interchanged, as rows in the presentation matrix, by a repeated application of the operation of adding a multiple of a row to another row (see [96; p. 118]). Therefore it is not surprising that the definition of Ω_L that appears to be the closest to being “conceptual” [94] proceeds by redoing everything in terms of determinants.

Problem 6.3. (a) Can Ω_L be constructed as an invariant of the $\mathbb{Z}[\pi_L/\pi'_L]$ -module π'_L/π''_L endowed with some additional structure?

(b) Is there a “conceptual” proof (as opposed to a verification of the axioms) that the two definitions of Ω_L in terms of the Fox calculus [37], [9; §2] and the third one in terms of Seifert surfaces [21] are all equivalent to each other?

7. SOME KNOWN TOOLS FOR COMPUTING Ω

Theorem 7.1 (Torres–Hartley [37]). Let $L = (K_1, \dots, K_m, K)$ be a PL link, where $m \geq 1$, and let $l_i = \text{lk}(K, K_i)$. Let $L' = (K_1, \dots, K_m)$. Then

$$(x_1^{l_1} \cdots x_m^{l_m} - x_1^{-l_1} \cdots x_m^{-l_m}) \Omega_{L'}(x_1, \dots, x_m) = \Omega_L(x_1, \dots, x_m, 1).$$

In particular, if all $l_i = 0$, then $\Omega_L(x_1, \dots, x_m, 1) = 0$; and if $m = 1$ and $l_1 = 1$, then $\Omega_L(x, 1) = \nabla_{K_1}(x - x^{-1})$.

In the case $m = 0$ we have

Theorem 7.2 (Alexander [2]). If K is a PL knot, then $\nabla_K(0) = 1$.

Proof. (I) implies that $\nabla_K(0)$ is the same for all knots K . By (H) and (Φ) $\nabla_{\bigcirc} = 1$. \square

Example 7.3. Let $L = (K_1, K_2)$ be a 2-component link and $n = \text{lk}(L)$. By Theorem 7.1

$$\Omega_L(x, 1) = \frac{x^n - x^{-n}}{x - x^{-1}} \nabla_{K_1}(x - x^{-1}).$$

Here

$$x^n - x^{-n} = \begin{cases} (x + x^{-1})F_n(x - x^{-1}) & \text{if } n \text{ is even,} \\ L_n(x - x^{-1}) & \text{if } n \text{ is odd,} \end{cases}$$

where $F_n(z)$ is the n th Fibonacci polynomial, defined for $n \geq 0$ by $F_0(z) = 0$, $F_1(z) = 1$ and $F_{n+1}(z) = zF_n(z) + F_{n-1}(z)$, and $L_n(z)$ is the n th Lucas polynomial, defined for $n \geq 0$ by the same recursion $L_{n+1}(z) = zL_n(z) + L_{n-1}(z)$ with different initial values $L_0(z) = 2$, $L_1(z) = z$. The definitions are extended for negative indexes by $F_{-n}(z) = (-1)^{n+1}F_n(z)$ and $L_{-n}(z) = (-1)^nL_n(z)$.

Corollary 7.4. If L is a 2-component link, then $\Omega(1, 1) = \text{lk}(L)$.

³³Namely, $H_1(\widetilde{S^3 \setminus L}, p^{-1}(b))$, where $b \in S^3 \setminus L$ and $p: \widetilde{S^3 \setminus L} \rightarrow S^3 \setminus L$ is the covering map. See [48] concerning the relation between the two modules.

Proof. $\frac{x^n - x^{-n}}{x - x^{-1}} = x^{-(n-1)} + x^{-(n-3)} + \dots + x^{n-1}$, which contains $n = \text{lk}(L)$ terms. \square

Let $L = (K_1, \dots, K_m, K)$ and $L' = (Q, Q_1, \dots, Q_n)$ be PL links in S^3 , where Q is unknotted. Let N and N' be regular neighborhoods of K and Q in the complements to the other components, and let E' be the solid torus $\overline{S^3 \setminus N'}$. Let $h: E' \rightarrow N$ be a homeomorphism that sends a longitude of E' (which is a meridian of N') onto a longitude of N . Then $\Lambda := (K_1, \dots, K_m, h(Q_1), \dots, h(Q_n))$ is called the *splice* of L and L' .

Theorem 7.5 (Cimasoni [22]). *Let Λ be the splice of $L = (K_1, \dots, K_m, K)$ and $L' = (Q, Q_1, \dots, Q_n)$, and let $\mu_i = \text{lk}(K, K_i)$ and $\nu_i = \text{lk}(Q, Q_i)$. Assume that $n \geq 1$, and either $m \geq 1$ or at least one of the ν_i is nonzero. Then*

$$\Omega_\Lambda(x_1, \dots, x_m, y_1, \dots, y_n) = \Omega_L(x_1, \dots, x_m, y_1^{\nu_1} \cdots y_n^{\nu_n}) \Omega_{L'}(x_1^{\mu_1} \cdots x_m^{\mu_m}, y_1, \dots, y_n).$$

Theorem 7.5 can be used to reprove the relation (Φ) , by representing the connected sum of a link L with the Hopf link as a splice of L with a connected sum of two Hopf links (whose Ω can be computed directly). In turn, (Φ) and Theorem 7.5 easily yield a formula for an arbitrary connected sum:

Corollary 7.6 (Cimasoni [22], [21]). *Let $\Lambda = (K_1, \dots, K_m, K \# Q, Q_1, \dots, Q_n)$ be the connected sum of PL links $L = (K_1, \dots, K_m, K)$ and $L' = (Q, Q_1, \dots, Q_n)$, where L' is disjoint from some 3-ball containing L , along an arbitrary band joining K and Q and otherwise disjoint from L and L' . Then*

$$\Omega_\Lambda(x_1, \dots, x_m, y, y_1, \dots, y_n) = (y - y^{-1}) \Omega_L(x_1, \dots, x_m, y) \Omega_{L'}(y, y_1, \dots, y_n).$$

In particular, when $n = 0$, we have $\Omega_\Lambda(x_1, \dots, x_n, y) = \Omega_L(x_1, \dots, x_n, y) \nabla_Q(y - y^{-1})$ and $\nabla_\Lambda(z) = \nabla_L(z) \nabla_Q(z)$.

Proof. Let $L'' = (B, Q, Q_1, \dots, Q_m)$, where B is the boundary of a small embedded disk intersecting Q transversely in one interior point with positive sign (that is, so that $\text{lk}(B, Q) = +1$) and otherwise disjoint from L' . By (Φ) $\Omega_{L''}(x, y, y_1, \dots, y_n, y) = (y - y^{-1}) \Omega_{L'}(y, y_1, \dots, y_n)$. On the other hand, Λ is easily seen to be the splice of L and L'' , and the assertion follows from Theorem 7.5. \square

Theorem 7.5 can also be used to obtain a formula for adding several push-offs of a component to the link (compare related formulas in [94], [22]):

Corollary 7.7. *Let $L = (K_1, \dots, K_m, K)$ be a PL link and let $l_i = \text{lk}(K, K_i)$. Let $L_+ = (K_1, \dots, K_m, K_+^1, \dots, K_+^n)$, where $n \geq 1$ and K_+^1, \dots, K_+^n are pairwise disjoint parallel pushoffs of K . Then*

$$\Omega_{L_+}(x_1, \dots, x_m, y_1, \dots, y_n) = (x_1^{l_1} \cdots x_m^{l_m} - x_1^{-l_1} \cdots x_m^{-l_m})^{n-1} \Omega_L(x_1, \dots, x_m, y_1 \cdots y_n).$$

Proof. Let $H = (C, B)$ be the Hopf link, and let $H_+ = (C, B_+^1, \dots, B_+^n)$, where B_+^1, \dots, B_+^n are pairwise disjoint parallel pushoffs of B . Then L_+ is the splice of L and H_+ , and $\Omega_{H_+}(u, v_1, \dots, v_n) = (u - u^{-1})^{n-1}$ by using (Φ) . \square

Let us note that for $n = 0$ Corollary 7.7 does not make sense as stated (because of a potential division by 0), but this case is addressed by Theorem 7.1.

Proposition 7.8 (Hartley [37]). *If L' is obtained from L by reversing the orientation of the first component, then $\Omega_{L'}(x_1, x_2, \dots, x_n) = -\Omega_L(x_1^{-1}, \dots, x_n)$.*

Corollary 7.7, Theorem 7.1 and Proposition 7.8 imply

Corollary 7.9. *Let $L = (K_1, \dots, K_m, K)$ be a PL link, let $L' = (K_1, \dots, K_m, K, -K_+)$, where $-K_+$ is a parallel pushoff of K with orientation reversed, and let $l_i = \text{lk}(K, K_i)$. Then*

$$\Omega_{L'}(x_1, \dots, x_m, y, y) = -(x_1^{l_1} \cdots x_m^{l_m} - x_1^{-l_1} \cdots x_m^{-l_m})^2 \Omega_{L \setminus K}(x_1, \dots, x_m).$$

Corollary 7.9 and Conway's identity I imply

Corollary 7.10. *Let $L = (K_1, \dots, K_m, K)$ be a PL link, let $L' = (K_1, \dots, K_m, W)$ be its possibly twisted Whitehead doubling along the last component, and let $l_i = \text{lk}(K, K_i)$. Then*

$$\Omega_{L'}(x_1, \dots, x_m, y) = \pm(y - y^{-1})(x_1^{l_1} \cdots x_m^{l_m} - x_1^{-l_1} \cdots x_m^{-l_m})^2 \Omega_{L \setminus K}(x_1, \dots, x_m).$$

Theorem 7.11 (Murasugi–...–Miyazawa [72]). *Let (B, K_1, \dots, K_m) be an $(m + 1)$ -component PL link, where B is the unknot, and let \tilde{K}_j denote the link $p_r^{-1}(K_j)$, where $p_r: (S^3, B) \rightarrow (S^3, B)$ is the r -fold cover ramified over B . Then*

$$\Omega_{(\tilde{K}_1, \dots, \tilde{K}_m)}(x_1, \dots, x_m) = i^{(r-1)(1-\lambda)} \Omega_{(K_1, \dots, K_m)}(x_1, \dots, x_m) \prod_{j=1}^{r-1} \Omega_{(B, K_1, \dots, K_m)}(\xi^j, x_1, \dots, x_m),$$

where $i = \sqrt{-1}$, $\xi = e^{\pi i/r}$ and $\lambda = \text{lk}(B, K_1) + \dots + \text{lk}(B, K_m)$.

Corollary 7.12. *Let $M = (J, B)$ be a 2-component link with linking number 1 and let $p_r: (S^3, B) \rightarrow (S^3, B)$ be the r -fold cover ramified over B . Let $J_r = p_r^{-1}(J)$ and $M_r = (J_r, B)$. Let $\xi = e^{\pi i/r}$. Then*

- (a) $\Omega_{M_r}(x, y) = \prod_{j=0}^{r-1} \Omega_M(x, y^{\frac{1}{r}} \xi^j)$;
- (b) $\Omega_{J_r}(x) = \frac{1}{x-x^{-1}} \prod_{j=0}^{r-1} \Omega_M(x, \xi^j)$.

Proof. (a). Let $M_+ = (J, B, B_+)$ be obtained from M by adding a parallel pushoff B_+ of B . By Corollary 7.7 $\Omega_{M_+}(x, y, z) = (x - x^{-1})\Omega_M(x, yz)$. Let M_r^+ be the 2-colored link (J_r, B_r) , where $B_r = p_r^{-1}(B_+)$. Then by Theorem 7.11 $\Omega_{M_r^+}(x, y) = \Omega_M(x, y) \prod_{j=1}^{r-1} \Omega_{M_+}(x, \xi^j, y)$. Hence $\Omega_{M_r^+}(x, y) = (x - x^{-1})^{r-1} \prod_{j=0}^{r-1} \Omega_M(x, y \xi^j)$. But clearly $B_r = (B_+^1, \dots, B_+^r)$, the link of r parallel pushoffs of B . Hence by Corollary 7.7 $\Omega_{M_r^+}(x, y) = (x - x^{-1})^{r-1} \Omega_{M_r}(x, y^r)$. Thus we get that $\Omega_{M_r}(x, y) = \prod_{j=0}^{r-1} \Omega_M(x, y^{\frac{1}{r}} \xi^j)$. \square

(b). Although (b) follows from (a) and Theorem 7.1, it is easier to prove it by observing that it is a direct consequence of Theorems 7.11 and 7.1. \square

Remark 7.13. Explicit computations may be aided by the following congruence [74; Proposition 4.2]. If $P(x, y) \in \mathbb{Z}[x^{\pm 1}, y^{\pm 1}]$ ³⁴ and $r = p^n m$, where p is a prime, $n > 0$ and $(m, p) = 1$, then $\prod_{j=0}^{r-1} P(x, e^{2\pi j/r}) \equiv \left(\prod_{k=0}^{m-1} P(x, e^{2\pi k/m}) \right)^{p^n} \pmod{p}$.

8. COMPUTING Ω FOR RAMIFIED COVERS OF THE FAKE MAZUR LINK

The following notation will be used throughout this section. Let $M = (J, B)$ be the fake Mazur link (see Figure 7) and let $p_r: (S^3, B) \rightarrow (S^3, B)$ be the r -fold cover ramified over B . Let $J_r = p_r^{-1}(J)$ and $M_r = (J_r, B)$.

Lemma 8.1. *Let $\zeta = e^{2i\pi/r}$. Then*

$$(a) \Omega_{M_r}(x, y) = \prod_{j=0}^{r-1} \left(1 + (xy)^{\frac{2}{r}} \zeta^j - x^{-1} y^{-\frac{2}{r}} \zeta^{-j} \right) (x - x^{-1}) - (x - x^{-1})^2;$$

$$(b) \Omega_{J_r}(x) = \frac{1}{x-x^{-1}} \prod_{j=0}^{r-1} \left(1 + (x\zeta^j - x^{-1}\zeta^{-j})(x - x^{-1}) - (x - x^{-1})^2 \right).$$

Proof. By Example 6.1

$$\Omega_M(x, y) = 1 + (xy + x^{-1}y^{-1})(x - x^{-1})(y - y^{-1}) = 1 + (xy^2 - x^{-1}y^{-2})(x - x^{-1}) - (x - x^{-1})^2.$$

Now the assertion follows from Corollary 7.12. \square

Proposition 8.2. (a) $\nabla_{J_r}(z) = r \prod_{k=1}^{r-1} \left(\frac{1}{\zeta^{2k} - \zeta^k} - z^2 \right)$, where $\zeta = e^{2\pi i/r}$.

$$(b) \Omega_{M_r}(x, y) = \nabla_{J_r}(x - x^{-1}) + (x - x^{-1})^r (y - y^{-1}) ((-1)^{r+1} x^r y + x^{-r} y^{-1}).$$

$$(c) x^r y + (-1)^{r+1} x^{-r} y^{-1} = (y - y^{-1}) F_{r+1}(x - x^{-1}) + (x^{-1} y + xy^{-1}) F_r(x - x^{-1}).$$

$$(d) \nabla_{M_r}(z) = z \nabla_{J_r}(z) + (-1)^{r+1} z^{r+2} L_{r+1}(z).$$

Here F_r and L_r are the Fibonacci and Lucas polynomials (see Example 7.3).

Proof. (a). We start from the expression of Lemma 8.1(b) and multiply out the factors corresponding to j and to $r - j$, for $j = 1, \dots, \lfloor \frac{r-1}{2} \rfloor$. There remain:

- an additional factor corresponding to $j = 0$, but it equals 1; and
- an additional factor corresponding to $\frac{r}{2}$ when r is even.

The result is:

$$\nabla_{J_r}(z) = \begin{cases} \prod_{k=1}^n (1 + 2z^2(1 + c_k - 2c_k^2) + 2z^4(1 - c_k)) & \text{if } r = 2n + 1; \\ (1 - 2z^2) \prod_{k=1}^{n-1} (1 + 2z^2(1 + c_k - 2c_k^2) + 2z^4(1 - c_k)) & \text{if } r = 2n, \end{cases}$$

where $c_k = \cos \frac{2\pi k}{r}$. Using the trigonometric identities $\cos(2\varphi) = 1 - 2\sin^2 \varphi$ and $\sin(3\varphi) = 3\sin \varphi - 4\sin^3 \varphi$ this formula can be simplified as follows:

$$\nabla_{J_r}(z) = \begin{cases} \prod_{k=1}^n (1 + 4\sin(k\alpha)\sin(3k\alpha)z^2 + 4\sin^2(k\alpha)z^4) & \text{if } r = 2n + 1; \\ (1 - 2z^2) \prod_{k=1}^{n-1} (1 + 4\sin(k\alpha)\sin(3k\alpha)z^2 + 4\sin^2(k\alpha)z^4) & \text{if } r = 2n, \end{cases}$$

where $\alpha = \pi/r$. Let $x = 2\sin(k\alpha)z^2$, $\lambda = \sin(3k\alpha)$. Then $1 + 2\lambda x + x^2 = (r_+ - x)(r_- + x)$, where $r_{\pm} = -\lambda \pm \sqrt{\lambda^2 - 1} = -\sin(3k\alpha) \pm i\cos(3k\alpha) = i(i\sin(3k\alpha) \pm \cos(3k\alpha))$. Thus

³⁴ P has no negative exponents in [74], but it is easy to see that this hypothesis is redundant.

$r_+ = ie^{3k\alpha i}$ and $r_- = ie^{(\pi-3k\alpha)i} = ie^{3(\pi-k\alpha)i} = ie^{3(r-k)\alpha i}$. On the other hand, $\sin(k\alpha) = \sin(\pi - k\alpha) = \sin((r-k)\alpha)$. It follows that

$$\nabla_{J_r}(z) = \prod_{k=1}^{r-1} (ie^{3k\alpha i} - 2\sin(k\alpha)z^2).$$

Now using the trigonometric identity³⁵ $\prod_{k=1}^{r-1} \sin(k\alpha) = \frac{r}{2^{r-1}}$ we get

$$\nabla_{J_r}(z) = r \prod_{k=1}^{r-1} \left(\frac{ie^{3k\alpha i}}{2\sin(k\alpha)} - z^2 \right).$$

Finally, $\frac{ie^{3k\alpha i}}{2\sin(k\alpha)} = \frac{-e^{3k\alpha i}}{e^{k\alpha i} - e^{-k\alpha i}} = \frac{1}{e^{-4k\alpha i} - e^{-2k\alpha i}}$. We have $e^{-2k\alpha i} = e^{(2\pi-2k\alpha)i} = e^{2(r-k)\alpha i}$ and similarly $e^{-4k\alpha i} = e^{4(r-k)\alpha i}$. Hence

$$\nabla_{J_r}(z) = r \prod_{k=1}^{r-1} \left(\frac{1}{e^{4k\alpha i} - e^{2k\alpha i}} - z^2 \right).$$

□

(b). Since $\text{lk}(M_r) = 1$, the exponent of y in every term of $\Omega_{M_r}(x, y)$ must be even [68; Lemma 2.1]. Each of the r factors of the product in Lemma 8.1(a) is a sum of monomials of three types: cx^k , $cx^k y^{\frac{2}{r}}$ and $cx^k y^{-\frac{2}{r}}$, where $k \in \mathbb{Z}$ and $c \in \mathbb{C}$. A product of r such monomials $m_0 \cdots m_{r-1}$ can have a nonzero even integer exponent of y only when all m_i are of the second type or all m_i are of the third type. Since $\zeta^0 \zeta^1 \cdots \zeta^{r-1} = \zeta^{r(r-1)/2} = (-1)^{r+1}$, the sum of all such products is $(-1)^{r+1} x^r y^2 (x - x^{-1})^r$ in the first case and $-x^{-r} y^{-2} (x - x^{-1})^r$ in the second case. Thus $\Omega_{M_r}(x, y) = ((-1)^{r+1} x^r y^2 - x^{-r} y^{-2}) (x - x^{-1})^r + L(x)$ for some $L(x) \in \mathbb{Z}[x^{\pm 1}]$. To eliminate $L(x)$ let us note that $\Omega_{M_r}(x, 1) = ((-1)^{r+1} x^r - x^{-r}) (x - x^{-1})^r + L(x)$. Hence we get $\Omega_{M_r}(x, y) - \Omega_{M_r}(x, 1) = (x - x^{-1})^r ((-1)^{r+1} x^r (y^2 - 1) + x^{-r} (1 - y^{-2})) = (x - x^{-1})^r (y - y^{-1}) ((-1)^{r+1} x^r y + x^{-r} y^{-1})$. Now the assertion follows from Theorem 7.1. □

(c). This follows from the formula $\{x^n M\} = F_n(\{x\})\{xM\} + F_{n-1}(\{x\})\{M\}$, which in turn follows from [68; Lemma 2.5(a)]. □

(d). This follows from (b) and (c) using that $L_{r+1}(z) = zF_{r+1}(z) + 2F_r(z)$. □

Corollary 8.3. $\nabla_{M_r} \neq z\nabla_{J_r}$.

Proof. It is easy to see from Proposition 8.2 that $\nabla_{M_r}(z)$ is a polynomial of degree $2r+3$, and $\nabla_{J_r}(z)$ is a polynomial of degree $2r-2$. □

Proposition 8.2(a) gives a description of the polynomial ∇_{J_r} in terms of its roots. But we are more interested in its coefficients, which do not seem to be easily obtainable from

³⁵Let $\zeta = e^{2\pi i/r}$ and $\xi = e^{\pi i/r}$. Then $\prod_{k=1}^{r-1} \sin(k\alpha) = \prod_{k=1}^{r-1} \frac{\xi^k - \xi^{-k}}{2i} = (2i)^{1-r} \xi^{r(r-1)/2} \prod_{k=1}^{r-1} (1 - \xi^{-2k}) = (2i)^{1-r} e^{(r-1)\pi i/2} \prod_{k=1}^{r-1} (1 - \zeta^{-k}) = 2^{1-r} \prod_{k=1}^{r-1} (1 - \zeta^k)$. On the other hand, by dividing both sides of $x^r - 1 = \prod_{k=0}^{r-1} (x - \zeta^k)$ by $x - 1$ we get $1 + x + \cdots + x^{r-1} = \prod_{k=1}^{r-1} (x - \zeta^k)$, and by substituting $x = 1$ we get $r = \prod_{k=1}^{r-1} (1 - \zeta^k)$.

this description (except for specific low values of r). To get closer to understanding the coefficients of ∇_{J_r} we will take a different approach.

Proposition 8.4.

$$\sum_{r=1}^{\infty} \nabla_{J_r}(z)x^r = \frac{x}{1+z^2(x-x^2)} \left(\frac{x}{1-x} + \frac{1}{1+z^2x} \right).$$

This result is not used in the present paper, but it might be useful for further attacks on the Rolfsen Problem.




FIGURE 9. Recursion for ∇_{A_n} .

Proof. Let $A_n = J_n$ and let B_n , C_n , D_n , E_n and F_n be its modifications shown in Figure 9. Here A_n involves n repeats of the Fox–Artin tangle:



and so is defined for $n \geq 0$; B_n and C_n each involve $n - 1$ repeats of the same tangle, and so are defined for $n \geq 1$; and D_n , E_n and F_n each involve $n - 2$ repeats of the same tangle, and so are defined for $n \geq 2$. From Conway’s first relation (I) and the relation (Φ) we obtain

$$\begin{aligned} \nabla_{A_n} &= \nabla_{B_n} - z\nabla_{C_n}, & n \geq 1 \\ \nabla_{B_n} &= \nabla_{B_{n-1}} - z\nabla_{D_n}, & n \geq 2 \\ \nabla_{C_n} &= \nabla_{D_n} - z\nabla_{E_n}, & n \geq 2 \\ \nabla_{D_n} &= \nabla_{F_n} + z\nabla_{B_{n-1}}, & n \geq 2 \\ \nabla_{E_n} &= \nabla_{A_{n-2}} + z\nabla_{C_{n-1}}, & n \geq 2 \\ \nabla_{F_n} &= -z\nabla_{B_{n-2}}, & n \geq 3. \end{aligned}$$

It is easy to see from Figure 10 that B_1 is the unknot (whence $\nabla_{B_1} = 1$), C_1 is the two-component unlink (whence $\nabla_{C_2} = 0$) and each of C_2 and D_2 is the Hopf link (whence $\nabla_{C_2} = z$ and $\nabla_{B_2} = \nabla_{B_1} - z\nabla_{D_2} = 1 - z^2$).




FIGURE 10. Beginning of the recursion.

Setting $a_n = \nabla_{A_n}$, $b_n = \nabla_{B_n}$ and $c_n = \nabla_{C_n}$, we obtain a system of recurrences

$$\begin{aligned} a_n &= b_n - zc_n, & n \geq 1 \\ b_n &= (1 - z^2)b_{n-1} + zb_{n-2}, & n \geq 3 \\ c_n &= z(b_{n-1} - 2b_{n-2}) - z^2(c_{n-1} - c_{n-2}), & n \geq 3 \end{aligned}$$

with initial conditions $b_1 = 1$, $b_2 = 1 - z^2$, $c_1 = 0$ and $c_2 = z$.³⁶ Solving the second and third equations, we obtain

$$\begin{aligned} b_n &= 1 - z^2 + z^4 - z^6 + \dots + (-z^2)^{n-1} \\ \sum_{n=1}^{\infty} c_n x^n &= \frac{1}{1 + z^2(x - x^2)} \sum_{n=2}^{\infty} z(b_{n-1} - 2b_{n-2})x^n, \end{aligned}$$

where $b_0 = 0$. Using that $2b_{n-2} - b_{n-1} = b_{n-1} - 2(-z^2)^{n-2}$ for $n \geq 2$, we get

$$\sum_{n=1}^{\infty} a_n x^n = \frac{1}{1 + z^2(x - x^2)} \left(\sum_{n=1}^{\infty} b_n x^n (1 + z^2(x - x^2)) + z^2 \sum_{n=2}^{\infty} (b_{n-1} - 2(-z^2)^{n-2}) x^n \right).$$

Now observe that $b_n + z^2 b_{n-1} = 1$ and $z^2 \sum_{n=1}^{\infty} b_n x^n (x - x^2) = z^2 \sum_{n=1}^{\infty} (b_n - b_{n-1}) x^{n+1} = z^2 \sum_{n=1}^{\infty} (-z^2)^{n-1} x^{n+1} = - \sum_{n=2}^{\infty} (-z^2)^{n-1} x^n$. Hence we get

$$\sum_{n=1}^{\infty} a_n x^n = \frac{1}{1 + z^2(x - x^2)} \left(\sum_{n=1}^{\infty} x^n - \sum_{n=2}^{\infty} (-z^2)^{n-1} x^n + 2 \sum_{n=2}^{\infty} (-z^2)^{n-1} x^n \right).$$

Thus

$$\sum_{n=1}^{\infty} \nabla_{J_n}(z) x^n = \frac{x}{1 + z^2(x - x^2)} \left(\frac{1}{1 - x} + \frac{1}{1 + z^2 x} - 1 \right).$$

□

9. FORMAL LIMITS OF RATIONAL POWER SERIES

The following is well-known, though not necessarily in the form of this particular assertion.

³⁶Note that the same system of recurrences with $z = x - x^{-1}$ and with a different set of initial conditions applies also to $a_n = \Omega_{M_n}(x, y)$.

Lemma 9.1. *A formal power series $R = r_0 + r_1x + r_2x^2 + \dots \in \mathbb{Z}[[x]]$ satisfies a linear recurrence of the form $r_k = q_1r_{k-1} + \dots + q_nr_{k-n}$ for each $k > m$, where $q_1, \dots, q_n \in \mathbb{Z}$ (here in the case $n > m + 1$ we set $r_{-1} = r_{-2} = \dots = r_{m+1-n} = 0$), if and only if $R = P/Q$, where $P, Q \in \mathbb{Z}[x]$, $\deg Q \leq n$ and $\deg P \leq m$.*

Proof. Suppose that R is of the form P/Q , where P and Q are as above. Since Q is invertible in $\mathbb{Z}[[x]]$, we have $Q(0) = \pm 1$; by changing the signs of P and Q we may assume that $Q(0) = 1$. Let us write $Q = 1 - q_1x - \dots - q_nx^n$. Then

$$P = QR = (1 - q_1x - \dots - q_nx^n)(r_0 + r_1x + \dots) = r_0 + (r_1 - q_1r_0)x + (r_2 - q_1r_1 - q_2r_0)x^2 + \dots + (r_n - q_1r_{n-1} - \dots - q_nr_0)x^n + (r_{n+1} - q_1r_{n+2} - \dots - q_nr_1)x^{n+1} + \dots \quad (*)$$

Since $\deg P \leq m$, the coefficients at x^i for $i > m$ must be zero, so we obtain the desired recurrence relation.

Conversely, suppose that the said recurrence relation is satisfied for $k > m$, and let $Q = 1 - q_1x - \dots - q_nx^n$. Then $P := QR$ is a polynomial of degree $\leq m$ (see (*)). \square

Theorem 9.2. *Let $P_1, Q_1, P_2, Q_2, \dots \in \mathbb{Z}[x]$, where each $Q_n \in \mathbb{Z}[[x]]^\times$ (in other words, $Q_n(0) = \pm 1$). Suppose that the P_i converge³⁷ to 0 in $\mathbb{Z}[[x]]$ but each $P_i \neq 0$. Then there exists a function $f: \mathbb{N} \rightarrow \mathbb{N}$ such that for any increasing sequence r_1, r_2, \dots satisfying $r_{i+1} \geq f(r_i)$ for each i ,*

$$(a) \text{ the infinite product}^{38} R := \left(1 + \frac{P_{r_1}}{Q_{r_1}}\right) \left(1 + \frac{P_{r_2}}{Q_{r_2}}\right) \dots;$$

$$(b) \text{ the infinite sum}^{39} R := \frac{P_{r_1}}{Q_{r_1}} + \frac{P_{r_2}}{Q_{r_2}} + \dots$$

is not a rational power series.

One cannot do without selecting a subsequence in general. For instance,

$$S := \prod_{n=1}^{\infty} \left(1 + \frac{x^{n-1} - x^{n+1}}{1 - x^{2n}}\right)$$

is a rational power series. (Indeed, $S = \prod_{n=1}^{\infty} \frac{(1+x^{n-1})(1-x^{n+1})}{(1-x^n)(1+x^n)} = \frac{2}{1-x}$.) Also, if $c_n = \sum_{d|n} \mu(d)2^{n/d}$, where $\mu(n)$ is the Möbius function⁴⁰, then

$$S := \sum_{n=1}^{\infty} \frac{c_n x^n}{1 - x^n}$$

³⁷In the usual topology on $\mathbb{Z}[[x]]$, that is, its topology as the inverse limit of the discrete rings $\mathbb{Z}[x]/(x^i)$. More explicitly, the condition $P_i \rightarrow 0$ is saying the for each k there exists an n such that for each $i > n$, P_i is divisible by x^k .

³⁸That is, the limit of the finite products in the same topology on $\mathbb{Z}[[x]]$. Its existence is clearly guaranteed by the condition $P_i/Q_i \rightarrow 0$.

³⁹That is, the limit of the finite sums in the same topology on $\mathbb{Z}[[x]]$. Its existence is clearly guaranteed by the condition $P_i/Q_i \rightarrow 0$.

⁴⁰ $\mu(n) = (-1)^k$ if n is a product of k distinct primes and $\mu(n) = 0$ if n divisible by a square of a prime.

is a rational power series. Indeed, by the Möbius inversion formula $\sum_{d|n} c_d = 2^n$, whence $S = \sum_{n=1}^{\infty} c_n(x^n + x^{2n} + x^{3n} + \dots) = 2x + 2^2x^2 + 2^3x^3 + \dots = \frac{2x}{1-2x}$.

Proof. We will prove (a) and (b) simultaneously. Let $n_i = \deg Q_1 \cdots Q_i$ and

$$m_i = \begin{cases} \deg(P_1 + Q_1) \cdots (P_i + Q_i) & \text{in (a),} \\ \max\{\deg Q_1 \cdots Q_{j-1} P_j Q_{j+1} \cdots Q_i \mid j = 1, \dots, i\} & \text{in (b).} \end{cases}$$

Let us define $f: \mathbb{N} \rightarrow \mathbb{N}$ as follows: let $f(i)$ be such that P_j is divisible by $x^{\max(2n_i, m_i^2)+1}$ for all $j \geq f(i)$. Let r_1, r_2, \dots be any increasing sequence satisfying $r_{i+1} \geq f(r_i)$ for each i . We have $R = \lim R_i$, where

$$R_i = \begin{cases} \left(1 + \frac{P_{r_1}}{Q_{r_1}}\right) \cdots \left(1 + \frac{P_{r_i}}{Q_{r_i}}\right) = \frac{(P_{r_1} + Q_{r_1}) \cdots (P_{r_i} + Q_{r_i})}{Q_{r_1} \cdots Q_{r_i}} & \text{in (a)} \\ \frac{P_{r_1}}{Q_{r_1}} + \dots + \frac{P_{r_i}}{Q_{r_i}} = \frac{P_{r_1} Q_{r_2} \cdots Q_{r_i} + \dots + Q_{r_1} \cdots Q_{r_{i-1}} P_{r_i}}{Q_{r_1} \cdots Q_{r_i}} & \text{in (b).} \end{cases}$$

Suppose that $R = P/Q$, where $\deg P = M$ and $\deg Q = N$. Choose i_0 so that $R - R_i$ is divisible by $x^{\max(2M, N^2)+1}$ for all $i \geq i_0$. Let us fix some $i \geq i_0$. For each $j > i$ we have $r_j \geq r_{i+1} \geq f(r_i)$ and hence P_{r_j} is divisible by $x^{\max(2n_i, m_i^2)+1}$. Let $n = \deg Q_{r_1} \cdots Q_{r_i}$ and let

$$m = \begin{cases} \deg(P_{r_1} + Q_{r_1}) \cdots (P_{r_i} + Q_{r_i}) & \text{in (a),} \\ \max\{\deg Q_{r_1} \cdots Q_{r_{j-1}} P_j Q_{r_{j+1}} \cdots Q_{r_i} \mid j = 1, \dots, i\} & \text{in (b).} \end{cases}$$

Then $n \leq n_{r_i}$ and $m \leq m_{r_i}$, and hence P_{r_j} is divisible by $x^{\max(2n, m^2)+1}$ for each $j > i$. It easily follows from this that $R - R_i$ is also divisible by $x^{\max(2n, m^2)+1}$. In more detail, in (a) we note that $\left(1 + \frac{P_{r_{i+1}}}{Q_{r_{i+1}}}\right) \cdots \left(1 + \frac{P_{r_j}}{Q_{r_j}}\right) - 1$ is divisible by $x^{\max(2n, m^2)+1}$ for each $j > i$, and hence so is $\left(1 + \frac{P_{r_{i+1}}}{Q_{r_{i+1}}}\right) \left(1 + \frac{P_{r_{i+2}}}{Q_{r_{i+2}}}\right) \cdots - 1 = \frac{R}{R_i} - 1 = \frac{R - R_i}{R_i}$. In (b), $\frac{P_{r_{i+1}}}{Q_{r_{i+1}}} + \dots + \frac{P_{r_j}}{Q_{r_j}}$ is divisible by $x^{\max(2n, m^2)+1}$ for each $j > i$, and hence so is $\frac{P_{r_{i+1}}}{Q_{r_{i+1}}} + \frac{P_{r_{i+2}}}{Q_{r_{i+2}}} + \dots = R - R_i$.

Thus $R - R_i$ is divisible by x^s , where $s = \max(2M, N^2, 2n, m^2) + 1$. Then $QR - QR_i$ is also divisible by x^s . Let us write $QR_i = a_0 + a_1x + a_2x^2 + \dots$. Since $QR = P$ is a polynomial of degree $M < s$, we obtain that $a_0 + a_1x + \dots + a_Mx^M = P$ and $a_{M+1} = \dots = a_{s-1} = 0$. On the other hand, by Lemma 9.1 QR_i satisfies a linear recurrence of the form $a_j = c_1a_{j-1} + \dots + c_na_{j-n}$ for each $j > Nm$. Since $s > \max(2M, 2n) \geq M + n$ and $s > \max(N^2, m^2) \geq Nm$, we obtain that $0 = a_s = a_{s+1} = \dots$. Thus $QR_i = P$. In other words, $R_i = P/Q = R$. Now i was an arbitrary integer such that $i \geq i_0$. Thus $R_{i_0} = R = R_{i_0+1}$. Hence $P_{i_0+1} = 0$, which contradicts the hypothesis. \square

10. PROOF OF THEOREM A

For a two-component PL link L it will be convenient to use the shifted Conway polynomial $\nabla_L(z) := z^{-1}\nabla_L(z)$. Thus $\nabla_L(x - x^{-1}) = \Omega_L(x, x)$. If we divide the shifted

Conway polynomial $\nabla_L(z)$ of $L = (K_1, K_2)$ by the product of the non-shifted Conway polynomials of the two components:

$$\bar{\nabla}_L(z) = \frac{\nabla_L(z)}{\nabla_{K_1}(z)\nabla_{K_2}(z)},$$

then we get an invariant of PL isotopy of L (see Corollary 7.6).⁴¹ Its coefficients are finite type invariants (see [67; Lemma 10.12(a)]). Hence by Theorem 1.2 it extends to an invariant of isotopy of topological links.

Theorem 10.1. *Let Λ be the splice of two-component PL links L and L' with $\text{lk}(L) = \text{lk}(L') = 1$. Then $\bar{\Omega}_\Lambda(x, y) = \bar{\Omega}_L(x, y)\bar{\Omega}_{L'}(x, y)$, where⁴²*

$$\bar{\Omega}_{(K, K')}(x, y) = \frac{\Omega_{(K, K')}(x, y)}{\nabla_K(x - x^{-1})\nabla_{K'}(y - y^{-1})}.$$

In particular, $\bar{\nabla}_\Lambda(z) = \bar{\nabla}_L(z)\bar{\nabla}_{L'}(z)$.

Proof. By Theorem 7.5 $\Omega_\Lambda(x, y) = \Omega_L(x, y)\Omega_{L'}(x, y)$. Let $\Lambda = (K_1, K_2)$, $L = (K_1, J)$ and $L' = (B, Q)$. The knot K_2 is itself the splice of J with (B, Q) . Then by Theorem 7.5 $\Omega_{K_2}(y) = \Omega_J(y)\Omega_{(B, Q)}(1, y)$. Also by Theorem 7.1 $\Omega_{(B, Q)}(1, y) = \nabla_Q(y - y^{-1})$. Hence $\nabla_{K_2}(z) = \nabla_J(z)\nabla_Q(z)$. Since $\nabla_B(z) = 1$, the assertion follows. \square

Let X be the Bing sling contained in the solid torus $T := S^1 \times D^2$. Let us assume that T is standardly embedded in $S^3 = S^1 \times D^2 \cup D^2 \times S^1$, and let B be a meridian of T and C the core circle of T .

The Bing sling X is defined as the limit (in the space of maps) of PL knots X_0, X_1, \dots , all contained in T , where $X_0 = C$ and each X_{i+1} is the image of a periodic PL knot $J_{r_{i+1}}$ contained in T under an untwisted homeomorphism h_i between T and a regular neighborhood T_i of X_i . Here the knots J_r are as in §8, and, as discussed in §1, the sequence r_1, r_2, \dots needs to be chosen appropriately in order for X to be homeomorphic to S^1 .

Step I. Let us first show that $Y := (B, X)$ is not isotopic to any PL link, provided that the r_i increase sufficiently rapidly.

⁴¹We do need to use both shifted and non-shifted Conway polynomials in the same fraction. If we use the shifted Conway polynomials in the denominator, then we do not get the PL isotopy invariance. If we use the non-shifted Conway polynomial in the numerator, then we do not get the formula of Theorem 10.1. Of course one can take the view that both the numerator and the denominator use the polynomial $z^{m-1}\nabla_L(z)$, where m is the number of components of L ; but this approach does not apply in the multi-variable setting, which will be used to prove Theorems A' and C.

⁴²Although $\bar{\nabla}_L(z)$ may be understood either as a rational function or a formal power series, $\bar{\Omega}_L(x, y)$ will be understood as a rational function only. One may, of course, identify $\bar{\Omega}_L(x, y)$ with a formal Laurent series $\Lambda(x, y)$, but this is not the only possibility. If $\Omega_L(x, y)$ happens to be a polynomial in $u := x - x^{-1}$ and $v := y - y^{-1}$, then another possibility is to identify $\bar{\Omega}_L$ with a formal power series $\Phi(u, v)$. In this case one cannot in general recover $\Lambda(x, y)$ from $\Phi(u, v)$ by substituting $x - x^{-1}$ for u and $y - y^{-1}$ for v . For instance, if $(1 - x + x^{-1})^{-1}$ is treated as a Laurent series, it equals $x(1 - x^2 + x)^{-1} = x(1 + (x^2 - x) + (x^2 - x)^2 + \dots)$, which cannot be obtained from $(1 - z)^{-1} = (1 + z + z^2 + \dots)$ by substituting $x - x^{-1}$ for z .

Let $Y_i = (B, X_i)$ and $L_i = (B, J_{r_i})$.⁴³ Each Y_{i+1} is clearly the splice of Y_i with L_{i+1} . Then by Theorem 10.1 we get $\bar{\nabla}_{Y_{i+1}}(z) = \bar{\nabla}_{Y_i}(z) \bar{\nabla}_{L_{i+1}}(z)$. Since $Y_0 = (B, C)$ is the Hopf link, $\bar{\nabla}_{Y_0} = 1$, and we get by induction that $\bar{\nabla}_{Y_n} = \bar{\nabla}_{L_1} \cdots \bar{\nabla}_{L_n}$.

Since the links Y_n converge to Y in the C^0 topology, their reduced shifted Conway polynomials $\bar{\nabla}_{Y_0}, \bar{\nabla}_{Y_1}, \dots$ converge to $\bar{\nabla}_Y$ in the ring of formal power series. Since each $\bar{\nabla}_{Y_n}$ is the finite product $\bar{\nabla}_{L_1} \cdots \bar{\nabla}_{L_n}$, their limit $\bar{\nabla}_Y$ is the infinite product $\bar{\nabla}_{L_1} \bar{\nabla}_{L_2} \cdots$ whose factors $\bar{\nabla}_{L_i}$ converge to 1.⁴⁴ By Corollary 8.3 $\bar{\nabla}_{L_i} \neq 1$, and hence by Theorem 9.2(a) this infinite product is not a rational power series, as long as the sequence r_i increases sufficiently rapidly. Hence Y is not isotopic to any PL link.

Step II. Next we prove Theorem A in the general case. Thus we are going to show that $Z := (K, X)$ is not isotopic to any PL link.

Since K lies in the complement to the Bing sling X , it is disjoint from one of the regular neighborhoods T_i of the knots X_i such that $X = \bigcap_{i=0}^{\infty} T_i$. Say $K \cap T_k = 0$. Then $Z_i := (K, X_i)$ is a PL link for each $i \geq k$. Moreover, each Z_{i+1} is the splice of Z_i and L_{i+1} for all $i \geq k$. Since $\text{lk } Z_i = \text{lk } Z = 1$, arguing as before we obtain that $\bar{\nabla}_{Z_i} = \bar{\nabla}_{Z_k} \bar{\nabla}_{L_{k+1}} \cdots \bar{\nabla}_{L_i}$, and consequently $\bar{\nabla}_Z = \bar{\nabla}_{Z_k} \bar{\nabla}_{L_{k+1}} \bar{\nabla}_{L_{k+2}} \cdots$. Like before, $\bar{\nabla}_{L_{k+1}} \bar{\nabla}_{L_{k+2}} \cdots$ is not a rational power series, as long as the sequence r_i increases sufficiently rapidly. Also $\bar{\nabla}_{Z_k}$ is a rational power series, and it is not identically zero due to $\text{lk } Z_k = \text{lk } Z = 1$. Hence the power series $\bar{\nabla}_Z$ cannot be rational and thus Z is not isotopic to any PL link.

Remark 10.2. Let us write down an explicit condition on the r_i that was used in the proof. We have $\bar{\nabla}_{L_i} = \frac{z^{-1} \nabla_{M_r}}{\nabla_{J_r}}$, where $r = r_i$. Here $\deg \nabla_{J_r} = 2r - 2$ and $\deg z^{-1} \nabla_{M_r} = 2r + 2$ (see the proof of Corollary 8.3). If n_i and m_i are as in the proof of Theorem 9.2(a), then $n_i = \sum_{j=1}^i (2j - 2) = i(i - 1)$ and $m_i = \sum_{j=1}^i (2j + 2) = i(i + 3)$, and hence $\max(2n_i, m_i^2) + 1 = i^2(i + 3)^2 + 1$. The polynomial P_i in the proof of Theorem 9.2(a) is $z^{-1} \nabla_{M_i} - \nabla_{J_i} = (-1)^{i+1} z^{i+1} L_{i+1}(z)$ (see Proposition 8.2(d)). The function f in the proof of Theorem 9.2(a) is defined by: $f(i)$ is such that P_j is divisible by $x^{\max(2n_i, m_i^2)+1}$ for all $j \geq f(i)$. Since L_{i+1} contains a constant term for odd i (and a linear term for even i , so we will not get much by omitting the odd i), the divisibility condition is saying that $j + 1 \geq i^2(i + 3)^2 + 1$. Thus $f(i) = i^2(i + 3)^2$. That is, the condition used in the proof that $\bar{\nabla}_{L_1} \bar{\nabla}_{L_2} \cdots$ is not rational is that each $r_{i+1} \geq r_i^2(r_i + 3)^2$. Since $\bar{\nabla}_{L_1} \cdots \bar{\nabla}_{L_k}$ is rational, to prove that $\bar{\nabla}_{L_{k+1}} \bar{\nabla}_{L_{k+2}} \cdots$ is not rational we may use the same condition.

⁴³Let us note that the unknotted component B comes first here, and therefore will correspond to the variable x ; whereas in §8 it came last in $M_r = (J_r, B)$ and therefore corresponded to the variable y . There are reasons for both choices, and it seems that reversing either of them would result in a greater inconvenience than the present inconvenience of switching from one choice to another.

⁴⁴There is also a more geometric way to see that $\bar{\nabla}_{L_i}$ converge to 1. The knots J_{r_1}, J_{r_2}, \dots , or rather their equivalent versions in thinner neighborhoods of C in T , have to converge to C in order for X_1, X_2, \dots to converge to a topological embedding of a circle. Since $Y_0 = (B, C)$ is the Hopf link, each $L_i = (B, J_{r_i})$ is k_i -quasi-isotopic to the Hopf link, where $k_i \rightarrow \infty$ as $i \rightarrow \infty$. Hence the coefficients of $\bar{\nabla}_{L_i}$ at degrees $\geq 2k_i$ are zero.

11. PROOF OF THEOREM B

Lemma 11.1. *Let X be a knot and let K and K' be PL knots each linking X with $\text{lk} = 1$. If there exists a PL solid torus T , disjoint from both K and K' and containing X so that $[X]$ is a generator of $H_1(T)$, then $\bar{\nabla}_{(K',X)} = R\bar{\nabla}_{(K,X)}$ for some rational power series R .*

Proof. Let X_0 be a core circle of T , and X_1, X_2, \dots be PL knots in the interior of T converging to X . Let h be a homomorphism between T and the standard unknotted solid torus $S^1 \times D^2 \subset S^3$. Let $M \subset S^3$ be a meridian $pt \times \partial D^2$ of the standard unknotted solid torus, $Q = h(X)$ and $Q_n = h(X_n)$. Then each link $Z_n := (K, X_n)$ is a splice of Z_0 and $L_n := (M, Q_n)$. Hence $\bar{\nabla}_{Z_n} = \bar{\nabla}_{Z_0} \bar{\nabla}_{L_n}$. Since $Z_n \rightarrow Z := (K, X)$ and $L_n \rightarrow L := (M, Q)$, we have $\bar{\nabla}_{Z_n} \rightarrow \bar{\nabla}_Z$ and $\bar{\nabla}_{L_n} \rightarrow \bar{\nabla}_L$. The latter clearly implies $\bar{\nabla}_{Z_0} \bar{\nabla}_{L_n} \rightarrow \bar{\nabla}_{Z_0} \bar{\nabla}_L$. Hence $\bar{\nabla}_Z = \bar{\nabla}_{Z_0} \bar{\nabla}_L$. Similarly $\bar{\nabla}_Y = \bar{\nabla}_{Y_0} \bar{\nabla}_L$, where $Y = (K', X)$ and $Y_0 = (K', X_0)$. Since $\text{lk}(Z_0) = \text{lk}(Z) = 1$ and $\text{lk}(Z_0)$ is the free term of $\bar{\nabla}_{Z_0}$, we have $\bar{\nabla}_{Z_0} \neq 0$. Then $\bar{\nabla}_Y = \bar{\nabla}_{Y_0} \bar{\nabla}_{Z_0}^{-1} \bar{\nabla}_Z$, where $\bar{\nabla}_{Y_0} \bar{\nabla}_{Z_0}^{-1}$ is a rational power series. \square

Lemma 11.2. *Let \mathcal{K} be a knot in S^3 which is the intersection of solid tori $T_1 \supset T_2 \supset \dots$. Then \mathcal{K} represents a generator of $H_1(T_i)$ for all but finitely many i .*

Proof. By the Alexander duality $H_1(S^3 \setminus \mathcal{K}) \simeq \mathbb{Z}$ and each $H_1(S^3 \setminus T_i) \simeq \mathbb{Z}$. Since $S^3 \setminus \mathcal{K} = \bigcup_i S^3 \setminus T_i$, we have $H_1(S^3 \setminus \mathcal{K}) \simeq \text{colim } H_1(S^3 \setminus T_i)$. It is easy to see that $\text{colim}(\mathbb{Z} \xrightarrow{f_1} \mathbb{Z} \xrightarrow{f_2} \dots) \not\simeq \mathbb{Z}$ if infinitely many of the f_i are not isomorphisms. Hence only finitely many of the f_i are not isomorphisms, and then without loss of generality all of them are isomorphisms. Then the inclusion induced maps $H_1(T_i) \rightarrow H_1(T_{i+1})$ are also isomorphisms. Then \mathcal{K} represents a generator of $H_1(T_i)$ for each i . \square

Theorem B is a consequence of the following

Theorem 11.3. *Suppose that the Bing sling X_0 is isotopic to a knot X_1 through knots X_t such that when t belongs to a dense subset S of $[0, 1]$, X_t is the intersection of a nested sequence of PL solid tori. Then X_1 is a wild knot.*

Proof. Let $0 = t_0 < \dots < t_n = 1$ and K_0, \dots, K_{n-1} be given by Lemma 3.1. By the proof of Theorem A, $\bar{\nabla}_{(K_0, X_0)}$ is not rational. Then it follows from Lemmas 11.2 and 11.1 that $\bar{\nabla}_{(K_{n-1}, X_1)}$ is also not rational. Hence X_1 is wild. \square

12. TWO-VARIABLE CONWAY POLYNOMIAL

For a 2-component PL link $L = (K_1, K_2)$ of linking number λ it is not hard to show that $\Omega_L(x, y)$ can be uniquely rewritten in the form

$$\nabla_L(u, v, w) := \sum_{i \equiv j \not\equiv \lambda \pmod{2}} a_{ij} u^i v^j + \sum_{i \equiv j \equiv \lambda \pmod{2}} a_{ij} u^i v^j w,$$

where $u = x - x^{-1}$, $v = y - y^{-1}$ and $w = xy^{-1} + x^{-1}y$ and each $a_{ij} \in \mathbb{Z}$ (see [68]).

Remark 12.1. Let $\bar{w} = xy + x^{-1}y^{-1}$. Then $w - \bar{w} = -uv$ and $w\bar{w} = u^2 + v^2 + 4$. So by Vieta's theorem w and $-\bar{w}$ are the roots of the equation $w^2 + uvw - (u^2 + v^2 + 4) = 0$. Thus $w = -\frac{uv}{2} + \sqrt{\frac{u^2v^2}{4} + u^2 + v^2 + 4}$, in the sense that $(w + \frac{uv}{2})^2 = \frac{u^2v^2}{4} + u^2 + v^2 + 4$ in $\mathbb{Q}[x^{\pm 1}, y^{\pm 1}]$. In fact, $w + \frac{uv}{2} = \frac{1}{2}(x + x^{-1})(y + y^{-1})$. Thus the \mathbb{Q} -subalgebra of $\mathbb{Q}[x^{\pm 1}, y^{\pm 1}]$ generated by u, v and w can be described as $\mathbb{Q}[u, v] \left[\sqrt{u^2v^2 + 4(u^2 + v^2 + 4)} \right]$, where $\sqrt{u^2v^2 + 4(u^2 + v^2 + 4)}$ can be identified with the element $(x + x^{-1})(y + y^{-1})$ of $\mathbb{Q}[x^{\pm 1}, y^{\pm 1}]$. (But there is no such description over \mathbb{Z} .) Of course, $\mathbb{Q}[x^{\pm 1}, y^{\pm 1}]$ itself is generated by its elements $u, v, x + x^{-1}$ and $y + y^{-1}$.

Let us note that upon equating x and y we get $u = v$ and $w = 2$, with $\nabla_L(z) = \nabla_L(z, z, 2)$. In general, we will call polynomials (resp. formal power series) in u, v, w that are linear in w and include only those monomials where the degrees of all three variables have the same parity *w-polynomials* (resp. *w-series*). Thus when $\text{lk}(L)$ is odd, ∇_L is a *w-polynomial*. When multiplying two or more *w-polynomials* we need to remember that there is an algebraic relation between u, v and w , namely, $w^2 = u^2 + v^2 + 4 - uvw$. Therefore a product of finitely many *w-polynomials* will again be a *w-polynomial* (if we take the product in $\mathbb{Z}[u, v, w]/(w^2 + uvw - u^2 - v^2 - 4)$), and a product of finitely many *w-series* will again be a *w-series* (if we take the product in $\mathbb{Z}[[u, v, w]]/(w^2 + uvw - u^2 - v^2 - 4)$). This also works for infinitely many *w-series* as long as they converge to 1 in the usual topology on $\mathbb{Z}[[u, v, w]]$. It is easy to see that the following are equivalent for a *w-series* R :

- (1) R is rational (as a power series in $\mathbb{Z}[[u, v, w]]$);⁴⁵
- (2) R is of the form $P(u, v, w)/Q(u, v)$, where P and Q are *w-polynomials* with Q constant in w and $Q(0, 0) = 1$;
- (3) R is of the form $R'(u, v) + wR''(u, v)$, where R' and R'' are rational power series in u and v .

13. PROOF OF THEOREM A'

For a 2-component PL link $L = (K_1, K_2)$ let

$$\bar{\nabla}_L(u, v, w) = \frac{\nabla_L(u, v, w)}{\nabla_{K_1}(u) \nabla_{K_2}(v)}.$$

Then $\bar{\nabla}_L(u, v, w)$ is a rational *w-series* if $\text{lk}(L)$ is odd. By [68; Theorem B(a)] each coefficient of $\bar{\nabla}_L$ assumes the same value on all sufficiently close PL approximations of a topological link; consequently $\bar{\nabla}_L$ extends to an invariant of isotopy of topological links (which is, of course, a *w-series* that no longer needs to be rational).

Remark 13.1. The rational function $\bar{\Omega}_L$ does not seem to have a reasonable extension to an invariant of topological links (with respect to some completion of the field of rational functions). On the other hand, if L and L' are topologically isotopic PL links,

⁴⁵A power series in $\mathbb{Z}[[x_1, \dots, x_n]]$ is called *rational* if it is of the form P/Q , where $P, Q \in \mathbb{Z}[x_1, \dots, x_n]$ and $Q \in \mathbb{Z}[[x_1, \dots, x_n]]^\times$, i.e. $Q(0, \dots, 0) = \pm 1$; compare [42].

then $\bar{\nabla}_L = \bar{\nabla}_{L'}$, and if we understand $\bar{\nabla}_L$ and $\bar{\nabla}_{L'}$ as fractions whose denominator does not involve w (rather than power series) and substitute $x - x^{-1}$, $y - y^{-1}$ and $xy^{-1} + x^{-1}y$ for u , v and w separately in the numerator and in the denominator, then we obtain that $\bar{\Omega}_L = \bar{\Omega}_{L'}$.

The definition of ∇_L along with the fact that the Conway polynomial of a knot has no odd exponents imply the following representation for $\bar{\nabla}_L$:

$$\bar{\nabla}_L(u, v, w) = \begin{cases} \sum_{i=0}^{\infty} u^{2i+1} v P_{2i+1}(v^2) + w \sum_{i=0}^{\infty} u^{2i} P_{2i}(v^2), & \text{if } \text{lk}(L) \text{ is even;} \\ \sum_{i=0}^{\infty} u^{2i} P_{2i}(v^2) + w \sum_{i=0}^{\infty} u^{2i+1} v P_{2i+1}(v^2), & \text{if } \text{lk}(L) \text{ is odd,} \end{cases}$$

where each P_k is a rational power series. It follows from Theorem 7.1 that $\bar{\nabla}_L(0, v, y + y^{-1}) = \bar{\Omega}_L(1, y) = \frac{y^{\text{lk}(L)} - y^{-\text{lk}(L)}}{y - y^{-1}}$. Therefore P_0 is a function of $\text{lk}(L)$.⁴⁶

Let us denote P_1 by C_L . When $\text{lk}(L) = 0$, $C_L(z) = \sum_{k=0}^{\infty} (-1)^k \beta_{k+1} z^{2k}$, where β_k are Cochran's derived invariants (see [68; Theorem C]). By the above, C_L extends to an invariant of isotopy of topological links and by [68; Theorem B(b)] the extension invariant under sufficiently close C^0 -perturbation of the first component.

Theorem 13.2. *Let Λ be the splice of two-component PL links L and L' with $\text{lk } L = \text{lk } L' = 1$. Then $C_{\Lambda}(z) = C_L(z) + C_{L'}(z)$.*

Proof. Let $L = (K_1, K_2)$. By Theorem 7.1 $\Omega_L(1, y) = \nabla_{K_2}(y - y^{-1})$ and $\nabla_{K_1}(0) = 1$. Hence $\bar{\Omega}_L(1, y) = 1$. Therefore $P_0 = 1$. Thus $\bar{\nabla}_L(u, v, w) \equiv 1 + uvw C_L(v) \pmod{(u^2)}$. And similarly for L' and Λ . By Theorem 10.1 $\bar{\nabla}_{\Lambda}(u, v, w) = \bar{\nabla}_L(u, v, w) \bar{\nabla}_{L'}(u, v, w)$, and the assertion follows. \square

Proof of Theorem A'. Suppose that \mathcal{K} is a topological knot in the complement of the Bing sling X such that $\text{lk}(\mathcal{K}, X) = 1$. Since C_L is invariant under sufficiently close C^0 -perturbation of the first component, $C_{(\mathcal{K}, X)} = C_{(K, X)}$ for any PL knot K sufficiently close to \mathcal{K} . Let $Z = (K, X)$. Since K lies in $S^3 \setminus X$, it is disjoint from one of the regular neighborhoods T_i of the knots X_i such that $X = \bigcap_{i=0}^{\infty} T_i$. Say $K \cap T_k = 0$. Then $Z_i := (K, J_{r_i})$ is a PL link for each $i \geq k$. Then similarly to the proof of the PL case of Theorem A, using Theorem 13.2 instead of Theorem 10.1, we get that $C_Z = C_{Z_k} + C_{L_{k+1}} + C_{L_{k+2}} + \dots$. By Proposition 8.2(b,c) $C_{L_i} = C_{(B, J_{r_i})} = z^r F_r(z) / \nabla_{J_r}(z) \neq 0$, where $r = r_i$. Hence by Theorem 9.2(b) we get that $C_{L_{k+1}} + C_{L_{k+2}} + \dots$ is not a rational power series, as long as the sequence r_i increases sufficiently rapidly. Also C_{Z_k} is a rational power series. Hence the power series $C_{(\mathcal{K}, X)} = C_Z$ cannot be rational. Since C_L is invariant under isotopy of L , we conclude that (K, X) is not isotopic to any PL link. \square

Remark 13.3. Let us write down an explicit condition on the r_i that was used in the proof. We have $C_{L_i} = z^r F_r(z) / \nabla_{J_r}(z)$, where $r = r_i$. Here $\deg \nabla_{J_r} = 2r - 2$ (see the proof of Corollary 8.3) and $\deg z^r F_r = 2r - 1$. If n_i and m_i are as in the proof of Theorem

⁴⁶In fact, it is easy to see from Example 7.3 that when $\text{lk}(L)$ is even, $P_0(v) = v^{-1} F_{\text{lk}(L)}(v)$; and when $\text{lk}(L)$ is odd, $P_0(v) = v^{-1} L_{\text{lk}(L)}(v)$.

9.2(b), then $n_i = \sum_{j=1}^i (2j - 2) = i(i - 1)$ and $m_i = n_i + 1 = i^2 - i + 1$, and hence $\max(2n_i, m_i^2) + 1 = (i^2 - i + 1)^2 + 1$. The polynomial P_i in the proof of Theorem 9.2(b) is $z^i F_i(z)$. The function f in the proof of Theorem 9.2(b) is defined by: $f(i)$ is such that P_j is divisible by $x^{\max(2n_i, m_i^2)+1}$ for all $j \geq f(i)$. Since F_i contains a constant term for odd i (and a linear term for even i , so we will not get much by omitting the odd i), the divisibility condition is saying that $j \geq (i^2 - i + 1)^2 + 1$. Thus $f(i) = (i^2 - i + 1)^2 + 1$. That is, the condition used in the proof that $C_{L_1} + C_{L_2} + \dots$ is not rational is that each $r_{i+1} \geq (r_i^2 - r_i + 1)^2 + 1$. Since $C_{L_1} + \dots + C_{L_k}$ is rational, to prove that $C_{L_{k+1}} + C_{L_{k+2}} + \dots$ is not rational we may use the same condition.

Lemma 13.4. *Let (K, X) and (K', X) be links with $\text{lk}(K, X) = \text{lk}(K', X) = 1$. Suppose that there exists a PL solid torus $T \subset S^3$, disjoint from both K and K' and containing X so that $H_1(T)$ is generated by $[X]$. Then $C_{(K, X)} - C_{(K', X)}$ is a rational power series.*

Proof. Similarly to the proof of Lemma 11.1. □

Let us call links (K, X) and (K', X) as in Lemma 13.4 *anti-F-isotopic*. Then we get

Theorem 13.5. *Let X be the Bing sling and K be a knot linking X with $\text{lk} = 1$. Suppose that (K, X) is related to a link (K', X') by the equivalence relation generated by isotopy and anti-F-isotopy. Then X' is a wild knot.*

14. A DISAPPOINTING EXAMPLE

The proofs of Theorem B and Theorem 13.5 raise the following two questions:

Problem 14.1. *Let X be a knot and let K and K' be PL knots each linking X with $\text{lk} = 1$. Is it true that*

- (a) $C_{(K', X)}(z)$ is a rational power series if and only if so is $C_{(K, X)}(z)$?
- (b) $\bar{\nabla}_{(K', X)}(z)$ is a rational power series if and only if so is $\bar{\nabla}_{(K, X)}(z)$?



A positive solution to either (a) or (b) would imply that the Bing sling is not isotopic to the unknot.

We will show, however, that (a) has a negative solution, and so does the two-variable version of (b). This does not yet provide a solution of (b), but there does not seem to be much hope that it can be positive.

Theorem 14.2. *There exist 2-component links $L' = (K', X)$ and $L^* = (K^*, X)$ such that K' and K^* are PL knots, $\text{lk}(L') = \text{lk}(L^*) = 1$ and*

- (a) $C_{L^*}(z)$ is rational but $C_{L'}(z)$ is not rational; also,
- (b) $\bar{\nabla}_{L^*}(u, v, w)$ is rational but $\bar{\nabla}_{L'}(u, v, w)$ is not rational.

Proof. Let $L = (K, X)$ be one of the links constructed in [65], with $\text{lk}(L) = 0$ and C_L not being rational (see Figure 11). Let K' and K'' be PL knots in the complement of X such that the links $L = (K, X)$, $L' := (K', X)$ and $L'' := (K'', X)$ are related as in the second Conway identity; see Figure 12.

FIGURE 11. The link $L = W_{n_1, n_2, n_3, \dots}$ where $(n_1, n_2, n_3) = (1, 0, -1)$.FIGURE 12. The links L' (top) and L'' (bottom).

Since K is unknotted in S^3 (see [65]), so are K' and K'' . If Q is a sufficiently close PL approximation of X , then $\Lambda := (K, Q)$, $\Lambda' := (K', Q)$ and $\Lambda'' := (K'', Q)$ are also related as in the second Conway identity, and so we have $\Omega_{\Lambda'}(x, y) + \Omega_{\Lambda''}(x, y) = (xy + x^{-1}y^{-1})\Omega_{\Lambda}(x, y)$. Writing $\Lambda^* = (K^*, Q)$, where $K^* = -K''$, that is, K'' with orientation reversed, by Proposition 7.8 we get $\Omega_{\Lambda'}(x, y) - \Omega_{\Lambda^*}(x^{-1}, y) = (xy + x^{-1}y^{-1})\Omega_{\Lambda}(x, y)$. Upon substituting x with x^{-1} we get $\Omega_{\Lambda'}(x^{-1}, y) - \Omega_{\Lambda^*}(x, y) = (x^{-1}y + xy^{-1})\Omega_{\Lambda}(x^{-1}, y)$ or equivalently $\nabla_{\Lambda'}(-u, v, w) - \nabla_{\Lambda^*}(u, v, w) = w\nabla_{\Lambda}(-u, v, w)$. Since K , K' and K^* are unknotted, we get $\bar{\nabla}_{\Lambda'}(-u, v, w) - \bar{\nabla}_{\Lambda^*}(u, v, w) = w\bar{\nabla}_{\Lambda}(-u, v, w)$. Since Q can be

any sufficiently close approximation of X , we obtain $\bar{V}_{L'}(-u, v, w) - \bar{V}_{L^*}(u, v, w) = w\bar{V}_L(-u, v, w)$, where $L^* = (K^*, X)$. Now since X is locally flat except at one point, it is easy to see that L^* is isotopic to the Hopf link. Hence $\bar{V}_{L^*}(u, v, w) = 1$, and in particular $C_{L^*} = 0$. Therefore $\bar{V}_{L'}(u, v, w) = w\bar{V}_L(u, v, w)$. Hence also $C_{L'}(z) = C_L(z)$, which is not rational. Then in particular $\bar{V}_{L'}(u, v, w)$ itself cannot be rational. \square

Remark 14.3. One might wonder whether the fact that $\bar{V}_{(K,X)}(z)$ and $\bar{V}_{(K,X)}(u, v, w)$ are represented as infinite products of rational power series for any PL knot K that links the Bing sling X with linking number 1 is remarkable as a property of the Bing sling. It is not. An arbitrary power series of the form $1 + a_1z + a_2z^2 + \dots$ factors in $\mathbb{Z}[[z]]$ into a product of polynomials of the form $(1 + b_1z)(1 + b_2z^2)(1 + b_3z^3) \dots$, where each b_n is uniquely determined, assuming that b_1, \dots, b_{n-1} have already been found, by equating the degree n term of $\prod_{i=1}^n (1 + b_i z^i)$ with that of the given power series. Similarly, every w -series $\sum_{i,j=0, i \equiv j \pmod{2}}^{\infty} a_{ij} u^i v^j w^{i \bmod 2}$ with $a_{00} = 1$ factors in $\mathbb{Z}[[u, v, w]]/(w^2 + uvw - u^2 - v^2 - 4)$ into a product of polynomials of the form $\prod_{i,j=0, i \equiv j \pmod{2}}^{\infty} (1 + b_{ij} u^i v^j w^{i \bmod 2})$ with $b_{00} = 0$.

15. PROOFS OF THEOREMS C AND D'

Theorem D' is a reformulation of the following

Theorem 15.1. *There exists a 2-component link \mathcal{L} in S^3 with $\text{lk}(\mathcal{L}) = 0$ and second component being PL, which is not equivalent to any PL link under the equivalence relation generated by (i) isotopy and (ii) replacing a link (Q, \mathcal{K}) by a link (Q', \mathcal{K}) , provided that Q and Q' are PL knots homotopic in $S^3 \setminus \mathcal{K}$ or more generally represent the same conjugacy class in the quotient of $\pi_1(S^3 \setminus \mathcal{K})$ by the second commutator subgroup.*

In fact, the proof also works for the following more general version (ii') of the transformation (ii): replace a link (Q, \mathcal{K}) by a link (Q', \mathcal{K}) , provided that Q and Q' represent the same conjugacy class in the quotient of $G := \pi_1(S^3 \setminus \mathcal{K})$ by the preimage in G' of the torsion submodule of the knot module G'/G'' .⁴⁷

Proof. The proof of Theorem 1.1 in [65] shows that for 2-component links L with $\text{lk}(L) = 0$ the Cochran power series C_L is invariant under isotopy of L , is rational for PL links, and is not rational for a certain link \mathcal{L} whose second component is PL. To complete the proof it suffices to show that for any transformation of the type (ii), $C_{(Q,\mathcal{K})} - C_{(Q',\mathcal{K})}$ is always rational. In fact we will prove that this difference is a polynomial. (For transformations of type (ii') this difference will only be rational, not polynomial.)

Since Q and Q' represent the same conjugacy class in the quotient of $\pi_1(S^3 \setminus \mathcal{K})$ by the second commutator subgroup, they admit lifts \tilde{Q}, \tilde{Q}' to the infinite cyclic cover \mathcal{X} of $S^3 \setminus \mathcal{K}$ such that the 1-cycle $\tilde{Q} - \tilde{Q}'$ bounds a 2-chain ζ in \mathcal{X} . The image of ζ in $S^3 \setminus \mathcal{K}$ as well as Q and Q' are disjoint from some polyhedral neighborhood N of \mathcal{K} . Hence ζ as well as \tilde{Q} and \tilde{Q}' lie in the preimage Y of $S^3 \setminus N$ in \mathcal{X} . This Y can also be described as

⁴⁷Equivalently, $Q' = Q \#_b R$ for some knot R in $S^3 \setminus \mathcal{K}$ such that the Kojima–Yamasaki λ -polynomial [53] $\lambda(R, \mathcal{K}) \neq 0$ and for some band $b \subset S^3 \setminus \mathcal{K}$.

the cover of $S^3 \setminus N$ corresponding to (the kernel of) the homomorphism $\pi_1(S^3 \setminus N) \rightarrow \mathbb{Z}$ given by $[Z] \mapsto \text{lk}(Z, \mathcal{K})$. (In other words, this homomorphism is determined by the class in $H^1(S^3 \setminus N)$ which is Alexander dual to $[\mathcal{K}] \in H_1(N)$.) If K is a sufficiently close PL approximation of \mathcal{K} , then it is homotopic to \mathcal{K} within N , and then the preimage of $S^3 \setminus N$ in the infinite cyclic cover X of $S^3 \setminus K$ is the same cover Y of $S^3 \setminus N$. Thus ζ as well as \tilde{Q} and \tilde{Q}' can be regarded as lying in X .

For PL links we have $C_L(z) = \eta_L(t)$, where η_L is Kojima's η -function, $t = x^2$ and $z = x - x^{-1}$. By definition $\eta_{(Q,K)}(t) = \langle \tilde{Q}, \tilde{Q}_+ \rangle$, where $\langle \cdot, \cdot \rangle$ is the Cochran pairing in the infinite cyclic cover X of $S^3 \setminus K$ and \tilde{Q} and \tilde{Q}_+ are nearby lifts of Q and of its parallel pushoff. By [68; Lemma 4.4(a)] $\langle \tilde{Q}, \tilde{Q}_+ \rangle - \langle \tilde{Q}', \tilde{Q}'_+ \rangle = \langle \partial\zeta, \tilde{Q}_+ \rangle + \langle \tilde{Q}', \partial\zeta_+ \rangle$, where $\partial\zeta = \tilde{Q} - \tilde{Q}'$ and $\partial\zeta_+ = \tilde{Q}_+ - \tilde{Q}'_+$. By [68; Lemma 4.4(c)] $\langle \tilde{Q}', \partial\zeta_+ \rangle(t) = \langle \partial\zeta_+, \tilde{Q}' \rangle(t^{-1})$. Thus

$$\eta_{(Q,K)}(t) - \eta_{(Q',K)}(t) = \langle \partial\zeta, \tilde{Q}_+ \rangle(t) + \langle \partial\zeta_+, \tilde{Q}' \rangle(t^{-1}) = \sum_{n=-\infty}^{\infty} (\zeta \cdot t^n \tilde{Q}_+ + \zeta_+ \cdot t^{-n} \tilde{Q}') t^n.$$

The right hand side is a Laurent polynomial (so, only finitely many summands are nonzero) which depends only on the mutual position of ζ and Q_+ and of ζ_+ and Q' in Y . Thus it does not depend on the choice of the PL approximation K , as long as it is homotopic to \mathcal{K} within N . Since $\eta_L(t^{-1}) = \eta_L(t)$, this Laurent polynomial is invariant under the involution $t \mapsto t^{-1}$. So it must be of the form $\sum_{n=0}^{\infty} a_n(t^n + t^{-n})$. It is not hard to check that $t^n + t^{-n} = L_{2n}(z)$ for $n \geq 0$, where L_n is the Lucas polynomial, defined by $L_0(z) = 2$, $L_1(z) = z$ and $L_{n+1}(z) = zL_n(z) + L_{n-1}(z)$. (Thus L_{2n} contains only even powers of z .) Since each coefficient of $C_L(z)$ is invariant under sufficiently close approximation [65; Theorem 2.3], $C_{(Q,K)}(z) - C_{(Q',K)}(z) = C_{(Q,K)}(z) - C_{(Q',K)}(z) = \sum_{n=0}^{\infty} a_n L_{2n}(z)$, which is a polynomial. \square

Theorem C follows by the proof of Theorem A' if we additionally use the following

Lemma 15.2. *Let \mathcal{K} be a knot in S^3 and let Q_1, Q_2 be PL knots in $S^3 \setminus \mathcal{K}$ such that $\text{lk}(Q_i, \mathcal{K}) = 1$. If Q_1 and Q_2 are homotopic in $S^3 \setminus \mathcal{K}$, then $C_{(Q_1, \mathcal{K})} - C_{(Q_2, \mathcal{K})}$ is a polynomial.*

Proof. Let N be a polyhedral neighborhood of \mathcal{K} such that the given homotopy takes place in $S^3 \setminus N$. Let K be a PL knot homotopic to \mathcal{K} within N . It suffices to consider the case where Q_1 is obtained from Q_2 by a single crossing change. Then for the 2-variable Conway polynomial of [68] we have

$$\nabla_{(Q_1, K)}(u, v) - \nabla_{(Q_2, K)}(u, v) = u \nabla_{(L, K)}(u, v),$$

where $L = (L_1, L_2)$ is the 2-component link obtained by smoothing the crossing. If we divide both sides of the equation by $\nabla_K(v)$, then differentiate both sides with respect to u and finally set $u = 0$, then we get

$$C_{(Q_1, K)}(v^2) - C_{(Q_2, K)}(v^2) = \frac{\nabla_{(L, K)}(0, v)}{\nabla_K(v)}.$$

Let us note that $\text{lk}(L_1, K) + \text{lk}(L_2, K) = \text{lk}(Q_i, K) = 1$. So precisely one of the integers $\text{lk}(L_1, K)$, $\text{lk}(L_2, K)$ is even, and if we denote it by n , then the other one equals $1 - n$. Then by Theorem 7.1 and Example 7.3

$$\Omega_{(L,K)}(1, y) = \frac{(y^n - y^{-n})(y^{1-n} - y^{n-1})}{y - y^{-1}} \nabla_K(y - y^{-1}) = (y + y^{-1})F_n(v) \frac{L_{1-n}(v)}{v} \nabla_K(v),$$

where $v = y - y^{-1}$. Then the part of $\Omega_{(L,K)}(x, y)$ which involves no powers of $u = x - x^{-1}$ must be $(x^{-1}y + xy^{-1})F_n(v) \frac{L_{1-n}(v)}{v} \nabla_K(v)$. Therefore $\nabla_{(L,K)}(0, v) = F_n(v) \frac{L_{1-n}(v)}{v} \nabla_K(v)$. Hence

$$C_{(Q_1,K)}(v^2) - C_{(Q_2,K)}(v^2) = F_n(v) \frac{L_{1-n}(v)}{v}.$$

Here the left hand side converges to $C_{(Q_1,\mathcal{K})} - C_{(Q_2,\mathcal{K})}$ as $K \rightarrow \mathcal{K}$ in C^0 topology, whereas the right side does not depend on K . Thus $C_{(Q_1,\mathcal{K})}(v^2) - C_{(Q_2,\mathcal{K})}(v^2) = F_n(v) \frac{L_{1-n}(v)}{v}$, which is a polynomial in v^2 . (Note that since n is even, $F_n(v)$ involves only even powers of v and L_{1-n} only odd ones.) \square

16. PROOF OF THEOREM D

Proposition 16.1. *Let \mathcal{K} be a knot in S^3 and Q a PL knot in $S^3 \setminus \mathcal{K}$ such that $\text{lk}(Q, \mathcal{K}) = 1$. Let S be a Seifert surface bounded by Q in S^3 and let F be an embedded PL surface in $S^3 \setminus \mathcal{K}$, disjoint from S and such that every knot in F is null-homologous in $S^3 \setminus \mathcal{K}$. Let (A_1, \dots, A_{2g}) be a symplectic basis for F , and let \tilde{A}_i be some lifts of A_i to the infinite cyclic cover \mathcal{X} of $S^3 \setminus \mathcal{K}$. Finally let $Q' = Q \#_b \partial F$, where b is a band in $S^3 \setminus \mathcal{K}$, connecting Q to ∂F and otherwise disjoint from S and F . If each $[\tilde{A}_{2i}]$ is annihilated by some Laurent polynomial in $H_1(\mathcal{X})$, then $C_{(Q,\mathcal{K})}(z) - C_{(Q',\mathcal{K})}(z)$ is a rational power series.*

Proof. By the hypothesis there exist $p_1, \dots, p_g \in \mathbb{Z}[t^{\pm 1}]$ such that each $p_i \tilde{A}_{2i}$ bounds a 2-chain ζ_i in \mathcal{X} . The images of the chains ζ_i in $S^3 \setminus \mathcal{K}$ as well as the surface F and the knot Q are disjoint from some polyhedral neighborhood N of \mathcal{K} . Hence the ζ_i and the \tilde{A}_{2i-1} lie in the preimage Y of $S^3 \setminus N$ in \mathcal{X} . This Y can also be described as the cover of $S^3 \setminus N$ corresponding to (the kernel of) the homomorphism $\pi_1(S^3 \setminus N) \rightarrow \mathbb{Z}$ given by $[Z] \mapsto \text{lk}(Z, \mathcal{K})$. If K is a sufficiently close PL approximation of \mathcal{K} , then it is homotopic to \mathcal{K} within N , and then the preimage of $S^3 \setminus N$ in the infinite cyclic cover X of $S^3 \setminus K$ is the same cover Y of $S^3 \setminus N$. Thus the ζ_i and the A_{2i-1} can be regarded as lying in X .

Let Σ be a Seifert surface bounded by K in S^3 . Since K is homotopic to \mathcal{K} within N , which is disjoint from F , $\text{lk}(R, K) = \text{lk}(R, \mathcal{K}) = 0$ for every knot R in F . Then by the proof of (3) \Rightarrow (1) for $g = 2$ in Theorem 17.10, we may assume that Σ is disjoint from F . If Σ intersects the band b , we can eliminate any such intersections by pushing them into S . Now by the proof of Lemma 3.5 we can eliminate all intersections between Σ and S except for one clasp arc J with one endpoint in K and another in Q . (This elimination consists of modifications of Σ and S which take place in a small neighborhood of $\Sigma \cup S$ and therefore create no intersections of either Σ or S with F . Also the innermost arc argument which is used to eliminate the intersections between Σ and Q can be carried out

in the arc $Q \setminus b$ rather than in the entire knot Q , and therefore will create no intersections of Σ with b .) Thus $S' := S \cup b \cup F$ is an embedded surface, and its intersection with Σ consists only of the arc $J \subset S$.

Now [68; Theorem D] applies to each of the pairs (Σ, S) and (Σ, S') , which yields

$$C_{(Q',K)}(s) - C_{(Q,K)}(s) = (-1)^g \sum_{n=1}^{\infty} (-1)^n \sum_{i=1}^g \beta_{\Sigma}^{n-1,n}(A_{2i}, A_{2i-1}) s^{n-1}.$$

The left hand side is a rational power series since both (Q, K) and (Q', K) are PL links. On the other hand, by [68; Theorem 6.6(a)] each $\beta_{\Sigma}^{n-1,n}(A_{2i}, A_{2i-1})$ depends only on $\langle A_{2i}, A_{2i-1} \rangle_{\Sigma}$, which in turn depends only on the mutual position of the ζ_i and the A_{2i-1} in Y . Thus it does not depend on the choice of the PL approximation K , as long as it is homotopic to \mathcal{K} within N . Therefore $C_{(Q',K)}(s) - C_{(Q,K)}(s)$ is rational. \square

Theorem 16.2. *Let \mathcal{K} be a knot in S^3 and let Q_1, Q_2 be PL knots in $S^3 \setminus \mathcal{K}$ such that $\text{lk}(Q_i, \mathcal{K}) = 1$. Let $G = \pi_1(S^3 \setminus \mathcal{K})$, let T_0 be the torsion submodule of the knot module G'/G'' , and let T be the preimage of T_0 in G' . If Q_1 and Q_2 represent the same conjugacy class of $G/[G', T]$, then $C_{(Q_1, \mathcal{K})} - C_{(Q_2, \mathcal{K})}$ is a rational power series.*

Theorem D follows from Theorem 16.2 and the proof of Theorem A'.

Proof. We may assume that Q_1 and Q_2 pass through the basepoint of $S^3 \setminus \mathcal{K}$ and hence determine some classes $a, b \in G$. Then by the hypothesis there exists a $g \in G$ such that $a = b^g$ modulo $[G', T]$. Hence $a^{-1}b^g = [c_1, t_1] \cdots [c_n, t_n]$ for some $c_i \in G'$ and some $t_i \in T$. Let O be a small 3-ball in $S^3 \setminus \mathcal{K}$ containing the basepoint of $S^3 \setminus \mathcal{K}$. We may represent c_1, \dots, c_n and t_1, \dots, t_n by PL knots A_1, \dots, A_n and B_1, \dots, B_n in $S^3 \setminus (\mathcal{K} \cup Q_1)$ which are all pairwise disjoint except that each A_i meets each B_i in a single point $p_i \in O$. We may assume that $\text{lk}(A_i, Q_1) = \text{lk}(B_i, Q_1) = 0$ for each i . Each of the wedges $A_i \vee B_i$ is a deformation retract of a punctured torus $\tau_i := \alpha_i \cup \beta_i$, where α_i and β_i are thin ribbons which deformation retract onto A_i and B_i and intersect in a 2-disk (a ‘‘square’’) which deformation retracts onto p_i . Thus each $[\partial\tau_i] = [c_i, t_i] \in G$. We may assume that the τ_i are pairwise disjoint and also are disjoint from \mathcal{K} and from Q_1 . For $i = 1, \dots, n-1$ let b_i be a band in O connecting $\partial\tau_i$ to $\partial\tau_{i+1}$ and let b_0 be a band in O connecting Q_1 to $\partial\tau_1$ so that the n bands are pairwise disjoint from each other and also from the τ_i and from Q_1 apart at their attaching arcs. These attaching arcs can be chosen so that the class of $Q'_1 := Q_1 \#_{b_0} \partial\tau_1 \#_{b_1} \cdots \#_{b_{n-1}} \partial\tau_n$ in G equals $a[c_1, t_1] \cdots [c_n, t_n]$. Since $\text{lk}(A_i, Q_1) = \text{lk}(B_i, Q_1) = 0$ for each i , it is easy to see that Q_1 bounds a Seifert surface S in S^3 disjoint from each τ_i . Also we can make S disjoint from the bands b_i by attaching to S punctured parallel copies of the surfaces ∂N_i , where N_n is a small regular neighborhood of τ_n and each N_i is a small regular neighborhood of $\tau_i \cup b_i \cup N_{i+1}$. In particular, S is disjoint from the surface $F := \tau_1 \cup b_1 \cup \tau_2 \cup \cdots \cup b_{n-1} \cup \tau_n$ and we have $Q'_1 = Q_1 \#_{b_0} \partial F$. Then by Proposition 16.1 $C_{(Q'_1, \mathcal{K})} - C_{(Q_1, \mathcal{K})}$ is a rational power series. On the other hand, since $b^g = a[c_1, t_1] \cdots [c_n, t_n]$, there is a free homotopy between Q'_1 and Q_2 in $S^3 \setminus \mathcal{K}$. Then by Lemma 15.2 $C_{(Q_2, \mathcal{K})} - C_{(Q'_1, \mathcal{K})}$ is a polynomial. \square

Corollary 16.3. *The Bing sling is not isotopic to any PL knot though knots whose knot module is a torsion module.*

Proof. Suppose that such an isotopy X_t exists. Let $0 = t_0 < \dots < t_n = 1$ and K_0, \dots, K_{n-1} be given by Lemma 3.1. Let $G_i = \pi_1(S^3 \setminus X_{t_i})$. By Theorem 3.2 K_{i-1} and K_i represent the same conjugacy class in $G_i/[G'_i, G'_i]$. Let $T_i \subset G'_i$ be the preimage of the torsion submodule of the knot module G'_i/G''_i . By the hypothesis $T_i = G'_i$. Hence K_{i-1} and K_i represent the same conjugacy class in $G_i/[G'_i, T_i]$. Then by Theorem 16.2 $C_{(K_{i-1}, X_{t_i})} - C_{(K_i, X_{t_i})}$ is a rational power series. By the proof of Theorem A each $C_{(K_i, X_{t_i})} = C_{(K_i, X_{t_{i+1}})}$, and $C_{(K_0, X_0)}$ is not rational. Then $C_{(K_{n-1}, X_1)}$ is not rational, contradicting our hypothesis that X_1 is a PL knot. \square

17. BOUNDARY-LINK-LIKE HANDLEBODIES

Lemma 17.1. *Let Σ be a closed orientable 2-manifold.*

(a) [69], [84], [10] (see also [31; §3], [77; §2]) *A nonzero class $\alpha \in H_1(\Sigma)$ is realizable by an embedded circle if and only if α is primitive.*⁴⁸

(b) [31] (see also [62; Corollary in §2]) *A linearly independent collection of classes $\alpha_1, \dots, \alpha_k \in H_1(\Sigma)$ is realizable by k pairwise disjoint embedded circles if and only if the subgroup $\langle \alpha_1, \dots, \alpha_k \rangle$ of $H_1(\Sigma)$ is its direct summand and $\alpha_i \cdot \alpha_j = 0$ for $i \neq j$.*

(c) [77; Lemma 6.2] *A collection of classes $\alpha_1, \dots, \alpha_k \in H_1(\Sigma)$ is realizable by k pairwise disjoint embedded circles whose union does not separate Σ if and only if they form a basis of a direct summand of $H_1(\Sigma)$ and $\alpha_i \cdot \alpha_j = 0$ for $i \neq j$.*

Let us note that (c) follows easily from (b), by applying the following lemma.

Lemma 17.2. *Let Σ be a closed 2-manifold and M its closed 1-submanifold. Then M does not separate Σ if and only if the classes of the connected components of M are linearly independent in $H_1(\Sigma)$.*

Proof. If $\Sigma \setminus M$ is disconnected, then the closure of any of its components, regarded as a 2-chain in M , provides a nontrivial linear dependence between the homology classes of (some of) the components of M . Conversely, if $\Sigma \setminus M$ is connected, then the restriction map $H^0(\Sigma) \rightarrow H^0(\Sigma \setminus M)$ is onto. Then by the Poincaré duality so is the homomorphism $H_2(\Sigma) \rightarrow H_2^\infty(\Sigma \setminus M) \simeq H_2(\Sigma, M)$. Then the boundary map $H_2(\Sigma, M) \rightarrow H_1(M)$ is zero, and hence the homomorphism $H_1(M) \rightarrow H_1(\Sigma)$ is injective. Therefore the homology classes of the components of M are linearly independent in $H_1(\Sigma)$. \square

By a *handlebody* in S^3 we mean a regular neighborhood of a PL embedded connected graph, or in other words a connected PL submanifold of S^3 which admits a handle decomposition with handles of indices 0 and 1 only.

Let $V \subset S^3$ be a handlebody and let $W = \overline{S^3 \setminus V}$. Let M_V and L_V be the kernels of the inclusion induced homomorphisms $\varphi: H_1(\partial V) \rightarrow H_1(V)$ and $\psi: H_1(\partial V) \rightarrow H_1(W)$.

⁴⁸An element of an abelian group is called *primitive* if it is not divisible by an integer greater than 1.

Lemma 17.3. (a) $H_i(W) \simeq H_i(V)$ (thus W is a “homology handlebody”);

(b) $H_1(\partial V) = M_V \oplus L$ and $\varphi|_{L_V}: L_V \rightarrow H_1(V)$ and $\psi|_{M_V}: M_V \rightarrow H_1(W)$ are isomorphisms;

(c) The intersection form $H_1(\partial V) \times H_1(\partial V) \rightarrow \mathbb{Z}$ vanishes on $L_V \times L_V$ and on $M_V \times M_V$.

Proof. (a). By the Alexander duality $\tilde{H}_i(W) \simeq \tilde{H}^{3-i-1}(V)$. Since V is a regular neighborhood of a connected graph, $\tilde{H}^{3-i-1}(V) = 0 = \tilde{H}_i(V)$ for $i \neq 1$ and $H^1(V) \simeq \text{Hom}(H_1(V), \mathbb{Z}) \simeq H_1(V)$. \square

(b). Since $H_1(S^3) = 0 = H_2(S^3)$, by the Mayer–Vietoris exact sequence the homomorphism $\varphi + \psi: H_1(\partial V) \rightarrow H_1(V) \oplus H_1(W)$ is an isomorphism. Hence the restriction $(\varphi + \psi)|_{L_V}$ is an isomorphism onto its image, but clearly it coincides with $\varphi|_{L_V}$ and so its image is $H_1(V)$. Thus $\varphi|_{L_V}: L_V \rightarrow H_1(V)$ is an isomorphism, and similarly so is $\psi|_{M_V}: M_V \rightarrow H_1(W)$. Also we get that the preimage of $H_1(V)$ under $\varphi + \psi$ is L_V , and similarly the preimage of $H_1(W)$ under $\varphi + \psi$ is M_V . Hence $H_1(\partial V) = M_V \oplus L_V$. \square

(c). Let $\alpha, \beta \in L_V$. These can be represented by elements of $\pi_1(\partial W)$, which in turn can be represented by loops $f, g: S^1 \rightarrow \partial W$ which are transverse to each other. Since f and g are null-homologous in W , they bound maps $F: \Sigma \rightarrow W$ and $G: \Sigma' \rightarrow W$ of some oriented surfaces, which may also be assumed to be transverse to each other. Then the oriented 0-manifold $f \cap g := \{(x, y) \in S^1 \times S^1 \mid f(x) = g(y)\}$ is the oriented boundary of the oriented 1-manifold $F \cap G := \{(x, y) \in \Sigma \times \Sigma' \mid F(x) = G(y)\}$. Hence $\alpha \cdot \beta = 0$. Similarly $\gamma \cdot \delta = 0$ for any $\gamma, \delta \in M_V$. \square

An oriented PL embedded circle in ∂V representing a nonzero element of L_V will be called a *longitude* of V . A collection of $\text{rk } L_V$ pairwise disjoint longitudes of V whose union is non-separating in ∂V will be called a *system of longitudes* for V .

Proposition 17.4. *Every basis of L_V can be represented by a system of longitudes of V , and every system of longitudes of V represents a basis of L_V .*

Of course, here one can speak of “bases of $H_1(V)$ ” instead of “bases of L_V ”.

Proof. This follows immediately from Lemmas 17.3 and 17.1. \square

Proposition 17.5. *Suppose that a basis of L_V is represented by a collection of $g := \text{rk } L_V$ pairwise disjoint embedded closed 1-manifolds $M_i \subset \partial V$ whose connected components are non-separating in ∂V . Then the following are equivalent:*

- (1) each M_i is connected;
- (2) $M_1 \cup \dots \cup M_g$ is non-separating in ∂V ;
- (3) (M_1, \dots, M_g) is a system of longitudes of V .

Proof. Since (3) is the conjunction of (1) and (2), it suffices to show that (1) \Leftrightarrow (2).

(1) \Rightarrow (2). This follows from Lemma 17.2.

(2) \Rightarrow (1). By Lemma 17.2 the classes of the components M_{ij} of the M_i are linearly independent in $H_1(\Sigma)$. On the other hand, since the M_{ij} are pairwise disjoint, they

have zero pairwise intersection numbers. Hence the M_{ij} form a basis of a subspace Λ of $H_1(\partial V)$ such that the intersection form vanishes on $\Lambda \times \Lambda$. Then $\text{rk } \Lambda \leq \frac{1}{2} \text{rk } H_1(\partial V) = g$, and therefore there are at most g of the M_{ij} . Hence each M_i is in fact connected. \square

Lemma 17.6. *Every longitude of V extends to a system of longitudes of V .*

Proof. By Lemma 17.1(a) the given longitude λ represents a primitive homology class. Hence $L_V / \langle \lambda \rangle$ is torsion free, and therefore (being finitely generated) is free abelian. Then the short exact sequence $0 \rightarrow \langle \lambda \rangle \rightarrow L_V \rightarrow L_V / \langle \lambda \rangle \rightarrow 0$ splits, and so $\langle \lambda \rangle$ is a direct summand of L_V . Let $\lambda_1, \dots, \lambda_{g-1}$ be a basis of a complementary summand. Then $\lambda_1, \dots, \lambda_{g-1}, \lambda$ is a basis of L_V . By Lemma 17.3 L_V is a direct summand of $H_1(\partial V)$ and the intersection form vanishes on $L_V \times L_V$. Then by [31; Proof of Proposition 4.2] $\lambda_1, \dots, \lambda_{g-1}$ can be represented by PL embedded circles in ∂V , pairwise disjoint from each other and disjoint from λ . By Lemma 17.2 these form a system of longitudes of V . \square

Lemma 17.7. *Let $m \in M_V$ be a nonzero primitive homology class. Then m can be represented by the boundary of a properly embedded disk in V .*

Proof. There is a collection of pairwise disjoint properly embedded disks D_1, \dots, D_g in V such that V cut open along these disks is the 3-ball. On the other hand, by Lemma 17.1(a) m can be realized by an embedded circle μ_1 in ∂V , and by the proof of Lemma 17.6 this extends to a collection of embedded circles μ_1, \dots, μ_g whose union is non-separating in ∂V and whose homology classes form a basis of M_V . Then it is easy to find embedded circles $\lambda_1, \dots, \lambda_g$ such that each λ_i meets μ_i transversely in a single point and is disjoint from μ_j for $j \neq i$. Arguing by induction, it is easy to see that the λ_i can be chosen pairwise disjoint. A regular neighborhood T_i of $\lambda_i \cup \mu_i$ is a punctured torus, and we may assume that the T_i are pairwise disjoint. By counting the Euler characteristic $\partial V \setminus T$ is a sphere with g punctures. If we repeat the same construction starting from the ∂D_i in place of the μ_i , we will again end up with punctured tori and a sphere with g punctures. Using this, it is easy to construct a homeomorphism φ of ∂V onto itself sending each ∂D_i onto μ_i .

In particular, we get a symplectic automorphism φ_* of $H_1(\partial V)$ sending each $[\partial D_i]$ onto $[\mu_i]$. Since $\varphi_*(M) = M$, by [13; Lemma 2.2] or [40; Theorem 2.1] φ_* is realized by a homeomorphism ψ of V onto itself. The latter sends ∂D_1 onto an embedded circle which represents the homology class m and bounds the disk $\psi(D_1)$. \square

Lemma 17.8. *If M is a 3-manifold (possibly noncompact) and Q is a smooth knot in ∂M which is null-homologous in M , then Q bounds a smooth embedded surface in M .*

Presumably this is well-known, but we include a proof for completeness.

Proof. Since $[Q] \in H_1(\partial M)$ maps to $0 \in H_1(M)$, it is the image of some $\alpha \in H_2(M, \partial M)$. Then the Poincaré dual⁴⁹ $D[Q] \in H_c^1(\partial M)$ is the image of $D(\alpha) \in H_c^1(M)$. The Pontryagin construction (see for instance [71]) yields a smooth map $f: \partial M \rightarrow S^1$ such that

⁴⁹See for instance [39; Theorem 3.35] for the required version of the Poincaré duality.

$f(U) = 1$ for some neighborhood of infinity U (=complement of a compact set) and $Q = f^{-1}(-1)$. Its homotopy class corresponds to $D[Q]$. Since $D[Q]$ is the image of α , f extends to a smooth map $\bar{f}: M \rightarrow S^1$ such that $\bar{f}(\bar{U}) = 1$ for some neighborhood of infinity \bar{U} . Then $\bar{f}^{-1}(-1)$ is the desired embedded surface. \square

Lemma 17.9. *Let $K = \bigvee_{i=1}^n K_i$ be a wedge of circles embedded in S^3 and Q be a knot in $S^3 \setminus K$. If Q is null-homologous in each $S^3 \setminus K_i$, then Q is null-homologous in $S^3 \setminus K$.*

Algebraic proof. By an inductive application of the Mayer–Vietoris exact sequence, using that $H_2(S^3 \setminus pt) = 0$. \square

Geometric proof. Q bounds a Seifert surface Σ which is disjoint from the wedge point and transverse to each K_i . Since $\text{lk}(Q, K_i) = 0$, we have $\Sigma \cdot K_i = 0$. Then K_i contains an arc J such that ∂J consists of two intersection points of Σ and K_i of opposite signs and J itself is disjoint from the other intersection points and from the wedge point. By attaching a tube to Σ going along J we remove the two intersection points. Proceeding in this fashion we obtain a new Seifert surface Σ' for Q which is disjoint from K . \square

Theorem 17.10. *Let $g = \text{rk } H_1(V)$. If $g = 2$, then the following are equivalent:*

- (1) *some system of longitudes of V bounds disjoint Seifert surfaces in W ;*
- (2) *every nonzero primitive class of L_V is represented by a longitude of V which bounds a Seifert surface Σ in W such that every knot in Σ is null-homologous in W ;*
- (3) *some longitude of V bounds a Seifert surface Σ in W such that every knot in Σ is null-homologous in W .*

For arbitrary g we still have (1) \Rightarrow (2) \Rightarrow (3).

The proof of (3) \Rightarrow (1) for $g = 2$ is found in [95; Theorem 3.3], of which the author was unaware when writing it up. The idea of proof of (1) \Rightarrow (2) comes from [46; proof of Theorem 5].

Proof. (1) \Rightarrow (2). Let $l \in L_V$ be the given primitive class. By Theorem 1.5 there exists a map f from W to a handlebody U which restricts to a homeomorphism $\partial W \rightarrow \partial U$. It is clear that $L_U = M_V$ and $M_U = L_V$ (see Lemma 17.3). Then by Lemma 17.7 $l \in L_V = M_U$ is realized by an embedded circle λ in $\partial W = \partial U$ which bounds a disk D in U . We may assume that f is transverse to D . Then $\Sigma := f^{-1}(D)$ is a Seifert surface for λ in W . Moreover, every knot $Q \subset \Sigma$ is sent by f into D , and hence represents a conjugacy class in the kernel of $f_*: \pi_1(W) \rightarrow \pi_1(U)$. Then by [46; Corollary to Theorem 1] Q represents a conjugacy class in $\gamma_\omega \pi_1(W)$, and in particular in $\gamma_2 \pi_1(W)$. Hence it is null-homologous in W .

(2) \Rightarrow (3). This is trivial.

(3) \Rightarrow (1) for $g = 2$. The hypothesis implies that each knot Q in the interior of Σ is null-homologous in $S^3 \setminus V$. On the other hand, by Lemma 17.6 λ extends to a system of longitudes λ, λ' . Since $\lambda' \subset V$, we have $\text{lk}(Q, \lambda') = 0$. Hence λ' is null-homologous in the complement to every knot in Σ . Now Σ deformation retracts onto a wedge of circles

$X = \bigvee_{k=1}^{2g_1} Q_j$. By Lemma 17.9 λ' is null-homologous in $S^3 \setminus X$. By Lemma 17.8 λ' bounds a Seifert surface F in $S^3 \setminus X$. We may assume that F intersects Σ transversely. It is easy to see that $\Sigma \setminus X$ is homeomorphic to $S^1 \times [0, 1)$. Hence every component C of $F \cap \Sigma$ bounds a surface $F_C \subset \Sigma$. If C is innermost on Σ among all such components, then by surgering F along F_C we can get rid of the component C . In the end we obtain a Seifert surface Σ' for λ' in $S^3 \setminus \Sigma$. \square

Remark 17.11. If V is of boundary type, it is not true that *every* longitude of V bounds a Seifert surface Σ in W such that every knot in Σ is null-homologous in W . For instance, let V be the unknotted handlebody of genus $g \geq 2$. Thus $\pi_1(\partial V) \simeq \langle m_1, \dots, m_g, l_1, \dots, l_g \mid [m_1, l_1] \cdots [l_g, m_g] = 1 \rangle$ and the map $\pi_1(\partial V) \rightarrow \pi_1(V)$ sends each m_i to 1 and each l_i to the i th free generator; and the map $\pi_1(\partial V) \rightarrow \pi_1(W)$ sends each l_i to 1 and each m_i to the i th free generator. Then any longitude of V representing $[m_i, m_j]l_k$, where $i \neq j$, is not going to be represented by an element of $\pi_1(W)''$. Hence any such longitude does not bound a Seifert surface Σ in W such that every knot in Σ is null-homologous in W .

Theorem 17.12. *Let V be a genus g PL handlebody in S^3 and let $W = \overline{S^3 \setminus V}$. Then the following conditions are equivalent to those of Theorem 1.5:*

- (1') *W admits a retraction onto a PL embedded wedge $Q \subset \partial W$ of g whiskered circles,⁵⁰ whose circles bound pairwise disjoint PL embedded disks in V ;*
- (2') *$\pi_1(W)$ admits a homomorphism to the free group F_g such that for some $Q \subset \partial W$ as in (1') the composition $\pi_1(Q) \rightarrow \pi_1(W) \rightarrow F_g$ is surjective;*
- (3') *$\pi_1(W)/\gamma_\omega \pi_1(W)$ is a free group and it is the image of $\pi_1(Q)$ for some $Q \subset \partial W$ as in (1');*
- (4') *W contains pairwise disjoint properly PL embedded surfaces $\Sigma_1, \dots, \Sigma_g$ and V contains pairwise disjoint properly PL embedded disks D_1, \dots, D_g such that each $\partial \Sigma_i$ is connected and meets ∂D_i transversely in a single point, and $\partial \Sigma_i \cap \partial D_j = \emptyset$ for $i \neq j$.*

Proof. (5) \Rightarrow (4'). This is similar to [8; proof of Theorem 7.3]. The given map $f: W \rightarrow U$ extends to $f \cup \text{id}_V: S^3 \rightarrow U \cup_h V$, where h is the homeomorphism $f|_{\partial W}: \partial V \rightarrow \partial U$. Clearly $M := U \cup_h V$ is a homology sphere (see Lemma 17.3). Since S^3 is simply-connected, f factors through the universal covering $p: \tilde{M} \rightarrow M$. Since M is orientable and f has degree 1, it induces an isomorphism on integral 3-homology, and hence so does p . It follows that M is simply-connected. Thus by Perelman's theorem $M \cong S^3$. Then $U \cup V$ is a Heegaard splitting of S^3 , and hence by Waldhausen's theorem (see [85]) it must be the standard one of genus g . Then V contains pairwise disjoint properly PL embedded 2-disks D_1, \dots, D_g and U contains pairwise disjoint properly PL embedded 2-disks $\Delta_1, \dots, \Delta_g$ such that each ∂D_i meets $\partial \Delta_i$ transversely in a single point and $\partial D_i \cap \partial \Delta_j = \emptyset$ for $i \neq j$. We may assume that the map $f: W \rightarrow U$ is transverse to each Δ_i , and then it remains to set $\Sigma_i = f^{-1}(\Delta_i)$.

⁵⁰A *whiskered circle* is the graph "letter P", endowed with a basepoint, which is its bottom point.

$(4') \Rightarrow (4)$. A regular neighborhood T_i of $\partial\Sigma_i \cup \partial D_i$ is a punctured torus, and we may assume that the T_i are pairwise disjoint. Then $T := T_1 \cup \dots \cup T_g$ is non-separating in ∂V (in fact, by counting the Euler characteristic $\partial V \setminus T$ is a sphere with g punctures). In particular, $\Sigma_1 \cup \dots \cup \Sigma_g$ is non-separating in ∂V .

$(1') \Leftrightarrow (2') \Leftrightarrow (3') \Leftrightarrow (4')$. Similarly to [89] (see also [46], [44]). \square

18. PROOF OF THEOREM F

Theorem 18.1. *Let \mathcal{K} be a knot in S^3 which is the intersection of a sequence of handlebodies $V_0 \supset V_1 \supset \dots$ and let $K_i \subset \partial V_i$ be knots such that each K_i is homologous to \mathcal{K} in V_i . Suppose that each K_i bounds a Seifert surface Σ_i in $W_i := S^3 \setminus \overline{V_i}$ such that each knot in Σ_i is null-homologous in W_i . Then the knot module of \mathcal{K} is a torsion module.*

Proof. The knot module of \mathcal{K} is the $\mathbb{Z}[t^{\pm 1}]$ -module $H_1(X)$, where X is the infinite cyclic cover of $S^3 \setminus K$. Every element of $H_1(X)$ can be represented by a knot $\tilde{Q}_0 \subset X$. We may assume that \tilde{Q}_0 projects to a knot Q_0 (i.e. Q_0 has no self-intersections) in $S^3 \setminus K$. Since Q_0 is null-homologous in $S^3 \setminus K$, it bounds a Seifert surface F_0 in $S^3 \setminus K$. Then F_0 is disjoint from some V_n . We may assume that F_0 intersects Σ_n transversely. Their intersection Q_1 is null-homologous in W_n , and hence bounds a Seifert surface F_1 in W_n . We may assume that F_1 intersects Σ_n transversely. Their intersection Q_2 is null-homologous in W_n . Continuing in this fashion, we obtain links $Q_1, Q_2, \dots \subset \Sigma_n$ which are null-homologous in W_n and in particular in $S^3 \setminus K$. Let $\tilde{Q}_1, \tilde{Q}_2, \dots$ be their lifts to X . They lie in the preimage \tilde{W}_n of W_n in X . This \tilde{W}_n can also be described as the cover of W_n corresponding to (the kernel of) the homomorphism $\pi_1(W_n) \rightarrow \mathbb{Z}$ given by $[Z] \mapsto \text{lk}(Z, \mathcal{K})$. (In other words, this homomorphism is determined by the class in $H^1(W_n)$ which is Alexander dual to $[\mathcal{K}] \in H_1(V_n)$.) Hence a fundamental domain of the action of \mathbb{Z} on \tilde{W}_n is a 3-manifold with boundary whose interior as a subset of \tilde{W}_n is homeomorphic to $W_n \setminus \Sigma_n$; two consecutive copies of the fundamental domain meet along a copy of Σ_n . Then each \tilde{Q}_i (starting from $i = 0$) is homologous in \tilde{W}_n to $(1-t)\tilde{Q}_{i+1}$. Namely, $\tilde{Q}_i - (1-t)\tilde{Q}_{i+1} = \partial \tilde{F}_i$, where \tilde{F}_i is a surface with boundary whose interior as a subset of \tilde{W}_n is homeomorphic to $F_i \setminus \Sigma_n$. Thus each $[\tilde{Q}_i] = (1-t)[\tilde{Q}_{i+1}] \in H_1(\tilde{W}_n)$. Since W_n is a compact 3-manifold, $H_1(\tilde{W}_n)$ is finitely generated over $\mathbb{Z}[t^{\pm 1}]$. Since $\mathbb{Z}[t^{\pm 1}]$ is Noetherian, so is $H_1(\tilde{W}_n)$. Hence its submodules $\langle [Q_0] \rangle \subset \langle [Q_1] \rangle \subset \dots$ stabilize at some point. Thus there exist a k and a Laurent polynomial p such that $[Q_k] = p[Q_{k-1}]$. Since $[Q_{k-1}] = (1-t)[Q_k]$, we get $[Q_k] = p(1-t)[Q_k]$, or $(p(t-1) - 1)[Q_k] = 0$. Hence also $(p(t-1) - 1)[Q_0] = (p(t-1) - 1)(1-t)^k[Q_k] = 0$. \square

Corollary 18.2. *If a knot $\mathcal{K} \subset S^3$ is the intersection of a nested sequence of boundary-link-like handlebodies, then the knot module of \mathcal{K} is a torsion module.*

Theorem F follows from Corollaries 16.3 and 18.2.

Proof. Let $V_1 \supset V_2 \supset \dots$ be the given handlebodies. Since $H_2(V_i) = 0$, we have $\lim^1 H_2(V_i) = 0$ and consequently by Milnor's short exact sequence $\lim H_1(V_i) \simeq H_1(\mathcal{K})$. Moreover, this isomorphism sends the thread consisting of the classes $[\mathcal{K}] \in H_1(V_i)$ to

the fundamental class in $H_1(\mathcal{K})$, which is nonzero. Hence $[\mathcal{K}] \in H_1(V_i)$ is nonzero for all sufficiently large i . By omitting a finite number of initial handlebodies we may assume that all of these classes are nonzero. Moreover, all of them are primitive, since so is the fundamental class of $H_1(\mathcal{K})$.

Let $W_i = \overline{S^3 \setminus V_i}$. Since $L_V = \ker(\psi: H_1(\partial V_i) \rightarrow H_1(W_i))$ is mapped isomorphically onto $H_1(V)$ by the restriction of $\varphi: H_1(\partial V_i) \rightarrow H_1(V_i)$ (see Lemma 17.3(b)), each $[\mathcal{K}] \in V_i$ is the image of some nonzero primitive class $l_i \in L_{V_i}$. By Lemma 17.10 this class l_i is represented by a longitude K_i of V which bounds a Seifert surface Σ in W such that every knot in Σ is null-homologous in W . Then by Theorem 18.1 the knot module of \mathcal{K} is a torsion module. \square

APPENDIX A. EVERY KNOT IS I -EQUIVALENT TO THE UNKNOT

This Appendix contains a proof of the following result, due largely to C. Giffen.

Theorem A.1. *Every topological knot is I -equivalent to the unknot.*

Theorem A.1 follows from Proposition A.3 and Theorem A.2.

An alternative proof of Theorem A.1 can be found in Ancel's paper [3].

In fact, the reason why I wrote down a proof of Theorem A.1 was that I had previously stated this result as known, with attribution to Giffen [65]. In August 2021 I was contacted by M. Freedman who mentioned among other things that he and R. Ancel are wondering about the details of the proof. In a few days I've sent him (Freedman) an email with a writeup of the proof of Theorem A.2 (essentially the same as below), suggesting that he forward it to Ancel. I've also sent the same proof to Ancel himself in response to his email in early September, but for some reason he did not get my reply. Eventually he got my email in December 2021, but by that time he already wrote down his own account [3].

A.1. F -isotopy implies I -equivalence.

Theorem A.2 (Giffen [33], [34]). *Every topological knot in $S^1 \times D^2$ that is homotopic to the core circle is I -equivalent to the core circle.*

Ancel's alternative exposition of what appears to be more or less the same construction [3] is more detailed and more thoughtful, but also much longer.

Proof. Let $k: S^1 \rightarrow S^1 \times D^2$ be the given knot. The identity $\text{id}: S^1 \times D^2 \rightarrow S^1 \times D^2 \times \{0\}$ and the core circle embedding $S^1 \times \{\infty\} \rightarrow S^1 \times D^2 \times \{1\}$ cobound an embedding $E: S^1 \times CD^2 \rightarrow S^1 \times D^2 \times [0, 1]$, where CD^2 is the cone over D^2 with apex ∞ . We may identify CD^2 with the one-point compactification of $D^2 \times [0, \infty)$, with ∞ being the added point. Let \tilde{w} denote the composition $S^1 \times \mathbb{R} \xrightarrow{k \times \text{id}_{\mathbb{R}}} S^1 \times D^2 \times \mathbb{R} \xrightarrow{\tilde{\varphi}} S^1 \times D^2 \times \mathbb{R}$, where $\tilde{\varphi}$ is the level-preserving homeomorphism given by $\tilde{\varphi}(\alpha, x, t) = (\alpha - [t], x, t)$; hereafter we identify S^1 with \mathbb{R}/\mathbb{Z} . The closure \bar{A} of $A := \tilde{w}(S^1 \times [0, \infty))$ in $S^1 \times CD^2$ is the union of A and $S^1 \times \{\infty\}$. We will show that \bar{A} is homeomorphic to $S^1 \times I$; this easily implies that $E(\bar{A})$ is the image of an I -equivalence between k and the core circle.

Let $K = k(S^1)$ and let $\tilde{W} = \tilde{w}(S^1 \times \mathbb{R})$. Let φ be the self-homeomorphism of $S^1 \times D^2 \times S^1$ given by $\varphi(\alpha, x, \beta) = (\alpha - \beta, x, \beta)$, and let $W = \varphi(K \times S^1)$. Then \tilde{W} is the preimage of W under the covering $\text{id}_{S^1 \times D^2} \times q: S^1 \times D^2 \times \mathbb{R} \rightarrow S^1 \times D^2 \times S^1$, where q is the quotient map $\mathbb{R} \rightarrow S^1$. On the other hand, let ψ be the self-homeomorphism of $S^1 \times D^2 \times S^1$ given by $\psi(\alpha, x, \beta) = (\alpha, x, \alpha - \beta)$. Then ψ takes $K \times S^1$ onto itself. Thus $W = \chi(K \times S^1)$, where $\chi = \varphi \circ \psi$. Let us note that $\chi(\alpha, x, \beta) = (\beta, x, \alpha - \beta)$.

Let \tilde{K} be the preimage of K under $q \times \text{id}_{D^2}: \mathbb{R} \times D^2 \rightarrow S^1 \times D^2$. The covering translation $\sigma: \mathbb{R} \times D^2 \rightarrow \mathbb{R} \times D^2$, given by $\sigma(t, x) = (t + 1, x)$, restricts to a homeomorphism $\sigma_K: \tilde{K} \rightarrow \tilde{K}$. Since σ is isotopic to $\text{id}_{\mathbb{R} \times D^2}$, its mapping torus $MT(\sigma)$ is homeomorphic to $\mathbb{R} \times D^2 \times S^1$. Writing $MT(\sigma)$ as $\mathbb{R} \times D^2 \times [0, 1]/(t, x, 0) \sim (t+1, x, 1)$, a specific homeomorphism $\tilde{\chi}: MT(\sigma) \rightarrow S^1 \times D^2 \times \mathbb{R}$ is given by $\tilde{\chi}([(t, x, s)]) = ([s], x, t - s)$. Since σ is a lift of $\text{id}_{S^1 \times D^2}$, there is a covering $p: MT(\sigma) \rightarrow MT(\text{id}_{S^1 \times D^2})$ and a commutative diagram

$$\begin{array}{ccc} MT(\sigma) & \xrightarrow{\tilde{\chi}} & S^1 \times D^2 \times \mathbb{R} \\ p \downarrow & & \text{id}_{S^1 \times D^2} \times q \downarrow \\ S^1 \times D^2 \times S^1 & \xrightarrow{\chi} & S^1 \times D^2 \times S^1. \end{array}$$

It is easy to see that $MT(\sigma_K) = p^{-1}(K \times S^1)$. Since $K \times S^1 = \chi^{-1}(W)$, we obtain that $MT(\sigma_K) = \tilde{\chi}^{-1}(\tilde{W})$.

We may regard the suspension ΣD^2 as the compactification of $\mathbb{R} \times D^2$ obtained by adding the two suspension points $\pm\infty$. Then σ uniquely extends to a homeomorphism $\bar{\sigma}: \Sigma D^2 \rightarrow \Sigma D^2$. In turn, $\bar{\sigma}$ restricts to a homeomorphism $\bar{\sigma}_K: \bar{K} \rightarrow \bar{K}$, where $\bar{K} = \tilde{K} \cup \{\pm\infty\}$. Since $\tilde{\chi}$ commutes with the ‘‘projection’’ $MT(\sigma) \rightarrow S^1$, $[(t, x, s)] \mapsto [s]$, and the projection $S^1 \times D^2 \times \mathbb{R} \rightarrow S^1$, it extends to a homeomorphism $\bar{\chi}: MT(\bar{\sigma}) \rightarrow S^1 \times \Sigma D^2$. (To be precise, ΣD^2 is regarded here as a compactification of $D^2 \times \mathbb{R}$ rather than $\mathbb{R} \times D^2$.) Clearly, $MT(\bar{\sigma}_K) = \bar{\chi}^{-1}(\bar{W})$, where $\bar{W} = \tilde{W} \cup S^1 \times \{\pm\infty\}$. On the other hand, since $\bar{\sigma}_K$ is isotopic to $\text{id}_{\bar{K}}$, its mapping torus is homeomorphic to $\bar{K} \times S^1$. Thus \bar{W} is homeomorphic to $S^1 \times [-1, 1]$. Let us note that this homeomorphism sends \tilde{W} onto $S^1 \times (-1, 1)$. Now $\bar{A} = \bar{W} \cap S^1 \times CD^2$, where $\bar{W} \cap S^1 \times D^2$ is a copy of S^1 contained in \tilde{W} , so it is clear that \bar{A} is homeomorphic to $S^1 \times [0, 1]$. \square

A.2. Every knot is F-isotopic to the unknot.

Proposition A.3. *Every oriented topological knot K in S^3 lies in the interior of an unknotted solid torus T such that the inclusion $K \rightarrow T$ is a homotopy equivalence.*

Ancel’s proof of Proposition A.3 in [3] is decidedly simpler than what follows. But perhaps the approach below might still be of some interest, as it answers some questions that naturally arise in connection with this statement.

Lemma A.4. *If K is a topological knot in S^3 , then every PL knot Q in $S^3 \setminus K$ is homotopic in $S^3 \setminus K$ to an unknot.*

It turns out that this lemma is proved in the literature: McMillan proved that if X is a compact 1-dimensional ANR in S^3 , then X is the intersection of a nested sequence

of handlebodies [60; Lemmas 2 and 3], and Bothe noted that this property implies that every knot Q in $S^3 \setminus X$ is homotopic in $S^3 \setminus X$ to a knot which is unknotted as a knot in S^3 [17; assertion “(A) implies (C)” on p. 176].

Proof. Let us pick some point in $S^3 \setminus (K \cup Q)$ and identify its complement in S^3 with \mathbb{R}^3 by a PL homeomorphism. By general position there exists a linear surjection $\pi: \mathbb{R}^3 \rightarrow \mathbb{R}^2$ whose restriction to Q has only finitely many double points and no triple points. If $J \subset Q$ is a small arc whose image does not contain any double points, and $f: Q \rightarrow \mathbb{R}$ is a PL map which is monotone on $Q \setminus J$, then $\pi \times f: Q \rightarrow \mathbb{R}^2 \times \mathbb{R}$ is easily seen to be a PL unknot. Hence we can unknot Q (in \mathbb{R}^3) by changing some of the crossings in its plane diagram $\pi(Q)$. Unfortunately, we may not assume that these crossings avoid $\pi(K)$.⁵¹ However, by a well-known theorem asserting that K is tame in the sense of Shtan’ko (or has embedding dimension 1 in another terminology) [60; Lemma 3], [16; Satz 4] (see also [87; Theorem 4]), there exists an ambient isotopy $h_t: \mathbb{R}^3 \rightarrow \mathbb{R}^3$, $h_0 = \text{id}$, keeping Q fixed and such that for every double point $\pi(p) = \pi(q)$, $p, q \in Q$, the straight line segment $[p, q]$ between p and q is disjoint from $h_1(K)$.⁵² Therefore the desired crossing changes can be realized by a PL homotopy K_t within $\mathbb{R}^3 \setminus h_1(K)$. The resulting PL knot K_1 is unknotted in \mathbb{R}^3 and homotopic to Q in $\mathbb{R}^3 \setminus h_1(K)$. Hence $h_1^{-1}(K_1)$, which is still an unknot, is homotopic to $h_1^{-1}(Q) = Q$ in $\mathbb{R}^3 \setminus K$. \square

Given disjoint oriented topological knots K and Q in S^3 , let $\text{lk}(K, Q)$ denote the integer $[Q] \in H_1(S^3 \setminus K) \xrightarrow{A} H^1(K) \xrightarrow{P} H_0(K) = \mathbb{Z}$, where A is the Alexander duality (involving Čech cohomology and assuming a fixed orientation of S^3) and P is the Poincaré duality $c \mapsto [K] \cap c$.

Lemma A.5. $\text{lk}(K, Q) = \text{lk}(Q, K)$.

Proof. It is easy and well-known that $\text{lk}(K, Q) = \text{lk}(Q, K)$ for PL links (K, Q) . Also it is easy to show that any two sufficiently close PL approximations of a given topological link (K, Q) are link homotopic (that is, homotopic by a homotopy that keeps the two components disjoint), and therefore have the same linking number. This yields an extension lk_{PL} of lk from PL to topological links, which is symmetric. It remains to show that $\text{lk} = \text{lk}_{PL}$.

Let T be some triangulation of S^3 and let N_i be the union of all simplexes of the i th derived subdivision $T^{(i)}$ that meet K . Then each N_i is a closed neighborhood of K which is a 3-manifold with boundary, $N_1 \supset N_2 \supset \dots$ and $\bigcap_i N_i = K$. The Alexander

⁵¹K. Borsuk, answering a question of R. H. Fox, constructed a knot in \mathbb{R}^m for each $m \geq 3$ whose projection to every hyperplane has a nonempty interior [15].

⁵²The construction of h_t is quite easy. Let $J_{pp'}$ (resp. $J_{qq'}$) be a small arc of Q with p (resp. q) as one of its endpoints. The join $J_{pp'} * J_{qq'}$ is a ball, and its boundary sphere Σ is not separated by $\Sigma \cap K$ since $\tilde{H}_0(\Sigma \setminus K) \simeq H^1(\Sigma \cap K) = 0$ due to $K \not\subset \Sigma$. Hence p and q are joined by a PL arc J_{pq} in $\Sigma \setminus K$. We may assume that J_{pq} meets $J_{pp'}$ only in p and $J_{qq'}$ only in q . Then there is a PL ambient isotopy of Σ keeping $J_{pp'} \cup J_{qq'}$ fixed from id_Σ to a homeomorphism sending J_{pq} onto $[p, q]$. It extends to a PL ambient isotopy h_t^{pq} of \mathbb{R}^3 with support in a small neighborhood of $J_{pp'} * J_{qq'}$ and keeping Q fixed. The composition $h_t^{p_1 q_1} h_t^{p_2 q_2} \dots$ of all such isotopies is the desired isotopy h_t .

duality isomorphism $A: H_1(S^3 \setminus K) \xrightarrow{\cong} H^1(K)$ is the direct limit of the Alexander duality isomorphisms $A_i: H_1(S^3 \setminus N_i) \xrightarrow{\cong} H^1(N_i)$. There exists an r such that $Q \cap N_r = \emptyset$. Let $i: K \rightarrow N_r$ and $i': Q \rightarrow S^3 \setminus N_r$ be the inclusion maps. As long as we fix homeomorphisms of K and Q with S^1 , it makes sense to say that i and i' are homotopic to PL embeddings $e: K \rightarrow N_r$ and $e': Q \rightarrow S^3 \setminus N_r$.

Now $\text{lk}(K, Q)$ is the image of $i'_*[Q] \in H_1(S^3 \setminus N_r)$ in $H_1(S^3 \setminus K) \xrightarrow{A} H^1(K) \xrightarrow{P} H_0(K) = \mathbb{Z}$. This is the same as the image of $A_r(i'_*[Q]) \in H^1(N_r)$ in $H^1(K) \xrightarrow{P} H_0(K) = \mathbb{Z}$. Thus $\text{lk}(K, Q) = i^*(A_r(i'_*[Q])) \frown [K] \in H_0(K) = \mathbb{Z}$. Writing $K' = e(K)$ and $Q' = e'(Q)$, we similarly have $\text{lk}(K', Q') = e^*(A_r(e'_*[Q])) \frown [K] \in H_0(K) = \mathbb{Z}$. But $i^* = e^*$ and $i'_* = e'_*$. Thus $\text{lk}(K, Q) = \text{lk}(K', Q')$, and it follows that $\text{lk}(K, Q) = \text{lk}_{PL}(K, Q)$. \square

Remark A.6. A direct algebraic (but rather long) proof of Lemma A.5 can be found in [66]. It shows that each of $\text{lk}(K, Q)$ and $\text{lk}(Q, K)$ is equal to $\text{lk}_S(K, Q) := \langle S[Q], i_*[K] \rangle$, where $i: K \rightarrow S^3 \setminus Q$ is the inclusion and $S: H_1(Q) \rightarrow H^1(S^3 \setminus Q)$ is the Steenrod duality. In more detail, the equality $\text{lk}(K, Q) = \text{lk}_S(K, Q)$ is an easy consequence of the Sitnikov duality (for arbitrary subsets of S^3 , not necessarily open or closed). To prove that $\text{lk}_S(K, Q) = \text{lk}(Q, K)$, one defines an evaluation pairing $H^k(X) \otimes H_k(X) \rightarrow \mathbb{Z}$ between Čech cohomology and Steenrod homology of the compactum X , and shows that when $X \subset S^n$, it is determined by the usual evaluation pairing $H^k(S^n \setminus X) \otimes H_k(S^n \setminus X) \rightarrow \mathbb{Z}$.

Lemma A.7. *If K is an oriented topological knot in S^3 , then there exists an oriented PL knot Q in $S^3 \setminus K$ such that $\text{lk}(Q, K) = 1$ and Q is unknotted (in S^3).*

Proposition A.3 is an easy consequence of Lemma A.7.

Proof. The positive generator of $H_1(S^3 \setminus K) \xrightarrow{A} H^1(K) \xrightarrow{P} H_0(K) = \mathbb{Z}$ can certainly be represented by a PL knot Q' in $S^3 \setminus K$. By Lemma A.4 Q' is homotopic to a PL unknot Q in $S^3 \setminus K$. Hence by Lemma A.5 $\text{lk}(Q, K) = \text{lk}(K, Q) = \text{lk}(K, Q') = 1$. \square

ACKNOWLEDGEMENTS

I would like to thank P. M. Akhmetiev, F. D. Ancel, M. H. Freedman, L. Funar, M. Il'insky and A. Zastrow for useful discussions and thoughtful remarks. Also I'm grateful to R. D. Edwards, M. Farber, J. Keesling and S. P. Novikov for stimulating my interest in isotopy of wild knots (around 2005).

REFERENCES

- [1] S. Akbulut, *A solution to a conjecture of Zeeman*, *Topology* **30** (1991), 513–515. [ScienceDirect](#). $\uparrow 1, 22$
- [2] J. W. Alexander, *Topological invariants of knots and links*, *Trans. Amer. Math. Soc.* **30** (1928), 275–306. $\uparrow 26, 7.2$
- [3] F. D. Ancel, *Semi-isotopic knots*. [arXiv:2112.13940](#). $\uparrow A, A.1, A.2$
- [4] ———, *Isotopies, semi-isotopies and thickenings of wild knots and links*, Video of talk at the meeting of the Moscow Geometric Topology Seminar dedicated to A. V. Chernavsky's 85th birthday (February 2023). https://www.mathnet.ru/php/seminars.phtml?&presentid=38111&option_lang=eng. $\uparrow 3$

- [5] F. D. Ancel and C. R. Guilbault, *Mapping swirls and pseudo-spines of compact 4-manifolds*, Topology Appl. **71** (1996), 277–293. [Journal](#). [↑](#)
- [6] J. Amundsen, E. Anderson, C. W. Davis, and D. Guyer, *The C-complex clasp number of links*, Rocky Mountain J. Math. **50** (2020), 839–850. [arXiv:1907.12472](#). [↑3.5](#)
- [7] G. Baumslag, F. B. Cannonito, and D. J. S. Robinson, *The algorithmic theory of finitely generated metabelian groups*, Trans. Amer. Math. Soc. **344** (1994), 629–648. [Journal](#). [↑3](#)
- [8] R. Benedetti and R. Frigerio, *Levels of knotting of spatial handlebodies*, Trans. Amer. Math. Soc. **365** (2013), 2099–2167. [arXiv:1101.2151](#). [↑1](#), [17](#)
- [9] M. Benheddi and D. Cimasoni, *Link Floer homology categorifies the Conway function*, Proc. Edinburgh Math. Soc. **59** (2016), 813–836. [arXiv:1408.3517](#). [↑6.2](#), [6.3](#)
- [10] D. Bennequin, *Sur les représentants lisses des classes d’homologie entière de codimension 1 des variétés*, C. R. Acad. Sci. Paris Sér. A–B **285** (1977), no. 4, A207–A210. [BnF Gallica](#). [↑17.1](#)
- [11] R. Bieri, *Mayer–Vietoris sequences for HNN-groups and homological duality*, Math. Z. **143** (1975), 123–130. [EuDML](#). [↑18](#)
- [12] R. H. Bing, *A simple closed curve that pierces no disk*, J. Math. Pures Appl. **35** (1956), 337–343. [↑1](#), [1](#), [5](#)
- [13] J. S. Birman, *On the equivalence of Heegaard splittings of closed, orientable 3-manifolds*, Knots, Groups, and 3-Manifolds, Ann. Math. Studies, vol. 84, Princeton Univ. Press, Princeton, NJ, 1975, pp. 137–164. [↑17](#)
- [14] W. A. Blankinship and R. H. Fox, *Remarks on certain pathological open subsets of 3-space and their fundamental groups*, Proc. Amer. Math. Soc. **1** (1950), 618–624. [Journal](#). [↑2.2](#)
- [15] K. Borsuk, *An example of a simple arc in space whose projection in every plane has interior points*, Fund. Math. **34** (1947), 272–277. [EuDML](#). [↑51](#)
- [16] H. G. Bothe, *Universalmengen bezüglich der Lage im E^n* , Fund. Math. **56** (1964), 203–212 (German). [EuDML](#). [↑7](#), [A.2](#)
- [17] ———, *A wildly embedded 1-dimensional compact set in S^3 each of whose components is tame*, Fund. Math. **99** (1978), 175–187. [EuDML](#). [↑7](#), [A.2](#)
- [18] M. Brin, *Curves isotopic to tame curves*, Continua, Decompositions, Manifolds, Univ. Texas Press, Austin, TX, 1983, pp. 163–166. [↑1](#), [1](#)
- [19] E. J. Brody, *On infinitely generated modules*, Quart. J. Math. Oxford **11** (1960), 141–150. [↑2](#), [1.6](#)
- [20] G. Burde and H. Zieschang, *Knots*, 2nd ed., De Gruyter Studies in Math., vol. 5, de Gruyter, Berlin, 2003. [↑2](#), [4.4](#)
- [21] D. Cimasoni, *A geometric construction of the Conway potential function*, Comment. Math. Helv. **79** (2004), 124–146. [arXiv:math/0406172](#). [↑6](#), [29](#), [6.2](#), [6.3](#), [7.6](#)
- [22] ———, *The Conway function of a splice*, Proc. Edinb. Math. Soc. **48** (2005), no. 1, 61–73. [arXiv:math/0407130](#). [↑7.5](#), [7.6](#), [7](#)
- [23] D. Cimasoni and V. Florens, *Generalized Seifert surfaces and signatures of colored links*, Trans. Amer. Math. Soc. **360** (2008), 1223–1264. [arXiv:math/0505185](#). [↑6.2](#)
- [24] T. D. Cochran, *Geometric invariants of link cobordism*, Comment. Math. Helv. **60** (1985), 291–311. [EuDML](#). [↑6](#)
- [25] T. D. Cochran, K. E. Orr, and P. Teichner, *Knot concordance, Whitney towers and L^2 -signatures*, Ann. Math. **157** (2003), 433–519. [arXiv:math.GT/9908117](#). [↑1](#)
- [26] J. H. Conway, *An enumeration of knots and links, and some of their algebraic properties*, Computational Problems in Abstract Algebra (Proc. Conf., Oxford, 1967), Pergamon, Oxford, 1970, pp. 329–358. [A. Ranicki’s archive](#). [↑26](#)
- [27] D. Cooper, *Signatures of surfaces with applications to knot and link cobordism*, Ph.D. thesis, University of Warwick, 1982. [Author’s homepage](#). [↑29](#)

- [28] D. S. Coram and P. F. Duvall, *Homotopy equivalences and mapping torus projections*, *Fund. Math.* **109** (1980), 1–7. [EuDML](#). [↑18](#)
- [29] R. J. Daverman, *Decompositions of Manifolds*, Pure Appl. Math., vol. 124, Academic Press, Orlando, FL, 1986. [↑](#)
- [30] R. J. Daverman and G. A. Venema, *Embeddings in Manifolds*, Grad. Studies in Math., vol. 106, Amer. Math. Soc., Providence, RI, 2009. [Venema’s webpage](#). [↑2.2](#), [5](#)
- [31] A. L. Edmonds, *Systems of curves on a closed orientable surface*, *Enseign. Math.* **42** (1996), 311–339. [E-Periodica](#). [↑17.1](#), [17](#)
- [32] R. H. Fox, *Some problems in knot theory*, *Topology of 3-Manifolds and Related Topics (Proc. of The Univ. of Georgia Institute, 1961)*, Prentice-Hall, Inc., Englewood Cliffs, NJ, 1961, pp. 168–176. [↑1](#), [24](#)
- [33] C. H. Giffen, *Disciplining dunce hats in 4-manifolds* (1976). Unpublished typeset manuscript. [↑A.2](#)
- [34] ———, *F-isotopy and I-equivalence of links* (1976). Unpublished handwritten manuscript. [↑A.2](#)
- [35] D. S. Gillman, *Sequentially 1-ULC tori*, *Trans. Amer. Math. Soc.* **111** (1964), 449–456. [Journal](#). [↑1](#), [1](#)
- [36] M. Hall, *The Theory of Groups*, The Macmillan Co., New York, 1959. [↑4.1](#)
- [37] R. Hartley, *The Conway potential function for links*, *Comment. Math. Helv.* **58** (1983), no. 3, 365–378. [EuDML](#). [↑26](#), [6](#), [29](#), [6.2](#), [6.3](#), [7.1](#), [7.8](#)
- [38] S. Harvey and E. Tweedy, *Co-rank of weakly parafree 3-manifold groups*, *Groups Geom. Dyn.* **14** (2020), 25–39. [arXiv:1711.10659](#). [↑1](#)
- [39] A. Hatcher, *Algebraic Topology*, Cambridge Univ. Press, 2002. [Author’s webpage](#). [↑2.2](#), [49](#)
- [40] S. Hirose, *The action of the handlebody group on the first homology group of the surface*, *Kyungpook Math. J.* **46** (2006), 399–408. [Journal](#). [↑17](#)
- [41] R. Holmes and N. Smythe, *Algebraic invariants of isotopy of links*, *Amer. J. Math.* **88** (1966), 646–654. [↑1](#)
- [42] Xiang-Dong Hou, S. R. López-Permouth, and B. R. Parra-Avila, *Rational power series, sequential codes and periodicity of sequences*, *J. Pure Appl. Algebra* **213** (2009), 1157–1169. [↑45](#)
- [43] J. F. P. Hudson and E. C. Zeeman, *On combinatorial isotopy*, *Inst. Hautes Études Sci. Publ. Math.* **19** (1964), 69–94. [EuDML](#). [↑2.4](#)
- [44] W. Jaco, *Nonretractable cubes-with-holes*, *Michigan Math. J.* **18** (1971), 193–201. [ProjectEuclid](#). [↑1](#), [17](#)
- [45] ———, *Geometric realizations for free quotients*, *J. Austral. Math. Soc.* **14** (1972), 411–418. [Cambridge Core](#). [↑1](#)
- [46] W. Jaco and D. R. McMillan Jr., *Retracting three-manifolds onto finite graphs*, *Illinois J. Math.* **14** (1970), 150–158. [ProjectEuclid](#). [↑1](#), [1](#), [17](#), [17](#)
- [47] Boju Jiang, *On Conway’s potential function for colored links*, *Acta Math. Sin. (Engl. Ser.)* **32** (2016), no. 1, 25–39. [arXiv:1407.3081](#). [↑30](#)
- [48] A. Kawauchi, *A Survey of Knot Theory*, Birkhäuser, Basel, 1996. Translated and revised from the 1990 Japanese original by the author. [↑3](#), [10](#), [33](#)
- [49] L. H. Kauffman, *The Conway polynomial*, *Topology* **20** (1981), 101–108. [↑26](#)
- [50] M. E. Kidwell, *On the two-variable Conway potential function*, *Proc. Amer. Math. Soc.* **98** (1986), 485–494. [Journal](#). [↑](#)
- [51] P. Kirk, C. Livingston, and Z. Wang, *The Gassner representation for string links*, *Comm. Contemp. Math.* **3** (2001), 87–136. [arXiv:math/9806035](#). [↑6.2](#)
- [52] A. Kirkor, *In S^3 every compact ANR with the homotopy type of S^1 is a deformation retract of a neighborhood*, *Proc. Int. Conf. on Geometric Topology (Warsaw, 1978)* (K. Borsuk and A. Kirkor, eds.), PWN, Warsaw, 1980, pp. 241–243. [↑1](#)

- [53] S. Kojima and M. Yamasaki, *Some new invariants of links*, *Invent. Math.* **54** (1979), 213–228. [EuDML](#). [↑3, 47](#)
- [54] M. Kreck and W. Lück, *Topological rigidity for non-aspherical manifolds*, *Pure Appl. Math. Q.* **5** (2009), 873–914. [arXiv:math.GT/0509238](#). [↑18](#)
- [55] H. W. Lambert, *Mapping cubes with holes onto cubes with handles*, *Illinois J. Math.* **13** (1969), 606–615. [ProjectEuclid](#). [↑1](#)
- [56] J. Levine, *A factorization of the Conway polynomial*, *Comment. Math. Helv.* **74** (1999), 27–52. [arXiv:q-alg/9711007](#). [↑6.2](#)
- [57] W. Magnus, A. Karrass, and D. Solitar, *Combinatorial Group Theory: Presentations of Groups in Terms of Generators and Relations*, 2nd ed., Dover, New York, 1976. [↑2.2, 2](#)
- [58] Y. Matsumoto, *Wild embeddings of piecewise linear manifolds in codimension two*, *Geometric Topology (Proc. Georgia Topology Conf.)*, Academic Press, New York, 1979, pp. 393–428. [↑](#)
- [59] B. Mazur, *A note on some contractible 4-manifolds*, *Ann. of Math.* **73** (1961), 221–228. [↑22](#)
- [60] D. R. McMillan Jr., *A criterion for cellularity in a manifold. II*, *Trans. Amer. Math. Soc.* **126** (1967), 217–224. [Journal](#). [↑1, 7, A.2](#)
- [61] J. M. McPherson, *On the nullity and enclosure genus of wild knots*, *Trans. Amer. Math. Soc.* **144** (1969), 545–555. [Journal](#). [↑2, 1.6](#)
- [62] W. H. Meeks III and J. Patrusky, *Representing homology classes by embedded circles on a compact surface*, *Illinois J. Math.* **22** (1978), 262–269. [ProjectEuclid](#). [↑17.1](#)
- [63] S. A. Melikhov and D. Repovš, *n-Quasi-isotopy. I. Questions of nilpotence*, *J. Knot Theory Ram.* **14** (2005), 571–602. [arXiv:math/0103113](#). [↑1, 2.7, 22](#)
- [64] S. A. Melikhov, *Steenrod homotopy*, *Russ. Math. Surv.* **64** (2009), 469–551; Russian transl., *Uspekhi Mat. Nauk* **64:3** (2009), 73–166. [arXiv:0812.1407](#). [↑3](#)
- [65] ———, *Topological isotopy and Cochran’s derived invariants*, *Topology, Geometry, and Dynamics — V. A. Rokhlin Memorial*, *Contemp. Math.*, vol. 772, Amer. Math. Soc., Providence, RI, 2021, pp. 249–266. [arXiv:2011.01409](#). [↑1.1, 1, 3, 17, 14, 14, 15, A](#)
- [66] S. A. Melikhov, *Topology of Metric Spaces* (18.11.2022). [ResearchGate](#). [↑3, A.6](#)
- [67] ———, *Topological isotopy and finite type invariants*. [arXiv:?](#). [↑1.2, 1, 10](#)
- [68] ———, *The two-variable Conway polynomial and Cochran’s derived invariants*. [arXiv:?](#). [↑1, 1, 3, 4, 6, 8, 12, 13, 13, 15, 15, 16](#)
- [69] M. D. Meyerson, *Representing homology classes of closed orientable surfaces*, *Proc. Amer. Math. Soc.* **61** (1976), 181–182 (1977). [Journal](#). [↑17.1](#)
- [70] J. Milnor, *Isotopy of links*, *Algebraic Geometry and Topology: A Symposium in Honor of S. Lefschetz*, Princeton Univ. Press, Princeton, NJ, 1957, pp. 280–306. [↑1](#)
- [71] J. W. Milnor, *Topology from the Differentiable Viewpoint*, Univ. Press of Virginia, Charlottesville, VA, 1965. [A. Ranicki’s web archive](#). [↑17](#)
- [72] Y. Miyazawa, *Conway polynomials of periodic links*, *Osaka J. Math.* **31** (1994), 147–163. [↑1, 7.11](#)
- [73] J. Murakami, *On local relations to determine the multi-variable Alexander polynomial of colored links*, *Knots 90* (Osaka, 1990), de Gruyter, Berlin, 1992, pp. 455–464. [↑30](#)
- [74] K. Murasugi, *On periodic knots*, *Comment. Math. Helv.* **46** (1971), 162–174. [EuDML](#). [↑7.13, 34](#)
- [75] Y. Nakanishi, *Three-variable Conway potential function of links*, *Tokyo J. Math.* **13** (1990), 163–177. [↑6, 30](#)
- [76] M. Ozawa and K. Shimokawa, *Dehn surgery and Seifert surface systems*, *Ann. Sc. Norm. Super. Pisa Cl. Sci.* **17** (2017), no. 1, 267–276. [arXiv:1406.6264](#). [↑1](#)
- [77] A. Putman, *Lectures on the Torelli group*, 2014. [Author’s webpage](#). [↑17.1](#)
- [78] D. Rolfsen, *Some counterexamples in link theory*, *Canadian J. Math.* **26** (1974), 978–984. [Journal](#). [↑1, 2.2](#)
- [79] ———, *Localized Alexander invariants and isotopy of links*, *Ann. Math.* **101** (1975), 1–19. [↑](#)

- [80] ———, *Knots and Links*, Math. Lecture Series, vol. 7, Publish or Perish, Inc., Berkeley, CA, 1976. [↑4.4](#)
- [81] ———, *Piecewise-linear I-equivalence of links*, Low-Dimensional Topology (Chelwood Gate, 1982), London Math. Soc. Lecture Note Ser., vol. 95, Cambridge Univ. Press, Cambridge, 1985, pp. 161–178. [↑1](#)
- [82] ———, *PL link isotopy, essential knotting and quotients of polynomials*, Canad. Math. Bull. **34** (1991), 536–541. [Journal](#). [↑](#)
- [83] N. Sato, *Cobordisms of semiboundary links*, Topology Appl. **18** (1984), 225–234. [Journal](#). [↑17](#)
- [84] J. A. Schafer, *Representing homology classes on surfaces*, Canad. Math. Bull. **19** (1976), 373–374. [Cambridge Core](#). [↑17.1](#)
- [85] J. Schultens, *Introduction to 3-Manifolds*, Grad. Studies in Math., vol. 151, Amer. Math. Soc., Providence, RI, 2014. [↑17](#)
- [86] M. A. Štan'ko, *Imbedding of compacta in euclidean space*, Mat. Sb. **83** (1970), 234–255 (Russian); English transl., Math. USSR-Sb. **12** (1970), 234–254. [MathNet](#). [↑7](#)
- [87] ———, *Imbedding of compacta in Euclidean space of dimension 5, 4 and 3*, Mat. Sb. **99** (1976), 626–633 (Russian); English transl., Math. USSR-Sb. **28** (1976), no. 4, 563–569. [MathNet](#). [↑7](#), [A.2](#)
- [88] D. S. Silver, W. Whitten, and S. G. Williams, *Knot groups with many killers*, Bull. Aust. Math. Soc. **81** (2010), 507–513. [arXiv:0909.3275](#). [↑2](#)
- [89] N. Smythe, *Boundary links*, Topology Seminar (Wisconsin, 1965) (R. H. Bing, ed.), Annals of Math. Studies, vol. 60, Princeton Univ. Press, 1966, pp. 69–72. [A. Ranicki's archive](#). [↑1](#), [1](#), [17](#)
- [90] S. Suzuki, *Alexander ideals of graphs in the 3-sphere*, Tokyo J. Math. **7** (1984). [ProjectEuclid](#). [↑1](#)
- [91] L. Traldi, *Conway's potential function and its Taylor series*, Kobe J. Math. **5** (1988), 233–263. [Lafayette Repository](#). [↑6.2](#)
- [92] Chichen M. Tsau, *Nonalgebraic killers of knot groups*, Proc. Amer. Math. Soc. **95** (1985), 139–146. [Journal](#). [↑2](#)
- [93] T. Tsukamoto and A. Yasuhara, *A factorization of the Conway polynomial and covering linkage invariants*, J. Knot Theory Ram. **16** (2007), 631–640. [arXiv:math/0405481](#). [↑6.2](#)
- [94] V. G. Turaev, *Reidemeister torsion in knot theory*, Uspekhi Mat. Nauk **41** (1986), no. 1, 97–147, 240, [MathNet](#); English transl., Russ. Math. Surveys **41** (1986), 119–182. [↑30](#), [6.2](#), [7](#)
- [95] A. H. Wright, *Boundary-preserving mappings of 3-manifolds onto cubes-with-handles*, Illinois J. Math. **20** (1976), 98–106. [ProjectEuclid](#). [↑17](#)
- [96] H. J. Zassenhaus, *The Theory of Groups*, 2nd ed., Chelsea, New York, 1958. [↑6.2](#)
- [97] A. Zastrow, *An example of a non-extendable isotopy of a knot* (2024), preprint. [↑1](#), [2.4](#)
- [98] ———, *An Embedded Circle into \mathbb{R}^3 Might Not Be Able to Escape Before an Isotoped Linked Circle* (June 6, 2022). Talk at [The 39th Annual \(Online\) Workshop in Geometric Topology](#). Video: https://www.youtube.com/watch?v=Zt_C_0vw4PQ. [↑1](#)

STEKLOV MATHEMATICAL INSTITUTE OF RUSSIAN ACADEMY OF SCIENCES, MOSCOW, RUSSIA
 Email address: melikhov@mi-ras.ru