

# VISIBILITY DOMAINS RELATIVE TO THE KOBAYASHI DISTANCE IN COMPLEX MANIFOLDS

RUMPA MASANTA

ABSTRACT. In this paper, we extend the notion of visibility relative to the Kobayashi distance to domains in arbitrary complex manifolds. Visibility here refers to a property resembling visibility in the sense of Eberlein–O’Neill for Riemannian manifolds. Since it is difficult, in general, to determine whether domains are Cauchy-complete with respect to the Kobayashi distance, we do not assume so here. We provide many sufficient conditions for visibility. We establish a Wolff–Denjoy-type theorem in a very general setting as an application. We also explore some connections between visibility and Gromov hyperbolicity for Kobayashi hyperbolic domains in the above setting.

## 1. INTRODUCTION AND STATEMENT OF RESULTS

In the past few years, Bharali–Zimmer introduced a property that may be viewed as a weak notion of negative curvature for domains  $\Omega \subsetneq \mathbb{C}^n$  viewed as metric spaces equipped with the Kobayashi distance  $K_\Omega$ —first for bounded domains [8] and later for unbounded domains [9]. This property of  $\Omega \subsetneq \mathbb{C}^n$  has been used to understand extensions and the behaviour of certain holomorphic mappings into  $\Omega$ , and also to study properties of  $K_\Omega$ : see, for instance, [8, 7, 13, 16, 9, 28, 26]. In [8, 9], the property referenced above was called the visibility property with respect to  $K_\Omega$  because it is reminiscent of visibility in the sense of Eberlein–O’Neill [19]. Very roughly, the visibility property is that all geodesics originating sufficiently close to and terminating sufficiently close to two distinct points in  $\partial\Omega$  must bend uniformly into  $\Omega$ .

Motivated by the usefulness of the visibility property, we extend this notion to Kobayashi hyperbolic domains  $\Omega \subsetneq X$  (i.e., domains on which the Kobayashi pseudodistance  $K_\Omega$  is a distance) for arbitrary complex manifolds  $X$ , and explore some of its consequences.

Given a Kobayashi hyperbolic domain  $\Omega$  in a complex manifold  $X$ , if  $\dim_{\mathbb{C}}(X) \geq 2$ , it is in general very difficult to determine whether the metric space  $(\Omega, K_\Omega)$  is Cauchy-complete—even when  $\Omega \subsetneq \mathbb{C}^n$ —and, so, whether  $(\Omega, K_\Omega)$  is a geodesic space. Thus, the formal definition of the visibility property requires some care. In the process, we get two closely related notions of visibility—the second notion being termed *weak visibility with respect to  $K_\Omega$* . We will formally define both these notions below. Functionally, these two notions were introduced to study continuous extensions to  $\overline{D}$  of holomorphic mappings between domains  $f : D \rightarrow \Omega$  when  $\Omega$  has the (weak) visibility property. To quote Bharali–Zimmer [9], for  $\Omega$  as above:

- weak visibility with respect to  $K_\Omega$  is the property that allows one to deduce continuous extension for isometries with respect to  $K_D$  and  $K_\Omega$ .
- visibility with respect to  $K_\Omega$  is the property that allows one to deduce continuous extension for continuous **quasi**-isometries with respect to  $K_D$  and  $K_\Omega$ .

That being said, in this paper we shall not discuss results on continuous extension since the proofs involving (target spaces that are) domains in  $\mathbb{C}^n$  extend verbatim to the more general setting of this paper. The results we shall present are (with perhaps one exception) such that their proofs are **not** routine extensions of the arguments used for domains in  $\mathbb{C}^n$ .

---

2020 *Mathematics Subject Classification*. Primary: 32F45, 32Q45, 53C23 ; Secondary: 32H50, 32Q05.  
*Key words and phrases*. Complex manifolds, hyperbolic imbedding, Kobayashi metric, visibility.

**1.1. General theorems on visibility.** We first formally define the notion of visibility discussed above. Before we do so, recall the discussion above on the difficulty of knowing whether  $(\Omega, K_\Omega)$  is Cauchy-complete (for brevity:  $\Omega$  is *Kobayashi complete*) and, consequently, whether any pair of distinct points admits a geodesic joining them. This difficulty is overcome by considering  $(\lambda, \kappa)$ -almost-geodesics (where  $\lambda \geq 1$  and  $\kappa \geq 0$ ), which will serve as substitutes for geodesics. A  $(\lambda, \kappa)$ -almost-geodesic is, essentially, a kind of “nice” quasi-geodesic: we refer the reader to Section 2.1 for the precise definition. The relevance of these curves is as follows: given any  $\lambda \geq 1$  and  $\kappa > 0$ , for any two distinct points  $x, y \in \Omega$ , there exists a  $(\lambda, \kappa)$ -almost-geodesic joining them. This is a consequence of Proposition 2.8. We can now give the following

**Definition 1.1.** Let  $X$  be a complex manifold and  $\Omega \subsetneq X$  be a Kobayashi hyperbolic domain.

- (a) Let  $p, q \in \partial\Omega$ ,  $p \neq q$ . We say that the pair  $(p, q)$  satisfies the *visibility condition with respect to  $K_\Omega$*  if there exist neighbourhoods  $U_p$  of  $p$  and  $U_q$  of  $q$  in  $X$  such that  $\overline{U_p} \cap \overline{U_q} = \emptyset$  and such that for each  $\lambda \geq 1$  and each  $\kappa \geq 0$ , there exists a compact set  $K \subsetneq \Omega$  such that the image of each  $(\lambda, \kappa)$ -almost-geodesic  $\sigma : [0, T] \rightarrow \Omega$  with  $\sigma(0) \in U_p$  and  $\sigma(T) \in U_q$  intersects  $K$ .
- (b) We say that  $\partial\Omega$  is *visible* if every pair of points  $p, q \in \partial\Omega$ ,  $p \neq q$ , satisfies the visibility condition with respect to  $K_\Omega$ .
- (c) Let  $p, q \in \partial\Omega$ ,  $p \neq q$ . We say that the pair  $(p, q)$  satisfies the *weak visibility condition with respect to  $K_\Omega$*  if the condition in (a) is satisfied **only** for  $\lambda = 1$ . We say that  $\partial\Omega$  is *weakly visible* if every pair of points  $p, q \in \partial\Omega$ ,  $p \neq q$ , satisfies the weak visibility condition with respect to  $K_\Omega$ .

We first explore the question whether, with  $\Omega$  as at the top of this paper,  $\partial\Omega$  being locally visible is the same as  $\partial\Omega$  being visible. Firstly, we clarify what “ $\partial\Omega$  being locally visible” means.

**Definition 1.2.** Let  $X$  be a complex manifold and  $\Omega \subsetneq X$  be a domain.

- (a) Let  $p \in \partial\Omega$ . We say that  $\partial\Omega$  is *locally visible at  $p$*  if there exists a neighbourhood  $U_p$  of  $p$  in  $X$  such that  $U_p \cap \Omega$  is a Kobayashi hyperbolic domain and there exists a neighbourhood  $V_p$  of  $p$  in  $X$ ,  $V_p \subseteq U_p$ , such that every pair of distinct points  $q_1, q_2 \in V_p \cap \partial\Omega$  satisfies the visibility condition with respect to  $K_{U_p \cap \Omega}$ .
- (b) We say that  $\partial\Omega$  is *locally visible* if  $\partial\Omega$  is locally visible at each boundary point  $p \in \partial\Omega$ .
- (c) Let  $p \in \partial\Omega$ . We say that  $\partial\Omega$  is *locally weakly visible at  $p$*  if the condition in (a) is satisfied **only** for  $\lambda = 1$ . We say that  $\partial\Omega$  is *locally weakly visible* if  $\partial\Omega$  is locally weakly visible at each boundary point  $p \in \partial\Omega$ .

The above question was first explored by Bracci *et al.* in [11] in the context of bounded Kobayashi complete domains in  $\mathbb{C}^n$ . We extend their analysis to domains in arbitrary complex manifolds that are not necessarily relatively compact and not necessarily Kobayashi complete. Our result, however, is **not** a routine extension of the analysis in [11]. To elaborate upon this point, let us first state a definition.

**Definition 1.3.** Let  $X$  be a complex manifold and let  $Z$  be a connected complex submanifold in  $X$ .  $Z$  is said to be *hyperbolically imbedded in  $X$*  if for each  $x \in \overline{Z}$  and for each neighbourhood  $V_x$  of  $x$  in  $X$ , there exists a neighbourhood  $W_x$  of  $x$  in  $X$ ,  $\overline{W_x} \subset V_x$ , such that

$$K_Z(\overline{W_x} \cap Z, Z \setminus V_x) > 0.$$

*Remark 1.4.* Clearly,  $Z$  is hyperbolically imbedded in  $X$  if and only if, for  $x, y \in \overline{Z}$  and  $x \neq y$ , there exist neighbourhoods  $V_x$  of  $x$  and  $V_y$  of  $y$  in  $X$  such that  $K_Z(V_x \cap Z, V_y \cap Z) > 0$ .

The notion of  $\partial\Omega$  being locally weakly visible at  $p \in \partial\Omega$  is the closer of the two notions in Definition 1.2 to local visibility at  $p \in \partial\Omega$  as considered in [11]. For either notion, the extension

of the arguments in [11] to our setting faces several obstacles. E.g., with  $X$  as above, it is not true in general that one can fix a Hermitian metric  $h$  on  $X$  such that the open balls given by the associated distance  $d_h$  are Kobayashi hyperbolic. If we assume that  $\partial\Omega$  is locally visible (resp., locally weakly visible), then, by augmenting this by the assumption that  $\Omega$  is hyperbolically imbedded, we are able to deal with the obstacles mentioned and deduce that  $\partial\Omega$  is visible (resp., weakly visible). But the additional assumption proves not to be unnatural because the converses of the latter statements hold true! This summarises the following two theorems.

**Theorem 1.5.** *Let  $X$  be a complex manifold and let  $\Omega \subsetneq X$  be a Kobayashi hyperbolic domain. Then, the following conditions are equivalent:*

- (a)  $\partial\Omega$  is locally visible and  $\Omega$  is a hyperbolically imbedded domain,
- (b)  $\partial\Omega$  is visible.

**Theorem 1.6.** *Let  $X$  be a complex manifold and let  $\Omega \subsetneq X$  be a Kobayashi hyperbolic domain. Then, the following conditions are equivalent:*

- (a)  $\partial\Omega$  is locally weakly visible and  $\Omega$  is a hyperbolically imbedded domain,
- (b)  $\partial\Omega$  is weakly visible.

The next concept that we will examine in this paper is introduced by the following definition.

**Definition 1.7.** Let  $X$  be a complex manifold and  $\Omega \subsetneq X$  be a Kobayashi hyperbolic domain. Let  $p \in \partial\Omega$ . We say that  $p$  is a *visible point* of  $\partial\Omega$  (resp.,  $p$  is a *weakly visible point* of  $\partial\Omega$ ) if there exists a neighbourhood  $U_p$  of  $p$  in  $X$  such that every pair of distinct points  $q_1, q_2 \in U_p \cap \partial\Omega$  satisfies the visibility condition (resp., the weak visibility condition) with respect to  $K_\Omega$ . Let  $\partial_{\text{visi}}\Omega$  denote the set of all visible points of  $\partial\Omega$  and  $\partial_{\text{wvisi}}\Omega$  denote the set of all weakly visible points of  $\partial\Omega$ .

The pair of sets defined above are important, but technical, objects that act along with more concrete notions to give us various sufficient conditions such that, given a domain  $\Omega \subsetneq X$ ,  $X$  and  $\Omega$  as in the above definition,  $\partial\Omega$  is visible (resp., weakly visible). The principal results that rely on Definition 1.7 are stated later in this section or in Section 4 and which will be proved in Sections 4 and 5. An easy consequence of the results in Section 4 is as follows.

**Proposition 1.8.** *Let  $X$  be a complex manifold and let  $\Omega \subsetneq X$  be a Kobayashi hyperbolic domain. Each of the sets  $\partial\Omega \setminus \partial_{\text{visi}}\Omega$  and  $\partial\Omega \setminus \partial_{\text{wvisi}}\Omega$  is either not totally disconnected or is empty.*

**1.2. Sufficient conditions for (weak) visibility.** Next, we show that domains  $\Omega \subsetneq X$  such that  $\partial\Omega$  is (weakly) visible are abundant.

**Definition 1.9.** Let  $X$  be a complex manifold and  $\Omega \subsetneq X$  be a Kobayashi hyperbolic domain. Let  $p \in \partial\Omega$ . We say that  $\Omega$  is *visible around  $p$*  (resp.,  $\Omega$  is *weakly visible around  $p$* ) if there exists a holomorphic chart  $(U, \varphi)$  of  $X$  centered at  $p$  such that  $\varphi(U)$  is bounded and  $\varphi(U \cap \Omega) \subsetneq \mathbb{C}^n$  is a domain, and such that  $\partial(\varphi(U \cap \Omega))$  is visible (resp., weakly visible). Let  $\partial_V\Omega$  denote the set of all points at which  $\Omega$  is visible around  $p$  and  $\partial_{WV}\Omega$  denote the set of all points at which  $\Omega$  is weakly visible around  $p$ .

Superficially, Definition 1.9 seems to be no different from Definition 1.2. However, the formulation of the former definition in terms of holomorphic charts allows us to construct large families of domains  $\Omega \subsetneq X$ ,  $X$  different from  $\mathbb{C}^n$ , having the visibility property by appealing to the many examples one knows in  $\mathbb{C}^n$ . This point will be clearer after we state our next theorem.

**Theorem 1.10.** *Let  $X$  be any one of the following:*

- (a) A Kobayashi hyperbolic complex manifold,
- (b) A Stein manifold,
- (c) A holomorphic fiber bundle  $\pi : X \rightarrow Y$  such that  $Y$  is either of (a) or (b) and the fibers are Kobayashi hyperbolic,
- (d) A holomorphic covering space  $\pi : X \rightarrow Y$  such that  $Y$  is either of (a) or (b).

Let  $\Omega \subsetneq X$  be a relatively compact domain.

- (i) Suppose the set  $\partial\Omega \setminus \partial_V\Omega$  is totally disconnected. Then,  $\partial\Omega$  is visible.
- (ii) Suppose the set  $\partial\Omega \setminus \partial_{WV}\Omega$  is totally disconnected. Then,  $\partial\Omega$  is weakly visible.

The above result can be interpreted as follows: if, around “most” points  $p \in \partial\Omega$ ,  $\Omega$  (locally) looks like a visibility domain in  $\mathbb{C}^{\dim(X)}$  —and, as stated above, we have **many** ways of identifying visibility in  $\mathbb{C}^n$ , thanks to Bharali–Zimmer [8, 9], Bharali–Maitra [7], Bracci *et al.* [13] —then  $\partial\Omega$  is visible.

Now, we will present a result that is a method for obtaining new domains with visible boundaries from old domains with the former property.

**Theorem 1.11.** *Let  $X$  be a complex manifold of dimension  $n$  and  $\Omega \subsetneq X$  be a Kobayashi hyperbolic domain. Let  $\partial\Omega$  be visible. Suppose  $S \subsetneq X$  is a closed, totally disconnected set such that  $\Omega \setminus S$  is connected, and such that  $\mathcal{H}^{2n-2}(S) = 0$ . Then,  $\partial(\Omega \setminus S)$  is visible.*

Here,  $\mathcal{H}^\alpha$  denotes the  $\alpha$ -dimensional Hausdorff measure for  $\alpha \geq 0$ .

To state our next result, we need a definition.

**Definition 1.12.** Let  $\Omega$  be a domain in  $\mathbb{C}$ . Let  $\widehat{\mathbb{C}}$  be the one-point compactification of  $\mathbb{C}$  and  $\partial_\infty\Omega$  be the boundary of  $\Omega$  viewed as a domain in  $\widehat{\mathbb{C}}$ .

- (a) We say that  $\partial_\infty\Omega$  is a *Jordan curve* if there exists a homeomorphism between  $S^1$  and  $\partial_\infty\Omega$ .
- (b) We say that  $\Omega$  is a *Jordan domain* if  $\partial_\infty\Omega$  is a Jordan curve.

We now present a new class of planar domains with the visibility property. To the best of my knowledge, there is no version of the following result in the literature for **unbounded** domains.

**Proposition 1.13.** *Let  $\Omega$  be a simply connected domain in  $\mathbb{C}$  such that  $\Omega$  is a Jordan domain. Then,  $\partial\Omega$  is visible.*

Experts will be familiar with the notion of “local Goldilocks points” introduced in [9] providing a sufficient condition for visibility for domains in  $\mathbb{C}^n$ . Is there an analogue of this in the more general setting of this paper? The answer to this is “Yes,” but since formulating this notion in a coordinate-independent way requires some work, we defer this discussion to Section 5.

**1.3. Compactifications, visibility domains, and a Wolff–Denjoy-type theorem.** The main notions studied in this paper —i.e., visibility and weak visibility of  $\partial\Omega$ —formally differ from the notion alluded to at the top of this section and which motivates this work: that of (weak) visibility domains with respect to  $K_\Omega$  for domains  $\Omega \subsetneq \mathbb{C}^n$ . We refer readers to [7, Definition 1.1] for the definition of the latter notion when  $\Omega$  is bounded and to [9, Definition 1.1] when  $\Omega$  is unbounded. The question arises: *are the two notions equivalent?* When  $\Omega \subsetneq \mathbb{C}^n$  is bounded, it is elementary to see that  $\partial\Omega$  is visible (resp., weakly visible) in the sense of Definition 1.1 if and only if  $\Omega$  is a visibility domain (resp., a weak visibility domain) in the sense of [7, 9]. However, for  $\Omega \subsetneq \mathbb{C}^n$  unbounded, the definition of a (weak) visibility domain involves the end compactification of  $\overline{\Omega}$ , and its relationship with Definition 1.1 is *a priori* unclear. The reasons that the end compactification of  $\overline{\Omega}$  forms a part of [9, Definition 1.1] are:

- (a) For many of the applications of visibility shown in the works cited above, an appropriate compactification of  $\Omega$  is needed, when  $\Omega$  is unbounded, for proofs to be valid — which is an issue we too must address in this paper.
- (b) The end compactification of  $\overline{\Omega}$  (see [9, Section 4] for a definition) suggests itself as a natural choice of a compactification because it arises naturally in results involving the Kobayashi distance; see, for instance, [12].

In the more general setting of this paper, we formalise the need suggested in (a) above as follows:

**Definition 1.14.** Let  $X$  be a complex manifold and let  $\Omega \subsetneq X$  be a Kobayashi hyperbolic domain. We say that  $\overline{\Omega}^\infty$  is an *admissible compactification of  $\overline{\Omega}$*  if  $\overline{\Omega}^\infty$  is a Hausdorff, sequentially compact topological space that admits a map  $\mathbf{j} : \overline{\Omega} \rightarrow \overline{\Omega}^\infty$  that is a homeomorphism onto its image, such that  $\mathbf{j}(\overline{\Omega})$  is open and dense in  $\overline{\Omega}^\infty$ , and such that  $\overline{\Omega}^\infty \setminus \mathbf{j}(\overline{\Omega})$  is totally disconnected.

In the present setting, no preferred compactification suggests itself. Hence the need for the above definition. In fact, for  $\Omega \subsetneq \mathbb{C}$ , it turns out to be more natural to take  $\overline{\Omega}^\infty$  to be the closure of  $\Omega$  in  $\widehat{\mathbb{C}}$ : the one-point compactification of  $\mathbb{C}$ . This will be seen in the results of a forthcoming work (and is hinted at by the proof of Proposition 1.13).

We can now propose a definition for a visibility domain. In the definition below, we will commit the following abuse of notation: for any point  $x \in \overline{\Omega}$ , we shall write  $\mathbf{j}(x)$  simply as  $x$ .

**Definition 1.15.** Let  $X$  be a complex manifold and  $\Omega \subsetneq X$  be a domain (not necessarily relatively compact) and let  $\overline{\Omega}^\infty$  be some admissible compactification of  $\overline{\Omega}$ . We say that  $\Omega$  is a *visibility domain subordinate to  $\overline{\Omega}^\infty$*  if  $\Omega$  is Kobayashi hyperbolic and has the following property:

- (\*) Given  $\lambda \geq 1$ ,  $\kappa \geq 0$ , distinct points  $p, q \in \partial_\infty \Omega$ , and  $\overline{\Omega}^\infty$ -open neighbourhoods  $V_p$  of  $p$  and  $V_q$  of  $q$  whose closures (in  $\overline{\Omega}^\infty$ ) are disjoint, there exists a compact set  $K \subsetneq \Omega$  such that the image of each  $(\lambda, \kappa)$ -almost-geodesic  $\sigma : [0, T] \rightarrow \Omega$  with  $\sigma(0) \in \mathbf{j}^{-1}(V_p \setminus \partial_\infty \Omega)$  and  $\sigma(T) \in \mathbf{j}^{-1}(V_q \setminus \partial_\infty \Omega)$  intersects  $K$ .

We say that  $\Omega$  is a *weak visibility domain subordinate to  $\overline{\Omega}^\infty$*  if the property (\*) holds true **only** for  $\lambda = 1$ . Here,  $\mathbf{j}$  is the embedding of  $\overline{\Omega} \hookrightarrow \overline{\Omega}^\infty$  that defines  $\overline{\Omega}^\infty$  and  $\partial_\infty \Omega := \overline{\Omega}^\infty \setminus \mathbf{j}(\overline{\Omega})$ .

The question raised above is answered as follows:

**Theorem 1.16** (see Corollaries 4.2, 4.3). *Let  $X$  be a complex manifold and  $\Omega \subsetneq X$  a Kobayashi hyperbolic domain. Then,  $\partial\Omega$  is visible (resp., weakly visible) if and only if  $\Omega$  is a visibility domain (resp., weak visibility domain) subordinate to any admissible compactification of  $\overline{\Omega}$ .*

We will not overtly focus on the property in Definition 1.15. In this paper, the latter property:

- is used to give a unified discussion — irrespective of whether or not  $\Omega \subsetneq X$  is relatively compact — of the connection between visibility of  $\partial\Omega$  and the local Goldilocks property that was alluded to above, and
- plays an absolutely vital role in establishing a Wolff–Denjoy-type theorem in the very general setting of this paper.

One of the goals of this paper is to show how ubiquitous the Wolff–Denjoy phenomenon is. The Wolff–Denjoy theorem (see Result 6.1 below) has been generalised in many ways in the literature but, till recently, these have featured convex domains. Since the history of generalisations of the Wolff–Denjoy theorem is extensive, and their contrast with our Wolff–Denjoy-type theorem (see Theorem 6.2) is significant, we shall introduce and discuss it in Section 6.

**1.4. Connections with Gromov hyperbolicity.** We will assume that the reader is familiar with Gromov hyperbolic metric spaces. Now, length metric spaces that are not geodesic spaces (and hence, by the Hopf–Rinow Theorem, are not Cauchy-complete) are less frequently

encountered in geometry. But, as we have emphasised above, we do not assume in this paper that  $(\Omega, K_\Omega)$  — for  $\Omega$  as above — is Cauchy-complete. The definition of Gromov hyperbolicity of  $(\Omega, K_\Omega)$  in the latter setting might be useful. We refer the reader to the definition in Section 2.3. Every Gromov hyperbolic metric space  $(X, d)$  has an abstract boundary called the *Gromov boundary*, denoted by  $\partial_G X$ . We define *Gromov bordification* to be the set  $\overline{X}^G := X \cup \partial_G X$  equipped with a topology with respect to which the inclusion  $X \hookrightarrow \overline{X}^G$  is a homeomorphism. This is called the *Gromov topology*, which we will discuss in Section 2.3. Here, we introduce the notation “ $\xrightarrow{G} \xi$ ” to denote the approach to  $\xi \in \partial_G X$  with respect to the Gromov topology.

When a length metric space  $(X, d)$  is Cauchy-complete and Gromov hyperbolic, then it is well known that  $\partial_G X$  is visible to geodesic lines (see [14, Part III, Lemma 3.2]). This property of  $\partial_G X$  is key to a range of applications of Gromov hyperbolicity. We now specialise to the metric space  $(\Omega, K_\Omega)$ ,  $\Omega$  is as in the theorems above. This, in view of Proposition 2.8, raises the natural question: *when  $(\Omega, K_\Omega)$  is **not** Cauchy-complete, then is  $\partial_G \Omega$  visible to almost-geodesics?* An answer to this question would, among other things, give us necessary conditions for when  $\overline{\Omega}$  embeds into  $\overline{\Omega}^G$  (which would open up many applications in the present setting). Firstly, we must clarify what is meant by the above question. To this end, we need a definition.

**Definition 1.17.** Let  $X$  be a complex manifold and  $\Omega \subsetneq X$  be a Kobayashi hyperbolic domain (that is not necessarily Kobayashi complete) such that  $(\Omega, K_\Omega)$  is Gromov hyperbolic. We say that  $\partial_G \Omega$  is *visible to almost-geodesics* if for any  $\xi, \eta \in \partial_G \Omega$ ,  $\xi \neq \eta$ , given any  $\kappa \geq 0$  and sequences  $(x_\nu)_{\nu \geq 1} \subset \Omega$  and  $(y_\nu)_{\nu \geq 1} \subset \Omega$  such that  $x_\nu \xrightarrow{G} \xi$ , and  $y_\nu \xrightarrow{G} \eta$ , there exists a compact set  $K \subsetneq \Omega$  such that for each  $\nu$ , the image of each  $(1, \kappa)$ -almost-geodesic joining  $x_\nu$  and  $y_\nu$  intersects  $K$ .

It turns out that the answer to the above question is in the affirmative in many cases.

**Theorem 1.18.** *Let  $X$  be a complex manifold and  $\Omega \subsetneq X$  be a Kobayashi hyperbolic domain such that  $(\Omega, K_\Omega)$  is Gromov hyperbolic. Suppose  $\overline{\Omega}^G$  is locally compact. Then,  $\partial_G \Omega$  is visible to almost-geodesics.*

The question now arises: *how natural is the assumption in Theorem 1.18 that  $\overline{\Omega}^G$  be locally compact when  $(\Omega, K_\Omega)$  is incomplete?* The final result of this section gives a construction of a large collection of domains that satisfy the conditions in Theorem 1.18.

**Proposition 1.19.** *Let  $D \subsetneq \mathbb{C}^n$  be a Kobayashi hyperbolic domain such that*

- (a)  $(D, K_D)$  is Gromov hyperbolic,
- (b)  $id_D$  extends to a homeomorphism between  $\overline{D}$  and  $\overline{D}^G$ .

*Let  $K \subsetneq D$  be a compact set such that  $\mathcal{H}^{2n-2}(K) = 0$ . Write  $\Omega := D \setminus K$ . Then,*

- (i)  $\Omega$  is Kobayashi hyperbolic and  $(\Omega, K_\Omega)$  is Gromov hyperbolic,
- (ii)  $\overline{\Omega}^G$  is homeomorphic to  $\overline{D} \setminus K$ .

*In particular  $\overline{\Omega}^G$  is non-compact and locally compact.*

The conditions (a) and (b) are observed rather frequently in this field. For instance, these conditions are satisfied by bounded strongly pseudoconvex domains [5] and by  $C^\infty$ -smoothly bounded convex domains of finite type [31].

## 2. METRICAL PRELIMINARIES AND ALMOST-GEODESICS

In this section, we elaborate upon certain metrical statements made in Section 1 for which the associated definitions and explanations had been deferred.

2.1.  **$(\lambda, \kappa)$ -almost-geodesics.** We present a couple of definitions.

**Definition 2.1.** Let  $X$  be a complex manifold of dimension  $n$ . A path  $\sigma : I \rightarrow X$ , where  $I \subseteq \mathbb{R}$  is an interval, is said to be *absolutely continuous* if for each  $t_0 \in I$  and each holomorphic chart  $(U, \varphi)$  around  $\sigma(t_0)$ , with  $I(\varphi, t_0)$  denoting any closed interval of  $\sigma^{-1}(U)$  containing  $t_0$ ,  $\varphi \circ \sigma|_{I(\varphi, t_0)}$  is absolutely continuous as a path in  $\mathbb{R}^{2n}$ .

*Remark 2.2.* From Definition 2.1 and the fact that every absolutely continuous path in  $\mathbb{R}^{2n}$  is almost-everywhere differentiable, it follows that the same is true in case of the paths defined in Definition 2.1. If  $\sigma$  is such a path and  $t \in I$  is such that  $D\sigma(t)$  exists, there is a canonical identification of the vector  $D\sigma(t)1 \in T_{\sigma(t)}X$  with a vector in  $T_{\sigma(t)}^{(1,0)}X$ . In this paper, we shall denote the latter by  $\sigma'(t)$ .

**Definition 2.3.** Let  $X$  be a complex manifold of dimension  $n$  and let  $\Omega \subseteq X$  be a Kobayashi hyperbolic domain. Let  $I \subseteq \mathbb{R}$  be an interval. For  $\lambda \geq 1$  and  $\kappa \geq 0$ , a curve  $\sigma : I \rightarrow \Omega$  is said to be a  $(\lambda, \kappa)$ -almost-geodesic if

(a) for all  $s, t \in I$

$$\frac{1}{\lambda}|s - t| - \kappa \leq K_{\Omega}(\sigma(s), \sigma(t)) \leq \lambda|s - t| + \kappa,$$

(b)  $\sigma$  is absolutely continuous (whence  $\sigma'(t)$  exists for almost every  $t \in I$ ), and for almost every  $t \in I$ ,  $k_{\Omega}(\sigma(t); \sigma'(t)) \leq \lambda$  (here  $k_{\Omega}$  denotes the Kobayashi pseudometric on  $\Omega$ ).

2.2. **General properties of Kobayashi hyperbolic domains.** With  $X$  and  $\Omega$  as above and  $\sigma : [a, b] \rightarrow \Omega$  an absolutely continuous path in  $\Omega$ , we introduce *the length of  $\sigma$  relative to the Kobayashi pseudometric*, denoted by  $l_{\Omega}(\sigma)$ . For this, we first observe that the Kobayashi pseudometric  $k_{\Omega}$  is an upper semicontinuous function on  $T^{(1,0)}\Omega := T^{(1,0)}X|_{\Omega}$ . Hence, from Remark 2.2,  $k_{\Omega}(\sigma(t); \sigma'(t))$  is integrable. Thus, the integral below makes sense and we define

$$l_{\Omega}(\sigma) := \int_a^b k_{\Omega}(\sigma(t); \sigma'(t)) dt.$$

For a Kobayashi hyperbolic domain  $\Omega \subseteq X$ , the Kobayashi pseudometric  $k_{\Omega}$  has the following properties:

**Result 2.4.** *Let  $X$  be a complex manifold of dimension  $n$  and  $\Omega \subseteq X$  be a Kobayashi hyperbolic domain.*

(a) [27, Theorem 1] *For any  $z, w \in \Omega$  we have*

$$K_{\Omega}(z, w) = \inf\{l_{\Omega}(\sigma) : \sigma : [a, b] \rightarrow \Omega \text{ is piecewise } \mathcal{C}^1 \text{ with } \sigma(a) = z \text{ and } \sigma(b) = w\}.$$

(b) [29, Theorem 3.1] *For any  $z, w$  we have*

$$K_{\Omega}(z, w) = \inf\{l_{\Omega}(\sigma) : \sigma : [a, b] \rightarrow \Omega \text{ is absolutely continuous with } \sigma(a) = z \text{ and } \sigma(b) = w\}.$$

**Result 2.5** (paraphrasing [27, Theorem 2]). *Let  $X$  be a complex manifold of dimension  $n$  and  $\Omega \subseteq X$  be a domain. Fix a Hermitian metric  $h$  on  $X$  and let  $d_h$  denote the distance induced by  $h$ . Then,  $\Omega$  is Kobayashi hyperbolic if and only if for each  $x \in \Omega$ , there exist constants  $c_x > 0$ ,  $r_x > 0$  such that  $k_{\Omega}(y; v) \geq c_x h_y(v)$  for every  $v \in T_y^{(1,0)}\Omega$  and every  $y \in B_{d_h}(x, r_x)$ .*

*Remark 2.6.* In the above statement,  $B_{d_h}(x, r)$  denotes the open  $d_h$ -ball centered at  $x$  and having radius  $r$ . In the above, and elsewhere in this paper, we abbreviate  $\sqrt{h_x(v, v)}$ ,  $v \in T_x^{(1,0)}X$ , as  $h_x(v)$ . The above result implies that when  $\Omega$  is Kobayashi hyperbolic, the Kobayashi pseudometric  $k_{\Omega}$  is a (Finsler) metric.

Now, we will prove that in the metric space  $(\Omega, K_\Omega)$  (where  $\Omega$  is a Kobayashi hyperbolic domain in a complex manifold  $X$ ), any two points in  $\Omega$  can be joined by a  $(1, \kappa)$ -almost-geodesic. To prove this result, we need a technical lemma.

**Lemma 2.7.** *Let  $X$  be a complex manifold and  $\Omega \subseteq X$  be a Kobayashi hyperbolic domain. Let  $\sigma : [a, b] \rightarrow \Omega$  be an absolutely continuous path. If*

$$l_\Omega(\sigma) \leq K_\Omega(\sigma(a), \sigma(b)) + \varepsilon,$$

for some  $\varepsilon > 0$ , then, whenever  $a \leq a' \leq b' \leq b$ , we have

$$l_\Omega(\sigma|_{[a', b']}) \leq K_\Omega(\sigma(a'), \sigma(b')) + \varepsilon.$$

In view of part (b) of Result 2.4, the above lemma follows from the definition of  $l_\Omega(\sigma)$  and the triangle inequality for the Kobayashi distance on  $\Omega$ .

**Proposition 2.8.** *Let  $X$  be a complex manifold and  $\Omega \subseteq X$  be a Kobayashi hyperbolic domain in  $X$ . For any  $\kappa > 0$  and  $x, y \in \Omega$ , there exists a  $(1, \kappa)$ -almost-geodesic  $\sigma : [a, b] \rightarrow \Omega$  with  $\sigma(a) = x$  and  $\sigma(b) = y$ .*

*Proof.* Fix a Hermitian metric  $h$  on  $X$  and let  $d_h$  be the distance induced by  $h$  on  $X$ . It is a classical result that

$$Top(\Omega) = Top(d_h|_{\Omega \times \Omega}),$$

where  $Top(\Omega)$  denotes the manifold topology on  $\Omega$ . Now, from part (a) of Result 2.4, there exists a piecewise  $\mathcal{C}^1$  curve  $\gamma : [0, T] \rightarrow \Omega$  such that  $\gamma(0) = x$ ,  $\gamma(T) = y$ , and

$$l_\Omega(\gamma) < K_\Omega(x, y) + \kappa. \quad (2.1)$$

Since  $\gamma([0, T])$  is compact, it can be covered by finitely many charts on  $\Omega$ . Also the Kobayashi metric  $k_\Omega$  is upper semicontinuous on  $T^{(1,0)}\Omega$ . Therefore, perturbing  $\gamma$  if needed, we can assume that  $\gamma$  is  $\mathcal{C}^1$  smooth and that  $h_{\gamma(t)}(\gamma'(t)) \neq 0$  for all  $t \in [0, T]$ . Now, we define the function  $f : [0, T] \rightarrow \mathbb{R}$  by

$$f(t) = \int_0^t k_\Omega(\gamma(s); \gamma'(s)) ds.$$

Since we know  $k_\Omega$  is upper semicontinuous on  $T^{(1,0)}\Omega$ , there exists a constant  $C_1 > 0$  such that for all  $t \in [0, T]$  and every tangent vector  $v \in T_{\gamma(t)}^{(1,0)}\Omega$  satisfying  $h_{\gamma(t)}(v) = 1$ , we have

$$\begin{aligned} k_\Omega(\gamma(t); v) &\leq C_1 \\ \implies k_\Omega(\gamma(t); \gamma'(t)) &\leq C_1 h_{\gamma(t)}(\gamma'(t)). \end{aligned} \quad (2.2)$$

Since  $\Omega$  is Kobayashi hyperbolic, from Result 2.5 it follows that for each  $t \in [0, T]$ , there exist constants  $r_t > 0$ ,  $\tilde{C}_t > 0$  such that for all  $y \in B_{d_h}(\gamma(t), r_t)$  and for all  $v \in T_y^{(1,0)}\Omega$

$$k_\Omega(y; v) \geq \tilde{C}_t h_y(v).$$

Since  $\gamma([0, T])$  is compact, there exist  $t_1, \dots, t_N \in [0, T]$  such that  $B_{d_h}(\gamma(t_1), r_{t_1}), \dots, B_{d_h}(\gamma(t_N), r_{t_N})$  cover  $\gamma([0, T])$ . Therefore, there exists a constant  $C_2 > 0$  such that, taking (2.2) into consideration, we have

$$C_2 h_{\gamma(t)}(\gamma'(t)) \leq k_\Omega(\gamma(t); \gamma'(t)) \leq C_1 h_{\gamma(t)}(\gamma'(t)),$$

for all  $t \in [0, T]$ . By assumption,  $h_{\gamma(t)}(\gamma'(t)) \neq 0$  for all  $t \in [0, T]$ . As  $h$  is continuous, there exists a constant  $\alpha > 1$  such that

$$\begin{aligned} (1/\alpha) &\leq C_2 h_{\gamma(t)}(\gamma'(t)) \leq k_\Omega(\gamma(t); \gamma'(t)) \leq C_1 h_{\gamma(t)}(\gamma'(t)) \leq \alpha \\ \implies (1/\alpha)|t_1 - t_2| &\leq |f(t_1) - f(t_2)| \leq \alpha|t_1 - t_2|, \end{aligned}$$



for all  $t_1, t_2 \in [0, T]$ . Hence,  $f$  is bi-Lipschitz and strictly increasing function. Therefore, we can define the function  $g = f^{-1} : [0, l_\Omega(\gamma)] \rightarrow [0, T]$ .

**Claim.**  $\tilde{\gamma} := \gamma \circ g$  is a  $(1, \kappa)$ -almost-geodesic joining  $x$  and  $y$ .

*Proof of Claim.* Since  $f$  is bi-Lipschitz,  $g$  is bi-Lipschitz. Now, fix  $t_0 \in [0, l_\Omega(\gamma)]$ , fix a holomorphic chart  $(U, \varphi)$  around  $\tilde{\gamma}(t_0)$ , and let  $I(\varphi, t_0)$  be a closed interval of  $\tilde{\gamma}^{-1}(U)$  containing  $t_0$ . So,  $g(I(\varphi, t_0))$  is a closed interval. Since  $\varphi$  is smooth, and  $\gamma$  is  $\mathcal{C}^1$  smooth on  $[0, T]$ ,  $\varphi \circ \gamma|_{g(I(\varphi, t_0))}$  is Lipschitz. Hence,  $\varphi \circ \tilde{\gamma}|_{I(\varphi, t_0)}$  is Lipschitz. Therefore,  $\varphi \circ \tilde{\gamma}|_{I(\varphi, t_0)}$  is an absolutely continuous path. So, by Definition 2.1,  $\tilde{\gamma}$  is absolutely continuous.

Since  $k_\Omega(\gamma(s); \gamma'(s))$  is bounded on  $[0, T]$ , it is integrable on  $[0, T]$ . By Lebesgue differentiation theorem, there exists a set  $E \subseteq [0, T]$  of full measure such that  $f$  is differentiable and  $f'(s) = k_\Omega(\gamma(s); \gamma'(s))$  for each  $s \in E$ . Since  $g$  is bi-Lipschitz,  $g^{-1}(E) \subseteq [0, l_\Omega(\gamma)]$  has full measure, hence on  $g^{-1}(E)$ ,  $g$  is differentiable, and for all  $s \in g^{-1}(E)$ :

$$g'(s) = 1/f'(g(s)).$$

At this stage we are reduced to examining functions defined on intervals in  $\mathbb{R}$ . No charts of  $X$  are now involved. Thus, in view of Lemma 2.7, the proof that  $\tilde{\gamma}$  has properties (a) and (b) given in Definition 2.1 now proceeds exactly as in the proof of [8, Proposition 4.4].  $\blacktriangleleft$

Since the above claim is true, the result follows.  $\square$

We conclude this section with a technical lemma. Its proof is trivial, despite its technical hypothesis, and we state it here because we shall invoke it several times in the later sections.

**Lemma 2.9.** *Let  $X$  be a complex manifold and  $\Omega \subsetneq X$  be a Kobayashi hyperbolic domain in  $X$ . Let  $h$  be a Hermitian metric on  $X$  and let  $(K_\nu)_{\nu \geq 1}$  be an exhaustion by compacts of  $\Omega$ . Fix  $\lambda \geq 1$ ,  $\kappa \geq 0$ , and  $p \in \partial\Omega$ . Let  $U$  and  $W$  be neighbourhoods of  $p$  in  $X$  such that  $U$  is relatively compact and  $W \Subset U$ . Let  $(x_\nu)_{\nu \geq 1} \subset W \cap \Omega$  and, for each  $\nu$ , let  $\gamma_\nu : [a_\nu, b_\nu] \rightarrow \Omega$  be a  $(\lambda, \kappa)$ -almost-geodesic such that  $\gamma_\nu(a_\nu) = x_\nu$  and such that*

$$\gamma_\nu([a_\nu, b_\nu]) \subset \Omega \setminus K_\nu \quad \text{and} \quad \gamma_\nu(b_\nu) \notin U \quad \forall \nu \geq 1.$$

*Given a neighbourhood  $V$  of  $p$  in  $X$  such that  $W \Subset V \Subset U$  and  $r_\nu \in (a_\nu, b_\nu)$  such that*

$$y'_\nu := \gamma_\nu(r_\nu) \in V \setminus \overline{W} \quad \text{and} \quad \gamma_\nu([a_\nu, r_\nu]) \subset U \quad \forall \nu \geq 1.$$

*Write  $\tilde{\gamma}_\nu := \gamma_\nu|_{[a_\nu, r_\nu]}$ . Then, there exist a point  $p' \in \overline{W} \cap \partial\Omega$  and a subsequence  $(\tilde{\gamma}_{\nu_k})_{k \geq 1}$  such that such that  $x_{\nu_k} \rightarrow p'$  and*

$$\lim_{k \rightarrow \infty} \sup_{z \in \text{image}(\tilde{\gamma}_{\nu_k})} d_h(z, \partial\Omega) = 0.$$

*Remark 2.10.* In the above statement,  $d_h$  denotes the distance induced by  $h$  on  $X$ . The existence of a neighbourhood  $V$  and numbers  $r_\nu \in (a_\nu, b_\nu)$ ,  $\nu \geq 1$ , with the properties stated in the hypothesis above is guaranteed by the condition  $\gamma_\nu([a_\nu, b_\nu]) \subset \Omega \setminus K_\nu$  for each  $\nu \geq 1$ .

The proof of Lemma 2.9 is a consequence of the condition  $\gamma_\nu([a_\nu, b_\nu]) \subset \Omega \setminus K_\nu$  for all  $\nu \geq 1$  and the fact that  $U$  is relatively compact, and is trivial. Thus, we shall skip the proof.

**2.3. Gromov hyperbolic metric spaces.** In Section 1.4 we mentioned Gromov hyperbolic metric spaces. Let us begin with the definition of Gromov hyperbolicity of a metric space  $(X, d)$  that does not require  $(X, d)$  to be a geodesic space. But first, for  $x, y, o \in X$ , we define the *Gromov product*  $(x|y)_o$  by  $(x|y)_o := (d(x, o) + d(y, o) - d(x, y))/2$ .

**Definition 2.11.** Let  $(X, d)$  be a metric space. We say that  $(X, d)$  is *Gromov hyperbolic* if there exists  $\delta \geq 0$  such that for any four points  $x, y, z, o \in X$ ,

$$\min \{ (x|y)_o, (y|z)_o \} \leq (x|z)_o + \delta. \quad (2.3)$$

Here, we will present some more notations and a useful result to prove Theorem 1.18. These are taken from [17, Section 3.4].

**Definition 2.12.** Let  $(X, d)$  be a metric space.

(a) We say that a sequence  $(x_\nu)_{\nu \geq 1} \subset X$  is a *Gromov sequence* if for some (hence any)  $o \in X$ ,

$$\lim_{\nu, l \rightarrow \infty} (x_l | x_\nu)_o = +\infty.$$

(b) Let  $(X, d)$  be Gromov hyperbolic. We say that two Gromov sequences  $(x_\nu)_{\nu \geq 1}$  and  $(y_\nu)_{\nu \geq 1}$  are *equivalent* if for some (and hence any)  $o \in X$ ,

$$\lim_{\nu, l \rightarrow \infty} (x_l | y_\nu)_o = +\infty.$$

If  $(X, d)$  is Gromov hyperbolic, it follows that the above relation (between two sequences of  $X$ ) is an equivalence relation.

We shall denote the equivalence class of Gromov sequences  $(x_\nu)_{\nu \geq 1}$  by  $[x_\nu]$ . We define the *Gromov boundary* of  $X$  to be the set of all equivalence classes of Gromov sequences in  $X$ . We shall denote the Gromov boundary of  $X$  by  $\partial_G X$ . Recall that the *Gromov bordification* of  $X$ , denoted by  $\overline{X}^G$ , as a set is  $\overline{X}^G := X \cup \partial_G X$ . We will not elaborate upon the construction of the topology on  $\overline{X}^G$ ; we refer the reader to [17, Section 3.4] for details about Gromov topology. Instead, we present the following result which is relevant for our paper and also gives some idea about the Gromov topology.

**Result 2.13.** *Let  $(X, d)$  be a Gromov hyperbolic metric space. Then,*

- (a) *The Gromov topology depends only on  $d$  and the inclusion  $X \hookrightarrow \overline{X}^G$  ( $X$  being equipped with the metric topology) is homeomorphic.*
- (b) *A sequence  $(x_\nu)_{\nu \geq 1} \subset X$  converges to a point  $\xi \in \partial_G X$  if and only if  $(x_\nu)_{\nu \geq 1}$  is a Gromov sequence and  $[x_\nu] = \xi$ .*

### 3. THE PROOFS OF THEOREMS 1.5 AND 1.6

To prove the above-mentioned theorems, we will need the following result:

**Result 3.1** (paraphrasing [27, Lemma 2 of §4]). *Let  $X$  be a complex manifold and  $\Omega \subseteq X$  be a Kobayashi hyperbolic domain. Let  $U$  be an open subset of  $X$  such that  $U \cap \Omega \neq \emptyset$  with  $U \cap \Omega$  connected. Then, for every  $v \in T_x^{(1,0)}(U \cap \Omega)$  and every  $x \in U \cap \Omega$ , we have*

$$k_\Omega(x; v) \leq k_{U \cap \Omega}(x; v) \leq \coth(K_\Omega(x, \Omega \setminus U)) k_\Omega(x; v).$$

In what follows, if  $\sigma : I \rightarrow \Omega$ ,  $I \subseteq \mathbb{R}$  an interval, is a curve, then we will denote the image of  $\sigma$  by  $\langle \sigma \rangle$ . In the proof that follows, we shall use the notation introduced in Section 2.

*The proof of Theorem 1.5.* Fix a Hermitian metric  $h$  on  $X$  and let  $d_h$  be the distance induced by  $h$  on  $X$ . First we will prove that (a) implies (b). In order to prove that  $\partial\Omega$  is visible, fix  $p, q \in \partial\Omega$  such that  $p \neq q$ . Since  $\partial\Omega$  is locally visible, there exists a neighbourhood  $U_p$  of  $p$  in  $X$  such that  $U_p \cap \Omega$  is a Kobayashi hyperbolic domain and there is a neighbourhood  $V_p$  of  $p$  in  $X$ ,  $V_p \subseteq U_p$ , such that every pair of distinct points  $q_1, q_2 \in V_p \cap \partial\Omega$  satisfies the visibility condition with respect to  $K_{U_p \cap \Omega}$ . Without loss of generality, we may assume that  $U_p$  is relatively compact.

We also fix  $\lambda \geq 1$  and  $\kappa \geq 0$ . Since  $\Omega$  is a hyperbolically imbedded domain in  $X$ , shrinking  $V_p$  if needed, we may assume without loss of generality that  $K_\Omega(V_p \cap \Omega, \Omega \setminus U_p) := \delta_p > 0$  and that  $q \notin \overline{V_p}$ . Therefore, there exists a neighbourhood  $V_q$  of  $q$  in  $X$  such that  $V_q$  is relatively compact, and such that  $\overline{V_p} \cap \overline{V_q} = \emptyset$ . Let  $W_p$  be a neighbourhood of  $p$  in  $X$  such that  $\overline{W_p} \subseteq V_p$ . For any pair  $x, y \in \Omega$  we define the set

$$AG_\Omega(x, y) := \{\gamma : [0, T] \longrightarrow \Omega : \gamma \text{ is a } (\lambda, \kappa)\text{-almost-geodesic joining } x \text{ and } y\}.$$

Here, the parameters  $\lambda$  and  $\kappa$  are those that we fixed above (we do not assume that  $AG_\Omega(x, y) \neq \emptyset$  when  $\kappa = 0$ ).

**Claim 1.** *There exists a compact set  $K \subset \Omega$  such that for every  $\gamma \in AG_\Omega(x, y)$ , every  $x \in W_p \cap \Omega$ , and every  $y \in V_q \cap \Omega$ , we have  $\langle \gamma \rangle \cap K \neq \emptyset$ .*

*Proof of Claim 1.* If possible let the claim be false. Let  $(K_\nu)_{\nu \geq 1}$  be an exhaustion by compacts of  $\Omega$ . Then, there exist sequences  $(x_\nu)_{\nu \geq 1} \subset W_p \cap \Omega$ ,  $(y_\nu)_{\nu \geq 1} \subset V_q \cap \Omega$ , and  $(\lambda, \kappa)$ -almost-geodesics  $\gamma_\nu$  in  $AG_\Omega(x_\nu, y_\nu)$ , for each  $\nu \geq 1$ , such that  $\langle \gamma_\nu \rangle \cap K_\nu = \emptyset$ . Fix neighbourhoods  $W_p^1, W_p^2$  of  $p$  such that  $W_p \Subset W_p^1 \Subset W_p^2 \Subset V_p$ . Let  $r_\nu > 0$  and  $y'_\nu = \gamma_\nu(r_\nu)$  be such that  $y'_\nu \in (V_p \setminus \overline{W_p^1}) \cap \Omega$ , and such that the image of  $\tilde{\gamma}_\nu := \gamma_\nu|_{[0, r_\nu]}$  is contained in  $W_p^2$ . By Lemma 2.9, and as we can relabel the subsequences given by it, there exist a point  $p' \in \overline{W_p} \cap \partial\Omega$  such that  $x_\nu \rightarrow p'$  and

$$\lim_{\nu \rightarrow \infty} \sup_{z \in \langle \tilde{\gamma}_\nu \rangle} d_h(z, \partial\Omega) = 0. \quad (3.1)$$

From Result 3.1 and the fact that  $K_\Omega(z, \Omega \setminus U_p) \geq \delta_p$  for all  $z \in V_p \cap \Omega$ , it follows that for every  $v \in T_z^{(1,0)}\Omega$  and every  $z \in V_p \cap \Omega$

$$k_\Omega(z; v) \leq k_{U_p \cap \Omega}(z; v) \leq \coth(K_\Omega(z, \Omega \setminus U_p))k_\Omega(z; v) \leq Ak_\Omega(z; v),$$

where  $A := \coth \delta_p (> 1)$ . Therefore, from part (a) of Result 2.4, we have

$$K_\Omega(z, w) \leq K_{U_p \cap \Omega}(z, w) \leq AK_\Omega(z, w),$$

for all  $z, w \in V_p \cap \Omega$ . Since  $\langle \tilde{\gamma}_\nu \rangle \subset V_p \cap \Omega$  and  $\gamma_\nu$  is a  $(\lambda, \kappa)$ -almost-geodesic for all  $\nu \geq 1$  and as  $A > 1$ , it follows from the last few inequalities that for all  $t, s \in [0, r_\nu]$

- (i)  $(A\lambda)^{-1}|t - s| - A\kappa \leq K_{U_p \cap \Omega}(\tilde{\gamma}_\nu(t), \tilde{\gamma}_\nu(s)) \leq A\lambda|t - s| + A\kappa$ ,
- (ii)  $k_{U_p \cap \Omega}(\tilde{\gamma}_\nu(t); \tilde{\gamma}'_\nu(t)) \leq A\lambda$  for a.e.  $t \in [0, r_\nu]$ .

Hence,  $\tilde{\gamma}_\nu$  is a  $(A\lambda, A\kappa)$ -almost-geodesic with respect to  $K_{U_p \cap \Omega}$  joining  $x_\nu$  to  $y'_\nu$ . As  $W_p^2$  is relatively compact, there exists a sequence  $(\nu_j)_{j \geq 1} \subset \mathbb{N}$  and a point  $q' \in \overline{W_p^2} \cap \partial\Omega \subseteq V_p \cap \partial\Omega$  such that  $\gamma_{\nu_j}(r_{\nu_j}) \rightarrow q'$ . By construction,  $p' \neq q'$ . Now, since  $V_p \cap \partial\Omega$  satisfies the visibility condition with respect to  $K_{U_p \cap \Omega}$ , there exists a compact set  $K \subset U_p \cap \Omega$  such that for every  $j \geq 1$  we can find  $z_j \in \langle \tilde{\gamma}_{\nu_j} \rangle$  with  $z_j \in K$ , which contradicts (3.1). Hence, the claim.  $\blacktriangleleft$

By this claim, we have established that  $\partial\Omega$  is visible.

Now to prove (b) implies (a), we only need to prove that  $\Omega$  is a hyperbolically imbedded domain. We shall establish the equivalent property stated in Remark 1.4. Let  $x, y \in \overline{\Omega}$  such that  $x \neq y$ . When at least one of  $x, y$  is in  $\Omega$ , this property follows easily from the fact that  $\Omega$  is Kobayashi hyperbolic, whereby  $Top(\Omega) = Top(K_\Omega)$ . Thus, consider the case when both  $x, y \in \partial\Omega$ . Since  $\partial\Omega$  is visible, there exist neighbourhoods  $V_x$  of  $x$  and  $V_y$  of  $y$  in  $X$  such that  $\overline{V_x} \cap \overline{V_y} = \emptyset$  and such that there exists a compact set  $K_0 \subset \Omega$  such that the image of each  $(1,1)$ -almost-geodesic  $\gamma : [0, T] \longrightarrow \Omega$  with  $\gamma(0) \in V_x$  and  $\gamma(T) \in V_y$  intersects  $K_0$ .

**Claim 2.**  $K_\Omega(V_x \cap \Omega, V_y \cap \Omega) > 0$

*Proof of Claim 2.* If possible let  $K_\Omega(V_x \cap \Omega, V_y \cap \Omega) = 0$ . Then, there exist sequences  $(z_\nu)_{\nu \geq 1} \subset V_x \cap \Omega$ ,  $(w_\nu)_{\nu \geq 1} \subset V_y \cap \Omega$  such that  $K_\Omega(z_\nu, w_\nu) \rightarrow 0$  as  $\nu \rightarrow \infty$ . By Proposition 2.8, we can

find  $\tau_\nu : [0, T_\nu] \rightarrow \Omega$ , a  $(1, 1/\nu)$ -almost-geodesic such that  $\tau(0) = z_\nu$  and  $\tau_\nu(T_\nu) = w_\nu$  for each  $\nu \geq 1$ . Clearly, each  $\tau_\nu$  is also a  $(1, 1)$ -almost-geodesic with respect to  $K_\Omega$ . Therefore, by assumption,  $\langle \tau_\nu \rangle \cap K_0 \neq \emptyset$  for all  $\nu \geq 1$ . Let  $s_\nu \in [0, T_\nu]$  be such that  $\tau_\nu(s_\nu) =: a_\nu \in \langle \tau_\nu \rangle \cap K_0$  and, by passing to a subsequence if needed, assume that  $a_\nu \rightarrow a \in K_0$  as  $\nu \rightarrow \infty$ . Since  $\tau_\nu$  is a  $(1, 1/\nu)$ -almost-geodesic passing through  $a_\nu$ , we have

$$\begin{aligned} K_\Omega(z_\nu, w_\nu) &\geq T_\nu - 1/\nu = (T_\nu - s_\nu + 1/\nu) + (s_\nu - 0 + 1/\nu) - 3/\nu \\ &\geq K_\Omega(z_\nu, a_\nu) + K_\Omega(a_\nu, w_\nu) - 3/\nu, \end{aligned}$$

for each  $\nu \geq 1$ . Therefore,  $K_\Omega(z_\nu, a_\nu) \rightarrow 0$  and  $K_\Omega(a_\nu, w_\nu) \rightarrow 0$  as  $\nu \rightarrow \infty$ . Since  $\Omega$  is a Kobayashi hyperbolic domain, from the discussion above it follows that  $z_\nu \rightarrow a$  and  $w_\nu \rightarrow a$  as  $\nu \rightarrow \infty$ , which contradicts the fact that  $z_\nu \in V_x \cap \Omega$ ,  $w_\nu \in V_y \cap \Omega$  and  $\overline{V_x} \cap \overline{V_y} = \emptyset$ . Hence, Claim 2 is true.  $\blacktriangleleft$

Combining this claim with the discussion that precedes it, the result follows.  $\square$

We shall now undertake the proof of Theorem 1.6. We will need the following lemmas to prove this theorem:

**Lemma 3.2.** *Let  $X$  be a complex manifold and  $\Omega \subseteq X$  be a Kobayashi hyperbolic domain. Let  $U$  be an open subset of  $X$  such that  $U \cap \Omega \neq \emptyset$  with  $U \cap \Omega$  connected. Then, for every  $V \subset U$  with  $V \cap \Omega \neq \emptyset$  and  $K_\Omega(V \cap \Omega, \Omega \setminus U) > 0$ , there exists a constant  $M > 0$  such that for every  $v \in T_x^{(1,0)}(V \cap \Omega)$  and every  $x \in V \cap \Omega$ ,*

$$k_\Omega(x; v) \leq k_{U \cap \Omega}(x; v) \leq (1 + Me^{-K_\Omega(x, \Omega \setminus U)})k_\Omega(x; v).$$

In view of Result 3.1, the argument for Lemma 3.2 is as in the proof of [28, Lemma 3.1].

**Lemma 3.3.** *Let  $X$  be a complex manifold and  $\Omega \subsetneq X$  be a Kobayashi hyperbolic domain. Let  $U, V, W$  be non-empty open subsets of  $X$  such that  $W \cap \Omega \neq \emptyset$ ,  $W \Subset V \Subset U$  with  $U \cap \Omega$  connected, and such that  $\delta := K_\Omega(V \cap \Omega, \Omega \setminus U) > 0$ . Let  $(U \cap \Omega, K_{U \cap \Omega})$  have the weak visibility property for every pair of distinct points in  $V \cap \partial\Omega$ . Let  $\kappa \geq 0$ . Then, there exists a constant  $\kappa_0 > 0$  such that for every  $x, y \in W \cap \Omega$  with  $x \neq y$ , and for  $\sigma : [0, T] \rightarrow \Omega$  a  $(1, \kappa)$ -almost-geodesic with respect to  $K_\Omega$  joining  $x$  and  $y$  such that the image of  $\sigma$  is contained in  $W$ ,*

$$l_{U \cap \Omega}(\sigma|_{[s_1, s_2]}) \leq K_{U \cap \Omega}(\sigma(s_1), \sigma(s_2)) + \kappa_0$$

for every  $s_1 \leq s_2 \in [0, T]$ .

Essentially, following the same argument as in the proof of [28, Lemma 3.5]—which is valid in the present setting in view of Lemma 3.2—up to the estimate [28, (3.6)] gives us the desired inequality.

*The proof of Theorem 1.6.* First we will prove that (a) implies (b). **Fix**  $\kappa \geq 0$ . As in the proof of Theorem 1.5, considering the **same** notations, we will prove Claim 1 occurring in its proof with  $\lambda = 1$ . As before, we will assume that the latter claim is false. This gives us  $\gamma_\nu$ ,  $\nu = 1, 2, 3, \dots$ ,  $(1, \kappa)$ -almost-geodesics with respect to  $K_\Omega$  joining  $x_\nu$  to  $y_\nu$ , where  $x_\nu \in W_p \cap \Omega$ ,  $y_\nu \in V_q \cap \Omega$ , having **all** the other properties listed in the proof of the above-mentioned Claim 1. Let  $r_\nu > 0$  and  $y'_\nu = \gamma_\nu(r_\nu)$  be such that  $y'_\nu \in (V_p \setminus \overline{W_p^1}) \cap \Omega$ , and such that the image of  $\tilde{\gamma}_\nu := \gamma_\nu|_{[0, r_\nu]}$  is contained in  $W_p^2$ . As indicated above,  $W_p^1$  and  $W_p^2$  are the same as in the proof of the above-mentioned Claim 1, and as argued in that proof using Lemma 2.9, we get  $p' \in \overline{W_p^1} \cap \partial\Omega$  such that  $x_\nu \rightarrow p'$  and

$$\lim_{\nu \rightarrow \infty} \sup_{z \in \langle \tilde{\gamma}_\nu \rangle} d_h(z, \partial\Omega) = 0. \quad (3.2)$$

By Lemma 3.3, there exists a constant  $\kappa_0 > 0$  such that for every  $s_1, s_2 \in [0, r_\nu]$  and every  $\nu \geq 1$  we have

$$l_{U_p \cap \Omega}(\tilde{\gamma}_\nu|_{[s_1, s_2]}) \leq K_{U_p \cap \Omega}(\tilde{\gamma}_\nu(s_1), \tilde{\gamma}_\nu(s_2)) + \kappa_0. \quad (3.3)$$

For each  $\nu$ , reparametrizing  $\tilde{\gamma}_\nu$  by (the Kobayashi–Royden) arc length and denoting this by  $\sigma_\nu$ , we conclude from (3.3) that  $\sigma_\nu : [0, r'_\nu] \rightarrow W_p^2$  is a  $(1, \kappa_0)$ -almost-geodesic with respect to  $K_{U_p \cap \Omega}$  joining  $x_\nu$  to  $y'_\nu$ . As  $W_p^2$  is relatively compact, there exists a sequence  $(\nu_j)_{j \geq 1} \subset \mathbb{N}$  and a point  $q' \in \overline{W_p^2} \cap \partial\Omega \subseteq V_p \cap \partial\Omega$  such that  $\gamma_{\nu_j}(r_{\nu_j}) = \sigma_{\nu_j}(r'_{\nu_j}) \rightarrow q'$ . By construction,  $p' \neq q'$ . Now, since  $V_p \cap \partial\Omega$  satisfies the weak visibility condition with respect to  $K_{U_p \cap \Omega}$ , there exists a compact set  $K \subset U_p \cap \Omega$  such that for every  $j \geq 1$  we can find  $z_j \in \langle \sigma_{\nu_j} \rangle = \langle \tilde{\gamma}_{\nu_j} \rangle$  with  $z_j \in K$ , which contradicts (3.2). Hence, our claim is true; therefore,  $\partial\Omega$  is weakly visible.

Now, to prove the converse, we only need to prove that  $\Omega$  is a hyperbolically imbedded domain. Note that, in the argument for (b) implies (a) in the proof of Theorem 1.5, we only need to work with  $(1, \kappa)$ -almost-geodesics. Therefore, that argument establishes the implication that we want. Hence, the result.  $\square$

#### 4. COMPACTIFICATIONS AND VISIBILITY DOMAINS

With  $X$  and  $\Omega \subsetneq X$  as in the discussions above, recall what an “admissible compactification of  $\overline{\Omega}$ ” means; see Definition 1.14. If  $\overline{\Omega}^\infty$  is an admissible compactification of  $\overline{\Omega}$  and  $\mathbf{j} : \overline{\Omega} \hookrightarrow \overline{\Omega}^\infty$  is the embedding that is part of the definition of  $\overline{\Omega}^\infty$ , then let us denote the set  $\overline{\Omega}^\infty \setminus \mathbf{j}(\Omega)$  by  $\partial_\infty\Omega$ .

We now present the central result of this section, which will play a key supporting role in the proofs presented in the subsequent sections. In what follows, given a set  $S \subseteq \overline{\Omega}^\infty$ ,  $\overline{S}^\infty$  will denote the closure of  $S$  in  $\overline{\Omega}^\infty$ . The sets  $\partial_{\text{visi}}\Omega$  and  $\partial_{\text{wvisi}}\Omega$  appearing in the result below are as defined in Definition 1.7.

**Theorem 4.1.** *Let  $X$  be a complex manifold and  $\Omega \subsetneq X$  be a Kobayashi hyperbolic domain. Let  $\overline{\Omega}^\infty$  be any admissible compactification of  $\overline{\Omega}$ .*

- (a) *Suppose the set  $\partial\Omega \setminus \partial_{\text{visi}}\Omega$  is totally disconnected. Then,  $\Omega$  is a visibility domain subordinate to  $\overline{\Omega}^\infty$ .*
- (b) *Suppose the set  $\partial\Omega \setminus \partial_{\text{wvisi}}\Omega$  is totally disconnected. Then,  $\Omega$  is a weak visibility domain subordinate to  $\overline{\Omega}^\infty$ .*

*Proof.* Fix a Hermitian metric  $h$  on  $X$  and let  $d_h$  be the distance induced by  $h$  on  $X$ . Fix  $p, q \in \partial_\infty\Omega$  such that  $p \neq q$ . Let us first prove (a). Therefore, fix  $\lambda \geq 1$  and  $\kappa \geq 0$ . Let  $V_p, V_q$  be two  $\overline{\Omega}^\infty$ -open neighbourhoods such that  $\overline{V_p}^\infty \cap \overline{V_q}^\infty = \emptyset$ . For any pair  $x, y \in \Omega$  we define the set

$$AG_\Omega(x, y) := \{\gamma : [0, T] \longrightarrow \Omega : \gamma \text{ is a } (\lambda, \kappa)\text{-almost-geodesic joining } x \text{ and } y\}.$$

Here, the parameters  $\lambda$  and  $\kappa$  are those that we fixed above.

We shall show that there exists a compact set  $K \subset \Omega$  such that for every  $\gamma \in AG_\Omega(x, y)$ , every  $x \in V_p \cap \Omega$ , and every  $y \in V_q \cap \Omega$ , we have  $\langle \gamma \rangle \cap K \neq \emptyset$ . If possible, let this assertion be false. Let  $(K_\nu)_{\nu \geq 1}$  be an exhaustion by compacts of  $\Omega$ . Then, there exist sequences  $(x_\nu)_{\nu \geq 1} \subset V_p \cap \Omega$ ,  $(y_\nu)_{\nu \geq 1} \subset V_q \cap \Omega$ , and  $(\lambda, \kappa)$ -almost-geodesics  $\gamma_\nu$  in  $AG_\Omega(x_\nu, y_\nu)$ , for each  $\nu \geq 1$ , such that  $\langle \gamma_\nu \rangle \cap K_\nu = \emptyset$ . Let  $\gamma_\nu : [0, T_\nu] \longrightarrow \Omega$ . Since  $S_\nu := \gamma_\nu([0, T_\nu])$  is a compact subset of  $X$ , passing to a subsequence and relabelling, if needed, we can assume that  $(S_\nu)_{\nu \geq 1}$  converges in the local

Hausdorff topology to a closed subset  $S \subseteq X$ . Since,  $\overline{V_p}^\infty \cap \overline{V_q}^\infty = \emptyset$  and  $\langle \gamma_\nu \rangle \cap K_\nu = \emptyset$  for all  $\nu \geq 1$ , we can conclude that  $\overline{S}^\infty$  contains at least two elements and  $S \subseteq \partial\Omega$ .

**Claim.**  $\overline{S}^\infty$  is connected.

*Proof of Claim.* If possible, let the claim be false. Then, there exist two disjoint  $\overline{\Omega}^\infty$ -open sets  $U, V$  such that  $\overline{S}^\infty \subseteq U \cup V$  and  $U \cap \overline{S}^\infty \neq \emptyset, V \cap \overline{S}^\infty \neq \emptyset$ . Therefore, as each  $S_\nu$  is connected, for sufficiently large  $\nu$ , there exists a point (recall our mild abuse of notation described in Section 1.3)  $z'_\nu \in \mathbf{j}(S_\nu) \setminus (U \cup V)$ . As  $\overline{\Omega}^\infty$  is sequentially compact, there exists a subsequence  $(z'_{\nu_k})_{k \geq 1}$  and a point  $\tilde{p} \in \overline{S}^\infty$  such that  $\mathbf{j}(z'_{\nu_k}) \rightarrow \tilde{p}$  with respect to the topology of  $\overline{\Omega}^\infty$ . Then  $\tilde{p} \in \overline{S}^\infty \setminus (U \cup V)$ , which is impossible. Hence, the claim.  $\blacktriangleleft$

Write  $\omega := \partial\Omega \setminus \partial_{\text{visi}}\Omega$ , a closed set in  $\partial\Omega$ . Since  $\overline{\Omega}^\infty \setminus \mathbf{j}(\overline{\Omega})$  is totally disconnected, it easily follows that  $\overline{\omega}^\infty$  is totally disconnected. In view of the above claim, and as  $S$  is non-empty, there exists  $\xi \in S \cap \partial_{\text{visi}}\Omega$ . Since  $\overline{V_p}^\infty \cap \overline{V_q}^\infty = \emptyset$ , without loss of generality we can assume that  $\xi \notin \overline{V_q}^\infty$ . Since  $\xi \in S$ , passing to a subsequence and relabelling, if needed, we may assume that for each  $\nu \geq 1$ , there exists  $r_\nu \in (0, T_\nu)$  such that  $\gamma_\nu(r_\nu) \rightarrow \xi$  as  $\nu \rightarrow \infty$ . Since  $\xi \in \partial_{\text{visi}}\Omega$  and  $\xi \notin \overline{V_q}^\infty$ , there exists a neighbourhood  $U_\xi$  of  $\xi$  such that  $U_\xi$  is relatively compact,  $\overline{U_\xi} \cap \overline{V_q}^\infty = \emptyset$ , and such that every pair of distinct points  $\xi_1, \xi_2 \in U_\xi \cap \partial\Omega$  satisfies the visibility condition with respect to  $K_\Omega$ . Also, fix neighbourhoods  $W_\xi, V_\xi$  of  $\xi$  such that  $W_\xi \Subset V_\xi \Subset U_\xi$ . We may assume that  $\gamma_\nu(r_\nu) \in W_\xi \cap \Omega$  for all  $\nu \geq 1$ . Let  $s_\nu > r_\nu$  and  $y'_\nu = \gamma_\nu(s_\nu)$  be such that  $y'_\nu \in (U_\xi \setminus \overline{W_\xi}) \cap \Omega$ , and such that the image of  $\tilde{\gamma}_\nu := \gamma_\nu|_{[r_\nu, s_\nu]}$  is contained in  $V_\xi$ . By Lemma 2.9 (and relabeling subsequences as usual) we have

$$\lim_{\nu \rightarrow \infty} \sup_{z \in \langle \tilde{\gamma}_\nu \rangle} d_h(z, \partial\Omega) = 0. \quad (4.1)$$

Therefore, there exists a sequence  $(\nu_j)_{j \geq 1} \subset \mathbb{N}$  and a point  $\eta \in \overline{V_\xi} \cap \partial\Omega \subseteq U_\xi \cap \partial\Omega$  such that  $y'_{\nu_j} \rightarrow \eta$ . By construction,  $\xi \neq \eta$ . Now, since  $U_\xi \cap \partial\Omega$  satisfies the visibility condition with respect to  $K_\Omega$ , there exists a compact set  $L \subseteq \Omega$  such that for every  $j \geq 1$  we can find  $\tilde{z}_{\nu_j} \in \langle \tilde{\gamma}_{\nu_j} \rangle$  with  $\tilde{z}_{\nu_j} \in L$ , which contradicts (4.1). This establishes (a).

It is evident that owing to our definitions, the entire argument above would be valid if we fix  $\lambda = 1$  and take  $\omega := \partial\Omega \setminus \partial_{\text{wvisi}}\Omega$ . Thus (b) follows.  $\square$

The above result gives us the following corollaries.

**Corollary 4.2.** *Let  $X$  be a complex manifold and  $\Omega$  be a Kobayashi hyperbolic domain. Then, the following are equivalent:*

- (a)  $\partial\Omega$  is visible.
- (b)  $\Omega$  is a visibility domain subordinate to any admissible compactification of  $\overline{\Omega}$ .
- (c)  $\Omega$  is a visibility domain subordinate to some admissible compactification of  $\overline{\Omega}$ .

*Proof.* Let  $\partial\Omega$  be visible. Then, every point in  $\partial\Omega$  is a visible point of  $\Omega$ . Therefore, (a) implies (b) follows from Theorem 4.1-(a). That (b) implies (c) is trivial. Finally, (c) implies (a) follows by Definition 1.1.  $\square$

**Corollary 4.3.** *Let  $X$  be a complex manifold and  $\Omega$  be a Kobayashi hyperbolic domain. Then, the following are equivalent:*

- (a)  $\partial\Omega$  is weakly visible.
- (b)  $\Omega$  is a weak visibility domain subordinate to any admissible compactification of  $\overline{\Omega}$ .
- (c)  $\Omega$  is a weak visibility domain subordinate to some admissible compactification of  $\overline{\Omega}$ .

The proof of the above is analogous to that of Corollary 4.2.

Note that, in view of the above corollaries, Theorem 1.16 follows immediately.

Finally, we can provide a proof of Proposition 1.8.

*The proof of Proposition 1.8.* If possible let  $\partial\Omega \setminus \partial_{\text{visi}}\Omega$  be a non-empty, totally disconnected set. Then, by Theorem 4.1 and Corollary 4.2, we have  $\partial\Omega$  is visible. Therefore, by Definition 1.7, it follows that  $\partial\Omega = \partial_{\text{visi}}\Omega$ , which contradicts the fact that  $\partial\Omega \setminus \partial_{\text{visi}}\Omega$  is non-empty. Similarly, we can prove that  $\partial\Omega \setminus \partial_{\text{wvisi}}\Omega$  cannot be a non-empty, totally disconnected set.  $\square$

## 5. SUFFICIENT CONDITIONS FOR VISIBILITY

In this section, we shall present several sufficient conditions for a domain  $\Omega \subsetneq X$ ,  $\Omega$  and  $X$  as in the previous sections, such that  $\partial\Omega$  is visible. We will first begin with a sufficient condition that is inspired by a construction in [9] — i.e., involving the notion of a “local Goldilocks point” alluded to in Section 1.2. Thereafter, we will provide proofs of the results stated in Section 1.2.

We first need a few definitions and observations.

**Definition 5.1.** Let  $X$  be a complex manifold and  $\Omega \subsetneq X$  be a Kobayashi hyperbolic domain. Fix  $p \in \partial\Omega$ . Let  $(U, \varphi)$  be a holomorphic chart of  $X$  centered at  $p$ . We define

$$M(r; \varphi, U) := \sup \left\{ \frac{1}{k_{\Omega}(x; v)} : v \in T_x^{(1,0)}(\Omega) \text{ such that } \|\varphi'(x)v\| = 1 \right. \\ \left. \text{and } x \in U \cap \Omega \text{ such that } d_{\text{Euc}}(\varphi(x), \varphi(U \cap \partial\Omega)) \leq r \right\}.$$

The function  $M(\cdot; \varphi, U)$  depends on the choice of charts, which could potentially pose difficulties in any analysis purportedly about  $\Omega$  but which relies on estimates for  $M(\cdot; \varphi, U)$ . The following lemma addresses this issue. In this paper, we abbreviate  $d_{\text{Euc}}(z, S)$  as  $\delta_S(z)$ , for any  $z \in \mathbb{C}^n$  and  $S \subseteq \mathbb{C}^n$ . The notation  $\|\cdot\|$  denotes the Euclidean norm on  $\mathbb{C}^n$ .

**Lemma 5.2.** *Let  $X$ ,  $\Omega$ , and  $p \in \partial\Omega$  be as in Definition 5.1. Let  $(U, \varphi)$  and  $(V, \psi)$  be two holomorphic charts of  $X$  centered at  $p$ . Then, there exists a neighbourhood  $W$  of  $p$  contained in  $U \cap V$  and constants  $R, \beta \geq 1$  which depend only on  $\varphi$  and  $\psi$  such that*

$$M(r; \varphi, W) \geq (1/\beta)M(Rr; \psi, V) \quad \text{and} \quad \delta_{\psi(V \cap \partial\Omega)}(\psi(x)) \leq R\delta_{\varphi(W \cap \partial\Omega)}(\varphi(x))$$

for every  $r > 0$  sufficiently small and for every  $x \in W \cap \Omega$ .

*Proof.* Let  $W$  be a neighbourhood of  $p$  such that  $\overline{W} \subseteq U \cap V$ , and such that  $\varphi(W)$  is a polydisk. Let  $x \in W \cap \Omega$  and  $r > 0$  be such that  $\delta_{\varphi(W \cap \partial\Omega)}(\varphi(x)) \leq r$ . Then, there exists  $\xi \in W \cap \partial\Omega$  such that  $\|\varphi(x) - \varphi(\xi)\| \leq r$ . Since  $\psi \circ (\varphi|_W)^{-1}$  is the restriction of a biholomorphic map on  $\varphi(U \cap V)$  and  $\overline{W}$  is compact, there exists a constant  $R \geq 1$  such that the operator norm  $\|\psi \circ ((\varphi|_W)^{-1})'(\zeta)\|_{\text{OP}} \leq R$  for all  $\zeta \in \varphi(W)$ . In what follows, we shall abbreviate  $\psi \circ (\varphi|_W)^{-1}$  to  $\psi \circ \varphi^{-1}$  for simplicity of notation. Since the line segment joining  $\varphi(x)$  and  $\varphi(\xi)$  lies in  $\varphi(W)$ , we have

$$\begin{aligned} \|\psi(x) - \psi(\xi)\| &= \|(\psi \circ \varphi^{-1})(\varphi(x)) - (\psi \circ \varphi^{-1})(\varphi(\xi))\| \\ &\leq \sup_{\zeta \in \varphi(W)} \|(\psi \circ \varphi^{-1})'(\zeta)\|_{\text{OP}} \|\varphi(x) - \varphi(\xi)\| \\ &\leq R\delta_{\varphi(W \cap \partial\Omega)}(\varphi(x)). \end{aligned} \tag{5.1}$$

It follows that  $\delta_{\psi(V \cap \partial\Omega)}(\psi(x)) \leq R\delta_{\varphi(W \cap \partial\Omega)}(\varphi(x))$ . Now, there exists a constant  $\beta \geq 1$  such that for any  $z \in W \cap \Omega$  and every  $v \in T_z^{(1,0)}\Omega$  that satisfies  $\|\varphi'(z)v\| = 1$ , we have  $1/\beta \leq \|\psi'(z)v\| \leq \beta$ .

As  $k_\Omega(z; c \cdot) = |c|k_\Omega(z; \cdot)$  on  $T_z^{(1,0)}(\Omega)$  for any  $z \in \Omega$  and any  $c \in \mathbb{C}$ , and as (5.1) holds true for any  $x$  in  $W \cap \Omega$  such that  $\delta_\varphi(W \cap \partial\Omega)(\varphi(x)) \leq r$ , it follows by definition that

$$M(r; \varphi, W) \geq (1/\beta)M(Rr; \psi, V).$$

Hence the proof.  $\square$

**Definition 5.3.** Let  $X$  be a complex manifold and  $\Omega \subsetneq X$  be a Kobayashi hyperbolic domain. A boundary point  $p \in \partial\Omega$  is called a *local Goldilocks point* if there exists a holomorphic chart  $(U, \varphi)$  centered at  $p$  and a neighbourhood  $U$  of  $p$  such that  $p \in U \subseteq \mathcal{U}$  such that

(a) for some  $\varepsilon > 0$  we have

$$\int_0^\varepsilon \frac{1}{r} M(r; \varphi, U) dr < \infty,$$

(b) for each  $z_0 \in \Omega$  there exist constants  $C, \alpha > 0$  (which depend on  $z_0, U, \varphi$ ) such that for all  $x \in U \cap \Omega$

$$K_\Omega(z_0, x) \leq C + \alpha \log \frac{1}{\delta_{\varphi(U \cap \partial\Omega)}(\varphi(x))}.$$

Let  $\partial_{\text{lg}}\Omega \subseteq \partial\Omega$  denote the set of all local Goldilocks points.

*Remark 5.4.* Note that the function  $M(\cdot; \varphi, U)$  is a non-negative, monotone increasing function, owing to which the integral in (a) makes sense. We see that, in view of Lemma 5.2, the status of whether a point  $p \in \partial\Omega$  is a local Goldilocks point is independent of the choice of chart occurring in Definition 5.3.

**Lemma 5.5.** Let  $X$  be a complex manifold and  $\Omega \subsetneq X$  be a Kobayashi hyperbolic domain. Let  $p \in \partial_{\text{lg}}\Omega$ . Let  $(U, \varphi)$  be a holomorphic chart centered at  $p$  that satisfies the conditions in Definition 5.3 and such that  $U$  is relatively compact. Fix  $\lambda \geq 1$  and  $\kappa \geq 0$ . Suppose for each  $\nu \geq 1$ ,  $\gamma_\nu : [0, r_\nu] \rightarrow \Omega$  is a  $(\lambda, \kappa)$ -almost-geodesic such that  $\gamma_\nu([0, r_\nu]) \subseteq U$ , and such that  $\gamma_\nu(0) \rightarrow p' \in U \cap \partial\Omega$ ,  $\gamma_\nu(r_\nu) \rightarrow q' \in U \cap \partial\Omega$  as  $\nu \rightarrow \infty$  with  $p' \neq q'$ . Let  $(\varphi \circ \gamma_\nu)_{\nu \geq 1}$  converge locally uniformly to a curve  $\sigma : [0, M] \rightarrow \overline{\varphi(U)}$ , for some  $M \in (0, \infty]$ . Then,  $\sigma$  is non-constant.

*Proof.* Define  $\sigma_\nu := \varphi \circ \gamma_\nu$  for every  $\nu \geq 1$  and  $Z := \varphi(U \cap \partial\Omega)$ . Fix  $z_0 \in U \cap \Omega$ . From part (b) of Definition 5.3 there exist constants  $C, \alpha > 0$  (which depend on  $z_0, U, \varphi$ ) such that for all  $x \in U \cap \Omega$

$$K_\Omega(z_0, x) \leq C + \alpha \log \frac{1}{\delta_{\varphi(U \cap \partial\Omega)}(\varphi(x))}.$$

Therefore, since each  $\gamma_\nu$  is a  $(\lambda, \kappa)$ -almost-geodesic, for every  $t \in [0, r_\nu]$ , every  $\nu \geq 1$ , we have

$$\begin{aligned} \frac{1}{\lambda}|t| - \kappa &\leq K_\Omega(\gamma_\nu(0), \gamma_\nu(t)) \leq K_\Omega(\gamma_\nu(0), z_0) + K_\Omega(z_0, \gamma_\nu(t)) \\ &\leq 2C + \alpha \log \frac{1}{\delta_Z(\sigma_\nu(0))\delta_Z(\sigma_\nu(t))}. \end{aligned}$$

Now, at this stage we can proceed exactly as in the proof of [9, Lemma 6.1] and hence, the result follows.  $\square$

We are now in a position to give a sufficient condition for visibility. This result is inspired by Theorem 1.4 in [9] and (in view of Corollary 4.2) is the exact analogue of the latter theorem.

**Theorem 5.6.** Let  $X$  be a complex manifold and  $\Omega \subsetneq X$  be a Kobayashi hyperbolic domain. Suppose the set  $\partial\Omega \setminus \partial_{\text{lg}}\Omega$  is totally disconnected. Then,  $\partial\Omega$  is visible.



*Proof.* Fix a Hermitian metric  $h$  on  $X$  and let  $d_h$  be the distance induced by  $h$  on  $X$ .

**Claim.** *Every local Goldilocks point is a visible point.*

*Proof of Claim.* Let  $p \in \partial_{\text{lg}}\Omega$ . Therefore, there exists a holomorphic chart  $(U, \varphi)$  centered at  $p$  that satisfies the conditions in Definition 5.3. Without loss of generality, we can assume that  $U$  is relatively compact and  $\tilde{U} := \varphi(U)$  is bounded. Fix  $\xi, \eta \in U \cap \partial\Omega$  such that  $\xi \neq \eta$ . We also fix  $\lambda \geq 1$  and  $\kappa \geq 0$ . Let  $V_\xi, V_\eta$  be two neighbourhoods of  $\xi$  and  $\eta$  respectively such that  $\overline{V_\xi} \cap \overline{V_\eta} = \emptyset$  and  $\overline{V_\xi} \subseteq U$ . Let  $W_\xi$  be a neighbourhood of  $\xi$  such that  $\overline{W_\xi} \subseteq V_\xi$ . For any pair  $x, y \in \Omega$ , let  $AG_\Omega(x, y)$  have the same meaning as in Section 3.

We will show that there exists a compact set  $K \subset \Omega$  such that for every  $\gamma \in AG_\Omega(x, y)$ , every  $x \in W_\xi \cap \Omega$ , and every  $y \in V_\eta \cap \Omega$ , we have  $\langle \gamma \rangle \cap K \neq \emptyset$ . If possible, let this assertion be false. Let  $(K_\nu)_{\nu \geq 1}$  be an exhaustion by compacts of  $\Omega$ . Then, there exist sequences  $(x_\nu)_{\nu \geq 1} \subset W_\xi \cap \Omega$ ,  $(y_\nu)_{\nu \geq 1} \subset V_\eta \cap \Omega$ , and  $(\lambda, \kappa)$ -almost-geodesics  $\gamma_\nu$  in  $AG_\Omega(x_\nu, y_\nu)$ , for each  $\nu \geq 1$ , such that  $\langle \gamma_\nu \rangle \cap K_\nu = \emptyset$ . Fix neighbourhoods  $W_\xi^1, W_\xi^2$  of  $\xi$  such that  $W_\xi \subseteq W_\xi^1 \subseteq W_\xi^2 \subseteq V_\xi$ . Let  $r_\nu > 0$  and  $y'_\nu = \gamma_\nu(r_\nu)$  be such that  $y'_\nu \in (V_\xi \setminus \overline{W_\xi^1}) \cap \Omega$ , and such that the image of  $\tilde{\gamma}_\nu := \gamma_\nu|_{[0, r_\nu]}$  is contained in  $W_\xi^2$ . By Lemma 2.9, and as we can relabel the subsequences given by it, there exist a point  $p' \in \overline{W_\xi} \cap \partial\Omega$  such that  $x_\nu \rightarrow p'$  and

$$\lim_{\nu \rightarrow \infty} \sup_{z \in \langle \tilde{\gamma}_\nu \rangle} d_h(z, \partial\Omega) = 0. \quad (5.2)$$

Therefore, there exists a sequence  $(\nu_j)_{j \geq 1} \subset \mathbb{N}$  and a point  $q' \in \overline{W_\xi^2} \cap \partial\Omega \subseteq V_\xi \cap \partial\Omega$  such that  $\gamma_{\nu_j}(r_{\nu_j}) \rightarrow q'$ . By construction,  $p' \neq q'$ . Define  $\sigma_j := \varphi \circ \tilde{\gamma}_{\nu_j}$  for every  $j \geq 1$  and let  $Z := \varphi(U \cap \partial\Omega)$ .

Since  $p$  is a local Goldilocks point, from part (a) of Definition 5.3 it follows that  $M(r; \varphi, U) \rightarrow 0$  as  $r \rightarrow 0^+$ . Let  $\delta > 0$  be such that whenever  $r \in (0, \delta)$ , we have  $0 \leq M(r; \varphi, U) < 1$ . Therefore, as  $k_\Omega(z; c \cdot) = |c|k_\Omega(z; \cdot)$  on  $T_z^{(1,0)}(\Omega)$  for  $z \in \Omega$  and  $c \in \mathbb{C}$ , for every  $v \in T_z^{(1,0)}(\Omega)$ , every  $z \in U \cap \Omega$  that satisfies  $\delta_Z(\varphi(z)) < \delta$ , we have  $k_\Omega(z; v) \geq \|\varphi'(z)v\|$ . Now, let  $V := U \cap \{z \in \Omega : \delta_Z(\varphi(z)) \geq \delta\}$ . Since  $U$  is relatively compact,  $\overline{V} \subseteq \Omega$  and  $\overline{V}$  is compact. Therefore, by Result 2.5, there exists a constant  $L_0 > 0$  such that for every  $v \in T_z^{(1,0)}(\Omega)$ , every  $z \in V$ , we have  $k_\Omega(z; v) \geq L_0 \|\varphi'(z)v\|$ . Let  $L := \min\{L_0, 1\}$ . Since  $\tilde{\gamma}_\nu$  is a  $(\lambda, \kappa)$ -almost-geodesic, from above it follows that

$$\|\sigma'_j(t)\| = \|\varphi'(\tilde{\gamma}_{\nu_j}(t))\tilde{\gamma}'_{\nu_j}(t)\| \leq \frac{1}{L}k_\Omega(\tilde{\gamma}_{\nu_j}(t); \tilde{\gamma}'_{\nu_j}(t)) \leq \lambda/L,$$

for almost every  $t \in [0, r_{\nu_j}]$ , every  $j \geq 1$ . Therefore, for every  $s, r \in [0, r_{\nu_j}]$ , every  $j \geq 1$  we have

$$\begin{aligned} \sigma_j(r) &= \sigma_j(s) + \int_s^r \sigma'_j(t) dt \\ \implies \|\sigma_j(r) - \sigma_j(s)\| &\leq \frac{\lambda}{L}|r - s|. \end{aligned}$$

Therefore,  $\sigma_j$  is  $\lambda/L$ -Lipschitz for every  $j \geq 1$ . Therefore, since  $\langle \sigma_j \rangle \subseteq \tilde{U}$ , by Arzelá–Ascoli Theorem, passing to a subsequence and relabelling, if needed, we can assume that  $\sigma_j$  converges locally uniformly to a curve  $\sigma : [0, M) \rightarrow \tilde{U}$  (where  $M \in (0, \infty]$ ). We will show that for almost every  $t \in [0, r_{\nu_j}]$ , every  $j \geq 1$

$$\|\sigma'_j(t)\| \leq \lambda M(\delta_Z(\sigma_j(t)); \varphi, U). \quad (5.3)$$

Now, (5.3) is true when  $\|\sigma'_j(t)\| = 0$ . Let  $\|\sigma'_j(t)\| \neq 0$ . Since  $\tilde{\gamma}_{\nu_j}$  is a  $(\lambda, \kappa)$ -almost-geodesic, for almost every  $t \in [0, r_{\nu_j}]$ , every  $j \geq 1$ , we have

$$\begin{aligned} k_{\Omega}(\tilde{\gamma}_{\nu_j}; \tilde{\gamma}'_{\nu_j}(t)) &\leq \lambda \\ \implies \|\sigma'_j(t)\| &\leq \frac{\lambda}{k_{\Omega}\left(\tilde{\gamma}_{\nu_j}(t); \frac{\tilde{\gamma}'_{\nu_j}(t)}{\|\sigma'_j(t)\|}\right)} \leq \lambda M(\delta_Z(\sigma_j(t)); \varphi, U). \end{aligned}$$

Now, by (5.2) we may assume, passing to a subsequence and relabelling, if needed, that  $\sup_{\zeta \in \langle \sigma_j \rangle} \delta_Z(\zeta) \searrow 0$  as  $j \rightarrow \infty$ . From this, it follows that  $\lim_{j \rightarrow \infty} \sup_{\zeta \in \langle \sigma_j \rangle} M(\delta_Z(\zeta); \varphi, U) = 0$  since  $M(\cdot; \varphi, U)$  is monotone decreasing. Hence, for every  $s, r \in [0, M]$  we have

$$\begin{aligned} \|\sigma(s) - \sigma(r)\| &= \lim_{j \rightarrow \infty} \|\sigma_j(s) - \sigma_j(r)\| \leq \limsup_{j \rightarrow \infty} \int_s^r \|\sigma'_j(t)\| dt \\ &\leq \frac{\lambda}{L} \limsup_{j \rightarrow \infty} \int_s^r M(\delta_Z(\sigma_j(t)); \varphi, U) dt = 0. \end{aligned}$$

Therefore,  $\sigma$  is constant, which contradicts Lemma 5.5. Hence, the claim.  $\blacktriangleleft$

By the above claim, since  $\partial\Omega \setminus \partial_{\mathbb{g}}\Omega$  is totally disconnected, we conclude that  $\partial\Omega \setminus \partial_{\text{visi}}\Omega$  is totally disconnected. Therefore, by Theorem 4.1 and Corollary 4.2, we have the result.  $\square$

At this stage, we can give proofs of some more sufficient conditions for visibility. These are the results stated in Section 1.2. We begin with proving Theorem 1.10.

*The proof of Theorem 1.10.* Fix a Hermitian metric  $h$  on  $X$  and let  $d_h$  be the distance induced by  $h$  on  $X$ . Since  $\Omega \subsetneq X$  is relatively compact, by [25, Result 3.1, Proposition 2.4], we can conclude that  $\Omega$  is a hyperbolically imbedded domain.

**Claim.** *Each point  $p$  in  $\partial_V\Omega$  is a visible point of  $\partial\Omega$ .*

*Proof of Claim.* Let  $p \in \partial_V\Omega$ . Therefore, there exists a holomorphic chart  $(U, \varphi)$  centered at  $p$  such that  $\varphi(U)$  is bounded,  $\varphi(U \cap \Omega)$  is a Kobayashi hyperbolic domain, and such that  $\partial(\varphi(U \cap \Omega))$  is visible. Let  $G := \varphi(U \cap \Omega)$ . Without loss of generality, we may assume that  $U$  is relatively compact. Fix  $\xi, \eta \in U \cap \partial\Omega$  such that  $\xi \neq \eta$ . Therefore,  $\varphi(\xi), \varphi(\eta) \in \partial G$  such that  $\xi' = \varphi(\xi) \neq \varphi(\eta) = \eta'$ . We also fix  $\lambda \geq 1$  and  $\kappa \geq 0$ . Since  $\partial G$  is visible, the pair  $(\xi', \eta')$  satisfies the visibility condition with respect to  $K_G$ . Therefore, since  $\varphi$  is a biholomorphism on  $U$  and  $\xi, \eta \in \partial(U \cap \Omega)$ , the pair  $(\xi, \eta)$  satisfies the visibility condition with respect to  $K_{U \cap \Omega}$ . Therefore,  $\partial\Omega$  is locally visible at  $p$ .

Since  $\Omega$  is a hyperbolically imbedded domain in  $X$ , there exists a neighbourhood  $V_{\xi} \subseteq U$  of  $\xi$  in  $X$  such that  $K_{\Omega}(V_{\xi} \cap \Omega, \Omega \setminus U) := \delta_{\xi} > 0$  and that  $\eta \notin \overline{V_{\xi}}$ . Since  $\eta \notin \overline{V_{\xi}}$ , there exists a neighbourhood  $V_{\eta}$  of  $\eta$  in  $X$  such that  $V_{\eta}$  is relatively compact, and such that  $\overline{V_{\eta}} \cap \overline{V_{\xi}} = \emptyset$ . Let  $W_{\xi}$  be a neighbourhood of  $\xi$  in  $X$  such that  $\overline{W_{\xi}} \subseteq V_{\xi}$ . For any pair  $x, y \in \Omega$  let  $AG_{\Omega}(x, y)$  have the same meaning as in Section 3. It suffices to show that there exists a compact set  $K \subset \Omega$  such that for every  $\gamma \in AG_{\Omega}(x, y)$ , every  $x \in W_{\xi} \cap \Omega$ , and every  $y \in V_{\eta} \cap \Omega$ , we have  $\langle \gamma \rangle \cap K \neq \emptyset$ . At this stage, the existence of such a  $K$  follows by the same argument as in the proof of Claim 1 of the proof of Theorem 1.5. Therefore,  $p$  is a visible point of  $\partial\Omega$  and hence, the claim.  $\blacktriangleleft$

By the above claim, since  $\partial\Omega \setminus \partial_V\Omega$  is totally disconnected, we conclude that  $\partial\Omega \setminus \partial_{\text{visi}}\Omega$  is totally disconnected. Therefore, by Theorem 4.1 and Corollary 4.2, part (i) follows.

It is evident that owing to our definitions, the entire argument above would be valid if we fix  $\lambda = 1$  and replace the reference to Theorem 1.5 with Theorem 1.6, and substitute  $\partial\Omega \setminus \partial_V\Omega$  with  $\partial\Omega \setminus \partial_{WV}\Omega$ . Thus, (ii) follows.  $\square$

Next, we will provide a proof of Theorem 1.11. This result is inspired by Banik's result [6, Theorem 3.1]. However, there are two differences between the hypotheses of our result and [6, Theorem 3.1]. Firstly, our setting is domains in arbitrary complex manifolds that are not necessarily relatively compact. Secondly, instead of removing a compact set contained in the "old" domain, we remove a closed (not necessarily compact) totally disconnected set. Consequently, there are significant differences between the proofs of Theorem 1.11 and [6, Theorem 3.1].

*The proof of Theorem 1.11.* Fix a Hermitian metric  $h$  on  $X$  and let  $d_h$  be the distance induced by  $h$  on  $X$ . Since  $\Omega$  is a Kobayashi hyperbolic domain and  $\Omega \setminus S$  is connected, it follows that  $\Omega \setminus S$  is a Kobayashi hyperbolic domain. Since  $S$  is closed and  $\mathcal{H}^{2n-2}(S) = 0$ , by [24, Theorem 3.2.22] we have

$$K_{\Omega \setminus S} = K_{\Omega}|_{(\Omega \setminus S) \times (\Omega \setminus S)}. \quad (5.4)$$

**Claim.** *Each point  $p$  in  $\partial\Omega \setminus S$  is a visible point of  $\partial(\Omega \setminus S)$ .*

*Proof of Claim.* Let  $p \in \partial\Omega \setminus S \subseteq \partial(\Omega \setminus S)$ . Since  $S$  is closed, there exists a neighbourhood  $U_p$  of  $p$  in  $X$  such that  $U_p$  is relatively compact and  $\overline{U_p} \cap S = \emptyset$ . Therefore,  $U_p \cap \partial(\Omega \setminus S) = U_p \cap \partial\Omega$ . Fix  $\lambda \geq 1$  and  $\kappa \geq 0$ . We also fix  $\xi, \eta \in U_p \cap \partial(\Omega \setminus S)$  such that  $\xi \neq \eta$ . There exist two neighbourhoods  $V_\xi, V_\eta$  of  $\xi, \eta$  respectively in  $X$  such that  $\overline{V_\xi} \cap \overline{V_\eta} = \emptyset$  and such that  $\overline{V_\xi}, \overline{V_\eta} \subseteq U_p$ . Let  $W_\xi$  be a neighbourhood of  $\xi$  in  $X$  such that  $\overline{W_\xi} \subseteq V_\xi$ . For any pair  $x, y \in \Omega \setminus S$ , we define the set

$$AG_{\Omega \setminus S}(x, y) := \{\gamma : [0, T] \longrightarrow \Omega \setminus S : \\ \gamma \text{ is a } (\lambda, \kappa)\text{-almost-geodesic with respect to } K_{\Omega \setminus S} \text{ joining } x \text{ and } y\},$$

where  $\lambda$  and  $\kappa$  are as fixed above.

We shall show that there exists a compact set  $K \subset \Omega \setminus S$  such that for every  $\gamma \in AG_{\Omega \setminus S}(x, y)$ , every  $x \in W_\xi \cap (\Omega \setminus S)$ , and every  $y \in V_\eta \cap (\Omega \setminus S)$ , we have  $\langle \gamma \rangle \cap K \neq \emptyset$ . If possible, let this assertion be false. Let  $(K_\nu)_{\nu \geq 1}$  be an exhaustion by compacts of  $\Omega \setminus S$ . Then, there exist sequences  $(x_\nu)_{\nu \geq 1} \subset W_\xi \cap (\Omega \setminus S)$ ,  $(y_\nu)_{\nu \geq 1} \subset V_\eta \cap (\Omega \setminus S)$ , and  $(\lambda, \kappa)$ -almost-geodesics  $\gamma_\nu$  in  $AG_{\Omega \setminus S}(x_\nu, y_\nu)$ , for each  $\nu \geq 1$ , such that  $\langle \gamma_\nu \rangle \cap K_\nu = \emptyset$ . At this point, we observe that by (5.4) and the fact that  $k_{\Omega \setminus S} \geq k_\Omega$ , for each  $\nu$

$$\gamma_\nu|_{I_\nu} \text{ is a } (\lambda, \kappa)\text{-almost-geodesic with respect to } K_\Omega,$$

where  $I_\nu$  is any interval contained in  $\text{dom}(\gamma_\nu)$ . Furthermore

$$U_p \cap \partial(\Omega \setminus S) = U_p \cap \partial\Omega \quad \text{and} \quad V_\eta \cap (\Omega \setminus S) = V_\eta \cap \Omega.$$

These two facts combined with an argument similar to the proof of the Claim made while proving Theorem 4.1 leads, essentially, to a similar contradiction. This proves the present claim.  $\blacktriangleleft$

Since  $S \subsetneq X$  is closed,  $\partial(\Omega \setminus S) \setminus (\partial\Omega \setminus S) \subseteq S$ . Therefore, since  $S$  is totally disconnected, from the above claim it follows that  $\partial(\Omega \setminus S) \setminus \partial_{\text{visi}}(\Omega \setminus S)$  is totally disconnected. Therefore, by Theorem 4.1 and Corollary 4.2, we can conclude that  $\partial(\Omega \setminus S)$  is visible.  $\square$

We conclude this section by presenting a proof of Proposition 1.13.

*The proof of Proposition 1.13.* Since  $\Omega$  is a simply connected domain, there exists a biholomorphic map  $f : \mathbb{D} \longrightarrow \Omega$ . Since  $\Omega$  is a Jordan domain, by [10, Theorem 4.3.3] we can conclude that  $f$  can be extended to a homeomorphism  $\tilde{f} : \overline{\mathbb{D}} \longrightarrow \overline{\Omega}^\infty$ , where  $\overline{\Omega}^\infty$  is the closure of  $\Omega$  viewed as a domain in  $\widehat{\mathbb{C}}$ . Fix  $p, q \in \partial\Omega \subseteq \partial_\infty\Omega$  such that  $p \neq q$ . We also fix  $\lambda \geq 1$  and  $\kappa \geq 0$ . Let  $p' = \tilde{f}^{-1}(p) \neq q' = \tilde{f}^{-1}(q)$ . So,  $p', q' \in \partial\mathbb{D}$ . Since  $\partial\mathbb{D}$  is visible,  $(p', q')$  satisfies

the visibility condition with respect to  $K_{\mathbb{D}}$ . Therefore, there exist neighbourhoods  $\widetilde{U}_{p'}$  of  $p'$  and  $\widetilde{U}_{q'}$  of  $q'$  in  $\mathbb{C}$  such that  $\overline{\widetilde{U}_{p'}} \cap \overline{\widetilde{U}_{q'}} = \emptyset$  and a compact set  $\widetilde{K}$  of  $\mathbb{D}$  such that the image of each  $(\lambda, \kappa)$ -almost-geodesic  $\tilde{\sigma} : [0, T] \rightarrow \mathbb{D}$  with  $\tilde{\sigma}(0) \in \widetilde{U}_{p'}$  and  $\tilde{\sigma}(T) \in \widetilde{U}_{q'}$  intersects  $\widetilde{K}$ . Consider two neighbourhoods  $U_p$  of  $p$  and  $U_q$  of  $q$  such that  $\overline{U_p} \cap \overline{U_q} = \emptyset$  and such that

$$\tilde{f}(\overline{\widetilde{U}_{p'}} \cap \overline{\mathbb{D}}) = \overline{U_p} \cap \overline{\Omega} \quad \text{and} \quad \tilde{f}(\overline{\widetilde{U}_{q'}} \cap \overline{\mathbb{D}}) = \overline{U_q} \cap \overline{\Omega}.$$

Since  $f$  is a biholomorphism, the pre-image (under  $f$ ) of each  $(\lambda, \kappa)$ -almost-geodesic in  $\Omega$  is a  $(\lambda, \kappa)$ -almost-geodesic in  $\mathbb{D}$ . Thus, by construction, the image of each  $(\lambda, \kappa)$ -almost-geodesic  $\sigma : [0, r] \rightarrow \Omega$  with  $\sigma(0) \in U_p$  and  $\sigma(r) \in U_q$  intersects the compact set,  $f(\widetilde{K})$ . Hence the result.  $\square$

## 6. AN APPLICATION: A WOLFF–DENJOY-TYPE THEOREM

Recall that the classical Wolff–Denjoy Theorem is as follows:

**Result 6.1** ([18, 30]). *Suppose  $f : \mathbb{D} \rightarrow \mathbb{D}$  is a holomorphic map then either:*

- (a)  *$f$  has a fixed point in  $\mathbb{D}$ ; or*
- (b) *there exists a point  $p \in \partial\mathbb{D}$  such that  $\lim_{\nu \rightarrow \infty} f^\nu(x) = p$  for any  $x \in \mathbb{D}$ , this convergence being uniform on compact subsets of  $\mathbb{D}$ .*

There is a rich history of generalisations of Result 6.1. However, until recently, these featured various convex domains in place of  $\mathbb{D}$ , for instance: unit ball in  $\mathbb{C}^n$  ( $n \geq 2$ ) [20], bounded strongly convex domains in  $\mathbb{C}^n$  [1], bounded convex domains in  $\mathbb{C}^n$  with progressively weaker assumptions [4], bounded strictly convex domains in a complex Banach space [15]. To the best of my knowledge, the main result in [21] is the first result generalising Result 6.1 to a class of contractible domains in  $\mathbb{C}^n$ ,  $n \geq 2$ , that includes domains that are not convex. Wolff–Denjoy-type theorems are now known for certain metric spaces where  $\partial\mathbb{D}$  is replaced by some abstract boundary — see, for instance [23, 22]. The metrical approaches to Wolff–Denjoy-type theorems motivated several generalisations of Result 6.1 to visibility domains (in the sense of [7, 9]): see, for instance, [8, 7, 9]. Our next theorem is in the spirit of the last three papers, extending the results therein to domains in arbitrary complex manifolds.

**Theorem 6.2.** *Let  $X$  be a complex manifold and  $\Omega \subsetneq X$  be a taut domain such that  $\partial\Omega$  is weakly visible. Let  $\overline{\Omega}^\infty$  be some admissible compactification of  $\overline{\Omega}$ . Suppose  $f : \Omega \rightarrow \Omega$  is a holomorphic map. Then, one of the following holds:*

- (a) *for every  $x \in \Omega$ , the set  $\{f^\nu(x) : \nu \geq 1\}$  is relatively compact in  $\Omega$ ; or*
- (b) *there exists  $p \in \partial_\infty\Omega$  such that for every  $x \in \Omega$ ,  $f^\nu(x)$  converges locally uniformly to  $p$  as  $\nu \rightarrow \infty$ .*

In the statement of the above theorem, we have implicitly used the fact that when  $\Omega$  is taut, then it is Kobayashi hyperbolic.

If  $\Omega$  is not contractible, there is no reason to expect  $f : \Omega \rightarrow \Omega$  as in the above theorem to have a fixed point. In fact, in [3], Abate–Heinzner construct *contractible* domains  $\Omega \subsetneq \mathbb{C}^n$ ,  $n \geq 2$ , and holomorphic maps  $f : \Omega \rightarrow \Omega$  that do not obey the dichotomy in Result 6.1. Thus, in the generality we are interested in, (a) of Theorem 6.2 is the natural substitute of (a) of Result 6.1.

In case Theorem 6.2 seems rather abstract, owing to the involvement of an abstract admissible compactification of  $\overline{\Omega}$ , let us state a theorem with a more explicit hypothesis. Wolff–Denjoy-type theorems are generally stated for bounded domains, but the following result shows that

the Wolff–Denjoy phenomenon extends to unbounded/non-relatively-compact domains (which is also seen in the metrical versions of the Wolff–Denjoy phenomenon mentioned above).

**Theorem 6.3.** *Let  $X$  be a non-compact complex manifold and  $\Omega \subsetneq X$  be a taut domain that is not relatively compact and such that  $\partial\Omega$  is weakly visible. Suppose  $f : \Omega \rightarrow \Omega$  is a holomorphic map. Then, one of the following holds:*

- (a) *for every  $x \in \Omega$ , the set  $\{f^\nu(x) : \nu \geq 1\}$  is relatively compact in  $\Omega$ ; or*
- (b) *either there exists  $p \in \partial\Omega$  such that for each  $x \in \Omega$ ,  $f^\nu(x) \rightarrow p$ , or, for each  $x \in \Omega$ , the sequence  $(f^\nu(x))_{\nu \geq 1}$  exits every compact subset of  $X$ . In the first case,  $(f^\nu)_{\nu \geq 1}$  converges locally uniformly to  $p$  while, in the second case,  $(f^\nu)_{\nu \geq 1}$  is compactly divergent viewed as a sequence of maps into  $X$ .*

Theorem 6.3 follows immediately from Theorem 6.2 by taking the admissible compactification of  $\bar{\Omega}$  to be the closure of  $\Omega$  in the one-point compactification of  $X$ .

The argument for Theorem 6.2 is similar to the argument in the proof of [7, Theorem 1.8]. However, as our setting is more general, we need two lemmas that are analogous to [7, Proposition 4.1] and [7, Theorem 4.3].

**Lemma 6.4.** *Let  $X$  be a complex manifold and  $\Omega \subsetneq X$  be a Kobayashi hyperbolic domain such that  $\partial\Omega$  is weakly visible. Let  $\bar{\Omega}^\infty$  be any admissible compactification of  $\bar{\Omega}$ . Suppose  $(f_\nu)_{\nu \geq 1}$  is a compactly divergent sequence in  $\text{Hol}(\Omega, \Omega)$ . Then, there exists  $p \in \partial_\infty\Omega$  and a subsequence  $(f_{\nu_j})_{j \geq 1}$  such that for all  $x \in \Omega$*

$$\lim_{j \rightarrow \infty} f_{\nu_j}(x) = p.$$

*Proof.* By Corollary 4.3, we deduce that  $\Omega$  is a weak visibility domain subordinate to  $\bar{\Omega}^\infty$ . At this stage, the argument proceeds exactly as in the proof of [9, Lemma 11.1] with  $\partial\bar{\Omega}^{\text{End}}$  replaced by  $\partial_\infty\Omega$ , which works because charts are irrelevant to the latter proof. The proof of [9, Lemma 11.1] relies on the sequential compactness of the specific admissible compactification  $\bar{\Omega}^{\text{End}}$  and on examining maps defined on intervals of  $\mathbb{R}$ , which carry over verbatim to our case.  $\square$

**Lemma 6.5.** *Let  $X$  be a complex manifold and  $\Omega \subsetneq X$  be a Kobayashi hyperbolic domain such that  $\partial\Omega$  is weakly visible. Let  $\bar{\Omega}^\infty$  be any admissible compactification of  $\bar{\Omega}$ . Let  $z_0 \in \Omega$ . Suppose  $f : \Omega \rightarrow \Omega$  is a holomorphic map that satisfies*

$$\limsup_{\nu \rightarrow \infty} K_\Omega(f^\nu(z_0), z_0) = \infty.$$

*Then, there exists  $p \in \partial_\infty\Omega$  such that for every sequence  $(\mu_k)_{k \geq 1} \subset \mathbb{N}$  that satisfies  $\lim_{k \rightarrow \infty} K_\Omega(f^{\mu_k}(z_0), z_0) = \infty$ , we have*

$$\lim_{k \rightarrow \infty} f^{\mu_k}(z_0) = p.$$

*Proof.* From Corollary 4.3 it follows that  $\Omega$  is a weak visibility domain subordinate to  $\bar{\Omega}^\infty$ . At this stage, we can essentially follow the same argument as in the proof of [9, Lemma 11.2] with  $\partial\bar{\Omega}^{\text{End}}$  replaced by  $\partial_\infty\Omega$  — which works because the latter proof relies on the sequential compactness of the specific admissible compactification  $\bar{\Omega}^{\text{End}}$  and on the behaviour of  $(1, 1)$ -almost-geodesics.  $\square$

**Result 6.6** ([2, Theorem 2.4.3]). *Let  $X$  be a complex manifold such that  $X$  is taut. Suppose  $f : X \rightarrow X$  is a holomorphic map. Then, either of the following holds:*

- (a)  *$(f^\nu)_{\nu \geq 1}$  is relatively compact in  $\text{Hol}(X, X)$ ; or*

(b)  $(f^\nu)_{\nu \geq 1}$  is compactly divergent.

Now, we are in a position to give a proof of Theorem 6.2.

*The proof of Theorem 6.2.* Fix a Hermitian metric  $h$  on  $X$  and let  $d_h$  be the distance induced by  $h$  on  $X$ . Since  $\Omega$  is taut, from Result 6.6 it follows that either for every  $x \in \Omega$ , the set  $\{f^\nu(x) : \nu \geq 1\}$  is relatively compact in  $\Omega$  or  $(f^\nu)_{\nu \geq 1}$  is compactly divergent. Therefore, we only need to show that (b) follows when  $(f^\nu)_{\nu \geq 1}$  is compactly divergent. Since  $\partial\Omega$  is weakly visible, by Corollary 4.3 we deduce that  $\Omega$  is a weak visibility domain subordinate to  $\overline{\Omega}^\infty$ . Therefore, at this stage, the proof proceeds in a similar way as the proof of [7, Theorem 1.8], with Lemma 6.4 serving in place of Montel's Theorem, Lemmas 6.4 and 6.5 replacing any reference to [7, Theorem 4.3] and [7, Proposition 4.1], respectively, and the words “ $\Omega$  is a weak visibility domain subordinate to  $\overline{\Omega}^\infty$ ” replacing the phrase “ $\Omega$  is a visibility domain”. This works because the proof of [7, Theorem 1.8] is metric-geometrical and charts are irrelevant to the latter proof. Essentially, the latter proof relies on  $(1, \kappa)$ -almost-geodesics. So, the only modifications that are required are the following:

- We need to define a neighbourhood of  $\xi$ , which will be the analogue of  $U_\delta$  appearing in the proof of [7, Theorem 1.8], when  $\xi \in \partial_\infty\Omega$ . Fix  $\tilde{z} \in \Omega$ . For  $\xi \in \overline{\Omega}^\infty \setminus \mathbf{j}(\overline{\Omega})$ , define

$$U_\delta := \text{the connected component of } \overline{\Omega}^\infty \setminus \mathbf{j}(\overline{B_{d_h}(\tilde{z}, \delta^{-1})} \cap \overline{\Omega}) \text{ that contains } \xi.$$

For  $\xi \in \partial\Omega$  define  $U_\delta := B_{d_h}(\xi, \delta) \cap \overline{\Omega}$ .

- Now, for arbitrary  $\delta > 0$  and  $\xi \in \partial_\infty\Omega$  define the function  $G_\delta$  by

$$G_\delta(x_1, x_2) := \inf\{K_\Omega^m(x_1, x_2) : m \in \mathbb{N}, f^m(x_1) \in U_\delta\}.$$

- For any arbitrary compact subset  $K \subset \Omega$  and  $\eta \in \partial_\infty\Omega$ , we need to show that

$$\varepsilon := \liminf_{z \rightarrow \eta} \inf_{k \in K} K_\Omega(k, z)$$

is in  $(0, \infty]$  (where  $\eta$  has the same meaning as in [7]). Since  $\Omega$  is a Kobayashi hyperbolic domain, by [24, Theorem 3.2.1] we have  $Top(\Omega) = Top(K_\Omega)$ , where  $Top(\Omega)$  denotes the manifold topology on  $\Omega$ . The argument to show that  $\varepsilon \in (0, \infty]$  in the proof of [9, Theorem 1.14] relies only on the latter fact and, so, the same argument applies here.

In view of the opening comments above, the result follows.  $\square$

## 7. CONNECTIONS WITH GROMOV HYPERBOLICITY

In this section, we shall follow the notation introduced in Sections 1.4 and 2.3. However, we need some more notation and concepts before we can prove Theorem 1.18. These are taken from [9] but, in the interests of exposition that is reasonably self-contained, we present them here.

**Definition 7.1.** Let  $X$  be a complex manifold and  $\Omega \subsetneq X$  be a Kobayashi hyperbolic domain. Let  $I \subset \mathbb{R}$  be an interval. For  $\kappa \geq 0$ , a path  $\sigma : I \rightarrow \Omega$  is said to be a  $(1, \kappa)$ -near-geodesic if  $\sigma$  is continuous and for all  $s, t \in I$ ,

$$|s - t| - \kappa \leq K_\Omega(\sigma(s), \sigma(t)) \leq |s - t| + \kappa.$$

**Definition 7.2.** Let  $X$  be a complex manifold and  $\Omega \subsetneq X$  be a Kobayashi hyperbolic domain. Fix  $\kappa \geq 0$ . We shall say that a family  $\mathcal{F}$  of  $(1, \kappa)$ -near-geodesics is a  $\kappa$ -admissible family if

- (a) for each pair of distinct points  $x, y \in \Omega$ , there exists a  $(1, \kappa)$ -near-geodesic  $\sigma \in \mathcal{F}$  joining  $x$  and  $y$ ,

(b) if a  $(1, \kappa)$ -near-geodesic  $\sigma : [0, T] \rightarrow \Omega$  is in  $\mathcal{F}$ , then so is  $\bar{\sigma}$ , where

$$\bar{\sigma}(t) := \sigma(T - t), \quad t \in [0, T].$$

For  $x, y \in \Omega$ , we shall use the notation  $\sigma|x \frown y$  to denote that  $\sigma$  is a  $(1, \kappa)$ -near-geodesic (for some  $\kappa \geq 0$ ) joining  $x$  and  $y$ . In view of Proposition 2.8, for any  $\kappa > 0$ , there exists a non-empty  $\kappa$ -admissible family. Therefore, the above discussion is naturally relevant to the ideas we will be discussing in this section.

We can now give the following definition, wherein we will commit the following abuse of notation: for any curve  $\sigma$ , we shall write the **image** of  $\sigma$  simply as  $\sigma$ .

**Definition 7.3.** Let  $X$  be a complex manifold and  $\Omega \subsetneq X$  be a Kobayashi hyperbolic domain. Fix  $\kappa \geq 0$ .

- (a) A  $\kappa$ -triangle is a triple of  $(1, \kappa)$ -near-geodesics  $\alpha|x \frown o$ ,  $\beta|y \frown o$ ,  $\gamma|x \frown y$ , where  $x, y, o \in \Omega$ . We call  $x, y, o$  the *vertices*, and  $\alpha, \beta, \gamma$  the *sides* of the  $\kappa$ -triangle.
- (b) Let  $\delta \geq 0$ . A  $\kappa$ -triangle  $\Delta$  is said to be  $\delta$ -slim if, for each side of  $\Delta$ ,

$$\sigma \subset \bigcup_{\Sigma \in \Delta \setminus \{\sigma\}} \{z \in \Omega : K_{\Omega}(z, \Sigma) \leq \delta\}.$$

- (c) Let  $\mathcal{F}$  be a  $\kappa$ -admissible family of  $(1, \kappa)$ -near-geodesics. Let  $\delta \geq 0$ . We say that  $\Omega$  satisfies  $(\mathcal{F}, \kappa, \delta)$ -Rips condition if every  $\kappa$ -triangle in  $\Omega$  whose sides are (the images of)  $(1, \kappa)$ -near-geodesics belonging to  $\mathcal{F}$  is  $\delta$ -slim.

The result that we need, and which the above notation serves is, the following:

**Result 7.4** (paraphrasing [9, Theorem 10.6]). *Let  $X$  be a connected complex manifold and  $\Omega \subsetneq X$  be a Kobayashi hyperbolic domain. Let  $\kappa \geq 0$  be such that there exists some  $\kappa$ -admissible family of  $(1, \kappa)$ -near-geodesics. Suppose  $(\Omega, K_{\Omega})$  is  $\delta$ -hyperbolic for some  $\delta \geq 0$ . Then, for any  $\kappa$ -admissible family  $\mathcal{F}$ ,  $\Omega$  satisfies the  $(\mathcal{F}, \kappa, 3\delta + 6\kappa)$ -Rips condition.*

The above result paraphrases [9, Theorem 10.6] in two ways. Firstly, [9, Theorem 10.6] has been stated for domains  $\Omega \subsetneq \mathbb{C}^n$ . Secondly, a partial converse of the above result, with  $X = \mathbb{C}^n$ , forms a part of [9, Theorem 10.6]. We have not stated (the relevant version of) the latter because it is not needed here. The proof of [9, Theorem 10.6] extends verbatim to the setting of Result 7.4 because this proof relies on viewing  $\Omega$  merely as a metric space, i.e.,  $(\Omega, K_{\Omega})$ , and on the fact that near-geodesics are continuous.

In the proof that follows:

- The symbol  $\mathbf{j}$  will denote the embedding  $\Omega \hookrightarrow \overline{\Omega}^G$  that is a part of the definition of  $\overline{\Omega}^G$ .
- $\lim$  will denote limits in all contexts other than limits with respect to the topology on  $\overline{\Omega}^G$ .
- Given  $x, y, o \in \Omega$ , the Gromov product with the underlying distance being  $K_{\Omega}$  will be denoted by  $(x|y)_o^{\Omega}$ .

Also, we shall commit the (mild) abuse of notation described in Section 1.3. With these words, we can now present

*The proof of Theorem 1.18.* Fix  $\xi, \eta \in \partial_G \Omega$  such that  $\xi \neq \eta$ . We also fix  $\kappa \geq 0$ . Consider sequences  $(x_{\nu})_{\nu \geq 1} \subset \Omega$  and  $(y_{\nu})_{\nu \geq 1} \subset \Omega$  such that  $x_{\nu} \xrightarrow{G} \xi$ , and  $y_{\nu} \xrightarrow{G} \eta$ . For any pair  $x, y \in \Omega$  we define the set (with  $\kappa$  understood to be fixed as above)

$$AG_{\Omega}(x, y) := \{\gamma : [0, T] \rightarrow \Omega : \gamma \text{ is a } (1, \kappa)\text{-almost-geodesic joining } x \text{ and } y\}.$$

We shall show that there exists a compact set  $K \subset \Omega$  such that for every  $\nu \geq 1$ , and every  $\gamma \in AG_\Omega(x_\nu, y_\nu)$ , we have  $\langle \gamma \rangle \cap K \neq \emptyset$ . If possible, let this assertion be false. Let  $(K_\nu)_{\nu \geq 1}$  be an exhaustion by compacts of  $\Omega$ . Then, passing to a subsequence and relabelling, if needed, we may assume, for each  $\nu \geq 1$ , there exists a  $(1, \kappa)$ -almost-geodesic  $\gamma_\nu \in AG_\Omega(x_\nu, y_\nu)$  such that  $\langle \gamma_\nu \rangle \cap K_\nu = \emptyset$ . Since  $\overline{\Omega}^G$  is locally compact, there exists a  $\overline{\Omega}^G$ -open neighbourhood  $V_\xi$  of  $\xi$  such that  $\overline{V_\xi}^G$  is compact. Fix neighbourhoods  $W_\xi, W_\xi^1, W_\xi^2$  of  $\xi$  such that  $W_\xi \subseteq W_\xi^1 \subseteq W_\xi^2 \subseteq V_\xi$ . Without loss of generality, we may assume that  $x_\nu \in W_\xi$  for all  $\nu \geq 1$ . Let  $r_\nu > 0$  and  $y'_\nu = \gamma_\nu(r_\nu)$  be such that  $y'_\nu \in \mathbf{j}^{-1}(V_\xi \setminus (\overline{W_\xi^1}^G \cup \partial_G \Omega))$ , and such that the image of  $\tilde{\gamma}_\nu := \gamma_\nu|_{[0, r_\nu]}$  is contained in  $W_\xi^2$ . Since  $\overline{V_\xi}^G$  is compact, by passing to a subsequence and relabelling, if needed, we may assume that there exists a point  $\eta' \in \overline{\Omega}^G$  such that  $y'_\nu \xrightarrow{G} \eta'$ . Since  $y'_\nu \notin K_\nu$  for each  $\nu \geq 1$ , we have  $\eta' \in \partial_G \Omega$ . By construction,  $\xi \neq \eta'$ . Therefore, by Part (b) of Result 2.13, it follows that  $(x_\nu)_{\nu \geq 1}$  and  $(y_\nu)_{\nu \geq 1}$  are Gromov sequences and  $\xi = [x_\nu] \neq [y_\nu] = \eta'$ . Therefore, by Definition 2.12, there exists a constant  $M > 0$  such that  $(x_\nu | y'_\nu)_o^\Omega < M$  for all  $\nu \geq 1$ . Let  $\kappa' > \kappa$ . Define  $\mathcal{F} := \{\tilde{\gamma}_\nu : \nu \geq 1\}$ . Since each  $\tilde{\gamma}_\nu$  is also a  $(1, \kappa')$ -almost-geodesic in  $\Omega$ ,  $\mathcal{F}$  is a subset of a  $\kappa'$ -admissible family, namely the collection of all  $(1, \kappa')$ -almost-geodesics in  $\Omega$ . Denote the latter family as  $\mathcal{F}$ .

Fix a point  $o \in \Omega$ . Since  $\kappa' > 0$ , in view of Proposition 2.8, for each  $\nu \geq 1$  we can fix  $\alpha_\nu$ , a  $(1, \kappa')$ -almost-geodesic joining  $x_\nu$  and  $o$ , and fix  $\beta_\nu$ , a  $(1, \kappa')$ -almost-geodesic joining  $y'_\nu$  and  $o$ . Let  $\Delta_\nu$  denote the  $\kappa'$ -triangle whose sides are  $\alpha_\nu, \beta_\nu$ , and  $\tilde{\gamma}_\nu$ . Let  $\delta \geq 0$  be such that  $(\Omega, K_\Omega)$  is  $\delta$ -hyperbolic. Let  $\delta' := 3\delta + 6\kappa'$ . By Result 7.4,  $\Omega$  satisfies  $(\mathcal{F}, \kappa', \delta')$ -Rips condition. Therefore, for each  $\nu \geq 1$

$$\tilde{\gamma}_\nu \subset \bigcup_{\Sigma \in \Delta_\nu \setminus \{\tilde{\gamma}_\nu\}} \{z \in \Omega : K_\Omega(z, \Sigma) \leq \delta'\}.$$

Since  $\tilde{\gamma}_\nu$  is connected, for each  $\nu$ , there exists  $z_\nu \in \tilde{\gamma}_\nu$  such that  $z_\nu \in \{z \in \Omega : K_\Omega(z, \alpha_\nu) \leq \delta'\} \cap \{z \in \Omega : K_\Omega(z, \beta_\nu) \leq \delta'\}$ . For each  $\nu \geq 1$ , fix  $a_\nu \in \alpha_\nu$  and  $c_\nu \in \beta_\nu$  such that  $K_\Omega(a_\nu, z_\nu), K_\Omega(c_\nu, z_\nu) \leq \delta'$ . By triangle inequality it follows that

$$K_\Omega(o, a_\nu) \geq K_\Omega(o, z_\nu) - \delta', \quad (7.1)$$

$$K_\Omega(o, c_\nu) \geq K_\Omega(o, z_\nu) - \delta'. \quad (7.2)$$

Therefore, from (7.1), and since  $\alpha_\nu$  is a  $(1, \kappa')$ -almost-geodesic, we have

$$\begin{aligned} K_\Omega(o, a_\nu) + K_\Omega(a_\nu, x_\nu) &\leq K_\Omega(o, x_\nu) + 3\kappa' \\ \implies K_\Omega(a_\nu, x_\nu) &\leq K_\Omega(o, x_\nu) - K_\Omega(o, z_\nu) + \delta' + 3\kappa'. \end{aligned}$$

Therefore, for each  $\nu \geq 1$ ,

$$\begin{aligned} K_\Omega(x_\nu, z_\nu) &\leq K_\Omega(a_\nu, x_\nu) + K_\Omega(a_\nu, z_\nu) \\ &\leq K_\Omega(o, x_\nu) - K_\Omega(o, z_\nu) + 2\delta' + 3\kappa'. \end{aligned}$$

Similarly, from (7.2), and since  $\beta_\nu$  is a  $(1, \kappa')$ -almost-geodesic, we have  $K_\Omega(y'_\nu, z_\nu) \leq K_\Omega(o, y'_\nu) - K_\Omega(o, z_\nu) + 2\delta' + 3\kappa'$ , for all  $\nu \geq 1$ . Combining last two inequalities, it follows that for all  $\nu \geq 1$ ,

$$2K_\Omega(o, z_\nu) \leq 2(x_\nu | y'_\nu)_o^\Omega + 4\delta' + 6\kappa' < 2M + 4\delta' + 6\kappa'. \quad (7.3)$$

Now, since  $(z_\nu)_{\nu \geq 1} \subset V_\xi$  and  $\overline{V_\xi}^G$  is compact, there exists a subsequence  $(\nu_j)_{j \geq 1}$  and  $\xi' \in \overline{V_\xi}^G$  such that  $z_{\nu_j} \xrightarrow{G} \xi'$ . Now, since  $z_{\nu_j} \notin K_{\nu_j}$  for all  $j \geq 1$ , we have  $\xi' \in \partial_G \Omega$ . Therefore, by Part (b) of Result 2.13,  $(z_{\nu_j})_{j \geq 1}$  is a Gromov sequence. Hence, from Definition 2.12, we have  $\lim_{j \rightarrow \infty} K_\Omega(o, z_{\nu_j}) = \infty$ , which contradicts (7.3). Hence the result.  $\square$



Now, we will give a proof of Proposition 1.19. The map  $\mathbf{j} : D \hookrightarrow \overline{D}^G$  is as explained above.

*The proof of Proposition 1.19.* Since  $D$  is Kobayashi hyperbolic, by the distance decreasing property for the Kobayashi distance of the inclusion map  $\Omega \hookrightarrow D$ , it follows that  $\Omega$  is a Kobayashi hyperbolic domain. Since  $K$  is compact and  $\mathcal{H}^{2n-2}(K) = 0$ , by [24, Theorem 3.2.22] we have

$$K_\Omega = K_D|_{\Omega \times \Omega}. \quad (7.4)$$

Since  $D$  is Gromov hyperbolic, there exists  $\delta \geq 0$  such that  $(D, K_D)$  is  $\delta$ -hyperbolic. Therefore, by Definition 2.11 and (7.4),  $(\Omega, K_\Omega)$  is  $\delta$ -hyperbolic. This proves part (i).

Now, to prove part (ii), since  $\overline{D}$  is homeomorphic to  $\overline{D}^G$ , it is enough to prove that  $\overline{\Omega}^G$  is homeomorphic to  $\overline{D}^G \setminus \mathbf{j}(K)$ . Since  $K$  is compact, as a set,

$$\overline{D}^G \setminus \mathbf{j}(K) = \Omega \cup \partial_G D. \quad (7.5)$$

From (7.4) and Definition 2.12, it follows that every equivalence class of Gromov sequences of  $\Omega$  is an equivalence class of Gromov sequences of  $D$  and the vice versa (since  $K$  is compact). Therefore, as a set  $\overline{\Omega}^G = \overline{D}^G \setminus \mathbf{j}(K)$ . Now, since the Gromov topology depends only on the Kobayashi distance, by Result 2.13, (7.5), and (7.4) we have that  $id_{D \setminus K}$  extends to a homeomorphism between  $\overline{\Omega}^G$  and  $\overline{D}^G \setminus \mathbf{j}(K)$ . Therefore, clearly,  $\overline{\Omega}^G$  is non-compact and locally compact.  $\square$

#### ACKNOWLEDGEMENTS

I would like to thank my thesis advisor, Prof. Gautam Bharali, for many invaluable discussions during course of this work. I am also grateful to him for his advice on the writing of this paper. This work is supported in part by a scholarship from the Prime Minister's Research Fellowship (PMRF) programme (fellowship no. 0201077) and by a DST-FIST grant (grant no. DST FIST-2021 [TPN-700661]).

#### REFERENCES

- [1] Marco Abate, *Horospheres and iterates of holomorphic maps*, Math. Z. **198**, no. 2, 225–238, 1988.
- [2] Marco Abate, *Iteration Theory of Holomorphic Maps on Taut Manifolds*, Research and Lecture Notes in Mathematics, Complex Analysis and Geometry, Mediterranean Press, Rende, 1989.
- [3] Marco Abate and Peter Heinzner, *Holomorphic actions on contractible domains without fixed points*, Math. Z. **211**, no. 4, 547–555, 1992.
- [4] Marco Abate and Jasmin Raissy, *Wolff–Denjoy theorems in nonsmooth convex domains*, Ann. Mat. Pura Appl. **4**, no. 193(5), 1503–1518, 2014.
- [5] Zoltán M. Balogh and Mario Bonk, *Gromov hyperbolicity and the Kobayashi metric on strictly pseudoconvex domains*, Comment. Math. Helv. **75**, no. 3, 504–533, 2000.
- [6] Annapurna Banik, *Visibility domains that are not pseudoconvex*, Bull. Sci. math. **193**, 2024.
- [7] Gautam Bharali and Anwoy Maitra, *A weak notion of visibility, a family of examples, and Wolff–Denjoy theorems*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. **22**, no. 1, 195–240, 2021.
- [8] Gautam Bharali and Andrew Zimmer, *Goldilocks domains, a weak notion of visibility, and applications*, Adv. Math. **310**, 377–425, 2017.
- [9] Gautam Bharali and Andrew Zimmer, *Unbounded visibility domains, the end compactification, and applications*, Trans. Amer. Math. Soc. **376**, no. 8, 5949–5988, 2023.
- [10] Filippo Bracci, Manuel D. Contreras, and Santiago Díaz-Madrigal, *Continuous Semigroups of Holomorphic Self-maps of the Unit Disc*, Springer Monographs in Mathematics, Springer Nature Switzerland AG, 2020.
- [11] Filippo Bracci, Hervé Gaussier, Nikolai Nikolov, and Pascal J. Thomas, *Local and global visibility and Gromov hyperbolicity of domains with respect to the Kobayashi distance*, Trans. Amer. Math. Soc. **377**, no. 1, 471–493, 2024.

- [12] Filippo Bracci, Hervé Gaussier, and Andrew Zimmer, *Homeomorphic extension of quasi-isometries for convex domains in  $\mathbb{C}^d$  and iteration theory*, Math. Ann. **379**, no. 1–2, 691–718, 2021.
- [13] Filippo Bracci, Nikolai Nikolov, and Pascal J. Thomas, *Visibility of Kobayashi geodesics in convex domains and related properties*, Math. Z. **301**, no. 2, 2011–2035, 2022.
- [14] Martin R. Bridson and André Haefliger, *Metric Spaces of Non-Positive Curvature*, Grundlehren der Mathematischen Wissenschaften **319**, Springer-Verlag, Berlin, 1999.
- [15] Monika Budzyńska, Tadeusz Kuczumow, and Simeon Reich, *A Denjoy–Wolf theorem for compact holomorphic mappings in complex Banach spaces*, Ann. Acad. Sci. Fenn. Math. **38**, no. 2, 747–756, 2013.
- [16] Vikramjeet Singh Chandel, Anwoy Maitra, and Amar Deep Sarkar, *Notions of visibility with respect to the Kobayashi distance: comparison and applications*, Ann. Mat. Pura Appl. **203**, 475–498, 2024.
- [17] Tushar Das, David Simmons, and Mariusz Urbański, *Geometry and Dynamics in Gromov Hyperbolic Metric Spaces*, Mathematical Surveys and Monographs **218**, American Mathematical Society, Providence, RI, 2017.
- [18] A. Denjoy, *Sur l’itération des fonctions analytiques*, C.R. Acad. Sci. Paris **182**, 255–257, 1926.
- [19] P. Eberlein and B. O’Neill, *Visibility manifolds*, Pacific J. Math. **46**, 45–109, 1973.
- [20] Michel Hervé, *Quelques propriétés des applications analytiques d’une boule à  $m$  dimensions dan elle-même*, J. Math. Pures Appl. **9**, no. 42, 117–147, 1963.
- [21] Xiao Jun Huang, *A non-degeneracy property of extremal mappings and iterates of holomorphic self-mappings*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. **4**, no. 21(3), 399–419, 1994.
- [22] Aleksandra Huczek and Andrzej Wiśnicki, *Theorems of Wolff–Denjoy type for semigroups of nonexpansive mappings in geodesic spaces* Math. Nachr. **296**, no. 8, 3387–3394, 2023.
- [23] Anders Karlsson, *Non-expanding maps and Busemann functions*, Ergodic Theory Dynam. Systems **21**, no. 5, 1447–1457, 2001.
- [24] Shoshichi Kobayashi, *Hyperbolic Complex Spaces*, Grundlehren Math. Wiss. **318**, Springer-Verlag, Berlin, 1998.
- [25] Rumpa Masanta, *Kobayashi complete domains in complex manifolds*, To appear in Int. J. Math., arXiv reference: [arXiv:2308.07181](https://arxiv.org/abs/2308.07181).
- [26] Nikolai Nikolov, Ahmed Yekta Ökten, and Pascal J. Thomas, *Local and global notions of visibility with respect to Kobayashi distance: a comparison*, Ann. Polon. Math. **132**, no. 2, 169–185, 2024.
- [27] H. L. Royden, *Remarks on Kobayashi metric*, Lecture Notes in Math. **185**, 125–137, Springer, Berlin, 1971.
- [28] Amar Deep Sarkar, *Localization of the Kobayashi distance for any visibility domain*, Geom. Anal. **33**, no. 5, paper no. 144, 16 pp, 2023.
- [29] Sergio Venturini, *Pseudodistances and pseudometrics on real and complex manifolds*, Ann. Mat. Pura Appl. **154**, no. 4, 385–402, 1989.
- [30] J. Wolff, *Sur une généralisation d’un théorème de Schwarz*, C.R. Acad. Sci. Paris **182**, 918–920, 1926.
- [31] Andrew M. Zimmer, *Gromov hyperbolicity and the Kobayashi metric on convex domains of finite type* Math. Ann. **365**, no. 3–4, 1425–1498, 2016.

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF SCIENCE, BANGALORE 560012, INDIA  
 Email address: [rumpamasanta@iisc.ac.in](mailto:rumpamasanta@iisc.ac.in)