

Wiener-Hopf indices of unimodular functions on the imaginary axis

A.E. Frazho, A.C.M. Ran and F. van Schagen

Abstract. This paper is concerned with the Wiener-Hopf indices of unimodular rational matrix functions on the imaginary axis. These indices play a role in the Fredholm theory for Wiener-Hopf integral operators. Our main result gives formulas for the Wiener-Hopf indices in terms of the matrices appearing in realizations of the factors in a Douglas-Shapiro-Shields factorization of the unimodular function. Two approaches to this problem are presented: one direct approach using operator theoretic methods, and a second approach using the Cayley transform which allows to use results for an analogous problem regarding unimodular functions on the unit circle and corresponding Toeplitz operators.

Mathematics Subject Classification (2010). 47A68, 47B35, 47A56, 47A53.

Keywords. Wiener-Hopf indices, unitary rational matrix functions, Wiener-Hopf operators.

1. Introduction

Wiener-Hopf factorization of matrix valued functions plays an important role in determining the Fredholm properties of several classes of operators, such as singular and Wiener-Hopf integral operators and Toeplitz operators, see e.g., [8, 9, 14, 15, 16] and [18]. To make this more explicit, let $R(s)$ be an $m \times m$ rational matrix valued function on the imaginary axis, which is continuous and takes invertible values for s on the imaginary axis. A factorization

$$R(s) = W_-(s) \operatorname{diag} \left(\left(\frac{1-s}{1+s} \right)^{\kappa_j} \right)_{j=1}^m W_+(s),$$

where W_- and its inverse are analytic on the closed left half of the complex plane, including infinity, and W_+ and its inverse are analytic on the closed right hand half plane, including infinity, and $\kappa_j \in \mathbb{Z}$ for $j = 1, \dots, m$, is called a (*right*) *Wiener-Hopf factorization* with respect to the imaginary axis. The

integers κ_j are uniquely determined by R and they are called the *Wiener-Hopf indices* of R .

In order to present the main results of this paper we recall some definitions and fix some notations. Throughout \mathcal{E} is a finite dimensional Hilbert space and $L_+^2(\mathcal{E})$ denotes the space of square integrable functions from $[0, \infty)$ to \mathcal{E} . The Laplace transform \mathfrak{L} is the unitary operator mapping $L_+^2(\mathcal{E})$ onto $H^2(\mathcal{E})$ defined by

$$(\mathfrak{L}f)(s) = \int_0^\infty e^{-st} f(t) dt \quad (f \in L_+^2(\mathcal{E})). \quad (1.1)$$

Here $H^2(\mathcal{E})$ is the Hardy space of \mathcal{E} -valued functions that are analytic in the right hand half complex plane $\mathbb{C}_+ = \{s \in \mathbb{C} : \Re(s) > 0\}$ and square integrable on the imaginary axis. Furthermore $H^\infty(\mathcal{E}, \mathcal{E})$ is the Hardy space of all functions Θ whose values are operators on \mathcal{E} and that are analytic and uniformly bounded in the open right half plane \mathbb{C}_+ , i.e.,

$$\|\Theta\|_\infty = \sup\{\|\Theta(s)\| : \Re(s) > 0\} < \infty.$$

(If $\mathcal{E} = \mathbb{C}$, then $H^\infty(\mathbb{C}, \mathbb{C})$ is denoted by H^∞ .) Similarly, $H^\infty(\mathcal{E})$ is the Hardy space consisting of the set of all \mathcal{E} -valued functions $\Psi(s)$ that are analytic in the open right half plane and such that

$$\|\Psi\|_\infty = \sup\{\|\Psi(s)\| : \Re(s) > 0\} < \infty.$$

Let $R(s)$ be a rational function taking unitary values on the finite dimensional space \mathcal{E} for values of s on the imaginary line. In particular, $R(\infty)$ is also a unitary operator. Let $r(t)$ be given via $R(i\omega) = R(\infty) + (\mathfrak{L}r)(i\omega)$.

Define T_R to be the Wiener-Hopf operator on $L_+^2(\mathcal{E})$ determined by R , that is,

$$(T_R f)(t) = R(\infty)f(t) + \int_0^\infty r(t - \tau)f(\tau)d\tau \quad (f \in L_+^2(\mathcal{E})). \quad (1.2)$$

For such an operator the image is denoted by $\text{Im } T_R$ and the kernel or null space by $\text{Ker } T_R$.

Throughout $\zeta(s) = \frac{1-s}{1+s}$ is the conformal mapping which maps the open right half plane \mathbb{C}_+ onto the open unit disc $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$. Let $-\kappa_1, -\kappa_2, \dots, -\kappa_p$ with $\kappa_1 \geq \kappa_2 \geq \dots \geq \kappa_p$ be the negative Wiener-Hopf indices of the function R . Then (see [15] Theorem XIII.3.2) we have that the dimension $\mathfrak{n}(T_R)$ of the null space $\text{Ker } T_R$ is given by

$$\mathfrak{n}(T_R) = \sum_{\kappa_j \geq 1} \kappa_j.$$

Consider the function R multiplied by $\zeta(s)^k$, which we denote by $\zeta^k R$. Define for $k = 1, 2, \dots$ the numbers μ_k by

$$\mu_k = \mathfrak{n}(T_{\zeta^{k-1}R}) - \mathfrak{n}(T_{\zeta^k R}).$$

Then (see Section 2 below)

$$\kappa_j = \#\{k \mid \mu_k \geq j\}.$$

Here $\#E$ denotes the number of elements of the set E .

Finding the Wiener-Hopf indices in terms of matrices in a realization of the function R is a problem that has already some history, see [3, 4, 5, 7, 18, 19]. An analogous problem for a unimodular function on the unit circle was studied in [20]. There, significant use was made of the Douglas-Shapiro-Shields factorization of R , that is, writing R as $R = VW^*$, where V and W are bi-inner. Note that in this case we say that the operator valued function Θ is a *bi-inner function* if Θ is analytic and uniformly bounded on \mathbb{D} , and $\Theta(e^{i\omega})$ is a unitary operator on \mathcal{E} for almost all $\omega \in \mathbb{R}$. The Wiener-Hopf indices were given in [20] in terms of realizations of V and W , based on earlier work in [12]. These results were extended to formulas for the Wiener-Hopf indices for any rational matrix valued function in [21].

Our aim in this paper is to obtain a result analogous to the result of [20] but for the imaginary axis replacing the unit circle and with a different method, which is more operator theoretic. This method is in parallel with our earlier results in [13] for the case of unimodular rational matrix functions on the unit circle and their related Toeplitz operators. Our approach leads to simple formulas for the Wiener-Hopf indices of a rational matrix function that takes unitary values on the imaginary axis.

Finally in this introduction we give a short description of the various sections of this paper. In Section 2, we introduce the functions and their realizations, and present the main result in Theorem 2.2. Section 3 is concerned with the Wiener-Hopf and Hankel operators corresponding to bi-inner rational matrix functions. Section 4 gives more detailed results on the unimodular function R , its Wiener-Hopf operator T_R and factorization. In subsection 4.1 we specify the results for the case when the bi-inner functions are scalar valued Blaschke products. We derive the main results for this special case. Section 5 is devoted to the proof of the main theorem. In Section 6 we treat the Cayley transform and the connection it gives between the realizations of rational matrix functions on the unit circle in the complex plane and the realizations of rational matrix functions on the imaginary axis in the complex plane. In Section 7 these relations are used to prove the equivalence of the main result, Theorem 2.2, of the paper with [13, Theorem 2.2]. In Subsection 7.1 we present an example.

2. The main result.

To present our method to compute the Wiener-Hopf indices, let us fix some notation. Recall that $\{A \text{ on } \mathcal{X}, B, C, D\}$ is a *realization of a function* $\Theta(s)$ if

$$\Theta(s) = D + C(sI - A)^{-1}B.$$

Here A is an operator on \mathcal{X} and B maps \mathcal{U} into \mathcal{X} , while C maps \mathcal{X} into \mathcal{Y} and D maps \mathcal{U} into \mathcal{Y} . Two state space realizations $\{A \text{ on } \mathcal{X}, B, C, D\}$ and $\{A_1 \text{ on } \mathcal{X}_1, B_1, C_1, D_1\}$ are *unitarily equivalent* if $D = D_1$ and there exists a

unitary operator U mapping \mathcal{X}_1 onto \mathcal{X} such that

$$AU = UA_1 \quad \text{and} \quad B = UB_1 \quad \text{and} \quad CU = C_1.$$

We say that an operator A on \mathcal{X} is *dissipative* if $A^* + A \leq 0$. An operator A on a finite dimensional space \mathcal{X} is *stable* if all the eigenvalues for A are contained in the open left hand plane $\mathbb{C}_- = \{s \in \mathbb{C} : \Re(s) < 0\}$.

Throughout we will be dealing with finite dimensional realizations, that is, realizations of the form $\{A \text{ on } \mathcal{X}, B, C, D\}$ where the state space \mathcal{X} is finite dimensional. In general we will call such a realization MIMO (multi input, multi output). In the particular case when \mathcal{U} and \mathcal{Y} are one-dimensional we say that the realization is SISO (single input, single output). The realization $\{A, B, C, D\}$ is *stable*, if A is stable. Next, we say that the realization $\{A, B, C, D\}$ is *stable and dissipative* if the following holds:

1. The operator A is stable and $A + A^* + C^*C = 0$;
2. the operator D is unitary;
3. $B = -C^*D$.

It is noted that the previous three conditions are equivalent to

- a) The operator A is stable and $A + A^* + BB^* = 0$;
- b) the operator D is unitary;
- c) $C = -DB^*$.

We are now ready to present the following classical result, compare, e.g., [2], [7], Section 17.5. The result is closely related to results on so called Livsic-Brodskii characteristic operator functions, see Section 1.2 in [6].

Theorem 2.1. *Let Θ be a rational function in $H^\infty(\mathcal{E}, \mathcal{E})$. Then Θ is bi-inner if and only if Θ admits a stable dissipative realization $\{A, B, C, D\}$. In this case, all stable dissipative realizations of Θ are unitarily equivalent.*

Let R take unitary values on the imaginary axis. Due to the Douglas-Shapiro-Shields factorization, when computing the Wiener-Hopf indices of R , without loss of generality, one can assume that $R = VW^*$ where V and W are two rational bi-inner functions in $H^\infty(\mathcal{E}, \mathcal{E})$. Here W^* denotes the function defined by $W^*(s) = (W(-\bar{s}))^*$ ($s \in \mathbb{C}$) and thus we have that $W^*(i\omega) = (W(i\omega))^*$ for all $\omega \in \mathbb{R}$. Let $\{A_v \text{ on } \mathcal{X}_v, B_v, C_v, D_v\}$ and $\{A_w \text{ on } \mathcal{X}_w, B_w, C_w, D_w\}$ be two stable dissipative realizations of V and W respectively. In particular,

$$V(s) = D_v + C_v(sI - A_v)^{-1}B_v \tag{2.1}$$

$$W(s) = D_w + C_w(sI - A_w)^{-1}B_w. \tag{2.2}$$

Let T_R be the Wiener-Hopf operator on $L^2_+(\mathcal{E})$ determined by R given by (1.2).

Let $-\kappa_1, -\kappa_2, \dots, -\kappa_p$ with $\kappa_1 \geq \kappa_2 \geq \dots \geq \kappa_p$ be the negative Wiener-Hopf indices of the function R and $R(s) = W_-(s)D(s)W_+(s)$ the Wiener-Hopf factorization of R . Then (see [15] Theorem XIII.3.2) we have that the

dimension $\mathbf{n}(T_R)$ of the null space of T_R , is given by

$$\mathbf{n}(T_R) = \sum_{\kappa_j \geq 1} \kappa_j.$$

Recall that $\zeta(s) = \frac{1-s}{1+s}$. Consider the function R multiplied by ζ^k , which we denote by $\zeta^k R$. Since $\zeta^k(s)R(s) = W_-(s)(\zeta^k(s)D(s))W_+(s)$, the Wiener-Hopf indices of $\zeta^k R$ are each k higher than the corresponding index of R . Therefore

$$\mathbf{n}(T_{\zeta^k R}) = \sum_{\kappa_j \geq k+1} (\kappa_j - k).$$

Define for $k = 1, 2, \dots$, the numbers μ_k by

$$\mu_k = \mathbf{n}(T_{\zeta^{k-1} R}) - \mathbf{n}(T_{\zeta^k R}) = \#\{j : \kappa_j \geq k\}. \quad (2.3)$$

Then (see [17] Proposition III.4.1)

$$\kappa_j = \#\{k : \mu_k \geq j\}. \quad (2.4)$$

The main result of the paper is the following theorem.

Theorem 2.2. *Assume that $R = VW^*$ where V and W are two bi-inner rational functions in $H^\infty(\mathcal{E}, \mathcal{E})$. Let $\{A_v, B_v, C_v, D_v\}$ and $\{A_w, B_w, C_w, D_w\}$ be stable dissipative realizations of V and W , respectively. Let Ω be the unique solution of the Lyapunov equation*

$$A_v \Omega + \Omega A_w^* + B_v B_w^* = 0. \quad (2.5)$$

Let C_\circ be the operator mapping \mathcal{X}_w into \mathcal{E} defined by

$$C_\circ = D_v B_w^* + C_v \Omega. \quad (2.6)$$

Finally, let Q be the unique solution to the Lyapunov equation

$$A_w Q + Q A_w^* + C_\circ^* C_\circ = 0. \quad (2.7)$$

Then the following holds:

1. The operator Q is a positive contraction.
2. The multiplicity of 1 as an eigenvalue of Q equals $\mathbf{n}(T_R)$. In other words, $\mathbf{n}(T_R) = \mathbf{n}(I - Q)$. Moreover, for $\zeta(s) = \frac{1-s}{1+s}$, we have

$$\mathbf{n}(T_{\zeta^k R}) = \dim(\text{Ker}(I - \zeta(-A_w)^k Q \zeta(-A_w)^{*k})). \quad (2.8)$$

3. For $k = 1, 2, \dots$, consider the integers

$$\mu_k = \mathbf{n}(I - \zeta(-A_w)^{k-1} Q (\zeta(-A_w)^*)^{k-1}) - \mathbf{n}(I - \zeta(-A_w)^k Q \zeta(-A_w)^{*k}). \quad (2.9)$$

Then the negative Wiener-Hopf indices $-\kappa_1, \dots, -\kappa_p$ of T_R are given by

$$\kappa_j = \#\{k : \mu_k \geq j\}, \quad (j = 1, \dots, p = \mu_1). \quad (2.10)$$

Notice that, once Parts 1 and 2 are proven, Part 3 follows from the equations (2.3) and (2.4).

The dual statement for the positive Wiener-Hopf indices is obtained by applying the above theorem to the function $R^*(s) = R(-\bar{s})^*$; see Corollary 5.3 below.

3. Unitary functions on the imaginary axis.

Let $\Theta \in H^\infty(\mathcal{E}, \mathcal{E})$ be such that $\Theta(s) - \Theta(\infty) = (\mathfrak{L}\theta)(s)$ and $(\mathfrak{L}\theta)(s)$ is the Laplace transform of a rational function $\theta(t)$ whose values are linear operators on \mathcal{E} and such that $\int_0^\infty \|\theta(t)\| dt < \infty$. (Here $\Theta(\infty)$ is the constant operator function on \mathcal{E} defined by $\Theta(\infty) = \lim_{s \rightarrow \infty} \Theta(s)$.) Then T_Θ is the *Wiener-Hopf operator* on $L_+^2(\mathcal{E})$ and H_Θ is the *Hankel operator* on $L_+^2(\mathcal{E})$ respectively defined by

$$(T_\Theta f)(t) = \Theta(\infty)f(t) + \int_0^t \theta(t - \tau)f(\tau)d\tau \quad (3.1)$$

$$(H_\Theta f)(t) = \int_0^\infty \theta(t + \tau)f(\tau)d\tau \quad (3.2)$$

where f is a function in $L_+^2(\mathcal{E})$. Throughout $\tilde{\Theta}$ is the function in $H^\infty(\mathcal{E}, \mathcal{E})$ defined by $\tilde{\Theta}(s) = \Theta(\bar{s})^*$ for all s in the open right half plane. In particular, $\{A, B, C, D\}$ is a realization for Θ if and only if $\{A^*, C^*, B^*, D^*\}$ is a realization for $\tilde{\Theta}$. Moreover, one readily obtains the following identity for Hankel operators: $H_{\tilde{\Theta}} = H_\Theta^*$. Finally, it is noted that $\theta(t) = Ce^{At}B$ for $t \geq 0$.

As before, let Θ be a bi-inner function in $H^\infty(\mathcal{E}, \mathcal{E})$. In this case, the corresponding Wiener-Hopf operator T_Θ is an isometry on $L_+^2(\mathcal{E})$. Indeed, $T_{\Theta^*}T_\Theta = T_{\Theta^*} = I$. Because $\tilde{\Theta}$ is also bi-inner, $T_{\tilde{\Theta}}$ is also an isometry on $L_+^2(\mathcal{E})$. Let $\mathfrak{H}(\Theta)$ and $\mathfrak{H}(\tilde{\Theta})$ denote the orthogonal complements of the ranges of T_Θ and $T_{\tilde{\Theta}}$, respectively, that is,

$$\mathfrak{H}(\Theta) = L_+^2(\mathcal{E}) \ominus T_\Theta L_+^2(\mathcal{E}) \quad \text{and} \quad \mathfrak{H}(\tilde{\Theta}) = L_+^2(\mathcal{E}) \ominus T_{\tilde{\Theta}} L_+^2(\mathcal{E}). \quad (3.3)$$

By consulting Equation (24) in Section XII.2 of [15] and with the fact that Θ is bi-inner, we see that

$$P_{\mathfrak{H}(\Theta)} = H_\Theta H_\Theta^* = I - T_\Theta T_\Theta^* \quad \text{and} \quad P_{\mathfrak{H}(\tilde{\Theta})} = H_{\tilde{\Theta}}^* H_{\tilde{\Theta}} = I - T_{\tilde{\Theta}} T_{\tilde{\Theta}}^* \quad (3.4)$$

are orthogonal operators on $\mathfrak{H}(\Theta)$ and $\mathfrak{H}(\tilde{\Theta})$, respectively. The second identity follows from the fact that $\tilde{\Theta}$ is also bi-inner. In particular, $P_{\mathfrak{H}(\Theta)} = H_\Theta H_\Theta^*$ and the range of H_Θ equals $\mathfrak{H}(\Theta)$. From the second equality in (3.4) we also have that $P_{\mathfrak{H}(\tilde{\Theta})} = H_{\tilde{\Theta}}^* H_{\tilde{\Theta}}$ and the range of the Hankel operator $H_{\tilde{\Theta}}^*$ equals $\mathfrak{H}(\tilde{\Theta})$. Therefore

$$\text{Im}(H_\Theta) = \mathfrak{H}(\Theta) \quad \text{and} \quad \text{Ker}(H_\Theta)^\perp = \mathfrak{H}(\tilde{\Theta}).$$

Using $P_{\mathfrak{H}(\tilde{\Theta})} = H_{\tilde{\Theta}}^* H_{\tilde{\Theta}}$, we see that there exists a unitary operator U mapping $\mathfrak{H}(\tilde{\Theta})$ onto $\text{Im}(H_\Theta) = \mathfrak{H}(\Theta)$ such that $UP_{\mathfrak{H}(\tilde{\Theta})} = H_\Theta$. Therefore the Hankel operator H_Θ can be viewed as a unitary operator mapping $\text{Ker}(H_\Theta)^\perp = \mathfrak{H}(\tilde{\Theta})$ onto $\text{Im}(H_\Theta) = \mathfrak{H}(\Theta)$. In particular, $\mathfrak{H}(\tilde{\Theta})$ and $\mathfrak{H}(\Theta)$ have the same dimension.

Let $\{A \text{ on } \mathcal{X}, B, C, D\}$ be any stable dissipative realization of a rational bi-inner function Θ in $H^\infty(\mathcal{E}, \mathcal{E})$. Then its *observability operator* Γ mapping \mathcal{X}

into $L_+^2(\mathcal{E})$ and *controllability operator* Υ mapping $L_+^2(\mathcal{E})$ into \mathcal{X} are defined by

$$\Gamma x = Ce^{At}x \quad (x \in \mathcal{X}, t \geq 0) \quad (3.5)$$

$$\Upsilon u = \int_0^\infty e^{At}Bu(t)dt \quad (u \in L_+^2(\mathcal{E})). \quad (3.6)$$

Because $A^* + A + C^*C = 0$ and A is stable, the observability operator Γ is an isometry. Likewise, since $A^* + A + BB^* = 0$ and A is stable, the controllability operator Υ is a co-isometry. Using the fact that $\Theta(s) - \Theta(\infty)$ is the Laplace transform of $\theta(t) := Ce^{At}B$, it follows that the Hankel operator H_Θ admits a factorization of the form:

$$H_\Theta = \Gamma\Upsilon. \quad (3.7)$$

Recall that H_Θ can be viewed as a unitary operator from $\mathfrak{H}(\tilde{\Theta})$ onto $\mathfrak{H}(\Theta)$. Since Γ is an isometry and Υ is a co-isometry, the equalities $\mathfrak{H}(\Theta) = \text{Im}(\Gamma)$ and $\mathfrak{H}(\tilde{\Theta}) = \text{Im}(\Upsilon^*)$ hold. The equation $H_\Theta = \Gamma\Upsilon$ with (3.4), readily implies that

$$P_{\mathfrak{H}(\Theta)} = \Gamma\Gamma^* \quad \text{and} \quad P_{\mathfrak{H}(\tilde{\Theta})} = \Upsilon^*\Upsilon. \quad (3.8)$$

4. The function $R = VW^*$.

Let V and W be two rational bi-inner functions in $H^\infty(\mathcal{E}, \mathcal{E})$. Let R be the rigid function in $L^\infty(\mathcal{E}, \mathcal{E})$ defined by

$$R(i\omega) = V(i\omega)W(i\omega)^* \quad (\text{for } -\infty < \omega < \infty). \quad (4.1)$$

(A function Ξ in $L^\infty(\mathcal{E}, \mathcal{E})$ is *rigid* if $\Xi(i\omega)$ is almost everywhere a unitary operator on \mathcal{E} .) Since V , W and R are rational we extend the definition of R to all but a finite number of values of $s \in \mathbb{C}$ by $R(s) = V(s)W(-\bar{s})^*$. Let T_R be the Wiener-Hopf operator on $L_+^2(\mathcal{E})$ determined by R , see (1.2). Because V and W are bi-inner, H_V is a unitary operator from $\mathfrak{H}(\tilde{V})$ onto $\mathfrak{H}(V)$, and H_W is a unitary operator from $\mathfrak{H}(\tilde{W})$ onto $\mathfrak{H}(W)$. Recall that $T_R = T_V T_W^* + H_V H_W^*$; see Equation (24) in Section XII.2 of [15]. This readily implies that

$$T_R = T_V T_W^* + H_V Y H_W^* \quad (4.2)$$

where Y is the contraction mapping $\mathfrak{H}(\tilde{W})$ into $\mathfrak{H}(\tilde{V})$ defined by

$$Y = P_{\mathfrak{H}(\tilde{V})} |\mathfrak{H}(\tilde{W}) : \mathfrak{H}(\tilde{W}) \rightarrow \mathfrak{H}(\tilde{V})|. \quad (4.3)$$

Since V and W are both bi-inner,

$$L_+^2(\mathcal{E}) = \text{Im}(T_V) \oplus \text{Im}(H_V) \quad \text{and} \quad L_+^2(\mathcal{E}) = \text{Im}(T_W) \oplus \text{Im}(H_W).$$

Using this with $T_R = T_V T_W^* + H_V Y H_W^*$, we see that T_R admits a "singular value type" decomposition of the form:

$$T_R = T_V T_W^* + H_V Y H_W^* = \begin{bmatrix} T_V & H_V \end{bmatrix} \begin{bmatrix} I & 0 \\ 0 & Y \end{bmatrix} \begin{bmatrix} T_W^* \\ H_W^* \end{bmatrix}. \quad (4.4)$$

Here

$$[T_V \ H_V] : \begin{bmatrix} L_+^2(\mathcal{E}) \\ \mathfrak{H}(\widetilde{V}) \end{bmatrix} \rightarrow L_+^2(\mathcal{E}) \quad \text{and} \quad \begin{bmatrix} T_W^* \\ H_W^* \end{bmatrix} : L_+^2(\mathcal{E}) \rightarrow \begin{bmatrix} L_+^2(\mathcal{E}) \\ \mathfrak{H}(\widetilde{W}) \end{bmatrix}$$

are both unitary operators. Moreover, the middle term

$$\begin{bmatrix} I & 0 \\ 0 & Y \end{bmatrix} = \begin{bmatrix} I & 0 \\ 0 & P_{\mathfrak{H}(\widetilde{V})} | \mathfrak{H}(\widetilde{W}) \end{bmatrix} : \begin{bmatrix} L_+^2(\mathcal{E}) \\ \mathfrak{H}(\widetilde{W}) \end{bmatrix} \rightarrow \begin{bmatrix} L_+^2(\mathcal{E}) \\ \mathfrak{H}(\widetilde{V}) \end{bmatrix} \quad (4.5)$$

is a contraction.

Due to the decomposition of T_R in (4.4), it follows that all the properties such as invertibility and Fredholmness of the operator T_R are the same as those of the contraction Y .

It is noted that x is in $\text{Ker}(Y)$ if and only if x is in $\mathfrak{H}(\widetilde{W})$ and $P_{\mathfrak{H}(\widetilde{V})} x = 0$, or equivalently, x is in $\mathfrak{H}(\widetilde{W})$ and x is in $\text{Im}(T_{\widetilde{V}}) = \mathfrak{H}(\widetilde{V})^\perp$. In other words,

$$\text{Ker}(Y) = \text{Im}(T_{\widetilde{V}}) \cap \mathfrak{H}(\widetilde{W}) \quad \text{and} \quad \text{Ker}(Y^*) = \text{Im}(T_{\widetilde{W}}) \cap \mathfrak{H}(\widetilde{V}),$$

where the second equality follows from a similar argument.

Recall that an operator T mapping \mathcal{X} into \mathcal{Y} admits a Moore-Penrose inverse T^{pinv} if the operator $T|_{\text{Ker}(T)^\perp}$ mapping $\text{Ker}(T)^\perp$ into the range of T is invertible. In this case, the Moore-Penrose inverse of T is given by $T^{pinv} = (T|_{\text{Ker}(T)^\perp})^{-1} P_{\text{Im}(T)}$. By consulting the form of T_R in (4.4), we obtain the following result.

Proposition 4.1. *Let $R = VW^*$ where V and W are both bi-inner functions in $H^\infty(\mathcal{E}, \mathcal{E})$. Moreover, let Y be the contraction mapping $\mathfrak{H}(\widetilde{W})$ into $\mathfrak{H}(\widetilde{V})$ defined by $Y = P_{\mathfrak{H}(\widetilde{V})} | \mathfrak{H}(\widetilde{W})$. Then the following hold.*

1. *The operator T_R is invertible if and only if Y is invertible. In this case,*

$$T_R^{-1} = T_W T_V^* + H_W Y^{-1} H_V^*. \quad (4.6)$$

2. *The subspaces $\text{Ker}(T_R)$ and $\text{Ker}(Y)$ have the same dimension. In fact,*

$$\text{Ker}(T_R) = H_W \text{Ker}(Y) \quad \text{and} \quad \text{Ker}(Y) = \text{Im}(T_{\widetilde{V}}) \cap \mathfrak{H}(\widetilde{W}). \quad (4.7)$$

3. *The subspaces $\text{Ker}(T_R^*)$ and $\text{Ker}(Y^*)$ have the same dimension. In particular,*

$$\text{Ker}(T_R^*) = H_V \text{Ker}(Y^*) \quad \text{and} \quad \text{Ker}(Y^*) = \text{Im}(T_{\widetilde{W}}) \cap \mathfrak{H}(\widetilde{V}). \quad (4.8)$$

4. *The subspaces $\text{Im}(T_R)^\perp$ and $\text{Im}(Y)^\perp$ have the same dimension. In fact,*

$$\text{Im}(T_R)^\perp = \text{Ker}(T_R^*) = H_V \text{Ker}(Y^*) = H_V \text{Im}(Y)^\perp. \quad (4.9)$$

5. *The operator T_R admits a Moore-Penrose restricted inverse if and only if Y admits a Moore-Penrose restricted inverse. In this case,*

$$T_R^{pinv} = T_W T_V^* + H_W Y^{pinv} H_V^*. \quad (4.10)$$

4.1. The Blaschke product case

In this section, to gain some insight into the general case, we will study the contraction $Y = P_{\mathfrak{H}(\varphi)}|_{\mathfrak{H}(m)}$ mapping $\mathfrak{H}(m)$ into $\mathfrak{H}(\varphi)$ when m and φ are scalar Blaschke products. We say that a function $b(s)$ is a *Blaschke product* if

$$b(s) = \rho \prod_{k=1}^n \frac{s + \bar{\alpha}_k}{s - \alpha_k} \quad (\text{where } \Re(\alpha_k) < 0 \text{ for all } k). \quad (4.11)$$

(Here ρ is a complex number on the unit circle.) Throughout $\tilde{b}(s) = \overline{b(\bar{s})}$. Finally, it is noted that $\tilde{b}(-s) = \frac{1}{b(s)}$.

Moreover, $n = \deg(b)$ is the *degree* of the Blaschke product. We will only consider Blaschke products of finite degree. So if we say that $b(s)$ is a Blaschke product, then we assume that $b(s)$ is a function of the form (4.11) and the degree of b is finite. It is well known that $b(s)$ is a rational inner function in H^∞ if and only if b is a Blaschke product (of finite degree). Furthermore, the SISO (single input single output) function $b(s)$ is a Blaschke product of degree n if and only if $b(s)$ admits a stable dissipative realization $\{A \text{ on } \mathcal{X}, B, C, D\}$ where n is the dimension of the state space \mathcal{X} . (See, eg., [7] Section 17.5.) In this case, the poles of $b(s)$ are precisely the eigenvalues of A .

Now assume that $\{A \text{ on } \mathcal{X}, B, C, D\}$ is a stable dissipative realization for a Blaschke product $b(s)$ of degree n . Let Γ mapping \mathcal{X} into L_+^2 be the observability operator formed by the pair $\{C, A\}$. Recall that Γ is an isometry. Moreover, the range of Γ equals $\mathfrak{H}(b)$. Because the dimension of the state space is n , it follows that the dimension of $\mathfrak{H}(b)$ (denoted by $\dim(\mathfrak{H}(b))$) equals n . Furthermore, the Laplace transform of the space $\mathfrak{H}(b)$ is given by

$$\mathfrak{L}(\mathfrak{H}(b)) = \{\mathfrak{L}(\Gamma x) : x \in \mathcal{X}\} = \{C(sI - A)^{-1}x : x \in \mathcal{X}\}. \quad (4.12)$$

Here $\mathfrak{L}(\mathfrak{H}(b))$ stands for the set of the Laplace transforms of the elements in $\mathfrak{H}(b)$. Because the pair $\{C, A\}$ is observable,

$$\mathfrak{L}(\mathfrak{H}(b)) = \left\{ \frac{p(s)}{\det[sI - A]} : p(s) \text{ is a polynomial of degree } < \dim(\mathcal{X}) \right\}. \quad (4.13)$$

(To see this, if $f(s) \in \mathfrak{L}(\mathfrak{H}(b))$, then $f(s) = \frac{p(s)}{\det[sI - A]}$, where $p(s)$ is a polynomial of degree less than $\dim(\mathcal{X})$. Moreover, the linear spaces in the right hand sides of (4.12) and (4.13) both have dimension n . Therefore they are equal.) In fact, $\det[sI - A] = \prod_{k=1}^n (s - \alpha_k)$. So if $b(s)$ is the Blaschke product of degree n given in (4.11), then

$$\mathfrak{L}(\mathfrak{H}(b)) = \left\{ \frac{p(s)}{\prod_{k=1}^n (s - \alpha_k)} : p(s) \text{ is a polynomial of degree } < n \right\}. \quad (4.14)$$

Let $G(s) = G(\infty) + (\mathfrak{L}g)(s)$ be a function in H^∞ where g is in L^1_+ . Let A be a stable operator on \mathcal{X} . Then

$$\begin{aligned}
(T_G^* e^{A(\cdot)})(t) &= \overline{G(\infty)} e^{At} + \int_t^\infty g(\tau - t)^* e^{A\tau} d\tau \\
&= \overline{G(\infty)} e^{At} + e^{At} \int_0^\infty g(v)^* e^{Av} dv \\
&= \overline{G(\infty)} e^{At} + e^{At} \left[\int_0^\infty g(\tau)^* e^{-s\tau} d\tau \right] \text{evaluated at } -A \\
&= \overline{G(\infty)} e^{At} + e^{At} (\mathfrak{L}(g(\cdot)^*))(-A) \\
&= e^{At} \tilde{G}(-A).
\end{aligned}$$

Hence

$$(T_G^* e^{A(\cdot)})(t) = e^{At} \tilde{G}(\cdot)(-A). \quad (4.15)$$

Let A be a stable dissipative operator on \mathcal{X} . Let C be any operator from \mathcal{X} onto \mathcal{E} such that $A^* + A + C^*C = 0$. Let Γ be the observability operator from \mathcal{X} into $L^2_+(\mathcal{E})$ defined by the pair $\{C, A\}$. Recall that Γ is an isometry. Let $\psi(s)$ be any function in H^∞ . Then

$$(T_\psi^* \Gamma)(t) = C(T_\psi^* e^{A(\cdot)})(t) = C e^{At} \tilde{\psi}(-A) = (\Gamma(t)) \tilde{\psi}(-A).$$

In other words,

$$T_\psi^* \Gamma = \Gamma \tilde{\psi}(-A) \quad \text{and thus} \quad \Gamma^* T_\psi^* \Gamma = \tilde{\psi}(-A). \quad (4.16)$$

Therefore if A is a stable dissipative operator and ψ is a function in H^∞ , then the evaluation $\psi(-A)$ of the function $\psi(s)$ at $-A$ is given by

$$\psi(-A) = \Gamma^* T_\psi^* \Gamma. \quad (4.17)$$

Using this along with (again) the fact that Γ is an isometry, we obtain

$$\|\psi(-A)\| = \|\Gamma^* T_\psi^* \Gamma\| \leq \|T_\psi^*\| = \|\psi\|_\infty.$$

This readily implies that

$$\|\psi(-A)\| \leq \|\psi\|_\infty \quad (\psi \in H^\infty \text{ and } A \text{ is stable and dissipative}). \quad (4.18)$$

In particular, if ψ is an inner function, then $\psi(-A)$ is a contraction. Finally, if ψ and θ are two functions in H^∞ , it follows that

$$(\theta\psi)(-A) = \theta(-A)\psi(-A)$$

Since ψ and θ commute, $\theta(-A)\psi(-A) = \psi(-A)\theta(-A)$.

Recall that if T is a contraction mapping \mathcal{X} into \mathcal{Z} , then D_T is the positive square root of $I - T^*T$, and \mathfrak{D}_T is the (closed) range of D_T . Finally, \mathfrak{d}_T is the dimension of \mathfrak{D}_T . This sets the stage for the following result.

Lemma 4.2. *Let $m(z)$ and $\varphi(z)$ be two finite Blaschke products in H^∞ . Consider the contraction $Y = P_{\mathfrak{H}(\varphi)}|_{\mathfrak{H}(m)}$ mapping $\mathfrak{H}(m)$ into $\mathfrak{H}(\varphi)$. Let $\{A, B, C, D\}$ be a stable dissipative realization for m . Then the following holds.*

1. *There exists a unitary operator Ψ mapping the range of Y onto $\mathfrak{D}_{\tilde{\varphi}(-A)}$ such that*

$$\Psi P_{\mathfrak{H}(\varphi)} \Gamma_m = D_{\tilde{\varphi}(-A)}. \quad (4.19)$$

(Here Γ_m is the observability operator formed by the state space realization $\{A \text{ on } \mathcal{X}, B, C, D\}$ for m .) In this case,

$$\mathfrak{d}_{\tilde{\varphi}(-A)} = \mathfrak{d}_{\varphi(-A)} = \min \{ \deg(\varphi), \deg(m) \} = \text{rank} (P_{\mathfrak{H}(\varphi)} | \mathfrak{H}(m)). \quad (4.20)$$

2. *In particular, $\deg(\varphi) \leq \deg(m)$ if and only if the range of the contraction $Y = P_{\mathfrak{H}(\varphi)} | \mathfrak{H}(m)$ equals $\mathfrak{H}(\varphi)$. In this case, $\mathfrak{d}_{\varphi(-A)} = \mathfrak{d}_{\tilde{\varphi}(-A)} = \deg(\varphi)$, and*

$$\dim(\text{Ker}(Y)) = \deg(m) - \deg(\varphi) \quad (\text{when } \deg(\varphi) \leq \deg(m)). \quad (4.21)$$

3. *The operator Y is one to one if and only if $\deg(m) \leq \deg(\varphi)$. In this case, $\mathfrak{d}_{\varphi(-A)} = \mathfrak{d}_{\tilde{\varphi}(-A)} = \deg(m)$,*

$$\dim(\text{Ker}(Y^*)) = \dim(\text{Im}(Y)^\perp) = \deg(\varphi) - \deg(m). \quad (4.22)$$

4. *If $\deg(\varphi) < \deg(m)$, then the Blaschke product*

$$\varphi(s) = \frac{C(sI - A)^{-1}x}{C(sI - A)^{-1}\tilde{\varphi}(-A)x} \quad (\text{if } 0 \neq x \in \mathfrak{D}_{\tilde{\varphi}(-A)}^\perp). \quad (4.23)$$

Recall that for a Blaschke product φ , we have $\tilde{\varphi}(-s) = \frac{1}{\varphi(s)}$. Hence

$$T_\varphi^* \Gamma_m = \Gamma_m \tilde{\varphi}(-A) = \Gamma_m \varphi^{-1}(A). \quad (4.24)$$

PROOF. Because $\{A \text{ on } \mathcal{X}, B, C, D\}$ is a stable dissipative realization of m , we have $\deg(m) = \dim(\mathcal{X})$. Recall that the Hankel operator $H_m = \Gamma_m \Upsilon_m$ where Γ_m maps \mathcal{X} into L_+^2 is the observability operator formed by $\{C, A\}$. Moreover, Υ_m mapping L_+^2 onto \mathcal{X} is the controllability operator determined by $\{A, B\}$. Furthermore, Γ_m is an isometry and Υ_m is a co-isometry. Since the range of H_m equals $\mathfrak{H}(m)$, it follows that the subspace $\mathfrak{H}(m)$ equals the range of Γ_m . Notice that h is in $\mathfrak{H}(m)$ if and only if $h = \Gamma_m x$ for some x in \mathcal{X} . In fact, this x is uniquely determined by h and given by $x = \Gamma_m^* h$. Using $P_{\mathfrak{H}(\varphi)} = I - T_\varphi T_\varphi^*$ with $T_\varphi^* \Gamma_m = \Gamma_m \tilde{\varphi}(-A)$ and T_φ is an isometry, we have

$$\begin{aligned} \|P_{\mathfrak{H}(\varphi)} \Gamma_m x\|^2 &= \|(I - T_\varphi T_\varphi^*) \Gamma_m x\|^2 = \|\Gamma_m x\|^2 - \|T_\varphi T_\varphi^* \Gamma_m x\|^2 \\ &= \|x\|^2 - \|T_\varphi^* \Gamma_m x\|^2 = \|x\|^2 - \|\Gamma_m \tilde{\varphi}(-A)x\|^2 \\ &= \|x\|^2 - \|\tilde{\varphi}(-A)x\|^2 = \langle x, (I - \tilde{\varphi}(-A)^* \tilde{\varphi}(-A))x \rangle \\ &= \|(I - \tilde{\varphi}(-A)^* \tilde{\varphi}(-A))^{\frac{1}{2}} x\|^2 = \|D_{\tilde{\varphi}(-A)} x\|^2. \end{aligned} \quad (4.25)$$

Hence there exists a unitary operator Ψ mapping the range of Y onto $\mathfrak{D}_{\tilde{\varphi}(-A)}$ such that

$$\Psi P_{\mathfrak{H}(\varphi)} \Gamma_m = D_{\tilde{\varphi}(-A)}. \quad (4.26)$$

This proves equation (4.19) in Part 1.

Now let us show that $\mathfrak{d}_{\tilde{\varphi}(-A)} = \min\{\deg(\varphi), \deg(m)\}$. To this end, first assume that $\deg(\varphi) \leq \deg(m)$. Then we claim that Y is onto $\mathfrak{H}(\varphi)$, and thus, the rank of Y equals $\dim(\mathfrak{H}(\varphi)) = \deg(\varphi)$. Assume that a vector $h \in \mathfrak{H}(\varphi)$ is orthogonal to the range of $Y = P_{\mathfrak{H}(\varphi)} | \mathfrak{H}(m)$. Then it follows from (4.8)

with $\tilde{V} = \varphi$ and $\tilde{W} = m$ that h is also a vector in the range of T_m , that is, $h \in \mathfrak{H}(\varphi) \cap T_m L_{\perp}^2$. By consulting (4.13) or (4.14), we see that the Laplace transform $\mathfrak{L}(\mathfrak{H}(\varphi))$ of the subspace $\mathfrak{H}(\varphi)$ consists of a set of rational functions, with at most $\deg(\varphi) - 1$ zeros. The Laplace transform of h is given by

$$\hat{h}(s) = (\mathfrak{L}h)(s) \in \left(\mathfrak{L}(\mathfrak{H}(\varphi)) \cap mH^2 \right).$$

Since $\deg(\varphi) \leq \deg(m)$, and m is a rational function with $\deg(m)$ zeros, the subspace $\left(\mathfrak{L}(\mathfrak{H}(\varphi)) \cap mH^2 \right) = \{0\}$. Therefore $h = 0$ and the operator Y is onto, whenever $\deg(\varphi) \leq \deg(m)$. This with $\Psi P_{\mathfrak{H}(\varphi)} \Gamma_m = D_{\tilde{\varphi}(-A)}$, implies that $\mathfrak{d}_{\tilde{\varphi}(-A)} = \deg(\varphi)$. Replacing φ with $\tilde{\varphi}$ shows that $\mathfrak{d}_{\varphi(-A)} = \deg(\varphi)$ when $\deg(\varphi) \leq \deg(m)$.

Now assume that $\deg(m) \leq \deg(\varphi)$. Clearly, Y and Y^* have the same rank. Notice that Y^* is the contraction determined by

$$Y^* = P_{\mathfrak{H}(m)} | \mathfrak{H}(\varphi) : \mathfrak{H}(\varphi) \rightarrow \mathfrak{H}(m).$$

So Y^* has the same form as Y , except m and φ interchange places. By our previous analysis $\text{rank}(Y^*) = \deg(m)$ and Y^* is onto $\mathfrak{H}(m)$. So Y is one to one. Recall that $\Psi P_{\mathfrak{H}(\varphi)} \Gamma_m = D_{\tilde{\varphi}(-A)}$. Because Y is one to one, $D_{\tilde{\varphi}(-A)}$ must also be one to one. Since $D_{\tilde{\varphi}(-A)}$ is one to one and $\dim(\mathcal{X}) = \deg(m)$, we see that $\mathfrak{d}_{\tilde{\varphi}(-A)} = \deg(m)$. This completes the proof of Part 1.

To prove Part 2, we showed that if $\deg(\varphi) \leq \deg(m)$ then Y is onto $\mathfrak{H}(\varphi)$. Moreover, $\dim(\text{Ker}(Y)) + \dim(\text{Im}(Y)) = \dim(\mathfrak{H}(m))$, which proves (4.21). On the other hand if $\deg(\varphi) > \deg(m)$, then $\text{rank } Y = \text{rank } Y^* = \deg m$ and hence Y is not onto $\mathfrak{H}(\varphi)$.

Part 3 is proven in the same way by replacing Y by Y^* .

To establish Part 4, assume that $\deg(\varphi) < \deg(m)$. Then there exists a nonzero x such that $D_{\tilde{\varphi}(-A)}x = 0$, or equivalently, $x = \tilde{\varphi}(-A)^* \tilde{\varphi}(-A)x$. Using

$$\Psi P_{\mathfrak{H}(\varphi)} \Gamma_m x = D_{\tilde{\varphi}(-A)} x = 0,$$

we have $P_{\mathfrak{H}(\varphi)} \Gamma_m x = 0$. By employing $P_{\mathfrak{H}(\varphi)} = I - T_{\varphi} T_{\varphi}^*$, we obtain

$$0 = P_{\mathfrak{H}(\varphi)} \Gamma_m x = (I - T_{\varphi} T_{\varphi}^*) \Gamma_m x = \Gamma_m x - T_{\varphi} \Gamma_m \tilde{\varphi}(-A)x.$$

In other words, $\Gamma_m x = T_{\varphi} \Gamma_m \tilde{\varphi}(-A)x$. By taking the Laplace transform, we arrive at

$$C(sI - A)^{-1}x = \varphi(s)C(sI - A)^{-1}\tilde{\varphi}(-A)x. \quad (4.27)$$

Since Γ_m is one to one and $\tilde{\varphi}(-A)x$ is nonzero, $C(sI - A)^{-1}\tilde{\varphi}(-A)x$ is a nonzero function in H^2 . Hence the rational function $C(sI - A)^{-1}\tilde{\varphi}(-A)x$ is nonzero. Dividing (4.27) by $C(sI - A)^{-1}\tilde{\varphi}(-A)x$, yields the formula that we have been looking for, that is,

$$\varphi(s) = \frac{C(sI - A)^{-1}x}{C(sI - A)^{-1}\tilde{\varphi}(-A)x}.$$

This completes the proof. □

Remark 4.3. Let R be the rational rigid function in L^∞ defined by $R(i\omega) = \varphi(i\omega)\overline{m(i\omega)}$, where φ and m are two Blaschke products. Let T_R be the Wiener-Hopf operator on L^2_+ determined by R . By consulting Proposition 4.1 and Lemma 4.2, we readily obtain the following.

1. The operator T_R is invertible if and only if $\deg(\varphi) = \deg(m)$, that is, the numbers of zeros and poles in \mathbb{C}_+ of $R(s)$ are equal.
2. The kernel of T_R is nonzero if and only if $\deg(\varphi) < \deg(m)$. In this case, the operator T_R is onto L^2_+ and $\dim(\text{Ker}(T_R)) = \deg(m) - \deg(\varphi)$. The number of poles in \mathbb{C}_+ of $R(s)$ is higher than the number of zeros.
3. The subspace $\text{Im}(T_R)^\perp$ is nonzero if and only if $\deg(m) < \deg(\varphi)$. In this case, the range of the operator T_R is closed, $\text{Ker}(T_R) = \{0\}$ and $\dim(\text{Im}(T_R)^\perp) = \deg(\varphi) - \deg(m)$.

Corollary 4.4. Let A be a stable dissipative operator on a finite dimensional space \mathcal{X} such that $A^* + A$ has rank one, and φ a rational Blaschke product in H^∞ . Then

$$\mathfrak{d}_{\varphi(-A)} = \mathfrak{d}_{\varphi(-A^*)} = \min\{\deg(\varphi), \dim(\mathcal{X})\}. \quad (4.28)$$

PROOF. Let C be any operator mapping \mathcal{X} into \mathbb{C} such that $A^* + A = -C^*C$. Set $B = -C^*$ mapping \mathbb{C} into \mathcal{X} and $D = 1$. Then $\{A, B, C, D\}$ is a dissipative realization for a Blaschke product $m(z)$ with degree $\dim(\mathcal{X})$; see Theorem 2.1. Applying Lemma 4.2 yields (4.28). \square

Proposition 4.5. Let A be a stable dissipative operator on \mathcal{X} such that the rank of $A^* + A$ equals 1. Let φ be a Blaschke product where $\deg(\varphi) < \dim(\mathcal{X})$. Then the inner function φ is given by

$$\varphi(s) = \frac{C(sI - A)^{-1}\varphi(-A^*)x}{C(sI - A)^{-1}x} \quad (\text{where } x = \varphi(-A^*)^*\varphi(-A^*)x \text{ and } x \neq 0). \quad (4.29)$$

Here C is any operator mapping \mathcal{X} into \mathbb{C} such that $A^* + A = -C^*C$. If θ is any function in H^∞ such that $\theta(-A) = \varphi(-A)$ and $\|\theta\|_\infty \leq 1$, then $\theta(s) = \varphi(s)$.

PROOF. Formula (4.29) will be derived from formula (4.23). To this end, recall that if Z is a contraction, then Z is a unitary operator from $\mathfrak{D}_{\frac{1}{Z}}$ onto $\mathfrak{D}_{\frac{1}{Z^*}}$. Hence $\tilde{\varphi}(-A)$ is a unitary operator from $\mathfrak{D}_{\tilde{\varphi}(-A)}^\perp$ onto $\mathfrak{D}_{\varphi(-A^*)}^\perp$. In particular,

$$\tilde{\varphi}(-A) : \mathfrak{D}_{\tilde{\varphi}(-A)}^\perp \rightarrow \mathfrak{D}_{\varphi(-A^*)}^\perp$$

is unitary. So for $x \neq 0$ in $\mathfrak{D}_{\tilde{\varphi}(-A)}^\perp$, we see that $y = \tilde{\varphi}(-A)x$ is in $\mathfrak{D}_{\varphi(-A^*)}^\perp$ and $x = \tilde{\varphi}(-A)^*y$. By interchanging the roles of x and y , we see that (4.23) yields (4.29) and vice-versa.

Let us show that if θ is a function H^∞ such that $\theta(-A^*) = \varphi(-A^*)$ and the H^∞ norm $\|\theta\|_\infty \leq 1$, then $\theta(s) = \varphi(s)$.

Assume that θ is a function in H^∞ such that $\tilde{\theta}(-A) = \tilde{\varphi}(-A)$ and $\|\theta\|_\infty \leq 1$. By taking the adjoint, we see that $\theta(-A^*) = \varphi(-A^*)$. In this case,

$$T_\theta^*\Gamma = \Gamma\tilde{\theta}(-A) = \Gamma\tilde{\varphi}(-A).$$

Since $T_\varphi^* \Gamma = \Gamma \tilde{\varphi}(-A)$, we also have $T_\varphi^* \Gamma = T_\theta^* \Gamma$, or equivalently, $\Gamma^* T_\varphi = \Gamma^* T_\theta$. Multiplying by Γ on both sides, we obtain with x as above

$$\Gamma \Gamma^* T_\theta \Gamma x = \Gamma \Gamma^* T_\varphi \Gamma x = \Gamma \varphi(-A^*) x.$$

Because $\Gamma \Gamma^*$ is an orthogonal projection, $\|\varphi(-A^*) x\|^2 = \|x\|^2$ and T_θ is a contraction, we see that

$$\|x\| \geq \|T_\theta \Gamma x\| \geq \|\Gamma \Gamma^* T_\theta \Gamma x\| = \|\Gamma \varphi(-A^*) x\| = \|\varphi(-A^*) x\| = \|x\|.$$

Therefore we have equality, and thus,

$$T_\theta \Gamma x = \Gamma \Gamma^* T_\theta \Gamma x = \Gamma \varphi(-A^*) x \quad (\text{when } 0 \neq x \in \mathfrak{D}_{\varphi(-A^*)}^\perp).$$

By taking the Laplace transform of both sides, and using (4.29), we obtain

$$\theta(s) = \frac{C(sI - A)^{-1} \varphi(-A^*) x}{C(sI - A)^{-1} x} = \varphi(s).$$

Therefore $\theta(s) = \varphi(s)$. In other words, if θ is a function in H^∞ such that $\tilde{\theta}(-A) = \tilde{\varphi}(-A)$ and $\|\theta\|_\infty \leq 1$, then $\theta(s) = \varphi(s)$. Replacing $\tilde{\varphi}$ by φ shows that if θ is a function in H^∞ such that $\theta(-A) = \varphi(-A)$ and $\|\theta\|_\infty \leq 1$, then $\tilde{\theta}(s) = \tilde{\varphi}(s)$, and thus, $\theta(s) = \varphi(s)$. \square

Finally, it is noted that this result is an application of the Sz.-Nagy-Foias commutant lifting theorem; see Corollary 2.7 page 142 of [11] and is also deeply connected to some of the results in [1]. See also [10].

Proposition 4.6. *Let A be a stable dissipative operator on a finite dimensional space \mathcal{X} and φ a Blaschke product. If $\deg(\varphi) \geq \dim(\mathcal{X})$, then the defect index for $\varphi(-A)$ equals $\dim(\mathcal{X})$, or equivalently, $\varphi(-A)^* \varphi(-A)$ does not have 1 as an eigenvalue.*

PROOF. Let C be any operator from \mathcal{X} onto \mathcal{E} such that $A^* + A = -C^* C$. Then the observability operator Γ formed by $\{C, A\}$ is an isometry from \mathcal{X} into $L_+^2(\mathcal{E})$. Recall that if $u(z)$ is any function in H^∞ , then $\|u(-A)\| \leq \|u\|_\infty$; see (4.18). Because φ is an inner function, $\tilde{\varphi}(-A)$ is a contraction. For x in \mathcal{X} , we have

$$\begin{aligned} \|(I - \tilde{\varphi}(-A)^* \tilde{\varphi}(-A))^{\frac{1}{2}} x\|^2 &= \langle (I - \tilde{\varphi}(-A)^* \tilde{\varphi}(-A)) x, x \rangle \\ &= \|x\|^2 - \|\tilde{\varphi}(-A) x\|^2 = \|\Gamma x\|^2 - \|\tilde{\varphi}(-A) x\|^2 \\ &= \|\Gamma x\|^2 - \|T_{\varphi I} T_{\varphi I}^* \Gamma x\|^2 = \|(I - T_{\varphi I} T_{\varphi I}^*) \Gamma x\|^2 = \|P_{\mathfrak{R}(\varphi I)} \Gamma x\|^2. \end{aligned}$$

Hence there exists a unitary operator Ψ from the range of $P_{\mathfrak{R}(\varphi I)} \Gamma$ onto the range of the defect operator $(I - \tilde{\varphi}(-A)^* \tilde{\varphi}(-A))^{\frac{1}{2}}$ such that

$$\Psi P_{\mathfrak{R}(\varphi I)} \Gamma = (I - \tilde{\varphi}(-A)^* \tilde{\varphi}(-A))^{\frac{1}{2}}. \quad (4.30)$$

In particular, the defect index $\mathfrak{d}_{\tilde{\varphi}(-A)}$ equals the rank of $P_{\mathfrak{R}(\varphi I)} \Gamma$.

Notice that x is in the kernel of $P_{\mathfrak{R}(\varphi I)} \Gamma$ if and only if x is a vector with eigenvalue 1 for $\tilde{\varphi}(-A)^* \tilde{\varphi}(-A)$. In this case,

$$0 = P_{\mathfrak{R}(\varphi I)} \Gamma x = \Gamma x - T_{\varphi I} T_{\varphi I}^* \Gamma x.$$

Hence

$$\Gamma x = T_{\varphi I} \Gamma \tilde{\varphi}(-A)x \quad (\text{for } x \in \mathfrak{D}_{\tilde{\varphi}(-A)}^{\perp}).$$

By taking the Laplace transform

$$C(sI - A)^{-1}x = \varphi(s)C(sI - A)^{-1}\tilde{\varphi}(-A)x \quad (\text{for } x \in \mathfrak{D}_{\tilde{\varphi}(-A)}^{\perp}). \quad (4.31)$$

Because $\varphi(s)$ has all its zeroes in the open right half plane, and the eigenvalues of A are in the open left half plane, the numerator of $C(sI - A)^{-1}x$ has at least $\deg(\varphi)$ zeros. However, the numerator of $C(sI - A)^{-1}x$ is a polynomial of degree at most $\dim(\mathcal{X}) - 1 < \deg(\varphi)$. Therefore $x = 0$, and $\tilde{\varphi}(-A)^* \tilde{\varphi}(-A)$ has no eigenvalue on the unit circle. The same argument applies by replacing $\tilde{\varphi}(s)$ by $\varphi(s)$. This completes the proof. \square

Winding numbers in the scalar case. Let R be the rigid function in L^{∞} defined by $R(i\omega) = \varphi(i\omega)\overline{m(i\omega)} = (\varphi m^*)(i\omega)$, where both φ and m are Blaschke products. The *winding number* for R is given by $\deg(\varphi) - \deg(m)$. As before, let Y be the contraction from $\mathfrak{H}(m)$ into $\mathfrak{H}(\varphi)$ given by $Y = P_{\mathfrak{H}(\varphi)}|_{\mathfrak{H}(m)}$. By consulting Lemma 4.2, we see that

1. The winding number for $R = \varphi m^*$ equals $\dim(\text{Im}(Y)^{\perp})$ when $\deg(m) \leq \deg(\varphi)$.
2. The winding number for $R = \varphi m^*$ equals $-\dim(\text{Ker}(Y))$ when $\deg(\varphi) \leq \deg(m)$.
3. Hence the winding number for $R = \varphi m^*$ equals $-\text{ind}(Y)$ where $\text{ind}(Y)$ denotes the Fredholm index ($\dim(\text{Ker}(Y)) - \dim(\text{Ker}(Y^*))$) for Y .

Recall that the Wiener-Hopf operator T_R admits a decomposition of the form:

$$T_R = T_{\varphi} T_m^* + H_{\varphi} Y H_m^*;$$

see (4.4). Here Y is the finite dimensional contraction from $\mathfrak{H}(\tilde{m})$ into $\mathfrak{H}(\tilde{\varphi})$ determined by $Y = P_{\mathfrak{H}(\tilde{\varphi})}|_{\mathfrak{H}(\tilde{m})}$. Hence T_R is Fredholm. Moreover, T_R and Y have the same Fredholm index. Hence

$$\text{ind}(T_R) = \dim(\text{Ker}(T_R)) - \dim(\text{Ker}(T_R^*)) = \text{ind}(Y) = \deg(m) - \deg(\varphi). \quad (4.32)$$

The Fredholm index of T_R equals minus the winding number of $R = \varphi m^*$.

We say that a polynomial $p(s)$ is *stable* if all the roots of $p(s)$ are contained in the open left hand plane \mathbb{C}_- . Let $p^{\sharp}(s)$ be the polynomial defined by

$$p^{\sharp}(s) = \overline{p(-\bar{s})}.$$

It is noted that if $p(s)$ is a stable polynomial of degree n , then $\varphi(s) = \frac{p^{\sharp}(s)}{p(s)}$ is a Blaschke product of order n . In particular, if A is a stable dissipative operator and $A^* + A$ has rank one, then

$$\varphi(-A) = p^{\sharp}(s)(-A)(p(-A))^{-1}$$

is a strict contraction; see Corollary 4.4. In other words, if $p(s)$ is a stable polynomial of degree n , then

$$p(-A)^* p(-A) - (p^{\sharp}(-A))^* p^{\sharp}(s)(-A)$$

is a strictly positive.

The following is a continuous time version of the Schur-Cohen stability test.

Remark 4.7. Assume that $p(s)$ is a polynomial of degree n and let

$$R(s) = \frac{p^\sharp(s)}{p(s)}.$$

Then the following statements are equivalent.

1. The polynomial $p(s)$ is stable.
2. The winding number for $R(s)$ equals n .
3. The Fredholm index for T_R equals $-n$.
4. If A is a stable dissipative operator on a n dimensional space and the rank of $A + A^*$ equals 1, then

$$p(-A)^*p(-A) - (p^\sharp(-A))^*p^\sharp(-A)$$

is a strictly positive.

Proof. First observe that if q is a stable polynomial of degree strictly less than n and $\theta(s) = \frac{q^\sharp(s)}{q(s)}$, then the norm of $\theta(-A)$ equals 1, or equivalently,

$$q(-A)^*q(-A) - (q^\sharp(-A))^*q^\sharp(-A)$$

is positive and singular; see Corollary 4.4.

Now let $p(s)$ be any polynomial of degree n . Then $p(s)$ admits a factorization of the form: $p(s) = q(s)u(s)$ where $q(s)$ is stable and $u(s)$ is unstable. Then

$$\begin{aligned} & p(-A)^*p(-A) - (p^\sharp(-A))^*p^\sharp(-A) = \\ & u(-A)^*q(-A)^*q(-A)u(-A) - (q^\sharp(-A))^*(u^\sharp(-A))^*q^\sharp(-A)u^\sharp(-A) \end{aligned}$$

Next we prove that item (1) implies item (4). Since R is a Blaschke product we know that $R(-A)$ is a contraction. (See text after formula (4.19)). Moreover equality (4.28) shows that the defect index of $R(-A)$ is equal to n . Thus we have that $R^*(-A)R(-A) < I$ or $I - R^*(-A)R(-A) > 0$. Multiplying the last inequality by $p^*(-A)p(-A)$ gives that

$$p(-A)^*p(-A) - (p^\sharp(-A))^*p^\sharp(-A)$$

is a strictly positive.

To prove that item (4) implies item (1) assume that p is not stable and $p(-A)^*p(-A) - (p^\sharp(-A))^*p^\sharp(-A) > 0$. We will show that this leads to an obvious contradiction. Write $p = p_1p_2^\sharp$ with p_1 and p_2 both stable polynomials. The assumption gives that p_2 is not a constant. Note that $p(-A)$ must be invertible. It follows that the zeros of p and the eigenvalues of $-A$ are disjoint sets. Write

$$B_i(s) = p_i^\sharp(s)p_i(s)^{-1} \quad (i = 1, 2), \text{ and } R(s) = B_1(s)B_2(s)^{-1}.$$

Since B_i is a Blaschke product, we have that $B_i(-A)$ is a contraction. Also our assumption gives that $B_1(-A)^*B_1(-A) < B_2(-A)^*B_2(-A)$. As the degree

of B_1 is strictly less than n , also the defect index, which is the minimum of n and the degree of B_1 , is strictly less than n . Therefore there exists a vector v such that

$$v^* B_1(-A)^* B_1(-A)v = v^* v. \quad (4.33)$$

On the other hand

$$v^* B_1(-A)^* B_1(-A)v < v^* B_2(-A)^* B_2(-A)v \leq v^* v. \quad (4.34)$$

The two equalities (4.33) and (4.34) contradict and therefore p_2 is a constant. We conclude that p is stable. \square

A discrete time version of part of Remark 4.7 with a different type proof is presented in [22].

Finally, for an example of a stable dissipative matrix A such that $A + A^*$ has rank one, consider the lower triangular Toeplitz matrix A on \mathbb{C}^n given by

$$A = \begin{bmatrix} -1 & 0 & 0 & 0 & \cdots & 0 \\ 2 & -1 & 0 & 0 & \cdots & 0 \\ -2 & 2 & -1 & 0 & \cdots & 0 \\ 2 & -2 & 2 & -1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ (-1)^n 2 & (-1)^{n-1} 2 & (-1)^{n-2} 2 & (-1)^{n-3} 2 & \cdots & -1 \end{bmatrix}$$

$$C = \sqrt{2} [1 \quad -1 \quad 1 \quad -1 \quad \cdots \quad (-1)^{n-1}].$$

Notice that A is the lower triangular Toeplitz matrix formed by

$$\{-1, 2, -2, 2, -2, \dots, 2(-1)^n\}.$$

Then A is a stable dissipative matrix and the rank of $A + A^*$ equals one. In fact, $A + A^* + C^*C = 0$. This matrix is motivated by equation (7.24) below.

5. Point evaluation and MIMO systems

Let us return to the MIMO case and provide a proof of one of our main results, Theorem 2.2.

As before, assume that V and W are two bi-inner rational functions in $H^\infty(\mathcal{E}, \mathcal{E})$. Let $\{A_v$ on $\mathcal{X}_v, B_v, C_v, D_v\}$ and $\{A_w$ on $\mathcal{X}_w, B_w, C_w, D_w\}$ be stable dissipative realizations of V and W , respectively. Recall that $R = VW^*$ and the Wiener-Hopf operator T_R on $L_+^2(\mathcal{E})$ admits a decomposition of the form:

$$T_R = T_V T_W^* + H_V Y H_W^*. \quad (5.1)$$

Here Y is the contraction defined by

$$Y = P_{\mathfrak{H}(\tilde{V})} | \mathfrak{H}(\tilde{W}) : \mathfrak{H}(\tilde{W}) \rightarrow \mathfrak{H}(\tilde{V}). \quad (5.2)$$

Recall that Proposition 4.1 shows that the dimensions of $\text{Ker}(Y)$ and $\text{Ker}(T_R)$ are equal. Therefore we are interested in calculating the dimension of $\text{Ker}(Y)$

in terms of the realizations of V and W . Since $H_W = \Gamma_w \Upsilon_w$, and the range of the Hankel operator H_W^* is given by $\text{Im}(H_W^*) = \mathfrak{H}(\widetilde{W})$, we have

$$\mathfrak{H}(\widetilde{W}) = \text{Im}(\Upsilon_w^*) \quad \text{where} \quad \Upsilon_w^* = B_w^* e^{A_w^* t} : \mathcal{X}_w \rightarrow L_+^2(\mathcal{E}).$$

Furthermore, Υ_w is a co-isometry and A_w^* is a stable dissipative operator on \mathcal{X}_w satisfying $A_w^* + A_w + B_w B_w^* = 0$, where B_w is an operator mapping \mathcal{E} into \mathcal{X}_w . Using the state space realization $V(s) = D_v + C_v(sI - A_v)^{-1}B_v$, we have

$$\begin{aligned} \left(T_{\widetilde{V}}^* \Upsilon_w^*\right)(t) &= (D_v \Upsilon_w^*)(t) + \int_t^\infty C_v e^{A_v(\tau-t)} B_v B_w^* e^{A_w^* \tau} d\tau \\ &= (D_v \Upsilon_w^*)(t) + \left(\int_0^\infty C_v e^{A_v \tau} B_v B_w^* e^{A_w^* \tau} d\tau\right) e^{A_w^* t} \\ &= \left(D_v B_w^* + C_v \int_0^\infty e^{A_v \tau} B_v B_w^* e^{A_w^* \tau} d\tau\right) e^{A_w^* t}. \end{aligned}$$

In other words,

$$\left(T_{\widetilde{V}}^* \Upsilon_w^*\right)(t) = \left(D_v B_w^* + C_v \int_0^\infty e^{A_v \tau} B_v B_w^* e^{A_w^* \tau} d\tau\right) e^{A_w^* t}. \quad (5.3)$$

Let C_o be the operator mapping \mathcal{X}_w into \mathcal{E} defined by

$$C_o = D_v B_w^* + C_v \int_0^\infty e^{A_v \tau} B_v B_w^* e^{A_w^* \tau} d\tau. \quad (5.4)$$

Because the operators A_v and A_w are both stable, the operator C_o is well defined. In fact, C_o can be computed by first solving the Lyapunov equation

$$A_v \Omega + \Omega A_w^* + B_v B_w^* = 0. \quad (5.5)$$

Because A_v and A_w are stable, the solution Ω to this Lyapunov equation is unique and given by

$$\Omega = \int_0^\infty e^{A_v \tau} B_v B_w^* e^{A_w^* \tau} d\tau. \quad (5.6)$$

Therefore

$$C_o = D_v B_w^* + C_v \Omega. \quad (5.7)$$

Let Γ_o be the operator from \mathcal{X}_w into $L_+^2(\mathcal{E})$ defined by $\Gamma_o = T_{\widetilde{V}}^* \Upsilon_w^*$. By consulting (5.3) with the definition of C_o in (5.4), we see that

$$(\Gamma_o x)(t) = \left(T_{\widetilde{V}}^* \Upsilon_w^* x\right)(t) = C_o e^{A_w^* t} x \quad (x \in \mathcal{X}_w). \quad (5.8)$$

Notice that Γ_o is the observability operator determined by the pair $\{C_o, A_w^*\}$.

It is emphasized that if $V = \varphi I$ where φ is an inner function in H^∞ , then $C_o = B_w^* \varphi(-A_w^*)$. In other words, in the scalar case

$$\Gamma_o = T_{\widetilde{\varphi}I}^* \Upsilon_w^* = \Upsilon_w^* \varphi(-A_w^*) \quad (5.9)$$

and the operator C_o plays the role of $B_w^* \varphi(-A_w^*)$ in the MIMO case. Note that in the general case, C_o is the left point evaluation of $-A_w^*$ with respect to B_w^* . For a further discussion on MIMO function evaluation with applications to H^∞ interpolation theory, see Section 1.2 page 15 of [11].

The next lemma with $\mathfrak{n}(T_R) = \mathfrak{n}(Y)$ (see Part 2 of Proposition 4.1) provides the proof of Part 1 and the first half of Part 2 of Theorem 2.2.

Lemma 5.1. *Let V and W be bi-inner rational functions in $H^\infty(\mathcal{E}, \mathcal{E})$. Consider the contraction Y mapping $\mathfrak{H}(\widetilde{W})$ into $\mathfrak{H}(\widetilde{V})$ determined by*

$$Y = P_{\mathfrak{S}(\widetilde{V})} | \mathfrak{H}(\widetilde{W}) : \mathfrak{H}(\widetilde{W}) \rightarrow \mathfrak{H}(\widetilde{V}). \quad (5.10)$$

Let C_\circ be the operator from \mathcal{X}_w into \mathcal{E} defined by (5.4) or (5.5) and (5.7). Let Q be the unique solution to the Lyapunov equation

$$A_w Q + Q A_w^* + C_\circ^* C_\circ = 0. \quad (5.11)$$

Then Q is a positive contraction. Moreover, Y and $I - Q$ have the same rank. In particular,

$$\text{Ker}(Y) = \Upsilon_w^* \text{Ker}(I - Q) \quad \text{and} \quad \dim(\text{Ker}(Y)) = \dim(\text{Ker}(I - Q)). \quad (5.12)$$

Furthermore, we have

$$\dim(\text{Ker}(Y^*)) = \dim(\mathfrak{H}(\widetilde{V})) - \text{rank}(I - Q). \quad (5.13)$$

If $V = \varphi I$ where φ is a Blaschke product, then $Q = \varphi(-A_w^*)^* \varphi(-A_w^*)$.

Proof. Notice that $Q = \Gamma_\circ^* \Gamma_\circ$ is the unique solution to the Lyapunov equation

$$A_w Q + Q A_w^* + C_\circ^* C_\circ = 0.$$

Recall that the orthogonal projection $P_{\mathfrak{S}(\widetilde{V})} = I - T_{\widetilde{V}} T_{\widetilde{V}}^*$. Moreover, $\mathfrak{H}(\widetilde{W})$ equals the range of the isometry Υ_w^* . So h is in $\mathfrak{H}(\widetilde{W})$ if and only if $h = \Upsilon_w^* x$ for some x in \mathcal{X}_w . In fact, $x = \Upsilon_w h$. Now observe that

$$\begin{aligned} \|P_{\mathfrak{S}(\widetilde{V})} \Upsilon_w^* x\|^2 &= \|(I - T_{\widetilde{V}} T_{\widetilde{V}}^*) \Upsilon_w^* x\|^2 = \|\Upsilon_w^* x\|^2 - \|T_{\widetilde{V}} T_{\widetilde{V}}^* \Upsilon_w^* x\|^2 \\ &= \|\Upsilon_w^* x\|^2 - \|T_{\widetilde{V}}^* \Upsilon_w^* x\|^2 = \|x\|^2 - \|\Gamma_\circ x\|^2 = \|x\|^2 - \langle Qx, x \rangle. \end{aligned}$$

(The fourth equality follows from (5.8).) Hence Q is a positive contraction, and thus,

$$\|P_{\mathfrak{S}(\widetilde{V})} \Upsilon_w^* x\|^2 = \|(I - Q)^{\frac{1}{2}} x\|^2 \quad (x \in \mathcal{X}).$$

So there exists a unitary operator Ψ mapping the range of $P_{\mathfrak{S}(\widetilde{V})} \Upsilon_w^*$ onto the range $(I - Q)^{\frac{1}{2}}$ such that

$$\Psi Y \Upsilon_w^* x = \Psi P_{\mathfrak{S}(\widetilde{V})} \Upsilon_w^* x = (I - Q)^{\frac{1}{2}} x \quad (\text{for } x \in \mathcal{X}_w). \quad (5.14)$$

In particular, Y and $I - Q$ have the same rank. Recall that $\mathfrak{H}(\widetilde{W})$ equals the range of Υ_w^* and that $Y = P_{\mathfrak{S}(\widetilde{V})} | \mathfrak{H}(\widetilde{W})$. Therefore, by (5.14)

$$\text{Ker}(Y) = \Upsilon_w^* \text{Ker}(I - Q) \quad \text{and} \quad \dim(\text{Ker}(Y)) = \dim(\text{Ker}(I - Q)).$$

In other words, $\dim(\text{Ker}(Y))$ equals the number of eigenvalues of Q equal to 1 counting multiplicities.

By applying the previous result to Y^* , we obtain

$$\dim(\text{Ker}(Y^*)) = \dim(\mathfrak{H}(\widetilde{V})) - \text{rank}(I - Q).$$

Finally, it is noted that if $V = \varphi I$ where φ is an inner function in H^∞ , then

$$\Gamma_\circ = \Upsilon_w^* \varphi(-A_w^*);$$

see (5.9). Because Υ_w is a co-isometry, we have

$$Q = \Gamma_\circ^* \Gamma_\circ = \varphi(-A_w^*)^* \Upsilon_w \Upsilon_w^* \varphi(-A_w^*) = \varphi(-A_w^*)^* \varphi(-A_w^*).$$

□

The following lemma implies Part 2 of Theorem 2.2.

Lemma 5.2. *Let V and W be two rational bi-inner functions in $H^\infty(\mathcal{E}, \mathcal{E})$, and ψ be an inner function in H^∞ . Let Y_ψ be the contraction from $\mathfrak{H}(\widetilde{W})$ into $\mathfrak{H}(\psi\widetilde{V})$ defined by $Y_\psi = P_{\mathfrak{H}(\psi\widetilde{V})} | \mathfrak{H}(\widetilde{W})$. Then $\dim(\text{Ker}(Y_\psi))$ is equal to the multiplicity of 1 as an eigenvalue of $\psi(-A_w)Q\psi(-A_w)^*$. Also*

$$\dim(\text{Ker}(Y_\psi^*)) = \deg(\psi) \dim \mathcal{E} + \dim(\mathfrak{H}(\widetilde{V})) - \text{rank}(I - \psi(-A_w)Q\psi(-A_w)^*).$$

*In particular, if $\zeta(s) = \frac{1-s}{1+s}$, then $\dim(\text{Ker}(Y_{\zeta^k}))$ is equal to the multiplicity of 1 as an eigenvalue of $\zeta(-A_w)^k Q \zeta(-A_w)^{*k}$. Also*

$$\dim(\text{Ker}(Y_{\zeta^k}^*)) = k \dim(\mathcal{E}) + \dim(\mathfrak{H}(\widetilde{V})) - \text{rank}(I - \zeta(-A_w)^k Q \zeta(-A_w)^{*k}).$$

Proof. Set $\Gamma = \Upsilon_w^*$. Then for x in \mathcal{X}_w we have

$$\|P_{\mathfrak{H}(\psi\widetilde{V})} \Gamma x\|^2 = \|(I - T_{\psi\widetilde{V}} T_{\psi\widetilde{V}}^*) \Gamma x\|^2 = \|\Gamma x\|^2 - \|T_{\psi\widetilde{V}} T_{\psi\widetilde{V}}^* \Gamma x\|^2.$$

Since $T_{\psi\widetilde{V}}$ is an isometry, and using (5.8), we have

$$\begin{aligned} \|P_{\mathfrak{H}(\psi\widetilde{V})} \Gamma x\|^2 &= \|\Gamma x\|^2 - \|T_{\psi\widetilde{V}}^* \Gamma x\|^2 = \|\Gamma x\|^2 - \|T_{\widetilde{V}}^* T_\psi^* \Gamma x\|^2 \\ &= \|\Gamma x\|^2 - \|T_{\widetilde{V}}^* \Gamma \psi(-A_w)^* x\|^2 \\ &= \|\Gamma x\|^2 - \|\Gamma_\circ \psi(-A_w)^* x\|^2. \end{aligned}$$

Recall that $\Gamma_\circ^* \Gamma_\circ = Q$. Therefore

$$\begin{aligned} \|P_{\mathfrak{H}(\psi\widetilde{V})} \Gamma x\|^2 &= \|x\|^2 - \langle \psi(-A_w)Q\psi(-A_w)^* x, x \rangle \\ &= \|(I - \psi(-A_w)Q\psi(-A_w)^*)^{\frac{1}{2}} x\|^2. \end{aligned}$$

So there exists a unitary operator Ψ mapping the range of $P_{\mathfrak{H}(\psi\widetilde{V})} \Gamma$ onto the range of $(I - \psi(-A_w)Q\psi(-A_w)^*)^{\frac{1}{2}}$ such that

$$\Psi P_{\mathfrak{H}(\psi\widetilde{V})} \Gamma x = (I - \psi(-A_w)Q\psi(-A_w)^*)^{\frac{1}{2}} x \quad (\text{for } x \in \mathcal{X}_w).$$

Recall that $\mathfrak{H}(\widetilde{W}) = \Gamma \mathcal{X}_w$. Therefore, when $Y_\psi = P_{\mathfrak{H}(\psi\widetilde{V})} | \mathfrak{H}(\widetilde{W})$, we have

$$\dim \left(\text{Ker} \left(P_{\mathfrak{H}(\psi\widetilde{V})} | \mathfrak{H}(\widetilde{W}) \right) \right) = \dim \left(\text{Ker} \left(I - \psi(-A_w)Q\psi(-A_w)^* \right) \right).$$

In other words, the dimension of $\text{Ker}(Y_\psi)$ is equal to the multiplicity of 1 as an eigenvalue of $\psi(-A_w)Q\psi(-A_w)^*$.

Using $\mathfrak{H}(\psi\widetilde{V}) = \mathfrak{H}(\psi I_{\mathcal{E}}) \oplus T_\psi \mathfrak{H}(\widetilde{V})$, we also have

$$\dim \left(\text{Ker}(Y_\psi^*) \right) = \deg(\psi) \dim(\mathcal{E}) + \dim(\mathfrak{H}(\widetilde{V})) - \text{rank}(I - \psi(-A_w)Q\psi(-A_w)^*).$$

□

The previous lemma with $\mathbf{n}(T_{\zeta^k R}) = \mathbf{n}(Y_{\zeta^k})$, yields

$$\mathbf{n}(T_{\zeta^k R}) = \dim (\text{Ker} (I - \zeta(-A_w)^k Q \zeta(-A_w)^{*k})).$$

This completes the proof of Part 2 of Theorem 2.2.

Since Part 3 of Theorem 2.2 follows from the Parts 1 and 2 the proof of Theorem 2.2 is complete now.

The next Corollary provides similar formulas for the positive Wiener-Hopf indices. First we define C_{o*} as follows. Solve the Lyapunov equation

$$A_w \Omega_* + \Omega_* A_v^* + B_w B_v^* = 0. \quad (5.15)$$

(Note that $\Omega_* = \Omega_*^*$.) Because A_v and A_w are stable the solution Ω_* to this Lyapunov equation is unique and given by

$$\Omega_* = \int_0^\infty e^{A_w t} B_w B_v^* e^{A_v t} dt. \quad (5.16)$$

Put

$$C_{o*} = D_w B_v^* + C_w \Omega_*. \quad (5.17)$$

Now let Q_* be the unique solution of the Lyapunov equation

$$A_v Q_* + Q_* A_v^* + C_{o*}^* C_{o*} = 0. \quad (5.18)$$

Corollary 5.3. *Let V and W be given by (2.1) and (2.2) and $R = WV^*$. Furthermore let $\zeta(s) = \frac{1-s}{1+s}$ and Q_* be defined by (5.18). Put*

$$\begin{aligned} \nu_k = & \dim(\text{Ker} (I - \zeta(-A_v)^{k-1} Q_* \zeta(-A_v)^{*k-1})) + \\ & - \dim(\text{Ker} (I - \zeta(-A_v)^k Q_* \zeta(-A_v)^{*k})). \end{aligned} \quad (5.19)$$

Then the positive Wiener-Hopf indices $\omega_1, \dots, \omega_q$ of T_R are given by

$$\omega_j = \#\{k \mid \nu_k \geq j\}, \quad (j = 1, \dots, q = \nu_1).$$

Proof. Notice $R^* = WV^*$. Recall $R^*(s)$ is defined by $R^*(s) = (R(-\bar{s}))^*$. If

$$R(s) = W_-(s) \text{diag} (\zeta(s)^{\kappa_j})_{j=1}^m W_+(s),$$

with W_+ and its inverse are analytic on the right half plane \mathbb{C}_+ and W_- and its inverse are analytic on the left half plane, \mathbb{C}_- , then

$$R^*(s) = W_+^*(s) \text{diag} (\zeta(s)^{-\kappa_j})_{j=1}^m W_-^*(s).$$

Moreover W_-^* and its inverse are analytic on \mathbb{C}_+ and W_+^* and its inverse are analytic on \mathbb{C}_- . This shows that $-\kappa_1, \dots, -\kappa_m$ are the Wiener-Hopf indices of R^* . The positive Wiener-Hopf indices of R are the opposite to the negative Wiener-Hopf indices of R^* . So the Corollary is immediate from applying Theorem 2.2 to R^* . \square

6. Rational matrix functions and the Cayley transform

We denote the open unit disk in the complex plane by \mathbb{D} , the unit circle by \mathbb{T} , the closed unit disk by $\overline{\mathbb{D}}$, the open right hand half plane by \mathbb{C}_+ , the imaginary axis by $i\mathbb{R}$ and the open left hand half plane by \mathbb{C}_- .

Let the conformal transformation $\zeta : \mathbb{C} \rightarrow \mathbb{C}$ be given by

$$z = \zeta(s) = \frac{1-s}{s+1} \quad (s \in \mathbb{C} \setminus \{-1\}).$$

Then also

$$s = \zeta^{-1}(z) = \frac{1-z}{z+1} \quad (z \in \mathbb{C} \setminus \{-1\}).$$

If $s \in \mathbb{C}_+$ then the distance of s to -1 is larger than the distance to 1 , or in formula $|s+1| > |1-s|$ or $|\zeta(s)| < 1$. So $\zeta(\mathbb{C}_+) = \mathbb{D}$. For $s \in i\mathbb{R}$ the distance of s to -1 is equal to the distance to 1 , or in formula $|s+1| = |1-s|$ or $|\zeta(s)| = 1$. So $\zeta(i\mathbb{R}) = \mathbb{T}$. Similarly one sees $\zeta(\mathbb{C}_-) = \mathbb{C} \setminus \overline{\mathbb{D}}$. Using the similarity of the formula of ζ^{-1} with the formula for ζ we see that $\zeta(\mathbb{D}) = \mathbb{C}_+$, $\zeta(\mathbb{T}) = i\mathbb{R}$ and $\zeta(\mathbb{C} \setminus \overline{\mathbb{D}}) = \mathbb{C}_-$.

The following result is well-known, compare Section 3.6 in [6]. The result is presented here for sake of completeness.

Lemma 6.1. *Let $\{A_d$ on $\mathcal{X}, B_d, C_d, D_d\}$ be a discrete time realization of the function Θ , that is,*

$$\Theta(z) = D_d + zC_d(I - zA_d)^{-1}B_d.$$

Then

$$\Theta(\zeta(s)) = D_c + C_c(sI - A_c)^{-1}B_c. \quad (6.1)$$

Here

$$A_c = (A_d - I)(I + A_d)^{-1}, \quad D_c = D_d - C_d(I + A_d)^{-1}B_d, \quad (6.2)$$

$$B_c = \sqrt{2}(I + A_d)^{-1}B_d, \quad C_c = \sqrt{2}C_d(I + A_d)^{-1}. \quad (6.3)$$

Proof. First note that

$$\begin{aligned} \Theta(\zeta(s)) &= D_d + \frac{1-s}{s+1}C_d \left(I - \frac{1-s}{s+1}A_d \right)^{-1} B_d \\ &= D_d + (1-s)C_d((s+1)I - (1-s)A_d)^{-1}B_d \\ &= D_d + (1-s)C_d((I - A_d) + s(I + A_d))^{-1}B_d. \end{aligned}$$

By employing $A_c = (A_d - I)(I + A_d)^{-1}$, we have

$$\Theta(\zeta(s)) = D_d + (1-s)C_d(sI - A_c)^{-1}(I + A_d)^{-1}B_d.$$

Since $(1-s)(sI - A_c)^{-1} = (I - A_c)(sI - A_c)^{-1} - I$, we obtain

$$\Theta(\zeta(s)) = D_d - C_d(I + A_d)^{-1}B_d + C_d(I - A_c)(sI - A_c)^{-1}(I + A_d)^{-1}B_d.$$

Finally, by using $I - A_c = 2(I + A_d)^{-1}$, we see that

$$\Theta(\zeta(s)) = D_d - C_d(I + A_d)^{-1}B_d + C_d\sqrt{2}(I + A_d)^{-1}(sI - A_c)^{-1}\sqrt{2}(I + A_d)^{-1}B_d.$$

We conclude that (6.1) holds. \square

We can also express the operators A_d, B_d, C_d and D_d in A_c, B_c, C_c, D_c . Indeed we have the following result.

Corollary 6.2. *With A_d, B_d, C_d, D_d and A_c, B_c, C_c, D_c as in the Lemma 6.1 we have*

$$\begin{aligned} A_d &= (I - A_c)^{-1}(A_c + I), & B_d &= \sqrt{2}(I - A_c)^{-1}B_c, \\ C_d &= \sqrt{2}C_c(I - A_c)^{-1}, & D_d &= D_c + C_c(I - A_c)^{-1}B_c. \end{aligned}$$

In particular, if $G(s)$ admits a continuous time realization of the form

$$G(s) = D_c + C_c(sI - A_c)^{-1}B_c,$$

then the corresponding discrete time realization is given by

$$G(\zeta^{-1}(z)) = D_d + zC_d(I - zA_d)^{-1}B_d,$$

and vice-versa.

Proof. This follows from a simple rewriting of (6.2) and (6.3). \square

From [23, 24] it is known that inner functions on the unit circle have a stable unitary realization, and conversely. The Cayley transform converts a stable unitary realization into a stable dissipative realization as is shown in the following lemma.

Recall that a discrete time transfer function $\Theta(z)$ is bi-inner if and only if $\Theta(z)$ admits a stable discrete time realization

$$\begin{aligned} \Theta(z) &= D_d + zC_d(I - zA_d)^{-1}B_d & \text{where} \\ \begin{bmatrix} A_d & B_d \\ C_d & D_d \end{bmatrix} : \begin{bmatrix} \mathcal{X}_d \\ \mathcal{U} \end{bmatrix} &\rightarrow \begin{bmatrix} \mathcal{X}_d \\ \mathcal{U} \end{bmatrix} & \text{is unitary.} \end{aligned} \quad (6.4)$$

Finally, a discrete time system $\{A_d, B_d, C_d, D_d\}$ is *stable* if all the eigenvalues for A_d are contained in \mathbb{D} .

Lemma 6.3. *Let $\{A_d$ on $\mathcal{X}, B_d, C_d, D_d\}$ be a discrete time realization of the function $\Theta(z)$, that is,*

$$\Theta(z) = D_d + zC_d(I - zA_d)^{-1}B_d.$$

Let $\{A_c$ on $\mathcal{X}, B_c, C_c, D_c\}$ be the corresponding continuous time realization of $\Theta(\zeta(s))$, where

$$\begin{aligned} A_c &= (A_d - I)(I + A_d)^{-1}, & D_c &= D_d - C_d(I + A_d)^{-1}B_d, \\ B_c &= \sqrt{2}(I + A_d)^{-1}B_d, & C_c &= \sqrt{2}C_d(I + A_d)^{-1}. \end{aligned}$$

Then $\{A_d$ on $\mathcal{X}, B_d, C_d, D_d\}$ is a stable unitary realization of $\Theta(z)$ if and only if $\{A_c$ on $\mathcal{X}, B_c, C_c, D_c\}$ is a stable dissipative realization of $\Theta(\zeta(s))$.

Proof. The continuous time realization $\{A_c$ on $\mathcal{X}, B_c, C_c, D_c\}$ being stable and dissipative means that A_c has all its eigenvalues in the left hand half plane \mathbb{C}_- and $A_c + A_c^* + C_c^*C_c = 0$, D_c is unitary, and $B_c = -C_c^*D_c$.

To start with the stability we suppose that $A_c x = \lambda x$. Then

$$A_d x = (A_c - I)^{-1}(A_c + I)x = \frac{1 + \lambda}{\lambda - 1}x.$$

Note that $\frac{1+\lambda}{\lambda-1}$ is an eigenvalue of A_d and this means $\frac{1+\lambda}{\lambda-1} \in \mathbb{D}$. In other words $|\lambda + 1| < |\lambda - 1|$ and this is equivalent to $\lambda \in \mathbb{C}_-$. We conclude that A_d is (discrete time) stable if and only if A_c is (continuous time) stable.

Recall that (6.4) being unitary means that

$$\begin{bmatrix} I & 0 \\ 0 & I \end{bmatrix} = \begin{bmatrix} A_d^* & C_d^* \\ B_d^* & D_d^* \end{bmatrix} \begin{bmatrix} A_d & B_d \\ C_d & D_d \end{bmatrix} = \begin{bmatrix} A_d^* A_d + C_d^* C_d & A_d^* B_d + C_d^* D_d \\ B_d^* A_d + D_d^* C_d & B_d^* B_d + D_d^* D_d \end{bmatrix}. \quad (6.5)$$

Notice that

$$C_c^* C_c = 2(I + A_d^*)^{-1} C_d^* C_d (I + A_d)^{-1} \quad (6.6)$$

and

$$\begin{aligned} A_c + A_c^* &= (A_d - I)(I + A_d)^{-1} + (I + A_d^*)^{-1}(A_d^* - I) \\ &= (I + A_d^*)^{-1}((A_d^* - I)(I + A_d) + (I + A_d^*)(A_d - I))(I + A_d)^{-1} \\ &= (I + A_d^*)^{-1}(2A_d^* A_d - 2I)(I + A_d)^{-1}. \end{aligned} \quad (6.7)$$

Adding the two equalities (6.6) and (6.7) we see that

$$A_c + A_c^* + C_c^* C_c = 2(I + A_d^*)^{-1}(A_d^* A_d + C_d^* C_d - I)(I + A_d)^{-1}.$$

We conclude that $A_c + A_c^* + C_c^* C_c = 0$ if and only if $A_d^* A_d + C_d^* C_d - I = 0$.

The following calculation shows that $B_c = -C_c^* D_c$ if and only if $A_d^* B_d + C_d^* D_d = 0$:

$$\begin{aligned} A_d^* B_d + C_d^* D_d &= \sqrt{2} \left[(I - A_c^*)^{-1} (I + A_c^*) (I - A_c)^{-1} B_c + (I - A_c^*)^{-1} C_c^* D_c + \right. \\ &\quad \left. + (I - A_c^*)^{-1} C_c^* C_c (I - A_c)^{-1} B_c \right] \\ &= \sqrt{2} (I - A_c^*)^{-1} \left[(I + A_c^*) (I - A_c)^{-1} B_c + C_c^* C_c (I - A_c)^{-1} B_c + C_c^* D_c \right] \\ &= \sqrt{2} (I - A_c^*)^{-1} \left[(I + A_c^*) (I - A_c)^{-1} B_c - (A_c + A_c^*) (I - A_c)^{-1} B_c + C_c^* D_c \right] \\ &= \sqrt{2} (I - A_c^*)^{-1} (B_c + C_c^* D_c). \end{aligned}$$

Finally, we check that, given $A_c + A_c^* + C_c^* C_c = 0$ and $B_c = -C_c^* D_c$, we have that $D_d^* D_d + B_d^* B_d = I$ if and only if $D_c^* D_c = I$, i.e., D_c is unitary. To this end, observe that

$$\begin{aligned} D_d^* D_d + B_d^* B_d &= [D_c^* + B^*(I - A_c^*)^{-1} C_c^*] [D_c + C_c (I - A_c^*)^{-1} B_c] + \\ &\quad + 2B_c^* (I - A_c^*)^{-1} (I - A_c)^{-1} B_c \\ &= D_c^* D_c + D_c^* C_c (I - A_c)^{-1} B_c + B_c^* (I - A_c^*)^{-1} C_c^* D_c \\ &\quad + B_c^* (I - A_c^*)^{-1} C_c^* C_c (I - A_c)^{-1} B_c + 2B_c^* (I - A_c^*)^{-1} (I - A_c)^{-1} B_c \\ &= D_c^* D_c + B_c^* (I - A_c^*)^{-1} \left[-(I - A_c^*) - (I - A_c) - A_c - A_c^* + 2I \right] (I - A_c)^{-1} B_c \\ &= D_c^* D_c, \end{aligned}$$

In the third equality we used that $B_c = -C_c^* D_c$ and $A_c + A_c^* + C_c^* C_c = 0$. Since $D_d^* D_d + B_d^* B_d = D_c^* D_c$ we conclude that $D_d^* D_d + B_d^* B_d = I$ if and only if $D_c^* D_c = I$.

Therefore $\{A_d \text{ on } \mathcal{X}, B_d, C_d, D_d\}$ is a (discrete time) stable unitary realization of $\Theta(z)$ if and only if $\{A_c \text{ on } \mathcal{X}, B_c, C_c, D_c\}$ is a stable dissipative (continuous time) realization of $\Theta(\zeta(s))$. \square

7. Equivalence of discrete time and continuous time theorems

Consider the functions $V(z)$ and $W(z)$ in $H_{\mathbb{D}}^{\infty}(\mathcal{E}, \mathcal{E})$. Let $V(\zeta(s))$ and $W(\zeta(s))$ in $H^{\infty}(\mathcal{E}, \mathcal{E})$ be their corresponding Cayley transforms. Moreover, consider their corresponding discrete and continuous time realizations given by

$$V(z) = D_{dv} + zC_{dv}(I - zA_{dv})^{-1}B_{dv}, \quad (7.1)$$

$$W(z) = D_{dw} + zC_{dw}(I - zA_{dw})^{-1}B_{dw}, \quad (7.2)$$

$$V(\zeta(s)) = D_{cv} + C_{cv}(sI - A_{cv})^{-1}B_{cv}, \quad (7.3)$$

$$W(\zeta(s)) = D_{cw} + C_{cw}(sI - A_{cw})^{-1}B_{cw}. \quad (7.4)$$

The discrete realization in (7.1) and (7.2), and their corresponding continuous time counter parts (7.3) and (7.4) are related by the transformations presented in Lemma 6.1. Recall from Corollary 6.2

$$\begin{aligned} A_{dv} &= (I - A_{cv})^{-1}(A_{cv} + I), & B_{dv} &= \sqrt{2}(I - A_{cv})^{-1}B_{cv}, \\ C_{dv} &= \sqrt{2}C_{cv}(I - A_{cv})^{-1}, & D_{dv} &= D_{cv} + C_{cv}(I - A_{cv})^{-1}B_{cv}, \end{aligned} \quad (7.5)$$

and

$$\begin{aligned} A_{dw} &= (I - A_{cw})^{-1}(A_{cw} + I), & B_{dw} &= \sqrt{2}(I - A_{cw})^{-1}B_{cw}, \\ C_{dw} &= \sqrt{2}C_{cw}(I - A_{cw})^{-1}, & D_{dw} &= D_{cw} + C_{cw}(I - A_{cw})^{-1}B_{cw}. \end{aligned} \quad (7.6)$$

Next we would like to develop a relationship between a special Stein equation in the discrete time and a corresponding Lyapunov equation in continuous time. To this end, consider the Stein equation:

$$\Omega = A_{dv}\Omega A_{dv}^* + B_{dv}B_{dv}^*. \quad (7.7)$$

By employing the corresponding transformations in (7.5) and (7.6), we obtain

$$\begin{aligned} \Omega &= (I - A_{cv})^{-1}(A_{cv} + I)\Omega(A_{cv}^* + I)(I - A_{cv}^*)^{-1} + \\ &\quad + (I - A_{cv})^{-1}B_{cv}2B_{cv}^*(I - A_{cv}^*)^{-1}. \end{aligned}$$

Multiplying by $I - A_{cv}$ on the left and by $I - A_{cv}^*$ on the right, we arrive at

$$(I - A_{cv})\Omega(I - A_{cv}^*) = (A_{cv} + I)\Omega(A_{cv}^* + I) + 2B_{cv}B_{cv}^*.$$

This simplifies to

$$A_{cv}\Omega + \Omega A_{cv}^* + B_{cv}B_{cv}^* = 0. \quad (7.8)$$

Therefore Ω is a solution of the Stein equation (7.7) if and only if Ω is a solution of the Lyapunov equation (7.8).

Next we establish the relation between $C_{c\circ}$ and $C_{d\circ}$ where

$$C_{d\circ} = D_{dv}B_{dv}^* + C_{dv}\Omega A_{dv}^*$$

and

$$C_{c\circ} = D_{cv}B_{cw}^* + C_{cv}\Omega.$$

We have

$$\begin{aligned} C_{d\circ} &= D_{dv}B_{dw}^* + C_{dv}\Omega A_{dw}^* \\ &= \sqrt{2} \left[[D_{cv} + C_{cv}(I - A_{cv})^{-1}B_{cv}]B_{cw}^* + \right. \\ &\quad \left. + C_{cv}(I - A_{cv})^{-1}\Omega(I + A_{cw}^*) \right] (I - A_{cw}^*)^{-1}. \end{aligned}$$

Now use (7.8) in

$$\begin{aligned} &C_{cv}(I - A_{cv})^{-1} [B_{cv}B_{cw}^* + \Omega(I + A_{cw}^*)] \\ &= C_{cv}(I - A_{cv})^{-1} [(I - A_{cv})\Omega] \\ &= C_{cv}\Omega. \end{aligned}$$

So we have

$$C_{d\circ} = \sqrt{2} \left[D_{cv}B_{cw}^*(I - A_{cw}^*)^{-1} + C_{cv}\Omega(I - A_{cw}^*)^{-1} \right].$$

We conclude that

$$C_{d\circ} = \sqrt{2}C_{c\circ}(I - A_{cw}^*)^{-1}. \quad (7.9)$$

Our aim is to develop a connection between [13, Theorem 2.2] in the discrete time setting and our Theorem 2.2. To this end, consider the equations

$$Q_d = A_{dw}Q_dA_{dw}^* + C_{d\circ}^*C_{d\circ} \quad (7.10)$$

and

$$A_{cw}Q_c + Q_cA_{cw}^* + C_{c\circ}^*C_{c\circ} = 0. \quad (7.11)$$

Because A_{dw} is discrete time stable and A_{cw} is continuous time stable, the solution to both of these equations is unique. We claim that $Q_d = Q_c$. Consider

$$\begin{aligned} Q &= A_{dw}QA_{dw}^* + C_{d\circ}^*C_{d\circ} \\ &= (I - A_{cw})^{-1}(A_{cw} + I)Q(A_{cw}^* + I)(I - A_{cw}^*)^{-1} \\ &\quad + 2(I - A_{cw})^{-1}C_{c\circ}^*C_{c\circ}(I - A_{cw}^*)^{-1}. \end{aligned}$$

Multiply on the left with $I - A_{cw}$ and on the right with $I - A_{cw}^*$, we have

$$(I - A_{cw})Q(I - A_{cw}^*) = (A_{cw} + I)Q(A_{cw}^* + I) + 2C_{c\circ}^*C_{c\circ}.$$

This simplifies to

$$A_{cw}Q + QA_{cw}^* + C_{c\circ}^*C_{c\circ} = 0.$$

We conclude that the equations (7.11) and (7.10) have the same solutions.

We will show the equivalence of the next two Theorems. The first is a rephrase of [13, Theorem 2.2] and the second rephrases Theorem 2.2 above. We denote with $H_{\mathbb{D}}^{\infty}(\mathcal{E}, \mathcal{E})$ the space consisting of the set of all operator valued functions $\Theta(s)$ on \mathcal{E} that are analytic in the open unit circle \mathbb{D} and such that

$$\|\Theta\|_{\infty} = \sup\{\|\Theta(z)\| : z \in \mathbb{D}\} < \infty.$$

In the remainder of the paper the boldface \mathbf{T}_R denotes the Toeplitz operator on $\ell_+^2(\mathcal{E})$ with symbol R in $L_\infty^\mathbb{T}(\mathcal{E}, \mathcal{E})$.

Theorem 7.1. *Assume that $R(e^{i\omega}) = V(e^{i\omega})W(e^{i\omega})^*$ where V and W are two bi-inner rational functions in $H_\mathbb{D}^\infty(\mathcal{E}, \mathcal{E})$. Let \mathbf{T}_R be the corresponding Toeplitz operator on $\ell_+^2(\mathcal{E})$. Let $\{A_{dv}, B_{dv}, C_{dv}, D_{dv}\}$ and $\{A_{dw}, B_{dw}, C_{dw}, D_{dw}\}$ be stable unitary realizations of $V(z)$ and $W(z)$, respectively. Let Ω_d be the unique solution of the Stein equation*

$$\Omega_d = A_{dv}\Omega_d A_{dw}^* + B_{dv}B_{dw}^*. \quad (7.12)$$

Let $C_{d\circ}$ be the operator mapping \mathcal{X}_w into \mathcal{E} defined by

$$C_{d\circ} = D_{dv}B_{dw}^* + C_{dv}\Omega_d A_{dw}^*. \quad (7.13)$$

Finally, let Q_d be the unique solution to the Stein equation

$$Q_d = A_{dw}Q_d A_{dw}^* + C_{d\circ}^* C_{d\circ}. \quad (7.14)$$

Then the following holds:

1. The operator Q_d is a positive contraction.
2. The multiplicity of 1 as an eigenvalue of Q_d equals $\mathbf{n}(\mathbf{T}_R)$. In other words, $\mathbf{n}(T_R) = \mathbf{n}(I - Q)$. Moreover, for $k = 0, 1, 2, \dots, p$, we have

$$\mathbf{n}(\mathbf{T}_{z^k R}) = \dim(\text{Ker}(I - A_{dw}^k Q A_{dw}^{*k})). \quad (7.15)$$

3. For $k = 1, 2, \dots$, consider the integers

$$\mu_k = \mathbf{n}(I - A_{dw}^{k-1} Q (A_{dw}^*)^{k-1}) - \mathbf{n}(I - A_{dw}^k Q A_{dw}^{*k}). \quad (7.16)$$

Then the negative Wiener-Hopf indices $-\kappa_1, \dots, -\kappa_p$ of the Toeplitz operator \mathbf{T}_R are given by

$$\kappa_j = \#\{k : \mu_k \geq j\}, \quad (j = 1, \dots, p = \mu_1). \quad (7.17)$$

Recall that the transformation $\zeta : \mathbb{C} \rightarrow \mathbb{C}$ is given by

$$\zeta(s) = \frac{1-s}{s+1} \quad (s \in \mathbb{C} \setminus \{-1\}).$$

Theorem 7.2. *Assume that $R(\zeta(i\omega)) = V(\zeta(i\omega))W(\zeta(i\omega))^*$ ($\omega \in \mathbb{R}$) where $V(\zeta(s))$ and $W(\zeta(s))$ are two bi-inner rational functions in $H^\infty(\mathcal{E}, \mathcal{E})$. Let*

$$\{A_{cv}, B_{cv}, C_{cv}, D_{cv}\} \quad \text{and} \quad \{A_{cw}, B_{cw}, C_{cw}, D_{cw}\}$$

be two stable dissipative realizations of $V(\zeta(s))$ and $W(\zeta(s))$, respectively. Let Ω_c be the unique solution of the Lyapunov equation

$$A_{cv}\Omega_c + \Omega_c A_{cw}^* + B_{cv}B_{cw}^* = 0. \quad (7.18)$$

Let $C_{c\circ}$ be the operator mapping \mathcal{X}_w into \mathcal{E} defined by

$$C_{c\circ} = D_{cv}B_{cw}^* + C_{cv}\Omega_c. \quad (7.19)$$

Finally, let Q_c be the unique solution to the Lyapunov equation

$$A_{cw}Q_c + Q_c A_{cw}^* + C_{c\circ}^* C_{c\circ} = 0. \quad (7.20)$$

Then the following holds:

1. The operator Q_c is a positive contraction.

2. The multiplicity of 1 as an eigenvalue of Q_c equals $\mathbf{n}(T_{R\zeta})$. In other words, $\mathbf{n}(T_{R\zeta}) = \mathbf{n}(I - Q_c)$. Moreover, for $\zeta(s) = \frac{1-s}{1+s}$, we have

$$\mathbf{n}(T_{\zeta^k(R\zeta)}) = \dim(\text{Ker}(I - \zeta(-A_{cw})^k Q_c \zeta(-A_{cw})^{*k})). \quad (7.21)$$

3. For $k = 1, 2, \dots$, consider the integers

$$\begin{aligned} \mu_k = & \mathbf{n}(I - \zeta(-A_{cw})^{k-1} Q_c (\zeta(-A_{cw})^*)^{k-1}) + \\ & - \mathbf{n}(I - \zeta(-A_{cw})^k Q_c \zeta(-A_{cw})^{*k}). \end{aligned} \quad (7.22)$$

Then the negative Wiener-Hopf indices $-\kappa_1, \dots, -\kappa_p$ of $T_{R\zeta}$ are given by

$$\kappa_j = \#\{k : \mu_k \geq j\}, \quad (j = 1, \dots, p = \mu_1). \quad (7.23)$$

Assume we have Theorem 7.1. Since $Q_c = Q_d$, Part 1 of Theorem 7.2 is proven. Notice that the Wiener-Hopf factorization of R with respect to \mathbb{T} immediately generates a Wiener-Hopf factorization of $R(\zeta(i\omega))$ with respect to $i\mathbb{R}$ with the same Wiener-Hopf indices. Indeed, if $R = W_+ DW_-$ then, after substitution of z by $\zeta(s)$, we have $R\zeta = (W_+ \zeta)(D\zeta)(W_- \zeta)$. So the dimensions of the null spaces of $\mathbf{T}_{z^k R}$ and of $T_{\zeta^k R\zeta}$ coincide. Together with $A_{dw} = \zeta(-A_{cw})$ the items 2 and 3 of Theorem 7.1 give the items 2 and 3 of Theorem 7.2.

Obviously, also Theorem 7.2 implies Theorem 7.1.

7.1. An example

As an illustration of Theorem 7.2 we present the following example which is a continuous version of the discrete example on page 706 in [20]. To this end, let

$$R_c(s) = \begin{bmatrix} \zeta(s)^{-4} & 0 & 0 & 0 & 0 \\ 0 & \zeta(s)^{-2} & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & \zeta(s)^3 & 0 \\ 0 & 0 & 0 & 0 & \zeta(s)^5 \end{bmatrix}.$$

Recall that $\zeta(s) = \frac{1-s}{1+s}$. Then $R_c(s)$ factors as $R_c(i\omega) = V_c(i\omega)W_c(i\omega)^*$, where

$$V_c(s) = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & \zeta(s)^3 & 0 \\ 0 & 0 & 0 & 0 & \zeta(s)^5 \end{bmatrix} \quad W_c(s) = \begin{bmatrix} \zeta(s)^4 & 0 & 0 & 0 & 0 \\ 0 & \zeta(s)^2 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

Let us first construct a stable dissipative realization $\{A_n \text{ on } \mathbb{C}^n, B_n, C_n, D_n\}$ for $\zeta(s)^n$. Notice that $p(z) = ze_1^T (I - zJ_n(0))^{-1} e_n = z^n$, where e_1 and e_n are the first and last unit vectors in \mathbb{C}^n and $J_n(0)$ is the upper triangular Jordan matrix with eigenvalue 0, or, in other words, the upward shift. Then

Lemma 6.1 provides the realization of $\zeta(s)^n = p(\zeta(s))$. To see this observe that

$$(I_n + J_n(0))^{-1} = \sum_{j=0}^{n-1} (-J_n(0))^j.$$

This with Lemma 6.1 readily implies that

$$\begin{aligned} A_n &= (J_n(0) - I_n)(I_n + J_n(0))^{-1} = -I_n + 2 \sum_{j=1}^{n-1} (-1)^{j+1} J_n(0)^j, \\ B_n &= \sqrt{2} (I_n + J_n(0))^{-1} e_n = \sqrt{2} [(-1)^{n-1} \quad (-1)^{n-2} \quad \dots \quad -1 \quad 1]^T, \\ C_n &= \sqrt{2} e_1^T (I_n + J_n(0))^{-1} = \sqrt{2} [1 \quad -1 \quad \dots \quad (-1)^{n-1}], \\ D_n &= 0 - e_1^T (I_n + J_n(0))^{-1} e_n = (-1)^n. \end{aligned} \tag{7.24}$$

Notice that A_n is an upper triangular Toeplitz matrix.

According to Lemma 6.3 and using that the realization of $p(z)$ is unitary, we have that $\{A_n, B_n, C_n, D_n\}$ is indeed a stable dissipative realization of $\zeta(s)^n$. Nevertheless let us verify that directly. Recall that the realization $\{A_n \text{ on } \mathcal{X}, B_n, C_n, D_n\}$ being stable and dissipative means that A_n has all its eigenvalues in the open left hand half plane and $A_n + A_n^* + C_n^* C_n = 0$, D_n is unitary, and $B_n = -C_n^* D_n$. First observe that -1 is the only eigenvalue of A_n , and thus, A_n is continuous time stable. Next check by direct calculation that $A_n^* + A_n + C_n^* C_n = 0$ and $A_n^* + A_n + B_n B_n^* = 0$. Since $D_n = (-1)^n$ is unitary and $B_n = -C_n^* D_n$ we are done.

Motivated by the previous realization, the factors $V_c(s)$ and $W_c(s)$ can be given by the following stable unitary realizations:

$$V_c(s) = D_v + C_v(sI - A_v)^{-1} B_v.$$

where

$$\begin{aligned} A_v &= \begin{bmatrix} A_3 & 0 \\ 0 & A_5 \end{bmatrix} : \begin{bmatrix} \mathbb{C}^3 \\ \mathbb{C}^5 \end{bmatrix} \rightarrow \begin{bmatrix} \mathbb{C}^3 \\ \mathbb{C}^5 \end{bmatrix}, \\ B_v &= \begin{bmatrix} 0_{3 \times 3} & B_3 & 0 \\ 0_{5 \times 3} & 0 & B_5 \end{bmatrix} : \begin{bmatrix} \mathbb{C}^3 \\ \mathbb{C} \\ \mathbb{C} \end{bmatrix} \rightarrow \begin{bmatrix} \mathbb{C}^3 \\ \mathbb{C}^5 \end{bmatrix}, \\ C_v &= \begin{bmatrix} 0_{3 \times 3} & 0_{3 \times 5} \\ C_3 & 0 \\ 0 & C_5 \end{bmatrix} : \begin{bmatrix} \mathbb{C}^3 \\ \mathbb{C}^5 \end{bmatrix} \rightarrow \begin{bmatrix} \mathbb{C}^3 \\ \mathbb{C} \\ \mathbb{C} \end{bmatrix} = \mathbb{C}^5, \\ D_v &= \begin{bmatrix} I_3 & 0 & 0 \\ 0 & D_3 & 0 \\ 0 & 0 & D_5 \end{bmatrix} : \begin{bmatrix} \mathbb{C}^3 \\ \mathbb{C} \\ \mathbb{C} \end{bmatrix} \rightarrow \begin{bmatrix} \mathbb{C}^3 \\ \mathbb{C} \\ \mathbb{C} \end{bmatrix}. \end{aligned}$$

In this case,

$$W_c(s) = D_w + C_w(sI - A_w)^{-1} B_w,$$

where

$$\begin{aligned}
A_w &= \begin{bmatrix} A_4 & 0 \\ 0 & A_2 \end{bmatrix} : \begin{bmatrix} \mathbb{C}^4 \\ \mathbb{C}^2 \end{bmatrix} \rightarrow \begin{bmatrix} \mathbb{C}^4 \\ \mathbb{C}^2 \end{bmatrix}, \\
B_w &= \begin{bmatrix} B_4 & 0 & 0_{4 \times 3} \\ 0 & B_2 & 0_{2 \times 3} \end{bmatrix} : \begin{bmatrix} \mathbb{C} \\ \mathbb{C} \\ \mathbb{C}^3 \end{bmatrix} \rightarrow \begin{bmatrix} \mathbb{C}^4 \\ \mathbb{C}^2 \end{bmatrix}, \\
C_w &= \begin{bmatrix} C_4 & 0 \\ 0 & C_2 \\ 0_{3 \times 4} & 0_{3 \times 2} \end{bmatrix} : \begin{bmatrix} \mathbb{C}^4 \\ \mathbb{C}^2 \end{bmatrix} \rightarrow \begin{bmatrix} \mathbb{C} \\ \mathbb{C} \\ \mathbb{C}^3 \end{bmatrix} = \mathbb{C}^5, \\
D_w &= \begin{bmatrix} D_4 & 0 & 0 \\ 0 & D_2 & 0 \\ 0 & 0 & I_3 \end{bmatrix} : \begin{bmatrix} \mathbb{C} \\ \mathbb{C} \\ \mathbb{C}^3 \end{bmatrix} \rightarrow \begin{bmatrix} \mathbb{C} \\ \mathbb{C} \\ \mathbb{C}^3 \end{bmatrix}.
\end{aligned}$$

It is noted that $B_v B_w^* = 0$. Therefore the unique solution to the Lyapunov equation $A_v \Omega + \Omega A_w^* + B_v B_w^* = 0$ is $\Omega = 0$. Hence

$$C_\circ = D_v B_w^* + C_v \Omega = D_v B_w^* = \sqrt{2} \begin{bmatrix} -1 & 1 & -1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

The unique solution to the Lyapunov equation $A_w Q + Q A_w^* + C_\circ^* C_\circ = 0$ is given by $Q = I$. Now observe that

$$\zeta(-A_w) = \begin{bmatrix} 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

It is noted that $\zeta(-A_w) = S_4 \oplus S_2$ on $\mathbb{C}^4 \oplus \mathbb{C}^2$ where S_k on \mathbb{C}^k is the upward shift, that is, all the entries of S_k immediately above the main diagonal are 1 and all the other entries are zero, that is $S_k = J_k(0)$. (See also Corollary 6.2.) Because $Q = I$, we readily see that

$$\begin{aligned}
\dim(\text{Ker}(I - Q)) &= 6 \\
\dim(\text{Ker}(I - \zeta(-A_w)Q\zeta(-A_w^*))) &= 4 \\
\dim(\text{Ker}(I - \zeta(-A_w)^2 Q \zeta(-A_w^*)^2)) &= 2 \\
\dim(\text{Ker}(I - \zeta(-A_w)^3 Q \zeta(-A_w^*)^3)) &= 1 \\
\dim(\text{Ker}(I - \zeta(-A_w)^k Q \zeta(-A_w^*)^k)) &= 0 \quad \text{for } k \geq 4.
\end{aligned}$$

In other words,

$$\mu_1 = 6 - 4 = 2, \quad \mu_2 = 4 - 2 = 2, \quad \mu_3 = 2 - 1 = 1 \quad \text{and} \quad \mu_4 = 1 - 0 = 0.$$

Using this we have

$$\kappa_2 = \#\{k : \mu_k \geq 1\} = 4 \quad \text{and} \quad \kappa_1 = \#\{k : \mu_k \geq 2\} = 2.$$

Therefore the negative Wiener-Hopf indices for R are $\{-4, -2\}$.

Acknowledgements

The work of the second author is supported in part by the National Research Foundation of South Africa (NRF), Grant Number 145688.

References

- [1] V.M. Adamjan, D.Z. Arov, M.G. Krein, Infinite Hankel block matrices and related extension problems, *Izv. Akad. Nauk SSSR. Ser. Math.* **6** (1971), pp. 87-112 (Russian); English Transl., *Amer. Math. Soc. Trans I.* **111** (1978), pp. 133-156.
- [2] D. Alpay, I. Gohberg, Unitary rational matrix functions. In *Topics in Interpolation Theory of Rational Matrix-valued Functions*. Oper. Theory Adv. Appl. 33, Birkhäuser Verlag, Basel, 1988, 175-222.
- [3] H. Bart, I. Gohberg, M.A. Kaashoek, The coupling method for solving integral equations. In: *Topics in operator theory, systems and networks*. Oper. Theory Adv. Appl. 12, Birkhäuser Verlag, Basel, 1984, 39-73.
- [4] H. Bart, I. Gohberg, M.A. Kaashoek, Explicit Wiener-Hopf factorization and realization. In: *Constructive methods of Wiener-Hopf factorization*. Oper. Theory Adv. Appl. 21, Birkhäuser Verlag, Basel, 1986, 235-316.
- [5] H. Bart, I. Gohberg, M.A. Kaashoek, Invariants for Wiener-Hopf equivalence of analytic operator functions. In: *Constructive methods of Wiener-Hopf factorization*. Oper. Theory Adv. Appl. 21, Birkhäuser Verlag, Basel, 1986, 317-355.
- [6] H. Bart, I. Gohberg, M.A. Kaashoek, A.C.M. Ran, *Factorization of matrix and operator functions: the state space method*. Oper. Theory Adv. Appl. 178, Birkhäuser Verlag, Basel, 2008.
- [7] H. Bart, I. Gohberg, M.A. Kaashoek, A.C.M. Ran, *A state space approach to canonical factorization with applications*. Oper. Theory Adv. Appl. 200, Birkhäuser, Basel, 2010.
- [8] A. Böttcher, B. Silbermann, *Analysis of Toeplitz Operators*. Second ed., Springer Verlag, Berlin, Heidelberg, 2006.
- [9] K. Clancey, I. Gohberg, *Factorization of matrix functions and singular integral operators*. Oper. Theory Adv. Appl. 3, Birkhäuser Verlag, Basel, 1981.
- [10] C. Foias and A. E. Frazho, *The Commutant Lifting Approach to Interpolation Problems*, Oper. Theory Adv. Appl., **44**, Birkhäuser-Verlag, Basel, 1990.
- [11] C. Foias, A.E. Frazho, I. Gohberg, and M.A. Kaashoek, *Metric constrained interpolation, commutant lifting and systems*, Oper. Theory Adv. Appl. 100, Birkhäuser Verlag, Basel, 1998.
- [12] A.E. Frazho, M.A. Kaashoek, Canonical factorization of rational matrix functions. A note on a paper by P. Dewilde. *Indag. Math.* 23 (2012), 1154-1164.

- [13] A.E. Frazho, M.A. Kaashoek, A.C.M. Ran, F. van Schagen, Wiener-Hopf indices of unimodular functions on the unit circle, revisited. *Linear Algebra Appl.* to appear.
- [14] I.C. Gohberg, I. A. Feldman, *Convolution equations and projection methods for their solution.* Transl. Math. Monogr., Vol. 41, Amer. Math. Soc., Providence, Rhode Island, 1974.
- [15] I. Gohberg, S. Goldberg, M.A. Kaashoek, *Classes of linear operators*, Vol. I. Oper. Theory Adv. Appl. 49, Birkhäuser Verlag, Basel, 1990.
- [16] I. Gohberg, S. Goldberg, M.A. Kaashoek, *Classes of linear operators*, Vol. II. Oper. Theory Adv. Appl. 63, Birkhäuser Verlag, Basel, 1993.
- [17] I. Gohberg, M.A. Kaashoek, F. van Schagen, *Partially Specified Matrices and Operators: Classification, Completion, Applications*, Oper. Theory Adv. Appl. 79, Birkhäuser Verlag, Basel, 1995, 340 pp.
- [18] I. Gohberg, M.A. Kaashoek, I.M. Spitkovsky, An overview of matrix factorization theory and operator applications. In: *Factorization and integrable systems.* Oper. Theory Adv. Appl. 141, Birkhäuser Verlag, Basel, 2003, 1-102.
- [19] I. Gohberg, L. Lerer, L. Rodman, Factorization indices for matrix polynomials. *Bull. Amer. Math. Soc.* 84 (1978), 275-277.
- [20] G. J. Groenewald, M. A. Kaashoek, A. C. M. Ran, Wiener-Hopf indices of unitary functions on the unit circle in terms of realizations and related results on Toeplitz operators. *Indag. Math.* 28, (2017), 649-710.
- [21] G. J. Groenewald, M. A. Kaashoek, A. C. M. Ran, Wiener-Hopf factorization indices of rational matrix functions with respect to the unit circle in terms of realization. *Indag. Math.* 34 (2023), 338-356.
- [22] V. Pták and N. J. Young, A generalization of the zero location theorem of Schur and Cohn, *IEEE trans. Automatic Control*, 25 (1980), 978-980.
- [23] B. Sz.-Nagy and C. Foias *Harmonic Analysis of Operators on Hilbert Space*, North Holland Publ. Co., Amsterdam-Budapest, 1970.
- [24] B. Sz.-Nagy, C. Foias, H. Bercovici and L. Kérchy *Harmonic Analysis of Operators on Hilbert Space: Second edition*, Springer, New York 2009.

A.E. Frazho

Department of Aeronautics and Astronautics, Purdue University, West Lafayette, IN 47907, USA.

e-mail: frazho@purdue.edu

A.C.M. Ran

Department of Mathematics, Faculty of Science, VU Amsterdam, De Boelelaan 1111, 1081 HV Amsterdam, The Netherlands and Research Focus: Pure and Applied Analytics, North-West University, Potchefstroom, South Africa

e-mail: a.c.m.ran@vu.nl

F. van Schagen

Department of Mathematics, Faculty of Science, VU Amsterdam, De Boelelaan 1111, 1081 HV Amsterdam

e-mail: f.van.schagen@vu.nl

(FvS) Department of Mathematics, Faculty of Science, VU Amsterdam, De Boelelaan 1111, 1081 HV Amsterdam