

Test particles in Kaluza-Klein models

João Baptista

June 2024

Abstract

Geodesics in general relativity describe the behaviour of test particles in a gravitational field. In 5D Kaluza-Klein, geodesics reproduce the Lorentz force motion of particles in an electromagnetic field. This paper studies geodesic motion on $M_4 \times K$ in the presence of general gauge fields and Higgs-like scalars. It shows that the classical mass and charge of a test particle become variable quantities when the geodesic traverses regions of spacetime with massive gauge fields, such as the weak force field, or with non-constant Higgs scalars. This agrees with the physical fact that interactions mediated by massive bosons can change the mass and charge of particles. The variation rates of mass and charge along a geodesic are given by natural geometric formulae. In regions where mass is preserved there are additional constants of motion, one for every abelian or simple summand in the Killing algebra of K . The last part of the paper discusses traditional difficulties of Kaluza-Klein models, such as the low q/m ratios in the 5D model. It suggests possible ways to circumvent them. It also advocates the naturalness of a model where elementary particles always travel at the speed of light in higher dimensions.

Keywords: Kaluza-Klein theories; geodesic motion; Riemannian submersions; test particles; momentum; rest mass; charge.

Contents

1	Introduction and overview	2
2	Motion in the vacuum and 4D inertial frames	10
3	Curves in Riemannian submersions	11
4	Geodesics among massless gauge fields	13
5	Geodesics in general Riemannian submersions	17
6	Higher-dimensional momentum and mass	21
7	Rest mass variation	23
8	Charges and charge variation	24
9	Circumventing a traditional difficulty	27
10	A unique speed in higher dimensions	31
11	Parameterizing the space of null geodesics	33
A	Auxiliary results	37
B	Parallel transport in Riemannian submersions	39
	References	40

1 Introduction and overview

This paper studies geodesics in the Kaluza-Klein framework. It works in a general setting where the higher-dimensional metric encodes both massless and massive 4D fields. The starting point is the notion that, in these conditions, a particle's classical mass and electromagnetic charge should not be strict constants of geodesic motion. In regions of spacetime with massive gauge fields or non-constant Higgs fields, physical particles can interact with massive bosons, so their own charge and mass can change. Thus, it would be natural to have definitions of mass and charge of a test particle that enabled such variations along a geodesic. This would also illustrate how a higher-dimensional, classical theory can exhibit qualitative features (particle charge and mass change) that are usually reserved for QFT calculations.

Thinking along those lines, the author was lead to consider how 4D proper time, denoted τ , evolves along an affine geodesic $\gamma(s)$ on the higher-dimensional spacetime $P = M_4 \times K$. In this setting, the covariantly conserved momentum vector is

$$p(s) = \sigma \frac{d\gamma}{ds} = \sigma \frac{d\tau}{ds} \frac{d\gamma}{d\tau} = m(s) \frac{d\gamma}{d\tau}, \quad (1.1)$$

where σ is a constant with dimension MT^{-1} and $m(s) = \sigma \frac{d\tau}{ds}$ is the particle's rest mass. So a non-constant rate of change $\frac{d\tau}{ds}$ is equivalent to a variable mass. It turns out that 4D proper time and the geodesic parameter s need not be proportional to each other in regions where the internal geometry is not covariantly constant. So $m(s)$ can vary in those regions, as desired.

It also turns out that the phenomenon of geodesic mass variation had been previously noticed in the Kaluza-Klein literature. Specifically, in studies of the 5D model, with $K = \text{U}(1)$ and mass variation coming from a non-constant size of the internal circle [1–5]. In that electromagnetic setting, however, it was mainly regarded as a difficulty of Kaluza-Klein theory. Rest mass variation is not observed in real electromagnetic interactions, so it was not easy to offer a physical understanding of the phenomenon. Arguments were proposed to avoid it or find alternative interpretations [6, 7].

A wider perspective seems to be useful in this instance. For higher-dimensional K it becomes possible to distinguish the separate effects on geodesics of electromagnetic-like gauge fields, non-abelian massless gauge fields, massive gauge fields and Higgs-like scalars. And when one recognizes that the only fields able to change the mass of a test particle are the massive gauge fields (such as the weak force field) and the Higgs-like scalars, a natural physical interpretation emerges. The phenomenon of geodesic mass variation seems to describe the physical fact that interactions mediated by massive bosons are able to change the mass of particles.

In the geodesic model all internal motion corresponds to mass in four dimensions, but electromagnetic charge is related to the component of internal momentum along a specific Killing direction inside K . So in this model there can be mass without charge, but not the other way around, which fits observation. The independent existence of mass and charge can occur only when the dimension of K is greater than one. Otherwise there is no meaningful distinction between the internal momentum and its component along the Killing field.

The higher-dimensional definition of charge of a test particle has three natural properties: *i)* charge is a constant of geodesic motion in regions of spacetime that only have massless gauge fields and constant Higgs scalars; *ii)* in those regions, geodesics project down to Lorentz force motions in four dimensions, as usual in 5D Kaluza-Klein; *iii)* in regions of spacetime with massive gauge fields, the test particle's charge can vary along a geodesic if and only if the bosons associated to the fields are charged. Thus, higher-dimensional geodesics give a fairly good account of the properties of charged particles, at least qualitatively. They model the physical fact that interactions mediated by charged gauge bosons, such as the Standard Model's W bosons, can modify the charge of particles, while interactions mediated by massless or by massive, neutral gauge bosons, cannot. This property of higher-dimensional geodesics apparently has not been remarked before.

Another relevant property of the geodesic model is that particles travelling at the speed of light on M_4 cannot interact directly with gauge fields or Higgs-like scalars, at least classically. More precisely, the projection to M_4 of the particle's motion on P is independent of the values of such fields. This property is incompatible with the motion of the old massless neutrinos, since they were thought to interact with the weak field and travel at the speed of light on Minkowski space. So the geodesic model disfavors the existence of massless neutrinos or similar particles. This also does not seem to have been remarked before. Of course, one should still bear in mind that geodesic motion is an approximation that disregards back-reaction and the fermionic nature of particles.

Our study of geodesic motion on $P = M_4 \times K$, for arbitrary K , relies on classical geometrical results about Riemannian submersions developed in [8–11] and presented in [12, 13], for example. Extended reviews of the Kaluza-Klein framework can be found in [14–21]. Some of the early original references are [22–31]. Geodesic motion has been studied since the very beginning of Kaluza-Klein literature. The main focus has been on the 5D case, where calculations are more explicit and differential geometric techniques less necessary (e.g. [2, 22, 28, 32–35]). This paper follows the notation in [36] and its treatment of massive gauge fields and Higgs-like fields. A shorter account of the results on rest mass variation is given in the companion paper [37]. More comments about the literature will be added below, as we give a brief overview of the main results and observations.

Rest mass variation

Consider a submersive metric g_P on the higher-dimensional spacetime $P = M_4 \times K$. As described in section 3, the assumptions on g_P that generalize the traditional Kaluza ansatz are: *i*) it projects down to a unique Lorentzian metric g_M on M_4 ; *ii*) the restriction of g_P to the slices $\{x\} \times K$ defines a family of Riemannian internal metrics $g_K(x)$ parametrized by the points in M_4 ; *iii*) the non-diagonal components of g_P encode gauge fields A^a on M_4 with values in the vector fields on K , so the gauge group is $\text{Diff}(K)$ or a subgroup. Notice that the gauge group need not act on K only through isometries of g_K .

Now let $\gamma(s)$ be a time-like or null geodesic on P representing the motion of a test particle. It is a curve satisfying $\nabla_{\dot{\gamma}}\dot{\gamma} = 0$. Denote by $\gamma_M(s)$ the projection of this curve to Minkowski space. The main result of section 7 says that the particle's rest mass, as defined in (1.1), changes according to

$$c^2 \frac{d}{ds} m^2(s) = - (d^A g_K)_{\dot{\gamma}_M} (p^\mathcal{Y}, p^\mathcal{Y}) . \quad (1.2)$$

Here $p^\mathcal{Y}$ denotes the vertical (internal) component of the particle's momentum vector, while $(d^A g_K)_{\dot{\gamma}_M}$ is a covariant derivative of the internal metric along the vector $\dot{\gamma}_M$ tangent to M_4 . So the rest mass of the particle is a constant of geodesic motion in regions where g_K is covariantly constant, but may change elsewhere.

The derivative $d^A g_K$ measures how g_K changes along M_4 up to diffeomorphisms of K . It is equivariant under $\text{Diff}(K)$ -gauge transformations. Geometrically, it can be identified with the second fundamental form of the fibres of P . As in [36], it can be expressed in terms of Lie derivatives and the gauge one-forms A^a as

$$(d^A g_K)_X(U, V) = (\mathcal{L}_X g_K)(U, V) + A^a(X) (\mathcal{L}_{e_a} g_K)(U, V) . \quad (1.3)$$

Here X is any vector in TM ; U and V are vertical vectors in TP ; and $\mathcal{L}_{e_a} g_K$ denotes the Lie derivative along the internal vector field e_a . Thus, the covariant derivative vanishes when the internal metric is constant along M_4 and, simultaneously, only gauge fields with values in the Killing vectors of g_K are non-zero. Those are precisely the massless gauge fields, since

$$\left(\text{Mass } A_\mu^a\right)^2 \propto \frac{\int_K \langle \mathcal{L}_{e_a} g_K, \mathcal{L}_{e_a} g_K \rangle \text{vol}_{g_K}}{2 \int_K g_K(e_a, e_a) \text{vol}_{g_K}} \quad (1.4)$$

for any (divergence-free) vector field e_a on a compact K [36]. Therefore, we conclude that when the geodesic traverses regions of spacetime with changing g_K or with massive gauge fields, such as the weak force field, the particle's classical rest mass $m(s)$ can indeed vary.

Rest mass variation can be understood as a transfer between 4D momentum and internal momentum. This transfer does not occur in regions with vacuum-type, product

geometries $g_M \times g_K$. There, the two momentum components are uncoupled and separately conserved. If massless gauge fields are turned on the geometry is no longer a product, but 4D motion is only lightly coupled to internal motion. The direction of 4D momentum can change but its magnitude is still conserved. These are Lorentz force-type motions. Internal and 4D momenta will just rotate within the vertical and horizontal subspaces, respectively. In regions where the higher-dimensional geometry is very different from a product, only the full momentum vector conserves its norm along general geodesics. The norms of the 4D and internal components can both change, with opposite signs, due to a rotation of the full momentum mixing those components. This process is perceived as a change of the particle's rest mass. The geometry distortions that produce it correspond to the presence of massive gauge fields or to a non-constant internal metric.

Note that having an internal metric that changes along M_4 is equivalent to having non-constant Higgs-like fields. If we regard the components of g_K as fields on M_4 , they play the exact role of Higgs fields in usual gauge theories. This is apparent from the decomposition of the higher-dimensional scalar curvature R_{g_P} in the Einstein-Hilbert action,

$$\int_P R_{g_P} \text{vol}_{g_P} = \int_P \left[R_{g_M} + R_{g_K} - \frac{1}{4} |F_A|^2 - \frac{1}{4} |d^A g_K|^2 + |d^A (\text{vol}_{g_K})|^2 \right] \text{vol}_{g_P}. \quad (1.5)$$

This formula extends the usual Kaluza-Klein result to the setting of general Riemannian submersions, where the geometry of the fibres can change. For more details see [36].

Charges and charge variation

Now let us consider particles' charges. Let ξ be an electromagnetic-like Killing vector field of g_K . It is an internal Killing field that commutes with all other Killing fields on K . If a particle's motion is parameterized by a geodesic $\gamma(s)$ on P , we define the particle's charge with respect to ξ as the scalar

$$q_\xi(s) := -g_P(\xi, p) = -g_K(\xi, p^\mathcal{Y}). \quad (1.6)$$

Section 4 shows that this is a constant of geodesic motion in regions where only massless gauge fields are present and the internal geometry does not change, independently of g_M . But there are more constants of motion in these regions. Essentially, there is one constant for every summand in the decomposition

$$\mathfrak{k} = \mathfrak{a}_1 \oplus \cdots \oplus \mathfrak{a}_m \quad (1.7)$$

of the Killing algebra of g_K as a sum of abelian or simple subalgebras. That constant of motion measures how orthogonal the derivative vector $\dot{\gamma}(s)$ is to the subspace of $T_{\gamma(s)}P$ spanned by the Killing fields in the corresponding summand. And this is independent of s . The Killing algebra of g_K , recall, determines the massless gauge fields in the model.

In regions where g_K is constant but some massive gauge fields are non-zero, the charges described above are well-defined but are not constants of motion anymore. For example, in section 8 it is shown that $q_\xi(s)$ evolves as

$$\frac{d}{ds} q_\xi(s) = A^a(\dot{\gamma}_M) g_P([\xi, e_a], p) \quad (1.8)$$

along a higher-dimensional geodesic $\gamma(s)$ with momentum $p(s)$. So the test particle's electromagnetic-like ξ -charge may vary when A^a is non-zero and the associated the gauge boson is ξ -charged (i.e. $[\xi, e_a] \neq 0$ as a vector field on K). This effect apparently has not been reported before. It agrees with the physical fact that interactions mediated by massive, charged gauge bosons, such as the Standard Model's W bosons, can modify the charge of particles.

The charge variation formula can be extended to the scalars associated to the simple summands \mathfrak{a}_r in the Killing algebra of g_K . This variation is different from the rotation of isospin along a geodesic found in [28, 38] and reviewed in [39]. More details in section 8.

Equations of geodesic motion

In section 5 we write down the equation of geodesic motion for a general submersive metric on $M_4 \times K$, as translated from [11]. The horizontal component of that equation says that the projection of the geodesic to four dimensions, denoted $\gamma_M(s)$, satisfies

$$g_M(\nabla_{\dot{\gamma}_M}^M \dot{\gamma}_M, X) = g_K(e_a, \dot{\gamma}^\mathcal{Y}) F_A^a(\dot{\gamma}_M, X) + \frac{1}{2} (d^A g_K)_X(\dot{\gamma}^\mathcal{Y}, \dot{\gamma}^\mathcal{Y}). \quad (1.9)$$

This is a generalization of the equation derived by Kerner when K is a Lie group with a constant bi-invariant metric [28]. It also extends the equations obtained when K is a circle of variable size [1–3, 5]. In regions where there are no massive gauge fields and the internal geometry is constant, the term $d^A g_K$ vanishes and the equation simplifies. It reduces to the usual Lorentz force equation when only an electromagnetic-like gauge field is present, as in the 5D calculation. This is verified in section 8 using the previous definitions of mass and charge. It justifies the interpretation of the scalar $q_\xi(s)$ as a charge.

Circumventing traditional difficulties

Kaluza-Klein models are often studied under strong simplifying assumptions, such as minimal 5D dimensions, or constant internal geometry, or the assumption that all relevant gauge fields are associated with internal isometries. Those simplifications facilitate the analysis but also create problems. In fact, we argue that some of the difficulties traditionally attributed to the Kaluza-Klein framework are due to those simplifications.

For example, the physical weak force field is usually associated to an $SU(2)$ -isometry of internal space. But its bosons are massive, even if light when compared to the Planck mass. So the mass formula (1.4) suggests that the weak field should not be associated with exact isometries of g_K . The Lie derivatives $\mathcal{L}_{e_a}g_K$ can be small yet non-zero. Abandoning the exact isometry assumption has an extra advantage, as described in [36]. It offers a possible way out of the main no-go arguments against having chiral fermions in Kaluza-Klein, such as the arguments based on the Atiyah-Hirzebruch theorem [40].

Another often-cited difficulty, in the 5D model, is that the Lorentz force equation of motion can be recovered from geodesics only in regions where the internal circle has constant size. And if this condition is granted, the full 5D equations of motion force the norm $|F_A|^2$ of the electromagnetic field strength to vanish in those same regions, which is not realistic [19, 20]. Moreover, even ignoring that problem, the range of 4D Lorentz force motions determined by 5D geodesics has severe limitations. Due to the normalization condition of the Killing field ξ , all 5D geodesics project down to Lorentz force motions on M_4 with q/m ratios that are way too low when compared to the physical ratios for elementary particles [2, 4]. So 5D geodesics cannot describe the motion of charged elementary particles.

Our first point is that these difficulties are less acute in higher-dimensional models. For example, for higher-dimensional K , after transforming the Lagrangian in (9.1) to the Einstein frame, the local constancy of internal volume only implies that²

$$\left(\frac{\kappa_P}{\kappa_M \text{Vol}_{g_K}}\right)^2 g_K(e_a, e_b) (F_A^a)^{\mu\nu} (F_A^b)_{\mu\nu} + R_{g_K} - \frac{2\Lambda k}{k+2} = 0, \quad (1.10)$$

where k denotes the dimension of K . So there is room for $|F_A|^2$ to vary, as long as those changes are compensated by variations of the internal metric g_K that change the scalar curvature R_{g_K} without affecting the total volume. For example TT-deformations of g_K . Moreover, these constraints are derived solely from the Einstein-Hilbert action on P , which should not tell the whole story for realistic models operating at different scales [36].

In section 9 we address the second difficulty, the small q/m ratios implied in 5D geodesics. On the one hand, it is shown that for higher-dimensional K the normalization of the Killing field no longer determines the value of $g_K(\xi, \xi)$, only the average value of that norm over K . This makes the problem less acute, because that average can be significantly different from the point value $g_K(\xi, \xi)$ that appears in the geodesic equation. Moreover, we note that the usual derivation of the normalization condition relies on a traditional but undue assumption. Namely, it assumes that the physical metric g_M on M_4 coincides with the projection \tilde{g}_M of the metric \tilde{g}_P that appears in the higher-dimensional

²This equation can be derived from the last equation in section 3.4 of [36] after expressing the normalized metric \bar{g}_K of that section in terms of the plain, un-normalized metric g_K .

Einstein-Hilbert action. And in general they should not coincide, only belong to the same conformal class. This is related to the well-known need to transform the dimensionally-reduced Lagrangian from the Jordan frame to the Einstein frame through a Weyl rescaling of g_M . And when this is taken into account, the normalization condition for ξ need not be problematic anymore.

A unique speed in higher dimensions

In section 10 we advocate the naturalness of the postulate that elementary particles always travel at the speed of light in higher dimensions. It is the projection of velocities to three dimensions that appears to produce speeds in the range $[0, c]$, as observed macroscopically. This is equivalent to saying that particles always follow null paths on P .

This hypothesis is not entirely unreasonable because null paths on P always project down to timelike or to null paths on M_4 , in the Kaluza-Klein framework. They never project down to spacelike paths. Higher-dimensional null paths can cover all types of causal motion on Minkowski space. Timelike paths on P do not seem necessary. Therefore, for simplicity's sake, it is natural to do away with timelike paths and investigate the consistency of a dynamical model entirely based on null paths on P .

In these conditions, all particles obey an energy-momentum relation similar to that of photons, but in higher dimensions. It projects down to the usual energy-momentum relation in four dimensions. For example, when $g_P = g_M + g_K$ is a simple product metric, the higher-dimensional momentum vector can be written in an inertial frame as $p = (E, \mathbf{p} + p_K)$, where E is the particle's energy, \mathbf{p} and p_K are its 3-momentum and internal momentum vectors, respectively, and we put $c = 1$. So when p is null with respect to g_P we get that

$$E^2 = g_P(\mathbf{p} + p_K, \mathbf{p} + p_K) = |\mathbf{p}|^2 + g_K(p_K, p_K).$$

The first equality is a photon-like energy-momentum relation in higher-dimensions. The second equality becomes the usual 4D energy-momentum relation if the particle's rest mass is identified with the norm of its internal momentum, $m^2 = g_K(p_K, p_K)$.

If the higher-dimensional speed is always c , as advocated, a particle at rest on 3D space is necessarily moving at full speed c along K . Then the associated kinetic energy is a natural source of the particle's energy at spatial rest, $E_0 = mc^2$. It is appealing to think that rest energy is simply the kinetic energy of internal motion, with no need for alternative mechanisms to store energy in a point-like mass.

The hypothesis of a unique speed in higher dimensions provides a natural origin for 3D rest energy. Conversely, the assumption of a fully kinetic origin of rest energy, if

granted, also implies that timelike geodesics in higher-dimensions should not be allowed to represent physical motions. Otherwise, a timelike particle moving with a 3-dimensional speed lower than c could very well have zero velocity along K , and hence have no rest energy or rest mass. And such particles have never been observed. Thus, when 3D rest energy is identified with the internal kinetic energy, only allowing null geodesics on P derives from the experimental fact that massless particles always travel at the speed of light on M_4 . Having a unique finite speed c for all elementary particles is also a mathematically attractive feature, simpler than having a closed interval $[0, c]$ of possible speeds. The inescapable price is having to work with a higher-dimensional spacetime, of course.

Spaces of null geodesics

Let \mathcal{N}_h^+ denote the space of null geodesics starting at a point h in P and moving forward in time. Each geodesic $\gamma(s)$ in this space is characterized by its null tangent vector $\dot{\gamma}(0)$ at h . In section 11 we describe two distinct parameterizations of \mathcal{N}_h^+ , one relying on the particles' momenta and the other on the celestial sphere of velocities.

Different geodesics in \mathcal{N}_h^+ represent the motion of particles with different masses and electromagnetic charges. Fixing the values of these constants carves out a smaller subspace $\mathcal{N}_h^+(m, q_\xi)$ inside \mathcal{N}_h^+ . In section 11 it is shown that

$$\mathcal{N}_h^+(m, q_\xi) \simeq \begin{cases} \emptyset & \text{if } |q_\xi| > m c |\xi| \\ \mathbb{R}^3 & \text{if } |q_\xi| = m c |\xi| \\ \mathbb{R}^3 \times S^{k-2} & \text{if } |q_\xi| < m c |\xi|. \end{cases} \quad (1.11)$$

Here $|\xi|$ denotes the Riemannian length $\sqrt{(g_K)_h(\xi, \xi)}$ of the Killing vector field at the point h . Thus, in this model, particles with a given classical mass cannot have arbitrarily strong charge. This is natural because, according to (6.6), mass is related to the norm of vertical momentum, while q_ξ measures the component of that same momentum along ξ .

Conceptual simplicity

Kaluza-Klein models strive for conceptual unification at the classical level, before thinking about quantization. Traditionally, they mainly deal with the unification of gauge fields and the 4D metric as components of a unique, higher-dimensional metric. Both in abelian and non-abelian gauge theories. It should also be possible to describe spontaneous symmetry breaking as a dynamical process of the internal metric, in which the isometry group of g_K is broken to generate the gauge bosons' mass according to formula (1.4) [36].

The main message of the present paper, in turn, is that conceptual unification can be taken farther in simple higher-dimensional models, hopefully without contradicting observation. At the level of test particles and geodesic motion, one can construct a model where massive and massless particles both travel at the speed of light in higher dimensions, satisfying a photon-like energy-momentum relation that projects down to the usual 4D relation on M_4 . A model where mass, charges and 4D momentum describe different aspects of a unique higher-dimensional momentum vector, which is covariantly conserved along geodesics. A model where the energy stored in 3D rest mass of classical particles is simply the kinetic energy of internal motion.

In this picture, the classical rest mass is not a constant attribute of a test particle. It is a dynamical quantity measuring the internal component of the particle's momentum. It can vary along geodesics if the background geometry is sufficiently distorted away from the vacuum configuration, since this enables transfers between the horizontal and vertical components of momentum. In particular, the geodesic model illustrates how a higher-dimensional, classical theory can exhibit qualitative features (particle charge and mass change) that are usually reserved for QFT calculations.

2 Motion in the vacuum and 4D inertial frames

To establish notation, we start by considering motion on $P = M_4 \times K$ equipped with a product metric $g_M \times g_K$. Consider an inertial frame on M_4 with coordinates (t, x^1, x^2, x^3) and write the Minkowski metric as

$$g_M = dx^1 \otimes dx^1 + dx^2 \otimes dx^2 + dx^3 \otimes dx^3 - c^2 dt \otimes dt . \quad (2.1)$$

Take arbitrary local coordinates y^j on the internal space K . In an inertial frame related by a boost with speed u along the x^1 -axis, the new coordinates satisfy the usual relations

$$\begin{aligned} dx'^1 &= \frac{dx^1 - u dt}{\sqrt{1 - u^2/c^2}} & dt' &= \frac{dt - u dx^1/c^2}{\sqrt{1 - u^2/c^2}} \\ dx'^n &= dx^n \quad \text{for } n = 2, 3 & dy'^j &= dy^j . \end{aligned} \quad (2.2)$$

All coordinates transversal to the boost remain invariant, including the internal ones. A particle moving on $M_4 \times K$ can be parameterized by a curve $\gamma(s) = (\gamma_M(s), \gamma_K(s))$. Taking the derivative with respect to s , we have the tangent vectors

$$\frac{d\gamma}{ds}(s) = \dot{\gamma} = \dot{\gamma}_M + \dot{\gamma}_K ,$$

with $\dot{\gamma}_M$ tangent to M_4 and $\dot{\gamma}_K$ tangent to K . They satisfy the relation

$$g_K(\dot{\gamma}_K, \dot{\gamma}_K) - g_P(\dot{\gamma}, \dot{\gamma}) = -g_M(\dot{\gamma}_M, \dot{\gamma}_M) = \left(c \frac{d\tau}{ds} \right)^2 , \quad (2.3)$$

where τ denotes the particle's 4D proper time. Since g_K is Riemannian by assumption, the term $g_K(\dot{\gamma}_K, \dot{\gamma}_K)$ is always non-negative. Thus, if γ is a timelike curve on P , the projection γ_M will also be timelike on M_4 . If γ is null on P , the projection γ_M will generically be timelike on M_4 , but can also be null when $\dot{\gamma}_K$ is zero. In principle, even some spacelike curves on P can project down to timelike curves on M_4 , if the norm $g_K(\dot{\gamma}_K, \dot{\gamma}_K)$ is sufficiently large. We will not consider that possibility here. In a Kaluza-Klein model the additional dimensions are interpreted to be physical as well, so subject to the same causal restrictions as the Minkowski dimensions. So we will only consider curves such that $g_P(\dot{\gamma}, \dot{\gamma})$ is negative or zero.

A particle's velocity in the inertial frame (t, x^1, x^2, x^3, y^j) is the derivative of its position with respect to the time coordinate. So the internal velocity of the test particle represented by $\gamma(s)$ is given by

$$v_K = \frac{d\gamma_K}{dt} = \frac{ds}{dt} \frac{d\gamma_K}{ds} = \frac{1}{\dot{\gamma}^0} \dot{\gamma}_K ,$$

which is a vector tangent to K at the point $\gamma_K(s)$. Similarly, the particle's 3D velocity in the frame is

$$\mathbf{v} = \frac{d\gamma^n}{dt} \frac{\partial}{\partial x^n} = \frac{1}{\dot{\gamma}^0} \dot{\gamma}^n \frac{\partial}{\partial x^n} ,$$

with an implicit sum over $n = 1, 2, 3$. The dot denotes derivation with respect to s .

An affine geodesic on P is a curve $\gamma(s)$ satisfying $\nabla_{\dot{\gamma}}\dot{\gamma} = 0$, where ∇ denotes the Levi-Civita connection of g_P . When g_P is a product metric, the projected curves $\gamma_M(s)$ and $\gamma_K(s)$ are also geodesics on the respective spaces. For more general g_P they are not. Although the geodesic equation is defined here using the Levi-Civita connection of g_P , any connection with totally anti-symmetric torsion would lead to the same equation, hence to the same geodesics on P .

3 Curves in Riemannian submersions

Take a Lorentzian metric g_P on the higher-dimensional space $P = M_4 \times K$ such that the projection $\pi : P \rightarrow M_4$ is a Riemannian submersion. As in the usual Kaluza-Klein framework, this metric determines three more familiar objects:

- i) through projection, a unique Lorentzian metric g_M on M_4 ;
- ii) through restriction to the fibres $\{x\} \times K$, a family of Riemannian metrics $g_K(x)$ on the internal spaces parameterized by the points in M_4 ;
- iii) gauge fields on spacetime, encapsulated in a one-form A on M_4 with values in the Lie algebra of vector fields on K .

The equations linking these objects to the higher-dimensional metric g_P are

$$\begin{aligned} g_P(U, V) &= g_K(U, V) \\ g_P(X, V) &= -g_K(A(X), V) \\ g_P(X, Y) &= g_M(X, Y) + g_K(A(X), A(Y)) \ , \end{aligned} \tag{3.1}$$

for all tangent vectors $X, Y \in TM$ and vertical vectors $U, V \in TK$. These relations generalize the usual the Kaluza ansatz for g_P . They show how to reconstruct the higher-dimensional metric from the data (g_M, A, g_K) . The correspondence between submersive metrics on P and that data is a bijection. This is described in more detail in [36].

Choosing a set $\{e_a\}$ of independent vector fields on K , the one-form on spacetime can be decomposed as a sum

$$A(X) = \sum_a A^a(X) e_a \ , \tag{3.2}$$

where the real-valued coefficients $A^a(X)$ are the traditional gauge fields on M_4 . For general submersive metrics on P this can be an infinite sum, with $\{e_a\}$ being a basis for the full space of vector fields on K , which coincides with the Lie algebra of the diffeomorphism group $\text{Diff}(K)$. The curvature F_A is a two-form on M_4 with values in the Lie algebra of vector fields on K . It can be defined by

$$F_A(X, Y) := (d_M A^a)(X, Y) e_a + A^a(X) A^b(Y) [e_a, e_b] \ ,$$

where the last term is just the Lie bracket $[A(X), A(Y)]$ of vector fields on K .

The tangent space to $P = M_4 \times K$ has a natural decomposition $TP = TM \oplus TK$. Since TK is the kernel of the projection $TP \rightarrow TM$, it is also called the vertical sub-bundle \mathcal{V} of TP . The higher-dimensional metric g_P determines an orthogonal complement $\mathcal{H} \simeq (TK)^\perp$, called the horizontal sub-bundle. So from g_P we get a second decomposition

$$TP = \mathcal{H} \oplus \mathcal{V} \ , \tag{3.3}$$

and every tangent vector $Z \in TP$ can be written as a sum $Z^{\mathcal{H}} + Z^{\mathcal{V}}$. The relation between the two decompositions of TP is quite simple in a Riemannian submersion. Writing $Z = Z_M + Z_K$ according to the decomposition $TP = TM \oplus TK$, we have

$$Z^{\mathcal{V}} = Z_K - A(Z_M) \quad Z^{\mathcal{H}} = Z_M + A(Z_M) \ . \tag{3.4}$$

So the information contained in the gauge one-form A on M is equivalent to the information contained in the horizontal distribution $\mathcal{H} \subset TP$. Geometrically, it is well-known that the curvature F_A is the obstruction to the integrability of the distribution \mathcal{H} , in the sense that it vanishes if and only if P can be foliated by horizontal submanifolds whose tangent space coincides with \mathcal{H} [12].

Now let $\gamma(s)$ be a null curve on P parameterized by s . Let $\gamma_M(s)$ and $\gamma_K(s)$ denote its projections onto the factors M_4 and K . The tangent vectors to P obtained by derivation with respect to s have two decompositions

$$\frac{d\gamma}{ds} = \dot{\gamma} = \dot{\gamma}_M + \dot{\gamma}_K = \dot{\gamma}^{\mathcal{V}} + \dot{\gamma}^{\mathcal{H}}. \quad (3.5)$$

According to (3.4) these are related by

$$\begin{aligned} \dot{\gamma}^{\mathcal{V}} &= \dot{\gamma}_K - A^a(\dot{\gamma}_M) e_a \\ \dot{\gamma}^{\mathcal{H}} &= \dot{\gamma}_M + A^a(\dot{\gamma}_M) e_a. \end{aligned} \quad (3.6)$$

Since g_P restricted to horizontal vectors projects down to g_M and the second decomposition is g_P -orthogonal, we have

$$-g_M(\dot{\gamma}_M, \dot{\gamma}_M) = -g_P(\dot{\gamma}^{\mathcal{H}}, \dot{\gamma}^{\mathcal{H}}) = g_P(\dot{\gamma}^{\mathcal{V}}, \dot{\gamma}^{\mathcal{V}}) - g_P(\dot{\gamma}, \dot{\gamma}). \quad (3.7)$$

The restriction of g_P to vertical vectors is the Riemannian metric g_K on the fibre. Hence the first term on the right-hand side is always non-negative. So is the second term when $\gamma(s)$ is a null or timelike curve on P . In that case, the projection $\gamma_M(s)$ is also a null or timelike curve on M_4 .

The particle's proper time on Minkowski space along the path $\gamma_M(s)$ is measured by

$$c[\tau(s_1) - \tau(s_2)] = \int_{s_1}^{s_2} \sqrt{-g_M(\dot{\gamma}_M, \dot{\gamma}_M)} ds = \int_{s_1}^{s_2} \sqrt{g_K(\dot{\gamma}^{\mathcal{V}}, \dot{\gamma}^{\mathcal{V}}) - g_P(\dot{\gamma}, \dot{\gamma})} ds. \quad (3.8)$$

The last integral depends on g_K and on the gauge fields, as is clear from (3.6).

4 Geodesics among massless gauge fields

The aim of this section is to identify constants of geodesic motion in certain regions of spacetime, namely regions where the Higgs-like scalars are constant and where all massive gauge fields vanish. In the physical world, this would allow an electromagnetic field but would exclude a non-zero weak field, for example. Under these conditions, the classical mass and charge of physical particles are constant. So the definitions of mass and charge adopted in the geodesic model should be searched among quantities that are constants of motion in these regions.

Consider a higher-dimensional submersive metric characterized by the equivalent data $g_P \simeq (g_M, A, g_K)$, as before. In this section we will assume that:

H1) The internal metrics g_K are the same for all fibres;

H2) The one-form $A(X)$ has values in the space of Killing vector fields on (K, g_K) .

According to [36] and the mass formula (1.4), these assumptions correspond to regions of M_4 where the Higgs-like scalars are constant and only massless gauge fields are present. In particular, the second fundamental form of the fibres, denoted S , which is equivalent to the covariant derivative $d^A g_K$ of (1.3), vanishes. With these assumptions, the equations of section 5 for higher-dimensional geodesics simplify considerably.

For example, since the norm $g_P(\dot{\gamma}, \dot{\gamma})$ is always a constant of geodesic motion, equations (3.7) and (5.2) imply that, for geodesics,

$$\frac{d}{ds} \left(c \frac{d\tau}{ds} \right)^2 = - \frac{d}{ds} g_M(\dot{\gamma}_M, \dot{\gamma}_M) = \frac{d}{ds} g_P(\dot{\gamma}^\mathcal{Y}, \dot{\gamma}^\mathcal{Y}) = 0 . \quad (4.1)$$

So in these regions the rate of change of proper time, $\frac{d\tau}{ds}$, is a constant of motion. In section 6 we will relate it to the mass of the test particle. Moreover, since S vanishes and γ is a geodesic by assumption, the two equations in (5.1) are simplified to

$$\begin{aligned} g_P(\nabla_{\dot{\gamma}} \dot{\gamma}^\mathcal{Y}, V) &= 0 \\ g_M(\nabla_{\dot{\gamma}_M}^M \dot{\gamma}_M, X) &= g_P(e_a, \dot{\gamma}) F_A^a(\dot{\gamma}_M, X) . \end{aligned} \quad (4.2)$$

Here V is any vertical vector on P and X is any vector on the base M_4 . The first equation describes the vertical component $\dot{\gamma}^\mathcal{Y}$ of the tangent $\dot{\gamma}$. It says that although the vectors $\dot{\gamma}^\mathcal{Y}$ are not parallelly transported along the geodesic, as the $\dot{\gamma}$ are, at least the vertical part of the covariant derivative $\nabla_{\dot{\gamma}} \dot{\gamma}^\mathcal{Y}$ vanishes. The second equation says that the projection of the geodesic to Minkowski space is a curve $\gamma_M(s)$ satisfying something similar to a Lorentz force law [41, sec. 4.3], but with more gauge fields involved. It reduces to the equation derived by Kerner when K is a Lie group and g_K is a bi-invariant metric [28]. The inner-products $g_P(e_a, \dot{\gamma})$ play the role of ‘‘charges’’, coupling the Minkowski motion $\gamma_M(s)$ to the curvature F_A^a of the background gauge fields. The next few paragraphs will investigate the extent to which these inner-products are constant along the geodesics, so that (4.2) truly resembles a Lorentz force equation of motion.

Constants of motion

Consider a general geodesic $\gamma(s)$ on the higher-dimensional P . As usual, if Z is a Killing vector field with respect to g_P , we have that

$$\frac{d}{ds} g_P(\dot{\gamma}, Z) |_{\gamma(s)} = g_P(\nabla_{\dot{\gamma}} \dot{\gamma}, Z) + g_P(\dot{\gamma}, \nabla_{\dot{\gamma}} Z) = 0 .$$

So the inner-product $g_P(\dot{\gamma}, Z)$ is constant along the geodesic $\gamma(s)$.

For a simple product metric, $g_P = g_M \times g_K$, the Killing fields of g_P are sums of Killing fields of g_M and g_K . So all the Killing fields of g_K determine constants of geodesic motion, besides those determined by the isometries of M_4 . However, these additional constants cannot be perceived from the 4D projection of motion. A product metric has vanishing gauge fields and constant internal geometry, so the second relation in (4.2) says that γ_M is a pure geodesic of g_M , uncoupled to the internal metric and to internal motion.

When the gauge fields are non-zero, some of the Killing fields of g_K may no longer preserve the higher-dimensional metric $g_P \simeq (g_M, A, g_K)$ on $M_4 \times K$. So the description of the constants of motion becomes less straightforward. It is still simple enough, however, if we assume that g_P satisfies conditions H1 and H2. Let us start by describing how the general formulae of section 5 are simplified under H1.

Lemma 4.1. *Let $g_P \simeq (g_M, A, g_K)$ be a submersive metric on P satisfying assumption H1. Let $\gamma(s)$ be a geodesic of g_P . If V is a Killing field of g_K , we have that*

$$\frac{d}{ds} g_P(V, \dot{\gamma}) = A^a(\dot{\gamma}_M) g_P([e_a, V], \dot{\gamma}) . \quad (4.3)$$

Proof. Assumption H1 says that the internal metric is the same for all fibres. So if V is a Killing field of g_K , the term with $\mathcal{L}_V g_K$ vanishes in (5.7). Moreover, if we regard V as a vector field on $M_4 \times K$ that is constant along the M_4 direction, the Lie bracket $[X, V]$ vanishes for arbitrary vector fields X on M_4 . This simplifies (5.8). The result follows from the combination of the simplified forms of (5.7) and (5.8). \square

Now, according to assumption H2, the gauge one-form $A(X)$ has values on the space of Killing fields of g_K . So when V commutes with all other Killing fields on K , formula (4.3) implies that $g_P(V, \dot{\gamma})$ has vanishing derivative with respect to the geodesic parameter s . Using the equivalence relations in proposition 5.2, we conclude that:

Lemma 4.2. *Let $g_P \simeq (g_M, A, g_K)$ be a submersive metric on P satisfying assumptions H1 and H2. Let ξ be a Killing vector field on (K, g_K) that commutes with all other Killing fields. Then ξ is also a Killing field on (P, g_P) and the inner-product $g_P(\xi, \dot{\gamma})$ is a constant of motion for every geodesic γ on P .*

If the vector field ξ does not commute with all other Killing fields of g_K , then in general ξ will not be Killing for g_P . However, it is still possible to extract a constant of motion from the subalgebra of the Killing algebra that contains V . To see how this comes about, we must describe the Killing algebra of g_K in more detail.

Since K is a compact manifold by assumption, the isometry group of g_K is a finite-dimensional, compact Lie group. The corresponding Lie algebra can be identified with

the algebra \mathfrak{k} of Killing vector fields on K . It admits a decomposition of the form

$$\mathfrak{k} = \mathfrak{a}_1 \oplus \cdots \oplus \mathfrak{a}_m , \quad (4.4)$$

where the \mathfrak{a}_r are either $\mathfrak{u}(1)$ lines or simple, non-abelian Lie algebras. To describe the constants of motion associated to the summands, let $\{\xi_{r,b} : 1 \leq b \leq \dim \mathfrak{a}_r\}$ denote a basis of \mathfrak{a}_r . Each $\xi_{r,b}$ is a Killing field of g_K . Since \mathfrak{k} is compact, it is possible to choose the basis so that the structure constants defined by $[\xi_{r,b}, \xi_{r,c}] = (f_r)_{bc}^d \xi_{r,d}$ are totally anti-symmetric in their three indices (see lemma A.5). With this choice, we have the following result.

Proposition 4.3. *Let $g_P \simeq (g_M, A, g_K)$ be a submersive metric on P satisfying assumptions H1 and H2. For each subalgebra \mathfrak{a}_r of the Killing algebra of g_K , the scalar*

$$\sum_b [g_P(\xi_{r,b}, \dot{\gamma})]^2$$

is a constant of motion for every geodesic $\gamma(s)$ on P .

Proof. To justify the assertion, observe that formula (4.3) implies that

$$\frac{d}{ds} [g_P(\xi_{r,b}, \dot{\gamma})]^2 = 2 g_P(\xi_{r,b}, \dot{\gamma}) \frac{d}{ds} g_P(\xi_{r,b}, \dot{\gamma}) = 2 g_P(\xi_{r,b}, \dot{\gamma}) g_P([A(\dot{\gamma}_M), \xi_{r,b}], \dot{\gamma}) .$$

By assumption H2, the contraction $A(\dot{\gamma}_M)$ is a Killing vector field on K , so a vector in the Killing algebra (4.4). Its component in the subspace \mathfrak{a}_r to which $\xi_{r,b}$ belongs is just $\sum_c A^c(\dot{\gamma}_M) \xi_{r,c}$. Since (4.4) is a Lie algebra decomposition, not just a vector space decomposition, this component is the only one that matters when calculating the bracket $[A(\dot{\gamma}_M), \xi_{r,b}]$. All other components commute with $\xi_{r,b}$. So we have

$$[A(\dot{\gamma}_M), \xi_{r,b}] = \sum_c A^c(\dot{\gamma}_M) [\xi_{r,c}, \xi_{r,b}] = \sum_{c,d} A^c(\dot{\gamma}_M) (f_r)_{cb}^d \xi_{r,d} .$$

Thus, we finally get that

$$\frac{d}{ds} \sum_b [g_P(\xi_{r,b}, \dot{\gamma})]^2 = 2 \sum_{b,c,d} g_P(\xi_{r,b}, \dot{\gamma}) A^c(\dot{\gamma}_M) (f_r)_{cb}^d g_P(\xi_{r,d}, \dot{\gamma}) = 0 ,$$

where the last equality follows from the anti-symmetry of the structure constants f_r in the indices b and d . \square

This proposition gives us constants of geodesic motion in regions of spacetime where all gauge fields are massless and the internal metric is constant. However, inspecting the calculation in the proof, one quickly realizes that some of the original assumptions may be relaxed. We can drop H2 and accept non-zero massive gauge fields as long as those fields commute with the Killing fields of g_K . More precisely, the previous proposition can be generalized by a similar calculation to the following result.

Proposition 4.4. *Let $g_P \simeq (g_M, A, g_K)$ be a submersive metric on P satisfying assumption H1. Suppose that the gauge one-form $A(X)$ has values in a subspace of vector fields on K of the form $\mathfrak{a}_r \oplus \mathfrak{h}$ with the commutation relation $[\mathfrak{a}_r, \mathfrak{h}] = 0$. Then the scalar*

$$\sum_b [g_P(\xi_{r,b}, \dot{\gamma})]^2$$

is a constant of motion for every geodesic $\gamma(s)$ on P .

The difference between the two propositions can be illustrated in the case where $K = G$ is a compact, simple Lie group and g_K is a generic left-invariant metric. For more details about these metrics see [36, 42], for example. The Killing algebra of g_K is the space of right-invariant vector fields on G , denoted \mathfrak{g}^R . If $\{\xi_b\}$ is a basis of $\text{Lie}(G)$ that is orthonormal with respect to the Killing form, then its right-invariant extension to G , $\{\xi_b^R\}$, is a basis of \mathfrak{g}^R with totally anti-symmetric structure constants. The left-invariant vector fields on G are not Killing, but their Lie bracket with the right-invariant fields does vanish. So we are in the conditions of Proposition 4.4 with $\mathfrak{a}_r = \mathfrak{g}^R$ and $\mathfrak{h} = \mathfrak{g}^L$. This means that the scalar $\sum_b [g_P(\xi_b^R, \dot{\gamma})]^2$ will be a constant of motion even in the presence of the massive gauge fields associated to the left-invariant vector fields on K .

A similar example shows that these constants of geodesic motion are not necessarily all independent of each other. To this end, still take K to be a compact, simple Lie group but now let g_K be a bi-invariant metric. Then the Killing algebra of g_K is the sum of the spaces of left-invariant and right-invariant vector fields, $\mathfrak{g}^L \oplus \mathfrak{g}^R$. If $\{\xi_b\}$ is a basis of $\text{Lie}(G)$ orthonormal with respect to the Killing form, then its left-invariant extension, denoted $\{\xi_b^L\}$, is a basis of \mathfrak{g}^L with totally anti-symmetric structure constants. Moreover, at every point of G , the vector fields ξ_b^L are g_K -orthogonal to each other and span the tangent space to the point. Similarly for the right-invariant extensions $\{\xi_b^R\}$. Then it is straightforward to check that the two constants of geodesic motion associated by proposition 4.3 to the summands \mathfrak{g}^L and \mathfrak{g}^R coincide with each other for all geodesics. Both are equal to the norm $g_K(\dot{\gamma}^\mathcal{V}, \dot{\gamma}^\mathcal{V})$, up to a constant factor.

5 Geodesics in general Riemannian submersions

Equations of motion

Let $\gamma(s)$ be a general curve on P and let ∇ denote the Levi-Civita connection on TP associated with the metric g_P . The covariant derivative $\nabla_{\dot{\gamma}} \dot{\gamma}$ determines the parallel transport of the tangent vector $\dot{\gamma}$ along the path $\gamma(s)$. When $P = M_4 \times K$ and the metric g_P defines a Riemannian submersion, one can ask how $\nabla_{\dot{\gamma}} \dot{\gamma}$ decomposes into horizontal

and vertical parts. Adapting the notation and using the properties of the tensors involved, as in [36], classic results of O'Neill [11, corollary 1] imply that

$$\begin{aligned} g_P(\nabla_{\dot{\gamma}}\dot{\gamma}, V) &= g_P(\nabla_{\dot{\gamma}}\dot{\gamma}^{\mathcal{V}}, V) - g_P(S_V \dot{\gamma}^{\mathcal{V}}, \dot{\gamma}^{\mathcal{H}}) \\ g_P(\nabla_{\dot{\gamma}}\dot{\gamma}, Z) &= g_M(\nabla_{\dot{\gamma}_M}^M \dot{\gamma}_M, \pi_*Z) + F_A^a(\pi_*Z, \dot{\gamma}_M) g_P(e_a, \dot{\gamma}) + g_P(S_{\dot{\gamma}^{\mathcal{V}}} \dot{\gamma}^{\mathcal{V}}, Z) \end{aligned} \quad (5.1)$$

for any curve $\gamma(s)$ on P , any vertical vector V and any horizontal vector Z in TP . The first equation determines the vertical component of $\nabla_{\dot{\gamma}}\dot{\gamma}$ and the second its horizontal part. The notation here is as in section 3 and in [36, sec. 2]. It differs from the conventional notation in the literature about Riemannian submersions because the latter clashes with the traditional physics notation (see the remark in appendix A). We use the decomposition $TP = \mathcal{H} \oplus \mathcal{V}$; the curvature F_A is a two-form on M with values on the vector fields on K ; the tensor $S : \mathcal{V} \times \mathcal{V} \rightarrow \mathcal{H}$ is the second fundamental form of the fibres of the projection $\pi : P \rightarrow M_4$, which can be identified with the covariant derivative $d^A g_K$ of (1.3) through formula (5.5); the symbol ∇^M denotes the Levi-Civita connection on (M, g_M) . The covariant derivative $\nabla_{\dot{\gamma}_M}^M \dot{\gamma}_M$ is a vector field along the curve $\gamma_M(s)$ on M_4 . It does not vanish in general, since the projected curve $\gamma_M = \pi \circ \gamma$ need not be a geodesic on the base (M, g_M) .

Now let $\gamma(s)$ be a geodesic curve on P satisfying $\nabla_{\dot{\gamma}}\dot{\gamma} = 0$. The first equation in (5.1) implies that the norm of the vertical component $\dot{\gamma}^{\mathcal{V}}$ evolves according to

$$\frac{d}{ds} g_P(\dot{\gamma}^{\mathcal{V}}, \dot{\gamma}^{\mathcal{V}}) = 2 g_P(\nabla_{\dot{\gamma}}\dot{\gamma}^{\mathcal{V}}, \dot{\gamma}^{\mathcal{V}}) = 2 g_P(S_{\dot{\gamma}^{\mathcal{V}}} \dot{\gamma}^{\mathcal{V}}, \dot{\gamma}^{\mathcal{H}}). \quad (5.2)$$

Using formula (3.7) and the fact that $g_P(\dot{\gamma}, \dot{\gamma})$ is always constant along a geodesic, this implies that

$$\frac{d}{ds} \left(c \frac{d\tau}{ds} \right)^2 = - \frac{d}{ds} g_M(\dot{\gamma}_M, \dot{\gamma}_M) = 2 g_P(S_{\dot{\gamma}^{\mathcal{V}}} \dot{\gamma}^{\mathcal{V}}, \dot{\gamma}^{\mathcal{H}}). \quad (5.3)$$

So the norm on M_4 of the tangent $\dot{\gamma}_M$ may not be constant as the parameter s varies. Note that the norm of the full tangent vector $\dot{\gamma}$ is always preserved along the geodesic. It is only the norm of the components $\dot{\gamma}^{\mathcal{V}}$ and $\dot{\gamma}^{\mathcal{H}}$ that may change, with opposite signs, in regions where the tensor S is non-zero, i.e. in regions where the fibres of P are not totally geodesic.

Now consider the second equation in (5.1). If $\gamma(s)$ is a geodesic on P , it implies that the projected curve $\gamma_M = \pi \circ \gamma$ on Minkowski space satisfies

$$g_M(\nabla_{\dot{\gamma}_M}^M \dot{\gamma}_M, \pi_*Z) = - g_P(e_a, \dot{\gamma}) F_A^a(\pi_*Z, \dot{\gamma}_M) - g_P(S_{\dot{\gamma}^{\mathcal{V}}} \dot{\gamma}^{\mathcal{V}}, Z) \quad (5.4)$$

for all horizontal vectors $Z \in TP$. The restriction of g_P to the fibres, denoted g_K , is a family of Riemannian metrics on K parameterized by the points in M_4 . The second

fundamental form S is equivalent to the covariant derivative of g_K along vectors on M_4 , as defined in (1.3). This follows from the identity

$$(d^A g_K)_{\pi_* Z}(U, V) = -2 g_P(S_U V, Z) \quad (5.5)$$

justified in [36, sec. 2]. Therefore, denoting by X the projection $\pi_* Z$ in TM and using that e_a is a vertical vector, the equation above can also be written as

$$g_M(\nabla_{\dot{\gamma}_M}^M \dot{\gamma}_M, X) = g_K(e_a, \dot{\gamma}^\mathcal{Y}) F_A^a(\dot{\gamma}_M, X) + \frac{1}{2} (d^A g_K)_X(\dot{\gamma}^\mathcal{Y}, \dot{\gamma}^\mathcal{Y}) \quad (5.6)$$

for any X in TM . This equation generalizes the Lorentz force equation of motion to regions of spacetime where $S \neq 0$, i.e. to regions where the internal metric g_K is not covariantly constant as one moves along M_4 . It generalizes the equation derived by Kerner when K is a Lie group and g_K is a bi-invariant metric [28], and also the equations for geodesic motion when $K = U(1)$ is a circle with variable size [1–3, 5]. It involves the changing metrics g_K and couples the motion on M_4 to the internal motion $\gamma_K(s)$ through the term $\dot{\gamma}^\mathcal{Y} = \dot{\gamma}_K - A^a(\dot{\gamma}_M) e_a$, described in (3.6).

Horizontal geodesics

Equations (5.1) imply that a horizontal curve $\gamma^{\text{hor}}(s)$ on P , i.e. a curve with vanishing $(\dot{\gamma}^{\text{hor}})^\mathcal{Y}$, satisfies

$$g_P(\nabla_{\dot{\gamma}^{\text{hor}}} \dot{\gamma}^{\text{hor}}, Z) = g_M(\nabla_{\dot{\gamma}_M^{\text{hor}}}^M \dot{\gamma}_M^{\text{hor}}, \pi_* Z)$$

for every tangent vector Z in TP . So the curve γ^{hor} is a geodesic on P if and only if its projection γ_M^{hor} is a geodesic on M_4 . So a direct consequence of (5.1) is that

Proposition 5.1 ([11]). *The horizontal geodesics on (P, g_P) are exactly the horizontal lifts of geodesics on (M_4, g_M) .*

In particular, if the gauge fields A^a and the internal metrics g_K change but the metric g_M on the base does not, then the horizontal geodesics will change as curves on P but their projection to M_4 will remain identical.

Now suppose that $\gamma(s)$ is a causal (timelike or null) curve on P that projects down to a null curve on M_4 . This means that the particle represented by γ is moving at the speed of light on Minkowski space. Using that $g_K(\dot{\gamma}^\mathcal{Y}, \dot{\gamma}^\mathcal{Y})$ and $-g_P(\dot{\gamma}, \dot{\gamma})$ are both non-negative, it follows from (3.7) that a zero $g_M(\dot{\gamma}_M, \dot{\gamma}_M)$ implies that the two latter terms must be zero as well. In other words, γ projects down to a null curve on M_4 if and only if γ itself is horizontal and null on P . Thus, we conclude that particles moving at the speed of light on M_4 are always represented by horizontal, null geodesics on P . Combining this with the previous observations about horizontal geodesics, we get that:

Physical implication. *In a causal, higher-dimensional geodesic model, particles moving at the speed of light on M_4 are always represented by horizontal, null geodesics on P . Their 4D motion is not affected by the configuration of gauge fields and internal metrics on P . They follow the null geodesics on M_4 determined by g_M alone.*

This is not the case for particles travelling at lower speeds on M_4 , since the corresponding geodesics on P can have a non-zero vertical component $\dot{\gamma}^\mathcal{V}$ that, according to (5.1), couples to the gauge fields and internal geometry through the non-vanishing tensors F_A and $d^A g_K$. As mentioned in the Introduction, this property of the geodesic model is incompatible with the motion of the old massless neutrinos, since they were thought to interact with the weak field and travel at the speed of light on Minkowski space. So a causal, geodesic model on $M_4 \times K$ disfavors the existence of massless neutrinos or similar particles. This does not seem to have been remarked before.

Vertical Killing fields

In the next few paragraphs we study the conditions necessary for a Killing field of g_K to be a Killing field of $g_P \simeq (g_M, A, g_K)$ as well. When this happens, we get additional constants of the higher-dimensional, geodesic motion.

Let $\gamma(s)$ be a geodesic for g_P and let V be a vertical vector field on P . In general, the inner-product $g_P(V, \dot{\gamma})$ is not constant along the geodesic. Using lemma A.3 in the appendix, its dependence on the parameter s is calculated to be

$$\begin{aligned}
2 \frac{d}{ds} g_P(V, \dot{\gamma}) &= 2 g_P(\nabla_{\dot{\gamma}} V, \dot{\gamma}) = (\mathcal{L}_V g_P)(\dot{\gamma}, \dot{\gamma}) \\
&= (\mathcal{L}_V g_P)(\dot{\gamma}^\mathcal{H}, \dot{\gamma}^\mathcal{H}) + 2 (\mathcal{L}_V g_P)(\dot{\gamma}^\mathcal{H}, \dot{\gamma}^\mathcal{V}) + (\mathcal{L}_V g_P)(\dot{\gamma}^\mathcal{V}, \dot{\gamma}^\mathcal{V}) \\
&= (\mathcal{L}_V g_K)(\dot{\gamma}^\mathcal{V}, \dot{\gamma}^\mathcal{V}) + 2 g_K([\dot{\gamma}^\mathcal{H}, V], \dot{\gamma}^\mathcal{V}) \\
&= (\mathcal{L}_V g_K)(\dot{\gamma}^\mathcal{V}, \dot{\gamma}^\mathcal{V}) + 2 g_K(d_{\dot{\gamma}^\mathcal{H}}^A V, \dot{\gamma}^\mathcal{V}).
\end{aligned} \tag{5.7}$$

In the last equality we have used the $\text{Diff}(K)$ -covariant derivative of a vertical field V on P along a vector field X on M . Using the Lie bracket of vector fields on P , it is defined through the expression

$$d_X^A V := [X^\mathcal{H}, V] = [X, V] + [A(X), V] = (dV^j)(X) \frac{\partial}{\partial y^j} + A^a(X) [e_a, V], \tag{5.8}$$

where the y^j are any coordinates on K . So $d_X^A V$ is another vertical field on P . These equalities use the fact that V is vertical; that the functions $A^a(X)$ do not depend on the coordinates y^j ; and that the Lie bracket $[X, \partial/\partial y^j]$ vanishes, since these are vector fields on different manifolds. The covariant derivative d^A is an interesting object of study. As

in [36, sec. 2.5], one can check that it is $C^\infty(M)$ -linear in both entries and is equivariant with respect to $\text{Diff}(K)$ -gauge transformations of V and the gauge one-form A .

Now, at a fixed point on P there are geodesics passing through with arbitrary vectors $\dot{\gamma}^\mathcal{Y}$ and $\dot{\gamma}_M$. So formula (5.7) implies the equivalence of conditions 2 and 3 below.

Proposition 5.2. *Let V be a vertical vector field on P and let $g_P \simeq (g_M, A, g_K)$ be a submersive metric. Then the following conditions are equivalent:*

1. V is a Killing vector field of g_P ;
2. The restriction of V to each fibre is Killing and $d_X^A V$ vanishes for all X in TM ;
3. The inner-products $g_P(V, \dot{\gamma})$ are constant for all geodesics $\gamma(s)$ on P .

The equivalence of conditions 1 and 2 follows from lemma A.3 in the appendix.

6 Higher-dimensional momentum and mass

The motion of test particles on the higher-dimensional space P is parameterized by time-like or null geodesics. Let $\gamma(s)$ be such a geodesic. The higher-dimensional momentum vector of the associated particle is the vector tangent to P defined simply by

$$p(s) := \sigma \frac{d\gamma}{ds} = \sigma \dot{\gamma}(s) . \quad (6.1)$$

Here σ is a positive constant with dimension MT^{-1} . Since γ is geodesic, the momentum vectors are covariantly constant along the curve, $\nabla_{\dot{\gamma}} p = 0$. The higher-dimensional momentum vector can be decomposed in two different ways:

$$p(s) = p^\mathcal{H} + p^\mathcal{Y} = p_M + p_K . \quad (6.2)$$

As in (3.5) and (3.6), writing the curve $\gamma(s)$ on $M_4 \times K$ as $(\gamma_M(s), \gamma_K(s))$, the different components of momentum are given by

$$\begin{aligned} p_M &= \sigma \dot{\gamma}_M \\ p_K &= \sigma \dot{\gamma}_K \\ p^\mathcal{H} &= \sigma \dot{\gamma}^\mathcal{H} = p_M + A^a(p_M) e_a \\ p^\mathcal{Y} &= \sigma \dot{\gamma}^\mathcal{Y} = p_K - A^a(p_M) e_a . \end{aligned} \quad (6.3)$$

For each value of s the vectors p_M and p_K can be regarded as tangent to M_4 and K , but they are not g_P -orthogonal to each other. The vectors $p^\mathcal{H}$ and $p^\mathcal{Y}$ are orthogonal on P , but they depend on the gauge fields and mix the components along M_4 and K .

Since γ is timelike or null, by (3.7) the same happens with its projection γ_M to M_4 . So the inner-product $g_M(\dot{\gamma}_M, \dot{\gamma}_M)$ is non-positive. The particle's 4D proper time τ is defined, up to an additive constant, by

$$c \frac{d\tau}{ds} = \sqrt{-g_M(\dot{\gamma}_M, \dot{\gamma}_M)}. \quad (6.4)$$

When this derivative is non-zero, τ can also be used as a parameter for the curve. Then the higher-dimensional momentum vector can be written as

$$p(s) = m(s) \frac{d\gamma}{d\tau}, \quad (6.5)$$

where we have defined the particle's rest mass function

$$m(s) := \sigma \frac{d\tau}{ds} = \frac{1}{c} \sqrt{-g_M(p_M, p_M)} = \frac{1}{c} \sqrt{g_P(p^\nu, p^\nu) - g_P(p, p)}. \quad (6.6)$$

The last equality uses (3.7). The mass function is independent of the 4D frame because it is defined in terms of proper time. It is positive for all geodesics on P that project down to timelike curves $\gamma_M(s)$ on M_4 . It vanishes only when $\gamma(s)$ projects to a null curve on M_4 . Combining (6.5) with formula (6.3) for the momentum components, it is clear that in 4D we also have

$$p_M = m(s) \frac{d\gamma_M}{d\tau}.$$

So as long as p_M is interpreted as the particle's 4-momentum on M_4 , this justifies the interpretation of the scalar $m(s)$ as its mass. In particular, given inertial coordinates (t, x^1, x^2, x^3) on M_4 , the energy E and 3-momentum \mathbf{p} of the particle parameterized by $\gamma(s)$ should be read from the projection p_M through the usual (coordinate-dependent) relation³

$$p_M = \frac{E}{c^2} \frac{\partial}{\partial t} + p^1 \frac{\partial}{\partial x^1} + p^2 \frac{\partial}{\partial x^2} + p^3 \frac{\partial}{\partial x^3}. \quad (6.7)$$

Now consider the case where γ projects down to a null curve on M_4 . By the discussion below proposition 5.1, γ itself must be a horizontal, null curve on P . Then relation (6.4) says that $d\tau/ds$ vanishes, and hence proper time cannot be used to parameterize $\gamma_M(s) = (\gamma^0(s), \gamma^n(s))$. In this case the relation between momentum and mass expressed in (6.5) is no longer valid, but the original definitions in (6.3) still are. Assuming that $\dot{\gamma}^0$ does not vanish, the projected momentum p_M can be written as

$$p_M = \sigma \frac{d\gamma_M}{ds} = \sigma \frac{dt}{ds} \frac{d\gamma_M}{dt} = \sigma \dot{\gamma}^0 \left(\frac{\partial}{\partial t} + \mathbf{v} \right), \quad (6.8)$$

³The energy of the particle measured by an observer at rest in the frame determined by (t, x^1, x^2, x^3) is $E = -g_M(p_M, \frac{\partial}{\partial t})$ [41, sec. 4.3]. So for the Minkowski metric (2.1) we get (6.7).

with $|\mathbf{v}| = c$ because $\gamma_M(s)$ is null on M_4 . Thus, to keep (6.7) valid, we must relate the (massless) particle's energy and 3-momentum to the components of p_M through

$$\frac{E}{c^2} = \sigma \dot{\gamma}^0 \quad p^n = \sigma \dot{\gamma}^0 v^n = \sigma \frac{dt}{ds} \frac{d\gamma^n}{dt} = \sigma \dot{\gamma}^n . \quad (6.9)$$

In particular, using Einstein's relation for photons, the frequency associated to the massless particle represented by the null curve $\gamma_M(s)$, in the frame (t, x^1, x^2, x^3) , is just $\nu = \sigma c^2 \dot{\gamma}^0 / h$.

Observe that, in this picture, there is no physical freedom to reparameterize affine geodesics on P . More precisely, if $\tilde{\gamma}(s) := \gamma(\alpha s)$ is a reparameterization of γ by a positive constant, then $\tilde{\gamma}(s)$ is still a geodesic on P , of course, but it now represents the motion of a test particle with a different higher-dimensional momentum. Evaluating the momentum at the point $\tilde{\gamma}(s)$ in P , we have

$$\tilde{p} |_{\tilde{\gamma}(s)} = \sigma \frac{d\tilde{\gamma}}{ds} |_{\tilde{\gamma}(s)} = \sigma \alpha \frac{d\gamma}{ds} |_{\gamma(\alpha s)} = \alpha p |_{\gamma(\alpha s)} = \alpha p |_{\tilde{\gamma}(s)} .$$

So $\tilde{p} = \alpha p$ as vector fields over the locus on P of the curve. Also, at the point $q = \tilde{\gamma}(s)$, the masses of the particles represented by $\tilde{\gamma}$ and γ , as written in (6.6), are related to each other by

$$\tilde{m}^2 |_q = -c^{-2} g_P(\tilde{p}^{\mathcal{H}}, \tilde{p}^{\mathcal{H}}) |_q = -c^{-2} \alpha^2 g_P(p^{\mathcal{H}}, p^{\mathcal{H}}) |_q = \alpha^2 m^2 |_q .$$

So $\tilde{\gamma}$ and γ parameterize the motion on P of different physical particles. In the case where $\tilde{\gamma}$ and γ are horizontal geodesics, both \tilde{m} and m vanish, of course. But a similar argument implies that the 4-momenta on M_4 of the respective particles are related by

$$\tilde{p}_M |_{\pi(q)} = \alpha p_M |_{\pi(q)} .$$

Hence the two geodesics are associated to massless particles with different energies in the frame (t, x^1, x^2, x^3) .

7 Rest mass variation

According to (5.3), the scalar $m(s)$ defined in (6.6) is a constant of motion when the geodesic traverses regions in P with totally geodesic fibres, i.e. regions where $S = d^A g_K = 0$. In those regions the geodesic parameterizes the motion of a test particle with fixed rest mass. In regions where the internal metric g_K is not covariantly constant, the same formula (5.3) implies that the mass function can vary according to

$$c^2 \frac{d}{ds} m^2(s) = 2 g_P(S_{p^\mathcal{V}} p^\mathcal{V}, \dot{\gamma}^{\mathcal{H}}) = - (d^A g_K)_{\dot{\gamma}_M} (p^\mathcal{V}, p^\mathcal{V}) . \quad (7.1)$$

The last equality uses (5.5). At the same time, identity (3.7) says that the different components of the momentum vector defined in (6.3) are related by

$$-g_M(p_M, p_M) = -g_P(p^{\mathcal{H}}, p^{\mathcal{H}}) = g_P(p^{\mathcal{V}}, p^{\mathcal{V}}) - g_P(p, p). \quad (7.2)$$

Since norm of the full momentum $g_P(p, p)$ is a constant of geodesic motion, it follows from (6.6) that

$$\frac{d}{ds} g_P(p^{\mathcal{V}}, p^{\mathcal{V}}) = -\frac{d}{ds} g_P(p^{\mathcal{H}}, p^{\mathcal{H}}) = c^2 \frac{d}{ds} m^2(s). \quad (7.3)$$

So the norms of the horizontal and vertical components of momentum may change with opposite signs along the motion, while the norm of the full momentum p remains constant. The picture is that in regions with non-vanishing $d^A g_K$ the higher-dimensional geometry is distorted, so forces a transfer between the horizontal and vertical components of momentum of particles in free fall along geodesics. Since internal momentum is seen in 4D physics as rest mass, in those regions we observe a phenomenon of rest mass variation.

For example, suppose that for $s < s_1$ a geodesic $\gamma(s)$ traverses a region of P with vanishing tensor $d^A g_K$; after that it enters a region where massive gauge fields are present and the internal scalars are not constant; then for $s > s_2$ it comes out into another region with vanishing $d^A g_K$. The test particle parameterized by this geodesic has a well-defined and constant mass inside the first and third regions. Those two masses may not coincide, because the geodesic traversed a region with changing internal geometry, which is able to transfer between vertical momentum and horizontal momentum. The classical mass change in the overall process is given by

$$m^2(s_2) - m^2(s_1) = -\frac{1}{c^2} \int_{s_1}^{s_2} (d^A g_K)_{\dot{\gamma}^M} (p^{\mathcal{V}}, p^{\mathcal{V}}) ds.$$

Qualitatively, this property of geodesics agrees with the physical fact that in regions where massive gauge fields are present, or where massive scalar fields (such as the Higgs field) are non-constant, particles may interact with the respective massive bosons and undergo a process of mass change. Of course, in this case we would not call the ingoing and outgoing object the same particle anymore, since, by convention, particles with different rest masses are called different names.

8 Charges and charge variation

Let ξ be a Killing vector field on (K, g_K) that commutes with all other Killing fields of g_K . Let $\gamma(s)$ be a geodesic on P parameterizing the motion of a test particle with momentum $p = \sigma \dot{\gamma}$. We define the particle's charge with respect to ξ to be the scalar function

$$q_\xi(s) := -g_P(\xi, p) = -g_K(\xi, p^{\mathcal{V}}). \quad (8.1)$$

So $q_\xi(s)$ measures the component of the particle's momentum along the vertical Killing field ξ . According to lemma 4.2, this scalar is a constant of motion when the geodesic traverses regions of P that satisfy conditions H1 and H2. Just like the particle's mass $m(s)$. Now suppose that in those regions the gauge form has values in the span of ξ , i.e. suppose that we can write $A(X) = A^\xi(X) \xi$ for all X tangent to M_4 . Then formula (4.2) says that the projection to M_4 of the geodesic $\gamma(s)$ satisfies

$$g_M(\nabla_{\dot{\gamma}_M}^M \dot{\gamma}_M, Y) = \frac{1}{\sigma} q_\xi(s) F_A^\xi(Y, \dot{\gamma}_M) .$$

But by definition of $m(s)$ we have

$$\dot{\gamma}_M = \frac{d\gamma_M}{ds} = \frac{m(s)}{\sigma} \frac{d\gamma_M}{d\tau} .$$

Thus, in terms of the 4-velocity $d\gamma_M/d\tau$ on Minkowski space, we get

$$g_M\left(\nabla_{\frac{d\gamma_M}{d\tau}}^M \frac{d\gamma_M}{d\tau}, Y\right) = \frac{q_\xi}{m} F_A^\xi\left(Y, \frac{d\gamma_M}{d\tau}\right) . \quad (8.2)$$

So the projection to M_4 of the particle's motion satisfies a Lorentz force equation [41, sec. 4.3], as long as we identify the scalar $q_\xi(s)$ with the particle's charge. This calculation justifies that interpretation of the conserved scalar. It is very familiar from 5D Kaluza-Klein [1, 2, 4, 22, 43]. So when the electromagnetic field is the only gauge field present, higher-dimensional Kaluza-Klein is similar to the 5D version. It is not quite the same though, and this can be used to mitigate some of the traditional difficulties of the 5D setting, as will be discussed in section 9.

A curious consequence of the definitions of charge and mass of a test particle is that

$$|q_\xi|^2 = |g_K(\xi, p^\mathcal{Y})|^2 \leq g_K(\xi, \xi) g_K(p^\mathcal{Y}, p^\mathcal{Y}) \leq g_K(\xi, \xi) m^2 c^2 . \quad (8.3)$$

The last inequality uses (6.6) and the fact that $g_P(p, p)$ is non-positive for any timelike or null geodesic on P . Therefore, in this geodesic model, test particles with a given classical mass at a point $\gamma(s)$ cannot have arbitrarily strong charge. There is a maximum limit determined by the norm of the Killing field at that point. The inequality above is saturated only when the particle's momentum p is null and, additionally, its vertical component is proportional to ξ at the point.

Now suppose that condition H1 of section 4 is satisfied but condition H2 is not. In other words, we are in a region of P where the internal metrics g_K are constant but there are both massive and massless gauge fields present. Let ξ denote the same Killing field of g_K as before. It is a direct consequence of lemma 4.1 and definition (8.1) that the charge of the test particle represented by a geodesic $\gamma(s)$ will evolve according to

$$\frac{d}{ds} q_\xi(s) = - \frac{d}{ds} g_P(\xi, p) = A^a(\dot{\gamma}_M) g_P([\xi, e_a], p) . \quad (8.4)$$

Now, by assumption ξ is an electromagnetic-like internal Killing field, i.e. it is a Killing field that commutes with all other Killing fields of g_K . So if only massless gauge fields are non-zero, the corresponding Lie brackets $[\xi, e_a]$ will vanish on K and charge is conserved. This is the content of lemma 4.2 of course. If there are massive gauge fields around, but the corresponding internal fields e_a are such that $[\xi, e_a] = 0$, then charge is still conserved. This is a special case of proposition 4.4. However, if we are in a region of spacetime with non-zero massive gauge fields A^a such that $[\xi, e_a]$ does not vanish, then the charge $q_\xi(s)$ may no longer be a constant of geodesic motion. It will indeed vary if the particle's internal momentum p^γ is not orthogonal to $[\xi, e_a]$ in the tangent space to K .

These properties of geodesic motion agree with the physical fact that gauge interactions mediated massive bosons will preserve a particle's charge if the bosons are neutral (i.e. if $[\xi, e_a]$ vanishes). This is illustrated by the Z boson of the Standard Model. But the interactions will not preserve the particle's charge if the gauge bosons are charged (i.e. if $[\xi, e_a]$ is non-zero), as in the case of the two W bosons. So there is a natural physical interpretation of the phenomenon of geodesic charge variation described by (8.4).

The fact that 4D electromagnetic charge may vary along a geodesic in higher dimensions, as in (8.4), apparently has not been reported before. That may be related to the fact that massive gauge fields are usually discarded in the Kaluza-Klein literature, after the quick remark that their bosons will have masses in the Planck scale. This is remark is not fully justified, in the author's view, having in mind the mass formula (1.4) (see the discussion in [36]). For point particles, the general relations between mass, charges and direction of internal motion also do not seem to be entirely clear in the literature. That may be related to the fact that, in the thoroughly studied 5D setting, it is not possible to distinguish between the direction of internal motion and the directions of the internal Killing vector fields.

As in section 4, under assumption H1 of a constant internal metric, it is possible to define a scalar associated to $\gamma(s)$ and each summand in decomposition (4.4) of the Killing algebra of K :

$$\alpha_{\mathfrak{a}_r}^2(s) := \sum_b \left[g_P(\xi_{r,b}, p) \right]^2 .$$

The scalar measures how orthogonal the particle's momentum vector $p(s)$ is to the subspace of $T_{\gamma(s)}P$ spanned by the Killing fields in the summand \mathfrak{a}_r . It is a constant of geodesic motion if only massless gauge fields are present, as in proposition 4.3. Massive gauge fields that satisfy the conditions of proposition 4.4 also do not spoil its conservation along geodesics. If other massive gauge fields are present, then the scalar evolves along a geodesic $\gamma(s)$ according to

$$\frac{d}{ds} \alpha_{\mathfrak{a}_r}^2(s) = 2 \sum_b g_P(\xi_{r,b}, p) g_P([A(\dot{\gamma}_M), \xi_{r,b}], p) . \quad (8.5)$$

The derivation of this formula is similar to that of (8.4). It is justified in the proof of proposition 4.4.

To end this section, we note that the charge variation described by (8.4) and (8.5) is different from the rotation of non-abelian isospin along a geodesic found in [28, 38] and reviewed in [39], for example. The rotation of isospin was derived in the conditions of constant internal metric g_K and massless gauge fields. It is trivial for abelian charges. In contrast, the charge variation described by (8.4) and (8.5) is due to massive gauge fields and is non-trivial for abelian charges. The relation between the two effects can be summarized by saying that the constants of motion of isospin rotation become variable quantities when massive gauge fields are introduced in the picture. Then (8.5) describes the variation along the geodesic of those former constants of motion.

9 Circumventing a traditional difficulty

Physical metric versus action metric

One manifestation of the Kaluza-Klein miracle, as it is sometimes dubbed, is the derivation of the Lorentz force equation of motion on M_4 from the geodesic equation on P . Another manifestation is the derivation of the Yang-Mills action on M_4 from dimensional reduction of the Einstein-Hilbert action on P . Both derivations start from a higher-dimensional metric satisfying certain restrictions, which generalize the Kaluza ansatz. Both calculations are time-honoured and very well known.

An important detail in the story, perhaps less frequently stated, is that the higher-dimensional metric appearing in these two derivations is not exactly the same. The metric g_P that appears in the geodesic equation, through the covariant derivative $\nabla_{\dot{\gamma}}\dot{\gamma}$, is the physical metric on P . It determines the path of test particles on this spacetime and, after projection to M_4 , determines the physical metric g_M in four dimensions. In contrast, the metric \tilde{g}_P that appears in the higher-dimensional Einstein-Hilbert action,

$$\mathcal{E}(\tilde{g}_P) := \frac{1}{2\kappa_P} \int_P (R_{\tilde{g}_P} - 2\Lambda) \text{vol}_{\tilde{g}_P} , \quad (9.1)$$

should not be directly interpreted as the physical metric on P . Nor should its projection \tilde{g}_M be taken as the physical metric on M_4 . This caution is necessary because dimensional reduction of (9.1) produces a 4D Lagrangian in the Jordan frame, not in the Einstein frame. This means that the gravity component of the 4D Lagrangian does not appear in the traditional guise $R_{\tilde{g}_M} \text{vol}_{\tilde{g}_M}$, but instead appears multiplied by a scalar field that depends on the M_4 -coordinates. This field is a hallmark of Kaluza-Klein theories. It is related to the volume $\text{Vol}_{\tilde{g}_K}$ of the internal spaces, which can be variable across M_4 .

Since the Einstein frame is preferred for a good physical interpretation of the 4D theory [44, 45], the recommended procedure in similar situations is to use a Weyl rescaling $\tilde{g}_M = e^{2\omega} g_M$ to transform the Lagrangian to the Einstein frame. To this end, one should find a function $\omega^2(\text{Vol}_{\tilde{g}_K})$ on M_4 with the property that, after the rescaling, the non-standard terms for \tilde{g}_M reduce to the standard GR term $R_{g_M} \text{vol}_{g_M}$ for g_M , plus derivatives of ω . Then one declares that g_M is the actual physical metric on M_4 , while the original \tilde{g}_M is a combination of g_M with internal scalar fields and has no direct physical meaning.

Let us review this in more detail. For a Riemannian submersion, the higher-dimensional scalar curvature $R_{\tilde{g}_P}$ can be decomposed as sum of natural geometric terms. These are the scalar curvatures $R_{\tilde{g}_M}$ of the base and $R_{\tilde{g}_K}$ of the fibres; the norm of F_A , which originates the Yang-Mills terms after dimensional reduction; the norms of the second fundamental form and mean curvature vector of the fibres. As described in [36], the latter two tensors can also be interpreted as the covariant derivatives along M_4 of the internal metric \tilde{g}_K and of the internal volume form $\text{vol}_{\tilde{g}_K}$. After discarding a total derivative term, the Einstein-Hilbert action for \tilde{g}_P can then be rewritten in the suggestive form

$$\mathcal{E}(\tilde{g}_P) = \frac{1}{2\kappa_P} \int_P \left[R_{\tilde{g}_M} - 2\Lambda + R_{\tilde{g}_K} - \frac{1}{4}|F_A|^2 - \frac{1}{4}|d^A \tilde{g}_K|^2 + |d^A(\text{vol}_{\tilde{g}_K})|^2 \right] \text{vol}_{\tilde{g}_P}. \quad (9.2)$$

This justifies the interpretation of the components of \tilde{g}_K as Higgs-like fields. Due to the appearance of the covariant derivative $d^A \tilde{g}_K$, they can produce non-zero mass terms for the gauge fields through the usual mechanism of spontaneous symmetry breaking. All this is described in more detail in [36]. Now, integration of the first term over the fibre K leads to

$$\int_P R_{\tilde{g}_M} \text{vol}_{\tilde{g}_P} = \int_M (\text{Vol}_{\tilde{g}_K}) R_{\tilde{g}_M} \text{vol}_{\tilde{g}_M}.$$

Since $\text{Vol}_{\tilde{g}_K}$ is in general a non-constant function on M_4 , the gravitational part of the 4D Lagrangian is manifestly not in the Einstein frame. To correct this situation, define a function ω and a new metric g_M on M_4 through the relations

$$\tilde{g}_M = e^{2\omega} g_M \quad \text{with} \quad e^{2\omega} := \frac{\kappa_P}{\kappa_M(\text{Vol}_{\tilde{g}_K})}.$$

Using the standard transformation rules of the Riemannian volume form and the scalar curvature under Weyl rescalings [36, 41], one then calculates that

$$\begin{aligned} \frac{1}{2\kappa_P} \int_P R_{\tilde{g}_M} \text{vol}_{\tilde{g}_P} &= \frac{1}{2\kappa_M} \int_M \frac{\kappa_M(\text{Vol}_{\tilde{g}_K})}{\kappa_P} e^{2\omega} \left[R_{g_M} - 6 \square_{g_M} \omega - 6 |d\omega|_{g_M}^2 \right] \text{vol}_{g_M} \\ &= \frac{1}{2\kappa_M} \int_M \left[R_{g_M} - 6 |d\omega|_{g_M}^2 \right] \text{vol}_{g_M} \end{aligned} \quad (9.3)$$

So the kinetic term of the metric g_M now appears in the precise GR form. It is accompanied by the kinetic term of the scalar field ω on M_4 , which measures the volume of the

internal space. This shows that dimensional reduction of the action $\mathcal{E}(\tilde{g}_P)$ produces a 4D Lagrangian in the Einstein frame as long as the physical metric on M_4 is not identified with \tilde{g}_M — the simple projection of \tilde{g}_P — but is instead identified with

$$g_M = \kappa_P^{-1} \kappa_M (\text{Vol}_{\tilde{g}_K}) \tilde{g}_M .$$

The scalar field will appear with the canonical normalization for Klein-Gordon fields in 4D if we define $\phi := \sqrt{6/\kappa_M} \omega + \text{const}$. This is the scalar field used in [36].

Let us now consider the Yang-Mills term of the action $\mathcal{E}(\tilde{g}_P)$. After dimensional reduction, it will not appear with the canonical normalization of 4D physics for a general, varying internal metric \tilde{g}_K . When \tilde{g}_K changes its isometry group can also change, and so will the structure of massive / massless gauge fields in the model. Thus, the observed structure of Yang-Mills terms in the Standard Model cannot correspond to a global property of the Kaluza-Klein model. Only to a local one, at best, valid in regions of M_4 where \tilde{g}_K is approximately constant and close to its vacuum value \tilde{g}_K^0 , which one assumes to be the present-time conditions. In short, the aim after dimensional reduction is to obtain 4D Yang-Mills terms with the canonical normalization only in regions of spacetime where $\tilde{g}_K \simeq \tilde{g}_K^0$. With this principle established, one observes that

$$\begin{aligned} \int_P |F_A|^2 \text{vol}_{\tilde{g}_P} &= \int_M \tilde{g}_M^{\mu\nu} \tilde{g}_M^{\sigma\rho} (F_A^a)_{\mu\sigma} (F_A^b)_{\nu\rho} \left[\int_K \tilde{g}_K(e_a, e_b) \text{vol}_{\tilde{g}_K} \right] \text{vol}_{\tilde{g}_M} \\ &= \int_M g_M^{\mu\nu} g_M^{\sigma\rho} (F_A^a)_{\mu\sigma} (F_A^b)_{\nu\rho} \left[\int_K \tilde{g}_K(e_a, e_b) \text{vol}_{\tilde{g}_K} \right] \text{vol}_{g_M} . \end{aligned}$$

The last equality reflects the invariance of the 4D Yang-Mills action under Weyl rescalings of g_M . Therefore, in the vacuum conditions $\tilde{g}_K \simeq \tilde{g}_K^0$, dimensional reduction of $\mathcal{E}(\tilde{g}_P)$ will produce terms with the canonical normalization $-\frac{1}{4}(F_A^a)^{\mu\nu}(F_A^a)_{\mu\nu}$ if and only if

$$\int_K \tilde{g}_K^0(e_a, e_b) \text{vol}_{\tilde{g}_K^0} = 2 \kappa_P \delta_{ab} . \quad (9.4)$$

This is a normalization condition for the fields e_a on K . It will be important below.

As in the case of the 4D metric \tilde{g}_M , the internal metric \tilde{g}_K does not need to be directly identified with the physical metric g_K that appears in the geodesic equation on $M_4 \times K$. One can imagine a similar rescaling relation $\tilde{g}_K = e^{2f} g_K$ for some function f on the base M_4 . For example a function dependent on $\text{Vol}_{\tilde{g}_K}$. Natural choices would be to have a constant f or to have $f = \pm \omega$. The $+$ choice corresponds to rescaling the full higher-dimensional metric as $\tilde{g}_P = e^{2\omega} g_P$. There could be more useful choices, however. The additional degree of freedom provided by the identification of the physical metric g_K should be used to facilitate the correspondence between the physics derived from the $\mathcal{E}(\tilde{g}_P)$ action and the observed physics in four dimensions.

Normalization of Killing fields and electromagnetic q/m ratios

One of the traditional difficulties attributed to the Kaluza-Klein framework comes from the normalization condition (9.4) in the 5D model [2]. In our view, it is caused by the lack of freedom in a unidimensional K and, perhaps more importantly, by not distinguishing between the action metric \tilde{g}_M and the physical metric g_M .

Most Kaluza-Klein literature does not make that distinction [44]. So does not perform the field redefinitions $\tilde{g}_M \rightarrow g_M$ to obtain the general, dimensionally-reduced Lagrangian in the Einstein frame. Instead, the standard procedure is to assume the internal vacuum condition $g_K \simeq g_K^0$ before dimensional reduction and then fix the constant κ_P to be $\kappa_M \text{Vol}_{g_K^0}$. This produces a 4D Lagrangian in the Einstein frame, but only in the case of a constant internal metric $g_K = g_K^0$. Moreover, fixing the constant κ_P to a vacuum-dependent value does not seem like a general approach. If the theory had several local vacua, associated to internal spaces of different sizes, one would run into an ambiguous choice of κ_P . A more general approach to obtaining an action in the Einstein frame is to keep κ_P unconstrained and, instead, redefine the physical metric g_M through the Weyl rescaling described above. This produces a Lagrangian in the Einstein frame in all circumstances, even when the internal metric g_K is not constant along M_4 .

After this lengthy preamble, let us describe the actual difficulty. If we take $\tilde{g}_P = g_P$ and set $\kappa_P = \kappa_M \text{Vol}_{g_K^0}$, as in [2], the normalization condition (9.4) would imply that

$$\frac{1}{\text{Vol}_{g_K^0}} \int_K g_K^0(e_a, e_b) \text{vol}_{g_K^0} = 2 \kappa_M \delta_{ab} . \quad (9.5)$$

In the 5D model, with g_K^0 equal to the round metric on $K = \text{U}(1)$ and $e_a = \xi$ equal to its standard Killing field, the function $g_K^0(\xi, \xi)$ is constant along the circle K . So the normalization condition for ξ reduces to

$$g_K^0(\xi, \xi) = 2 \kappa_M .$$

This creates a problem when comparing to the Lorentz force motions coming from the geodesic picture. For instance, formula (8.3) says that for non-trivial geodesics on $M_4 \times \text{U}(1)$ we must have

$$\frac{q\xi}{m} \leq c \sqrt{g_K^0(\xi, \xi)} = c \sqrt{2 \kappa_M} ,$$

while the charge to mass ratio of the electron is many orders of magnitude above this value. This is the often-mentioned difficulty described in [2]. See also [4].

As described above, it only arises if we assume that the action metric \tilde{g}_P coincides with the physical metric g_P and fix the value $\kappa_P = \kappa_M \text{Vol}_{g_K^0}$. With the more general transformation of \tilde{g}_P to the Einstein frame, the constant κ_P remains unconstrained and

the normalization condition (9.4) need not be problematic. Furthermore, even if we insist on setting $\kappa_P = \kappa_M \text{Vol}_{g_K^0}$, it is only for unidimensional fibres that (9.5) actually fixes the value of $g_K^0(\xi, \xi)$. In higher dimensions only the average of $g_K^0(\xi, \xi)$ over the manifold (K, g_K^0) is fixed. And in principle that average can be very different from the value of $g_K^0(\xi, \xi)$ at a given point. Finally, one could also use the freedom in the identification of the physical internal metric g_K , as opposed to the action metric \tilde{g}_K , to mitigate problems created by the normalization condition (9.4). So this specific traditional difficulty seems to be circumventable. For observations about other difficulties, such as the compatibility of Kaluza-Klein with chiral fermions, see [36].

10 A unique speed in higher dimensions

Kaluza-Klein models strive for conceptual simplicity at classical level, before thinking about quantization. Following this motto, in this section we advocate the naturalness of the postulate that all elementary particles travel at the speed of light in higher dimensions. It is the projection of velocities to three dimensions that appears to produce speeds in the range $[0, c]$, as observed macroscopically. Under this assumption the previous geodesic model is simplified, as we will see. Both massless and massive particles satisfy a photon-like energy-momentum relation in higher dimensions, which projects down to the usual energy-momentum relation in 4D. The energy stored in 3D rest mass of classical particles becomes entirely due to the kinetic energy of internal motion.

Let us then assume that test particles always follow null geodesics on P . In a Riemannian submersion, we know that tangent vectors to a null path $\gamma(s)$ in higher dimensions satisfy

$$-g_M(\dot{\gamma}_M, \dot{\gamma}_M) = -g_P(\dot{\gamma}^{\mathcal{H}}, \dot{\gamma}^{\mathcal{H}}) = g_P(\dot{\gamma}^{\mathcal{V}}, \dot{\gamma}^{\mathcal{V}}) . \quad (10.1)$$

This is just a simplification of (3.7). Since $\dot{\gamma}^{\mathcal{V}}$ is tangent to the fibres and the restriction of g_P to those fibres are the Riemannian metrics g_K , the right-hand side is always non-negative. It is zero only if $\dot{\gamma}^{\mathcal{V}}$ vanishes. So the four-dimensional projection $\gamma_M(s)$ of the null path is timelike on M_4 when γ has a vertical component and is null when γ is completely horizontal. The projection of a null path on P can never be spacelike on M_4 . Thus, higher-dimensional null paths on P can describe all types of causal motion on M_4 and never correspond to acausal ones. In the name of simplicity, it is natural to investigate the consistency of a dynamical model entirely based on higher-dimensional null paths.

From (3.8), the 4D proper time of particle moving along a null curve on P satisfies the simplified relation

$$c[\tau(s_1) - \tau(s_2)] = \int_{s_1}^{s_2} \sqrt{g_K(\dot{\gamma}^{\mathcal{V}}, \dot{\gamma}^{\mathcal{V}})} ds .$$

When there are no gauge fields around, we have $\dot{\gamma}^{\mathcal{Y}} = \dot{\gamma}_K$ and proper time is just a measure of the Riemannian distance travelled by the particle in the internal space.

Regarding mass, for a null path in higher dimensions relation (6.6) reduces to

$$m(s) = \frac{1}{c} \sqrt{g_P(p^{\mathcal{Y}}, p^{\mathcal{Y}})} . \quad (10.2)$$

So the test particle's rest mass is simply the norm of its vertical momentum. Internal motion is the sole source of rest energy. Mass vanishes for particles travelling along horizontal, null paths on P , which of course also project down to null paths on M_4 . So mass vanishes if and only if the particle has speed c on Minkowski space. Rest mass will vary if there is a transfer between the horizontal and vertical components of momentum. This can happen when the geometry of P is sufficiently distorted in comparison to the vacuum geometry $g_M \times g_K$. More precisely, when $d^A g_K$ does not vanish, as described in section 7. The total momentum $p(s)$ is always covariantly conserved along a higher-dimensional geodesic.

For a null curve on P , relation (10.1) implies that the momentum components satisfy

$$-g_M(p_M, p_M) = -g_P(p^{\mathcal{H}}, p^{\mathcal{H}}) = g_P(p^{\mathcal{Y}}, p^{\mathcal{Y}}) = c^2 m^2 . \quad (10.3)$$

So the horizontal and vertical momenta have the same norm but with opposite signs. This follows from the vanishing of $g_P(p, p)$, of course. Now suppose that g_M is the Minkowski metric (2.1), choose coordinates (t, x^1, x^2, x^3) on M_4 and decompose the 4-momentum as

$$p_M = \frac{E}{c^2} \frac{\partial}{\partial t} + \mathbf{p} ,$$

as in (6.7). Then the momentum relation (10.3) can be rewritten as the 4D relation

$$E^2 = c^2 |\mathbf{p}|^2 + m^2 c^4 ,$$

or, alternatively, as the higher-dimensional relation

$$E^2 = c^2 g_P(\mathbf{p}^{\mathcal{H}} + p^{\mathcal{Y}}, \mathbf{p}^{\mathcal{H}} + p^{\mathcal{Y}}) . \quad (10.4)$$

The former is the usual 4D energy-momentum relation. The latter is similar to the energy-momentum relation of photons, but in higher dimensions. This is not surprising, because we are assuming that test particles follow null paths on P , just as photons do in 4D.

Dividing equation (10.4) by $(\sigma c \dot{\gamma}^0)^2$ and using (6.8) and (6.9), we obtain the relation for velocities in the chosen frame,

$$c^2 = g_P(\mathbf{v}^{\mathcal{H}} + v^{\mathcal{Y}}, \mathbf{v}^{\mathcal{H}} + v^{\mathcal{Y}}) = g_M(\mathbf{v}, \mathbf{v}) + g_K(v^{\mathcal{Y}}, v^{\mathcal{Y}}) . \quad (10.5)$$

Thus, for non-trivial, null geodesics on P , the norm of the velocity vector in $\mathbb{R}^3 \times K$ is always c . Test particles always travel at the speed of light in higher dimensions, as expected. The projection of velocities to three dimensions is the sole reason for the appearance of speeds in the range $[0, c]$. This generalizes the discussion in the last appendix of [46].

11 Parameterizing the space of null geodesics

Motivated by the model described in the last section, we will now have a closer look at null geodesics on P . Denote by \mathcal{N}_h the space of non-trivial, null geodesics $\gamma(s)$ passing through a point h in P at the parameter value $s = 0$. The trivial geodesic we are excluding is the one with image h for all s . Standard properties of the geodesic equation say that \mathcal{N}_h can be parameterized by the non-zero, null vectors $\dot{\gamma}(0)$ in the tangent space $T_h P$. Each of these vectors is parallelly transported along the geodesic it determines, so if $\dot{\gamma}$ is non-zero and null at $s = 0$, it will have the same properties for every value of s .

Now pick a local coordinate system (t, x^n, y^j) on $M_4 \times K$, where the x^n and y^j are the coordinates in \mathbb{R}^3 and K , respectively. Write the geodesic as

$$\gamma(s) = \left(\gamma_M(s), \gamma_K(s) \right) = \left(\gamma^0(s), \gamma^n(s), \gamma^j(s) \right).$$

Since $\dot{\gamma}(s)$ is null and non-zero, the time component $\dot{\gamma}^0(s)$ must non-zero for all s . In particular, it is always positive or always negative. So the function $s \mapsto \gamma^0(s)$ is strictly monotonous. According to the sign of $\dot{\gamma}^0$ we can divide the space of geodesics \mathcal{N}_h into its components \mathcal{N}_h^+ and \mathcal{N}_h^- , corresponding to particles moving forward and backward in time, respectively.

Each of these components is isomorphic to $\mathbb{R}^{3+k} \setminus \{0\}$, where k denotes the dimension of K . For example, thinking of \mathcal{N}_h^+ as a subset of $T_h P$, there is a bijection

$$\varphi_1 : \mathbb{R}^{3+k} \setminus \{0\} \longrightarrow \mathcal{N}_h^+ \quad (u^1, u^2, u^3, w^j) \longmapsto (u^{\mathcal{H}}, w), \quad (11.1)$$

where we have defined

$$\begin{aligned} \mathbf{u} &:= \sum_{n=1}^3 u^n \frac{\partial}{\partial x^n} & w &:= \sum_j w^j \frac{\partial}{\partial y^j} \\ u &:= \mathbf{u} + c^{-1} \sqrt{|\mathbf{u}|^2 + g_K(w, w)} \frac{\partial}{\partial t} \end{aligned} \quad (11.2)$$

and the horizontal lift $u^{\mathcal{H}}$ is taken according to (3.4). By construction, $\varphi_1(u, w)$ is always a non-zero, null vector in $T_h P$.

The bijection φ_1 parameterizes all the null geodesics starting at h and moving forward in time. These represent the motion on P of particles with different masses. Fixing the mass of the particle means restricting to a subset $\mathcal{N}_h^+(m)$ of the full space \mathcal{N}_h^+ . According to (6.6), after the simplification $g_P(p, p) = 0$ the mass of a particle at h is just $c^{-1} \sqrt{g_P(p^{\mathcal{V}}, p^{\mathcal{V}})}$. So it essentially corresponds to the norm of the vertical vector w in (11.2). Thus, under the bijection φ_1 , the space of null geodesics at h of mass m corresponds to the subset

$$\mathcal{N}_h^+(m) \simeq \left\{ (u^1, u^2, u^3, w^j) \in \mathbb{R}^{3+k} \setminus \{0\} : (g_K|_h)_{ij} w^i w^j = \sigma^{-2} c^2 m^2 \right\} \quad (11.3)$$

of the full $\mathbb{R}^{3+k} \setminus \{0\}$. Topologically, it is clear that $\mathcal{N}_h^+(m)$ is isomorphic to $\mathbb{R}^3 \times S^{k-1}$ for non-zero m and to $\mathbb{R}^3 \setminus \{0\}$ for vanishing mass.

When the internal metric g_K has isometries in the region around the point h , we can also consider the space of null geodesics representing particles with given charges, besides a given mass. For example, one can consider the subset $\mathcal{N}_h^+(q_\xi) \subset \mathcal{N}_h^+$ representing particles with electromagnetic charge q_ξ with respect to a Killing field ξ of g_K . The charge condition (8.1) imposes the constraint

$$(g_K|_h)_{ij} \xi^i w^j = -\sigma^{-1} q_\xi \quad (11.4)$$

on the coordinates w^j of $\mathbb{R}^{3+k} \setminus \{0\}$. So the subspaces $\mathcal{N}_h^+(q_\xi)$ are isomorphic to $\mathbb{R}^{2+k} \setminus \{0\}$.

Fixing mass and electromagnetic charge simultaneously defines the smaller subspaces $\mathcal{N}_h^+(m, q_\xi) = \mathcal{N}_h^+(m) \cap \mathcal{N}_h^+(q_\xi)$ inside the space of null geodesics \mathcal{N}_h^+ . Adding the mass condition in (11.3) to the charge condition (11.4), it is clear that these spaces are isomorphic to

$$\mathcal{N}_h^+(m, q_\xi) \simeq \begin{cases} \emptyset & \text{if } |q_\xi| > m c |\xi| \\ \mathbb{R}^3 & \text{if } |q_\xi| = m c |\xi| \\ \mathbb{R}^3 \times S^{k-2} & \text{if } |q_\xi| < m c |\xi|. \end{cases} \quad (11.5)$$

Here $|\xi|$ denotes the Riemannian length $\sqrt{(g_K)_h(\xi, \xi)}$ of the Killing vector field at the point h . The upper bound for the charge of a test particle at the point h , given the value of its mass, is of course the same as in formula (8.3) of section 8.

The higher-dimensional momenta of particles with the strongest possible charge, $|q_\xi| = m c |\xi|$, have fewer degrees of freedom. This happens because the vertical component of momentum must be completely aligned with ξ . Hence, the motion of a particle of maximum charge is completely determined by an initial point in P and by the 3D momentum vector at that point. For particles with weaker charges the situation is different. One also needs to choose the components of the vertical momentum at the initial point, and there is an S^{k-2} of possible choices compatible with the mass and charge of the particle.

If we fix more constants of motion of the particle, besides mass and electromagnetic charge, for example the constants described in proposition 4.3, then the space of representative null geodesics at h will be further constrained.

Celestial spheres

We will now describe a second parameterization of the space of null geodesics \mathcal{N}_h^+ . It uses physical velocities instead of momenta. Its construction relies on the fact that the time

component $\dot{\gamma}^0$ is always positive for geodesics in \mathcal{N}_h^+ . Given a null geodesic, define

$$v := \frac{1}{\dot{\gamma}^0} \dot{\gamma} \quad (11.6)$$

as a tangent vector in $T_h P$. Recalling that $\gamma^0(s)$ is just the time coordinate of the geodesic, the projection of v to the 4D tangent space $T_{\pi(h)} M_4$ can be written as

$$\pi_*(v) = \left(\frac{d\gamma^0}{ds} \right)^{-1} \sum_{n=1}^4 \frac{d\gamma^n}{ds} \frac{\partial}{\partial x^n} = \frac{\partial}{\partial t} + \sum_{n=1}^3 \frac{d\gamma^n}{dt} \frac{\partial}{\partial x^n} = \frac{\partial}{\partial t} + \mathbf{v} .$$

Here \mathbf{v} is derivative of position in \mathbb{R}^3 with respect to time, so it is the 3-dimensional velocity vector of the particle in the coordinates (t, x^1, x^2, x^3) . Decomposing v into its horizontal and vertical components, we then have

$$v = v^{\mathcal{H}} + v^{\mathcal{V}} = \left(\frac{\partial}{\partial t} + \mathbf{v} \right)^{\mathcal{H}} + v^{\mathcal{V}} .$$

The vector $\mathbf{v}^{\mathcal{H}} + v^{\mathcal{V}}$ is the derivative of position in $\mathbb{R}^3 \times K$ with respect to time. So it is the higher-dimensional velocity vector of the particle represented by $\gamma(s)$, in the coordinate system. Since $\dot{\gamma}$ and v are null vectors at h , we have $g_P(v, v) = 0$ and hence

$$\begin{aligned} c^2 &= -g_M \left(\frac{\partial}{\partial t}, \frac{\partial}{\partial t} \right) = -g_P \left(\left(\frac{\partial}{\partial t} \right)^{\mathcal{H}}, \left(\frac{\partial}{\partial t} \right)^{\mathcal{H}} \right) \\ &= g_P \left(\mathbf{v}^{\mathcal{H}} + v^{\mathcal{V}}, \mathbf{v}^{\mathcal{H}} + v^{\mathcal{V}} \right) = g_M(\mathbf{v}, \mathbf{v}) + g_K(v^{\mathcal{V}}, v^{\mathcal{V}}) . \end{aligned} \quad (11.7)$$

Here we have also used the form (2.1) of the Minkowski metric and the fact that g_P applied to horizontal vectors coincides with g_M applied to the 4D-projection of those vectors. Thus, the velocity vectors $\mathbf{v}^{\mathcal{H}} + v^{\mathcal{V}}$ associated to null geodesics at h define a Euclidean sphere S_c^{k+2} of dimension $k+2$ and radius c inside the tangent space $T_h P$. Generalizing the terminology of the 4D case, this will be called the celestial sphere at h .

It is clear from this discussion that the correspondence between geodesics in \mathcal{N}_h^+ and points in the celestial sphere of velocities is surjective but not injective. A reparameterization of the geodesic produces a constant factor in $\dot{\gamma}$ that cancels out in the quotient (11.6), so leads to the same velocity vector v . Such constant factors can be explicitly controlled by the time component $\dot{\gamma}^0$ of the derivative of $\gamma(s)$ at h . But, according to (6.9), this component is proportional to the energy of the particle in the coordinate system,

$$E = \sigma c^2 \dot{\gamma}^0 . \quad (11.8)$$

This means that the correspondence that takes a non-zero, null geodesic at h to the velocity vector plus the energy of the particle it represents,

$$\varphi_2 : \mathcal{N}_h^+ \longrightarrow S_c^{k+2} \times \mathbb{R}^+ \quad \dot{\gamma}_h \longmapsto \left(\mathbf{v}^{\mathcal{H}} + v^{\mathcal{V}}, E \right)$$

defines a second bijection that can be used to parameterize \mathcal{N}_h^+ . Of course, topologically, $S_c^{k+2} \times \mathbb{R}^+$ is isomorphic to the parameter space $\mathbb{R}^{3+k} \setminus \{0\}$ used in bijection (11.1). The virtue of the second parameterization is that it is expressed in terms of clear physical quantities, namely the velocity and energy of the particle represented by the null geodesic.

The next question is how the subsets of geodesics with fixed mass, previously denoted by $\mathcal{N}_h^+(m)$, look like under the parameterization φ_2 . In other words, how are they carved out from the full parameter space $S_c^{k+2} \times \mathbb{R}^+$. Using definition (6.6) for the mass of the particle associated to a geodesic, on the one hand, and using relation (6.9) for the energy associated to the particle in the frame, on the other hand, it is not difficult to recognize that under the parameterization φ_2 we have:

$$\mathcal{N}_h^+(m) \simeq \left\{ (\mathbf{v}^{\mathcal{H}} + v^{\mathcal{Y}}, E) \in S_c^{k+2} \times \mathbb{R}^+ : E^2 g_K(v^{\mathcal{Y}}, v^{\mathcal{Y}}) = m^2 c^6 \right\}. \quad (11.9)$$

Thus, when $m = 0$, the mass condition is satisfied if and only if $v^{\mathcal{Y}}$ vanishes. When $m > 0$, the mass condition only allows points in the celestial sphere with non-zero $v^{\mathcal{Y}}$, and then the component E in \mathbb{R}^+ is fully determined by the norm of $v^{\mathcal{Y}}$. So for $m > 0$ the space $\mathcal{N}_h^+(m)$ is parameterized by the sphere S_c^{k+2} after the exclusion of an S^2 corresponding to the points with vanishing $v^{\mathcal{Y}}$. The excluded points correspond to the 4D motions at the speed of light, of course.

Combining (11.6), (11.8) and the mass condition in (11.9), one can check that the inverse of the parameterization φ_2 satisfies

$$\begin{aligned} \varphi_2^{-1} : S_c^{k+2} \setminus S^2 &\longrightarrow \mathcal{N}_h^+(m) & (11.10) \\ \mathbf{v}^{\mathcal{H}} + v^{\mathcal{Y}} &\longmapsto \dot{\gamma}(0) = \frac{m c}{\sigma \sqrt{g_K(v^{\mathcal{Y}}, v^{\mathcal{Y}})}} \left[\left(\frac{\partial}{\partial t} + \mathbf{v} \right)^{\mathcal{H}} + v^{\mathcal{Y}} \right]. \end{aligned}$$

Finally, it is relevant to mention that the parameterizations φ_1 and φ_2 of \mathcal{N}_h^+ are not canonical, in the sense that they both rely on a choice of coordinates on Minkowski space.

Acknowledgements

It is a pleasure to thank Nick Manton and Nuno Romão for helpful comments on an earlier version of this paper.

Appendices

A Auxiliary results

Classical results of Ehresmann [8] and Hermann [9] imply that:

Proposition A.1. *Let $\pi : (P, g_P) \rightarrow (M, g_M)$ be a Riemannian submersion. If the metric g_P is complete, then π defines a locally trivial fibration. In this case, given a path $\gamma_M(s)$ on M starting at a point x and given a point p in P such that $\pi(p) = x$, there exist a unique path $\gamma(s)$ on P starting at p such that $\gamma_M = \pi \circ \gamma$ and the derivatives $\dot{\gamma}$ are always horizontal vectors on P . This γ is called a horizontal lift of γ_M .*

Regarding horizontal geodesics, O'Neill has the following result:

Proposition A.2 ([11]). *Let $\pi : (P, g_P) \rightarrow (M, g_M)$ be a Riemannian submersion and let $\gamma(s)$ be a geodesic on P . If the derivative $\dot{\gamma}$ is a horizontal vector at some point, then it is always horizontal. Moreover, in this case the projection $\gamma_M(s) = \pi \circ \gamma(s)$ is a geodesic on (M, g_M) .*

Using basic properties of Riemannian submersions (e.g. see [12, sec. 9]), we show that:

Lemma A.3. *Let $\pi : P \rightarrow M$ be a Riemannian submersion with metric g_P determined by the equivalent data (g_M, g_K, A) . Let U and V be vertical vector fields and let Z be a basic vector field on P . Then*

$$\begin{aligned} (\mathcal{L}_V g_P)(U, U) &= (\mathcal{L}_V g_K)(U, U) & (\mathcal{L}_V g_P)(Z, Z) &= 0 \\ (\mathcal{L}_V g_P)(Z, U) &= g_K([Z, V], U) . \end{aligned} \tag{A.1}$$

In particular, V is Killing with respect to g_P if and only if the restriction of V to each fibre is Killing with respect to g_K and, additionally, the Lie bracket $[V, Z]$ vanishes for every basic vector field Z on P .

Proof. The horizontal basic field Z is π -related to a well-defined vector field $\pi_* Z$ on the base M . Since V is vertical, we have $\pi_* [V, Z] = [\pi_* V, \pi_* Z] = 0$, and the bracket $[V, Z]$ is also vertical. So by definition of Lie derivative:

$$(\mathcal{L}_V g_P)(Z, Z) = \mathcal{L}_V(g_P(Z, Z)) - 2g_P([V, Z], Z) = \mathcal{L}_V(g_M(\pi_* Z, \pi_* Z)) = 0 .$$

Using that U and $[V, U]$ are both vertical, so are both orthogonal to Z , we have

$$(\mathcal{L}_V g_P)(Z, U) = \mathcal{L}_V(g_P(Z, U)) - g_P([V, Z], U) - g_P(Z, [V, U]) = g_P([Z, V], U) .$$

Finally, since U, V and $[U, V]$ are all vertical and g_K just denotes the restriction of g_P to the fibres, it is clear that

$$\begin{aligned} (\mathcal{L}_V g_P)(U, U) &= \mathcal{L}_V(g_P(U, U)) - 2g_P([V, U], U) \\ &= \mathcal{L}_V(g_K(U, U)) - 2g_K([V, U], U) = (\mathcal{L}_V g_K)(U, U) . \end{aligned} \tag{A.2}$$

Now suppose that $\mathcal{L}_V g_P$ vanishes. Then by (A.1) we get that $(\mathcal{L}_V g_K)(U, U)$ vanishes for all U . Since $\mathcal{L}_V g_K$ is symmetric in both entries, it must also vanish. Moreover, also by (A.1), the vanishing of $(\mathcal{L}_V g_P)(Z, U)$ for all U and the non-degeneracy of g_K imply that $[Z, V]$ vanishes for all basic Z . This confirms the ‘‘only if’’ part of the last statement, while the converse is clear. \square

The following observation, stated in [12, sec. 9F], is a consequence of results in [9, 10].

Proposition A.4. *Let the metric $g_P \simeq (g_M, A, g_K)$ define a Riemannian submersion on P with totally geodesic fibres, i.e. with $d^A g_K = 0$. Then the curvature form $F_A(X, Y)$ has values in the Killing vector fields of g_K for every $X, Y \in TM$.*

Remark. We should also mention that the notation used in this paper differs significantly from the conventional notation in the literature about Riemannian submersions. This is because the latter clashes with the traditional notation in physics. So the tensor called A in [10–13] is essentially what we call here F_A , since it represents the physical gauge fields strength. The tensor called T in [10–13] is what we call here $d^A g_K$ (or S less often). This avoids confusion with torsion or with the energy-momentum tensor. It also emphasizes its physical role as a covariant derivative of Higgs-like fields.

In section 4 we use the following standard result.

Lemma A.5. *Let $\{u_j\}$ be a basis of the space of Killing vector fields on the compact manifold (K, g_K) . Define the structure constants with respect to this basis by the relations $[u_i, u_j] = f_{ij}^l u_l$. If the basis is orthonormal with respect to the L^2 -inner product on (K, g_K) , then the f_{ij}^l are totally anti-symmetric in their three indices.*

Proof. The Lie derivatives of the metric g_K and volume form vol_{g_K} both vanish along Killing vector fields. Using this fact, the definition of the L^2 -inner product of vector fields and Stokes’ theorem, we can write

$$\begin{aligned} \sum_l f_{ij}^l \langle u_l, u_k \rangle_{L^2} &= \langle [u_i, u_j], u_k \rangle_{L^2} = \int_K g_K([u_i, u_j], u_k) \text{vol}_{g_K} \\ &= \int_K \left\{ \mathcal{L}_{u_i} [g_K(u_j, u_k)] - g_K(u_j, [u_i, u_k]) \right\} \text{vol}_{g_K} \\ &= \int_K \mathcal{L}_{u_i} [g_K(u_j, u_k) \text{vol}_{g_K}] - \int_K g_K(u_j, [u_i, u_k]) \text{vol}_{g_K} \\ &= - \int_K g_K(u_j, [u_i, u_k]) \text{vol}_{g_K} = - \sum_l f_{ik}^l \langle u_j, u_l \rangle_{L^2} . \end{aligned}$$

Thus, when the basis $\{u_j\}$ is L^2 -orthonormal, we get that

$$\sum_l f_{ij}^l \delta_{lk} + f_{ik}^l \delta_{jl} = 0 \quad \iff \quad f_{ij}^k = -f_{ik}^j .$$

The anti-symmetry of f_{ij}^l in the lower two indices follows from its definition and is true for any basis. \square

B Parallel transport in Riemannian submersions

For reference, in this appendix we write down the equations of parallel transport of vectors along general curves $\gamma(s)$ on the higher-dimensional space $P = M \times K$. That space is assumed to be equipped with a submersive metric $g_P \simeq (g_M, A, g_K)$. Parallel transport is determined by its Levi-Civita connection ∇ . As in section 3, a vector E in TP can be decomposed into horizontal and vertical components, $E = E^{\mathcal{H}} + E^{\mathcal{V}}$, defined through (3.4). The projection of the curve $\gamma(s)$ to the base M is denoted by $\gamma_M(s)$, so that $\gamma_M = \pi \circ \gamma$. The Levi-Civita connection on M determined by the projected metric g_M is denoted by ∇^M .

With this notation, the equations in [11] for the horizontal and vertical components of the covariant derivative $\nabla_{\dot{\gamma}} E$ along $\gamma(s)$ can be written as:

$$\begin{aligned} g_P(\nabla_{\dot{\gamma}} E, V) &= g_P(\nabla_{\dot{\gamma}} E^{\mathcal{V}}, V) + \frac{1}{2} F_A^a(\dot{\gamma}_M, E_M) g_P(e_a, V) - g_P(S_V \dot{\gamma}^{\mathcal{V}}, E^{\mathcal{H}}) \\ g_P(\nabla_{\dot{\gamma}} E, Z) &= g_M(\nabla_{\dot{\gamma}_M}^M E_M, \pi_* Z) + \frac{1}{2} F_A^a(\pi_* Z, \dot{\gamma}_M) g_P(e_a, E^{\mathcal{V}}) \\ &\quad + \frac{1}{2} F_A^a(\pi_* Z, E_M) g_P(e_a, \dot{\gamma}) + g_P(S_{\dot{\gamma}^{\mathcal{V}}} E^{\mathcal{V}}, Z) \end{aligned} \quad (\text{B.1})$$

Here Z and V are any horizontal and vertical vectors in TP , respectively. We have also translated O'Neill's notation to the one used in this paper, as described in the remark of appendix A. As in section 5, the tensor S is the second fundamental form of the fibres of P . It essentially coincides with our $d^A g_K$, as expressed in (5.5).

The general equations B.1 reduce to (5.1) when we take $E = \dot{\gamma}$. The equations satisfied by a parallelly transported vector field $E_{\gamma(s)}$ can be obtained from B.1 by setting $\nabla_{\dot{\gamma}} E = 0$ on the left-hand sides.

References

- [1] D. Kovacs: The geodesic equation in five-dimensional relativity theory of Kaluza-Klein, *Gen. Relat. Gravit.* **16** (1984), 645–655.
- [2] J. Gegenberg and G. Kunstatter: The motion of charged particles in Kaluza-Klein space-time, *Phys. Letters* **106A** (1984), 410–414.
- [3] Y. Cho and D. Park: Higher-dimensional unification and fifth force, *Il Nuovo Cimento*, **105 B** (1990), 817–829.
- [4] R. Coquereaux and G. Esposito-Farese: The theory of Kaluza-Klein-Jordan-Thiry revisited, *Ann. Inst. H. Poincaré* **52** (1990), 113–150.
- [5] H. Liu and B. Mashhoon: Spacetime measurements in Kaluza-Klein gravity, *Phys. Letters* **272A** (2000), 26–31.
- [6] S. Seahra and P. Wesson: Null geodesics in five-dimensional manifolds, *Gen. Relat. Gravit.*, **33** (2001), 1731–1752.
- [7] E. Minguzzi: Proper time and conformal problem in Kaluza-Klein theory, *Int. J. Geom. Meth. Mod. Phys.* **12** (2015), 1550063.
- [8] C. Ehresmann: Les connexions infinitésimales dans un espace fibré différentiable, *Colloque de Topologie* (1950), 29–55.
- [9] R. Hermann: A sufficient condition that a mapping of Riemannian manifolds be a fibre bundle, *Proc. Amer. Math. Soc.* **11** (1960), 236–242.
- [10] B. O’Neill: The fundamental equations of a submersion, *Michigan Math. J.* **13** (1966), 459–469.
- [11] B. O’Neill: Submersions and geodesics, *Duke Math. J.* **34** (1967), 363–373.
- [12] A. Besse: *Einstein manifolds*, Classics in Mathematics, Springer-Verlag, 1987.
- [13] M. Falcitelli, S. Ianus and A. Pastore: *Riemannian submersions and related topics*, World Scientific Publishing, 2004.
- [14] D. Bailin and A. Love: Kaluza-Klein theories, *Rep. Prog. Phys.* **50** (1987), 1087–1170.
- [15] D. Bleecker: *Gauge theory and variational principles*, Addison-Wesley, 1981.
- [16] J. Bourguignon: A mathematician’s visit to Kaluza-Klein theory, *Rend. Sem. Mat. Univ. Politec. Torino* (1989), 143–163.

- [17] L. Castellani, P. Fré and R. D’Auria: *Supergravity and superstrings: a geometric perspective*, Vol. 2, Part five, World Scientific Publishing, 1991.
- [18] R. Coquereaux and A. Jadczyk: *Riemannian geometry, fiber bundles, Kaluza-Klein theories and all that....*, World Scientific Publishing, 1988.
- [19] M. Duff, B. Nilsson and C. Pope: Kaluza-Klein supergravity, *Phys. Reports* **130** (1986), 1–142.
- [20] J. Overduin and P. Wesson: Kaluza-Klein gravity, *Phys. Reports* **283** (1997), 303–380.
- [21] E. Witten: Search for a realistic Kaluza-Klein theory, *Nucl. Phys.* **B186** (1981), 412–428.
- [22] T. Kaluza: Zum Unittäsproblem in der Physik, *Sitzungsber. Preuss. Akad. Wiss. Berlin Math. Phys.* **K1** (1921), 966–972.
- [23] O. Klein: Quantentheorie und fünfdimensionale Relativitätstheorie, *Zeitschrift Phys.* **37** (1926), 895–906.
- [24] A. Einstein and P. Bergmann: On a generalization of Kaluza’s theory of electricity, *Annals Math.* **39** (1938), 683–701.
- [25] P. Jordan: Relativistische Gravitationstheorie mit variabler Gravitationskonstante, *Naturwissenschaften* **33** (1946), 250–251.
- [26] Y. Thiry: Les équations de la théorie unitaire de Kaluza, *Comptes Rendus Acad. Sci. Paris* **226** (1948), 216–218.
- [27] B. DeWitt: Dynamical theory of groups and fields, in *Lectures at 1963 Les Houches School*, Gordon and Breach, 1964, 585–820.
- [28] R. Kerner: Generalization of the Kaluza-Klein theory for an arbitrary non-abelian gauge group, *Ann. Inst. H. Poincaré* **9** (1968), 143–152.
- [29] A. Trautman: Fiber bundles associated with space-time, *Rep. Math. Phys.* **1** (1970), 29–62.
- [30] Y. Cho: Higher-dimensional unifications of gravitation and gauge theories, *J. Math. Phys.* **16** (1975), 2029–2035.
- [31] Y. Cho and P. Freund: Non-Abelian gauge fields as Nambu-Goldstone fields, *Phys. Rev.* **D12** (1975) 1711–1720.

- [32] E. Leibowitz and N. Rosen: Five-dimensional relativity theory, *Gen. Relat. Gravit.* **4** (1973), 449–474.
- [33] L. Fehér and P. Horvathy: Dynamical symmetry of the Kaluza-Klein monopole, in *Proc. Symmetries in Science III*, Plenum, 1989, 399–417.
- [34] P. Wesson and J. Leon: The equations of motion in Kaluza-Klein cosmology and its implications for astrophysics, *Astron. Astrophys.* **294** (1995), 1–7.
- [35] V. Lacquaniti, G. Montani and D. Pugliese: Massive test particle motion in 5-dimensional electromagnetic-free Kaluza-Klein theory, *Gen. Relat. Gravit.* **43** (2011), 1103–1120.
- [36] J. Baptista: Internal symmetries in Kaluza-Klein models, *J. High Energ. Phys.* **2024** (2024), 178.
- [37] J. Baptista: Rest mass variation in Kaluza-Klein models, [arXiv:2406.09503v1](https://arxiv.org/abs/2406.09503) [[hep-th](https://arxiv.org/abs/2406.09503v1)].
- [38] S. Wong: Field and particle equations for the classical Yang-Mills field and particles with isotopic spin, *Nuovo Cimento* **65** (1970), 689–694.
- [39] P. Horvathy and P. Zhang: Kerner equation for motion in a non-abelian gauge field, *Universe* **9** (2023), 519.
- [40] E. Witten: Fermion quantum numbers in Kaluza-Klein theory, in *Shelter Island II, Proceeding of the 1983 Shelter Island conference*, MIT Press, 1985, 227–277.
- [41] R. Wald: *General relativity*, Chicago Univ. Press, 1984.
- [42] J. Milnor: Curvature of left invariant metrics on Lie groups, *Adv. Math.* **21** (1976), 293–329.
- [43] R. Kerner, J. Martin, S. Mignemi and J. Holten: Geodesic deviation in Kaluza-Klein theories, *Phys. Rev.* **D63** (2000), 027502.
- [44] V. Faraoni, E. Gunzig and P. Nardone: Conformal transformations in classical gravitational theories and in cosmology, *Fund. Cosmic Phys.* **20** (1999), 121–175.
- [45] Y. Cho: Unified cosmology, *Phys. Rev.* **D41** (1990) 2462–2471.
- [46] J. Baptista: Higher-dimensional routes to the Standard Model fermions, [arXiv:2105.02901](https://arxiv.org/abs/2105.02901) [[hep-th](https://arxiv.org/abs/2105.02901)].