MULTIGRADED STILLMAN'S CONJECTURE

JOHN COBB, NATHANIEL GALLUP, AND JOHN SPOERL

ABSTRACT. We resolve Stillman's conjecture for families of polynomial rings that are graded by any abelian group under mild conditions. Conversely, we show that these conditions are necessary for the existence of a Stillman bound. This has applications even for the wellknown standard graded case.

1. INTRODUCTION

Let k be a field and $S = k[x_1, \ldots, x_n]$ be the *n*-variable polynomial ring. Famously, the Hilbert syzygy theorem says that any finitely generated S-module has projective dimension less than n. In 2000, Mike Stillman conjectured that the projective dimension of a homogeneous ideal in a polynomial ring can be bounded just in terms of the number and degrees of the generators provided that S is given the standard grading [PS09]. Importantly, this bound is independent of the number of variables of S. This conjecture was proven by Ananyan and Hochster [AH16, Theorem C] in 2016, and subsequently reproven by Erman, Sam, and Snowden [ESS19] and Draisma, Lasoń, and Leykin [DLL19].

The property that the projective dimension of finitely generated ideals are bounded only in terms of the number and degrees of their generators is sometimes called being *Stillman bounded* and is part of a larger program of similar phenomena of Stillman uniformity [ESS18]. For instance, Caviglia proved that a Stillman bound on projective dimension is equivalent to one on regularity [Pee10, Theorem 29.5] which links to work on the Eisenbud–Goto conjecture [BMNB⁺11, MP18]. More generally, "projective dimension" can be swapped out with a host of other ideal invariants [ESS21]. Due mostly to work in toric geometry, there has been great interest in understanding analogs of such results for gradings by other abelian groups [HS02, MS04, HSS06, CMR06, Hà07, LS12, BES20, Yan19, CN20, BS22, CH22, BHS22, BPC22, Cob24]. Since Stillman uniformity requires finding a bound independent of the number of variables, it is inherently a property of a *family* of graded polynomial rings (e.g. all standard graded polynomial rings) and not of any particular fixed *S*. Ananyan and Hochster established that the family of \mathbb{Z}_+ -graded polynomial rings has Stillman bounded projective dimension [AH16, page 12] for possibly inhomogeneous ideals. It is our goal to understand which graded families have Stillman uniformity.

The following example demonstrates that allowing the grading group to have infinite descending chains (i.e. the divisibility order is not well-founded) permits counterexamples to a multigraded Stillman bound on projective dimension.

Example 1.1. Let $S_n = k[x, y, z_1, ..., z_n]$ be a \mathbb{Q}_+ -graded polynomial ring with grading given by $\deg(x) = \deg(y) = \deg(z_i) = 1/n$. The homogeneous ideal

$$I_n = \left\langle x^n, y^n, x^{n-1}z_1 + x^{n-2}yz_2 + \dots + xy^{n-2}z_{n-1} + y^{n-1}z_n \right\rangle$$

is generated by 3 degree 1 elements. It is shown in [McC11] that the projective dimension of S_n/I_n is n + 2. Therefore the family $\{S_n\}$ of \mathbb{Q}_+ -graded polynomial rings cannot have a Stillman bound since we can make n arbitrarily large while I_n is generated in degree 1. \diamond On the other hand, there is an obvious condition on the grading which is sufficient for the existence of a Stillman bound. If there is a sufficiently nice "flattening" map from the grading group to \mathbb{Z} , we can regrade our family by \mathbb{Z}_+ and apply Ananyan and Hochster's result, since changing the grading does not change projective dimension. However, not all grading groups admit such a map (see Example 4.2). Our main result gives a condition on the grading group that is weaker than having a flattening map and stronger than well-foundedness which is equivalent to the existence of a Stillman bound for any graded family. To be more precise, we need a few definitions.

A grading of a polynomial ring S = k[X] over a (possibly infinite) set of variables X by a (possibly infinitely generated) abelian group Γ is a decomposition of S into k-submodules

$$S = \bigoplus_{g \in \Gamma} S_g, \quad \text{with } S_g \cdot S_h \subset S_{g+h}$$

such that for all variables $x \in X$, $x \in S_g$ for some $g \in \Gamma$. If $s \in S_g$ we say that s is homogeneous of degree g and write deg(s) = g. We say that S is connected if $S_0 = k$. The (grading) support of S is the submonoid of Γ generated by the degrees of all the monomials along with the identity. We say a (possibly inhomogeneous) ideal has degree sequence bounded by $\mathbf{d} = (d_1, \ldots, d_n) \in \Gamma^n$ if the ideal can be generated by elements f_1, \ldots, f_n whose monomials have degree less than d_1, \ldots, d_n . Any submonoid $\Lambda \subseteq \Gamma$ has bounded factorization if it is impossible to express an element in Λ as an arbitrarily large sum of other elements in Λ . Our main theorem shows that for a fixed $\Lambda \subseteq \Gamma$, any family of connected Γ -graded polynomial rings with support contained in Λ has Stillman bounded projective dimension if and only if Λ has bounded factorization.

Theorem 1.2. For any degree sequence d from Λ , there is a number $N(\Lambda, d)$ depending only on Λ and d bounding the projective dimension of any ideal with degree sequence bounded by d in any connected Γ -graded polynomial ring with support contained in Λ if and only if Λ has bounded factorization.

After fixing any particular polynomial ring S and ideal I, the projective dimension of S/I does not change as one varies the grading. Hilbert's syzygy theorem guarantees that the projective dimension of *every* ideal in S is bounded by the (hopefully finite) number of variables in S. The classical Stillman's conjecture ensures a bound shared across the entire family of standard graded polynomial rings at the cost of restricting the desired class of ideals I to those with a particular degree sequence. Theorem 1.2 asserts that if you also allow your grading to vary, under mild conditions you must only further fix Λ for Stillman uniformity. Example 1.1 fails to have Stillman bounded projective dimension exactly because the supports of the family fail to lie in a bounded factorization monoid. Letting deg(x) = deg(y) = 0 and deg $(z_i) = d$ in the same example shows that non-connected gradings do not have a Stillman bound. We emphasize the fact that the bound in Theorem 1.2 works over families of different gradings with the following example.

Example 1.3. Let $\Gamma = \mathbb{Z} \times \mathbb{Z}$ and let $\Lambda = (\mathbb{Z} \times \mathbb{Z}_{>0}) \cup \{(0,0)\}$. Then despite being infinitely generated, Λ has bounded factorization.¹ Now for any n, let $S_n = k[x_1, \ldots, x_n, y_1, \ldots, y_n]$ and give S_n a Γ -grading by setting deg $(x_n) = (-n, 1)$ and deg $(y_n) = (n, 1)$. Theorem 1.2

¹This is true since we have a height function $\ell : \Lambda \to \mathbb{Z}_{\geq 0}$ defined by $\ell(n,m) = m$, meaning ℓ satisfies $\ell(g) = 0$ iff g = 0 and $\ell(g+h) \ge \ell(g) + \ell(h)$ for all $g, h \in \Lambda$. A pointed monoid has bounded factorization if and only if it has such a height function [GV23].

3

says that the family $\{S_n\}$ has Stillman uniformity because for all n, the support of S_n is contained in Λ which has bounded factorization. In fact we could have assigned the degrees of x_n and y_n to be anything inside of Λ and the same Stillman bounds would hold. \diamond

After finding the right condition on Λ , the proof of Theorem 1.2 ends up being very short. The key intuition is to leverage the known Stillman bound for the \mathbb{Z}_+ -graded case by using bounded factorization to guarantee a nice regrading by \mathbb{Z}_+ for the "if" direction, and to construct an explicit counterexample using only non-bounded factorization for the "only if" direction. Even in the $\Lambda = \mathbb{Z}_+$ case proven by Ananyan and Hochster [AH16], Theorem 1.2 says something new; the same Stillman bound holds for families of varying \mathbb{Z}_+ -gradings, with the bounds potentially getting tighter for polynomial rings whose support refines \mathbb{Z}_+ . That is, by equipping a ring with a finer-graded structure, we can get better bounds for a given ideal, as shown in the following example.

Example 1.4. Let $S = k[x_1, x_2, ...]$ be an infinite polynomial ring, and suppose we want to bound the projective dimension of a particular ideal I generated by $f = x_1x_4x_7 + x_{10}x_{13}x_{16}$, $g = x_2x_5x_8 + x_{11}x_{14}x_{17}$, and $h = x_3x_6x_9 + x_{12}x_{15}x_{18}$. Of course, we get the bound of 18 from Hilbert's syzygy theorem since f, g, and h only involve 18 variables, but we can do better. Under the standard grading, $\langle f, g, h \rangle$ has the degree sequence d = (3, 3, 3) whose tight Stillman bound has been computed to be 5 [MM19]. Consider the \mathbb{Z}^3 -grading on S by

$$\deg(x_i) = \begin{cases} (1,0,0) & \text{if } i = 1 \mod 3, \\ (0,1,0) & \text{if } i = 2 \mod 3, \\ (0,0,1) & \text{if } i = 0 \mod 3. \end{cases}$$

Under this new grading, the degree sequence of $\langle f, g, h \rangle$ is $\boldsymbol{e} = \{(3, 0, 0), (0, 3, 0), (0, 0, 3)\}$. This grading is *finer* in the sense that the family of ideals that are degree \boldsymbol{e} is strictly contained in the larger family of ideals that are degree $\varphi(\boldsymbol{e}) = \boldsymbol{d}$, where φ is the natural flattening map summing the entries. Therefore one would expect that $N(\boldsymbol{e}) < N(\boldsymbol{d})$ and in this case, this is true; the Stillman bound under this grading is exactly the projective dimension of 3.

In Example 1.4, the \mathbb{Z}^3 -grading reveals that the variables showing up among f, g, and h are disjoint and therefore f, g, h form a regular sequence. In general, if the bound in Theorem 1.2 were made effective (e.g. as in the standard graded case [KZ20]), one could find better Stillman bounds for a fixed polynomial ring S and ideal I by varying the grading. That is, given a fixed \mathbb{Z}_+ -graded S and ideal with degree sequence d, you may consider equipping S with various positive Γ -gradings with flattening maps $\varphi : \Lambda \to \mathbb{Z}_+$ to obtain the bound

(1)
$$\operatorname{pdim}_{k}(S/I) \leq \min_{\Lambda} \{ N(\varphi(\Lambda), \varphi(\boldsymbol{e})) \mid \Lambda \text{ grades } S \text{ s.t. } \varphi(\boldsymbol{e}) = \boldsymbol{d} \}$$

which Example 1.4 shows can be tight.

Open problems. Here are some open problems raised by our work:

- Are there similar bounds for a larger class of ideal invariants such as multigraded regularity as shown in [ESS21]?
- When is the bound in Equation (1) tight?

Acknowledgements. Thanks to Daniel Erman, Jason McCullough, Christopher O'Neill, and Ivan Aidun for invaluable discussions. Cobb acknowledges the support of the National Science Foundation Grant DMS-2402199.

2. Γ -graded Commutative Algebra

Let Γ be an abelian group, k a field, and S a Γ -graded polynomial ring with graded support Λ . The graded structure (and thus many of the algebraic properties) of S is entirely determined by Λ . In fact, we could eliminate mentions of Γ as long as we assume that Λ is commutative and *cancellative*, which means that g + h = g + h' implies h = h' for all $g, h, h' \in \Lambda$. This comes for free when Λ is a submonoid of a group, otherwise, it allows us to construct a unique group (its Grothendieck group) containing Λ by adding formal inverses. We will nearly always assume that Λ is *pointed*, which means that q + q' = 0implies q = q' = 0 for $q, q' \in \Lambda$. Note that Λ must be torsion-free in order to be pointed. If Λ is pointed and cancellative, it has a natural partial order \leq_{Λ} :

 $g \leq_{\Lambda} h \iff g + q = h$ for some $q \in \Lambda$.

This provides a notion of positivity in S by letting $S_+ = \{s \in S \mid \deg(s) >_{\Lambda} 0\}$ be the *positive degree* elements of S. By assuming that Λ is commutative and cancellative we can work directly with Λ -graded rings without mention of Γ . The following lemma shows that for Γ -graded polynomial rings, connected implies pointed:

Lemma 2.1. If S is a connected Λ -graded polynomial ring then the support of S is pointed.

Proof. Let S = k[X] and suppose that $g, h \in \Lambda$ are such that g + h = 0. Then by definition of Λ there exists $r, s \in S \setminus \{0\}$ such that $\deg(r) = g$ and $\deg(s) = h$. Then $\deg(rs) = \deg(r) + \deg(s) = g + h = 0$, so by connectedness, we have that $rs \in k$. Since S is a polynomial ring over k, we have $r \in k$ and $s \in k$, hence $g = \deg(r) = 0$ and $h = \deg(s) = 0$. \Box

We say that Λ (or S):

- is well-founded if there are no infinite decreasing chains in Λ under the relation \leq_{Λ} ,
- has bounded factorization if for all $g \in \Lambda$ there exists some N such that for every expression $g = g_1 + \ldots + g_s$ for $g_1, \ldots, g_s \in \Lambda \setminus \{0\}$, we have $s \leq N$,

If Λ is pointed, having bounded factorization allows a well-defined height function that assigns any element d in Λ to a natural number |d| satisfying a triangle inequality (see [GV23]). Although bounded factorization implies well-foundedness, the following example shows that there exist well-founded monoids which do not have bounded factorization.

Example 2.2 (Well-founded but no bounded factorization). Let Λ be the submonoid of $\mathbb{Q}_{\geq 0}$ generated by $\{\frac{1}{p} \mid p \in \mathbb{N} \text{ is prime}\}$. Λ is pointed since it is contained in the pointed monoid $\mathbb{Q}_{\geq 0}$, but it does not have bounded factorization since 1 can be written as a sum of p copies of 1/p for all primes p. However, we claim that Λ is well-founded. Since all elements in Λ are built by adding up reciprocals of primes, any element h in Λ can be written uniquely as $h_1 + \sum_{k=2}^{\ell} \frac{h_k}{p_k}$, where $0 < h_k < p_k$. We wish to show that any decreasing path from h is finite. Suppose we start such a path, $h \geq_{\Lambda} g$. Then there must exist some $g' \in \Lambda$ with g + g' = h. Writing each of these in the form mentioned above, it must be that $g_k + g'_k = h_k$ and thus that $g_k \leq h_k$. So any chain of elements strictly descending from h can have at most $h_1 + \ldots + h_\ell$ elements.

MULTIGRADED STILLMAN'S CONJECTURE

3. Main Results and Corollaries

In this section, Λ will always be commutative and cancellative. Consider a monomial of degree $d \in \Lambda$ in a connected Λ -graded polynomial ring k[X]. If we completely forget the Λ -grading and give k[X] the standard grading (i.e. "flatten" it), what degree will f be? The following theorem says that having bounded factorization is exactly what is needed to bound the new degree of f in the standard grading.

Lemma 3.1. Suppose k[X] is a connected Λ -graded polynomial ring with bounded factorization. For all $g \in \Lambda$, there exists $N \in \mathbb{N}$ such that if $x_1^{e_1} \dots x_m^{e_m}$ has degree $\leq_{\Lambda} g$ (for $x_i \in X$) then $e_1 + \dots + e_m \leq N$.

Proof. By definition of having bounded factorization, there exists $N \in \mathbb{N}$ such that if $g_1 + \ldots + g_s = g$ is a non-trivial factorization of g (i.e. $g_i \neq 0$) then $s \leq N$. If $\deg(x_1^{e_1} \ldots x_m^{e_m}) \leq_{\Lambda} g$ then there exists some $h \in \Lambda$ with $e_1 \deg(x_1) + \ldots + e_m \deg(x_m) + h = g$. Since $\deg(x_j) >_{\Lambda} 0$ by connectedness, it must be that $e_1 + \ldots + e_m \leq N$ as desired. \Box

The forward direction of Theorem 1.2 comes from constructing a specific counterexample only from the knowledge that Λ is not a bounded factorization monoid. Consider the following example, due to Burch in the local case [Bur68], Kohn in the global case [Koh72], and translated to the language of polynomial rings by McCullough and Seceleanu [MS12].

Example 3.2. Let $S = k[x_1, \ldots, x_n, y_1, \ldots, y_n]$, and let I be the ideal generated by the three elements $f_1 = \prod_{i=1}^n x_i$, $f_2 = \prod_{i=1}^n y_i$, and $f_3 = \sum_{i=1}^n \prod_{j \neq i} x_j y_j$. Then $\text{pdim}_S(S/I) = n+2$.

Proof of Theorem 1.2. For the forward direction, suppose that Λ does not have bounded factorization. This means there exists some $d \in \Lambda$ such that for every $b \in \mathbb{N}$ we can find another number B larger than b that is the length of a factorization $d = d_1 + \cdots + d_B$, where d_1, \ldots, d_B are nonzero degrees in Λ . Grouping terms arbitrarily, we can take B = b. For every b we can define a polynomial ring $S_b = k[x_1, \ldots, x_b, y_1, \ldots, y_b]$ with Λ -grading defined by setting deg $(x_i) = deg(y_i) = d_i$. Then with f_1, f_2, f_3 defined as in Example 3.2, we have that f_1 and f_2 are homogeneous of degree d, while f_3 is inhomogeneous but with the degree of each monomial bounded by 2d. Letting $I_b = \langle f_1, f_2, f_3 \rangle$, Example 3.2 confirms that pdim_{St} $(S_b/I_b) = b + 2$ can grow arbitrarily big, so there cannot be Stillman bounded.

For the backward direction, let S be any connected Λ -graded polynomial ring with bounded factorization and let $I = \langle f_1, \ldots, f_n \rangle$ be an ideal with degree sequence bounded by d. By Lemma 3.1, we can find a number B bounding the standard degree of all monomials appearing among f_1, \ldots, f_n after giving S the standard grading. Now, [AH16, Theorem C] provides a Stillman bound for I.

Corollary 3.3 shows that any connected polynomial ring with support $\Lambda' \subset \Lambda$ has the same Stillman bound, i.e. $N(\Lambda', d) \leq N(\Lambda, d)$.

Corollary 3.3. If S' and S have nested support $\Lambda' \subset \Lambda$ then $N(\Lambda', d) \leq N(\Lambda, d)$.

Proof. Given such an S with support $\Lambda' \subseteq \Lambda$ and, let $Y = \{y_g \mid g \in \Lambda \smallsetminus \Lambda'\}$ be a set of variables in bijection with $\Lambda \smallsetminus \Lambda'$. We give $T := k[X \sqcup Y]$ a Λ -grading by setting $\deg_T(x) = \deg_S(x)$ for all $x \in X$ and $\deg_T(y_g) = g$. Then T is a connected Λ -graded polynomial ring with S as a subring, hence the grading has bounded factorization and has the bound $N(\Lambda, \mathbf{d})$ as in Theorem 1.2.

4. Applications and Examples

If S is a connected Γ -graded polynomial ring, then the following example shows how a choice of Λ is equivalent to a choice of an effective cone in toric geometry.

Example 4.1. The Cox ring of the Hirzebruch \mathcal{H}_2 is $S = k[x_0, x_1, x_2, x_3]$ with the \mathbb{Z}^2 -grading given by deg $(x_0) = deg(x_2) = (1, 0)$ and deg $(x_1) = (-2, 1)$ and deg $(x_3) = (0, 1)$. Since S is connected, the effective cone $\{d \in \mathbb{Z}^2 \mid S_d \neq 0\}$ of \mathcal{H}_2 is exactly the support Λ . Since Λ has bounded factorization, there is a number $N(\Lambda, d)$ bounding the projective dimension of all ideals whose degree sequence is less than d in any variety whose effective cone sits inside that of \mathcal{H}_2 .

As discussed in the introduction, some Γ -gradings under consideration are exotic enough to fail to admit a flattening map $\Lambda \to \mathbb{Z}_+$. The following is one such example with Stillman bounded projective dimension:

Example 4.2. Let $S = k[x_0, x_1, x_2, \ldots]$ and define p_n , where $n \in \mathbb{N}$, to be the *n*th prime natural number. Define the degree of any constant to be 0, $\deg(x_0) = 1 \in \mathbb{Q}$, and $\deg(x_n) = n + \frac{1}{p_n} \in \mathbb{Q}$. Let $\Lambda \subseteq \mathbb{Q}$ be the \mathbb{Q} -support of this grading and then let Γ be the subgroup of \mathbb{Q} spanned by Λ . Then the grading on S is connected and thus Λ is pointed. Given any $q \in \mathbb{Q}$ it is clear that $\{n \mid n + \frac{1}{p_n} \leq q\}$ is finite, so Λ has bounded factorization. Therefore due to Theorem 1.2, S has Stillman bounded projective dimension. Interestingly, there is no flattening homomorphism $\Lambda \to \mathbb{Z}_+$. Indeed if φ is such a map, since 1 and $\frac{1}{p_n}$ are contained in Γ for all n, we have that $\varphi(1) = p_n \varphi(\frac{1}{p_n})$ for all n, which is impossible as φ takes values in the integers.

References

- [AH16] Tigran Ananyan and Melvin Hochster, Small subalgebras of polynomial rings and stillman's conjecture, J. Amer. Math. Soc. 33 (2016), no. 1, 291–309. arXiv:1610.09268. ↑1, 3, 5
- [BES20] Christine Berkesch, Daniel Erman, and Gregory G. Smith, Virtual resolutions for a product of projective spaces, Algebr. Geom. 7 (2020), no. 4, 460–481. arXiv:1703.07631. ↑1
- [BHS22] Juliette Bruce, Lauren Cranton Heller, and Mahrud Sayrafi, Bounds on multigraded regularity (2022). arXiv:2208.11115. ↑1
- [BMNB⁺11] Jesse S. Beder, Jason McCullough, Luis Núñez-Betancourt, Alexandra Seceleanu, Bart Snapp, and Branden Stone, *Ideals with larger projective dimension and regularity*, J. Symb. Comput. 46 (2011), 1105–1113. arXiv:1101.3368. ↑1
 - [BPC22] Caitlyn Booms-Peot and John Cobb, Virtual criterion for generalized eagon-northcott complexes, J. Pure. Appl. Algebra 226 (2022), no. 12. arXiv:2111.13773. ↑1
 - [BS22] Michael K. Brown and Mahrud Sayrafi, A short resolution of the diagonal for smooth projective toric varieties of picard rank 2 (2022). arXiv:2208.00562. ↑1
 - [Bur68] Lindsay Burch, A note on the homology of ideals generated by three elements in local rings, Math. proc. camb. philos. soc., 1968, pp. 949–952. ↑5
 - [CH22] Marc Chardin and Rafael Holanda, Multigraded tor and local cohomology (2022). arXiv:2211.14357. ↑1
 - [CMR06] Laura Costa and Rosa María Miró-Roig, m-blocks collections and castelnuovo-mumford regularity in multiprojective spaces, Nagoya Math. J. 186 (2006), 119 –155. arXiv:0609560. ↑1
 - [CN20] Marc Chardin and Navid Nemati, Multigraded regularity of complete intersections (2020). arXiv:2012.14899. ↑1
 - [Cob24] John Cobb, Syzygies of curves in products of projective spaces, Math. Z. 307 (2024), no. 27. arXiv:2301.05979. ↑1
 - [DLL19] Jan Draisma, Michal Lasón, and Anton Leykin, Stillman's conjecture via generic initial ideals, Comm. Algebra 47 (2019), 2384–2395. arXiv:1802.10139. ↑1

- [ESS18] Daniel Erman, Steven Sam, and Andrew Snowden, Cubics in 10 variables vs. cubics in 1000 variables: Uniformity phenomena for bounded degree polynomials, Bull. Am. Math. Soc. 56 (2018), no. 1, 87–114. arXiv:1809.09402. ↑1
- [ESS19] _____, Big polynomial rings and stillman's conjecture, Invent. Math. 218 (2019), no. 2, 413–439. arXiv:1801.09852. ↑1
- [ESS21] _____, Generalizations of stillman's conjecture via twisted commutative algebra, Int. Math. Res. Not. 2021 (2021), no. 16, 12281–12304. arXiv:1804.09807. ↑1, 3
- [GV23] Felix Gotti and Joseph Vulakh, On the atomic structure of torsion-free monoids, Semigr. Forum 107 (2023), no. 2, 402–423. arXiv:2212.08347. [↑]2, 4
- [Hà07] Huy Tài Hà, Multigraded regularity, a^{*}-invariant and the minimal free resolution, J. Algebra 310 (2007), no. 1, 156–179. arXiv:0501479. ↑1
- [HS02] Mark D. Haiman and Bernd Sturmfels, *Multigraded hilbert schemes*, J. Algebr. Geom. **13** (2002), 725–769. arXiv:0201271. ↑1
- [HSS06] Milena Hering, Hal Schenck, and Gregory G. Smith, Syzygies, multigraded regularity and toric varieties, Compos. Math. 142 (2006), no. 6, 1499–1506. arXiv:0502240. ↑1
- [Koh72] Peter Kohn, Ideals generated by three elements, Proc. Amer. Math. Soc. **35** (1972), no. 1, 55–58. $\uparrow 5$
- [KZ20] David Kazhdan and Tamar Ziegler, Properties of high rank subvarieties of affine spaces, Geom. Funct. Anal. 30 (2020), no. 4, 1063–1096. arXiv:1902.00767. ↑3
- [LS12] Victor Lozovanu and Gregory G. Smith, Vanishing theorems and the multigraded regularity of nonsingular subvarieties (2012). arXiv:1208.0484. ↑1
- [McC11] Jason McCullough, A family of ideals with few generators in low degree and large projective dimension, Proc. Amer. Math. Soc. 139 (2011), no. 6, 2017–2023. arXiv:1005.3361. ↑1
- [MM19] Paolo Mantero and Jason McCullough, The projective dimension of three cubics is at most 5, J. Pure. Appl. Algebra 223 (2019), no. 4, 1383–1410. arXiv:1801.08195. [↑]3
- [MP18] Jason McCullough and Irena Peeva, Counterexamples to the eisenbud–goto regularity conjecture, J. Amer. Math. Soc. 31 (2018), no. 2, 473–496. ↑1
- [MS04] Diane Maclagan and Gregory G. Smith, Multigraded castelnuovo-mumford regularity, J. Reine Angew. Math. 571 (2004), 179–212. arXiv:0305214. ↑1
- [MS12] Jason McCullough and Alexandra Seceleanu, Bounding projective dimension, Commutative algebra: Expository papers dedicated to david eisenbud on the occasion of his 65th birthday, 2012, pp. 551–576. ↑5
- [Pee10] Irena Peeva, Graded syzygies, Vol. 14, Springer Science & Business Media, 2010. ↑1
- [PS09] Irena Peeva and Mike Stillman, Open problems on syzygies and hilbert functions, J. Commut. Algebra 1 (2009), no. 1, 159–195. ↑1
- [Yan19] Jay Yang, Virtual resolutions of monomial ideals on toric varieties, Proc. Amer. Math. Soc. Ser. B (2019). arXiv:1906.00508. ↑1

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN, MADISON, WI Email address: jcobb2@math.wisc.edu URL: https://johndcobb.github.io

Email address: npgallup@ucdavis.edu
URL: https://sites.google.com/view/nathanielgallup/home

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN, MADISON, WI *Email address*: john.spoerl@wisc.edu