

Complex Symplectic Contractions and 3d Mirrors

Andrew Dancer,^{a,c} Julius F. Grimminger,^a Johan Martens,^b and Zhenghao Zhong^a

^a*Mathematical Institute, University of Oxford,*

Andrew Wiles Building, Woodstock Road, Oxford, OX2 6GG, UK

^b*School of Mathematics and Maxwell Institute, University of Edinburgh, James Clerk Maxwell Building, Edinburgh EH9 3FD, Scotland, UK*

^c*Jesus College, Oxford OX1 3DW, UK*

E-mail: dancer@maths.ox.ac.uk, julius.grimminger@maths.ox.ac.uk,
johan.martens@ed.ac.uk, zhenghao.zhong@maths.ox.ac.uk

ABSTRACT: We propose magnetic quivers for the complex-symplectic contraction spaces, which are related to implosions and have a natural interpretation in terms of the Moore-Tachikawa category. We use 3-d mirrors to provide computational checks.

Contents

1	Introduction and Summary	1
2	Magnetic Quivers for Contraction Spaces	2
2.1	The Moore-Tachikawa Category	3
2.2	Implosion and Contraction Spaces	3
3	3d Mirrors	7
4	Ortho-Symplectic Contractions (D-type)	10
A	Unframed Quivers and Choice of Gauge Group	13
B	Hyper-Kähler Quotients of Coulomb Branch Symmetries	14
B.1	Hyper-Kähler Quotient by Diagonal Symmetry of two Legs	14
B.2	Hyper-Kähler quotient of $U(1)_{\text{top}}$ of a single node.	15
B.3	Illustrative Examples	17

1 Introduction and Summary

Complex-symplectic varieties, and in particular symplectic singularities [1], have been at the heart of geometric representation theory for many years. In physics these spaces show up prominently as Higgs branches [2] of supersymmetric quantum field theories (SQFTs) with 8 supercharges in various dimensions (e.g. 3d $\mathcal{N} = 4$, 4d $\mathcal{N} = 2$, 5d $\mathcal{N} = 1$, or 6d $\mathcal{N} = (1, 0)$ SQFTs), or as Coulomb branches of 3d $\mathcal{N} = 4$ SQFTs [3–7].

The construction of such a space as the 3d $\mathcal{N} = 4$ Coulomb branch of a *quiver gauge theory* is extremely useful, as it allows us to compute a range of its properties via combinatorial operations on its quiver. From now on we will refer to such a quiver \mathbf{Q} , whose 3d $\mathcal{N} = 4$ Coulomb branch $\mathcal{C}(\mathbf{Q})$ is a variety X , as a *magnetic quiver* for X . We use $\mathcal{H}(\mathbf{Q})$ to denote the Higgs branch of the quiver, which is the symplectic dual [8] of $\mathcal{C}(\mathbf{Q})$.

On the other hand, we may also be able to construct our space in question directly as a Higgs branch of a quiver, which we then refer to as an *electric quiver* for our space. Two quivers \mathbf{Q} and \mathbf{Q}^\vee are called a *3d mirror pair*, if among other things (see [9]) the Higgs branch of one is the Coulomb branch of the other and vice versa.

In previous work [10–12] we investigated the magnetic quivers of complex-symplectic implosion spaces [13]. We remark here that in the A_n case such spaces have a stratified hyper-Kähler structure and this is expected to be true in general, hence they are often referred to as hyper-Kähler implosions.

In the current paper we extend this to the related construction of complex-symplectic contraction [14]. The contraction of a complex-symplectic space has the same dimension

as the original space but the symmetry is enhanced by an abelian group (in the simplest case by a maximal torus factor). As for implosion, the complex-symplectic contraction operation is the complex-symplectic analogue of operations in equivariant algebraic [15] and real symplectic geometry [16]. Contractions may be understood in terms of a universal example, which is a complex-symplectic quotient of the square of the universal implosion. In the Moore-Tachikawa category of complex-symplectic varieties [17] the contraction morphism is the composition of two implosion morphisms. A complex-symplectic quotient by abelian factors which are diagonal in the symmetry of two spaces (which is the case in this construction) leaves behind an abelian factor. Hence the global symmetry of contraction spaces are larger than ‘usual’.

We construct the magnetic quivers of contraction spaces as an operation on the implosion magnetic quivers, realising the complex-symplectic quotient on the Coulomb branch. We give background on these quotients in Appendix B. We note that these are complex-symplectic quotients by a complex reductive group, so can be interpreted as hyper-Kähler quotients.

To conclude the introduction we mention some advantages of having a magnetic quiver description Q of a variety X :

1. The stratification into symplectic leaves [18, 19] (and their transverse slices) can be obtained from a quiver subtraction algorithm [20–23] and more recently from a ‘decay and fission’ algorithm [24, 25].
2. Computing the Higgs branch of the magnetic quiver Q (which is well defined if the quiver is simply-laced) provides the symplectic dual X^\vee of the variety X .
3. We see discrete operations, like the quotient by the Weyl group.
4. Deformations (and the Namikawa Weyl group) of the symplectic dual X^\vee can be studied using a subtraction algorithm on the magnetic quiver [26].
5. By construction, if $X = \mathcal{C}(Q)$, X has only symplectic singularities [27].

Plan of the paper

The paper is organized as follows. In Section 2 we review the Moore-Tachikawa category and how the implosion and contraction constructions fit into this framework. We provide magnetic quivers of contraction spaces and explore the symplectic duals of the contraction spaces. In Section 3 we discuss 3d mirror pairs, using the quiver description of the $SU(n)$ implosion, and provide computational confirmation of our proposals for magnetic quivers. In Section 4 we discuss D-type contractions. In the Appendices we provide background material.

2 Magnetic Quivers for Contraction Spaces

In this section we first review some basics of the Moore-Tachikawa category and how implosion and contraction spaces show up as morphisms. We then construct explicit magnetic quivers for contraction spaces, and provide a variety of checks.

2.1 The Moore-Tachikawa Category

In this section we study certain morphisms in the Moore-Tachikawa category [17],¹ whose objects are reductive groups, and morphisms are complex-symplectic varieties whose global symmetry includes the source and target:

$$G_1 \xrightarrow{X_{12}} G_2. \quad (2.1)$$

Composition of morphisms is given via a complex-symplectic quotient:

$$\begin{array}{ccccc} G_1 & \xrightarrow{X_{12}} & G_2 & \xrightarrow{X_{23}} & G_3, \\ & \searrow & & \nearrow & \\ & & X_{13} = X_{23} \circ X_{12} & & \end{array} \quad (2.2)$$

with

$$X_{13} = X_{23} \circ X_{12} = (X_{12} \times X_{23}) // \Delta G_2, \quad (2.3)$$

where ΔG_2 denotes the diagonally embedded G_2 in the symmetry group $G_1 \times G_2 \times G_2 \times G_3$ of $X_{12} \times X_{23}$.

In general, this quotient will have a residual action of $G_1 \times G_3$, and hence be a morphism from G_1 to G_3 as required. However, if we allow the groups G to have nontrivial centers, then we may acquire more symmetries. In particular, if the middle group G_2 is Abelian then X_{13} actually has an action of $G_1 \times G_2 \times G_3$ because the anti-diagonally embedded G_2 commutes with the diagonally embedded G_2 and hence descends to the quotient.

Remark 1 *We note that there is a question in general as to how to interpret the quotient (2.3). We shall view this as the Geometric Invariant Theory quotient by ΔG_2 of the zero level set for the complex-symplectic moment map for this group. In cases of physical interest, the varieties are expected to be Coulomb branches and so affine varieties, in which case the quotient will exist as an affine variety since we are taking our groups to be reductive.*

Remark 2 *It may be possible to construct higher morphisms in the Moore-Tachikawa picture, with the role of 2-morphisms between morphisms X and Y in $\text{Hom}(G_1, G_2)$ being played by Lagrangian correspondences $L \subset X^- \times Y$ with a $G_1 \times G_2$ action where X^- refers to X equipped with the opposite symplectic structure. Floer homology groups $HF(L_1, L_2)$ might be used to define higher morphisms. We refer the reader to Cazassus [28] for a further discussion of this topic.*

2.2 Implosion and Contraction Spaces

Universal Implosion Let G be a simple group. The universal (right) implosion is a space Q_G of (complex) dimension $\dim G + \text{rank } G$ with an action of $G \times T$, where T is

¹The objects in the category which Moore and Tachikawa call *HS* are semi-simple Lie groups. Here we allow for any reductive group, in particular abelian factors, but still refer to it as Moore-Tachikawa category.

the maximal complex torus in G . Complex-symplectic reduction by T at level zero gives the nilpotent cone, with a residual action of G . Reductions at general levels give Kostant varieties, which are closures of complex coadjoint orbits.

This is universal in the following sense. Given a complex-symplectic space M with a G action, we may form its implosion

$$M_{\text{impl}} = (M \times \mathbb{Q}_G) // \Delta G \quad (2.4)$$

which now has a T action. In terms of the Moore-Tachikawa category (enlarged to allow reductive groups), the universal implosion is an element of $\text{Hom}(G, T)$ and a space with a G action is viewed as living in $\text{Hom}(1, G)$. The implosion operation is now just composition with the universal implosion to obtain an element M_{impl} of $\text{Hom}(1, T)$.

We may think of the universal right implosion \mathbb{Q}_G as being obtained from T^*G , which has a $G \times G$ action, by breaking the right action to a T action. Explicitly

$$\mathbb{Q}_G = \{(g, v) \in G \times \mathfrak{u}^\circ\} // U \quad (2.5)$$

where U denotes the maximal unipotent subgroup of G acting by $(g, v) \mapsto (gu^{-1}, \text{Ad}(u)v)$, and $\mathfrak{u} = \text{Lie}(U)$. Equivalently, the right implosion is the complex-symplectic quotient (in the Geometric Invariant Theory sense) of T^*G by the right action of the maximal unipotent U . As a morphism in the enlarged Moore-Tachikawa category we have:

$$G \xrightarrow{\mathbb{Q}_G} T. \quad (\text{Universal Implosion})$$

We also have an equivalent notion, denoted by ${}_G\mathbb{Q}$, of left implosion with a $T \times G$ action, so representing an element of $\text{Hom}(T, G)$. In the $G = SL_n$ case we have an explicit quiver description of the implosion [13].

A special feature of the implosion in complex symplectic geometry (not present for implosion in real symplectic geometry) is the presence of an extra symmetry by the Weyl group W_G of G : there is an action on \mathbb{Q}_G of $G \times (W_G \times T)$. A trace of this can be seen on the total space of the Grothendieck-Springer resolution of G , which is the quotient of an open subvariety of \mathbb{Q}_G by T . The Weyl group only acts on the regular part of the Grothendieck-Springer resolution though, whereas it acts on all of \mathbb{Q}_G .

In [10] magnetic quivers for the implosion were introduced. Let $G = SL_{n+1}$ (so $T = GL_1^n$), then, denoting as above the Coulomb branch by \mathcal{C} , we have:

$$\mathbb{Q}_G = \mathcal{C} \left(\begin{array}{ccccccc} & & & & & & \circ 1 \\ & & & & & & \vdots \\ & & & & & & \circ 1 \\ & & & & & & \circ 1 \\ & & & & & & \circ 1 \\ & & & & & & \circ 1 \\ \circ 1 & \text{---} & \circ 2 & \text{---} & \cdots & \text{---} & \circ n-1 & \text{---} & \circ n & \begin{array}{l} \text{---} \circ 1 \\ \text{---} \circ 1 \\ \text{---} \circ 1 \end{array} \\ & & & & & & & & & \left. \vphantom{\begin{array}{l} \circ 1 \\ \vdots \\ \circ 1 \\ \circ 1 \\ \circ 1 \end{array}} \right\} n+1 \end{array} \right). \quad (2.6)$$

The flavour symmetry of $X = \mathbb{Q}_{SL_{n+1}}$ is $F_X = SU(n+1) \times U(1)^n$. Gannon and Williams [29] have recently given an algebro-geometric proof that this is the correct magnetic quiver, and observed that the Weyl group action on the implosion is manifest as the symmetry permuting the Abelian bouquet nodes emanating from the n -dimensional node. It is interesting to note that the Weyl action is much less obvious in the quiver Higgs branch description of the implosion given in [13] (for an explicit description of the action in this picture see Wang [30]).

Partial Implosion We also have a notion of partial implosions, where we take the complex-symplectic quotient in the GIT sense of T^*G by the maximal unipotent U_P of a parabolic subgroup P of G . The universal contraction is

$$\mathbb{Q}_{G,P} = (G \times \mathfrak{u}_P^\circ) // U_P. \quad (2.7)$$

This has an action of $G \times L_P$ where L_P is the Levi subgroup of P , as P is the semidirect product of U_P and L_P , so L_P normalises U_P . The case when P is the Borel recovers the ordinary implosion. In general we have:

$$G \xrightarrow{\mathbb{Q}_{G,P}} L_P. \quad (\text{Partial Implosion})$$

where L_P is a Levi of G .

When $G = SL_{n+1}$ and $L_P = S(\prod_{i=1}^l GL_{k_i})$ (where $\sum_{i=1}^l k_i = n+1$),

$$\mathbb{Q}_{G,P} = \mathcal{C} \left(\begin{array}{ccccccc} & & & & & k_1 & \dots & 1 \\ & & & & & \circ & \dots & \circ \\ & & & & & \vdots & & \\ & & & & & k_{l-1} & \dots & 1 \\ & & & & & \circ & \dots & \circ \\ & & & & & k_l & \dots & 1 \\ 1 & 2 & \dots & n-1 & n & & & \end{array} \right). \quad (2.8)$$

$\mathbb{Q}_{G,P}$ is singular and the flavour symmetry of $\mathbb{Q}_{G,P}$ is $F_{\mathbb{Q}_{G,P}} = SU(n+1) \times S(\prod_{i=1}^l U(k_i))$, unless $k_1 = n$ when $\mathbb{Q}_{G,P} = \mathbb{H}^{n(n+1)}$ and $F_{\mathbb{Q}_{G,P}} = Sp(n(n+1))$, or $k_1 = n+1$ when $\mathbb{Q}_{G,P} = T^*SL_{n+1}$ and $F_{\mathbb{Q}_{G,P}} = SU(n+1)^2$.

Universal Contraction The universal contraction space corresponds to composing the left and right implosion morphisms in the enlarged Moore-Tachikawa category

$$\begin{array}{ccc} G & \xrightarrow{\mathbb{Q}_G} & T \xrightarrow{G\mathbb{Q}} G, \\ & \searrow & \nearrow \\ & & R_G = G\mathbb{Q} \circ \mathbb{Q}_G \end{array} \quad (2.9)$$

A standard example of a 3d mirror pair (in fact an example of a self-mirror) is the following

$$T[SU(n)] = \begin{array}{c} \circ - \circ - \dots - \circ - \circ - \square \\ 1 \quad 2 \quad \quad \quad n-1 \quad n \quad n+1 \end{array} \xleftrightarrow{3d \text{ MS}} \begin{array}{c} \square - \circ - \circ - \dots - \circ - \circ \\ n+1 \quad n \quad n-1 \quad \quad \quad 2 \quad 1 \end{array} \quad (3.1)$$

We have the following 3d mirror pair involving the universal implosion quiver:

$$\begin{array}{c} \circ - \circ - \dots - \circ - \circ \\ 1 \quad 2 \quad \quad \quad n-1 \quad n \end{array} \begin{array}{c} \circ \\ \vdots \\ \circ \\ \circ \end{array} \xleftrightarrow{3d \text{ MS}} \begin{array}{c} \square - \circ - \circ - \dots - \circ - \circ \\ n+1 \quad SU(n) \quad SU(n-1) \quad \quad \quad SU(2) \quad SU(1), \end{array} \quad (3.2)$$

where the theory on the left is $T[SU(n)]$, but with the $n+1$ flavour node ‘exploded’ into $n+1$ $U(1)$ gauge nodes, i.e. we gauge the $U(1)^n$ torus inside the flavour symmetry of the Higgs branch; and the theory on the right is $T[SU(n)]$, but with all the unitary gauge groups replaced with special unitary gauge groups, i.e. we gauge the $U(1)^n$ torus inside the flavour symmetry of the Coulomb branch.

The universal contraction quiver can be obtained by taking two copies of $T[SU(n)]$ and gauging the diagonal $U(1)^n$ of the tori of their Higgs branches. Therefore we expect that the 3d mirror is obtained by taking two copies of $T[SU(n)]$ and gauging the diagonal $U(1)^n$ of the tori of their Coulomb branches.

$$\begin{array}{c} \circ \\ 1 \\ \vdots \\ \circ \\ \vdots \\ \circ \\ 1 \end{array} \begin{array}{c} \circ - \circ - \dots - \circ - \circ \\ 1 \quad 2 \quad \quad \quad n-1 \quad n \end{array} \begin{array}{c} \circ - \circ - \dots - \circ - \circ \\ n \quad n-1 \quad \quad \quad 2 \quad 1 \end{array} \xleftrightarrow{3d \text{ MS}} \begin{array}{c} \square - \circ - \circ - \dots - \circ - \circ \\ n+1 \quad S[U(n)]S[U(n-1)] \quad \quad \quad S[U(2)] \quad S[U(1)] \\ n+1 \quad U(n) \quad U(n-1) \quad \quad \quad U(2) \quad U(1) \end{array} \quad (3.3)$$

The colours represent gauge groups where a diagonal $U(1)$ is decoupled from them. This result has been verified through explicit computation of Hilbert series of both quivers.

The more non-trivial side of the 3d mirror is to check the Higgs branch of the contracted quiver with the Coulomb branch of the product of two linear quivers. To do this, we compute the Hilbert series for the two coordinate rings (equivalently, the Higgs and Coulomb branch chiral ring from a physics point of view).

Note that in the contracted quiver, although there are no flavour nodes, we expect a $U(1)^{n-1}$ symmetry from the loops in the graph. We can compare this with the case of Asymptotically Locally Euclidean hyper-Kähler manifolds of quaternionic dimension one, which are known to arise from the extended Dynkin diagrams of ADE type. In the A_n case, uniquely, the extended diagram is a loop and this gives rise to the $U(1)$ -triholomorphic symmetry that occurs in this case but not in the D_n or E_n cases where the extended diagrams are trees.

The corresponding deformation parameters in the Coulomb branch are the choice of level set in the description of the contraction as the hyper-Kähler quotient by $T \cong U(1)^{n-1}$ of the product of two implosions.

For $n = 3$, the resulting Hilbert series is:

$$\begin{aligned} HS_{\text{Higgs}}^{\text{Contraction}}(t) &= \frac{1 + 6t^4 + 12t^6 + 18t^8 + 24t^{10} + 34t^{12} + 24t^{14} + 18t^{16} + 12t^{18} + 6t^{20} + t^{24}}{(-1+t)^8(1+t)^8(1+t^2)^4(1-t+t^2)^2(1+t+t^2)^2} \\ &= 1 + 2t^2 + 13t^4 + 38t^6 + 115t^8 + 284t^{10} + 666t^{12} + 1392t^{14} + O(t^{15}) \end{aligned} \quad (3.4)$$

which is the same as the Coulomb branch Hilbert series of the product quivers. In the second line, we expressed the result as a Taylor expansion in t . The coefficient of the leading term $2t^2$ is in accordance with our observation above of the $U(1)^2$ Higgs branch global symmetry group of the contracted quiver. The exact equivalence of the two Hilbert series gives very strong indication that the two moduli spaces are the same.

The other side of the duality can also be checked, but due to the complication of the computation, we only provide the first few terms:

$$HS_{\text{Coulomb}}^{\text{Contraction}}(t) = 1 + 18t^2 + 223t^4 + 1966t^6 + 13444t^8 + 74590t^{10} + O(t^{12}). \quad (3.5)$$

This is the same with the Higgs branch Hilbert series of the product quivers. The coefficient of the leading term $18t^2$ is consistent with the expectation that Coulomb branch global symmetry group of the contracted quivers is $SU(3)^2 \times U(1)^2$. The computations (3.4) and (3.5) give strong indications that the mirror pair in (2.15) is correct for $n = 3$. Recall [13],[10] that uniquely in the $n = 3$ case the symmetry group of the implosion is enhanced from $SU(3) \times U(1)^2$ to $SO(8)$, reflecting the fact that the Abelian bouquet nodes are balanced in this case. However when we form the contraction only the $SU(3)^2 \times U(1)^2$ survives. In terms of the quiver diagram, the bouquet nodes become unbalanced again when we fuse the implosion quivers together.

For $n = 4$, the resulting Hilbert series is:

$$\begin{aligned}
HS_{\text{Higgs}}^{\text{Contraction}}(t) &= \frac{1 - t^2 + 16t^4 + 41t^6 + 164t^8 + 499t^{10} + 1437t^{12} + 3336t^{14} + 7492t^{16} + \\
&14524t^{18} + 26484t^{20} + 43272t^{22} + 66107t^{24} + 91941t^{26} + 119980t^{28} + \\
&143407t^{30} + 160831t^{32} + 166042t^{34} + 160831t^{36} + 143407t^{38} + \\
&119980t^{40} + 91941t^{42} + 66107t^{44} + 43272t^{46} + 26484t^{48} + 14524t^{50} + \\
&7492t^{52} + 3336t^{54} + 1437t^{56} + 499t^{58} + 164t^{60} + 41t^{62} + 16t^{64} - t^{66} + t^{68}}{(-1+t)^{18}(1+t)^{18}(1+t^2)^9(1-t+t^2)^5(1+t+t^2)^5(1+t^4)^3} \\
&= 1 + 3t^2 + 28t^4 + 138t^6 + 674t^8 + 2808t^{10} + 10781t^{12} + 37479t^{14} + O(t^{15})
\end{aligned} \tag{3.6}$$

which is the same as the Coulomb branch Hilbert series of the product quivers.

The other side of the duality can also be checked with the first few terms being:

$$HS_{\text{Coulomb}}^{\text{Contraction}}(t) = 1 + 33t^2 + 559t^4 + 6685t^6 + 61900t^8 + 491584t^{10} + O(t^{12}). \tag{3.7}$$

Again, we see the t^2 coefficient gives the dimension of the global symmetry group $SU(4)^2 \times U(1)^3$ of the contraction.

Note that, like in the case of implosion, the action of the Weyl group is not reflected in the Hilbert series, related to the fact that it does not respect the \mathbb{C}^* -action on the universal implosion or contraction spaces.

These results have interesting implications. The study of linear quivers with mixed unitary and special unitary gauge groups is explored in great detail in [33]. However, a natural question there is whether one can have gauge groups where a diagonal $U(1)$ is decoupled, e.g. $S[\prod_i U(n_i)]$. Here, we provided an explicit example where this action will lead to a non-linear 3d mirror quiver. The application of removing diagonal $U(1)$ s can help us expand the landscape of 3d mirror pairs, especially when one side is a non-linear quiver. In particular, it is well known that many non-linear quivers (e.g star-shaped quivers) do not have a 3d mirror with a Lagrangian description (quiver description). Perhaps a 3d mirror description can be obtained through decoupling a diagonal subgroup of a Lagrangian (quiver) theory.

4 Ortho-Symplectic Contractions (D-type)

Following the orthosymplectic implosion constructions in [11] and partial implosion in [12], one can perform the same contractions for these quivers as well.

For simplicity we only focus on $G = SO(2n)$.

Universal Implosion The magnetic quiver construction for the universal implosion Q_G for $G = SO(2n)$ (with $T = U(1)^n$) is [11]:

$$Q_G = \mathcal{C} \left(\begin{array}{c} \left(\begin{array}{ccccccc} \bullet & \bullet & \bullet & \bullet & \dots & \bullet & \begin{array}{c} \circ \\ \vdots \\ \circ \end{array} \\ \left. \begin{array}{l} 2 \\ 2 \\ 4 \\ 4 \\ \dots \\ 2n-2 \end{array} \right\} n \end{array} \right) \Bigg/ \mathbb{Z}_2 \end{array} \right). \tag{4.1}$$

has magnetic quiver construction

$$\mathbb{R}_{G,P} = \mathcal{C} \left(\begin{array}{c} \text{---} \left(\text{red } 2 \text{---} \text{blue } 2 \text{---} \text{red } 4 \text{---} \text{blue } 4 \text{---} \dots \text{---} \text{blue } 2n-2 \right) \begin{array}{c} \text{---} \text{white } k_1 \\ \text{---} \vdots \\ \text{---} \text{white } k_l \\ \text{---} \text{red } 2k \end{array} \left(\text{---} \text{blue } 2n-2 \text{---} \dots \text{---} \text{blue } 4 \text{---} \text{red } 4 \text{---} \text{blue } 2 \text{---} \text{red } 2 \text{---} \right) \text{---} \\ \text{---} \mathbb{Z}_2 \end{array} \right). \quad (4.8)$$

The flavour symmetry of $\mathbb{R}_{G,P}$ is $F_{\mathbb{R}_{G,P}} = SO(2n) \times U(1)^l \times SO(2n)$. The reduction by $U(1)^l$ is

$$\mathbb{R}_{G,P} = \mathcal{C} \left(\begin{array}{c} \text{---} \left(\text{red } 2 \text{---} \text{blue } 2 \text{---} \text{red } 4 \text{---} \text{blue } 4 \text{---} \dots \text{---} \text{blue } 2n-2 \right) \begin{array}{c} \text{---} \text{white } SU(k_1) \\ \text{---} \vdots \\ \text{---} \text{white } SU(k_l) \\ \text{---} \text{red } 2k \end{array} \left(\text{---} \text{blue } 2n-2 \text{---} \dots \text{---} \text{blue } 4 \text{---} \text{red } 4 \text{---} \text{blue } 2 \text{---} \text{red } 2 \text{---} \right) \text{---} \\ \text{---} \mathbb{Z}_{\text{gcd}(2,k_i)} \end{array} \right). \quad (4.9)$$

Acknowledgements

We would like to thank Antoine Bourget for discussions. AD and JM would like to thank the Isaac Newton Institute for Mathematical Sciences, Cambridge, for support and hospitality during the programme *New equivariant methods in algebraic and differential geometry* where work on this paper was undertaken. This work was supported by EPSRC grant no EP/R014604/1. JFG is supported by the EPSRC Open Fellowship (Schafer-Nameki) EP/X01276X/1 and the ‘‘Simons Collaboration on Special Holonomy in Geometry, Analysis and Physics’’. ZZ is supported by the ERC Consolidator Grant # 864828 ‘‘Algebraic Foundations of Supersymmetric Quantum Field Theory’’ (SCFTAlg).

A Unframed Quivers and Choice of Gauge Group

In this paper we deal mostly with unframed unitary or (special) unitary quivers, and some orthosymplectic quivers. We assume the reader is familiar with these concepts, however it is useful to recall some basic notions, see for example [34].

Quiver Notation. We use square nodes for flavour nodes, round nodes for gauge nodes, white for unitary groups (unless specified to be special unitary), red for special orthogonal gauge groups and blue for compact symplectic groups.

$$\begin{aligned}
 \begin{array}{c} \circ \\ n \end{array} &= \begin{array}{c} \circ \\ U(n) \end{array} \\
 \\
 \begin{array}{c} \circ \\ SU(n) \end{array} &= \begin{array}{c} \circ \\ SU(n) \end{array} \\
 \\
 \begin{array}{c} \bullet \\ n \end{array} &= \begin{array}{c} \bullet \\ SO(n) \end{array} \\
 \\
 \begin{array}{c} \bullet \\ n \end{array} &= \begin{array}{c} \bullet \\ USp(n) \end{array} .
 \end{aligned} \tag{A.1}$$

Unitary Quivers. Take a (connected) unframed unitary quiver \mathcal{Q} with N nodes of ranks $(r_i)_{i=1,\dots,N}$. Call $\tilde{G} = \prod_i U(r_i)$ the naive gauge group. There is a $U(1) \subset \tilde{G}$ which acts trivially on the hypermultiplets. We take the actual gauge group of the quiver to be $G = \tilde{G}/U(1)$, decoupling the trivially acting $U(1)$ ³. One would naively associate to every $U(r_i)$ factor in \tilde{G} a $U(1)$ factor in the torus of the Coulomb branch global symmetry. However, since we decoupled the overall $U(1)$, there are only $N - 1$ $U(1)$ factors in the torus of the Coulomb branch global symmetry. Where helpful, we will explicitly denote the decoupling of the $U(1)$ as

$$\mathcal{Q} \Big/ U(1). \tag{A.2}$$

Importantly, when we consider the union of two disjoint quivers, then each quiver has its own $U(1)$ in need of decoupling.

(Special) Unitary Quivers. Now consider a (connected) unframed quiver \mathcal{Q}^{su} with both N unitary and $M > 0$ special unitary nodes. Let the ranks of the special unitary nodes be $(r_j^{\text{su}} - 1)_{j=1,\dots,M}$. Again we may call $\tilde{G} = \prod_i U(r_i) \times \prod_j SU(r_j^{\text{su}})$ the naive gauge group. Let $g = \text{gcd}(r_j^{\text{su}})$. There is a $\mathbb{Z}_g \subset \tilde{G}$ which acts trivially on the hypermultiplets. We may *pick* any divisor $k|g$ and specify the actual gauge group to be $G^k = \tilde{G}/\mathbb{Z}_k$.⁴ This

³In the maths literature this is sometimes referred to as the Crawley-Boevey trick referring to an observation in [35].

⁴ \mathbb{Z}_g is called a 1-form symmetry in the physics literature, and changing the gauge group from \tilde{G} to \tilde{G}/\mathbb{Z}_k is achieved by gauging the subgroup $\mathbb{Z}_k \subset \mathbb{Z}_g$ of the 1-form symmetry.

Unitary Quiver. First let's start with a purely unitary quiver with a node of rank n

The diagram shows a quiver with a top node labeled $u(1)$ in red, which is circled with a red dashed line. Below it is a node labeled n . Ellipses \dots indicate intermediate nodes. The bottom node is labeled Q . A diagonal slash is drawn below the quiver, with $U(1)$ written to its right.

where we associate a $u(1)$ Coulomb branch symmetry to the node in question, keeping in mind that one decouples an overall $U(1)$.

We can turn this quiver in an unframed (special) unitary quiver

The diagram shows a quiver with a top node labeled $SU(n)$. Below it is a node labeled n . Ellipses \dots indicate intermediate nodes. The bottom node is labeled Q . A diagonal slash is drawn below the quiver, with \mathbb{Z}_n written to its right.

which has the same Higgs and Coulomb branch. We refer to this operation as a hyper-Kähler quotient by $SU(1)$, which is of course not meaningful as a geometric operation, but it is still a non-trivial operation on the quiver.

(Special) Unitary Quiver. Now take a (special) unitary quiver with $g = \gcd(r_j^{\text{su}})$, and a unitary node of rank n

The diagram shows a quiver with a top node labeled $u(1)$ in red, which is circled with a red dashed line. Below it is a node labeled n . Ellipses \dots indicate intermediate nodes. The bottom node is labeled Q^{su} . A diagonal slash is drawn below the quiver, with \mathbb{Z}_g written to its right.

where the node in question carries a $u(1)$ Coulomb branch symmetry, and there is no decoupling of an overall $U(1)$.

We can perform a hyper-Kähler quotient by $U(1)$, resulting in the (special) unitary

quiver

(B.9)

The quaternionic dimension of the Coulomb branch is decreased by 1 as expected. Furthermore, the centre of the gauge group remains unchanged, due to the quotient by $\mathbb{Z}_{\gcd(g,n)}$ in the latter quiver.

B.3 Illustrative Examples

In Figure 1 we give the simplest example relevant our paper of the hyper-Kähler quotients mentioned in Appendix B.1. In Figure 2 we give the simplest example relevant to our paper of the hyper-Kähler quotients mentioned in Appendix B.2.




Figure 1. Hyper-Kähler quotient of product of partial implosions by the diagonal of the Levy to reach the partial contraction, on the level of magnetic quivers and their 3d mirrors.

References

- [1] A. Beauville, *Symplectic singularities*, *Invent. Math.* **139** (2000) 541–549, [[math/9903070](#)].
- [2] N. J. Hitchin, A. Karlhede, U. Lindstrom and M. Rocek, *Hyperkahler Metrics and Supersymmetry*, *Commun. Math. Phys.* **108** (1987) 535.
- [3] S. Cremonesi, A. Hanany and A. Zaffaroni, *Monopole operators and Hilbert series of Coulomb branches of 3d $\mathcal{N} = 4$ gauge theories*, *JHEP* **01** (2014) 005, [[1309.2657](#)].
- [4] M. Bullimore, T. Dimofte and D. Gaiotto, *The Coulomb Branch of 3d $\mathcal{N} = 4$ Theories*, *Commun. Math. Phys.* **354** (2017) 671–751, [[1503.04817](#)].
- [5] H. Nakajima, *Towards a mathematical definition of Coulomb branches of 3-dimensional $\mathcal{N} = 4$ gauge theories, I*, *Adv. Theor. Math. Phys.* **20** (2016) 595–669, [[1503.03676](#)].
- [6] A. Braverman, M. Finkelberg and H. Nakajima, *Towards a mathematical definition of Coulomb branches of 3-dimensional $\mathcal{N} = 4$ gauge theories, II*, *Adv. Theor. Math. Phys.* **22** (2018) 1071–1147, [[1601.03586](#)].
- [7] A. Braverman, M. Finkelberg and H. Nakajima, *Coulomb branches of 3d $\mathcal{N} = 4$ quiver gauge theories and slices in the affine Grassmannian*, *Adv. Theor. Math. Phys.* **23** (2019) 75–166, [[1604.03625](#)].
- [8] T. Braden, A. Licata, N. Proudfoot and B. Webster, *Quantizations of conical symplectic resolutions II: category \mathcal{O} and symplectic duality*, *Astérisque* (2016) 75–179, [[1407.0964](#)].
- [9] K. A. Intriligator and N. Seiberg, *Mirror symmetry in three-dimensional gauge theories*, *Phys. Lett.* **B387** (1996) 513–519, [[hep-th/9607207](#)].
- [10] A. Dancer, A. Hanany and F. Kirwan, *Symplectic duality and implosions*, *Adv. Theor. Math. Phys.* **25** (2021) 1367–1387, [[2004.09620](#)].
- [11] A. Bourget, A. Dancer, J. F. Grimminger, A. Hanany, F. Kirwan and Z. Zhong, *Orthosymplectic implosions*, *JHEP* **08** (2021) 012, [[2103.05458](#)].
- [12] A. Bourget, A. Dancer, J. F. Grimminger, A. Hanany and Z. Zhong, *Partial implosions and quivers*, *JHEP* **07** (2022) 049, [[2112.10825](#)].
- [13] A. Dancer, F. Kirwan and A. Swann, *Implosion for hyperkähler manifolds*, *Compos. Math.* **149** (2013) 1592–1630, [[1209.1578](#)].
- [14] A. Dancer, F. Kirwan and J. Martens, *Implosion, contraction and Moore-Tachikawa*, *Int. J. Math.* (2024) to appear, [[2401.07920](#)].
- [15] V. L. Popov, *Contractions of actions of reductive algebraic groups*, *Mat. Sb. (N.S.)* **130(172)** (1986) 310–334, 431.
- [16] J. Hilgert, C. Manon and J. Martens, *Contraction of Hamiltonian K -spaces*, *Int. Math. Res. Not. IMRN* (2017) 6255–6309, [[1509.06406](#)].
- [17] G. W. Moore and Y. Tachikawa, *On 2d TQFTs whose values are holomorphic symplectic varieties*, *Proc. Symp. Pure Math.* **85** (2012) 191–208, [[1106.5698](#)].
- [18] H. Kraft and C. Procesi, *Minimal singularities in GL_n* , *Inventiones mathematicae* **62** (1980) 503–515.
- [19] D. Kaledin, *Symplectic singularities from the poisson point of view*, *Journal für die reine und angewandte Mathematik (Crelles Journal)* **2006** (2006) 135–156.

- [20] S. Cabrera and A. Hanany, *Quiver Subtractions*, *JHEP* **09** (2018) 008, [[1803.11205](#)].
- [21] A. Bourget, S. Cabrera, J. F. Grimminger, A. Hanany, M. Sperling, A. Zajac et al., *The Higgs mechanism — Hasse diagrams for symplectic singularities*, *JHEP* **01** (2020) 157, [[1908.04245](#)].
- [22] A. Bourget, J. F. Grimminger, A. Hanany and Z. Zhong, *The Hasse Diagram of the Moduli Space of Instantons*, [2202.01218](#).
- [23] A. Bourget and J. F. Grimminger, *Fibrations and Hasse diagrams for 6d SCFTs*, *JHEP* **12** (2022) 159, [[2209.15016](#)].
- [24] A. Bourget, M. Sperling and Z. Zhong, *Decay and Fission of Magnetic Quivers*, *Phys. Rev. Lett.* **132** (2024) 221603, [[2312.05304](#)].
- [25] A. Bourget, M. Sperling and Z. Zhong, *Higgs branch RG-flows via Decay and Fission*, [2401.08757](#).
- [26] A. Bourget, S. Giacomelli and J. F. Grimminger, *FI-flows of 3d $N=4$ Theories*, [2302.03698](#).
- [27] A. Weekes, *Quiver gauge theories and symplectic singularities*, *arXiv e-prints* (May, 2020) [arXiv:2005.01702](#), [[2005.01702](#)].
- [28] G. Cazassus, *A two-category of Hamiltonian manifolds, and a $(1+1+1)$ field theory*, *Indiana Univ. Math. J.* **72** (2023) 2101–2144, [[1903.10686](#)].
- [29] T. Gannon and H. Williams, *Differential operators on the base affine space of SL_n and quantised Coulomb branches*, [2312.10278](#).
- [30] X. Wang, *A new Weyl group action related to the quasi-classical Gelfand-Graev action*, *Selecta Math. (N.S.)* **27** (2021) Paper No. 38, 44, [[1912.04539](#)].
- [31] B. Jia, *The geometry of the affine closure of $T^*(SL_n/U)$* , [2112.08649](#).
- [32] T. Gannon, *Proof of the Ginzburg-Kazhdan conjecture*, *Advances in Mathematics* **448** (2024) 109701, [[2302.11160](#)].
- [33] A. Bourget, J. F. Grimminger, A. Hanany, R. Kalveks and Z. Zhong, *Higgs branches of U/SU quivers via brane locking*, *JHEP* **08** (2022) 061, [[2111.04745](#)].
- [34] A. Bourget, J. F. Grimminger, A. Hanany, R. Kalveks, M. Sperling and Z. Zhong, *Magnetic Lattices for Orthosymplectic Quivers*, *JHEP* **12** (2020) 092, [[2007.04667](#)].
- [35] W. Crawley-Boevey, *Geometry of the moment map for representations of quivers*, *Compos. Math.* **126** (May, 2001) 257–293.
- [36] A. Hanany, R. Kalveks and G. Kumaran, *Quotient quiver subtraction*, [2308.05853](#).
- [37] F. Benini, Y. Tachikawa and D. Xie, *Mirrors of 3d Sicilian theories*, *JHEP* **09** (2010) 063, [[1007.0992](#)].