Convergence analysis of a regularized Newton method with generalized regularization terms for convex optimization problems

Yuya Yamakawa[∗]and Nobuo Yamashita†

July 10, 2024

Abstract. This paper presents a regularized Newton method (RNM) with generalized regularization terms for unconstrained convex optimization problems. The generalized regularization includes quadratic, cubic, and elastic net regularizations as special cases. Therefore, the proposed method serves as a general framework that includes not only the classical and cubic RNMs but also a novel RNM with elastic net regularization. We show that the proposed RNM has the global $\mathcal{O}(k^{-2})$ and local superlinear convergence, which are the same as those of the cubic RNM.

Keywords. unconstrained convex optimization, regularized Newton method, generalized regularization, global $\mathcal{O}(k^{-2})$ convergence, superlinear convergence, local convergence

1 Introduction

We consider the following unconstrained convex optimization problem:

$$
\underset{x \in \mathbb{R}^n}{\text{minimize}} \quad f(x),\tag{1}
$$

where the function f is twice continuously differentiable and convex on \mathbb{R}^n .

Newton's method is one of the most well-known and basic iterative methods for solving unconstrained convex optimization problems. Each iteration computes a search direction d_k , which is a solution of the following subproblem:

$$
\underset{d \in \mathbb{R}^n}{\text{minimize}} \quad \langle \nabla f(x_k), d \rangle + \frac{1}{2} \langle \nabla^2 f(x_k) d, d \rangle,
$$

[∗]Graduate School of Informatics, Kyoto University, Yoshidahommachi, Sakyo-ku, Kyoto-shi, Kyoto 606-8501, Japan, E-mail: yuya@i.kyoto-u.ac.jp

[†]Graduate School of Informatics, Kyoto University, Yoshidahommachi, Sakyo-ku, Kyoto-shi, Kyoto 606-8501, Japan, Email: nobuo@kyoto-u.ac.jp

and updates the current point x_k as $x_{k+1} := x_k + t_k d_k$, where $t_k > 0$ denotes a step size. It converges rapidly thanks to the use of the second-order information, that is, $\nabla^2 f(x_k)$, of the objective function. However, it requires that $\nabla^2 f(x_k)$ is nonsingular at each iteration. Even if the Hessian is nonsingular, the convergence rate may be reduced to linear when the Hessian is close to singular. Several variants of Newton's method have been proposed to overcome these drawbacks including regularized Newton methods (RNMs) [\[1,](#page-14-0) [2,](#page-14-1) [3,](#page-14-2) [4,](#page-14-3) [5\]](#page-14-4), cubic RNMs [\[6,](#page-15-0) [7,](#page-15-1) [8\]](#page-15-2), and so forth [\[9,](#page-15-3) [10,](#page-15-4) [11,](#page-15-5) [12,](#page-15-6) [13,](#page-15-7) [14,](#page-15-8) [15,](#page-15-9) [16,](#page-15-10) [17,](#page-15-11) [18\]](#page-16-0).

RNMs can be considered modifications of Newton's method because they improve the subproblem of Newton's method such that it can be solved even if the Hessian matrix is singular. More precisely, RNMs iteratively solve the following subproblem to find a search direction d_k :

$$
\underset{d\in\mathbb{R}^n}{\text{minimize}} \quad \langle \nabla f(x_k), d \rangle + \frac{1}{2} \langle \nabla^2 f(x_k) d, d \rangle + \frac{\mu_k}{2} ||d||^2,
$$

where $\mu_k > 0$ denotes a parameter. Since the objective function is strongly convex, the subproblem has a unique optimum. Nesterov and Polyak [\[6\]](#page-15-0) proposed an RNM with the cubic regularization $\frac{\mu_k}{6} ||d||^3$. The proposed method is called the cubic RNM and iteratively solves the following subproblem:

$$
\underset{d\in\mathbb{R}^n}{\text{minimize}} \quad \langle \nabla f(x_k), d \rangle + \frac{1}{2} \langle \nabla^2 f(x_k) d, d \rangle + \frac{\mu_k}{6} ||d||^3,
$$

For classical and cubic RNMs, global $\mathcal{O}(k^{-2})$ and local superlinear convergence were proven in $[2, 4, 6, 5]$ $[2, 4, 6, 5]$ $[2, 4, 6, 5]$ $[2, 4, 6, 5]$.

For least squares problems, Ariizumi, Yamakawa, and Yamashita [\[19\]](#page-16-1) recently proposed a Levenberg-Marquardt method (LMM) equipped with a generalized regularization term and showed its global and local superlinear convergence. Although a subproblem of the classical LMM has a quadratic regularization term $\frac{\mu_k}{2} ||d||^2$, they generalized the regularization term such that another regularization can be adopted, such as the $L¹$ and elastic-net regularization. Moreover, they reported numerical experiments in which their LMM with the elastic-net regularization worked well for certain examples, owing to the sparsity of the search direction.

Inspired by Ariizumi, Yamakawa, and Yamashita [\[19\]](#page-16-1), we propose a generalized RNM (GRNM) for solving problem [\(1\)](#page-0-0). We adopt new regularization terms provided as $\frac{\mu_k}{p} ||d||_2^p + \rho_k ||d||_1$, where $p \in (1, 3]$ is a pre-fixed parameter. With the addition of the new regularization terms, the GRNM includes classical and cubic RNMs as well as novel RNMs with other regularizations, such as the elastic net and so forth. More precisely, if $p = 2$ and $\rho_k = 0$, the GRNM is reduced to the classical RNM; if $p = 3$ and $\rho_k = 0$, it is equivalent to the cubic RNM. Moreover, if $p = 2$ and $\rho_k > 0$, it can be regarded as a novel RNM with the elastic-net regularization.

The contributions of this study are as follows. This study provides

- (i) the generalized RNM stated above;
- (ii) sufficient conditions of p, μ_k , and ρ_k for which the GRNM has the global $\mathcal{O}(k^{-2})$ convergence;
- (iii) local superlinear convergence under the local error bound condition.

Hence, these contributions include classical and cubic RNMs as special cases and provide a framework for new RNMs such as the elastic-net RNM.

The remainder of this paper is organized as follows. Section [2](#page-2-0) provides a general proposition that plays an important role in the analysis of global $\mathcal{O}(k^{-2})$ convergence. Section [3](#page-5-0) describes the proposed method. Sec-tion [4](#page-6-0) presents global $\mathcal{O}(k^{-2})$ convergence of the proposed method. Section [5](#page-11-0) proves local and superlinear convergence. Finally, concluding remarks are presented in Section [6.](#page-14-5)

Throughout the paper, we use the following mathematical notation. Let N be the set of natural numbers (positive integers). For $p \in \mathbb{N}$ and $q \in \mathbb{N}$, the set of real matrices with p rows and q columns is denoted by $\mathbb{R}^{p \times q}$. Note that $\mathbb{R}^{p\times 1}$ is equal to the set of p-dimensional real vectors, that is, $\mathbb{R}^{p\times 1} = \mathbb{R}^p$, and note that \mathbb{R}^1 represents the set of real numbers, namely, $\mathbb{R}^1 = \mathbb{R}$. For any $w \in \mathbb{R}^p$, the transposition of w is represented as $w^{\top} \in \mathbb{R}^{1 \times p}$. For $u \in \mathbb{R}^p$ and $v \in \mathbb{R}^p$, the inner product of u and v is defined by $\langle u, v \rangle := u^\top v$. We denote by I the identity matrix, where these dimensions are defined by the context. For each $w \in \mathbb{R}^p$, the Euclidean and L^1 norms of w are respectively defined by $||w|| := \sqrt{\langle w, w \rangle}$ and $||w||_1 := |[w]_1| + |[w]_2| + \cdots + |[w]_p|$, where $[w]_j$ represents the j-th element of w. For $W \in \mathbb{R}^{p \times q}$, we denote by $||W||$ the operator norm of W, that is, $||W|| := \sup{||W u||, ||u|| \leq 1}$. Let φ be a function from \mathbb{R}^p to \mathbb{R} . The gradient of φ at $w \in \mathbb{R}^p$ is represented as $\nabla \varphi(w)$. The Hessian of φ at $w \in \mathbb{R}^p$ is denoted by $\nabla^2 \varphi(w)$. For a convex function $\phi: \mathbb{R}^p \to \mathbb{R}$, we denote by $\partial \phi(w)$ the subdifferential of ϕ at w. For $\eta \in \mathbb{R}^p$ and $r > 0$, we define $B(\eta, r) := \{ \mu \in \mathbb{R}^p; ||\mu - \eta|| \leq r \}$. For infinite sequences ${a_k} \subset \mathbb{R}$ and ${b_k} \subset \mathbb{R}$, we write $a_k = \mathcal{O}(b_k)$ $(k \to \infty)$ if there exist $c > 0$ and $n \in \mathbb{N}$ such that $|a_k| \leq c|b_k|$ for all $k \geq n$.

2 Preliminaries

This section presents a general proposition that provides sufficient conditions under which arbitrary sequences generated by optimization methods have global $\mathcal{O}(k^{-2})$ convergence. This proposition plays a critical role in Section [4.](#page-6-0) The proof of the proposition is inspired by the technique presented in [\[5,](#page-14-4) Theorem 1].

Proposition 1. Let $f: \mathbb{R}^n \to \mathbb{R}$ be a continuously differentiable convex function with a minimum $x^* \in \mathbb{R}^n$. Let $\{x_k\}$ be an infinite sequence in \mathbb{R}^n . Suppose that

- (i) $f(x_{k+1}) \leq f(x_k)$ for all $k \in \mathbb{N} \cup \{0\};$
- (ii) there exists $\gamma > 0$ such that $\|\nabla f(x_{k+1})\| \leq \gamma \|\nabla f(x_k)\|$ for all $k \in$ $\mathbb{N} \cup \{0\}$:
- (iii) there exists $\delta > 0$ such that $f(x_k) f^* \leq \delta \|\nabla f(x_k)\|$ for all $k \in \mathbb{N} \cup \{0\}$;
- (iv) there exist $\theta \in (0,1)$, $\nu > 0$, and $\ell \in \mathbb{N}$ such that $\theta^k \leq \nu k^{-2}$ and $(\gamma \theta)^{\frac{k}{2}} \leq \nu k^{-2}$ for all $k \geq \ell$, where γ is given in (ii).

Let $\mathcal{I}(\theta) := \{i \in \mathbb{N} \cup \{0\}; \theta \|\nabla f(x_i)\| \leq \|\nabla f(x_{i+1})\|\}$. Suppose also that

(v) there exists $\tau > 0$ such that $f(x_{k+1}) - f(x_k) \leq -\tau (f(x_k) - f^*)^{\frac{3}{2}}$ for all $k \in \mathcal{I}(\theta)$.

Then, one of the following statements holds:

(a) If $|\mathcal{I}(\theta)| < \infty$ holds, then

$$
f(x_k) - f^* \le \frac{\theta^{-(\widehat{i}+1)} \nu \delta \|\nabla f(x_{\widehat{i}+1})\|}{k^2} \quad \forall k \ge \max\{\ell, \widehat{i}+2\},
$$

where \hat{i} is the largest element of $\mathcal{I}(\theta)$.

(b) If $|\mathcal{I}(\theta)| = \infty$ holds, then

$$
f(x_k) - f^* \le \max\left\{\frac{36\tau^{-2}}{(k+4)^2}, \frac{\nu\delta \|\nabla f(x_0)\|}{k^2}\right\} \quad \forall k \ge \ell.
$$

Proof. Let $i_{\ell} \in \mathbb{N}$ be the ℓ -th smallest element of $\mathcal{I}(\theta)$, that is, $\mathcal{I}(\theta)$ = ${i_1, i_2, \ldots, \hat{i}}$ with $i_\ell < i_{\ell+1}$. Moreover, regarding assumption (ii), we suppose $\gamma \geq 1$ without loss of generality.

To begin with, we consider case (a), that is, $|\mathcal{I}(\theta)| < \infty$ is satisfied. Let $\mathcal{J}(\theta) \coloneqq \{i \in \mathbb{N} \cup \{0\}; \theta \|\nabla f(x_i)\| > \|\nabla f(x_{i+1})\|\}.$ Note that $\widehat{i} = i_{|\mathcal{I}(\theta)|}$. We can easily observe that

$$
k \in \mathcal{J}(\theta) \quad \forall k > \hat{i}.\tag{2}
$$

For every $k \geq \hat{i} + 2$, it follows from [\(2\)](#page-3-0) that $j \in \mathcal{J}(\theta)$ for any $j \in \{\hat{i} + 1, \hat{i} + \hat{j}\}$ $2, \ldots, k-1\}$, that is,

$$
\|\nabla f(x_k)\| < \theta \|\nabla f(x_{k-1})\| < \dots < \theta^{k - (\hat{i} + 1)} \|\nabla f(x_{\hat{i} + 1})\|.\tag{3}
$$

Let us take $k \ge \max{\{\ell, \hat{i} + 2\}}$ arbitrarily. By assumption (iii) and [\(3\)](#page-3-1), we obtain

$$
f(x_k) - f^* \le \delta \|\nabla f(x_k)\| < \theta^{k - (\hat{i} + 1)} \delta \|\nabla f(x_{\hat{i} + 1})\|.
$$
 (4)

Recall that $\theta^k \leq \nu k^{-2}$ from assumption (iv). Thus, the desired inequality is derived from (4) .

Next, we discuss the case where $|\mathcal{I}(\theta)| = \infty$ holds. Let $k \in \mathbb{N} \cup \{0\}$ and define $\psi_k := \tau^2(f(x_{i_k}) - f^*)$. Combining assumption (i) and $i_{k+1} \geq i_k + 1$ yields

$$
\psi_{k+1} = \tau^2(f(x_{i_{k+1}}) - f^*) \le \tau^2(f(x_{i_k+1}) - f^*). \tag{5}
$$

Since assumption (v) implies that $f(x_{i_k+1}) - f(x_{i_k}) \leq -\tau (f(x_{i_k}) - f^*)^{\frac{3}{2}}$,

$$
\tau^{2}(f(x_{i_{k}+1}) - f^{*}) \leq \tau^{2}(f(x_{i_{k}}) - f^{*}) - \tau^{3}(f(x_{i_{k}}) - f^{*})^{\frac{3}{2}} = \psi_{k} - \psi_{k}^{\frac{3}{2}}.
$$
 (6)

Exploiting [\(5\)](#page-4-0) and [\(6\)](#page-4-1) derives $\psi_{k+1} \leq \psi_k - \psi_k^{\frac{3}{2}} \leq \psi_k - \frac{2}{3}$ $\frac{2}{3}\psi_k^{\frac{3}{2}}$. It then follows from [\[5,](#page-14-4) Proposition 1] that $\psi_k \leq 9(k+2)^{-2}$, that is,

$$
f(x_{i_k}) - f^* \le \frac{9\tau^{-2}}{(k+2)^2} \quad \forall k \in \mathbb{N} \cup \{0\}.
$$
 (7)

Let $\mathcal{I}_k := \{i \in \mathcal{I}(\theta); i \leq k\}$. In the following, we assume $k \geq \ell$. There are two possible cases: Case (1) $|\mathcal{I}_k| \geq \frac{k}{2}$ and Case (2) $|\mathcal{I}_k| < \frac{k}{2}$.

Case (1): The largest element of \mathcal{I}_k can be represented by $i_{|\mathcal{I}_k|}$, and thus $i_{\vert \mathcal{I}_k \vert} \leq k$. This fact, assumption (i), and [\(7\)](#page-4-2) imply that

$$
f(x_k) - f^* \le f(x_{i_{|\mathcal{I}_k|}}) - f^* \le \frac{9\tau^{-2}}{(|\mathcal{I}_k| + 2)^2} \le \frac{36\tau^{-2}}{(k+4)^2}.
$$

Case (2): From assumption (ii), each $j \in \{0, 1, \ldots, k-1\}$ satisfies

$$
\|\nabla f(x_{j+1})\| \leq \gamma \|\nabla f(x_j)\| \quad \text{if } j \in \mathcal{I}_{k-1},
$$

$$
\|\nabla f(x_{j+1})\| < \theta \|\nabla f(x_j)\| \quad \text{if } j \notin \mathcal{I}_{k-1}.
$$
 (8)

Combining assumption (iii) and [\(8\)](#page-4-3) derives

$$
f(x_k) - f^* \le \delta \|\nabla f(x_k)\| \le \delta \gamma^{|\mathcal{I}_{k-1}|} \theta^{k - |\mathcal{I}_{k-1}|} \|\nabla f(x_0)\|.
$$
 (9)

Note that $\gamma \geq 1, \theta \in (0,1)$, and $(\gamma \theta)^{\frac{k}{2}} \leq \nu k^{-2}$ hold from assumption (iv) and $k \geq \ell$. It then follows from $|\mathcal{I}_{k-1}| \leq |\mathcal{I}_k| < \frac{k}{2}$ that

$$
\gamma^{\left|\mathcal{I}_{k-1}\right|}\theta^{k-\left|\mathcal{I}_{k-1}\right|} \leq \left(\gamma\theta\right)^{\frac{k}{2}} \leq \frac{\nu}{k^2}.\tag{10}
$$

We have from [\(9\)](#page-4-4) and [\(10\)](#page-4-5) that

$$
f(x_k) - f^* \le \frac{\nu \delta \|\nabla f(x_0)\|}{k^2}.
$$

Cases (1) and (2) guarantee that the desired inequality holds when $|\mathcal{I}(\theta)| = \infty$. Therefore, the assertion is proven ∞. Therefore, the assertion is proven.

Remark 1. We discuss sufficient conditions for assumptions (i)–(iv) of Proposition [1](#page-2-1). Assumptions (i) and (ii) would be satisfied for any sequence generated by descent methods. Note that γ of item (ii) is allowed to be greater than or equal to 1. Assumption (iii) is satisfied when $\{x_k\}$ is bounded. Assumption (iv) holds if $\theta \in (0, \gamma^{-1})$ because it implies $\theta^k = \mathcal{O}(k^{-2})$ and $(\gamma \theta)^{\frac{k}{2}} = \mathcal{O}(k^{-2})$ as $k \to \infty$. From these discussions, we can see that assumption (v) is the key to global $\mathcal{O}(k^{-2})$ convergence.

3 An RNM with generalized regularization terms

In this paper, we consider an RNM with generalized regularization terms that iteratively solves the following subproblem:

$$
\underset{d \in \mathbb{R}^n}{\text{minimize}} \quad \langle \nabla f(x_k), d \rangle + \frac{1}{2} \langle \nabla^2 f(x_k) d, d \rangle + \frac{\mu_k}{p} ||d||^p + \rho_k ||d||_1,\tag{11}
$$

where $\mu_k > 0$ and $\rho_k \geq 0$ are parameters, and $p \in (1,3]$ is a pre-fixed constant. The proposed method obtains a search direction d_k by solving subproblem [\(11\)](#page-5-1), and the point x_k is updated as $x_{k+1} := x_k + t_k d_k$, where t_k is a step size.

Now, we denote by φ_k the objective function of subproblem [\(11\)](#page-5-1), namely,

$$
\varphi_k(d) \coloneqq \langle \nabla f(x_k), d \rangle + \frac{1}{2} \langle \nabla^2 f(x_k) d, d \rangle + \frac{\mu_k}{p} ||d||^p + \rho_k ||d||_1.
$$

Since the objective function φ_k has generalized regularization terms $\frac{\mu_k}{p}||d||^p$ and $\rho_k||d||_1$, we call the proposed method a generalized RNM (GRNM).

Remark 2. The GRNM includes the quadratic, cubic, and elastic net regularization as special cases. Moreover, it includes a novel regularization in addition to the aforementioned regularization.

Remark 3. The proposed GRNM can adopt the L^1 regularization term, that is, $\mu_k = 0$ and $\rho_k > 0$. However, subproblem [\(11\)](#page-5-1) with $\mu_k = 0$ might have no global optimum when $\nabla^2 f(x_k)$ is not positive definite. Conversely, if ρ_k is sufficiently large, the solution becomes 0. We provide sufficient conditions under which [\(11\)](#page-5-1) has nonzero solutions.

Lemma 1. Let $x_k \in \mathbb{R}^n$, $\mu_k > 0$, and $\rho_k \ge 0$ be given. If $\rho_k < ||\nabla f(x_k)||_{\infty}$, then problem [\(11\)](#page-5-1) has a unique global optimum $d_k \neq 0$ that satisfies

$$
\nabla f(x_k) + (\nabla^2 f(x_k) + \mu_k ||d_k||^{p-2} I) d_k + \rho_k \eta_k = 0
$$

for some $\eta_k \in \partial ||d_k||_1$. Moreover, d_k is the descent direction of f at x_k , that is, $\langle \nabla f(x_k), d_k \rangle < 0$.

Proof. First, we show the solvability of [\(11\)](#page-5-1). Recall that φ_k is closed, proper, and coercive. Hence, by using [\[20,](#page-16-2) Proposition 3.2.1], problem [\(11\)](#page-5-1) has a global optimum d_k . The uniqueness of d_k is derived from the strict convexity of φ_k .

Hereafter, we show that $d_k \neq 0$ is satisfied. We assume to the contrary that $d_k = 0$ holds. As $d_k = 0$ satisfies the first-order optimality condition of [\(11\)](#page-5-1), there exists $\eta_k \in \partial ||d_k||_1$ such that $\nabla f(x_k) + \rho_k \eta_k = 0$. It then follows from $\rho_k < ||\nabla f(x_k)||_{\infty}$ that $||\nabla f(x_k)||_{\infty} = \rho_k < ||\nabla f(x_k)||_{\infty}$. This result contradicts, that is, $d_k \neq 0$.

Finally, the first-order optimality condition of [\(11\)](#page-5-1) leads to the desired equality, and it yields

$$
\langle \nabla f(x_k), d_k \rangle = -\langle (\nabla^2 f(x_k) + \mu_k ||d_k||^{p-2} I) d_k, d_k \rangle - \rho_k ||d_k||_1 < 0,
$$

where note that $\langle d_k, \eta_k \rangle = ||d_k||_1$ and $d_k \neq 0$. This completes the proof. \Box

Remark 4. By utilizing the line search strategy or an appropriate choice of μ_k and ρ_k , we can prove the global convergence of Algorithm [1](#page-6-1). However, because $\mathcal{O}(k^{-2})$ convergence implies global convergence, we omit discussions on the line search.

We provide a formal description of the proposed method in Algorithm [1.](#page-6-1)

Algorithm 1 (GRNM)

1: Choose $p \in (1,3], x_0 \in \mathbb{R}^n, \varepsilon > 0$, and set $k \coloneqq 0$.

2: If $\|\nabla f(x_k)\| \leq \varepsilon$, then stop.

- 3: Set parameters $\mu_k > 0$ and $\rho_k \geq 0$, and find a global optimum d_k of [\(11\)](#page-5-1).
- 4: Set $x_{k+1} := x_k + d_k$.
- 5: Set $k \leftarrow k + 1$, and go to Line [2.](#page-6-1)

4 Global $\mathcal{O}(k^{-2})$ convergence of Algorithm [1](#page-6-1)

This section shows that Algorithm [1](#page-6-1) globally converges with the $\mathcal{O}(k^{-2})$ rate. From now on, we denote by $x^* \in \mathbb{R}^n$ an optimal solution of problem [\(1\)](#page-0-0), and use the following notation: $f^* := f(x^*)$ and $\mathcal{S} := \{x \in \mathbb{R}^n; f(x) \leq f(x_0)\}.$

We will show global $\mathcal{O}(k^{-2})$ convergence of Algorithm [1](#page-6-1) by showing that assumptions (i)–(v) in Proposition [1](#page-2-1) hold for a sequence $\{x_k\}$ generated by Algorithm [1.](#page-6-1)

In the subsequent argument, we suppose that $\varepsilon = 0$ and Algorithm [1](#page-6-1) generates an infinite sequence $\{x_k\}$ satisfying $\nabla f(x_k) \neq 0$ for each $k \in$ ^N ∪ {0}. Moreover, we make the following assumptions.

(A1) There exists $L > 0$ such that for any $x, y \in \mathbb{R}^n$,

$$
\begin{aligned} \|\nabla f(x) - \nabla f(y) - \nabla^2 f(y)(x - y)\| &\le L \|x - y\|^2, \\ |f(x) - f(y) - \langle \nabla f(y), x - y \rangle - \frac{1}{2} \langle \nabla^2 f(y)(x - y), x - y \rangle| &\le \frac{L}{3} \|x - y\|^3. \end{aligned}
$$

(A2) The parameters μ_k and ρ_k are set as follows: For all $k \in \mathbb{N} \cup \{0\},$

$$
\mu_k := c_1^{\frac{p-1}{2}} \|\nabla f(x_k)\|^{\frac{3-p}{2}}, \ \rho_k := \min\left\{\frac{q}{\sqrt{n}}\|\nabla f(x_k)\|, c_2\|\nabla f(x_k)\|^{\frac{p+1}{2}}\right\},\
$$

where $c_1 \geq L, c_2 \in (0, 1)$, and $q \geq 0$.

(A3) There exists $R > 0$ such that $S \subset B(0, R)$.

Note that subproblem [\(11\)](#page-5-1) has a global optimum $d_k \neq 0$ because (A2) satisfies the condition of Lemma [1.](#page-5-2) Note also that several basic properties of linear algebra derive

$$
\left\| (\nabla^2 f(x_k) + \mu_k \|d_k\|^{p-2} I)^{-1} \right\| \le \mu_k^{-1} \|d_k\|^{2-p},\tag{12}
$$

$$
\left\| (\nabla^2 f(x_k) + \mu_k \|d_k\|^{p-2} I)^{-1} \nabla^2 f(x_k) \right\| \le 1.
$$
 (13)

We now show that assumptions (i) and (ii) in Proposition [1](#page-2-1) hold.

Lemma 2. Suppose that (A1) and (A2) are satisfied. Suppose also that $3 - (1+q)^{\frac{3-p}{p-1}} > 0$ where q is a constant in (A2). For any $k \in \mathbb{N} \cup \{0\}$, the following inequalities hold:

(a)
$$
f(x_{k+1}) - f(x_k) \le -\frac{3 - (1 + q)^{\frac{3-p}{p-1}}}{3} \mu_k \|d_k\|^p < 0
$$
,
\n(b) $\|\nabla f(x_{k+1})\| \le \left(1 + (1 + q)^{\frac{3-p}{p-1}}\right) \mu_k \|d_k\|^{p-1}$
\n $+ \min \left\{q \|\nabla f(x_k)\|, \sqrt{n}c_2 \|\nabla f(x_k)\|^{\frac{p+1}{2}}\right\},$
\n(c) $\|\nabla f(x_{k+1})\| \le \left(1 + 2q + (1 + q)^{\frac{2}{p-1}}\right) \|\nabla f(x_k)\|.$

Proof. It follows from Lemma [1](#page-5-2) and [\(12\)](#page-7-0) that

$$
||d_k|| \le ||(\nabla^2 f(x_k) + \mu_k ||d_k||^{p-2} I)^{-1} (\nabla f(x_k) + \rho_k \eta_k) ||
$$

\n
$$
\le \mu_k^{-1} ||d_k||^{2-p} (||\nabla f(x_k)|| + \sqrt{n} \rho_k)
$$

\n
$$
\le \mu_k^{-1} ||d_k||^{2-p} (1+q) ||\nabla f(x_k)||,
$$

that is,

$$
||d_k|| \le \frac{(1+q)^{\frac{1}{p-1}}}{\sqrt{c_1}} \sqrt{||\nabla f(x_k)||}.
$$
 (14)

By (14) and the first equality of $(A2)$, we have

$$
L||d_k||^2 \le c_1||d_k||^{3-p} \cdot ||d_k||^{p-1}
$$

\n
$$
\le (1+q)^{\frac{3-p}{p-1}} \cdot c_1^{\frac{p-1}{2}} ||\nabla f(x_k)||^{\frac{3-p}{2}} \cdot ||d_k||^{p-1}
$$

\n
$$
= (1+q)^{\frac{3-p}{p-1}} \mu_k ||d_k||^{p-1}.
$$
\n(15)

Recall that $x_{k+1} = x_k + d_k$ and $\langle \eta_k, d_k \rangle = ||d_k||_1$. Combining the second inequality of (A1), Lemma [1,](#page-5-2) and [\(15\)](#page-8-0) yields

$$
f(x_{k+1}) - f(x_k) \le \langle \nabla f(x_k) + \nabla^2 f(x_k) d_k, d_k \rangle - \frac{1}{2} \langle \nabla^2 f(x_k) d_k, d_k \rangle + \frac{L}{3} ||d_k||^3
$$

\n
$$
\le -\mu_k ||d_k||^p - \rho_k ||d_k||_1 + \frac{1}{3} ||d_k|| \cdot L ||d_k||^2
$$

\n
$$
\le -\frac{3 - (1+q)^{\frac{3-p}{p-1}}}{3} \mu_k ||d_k||^p.
$$

From $x_{k+1} = x_k + d_k$ and Lemma [1,](#page-5-2) we obtain

$$
\nabla f(x_{k+1}) = \nabla f(x_k + d_k) - \nabla f(x_k) - (\nabla^2 f(x_k) + \mu_k ||d_k||^{p-2} I) d_k - \rho_k \eta_k.
$$

Subsequently, exploiting the first inequality of (A1), the second equality of $(A2)$, and (15) derives

$$
\|\nabla f(x_{k+1})\| \le \|\nabla f(x_k + d_k) - \nabla f(x_k) - \nabla^2 f(x_k) d_k\| + \mu_k \|d_k\|^{p-1} + \sqrt{n}\rho_k
$$

\n
$$
\le L \|d_k\|^2 + \mu_k \|d_k\|^{p-1}
$$

\n
$$
+ \min \left\{ q \|\nabla f(x_k)\|, \sqrt{n}c_2 \|\nabla f(x_k)\|^{\frac{p+1}{2}} \right\}
$$

\n
$$
\le \left(1 + (1+q)^{\frac{3-p}{p-1}}\right) \mu_k \|d_k\|^{p-1}
$$

\n
$$
+ \min \left\{ q \|\nabla f(x_k)\|, \sqrt{n}c_2 \|\nabla f(x_k)\|^{\frac{p+1}{2}} \right\}, \qquad (16)
$$

namely, item (b) is verified. Meanwhile, from [\(14\)](#page-7-1) and the first equality of (A2), we have $\mu_k ||d_k||^{p-1} \leq (1+q) ||\nabla f(x_k)||$. Utilizing this result, [\(16\)](#page-8-1), and $\min\{q\|\nabla f(x_k)\|, \sqrt{n}c_2\|\nabla f(x_k)\|^{\frac{p+1}{2}}\} \leq q\|\nabla f(x_k)\|$ means

$$
\|\nabla f(x_{k+1})\| \le \left(1 + 2q + (1+q)^{\frac{2}{p-1}}\right) \|\nabla f(x_k)\|.
$$

Therefore, the desired inequalities are obtained.

Now, we provide the global $\mathcal{O}(k^{-2})$ convergence property of Algorithm [1](#page-6-1) by showing assumptions (iii)- (v) in Proposition [1.](#page-2-1)

Theorem 1. Suppose that $(A1)$ – $(A3)$ hold. Moreover, suppose that the following assumptions $(A4)–(A6)$ hold:

 \Box

- (A4) $3 (1+q)^{\frac{3-p}{p-1}} > 0;$
- (A5) $1 + 2q + (1 + q)^{\frac{2}{p-1}} \geq 1;$
- (A6) there exist $\theta \in (q, 1)$, $\nu > 0$, and $\ell \in \mathbb{N}$ such that

$$
\theta^k \le \nu k^{-2}, \quad \left(\left(1 + 2q + (1+q)^{\frac{2}{p-1}} \right) \theta \right)^{\frac{k}{2}} \le \nu k^{-2} \quad \forall k \ge \ell.
$$

Let $\mathcal{I}(\theta) \coloneqq \{i \in \mathbb{N} \cup \{0\}; \theta \|\nabla f(x_i)\| \leq \|\nabla f(x_{i+1})\|\}.$ Let D and τ be defined as

$$
D := R + \|x^*\|, \quad \tau := \frac{(\theta - q)^{\frac{p}{p-1}} \left(3 - (1+q)^{\frac{3-p}{p-1}}\right)}{3\sqrt{c_1}D^{\frac{3}{2}} \left(1 + (1+q)^{\frac{3-p}{p-1}}\right)^{\frac{p}{p-1}}}.
$$

Then, a sequence $\{x_k\}$ generated by Algorithm [1](#page-6-1) satisfies one of the following statements:

(a) If $|\mathcal{I}(\theta)| < \infty$, then

$$
f(x_k) - f^* \le \frac{\theta^{-(\widehat{i}+1)} \nu D \|\nabla f(x_{\widehat{i}+1})\|}{k^2} \quad \forall k \ge \max\{\ell, \widehat{i}+2\},
$$

where \widehat{i} is the largest element of $\mathcal{I}(\theta)$.

(b) If $|\mathcal{I}(\theta)| = \infty$, then

$$
f(x_k) - f^* \le \max\left\{\frac{36\tau^{-2}}{(k+4)^2}, \frac{\nu D \|\nabla f(x_0)\|}{k^2}\right\} \quad \forall k \ge \ell.
$$

Proof. If items (i)–(v) in Proposition [1](#page-2-1) hold, then the desired result can be obtained. Item (i) directly follows from (a) of Lemma [2.](#page-7-2) Let us define $r := 1+2q+(1+q)^{\frac{2}{p-1}}$. Recall that (A5) ensures $r \geq 1$. Thus, item (ii) holds from (c) of Lemma [2.](#page-7-2) The definition of r and (A6) imply that item (iv) is satisfied. Thus, it is sufficient to show items (iii) and (v).

Since $\{x_k\} \subset \mathcal{S}$ holds, we have from (A3) that $||x_k|| \leq R$ for $k \in \mathbb{N} \cup \{0\}$. Let us take $k \in \mathbb{N} \cup \{0\}$ arbitrarily. Then, it is clear that $||x_k - x^*|| \leq D$, and hence the convexity of f yields $f(x_k) - f^* \le \langle \nabla f(x_k), x_k - x^* \rangle \le D || \nabla f(x_k) ||$. This fact implies that item (iii) holds, and

$$
\left(\frac{f(x_k) - f^*}{D}\right)^{\frac{3}{2}} \le \|\nabla f(x_k)\|^{\frac{3}{2}}.
$$
\n(17)

Now, we take arbitrary $k \in \mathcal{I}(\theta)$. The definition of μ_k and (b) of Lemma [2](#page-7-2) derive

$$
\theta \|\nabla f(x_k)\| \le \left(1 + (1+q)^{\frac{3-p}{p-1}}\right) c_1^{\frac{p-1}{2}} \|\nabla f(x_k)\|^{\frac{3-p}{2}} \|d_k\|^{p-1} + q \|\nabla f(x_k)\|,
$$

which implies

$$
\frac{(\theta - q)^{\frac{p}{p-1}}}{c_1^{\frac{p}{2}} \left(1 + (1 + q)^{\frac{3-p}{p-1}}\right)^{\frac{p}{p-1}}} \|\nabla f(x_k)\|^{\frac{p}{2}} \le \|d_k\|^p.
$$

Multiplying both sides of this inequality by $\mu_k = c_1^{\frac{p-1}{2}} \|\nabla f(x_k)\|^{\frac{3-p}{2}}$ yields

$$
\frac{(\theta - q)^{\frac{p}{p-1}}}{\sqrt{c_1} \left(1 + (1 + q)^{\frac{3-p}{p-1}}\right)^{\frac{p}{p-1}}} \|\nabla f(x_k)\|^{\frac{3}{2}} \le \mu_k \|d_k\|^p. \tag{18}
$$

 \Box

Using item (a) of Lemma [2](#page-7-2) and [\(18\)](#page-10-0), we obtain

$$
f(x_{k+1}) - f(x_k) \le -\frac{(\theta - q)^{\frac{p}{p-1}} \left(3 - (1+q)^{\frac{3-p}{p-1}}\right)}{3\sqrt{c_1} \left(1 + (1+q)^{\frac{3-p}{p-1}}\right)^{\frac{p}{p-1}}} \|\nabla f(x_k)\|^{\frac{3}{2}}.
$$
 (19)

Moreover, from (17) and (19) ,

$$
f(x_{k+1}) - f(x_k) \leq -\tau (f(x_k) - f^*)^{\frac{3}{2}} \quad \forall k \in \mathcal{I}(\theta).
$$

Therefore, we can verify that item (v) of Proposition [1](#page-2-1) holds.

From Theorem [1,](#page-8-2) we have to indicate the existence of q and θ satisfying $(A4)$ – $(A6)$ to show global $\mathcal{O}(k^{-2})$ convergence of Algorithm [1.](#page-6-1) Although the existence of these parameters cannot be ensured for all $p > 1$, we can show their existence for specific $p \in (1,3]$. Two examples of these concrete parameters are presented.

Example 1.

$$
q \coloneqq 0, \quad \theta \coloneqq \frac{3}{8}.
$$

Example 2.

$$
q \coloneqq \min\left\{\frac{1}{10}(2^{\frac{p-1}{3-p}} - 1), \frac{1}{20}2^{\frac{3-p}{p-1}}\right\}, \quad \theta \coloneqq \frac{1}{5}.
$$

The parameters q and θ described in Examples [1](#page-10-2) and [2](#page-10-3) satisfy conditions (A4)–(A6). For details, see Appendix [A.](#page-16-3)

Remark 5. When $(p,q,\theta) = (2,0,4^{-1})$, we can easily verify that $\tau =$ $(96D^{3/2}\sqrt{c_1})^{-1}$. This value coincides with that of Mishchenko [\[5\]](#page-14-4), implying that the proposed method is a generalization of [\[5\]](#page-14-4).

Remark 6. Parameter θ described in Theorem [1](#page-8-2) is only required for the proof and is unrelated to problem [\(1\)](#page-0-0) and Algorithm [1](#page-6-1). Hence, it should be selected to provide a good coefficient regarding the convergence rate. Since the coefficients are determined by

$$
\frac{1}{\theta^{(\hat{i}+1)}}, \quad \left[\frac{3\sqrt{c_1}D^{\frac{3}{2}}\left(1+(1+q)^{\frac{3-p}{p-1}}\right)^{\frac{p}{p-1}}}{(\theta-q)^{\frac{p}{p-1}}\left(3-(1+q)^{\frac{3-p}{p-1}}\right)}\right]^2,
$$

we should take θ as large as possible.

5 Local superlinear convergence of Algorithm [1](#page-6-1)

This section aims to show local and superlinear convergence of Algorithm [1.](#page-6-1) Throughout this section, the set of optimal solutions is denoted by $X^* \subset \mathbb{R}^n$. We first make an additional assumption.

(A7) There exist $r_1 > 0$ and $m_1 > 0$ such that $dist(x, X^*) \leq m_1 ||\nabla f(x)||$ for each $x \in B(x^*, r_1)$.

For a given point $x \in \mathbb{R}^n$, let \widehat{x} be a point satisfying

 $\hat{x} \in X^*, \quad \|\hat{x} - x\| = \text{dist}(x, X^*).$

Some important inequalities for local convergence are as follows:

Lemma 3. Suppose that $(A1)$, $(A2)$, and $(A7)$ hold. Then, there exist $r_2 > 0$, $m_2 > 0$, $m_3 > 0$, and $m_4 > 0$ such that

- (a) $||d_k|| \leq m_2 \text{dist}(x_k, X^*)$ and $||d_k|| \geq m_3 \text{dist}(x_k, X^*)$ for $x_k \in B(x^*, r_2)$;
- (b) dist $(x_{k+1}, X^*) \leq m_4 \text{dist}(x_k, X^*)^{\frac{p+1}{2}}$ for $x_k, x_{k+1} \in B(x^*, r_2)$.

Proof. To begin with, we define u_1 and u_2 as

$$
u_1 := \sup \{ ||\nabla^2 f(z)||; z \in B(x^*, r_1) \}, \quad u_2 := u_1 + \frac{c_1 r_1}{2},
$$

respectively, and will show the following inequality:

$$
\|\nabla f(x)\| \le u_2 \text{dist}(x, X^*) \quad \forall x \in B(x^*, r_1). \tag{20}
$$

We have $\|\hat{x} - x\| \leq \|x - x^*\| \leq r_1$. Hence, the first inequality of (A1) guarantees $\|\nabla f(\hat{x}) - \nabla f(x) - \nabla^2 f(x)(\hat{x} - x)\| \leq \frac{c_1}{2} \text{dist}(x, X^*)^2$. It then follows from $\nabla f(\hat{x}) = 0$ and $dist(x, X^*) \le ||x - x^*|| \le r_1$ that $||\nabla f(x)||$ u_1 dist $(x, X^*) \leq \frac{c_1 r_1}{2}$ $\frac{r_1}{2}$ dist (x, X^*) . Thus, inequality [\(20\)](#page-11-1) holds.

We show item (a). Define r_2 as follows:

$$
r_2 := \min \left\{ r_1, \left(2\sqrt{n} c_2 u_2^{\frac{p-1}{2}} \right)^{-\frac{2}{p-1}} \right\}.
$$

Let $x_k \in B(x^*, r_2)$. The definition of $dist(x_k, X^*)$ and inequality [\(20\)](#page-11-1) ensure

$$
dist(x_k, X^*) \le \|x_k - x^*\| \le r_2
$$
\n(21)

$$
\|\nabla f(x_k)\| \le u_2 \text{dist}(x_k, X^*) \le u_2 \|x_k - x^*\| \le u_2 r_2. \tag{22}
$$

Lemma [1](#page-5-2) and [\(12\)](#page-7-0) lead to $||d_k|| \le ||(\nabla^2 f(x_k) + \mu_k||d_k||^{p-2}I)^{-1}\nabla f(x_k)|| +$ $\sqrt{n}\rho_k$ $\frac{\sqrt{n\rho_k}}{\mu_k||d_k||^{p-2}}$. It then follows from (A2) and [\(20\)](#page-11-1) that

$$
||d_k|| \le ||(\nabla^2 f(x_k) + \mu_k ||d_k||^{p-2} I)^{-1} \nabla f(x_k)||
$$

$$
+ \frac{\sqrt{n}c_2 u_2^{\frac{p+1}{2}}}{\mu_k ||d_k||^{p-2}} \text{dist}(x_k, X^*)^{\frac{p+1}{2}}.
$$
 (23)

Now, we notice that $\nabla f(x_k) = -(\nabla f(\widehat{x}_k) - \nabla f(x_k) - \nabla^2 f(x_k)(\widehat{x}_k - x_k))$ $\nabla^2 f(x_k)(\hat{x}_k - x_k)$ holds from $\nabla f(\hat{x}_k) = 0$. Then, combining (A1), [\(12\)](#page-7-0), and [\(13\)](#page-7-3) implies

$$
\|(\nabla^2 f(x_k) + \mu_k \|d_k\|^{p-2} I)^{-1} \nabla f(x_k)\|
$$

\n
$$
\leq \|(\nabla^2 f(x_k) + \mu_k \|d_k\|^{p-2} I)^{-1} \| \|\nabla f(\widehat{x}_k) - \nabla f(x_k) - \nabla^2 f(x_k)(\widehat{x}_k - x_k)\|
$$

\n
$$
+ \|(\nabla^2 f(x_k) + \mu_k \|d_k\|^{p-2} I)^{-1} \nabla^2 f(x_k) \| \|\widehat{x}_k - x_k\|
$$

\n
$$
\leq \frac{c_1}{2\mu_k \|d_k\|^{p-2}} \text{dist}(x_k, X^*)^2 + \text{dist}(x_k, X^*).
$$
 (24)

By [\(21\)](#page-12-0), [\(23\)](#page-12-1) and [\(24\)](#page-12-2), we get $||d_k|| \leq \frac{u_3}{\mu_k ||d_k||^{p-2}} \text{dist}(x_k, X^*)^{\frac{p+1}{2}} + \text{dist}(x_k, X^*),$ where $u_3 \coloneqq \frac{1}{2}$ $\frac{1}{2}c_1r_2^{\frac{3-p}{2}} + \sqrt{n}c_2u_2^{\frac{p+1}{2}}$. This inequality can be reformulated as $\mu_k ||d_k||^{p-1} \leq 2 \max\{u_3 \text{dist}(x_k, X^*)^{\frac{p+1}{2}}, \mu_k ||d_k||^{p-2} \text{dist}(x_k, X^*)\}.$ There are two possible cases: (i) $\mu_k ||d_k||^{p-1} \leq 2u_3 \text{dist}(x_k, X^*)^{\frac{p+1}{2}}$; (ii) $\mu_k ||d_k||^{p-1} \leq$ $2\mu_k ||d_k||^{p-2}$ dist (x_k, X^*) . In case (i), utilizing (A2) and (A7) derives

$$
\frac{c_1^{\frac{p-1}{2}}\text{dist}(x_k, X^*)^{\frac{3-p}{2}}}{m_1^{\frac{3-p}{2}}} \|d_k\|^{p-1} \le \mu_k \|d_k\|^{p-1} \le 2u_3\text{dist}(x_k, X^*)^{\frac{p+1}{2}}.
$$

Hence, we have $||d_k|| \leq c_1^{-\frac{1}{2}} (2u_3 m_1^{\frac{3-p}{2}})^{\frac{1}{p-1}} \text{dist}(x_k, X^*)$. Meanwhile, case (ii) leads to $||d_k|| \le 2$ dist (x_k, X^*) . Thus, there exists $m_2 > 0$ such that $||d_k|| \le$ $m_2\mathrm{dist}(x_k,X^*)$.

Now, Lemma [1,](#page-5-2) (A2), and [\(22\)](#page-12-3) yield

$$
||d_k|| \ge \frac{||\nabla f(x_k) + \rho_k \eta_k||}{||\nabla^2 f(x_k) + \mu_k||d_k||^{p-2}I||}
$$

\n
$$
\ge \frac{1 - \sqrt{n}c_2(u_2r_2)^{\frac{p-1}{2}}}{u_1 + \mu_k||d_k||^{p-2}} ||\nabla f(x_k)||
$$

\n
$$
\ge \frac{1}{2(u_1 + \mu_k||d_k||^{p-2})} ||\nabla f(x_k)||,
$$
\n(25)

where the last inequality follows from the definition of r_2 . Exploiting (A2) and $||d_k|| \leq m_2 \text{dist}(x_k, X^*)$ derives

$$
\mu_k \|d_k\|^{p-2} \le c_1^{\frac{p-1}{2}} \|\nabla f(x_k)\|^{\frac{3-p}{2}} m_2^{p-2} \text{dist}(x_k, X^*)^{p-2}.
$$

It then follows from [\(21\)](#page-12-0) and [\(22\)](#page-12-3) that

$$
\mu_k \|d_k\|^{p-2} \le c_1^{\frac{p-1}{2}} u_2^{\frac{3-p}{2}} m_2^{p-2} r_2^{\frac{p-1}{2}}.
$$
\n(26)

By [\(25\)](#page-13-0), [\(26\)](#page-13-1), and (A7), there exists $m_3 > 0$ satisfying $||d_k|| \geq m_3 \text{dist}(x_k, X^*)$.

Next, we prove item (b). Let $x_{k+1} = x_k + d_k \in B(x^*, r_2)$. Lemma [1](#page-5-2) implies

$$
\nabla f(x_{k+1}) = (\nabla f(x_k + d_k) - \nabla f(x_k) - \nabla^2 f(x_k) d_k) - (\mu_k ||d_k||^{p-2} d_k + \rho_k \eta_k) + (\nabla f(x_k) + (\nabla^2 f(x_k) + \mu_k ||d_k||^{p-2} I) d_k + \rho_k \eta_k) = (\nabla f(x_k + d_k) - \nabla f(x_k) - \nabla^2 f(x_k) d_k) - (\mu_k ||d_k||^{p-2} d_k + \rho_k \eta_k).
$$

Then, we have from $(A1)$, $(A2)$, and $(A7)$ that

$$
\begin{split} &\text{dist}(x_{k+1}, X^*)\\ &\leq m_1 \|\nabla f(x_k + d_k)\|\\ &\leq m_1 \|\nabla f(x_k + d_k) - \nabla f(x_k) - \nabla^2 f(x_k) d_k\| + m_1 \mu_k \|d_k\|^{p-1} + \sqrt{n} m_1 \rho_k\\ &\leq \frac{c_1 m_1}{2} \|d_k\|^2 + c_1^{\frac{p+1}{2}} m_1 \|\nabla f(x_k)\|^{\frac{3-p}{2}} \|d_k\|^{p-1} + \sqrt{n} c_2 m_1 \|\nabla f(x_k)\|^{\frac{p+1}{2}}. \end{split} \tag{27}
$$

Now, recall that $||d_k|| \leq m_2 \text{dist}(x_k, X^*)$, $||\nabla f(x_k)|| \leq u_2 \text{dist}(x_k, X^*)$, and $dist(x_k, X^*) \leq r_2$, where the second and third inequalities follow from [\(20\)](#page-11-1) and [\(21\)](#page-12-0), respectively. Thus, we can easily verify that

$$
\frac{c_1 m_1}{2} \|d_k\|^2 + c_1^{\frac{p+1}{2}} m_1 \|\nabla f(x_k)\|^{\frac{3-p}{2}} \|d_k\|^{p-1} + \sqrt{n} c_2 m_1 \|\nabla f(x_k)\|^{\frac{p+1}{2}}
$$

$$
\leq \left(\frac{c_1 m_1 m_2^2 r_2^{\frac{3-p}{2}}}{2} + c_1^{\frac{p+1}{2}} u_2^{\frac{3-p}{2}} m_2^{p-1} + \sqrt{n} c_2 m_1 u_2^{\frac{p+1}{2}}\right) \text{dist}(x_k, X^*)^{\frac{p+1}{2}}. (28)
$$

Therefore, combining [\(27\)](#page-13-2) and [\(28\)](#page-13-3) guarantees the existence of $m_4 > 0$ satisfying dist $(x_{k+1}, X^*) \leq m_4 \text{dist}(x_k, X^*)^{\frac{p+1}{2}}$. \Box Finally, we establish local and superlinear convergence of Algorithm [1.](#page-6-1) Although we can prove the theorem using the above lemmas in a manner similar to [\[2,](#page-14-1) Theorem 3.2], the proof is given in Appendix [B](#page-17-0) for completeness of the paper.

Theorem 2. Suppose that (A1), (A2), and (A7) hold. If an initial point x_0 is chosen sufficiently close to x^* , then any sequence $\{x_k\}$ generated by Algo-rithm [1](#page-6-1) converges to some global optimum $\bar{x} \in X^*$ superlinearly. Moreover, if $p = 3$ is satisfied, then $\{x_k\}$ converges to $\bar{x} \in X^*$ quadratically.

6 Concluding remarks

In this paper, we have proposed Algorithm [1,](#page-6-1) which is an RNM with generalized regularization terms. The proposed method is based on the RNM proposed by Mishchenko [\[5\]](#page-14-4), but it is a generalization of the existing one regarding regularization. Therefore, not only the quadratic and cubic RNMs but also novel RNMs with other regularization, such as the elastic net, are included in Algorithm [1.](#page-6-1) We have proven global $\mathcal{O}(k^{-2})$ and local superlinear convergence of Algorithm [1.](#page-6-1)

One of future research is to propose an accelerated GRNM that globally converges in the order of $\mathcal{O}(k^{-3})$.

Declarations

Conflict of interest The authors declare no conflicts of interest.

References

- [1] N. Doikov, K. Mishchenko, Y. Nesterov, Super-universal regularized Newton method, SIAM Journal on optimization 34 (2024) 27–56.
- [2] D. Li, M. Fukushima, L. Qi, N. Yamashita, Regularized Newton methods for convex minimization problems with singular solutions, Computational Optimization and Applications 28 (2004) 131–147.
- [3] Y.-J. Li, D.-H. Li, Truncated regularized Newton method for convex minimizations, Computational Optimization and Applications 43 (2009) 119–131.
- [4] R. A. Polyak, Regularized Newton method for unconstrained convex optimization, Mathematical Programming 120 (2009) 125–145.
- [5] K. Mishchenko, Regularized Newton method with global $\mathcal{O}(1/k^2)$ convergence, SIAM Journal on optimization 33 (2023) 1440–1462.
- [6] Y. Nesterov, B. T. Polyak, Cubic regularization of Newton method and its global performance, Mathematical Programming 108 (2006) 177– 205.
- [7] Y. Nesterov, Accelerating the cubic regularization of newton's method on convex problems, Mathematical Programming 112 (2008) 159–181.
- [8] M.-C. Yue, Z. Zhou, A. M.-C. So, On the quadratic convergence of the cubic regularization method under a local error bound condition, SIAM Journal on Optimization 29 (2019) 90–932.
- [9] D. Goldfarb, Y. Ren, A. Bahamou, Practical quasi-Newton methods for training deep neural networks, in: Advances in Neural Information Processing Systems, 2020, pp. 2386–2396.
- [10] L. Grippo, F. Lampariello, S. Luclidi, A nonmonotone line search technique for Newton's method, SIAM Journal on Numerical Analysis 23 (1986) 707–716.
- [11] A. Rodomanov, Y. Nesterov, Greedy quasi-Newton methods with explicit superlinear convergence, SIAM Journal on Optimization 31 (2021) 785–811.
- [12] R. Crane, F. Roosta, DINO: Distributed Newton-type optimization method, in: Proceedings of the 37th International Conference on Machine Learning, 2020, pp. 2174–2184.
- [13] N. Doikov, P. Richtárik, Randomized block cubic Newton method, in: Proceedings of the 35th International Conference on Machine Learning, PMLR, 2018, pp. 1290–1298.
- [14] R. M. Gower, D. Kovalev, F. Lieder, P. Richtárik, RSN: Randomized subspace Newton, in: Advances in Neural Information Processing Systems, 2019, pp. 616–625.
- [15] F. Hanzely, N. Doikov, P. Richtárik, Y. Nesterov, Stochastic subspace cubic Newton method, in: Proceedings of the 37th International Conference on Machine Learning, 2020, pp. 4027–4038.
- [16] U. Marteau-Ferey, F. Bach, A. Rudi, Globally convergent Newton methods for ill-conditioned generalized self-concordant losses, in: Advances in Neural Information Processing Systems, 2019, pp. 7636–7646.
- [17] A. Rodomanov, D. Kropotov, A superlinearly-convergent proximal Newton-type method for the optimization of finite sums, in: Proceedings of the International Conference on Machine Learning, PRML, 2016, pp. 2597–2605.
- [18] S. Soori, K. Mishchenko, A. Mokhtari, M. M. Dehnavi, M. Gürbüzbalaban, DAve-QN: A distributed averaged quasi-Newton method with local superlinear convergence rate, in: Proceedings of the International Conference on Artificial Intelligence and Statistics, 2020, pp. 1965–1976.
- [19] S. Ariizumi, Y. Yamakawa, N. Yamashita, Convergence properties of Levenberg-Marquardt methods with generalized regularization terms, Applied Mathematics and Computation 463 (2024) 128365.
- [20] D. P. Bertsekas, Convex Optimization Algorithms, Athena Scientific, Massachusetts, 2015.

A

In this appendix, we show that the parameters described in Examples [1](#page-10-2) and [2](#page-10-3) satisfy assumptions $(A4)$ – $(A6)$ stated in Theorem [1.](#page-8-2)

We now discuss Example [1.](#page-10-2) It can be verified that

$$
q = 0, \quad 3 - (1 + q)^{\frac{3-p}{p-1}} = 3, \quad 1 + 2q + (1 + q)^{\frac{2}{p-1}} = 2,
$$

$$
\left(1 + 2q + (1 + q)^{\frac{2}{p-1}}\right)\theta = \frac{3}{4}.
$$

Thus, we can easily verify that assumptions (A4) and (A5) hold. Moreover, assumption (A6) is obtained from

$$
\theta^k = \left(\frac{3}{8}\right)^k = \mathcal{O}(k^{-2}) \quad (k \to \infty),
$$

$$
\left(\left(1 + 2q + (1+q)^{\frac{2}{p-1}}\right)\theta\right)^{\frac{k}{2}} = \left(\frac{3}{4}\right)^{\frac{k}{2}} = \mathcal{O}(k^{-2}) \quad (k \to \infty).
$$

Next, we consider Example [2.](#page-10-3) In this case, the parameter q depends on $p \in (1,3]$, and hence we denote $q = q(p)$. Moreover, we use the following notation:

$$
s(p) \coloneqq 3 - (1 + q(p))^{\frac{3-p}{p-1}}, \quad t(p) \coloneqq 1 + 2q(p) + (1 + q(p))^{\frac{2}{p-1}}.
$$

For any $x \in (1, 3)$, we define

$$
q_1(x) := \frac{1}{10} (2^{\frac{x-1}{3-x}} - 1) > 0, \t t_1(x) := 1 + 2q_1(x) + (1 + q_1(x))^{\frac{2}{x-1}},
$$

$$
q_2(x) := \frac{1}{20} 2^{\frac{3-x}{x-1}} > 0, \t t_2(x) := 1 + 2q_2(x) + (1 + q_2(x))^{\frac{2}{x-1}}.
$$

Recall that $q(p)$ and $t(p)$ can be represented as follows:

$$
q(p) = \begin{cases} q_1(p) & \text{if } p \in (1,2], \\ q_2(p) & \text{if } p \in (2,3], \end{cases} \qquad t(p) = \begin{cases} t_1(p) & \text{if } p \in (1,2], \\ t_2(p) & \text{if } p \in (2,3]. \end{cases} (A.1)
$$

By the definitions of q_1 and q_2 , we obtain

$$
\frac{d}{dp}q_1(p) = \frac{1}{10(3-p)^2} 2^{\frac{2}{3-p}} \log 2 > 0 \quad \forall p \in (1,2],
$$
\n
$$
\frac{d}{dp}q_2(p) = -\frac{1}{20(p-1)^2} 2^{\frac{2}{p-1}} \log 2 < 0 \quad \forall p \in (2,3].
$$
\n(A.2)

Thus, the first equality of $(A.1)$ implies that q is monotonically increasing for $p \in (1, 2]$ and is monotonically decreasing for $p \in (2, 3]$. Thus, using $\theta = \frac{1}{5}$ yields

$$
0 < q(p) \le q(2) = \frac{1}{10} < \theta < 1 \quad \forall p \in (1, 3]. \tag{A.3}
$$

Since $q(p) = \min\{q_1(p), q_2(p)\} \le q_1(p) = \frac{1}{10}(2^{\frac{p-1}{3-p}} - 1) < 2^{\frac{p-1}{3-p}} - 1$ for $p \in (1,3]$, we have

$$
s(p) = 3 - (1 + q(p))^{\frac{3-p}{p-1}} > 3 - 2 = 1 > 0 \quad \forall p \in (1, 3].
$$
 (A.4)

Noting $\frac{2}{p-1} \ge 1$ and $q(p) > 0$ derives

$$
t(p) = 1 + 2q(p) + (1 + q(p))^{\frac{2}{p-1}} \ge 2 + 3q(p) \ge 1 \quad \forall p \in (1, 3]. \tag{A.5}
$$

Utilizing [\(A.2\)](#page-17-1) implies

$$
\frac{d}{dp}t_1(p) = \left(2 + \frac{2}{p-1}(1+q_1(p))^{\frac{3-p}{p-1}}\right) \frac{d}{dp}q_1(p) > 0 \quad \forall p \in (1,2],
$$

$$
\frac{d}{dp}t_2(p) = \left(2 + \frac{2}{p-1}(1+q_2(p))^{\frac{3-p}{p-1}}\right) \frac{d}{dp}q_2(p) < 0 \quad \forall p \in (2,3].
$$

Hence, the second equality of [\(A.1\)](#page-16-4) derives $t(p) \leq t(2) = \frac{241}{100}$ for $p \in (1,3]$. It then follows from [\(A.5\)](#page-17-2) and $\theta = \frac{1}{5}$ $\frac{1}{5}$ that

$$
0 < \frac{1}{5} \le \theta t(p) \le \frac{241}{500} < 1. \tag{A.6}
$$

Therefore, the assumptions are ensured by $(A.3)$, $(A.4)$, $(A.5)$, and $(A.6)$.

B

This appendix provides the proof of Theorem [2.](#page-14-6)

Proof. We define r_3 and r_4 as follows:

$$
r_3 \coloneqq \min\left\{r_2, \left(\frac{m_3}{3m_2m_4}\right)^{\frac{2}{p-1}}\right\}, \quad r_4 \coloneqq \frac{1}{2+m_2}\min\left\{r_3, \left(\frac{m_3}{3m_2m_4}\right)^{\frac{2}{p-1}}\right\},\
$$

where r_2 , m_2 , m_3 , and m_4 are positive constants described in Lemma [3.](#page-11-2) Assume that the initial point x_0 is selected from $B(x^*, r_4)$, thus it satisfies $||x_0 - x^*|| \leq r_4.$

The proof is divided into two parts: The former part will prepare two inequalities regarding $||d_k||$, and the latter part will prove fast convergence of $\{x_k\}$ using those inequalities. We first show

$$
||d_k|| \le \frac{1}{3}||d_{k-1}||, \quad ||d_k|| \le \frac{r_4m_2}{3^k} \quad \forall k \in \mathbb{N}.
$$
 (B.1)

Let $k \in \mathbb{N}$ be arbitrary. Using (a) and (b) of Lemma [3](#page-11-2) yields

$$
||d_k|| \le m_2 m_4 \text{dist}(x_{k-1}, X^*)^{\frac{p+1}{2}} \le \frac{m_2 m_4}{m_3} r_3^{\frac{p-1}{2}} ||d_{k-1}|| \le \frac{1}{3} ||d_{k-1}||, \quad (B.2)
$$

where note that the last inequality follows from the definition of r_3 . Now, let us show that

$$
x_{\ell} \in B(x^*, r_3) \quad \forall \ell \in \{0, 1, \dots, k\} \implies ||d_{\ell}|| \le \frac{r_4 m_2}{3^{\ell}} \quad \forall \ell \in \{0, 1, \dots, k\}.
$$
\n(B.3)

From [\(B.2\)](#page-18-0) and (a) of Lemma [3,](#page-11-2) we have

$$
||d_0|| \le m_2 \text{dist}(x_0, X^*) \le m_2 ||x_0 - x^*|| \le r_4 m_2,
$$

$$
||d_\ell|| \le \frac{1}{3} ||d_{\ell-1}|| \le \dots \le \frac{1}{3^{\ell}} ||d_0|| \le \frac{r_4 m_2}{3^{\ell}} \quad \forall \ell \in \{1, 2, \dots, k\},
$$

namely, [\(B.3\)](#page-18-1) can be verified.

From now on, we prove by mathematical induction that $x_k \in B(x^*, r_3)$ for all $k \in \mathbb{N}$. Let us consider the case where $k = 1$. Item (a) of Lemma [3](#page-11-2) implies $||x_1 - x^*|| \le ||x_0 - x^*|| + ||d_0|| \le r_4 + m_2 \text{dist}(x_0, X^*) \le r_4(1 + m_2) \le$ r₃. Next, let $k \in \mathbb{N}$ be arbitrary, and we assume that $x_j \in B(x^*, r_3)$ for $j \in \{0, 1, \ldots, k\}$. By [\(B.3\)](#page-18-1) and item (a) of Lemma [3,](#page-11-2) we obtain

$$
||x_{k+1} - x^*|| \le ||x_k - x^*|| + ||d_k|| \le \dots \le ||x_0 - x^*|| + \sum_{\ell=0}^k ||d_\ell||
$$

$$
\le r_4 + \frac{r_4 m_2}{2} \left(1 - \frac{1}{3^{k+1}}\right) \le \frac{2 + m_2}{2} r_4 \le r_3.
$$

Thus, we verify that $x_k \in B(x^*, r_3)$ for $k \in \mathbb{N}$. It then follows from [\(B.3\)](#page-18-1) that $||d_k|| \leq \frac{r_4m_2}{3^k}$ for $k \in \mathbb{N}$, namely, the disired inequalities of [\(B.1\)](#page-18-2) are proven.

The second part shows the local fast convergence of $\{x_k\}$. We arbitrarily

take $i \in \mathbb{N}$ and $j \in \mathbb{N}$ with $i \gg j$. Using [\(B.1\)](#page-18-2) yields

$$
||x_i - x_j|| \le ||x_{i-1} - x_j|| + ||d_{i-1}|| \le ||x_{i-2} - x_j|| + \sum_{\ell=i-2}^{i-1} ||d_{\ell}||
$$

$$
\le \cdots \le \sum_{\ell=j}^{i-1} ||d_{\ell}|| \le r_4 m_2 \sum_{\ell=j}^{i-1} \frac{1}{3^{\ell}} = r_4 m_2 \left(\frac{1}{3^{j-1}} - \frac{1}{3^{i-1}}\right) \le \frac{r_4 m_2}{3^{j-1}}.
$$

This fact implies that $\{x_k\}$ is a Cuachy sequence, that is, there exists $\bar{x} \in \mathbb{R}^n$ such that $x_k \to \bar{x}$ as $k \to \infty$. Meanwhile, it follows from (a) of Lemma [3](#page-11-2) and [\(B.1\)](#page-18-2) that $\{\text{dist}(x_k, X^*)\}$ converges to zero. We note that $\|\widehat{x}_k\| \le$ $\|\widehat{x}_k - x_k\| + \|x_k\| = \text{dist}(x_k, X^*) + \|x_k\|$, namely, $\{\widehat{x}_k\}$ is bounded. Hence, there exist $\widetilde{x} \in X^*$ and $\mathcal{K} \subset \mathbb{N}$ such that $\widehat{x}_k \to \widetilde{x}$ as $\mathcal{K} \ni k \to \infty$. These facts imply that $\|\bar{x}-\tilde{x}\| \leq \|x_k-\bar{x}\|+\text{dist}(x_k, X^*)+\|\hat{x}_k-\tilde{x}\| \to 0$ as $\mathcal{K} \ni k \to \infty$, that is, $\{x_k\}$ converges to some global optimum $\bar{x} = \tilde{x} \in X^*$.

Hereinafter, we show that $\{x_k\}$ converges to \bar{x} superlinearly. Combining items (a) and (b) of Lemma [3](#page-11-2) derives

$$
||d_{j+1}|| \le m_2 \text{dist}(x_{j+1}, X^*) \le m_2 m_4 \text{dist}(x_j, X^*)^{\frac{p+1}{2}} \le \frac{m_2 m_4}{m_3^{\frac{p+1}{2}}} ||d_j||^{\frac{p+1}{2}}.
$$
\n(B.4)

Since the first inequality of [\(B.1\)](#page-18-2) holds, it can be verfied that

$$
||d_{\ell}|| \leq \frac{1}{3^{\ell-j-1}}||d_{j+1}|| \quad \forall \ell \in \{j+1, j+2, \dots, i-1\},
$$

$$
||d_{\ell}|| \leq \frac{1}{3^{\ell-j}}||d_j|| \quad \forall \ell \in \{j+1, j+2, \dots, i-1\}.
$$

By these inequalities, we obtain

$$
||x_{j+1} - x_i|| = \left\| \sum_{\ell=j+1}^{i-1} d_{\ell} \right\| \le \sum_{\ell=j+1}^{i-1} ||d_{\ell}||
$$

$$
\sum_{\ell=j+1}^{i-1} \frac{1}{3^{\ell-j-1}} ||d_{j+1}|| = \frac{3}{2} \left(1 - \frac{1}{3^{i-j-1}} \right) ||d_{j+1}|| \le 2||d_{j+1}||,
$$

and

$$
||x_j - x_i|| = \left\| \sum_{\ell=j}^{i-1} d_{\ell} \right\| \ge ||d_j|| - \sum_{\ell=j+1}^{i-1} ||d_{\ell}||
$$

$$
\ge \left(1 - \sum_{\ell=j+1}^{i-1} \frac{1}{3^{\ell-j}}\right) ||d_j|| = \frac{1}{2} \left(1 + \frac{1}{3^{i-j-1}}\right) ||d_j|| \ge \frac{1}{2} ||d_j||.
$$

Hence taking the limit $i \to \infty$ implies

$$
||x_{j+1} - \bar{x}|| \le 2||d_{j+1}||, \quad ||d_j|| \le 2||x_j - \bar{x}||. \tag{B.5}
$$

Exploiting [\(B.4\)](#page-19-0) and [\(B.5\)](#page-20-0) derives

$$
||x_{j+1}-\bar{x}|| \leq \frac{2^{\frac{p+3}{2}}m_2m_4}{m_3^{\frac{p+1}{2}}}||x_j-\bar{x}||^{\frac{p+1}{2}}.
$$

Therefore, from $\frac{p+1}{2} \in (1,2]$, the sequence $\{x_k\}$ converges to \bar{x} superlinearly. Moreover, if $p = 3$ holds, then $\frac{p+1}{2} = 2$, that is, the rate of convergence is quadratic.