HYPERBOLIC ENTROPY FOR HARMONIC MEASURES ON SINGULAR HOLOMORPHIC FOLIATIONS

FRANÇOIS BACHER

ABSTRACT. Let $\mathscr{F}=(M,\mathscr{L},E)$ be a Brody-hyperbolic singular holomorphic foliation on a compact complex manifold M. Suppose that \mathscr{F} has isolated singularities and that its Poincaré metric is complete. This is the case for a very large class of singularities, namely, non-degenerate and saddle-nodes in dimension 2. Let μ be an ergodic harmonic measure on \mathscr{F} . We show that the upper and lower local hyperbolic entropies of μ are leafwise constant almost everywhere. Moreover, we show that the entropy of μ is at least 2.

1. Introduction

The dynamical theory for laminations by Riemann surfaces has recently received much attention. In particular, a lot of progress have been focused on the case of hyperbolic leaves. For singular holomorphic foliations on \mathbb{P}^n , this is a typical setup. Indeed, every polynomial vector field on \mathbb{C}^n induces a singular holomorphic foliation that can be compactified onto \mathbb{P}^n . This foliation is always singular. For $d, n \in \mathbb{N}$ with $n \geq 2$, denote by $\mathscr{F}_d(\mathbb{P}^n)$ the space of singular holomorphic foliations of degree d on \mathbb{P}^n . As is shown by Lins Neto and Soares [15], a generic foliation $\mathscr{F} \in \mathscr{F}_d(\mathbb{P}^n)$ has only non-degenerate singularities. This result is based on a previous analysis of the Jouanolou foliations [12]. By the results of Lins Neto [13, 14] and Glutsyuk [11], such a foliation is hyperbolic if $d \geq 2$ and even Brody-hyperbolic in the sense of [7]. In the case n = 2, Nguyên also uses the integrability of the holonomy cocycle [19] to compute the Lyapunov exponent [20] of a generic foliation in $\mathscr{F}_d(\mathbb{P}^2)$. We refer the reader to the survey articles [8, 9, 17, 18] for more details about these questions.

Solving heat equations with respect to a harmonic current, Dinh, Nguyên and Sibony prove in [5] ergodic theorems for laminations. In particular, they obtain an effective and geometric analog of Birkhoff's Theorem. Somehow, the comparison relies on considering the time to be the hyperbolic distance in a uniformization of a leaf by the Poincaré disk \mathbb{D} . Therefore, in a series of two articles [6, 7], the three authors study the modulus of continuity of the Poincaré metric of compact laminations and of singular holomorphic foliations with linearizable singularities. They also introduce various notions of entropy for hyperbolic laminations. These concepts, together with Lyapunov exponents [16], could bring a fruitful bridge between the dynamics of foliations and the dynamics of maps. Our works [1, 2, 3] are devoted to generalize some of their results about the

Université de Lille, Laboratoire de mathématiques Paul Painlevé, CNRS U.M.R. 8524, 59655 Villeneuve d'Ascq Cedex, France.

E-mail address: francois.bacher@univ-lille.fr.

Date: June 17, 2024.

Key words and phrases. Singular holomorphic foliation; Hyperbolic entropy; Ergodic Theory; Poincaré metric; Harmonic measures.

Poincaré metric, the heat diffusions and the topological entropy to foliations with nondegenerate singularities. Here, we are interested in the measure-theoretic hyperbolic entropy.

Let us introduce some notations to state our results. Let $\mathscr{F}=(M,\mathscr{L},E)$ be a singular holomorphic foliation. Here, M is a compact complex manifold, \mathscr{L} is a foliated atlas of $M\backslash E$ and E is the singular set of \mathscr{F} . Suppose that the leaves of \mathscr{F} are hyperbolic Riemman surfaces and for $x\in M\backslash E$, fix $\phi_x\colon \mathbb{D}\to L_x$ a uniformization of the leaf L_x . Let g_M be a Hermitian metric on M and denote by d the induced distance on M. Since we consider the hyperbolic distance in \mathbb{D} to be a time parametrizing L_x ,

$$d_R(x,y) = \inf_{\theta \in \mathbb{R}} \sup_{\xi \in \overline{\mathbb{D}}_R} d(\phi_x(\xi), \phi_y(e^{i\theta}\xi)), \quad x, y \in M \backslash E,$$

for R>0, can be thought as a *Bowen distance* up to time R [6, pp. 581–582]. Here, \mathbb{D}_R denotes the disk of center 0 and hyperbolic radius R. The infimum over θ means that we consider distances between the two closest parametrizations of the leaves L_x and L_y . That is, we consider the distance up to reparametrization. Therefore, we obtain a notion of Bowen balls and of *topological entropy* $h(\mathscr{F})$, which is a number that measures heuristically the exponential growth with R of the amount of Bowen balls needed to cover $M\backslash E$. The more precise definition of $h(\mathscr{F})$ will be given in Section 2. There are some cases for which it is known the topological entropy is finite. Namely, for smooth compact laminations without singularities [6] and foliations with only non-degenerate singularities on compact complex surfaces [7, 3]. Here, we show a lower bound for the entropy.

If moreover \mathscr{F} is endowed with a harmonic measure μ , one can ask more weakly how many balls are needed to cover a subset of measure at least $1-\delta$, for $\delta \to 0$, and we obtain a measure-theoretic entropy $h(\mu)$. It is clear that $h(\mu) \leqslant h(\mathscr{F})$. To study these numbers, Dinh, Nguyên and Sibony [6] introduce other notions of entropy, and in particular local upper and lower entropies $h^{\pm}(\mu,x)$, that measure the exponential decay with R of the μ -measure of the Bowen balls centered in x. In the case without singularities, they can show the following.

Theorem 1.1 (Dinh–Nguyên–Sibony [6, Theorem 4.2]). Let $\mathscr{F}=(M,\mathscr{L})$ be a smooth compact lamination by hyperbolic Riemann surfaces and μ be a harmonic measure on \mathscr{F} . Then, the local upper and lower entropies $x\mapsto h^\pm(\mu,x)$ are leafwise constant. In particular, if μ is ergodic, they are constant μ -almost everywhere and we denote them by $h^\pm(\mu)$. Moreover, $h(\mu) \geqslant 2$.

Here, we improve their results for singular holomorphic foliations.

Theorem 1.2. Let $\mathscr{F}=(M,\mathscr{L},E)$ be a Brody-hyperbolic singular holomorphic foliation on a compact complex manifold M. Let μ be an ergodic harmonic measure on \mathscr{F} . Suppose that the singularities of \mathscr{F} are isolated and that the Poincaré metric of \mathscr{F} is complete. Then, $x\mapsto h^\pm(\mu,x)$ are constant μ -almost everywhere. Moreover, $h(\mu)\geqslant 2$.

As we show in Section 2, if \mathscr{F} is Brody-hyperbolic, the assumption on the Poincaré metric is only local near the singularities. Therefore, by local works of Canille Martins and Lins Neto [4], our result can be applied to a large class of singularities.

Corollary 1.3. Let $\mathscr{F} = (M, \mathscr{L}, E)$ be a Brody-hyperbolic singular holomorphic foliation on a compact complex manifold M. Let μ be an ergodic harmonic measure on \mathscr{F} . Suppose that all the singularities $p \in E$ of \mathscr{F} are of one of the following type.

- (1) In a chart centered at p, \mathscr{F} is generated by a vector field $X = \sum_{j=k}^{+\infty} X_j$, with the X_j homogeneous vector fields of degree j, for $j \ge k$, and X_k admitting an isolated singularity. In particular, if k = 1, p is a non-degenerate singularity.
- (2) The point p is a saddle-node singularity of \mathscr{F} (in ambient dimension 2).

Then, the conclusions of Theorem 1.2 hold.

Let us explain the method of our proof. First, we follow the idea of Dinh, Nguyên and Sibony of considering transversal entropies $\tilde{h}^{\pm}(\mu, x)$. The main advantage is that it is not so difficult to show that they are leafwise constant using the quasi-invariance of μ . As in the non-singular case, we reduce our problem to proving $h^{\pm} = h^{\pm} + 2$. Somehow, this means showing that the plaquewise entropy is always 2. That is, we have to determine the size of the intersection of a Bowen ball with a plaque. In the non-singular case, estimates on the distance between the identity and $\tau \in \operatorname{Aut}(\mathbb{D})$ on \mathbb{D}_R show that this intersection is essentially a disk of radius $\simeq e^{-R}$. Indeed, if there is a $\xi \in \mathbb{D}_R$ such that $d_P(\xi, \tau(\xi))$ is big, then there are intermediate points where it is not so big to be in the same plaque, and still bigger than ε . In that case, the Poincaré distance and the distance in M are comparable. This simple argument does not work anymore in our context because of the singularities. When the above distance is not so big, we could be near the singularity. In that case, we can not conclude anything. On the other hand, when we are outside neighbourhoods of the singularities, the Poincaré distance between two points of the same leaf could be big and the distance in M very small if they do not belong to the same plaque.

Let us describe how we overcome this difficulty. First, we use Dinh, Nguyên and Sibony's Birkhoff type Theorem [5] to control the time spent near the singularities. Next, we prove thorough estimates on the distance between the identity and an automorphism τ of $\mathbb D$ and find a large amount of points where the distances between them is big enough but not too big. Then, this enables to show that the distances between two points of the intersection of a Bowen ball with a plaque correspond in some sense to large subsets of the disk which are spent near the singular set. Since this time is small, this should not happen often. More precisely, we show somehow that this intersection contains at most $e^{\delta R}$ disks of radius e^{-R} , for δ comparable to the proportion of time spent near the singular set. Since this δ can be chosen arbitrarily small, we have our result. The last argument relies on studying what we call *subsets of* \mathbb{R}^n *with prescribed steps*. That is, subsets for which it is known that the distance between any two points is in some small union of intervals. These look like what we obtain when constructing the Cantor ternary set by induction.

The article is organized as follows. In Section 2, we begin by recalling some basic facts about the Poincaré metric. We show that Corollary 1.3 is indeed a consequence of Theorem 1.2. Then, we define the different entropies we shall use. We recall some results of the three authors to reduce to studying the plaquewise entropy and showing Proposition 2.7. In Section 3, we prove this proposition. We begin by reducing to two lemmas that will allow to work in one single plaque. Next, we study briefly subsets of \mathbb{R}^n

with prescribed steps. We finish by introducing a fine geometric setup and proving the two lemmas.

Notation. Throughout this paper, we denote by \mathbb{D} the unit disk of \mathbb{C} , and $r\mathbb{D}$ the open disk of radius $r \in \mathbb{R}_+^*$ for the standard Euclidean metric of \mathbb{C} . For $R \in \mathbb{R}_+^*$, we also denote by \mathbb{D}_R the open disk of hyperbolic radius R in \mathbb{D} , so that $\mathbb{D}_R = r\mathbb{D}$ with $r = \frac{e^R - 1}{e^R + 1}$, or if $r \in [0, 1)$, with $R = \ln \frac{1+r}{1-r}$.

If $\mathscr{F}=(M,\mathscr{L},E)$ is a singular holomorphic foliation and $x\in M\backslash E$, we denote by L_x the leaf of \mathscr{F} through x. Moreover, if L_x is hyperbolic, we denote by $\phi_x\colon \mathbb{D}\to L_x$ a uniformization of L_x such that $\phi_x(0)=x$. Given a Hermitian metric g_M on M, we denote by d the distance induced by g_M . Similarly, we denote by d_P the Poincaré distance if one is given a Poincaré metric denoted g_P on the disk or on a leaf. We are also given Bowen distances d_R on $M\backslash E$, and denote by $B_R(x,\varepsilon)$ the ball of center x and radius ε for the distance d_R .

Finally, we denote by Leb(B) the Lebesgue measure of a Borel set $B \subset \mathbb{R}^n$. We use C, C', C'', etc. to denote positive constants which may change from a line to another.

Acknowledgments. The author is supported by the Labex CEMPI (ANR-11-LABX-0007-01) and by the project QuaSiDy (ANR-21-CE40-0016).

2. Hyperbolic entropy for foliations

2.1. Leafwise Poincaré metric. In all this section, we let $\mathscr{F} = (M, \mathscr{L}, E)$ be a singular holomorphic foliation on a compact complex manifold M. Suppose that M is endowed with a Hermitian metric g_M and for $x \in M \setminus E$, consider

$$(2.1) \eta(x) = \sup \left\{ \|\alpha'(0)\|_{g_M}; \ \alpha \colon \mathbb{D} \to L_x \text{ holomorphic such that } \alpha(0) = x \right\}.$$

Above, $\|v\|_{g_M}$ is the norm of a vector $v \in T_x L_x$ with respect to the Hermitian metric g_M . That is, $\|v\|_{g_M} = (g_{M,x}(v,v))^{1/2}$. The map η was introduced by Verjovsky in [21]. It is designed to satisfy the following facts.

Proposition 2.1. (1) For $x \in M \setminus E$, $\eta(x) < +\infty$ if and only if the leaf L_x is hyperbolic, that is, it is uniformized by the Poincaré disk \mathbb{D} .

- (2) If L_x is hyperbolic, we have $\eta(x) = \|\phi'(0)\|_{g_M}$, where $\phi \colon \mathbb{D} \to L_x$ is any uniformization of L_x such that $\phi(0) = x$.
- (3) If L_x is hyperbolic, then $\frac{4g_M}{\eta^2}$ induces the Poincaré metric on L_x .

Here, we are interested in the case of hyperbolic leaves. We need to specify our global setting. Let us introduce some terminology, for which we follow [7, Definition 3.1].

Definition 2.2. If all the leaves of \mathscr{F} are hyperbolic, we say that \mathscr{F} is *hyperbolic*. If moreover there exists a constant $c_0 > 0$ such that $\eta(x) < c_0$ for all $x \in M \setminus E$, we say that \mathscr{F} is *Brody-hyperbolic*.

We say the Poincaré metric of \mathscr{F} is complete if \mathscr{F} is hyperbolic and $\frac{4g_M}{\eta^2}$ is complete as a Hermitian metric on $M \setminus E$. We refer the reader to [4, Property **P.2.**] for more details.

As mentioned before, there are cases when we know the Poincaré metric is complete.

Theorem 2.3 (Canille Martins–Lins Neto [4, Theorems 1 and 2]). Let \mathscr{F} be a local singular holomorphic foliation near $0 \in \mathbb{C}^n$, with singularity at 0 of any type listed in Corollary 1.3. Then, the Poincaré metric of \mathscr{F} is complete at 0.

Therefore, Corollary 1.3 is a consequence of the following, the proof of which is essentially the same as [7, Proposition 3.3].

Lemma 2.4. Let $\mathscr{F} = (M, \mathscr{L}, E)$ be a Brody-hyperbolic singular holomorphic foliation on a compact complex manifold. Suppose that all $a \in E$ admit a neighbourhood U_a on which the Poincaré metric of $\mathscr{F}|_{U_a}$ is complete at a. Then, the Poincaré metric of \mathscr{F} is complete.

Proof. Let η_a denote the η map of the foliation $\mathscr{F}|_{U_a}$. We will show that there exists a constant $C_a>0$ such that $\eta\leqslant C_a\eta_a$ on some neighbourhood $V_a\subset U_a$ of a. Since M is compact and \mathscr{F} is Brody-hyperbolic, it is not difficult to see that it implies the result. Let $\rho>0$ be such that the ball $B(a,\rho)$ is contained in U_a and consider $V_a=B\left(a,\frac{\rho}{2}\right)$. Fix $x\in V_a$ and $\phi_x\colon \mathbb{D}\to L_x$ a uniformization of L_x such that $\phi_x(0)=x$. That way, $\eta(x)=\|\phi_x'(0)\|_{g_M}$. Since \mathscr{F} is Brody-hyperbolic, there exists a radius $r_0\in(0,1)$, independent on x, such that $\phi_x(r_0\mathbb{D})\subset U_a$. More precisely, the hyperbolic radius $R_0=\ln\frac{1+r_0}{1-r_0}$ corresponding to r_0 can be chosen to be equal to $\frac{\rho}{c_0}$, for $c_0>0$ as in Definition 2.2. By the extremal condition (2.1) of η_a , it follows that $\eta_a(x)\geqslant r_0\|\phi_x'(0)\|_{g_M}=r_0\eta(x)$.

2.2. Bowen distance and various notions of entropy. From now on, suppose that \mathscr{F} is hyperbolic. Also, denote by d the distance on M induced by the ambient Hermitian metric g_M . For $x \in M \setminus E$, fix $\phi_x \colon \mathbb{D} \to L_x$ a uniformization of L_x such that $\phi_x(0) = x$. The idea of Dinh, Nguyên and Sibony [6, pp. 581–582] is to consider the Poincaré distance in \mathbb{D} to be a canonical time. More precisely, for $R \geqslant 0$, consider the Bowen distance

$$d_R(x,y) = \inf_{\theta \in \mathbb{R}} \sup_{\xi \in \overline{\mathbb{D}}_R} d(\phi_x(\xi), \phi_y(e^{i\theta}\xi)), \quad x, y \in M \backslash E.$$

It measures the distance between the orbits of x and y up to time R. It is clear that it is independent on the choice of ϕ_x . This enables us to define various notions of entropy. For $x \in M \setminus E$, $R \geqslant 0$ and $\varepsilon > 0$, denote by $B_R(x,\varepsilon) = \{y \in M \; ; \; d_R(x,y) < \varepsilon\}$ the Bowen ball of radius ε and center x up to time R. For $Y \subset M \setminus E$, $R \geqslant 0$, $\varepsilon > 0$ and $F \subset Y$, we say that F is (R,ε) -dense in Y if $Y \subset \cup_{x \in F} B_R(x,\varepsilon)$. Denote by $N(Y,R,\varepsilon)$ the minimal cardinality of an (R,ε) -dense subset in Y. The topological hyperbolic entropy of Y is defined as

$$h(Y) = \sup_{\varepsilon > 0} \limsup_{R \to +\infty} \frac{1}{R} \log N(Y, R, \varepsilon).$$

For $Y = M \setminus E$, we denote it by $h(\mathscr{F})$. If M is compact, it is not difficult to prove that it does not depend on the choice of g_M . A similar and equivalent definition can be made with maximal (R, ε) -separated sets, but we do not need it. The interested reader can see [6, Section 3] for more details.

Now, we introduce some entropies for harmonic measures following [6, Section 4]. Basically, harmonic measures are quasi-invariant measures by the foliation \mathscr{F} . Here, we only use (2.2) below, which can be seen as a local expression of this quasi-invariance. That is, a harmonic measure can be locally disintegrated into a transverse Radon measure times a leafwise harmonic density. We refer the reader to [5, 9, 10, 18] for a definition and a link with dd^c -closed currents.

To define measure-theoretic entropy, the idea is to consider analogous quantities as those emerging from Brin–Katok Theorem. Let μ be a harmonic probability measure on \mathscr{F} . All our harmonic measures will be supposed to have total mass 1. For $R \ge 0$, $\varepsilon > 0$ and $\delta \in (0,1)$, let $N(R,\varepsilon,\delta)$ be the smallest integer N such that there exists $x_1,\ldots,x_N\in M\backslash E$ with $\mu\left(\bigcup_{i=1}^N B_R\left(x_i,\varepsilon\right)\right)\ge 1-\delta$. Define the *hyperbolic entropy of* μ as

$$h(\mu) = \sup_{\delta \in (0,1)} \sup_{\varepsilon > 0} \limsup_{R \to +\infty} \frac{1}{R} \log N(R, \varepsilon, \delta).$$

To study this quantity, Dinh, Nguyên and Sibony consider *local upper and lower hyperbolic* entropies $h^{\pm}(\mu, x)$, for $x \in M \setminus E$. Namely,

$$h^{+}(\mu, x, \varepsilon) = \limsup_{R \to +\infty} -\frac{1}{R} \log \mu \left(B_{R}(x, \varepsilon) \right), \qquad h^{+}(\mu, x) = \sup_{\varepsilon > 0} h^{+}(\mu, x, \varepsilon);$$
$$h^{-}(\mu, x, \varepsilon) = \liminf_{R \to +\infty} -\frac{1}{R} \log \mu \left(B_{R}(x, \varepsilon) \right), \qquad h^{-}(\mu, x) = \sup_{\varepsilon > 0} h^{-}(\mu, x, \varepsilon).$$

In the context of a non-singular foliation, they prove that these are leafwise constant (see Theorem 1.1). This enables them to obtain a link between all these entropies.

Theorem 2.5 (Dinh–Nguyên–Sibony [6, Proposition 4.5]). Let $\mathscr{F} = (M, \mathscr{L})$ be a (non-singular) hyperbolic holomorphic foliation on a complex manifold M (we do not assume that M is compact). Let μ be a harmonic measure on \mathscr{F} . If the quantities $x \mapsto h^{\pm}(\mu, x)$ are constant μ -almost everywhere, then,

$$h^{-}(\mu) \leqslant h(\mu) \leqslant h^{+}(\mu) \leqslant h(\mathscr{F}).$$

Under these hypotheses, the above theorem is in fact implicitly proved, for the authors state it in the setup of Theorem 1.1. However, they only use that $h^{\pm}(\mu,x)$ are constant. Let us define another notion of entropy that they introduce to prove their results.

2.3. Local transversal entropy. Let $x \in M \setminus E$. If $\varepsilon > 0$ is sufficiently small, then there exists a flow box $U \simeq \mathbb{D} \times \mathbb{T}$ such that $B(x, \varepsilon) \subset U$. Denote by $\pi_{\mathbb{T}} \colon U \to \mathbb{T}$ the projection on the second coordinate. Consider a disintegration of the measure μ in U. That is,

(2.2)
$$\mu = \int_{\mathbb{T}} \left(\int_{\mathbb{D}} f_t(z) g_P(z, t) \right) d\nu(t),$$

where ν is a finite Radon measure on \mathbb{T} and the f_t , for $t \in \mathbb{T}$, are harmonic functions on \mathbb{D} with $f_t(0) = 1$. Such a decomposition is unique up to changing the f_t on a ν -negligible set and exists due to [5, Propositions 2.3 and 5.1]. Define

$$\widetilde{h}^{+}(\mu, x, \varepsilon) = \limsup_{R \to +\infty} -\frac{1}{R} \log \nu \left(\pi_{\mathbb{T}} \left(B_{R}(x, \varepsilon) \right) \right), \qquad \widetilde{h}^{+}(\mu, x) = \sup_{\varepsilon > 0} \widetilde{h}^{+}(\mu, x, \varepsilon);$$

$$\widetilde{h}^{-}(\mu, x, \varepsilon) = \liminf_{R \to +\infty} -\frac{1}{R} \log \nu \left(\pi_{\mathbb{T}} \left(B_{R}(x, \varepsilon) \right) \right), \qquad \widetilde{h}^{-}(\mu, x) = \sup_{\varepsilon > 0} \widetilde{h}^{-}(\mu, x, \varepsilon).$$

It is not difficult to show that $\widetilde{h}^{\pm}(\mu,x)$ do not depend on the choice of the flow box U. The following is implicitly proved by the three authors. They state it with more restrictive hypotheses but their proof still works without changing anything.

Theorem 2.6 (Dinh–Nguyên–Sibony [6, Theorem 4.2]). Let $\mathscr{F} = (M, \mathscr{L})$ be a Brody-hyperbolic holomorphic foliation on a complex manifold M and μ be a harmonic measure

on \mathscr{F} . Then, the quantities $x \mapsto \widetilde{h}^{\pm}(\mu, x)$ are leafwise constant. In particular, if μ is ergodic, they are constant μ -almost everywhere and we denote them by $\widetilde{h}^{\pm}(\mu)$.

Note that there is no compactness assumption on M. Therefore, we can apply it to $(M \setminus E, \mathcal{L})$ in the singular case. Now, Theorem 1.2 wil be a consequence of Theorems 2.5 and 2.6, combined with the following, the proof of which occupies the next section and needs some detours.

Proposition 2.7. Let $\mathscr{F}=(M,\mathscr{L},E)$ be a Brody-hyperbolic singular holomorphic foliation on a compact complex manifold M. Let also μ be an ergodic harmonic measure on \mathscr{F} . Suppose that the singularities of \mathscr{F} are isolated and that the Poincaré metric of \mathscr{F} is complete in the sense of Definition 2.2. Then, for μ -almost every $x \in M \setminus E$,

$$h^{\pm}(\mu, x) = \tilde{h}^{\pm}(\mu, x) + 2.$$

In particular, $x \mapsto h^{\pm}(\mu, x)$ are constant μ -almost everywhere and $h^{-}(\mu) \geqslant 2$.

3. Proof of Proposition 2.7

3.1. **First reduction.** Let $\mathscr{F}=(M,\mathscr{L},E)$ be a Brody-hyperbolic singular holomorphic foliation on a compact complex manifold M and μ be an ergodic harmonic measure on \mathscr{F} . Suppose that the singularities of \mathscr{F} are isolated and that the Poincaré metric of \mathscr{F} is complete. First, we intend to reduce the proof to two lemmas, which together will give estimates to the measure of plaquewise Bowen balls. More precisely, we estimate the quantity of automorphisms τ of the disk, such that $\phi \circ \tau$ is close to ϕ , for some given ϕ uniformization of a leaf. These estimates will take some time to be proven, but we already show how to obtain Proposition 2.7 from them.

Lemma 3.1. For $\delta > 0$ and μ -almost every $x \in M \setminus E$, there exists $\varepsilon_0 > 0$ satisfying the following. For $\varepsilon > 0$ sufficiently small, there exists C > 0 such that for all sufficiently large R > 0 and $y \in B_R(x, \varepsilon_0)$,

$$C^{-1}e^{-R} \leq \text{Leb}\left(\left\{\theta \in [-\pi, \pi]; \ d_{\overline{\mathbb{D}}_R}(\phi_y \circ r_\theta, \phi_y) < \varepsilon\right\}\right) \leq Ce^{-(1-\delta)R}.$$

Here, $r_{\theta} \in Aut(\mathbb{D})$ *denotes the rotation of angle* θ .

We need a similar result for a general automorphism in $\operatorname{Aut}(\mathbb{D})$. For $\zeta \in \mathbb{D}$, denote by $\tau_{\zeta} \in \operatorname{Aut}(\mathbb{D})$ defined by $\tau_{\zeta}(\xi) = \frac{\xi + \zeta}{1 + \overline{\zeta} \xi}$. We need some preparation.

Lemma 3.2. Given $x \in M \setminus E$, there exist flow boxes $\mathbb{D} \times \mathbb{T} \simeq U \subset V$ containing x and $r_0 \in (0,1)$ such that

$$U \subset \bigcup_{t \in \mathbb{T}} \phi_t \left(\frac{r_0}{3} \mathbb{D} \right) \quad \textit{and} \quad \bigcup_{y \in U} \phi_y(r_0 \mathbb{D}) \subset V.$$

Proof. First fix a flow box V containing x. Since η is bounded above, we can find $r_0 \in (0,1)$ and U sufficiently small such that the map

$$r_0 \mathbb{D} \times U \to M, \qquad (\zeta, y) \mapsto \phi_y(\zeta)$$

is injective and with values in V. Now, since η is bounded from below near x, shrinking U if necessary, we can suppose $U \subset \bigcup_{t \in \mathbb{T}} \phi_t \left(\frac{r_0}{3} \mathbb{D} \right)$.

Lemma 3.3. For $\delta > 0$ and μ -almost every $x \in M \setminus E$, there exists $\varepsilon_0 > 0$ satisfying the following. For $\varepsilon > 0$ sufficiently small, there exists C > 0 such that for all sufficiently large R > 0 and $y \in B_R(x, \varepsilon_0)$,

$$C^{-1}e^{-3R} \leqslant \operatorname{Leb}\left(\left\{ (\zeta,\theta) \in r_0 \mathbb{D} \times [-\pi,\pi] \; ; \; d_{\overline{\mathbb{D}}_R}(\phi_y \circ \tau_\zeta \circ r_\theta,\phi_y) < \varepsilon \right\} \right) \leqslant Ce^{-3(1-\delta)R}.$$

Here, r_0 is given by the previous lemma.

Proof of Proposition 2.7. We take for granted Lemmas 3.1 and 3.3 and show how they give us our result. Consider $x \in M \setminus E$ satisfying the conclusions of the lemmas and fix $\delta > 0$. Let $r_0 \in (0,1)$, $\mathbb{D} \times \mathbb{T} \simeq U \subset V$ be given by Lemma 3.2 and $0 < \varepsilon < \frac{\varepsilon_0}{3}$ sufficiently small to have the conclusions of both other lemmas. Using (2.2), we get

$$\mu(B_R(x,\varepsilon)) = \int_{\mathbb{T}} \left(\int_{\mathbb{D}} f_t(z) \chi_{B_R(x,\varepsilon)}(z,t) g_P(z,t) \right) d\nu(t),$$

where χ_B denotes the characteristic function of a Borel set B. Now, since $x \in M \backslash E$ is still far from the singular set, g_P is equivalent to the Lebesgue measure and by the Harnack inequality, there exists c > 0 such that $c^{-1} \leqslant f_t \leqslant c$ on U. Therefore, there exists c' > 1 with

$$c'^{-1}\mu(B_R(x,\varepsilon)) \leqslant \int_{\pi_{\mathbb{T}}(B_R(x,\varepsilon))} \operatorname{Leb}\left(B_R(x,\varepsilon) \cap (\mathbb{D} \times \{t\})\right) d\nu(t) \leqslant c'\mu(B_R(x,\varepsilon)).$$

Here, the Lebesgue measure is the one of $\mathbb D$ in the flow box U. For $t \in \pi_{\mathbb T}(B_R(x,\varepsilon))$, choose $y_t \in B_R(x,\varepsilon) \cap (\mathbb D \times \{t\})$. Then, $B_R(x,\varepsilon) \cap (\mathbb D \times \{t\}) \subset B_R(y_t,2\varepsilon) \cap (\mathbb D \times \{t\})$. Moreover, for $y_t \in B_R\left(x,\frac{\varepsilon}{2}\right)$, $B_R\left(y_t,\frac{\varepsilon}{2}\right) \cap (\mathbb D \times \{t\}) \subset B_R(x,\varepsilon) \cap (\mathbb D \times \{t\})$. Hence,

(3.1)
$$\mu(B_R(x,\varepsilon)) \leqslant c' \nu(\pi_{\mathbb{T}}(B_R(x,\varepsilon))) \sup_{t \in \pi_{\mathbb{T}}(B_R(x,\varepsilon))} \text{Leb}(B_R(y_t,2\varepsilon) \cap (\mathbb{D} \times \{t\}));$$

$$\mu(B_R(x,\varepsilon)) \geqslant c'^{-1} \nu\left(\pi_{\mathbb{T}}\left(B_R\left(x,\varepsilon/2\right)\right)\right) \inf_{t \in \pi_{\mathbb{T}}(B_R(x,\varepsilon/2))} \text{Leb}\left(B_R\left(y_t,\varepsilon/2\right) \cap (\mathbb{D} \times \{t\})\right).$$

So, changing ε if necessary, it will be sufficient to bound above and below the Lebesgue measure of $B_R(y,\varepsilon) \cap (\mathbb{D} \times \{t\})$, for $y \in B_R(x,\varepsilon_0) \cap (\mathbb{D} \times \{t\})$. Let $y' \in B_R(y,\varepsilon) \cap (\mathbb{D} \times \{t\})$. Since y and y' are in the same plaque, there is $\zeta \in r_0\mathbb{D}$ such that $y' = \phi_y(\zeta)$. Thus,

$$B_R(y,\varepsilon) \cap (\mathbb{D} \times \{t\}) = \left\{ \phi_y(\zeta); \zeta \in r_0 \mathbb{D}, \exists \theta \in [-\pi, \pi], d_{\overline{\mathbb{D}}_R}(\phi_y \circ \tau_\zeta \circ r_\theta, \phi_y) < \varepsilon \right\}.$$

Denote by

$$A_{\varepsilon}(y) = \left\{ \zeta \in r_{0} \mathbb{D}, \exists \theta \in [-\pi, \pi], d_{\overline{\mathbb{D}}_{R}}(\phi_{y} \circ \tau_{\zeta} \circ r_{\theta}, \phi_{y}) < \varepsilon \right\},$$

$$B_{\varepsilon}(y) = \left\{ (\zeta, \theta) \in r_{0} \mathbb{D} \times [-\pi, \pi], d_{\overline{\mathbb{D}}_{R}}(\phi_{y} \circ \tau_{\zeta} \circ r_{\theta}, \phi_{y}) < \varepsilon \right\},$$

$$C_{\varepsilon}(y') = \left\{ \theta \in [-\pi, \pi], d_{\overline{\mathbb{D}}_{R}}(\phi_{y'} \circ r_{\theta}, \phi_{y'}) < \varepsilon \right\}, \quad y' \in B_{R}(y, \varepsilon) \cap (\mathbb{D} \times \{t\}).$$

Since the Poincaré metric in the flow box is equivalent to the Lebesgue measure, we have

(3.2)
$$c^{-1} \operatorname{Leb} (A_{\varepsilon}(y)) \leq \operatorname{Leb} (B_R(y, \varepsilon) \cap (\mathbb{D} \times \{t\})) \leq c \operatorname{Leb} (A_{\varepsilon}(y)).$$

Moreover,

$$\operatorname{Leb}(B_{\varepsilon}(y)) \leqslant \operatorname{Leb}(A_{\varepsilon}(y)) \sup_{\xi \in \pi_{\zeta}(B_{\varepsilon}(y))} \operatorname{Leb}(C_{2\varepsilon}(\phi_{y}(\xi))),$$

$$\operatorname{Leb}(B_{2\varepsilon}(y)) \geqslant \operatorname{Leb}(A_{\varepsilon}(y)) \inf_{\xi \in \pi_{\zeta}(B_{\varepsilon}(y))} \operatorname{Leb}(C_{\varepsilon}(\phi_{y}(\xi))),$$

where π_{ζ} is the projection on the disk from $\mathbb{D} \times [-\pi, \pi]$. Now, applying Lemmas 3.1 and 3.3 and coming back to (3.2), we get

$$C^{-1}e^{-(2+\delta)R} \leq \text{Leb}\left(B_R(y,\varepsilon) \cap (\mathbb{D} \times \{t\})\right) \leq Ce^{-(2-3\delta)R}$$
.

By (3.1), we obtain

$$(c'C)^{-1}\mu(B_R(x,\varepsilon))e^{(2-3\delta)R} \leqslant \nu(\pi_{\mathbb{T}}(B_R(x,\varepsilon))) \leqslant c'Ce^{(2+\delta)R}\mu(B_R(x,\varepsilon)).$$

Letting R go to infinity and noting that ε can be chosen arbitrarily small, this implies

$$h^{\pm}(\mu, x) - 2 - \delta \leqslant \widetilde{h}^{\pm}(\mu, x) \leqslant h^{\pm}(\mu, x) - 2 + 3\delta.$$

Since δ was chosen arbitrarily, we conclude the proof.

3.2. **Subsets of** \mathbb{R}^n **with prescribed steps.** This subsection is devoted to studying briefly subsets of \mathbb{R}^n with distances in a pre-defined set. As will appear in the next subsection, the sets appearing in Lemmas 3.1 and 3.3 will be of that kind (for n = 1 or 3) and we are especially interested in estimating their Lebesgue measure. The sets we study look like the union of intervals we obtain when constructing Cantor sets by induction. Let us be more precise.

Definition 3.4. Let $N \in \mathbb{N}$ and for $i \in [0, N]$, $j \in \{1, 2\}$, take $\varepsilon_{i,j} \ge 0$. Suppose that

$$0<\varepsilon_{i+1,2}<\varepsilon_{i,1}<\varepsilon_{i,2}, \qquad i\in \llbracket 0,N-1\rrbracket\,,$$

and $\varepsilon_{N,1}=0$. A Borel subset $A\subset\mathbb{R}^n$ is said to have *prescribed* $(\varepsilon_{i,j})_{i,j}$ -steps if for any $x,y\in A$, there exists $i\in[0,N]$ such that $||x-y||\in[\varepsilon_{i,1},\varepsilon_{i,2}]$.

In particular, it should be noted that $diam(A) \leq \varepsilon_{0,2}$. The following gives an upper bound to the Lebesgue measure of those sets. The bound is loose but will be sufficient.

Lemma 3.5. Let A be a Borel subset of \mathbb{R}^n with prescribed $(\varepsilon_{i,j})_{i \in \llbracket 0,N \rrbracket, j \in \{1,2\}}$ -steps. Suppose that for $i \in \llbracket 1,N \rrbracket$, $\varepsilon_{i,2} < \frac{\varepsilon_{i-1,1}}{2}$. Then, if $V_n = \operatorname{Vol}(B(0,1))$ in \mathbb{R}^n ,

Leb(A)
$$\leq V_n \varepsilon_{N,2}^n 3^{nN} \times \prod_{i=0}^{N-1} \left(\frac{\varepsilon_{i,2}}{\varepsilon_{i,1}}\right)^n$$
.

Proof. Let us show by induction on N that such an A is contained in the union of at most $p_N=3^{nN}\times\prod_{i=0}^{N-1}\left(\frac{\varepsilon_{i,2}}{\varepsilon_{i,1}}\right)^n$ balls of radius $\varepsilon_{N,2}$. If N=0, this is trivial. Suppose that this was proven at rank N-1. In particular, each of the at most p_{N-1} balls of radius $\varepsilon_{N-1,2}$ has prescribed $(\varepsilon_{i,j})_{i\in\{N-1,N\},j\in\{1,2\}}$ -steps. Here, we use that $\varepsilon_{N-1,2}<\frac{\varepsilon_{N-2,1}}{2}$. So, this is enough to prove it for N=1. This means that for $x,y\in A$, either $\|x-y\|\leqslant \varepsilon_{1,2}$, or $\|x-y\|\in [\varepsilon_{0,1},\varepsilon_{0,2}]$. Let x_1,\ldots,x_r be a maximal $\varepsilon_{1,2}$ -separated subset of A. Since the x_i are $\varepsilon_{1,2}$ -separated, $\|x_i-x_j\|>\varepsilon_{1,2}$ for $i\neq j$ and the prescribed steps imply that $\|x_i-x_j\|\geqslant \varepsilon_{0,1}$. Therefore, the balls $\left(B\left(x_i,\frac{\varepsilon_{0,1}}{2}\right)\right)_{i\in [\![1,r]\!]}$ are disjoint. Since $\mathrm{diam}(A)\leqslant \varepsilon_{0,2}$, they are also all included in $B\left(x_1,\varepsilon_{0,2}+\frac{\varepsilon_{0,1}}{2}\right)$. Hence,

$$\operatorname{Vol}\left(B\left(x_{1},\varepsilon_{0,2}+\frac{\varepsilon_{0,1}}{2}\right)\right)=V_{n}\left(\varepsilon_{0,2}+\frac{\varepsilon_{0,1}}{2}\right)^{n}\geqslant\sum_{i=1}^{r}\operatorname{Vol}\left(B\left(x_{i},\frac{\varepsilon_{0,1}}{2}\right)\right)=rV_{n}\left(\frac{\varepsilon_{0,1}}{2}\right)^{n}.$$

It follows that $r \leqslant \left(\frac{2\varepsilon_{0,2}+\varepsilon_{0,1}}{\varepsilon_{0,1}}\right)^n \leqslant \left(3\frac{\varepsilon_{0,2}}{\varepsilon_{0,1}}\right)^n$. Moreover, by maximality of the x_1,\ldots,x_r , $A \subset \bigcup_{i=1}^r B(x_i,\varepsilon_{1,2})$. The induction is complete.

3.3. **Geometric setup.** In this subsection, we prepare some geometric and dynamical ground. In particular, we need Dinh, Nguyên and Sibony's Birkhoff-type Theorem. Let $\mathscr{F} = (M, \mathscr{L}, E)$ be a Brody-hyperbolic singular holomorphic foliation on a compact complex manifold. Suppose that \mathscr{F} has isolated singularities. For $x \in M \setminus E$, fix $\phi_x \colon \mathbb{D} \to L_x$ a uniformization of L_x . For $r \in (0,1)$ and the corresponding $R = \ln \frac{1+r}{1-r}$, define

$$m_{x,R} = \frac{1}{M_R} (\phi_x)_* \left(\log^+ \frac{r}{|\zeta|} g_P \right),$$

where \log^+ stands for $\max(\log, 0)$, g_P denotes the Poincaré metric on the disk and

$$M_R = \int_{\mathbb{D}} \log^+ \frac{r}{|\zeta|} g_P = -2\pi \log(1 - r^2) \sim_{R \to +\infty} 2\pi R.$$

Theorem 3.6 (Dinh–Nguyên–Sibony [5, Theorem 7.1] (see also [18, Theorem 5.36])). Keep the above notations and hypotheses. Let μ be an ergodic harmonic measure on \mathscr{F} . Then, for μ -almost every $x \in M \setminus E$, $m_{x,R}$ converges weakly to μ when R tends to infinity.

Now, fix an $x \in M \setminus E$ such that $m_{x,R} \to \mu$. We use this convergence to control the time that L_x spends near the singular set. More precisely, we need the following.

Lemma 3.7. Suppose moreover that the Poincaré metric of \mathscr{F} is complete. Fix $\delta > 0$. There exist $0 < \rho_1 < \rho_2 < \rho_3$ satisfying the following. Denote by $U_i^{\text{sing}} = \{y \in M \setminus E; d(y, E) \leq \rho_i\}$, for $j \in \{1, 2, 3\}$.

- (i) If $y \in M \setminus E$ and $j \in \{1,2\}$ are such that $d(y,E) \leq \rho_j$ and $y' \in L_y$ is such that $d(y',E)\geqslant
 ho_{j+1}$, then $d_P(y,y')\geqslant 4$.

 (ii) $\mu\left(U_3^{\mathrm{sing}}\right)<\frac{\delta}{2}$. In particular, for all large enough R>0, $m_{x,R}\left(U_3^{\mathrm{sing}}\right)<\delta$.

 (iii) $x\notin U_3^{\mathrm{sing}}$.

Proof. Choose first $\rho_3 > 0$ satisfying (ii) and (iii). Clearly we can do so since $\mu(E) = 0$ and $m_{x,R} \to \mu$. Then, since the Poincaré metric is complete, we can choose $\rho_2 \in (0, \rho_3)$ and after that $\rho_1 \in (0, \rho_2)$ satisfying (i).

Lemma 3.8. With the notations of Lemma 3.7, there exist c > 1 and $\varepsilon_0 > 0$ such that the following holds. Let $y \notin \frac{1}{2}U_1^{\text{sing}}$, $y' \in L_y$ be such that $d_P(y, y') \leqslant \varepsilon_0$. Then,

$$c^{-1}d_P(y,y') \leqslant d(y,y') \leqslant cd_P(y,y').$$

Above, $\frac{1}{2}U_1^{\text{sing}} = \{y \in M \setminus E; \ d(y, E) \leqslant \frac{\rho_1}{2}\}.$

Proof. Given ρ_1 , we can cover $M \setminus \frac{1}{4} U_1^{\text{sing}}$ by a finite number of flow boxes. It follows that η is bounded by $c^{-1} < \eta < c$ on $M \setminus \frac{1}{2} U_1^{\text{sing}}$. Moreover, we find an $\varepsilon_0 > 0$ such that $y \notin \frac{1}{4}U_1^{\text{sing}}$ implies that $B(y, c\varepsilon_0)$ is contained in a flow box. For y and y' satisfying the above conditions, we get

$$\frac{c\varepsilon_0}{2} \geqslant \frac{c}{2} d_P(y, y') \geqslant d_{L_y}(y, y'),$$

where d_{L_y} denotes the distance on L_y induced by the restriction of g_M to L_y . This implies that y and y' are in the same plaque and the bounds on η give us both needed inequalities.

The following describes circles that are in a large part mapped to the singular set. These $R_{i,j}$ will prove to give prescribed steps of the sets involved in Lemmas 3.1 and 3.3. **Lemma 3.9.** With the notations of Lemma 3.7, take R > 0 sufficiently large. Denote by

$$I_j^{\text{sing}} = \left\{ R' \in [0, R] ; \text{ Leb } \left\{ \xi \in \partial \mathbb{D}_{R'}; \phi_x(\xi) \in U_j^{\text{sing}} \right\} > \frac{\pi}{3} \right\}, \quad j \in \{1, 2, 3\}.$$

Here, Leb of the circles $\partial \mathbb{D}_{R'}$ are normalized to have mass 2π . There exists $(R_{i,j})_{i \in \llbracket 0,N \rrbracket, j \in \{1,2\}}$, such that $4 < R_{i,2} < R_{i,1} < R_{i+1,2}$, $i \in \llbracket 0,N-1 \rrbracket$, $R \geqslant R_{N,2}$ and $R_{N,1} = +\infty$, with

- (1) $R_{i,1} R_{i,2} \geqslant 4$, $R_{i+1,2} R_{i,1} \geqslant 4$, for $i \in [0, N-1]$.
- (2) $I_1^{\text{sing}} \subset \bigcup_{i=0}^N (R_{i,2}, R_{i,1}).$
- (3) $[0,R] \cap \left(\cup_{i=0}^{N} (R_{i,2},R_{i,1}) \right) \subset I_3^{\text{sing}}$.

Moreover, $\sum_{i=0}^{N-1} R_{i,1} - R_{i,2} + (R - R_{N,2}) \leq 12\delta R$ and $N \leq 3\delta R$.

Big R will correspond to small ε , that is why the order is reversed compared to sets with prescribed steps.

Proof. This is where we use our weird conditions on ρ_1, ρ_2, ρ_3 . The $I_j^{\rm sing}$ are open subsets of [0,R]. Hence, they are an at most countable union of intervals. For each C_3 connected component of $I_3^{\rm sing}$, if $C_3 \cap I_1^{\rm sing} \neq \emptyset$, consider I_{C_3} to be the convex hull of the union of all C_2 connected components of $I_2^{\rm sing} \cap C_3$. Define I to be the union of the I_{C_3} , for all the C_3 . Since each I_{C_3} contains a point of $I_1^{\rm sing}$ and all points of $I_2^{\rm sing}$ in C_3 , condition (i) implies that the length of I_{C_3} is at least 4. Indeed, points in the part of the circle of hyperbolic radius $R_1 \in I_1^{\rm sing}$ which are in $U_1^{\rm sing}$ must go out of $U_2^{\rm sing}$ so that we leave I_{C_3} . Moreover, since we work component by component in $I_3^{\rm sing}$, the same condition (i) implies that the I_{C_3} are pairwise distant of at least 4. Indeed, points in the part of the circle of hyperbolic radius $R_2 \in I_{C_3}$ which are in $U_2^{\rm sing}$ must go out of $U_3^{\rm sing}$ so that we enter a new I_{C_3} . If we denote the I_{C_3} by $(R_{i,2},R_{i,1})$, points (1), (2) and (3) are then clear. To have $R_{0,2} > 4$, we use (iii). The bounds on $R_{N,j}$ are just technical for further notations. Here actually, it is only asked that $R_{N,2} - R_{N-1,1} \geqslant 4$. If this is not satisfied for $R_{N,2} = R$, then we can just put $R_{N-1,1} = +\infty$ and still get all points of the lemma.

For the estimate $\sum_{i=0}^{N-1} R_{i,1} - R_{i,2} + R - R_{N,2} \le 12\delta R$, we need the following computation.

Lemma 3.10. Let $0 < r_2 < r_1 < r$ and $R_j = \ln \frac{1+r_j}{1-r_j}$ (or $r_j = \frac{e^{R_j}-1}{e^{R_j}+1}$), $j \in \{1,2\}$ be the corresponding hyperbolic radii. Then

$$\int_{r_2}^{r_1} \log \left(\frac{r}{\rho}\right) \frac{4\rho}{(1-\rho^2)^2} d\rho = R_1 - R_2 + o(1),$$

where the o(1) stands for R_2 goes to infinity.

End of proof of Lemma 3.9. Taking for granted the above estimate, let us finish our proof. If R is sufficiently large, we have $m_{x,R}\left(U_3^{\mathrm{sing}}\right)<\delta$ by (ii). We get

$$M_R \delta > \int_{\mathbb{D}} \chi_{\left\{\phi_x \in U_3^{\text{sing}}\right\}}(\xi) \log^+ \left(\frac{r}{|\xi|}\right) g_P(\xi),$$

where as usual, χ_B is the characteristic function of B. Let us continue our computation.

$$M_{R}\delta > \int_{0}^{r} \left(\int_{0}^{2\pi} \chi_{\left\{\phi_{x} \in U_{3}^{\text{sing}}\right\}} (\rho e^{i\theta}) d\theta \right) \log \left(\frac{r}{\rho}\right) \frac{4\rho d\rho}{(1-\rho^{2})^{2}}$$

$$M_{R}\delta > \frac{\pi}{3} \sum_{i=0}^{N-1} \int_{r_{i,2}}^{r_{i,1}} \log \left(\frac{r}{\rho}\right) \frac{4\rho d\rho}{(1-\rho^{2})^{2}} + \frac{\pi}{3} \int_{r_{N,2}}^{r} \log \left(\frac{r}{\rho}\right) \frac{4\rho d\rho}{(1-\rho^{2})^{2}}$$

$$M_{R}\delta > \frac{\pi}{3} \sum_{i=0}^{N-1} (R_{i,1} - R_{i,2}) + \frac{\pi}{3} (R - R_{N,2}) + o(R).$$

Here, we have denoted $r_{i,j} = \frac{e^{R_{i,j}}-1}{e^{R_{i,j}}+1}$ as usual and we have used (3). For the last inequality, we applied Lemma 3.10 and Cesàro Theorem. Note that (1) implies $R_{i,2} > 8i + 4$, which goes to infinity with $i \leq N = O(R)$. Now, since $M_R \sim 2\pi R$, we get our result. \square

Proof of Lemma 3.10. This is only a silly computation. Integrating by parts,

$$\int_{r_2}^{r_1} \log \left(\frac{r}{\rho}\right) \frac{4\rho}{(1-\rho^2)^2} d\rho = 2\left[\frac{1}{1-\rho^2} \log \left(\frac{r}{\rho}\right)\right]_{r_2}^{r_1} + 2\int_{r_2}^{r_1} \frac{d\rho}{\rho(1-\rho^2)}.$$

Since $\frac{1}{\rho(1-\rho^2)} = \frac{1}{2} \left(\frac{1}{1-\rho} - \frac{1}{1+\rho} + \frac{2}{\rho} \right)$, we obtain

$$\int_{r_2}^{r_1} \log \left(\frac{r}{\rho}\right) \frac{4\rho}{\left(1-\rho^2\right)^2} d\rho = 2\left[\frac{\rho^2}{1-\rho^2} \log \left(\frac{r}{\rho}\right)\right]_{r_2}^{r_1} - \log(1-r_1^2) + \log(1-r_2^2).$$

Now, we translate into hyperbolic language. For $j \in \{1,2\}$, $\frac{r_j^2}{1-r_j^2} = \frac{1}{4}e^{R_j} + o\left(e^{R_j}\right)$. Moreover, $\log \frac{r}{r_j} = 2e^{-R_j} - 2e^{-R} + o\left(e^{-R_j}\right)$. So the first bracket above is o(1). The other terms satisfy $1 - r_j^2 = e^{-R_j}(4 + o(1))$. We get the estimate wanted.

3.4. **End of proof of Lemmas 3.1 and 3.3.** Our preparation is soon to be over and we move on to the end of our proof. We begin by Lemma 3.1, for it involves less computation and already explains the ideas of the proof of Lemma 3.3. First, let us study the distance to identity of a rotation in $\operatorname{Aut}(\mathbb{D})$.

Lemma 3.11. Let $0 < \varepsilon_1 < \varepsilon_2$ be sufficiently small with $\varepsilon_1 \leq \frac{1}{16}\varepsilon_2$, $\theta \in [-\pi, \pi]$ and $\xi \in \mathbb{D}$. Then, if $|\xi| \geq \frac{1}{4}$,

$$d_P\left(\xi, e^{i\theta}\xi\right) \in \left[\varepsilon_1, \varepsilon_2\right], \quad \text{for} \quad 8\varepsilon_2^{-1} \left|\sin(\theta/2)\right| \leqslant 1 - \left|\xi\right|^2 \leqslant \frac{1}{2}\varepsilon_1^{-1} \left|\sin(\theta/2)\right|.$$

Proof. Denote by $\tanh^{-1}(x) = \frac{1}{2} \ln \frac{1+x}{1-x}$, $x \in (-1,1)$ and compute.

$$d_{P}\left(\xi, e^{i\theta}\xi\right) = 2 \tanh^{-1} \left| \frac{\xi(1 - e^{i\theta})}{1 - e^{i\theta} |\xi|^{2}} \right| = 2 \tanh^{-1} \left| \frac{2\xi \sin(\theta/2)}{\cos(\theta/2) \left(1 - |\xi|^{2}\right) - i \sin(\theta/2) \left(1 + |\xi|^{2}\right)} \right|$$

$$= 2 \tanh^{-1} \left(\frac{2 |\xi| |\sin(\theta/2)|}{\left(\left(1 - |\xi|^{2}\right)^{2} + 4 |\xi|^{2} \sin^{2}(\theta/2)\right)^{1/2}} \right) = f\left(\frac{1 - |\xi|^{2}}{2 |\xi| |\sin(\theta/2)|}\right),$$

for $f : \mathbb{R}_+^* \to \mathbb{R}_+^*$ defined by $f(x) = 2 \tanh^{-1} \left(\frac{1}{\sqrt{1+x^2}}\right)$. The function f is strictly decreasing and has inverse $f^{-1} : y \mapsto \sqrt{\frac{1}{\tanh^2(y/2)} - 1} = \frac{2}{y} + O_{y \to 0}(1)$, as is shown by a straightforward

computation. It follows, that if $\varepsilon_1, \varepsilon_2$ are sufficiently small, and if $x \in [4\varepsilon_2^{-1}, \varepsilon_1^{-1}]$, then $x \in [f^{-1}(\varepsilon_2), f^{-1}(\varepsilon_1)]$ and $f(x) \in [\varepsilon_1, \varepsilon_2]$. The bounds on $1 - |\xi|^2$ exactly give

$$\frac{1-\left|\xi\right|^{2}}{2\left|\xi\right|\left|\sin(\theta/2)\right|} \in \left[4\varepsilon_{2}^{-1}, \varepsilon_{1}^{-1}\right],$$

for $|\xi| \leq \frac{1}{4}$, so we get our result.

Lemma 3.12. Take the notations of Lemmas 3.8 and 3.9 and let $\varepsilon > 0$ be sufficiently small, R > 0 be sufficiently large. Define $\varepsilon_{i,1} = 2\arcsin\left(8c\varepsilon e^{-R_{i,1}}\right)$ and $\varepsilon_{i,2} = 2\arcsin\left(16c\varepsilon e^{-R_{i,2}}\right)$, for $i \in [0, N]$. Note that $\varepsilon_{N,1} = 0$. Then, for $y \in B_R(x, \varepsilon_0)$,

$$A = \left\{ \theta \in \left[-\pi, \pi \right]; \ d_{\overline{\mathbb{D}}_R}(\phi_y \circ r_\theta, \phi_y) < \varepsilon \right\}$$

has prescribed $(\varepsilon_{i,j})_{i,j}$ -steps.

Proof. Let $y \in B_R(x, \varepsilon_0)$, ε be sufficiently small to determine and $\theta_1, \theta_2 \in A$. We have to show that $|\theta_1 - \theta_2| \in \bigcup_{i=0}^N [\varepsilon_{i,1}, \varepsilon_{i,2}]$. Since $\theta_1, \theta_2 \in A$, we have

$$(3.3) d_{\overline{\mathbb{D}}_R}(\phi_y \circ r_{\theta_1}, \phi_y \circ r_{\theta_2}) = d_{\overline{\mathbb{D}}_R}(\phi_y \circ r_{|\theta_1 - \theta_2|}, \phi_y) < 2\varepsilon.$$

Denote by $\theta=|\theta_1-\theta_2|$. With the notations of Lemma 3.8, set $\varepsilon_2<\varepsilon_0,\ \varepsilon_1=\frac{1}{16}\varepsilon_0$ sufficiently small, and $\varepsilon=\frac{1}{2}c^{-1}\varepsilon_1$. Fix ϕ_y a uniformization of L_y with $d_{\overline{\mathbb{D}}_R}(\phi_x,\phi_y)<\varepsilon_0$. By definition, it is clear that $\varepsilon_0<\frac{\rho_1}{2}$. Hence, if $\xi\in\overline{\mathbb{D}}_R$ satisfies $d(\phi_x(\xi),E)>\rho_1$, then $d(\phi_y(\xi),E)>\frac{\rho_1}{2}$. Take $\xi\in\mathbb{D}$ such that

$$1 - |\xi|^2 = 8\varepsilon_2^{-1} |\sin(\theta/2)| = \frac{1}{2}\varepsilon_1^{-1} |\sin(\theta/2)|.$$

Since y is still far from the singular set, it is clear by Lemma 3.8 that $A \subset [-C\varepsilon, C\varepsilon]$, for some constant C > 1. Shrinking ε if necessary, we obtain that $|\xi| \ge \frac{1}{4}$. By Lemma 3.11,

$$2c\varepsilon = \varepsilon_1 \leqslant d_P\left(\xi, e^{i\theta}\xi\right) \leqslant \varepsilon_2 < \varepsilon_0.$$

Hence, (3.3) together with Lemma 3.8 imply that either $d_P(0,\xi) > R$, or $\phi_y(\xi) \in \frac{1}{2}U_1^{\text{sing}}$. This should hold for every ξ with same modulus, so we deduce by Lemma 3.9(2) that $d_P(0,\xi) \in \bigcup_{i=0}^N (R_{i,1},R_{i,2})$. Therefore, $1-|\xi|^2 \in \bigcup_{i=0}^N \left(2e^{-R_{i,2}},4e^{-R_{i,1}}\right)$ and

$$\sin(\theta/2) \in \bigcup_{i=0}^{N} \left(8c\varepsilon e^{-R_{i,1}}, 16c\varepsilon e^{-R_{i,2}}\right).$$

This concludes the proof.

End of proof of Lemma 3.1. For the upper bound, we apply Lemmas 3.12 and 3.5. Note that $\varepsilon_{i,2} < \frac{\varepsilon_{i-1,1}}{2}$, because $R_{i,2} - R_{i-1,1} > 4$ in Lemma 3.9. We obtain

Leb(A)
$$\leq 2\varepsilon_{N,2}3^N \prod_{i=0}^{N-1} \frac{\varepsilon_{i,2}}{\varepsilon_{i,1}} \leq 32c\varepsilon e^{-R}e^{R-R_{N,2}}12^N \prod_{i=0}^{N-1} e^{R_{i,1}-R_{i,2}}.$$

The last statement of Lemma 3.9 now gives

$$Leb(A) \leq 32c\varepsilon e^{\delta(3\log(12)+12)R}e^{-R}$$
.

Since δ was chosen arbitrarily, we deduce the upper bound wanted. For the lower bound, we will use notations of Lemma 3.11. Recall also that $c_0 > 0$ is such that $\eta \leqslant c_0$ on $M \setminus E$. If θ satisfies $|\sin(\theta/2)| \leqslant c_0^{-1} \frac{\varepsilon}{4} e^{-R}$ and $\xi \in \overline{\mathbb{D}}_R$, then $\frac{1-|\xi|^2}{2|\xi||\sin(\theta/2)|} \geqslant \frac{4c_0}{\varepsilon}$. Thus,

$$d_P\left(\xi, e^{i\theta}\xi\right) = f\left(\frac{1 - |\xi|^2}{2|\xi| |\sin(\theta/2)|}\right) \leqslant f\left(\frac{4}{c_0\varepsilon}\right) \leqslant c_0^{-1}\varepsilon,$$

where f is defined in Lemma 3.11. Since ξ is arbitrary in $\overline{\mathbb{D}}_R$, we get that $\theta \in A$. It follows that $\operatorname{Leb}(A) \geqslant c_0^{-1} \frac{\varepsilon}{2} e^{-R}$. Such an estimate can also be thought as Lemma 3.11 for $\varepsilon_2 = c_0^{-1} \varepsilon$ and $\varepsilon_1 = 0$.

We argue similarly for Lemma 3.3, with analogous steps. But first, let us see how to compose the automorphisms of the disk.

Lemma 3.13. Let $\zeta_1, \zeta_2 \in \operatorname{Aut}(\mathbb{D})$ and $\theta_1, \theta_2 \in \mathbb{R}$. If ζ_1, ζ_2 are sufficiently small, there exist $\zeta \in \mathbb{D}$ and $\theta \in \mathbb{R}$, with

$$\frac{1}{2} \| (\zeta_1 - \zeta_2, \theta_1 - \theta_2) \|_{\infty} \le \| (\zeta, \theta) \|_{\infty} \le 3 \| (\zeta_1 - \zeta_2, \theta_1 - \theta_2) \|_{\infty},$$

satisfying

$$(\tau_{\zeta_1} \circ r_{\theta_1})^{-1} \circ (\tau_{\zeta_2} \circ r_{\theta_2}) = \tau_{\zeta} \circ r_{\theta}.$$

Here, $\|(\zeta, \theta)\|_{\infty} = \max(|\zeta|, |\theta|)$.

Proof. This is just computation, that we partly leave to the reader. We have the following composition rules.

$$r_{\theta} \circ \tau_{\zeta} = \tau_{e^{i\theta}\zeta} \circ r_{\theta}, \qquad \tau_{\zeta_1} \circ \tau_{\zeta_2} = r_{\theta(\zeta_1,\zeta_2)} \circ \tau_{\zeta(\zeta_1,\zeta_2)},$$

for $\zeta(\zeta_1,\zeta_2)=\tau_{\zeta_2}(\zeta_1)$ and $\theta(\zeta_1,\zeta_2)$ satisfying $e^{i\theta(\zeta_1,\zeta_2)}=\frac{1+\zeta_1\overline{\zeta_2}}{1+\overline{\zeta_1}\zeta_2}$. Playing with these rules, we get the form wanted, with $|\zeta|=|\tau_{\zeta_2}(-\zeta_1)|$. So indeed, $d_P(0,\zeta)=d_P(\zeta_1,\zeta_2)$ and $2|\zeta_1-\zeta_2|\leqslant |\zeta|\leqslant 3|\zeta_1-\zeta_2|$. Moreover, $\theta=\theta_2-\theta_1+\theta(-\zeta_1,\zeta_2)$. Therefore, we just need to show that $|\theta(-\zeta_1,\zeta_2)|\leqslant |\zeta_1-\zeta_2|$. We have

$$\left|1 - e^{i\theta(-\zeta_1,\zeta_2)}\right| = \left|\frac{2i\Im\left(\zeta_1\overline{\zeta_2}\right)}{1 - \overline{\zeta_1}\zeta_2}\right| = \left|\frac{2\Im\left((\zeta_1 - \zeta_2)\overline{\zeta_2}\right)}{1 - \overline{\zeta_1}\zeta_2}\right| \leqslant \frac{1}{2}\left|\zeta_1 - \zeta_2\right|,$$

if ζ_1 and ζ_2 are sufficiently small. Above, $\Im(z)$ denotes the imaginary part of a complex number z. We get the estimate wanted.

Lemma 3.14. There exist constants $C_1, C_2 > 1$ satisfying the following. Let $0 < \varepsilon_1 < \varepsilon_2$ be sufficiently small with $\varepsilon_1 \leq (C_1C_2)^{-1}\varepsilon_2$, $\theta \in [-\pi, \pi]$ and $\zeta \in r_0\mathbb{D}$. There exists a subset $\Lambda \subset [-\pi, \pi]$, with $\text{Leb}(\Lambda) = \frac{\pi}{2}$, such that if $\xi = \rho e^{i\lambda}$, $\rho \geqslant \frac{1}{4}$, $\lambda \in \Lambda$ and

$$C_2 \varepsilon_2^{-1} \| (\zeta, \theta) \|_{\infty} \leq 1 - |\xi|^2 \leq C_1^{-1} \varepsilon_1^{-1} \| (\zeta, \theta) \|_{\infty},$$

then $d_P\left(\xi, \tau_{\zeta}\left(e^{i\theta}\xi\right)\right) \in [\varepsilon_1, \varepsilon_2].$

Proof. First, let us compute a formula for the distance

$$d_{P}\left(\xi, \tau_{\zeta}\left(e^{i\theta}\xi\right)\right) = 2 \tanh^{-1} \left| \left(\xi - \frac{\zeta + e^{i\theta}\xi}{1 + \overline{\zeta}e^{i\theta}\xi}\right) \left(1 - \overline{\xi} \frac{\zeta + e^{i\theta}\xi}{1 + \overline{\zeta}e^{i\theta}\xi}\right)^{-1} \right|$$

$$= 2 \tanh^{-1} \left| \frac{\xi(1 - e^{i\theta}) - \zeta + \overline{\zeta}e^{i\theta}\xi^{2}}{1 - e^{i\theta} + e^{i\theta}\left(1 - |\xi|^{2}\right) + \overline{\zeta}e^{i\theta}\xi - \zeta\overline{\xi}} \right|$$

$$= 2 \tanh^{-1} \left| \frac{2i\xi \sin(\theta/2) + \zeta e^{-i\theta/2} - \overline{\zeta}e^{i\theta/2}\xi^{2}}{2i\left(\sin(\theta/2) + \Im\left(\zeta\overline{\xi}e^{-i\theta/2}\right)\right) - e^{i\theta/2}\left(1 - |\xi|^{2}\right)} \right|.$$

Now, take $\xi = \sigma i \frac{\zeta}{|\zeta|} e^{-i\theta/2} \rho e^{i\lambda}$, with $\sigma = \pm 1$ chosen such that $\sigma \sin(\theta/2) = -|\sin(\theta/2)|$. If $\zeta = 0$, then any complex number of modulus 1 can replace $\frac{\zeta}{|\zeta|}$. The above formula becomes

$$d_P\left(\xi, \tau_{\zeta}\left(e^{i\theta}\xi\right)\right) = 2\tanh^{-1}\left|\frac{2\rho e^{i\lambda}\left|\sin(\theta/2)\right| + \left|\zeta\right|\left(1 + \rho^2 e^{2i\lambda}\right)}{2i\left(\sin(\theta/2) - \sigma\rho\left|\zeta\right|\cos(\lambda)\right) - e^{i\theta/2}\left(1 - \rho^2\right)}\right|.$$

Next, suppose that $|\lambda| \leqslant \frac{\pi}{4}$. This gives indeed a subset Λ of angles, of Lebesgue measure $\frac{\pi}{2}$. It should be noted that the three terms in the numerator have non-negative real part. Moreover, if ρ satisfies the conditions of the lemma with ε_1 and ε_2 sufficiently small, it is clear that the term $1-\rho^2$ in the denominator dominates all the others. Therefore,

$$2\tanh^{-1}\left(\frac{\sqrt{2}/(4\pi)\|(\zeta,\theta)\|_{\infty}}{2C_{1}^{-1}\varepsilon_{1}^{-1}\|(\zeta,\theta)\|_{\infty}}\right) \leqslant d_{P}\left(\xi,\tau_{\zeta}\left(e^{i\theta}\xi\right)\right) \leqslant 2\tanh^{-1}\left(\frac{3\|(\zeta,\theta)\|_{\infty}}{C_{2}\varepsilon_{2}^{-1}/2\|(\zeta,\theta)\|_{\infty}}\right)$$
$$\varepsilon_{1} = \frac{\sqrt{2}}{8\pi}C_{1}\varepsilon_{1} \leqslant d_{P}\left(\xi,\tau_{\zeta}\left(e^{i\theta}\xi\right)\right) \leqslant 24C_{2}^{-1}\varepsilon_{2} = \varepsilon_{2},$$

for $\varepsilon_1, \varepsilon_2$ sufficiently small. Here we have defined the constants C_1, C_2 to get the extreme identities. The lemma is proven.

Lemma 3.15. Take the notations of Lemmas 3.8 and 3.9 and let $\varepsilon > 0$ be sufficiently small, R > 0 be sufficiently large. Define $\varepsilon_{i,1} = \frac{4}{3}cC_1\varepsilon e^{-R_{i,1}}$ and $\varepsilon_{i,2} = 16cC_1\varepsilon e^{-R_{i,2}}$, for $i \in [0, N]$. Note that $\varepsilon_{N,1} = 0$. Then, for $y \in B_R(x, \varepsilon_0)$,

$$A = \left\{ (\zeta, \theta) \in r_0 \mathbb{D} \times [-\pi, \pi] \; ; \; d_{\overline{\mathbb{D}}_R} (\phi_y \circ \tau_\zeta \circ r_\theta, \phi_y) < \varepsilon \right\}$$

has prescribed $(\varepsilon_{i,j})_{i,j}$ -steps.

Proof. We argue similarly to Lemma 3.12. Fix $\varepsilon_2 < \varepsilon_0$ sufficiently small, $\varepsilon_1 = (C_1 C_2)^{-1} \varepsilon_2$ and $\varepsilon = (2c)^{-1} \varepsilon_1$. Take $(\zeta_1, \theta_1), (\zeta_2, \theta_2) \in A$ and (ζ, θ) given by Lemma 3.13. That way,

$$2\varepsilon > d_{P,\overline{\mathbb{D}}_R}(\phi_y \circ \tau_{\zeta_1} \circ r_{\theta_1}, \phi_y \circ \tau_{\zeta_2} \circ r_{\theta_2}) \geqslant d_{P,\overline{\mathbb{D}}_{R-3|\zeta_1|}}(\phi_y \circ \tau_{\zeta} \circ r_{\theta}, \phi_y).$$

Finally, take $\xi = \rho e^{i\lambda}$, with $\lambda \in \Lambda$ given by Lemma 3.14 and

$$1 - \rho^2 = C_1^{-1} \varepsilon_1^{-1} \| (\zeta, \theta) \|_{\infty} = C_2 \varepsilon_2^{-1} \| (\zeta, \theta) \|_{\infty}.$$

As in Lemma 3.12, because $\operatorname{Leb}(\Lambda) = \frac{\pi}{2}$, we obtain $1 - \rho^2 \in \cup_{i=0}^N \left(2e^{-R_i,1}, 4e^{-R_{i,2}}\right)$. Here, we work with Lemma 3.9 for R-3 $|\zeta_1|$ but this does not change anything to the estimates. This gives $\|(\zeta,\theta)\|_{\infty} \in \left(4cC_1\varepsilon e^{-R_{i,1}}, 8c\varepsilon C_1e^{-R_{i,2}}\right)$. Lemma 3.13 then implies

$$\|(\zeta_1 - \zeta_2, \theta_1 - \theta_2)\|_{\infty} \in \bigcup_{i=0}^{N} \left(\frac{4}{3}cC_1\varepsilon e^{-R_{i,1}}, 16cC_1\varepsilon e^{-R_{i,2}}\right). \quad \Box$$

End of proof of Lemma 3.3. The proof is essentially the same as of Lemma 3.1. Note that $\varepsilon_{i,2} < \frac{\varepsilon_{i-1,1}}{2}$ because $R_{i,2} - R_{i-1,1} > 4$. By Lemmas 3.5 and 3.15, there exist universal constants C, C' such that the Lebesgue measure of the Borel set involved is lower than $C\varepsilon^3 e^{(C'\delta-3)R}$. For the lower bound, the computation of Lemma 3.14 easily shows that for $\|(\zeta,\theta)\|_{\infty} \leqslant C\varepsilon e^{-R}$ and $\xi \in \overline{\mathbb{D}}_R$, $d_P\left(\xi,\tau_{\zeta}\left(e^{i\theta}\xi\right)\right) \leqslant c_0^{-1}\varepsilon$. We conclude the same using that \mathscr{F} is Brody-hyperbolic.

REFERENCES

- [1] Bacher, François: Poincaré metric of holomorphic foliations with non-degenerate singularities. *Internat. J. Math.* **34** (2023), no.10, Paper No. 2350059.
- [2] Bacher, François: Heat diffusions on holomorphic foliations with non-degenerate singularities. *J. Geom. Anal.* **34**, 31 (2024). https://doi.org/10.1007/s12220-023-01485-6.
- [3] Bacher, François: Finiteness of hyperbolic entropy for holomorphic foliations with non-degenerate singularities. arXiv:2311.17236, (2023).
- [4] Canille Martins, Júlio Cesar; Lins Neto, Alcides: Hermitian metrics inducing the Poincaré metric, in the leaves of a singular holomorphic foliation by curves. *Trans. Amer. Math. Soc.* **356** (2004), no. 7, 2963–2988.
- [5] Dinh, Tien-Cuong; Nguyên, Viêt-Anh; Sibony, Nessim: Heat equation and ergodic theorems for Riemann surface laminations. *Math. Ann.* **354** (2012), no. 1, 331–376.
- [6] Dinh, Tien-Cuong; Nguyên, Viêt-Anh; Sibony, Nessim: Entropy for hyperbolic Riemann surface laminations I. *Frontiers in Complex Dynamics: a volume in honor of John Milnor's 80th birthday,* (A. Bonifant, M. Lyubich, S. Sutherland, editors), 569–592, (2014), Princeton University Press.
- [7] Dinh, Tien-Cuong; Nguyên, Viêt-Anh; Sibony, Nessim: Entropy for hyperbolic Riemann surface laminations II. *Frontiers in Complex Dynamics: a volume in honor of John Milnor's 80th birthday,* (A. Bonifant, M. Lyubich, S. Sutherland, editors), 593–622, (2014), Princeton University Press.
- [8] Dinh, Tien-Cuong; Sibony, Nessim: Some open problems on holomorphic foliation theory. *Acta Math. Vietnam.* **45** (2020), no. 1, 103–112.
- [9] Fornæss, John Erik; Sibony, Nessim: Riemann surface laminations with singularities. *J. Geom. Anal.* **18** (2008), no. 2, 400–442.
- [10] Garnett, Lucy: Foliations, the ergodic theorem and Brownian motion. *J. Functional Analysis* **51** (1983), no. 3, 285–311.
- [11] Glutsyuk, Alexey: Hyperbolicity of the leaves of a generic one-dimensional holomorphic foliation on a nonsingular projective algebraic variety. (Russian) *Tr. Mat. Inst. Steklova* **213** (1997), Differ. Uravn. s Veshchestv. i Kompleks. Vrem., 90–111; *translation in Proc. Steklov Inst. Math.* 1996, **213**(2), 83–103.
- [12] Jouanolou, Jean-Pierre: Équations de Pfaff algébriques. (French) [Algebraic Pfaffian equations] Lecture Notes in Mathematics, **708**. *Springer, Berlin,* 1979. v+255 pp.
- [13] Lins Neto, Alcides: Simultaneous uniformization for the leaves of projective foliations by curves. *Bol. Soc. Brasil. Mat. (N.S.)* **25** (1994), no.2, 181–206.
- [14] Lins Neto, Alcides: Uniformization and the Poincaré metric on the leaves of a foliation by curves. *Bol. Soc. Brasil. Mat. (N.S.)* **31** (2000), no. 3, 351–366.
- [15] Lins Neto, Alcides; Soares, Márcio G.: Algebraic solutions of one-dimensional foliations. *J. Differential Geom.* **43** (1996), no. 3, 652–673.
- [16] Nguyên, Viêt-Anh: Oseledec multiplicative ergodic theorem for laminations. *Mem. Amer. Math. Soc.* **246** (2017), no. 1164, ix+174 pp. ISBN: 978-1-4704-2253-0; 978-1-4704-3637-7.
- [17] Nguyên, Viêt-Anh: Ergodic theory for Riemann surface laminations: a survey. *Geometric complex analysis*, 291–327, Springer Proc. Math. Stat., **246**, *Springer, Singapore*, 2018.
- [18] Nguyên, Viêt-Anh: Ergodic theorems for laminations and foliations: recent results and perspectives. *Acta Math. Vietnam.* **46** (2021), no. 1, 9–101.
- [19] Nguyên, Viêt-Anh: Singular holomorphic foliations by curves I: integrability of holonomy cocycle in dimension 2. *Invent. Math.* **212** (2018), no. 2, 531–618.

- [20] Nguyên, Viêt-Anh: Singular holomorphic foliations by curves II: Negative Lyapunov exponent. *J. Geom. Anal.* **33** (2023), no.10, Paper No. 315, 53 pp.
- [21] Verjovsky, Alberto: A uniformization theorem for holomorphic foliations. *The Lefschetz centennial conference, Part III (Mexico City, 1984)*, 233–253, Contemp. Math., **58**, III, *Amer. Math. Soc., Providence, RI*, 1987.