

Regularising Spectral Curves for Homogeneous Yang-Baxter strings

Sibylle Driezen^{1,*} and Niranjan Kamath^{1,2,†}

¹*Institut für Theoretische Physik, ETH Zürich,*

Wolfgang-Pauli-Strasse 27, 8093 Zürich, Switzerland

²*Theoretische Natuurkunde, Vrije Universiteit Brussel (VUB) and The International Solvay Institutes, Pleinlaan 2, B-1050 Brussels, Belgium*

Abstract

In this Letter, we study the semi-classical spectrum of integrable worldsheet σ -models using the Spectral Curve. We consider a Homogeneous Yang-Baxter deformation of the $AdS_5 \times S^5$ superstring, understood as the composition of a Jordanian with a “non-diagonal” TsT deformation. We derive its type IIB supergravity solution, whose isometry algebra features zero supercharges and a non-relativistic conformal algebra in $0 + 1$ dimensions. While the Spectral Curves of non-diagonal TsT models are ill-defined, we demonstrate that the composition with a Jordanian model regularises this issue. From the regularised Curve, we derive the one-loop shift of the classical energy and the semi-classical spectrum of excitations of a point-like string. In the TsT limit, the one-loop shift vanishes despite the loss of supersymmetry. Our results suggest that it may be possible to use standard Bethe Ansatz on spin chain pictures of deformed $\mathcal{N} = 4$ Super-Yang-Mills theory dual to non-diagonal TsT models.

One of the earliest and most robust validations of the AdS/CFT correspondence is the match between the energy spectrum of the $AdS_5 \times S^5$ σ -model and the scaling dimensions for various operators of planar $\mathcal{N} = 4$ Super-Yang-Mills (SYM) theory. Crucial for this achievement was the use of an underlying integrable model and the development of new integrability techniques to solve for the exact spectrum in certain sectors. Particularly well-understood is the spectrum of long operators dual to semi-classical string configurations which are amenable to the algebraic Bethe Ansatz and the semi-Classical Spectral Curve (s-CSC) respectively, as reviewed in [1].

The success of AdS/CFT integrability has spurred, among other factors, significant interest in integrable deformations of string σ -models, cf. [2] for a review. These deformations preserve classical integrability but break many Noether symmetries, which encourages the development of new exact techniques for non-maximally supersymmetric generalisations of AdS/CFT. While numerous examples have now been developed, applying integrable methods to these models has proven challenging. Notable exceptions are “diagonal” TsT-transformations [3] and inhomogeneous Yang-Baxter deformations [4], where techniques similar to those used in the undeformed cases are applicable [5]. This can be attributed to the fact that these models preserve the Cartan subalgebra of the original symmetries.

In contrast, generic Homogeneous Yang-Baxter (HYB) deformations [6], which include all TsT-transformations [7] as well as non-abelian generalisations, generally break the Cartan subalgebra, rendering existing exact techniques challenging. Nonetheless, progress has been made using the fact that HYB deformations are on-shell equivalent to the undeformed σ -model with twisted boundary conditions [8]. This allowed the development of the s-CSC for particular point-like string solutions of a non-diagonal TsT [9] and a non-abelian HYB of Jordanian type [10]. However, unlike Jordanian models, the twist

for non-diagonal TsT is non-diagonalisable. This fact makes the asymptotics of the curve that holds the energy spectrum of non-diagonal TsT non-polynomial [8] and thereby turns the reconstruction of more generic (finite-gap) solutions and their spectra ill-defined. This limitation is reflected on the field theory side by rendering usual Bethe Ansatz techniques inapplicable and necessitating more complex methods [11] [12].

In this Letter, we consider a Jordanian deformation of $AdS_5 \times S^5$ that combines a *minimal* Jordanian with a non-diagonal TsT deformation. We show that this composition regularises the issues related to the non-diagonalisable twist and non-polynomial asymptotics of the non-diagonal TsT. We derive the spectrum of excitations and the one-loop shift of the classical energy of a point-like string solution from this “regularised” curve. In the TsT limit, the one-loop shift vanishes, while the degeneracy of excitations depends on the TsT parameter.

For the corresponding σ -models, we derive the type IIB supergravity solution explicitly from the worldsheet and show that the deformed metric is supported by NSNS, $F^{(3)}$ and $F^{(5)}$ fluxes, along with a constant dilaton. The minimal Jordanian model preserves 12 supersymmetries, the maximum found in the classification of Jordanian deformations of $AdS_5 \times S^5$ superstrings [13], while the non-diagonal TsT and their combination do not preserve any supersymmetry. Interestingly, the non-compact sector of the background exhibits non-relativistic conformal isometries encoded by a Schrödinger algebra in zero spatial dimension [14]. This is the non-relativistic analogue of the conformal algebra relevant for the SYK model [15] and AdS_2/CFT_1 holography [16]. In the non-diagonal TsT limit, the background further simplifies significantly and recovers the isometries of the round S^5 sphere.

With the semi-classical spectrum of a point-like string, our work provides concrete results at the gravity side, which we hope could be matched in the future with a holographic description of a HYB deformation of $\mathcal{N} = 4$

SYM or a potential anisotropic QFT obtained after dimensional reductions. Much progress is being made on the former [17], where they are understood as noncommutative deformations of SYM defined through twisted field products. However, a concrete construction for the Jordanian cases is yet to be developed.

Using the regularisation of a non-diagonal TsT with a Jordanian model, we anticipate the usage of traditional Bethe Ansatz techniques in the spin chain representation of QFTs dual to non-diagonal TsT models, such as e.g. the dipole deformations [11, 18] [19]. The non-diagonal TsT regularisation presented in this Letter can in fact be extended to other examples in the classification of Jordanian deformations of $AdS_5 \times S^5$ [13] [20]. It would therefore be valuable to understand and classify the space of possible regularisations of non-diagonal TsT's within [13] and to examine, for example, when there are nontrivial one-loop shifts in the energy more systematically. Another particular interesting example obtained by non-diagonal TsT is the Hashimoto-Itzhaki/Maldacena-Russo background [21], understood as the Groenewold-Moyal noncommutative deformation of SYM [17, 22]. In this case, taking the non-diagonal TsT limit directly on the twist will yield a non-diagonalisable result, thus requiring Jordanian regularisation at each stage of the computations.

Extending our s-CSC results to a Quantum Spectral Curve (QSC) description would also be very compelling in order to obtain and match with the exact spectrum of the underlying integrable models of the deformed duals. At the CSC level, we find that the deformations only affect the asymptotics of the curve, which aligns with known QSC descriptions of other deformations, e.g. [23].

Interestingly, the non-diagonal TsT we consider uses the same deformation operator as in [24], where it acts on the string σ -model in flat space instead of $AdS_5 \times S^5$, which allowed the authors to obtain the exact energy spectrum. Understanding a sensible flat space limit of our model and matching our results with theirs in the semi-classical limit would thus be very interesting [25].

Another intriguing possibility that we wish to understand further is the possible applications of our work to non-relativistic versions of AdS_2/CFT_1 .

The Jordanian string. Homogeneous Yang-Baxter (HYB) deformations of semi-symmetric space σ -models are realised by the action [4]

$$S = -\frac{\sqrt{\lambda}}{4\pi} \int d^2\sigma \operatorname{str}(\Pi^{\alpha\beta} J_\alpha \hat{d}_- (1 - \eta R_g \hat{d}_-)^{-1} J_\beta), \quad (1)$$

where $\frac{\sqrt{\lambda}}{4\pi}$ denotes the string tension, $\Pi^{\alpha\beta} = \frac{1}{2}(\gamma^{\alpha\beta} - \epsilon^{\alpha\beta})$ with $\gamma^{\alpha\beta}$ the unit worldsheet metric, $\epsilon^{\tau\sigma} = -\epsilon^{\sigma\tau} = -1$, str the supertrace of a \mathbb{Z}_4 -graded superalgebra $\mathfrak{g} = \operatorname{Lie}(G)$, $J = g^{-1}dg$ with $g(\tau, \sigma) \in G$, and $\hat{d}_\pm = \mp \frac{1}{2}P^{(1)} + P^{(2)} \pm \frac{1}{2}P^{(3)}$ with $P^{(0,1,2,3)}$ projectors on the

\mathbb{Z}_4 -eigenspaces. The deformation is induced by $R_g = \operatorname{Ad}_g^{-1}R\operatorname{Ad}_g$, with $\operatorname{Ad}_g x = gxg^{-1}$ for $x \in \mathfrak{g}$ and $R : \mathfrak{g} \rightarrow \mathfrak{g}$ a linear operator, and $\eta \in \mathbb{R}$ is the deformation parameter. When R is antisymmetric with respect to str and solves the classical Yang-Baxter equation (CYBE) the σ -model (1) is integrable [4]. When R is also unimodular with respect to \mathfrak{g} , the σ -model (1) will give rise to a type IIB supergravity solution if the $\eta = 0$ point does [26].

Introducing a basis T_A for \mathfrak{g} , with $A = 1, \dots, \dim \mathfrak{g}$, we can write $R T_A = R^B{}_A T_B$ and $K_{AB} = \operatorname{str}(T_A T_B)$. The R -operator is often written as a 2-fold (graded) wedge product $r = -\frac{1}{2}R^{AB}T_A \wedge T_B$ where $R^B{}_A = K_{AC}R^{CB}$. Unimodular Jordanian R -operators of rank-2 are HYB deformations for which [13, 26–28]

$$r = \mathfrak{h} \wedge \mathfrak{e} - \frac{i}{2}(\mathbb{Q}_1 \wedge \mathbb{Q}_1 + \mathbb{Q}_2 \wedge \mathbb{Q}_2), \quad (2)$$

with $\mathfrak{h}, \mathfrak{e}$ bosonic and $\mathbb{Q}_1, \mathbb{Q}_2$ fermionic generators satisfying $[\mathfrak{h}, \mathfrak{e}] = \mathfrak{e}$, $[\mathbb{Q}_i, \mathfrak{e}] = 0$, $[\mathfrak{h}, \mathbb{Q}_i] = \frac{1}{2}(\mathbb{Q}_i - \epsilon_{ij}a\mathbb{Q}_j)$, $\{\mathbb{Q}_i, \mathbb{Q}_j\} = -i\delta_{ij}\mathfrak{e}$, and $a \in \mathbb{C}$ a free parameter. In this paper, we take $\mathfrak{g} = \mathfrak{psu}(2, 2|4)$ and the R -operator \bar{R}_1 of the classification of [13] on $b = -1/2$, i.e.

$$\begin{aligned} \mathfrak{h} &= \frac{D - J_{03}}{2} + aJ_{12}, & \mathfrak{e} &= \mathfrak{p}_0 + \mathfrak{p}_3, \\ \mathbb{Q}_1 &= \frac{1}{\sqrt{2}}\mathcal{Q}_+^{21}, & \mathbb{Q}_2 &= \frac{i}{\sqrt{2}}\mathcal{Q}_-^{21}. \end{aligned} \quad (3)$$

with D the dilatation operator, J the Lorentz and \mathfrak{p} the translation generators of the conformal subalgebra, and \mathcal{Q}_\pm supercharges. For our conventions and superalgebra realisation, we refer to app. B and eq. (2.2) of [13].

The non-diagonal TsT limit and regularisation.

An interesting limit can be obtained by sending $\eta \rightarrow 0$ while keeping $\eta_{\text{TsT}} \equiv \eta a$ constant. We will call this the *non-diagonal TsT limit*. In fact, the R -matrix (2),(3) can be interpreted as the composition of two HYB deformations $R = R_J + R_{\text{TsT}}$; the “minimal” Jordanian deformation $R_J = R(a = 0)$ followed by the abelian deformation R_{TsT} along the commuting J_{12} and $\mathfrak{p}_0 + \mathfrak{p}_3$ residual isometries of R_J . The latter is equivalent to the sequence of abelian T-duality–shift–T-duality (TsT) transformations [7] along those isometries. In the TsT limit, due to the multiplication of R by η in the action, R_J is eliminated while R_{TsT} survives. Interestingly, our R_{TsT} is of non-diagonal type: at least one of the commuting generators (here $\mathfrak{p}_0 + \mathfrak{p}_3$) is non-diagonalisable. In this case, the on-shell equivalent twisted model has a non-diagonalisable twist and, therefore, a CSC with non-polynomial asymptotics for which it is unknown whether the full curve can be reconstructed for all finite-gap solutions [8]. Generically, this can in turn limit the reconstruction of the s-CSC fluctuations of a specified finite-gap solution. However, we will see that this issue can be regularised by considering the Jordanian composition $R = R_J + R_{\text{TsT}}$

whose twist is always diagonalisable. One can then take the non-diagonal TsT limit on the results.

Type IIB supergravity and isometries. The deformed target space will be manifestly isometric under the subalgebra $\{\mathbb{T}_{\bar{A}}\} \subset \mathfrak{psu}(2, 2|4)$ satisfying $\text{ad}_{\mathbb{T}_{\bar{A}}} R = \text{Rad}_{\mathbb{T}_{\bar{A}}}$ [29]. This can be divided into sets of bosonic generators of the conformal algebra $\mathfrak{t}_{\mathfrak{a}} \subset \mathfrak{so}(2, 4)$ and R-symmetry algebra $\mathfrak{t}_{\mathfrak{s}} \subset \mathfrak{so}(6)$, and of fermionic supercharges $\mathfrak{t}_{\mathfrak{q}}$. For the unimodular R -operator (3) with generic η, a we have [13]

$$\begin{aligned} \mathfrak{t}_{\mathfrak{a}} &= \text{span}(\mathbb{D} + \mathbb{J}_{03}, \mathbb{k}_0 + \mathbb{k}_3, \mathbb{p}_0, \mathbb{p}_3, \mathbb{J}_{12}) \cong \mathfrak{sl}(2, R) \oplus \mathfrak{u}(1)^2, \\ \mathfrak{t}_{\mathfrak{s}} &= \text{span}(\mathbb{R}_{16} - \mathbb{R}_{24}, \mathbb{R}_{14} + \mathbb{R}_{26}, \mathbb{R}_{36} + \mathbb{R}_{45}, \\ &\quad \mathbb{R}_{34} + \mathbb{R}_{56}, \mathbb{R}_{13} + \mathbb{R}_{25}, \mathbb{R}_{15} - \mathbb{R}_{23}, \\ &\quad \mathbb{R}_{12}, \mathbb{R}_{35}, \mathbb{R}_{46}) \cong \mathfrak{su}(3) \oplus \mathfrak{u}(1), \end{aligned} \quad (4)$$

and no supercharges $\mathfrak{t}_{\mathfrak{q}} = \emptyset$. On the special point $a = 0$, studied in [10, 28], $\mathfrak{t}_{\mathfrak{q}}$ enhances to 12 supercharges, while $\mathfrak{t}_{\mathfrak{a}}$ and $\mathfrak{t}_{\mathfrak{s}}$ are as in (4). In the non-diagonal TsT limit $\mathfrak{t}_{\mathfrak{s}}$ enhances to $\mathfrak{so}(6)$, while $\mathfrak{t}_{\mathfrak{q}} = \emptyset$ and $\mathfrak{t}_{\mathfrak{a}}$ is as in (4). In the undeformed limit one of course restores the full $\mathfrak{psu}(2, 2|4)$ with the maximal 32 supercharges.

The isometry algebra $\mathfrak{t}_{\mathfrak{a}}$ corresponds to the Schrödinger algebra in zero spatial dimensions, which is the non-relativistic analogue of zero-dimensional conformal symmetry, extended with the central element \mathbb{J}_{12} , which is a remnant of $\mathfrak{so}(4, 2)$. The $\mathfrak{sl}(2, R) = \text{span}(\mathbb{D} + \mathbb{J}_{03}, \mathbb{k}_0 + \mathbb{k}_3, \mathbb{p}_0 - \mathbb{p}_3)$ subalgebra of the Schrödinger algebra (in any spatial dimension) is central in defining non-relativistic scaling dimensions, primary operators and a state-operator map and, consequently, non-relativistic holography [30], while the central element $\mathbb{p}_0 + \mathbb{p}_3$ has the interpretation of non-relativistic mass.

Let us now extract the IIB supergravity background of the σ -model (1) with the unimodular Jordanian r -matrix (2) and (3) by following the methods of [26]. For this purpose, we can take a bosonic coset representative parametrised as $g = g_{\mathfrak{a}} g_{\mathfrak{s}}$ with $g_{\mathfrak{a}} \in SO(2, 4)$ and $g_{\mathfrak{s}} \in SO(6)$. As the bosonic part of the R -operator only affects the AdS space, we will take $g_{\mathfrak{a}}$ in such a way that the three Cartan generators of the residual $\mathfrak{t}_{\mathfrak{a}}$ isometries are realised as shifts through global left-acting transformations $g \rightarrow g_L g$ with $g_L \in G$ constant. We take

$$g_{\mathfrak{a}} = e^{T\mathbb{H}_T + V\mathbb{H}_V + \Theta\mathbb{H}_{\Theta}} e^{P\mathbb{p}_1} e^{\log(Z)\mathbb{D}}, \quad (5)$$

with $\mathbb{H}_T, \mathbb{H}_V$ and \mathbb{H}_{Θ} the Cartan generators given by $\mathbb{H}_T = \frac{1}{\sqrt{2}}(\mathbb{p}_0 - \mathbb{p}_3 - \frac{1}{2}(\mathbb{k}_0 + \mathbb{k}_3))$, $\mathbb{H}_V = \frac{1}{\sqrt{2}}(\mathbb{p}_0 + \mathbb{p}_3)$ and $\mathbb{H}_{\Theta} = \mathbb{J}_{12}$. \mathbb{H}_T is, up to conjugation, the unique time-like Cartan generator of $\mathfrak{t}_{\mathfrak{a}}$ [10]. The background is then invariant under shifts of the coordinates T, V , and Θ . P and Z are the remaining $SO(2, 4)/SO(1, 4)$ coordinates. We parametrise $g_{\mathfrak{s}}$ as in app. C of [31], with the $SO(6)/SO(5)$ coordinates labelled as $(\phi_1, \phi_2, \phi_3, \xi \equiv \arcsin \omega, r)$. We will denote the collection of all coordinates by X .

Following [26], we then take the operators $O_{\pm} = 1 \pm \eta R_g \hat{d}_{\pm}$ and calculate the one-forms $A_{\pm} = O_{\pm}^{-1} J$ [32]. Observing that for $M = O_{-}^{-1} O_{+}$ one has $M^T P^{(2)} M = P^{(2)}$ [33], shows that $P^{(2)} M P^{(2)}$ implements a local Lorentz transformation on the grade-2 subspace of \mathfrak{g} . This can be realised as $P^{(2)} M P^{(2)} = \text{Ad}_h^{-1} P^{(2)} = P^{(2)} \text{Ad}_h^{-1}$ for an element $h \in G^{(0)} = \exp(P^{(0)} \mathfrak{g})$. Hence, we can write

$$(P^{(2)} A_{+}) h = h (P^{(2)} A_{-}). \quad (6)$$

Next, we introduce the bosonic vielbeins of the deformed and undeformed models as $E = P^{(2)} A_{+}$ and $e = P^{(2)} J$ respectively. Defining the subsets $\mathbb{T}_{\mathfrak{a}} = P^{(2)} \mathbb{T}_{\bar{A}}$, $\mathbb{T}_{\alpha_1} = P^{(1)} \mathbb{T}_{\bar{A}}$, and $\mathbb{T}_{\alpha_2} = P^{(3)} \mathbb{T}_{\bar{A}}$, with $\mathfrak{a} = 0, \dots, 9$, and $\alpha_{1,2} = 1, \dots, 16$, we can then write the metric, B-field, and dilaton as $ds^2 = E^{\mathfrak{a}} E^{\mathfrak{b}} K_{\mathfrak{a}\mathfrak{b}}$, $B = \frac{1}{2} (O_{-}^{-1})_{\mathfrak{a}\mathfrak{b}} e^{\mathfrak{a}} \wedge e^{\mathfrak{b}}$ and $e^{\phi} = (\det O_{+})^{-1/2}$ respectively, while the RR-fluxes can be obtained from projecting the RR-bispinor $\mathcal{F}^{\alpha_1 \beta_2} = 8i (\text{Ad}_h (3 - 4O_{+}^{-1}))^{\alpha_1}_{\gamma_1} K^{\gamma_1 \beta_2}$ on the relevant basis elements of the ten-dimensional Clifford algebra; see [26] for more details. For the latter calculation, one can compute Ad_h by means of the formula (6.8) of [26]. This is, however, a heavy evaluation; we found it more efficient to construct a generic matrix h and solve for its elements by pulling the linear equation (6) onto the target-space basis one-forms. Demanding that the result is an element of $G^{(0)}$ then exhibits a unique expression for h . We find

$$\begin{aligned} ds^2 &= - \frac{2Z^4(Z^2 + P^2) + \eta^2(Z^2 + (1 + 4a^2)P^2)}{2Z^6} dT^2 \\ &\quad + \frac{dZ^2 + dP^2 + P^2 d\Theta^2 - 2dTdV}{Z^2} + ds_{S^5}^2, \\ B &= \frac{\eta}{\sqrt{2}} \left(\frac{PdP \wedge dT + 2aP^2 d\Theta \wedge dT + Z dZ \wedge dT}{Z^4} \right), \\ F^{(3)} &= F_{a=0}^{(3)} + e^{-\phi_0} 4\sqrt{2}\eta a \frac{PdP \wedge dT \wedge dZ}{Z^5} \\ &= e^{-\phi_0} \frac{\sqrt{2}\eta}{Z^5} dT \wedge [2P^2 d\Theta \wedge dZ + ZPdP \wedge d\Theta \\ &\quad - 4aPdP \wedge dZ + Z^2 d\phi_3 \wedge ((1 - r^2)dZ + rZdr) \\ &\quad + r\omega Z^2 d\phi_2 \wedge (\omega Zdr - r\omega dZ + rZd\omega) \\ &\quad + rZ^2(1 - \omega^2)d\phi_1 \wedge (Zdr - rdZ) \\ &\quad - r^2 Z^3 \omega d\phi_1 \wedge d\omega], \\ F^{(5)} &= F_{\eta=0}^{(5)} = e^{-\phi_0} (1 + \star) \frac{4PdT \wedge dV \wedge dZ \wedge dP \wedge d\Theta}{Z^5}, \end{aligned} \quad (7)$$

$F^{(1)} = 0$, and the dilaton $\phi = \phi_0$ constant [34]. In the minimal Jordanian limit $a \rightarrow 0$, the background must coincide with eq. (44) of [28] up to redefining η and after performing the (inverse) coordinate transformation (2.20) of [10] [35]. In the non-diagonal TsT-limit, the $F^{(3)}$ flux remains non-vanishing but simplifies significantly (with only legs in AdS). On $\eta \rightarrow 0$ we naturally find the undeformed $AdS_5 \times S^5$ spacetime with $F^{(3)} = 0$.

It is known that the deformed AdS_5 metric of (7) is geodesically complete on the (formal) parameter surface

$(1 + 4a^2) = 0$ (corresponding to the Schrödinger space-time Sch_2) [36] and on $a = 0$ [10]. We have checked that this remains to be the case for generic (η, a) . In fact, only the geodesic equation for the isometric coordinate V is modified, which does not affect the behaviour around the potential pathological points $Z, P = \{0, \infty\}$. The isometric coordinate T is thus a *global* time-like coordinate.

Let us now consider a point-like string solution of the σ -model in the target space (7), which is the analog of the BMN solution in undeformed $AdS_5 \times S^5$, on which we will apply the s-CSC techniques. We can take the ‘‘BMN-like’’ solution of [10, 11] which is trivial in the P -direction, and therefore also valid on non-trivial a ;

$$T = a_T \tau, \quad V = -\frac{\eta^2 a_T}{2b_Z^2} \tau, \quad Z = b_Z, \quad \phi_3 = a_\phi \tau, \quad (8)$$

with a_T, b_Z, a_ϕ real constants, $\gamma^{\alpha\beta} = \text{diag}(-1, +1)$, and all other fields (bosonic and fermionic) vanishing. The Virasoro constraints are solved on $a_\phi = \sqrt{1 - \frac{\eta^2}{2b_Z^4}} a_T$. We will from now on set $b_Z^4 = 1/2$ so that they require $-1 < \eta < 1$. The Noether Cartan charges $Q_\bullet = \frac{\sqrt{\lambda}}{2\pi} \int_0^{2\pi} d\sigma \text{str}(\mathbf{H}_\bullet \cdot \text{Ad}_g A_\tau^{(2)})$ associated to the \mathfrak{t}_a symmetries of the BMN-like solution (8) evaluate to [37]

$$Q_T = -\sqrt{\lambda} a_T, \quad Q_V = -\sqrt{2\lambda} a_T, \quad Q_\Theta = 0. \quad (9)$$

Twisted formulation. In the previous sections, we described the deformed *periodic* σ -model. We will now employ the fact that, on-shell, HYB models are classically equivalent to the undeformed model with twisted boundary conditions [8]. For a review for *Jordanian* HYB models we refer to sec. 4 and 7 of [10]. In the following, we will use tildes to denote objects related to the twisted variables. Note that, for the R -operator (2)–(3), the map between the deformed periodic variables $g(X)$ to the undeformed twisted variables $\tilde{g}(\tilde{X})$ is only non-trivial in the AdS-sector. On the solution (8) it results in

$$\tilde{T} = a_T \tau, \quad \tilde{Z} = \exp(\eta a_T \sigma) b_Z, \quad \phi_3 = a_\phi \tau, \quad (10)$$

and all other fields vanishing. This solution satisfies the following twisted boundary conditions [38]

$$\tilde{G}(\tau, 2\pi) = W \tilde{G}(\tau, 0) W^t, \quad W = \exp(4\pi \eta a_T \mathfrak{h}), \quad (11)$$

which are written in terms of the gauge-invariant collection of fields $\tilde{G} = \tilde{g} K \tilde{g}^t$ with K the $SO(1, 4) \times SO(5)$ invariant. On $\eta = 0$, $W = 1$ and the boundary conditions become periodic. For generic σ -model solutions, the twist of the bosonic undeformed fields reads

$$W = \exp(\mathbf{Q}(\mathfrak{h} - \mathbf{q}\mathfrak{e})), \quad (12)$$

with the expressions for \mathbf{Q} and \mathbf{q} in terms of $\tilde{g}(\tilde{X})$ given in eqs. (4.5) and (4.6) of [10]. Our (twisted) BMN-like solution is thus characterised by $\mathbf{Q} = 4\pi \eta a_T$ and $\mathbf{q} = 0$.

Since on-shell \mathbf{Q} and \mathbf{q} are constant, and in particular time-independent, these objects correspond to conserved quantities of the twisted model [8]. Yet, their existence is not apparent from a continuous symmetry of the action, in the traditional sense of Noether’s theorem. Importantly, however, for solutions with $\mathbf{Q} \neq 0$, the object \mathbf{q} can be removed via a suitable field redefinition of \tilde{g} [10], while \mathbf{Q} remains a physical ‘‘charge’’; In fact, we will see that \mathbf{Q} characterises the spectrum of the twisted model. We will henceforth refer to \mathbf{Q} as the ‘‘twist charge’’.

The Noether symmetries of the twisted model, on the other hand, are generated in the AdS sector by

$$\tilde{\mathfrak{t}}_a = \{\mathbf{D} + \mathbf{J}_{03}, \mathbf{k}_0 + \mathbf{k}_3, \mathbf{p}_0 - \mathbf{p}_3, \mathbf{J}_{12}\} \cong \mathfrak{sl}(2, R) \oplus \mathfrak{u}(1). \quad (13)$$

This is generally a subset of the symmetry algebra \mathfrak{t}_a of the deformed model [10]. In particular, the rank is reduced: the Cartan of \mathfrak{t}_a is spanned by $\{\mathbf{H}_T, \mathbf{H}_\Theta\}$, while the Cartan of $\tilde{\mathfrak{t}}_a$ is spanned by $\{\mathbf{H}_T, \mathbf{H}_\Theta, \mathbf{H}_V\}$. Morally, as made more precise later, the third Cartan charge Q_V of the deformed model is thus replaced by the twist charge \mathbf{Q} of the twisted model. The remaining Cartan charges $\tilde{Q}_\bullet = \frac{\sqrt{\lambda}}{2\pi} \int_0^{2\pi} d\sigma \text{str}(\mathbf{H}_\bullet \cdot \text{Ad}_{\tilde{g}} \tilde{J}_\tau^{(2)})$ coincide on-shell with those of the deformed model [39]; i.e. $\tilde{Q}_T = Q_T = -\sqrt{\lambda} a_T$ and $\tilde{Q}_\Theta = Q_\Theta = 0$.

While the solution and its charges are independent of a , the generic twist W (12) is a -dependent through \mathfrak{h} . Interestingly, in the non-diagonal TsT-limit, W remains diagonalisable and does not coincide with the non-diagonalisable twist W_{TsT} of [8] that one would obtain if calculated directly for the non-diagonal TsT model, i.e. $W_{\text{TsT}} = \exp(\frac{\eta \mathbf{T}_{\text{ST}}}{\sqrt{\lambda}} (\tilde{Q}_{\mathbf{J}_{12}} \mathbf{e} - \tilde{Q}_e \mathbf{J}_{12}))$. Instead $W_{\text{TsT, reg}} \equiv \lim_{\eta \rightarrow 0, a \rightarrow \eta \mathbf{T}_{\text{ST}}/\eta} W = \exp(-\frac{\eta \mathbf{T}_{\text{ST}}}{\sqrt{\lambda}} \tilde{Q}_e \mathbf{J}_{12})$, which is indeed diagonalisable [40]. Note, furthermore, that for TsT (i.e. abelian HYB) models $Q_{T_{\tilde{A}}} = \tilde{Q}_{T_{\tilde{A}}}$ [8] and since \mathbf{e} is proportional to \mathbf{H}_V , the Noether charge Q_V reinstates itself explicitly through the twist charge $\mathbf{Q}_{\text{TsT, reg}}$.

Semi-Classical Spectral Curve. Both the deformed and twisted σ -model are classically integrable; each have a flat Lax connection that coincides on the on-shell map between the models. We can therefore employ the methods of the Classical Spectral Curve (CSC) to analyse the spectrum of infinite charges. In contrast to the deformed model, in the twisted variables the CSC can be fully reconstructed in terms of *local* conserved charges which include the target-space energy $E \equiv \tilde{Q}_T$. To obtain the semi-classical quantum fluctuations and the one-loop shift to the energy of the BMN-like vacuum we will thus work in the twisted variables.

The twisted CSC is obtained from the conserved eigenvalues $\lambda(z)$ of the twisted monodromy matrix [8]

$$\Omega_W(z) = W^{-1} \mathcal{P} \exp\left(-\int_0^{2\pi} d\sigma \mathcal{L}_\sigma^{\tilde{g}}(z)\right), \quad (14)$$

with $\mathcal{L}^{\tilde{g}}(z) = \tilde{g} \mathcal{L}(z) \tilde{g}^{-1} - d\tilde{g} \tilde{g}^{-1}$ the gauge-transformed

Lax connection of the undeformed supercoset model

$$\mathcal{L}(z) = \tilde{J}^{(0)} + (z_+ + z_- \star) \tilde{J}^{(2)} + z \tilde{J}^{(1)} + z^{-1} \tilde{J}^{(3)}, \quad (15)$$

with $\tilde{J}^{(i)} \equiv P^{(i)} \tilde{J}$ and $z_{\pm} \equiv \frac{1}{2}(z^2 \pm z^{-2})$ [41]. Since $\mathcal{L}(z)$ is flat $\forall z \in \mathbb{C}$ and periodic also in the twisted variables [8, 10], the eigenvalues of $\Omega_W(z)$ give rise to infinite towers of conserved charges by expanding around fixed values of z . Generically, these charges are non-local, but since $\mathcal{L}_0^{\tilde{g}}(z=1) = 0$, the leading asymptotics around $z=1$ are local charges. Redefining the spectral parameter as $z = \sqrt{\frac{x+1}{x-1}}$ [42] this point in \mathbb{C} corresponds to $x = \infty$ around which the monodromy matrix reads

$$\Omega_W(x) = W^{-1}(1 + 4\pi x^{-1} O_W) + \mathcal{O}(x^{-2}), \quad (16)$$

with $O_W = \frac{1}{2\pi} \int_0^{2\pi} d\sigma \text{Ad}_{\tilde{g}}(\tilde{J}_\tau^{(2)} - \frac{1}{2}(\tilde{J}_\sigma^{(1)} - \tilde{J}_\sigma^{(3)}))$. Note that in the AdS-sector O_W is only conserved in the projections on $\tilde{\mathfrak{t}}_{\mathfrak{a}}$ [43]. However, after diagonalisation and a possible conjugation to the Cartan, we will precisely uncover the conserved charges only. In fact, in terms of the AdS *quasimomenta* $p_{\tilde{1}-\tilde{4}}(x)$ obtained from the AdS eigenvalues $\lambda_{\tilde{i}}(x) = e^{ip_{\tilde{i}}(x)}$ of $\Omega_W(x)$, we obtain [44]

$$\begin{pmatrix} p_{\tilde{1}} \\ p_{\tilde{2}} \\ p_{\tilde{3}} \\ p_{\tilde{4}} \end{pmatrix} \sim \frac{\mathbf{Q}}{2} \begin{pmatrix} a \\ i-a \\ -i-a \\ a \end{pmatrix} + \frac{2\pi}{x\sqrt{\lambda}} \begin{pmatrix} \tilde{Q}_{\Theta} - E \\ (2ia-1)\tilde{Q}_{\Theta} \\ (2ia+1)\tilde{Q}_{\Theta} \\ \tilde{Q}_{\Theta} + E \end{pmatrix}. \quad (17)$$

It is interesting to compare this situation to the undeformed periodic model ($W=1$), where these asymptotics are dictated by the three Noether Cartan charges of the AdS_5 isometries. In contrast, in the twisted case the Noether Cartan algebra is only two-dimensional, but the asymptotics of $p_{\tilde{i}}(x)$ are still determined by three conserved quantities: the Noether charges \tilde{Q}_{Θ} and $E = \tilde{Q}_T$ and, crucially, the twist charge \mathbf{Q} , similar as in [8, 10].

The quasimomenta of the BMN-like solution (8), or equivalently (10), can be evaluated on the full complex plane. In the AdS sector, we obtain

$$\begin{aligned} \frac{p_{\tilde{1}}(x)}{2\pi a_T} - a\eta &= -\frac{p_{\tilde{4}}(x)}{2\pi a_T} + a\eta = \frac{\sqrt{x^2 - \eta^2}}{x^2 - 1}, \\ \frac{p_{\tilde{2}}(x)}{2\pi a_T} + a\eta &= -\frac{p_{\tilde{3}}(x)}{2\pi a_T} - a\eta = \frac{x\sqrt{1 - \eta^2 x^2}}{x^2 - 1}, \end{aligned} \quad (18)$$

which corresponds to a finite gap solution with branch-cuts on $\mathcal{C}_{\tilde{1},\tilde{4}} \equiv [-\eta, \eta]$ and $\mathcal{C}_{\tilde{2},\tilde{3}} \equiv (-\infty, -\eta^{-1}] \cup [\eta^{-1}, \infty)$ in \mathbb{C} . Matching (18) with the (off-shell) asymptotics (17) is consistent with the values of the conserved quantities $E = \tilde{Q}_T, \tilde{Q}_{\Theta}$ and \mathbf{Q} given in the previous section. The quasimomenta of the sphere, which we denote by $p_{\tilde{1}-\tilde{4}}(x)$, are unaffected by the deformation and have no cuts [45]

$$p_{\tilde{1}}(x) = p_{\tilde{2}}(x) = -p_{\tilde{3}}(x) = -p_{\tilde{4}}(x) = \frac{2\pi a_{\phi} x}{x^2 - 1}. \quad (19)$$

Semi-classical quantum fluctuations on top of the BMN-like solution can be obtained at the level of the (twisted) Spectral Curve by introducing microscopic cuts between the Riemann sheets (18)–(19) in all possible ways [10, 45, 46]. Heuristically, since a branch cut can be viewed as a “condensation” of poles, the microscopic cut or excitation can be treated as a single pole singularity. The excitation is bosonic when this cut connects two AdS sheets $\hat{i}-\hat{j}$ or two sphere sheets $\tilde{i}-\tilde{j}$, while it is fermionic when it connects an AdS sheet \hat{i} with a sphere sheet \tilde{j} , see e.g. [47]. The backreaction from such excitations results in shifts of the classical (background) quasimomenta as $p_i \rightarrow p_i + \delta p_i$, $i \in \{\hat{i}, \tilde{i}\}$, which must satisfy a number of analytic properties coming from the BMN-like CSC as well as from $\mathfrak{psu}(2, 2|4)$. These properties are so restrictive that they fully determine δp_i through a simple linear problem which we will now summarise.

First, the corrections must not alter the gluing conditions of the classical macroscopic cuts, i.e.

$$\delta p_i(x + i\epsilon) - \delta p_j(x - i\epsilon) = 0, \quad x \in \mathcal{C}_{i,j}, \quad (20)$$

for infinitesimal ϵ . Similarly, on the location $x_n^{ij} \in \mathbb{C}$ of the new microscopic cuts between the sheets (i, j) and with mode number n we have, to leading order,

$$p_i(x_n^{ij}) - p_j(x_n^{ij}) = 2\pi n, \quad n \in \mathbb{Z}. \quad (21)$$

This condition fixes the positions x_n^{ij} . The number N_n^{ij} of such excitations is furthermore constrained by the level-matching condition $\sum_n n \sum_{\text{all } ij} N_n^{ij} = 0$. Due to the \mathbb{Z}_4 -grading of $\mathfrak{psu}(2, 2|4)$, the quasimomenta should in addition satisfy “inversion symmetry”

$$\delta p_i(x) = -\delta p_{i'}(x^{-1}), \quad \delta p_{\tilde{i}}(x) = -\delta p_{\tilde{i}'}(x^{-1}), \quad (22)$$

with $i = (1, 2, 3, 4)$ and $i' = (2, 1, 4, 3)$, and a “synchronisation” around the poles $x = \pm 1$ of the Lax connection

$$\text{Res}_{x=\pm 1} \delta p_i = \text{Res}_{x=\pm 1} \delta p_{i'} = \text{Res}_{x=\pm 1} \delta p_{\tilde{i}} = \text{Res}_{x=\pm 1} \delta p_{\tilde{i}'}, \quad (23)$$

which stems, in addition, from the supertracelessness of the Lax and the Virasoro constraints. For bosonic excitations there is a further simplification $\text{Res}_{x=\pm 1} \delta p_{\tilde{1}} = -\text{Res}_{x=\pm 1} \delta p_{\tilde{3}}$, and related implications from (23), due to the tracelessness of $\mathfrak{su}(2, 2)$ and $\mathfrak{su}(4)$ separately. At last, because of (17), we demand that the next-to-leading order asymptotics of $\delta p_i(x)$ around zeroes of the (gauge-transformed) Lax connection, here $x = \infty$, takes the form

$$\begin{aligned} \begin{pmatrix} \delta p_{\tilde{1}} \\ \delta p_{\tilde{2}} \\ \delta p_{\tilde{3}} \\ \delta p_{\tilde{4}} \end{pmatrix} &\sim \frac{\delta \mathbf{Q}}{2} \begin{pmatrix} a \\ i-a \\ -i-a \\ a \end{pmatrix} + \frac{4\pi}{x\sqrt{\lambda}} \begin{pmatrix} \frac{\delta \Delta}{2} + \sum_{i=\tilde{3},\tilde{4},\tilde{3},\tilde{4}} N_{\tilde{1}i} \\ \sum_{i=\tilde{3},\tilde{4},\tilde{3},\tilde{4}} N_{\tilde{2}i} \\ -\sum_{i=\tilde{1},\tilde{2},\tilde{1},\tilde{2}} N_{\tilde{3}\tilde{i}} \\ -\frac{\delta \Delta}{2} - \sum_{i=\tilde{1},\tilde{2},\tilde{1},\tilde{2}} N_{\tilde{4}\tilde{i}} \end{pmatrix} \\ \begin{pmatrix} \delta p_{\tilde{1}} \\ \delta p_{\tilde{2}} \\ \delta p_{\tilde{3}} \\ \delta p_{\tilde{4}} \end{pmatrix} &\sim \frac{4\pi}{x\sqrt{\lambda}} \begin{pmatrix} -\sum_{i=\tilde{3},\tilde{4},\tilde{3},\tilde{4}} N_{\tilde{1}i} \\ -\sum_{i=\tilde{3},\tilde{4},\tilde{3},\tilde{4}} N_{\tilde{2}i} \\ \sum_{i=\tilde{1},\tilde{2},\tilde{1},\tilde{2}} N_{\tilde{3}\tilde{i}} \\ \sum_{i=\tilde{1},\tilde{2},\tilde{1},\tilde{2}} N_{\tilde{4}\tilde{i}} \end{pmatrix}, \end{aligned} \quad (24)$$

where $\delta\Delta$ is the anomalous correction to the energy E , $\delta\mathbf{Q}$ the correction to the twist charge \mathbf{Q} , and $N_{ij} = \sum_n N_n^{ij}$ is the total number of excitations connecting sheets (i, j) [48]. If we denote each contribution to $\delta\Delta$ from the excitation N_n^{ij} as $\Omega^{ij}(x_n)$, we can write

$$\delta\Delta = \sum_n \Omega^{ij}(x_n) N_n^{ij}. \quad (25)$$

In analogy with the harmonic oscillator, we will call $\Omega^{ij}(x_n)$ the *frequencies*. For more details on the origin of the properties (20)–(24) we refer to [45–47] in the periodic case and [10] in the twisted case. Note that the only difference between the periodic and twisted models lies in the asymptotics (17) and (24)–(24) and, consequently, the identification of local charges in the spectral curve.

By combining the constraints from (20)–(24), we can now calculate all the frequencies $\Omega^{ij}(x_n)$ as well as $\delta\mathbf{Q}$ for our model. A significant advantage of employing the s-CSC over standard semiclassical quantisation methods based on effective actions is that, for a large class of classical solutions, the full spectrum of frequencies can be completely determined from a single sphere and a single *AdS* frequency (the “frequency basis”). This was explicitly proven in [46] for solutions with pairwise symmetric quasimomenta, i.e. $p_{\hat{1},\hat{2},\hat{1},\hat{2}} = -p_{\hat{4},\hat{3},\hat{4},\hat{3}}$, using inversion symmetry as well as the composition of off-shell frequencies which share poles of opposite residues. In our case, the (constant) a -dependent terms in the *AdS* quasimomenta (18) spoil this assumption; however, they only shift the reference values on the sheets and since the frequencies are derived from the x -dependent terms, which are pairwise symmetric, the proof in [46] readily goes through. Consequently, using table (B.1) therein, the spectrum of frequencies can be entirely determined through, e.g., $\Omega^{\hat{2}\hat{3}}$ and $\Omega^{\hat{2}\hat{3}}$ only.

To compute the sphere frequency $\Omega^{\hat{2}\hat{3}}$, we need to turn on only the bosonic excitation $N_n^{\hat{2}\hat{3}}$. The pole structure, incl. residues, at $x = \pm 1, x_n^{\hat{2}\hat{3}}$ for e.g. $\delta p_{\hat{2}}(x)$ is then easily obtained using inversion symmetry and comparison with the asymptotics (24), similar as done e.g. in [13, 45]. The backreaction of this excitation on the *AdS* sheets is expected to slightly shift the branch points connecting $\hat{1} - \hat{4}$ and $\hat{2} - \hat{3}$, justifying the ansatz

$$\delta p_{\hat{1}}(x) = f(x) + \frac{g(x)}{K(\frac{1}{x})}, \quad \delta p_{\hat{4}}(x) = f(x) - \frac{g(x)}{K(\frac{1}{x})}, \quad (26)$$

with $K(x) = \sqrt{1 - \eta^2 x^2}$ and $f(x), g(x)$ arbitrary functions. Using the synchronisation of the poles at $x = \pm 1$ for this bosonic excitation and the asymptotics (24), we find $f(x) = a\delta\mathbf{Q}/2$ using Liouville’s theorem, and

$$\delta\Delta = \sum_n \Omega^{\hat{2}\hat{3}}(x_n^{\hat{2}\hat{3}}) N_n^{\hat{2}\hat{3}}, \quad \Omega^{\hat{2}\hat{3}}(x_n^{\hat{2}\hat{3}}) = \frac{2K(1)}{(x_n^{\hat{2}\hat{3}})^2 - 1}. \quad (27)$$

For any *AdS* excitation, e.g. $N_n^{\hat{2}\hat{3}}$, there will be no correction to the quasimomenta of the sphere, $\delta p_i(x) = 0$,

which can be verified from (24) and the synchronisation of the poles at $x = \pm 1$ for bosonic excitations. We can furthermore assume the same ansatz (26) for $\delta p_{\hat{1}}(x)$ and $\delta p_{\hat{4}}(x)$. Liouville’s theorem again implies that $f(x) = a\delta\mathbf{Q}/2$ while using inversion symmetry we can write

$$g(x) = - \sum_n \frac{K(x_n^{\hat{2}\hat{3}}) \alpha((x_n^{\hat{2}\hat{3}})^{-1})}{x - (x_n^{\hat{2}\hat{3}})^{-1}} N_n^{\hat{2}\hat{3}} + \text{reg.}, \quad (28)$$

where $\alpha(x) = \frac{4\pi}{\sqrt{\lambda}} \frac{x^2}{x^2 - 1}$. Matching with (24) we find

$$\delta\Delta = \sum_n \Omega^{\hat{2}\hat{3}}(x_n^{\hat{2}\hat{3}}) N_n^{\hat{2}\hat{3}}, \quad \Omega^{\hat{2}\hat{3}}(x_n^{\hat{2}\hat{3}}) = \frac{2K(x_n^{\hat{2}\hat{3}})}{(x_n^{\hat{2}\hat{3}})^2 - 1} \quad (29)$$

As argued, using inversion symmetry and composition of the poles, the other frequencies can be easily extracted from (B.1) of [46]. Note that their form as functions of the pole positions x_n^{ij} is independent from the parameter a and thus coincides with (6.18)–(6.23) of [10]. The expressions for x_n^{ij} themselves, however, will receive a -contributions. After solving (21) for every excitation and inserting the obtained solutions for x_n^{ij} in $\Omega^{ij}(x_n^{ij})$ we find

$$\begin{aligned} \Omega^{\hat{1}\hat{3}} &= \Omega^{\hat{1}\hat{4}} = \Omega^{\hat{2}\hat{3}} = \Omega^{\hat{2}\hat{4}} = -\sqrt{1 - \eta^2} + \sqrt{1 - \eta^2 + n^2 a_T^{-2}}, \\ \Omega^{\hat{1}\hat{4}} &= -2 + \sqrt{2 + n^2 a_T^{-2} + 2\sqrt{1 + (1 - \eta^2)n^2 a_T^{-2}}}, \\ \Omega^{\hat{2}\hat{3}} &= \sqrt{2 + n^2 a_T^{-2} - 2\sqrt{1 + (1 - \eta^2)n^2 a_T^{-2}}}, \\ \Omega^{\hat{1}\hat{3}} &= -1 + \sqrt{1 + \eta^2 + (na_T^{-1} - 2a\eta)^2}, \\ \Omega^{\hat{2}\hat{4}} &= -1 + \sqrt{1 + \eta^2 + (na_T^{-1} + 2a\eta)^2} \quad (30) \\ \Omega^{\hat{1}\hat{3}} &= \Omega^{\hat{1}\hat{4}} = -1 + \sqrt{1 + (na_T^{-1} - a\eta)^2}, \\ \Omega^{\hat{1}\hat{4}} &= \Omega^{\hat{2}\hat{4}} = -1 + \sqrt{1 + (na_T^{-1} + a\eta)^2}, \\ \Omega^{\hat{2}\hat{3}} &= \Omega^{\hat{2}\hat{4}} = -\sqrt{1 - \eta^2} + \sqrt{1 + (na_T^{-1} + a\eta)^2}, \\ \Omega^{\hat{1}\hat{3}} &= \Omega^{\hat{2}\hat{3}} = -\sqrt{1 - \eta^2} + \sqrt{1 + (na_T^{-1} - a\eta)^2}. \end{aligned}$$

In the undeformed limit, all of the above energy frequencies degenerate to the single BMN frequency [49] (then, on the Virasoro constraint, $a_T = a_\phi$). On $a = 0$ they degenerate to six independent contributions, incl. two sets of 4-fold fermionic frequencies, that coincide with [10]. Interestingly, we see that turning on a breaks the degeneracy. This is expected given that we break all the 12 supersymmetries of the $a = 0$ model [13]. In the non-diagonal TsT limit, this breaking lifts back to five independent frequencies, incl. two sets of 4-fold fermionic frequencies distinct from the $a = 0$ case, which can now be attributed to the restoration of the $\mathfrak{so}(6)$ symmetries.

As a non-trivial consistency check of both the twisting and regularisation procedure, we verified the expressions of $\Omega^{ij}(x_n)$ from the periodic and deformed viewpoint through an independent calculation of the effective action of small fluctuations around the solution (8).

Let us now consider $\delta\mathbf{Q}$. First, note that we can generically write $\delta p_1(x) = A + h(x)$ with $A \equiv \delta p_1(\infty)$ a constant and $h(x)$ a function capturing all the $\mathcal{O}(x^{-1})$ corrections around $x = \infty$, i.e. $h(\infty) = 0$. From the previous discussion on the $\hat{2}\hat{3}$ and $\hat{3}\hat{3}$ excitations, we thus have $A = a\delta\mathbf{Q}/2$ and $h(x) = g(x)/K(x^{-1})$. Therefore, in both cases, $h(0) = 0$. By inversion symmetry, this implies $\delta p_2(x) = -A + \mathcal{O}(x^{-1})$, which is consistent with (24) only when $\delta\mathbf{Q} = 0$. This can be readily generalised to all the other excitations, as we can expect from their composition rules. We thus obtain no anomalous correction to the twist charge, which matches with the $a = 0$ Jordanian string [10] and the β -deformation [5].

At last, we can compute the one-loop correction to the vacuum energy of our classical string [45, 50]

$$E \approx Q_T + E_{1\text{-loop}} = Q_T + \frac{1}{2} \sum_{n \in \mathbb{Z}} \sum_{ij} (-)^{F_{ij}} \Omega_n^{ij}, \quad (31)$$

with $F_{ij} = 0$ for bosonic and $F_{ij} = 1$ for fermionic excitations. After approximating the sums as integrals by assuming (without loss of generality) $a_T \gg 1$, and using similar integration tricks as in [10], we find that the a -dependency drops out in the result after a number of non-trivial cancellations. Hence, we obtain

$$E_{1\text{-loop}} = a_T \left(\eta(1 - \eta) - (3 - \eta^2) \log \sqrt{1 - \eta^2} - (1 + \eta^2) \log \sqrt{\frac{(1 + \eta)(1 + \eta^2)}{1 - \eta}} \right), \quad (32)$$

as for the $a = 0$ Jordanian string [10]. Interestingly, this means that in the non-diagonal TsT limit $E_{1\text{-loop}}$ vanishes despite the broken supersymmetry.

Acknowledgements: We thank Niklas Beisert, Martí Berenguer Mimó, Sylvain Lacroix, Stijn Van Tongeren, and especially Riccardo Borsato, for useful discussions and comments on the draft. This work is partly based on the Master thesis of one of the authors (NK) prepared at ETH Zürich. SD is supported by the Swiss National Science Foundation through the NCCR SwissMAP. NK is supported in part by the FWO Vlaanderen through the project G006119N, as well as by the Vrije Universiteit Brussel through the Strategic Research Program “High-Energy Physics”.

* sdriezen@phys.ethz.ch

† niranjannkamath@gmail.com

- [1] N. Beisert *et al.*, *Lett. Math. Phys.* **99**, 3 (2012), [arXiv:1012.3982](https://arxiv.org/abs/1012.3982) [hep-th]; M. Staudacher, *Lett. Math. Phys.* **99**, 191 (2012), [arXiv:1012.3990](https://arxiv.org/abs/1012.3990) [hep-th]; S. Schafer-Nameki, *Lett. Math. Phys.* **99**, 169 (2012), [arXiv:1012.3989](https://arxiv.org/abs/1012.3989) [hep-th].

- [2] B. Hoare, *J. Phys. A* **55**, 093001 (2022), [arXiv:2109.14284](https://arxiv.org/abs/2109.14284) [hep-th].
- [3] O. Lunin and J. M. Maldacena, *JHEP* **0505**, 033 (2005), [arXiv:hep-th/0502086](https://arxiv.org/abs/hep-th/0502086) [hep-th]; S. Frolov, *JHEP* **0505**, 069 (2005), [arXiv:hep-th/0503201](https://arxiv.org/abs/hep-th/0503201) [hep-th]; S. A. Frolov, R. Roiban, and A. A. Tseytlin, *JHEP* **07**, 045 (2005), [arXiv:hep-th/0503192](https://arxiv.org/abs/hep-th/0503192) [hep-th].
- [4] C. Klimcik, *J.Math.Phys.* **50**, 043508 (2009), [arXiv:0802.3518](https://arxiv.org/abs/0802.3518) [hep-th]; F. Delduc, M. Magro, and B. Vicedo, *JHEP* **1311**, 192 (2013), [arXiv:1308.3581](https://arxiv.org/abs/1308.3581) [hep-th]; *Phys.Rev.Lett.* **112**, 051601 (2014), [arXiv:1309.5850](https://arxiv.org/abs/1309.5850) [hep-th].
- [5] L. F. Alday, G. Arutyunov, and S. Frolov, *JHEP* **0606**, 018 (2006), [arXiv:hep-th/0512253](https://arxiv.org/abs/hep-th/0512253) [hep-th]; N. Beisert and R. Roiban, *JHEP* **08**, 039 (2005), [arXiv:hep-th/0505187](https://arxiv.org/abs/hep-th/0505187) [hep-th]; M. de Leeuw and S. J. van Tongeren, *Nucl. Phys.* **B860**, 339 (2012), [arXiv:1201.1451](https://arxiv.org/abs/1201.1451) [hep-th]; V. Kazakov, *Ludwig Faddeev Memorial Volume*, 293 (2018), [arXiv:1802.02160](https://arxiv.org/abs/1802.02160) [hep-th]; G. Arutyunov, M. de Leeuw, and S. J. van Tongeren, *Teor. Mat. Fiz.* **182**, 28 (2014), [arXiv:1403.6104](https://arxiv.org/abs/1403.6104) [hep-th]; R. Klabbers and S. J. van Tongeren, *Nucl. Phys. B* **925**, 252 (2017), [arXiv:1708.02894](https://arxiv.org/abs/1708.02894) [hep-th].
- [6] I. Kawaguchi, T. Matsumoto, and K. Yoshida, *JHEP* **04**, 153 (2014), [arXiv:1401.4855](https://arxiv.org/abs/1401.4855) [hep-th]; S. J. van Tongeren, *JHEP* **06**, 048 (2015), [arXiv:1504.05516](https://arxiv.org/abs/1504.05516) [hep-th].
- [7] D. Osten and S. J. van Tongeren, *Nucl. Phys.* **B915**, 184 (2017), [arXiv:1608.08504](https://arxiv.org/abs/1608.08504) [hep-th].
- [8] R. Borsato, S. Driezen, and J. L. Miramontes, *JHEP* **04**, 053 (2022), [arXiv:2112.12025](https://arxiv.org/abs/2112.12025) [hep-th].
- [9] H. Ouyang, *JHEP* **12**, 126 (2017), [arXiv:1709.06844](https://arxiv.org/abs/1709.06844) [hep-th].
- [10] R. Borsato, S. Driezen, J. M. Nieto García, and L. Wyss, *Phys. Rev. D* **106**, 066015 (2022), [arXiv:2207.14748](https://arxiv.org/abs/2207.14748) [hep-th].
- [11] M. Guica, F. Levkovich-Maslyuk, and K. Zarembo, *J. Phys.* **A50**, 394001 (2017), [arXiv:1706.07957](https://arxiv.org/abs/1706.07957) [hep-th].
- [12] Despite this, more intricate methods based on the Baxter equation have been successfully employed in [11] resulting in the spectrum of a twisted spin chain with which the semi-classical spectrum of a specific point-like string configuration matched at large R-charge [9].
- [13] R. Borsato and S. Driezen, *SciPost Phys.* **14**, 160 (2023), [arXiv:2212.11269](https://arxiv.org/abs/2212.11269) [hep-th].
- [14] Models with Schrödinger symmetries in general dimensions have attracted also much attention as possible holographic descriptions of anisotropic condensed matter systems [30].
- [15] S. Sachdev and J. Ye, *Phys. Rev. Lett.* **70**, 3339 (1993), [arXiv:cond-mat/9212030](https://arxiv.org/abs/cond-mat/9212030); A. Kitaev, Talk at KITP, April 7, 2015, <https://online.kitp.ucsb.edu/online/entangled15/kitaev/> (2015); Talk at KITP, May 27, 2015, <https://online.kitp.ucsb.edu/online/entangled15/kitaev2/> (2015).
- [16] J. Maldacena and D. Stanford, *Phys. Rev. D* **94**, 106002 (2016), [arXiv:1604.07818](https://arxiv.org/abs/1604.07818) [hep-th]; G. Sárosi, *PoS Modave2017*, 001 (2018),

- arXiv:1711.08482 [hep-th].
- [17] S. J. van Tongeren, *Nucl. Phys.* **B904**, 148 (2016), arXiv:1506.01023 [hep-th]; *Phys. Lett.* **B765**, 344 (2017), arXiv:1610.05677 [hep-th]; T. Meier and S. J. van Tongeren, *Phys. Rev. Lett.* **131**, 121603 (2023), arXiv:2301.08757 [hep-th]; *JHEP* **12**, 045 (2023), arXiv:2305.15470 [hep-th].
- [18] T. Matsumoto and K. Yoshida, *JHEP* **04**, 180 (2015), arXiv:1502.00740 [hep-th].
- [19] It would be interesting to verify compatibility with the results of [9].
- [20] One can include Jordanian models that do not admit a unimodular extension, as this feature does not play a role in the non-diagonal TsT limit as explained later.
- [21] A. Hashimoto and N. Itzhaki, *Phys. Lett.* **B465**, 142 (1999), arXiv:hep-th/9907166 [hep-th]; J. M. Maldacena and J. G. Russo, *JHEP* **9909**, 025 (1999), arXiv:hep-th/9908134 [hep-th].
- [22] T. Matsumoto and K. Yoshida, *JHEP* **06**, 163 (2014), arXiv:1404.3657 [hep-th].
- [23] N. Gromov and F. Levkovich-Maslyuk, *JHEP* **04**, 134 (2016), arXiv:1510.02098 [hep-th]; V. Kazakov, S. Leurent, and D. Volin, *JHEP* **12**, 044 (2016), arXiv:1510.02100 [hep-th]; N. Gromov, V. Kazakov, G. Korchemsky, S. Negro, and G. Sizov, *JHEP* **01**, 095 (2018), arXiv:1706.04167 [hep-th].
- [24] K. Idiab and S. J. van Tongeren, (2024), arXiv:2403.07365 [hep-th].
- [25] We thank Stijn Van Tongeren for pointing out this connection.
- [26] R. Borsato and L. Wulff, *JHEP* **10**, 045 (2016), arXiv:1608.03570 [hep-th].
- [27] V. N. Tolstoy, ArXiv Mathematics e-prints (2004), math/0402433.
- [28] S. J. van Tongeren, *SciPost Phys.* **7**, 011 (2019), arXiv:1904.08892 [hep-th].
- [29] Following [24], we refer to *manifest* isometries as those corresponding to Noether symmetries that leave the R operator invariant and therefore the Lagrangian manifestly invariant, without compensating total derivatives. There may, however, be more general “enhanced” symmetries that leave the Lagrangian invariant up to total derivatives. For realisations of the latter see [24].
- [30] Y. Nishida and D. T. Son, *Phys. Rev. D* **76**, 086004 (2007), arXiv:0706.3746 [hep-th]; D. T. Son, *Phys. Rev. D* **78**, 046003 (2008), arXiv:0804.3972 [hep-th].
- [31] R. Borsato, *Integrable strings for AdS/CFT*, Ph.D. thesis, Utrecht U. (2015), arXiv:1605.03173 [hep-th].
- [32] Note that here \pm is not a worldsheet index.
- [33] Note that $\hat{d}_+ = (\hat{d}_-)^T$.
- [34] The last term in the B -field may be removed by the gauge transformation $B \rightarrow B + \frac{\eta}{2\sqrt{2}} d(Z^{-2} dT)$.
- [35] We find, however, a difference in the expressions of $F_{z\phi_2 x^-}^{(3)}$ and $F_{z\theta x^-}^{(3)}$ which turn out to be typographical errors in eq. (44) of [28], and we thank Stijn Van Tongeren for confirming this. We have checked that (7) solves the type IIB field equations and Bianchi identities.
- [36] M. Blau, J. Hartong, and B. Rollier, *JHEP* **07**, 027 (2009), arXiv:0904.3304 [hep-th].
- [37] The full family of Noether (super)charges is given by $Q_{T_{\tilde{A}}} = \frac{\sqrt{\tilde{\lambda}}}{2\pi} \int_0^{2\pi} d\sigma \operatorname{str} \left[T_{\tilde{A}} \cdot \operatorname{Ad}_{\tilde{g}}(A_{\tilde{\tau}}^{(2)} - \frac{1}{2}(A_{\tilde{\sigma}}^{(1)} - A_{\tilde{\sigma}}^{(3)})) \right]$ with $T_{\tilde{A}} \in \{\mathfrak{t}_a, \mathfrak{t}_s, \mathfrak{t}_q\}$.
- [38] In terms of the group element, the twisted boundary conditions read as $\tilde{g}(\tau, 2\pi) = W\tilde{g}(\tau, 0)h$ with $h \in G^{(0)}$ a possible right-acting gauge ambiguity, for more details see [10].
- [39] The full family of Noether charges of the twisted model is given by $\tilde{Q}_{T_{\tilde{A}}} = \frac{\sqrt{\tilde{\lambda}}}{2\pi} \int_0^{2\pi} d\sigma \operatorname{str} \left[T_{\tilde{A}} \cdot \operatorname{Ad}_{\tilde{g}}(\tilde{J}_{\tilde{\tau}}^{(2)} - \frac{1}{2}(\tilde{J}_{\tilde{\sigma}}^{(1)} - \tilde{J}_{\tilde{\sigma}}^{(3)})) \right]$ with $T_{\tilde{A}} \in \mathfrak{psu}(2, 2|4)$ satisfying $WT_{\tilde{A}}W^{-1} = T_{\tilde{A}}$ [8, 10].
- [40] This is, however, a special feature of the model under consideration, particularly when the non-diagonal TsT model has *at most* one non-diagonalisable generator in the R -operator. For non-diagonal TsT models with both non-diagonalisable generators in the R -operator, such as the one corresponding to the Maldacena-Russo background [22], the non-diagonal TsT limit would still result in a non-diagonalisable twist. In that case, we believe that the TsT limit should be taken only on physical quantities such as the energy spectrum.
- [41] The Hodge star on one-forms is defined on the worldsheet Σ as $\star J = J_{\alpha} \epsilon^{\alpha\beta\gamma} \gamma_{\beta\gamma} d\sigma^{\gamma}$.
- [42] Note that there is a typographical error in [10] in the redefinition from z to x after eq. (5.3), where $z = \sqrt{\frac{1+x}{1-x}}$ was written instead of $z = \sqrt{\frac{x+1}{x-1}}$.
- [43] This is in contrast to the undeformed periodic model ($W = 1$) where all projections on the $\mathfrak{psu}(2, 2|4)$ generators are conserved.
- [44] Note that the quasimomenta have a $2\pi\mathbb{Z}$ ambiguity.
- [45] N. Gromov and P. Vieira, *Nucl. Phys. B* **789**, 175 (2008), arXiv:hep-th/0703191.
- [46] N. Gromov, S. Schafer-Nameki, and P. Vieira, *JHEP* **12**, 013 (2008), arXiv:0807.4752 [hep-th].
- [47] N. Beisert, V. Kazakov, K. Sakai, and K. Zarembo, *Commun. Math. Phys.* **263**, 659 (2006), arXiv:hep-th/0502226.
- [48] We assume that only the twist charge and the energy can receive corrections, while the spin charge of angular momentum \tilde{Q}_{Θ} does not, as it sits in a representation of a compact group.
- [49] D. E. Berenstein, J. M. Maldacena, and H. S. Nastase, *JHEP* **0204**, 013 (2002), arXiv:hep-th/0202021 [hep-th].
- [50] S. Frolov and A. A. Tseytlin, *JHEP* **06**, 007 (2002), arXiv:hep-th/0204226.