

Computation of Robust Option Prices via Structured Multi-Marginal Martingale Optimal Transport

Linn Engström* Sigrid Källblad† Johan Karlsson‡

Abstract

We introduce an efficient computational framework for solving a class of multi-marginal martingale optimal transport problems, which includes many robust pricing problems of large financial interest. Such problems are typically computationally challenging due to the martingale constraint, however, by extending the state space we can identify them with problems that exhibit a certain sequential martingale structure. Our method exploits such structures in combination with entropic regularisation, enabling fast computation of optimal solutions and allowing us to solve problems with a large number of marginals. We demonstrate the method by using it for computing robust price bounds for different options, such as lookback options and Asian options.

Keywords: *Martingale Optimal Transport, Multi-Marginal Optimal Transport, Entropic Regularisation, Robust Finance, Numerical Methods*

1 Introduction

A fundamental problem in mathematical finance is to find fair prices of financial claims. The main goal of this article is to compute robust and model independent bounds on prices of exotic options when there is uncertainty about the true underlying market model. More specifically, we develop computationally efficient methods for obtaining such bounds when restricting to market models that are consistent with given market data and respect fundamental market modelling principles. For market data consisting of prices of liquidly traded call options for multiple maturities, the problem can be formulated as a multi-marginal martingale optimal transport (MOT) problem, a problem which has been extensively studied over the last decade. The approach taken here leverages recent computational results for solving multi-marginal optimal transport problems using entropic regularisation. More pertinently, we will combine entropic regularisation with methods utilising specific structures in the payoff function so as to obtain an efficient method for MOT problems in the presence of multiple marginal constraints.

The classical approach to pricing of financial derivatives is to start by postulating a market model. A model effectively consists of a filtered probability space, say $(\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$, supporting a stochastic process S modelling the underlying future stock prices. We here restrict to discrete time and suppose that the risk-free interest rate is zero. Such a model is then a valid model, in the sense that it does not allow for arbitrage-opportunities, if there exists a so-called risk-neutral measure \mathbb{Q} rendering the price process S a martingale. For an exotic option ensuring the holder a payoff given by $\phi(S_0, \dots, S_T)$, for some function $\phi : \mathbb{R}^{T+1} \rightarrow \mathbb{R}$, it turns out that any fair price thereof — that is, any price which does not introduce arbitrage opportunities to the market — must be given by

$$\mathbb{E}_{\mathbb{Q}}[\phi(S_0, \dots, S_T)] \tag{1}$$

*KTH Royal Institute of Technology; linneng@kth.se

†KTH Royal Institute of Technology; sigrid.kallblad@math.kth.se

‡KTH Royal Institute of Technology; johan.karlsson@math.kth.se

for such a risk-neutral pricing measure \mathbb{Q} . Fair prices are thus highly dependent on the original choice of market model, as well as on the choice of pricing measure. In reality, however, neither of the two is known. The acknowledgment of this fact has led to an intense stream of research aiming at quantifying the impact of the modelling assumptions.

A natural question to ask is whether it is possible to obtain robust bounds on fair prices. Naïvely, such bounds could be obtained by optimising (1) over all possible market models and pricing measures; that is, over all probability spaces $(\Omega, \mathcal{F}, \mathbb{Q})$ supporting a stochastic process S which is a martingale in its own filtration under \mathbb{Q} . Such a procedure leads however to bounds which are too wide to be of any practical use. Consensus has therefore been reached on also acknowledging that certain financial products are traded to such a great extent that their market prices can be viewed as ‘correct’ and should contain information about the market participants’ beliefs about the future. Put differently, the prices of liquidly traded options carry information about the pricing measure effectively used by the market. It is thus sensible to further restrict to market models and pricing measures for which the computed prices of such liquidly traded options agree with actual market prices. The most common choice in the literature is to strive for consistency with market prices of European call options, the perhaps most commonly traded option. Breeden and Litzenberger [9] observed that if one has access to call prices for a continuum of strikes, at given maturities $\mathcal{T} \subseteq \{0, 1, \dots, T\}$, one may recover the marginal distributions of the underlying price process at these dates. The thus obtained family of marginal distributions, say $\{\mu_t\}_{t \in \mathcal{T}}$, may therefore be considered to be known and fixed; it reflects what we know about the market-implied pricing measure. The problem of solving for the associated price bound, here formulated in terms of a lower bound, takes the following form:

$$\inf_{(\Omega, \mathcal{F}, \mathbb{Q}, S)} \mathbb{E}_{\mathbb{Q}}[\phi(S_0, \dots, S_T)] \quad (2a)$$

$$\text{subject to } \mathbb{E}_{\mathbb{Q}}[S_t | \sigma(S_0, \dots, S_{t-1})] = S_{t-1}, \quad t \in \{1, \dots, T\} \quad (2b)$$

$$S_t \sim_{\mathbb{Q}} \mu_t, \quad t \in \mathcal{T}. \quad (2c)$$

Note that if $\{\mu_t\}_{t \in \mathcal{T}}$ are obtained from consistent call prices, they are marginal distributions of a martingale and are thus in convex order. By Strassen [58], it turns out that the latter is both a necessary and sufficient condition for problem (2) to be well posed. The robust pricing problem was first introduced by Hobson [38]. Subsequently a rigorous duality theory has been established motivating the problem also from a pricing-hedging perspective; see Cheridito, Kupper and Tangpi [13] and Dolinsky and Soner [19].

About a decade ago, the insightful observation was made that problem (2) can be viewed as an optimal transport problem with an additional martingale constraint — an MOT problem; see Beiglböck, Henry-Labordère and Penkner [6] and Galichon, Henry-Labordère and Touzi [26]. The optimal transport (OT) problem — that is, problem (2) without condition (2b) — is a classical problem that has received renewed attention in recent years; we refer to Villani [60] for an overview. It was originally formulated for two marginals, and then amounts to finding a transport plan moving the mass of one distribution to another while minimising the associated cost, but has since been extended to transportation problems over multiple marginals; see, e.g., Gangbo and Świąch [27], Rüschendorf [53] and Pass [50]. Equipped with the additional martingale constraint, the MOT problem turned out to be an intriguing problem which triggered intense research and led to many new developments; for multi-marginal results, we refer to Nutz, Stebegg and Tan [48] and Sester [55]. Prior to the contributions of [6] and [26], the robust pricing problem was typically addressed via the so-called Skorokhod embedding problem (SEP). This approach was initiated by [38], who considered the problem with one marginal constraint and a claim depending on the past maximum of a *continuous* price process, and linked it to the well-known Azéma–Yor solution of the SEP. The connection to the SEP remained for long the predominant approach for addressing (continuous) robust pricing problems and extensive research in this area led to new contributions in terms of both robust price bounds and solutions of the SEP; for multi-marginal results in this direction, we refer, e.g., to Beiglböck, Cox and Huesmann [5], Cox, Oblój and Touzi [15], and Henry-Labordère et al. [35], see also [40].

Recently, there has been a rapid development of computational theory and algorithms for OT problems. The OT problem can be formulated as a linear programming problem when restricting

the marginals to have support on a pre-specified finite grid. However, the number of variables is often large, making the problem computationally intractable for many problems of practical interest. In particular, this is the case for multi-marginal problems, since the number of variables grows exponentially in the number of marginals. One of the dominant approaches for handling this has been to utilise entropic regularisation, inspired by Cuturi [16] and Benamou et al. [8]. The regularised problem can then be addressed by use of duality theory. In particular, the dual problem contains significantly fewer variables than the primal problem, since the number of dual variables is linear in the number of marginals. Further, the dual problem has a certain decomposable structure, which allows for obtaining explicit expressions characterising the optimal dual variables. These expressions allow for deriving efficient update formulas for blocks of dual variables, which makes coordinate dual ascent — that is, to cyclically optimise over subsets of the dual variables — a particularly suitable method for solving the regularised problem. In the context of OT, this method is often referred to as Sinkhorn’s algorithm for the bi-marginal problem. It was originally utilised for matrix balancing in Sinkhorn [56], but has since been reinvented several times. It can be derived, for example, using Bregman projections or Dykstra’s algorithm; see [8] and Peyré and Cuturi [51] for further details. It can be shown that the dual iterates generated by this method converge linearly, see Luo and Tseng [45] (cf. Franklin [25] for the bi-marginal case), and for bi-marginal OT problems, entropic regularisation thus yields an efficient method for solving the problem. For multi-marginal problems, however, the computational complexity also increases exponentially as a function of the number of marginals, see Lin et al. [44], and entropic regularisation alone is thus not enough for efficient computation of the problem. Fortunately, for many multi-marginal problems the cost function has graph-structures that can be exploited in combination with entropic regularisation. These are referred to as *graph-structured multi-marginal optimal transport problems* and have been studied, e.g., in [8, 23, 30, 31, 33, 46]; see also Altschuler and Boix-Adserà [2] and Beier et al. [4]. In order to carry out the coordinate dual ascent method, one effectively needs to compute certain matrix projections; the key point utilised in those papers is that by suitably exploiting the additional structure of the cost function, said projections can be computed rapidly, thus rendering an efficient algorithm also for very large numbers of marginal constraints.

In this article, we combine entropic regularisation with the exploitation of structures inherent in the problem, in order to computationally solve the multi-marginal MOT problem (2). In line with the above-mentioned literature on graph-structured multi-marginal OT problems, the idea is to exploit specific structures in the cost function. We therefore restrict ourselves to financial derivatives whose payoff is of the form

$$\phi(S_0, \dots, S_T) = \sum_{t=1}^T \phi_t(S_{t-1}, X_{t-1}, S_t, X_t), \quad (3)$$

where $X_0 = h_0(S_0)$ and $X_t = h_t(S_t, S_{t-1}, X_{t-1})$, for some functions $h_0 : \mathbb{R} \rightarrow \mathbb{R}$, $h_t : \mathbb{R}^3 \rightarrow \mathbb{R}$ and $\phi_t : \mathbb{R}^4 \rightarrow \mathbb{R}$, $t \in \{1, \dots, T\}$. We utilise this structure in a twofold manner. First, leveraging on the fact that for a given stochastic process, one can construct a Markov process with the same pair-wise joint marginal distributions onto adjacent marginals, for payoffs of the form (3), we manage to relax condition (2b) to $\mathbb{E}_{\mathbb{Q}}[S_t | \sigma(S_{t-1}, X_{t-1})] = S_{t-1}$, $t \in \{1, \dots, T\}$. In effect, we are facing a problem where each constraint is only connecting adjacent marginals and where the cost function has the same structure as in the above-mentioned works on graph-structured problems. In turn, we may thus employ methods from the theory on graph-structured multi-marginal OT problems to find an efficient algorithm for solving problem (2). Our first main result establishes the equivalence between the relaxed and the original problem formulation, while our second main result provides an efficient method for solving the corresponding entropy regularised problem. We illustrate the usefulness of our method by performing explicit computations in a number of examples: First, we apply our algorithm to problems for which the theoretical solution is known; we use it to recover the left-monotone coupling, late and early transport behaviour, as well as optimal models for digital options and lookback options. We then use it to compute the robust price bound for an Asian option, subject to several marginal constraints, where as far as we know, no theoretical solution is known.

For MOT problems, entropic regularisation was first introduced by de March [18], who derived a

method based on Sinkhorn’s algorithm and Newton’s method for solving bi-marginal problems. As for early numerical results on robust pricing problems, we also mention Henry-Labordere [34], Davis, Obłój and Raval [17] and Tan and Touzi [59]. Computationally solving multi-marginal MOT problems has since been addressed with neural network approaches in Eckstein and Kupper [21] and Eckstein et al. [20], and by linear programming approaches in [20] and Guo and Obłój [29], where the latter relies on a relaxation of the martingale constraint. However, due to the curse of dimensionality, it is computationally challenging to utilise these methods for problems with a large number of marginals. Markovian assumptions in a multi-marginal MOT context were introduced in Henry-Labordere [37, Remark 2.7] and further discussed in Sester [54], who considered a problem subject to an explicit Markov constraint. Such problems are not convex, which motivated Eckstein and Kupper [22] to consider multi-marginal MOT problems under a time-homogeneity assumption. While Alfonsi et al. [1] and Lindheim and Steidl [61] have developed methods for OT problems featuring constraints closely related to the martingale constraint, to the best of our knowledge, there are no other works using entropic regularisation methods for multi-marginal MOT problems specifically.

The rest of the article is organised as follows: In Section 2, we characterise our class of multi-marginal MOT problems in terms of properties inherent in the payoff function; we also provide examples of derivatives of practical financial interest belonging to this class, and provide some background on structured OT. Section 3 is devoted to the reformulation of the problem onto a form that allows for efficient computations: We show that for our class of payoff functions, the MOT problem is equivalent to an OT problem where the complexity of the martingale constraint has been reduced to involve only adjacent marginals; the latter problem is then discretised and formulated on tensor form. In Section 4, relying on the structure inherent in our reformulated problem, we derive efficient formulas for the optimal dual variables corresponding to its entropy regularised version. In Section 5, we apply our computational framework in a number of different examples and compare our results with existing analytical solutions.

Notation

We let \odot and \oslash denote the elementwise product and division of two tensors (or vectors/matrices), while $\exp(\cdot)$ denotes elementwise application of the exponential function. The Kronecker product is indicated by \otimes . The notation $\text{diag}(v)$ represents a diagonal matrix with the vector v on the main diagonal, while $\mathbf{1}_n$ and $\mathbf{0}_n$ are the vectors of length $n \in \mathbb{N}$ consisting of ones and zeros, respectively. The one-vector should not be mixed up with the indicator function $\chi_A : \mathbb{R}^n \rightarrow \mathbb{R}$ associated with some set $A \subset \mathbb{R}^n$,

$$\chi_A(x) := \begin{cases} 1, & x \in A \\ 0, & x \notin A. \end{cases}$$

We let $[m] := \{0, \dots, m\}$ for any $m \in \mathbb{N}$ and write $[m] \setminus k$ for short for $[m] \setminus \{k\}$ for $k \in [m]$. The inner product $\langle \cdot, \cdot \rangle$ refers to the Frobenius inner product, that is,

$$\langle A, B \rangle = \sum_{i_0, \dots, i_n \in [m]} A(i_0, \dots, i_n) B(i_0, \dots, i_n), \quad A, B \in \mathbb{R}^{n^m},$$

and we let $P_j : \mathbb{R}^{n^m} \rightarrow \mathbb{R}^n$ denote the projection operator that projects a tensor onto its j :th dimension, defined by

$$P_j(A)(i_j) := \sum_{i_\ell : \ell \in [m] \setminus j} A(i_0, \dots, i_m), \quad i_j = 1, \dots, n, \quad \text{where } A \in \mathbb{R}^{n^m}.$$

Similarly, we let $P_{j,k} : \mathbb{R}^{n^m} \rightarrow \mathbb{R}^{n^2}$ be the projection operator that projects the tensor jointly onto the two dimensions j and k (see [23, p. 7]). Every real-valued function is assumed to be Borel measurable.

2 Problem formulation and background

In this section we describe our main problem of interest, along with some examples demonstrating its relevance. We then provide a brief introduction to optimal transport with emphasis on the computational aspects of the problem; in particular, we review how a certain type of structure inherent in some transportation problems enables fast computation of optimal solutions.

2.1 Problem formulation for structured payoff functions

We will now specify our class of problems, which consists of MOT problems where the payoff function is of a certain form. Let a probability space $(\Omega, \mathcal{F}, \mathbb{Q})$ and a price process $S : \Omega \times [T] \rightarrow \mathbb{R}$ be given such that S is a martingale under \mathbb{Q} . Then define a second stochastic process $X : \Omega \times [T] \rightarrow \mathbb{R}$ by

$$X_t = \begin{cases} h_0(S_0), & t = 0 \\ h_t(S_t, S_{t-1}, X_{t-1}), & t \in [T] \setminus 0 \end{cases} \quad (4)$$

where $h_0 : \mathbb{R} \rightarrow \mathbb{R}$ and $h_t : \mathbb{R}^3 \rightarrow \mathbb{R}$ for $t \in [T] \setminus 0$. By defining X in this way we are able to isolate important features of the path of S , thus enabling the pricing of many exotic options. We are here interested in pricing path-dependent derivatives whose payoff function can be pairwise decoupled, i.e.,

$$\phi(S_0, \dots, S_T) = \sum_{t \in [T] \setminus 0} \phi_t(S_{t-1}, X_{t-1}, S_t, X_t), \quad (5)$$

where $\phi_t : \mathbb{R}^4 \rightarrow \mathbb{R}$ for $t \in [T] \setminus 0$ and X is obtained from (4) via a given family of functions h_t , $t \in [T]$. For payoff functions of this form the MOT problem (2) becomes

$$\inf_{(\Omega, \mathcal{F}, \mathbb{Q}, S)} \mathbb{E}_{\mathbb{Q}} \left[\sum_{t \in [T] \setminus 0} \phi_t(S_{t-1}, X_{t-1}, S_t, X_t) \right] \quad (6a)$$

$$\text{subject to } \mathbb{E}_{\mathbb{Q}}[S_t | \sigma(S_0, \dots, S_{t-1})] = S_{t-1}, \quad t \in [T] \setminus 0 \quad (6b)$$

$$S_t \sim_{\mathbb{Q}} \mu_t, \quad t \in \mathcal{T}. \quad (6c)$$

Here we assume that the given marginals μ_t , for $t \in \mathcal{T} \subseteq \{0, 1, \dots, T\}$, are in convex order. Such problems are well posed, since it follows by Strassen's theorem that their feasible sets are nonempty [58].

The contribution of this work is the development of a computational method that allows for approximately solving this class of multi-marginal MOT problems for T large. As we will see, many problems of practical financial interest belong to this class.

2.2 Financial examples

Below follows a few examples of financial derivatives whose payoff functions satisfy the assumptions of our framework; the first four of them will be revisited later in Section 5, while the remaining ones illustrate the versatility of our class of problems.

Example 2.1 (Lookback and barrier options). Consider the case when X is the maximum process of the price process S , that is, when $X_t = \max_{r \in [t]} S_r$ for $t \in [T]$; since it can be written

$$X_t = \begin{cases} S_0, & t = 0 \\ \max\{S_t, X_{t-1}\}, & t \in [T] \setminus 0 \end{cases}$$

the maximum process is of the form (4) (the minimum process can be considered analogously). Related derivatives are single asset, single period maximum lookback options such as the *floating strike lookback put*, whose payoff function is $\phi(S_T, X_T) = (X_T - S_T)^+$ (see, for example, [39, pp. 623–525]). Certain barrier options can also be priced by considering the maximum process, for example

the *up-and-in barrier call* with barrier b and strike K , which at maturity pays out $\phi(S_T, X_T) = \chi_{[b, \infty)}(X_T)(S_T - K)^+$ (see, for example, [12, pp. 143–150]). In Section 5 we will solve an upper bound MOT problem where the payoff is equal to the terminal value of the maximum.

Example 2.2 (Asian options). Let the process X be the arithmetic mean of the price process S , that is, let $X_t = (t+1)^{-1} \sum_{j \in [t]} S_j$ for $t \in [T]$. Since an equivalent way of defining X is

$$X_t = \begin{cases} S_0, & t = 0 \\ \frac{1}{t+1} S_t + \left(1 - \frac{1}{t+1}\right) X_{t-1}, & t \in [T] \setminus 0, \end{cases}$$

it is of the form (4). Derivatives whose payoff is a function of the arithmetic mean are Asian options with arithmetic averaging, such as the *average price call* and the *average strike call*. Their payoffs are $\phi(X_T) = (X_T - K)^+$ and $\phi(S_T, X_T) = (S_T - X_T)^+$, respectively, where $K > 0$ denotes a fixed strike price (see, for example, [39, pp. 626–627]). In Section 5 we will provide an example where we compute the robust fair price of an *Asian straddle*, given two, three and four known marginals.

Example 2.3 (Barrier options). Let $A_0, \dots, A_T \subseteq \mathbb{R}$ be given subsets and define a stochastic process via $X_t = \chi_{A_0 \times \dots \times A_t}(S_0, \dots, S_t)$ for $t \in [T]$; this X can be written

$$X_t = \begin{cases} \chi_{A_0}(S_0), & t = 0 \\ \chi_{A_t}(S_t) X_{t-1}, & t \in [T] \setminus 0. \end{cases}$$

Its terminal value, X_T , indicates whether the price S has passed through all the sets A_0, \dots, A_T during the duration of the contract. By setting all but two of the sets A_0, \dots, A_T equal to \mathbb{R} , it is possible to consider *second-order binary options*. Indeed, let $T_0 \in [T-1] \setminus 0$ and define $A_{T_0} := \{x \in \mathbb{R} : x \geq b_0\}$ and $A_T := \{x \in \mathbb{R} : x \geq b\}$ for some barrier levels b_0 and b . Let $A_t = \mathbb{R}$ for $t \in [T] \setminus \{T_0, T\}$. Then the payoff of an up-up type second-order asset binary option can be written $\phi(S_T, X_T) = S_T X_T$ (see, for example, [12, pp. 109–110]). A *double knockout option*, on the other hand, expires worthless if the price of the underlying crosses any of the two barriers $b_l < S_0 < b_u$ (see, for example, [49, p. 274]). Defining $A := \{x \in \mathbb{R} : b_l < x < b_u\}$ and setting $A_t = A$ for all $t \in [T]$ allows us to also consider such options. In Section 5 we will compute the robust price of a *digital option*.

Example 2.4 (Variance swaps). Consider the case when the process X is the realised variance of the price process S , that is, when $X_t = t^{-1} \sum_{j=1}^t (\log(S_j/S_{j-1}))^2$ for $t \in [T] \setminus [0]$. Set $X_0 := 0$; then X can be written as

$$X_t = \begin{cases} 0, & t = 0 \\ \left(1 - \frac{1}{t}\right) X_t + \frac{1}{t} \left(\log\left(\frac{S_t}{S_{t-1}}\right)\right)^2, & t \in [T] \setminus 0, \end{cases}$$

which shows that the realised variance is indeed of the form (4). The payoff of a *variance swap* is then given by $\phi(X_T) = X_T - \sigma_{\text{fixed}}^2$, where σ_{fixed}^2 denotes the fixed variance (see, for example, [39, p. 629]). Needless to say, the realised volatility of the price S can easily be obtained as $\sqrt{X_t}$ at any time point $t \in [T]$. We conclude this example by noting that the optimal solution of the corresponding MOT problem is known in the case when $T = 1$; it is the left-monotone transport plan [7, 36]. We will revisit this solution in Section 5.

Example 2.5 (Parisian options). In a similar fashion to Example 2.3, the payoff can be defined in terms of the duration during which the process is in a given interval; let $A \subset \mathbb{R}$ and set $X_t = \sum_{j \in [t]} \chi_A(S_j)$ for $t \in [T]$. This process can also be written

$$X_t = \begin{cases} \chi_A(S_0), & t = 0 \\ \chi_A(S_t) + X_{t-1}, & t \in [T] \setminus 0 \end{cases}$$

and its terminal value, X_T , corresponds to the number of, say, days that the price S has spent within A during the duration of the contract, a property that allows for pricing *Parisian options*. In order for such an option — written with a $D \in \mathbb{N}$ day window — to activate, the price of the underlying must spend at least D days within A (see, for example, [39, p. 622]). As an example, an up-and-in Parisian put with a D day window, a strike price K and a barrier b has a payoff that can be written $\phi(S_T, X_T) = (K - S_T)^+ \chi_{[D, \infty)}(X_T)$, with X_T defined as above for $A = \{x \in \mathbb{R} : x \geq b\}$.

Example 2.6 (Dual expiry options). Consider a setting with two times of expiry, T and $T_0 \in [T-1] \setminus 0$, and claims whose payoff depend on the value of the underlying at those two specific times only. By defining the process X as

$$X_t = \begin{cases} 0, & t \in [T_0 - 1] \\ S_{T_0}, & t \in [T] \setminus [T_0 - 1] \end{cases}$$

we are able to ‘remember’ the value of the underlying price process S at time T_0 throughout the duration of the contract; an equivalent definition is

$$X_t = \begin{cases} 0, & t = 0 \\ X_{t-1} \chi_{[T] \setminus T_0}(t) + S_t \chi_{\{T_0\}}(t), & t \in [T] \setminus 0, \end{cases}$$

which is of the form (4). This choice of X allows for pricing dual expiry options — the payoff of a *forward start call* is given by $\phi(S_T, X_T) = (S_T - X_T)^+$, while a *ratchet call option* with strike price K has a payoff $\phi(S_T, X_T) = \max\{(S_T - K)^+, (X_T - K)^+\}$ (see, for example, [12, pp. 107, 119]).

Example 2.7 (Cliquet options). Consider the sum of the truncated relative returns; let $C_l > 0$ be the local cap and let $X_t = \sum_{j=1}^t \max\{\min\{(S_j - S_{j-1})/S_{j-1}, C_l\}, 0\}$ for $t \in [T] \setminus 0$. Set $X_0 = 0$. Then X_t can be written on the form from equation (4); indeed,

$$X_t = \begin{cases} 0, & t = 0 \\ \max\left\{\min\left\{\frac{S_t - S_{t-1}}{S_{t-1}}, C_l\right\}, 0\right\} + X_{t-1}, & t \in [T] \setminus 0. \end{cases}$$

The payoff function of a globally floored, locally capped *cliquet option* is then given by $\phi(X_T) = \max\{X_T, F_g\}$, where F_g is the global floor (see, for example, [62, p. 379]).

2.3 Structured multi-marginal optimal transport

The OT problem was introduced in Section 1 as the problem of minimising the total cost for transporting the mass of one distribution to another one. The transport can occur in one or several steps — we refer to the former as bi-marginal transport, while the latter is multi-marginal transport. In order to formalise this, consider $m + 1$ probability spaces $(\mathcal{X}_0, \mathcal{F}_{\mathcal{X}_0}, \mu_0), \dots, (\mathcal{X}_m, \mathcal{F}_{\mathcal{X}_m}, \mu_m)$ and let $\mathcal{M}(\mathcal{X}_j)$ denote the set of probability measures on \mathcal{X}_j for $j \in [m]$. Similarly, let $\mathcal{M}(\mathcal{X})$ denote the set of probability measures on the product space $\mathcal{X} := \mathcal{X}_0 \times \dots \times \mathcal{X}_m$ equipped with the product σ -algebra. Let the projection operators $\mathcal{P}_j : \mathcal{M}(\mathcal{X}) \rightarrow \mathcal{M}(\mathcal{X}_j)$ be defined as $\mathcal{P}_j(\pi) := \int_{\mathcal{X}_0} \dots \int_{\mathcal{X}_{j-1}} \int_{\mathcal{X}_{j+1}} \dots \int_{\mathcal{X}_m} d\pi$, for $\pi \in \mathcal{M}(\mathcal{X})$ and $j \in [m]$. A measure $\pi \in \mathcal{M}(\mathcal{X})$ is said to be a coupling of (μ_0, \dots, μ_m) if it satisfies $\mathcal{P}_j(\pi) = \mu_j$ for $j \in [m]$; in the context of optimal transport, a coupling is often referred to as a transport plan. Given a cost function $c : \mathcal{X} \rightarrow \mathbb{R}$, where $c(x_0, \dots, x_m)$ is the cost associated with the path $(x_0, \dots, x_m) \in \mathcal{X}$, the optimal transport problem is the problem of finding a coupling that minimises the total associated cost, i.e. that attains

$$\inf_{\pi \in \mathcal{M}(\mathcal{X})} \int_{\mathcal{X}_0 \times \dots \times \mathcal{X}_m} c(x_0, \dots, x_m) d\pi(x_0, \dots, x_m) \quad (7a)$$

$$\text{subject to } \mathcal{P}_j(\pi) = \mu_j, \quad j \in [m]. \quad (7b)$$

A coupling of (μ_0, \dots, μ_m) that attains the minimum in problem (7) is said to be an optimal coupling or an optimal transport plan. Problem (7) can be generalised by relaxing the constraint (7b) to only hold for a subset $\Gamma \subseteq [m]$ of the marginals or by introducing unbalanced problems [4, 14, 23] (cf. bi-marginal problems [28, 42, 43, 52]).

In order to solve problems on the form (7) computationally, we assume that each of the sets $\mathcal{X}_0, \dots, \mathcal{X}_m$ is finite, that is, $\cup_{j \in [m]} \mathcal{X}_j$ is contained within $n \in \mathbb{N}$ points. Then the marginals can be represented by vectors $\mu_j \in \mathbb{R}^n$, $j \in [m]$, and the cost function and the transport plan by tensors $C \in \mathbb{R}^{n^{m+1}}$ and $Q \in \mathbb{R}_+^{n^{m+1}}$, respectively. The discrete analogue to problem (7) is then

$$\min_{Q \in \mathbb{R}_+^{n^{m+1}}} \langle C, Q \rangle \quad (8a)$$

$$\text{subject to } P_j(Q) = \mu_j, \quad j \in [m]. \quad (8b)$$

Problem (8) is a linear programming problem in n^{m+1} variables. Since the number of variables grows exponentially as a function of the number of marginals, the problem quickly becomes intractable for standard methods — even problems with two marginals ($m = 2$) are not straightforward to solve when the number of points n is large. One way to alleviate this difficulty is to regularise the problem with an entropy term, as proposed in [8, 16, 51]; we will see that this leads to a dual problem in $n(m+1)$ variables and explicit formulas for the optimal choice thereof.

The entropy regularised problem is given by

$$\min_{Q \in \mathbb{R}_+^{n^{m+1}}} \langle C, Q \rangle + \varepsilon D(Q) \quad (9a)$$

$$\text{subject to } P_j(Q) = \mu_j, \quad j \in [m], \quad (9b)$$

where $\varepsilon > 0$ is a regularisation parameter and

$$D(Q) := \sum_{i_0, \dots, i_m \in [n]} (Q(i_0, \dots, i_m) \log Q(i_0, \dots, i_m) - Q(i_0, \dots, i_m) + 1)$$

is an entropy term. It is solved by utilising duality theory. The Lagrangian function corresponding to problem (9) reads

$$\mathcal{L}(Q, \lambda) = \langle C, Q \rangle + \varepsilon D(Q) + \sum_{j \in [m]} \lambda_j^\top (\mu_j - P_j(Q)),$$

where $\lambda = \{\lambda_j\}_{j \in [m]}$ are Lagrangian multipliers in \mathbb{R}^n corresponding to the constraints (9b). The Lagrangian multipliers are also referred to as dual variables, while the dual functional of problem (9) is defined as $\varphi(\lambda) := \min\{\mathcal{L}(Q, \lambda) : Q \in \mathbb{R}_+^{n^{m+1}}\}$.

Since the regularisation term $D(Q)$ ensures that the Lagrangian function is strictly convex in Q , the minimum is attained by the unique $Q > 0$ for which the gradient of the Lagrangian function is zero; that is, for the transport plan Q satisfying

$$0 = \partial_{Q(i_0, \dots, i_m)} \mathcal{L}(Q, \lambda) = C(i_0, \dots, i_m) + \varepsilon \log Q(i_0, \dots, i_m) - \sum_{j \in [m]} \lambda_j(i_j)$$

for $i_0, \dots, i_m \in \{1, \dots, n\}$. Solving for Q yields that the primal solution is given by $Q_\lambda = K \odot U_\lambda$, where

$$K(i_0, \dots, i_m) := e^{-C(i_0, \dots, i_m)/\varepsilon}, \quad U_\lambda(i_0, \dots, i_m) := \prod_{j \in [m]} e^{\lambda_j(i_j)/\varepsilon}, \quad i_0, \dots, i_m = 1, \dots, n$$

— see this by noting that $\langle C, Q_\lambda \rangle + \varepsilon \langle Q_\lambda, \log Q_\lambda \rangle - \sum_{j \in [m]} \lambda_j^\top P_j(Q_\lambda) = 0$. Inserting this solution Q_λ into the Lagrangian function yields the dual functional φ . The dual of problem (9) becomes

$$\max_{\lambda} \varphi(\lambda) = \max_{\lambda} \sum_{j \in [m]} \lambda_j^\top \mu_j - \varepsilon \langle K, U_\lambda \rangle + \varepsilon n^{m+1}. \quad (10)$$

Note that the dual variables are unconstrained since the corresponding constraints of the primal problem are equality constraints (see, for example, [47]). One can show that for a primal-dual pair like the one above, strong duality holds (see, for example, [47]), meaning that the respective problem values of problems (9) and (10) coincide. Moreover, the optimal solution of the primal problem (9) is Q_{λ^*} , where λ^* is the optimal solution of the dual problem (10). The optimal solution Q_{λ^*} of the primal problem (9) can therefore be obtained by first solving the dual problem (10).

Since the dual functional φ is concave, the dual problem is a convex problem. Its maximisers are thus found as stationary points of φ . The optimal λ therefore satisfies $0 = \partial_{\lambda_j(i_j)}\varphi(\lambda) = \mu_j(i_j) - P_j(K \odot U_\lambda)(i_j)$ for $i_j \in \{1, \dots, n\}$ and $j \in [m]$. In order to simplify the expressions, the solution is reparameterised in terms of $u = \{u_j\}_{j \in [m]}$ where $u_j := \exp(\lambda_j/\varepsilon)$ for $j \in [m]$. It follows that the tensor U_λ can be written $U_\lambda(i_0, \dots, i_m) = \prod_{\ell \in [m]} u_\ell(i_\ell)$, for $i_0, \dots, i_m \in \{1, \dots, n\}$. Since U_λ is a rank one tensor, it holds that $P_j(K \odot U_\lambda) = u_j \odot P_j(K \odot U_\lambda^{-j})$, where $U_\lambda^{-j}(i_0, \dots, i_m) := \prod_{\ell \in [m] \setminus j} u_\ell(i_\ell)$. The optimal u therefore satisfies

$$u_j = \mu_j \odot P_j(K \odot U_\lambda^{-j}), \quad j \in [m]. \quad (11)$$

The computational bottleneck of the formula (11) is evaluating the projection, which requires manipulating an n^{m+1} -tensor. Consequently, the computational complexity increases exponentially as the number of marginals $m+1$ grows [44], and entropic regularisation alone is thus not enough for efficient computation of the problem. Fortunately, for many such multi-marginal problems the cost function has graph-structures that can be exploited in combination with entropic regularisation [2, 4, 8, 23, 30, 31, 33, 46]. These are referred to as *graph-structured multi-marginal optimal transport problems*. For cost functions that decouples sequentially, i.e. are of the form

$$C(i_0, \dots, i_m) = \sum_{j \in [m] \setminus 0} C_j(i_{j-1}, i_j), \quad i_0, \dots, i_m = 1, \dots, n \quad (12)$$

for some matrices $C_1, \dots, C_m \in \mathbb{R}^{n \times n}$, the projection can be computed efficiently using matrix-vector multiplications. This is described in the following proposition.¹

Proposition 2.8 ([23], Proposition 2). *Let the elements of the tensors K and U be of the form*

$$K(i_0, \dots, i_m) = \prod_{j \in [m] \setminus 0} K_j(i_{j-1}, i_j) \quad \text{and} \quad U(i_0, \dots, i_m) = \prod_{j \in [m]} u_j(i_j)$$

for matrices $K_j \in \mathbb{R}^{n \times n}$, $j \in [m] \setminus 0$, and vectors $u_j \in \mathbb{R}^n$, $j \in [m]$. Then,

$$P_j(K \odot U) = \left(u_0^\top K_1 \text{diag}(u_1) K_2 \dots K_{j-1} \text{diag}(u_{j-1}) K_j \right)^\top \odot u_j \\ \odot \left(K_{j+1} \text{diag}(u_{j+1}) K_{j+2} \dots K_{m-1} \text{diag}(u_{m-1}) K_m u_m \right), \quad j \in [m].$$

In Section 4 we will use this and similar results to develop an efficient method for solving our class of multi-marginal MOT problems.

3 Reformulation as a structured OT problem

We now show how each MOT problem of the form (6) can be identified with a structured multi-marginal OT problem whose optimal solution can be computed efficiently. This is carried out in two main steps, where the first one is to reduce the path dependency of the martingale constraint. Once this is done, we will transform the resulting problem to tensor form. A key insight throughout is to leverage on the sequential structure inherent in the problem.

¹In [23], Proposition 2.8 was stated in the special case when the matrices K_j , $j \in [m] \setminus 0$, do not depend on the index j . However, it is straightforward to generalise it to the version given here.

3.1 Markovian reformulation of the problem

When comparing the form (5) of our class of payoff functions with the form (12) of the cost functions characterising the sequentially structured OT problems, we note that they are consistent. Unfortunately, the presence of the martingale constraint introduces a dependency of each marginal to all previous marginals, ruining the structure of the problem. Therefore, our class of MOT problems cannot directly be solved via the techniques mentioned in Section 2.3, despite the payoff function being of the appropriate form. It can be noted though that when the price process S is a Markov process under the pricing measure \mathbb{Q} , the martingale constraint simplifies to only connecting adjacent marginals to each other. This recovers the structure of the problem, so that the projections can be computed efficiently. Leveraging on these insights, consider the modified multi-marginal OT problem

$$\inf_{(\Omega, \mathcal{F}, \mathbb{Q}, S)} \mathbb{E}_{\mathbb{Q}} \left[\sum_{t \in [T] \setminus 0} \phi_t(S_{t-1}, X_{t-1}, S_t, X_t) \right] \quad (13a)$$

$$\text{subject to } \mathbb{E}_{\mathbb{Q}}[S_t | \sigma(S_{t-1}, X_{t-1})] = S_{t-1}, \quad t \in [T] \setminus 0 \quad (13b)$$

$$S_t \sim_{\mathbb{Q}} \mu_t, \quad t \in \mathcal{T} \quad (13c)$$

where X is given in equation (4). Note that since the constraints (13b) and (13c) are linear in \mathbb{Q} , the feasible set of problem (13) is convex. We now show that any optimal solution to problem (13) can be used to find an optimal solution to problem (6).

Theorem 3.1. *The martingale optimal transport problem (6) and the optimal transport problem (13) are equivalent in the sense that any optimal solution of problem (6) is also an optimal solution of problem (13), while any optimal solution of problem (13) can be used to construct an optimal solution to problem (6). Moreover, the optimal objective values coincide.*

The key idea behind the above result is that if a model $(\Omega, \mathcal{F}, \mathbb{Q}, S)$ satisfies equations (13b) and (13c), where the process X is given by equation (4), then there exists a Markov process (\tilde{S}, \tilde{X}) , supported on some probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$, such that equations (13b) and (13c) hold, such that (\tilde{S}, \tilde{X}) is of the form (4), and such that the corresponding value of the objective function (13a) is the same as for the original model $(\Omega, \mathcal{F}, \mathbb{Q}, S)$. In order to prove Theorem 3.1, we first establish a few intermediate results.

Fix any tuple $(\Omega, \mathcal{F}, \mathbb{P}, S)$, consisting of a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ and a stochastic process S on that space, and define a process X out of the process S as in equation (4). Let $\nu : \mathcal{B}(\mathbb{R}^2) \rightarrow [0, 1]$ and $\kappa_t : \mathbb{R}^2 \times \mathcal{B}(\mathbb{R}^2) \rightarrow [0, 1]$, for $t \in [T] \setminus 0$, be defined by

$$\nu(A) := \mathbb{P}((S_0, X_0) \in A), \quad A \in \mathcal{B}(\mathbb{R}^2)$$

and

$$\kappa_t(y, A) := \mathbb{P}((S_t, X_t) \in A | (S_{t-1}, X_{t-1}) = y), \quad A \in \mathcal{B}(\mathbb{R}^2), \quad y \in \mathbb{R}^2,$$

respectively, where we let the regular conditional distribution be equal to some real-valued constant if conditioning on a null event. Then ν is the initial distribution of the joint process (S, X) on \mathbb{R}^2 , while $\{\kappa_t\}_{t=1}^T$ is a family of kernels on $(\mathbb{R}^2, \mathcal{B}(\mathbb{R}^2))$. Indeed, since for every $t \in [T]$ the random element (S_t, X_t) takes values in \mathbb{R}^2 and $(\mathbb{R}^2, \mathcal{B}(\mathbb{R}^2))$ is a Borel space, the regular conditional distribution given above exists and is almost everywhere equal to a kernel (see, for example, [41, Theorem 8.5]). The Kolmogorov extension theorem then guarantees the existence of an \mathbb{R}^2 -valued Markov process (\tilde{S}, \tilde{X}) , defined on some probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$, with initial distribution ν and transition kernels $\{\kappa_t\}_{t=1}^T$ (see, for example, [41, Theorem 11.4]); that is,

$$\tilde{\mathbb{P}}((\tilde{S}_0, \tilde{X}_0) \in A) = \nu(A), \quad A \in \mathcal{B}(\mathbb{R}^2) \quad (14a)$$

$$\tilde{\mathbb{P}}((\tilde{S}_t, \tilde{X}_t) \in A | (\tilde{S}_{t-1}, \tilde{X}_{t-1}) = y) = \kappa_t(y, A), \quad A \in \mathcal{B}(\mathbb{R}^2), \quad y \in \mathbb{R}^2, \quad t \in [T] \setminus 0. \quad (14b)$$

It can be taken as the canonical process on the probability space given by $\tilde{\Omega} = (\mathbb{R}^2)^{T+1}$, $\tilde{\mathcal{F}} = \sigma((\tilde{S}_0, \tilde{X}_0), \dots, (\tilde{S}_T, \tilde{X}_T))$ and $\tilde{\mathbb{P}} = \nu \otimes \kappa_1 \otimes \dots \otimes \kappa_T$.

Lemma 3.2. *Let the stochastic processes (S, X) and (\tilde{S}, \tilde{X}) be as defined above. Then the law of $((\tilde{S}_{t-1}, \tilde{X}_{t-1}), (\tilde{S}_t, \tilde{X}_t))$ is equal to the law of $((S_{t-1}, X_{t-1}), (S_t, X_t))$ for all $t \in [T] \setminus 0$.*

Proof. It is immediate from equation (14a) that $\text{Law}((\tilde{S}_0, \tilde{X}_0)) = \nu = \text{Law}((S_0, X_0))$. Then fix any $t \in [T] \setminus 0$ and assume that

$$\text{Law}((\tilde{S}_{t-1}, \tilde{X}_{t-1})) = \text{Law}((S_{t-1}, X_{t-1})).$$

Let $F_{(\tilde{S}_{t-1}, \tilde{X}_{t-1})}$ and $F_{(S_{t-1}, X_{t-1})}$ denote the joint cumulative distribution functions corresponding to $\text{Law}((\tilde{S}_{t-1}, \tilde{X}_{t-1}))$ and $\text{Law}((S_{t-1}, X_{t-1}))$, respectively. Then $F_{(\tilde{S}_{t-1}, \tilde{X}_{t-1})} = F_{(S_{t-1}, X_{t-1})}$, so for any $A \in \mathcal{B}(\mathbb{R}^2)$,

$$\begin{aligned} \tilde{\mathbb{P}}((\tilde{S}_t, \tilde{X}_t) \in A) &= \int_{\mathbb{R}^2} \kappa_t(y, A) dF_{(\tilde{S}_{t-1}, \tilde{X}_{t-1})}(y) \\ &= \int_{\mathbb{R}^2} \kappa_t(y, A) dF_{(S_{t-1}, X_{t-1})}(y) = \mathbb{P}((S_t, X_t) \in A), \end{aligned}$$

where the second equality follows from equation (14b). Since the above holds for any $t \in [T]$, the assertion follows from this in combination with equation (14b). \square

Remark 3.3. In case the joint process (S, X) is a Markov process under \mathbb{P} , it holds that

$$\text{Law}((\tilde{S}_0, \tilde{X}_0), \dots, (\tilde{S}_T, \tilde{X}_T)) = \text{Law}((S_0, X_0), \dots, (S_T, X_T)).$$

We now show that the Markov process (\tilde{S}, \tilde{X}) is almost surely of the form given in equation (4).

Lemma 3.4. *The process (\tilde{S}, \tilde{X}) satisfies equation (4) almost surely, that is,*

$$\tilde{X}_t = \begin{cases} h_0(\tilde{S}_0), & t = 0 \\ h_t(\tilde{S}_{t-1}, \tilde{S}_t, \tilde{X}_{t-1}), & t = 1, \dots, T \end{cases} \quad a.s.$$

Proof. The assertion for $t = 0$ follows from Lemma 3.2 after noting that

$$\{\tilde{X}_0 = h_0(\tilde{S}_0)\} = \{(h_0(\tilde{S}_0), \tilde{X}_0) \in \{(x, x) : x \in \mathbb{R}\}\}$$

and similarly for the event $\{X_0 = h_0(S_0)\}$. Indeed,

$$\begin{aligned} \tilde{\mathbb{P}}((h_0(\tilde{S}_0), \tilde{X}_0) \in \{(x, x) : x \in \mathbb{R}\}) &= \tilde{\mathbb{P}}((\tilde{S}_0, \tilde{X}_0) \in \{(h_0^{-1}(x), x) : x \in \mathbb{R}\}) \\ &= \mathbb{P}((S_0, X_0) \in \{(h_0^{-1}(x), x) : x \in \mathbb{R}\}) \\ &= \mathbb{P}((h_0(S_0), X_0) \in \{(x, x) : x \in \mathbb{R}\}), \end{aligned}$$

where the right-hand side is equal to one since by definition $X_0 = h_0(S_0)$. By similar arguments for $t \in [T] \setminus 0$, $\tilde{\mathbb{P}}(\tilde{X}_t = h_t(\tilde{S}_{t-1}, \tilde{S}_t, \tilde{X}_{t-1})) = \mathbb{P}(X_t = h_t(S_{t-1}, S_t, X_{t-1}))$. The assertion for $t \in [T] \setminus 0$ follows, which completes the proof. \square

We are now ready to prove Theorem 3.1.

Proof of Theorem 3.1. Let a tuple $(\Omega, \mathcal{F}, \mathbb{Q}, S)$ refer to a probability space $(\Omega, \mathcal{F}, \mathbb{Q})$ supporting a stochastic process $S : \Omega \times [T] \rightarrow \mathbb{R}$. Define two sets of tuples \mathfrak{F} and \mathfrak{F}' as

$$\mathfrak{F} := \{(\Omega, \mathcal{F}, \mathbb{Q}, S) : S_t \sim_{\mathbb{Q}} \mu_t \quad \forall t \in \mathcal{T} \quad \text{and} \quad \mathbb{E}_{\mathbb{Q}}[S_t | \sigma(S_0, \dots, S_{t-1})] = S_{t-1} \quad \forall t \in [T] \setminus 0\}$$

and

$$\mathfrak{F}' := \{(\Omega, \mathcal{F}, \mathbb{Q}, S) : S_t \sim_{\mathbb{Q}} \mu_t \quad \forall t \in \mathcal{T} \quad \text{and} \quad \mathbb{E}_{\mathbb{Q}}[S_t | \sigma(S_{t-1}, X_{t-1})] = S_{t-1} \quad \forall t \in [T] \setminus 0\},$$

respectively, where X is defined out of S as in equation (4). We note that \mathfrak{F} is the feasible set of problem (6) and that \mathfrak{F}' is the feasible set of problem (13). Take any tuple $(\Omega, \mathcal{F}, \mathbb{Q}, S) \in \mathfrak{F}$. Then, by the Tower property, for $t \in [T] \setminus 0$,

$$\begin{aligned} \mathbb{E}_{\mathbb{Q}}[S_t | \sigma(S_{t-1}, X_{t-1})] &= \mathbb{E}_{\mathbb{Q}}[\mathbb{E}_{\mathbb{Q}}[S_t | \sigma(S_0, \dots, S_{t-1})] | \sigma(S_{t-1}, X_{t-1})] \\ &= \mathbb{E}_{\mathbb{Q}}[S_{t-1} | \sigma(S_{t-1}, X_{t-1})] = S_{t-1} \quad \text{a.s.} \end{aligned}$$

It follows that $(\Omega, \mathcal{F}, \mathbb{Q}, S) \in \mathfrak{F}'$. Therefore $\mathfrak{F} \subseteq \mathfrak{F}'$.

Take any tuple $(\Omega, \mathcal{F}, \mathbb{Q}, S) \in \mathfrak{F}'$ and define X out of S as given in equation (4). Then note that there exists an \mathbb{R}^2 -valued Markov process (\tilde{S}, \tilde{X}) , supported on a probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{Q}})$, that has the same initial distribution and transition kernels as the joint process (S, X) (cf. (14)). Since by Lemma 3.2 the law of $(\tilde{S}_{t-1}, \tilde{S}_t, \tilde{X}_{t-1}, \tilde{X}_t)$ under $\tilde{\mathbb{Q}}$ is equal to the law of $(S_{t-1}, S_t, X_{t-1}, X_t)$ under \mathbb{Q} for every $t \in [T] \setminus 0$, it follows that the law of \tilde{S}_t under $\tilde{\mathbb{Q}}$ is equal to the law of S_t under \mathbb{Q} for every $t \in [T]$. Therefore $\tilde{S}_t \sim_{\tilde{\mathbb{Q}}} \mu_t$ for $t \in \mathcal{T}$. By similar arguments,

$$\mathbb{E}_{\tilde{\mathbb{Q}}}[\tilde{S}_t | \tilde{S}_{t-1} = s, \tilde{X}_{t-1} = x] = \mathbb{E}_{\mathbb{Q}}[S_t | S_{t-1} = s, X_{t-1} = x], \quad s, x \in \mathbb{R}, \quad t \in [T] \setminus 0.$$

Then note that by definition of \mathfrak{F}' , the joint process (S, X) satisfies $\mathbb{E}_{\mathbb{Q}}[S_t | S_{t-1} = s, X_{t-1} = x] = s$ for every $s, x \in \mathbb{R}$ and $t \in [T] \setminus 0$. Combining this with the above and with the Markov property of (\tilde{S}, \tilde{X}) yields that \tilde{S} is a martingale with respect to its own filtration under $\tilde{\mathbb{Q}}$. Therefore, $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{Q}}, \tilde{S}) \in \mathfrak{F}$. Moreover, it follows from Lemma 3.2 that

$$\mathbb{E}_{\tilde{\mathbb{Q}}} \left[\sum_{t \in [T] \setminus 0} \phi_t(\tilde{S}_{t-1}, \tilde{X}_{t-1}, \tilde{S}_t, \tilde{X}_t) \right] = \mathbb{E}_{\mathbb{Q}} \left[\sum_{t \in [T] \setminus 0} \phi_t(S_{t-1}, X_{t-1}, S_t, X_t) \right],$$

where the left-hand side is the payoff associated with the model $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{Q}}, \tilde{S})$ due to Lemma 3.4. We have thus shown that for any $(\Omega, \mathcal{F}, \mathbb{Q}, S) \in \mathfrak{F}'$ we can find $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{Q}}, \tilde{S}) \in \mathfrak{F}$ such that their corresponding values of the objective function are equal. This implies that the values of the two problems coincide.

It follows that if $(\Omega^*, \mathcal{F}^*, \mathbb{Q}^*, S^*) \in \mathfrak{F}'$ is an optimal solution of problem (13) there exists $(\tilde{\Omega}^*, \tilde{\mathcal{F}}^*, \tilde{\mathbb{Q}}^*, \tilde{S}^*) \in \mathfrak{F}$ that is an optimal solution of problem (6). Assume on the other hand that $(\Omega^*, \mathcal{F}^*, \mathbb{Q}^*, S^*) \in \mathfrak{F}$ is an optimal solution of problem (6). Since the two problem values are equal and $\mathfrak{F} \subseteq \mathfrak{F}'$, it immediately follows that then $(\Omega^*, \mathcal{F}^*, \mathbb{Q}^*, S^*)$ is an optimal solution also of problem (13). This completes the proof. \square

Remark 3.5. Note that when (S, X) is not a Markov process under \mathbb{Q} , it does not necessarily hold that

$$\mathbb{E}_{\mathbb{Q}}[S_t | \sigma(S_0, \dots, S_{t-1})] = \mathbb{E}_{\mathbb{Q}}[S_t | \sigma(S_{t-1}, X_{t-1})] \quad \text{a.s.,} \quad t \in [T] \setminus 0. \quad (15)$$

Therefore, in general $(\Omega, \mathcal{F}, \mathbb{Q}, S) \notin \mathfrak{F}$ when $(\Omega, \mathcal{F}, \mathbb{Q}, S) \in \mathfrak{F}'$. Consequently $\mathfrak{F} \neq \mathfrak{F}'$. From this it can be noted that \mathfrak{F} can be characterised as the subset of tuples $(\Omega, \mathcal{F}, \mathbb{Q}, S) \in \mathfrak{F}'$ such that equation (15) holds. We immediately realise that every tuple $(\Omega, \mathcal{F}, \mathbb{Q}, S) \in \mathfrak{F}'$ that is such that (S, X) is a Markov process under \mathbb{Q} also belongs to \mathfrak{F} . On the other hand, note that every tuple in \mathfrak{F}' does not correspond to an \mathbb{R}^2 -valued Markov process. On a related note, one can show that the multi-marginal MOT problem that is subject an additional Markov constraint is equivalent to problem (13) in the same way as we have shown that problem (6) is equivalent to problem (13). This means that there are in total three problems that are all equivalent to each other, given that the payoff function is of the form (5). This holds despite the fact that this third problem is not convex.

Remark 3.6. If we know that the optimal solution to a MOT problem is such that the price process is a Markov process, the process X can be omitted; see this by letting $X_t = S_t$ for $t \in [T]$ in the above results. This reduces the size of the problem. This holds in particular for all bi-marginal problems since a stochastic process defined over two discrete time points trivially is a Markov process. Along the same lines, we emphasise that MOT problems characterised by a payoff of the form $\phi(S_0, \dots, S_T) = \sum_{t \in [T] \setminus 0} \phi_t(S_{t-1}, S_t)$, $\phi_t : \mathbb{R}^2 \rightarrow \mathbb{R}$, can be shown to be equivalent with an OT problem subject to the constraint $\mathbb{E}_{\mathbb{Q}}[S_t | \sigma(S_{t-1})] = S_{t-1}$ for $t \in [T] \setminus 0$.

3.2 Formulation as a linear programming problem

We have now shown that the MOT problem (6) is equivalent to the OT problem (13), so that instead of computationally solving problem (6) we can solve problem (13), which has the sequential structure required for obtaining an efficient algorithm. In order to develop a computational method, we now consider the case when the support of the marginals is concentrated on a finite number of points — the objective function as well as the constraints of problem (13) can then be expressed in terms of tensors. We will now formalise this. With slight abuse of terminology, we refer to equation (13b) as ‘martingale constraints’ from now on.

Suppose that the support of the given marginals $\{\mu_t\}_{t \in \mathcal{T}}$ of the price process is concentrated on a finite number of points. This is the situation for information given by a real-world financial market. Let the support of the given marginals be contained within the discrete set $\mathcal{S} \subset \mathbb{R}$ given by

$$\mathcal{S} = \{s_1, \dots, s_{n_S}\}$$

where $n_S \in \mathbb{N}$ and $s_1, \dots, s_{n_S} \in \mathbb{R}$. Let $s \in \mathbb{R}^{n_S}$ be the vector consisting of the elements in \mathcal{S} , that is

$$s(i) := s_i, \quad i = 1, \dots, n_S.$$

For each given marginal μ_t we define a vector $\mu_t \in \mathbb{R}^{n_S}$ where each element is given by

$$\mu_t(i) := \mu_t(s(i)), \quad i = 1, \dots, n_S. \quad (16)$$

For practical convenience, we limit ourselves to only consider models $(\Omega, \mathcal{F}, \mathbb{Q}, S)$ where the support of S_t is contained in \mathcal{S} also for $t \in [T] \setminus \mathcal{T}$. Let X be defined in terms of the price process S as in equation (4) and let its (finite) range in \mathbb{R} be denoted by \mathcal{X} . Then \mathcal{X} is of the form

$$\mathcal{X} = \{x_1, \dots, x_{n_X}\}$$

for some $n_X \in \mathbb{N}$ and $x_1, \dots, x_{n_X} \in \mathbb{R}$. Let $x \in \mathbb{R}^{n_X}$ be the vector consisting of the elements of \mathcal{X} , that is

$$x(i) := x_i, \quad i = 1, \dots, n_X.$$

Note that depending on the choice of $h_t, t \in [T]$, we could have that $\mathcal{X} \equiv \mathcal{S}$ — this is the case for Examples 2.1 and 2.6.

Let us now take a model $(\Omega, \mathcal{F}, \mathbb{Q}, S)$ satisfying the above assumptions. For each $t \in [T]$ and almost all $\omega \in \Omega$ the price S then takes one of the values represented by s ,

$$S_t(\omega) = s(\omega_t^S),$$

for some $\omega_t^S \in \{1, \dots, n_S\}$. Note that ω_t^S denotes an index running from 1 to n_S . Also note that each time point t is characterised by its own index so that in total we have $T + 1$ indices $\omega_0^S, \dots, \omega_T^S$, all running from 1 to n_S . Then the index tuple $(\omega_0^S, \dots, \omega_T^S)$ represents the outcome $\omega \in \Omega$. Similarly, the process X at time $t \in [T]$ and for almost all $\omega \in \Omega$ takes one of the values represented by x ,

$$X_t(\omega) = x(\omega_t^X)$$

where $\omega_t^X \in \{1, \dots, n_X\}$. Here ω_t^X is an index running from 1 to n_X . By constructing a joint index

$$\omega_t := (\omega_t^S, \omega_t^X), \quad t = 0, \dots, T,$$

a joint index tuple

$$\boldsymbol{\omega} := (\omega_0, \dots, \omega_T) = ((\omega_0^S, \omega_0^X), \dots, (\omega_T^S, \omega_T^X))$$

and a joint index set

$$\boldsymbol{\Omega} := (\{1, \dots, n_S\} \times \{1, \dots, n_X\})^{T+1},$$

we can identify each sample $\omega \in \Omega$ with an element $\boldsymbol{\omega} \in \boldsymbol{\Omega}$ via

$$\left((S_0(\omega), X_0(\omega)), \dots, (S_T(\omega), X_T(\omega)) \right) = \left((s(\omega_0^S), x(\omega_0^X)), \dots, (s(\omega_T^S), x(\omega_T^X)) \right).$$

The index tuple $\boldsymbol{\omega} \in \boldsymbol{\Omega}$ thus represents one possible trajectory for the joint process (S, X) .

We will now have a look at how the payoff can be represented within this framework. Let $\Phi : \boldsymbol{\Omega} \rightarrow \mathbb{R}$ be given by

$$\Phi(\boldsymbol{\omega}) := \phi(s(\omega_0^S), \dots, s(\omega_T^S)), \quad \boldsymbol{\omega} \in \boldsymbol{\Omega}.$$

Then $\Phi(\boldsymbol{\omega})$ represents the payoff associated with the trajectory $\boldsymbol{\omega} \in \boldsymbol{\Omega}$. Also define Φ_t for $t \in [T] \setminus 0$ by

$$\Phi_t(\omega_{t-1}, \omega_t) := \phi_t(s(\omega_{t-1}^S), x(\omega_{t-1}^X), s(\omega_t^S), x(\omega_t^X)), \quad \omega_{t-1}, \omega_t \in \{1, \dots, n_S\} \times \{1, \dots, n_X\}.$$

Then

$$\Phi(\boldsymbol{\omega}) = \sum_{t \in [T] \setminus 0} \Phi_t(\omega_{t-1}, \omega_t), \quad \boldsymbol{\omega} \in \boldsymbol{\Omega}, \quad (17)$$

analogously to equation (5). In order to enforce a zero probability for every trajectory $\boldsymbol{\omega} \in \boldsymbol{\Omega}$ that does not respect equation (4), define a penalty term by

$$I(\boldsymbol{\omega}) := \sum_{t \in [T] \setminus 0} I_t(\omega_{t-1}, \omega_t), \quad \boldsymbol{\omega} \in \boldsymbol{\Omega},$$

with

$$I_t(\omega_{t-1}, \omega_t) := \begin{cases} 0 & \text{if } x(\omega_t^X) = h_t(s(\omega_t^S), s(\omega_{t-1}^S), x(\omega_{t-1}^X)) \\ \infty & \text{else} \end{cases}, \quad t = 2, \dots, T$$

and

$$I_1(\omega_0, \omega_1) := \begin{cases} 0 & \text{if } x(\omega_0^X) = h_0(s(\omega_0^S)) \text{ and } x(\omega_1^X) = h_1(s(\omega_1^S), s(\omega_0^S), x(\omega_0^X)) \\ \infty & \text{else.} \end{cases}$$

By adding the penalty term I to the payoff, we penalise forbidden trajectories $\boldsymbol{\omega} \in \boldsymbol{\Omega}$. Therefore, define C_t for $t \in [T] \setminus 0$ by

$$C_t(\omega_{t-1}, \omega_t) := \Phi_t(\omega_{t-1}, \omega_t) + I_t(\omega_{t-1}, \omega_t), \quad \omega_{t-1}, \omega_t \in \{1, \dots, n_S\} \times \{1, \dots, n_X\},$$

and let $C(\boldsymbol{\omega}) := \sum_{t \in [T] \setminus 0} C_t(\omega_{t-1}, \omega_t)$ for $\boldsymbol{\omega} \in \boldsymbol{\Omega}$. Note that C inherits the structure of Φ and I .

We have now presented the notation needed for formulating problem (13) as a linear programming problem; before doing so, we present some alternative notation that will simplify the expressions. By ordering the joint indices $\{1, \dots, n_S\} \times \{1, \dots, n_X\}$ according to

$$(1, 1), \dots, (n_S, 1), (1, 2), \dots, (n_S, 2), \dots, (1, n_X), \dots, (n_S, n_X), \quad (18)$$

Φ, I and C can be identified with $T + 1$ -dimensional tensors in $\mathbb{R}^{(n_S n_X)^{T+1}}$. Similarly Φ_t, I_t and C_t , $t \in [T] \setminus 0$, can be represented by matrices in $\mathbb{R}^{n_S n_X \times n_S n_X}$. Define the matrix $\Delta \in \mathbb{R}^{n_S n_X \times n_S n_X}$ as

$$\Delta(i, j) := (\mathbf{1}_{n_X} \otimes s)(j) - (\mathbf{1}_{n_X} \otimes s)(i), \quad i, j \in \{1, \dots, n_S\} \times \{1, \dots, n_X\}, \quad (19)$$

where it is assumed that the indices are ordered along each axis according to the rule given by (18). Recall the definitions of the projection operators P_t , for $t \in [T]$, and P_{t_1, t_2} , for $t_1, t_2 \in [T]$. They can be identified with maps from $\mathbb{R}_+^{(n_S n_X)^{T+1}}$ to $\mathbb{R}_+^{n_S n_X}$ and $\mathbb{R}_+^{n_S n_X \times n_S n_X}$, respectively, when the indices are ordered according to (18). Define $P^S : \mathbb{R}_+^{n_S n_X} \rightarrow \mathbb{R}_+^{n_S}$ as

$$P^S(v)(\omega_t^S) := \sum_{\omega_t^X=1}^{n_X} v((\omega_t^S, \omega_t^X)), \quad \omega_t^S = 1, \dots, n_S$$

where $v \in \mathbb{R}_+^{n_S n_X}$ and let $P_t^S := P^S \circ P_t$ for $t \in [T]$.

Given the framework introduced above, consider the linear programming problem

$$\min_{Q \in \mathbb{R}_+^{(n_S n_X)^{T+1}}} \langle C, Q \rangle \quad (20a)$$

$$\text{subject to } (P_{t-1,t}(Q) \odot \Delta) \mathbf{1}_{n_S n_X} = \mathbf{0}_{n_S n_X}, \quad t \in [T] \setminus 0 \quad (20b)$$

$$P_t^S(Q) = \mu_t, \quad t \in \mathcal{T}. \quad (20c)$$

We now show that problems (13) and (20) are equivalent.

Proposition 3.7. *Suppose that we restrict problem (13) to models such that the support of the price process at each time point is contained within \mathcal{S} . Then problems (13) and (20) are equivalent.*

Proof. Let $(\Omega, \mathcal{F}, \mathbb{Q}, S)$ be a model feasible to problem (13) and note that we can define a probability measure Q on $(\Omega, \sigma(\Omega))$ by

$$Q(\omega) := \mathbb{Q} \left(\bigcap_{t=0}^T \left\{ S_t = s(\omega_t^S), X_t = x(\omega_t^X) \right\} \right), \quad \omega \in \Omega. \quad (21)$$

It can be identified with a tensor, indexed by (18), in $\mathbb{R}_+^{(n_S n_X)^{T+1}}$. Starting with the objective function, it holds that

$$\mathbb{E}_{\mathbb{Q}} \left[\sum_{t \in [T] \setminus 0} \phi(S_{t-1}, X_{t-1}, S_t, X_t) \right] = \sum_{t \in [T] \setminus 0} \langle \Phi_t, P_{t-1,t}(Q) \rangle = \langle \Phi, Q \rangle = \langle C, Q \rangle,$$

where the last equality follows from the fact that $\langle I, Q \rangle = 0$ for any tensor Q defined as in equation (21). As for the marginal constraints (13c), it follows immediately from the definition of Q , for $t \in [T]$, that

$$\mathbb{Q}(S_t = s(i)) = P_t^S(Q)(i), \quad i = 1, \dots, n_S.$$

Combining this elementwise with equation (16) and the constraint (13c) yields the constraint (20c). Similarly, for $t \in [T] \setminus 0$, it holds that

$$\mathbb{E}_{\mathbb{Q}} [S_t | S_{t-1} = s(\omega_{t-1}^S), X_{t-1} = x(\omega_{t-1}^X)] = \frac{(P_{t-1,t}(Q)(\mathbf{1}_{n_X} \otimes s))(\omega_{t-1})}{P_{t-1}(Q)(\omega_{t-1})}$$

for $\omega_{t-1} \in \{1, \dots, n_S\} \times \{1, \dots, n_X\}$, since $s(\omega_t^S) = (\mathbf{1}_{n_X} \otimes s)(\omega_t)$ when $\omega_t \in \{1, \dots, n_S\} \times \{1, \dots, n_X\}$ is ordered according to (18). By combining the martingale constraint (13b) elementwise with this we obtain

$$P_{t-1,t}(Q)(\mathbf{1}_{n_X} \otimes s) = (\mathbf{1}_{n_X} \otimes s) \odot P_{t-1}(Q), \quad t = 1, \dots, T.$$

This is the martingale constraint (13b) formulated on vector form. By subtracting the right-hand side from the left-hand side and identifying the matrix $\Delta \in \mathbb{R}^{n_S n_X \times n_S n_X}$ the constraint (20b) follows.

Conversely, let $Q \in \mathbb{R}_+^{(n_S n_X)^{T+1}}$ be feasible to problem (20). Then $(\Omega, \sigma(\Omega), Q)$ is a probability space, on which we can identify Φ as a random variable. Assume that Q is such that $\langle I, Q \rangle = 0$. Then

$$\langle C, Q \rangle = \langle \Phi, Q \rangle = \mathbb{E}_Q[\Phi],$$

which means that the values of the objective functions (20a) and (13a) coincide. Also note that the model $(\Omega, \sigma(\Omega), Q, s^{T+1})$, induced by the tensor Q satisfying the constraints (20b) and (20c), satisfies the constraints (13b) and (13c). It remains to argue that there always exists a Q such that $\langle I, Q \rangle = 0$ whenever the feasible set of problem (20) is nonempty. Do so by noting that the feasible set of problem (13) is nonempty and that every feasible model $(\Omega, \mathcal{F}, \mathbb{Q}, S)$ induces Q such that $\langle I, Q \rangle = 0$. This completes the proof. \square

4 Solving the structured problem via regularisation

We have now arrived at a version of the problem that can be approached computationally. We therefore proceed by deriving a generalisation of the algorithm described in Section 2.3, modified to also take the martingale constraint into account. As in Section 2.3, this is done by adding a regularising entropy term $D(Q) := \sum_{\omega \in \Omega} (Q(\omega) \log Q(\omega) - Q(\omega) + 1)$, scaled by some small number $\varepsilon > 0$, to the objective function of problem (20). The entropy regularised version of problem (13) written on tensor form thus becomes

$$\min_{Q \in \mathbb{R}_+^{(n_S n_X)^{T+1}}} \langle C, Q \rangle + \varepsilon D(Q) \quad (22a)$$

$$\text{subject to } (P_{t-1,t}(Q) \odot \Delta) \mathbf{1}_{n_S n_X} = \mathbf{0}_{n_S n_X}, \quad t \in [T] \setminus 0 \quad (22b)$$

$$P_t^S(Q) = \mu_t, \quad t \in \mathcal{T}, \quad (22c)$$

where we recall that the tensor C is of the form $C(\omega) = \sum_{t \in [T] \setminus 0} C_t(\omega_{t-1}, \omega_t)$, $\omega \in \Omega$, for matrices $C_t \in \mathbb{R}^{n_S n_X \times n_S n_X}$. Note that the above problem is no longer linear, but rather strictly convex, due to the entropy term $D(Q)$ being strictly convex in Q . Since it is a convex problem whose feasible set has non-empty relative interior, one can show (see, for example, [47]) that strong duality holds. We can therefore solve (22) by considering the dual problem. We will see that the optimality conditions for the dual variables correspond to relatively simple equations, where explicit expressions can be obtained for the dual variables corresponding to marginal constraints, while for the dual variables corresponding to martingale constraints it turns out that T sets of equations must be solved numerically. This is a generalisation of the strategy used by [18] for the bi-marginal MOT problem. We then exploit the structure inherent in the problem to simplify the equations obtained; the resulting algorithm is a multi-marginal version of Sinkhorn's algorithm where some of the variables are updated by using Newton's method.

4.1 Lagrangian relaxation and coordinate dual ascent

In order to derive the dual of problem (22), we start by forming the corresponding Lagrangian function. It reads

$$\mathcal{L}(Q, \lambda, \gamma) = \langle C, Q \rangle + \varepsilon D(Q) + \sum_{t \in \mathcal{T}} [\lambda_t^\top (\mu_t - P_t^S(Q))] - \sum_{t \in [T] \setminus 0} [\gamma_{t-1}^\top (P_{t-1,t}(Q) \odot \Delta) \mathbf{1}_{n_S n_X}],$$

where $\lambda = \{\lambda_t\}_{t \in \mathcal{T}}$ are Lagrangian multipliers (dual variables) corresponding to the marginal constraints (22c) and $\gamma = \{\gamma_t\}_{t=0}^{T-1}$ are Lagrangian multipliers corresponding to the martingale constraints (22b). Note that each λ_t is a vector of length n_S , while each γ_t is a vector of length $n_S n_X$. The dual of problem (22) is then

$$\max_{\lambda, \gamma} \varphi(\lambda, \gamma),$$

where the dual functional is defined as $\varphi(\lambda, \gamma) := \min\{\mathcal{L}(Q, \lambda, \gamma) : Q \in \mathbb{R}_+^{(n_S n_X)^{T+1}}\}$. We proceed by finding the minimising $Q_{\lambda, \gamma}$ for given dual variables λ, γ . Since the Lagrangian function \mathcal{L} is strictly convex in Q and the entropy term $D(Q)$ prevents optimal solutions at the boundary, the minimising transport plan satisfies $\partial_{Q(\omega)} \mathcal{L}(Q, \lambda, \gamma) = 0$ for $\omega \in \Omega$. By differentiating the Lagrangian function we obtain that for any $\omega \in \Omega$

$$\partial_{Q(\omega)} \mathcal{L}(Q, \lambda, \gamma) = \varepsilon \log(Q(\omega)) - \sum_{t \in \mathcal{T}} \lambda_t(\omega_t^S) + \sum_{t \in [T] \setminus 0} (C_t(\omega_{t-1}, \omega_t) - \gamma_{t-1}(\omega_{t-1}) \Delta(\omega_{t-1}, \omega_t))$$

which yields a minimising transport plan

$$Q_{\lambda, \gamma}(\omega) = \left(\prod_{t \in \mathcal{T}} e^{\lambda_t(\omega_t^S)/\varepsilon} \right) \left(\prod_{t \in [T] \setminus 0} e^{-C_t(\omega_{t-1}, \omega_t)/\varepsilon} e^{\gamma_{t-1}(\omega_{t-1}) \Delta(\omega_{t-1}, \omega_t)/\varepsilon} \right), \quad \omega \in \Omega, \quad (23)$$

for which the first term of the scaled entropy $\varepsilon D(Q_{\lambda,\gamma})$ equals

$$\begin{aligned} & \varepsilon \sum_{\omega \in \Omega} Q_{\lambda,\gamma}(\omega) \log(Q_{\lambda,\gamma}(\omega)) \\ &= \sum_{\omega \in \Omega} Q_{\lambda,\gamma}(\omega) \left(\sum_{t \in \mathcal{T}} \lambda_t(\omega_t^S) - \sum_{t \in [T] \setminus 0} (C_t(\omega_{t-1}, \omega_t) - \gamma_{t-1}(\omega_{t-1}) \Delta(\omega_{t-1}, \omega_t)) \right) \\ &= \sum_{t \in \mathcal{T}} \lambda_t^\top P_t^S(Q_{\lambda,\gamma}) - \langle C, Q_{\lambda,\gamma} \rangle + \sum_{t \in [T] \setminus 0} \gamma_{t-1}^\top (P_{t-1,t}(Q_{\lambda,\gamma}) \odot \Delta) \mathbf{1}_{n_S n_X}. \end{aligned}$$

Inserting $Q_{\lambda,\gamma}$ into $\mathcal{L}(\cdot, \lambda, \gamma)$ thus yields that the objective of the dual of problem (22) becomes

$$\begin{aligned} & \varphi(\lambda, \gamma) \\ &= \sum_{t \in \mathcal{T}} \lambda_t^\top \mu_t + \varepsilon \sum_{\omega \in \Omega} \left[1 - \left(\prod_{t \in \mathcal{T}} e^{\lambda_t(\omega_t^S)/\varepsilon} \right) \left(\prod_{t \in [T] \setminus 0} e^{-C_t(\omega_{t-1}, \omega_t)/\varepsilon} e^{\gamma_{t-1}(\omega_{t-1}) \Delta(\omega_{t-1}, \omega_t)/\varepsilon} \right) \right]. \end{aligned}$$

At this point, we note that φ can be expressed in a more compact format — a format that reduces the number of direct applications of the exponential function later on — by reparameterising the dual variables. Therefore, let u_t for $t \in \mathcal{T}$ be a family of vectors in $\mathbb{R}^{n_S n_X}$ given by

$$u_t(\omega_t^S) := e^{\lambda_t(\omega_t^S)/\varepsilon}, \quad \omega_t^S = 1, \dots, n_S, \quad (24)$$

and let K_t, G_t , for $t \in [T] \setminus 0$, be two families of matrices in $\mathbb{R}^{n_S n_X \times n_S n_X}$, defined by

$$K_t(\omega_{t-1}, \omega_t) := e^{-C_t(\omega_{t-1}, \omega_t)/\varepsilon}, \quad G_t(\omega_{t-1}, \omega_t) := e^{\gamma_{t-1}(\omega_{t-1}) \Delta(\omega_{t-1}, \omega_t)/\varepsilon}, \quad (25)$$

for $\omega_{t-1}, \omega_t \in \{1, \dots, n_S\} \times \{1, \dots, n_X\}$. Then $u = \{u_t\}_{t \in \mathcal{T}}$ are variables associated with the marginal constraints (22c), while G_t is a matrix representing the t^{th} martingale constraint (22b). It connects the $(t-1)^{\text{th}}$ and t^{th} marginals to each other. The matrix K_t represents the cost associated with moving from the $(t-1)^{\text{th}}$ to the t^{th} marginal. Let $K, U_\lambda, G_\gamma \in \mathbb{R}^{(n_S n_X)^{T+1}}$ be tensors defined by

$$K(\omega) := \prod_{t \in [T] \setminus 0} K_t(\omega_{t-1}, \omega_t), \quad U_\lambda(\omega) := \prod_{t \in \mathcal{T}} u_t(\omega_t^S), \quad G_\gamma(\omega) := \prod_{t \in [T] \setminus 0} G_t(\omega_{t-1}, \omega_t), \quad (26)$$

for $\omega \in \Omega$. Also define

$$U_\lambda^{-r}(\omega) := \prod_{t \in \mathcal{T} \setminus r} u_t(\omega_t^S), \quad r \in \mathcal{T}, \quad G_\gamma^{-q}(\omega) := \prod_{t \in [T] \setminus \{0, q\}} G_t(\omega_{t-1}, \omega_t), \quad q \in [T] \setminus 0, \quad \omega \in \Omega.$$

In this notation, the expression for $Q_{\lambda,\gamma}$ in equation (23) becomes

$$Q_{\lambda,\gamma} = K \odot U_\lambda \odot G_\gamma. \quad (27)$$

This is the minimising primal variable Q , given dual variables λ, γ . The dual functional can be written as $\varphi(\lambda, \gamma) = \sum_{t \in \mathcal{T}} \lambda_j^\top \mu_j - \varepsilon \langle K, U_\lambda \odot G_\gamma \rangle + \varepsilon (n_S n_X)^{T+1}$ using the new notation; the dual problem thus reads

$$\max_{\lambda, \gamma} \sum_{t \in \mathcal{T}} \lambda_j^\top \mu_j - \varepsilon \langle K, U_\lambda \odot G_\gamma \rangle + \varepsilon (n_S n_X)^{T+1}. \quad (28)$$

It remains to find λ, γ such that the concave functional φ is maximised — that is, that satisfies the following set of equations

$$\partial_{\lambda_{t(i)}} \varphi(\lambda, \gamma) = 0, \quad t \in \mathcal{T}, \quad i \in \{1, \dots, n_S\} \quad (29a)$$

$$\partial_{\gamma_{t(i)}} \varphi(\lambda, \gamma) = 0, \quad t \in [T-1], \quad i \in \{1, \dots, n_S\} \times \{1, \dots, n_X\}. \quad (29b)$$

Once they are known, they can be used to recover the optimal transport plan $Q_{\lambda,\gamma}$ via equations (24) – (27).

Starting with the dual variables corresponding to marginal constraints in the primal problem, we differentiate and obtain

$$\partial_{\lambda_t(i)}\varphi(\lambda,\gamma) = \mu_t(i) - \sum_{\substack{\omega \in \Omega: \\ \omega_t^S = i}} K(\omega)G_\gamma(\omega) \prod_{r \in \mathcal{T}} e^{\lambda_r(\omega_r^S)/\varepsilon}, \quad t \in \mathcal{T}, \quad i \in \{1, \dots, n_S\}.$$

Solving for $u_t = \exp(\lambda_t/\varepsilon)$ thus yields

$$u_t = \mu_t \odot P_t^S(K \odot U_\lambda^{-t} \odot G_\gamma), \quad t \in \mathcal{T}. \quad (30)$$

This is the equation characterising the representation u of the optimal dual variables corresponding to marginal constraints in problem (22). Note that the right-hand side of equation (30) can be written as $(\mu_t \odot u_t) \odot P_t^S(K \odot U_\lambda \odot G_\gamma)$. Equation (30) thus requires that u_t should be chosen such that the corresponding $Q_{\lambda,\gamma}$ satisfies $P_t^S(Q_{\lambda,\gamma}) = \mu_t$, which we recognise as the constraint (22c).

Moving on to the dual variables corresponding to martingale constraints in the primal problem, we differentiate and get

$$\partial_{\gamma_t(j)}\varphi(\lambda,\gamma) = - \sum_{\substack{\omega \in \Omega: \\ \omega_t = j}} \Delta(j, \omega_{t+1})K(\omega)U_\lambda(\omega) \prod_{t \in [T] \setminus 0} e^{\gamma_{r-1}(\omega_{r-1})\Delta(\omega_{r-1}, \omega_r)/\varepsilon},$$

for $t \in [T-1]$ and $j \in \{1, \dots, n_S\} \times \{1, \dots, n_X\}$. When solving for γ such that $\partial_{\gamma_t(j)}\varphi(\lambda,\gamma) = 0$ for all t and j as given above, we do not get any explicit formulas. Instead we obtain for every $t \in [T-1]$, $n_S n_X$ equations in $\gamma_t(j)$, $j \in \{1, \dots, n_S\} \times \{1, \dots, n_X\}$. They can be written on vector form as

$$(P_{t,t+1}(K \odot U_\lambda \odot G_\gamma) \odot \Delta) \mathbf{1}_{n_S n_X} = \mathbf{0}_{n_S n_X}, \quad t \in [T-1], \quad (31)$$

where the above should be solved with respect to γ_t , noting that $G_\gamma = \exp(\text{diag}(\gamma_t)\Delta/\varepsilon) \odot G_\gamma^{-(t+1)}$. We recognise equation (31) as constraint (22b). We emphasise that there are no cross-dependencies between the elements of γ_t in equation (31); this will turn out to be useful when deriving a Newton scheme later on.

At this point, we can formulate a high-level method for solving problem (22). It is summarised in Algorithm 1.

Algorithm 1 High-level method for solving problem (22).

```

Initialise:  $u_t \leftarrow \mathbf{1}_{n_S}$  for  $t \in [T]$ 
            $\gamma_t \leftarrow \mathbf{1}_{n_S n_X}$  for  $t \in [T-1]$ 
while not converged do
  for  $t \in \mathcal{T}$  do
     $u_t \leftarrow \mu_t \odot P_t^S(K \odot U_\lambda^{-t} \odot G_\gamma)$ 
  end for
  for  $t = T-1, \dots, 0$  do
    find  $\hat{\gamma}_t$  such that  $(P_{t,t+1}(K \odot U_\lambda \odot G_\gamma) \odot \Delta) \mathbf{1}_{n_S n_X} = \mathbf{0}_{n_S n_X}$ 
     $\gamma_t \leftarrow \hat{\gamma}_t$ 
  end for
end while
 $Q_{\lambda,\gamma} \leftarrow K \odot U_\lambda \odot G_\gamma$ 
return  $Q_{\lambda,\gamma}$ 

```

4.2 Exploiting the structure for computing the projections

The computational bottleneck of Algorithm 1 is the evaluation of the projections; this is because the number of elements in the tensors grows exponentially in the number of marginals. A key ingredient for efficiently computing the optimal dual variables is therefore to reduce the work required for this part. We will now show how this can be done by exploiting the structure inherent in the problem, hence simplifying equations (30) and (31).

Our key result is summarised in the following theorem.

Theorem 4.1. *Let $\lambda = \{\lambda_t\}_{t \in \mathcal{T}}$ and $\gamma = \{\gamma_t\}_{t=0}^{T-1}$ be given families of vectors in \mathbb{R}^{n_S} and $\mathbb{R}^{n_S n_X}$, respectively. Let $u_t = \exp(\lambda_t/\epsilon)$ for $t \in \mathcal{T}$ and let $K_t = \exp(-C_t/\epsilon)$ and $G_t = \exp(\text{diag}(\gamma_t)\Delta/\epsilon)$ for $t \in [T] \setminus 0$, where the matrices C_t , $t \in [T] \setminus 0$, are as in problem (22) and the matrix Δ is as given in equation (19). Define two families of vectors, $\hat{\psi} = \{\hat{\psi}_t\}_{t=0}^T$ and $\psi = \{\psi_t\}_{t=0}^T$, via the recursions*

$$\hat{\psi}_t = \begin{cases} \mathbf{1}_{n_S n_X}, & t = 0 \\ (K_t \odot G_t)^\top (\hat{\psi}_{t-1} \odot (\mathbf{1}_{n_X} \otimes u_{t-1})), & (t-1) \in \mathcal{T}, \quad t \in [T] \setminus 0 \\ (K_t \odot G_t)^\top \hat{\psi}_{t-1}, & (t-1) \notin \mathcal{T}, \quad t \in [T] \setminus 0 \end{cases} \quad (32)$$

and

$$\psi_t = \begin{cases} \mathbf{1}_{n_S n_X}, & t = T \\ (K_{t+1} \odot G_{t+1}) (\psi_{t+1} \odot (\mathbf{1}_{n_X} \otimes u_{t+1})), & (t+1) \in \mathcal{T}, \quad t \in [T-1] \\ (K_{t+1} \odot G_{t+1}) \psi_{t+1}, & (t+1) \notin \mathcal{T}, \quad t \in [T-1]. \end{cases} \quad (33)$$

Then λ and γ are optimal variables for the dual problem (28) if and only if the following equations hold

$$u_t = \mu_t \oslash P^S(\hat{\psi}_t \odot \psi_t), \quad t \in \mathcal{T},$$

and

$$\begin{aligned} \hat{\psi}_t \odot (\mathbf{1}_{n_X} \otimes u_t) \odot (K_{t+1} \odot G_{t+1} \odot \Delta) (\psi_{t+1} \odot (\mathbf{1}_{n_X} \otimes u_{t+1})) &= \mathbf{0}_{n_S n_X}, & t \in \mathcal{T}, t+1 \in \mathcal{T} \\ \hat{\psi}_t \odot (K_{t+1} \odot G_{t+1} \odot \Delta) (\psi_{t+1} \odot (\mathbf{1}_{n_X} \otimes u_{t+1})) &= \mathbf{0}_{n_S n_X}, & t \notin \mathcal{T}, t+1 \in \mathcal{T} \\ \hat{\psi}_t \odot (\mathbf{1}_{n_X} \otimes u_t) \odot (K_{t+1} \odot G_{t+1} \odot \Delta) \psi_{t+1} &= \mathbf{0}_{n_S n_X}, & t \in \mathcal{T}, t+1 \notin \mathcal{T} \\ \hat{\psi}_t \odot (K_{t+1} \odot G_{t+1} \odot \Delta) \psi_{t+1} &= \mathbf{0}_{n_S n_X}, & t \notin \mathcal{T}, t+1 \notin \mathcal{T}. \end{aligned}$$

Remark 4.2. According to the above theorem, we can compute dual variables satisfying the optimality conditions from equations (29a) and (29b), by simply performing a number of matrix-vector products. This greatly reduces the computational work required, especially since the help vectors $\hat{\psi}$ (resp. ψ) can be computed inductively (resp. recursively); therefore, we do not have to recalculate the full chains for each individual variable update. We also emphasise that according to the above theorem, there is no need to explicitly form the tensors K , U_λ and G_γ .

In order to prove Theorem 4.1, we will need the following lemma.

Lemma 4.3. *Let $\lambda = \{\lambda_t\}_{t \in \mathcal{T}}$ and $\gamma = \{\gamma_t\}_{t=0}^{T-1}$ be given families of vectors in \mathbb{R}^{n_S} and $\mathbb{R}^{n_S n_X}$, respectively. Define u_t for $t \in \mathcal{T}$ and K_t and G_t for $t \in [T] \setminus 0$ as in Theorem 4.1 and let $U_\lambda(\omega) = \sum_{t \in \mathcal{T}} u_t(\omega_t^S)$ and $G_\gamma(\omega) = \sum_{t \in [T] \setminus 0} G_t(\omega_{t-1}, \omega_t)$ for $\omega \in \Omega$. Then*

$$P_t(K \odot U_\lambda \odot G_\gamma) = \hat{\psi}_t \odot \bar{u}_t \odot \psi_t, \quad t \in [T],$$

where $\hat{\psi}$ and ψ are functions of λ and γ as given in equations (32) and (33), respectively, and $\bar{u}_t \in \mathbb{R}^{n_S n_X}$ is given by

$$\bar{u}_t = \begin{cases} \mathbf{1}_{n_X} \otimes u_t, & t \in \mathcal{T} \\ \mathbf{1}_{n_S n_X}, & t \in [T] \setminus \mathcal{T}. \end{cases} \quad (34)$$

Moreover, for $t_1, t_2 \in [T]$ such that $t_1 < t_2$,

$$\begin{aligned} & P_{t_1, t_2}(K \odot U_\lambda \odot G_\gamma) \\ &= \text{diag}(\hat{\psi}_{t_1} \odot \bar{u}_{t_1}) \left((K_{t_1+1} \odot G_{t_1+1}) \text{diag}(\bar{u}_{t_1+1}) \dots (K_{t_2} \odot G_{t_2}) \text{diag}(\bar{u}_{t_2}) \right) \text{diag}(\psi_{t_2}). \end{aligned}$$

Proof. Start by noting that U_λ is of the form

$$U_\lambda(\omega_0, \dots, \omega_T) = \prod_{t \in [T]} \bar{u}_t(\omega_t), \quad \omega_0, \dots, \omega_T \in \{1, \dots, n_S\} \times \{1, \dots, n_X\}$$

since, when $\omega_t \in \{1, \dots, n_S\} \times \{1, \dots, n_X\}$ is ordered according to (18), $\bar{u}_t(\omega_t) = u_t(\omega_t^S)$ for $t \in \mathcal{T}$. Also note that $K \odot G_\gamma$ inherits the structure of K and G_γ ; indeed,

$$(K \odot G_\gamma)(\omega_0, \dots, \omega_T) = \prod_{t \in [T] \setminus 0} (K_t \odot G_t)(\omega_{t-1}, \omega_t), \quad \omega_0, \dots, \omega_T \in \{1, \dots, n_S\} \times \{1, \dots, n_X\}.$$

The requirements of Proposition 2.8 are thus satisfied and it yields expressions for the projections. Specifically, taking

$$\hat{\psi}_t := \begin{cases} \mathbf{1}_{n_S n_X}, & t = 0 \\ (K_t \odot G_t)^\top \text{diag}(\bar{u}_{t-1}) \dots (K_1 \odot G_1)^\top \bar{u}_0, & t \in [T] \setminus 0 \end{cases}$$

and

$$\psi_t := \begin{cases} \mathbf{1}_{n_S n_X}, & t = T \\ (K_{t+1} \odot G_{t+1}) \text{diag}(\bar{u}_{t+1}) \dots (K_T \odot G_T) \bar{u}_T, & t \in [T-1] \end{cases}$$

we get

$$P_t(K \odot U_\lambda \odot G_\gamma) = \hat{\psi}_t \odot \bar{u}_t \odot \psi_t, \quad t \in [T].$$

It is easily verified that this choice of $\hat{\psi}$ and ψ are given in equations (32) and (33), respectively, given the definition from equation (34).

Let $t_1, t_2 \in [T]$ such that $t_1 < t_2$. Applying the results from [23, Lemma 2 and proof of Proposition 2] and making the same arguments as for the one-marginal projection gives that

$$P_{t_1, t_2}(K \odot U_\lambda \odot G_\gamma) = \text{diag}(\hat{\psi}_{t_1} \odot \bar{u}_{t_1}) M_{t_1, t_2} \text{diag}(\bar{u}_{t_2} \odot \psi_{t_2})$$

where

$$M_{t_1, t_2} = (K_{t_1+1} \odot G_{t_1+1}) \text{diag}(\bar{u}_{t_1+1}) \dots (K_{t_2-1} \odot G_{t_2-1}) \text{diag}(\bar{u}_{t_2-1}) (K_{t_2} \odot G_{t_2}),$$

from where the result follows. \square

We now prove Theorem 4.1.

Proof of Theorem 4.1. It was shown in Section 4.1 that the (representation of) the optimal dual variables u and γ should satisfy the optimality conditions in equations (30) and (31). Start by considering the former equation, whose right-hand side is equal to

$$(\mu_t \odot u_t) \odot P_t^S(K \odot U_\lambda \odot G_\gamma), \quad t \in \mathcal{T}. \quad (35)$$

We must find an expression for the projection of the tensor $K \odot U_\lambda \odot G_\gamma$. Fix some $t \in \mathcal{T}$ and start by recalling that $P_t^S(K \odot U_\lambda \odot G_\gamma) = (P^S \circ P_t)(K \odot U_\lambda \odot G_\gamma)$, where Lemma 4.3 then yields an expression for $P_t(K \odot U_\lambda \odot G_\gamma)$. Therefore

$$P_t^S(K \odot U_\lambda \odot G_\gamma) = P^S(\hat{\psi}_t \odot (\mathbf{1}_{n_X} \otimes u_t) \odot \psi_t) = P^S(\hat{\psi}_t \odot \psi_t) \odot u_t \quad (36)$$

where the second equality follows from the fact that $(\mathbf{1}_{n_X} \otimes u_t)(\omega_t) = u_t(\omega_t^S)$ for every $\omega_t \in \{1, \dots, n_S\} \times \{1, \dots, n_X\}$ ordered by (18). Inserting equation (36) into equation (35) yields the expression stated in the theorem.

Moving on to equation (31), we apply Lemma 4.3 to obtain an expression for the bi-marginal projections,

$$P_{t,t+1}(K \odot U_\lambda \odot G_\gamma) = \text{diag}(\hat{\psi}_t \odot \bar{u}_t)(K_{t+1} \odot G_{t+1}) \text{diag}(\psi_{t+1} \odot \bar{u}_{t+1}), \quad t \in [T-1].$$

The optimality condition from equation (31) thus becomes

$$\hat{\psi}_t \odot \bar{u}_t \odot (K_{t+1} \odot G_{t+1} \odot \Delta)(\psi_{t+1} \odot \bar{u}_{t+1}) = \mathbf{0}_{n_S n_X}, \quad t \in [T-1].$$

Inserting the definition of \bar{u}_t and \bar{u}_{t+1} from Lemma 4.3 into the above expression proves the second part of the assertion and thus completes the proof. \square

We end this section by noting that Lemma 4.3 provides an expression that allows for fast computation of the sub-transport between any two marginals, without explicitly having to form the full transport Q . One type of sub-transport that is of particular interest is transportation between adjacent marginals; later we will see that such sub-transports allow for recovering the price without explicitly forming the full transport Q .

Corollary 4.4. *Let $\lambda = \{\lambda_t\}_{t \in \mathcal{T}}$ and $\gamma = \{\gamma_t\}_{t=0}^{T-1}$ be given families of vectors in \mathbb{R}^{n_S} and $\mathbb{R}^{n_S n_X}$, respectively. Define u_t for $t \in \mathcal{T}$ and K_t and G_t for $t \in [T] \setminus 0$ as in Theorem 4.1 and let the corresponding transport plan be denoted $Q_{\lambda, \gamma}$. Then the bi-marginal sub-transport $P_{t-1,t}(Q_{\lambda, \gamma})$ for $t \in [T] \setminus 0$ is given by*

$$\begin{cases} \text{diag}(\hat{\psi}_{t-1} \odot (\mathbf{1}_{n_X} \otimes u_{t-1}))(K_t \odot G_t) \text{diag}(\psi_t \odot (\mathbf{1}_{n_X} \otimes u_t)), & t-1 \in \mathcal{T}, \quad t \in \mathcal{T} \\ \text{diag}(\hat{\psi}_{t-1} \odot (\mathbf{1}_{n_X} \otimes u_{t-1}))(K_t \odot G_t) \text{diag}(\psi_t), & t-1 \in \mathcal{T}, \quad t \notin \mathcal{T} \\ \text{diag}(\hat{\psi}_{t-1})(K_t \odot G_t) \text{diag}(\psi_t \odot (\mathbf{1}_{n_X} \otimes u_t)), & t-1 \notin \mathcal{T}, \quad t \in \mathcal{T} \\ \text{diag}(\hat{\psi}_{t-1})(K_t \odot G_t) \text{diag}(\psi_t), & t-1 \notin \mathcal{T}, \quad t \notin \mathcal{T}, \end{cases}$$

where $\hat{\psi}$ and ψ are given as functions of λ and γ by equations (32) and (33).

Proof. Fix some $t \in [T] \setminus 0$ and note that by equation (27), $P_{t-1,t}(Q_{\lambda, \gamma}) = P_{t-1,t}(K \odot U_\lambda \odot G_\gamma)$, where

$$P_{t-1,t}(K \odot U_\lambda \odot G_\gamma) = \text{diag}(\hat{\psi}_{t-1} \odot \bar{u}_{t-1})(K_t \odot G_t) \text{diag}(\psi_t \odot \bar{u}_t)$$

follows from application of Lemma 4.3 with $t_1 = t-1$ and $t_2 = t$. Here \bar{u}_{t-1} and \bar{u}_t are defined as in equation (34). This proves the assertion. \square

4.3 Summary of the full algorithm

We have now arrived at a form of the optimality conditions that allows for efficient computation of the optimal dual variables. We can therefore construct a coordinate dual ascent method by cyclically fixing all but one variable and optimise over the remaining variable by selecting it so that it satisfies the corresponding optimality condition. We then move on to the next variable until all variables have been optimised over; one such cycle defines one iteration in the method.

Each iteration starts with the update of the variables u corresponding to the marginal constraints (22c). The closed-form formula for doing so is given by Theorem 4.1 as

$$u_t \leftarrow \mu_t \odot P^S(\hat{\psi}_t \odot \psi_t), \quad t \in \mathcal{T}.$$

From the above we note that it suffices to manipulate vectors in order to update u , something that enables fast computation.

We then proceed to updating the dual variables γ representing the martingale constraints (22b). This time, no closed-form formula exists for γ — the optimality condition thus has to be solved numerically. Theorem 4.1 provides a form of the equations that allows for efficiently doing so by

application of Newton's method. In order to do so, define a family of functions $\alpha_t : \mathbb{R}^{n_S n_X} \rightarrow \mathbb{R}^{n_S n_X}$ for $t \in [T] \setminus 0$ via

$$\alpha_t(z) := \begin{cases} (K_t \odot \exp(\text{diag}(z)\Delta/\varepsilon) \odot \Delta)(\psi_t \odot (\mathbf{1}_{n_X} \otimes u_t)), & t \in \mathcal{T} \\ (K_t \odot \exp(\text{diag}(z)\Delta/\varepsilon) \odot \Delta)\psi_t, & t \notin \mathcal{T}. \end{cases}$$

Then equation (31) can be written

$$\begin{cases} \hat{\psi}_t \odot (\mathbf{1}_{n_X} \otimes u_t) \odot \alpha_{t+1}(\gamma_t) = \mathbf{0}_{n_S n_X}, & t \in [T-1] \cap \mathcal{T} \\ \hat{\psi}_t \odot \alpha_{t+1}(\gamma_t) = \mathbf{0}_{n_S n_X}, & t \in [T-1] \setminus \mathcal{T}. \end{cases}$$

Application of Newton's method to the above equation requires iterating according to

$$\gamma_t^{(k+1)} \leftarrow \gamma_t^{(k)} - \theta^{(k)} \odot J_{\alpha_{t+1}}(\gamma_t^{(k)})^{-1} \alpha_{t+1}(\gamma_t^{(k)}), \quad k = 1, 2, \dots, \quad (37)$$

where $\theta^{(k)} \in \mathbb{R}^{n_S n_X}$ is a vector containing the respective step lengths to be used for each variable, such that $\theta^{(k)}(i) \in [0, 1]$ for $i \in \{1, \dots, n_S n_X\}$. It is determined by vectorised line search. In the above, $J_{\alpha_{t+1}}$ denotes the Jacobian matrix associated with α_{t+1} ; note that α_{t+1} is a vector-valued function whose i^{th} component is independent of any other components of γ_t than $\gamma_t(i)$, $i \in \{1, \dots, n_S\} \times \{1, \dots, n_X\}$. There are no cross dependencies. Therefore, the Jacobian is a diagonal matrix and given by $J_{\alpha_t}(\gamma_{t-1}) = \text{diag}(\beta_t(\gamma_{t-1}))$, where $\beta_t : \mathbb{R}^{n_S n_X} \rightarrow \mathbb{R}^{n_S n_X}$ for $t \in [T] \setminus 0$ is defined as

$$\beta_t(z) := \begin{cases} \frac{1}{\varepsilon}(K_t \odot \exp(\text{diag}(z) \odot \Delta/\varepsilon) \odot \Delta \odot \Delta)(\psi_t \odot (\mathbf{1}_{n_X} \otimes u_t)), & t \in \mathcal{T} \\ \frac{1}{\varepsilon}(K_t \odot \exp(\text{diag}(z) \odot \Delta/\varepsilon) \odot \Delta \odot \Delta)\psi_t, & t \notin \mathcal{T}. \end{cases}$$

The iterative scheme (37) can therefore be simplified to

$$\gamma_t^{(k+1)} \leftarrow \gamma_t^{(k)} - \theta^{(k)} \odot \alpha_{t+1}(\gamma_t^{(k)}) \oslash \beta_{t+1}(\gamma_t^{(k)}), \quad k = 1, 2, \dots$$

The inversion of the Jacobian and the matrix-vector multiplication used in (37) are thus avoided and replaced by the elementwise division of one vector with another.

Putting things together, we have completed the derivation of our algorithm — see Algorithm 2 for the full method. Convergence of the algorithm can be shown given that there exists a feasible solution that is positive for all variables that are not directly constrained to be zero by the constraints (cf. [32, Theorem 4.1 and Proposition 1]). In this case, the variables constrained to be zeros can be removed (following [32]); convergence of the coordinate dual ascent then follows from [45, Theorem 2 and Section 5].

Remark 4.5. Storing the tensor Q quickly becomes impossible as the number of marginals grows and consequently we cannot access the full transport Q in practice. Fortunately, it is often enough to obtain a set of sub-transport of the form $P_{t_1, t_2}(Q)$, where $t_1, t_2 \in [T]$ are such that $t_1 < t_2$. For example, the corresponding problem value — the lower bound on the price — can be obtained without explicitly forming and storing the full tensor. See this by recalling that

$$\langle \Phi, Q \rangle = \sum_{t \in [T] \setminus 0} \langle \Phi_t, P_{t-1, t}(Q) \rangle,$$

where the sub-transport $P_{t-1, t}(Q)$ for $t \in [T] \setminus 0$ are given by Corollary 4.4.

5 Computational results

In this section, we apply our computational framework to solve a number of MOT problems: First, we consider problems for which the optimal solution is known analytically; comparison of the computed solution with the known solutions allows for verifying that the method works as intended. We then conclude by solving a MOT problem where the optimal solution is not known.

Algorithm 2 Method for solving problem (22).

Given: matrices $\{K_t\}_{t=1}^T$ and Δ , index set \mathcal{T} , vectors $\{\mu_t\}_{t \in \mathcal{T}}$, scalar $\varepsilon > 0$
Initialise: $u_t \leftarrow \mathbf{1}_{n_S}$ for $t \in [T]$
 $\gamma_t \leftarrow \mathbf{1}_{n_S n_X}$ for $t \in [T-1]$
 $G_t \leftarrow \exp(\text{diag}(\gamma_{t-1})\Delta/\varepsilon)$ for $t \in [T] \setminus 0$
 $\hat{\psi}_1, \psi_T \leftarrow \mathbf{1}_{n_S n_X}$
 $\psi_t \leftarrow (K_{t+1} \odot G_{t+1})(\psi_{t+1} \odot (\mathbf{1}_{n_X} \otimes u_{t+1}))$ for $t \in [T-1]$
while Sinkhorn not converged **do**
 #Update marginal constraints:
 if $0 \in \mathcal{T}$ **then**
 $u_0 \leftarrow \mu_0 \oslash P^S(\psi_0)$
 end if
 for $t = 1, \dots, T-1$ **do**
 $\hat{\psi}_t \leftarrow (K_t \odot G_t)^\top (\hat{\psi}_{t-1} \odot (\mathbf{1}_{n_X} \otimes u_{t-1}))$
 if $t \in \mathcal{T}$ **then**
 $u_t \leftarrow \mu_t \oslash P^S(\hat{\psi}_t \odot \psi_t)$
 end if
 end for
 $\hat{\psi}_T \leftarrow (K_T \odot G_T)^\top (\hat{\psi}_{T-1} \odot (\mathbf{1}_{n_X} \otimes u_{T-1}))$
 if $T \in \mathcal{T}$ **then**
 $u_T \leftarrow \mu_T \oslash P^S(\hat{\psi}_T)$
 end if
 #Update martingale constraints:
 for $t = T, \dots, 1$ **do**
 if $t \neq T$ **then**
 $\psi_t \leftarrow (K_{t+1} \odot G_{t+1})(\psi_{t+1} \odot (\mathbf{1}_{n_X} \otimes u_{t+1}))$
 end if
 $k \leftarrow 0$
 while Newton not converged **do**
 $k \leftarrow k + 1$
 $\theta^{(k)} \leftarrow 1$
 $\gamma_{t-1}^{(k+1)} \leftarrow \gamma_{t-1}^{(k)} - \theta^{(k)} \odot \alpha_t(\gamma_{t-1}^{(k)}) \oslash \beta_t(\gamma_{t-1}^{(k)})$
 while $\gamma_{t-1}^{(k+1)} > \gamma_{t-1}^{(k)}$ **do**
 $\theta^{(k)} \leftarrow \theta^{(k)}/2$
 $\gamma_{t-1}^{(k+1)} \leftarrow \gamma_{t-1}^{(k)} - \theta^{(k)} \odot \alpha_t(\gamma_{t-1}^{(k)}) \oslash \beta_t(\gamma_{t-1}^{(k)})$
 end while
 $G_t \leftarrow \exp(\text{diag}(\gamma_{t-1}^{(k+1)})\Delta/\varepsilon)$
 end while
 end for
 $\psi_0 \leftarrow (K_1 \odot G_1)(\psi_1 \odot (\mathbf{1}_{n_X} \otimes u_1))$
end while
return $u, \hat{\psi}, \psi, \{G_t\}_{t=1}^T$




Figure 1: Optimal solutions to the upper bound bi-marginal MOT and OT problems corresponding to the payoff of a variance swap when the given marginals are uniform, as in Example 5.1. The computed solution of the MOT problem is displayed in Figure 1a; it exhibits the shape of the left-monotone transport. For comparison, the computed solution of the corresponding OT problem is displayed in Figure 1b; it displays the shape of the anti-monotone coupling.

5.1 Monotone transport

A well-known example from classical OT theory is the monotone transport plan, which is concentrated on the graph of an increasing function (see, for example, [60, p. 75]). It is also known as the Hoeffding-Fréchet coupling and solves the upper bound bi-marginal OT problem when the cost function $\phi : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ satisfies certain conditions² — in particular, the cross derivative $\partial_{s_0 s_1} \phi(s_0, s_1)$ should exist and be strictly positive [36, Theorem 2.2]. Similarly, the anti-monotone transport plan is concentrated on the graph of a decreasing function; it solves the corresponding lower bound OT problem. There exists a martingale analogue to the monotone coupling; the *left-monotone transport plan* was introduced in [7, Definition 1.4] and shown to solve the bi-marginal MOT problem for two types of cost functions [7, Theorems 6.1 and 6.3]. It is concentrated on the graphs of two measurable functions and exhibits a V-shape when visualised in the plane. It was shown in [36, Theorem 5.1] that the left-monotone coupling solves the upper bound bi-marginal MOT problem, given that the derivative $\partial_{s_0 s_1} \phi(s_0, s_1)$ exists and is strictly positive. Since the monotone couplings are visually easily recognised, they serve as a clear first demonstrating example.

Example 5.1 (Robust pricing of a variance swap). Let $\phi(s_0, s_1) = (\log(s_1/s_0))^2$ and let the initial marginal μ_0 be uniform with support on $n_{S_0} = 600$ atoms, evenly distributed over the interval $[1.25, 1.75]$. Similarly, let the terminal marginal μ_1 be uniform with support on $n_{S_1} = 1200$ atoms, evenly distributed over the interval $[1, 2]$. It is easily verified that $\partial_{s_0 s_1} \phi > 0$ and $\partial_{s_0 s_1} \phi < 0$ on the supports of μ_0 and μ_1 and that the marginals are in convex order; the optimal solutions for the corresponding upper-bound MOT and OT problems are thus the left-monotone and the anti-monotone couplings, respectively. Figure 1 shows the computed optimal solutions obtained for $\varepsilon = 4.5 \cdot 10^{-4}$. A dark blue colour indicates a close-to-zero probability for that specific transport to occur, whilst a yellow or green colour indicates a non-zero probability that this transport occurs. Note the V-shape of the optimal martingale coupling.

²The cost function ϕ should be upper semi-continuous and of linear growth.

5.2 Late and early transports

Next we consider a multi-marginal ($T > 1$) example where the payoff function is of the form

$$\phi(s_0, \dots, s_T) = \frac{1}{T+1} \sum_{t \in [T]} f(s_t), \quad s_0, \dots, s_T \in \mathcal{S}, \quad (38)$$

for some convex function $f : \mathbb{R} \rightarrow \mathbb{R}$, and where the initial and terminal marginals μ_0 and μ_T are the only given marginals. As always, it is assumed that μ_0 and μ_T are in convex order. Two particular martingale couplings are of interest here, the coupling that corresponds to not moving at all before $t = T - 1$, then performing the full transport between $t = T - 1$ and $t = T$, and the coupling that corresponds to performing the full transport between $t = 0$ and $t = 1$, then keeping the process constant until $t = T$. We refer to these couplings as late and early transport, respectively. The below result follows immediately from linearity of the expected value, since for any model $(\Omega, \mathcal{F}, \mathbb{Q}, S)$,

$$\mathbb{E}_{\mathbb{Q}} [f(S_T)] \geq \mathbb{E}_{\mathbb{Q}} [f(S_{T-1})] \geq \dots \geq \mathbb{E}_{\mathbb{Q}} [f(S_1)] \geq \mathbb{E}_{\mathbb{Q}} [f(S_0)],$$

by convexity of f . See [57] for a similar result.

Proposition 5.2. *Let $T > 1$ and consider a payoff function $\phi : \mathbb{R}^{T+1} \rightarrow \mathbb{R}$ of the form (38). Let μ_0 and μ_T be two given probability measures on $(\mathbb{R}, \mathcal{B}(\mathbb{R}))$ and assume that they are in convex order. Then the optimal solution of the associated lower bound MOT problem corresponds to the late transport defined above, whilst the early transport solves the upper bound MOT problem.*

Example 5.3. Let $f(x) = x^2$, $T = 50$ and let the two given marginal distributions μ_0 and μ_{50} be as in Figure 2; they are in convex order. Since the payoff function is of the form $\phi(s_0, \dots, s_T) = \phi_1(s_0, s_1) + \sum_{t \in [T] \setminus \{0,1\}} \phi_t(s_t)$ for

$$\begin{aligned} \phi_1(s_0, s_1) &= \frac{1}{T+1} (f(s_0) + f(s_1)), \quad s_0, s_1 \in \mathcal{S}, \\ \phi_t(s_t) &= \frac{1}{T+1} f(s_t), \quad s_t \in \mathcal{S}, \quad t = 2, \dots, T, \end{aligned}$$

it is by Remark 3.6 possible to omit the memory process X from the representation of the problem.

A subset of the marginal distributions of the computed optimal martingale transport plans are displayed in Figure 2, along with the given marginals μ_0 and μ_{50} . The lower bound solution (computed using $\varepsilon = 0.0065$) is shown in Figure 2a; since the computed intermediate marginals are close to the initial marginal this solution corresponds to a late transport, as predicted by Proposition 5.2. The upper bound solution (computed using $\varepsilon = 0.02$) is shown in Figure 2b; for this solution, we note that the intermediate marginals are close to the terminal marginal. It is therefore an example of early transport, as given by Proposition 5.2. For both solutions, the small deviations of the intermediate marginals from the initial or terminal marginal are due to the regularisation.

5.3 The maximum of the maximum

Suppose that μ_T is a given marginal distribution, centered in $s_0 \in \mathbb{R}$, and assume that $\mu_0 = \delta_{s_0}$. Let the process X be the rolling maximum of the price process S — that is, let X be as in Example 2.1. We here consider the problem of finding the martingale model that maximises the maximum while respecting the terminal marginal μ_T , that is, that solves the upper bound MOT problem with $\phi(S_0, \dots, S_T) = X_T$ and $\mathcal{T} = \{0, T\}$. Note that this payoff is of the form (5); the problem can thus be addressed within our framework.

Before we compute its optimal solution for a specific choice of μ_T , we will have a look at the corresponding continuous-time solution. Fix a probability space $(\Omega, \mathcal{F}, \mathbb{Q})$ and let S^{AY} denote the martingale on $[0, T]$ whose maximum has a first-order stochastic dominance over the maximum of any other continuous martingale with terminal marginal μ_T . It exists, and is related to the martingale constructed by Azéma and Yor [3] as a solution of the Skorokhod embedding problem with terminal




Figure 2: Marginal distributions corresponding to the optimal solutions of the upper- and lower bound MOTs from Example 5.3. The given initial marginal μ_0 is displayed in light blue bars, while the given terminal marginal μ_T is shown in light red bars. The marginals $P_t(Q)$ corresponding to the computed transport plan Q are shown for $t \in \{0, 10, 20, 30, 40, 50\}$ with solid lines. Figure 2a corresponds to the lower bound martingale transport plan while Figure 2b corresponds to the upper bound martingale transport plan.

marginal μ_T [10, Lemma 2.2]; we will refer to this martingale as the Azéma–Yor martingale. It follows that for the continuous-time case, an optimal solution of the above MOT problem is given by the Azéma–Yor martingale S^{AY} [38]. A solution of the problem in continuous-time is thus known. It follows from [11, Lemma 2.4] that said distribution can be obtained as

$$\mathbb{Q}(X_T^{\text{AY}} \geq B) = \min_{0 \leq y \leq B} f_B(y), \quad B > s_0, \quad (39)$$

where $f_B : [0, B] \rightarrow \mathbb{R}$ is given by $f_B(y) := (B - y)^{-1} \int (s - y)^+ d\mu_T(s)$ for $B > s_0$. Equation (39) thus directly provides the law of the maximum X_T that corresponds to an optimal solution of the problem, when an ‘infinite number of time steps’ is used. We will now compute this law for a specific choice of terminal marginal μ_T . We then solve the problem computationally for T increasing and compute the distribution of X_T corresponding to an optimal solution. We will see that the resulting distribution approaches the distribution obtained from equation (39) as T increases.

Example 5.4 (Comparison with the Azéma–Yor martingale). Let the terminal marginal μ_T be as in Figure 3a with $s_0 = 0.5$, see Figure 3b for examples of the function f_B corresponding to μ_T for a few different values of B — note that its minimum exists for each B . The probability $\mathbb{Q}(X_T^{\text{AY}} \geq B)$ is obtained by minimising f_B over $[0, B]$. By doing this for all $B \in (0.5, 1]$ we are able to recover the distribution of X_T^{AY} . It is displayed using a bold red line in Figure 4a. The corresponding problem value, the robust price, is $\mathbb{E}_{\mathbb{Q}}[X_T^{\text{AY}}] = 0.6133$.

We now solve the corresponding upper bound MOT problem for different values of T , ranging from 2 to 29, by application of our framework with $\varepsilon = 0.01$. The corresponding law of the maximum of the price process is displayed in Figure 4a. The corresponding robust prices, $\mathbb{E}_{\mathbb{Q}}[X_T]$, are shown in Figure 4b as a function of the number of marginals used. We note that the computed distribution approaches the known continuous-time solution and that the cumulative distribution function decreases as the number of time steps used increases. Similarly the computed problem values, the robust prices, approaches the problem value of the corresponding continuous-time problem.




Figure 3: The given terminal marginal used in Example 5.4 is displayed in Figure 3a. It is centered in 0.5 and its support is contained within $[0, 1]$. The function f_B is shown in Figure 3b for a few different values of B .




Figure 4: The distribution of the maximum of the maximum from Example 5.4, solved approximately for an increasing number of marginals $T + 1$, is shown in Figure 4a. The corresponding distribution for the continuous-time case is included for reference (bold red). We note that the discrete-time computed solutions approach the continuous-time solution as the number of time steps used increases, and that the cumulative distribution function decreases. The corresponding problem values are shown as a function of the number of marginals $T + 1$ in Figure 4b; the problem value for the continuous-time case is included for comparison (bold red).

5.4 The robust price of a digital option

An example related to the maximum of the maximum is the robust upper bound on the fair price of a *digital option* — a claim that pays one unit of money if the underlying price process S has exceeded the barrier B before maturity, else it expires worthless. Yet again, let μ_T be a given terminal distribution, centered in $s_0 \in \mathbb{R}$, and assume that the initial distribution is given by $\mu_0 = \delta_{s_0}$. Fix some $B > s_0$, let $T > 1$ and let X be the indicator process keeping track of whether the price process S has exceeded the barrier B so far, that is, let $X_t := \chi_{[B, \infty)}(\max_{r \in [t, T]} S_r)$ for $t \in [T]$. Then the payoff of a digital option with barrier B is given by $\phi(S_0, \dots, S_T) = X_T$; we are interested in solving the corresponding upper bound MOT problem.

We have seen that in continuous-time there exists a martingale S^{AY} that optimises all digital options simultaneously, that is, the Azéma–Yor martingale S^{AY} solves the problem for any barrier $B > s_0$. Indeed, since $\chi_{[B, \infty)}(\cdot)$ is an increasing function and the maximum of the martingale S^{AY} has a first-order stochastic dominance over the maximum of any other continuous martingale S with marginals μ_0 and μ_T ,

$$\mathbb{E}_{\mathbb{Q}} \left[\chi_{[B, \infty)} \left(\max_{0 \leq t \leq T} S_t^{\text{AY}} \right) \right] \geq \mathbb{E}_{\mathbb{Q}} \left[\chi_{[B, \infty)} \left(\max_{0 \leq t \leq T} S_t \right) \right], \quad B > s_0.$$

For a specific digital option though — that is, for B fixed — one can find an (optimal) coupling with the same objective value by using just one intermediate time step [24, pp.416–419]. In the below examples, we solve problems with $T = 2$ computationally; we will recover the coupling constructed in [24] as well as the distribution of the maximum of the Azéma–Yor martingale.

Example 5.5 (Recovering the optimal coupling from [24] when $T = 2$). Let $s_0 = 0.5$ and $\mu_T = 0.5(\delta_0 + \delta_1)$, then it follows from equation (39) that the the robust price of a digital option with barrier $B \in (0.5, 1]$ for the continuous-time problem is $(2B)^{-1}$. On the other hand, [24, Theorem 7.27] yields the same value when taking only discrete time models with $T \geq 2$ into account. Repeating the construction from the proof of the theorem allows for constructing a martingale coupling that attains this upper bound. For this example with $T = 2$, it corresponds to take at $t = 1$ any of the two values 0 or B with probability $1 - (2B)^{-1}$ and $(2B)^{-1}$, respectively, before moving to the terminal marginal. We note that the three marginals fully specifies a martingale coupling.

We approximately solve the corresponding tri-marginal MOT problem by application of our framework for $B = 0.75$, $\varepsilon = 0.02$ and $n_S = 100$. The corresponding intermediate marginal distribution ($t = 1$) of the price process is $0.27\delta_0 + 0.05\delta_{0.01} + 0.01\delta_{0.02} + 0.67\delta_{0.75}$; this can be compared with $0.33\delta_0 + 0.67\delta_{0.75}$, which is given by the theoretical construction. The computed distribution is subject to some smoothing around zero, which is due to regularisation.

We then repeat the computations for 2, 3, 5, 10 and 15 marginals and investigate what happens with the problem value, the robust price, as the number of marginals increases. For $T = 1$, the computed price is 0.5, while for larger T it is 0.66 — this confirms that it is not possible to increase the problem value by adding further time steps when $T \geq 2$.

Example 5.6 (The law of the maximum of the Azéma–Yor martingale via digital options). Let $T = 2$ and let the initial and terminal marginals be as in Example 5.4. We solve the corresponding MOT problem computationally for a set of barriers ranging from 0.5 to 0.92; a regularisation parameter of $\varepsilon = 0.01$ was used. By doing so we obtain the maximum of the value $\mathbb{E}_{\mathbb{Q}}[\chi_{[B, \infty)}(\max_{t \in [T]} S_t)] = \mathbb{Q}(\max_{t \in [T]} S_t \geq B)$ for each such barrier B ; the result is shown as a function of the barrier in Figure 5, along with the cumulative distribution function of the maximum of S^{AY} (cf. Example 5.4). We see that by successively solving for the upper bounds on the probability that digital options with barriers $B \in (0.5, 1]$ are not activated, we can approximately reproduce the cumulative distribution function of the maximum of the Azéma–Yor martingale.

5.5 Robust pricing of an Asian option

We conclude by considering an example where the optimal solution is not known, namely, the robust pricing of an Asian option subject to several marginal constraints. Asian options are a class of financial




Figure 5: The computed upper bound on the probability that the digital option with barrier B is not activated as a function of the barrier level B , as given in Example 5.6. The cumulative distribution function of the maximum of the corresponding Azéma–Yor martingale is displayed in bold red for reference. The grey vertical lines illustrate the support of the price process contained within $[0.5, 0.92]$. Note that the resolution of the computed approximations follow this grid.


derivatives that are of great practical financial interest; they are characterised by their payoff being of the form

$$\phi(S_0, \dots, S_T) = f\left(\frac{1}{T+1} \sum_{t \in [T]} S_t\right)$$


for some function $f : \mathbb{R} \rightarrow \mathbb{R}$. The dependence on the arithmetic mean makes the corresponding MOT problem a difficult one to solve analytically — this was discussed in [57], who provided the optimal solutions for a problem with two or three given marginals and f convex. As far as we know, it is still unclear what the optimal solution would look like for a MOT problem subject to more than four marginal constraints. By letting the stochastic process X be as given in Example 2.2, we can however solve these problems computationally. We will now do so for an example where we successively increase the number of marginal constraints. In order to provide visually clear results, all given marginals will be rather simple.

Example 5.7 (Pricing a straddle). Let the payoff function ϕ be as given above with $f(x) = |x - 30|$. Then the payoff corresponds to the payoff of an Asian straddle with strike 30. Let $T = 11$. Consider a trivial initial marginal $\mu_0 = \delta_{30}$ and a uniformly distributed terminal marginal μ_{11} whose support is contained within $n_S = 41$ atoms, evenly distributed over the interval $[25, 35]$. In order not to run into problems with rounding errors, we will here use an increasing grid for representing the support of the process X ; it is straightforward to generalise the results from Section 4 to this setting.


The optimal solution of the corresponding lower bound MOT problem, subject to marginal constraints on the initial and the terminal marginals, was computed using $\varepsilon = 0.006$. The marginals of the computed optimal coupling are displayed in Figure 6a, where the marginals that are subject to a constraint are marked in green. We note that it corresponds to a late transport, as given by [57]. The smoothing is due to regularisation. A third marginal constraint was then added on the fourth marginal, by requiring that $\mu_4 = \frac{1}{2}(\delta_{29} + \delta_{31})$. The problem was then solved again; the marginals of the computed optimal coupling are shown in Figure 6b. We note that the transport from the initial marginal μ_0 to the given intermediate marginal μ_4 corresponds to a late transport, as conjectured by [57]. The problem was then solved a third time with a fourth marginal constraint imposed on the eighth marginal, with $\mu_8 = \frac{1}{4}(\delta_{28} + 2\delta_{30} + \delta_{32})$. The marginals of the optimal coupling are shown in Figure 6c.



(a) $\mathcal{T} = \{0, 11\}$



(b) $\mathcal{T} = \{0, 4, 11\}$



(c) $\mathcal{T} = \{0, 4, 8, 11\}$

Figure 6: The marginals of the optimal couplings corresponding to the lower bound MOT problem of Example 5.7, subject to marginal constraints on the marginals belonging to \mathcal{T} . Marginals subject to constraints are marked with a green background.

6 Summary and future work

We have proposed a method for approximately solving a class of multi-marginal MOT problems computationally and demonstrated its utility on a number of different examples. Our examples show that we are able to solve problems over a large number of marginals and that the computed solutions are aligned with theoretical results. We have not seen other examples of methods that are able to handle this many marginals in a martingale transport context — our method is thus, to the best of our knowledge, unique in this sense. One bottleneck is to numerically solve the equations characterising the optimal dual variables corresponding to martingale constraints in the primal problem. It is therefore our belief that the method could be made faster by amending this sub-routine. Another, related, concern is the propagation of the non-zero errors in the constraints. Since the number of martingale constraints are typically much larger than the number of marginal constraints, an important question is how the residuals in Newton’s method accumulates as the number of time steps increases. Our numerical experiments suggests that 30 marginals can be used without this being an issue, but the matter should be investigated further. Lastly, this paper constitutes (as far as we know) the first work applying entropic regularisation to MOT problems over many marginals. Several theoretical matters remains unaddressed, such as showing convergence of optimal solutions to the regularised problem as the regularisation parameter ε vanishes. Our computational results indicate that we are able to obtain a good approximate solution when ε is small, however, obtaining such results would be an important contribution to the field.

References

- [1] A. ALFONSI, R. COYAUD, V. EHRLACHER, AND D. LOMBARDI, *Approximation of optimal transport problems with marginal moments constraints*, Math. Comp., 90 (2021), pp. 689–737.
- [2] J. M. ALTSCHULER AND E. BOIX-ADSERÀ, *Polynomial-time algorithms for multimarginal optimal transport problems with structure*, Math. Program., 199 (2023), pp. 1107–1178.
- [3] J. AZÉMA AND M. YOR, *Une solution simple au probleme de Skorokhod*, in Séminaire de Probabilités XIII: Université de Strasbourg 1977/78, Springer, 2006, pp. 90–115.
- [4] F. BEIER, J. VON LINDHEIM, S. NEUMAYER, AND G. STEIDL, *Unbalanced multi-marginal optimal transport*, J. Math. Imaging Vision, 65 (2023), pp. 394–413.
- [5] M. BEIGLBÖCK, A. M. G. COX, AND M. HUESMANN, *The geometry of multi-marginal Skorokhod embedding*, Probab. Theory Related Fields, 176 (2020), pp. 1045–1096.
- [6] M. BEIGLBÖCK, P. HENRY-LABORDÈRE, AND F. PENKNER, *Model-independent bounds for option prices—a mass transport approach*, Finance Stoch., 17 (2013), pp. 477–501.
- [7] M. BEIGLBÖCK AND N. JUILLET, *On a problem of optimal transport under marginal martingale constraints*, Ann. Probab., 44 (2016), pp. 42–106.
- [8] J.-D. BENAMOU, G. CARLIER, M. CUTURI, L. NENNA, AND G. PEYRÉ, *Iterative Bregman projections for regularized transportation problems*, SIAM J. Sci. Comput., 37 (2015), pp. A1111–A1138.
- [9] D. T. BREEDEN AND R. H. LITZENBERGER, *Prices of state-contingent claims implicit in option prices*, J. Bus., 51 (1978), pp. 621–651.
- [10] H. BROWN, D. HOBSON, AND L. C. G. ROGERS, *The maximum maximum of a martingale constrained by an intermediate law*, Probab. Theory Related Fields, 119 (2001), pp. 558–578.
- [11] ———, *Robust hedging of barrier options*, Math. Finance, 11 (2001), pp. 285–314.
- [12] P. BUCHEN, *An introduction to exotic option pricing*, Chapman & Hall/CRC Financial Mathematics Series, CRC Press, Boca Raton, FL, 2012.

- [13] P. CHERIDITO, M. KUPPER, AND L. TANGPI, *Duality formulas for robust pricing and hedging in discrete time*, SIAM J. Financial Math., 8 (2017), pp. 738–765.
- [14] L. CHIZAT, G. PEYRÉ, B. SCHMITZER, AND F.-X. VIALARD, *Scaling algorithms for unbalanced optimal transport problems*, Math. Comp., 87 (2018), pp. 2563–2609.
- [15] A. M. G. COX, J. OBLÒJ, AND N. TOUZI, *The Root solution to the multi-marginal embedding problem: an optimal stopping and time-reversal approach*, Probab. Theory Related Fields, 173 (2019), pp. 211–259.
- [16] M. CUTURI, *Sinkhorn distances: Lightspeed computation of optimal transport*, in Advances in Neural Information Processing Systems, Curran Associates, 2013, pp. 2292–2300.
- [17] M. DAVIS, J. OBLÒJ, AND V. RAVAL, *Arbitrage bounds for prices of weighted variance swaps*, Math. Finance, 24 (2014), pp. 821–854.
- [18] H. DE MARCH, *Entropic approximation for multi-dimensional martingale optimal transport*, arXiv:1812.11104, (2018).
- [19] Y. DOLINSKY AND H. M. SONER, *Martingale optimal transport and robust hedging in continuous time*, Probab. Theory Related Fields, 160 (2014), pp. 391–427.
- [20] S. ECKSTEIN, G. GUO, T. LIM, AND J. OBLÒJ, *Robust pricing and hedging of options on multiple assets and its numerics*, SIAM J. Financial Math., 12 (2021), pp. 158–188.
- [21] S. ECKSTEIN AND M. KUPPER, *Computation of optimal transport and related hedging problems via penalization and neural networks*, Appl. Math. Optim., 83 (2021), pp. 639–667.
- [22] ———, *Martingale transport with homogeneous stock movements*, Quant. Finance, 21 (2021), pp. 271–280.
- [23] F. ELVANDER, I. HAASLER, A. JAKOBSSON, AND J. KARLSSON, *Multi-marginal optimal transport using partial information with applications in robust localization and sensor fusion*, Signal Process., 171 (2020), p. 107474.
- [24] H. FÖLLMER AND A. SCHIED, *Stochastic finance : An introduction in discrete time*, De Gruyter, Berlin, 4th ed., 2016.
- [25] J. FRANKLIN AND J. LORENZ, *On the scaling of multidimensional matrices*, Linear Algebra Appl., 114 (1989), pp. 717–735.
- [26] A. GALICHON, P. HENRY-LABORDÈRE, AND N. TOUZI, *A stochastic control approach to no-arbitrage bounds given marginals, with an application to lookback options*, Ann. Appl. Probab., 24 (2014), pp. 312–336.
- [27] W. GANGBO AND A. ŚWIĘCH, *Optimal maps for the multidimensional Monge-Kantorovich problem*, Comm. Pure Appl. Math., 51 (1998), pp. 23–45.
- [28] T. T. GEORGIU, J. KARLSSON, AND M. S. TAKYAR, *Metrics for power spectra: an axiomatic approach*, IEEE Transactions on Signal Processing, 57 (2008), pp. 859–867.
- [29] G. GUO AND J. OBLÓJ, *Computational methods for martingale optimal transport problems*, Ann. Appl. Probab., 29 (2019), pp. 3311–3347.
- [30] I. HAASLER, J. KARLSSON, AND A. RINGH, *Control and estimation of ensembles via structured optimal transport*, IEEE Control Syst., 41 (2021), pp. 50–69.
- [31] I. HAASLER, A. RINGH, Y. CHEN, AND J. KARLSSON, *Multimarginal optimal transport with a tree-structured cost and the Schrödinger bridge problem*, SIAM J. Control Optim., 59 (2021), pp. 2428–2453.
- [32] ———, *Scalable computation of dynamic flow problems via multimarginal graph-structured optimal transport*, Math. Oper. Res., In Press (2023).
- [33] I. HAASLER, R. SINGH, Q. ZHANG, J. KARLSSON, AND Y. CHEN, *Multi-marginal optimal transport and probabilistic graphical models*, IEEE Trans. Inform. Theory, 67 (2021), pp. 4647–4668.

- [34] P. HENRY-LABORDÈRE, *Automated option pricing: numerical methods*, Int. J. Theor. Appl. Finance, 16 (2013), pp. 1350042, 27.
- [35] P. HENRY-LABORDÈRE, J. OBLÓJ, P. SPOIDA, AND N. TOUZI, *The maximum maximum of a martingale with given n marginals*, Ann. Appl. Probab., 26 (2016), pp. 1–44.
- [36] P. HENRY-LABORDÈRE AND N. TOUZI, *An explicit martingale version of the one-dimensional Brenier theorem*, Finance Stoch., 20 (2016), pp. 635–668.
- [37] P. HENRY-LABORDÈRE, *Model-free Hedging: A Martingale Optimal Transport Viewpoint*, Chapman and Hall/CRC, New York, 1st ed., 2017.
- [38] D. G. HOBSON, *Robust hedging of the lookback option*, Finance Stoch., 2 (1998), pp. 329–347.
- [39] J. C. HULL, *Options, Futures and Other Derivatives*, Pearson Education Ltd., Harlow, 11th ed., 2022.
- [40] S. KÄLLBLAD, X. TAN, AND N. TOUZI, *Optimal Skorokhod embedding given full marginals and Azéma-Yor peacocks*, Ann. Appl. Probab., 27 (2017), pp. 686–719.
- [41] O. KALLENBERG, *Foundations of modern probability*, vol. 99 of Probability Theory and Stochastic Modelling, Springer, New York, 3rd ed., 2021.
- [42] J. KARLSSON AND A. RINGH, *Generalized Sinkhorn iterations for regularizing inverse problems using optimal mass transport*, SIAM J. Imaging Sci., 10 (2017), pp. 1935–1962.
- [43] M. LIERO, A. MIELKE, AND G. SAVARÉ, *Optimal entropy-transport problems and a new Hellinger-Kantorovich distance between positive measures*, Invent. Math., 211 (2018), pp. 969–1117.
- [44] T. LIN, N. HO, M. CUTURI, AND M. I. JORDAN, *On the complexity of approximating multi-marginal optimal transport*, J. Mach. Learn. Res., 23 (2022), pp. 1–43.
- [45] Z. Q. LUO AND P. TSENG, *On the convergence of the coordinate descent method for convex differentiable minimization*, J. Optim. Theory Appl., 72 (1992), pp. 7–35.
- [46] M. MASCHERPA, I. HAASLER, B. AHLGREN, AND J. KARLSSON, *Estimating pollution spread in water networks as a Schrödinger bridge problem with partial information*, Eur. J. Control, 74 (2023), pp. 1–8.
- [47] S. G. NASH AND A. SOFER, *Linear and nonlinear programming*, McGraw-Hill series in industrial engineering and management science, McGraw-Hill, New York, 1996.
- [48] M. NUTZ, F. STEBEGG, AND X. TAN, *Multiperiod martingale transport*, Stochastic Process. Appl., 130 (2020), pp. 1568–1615.
- [49] H. H. PANJER, P. P. BOYLE, S. H. COX, D. DUFRESNE, H. GERBER, H. MUELLER, H. PEDERSEN, S. PLISKA, M. SHERRIS, E. SHIU, ET AL., *Financial economics : With applications to investments, insurance, and pensions*, Society of Actuaries Foundation, Schaumburg, Ill., 2001.
- [50] B. PASS, *Multi-marginal optimal transport: theory and applications*, ESAIM Math. Model. Numer. Anal., 49 (2015), pp. 1771–1790.
- [51] G. PEYRÉ AND M. CUTURI, *Computational optimal transport: With applications to data science*, Found. Trends Mach. Learn., 11 (2019), pp. 355–607.
- [52] B. PICCOLI AND F. ROSSI, *Generalized Wasserstein distance and its application to transport equations with source*, Arch. Ration. Mech. Anal., 211 (2014), pp. 335–358.
- [53] L. RÜSCHENDORF, *Optimal solutions of multivariate coupling problems*, Appl. Math. (Warsaw), 23 (1995), pp. 325–338.
- [54] J. SESTER, *Robust bounds for derivative prices in Markovian models*, Int. J. Theor. Appl. Finance, 23 (2020), pp. 2050015, 39.
- [55] ———, *A multi-marginal c -convex duality theorem for martingale optimal transport*, Statist. Probab. Lett., 210 (2024), p. 110112.

- [56] R. SINKHORN, *Diagonal equivalence to matrices with prescribed row and column sums*, Amer. Math. Monthly, 74 (1967), pp. 402–405.
- [57] F. STEBEGG, *Model-independent pricing of Asian options via optimal martingale transport*, arXiv:1412.1429, (2014).
- [58] V. STRASSEN, *The existence of probability measures with given marginals*, Ann. Math. Statist., 36 (1965), pp. 423–439.
- [59] X. TAN AND N. TOUZI, *Optimal transportation under controlled stochastic dynamics*, Ann. Probab., 41 (2013), pp. 3201–3240.
- [60] C. VILLANI, *Topics in optimal transportation*, vol. 58 of Graduate Studies in Mathematics, American Mathematical Society, Providence, RI, 2003.
- [61] J. VON LINDHEIM AND G. STEIDL, *Generalized iterative scaling for regularized optimal transport with affine constraints: Application examples*, arXiv:2305.07071, (2023).
- [62] P. WILMOTT, *The best of Wilmott*, vol. 1, John Wiley & Sons Inc., Chichester, 2005.