

The work of Walter Bergweiler in value distribution of meromorphic functions

Alexandre Eremenko

June 17, 2024

Abstract

This is the colloquium talk in CAU Kiel delivered on June 7, 2024 on the occasion of Walter Bergweiler's retirement. It contains a survey of his work in the value distribution theory of meromorphic functions of one complex variable. The talk was aimed at a broad audience, so a brief introduction to Nevanlinna theory is included.

2010 MSC: 30D35. Keywords: meromorphic function, differential polynomial, value distribution, Nevanlinna theory, Bloch's Principle.

Value distribution theory begins with the celebrated theorem of Emile Picard:

If an entire function omits two values, then it is constant.

This theorem (and its two-lines proof) published in 1879 made an enormous impression, and they say that all seminar talks in Paris in the early 20th century began with the words: "Picard proved...".

One can restate it as follows *if a meromorphic function in the plane omits three values then it is constant*. The generalization known as the Great Picard's theorem says that *If a meromorphic function in a punctured neighborhood of a point omits three values in $\overline{\mathbb{C}}$ then this point is a pole or removable singularity*.

Schottky [31] proved the corresponding result for the unit disk. We state it in the form proposed by Montel: *The family of all meromorphic functions in the unit disk which omit three values, say $0, 1, \infty$ is normal*. I recall that a family F of meromorphic functions in a region D is called *normal* if from any sequence (f_n) in F one can select a subsequence which converges uniformly

(with respect to the spherical metric in the image) on every compact subset of D .

One can show that the theorems of Picard and Montel can be simply derived from each other, and this is the basis of the famous “Bloch’s Principle”: *If some conditions imply that a meromorphic function in the plane is constant, then the same conditions imposed on a family of meromorphic functions in any region imply that this family is normal, and conversely.*

This heuristic principle turned out to be very fruitful, see, for example Walter’s survey [5]. It is equally important in the theory of holomorphic curves, [21], [34].

The condition that some value a is omitted can be usually replaced by the condition that all solutions of $f(z) = a$ have high multiplicity. Nevanlinna proved the following generalization of Picard’s theorem: *If f is a non-constant meromorphic function in \mathbf{C} , and a_j have the property that all solutions of $f(z) = a_j$ have multiplicities at least m_j , then*

$$\sum_j \left(1 - \frac{1}{m_j}\right) \leq 2, \tag{1}$$

where $m_j = \infty$ if a_j is omitted. This result was first obtained by Nevanlinna, as a consequence of his deep *Nevanlinna theory* which can be considered as a far-reaching quantification of Picard’s theorems. Since its inception it became the main part of the theory of meromorphic functions, and I will briefly introduce the main notions and results. Let $f : \mathbf{C} \rightarrow \overline{\mathbf{C}}$ be a meromorphic function, and

$$A(r, f) = \frac{1}{\pi} \int \int_{x^2+y^2 \leq r^2} \frac{|f'(x+iy)|^2}{(1+|f(x+iy)|^2)^2} dx dy,$$

the area of the image of the disk under f with respect to the spherical metric, divided by the area of the sphere. So for a rational function of degree d we have $A(r, f) \rightarrow d, r \rightarrow \infty$. Then the Nevanlinna characteristic is defined as an average of $A(r, f)$ with respect to r :

$$T(r, f) = \int_0^r \frac{A(t, f)}{t} dt.$$

This function measures the growth of a meromorphic function. For a rational function $T(r, f) \sim d \log r$, and for transcendental meromorphic functions we have $T(r, f)/\log r \rightarrow \infty$. Nevanlinna characteristic has formal algebraic

properties similar to the degree of a rational function, and this is one reason why we averaged with respect to r . For example,

$$T(r, f) = T(r, 1/f), \quad T(r, f^n) = nT(r, f), \quad T(r, fg) \leq T(r, f) + T(r, g) + O(1),$$

and $T(r, f + g) \leq T(r, f) + T(r, g) + O(1)$.

The order of a meromorphic function is defined as

$$\rho := \limsup_{r \rightarrow \infty} \frac{\log T(r, f)}{\log r} \in [0, \infty].$$

So for example, e^z and $\sin z$ are of order 1, e^{z^p} is of order p , $\wp(z)$ has order 2, and e^{e^z} is of infinite order.

For every $a \in \overline{\mathbf{C}}$ we define $n(r, a, f)$ as the number (counting multiplicity) of solutions of $f(z) = a$ in the disk $|z| \leq r$, and then apply the averaging in r as above:

$$N(r, a, f) = \int_0^r (n(t, a, f) - n(0, a, f)) \frac{dt}{t}.$$

Then the First Main Theorem of Nevanlinna says that for all $a \in \overline{\mathbf{C}}$ we have

$$N(r, a, f) \leq T(r, f) + O(1), \quad r \rightarrow \infty,$$

and the Second Main Theorem says that for any finite set $\{a_1, \dots, a_q\}$

$$\sum_{j=1}^q N(r, a_j, f) \geq (q - 2 + o(1))T(r, f) + N_1(r, f),$$

when $r \rightarrow \infty$ avoiding a set of finite measure. Here $N_1(r, f)$ is the ramification term: $n_1(r, f)$ is the number of critical points (counting multiplicity) in the disk $|z| \leq r$, and N_1 is obtained from n_1 by the averaging as above. The meaning of the Second Main Theorem can be made clearer if we define the deficiency

$$\delta(a, f) = 1 - \limsup_{r \rightarrow \infty} \frac{N(r, a, f)}{T(r, f)}.$$

So $0 \leq \delta(a, f) \leq 1$, and if the value a is omitted then $\delta(a, f) = 1$. Then the Second Main Theorem implies that $\delta(a, f) > 0$ for at most countable set of a (they are called deficient), and

$$\sum_{a \in \overline{\mathbf{C}}} \delta(a, f) \leq 2. \tag{2}$$

This is a generalization of Picard's theorem. If one takes multiplicities into account, then (1) follows.

Returning to Picard's theorem, it inspired a lot of variations and generalizations in 20th century. Our main subject here will be the analogs of Picard's theorem involving derivatives.

Among the earliest results on this subject, I mention a theorem of Milloux [22] *If an entire function omits some finite value, and its derivative omits some non-zero value, then f is constant.* The corresponding normality criterion was proved by Miranda [23] 5 years earlier. Example e^z shows that f' can indeed omit zero. This shows that the value 0 is special for the derivative.

On the other hand, Csillag [13] proved that *If f is entire and $f, f^{(m)}, f^{(n)}$ omit zero for some $0 < m < n$, then $f(z) = e^{az+b}$.* The case $(m, n) = (1, 2)$ was earlier proved by Saxer.

A great breakthrough in these questions was made by Hayman [16]. The main result of his article is the so-called

Hayman's Alternative. *If f is a transcendental meromorphic function in the plane and n is a positive integer, then either f assumes every finite value infinitely many times, or $f^{(n)}$ assumes every finite non-zero value infinitely many times.*

It is noteworthy that this theorem gives only two conditions on omitted values which imply that the function is constant. Picard, Schottky, Milloux and Csillag theorems when stated for meromorphic functions require 3 conditions (they all assume that ∞ is omitted).

Hayman's result generalizes the theorem of Milloux to meromorphic functions.

In this paper Hayman proved many other results, but he assumed that they are not best possible, and stated several conjectures. The most outstanding conjectures are these two:

Hayman's Conjecture 1. *If f is a transcendental meromorphic function in \mathbf{C} , and $n \geq 2$, then $(f^n)'$ assumes all non-zero complex values infinitely many times.*

He himself proved this for $n \geq 4$, and later Mues extended this to $n = 3$, and Clunie proved this conjecture for entire functions.

Hayman's Conjecture 2. *If f is a meromorphic function in \mathbf{C} and both*

f and $f^{(n)}$ are free of zeros for some $n \geq 2$, then $f(z) = e^{az+b}$ or $f(z) = (az + b)^{-m}$ with some integer m .

This was proved by Frank for $n \geq 3$ and by Mues for functions of finite order. The final result with $n = 2$ is due to Langley [17] who developed the methods of Mues and Frank in a very subtle way.

I met Walter in winter 1992/3 in a conference in Joensuu. For me this was probably the first international conference that I was able to attend (I moved from the Soviet Union to the US in 1991). We shared a 3-bedroom apartment with my adviser A. A. Goldberg and for him this was also one of the first trips abroad. We had no common language for all three of us: Goldberg did not speak English, so he had to communicate with Walter in German and with me in Russian. And I translated his conference talk into English.

Next summer I was invited by Walter to stay with him and his wife in Aachen, where he had his first position. When meeting me in the train station, he explained to me his proposal to prove Hayman's Conjecture 1 in the remaining case $n = 2$ for functions of finite order.

The result was the paper [6] which we wrote that summer and submitted to the journal. The proof was unusual for the theory of meromorphic functions since it used holomorphic dynamics. Holomorphic dynamics which was originated by Fatou (with some result obtained independently by Julia, Lattès and Ritt) in the beginning of 20th century experienced a strong revival since 1982.

We were both interested in holomorphic dynamics at that time: I was deeply impressed with the work of Misha Lyubich in the early 1980s, and we wrote several joint papers in Kharkiv on dynamics of transcendental functions. Walter started thinking about holomorphic dynamics under the influence of J. Hubbard, when he visited Cornell University as a postdoc in 1987-89. By 1993 we both wrote surveys on holomorphic dynamics: my survey was joint with Lyubich, and Walter's survey of dynamics of transcendental meromorphic function is to this time the best introductory survey; it accumulated 417 citations on MSN by the time I write this.

The idea that Walter explained to me in the train station was the following: if f has finitely many zeros, then Hayman's alternative is applicable, and we are done. So assume that it has infinitely many zeros. They are all

multiple zeros of f^n . Then the function

$$g(z) = z - f^n(z)$$

has infinitely many fixed points, and these fixed points are neutral, with multiplier 1, that is $g'(a) = 1$ at all these fixed points a . To each such fixed point an immediate domain of attraction P is associated (the so called Leau's petal). This domain is invariant under g , and according to the fundamental theorem of Fatou, P must contain either critical or asymptotic value of g . (We say that $a \in \overline{\mathbf{C}}$ is an asymptotic value of a function g meromorphic in \mathbf{C} if there is a curve tending to ∞ such that $g(z) \rightarrow a$ along this curve.) If this singular value which exists by Fatou's theorem is critical then $g'(z) = 0$ at some point z in the domain of attraction of P , and since these domains are disjoint for different neutral fixed points, we obtain infinitely many critical points, which are solutions of $(f^n)' = 1$.

So it remains to deal with the possibility that g has infinitely many asymptotic values but only finitely many critical values. Such meromorphic functions exist, an example can be found in [26].

But it turns out that such thing is impossible for functions of *finite order*. This was a great insight of Walter which he explained to me during that conversation in the train station. It took few weeks to work out the details, and the final result was the following

Theorem 1. *Let f be a meromorphic function of finite order ρ . Then every asymptotic value of f , except at most 2ρ of them, is a limit point of critical values different from this asymptotic value.*

By using the dynamics argument outlined above, we obtain

Theorem 2. *Let f be a transcendental meromorphic function of finite order, and assume that it has infinitely many multiple zeros. Then f' assumes every finite non-zero value infinitely often.*

Unfortunately, this theorem fails for functions of infinite order, so it seemed that we could not prove Hayman's Conjecture 1 in full generality with this method.

I knew from the work of Zalcman [35] and Brody that there exists a marvelous device to reduce Picard-type theorems to their special cases for functions of finite order. This is called the Zalcman Rescaling Lemma to the specialists in classical function theory, and Brody's Rescaling Lemma to

the specialists in holomorphic curves. Actually this argument goes back to the work of Bloch and Valiron in the beginning of 20th century. However this rescaling argument did not seem to be appropriate for working with derivatives.

Only next summer (1994) when browsing the book by Schiff on normal families, I learned about the paper by X. Pang [27] of 1989 where a suitable generalization of Zalcman's Lemma was proved¹. I immediately informed Walter (and Zalcman) and since our paper was not yet published, we managed to revise it, and obtain the full proof of Hayman's conjecture [6]. Our final result was

Theorem 3. *For every transcendental meromorphic function f , and every positive integers $k < n$, the function $(f^n)^{(k)}$ takes every finite non-zero value infinitely often.*

Meanwhile, the statement of our Theorem 2 reached China (this was before the advent of the arXiv): Walter gave a conference talk in Nanjing in June 1994, and the consequence was an "independent" proof of Hayman's conjecture published by Huaihui Chen and Mingliang Fang [12]. According to a later personal communication from X. Pang, he knew how to reduce Hayman's conjecture to its special case of finite order even before our paper was written, but Pang was not present at Walter's talk in Nanjing.

In a later paper [4], Walter applied Pang's lemma to give a new simple proof of Langley's theorem stated above. The corresponding normality criterion is

Let F be a family of meromorphic functions f such that ff'' omit zero. Then the family $\{f'/f : f \in F\}$ is normal.

Since then our Theorem 3 was very much used and generalized, paper [6] has 230 citations on MSN and 550 on Google Scholar.

The ultimate generalization for the first derivative was conjectured by Walter and proved by Jianming Chang [11], who used a deep intermediate result of Nevo, Pang and Zalcman [25]:

Suppose that for a transcendental meromorphic function f , $|f'|$ is bounded on the zero set of f . Then f takes every finite non-zero value infinitely often.

There is an essential difference between this and Theorem 2: (which is

¹Independently of Pang, similar arguments were used in Schwick's thesis [32].

not true for functions of infinite order): In Theorem 2 the assumption is that there are infinitely many multiple zeros, while in the theorem of Chang the assumption of boundedness of $|f'|$ applies to *all* zeros.

Despite many variations and generalizations, no new proof of Theorem 3 was found until 2020: all variants and generalizations used our Theorem 3. I always felt that our proof was unsatisfactory from the philosophical point of view. One reason for this was the use of dynamics: why a non-dynamical theorem should use dynamics in its proof? Walter showed me how dynamics can be eliminated (see [14] where his argument is recorded). It uses instead the so-called Logarithmic Change of the Variable, which is a common tool in transcendental dynamics, but formally does not belong to it. (It was invented by Teichmüller for a problem of value distribution theory).

More important philosophical objection is related to the weird overall scheme: we prove Theorem 2 for functions of finite order, (*and it is not true without this restriction!*). And then by some magic trick we derive Theorem 3 which applies to all meromorphic functions.

The new proof, which uses neither our arguments nor Pang's lemma, was obtained recently in [1] as a corollary of the very deep result of Yamanoi [33], and I briefly describe his result.

Erwin Mues conjectured in 1971 that the defect relation for the derivative of a meromorphic function can be improved to

$$\sum_{a \neq \infty} \delta(a, f') \leq 1.$$

In his attempts to prove this conjecture Goldberg made a new one *For all meromorphic functions*

$$\overline{N}(r, \infty, f) \leq N(r, 0, f'') + o(T(r, f)), \quad (3)$$

(as $r \rightarrow \infty$ avoiding a small exceptional set), where \overline{N} is the averaged counting function of distinct poles (that is counted without multiplicity). Goldberg showed that this implies Mues's conjecture, and proved (3) for the case when all poles of f are simple. Actually, both the Mues conjecture and Goldberg's conjecture for the case of simple poles can be derived from the arguments of Frank and Weissenborn [15].

It is amazing that no perturbation argument seems to derive the Mues and Goldberg conjectures from this special "generic" case.

Langley [19] proved that for meromorphic functions of finite order the condition that $f^{(k)}$ for some $k \geq 2$ has finitely many zeros implies that f

has finitely many poles. However he constructed an example of a meromorphic function of arbitrarily slow infinite order of growth for which f'' has no zeros but f has infinitely many poles. So there is no Picard-type theorem corresponding to Goldberg's conjecture!

Goldberg's conjecture was proved by Yamanoi [33] in the following stronger form

$$k\overline{N}(r, \infty, f) + \sum_{a \in A} N_1(r, a, f) \leq N(r, 0, f^{(k+1)}) + o(T(r, f))$$

outside a set of r of zero logarithmic density, where k is any positive integer and A is any finite set. This 77 pages paper is a real tour de force; this is probably the most difficult of all results in value distribution of meromorphic functions in dimension one. In the recent decades, the main interests in Nevanlinna theory shifted from meromorphic functions to holomorphic curves, but this result of Yamanoi (and his other results) on one-dimensional Nevanlinna theory show that it is still very much alive.

I make few comments on Theorem 1, which is of independent interest. It is related to the famous Denjoy–Carleman–Ahlfors theorem, and Iversen's classification of singularities of inverse functions. If a is an asymptotic value of a meromorphic function f , and $D(a, r)$ is an arbitrarily small (spherical) disk centered at a , then the preimage $f^{-1}(D(a, r))$ has an unbounded component V , the one that contains the asymptotic curve. If for some $r > 0$ the equation $f(z) = a$ has no solutions in V , we say that the singularity of f^{-1} over a is direct. Otherwise it is indirect. For example $\log z$ has two direct singularities: one over 0 another over ∞ . And $(\sin z)/z$ has two indirect singularities over 0 and two direct ones over ∞ .

The famous Denjoy–Carleman–Ahlfors theorem says that for a meromorphic function of order ρ , the inverse has at most 2ρ direct singularities. And the more precise form of our Theorem 1 is:

If f is a meromorphic function of finite order, then every neighborhood of an indirect singularity over a (that is every component V in the above explanation) contains infinitely many critical points with critical values distinct from a .

Iversen's classification was rarely used in value distribution theory before our paper, but since its publication it became a common tool in value distribution and transcendental holomorphic dynamics.

Returning to Walter's work, I would like to mention here three his other contributions.

In his paper [2] he proved the following

Theorem 4. *Let f be a meromorphic function of finite order, and $a \in \mathbf{C}$. If $a \neq 1$ and $a \neq (n+1)/n$ for all $n \in \mathbf{N}$, then $ff'' - a(f')^2$ has infinitely many zeros, unless $f = Re^P$ with rational R and polynomial P .*

He conjectured that the assumption of finite order can be removed in this theorem; this would give a generalization of Langley's proof of Hayman's Conjecture 2, which corresponds to $a = 0$. Walter's conjecture was proved by Langley [18], without use of Pang's lemma. A simple derivation based on Pang's Lemma appeared in [20].

In the joint paper [8] Walter and Jim study the question whether in Hayman's alternative one can replace omitted values by high multiplicity. As expected, they show that this is so, but probably their results are not best possible. Improving a previous result by Zhen Hua Chen, they show that a meromorphic function f whose zeros have multiplicity $\geq m$ while 1-points of $f^{(k)}$ have multiplicity $\geq n$ must be constant if

$$\left(2k + 3 + \frac{2}{k}\right)/m + \left(2k + 2 + \frac{2}{k}\right)/n < 1.$$

So when $k = 1$, we have the condition

$$\frac{7}{m} + \frac{8}{n} < 1.$$

They also constructed an example of a non-constant meromorphic function with $m = 2$ and $n = 3$. On the other hand, Nevo, Pang and Zalcman proved that

The derivative of a transcendental meromorphic function whose almost all zeros are multiple takes every non-zero value infinitely often.

Notice that the rational function

$$f(z) = \frac{(z-a)^2}{(z-b)}$$

has only one zero in \mathbf{C} , this zero is multiple, and f' omits 1.

So it is an open problem to obtain a best possible result about multiplicities in Hayman's alternative. The statement about normality corresponding to this via Bloch's Principle also holds in this setting.

In his influential paper [10] Bloch mentioned two philosophical principles: one that we stated in the beginning and another the “Principle of topological continuity”. To explain this second principle, consider a disk $V \subset \overline{\mathbf{C}}$. A component of preimage $f^{-1}(V)$ under a meromorphic function f is called an *island* over V if this component is bounded. In this case the restriction of f on this component is a ramified covering and thus has a *degree* which is a positive integer. This degree is called the multiplicity of an island. If $a \in V$ is a point, and f has an a -point of multiplicity m in an island over V , then the multiplicity of the island is at least m . Bloch proposed the following generalization of (1):

If V_j are disks with disjoint closures, and the multiplicity of all islands over V_j of a non-constant meromorphic function f in \mathbf{C} are at least m_j , then (1) holds.

If there are no islands over V_j we set $m_j = \infty$.

Bloch’s conjecture was proved as a consequence of very deep “Überlagerungsflächentheorie” of Ahlfors for which he obtained one of the first two Fields medals in 1936. This theory is very technical, and it does not use the Nevanlinna’s result (1). A modern proof of (1) can be obtained without Nevanlinna theory, a simple proof was obtained by Robinson.

Several simplified derivations of Ahlfors theory also exist, but none of them can be called really simple.

Walter found in [3] a simple derivation of the Ahlfors island result from (1), in complete agreement with Bloch’s vision. So this is by far the simplest available proof of the Ahlfors’s islands theorem.

The “Principle of Topological Continuity” as stated by Bloch, is not a theorem but a philosophical principle, and in our most recent paper with Walter [7] we investigate the cases when it can be converted into theorems. For example, one simple consequence of Nevanlinna theory (proved by H. Selberg) is

Let f be a non-constant entire function of order less than 1, and a, b are two points in \mathbf{C} . Then f one of these two points has a simple preimage.

Notice that $\cos \sqrt{z}$ is of order $1/2$, and has only one simple preimage of the pair of points $\{\pm 1\}$.

Bloch stated that to this should correspond:

Let f be a non-constant entire function of order less than 1, and A, B are two discs in \mathbf{C} with disjoint closures. Then over one of these disks there is

a simple island.

We proved this in [7]. But some other similar statements involving meromorphic functions and simple islands over one of the 4 disks with disjoint closures turn out to be wrong.

References

- [1] Ta Thi Hoai An and Nguyen Viet Phuong, A note on Hayman's conjecture, *Internat. J. Math.* 31 (2020), no. 6, 2050048, 10 pp.
- [2] W. Bergweiler, On the zeros of certain homogeneous differential polynomials, *Arch. Math. (Basel)* 64 (1995), no. 3, 199–202.
- [3] W. Bergweiler, A new proof of the Ahlfors five islands theorem, *J. Anal. Math.* 76 (1998), 337–347.
- [4] W. Bergweiler, Normality and exceptional values of derivatives, *Proc. AMS*, 129 (2000) 1, 121–129.
- [5] W. Bergweiler, Bloch's principle, *Comput. Methods Funct. Theory* 6 (2006), no. 1, 77–108.
- [6] W. Bergweiler and A. Eremenko, On the singularities of the inverse to a meromorphic function of finite order, *Rev. Mat. Iberoamericana* 11 (1995), no. 2, 355–373.
- [7] W. Bergweiler and A. Eremenko, On Bloch's Principle of Topological Continuity, arXiv:2309.14749 to appear in *CMFT*.
- [8] W. Bergweiler and J. Langley, Multiplicities in Hayman's alternative, *J. Aust. Math. Soc.* 78 (2005), no. 1, 37–57.
- [9] W. Bergweiler and X. Pang, On the derivative of meromorphic functions with multiple zeros, *J. Math. Anal. Appl.* 278 (2003), no. 2, 285–292.
- [10] A. Bloch, La conception actuelle de la théorie des fonctions entières et méromorphes, *Enseignement Math.* 25 (1926), 83–103.

- [11] J. Chang, On meromorphic functions whose first derivative has finitely many zeros, *Bull. LMS*, 44 (2012) 703–715.
- [12] H. H. Chen and M. L. Fang, On the value distribution of $f^n f'$, *Science in China*, 38, 7 (1995) 789–798.
- [13] P. Csillag, Über ganze Funktionen, welche drei nicht verschwindende Ableitungen besitzen, *Math. Ann.* 110 (1935), 745–752.
- [14] A. Eremenko, Interactions between holomorphic dynamics and function theory, arXiv:2007.02799.
- [15] G. Frank and Weissenborn, Rational deficient functions of meromorphic functions. *Bull. London Math. Soc.* 18 (1986), no. 1, 29–33.
- [16] W. Hayman, Picard values of meromorphic functions and their derivatives, *Ann. of Math. (2)* 70 (1959), 9–42.
- [17] J. Langley, Proof of a conjecture of Hayman concerning f and f'' , *J. London Math. Soc. (2)* 48 (1993), no. 3, 500–514.
- [18] J. Langley, A lower bound for the number of zeros of a meromorphic function and its second derivative, *Proc. Edinburgh Math. Soc. (2)* 39 (1996), no. 1, 171–185.
- [19] J. Langley, The second derivative of a meromorphic function of finite order, *Bull. London Math. Soc.* 35 (2003), no. 1, 97–108.
- [20] Wei-Chuan Lin and Hong-Xun Yi, On a conjecture of W. Bergweiler, *Proc. Japan Acad. Ser. A Math. Sci.* 79 (2003), no. 2, 23–27
- [21] M. McQuillan, The Bloch Principle, arxiv.org:1209.5402
- [22] H. Milloux, *Sur les fonctions méromorphes et leurs dérivées*, Paris, 1940.
- [23] C. Miranda, Sur un nouveau critère de normalité pour les familles de fonctions holomorphes, *Bull. Soc. Math. Fr.* 63, (1935) 185–196 (1935).
- [24] P. Montel, Sur les fonctions analytiques qui admettent deux valeurs exceptionnelles dans un domaine, *C. R.* 153 (1911) 996–998.

- [25] S. Nevo, X. Pang, and L. Zalcman, Quasinormality and meromorphic functions with multiple zeros, *Journal d'Analyse*, 101 (2007) 1–23. *Electron. Res. Announc. Amer. Math. Soc.* 12 (2006), 37–43.
- [26] L. I. Volkovyskii, Research on the Type Problem of a Simply Connected Riemann Surface, *Proc. Steklov Math. Inst.* 34, Acad. Sci. USSR, Moscow, 1950.
- [27] Xue Cheng Pang, Bloch's principle and normal criterion, *Science in China* 32 (1989), 782–791.
- [28] E. Picard, Sur les fonctions entières, *C. R.* 89 (1880) 662–665.
- [29] E. Picard, Sur les fonctions analytiques uniformes dans le voisinage d'un point singulier essentiels, *C. R.* 88 (1879) 745–747.
- [30] E. Picard, Sur une propriété des fonctions entières, *C. R.* 88 (1879) 1024–1027.
- [31] F. Schottky, Über den Picardschen Satz und die Borelschen Ungleichungen, *Sitzungsber. Preuss. Acad.* (1904), 1244–1263.
- [32] W. Schwick, Normality criteria for families of meromorphic functions, *J. d'Analyse*, 52 (1989) 241–289.
- [33] K. Yamanoi, Zeros of higher derivatives of meromorphic functions in the complex plane, *Proc. Lond. Math. Soc.* (3) 106 (2013), no. 4, 703–780.
- [34] K. Yamanoi, Bloch's Principle for holomorphic maps into subvarieties of semi-Abelian varieties, arXiv:2304.057
- [35] L. Zalcman, A heuristic principle in complex function theory, *Amer. Math. Monthly* 82 (1975), 813–817.

Department of Mathematics
Purdue University
West Lafayette, IN 47907
eremenko@purdue.edu