OPTIMAL RATES FOR FUNCTIONAL LINEAR REGRESSION WITH GENERAL REGULARIZATION

NAVEEN GUPTA, S. SIVANANTHAN, AND BHARATH K. SRIPERUMBUDUR

ABSTRACT. Functional linear regression is one of the fundamental and well-studied methods in functional data analysis. In this work, we investigate the functional linear regression model within the context of reproducing kernel Hilbert space by employing general spectral regularization to approximate the slope function with certain smoothness assumptions. We establish optimal convergence rates for estimation and prediction errors associated with the proposed method under a Hölder type source condition, which generalizes and sharpens all the known results in the literature.

1. INTRODUCTION

Functional data analysis (FDA) deals with analyzing and extracting information from curves, known as functional data, that include growth curves, weather data, health data, etc. The functional linear regression (FLR) model introduced by Ramsay and Dalzell [28] has emerged as one of the mainstays for analyzing functional data [29, 27, 19, 30, 23, 37]. Formally, the FLR model is stated as

$$Y = \int_{S} X(t)\beta^{*}(t) dt + \epsilon, \qquad (1.1)$$

where $Y(\omega)$ is a real-valued random variable, $(X(\omega, t); t \in S)$ is a continuous time process, ϵ is a zero mean random noise, independent of X, with finite variance σ^2 and β^* is an unknown slope function. Throughout the paper, we assume that X and β^* are in $L^2(S)$, $\mathbb{E}[||X||_{L^2}^2]$ is finite, and S is a compact subset of \mathbb{R}^d . From (1.1), it is easy to note that for any $s \in S$, $\mathbb{E}[YX(s)] = \mathbb{E}[\langle X, \beta^* \rangle X(s)] + \mathbb{E}[\epsilon X(s)]$ which implies $\mathbb{E}[XY](s) = (C\beta^*)(s)$, where $C := \mathbb{E}[X \otimes X]$ is the covariance operator. Therefore, the slope function β^* satisfies the operator equation $\mathbb{E}[XY] = C\beta^*$ which can be alternately expressed as

$$\beta^* := \arg\min_{\beta \in L^2(S)} \mathbb{E}[Y - \langle X, \beta \rangle_{L^2}]^2.$$
(1.2)

The goal is to construct an estimator $\hat{\beta}$ to estimate the unknown slope function β^* using observed empirical data $\{(X_1, Y_1), (X_2, Y_2), \ldots, (X_n, Y_n)\}$. Here, X_i 's are i.i.d. copies of the random process $X(\omega, \cdot)$, and Y_i 's are scalar responses. A general study by taking Y_i 's to be functional responses has been carried out in [35, 22, 4, 12].

One of the initial methodologies developed for estimating the slope function is functional principal component analysis (FPCA) [7, 24, 5, 20, 40], which relies on representing the estimator of β^* with respect to a basis, for example, a B-spline basis as in [8], or more popularly the eigenfunctions of the covariance operator. Under a smoothness condition on the slope function, [14] provided an optimal convergence rate for the FPCA method using covariance eigenfunctions. Recently, Holzleitner and Pereverzyev [16] explored the FPCA method under general regularization—the aforementioned works use spectral cutoff as the regularizer—, for the functional polynomial regression model with general source

Key words and phrases. Functional linear regression, reproducing kernel Hilbert space, regularization, minimax rates, integral operator, covariance operator.

condition. We refer the reader to [37, 15] and references therein for a detailed account of the FPCA approach.

Another approach for FLR that gained popularity is the method of regularization in a reproducing kernel Hilbert space (RKHS), i.e., an estimator of β^* is constructed by restricting it to an RKHS, \mathcal{H} (see Section 2 for details about this approach). The goal of this paper is to investigate this approach, obtain optimality results, and close the gaps in the literature, thereby obtaining a closure on this approach. Before discussing our contributions, we first establish the context by discussing the results in the literature.

The RKHS approach for FLR was first proposed and investigated by Yuan and Cai [38], wherein they developed an estimator of β^* based on Tikhonov regularization and established its optimality under the commutativity of the integral operator, T associated with the reproducing kernel and the covariance operator, C, assuming $\beta^* \in \mathcal{H}$. Under the same assumption of $\beta^* \in \mathcal{H}$, [6, 34] established optimal convergence rates when the integral and covariance operators do not commute. On the other hand, Zhang et al. [39] considered a different smoothness assumption of $\beta^* \in \mathcal{R}(T^{\frac{1}{2}}(T^{\frac{1}{2}}CT^{\frac{1}{2}})^s)$ and derived optimal convergence rates for the estimation and prediction errors associated with the estimator for $s \in (0,1]$ and $s \in (0,\frac{1}{2}]$, respectively, where $\mathcal{R}(A)$ denotes the range space of the operator A. Balasubramanian et al. [2] considered a non-linear extension to the FLR model, called the functional single-index model $(Y = g(\langle X, \beta^* \rangle_{L^2}) + \epsilon)$ where g is a non-linear function, and showed that if X is a Gaussian process with covariance operator C, then under appropriate regularity condition on g, the slope function β^* of the singleindex model satisfies the operator equation, $\mathbb{E}[XY] = C\beta^*$, which is also satisfied by the slope function of the FLR model. Therefore, they analyzed the functional single-index model by analyzing the FLR model, wherein they considered a least squares approach with Tikhonov regularization (similar to [38]) to obtain an estimator of β^* . Under the source condition of $\beta^* \in \mathcal{R}(T^{\alpha}) \subset L^2(S) \setminus \mathcal{H}, \alpha \in (0, \frac{1}{2})$ —note that this is weaker than that used in [38] which uses $\alpha = \frac{1}{2}$, i.e., $\beta^* \in \mathcal{R}(T^{\frac{1}{2}}) = \mathcal{H}$, [2] established convergence rates for estimation and prediction errors in the commutative setup, extending the results of [38]. [2] also obtained convergence rates for the estimation and prediction errors in the non-commutative setting under the assumption that $\beta^* \in \mathcal{R}(T^{\frac{1}{2}}(T^{\frac{1}{2}}CT^{\frac{1}{2}})^s), 0 \leq s \leq 1$, which partially extends the results of [6, 34, 39].

In the context of the above-mentioned results, there are many open questions related to the RKHS approach for FLR that need to be addressed: (i) the rates for a large range of smoothness, i.e., $\alpha > \frac{1}{2}$ (in the commutative setting) and s > 1 (in the noncommutative setting) is not known, (ii) the optimality of the rates is not known in the commutative setting except for $\alpha = \frac{1}{2}$, and in the non-commutative setting for s > 1, and (iii) the impact of qualification of a general regularization family on the convergence rate is not known since all the above-mentioned works employ Tikhonov regularization. The main contribution of this paper is to address these limitations and provide closure on the RKHS approach for FLR. To this end, we undertake a comprehensive examination of this approach by using a general regularization scheme (subsuming Tikhonov regularization) that caters to a wide range of smoothness and obtain optimal convergence rates for the estimation and prediction errors in the commutative and non-commutative settings. The details of these contributions are elaborated below.

1.1. **Contributions.** (i) We propose an estimator of β^* based on generalized regularization that subsumes Tikhonov regularization (which was already considered in the literature) so as to understand the impact of the source condition and the qualification (of the regularization) on the convergence rates.

(ii) Under the smoothness assumption, $\beta^* \in \mathcal{R}(T^{\alpha})$, $\alpha > 0$, in Theorems 3.1 and 3.4, we present the convergence rates for the estimation and prediction errors, respectively, in the commutative setting (i.e., T and C commute). These results match the optimal results of [38] when $\alpha = \frac{1}{2}$ and improve upon the rates of [2] for $0 < \alpha < \frac{1}{2}$ while providing convergence rates for $\alpha > \frac{1}{2}$ but up to a smoothness index determined by the qualification of the general regularization. If the qualification is infinity, then Theorems 3.1 and 3.4 provide convergence rates for all orders of smoothness, i.e., $\alpha > 0$. We would like to highlight that until this work, the convergence rates are known only for $0 < \alpha \leq \frac{1}{2}$ and the optimality of these rates are known only for $\alpha = \frac{1}{2}$. In this work, we not only resolve the issue of catering to $\alpha > \frac{1}{2}$ but also establish the optimality of the rates in the commutative setting for $\alpha > 0$ by deriving lower bounds in Theorem 4.3. A summary of our results in relation to those in the literature is captured in Table 1.

Results (Regularization)	Assumption	Estimation error $\ eta^* - \hat{eta}\ _{L^2}$	Prediction error $\ C^{\frac{1}{2}}(\beta^* - \hat{\beta})\ _{L^2}^2$
Cai and Yuan[38] (Tikhonov)	$\beta^* \in \mathcal{R}(T^{\frac{1}{2}})$	$\asymp n^{-\frac{t}{2(1+t+c)}}$	$\simeq n^{-\frac{t+c}{t+c+1}}$
Balasubramanian et al. [2] (Tikhonov)	$\beta^* \in \mathcal{R}(T^{\alpha})$ $(0 < \alpha \le \frac{1}{2})$	$\lesssim n^{-rac{lpha t}{1+c+2t(1-lpha)}}$	$\lesssim n^{-\frac{2\alpha t+c}{1+c+2t(1-\alpha)}}$
This paper Theorems 3.1, 3.4 (General)	$\beta^* \in \mathcal{R}(T^{\alpha})$	$\lesssim n^{-\frac{rt}{1+c+2rt}}$ $r = \alpha \wedge \left(\nu + \frac{c}{t}(\nu - \frac{1}{2})\right)$	$\lesssim n^{-\frac{2rt+c}{1+c+2rt}}$ $r = \alpha \wedge \left(\nu + \frac{1}{2} + \nu \frac{c}{t}\right)$
This paper Theorem 4.3	$\beta^* \in \mathcal{R}(T^{\alpha})$ $(0 < \alpha)$	$\gtrsim n^{-\frac{\alpha t}{1+c+2\alpha t}}$	$\gtrsim n^{-\frac{2\alpha t+c}{1+c+2\alpha t}}$

Table 1.	Conve	ergence rat	es for F	LR in the	e RKHS	frame	work whe	n the
kernel int	egral o	perator T	and the	e covarian	ce opera	ator C	commute.	Here
ν denotes	s the qu	alification	of the	regulariza	tion and	$d a \wedge b$	$:= \min\{a$	$,b\}.$

(iii) We provide convergence rates for the estimation and prediction errors in the noncommutative setting (i.e., without assuming the commutativity of T and C) in Theorems 3.3 and 3.5, respectively, under the source condition $\beta^* \in \mathcal{R}(T^{\frac{1}{2}}(T^{\frac{1}{2}}CT^{\frac{1}{2}})^s)$ where s > 0. These results generalize those of [2, 6, 39] by investigating rates for smoothness index, s > 1 since the prior results are known only for $0 \le s \le 1$ for the estimation error and $0 \le s \le \frac{1}{2}$ for the prediction error. Moreover, the optimality is known only for the prediction error for $0 \le s \le \frac{1}{2}$. We address this issue by showing the rates of Theorems 3.3 and 3.5 to be optimal by deriving lower bounds in Theorem 4.4. A summary of our contributions in the non-commutative setting in contrast to the existing results in the literature is provided in Table 2.

(*iv*) Since the slope function of a single-index model can be estimated using the FLR estimator if X is Gaussian (as shown in [2]), our results discussed in (i)-(iii) cater to the single-index model as well, thereby sharpening and generalizing the existing results in the literature (see Remark 3.6).

Results (Regularization)	Assumption	Estimation error $\ \beta^* - \hat{\beta}\ $	Prediction error $\ C^{\frac{1}{2}}(\beta^* - \hat{\beta})\ _{L^2}^2$	
Yuan and Cai [6] (Tikhonov)	$\beta^* \in \mathcal{R}(T^{\frac{1}{2}})$	N.A.	$\approx n^{-\frac{b}{b+1}}$	
Tong and Ng[34] (Tikhonov)	$\beta^* \in \mathcal{R}(T^{\frac{1}{2}})$	N.A.	$\lesssim n^{-rac{b}{b+1}}$ without Assumption 5	
Zhang et al.[39]	$\beta^* \in \mathcal{R}(T^{\frac{1}{2}}(T^{\frac{1}{2}}CT^{\frac{1}{2}})^s)$	$\asymp n^{-\frac{sb}{1+b+2sb}}$	$symp n^{-rac{(b+2sb)}{1+b+2sb}}$	
(Tikhonov)	$(0 < s \le 1)$	(RKHS norm)	$(0 \le s \le \frac{1}{2})$	
Balasubramanian et al.[2]	$\beta^* \in \mathcal{R}(T^{\frac{1}{2}}(T^{\frac{1}{2}}CT^{\frac{1}{2}})^s)$	$\lesssim n^{-rac{sb}{1+b+2sb}}$	$\lesssim n^{-\frac{b}{1+b}}; \beta^* \in \mathcal{R}(T^{\frac{1}{2}})$	
(Tikhonov)	$(0 < s \le 1)$	$(L^2 \text{ norm})$		
This paper Theorems 3.3, 3.5 (General)	$\beta^* \in \mathcal{R}(T^{\frac{1}{2}}(T^{\frac{1}{2}}CT^{\frac{1}{2}})^s)$ $(0 < s \le \nu)$	$ \begin{array}{l} \lesssim n^{-\frac{rb}{1+b+2rb}} \\ (\text{RKHS norm}) \\ r = s \wedge \nu \end{array} $	$ \lesssim n^{-\frac{(b+2rb)}{1+b+2rb}} r = s \wedge (\nu - \frac{1}{2}) $	
This paper Theorem 4.4	$\beta^* \in \mathcal{R}(T^{\frac{1}{2}}(T^{\frac{1}{2}}CT^{\frac{1}{2}})^s) \\ (s > 0)$	$\gtrsim n^{-rac{sb}{1+b+2sb}}$ (RKHS norm)	$\gtrsim n^{-rac{(b+2sb)}{1+b+2sb}}$	

Table 2. Convergence rates for FLR in the RKHS framework when the kernel integral operator T and the covariance operator C do not commute. Here ν denotes the qualification of the regularization and $a \wedge b := \min\{a, b\}$.

1.2. Organization. The paper is organized as follows. Section 2 presents the necessary background for the regularization method and the FLR model in the RKHS setting and provides some elementary results that are necessary for analysis. In Section 3, we discuss our assumptions and provide convergence rates for the estimation and prediction errors in both commutative and non-commutative settings for the general regularization method. We provide matching lower bounds for these errors in Section 4, which establish the

optimality of the general regularization method in the FLR setting. The proofs of all the results are presented in Section 5, while supplementary results needed to prove the main results are relegated to Appendix A.

1.3. Notations. $L^2(S)$ denotes the space of all real-valued square-integrable functions defined on S. For $f,g \in L^2(S)$, L^2 inner product and norm are defined as $\langle f,g \rangle_{L^2} = \int_S f(x)g(x) dx$ and $\|f\|_{L^2}^2 = \langle f, f \rangle_{L^2}$. The inner product and norm associated with the RKHS, \mathcal{H} are denoted as $\langle \cdot, \cdot \rangle_{\mathcal{H}}$ and $\|\cdot\|_{\mathcal{H}}$, respectively. For an operator A, $\mathcal{R}(A)$ denotes the range of operator A. For an operator $A : L^2 \to L^2$, the operator norm is defined as $\|A\|_{L^2 \to L^2} = \sup\{\|Af\|_{L^2}|f \in L^2, \|f\|_{L^2} = 1\}$. Similarly, for an operator $A : L^2 \to \mathcal{H}$, the operator norm is defined as $\|A\|_{L^2 \to \mathcal{H}} = \sup\{\|Af\|_{\mathcal{H}} : f \in L^2, \|f\|_{L^2} = 1\}$. For two positive numbers a and $b, a \lesssim b$ means $a \leq cb$ for some positive constant c. For positive sequences $(a_k)_k$ and $(b_k)_k, a_k \asymp b_k$ means $a_k \lesssim b_k \lesssim a_k$ for all k. For a random variable W with law P and a constant $b, W \lesssim_p b$ denotes that for any $\delta > 0$, there exists a positive constant $c_{\delta} < \infty$ such that $P(W \leq c_{\delta}b) \geq \delta$. For notational convenience, we define $\Lambda := T^{\frac{1}{2}}CT^{\frac{1}{2}}$ and $\hat{\Lambda}_n := T^{\frac{1}{2}}\hat{C}_nT^{\frac{1}{2}}$, where \hat{C}_n is an empirical estimator of C (see Section 2 for the definition of \hat{C}_n).

2. Preliminaries

A Hilbert space \mathcal{H} of real-valued functions on S is called an RKHS, if the pointwise evaluation functional is linear and continuous. There is a one-to-one correspondence between an RKHS \mathcal{H} and a reproducing kernel k which is a symmetric and positive definite function from $S \times S$ to \mathbb{R} such that $k(s, \cdot) \in \mathcal{H}$ and $f(s) = \langle k(s, \cdot), f \rangle_{\mathcal{H}}, \forall f \in \mathcal{H}$. We assume that k is measurable and $\sup_{x \in S} k(x, x) \leq \kappa$, where κ is a positive constant. Then the elements in the associated RKHS \mathcal{H} are measurable and bounded functions on S. It can be easily seen that \mathcal{H} is continuously embedded in $L^2(S)$ [11]. We refer the reader to [1, 32, 25, 26] for more details on RKHS.

We define the integral operator $T := JJ^* : L^2(S) \to L^2(S)$, where $J : \mathcal{H} \to L^2(S)$ is the inclusion operator defined as $(Jf)(x) = \langle k(x, \cdot), f \rangle_{\mathcal{H}}, \ \forall x \in S \text{ and } J^* : L^2(S) \to \mathcal{H}$ is the adjoint of the inclusion operator given as $J^*g = \int_S k(\cdot, t)g(t) dt$. It is easy to verify that $T^{\frac{1}{2}}$ is a partial isometry from $L^2(S)$ to \mathcal{H} , consequently $\mathcal{R}(T^{\frac{1}{2}}) = \mathcal{H}$. Moreover, for $\alpha > \frac{1}{2}$, $\mathcal{R}(T^{\alpha}) \subset \mathcal{H}$. In other words, coefficients of a representation of a function, $f \in \mathcal{R}(T^{\alpha})$ in terms of eigenfunctions of the operator T, will have a faster decay for larger value of α .

Motivated by the observation that β^* solves (1.2), using the given data $\{(X_i, Y_i)\}_{i=1}^n$, we consider the following estimator of β^* :

$$\hat{\beta} = \arg\min_{\beta \in \mathcal{H}} \frac{1}{n} \sum_{i=1}^{n} [Y_i - \langle \beta, X_i \rangle_{L^2}]^2 = \arg\min_{\beta \in \mathcal{H}} \frac{1}{n} \sum_{i=1}^{n} [Y_i - \langle J\beta, X_i \rangle_{L^2}]^2$$

$$= \arg\min_{\beta \in \mathcal{H}} \frac{1}{n} \sum_{i=1}^{n} [Y_i - \langle \beta, J^* X_i \rangle_{L^2}]^2.$$
(2.1)

Let us define

$$\hat{C}_n := \frac{1}{n} \sum_{i=1}^n X_i \otimes X_i$$
 and $\hat{R} := \frac{1}{n} \sum_{i=1}^n Y_i X_i$

It can be easily verified that the solution of the equation (2.1) can be given by solving the operator equation

$$J^* \hat{C}_n J \hat{\beta} = J^* \hat{R}. \tag{2.2}$$

Equation (2.2) can be seen as a discretized form of $J^*CJ\beta^* = J^*\mathbb{E}[XY]$, which is an illconditioned problem. Regularization techniques have been well explored in the learning theory framework to solve such ill-posed problems [3]. A detailed study on the Tikhonov regularization technique to solve (2.2) has been carried out in [6, 39, 2, 34, 33]. In this paper, we use the general regularization techniques from the theory of ill-posed inverse problems to solve (2.2), yielding the regularized estimator

$$\hat{\beta}_{\lambda} = g_{\lambda} (J^* \hat{C}_n J) J^* \hat{R}_{\lambda}$$

where $\lambda > 0$ is the regularization parameter. For simplicity, we remove the subscript λ and write $\hat{\beta}_{\lambda}$ as $\hat{\beta}$. Here $g_{\lambda} : [0, \eta] \to \mathbb{R}, 0 < \lambda \leq \eta$, is the regularization family satisfying the following conditions:

• There exists a constant A > 0 such that

$$\sup_{0 < \sigma \le \eta} |\sigma g_{\lambda}(\sigma)| \le A.$$
(2.3)

• There exists a constant B > 0 such that

$$\sup_{0 < \sigma \le \eta} |g_{\lambda}(\sigma)| \le \frac{B}{\lambda}.$$
(2.4)

• There exists a constant D > 0 such that

$$\sup_{0<\sigma\leq\eta} |1 - g_{\lambda}(\sigma)\sigma| = \sup_{0<\sigma\leq\eta} |r_{\lambda}(\sigma)| \leq D.$$
(2.5)

• The maximal *p* such that

$$\sup_{0<\sigma\leq\eta}|1-g_{\lambda}(\sigma)\sigma|\sigma^{p} = \sup_{0<\sigma\leq\eta}|r_{\lambda}(\sigma)|\sigma^{p}\leq\omega_{p}\lambda^{p},$$
(2.6)

is called the qualification of the regularization family g_{λ} , where the constant ω_p does not depend on λ .

Examples of regularization families include Tikhonov $(g_{\lambda}(\sigma) = (\sigma + \lambda)^{-1})$, spectral cut-off $(g_{\lambda}(\sigma) = \sigma^{-1} \mathbb{1}_{\sigma \geq \lambda})$, Showalter $(g_{\lambda}(\sigma) = \sigma^{-1}(1 - e^{-\sigma/\lambda})\mathbb{1}_{\sigma \neq 0} + \lambda^{-1}\mathbb{1}_{\sigma = 0})$, and Landweber iteration $(g_t(\sigma) = \sum_{i=1}^{t-1} (1 - \sigma)^i$ where λ is identified as t^{-1} , $t \in \mathbb{N}$), with qualification 1 for Tikhonov and ∞ for the rest, where $\mathbb{1}_{x \in \Omega} = 1$ if $x \in \Omega$, and 0, otherwise. We refer the reader to [21, 13] for more details about the regularization method.

3. MAIN RESULTS

In this section, we present the convergence rates of the general regularization method in the FLR model for both estimation and prediction errors. First, we begin with a list of assumptions required for the analysis.

Assumption 1. There exists a $h \in L^2(S)$ such that $\beta^* = T^{\alpha}h$, where $\alpha > 0$, i.e., $\beta^* \in \mathcal{R}(T^{\alpha})$.

Assumption 2. There exists a $h \in L^2(S)$ such that $\beta^* = T^{\frac{1}{2}} (T^{\frac{1}{2}} C T^{\frac{1}{2}})^s h$, where s > 0, *i.e.*, $\beta^* \in \mathcal{R}(T^{\frac{1}{2}} (T^{\frac{1}{2}} C T^{\frac{1}{2}})^s)$.

Assumptions 1 and 2, called the source condition, ensure the smoothness of β^* in commutative (*T* and *C* commute) and non-commutative (*T* and *C* do not commute) cases, respectively. Since $\mathcal{R}(T^{\frac{1}{2}}) = \mathcal{H}$ [11], it is clear that Assumption 2 is more stringent than Assumption 1. We would like to highlight that while both the commutative and non-commutative cases can be analyzed in a unified manner under Assumption 2, [2] demonstrated that sharper bounds can be obtained for the commutative case even under a weaker assumption (i.e., Assumption 1). Hence, we analyze our estimator under two separate settings of commutative and non-commutative, using the appropriate assumptions.

$$i^{-t} \lesssim \mu_i \lesssim i^{-t}$$
 and $i^{-c} \lesssim \xi_i \lesssim i^{-c}$ $\forall i \in \mathbb{N}$,

where $(\mu_i, \phi_i)_{i \in \mathbb{N}}, (\xi_i, \phi_i)_{i \in \mathbb{N}}$ are simple eigenvalue-eigenfunction pairs of operators T and C, respectively.

Assumption 4. For some b > 1,

$$i^{-b} \lesssim \tau_i \lesssim i^{-b} \quad \forall i \in \mathbb{N},$$

where $(\tau_i, e_i)_{i \in \mathbb{N}}$ is the eigenvalue-eigenfunction pair of operator $T^{\frac{1}{2}}CT^{\frac{1}{2}}$.

Assumptions 3 and 4 capture the decay rate of eigenvalues of operators, T, C and $T^{\frac{1}{2}}CT^{\frac{1}{2}}$, which are employed to obtain convergence rates in the commutative and non-commutative cases, respectively. Note that when T and C commute, then b = t + c.

Assumption 5. There exists a constant $d_1 > 0$ such that

$$\mathbb{E}[\langle X, f \rangle_{L^2}^4] \le d_1[\mathbb{E}\langle X, f \rangle_{L^2}^2]^2, \ \forall f \in L^2(S).$$

In Assumption 5, it is posited that the fourth moment is constrained not to exceed the square of the second moment. An instance of data that adheres to this inequality is the one that follows a Gaussian distribution.

3.1. Estimation error. We provide error bounds for estimating the slope function, β^* under commutative and non-commutative settings in Theorems 3.1 (proved in Sections 5.1 and 5.2) and 3.3 (proved in Section 5.3) under the source conditions mentioned in Assumptions 1 and 2, respectively.

Theorem 3.1 (Estimation error–Commutative). Suppose T and C commute, and Assumptions 1, 3 and 5 hold. Let ν be the qualification of the regularization family and $\nu \geq \frac{1}{2}$. Then, for $\lambda = n^{-\frac{t+c}{1+c+2tr}}$, we have

 $\begin{array}{l} (a) \ \|\beta^* - \hat{\beta}\|_{L^2} \lesssim_p n^{-\frac{rt}{1+c+2tr}}, \quad where \quad r = \min\{\alpha, \nu + \frac{c}{t}(\nu - \frac{1}{2})\};\\ (b) \ \|\beta^* - \hat{\beta}\|_{\mathcal{H}} \lesssim_p n^{-\frac{t(r-\frac{1}{2})}{1+c+2tr}}, \quad where \quad r = \min\{\alpha, \nu + \frac{c}{t}\nu + \frac{1}{2}\} \text{ for } \alpha \ge \frac{1}{2}. \end{array}$

Remark 3.2. (i) The assumption that $\beta^* \in \mathcal{R}(T^{\alpha})$ enforces a smoothness condition that influences the properties of the functional space to which the slope function belongs. For $\alpha \geq \frac{1}{2}$, $\mathcal{R}(T^{\alpha}) \subset \mathcal{H}$, while for $0 < \alpha < \frac{1}{2}$, $\mathcal{R}(T^{\alpha}) \subset L^2(S) \setminus \mathcal{H}$.

(ii) The results on L^2 estimation error in the literature mostly deal with $\alpha = \frac{1}{2}$ [38] and $0 < \alpha \leq \frac{1}{2}$ [2]. When $\alpha = \frac{1}{2}$, our L^2 -error rates align with the minimax rate established by [38] (also see Theorem 4.3), while for $0 < \alpha < \frac{1}{2}$, our rates improve those provided in [2] from $n^{-\frac{\alpha t}{1+c+2t(1-\alpha)}}$ to $n^{-\frac{\alpha t}{1+c+2t\alpha}}$. In fact, in Theorem 4.3, we show this improved rate to be minimax optimal. Moreover, the results presented in Theorem 3.1 cater to smoothness index $\alpha > \frac{1}{2}$, which means faster convergence rates are obtained for smoother slope function but with the rate saturating at a level that depends on the qualification of the regularization method.

(iii) As can be seen from Theorem 3.1, we can allow α to go beyond the qualification ν , and still the algorithm can achieve faster convergence rates. In particular, while using Tikhonov regularization for the commutative setting, we can get faster convergence rates for our algorithm even if $\alpha > 1$. This happens since the regularization is applied to the operator Λ while the source condition on β^* is enforced through the operator T.

The following result (proved in Section 5.2) presents the estimation error in the RKHS norm when T and C do not commute.

Theorem 3.3 (Estimation error–Non-commutative). Suppose Assumptions 2, 4 and 5 hold. Let ν be the qualification of the regularization family and $\nu \geq \frac{1}{2}$. Then, for $\lambda = n^{-\frac{b}{1+b+2rb}}$, we have

$$\|\beta^* - \hat{\beta}\|_{\mathcal{H}} \lesssim_p n^{-\frac{or}{1+b+2rb}}, \quad where \quad r = \min\{s, \nu\}.$$

The rate in Theorem 3.3 not only matches with the existing rates of [2, 39] when $0 < s \leq 1$, but also addresses the case of smoothness index s beyond 1 (but with the rate saturating at $s = \nu$, the qualification of the regularization family). Moreover, in Theorem 4.4, we show the bound of Theorem 3.3 to be minimax optimal.

3.2. **Prediction error.** We present the prediction error bounds in commutative and non-commutative settings in Theorems 3.4 (proved in Section 5.4) and 3.5 (proved in Section 5.5), respectively.

Theorem 3.4 (Prediction error–Commutative). Suppose T and C commute, and Assumptions 1, 3 and 5 hold. Let ν be the qualification of the regularization family and $\nu \geq \frac{1}{2}$. Then, for $\lambda = n^{-\frac{t+c}{1+c+2tr}}$, we have

$$\|C^{\frac{1}{2}}(\beta^* - \hat{\beta})\|_{L^2}^2 \lesssim_p n^{-\frac{2rt+c}{1+c+2tr}}, \quad where \quad r = \min\left\{\alpha, \nu + \frac{1}{2} + \nu \frac{c}{t}\right\}.$$

The above result matches the results of [2, 38], wherein [38] considers only $\alpha = \frac{1}{2}$, while [2] extended the result of [38] under Assumption 1 for $0 < \alpha \leq \frac{1}{2}$. Moreover, Theorem 3.4 obtains a rate for α beyond the above-mentioned ranges, and we show this rate to be minimax optimal in Theorem 4.3.

The following result, which considers the prediction error when T and C do not commute, is an extension of the results of [2, 6, 34, 39] in terms of the smoothness condition. [2, 6, 34] have considered the prediction error for non-commutative case with $\beta^* \in \mathcal{H}$. In [39], authors have extended these results with Assumption 3 but for a restricted range of $0 < s \leq \frac{1}{2}$. We extend all these results for all positive ranges of s in Assumption 3.

Theorem 3.5 (Prediction error–Non-commutative). Suppose Assumptions 2, 4 and 5 hold. Let ν be the qualification of the regularization family and $\nu \geq \frac{1}{2}$. Then, for $\lambda = n^{-\frac{b}{1+b+2rb}}$, we have

$$\|C^{\frac{1}{2}}(\beta^* - \hat{\beta})\|_{L^2}^2 \lesssim_p n^{-\frac{b(2r+1)}{1+b+2rb}}, \quad where \quad r = \min\left\{s, \nu - \frac{1}{2}\right\}.$$

Remark 3.6. For an index function $g : \mathbb{R} \to \mathbb{R}$, the single index model is defined as

$$Y = g\left(\int_{S} X(t)\beta^{*}(t)dt\right) + \epsilon.$$

A detailed study of estimating β^* in a single-index model has been carried out in [24, 17, 9, 31, 18]. In [2], the authors showed that if X is sampled from a Gaussian process, then under some regularity conditions on g, β^* in the single-index model satisfies the operator equation of the FLR model, i.e., $\mathbb{E}[XY] = C\beta^*$. This means the results obtained in this paper will address the question of estimating β^* in a single-index model if X is sampled from a Gaussian process, thereby sharpening and generalizing the results of [2].

4. Lower Bounds

In this section, we establish lower bounds for each of the cases discussed in the previous section and show them to match the upper bounds, thereby confirming the optimality of the proposed regularized estimator of β^* . The idea of the proof is inspired by the work of Cai and Yuan [6]. To derive the lower bounds, we need to discuss the divergence between two probability measures P_1 and P_2 defined on a measurable space $(\mathcal{X}, \mathcal{A})$. In our analysis, we use **Kullback-Leibler divergence** which is defined as :

$$\mathcal{K}(P_1, P_2) := \begin{cases} \int \log\left(\frac{dP_1}{dP_2}\right) dP_1 & \text{if } P_1 \ll P_2; \\ \mathcal{X} \\ +\infty & \text{otherwise,} \end{cases}$$

where $P_1 \ll P_2$ denotes P_1 is absolutely continuous with respect to P_2 .

The key concept to derive the lower bounds follows from [36, Chapter 2], where the major task is to find N + 1 elements $\{\theta'_0, \ldots, \theta'_N\}$ from a hypothesis space such that the distance between any two of them is at least 2r. Furthermore, associated with each θ'_j , we have a distribution $P_{\theta'_j}$, and the mean of the Kullback-Leibler divergence between distributions $P_{\theta'_j}$ and $P_{\theta'_0}$ should grow at most logarithmically. Consequently, we get that for large N, with high probability, the distance between θ'_j and any estimator $\hat{\theta}$ will be at least r. For easy reference, we state the key theorem [36, Theorem 2.5] below.

Let Θ' be a non-empty set. Suppose $d : \Theta' \times \Theta' \to [0, \infty)$ be a semi-distance function and $\mathcal{P} = \{P_{\theta} : \theta \in \Theta'\}$ be the set of probability measures on $(\mathcal{X}, \mathcal{A})$ indexed by Θ' .

Theorem 4.1. [36, Theorem 2.5] Assume that N > 2 and suppose that the hypothesis space Θ' contains elements $\theta'_0, \theta'_1, \ldots, \theta'_N$ such that:

(1) $d(\theta'_{j}, \theta'_{k}) \ge 2r > 0 \quad \forall \ 0 \le j < k \le N;$ (2) $P_{\theta'_{j}} \ll P_{\theta'_{0}} \quad \forall j = 1, 2, \dots, N, and$

$$\frac{1}{N}\sum_{j=1}^{N}\mathcal{K}(P_{\theta'_{j}}, P_{\theta'_{0}}) \le u\log N$$

for some $0 < u < \frac{1}{8}$. Then

$$\inf_{\hat{\theta}} \sup_{\theta \in \Theta'} P_{\theta}(d(\theta, \hat{\theta}) \ge r) \ge \frac{\sqrt{N}}{1 + \sqrt{N}} \left(1 - 2u - \sqrt{\frac{2u}{\log N}} \right),$$

where the infimum is taken over all estimators $\hat{\theta}$ of $\theta \in \Theta'$.

The next lemma, which ensures the existence of N+1 elements in $\{0, 1\}^M$ with Hamming distance at least $\frac{M}{8}$, will be helpful to construct N+1 elements in a hypothesis space with the desired distance.

Lemma 4.2 (Varshamov-Gilbert bound [36]). Let $M \ge 8$. Then there exists a subset $\Theta = \{\theta^{(0)}, \ldots, \theta^{(N)}\} \subset \{0, 1\}^M$ such that $\theta^{(0)} = (0, \cdots, 0)$,

$$H(heta, heta^{'}) > rac{M}{8}, \quad \forall \ heta
eq heta^{'} \in \Theta,$$

where $H(\theta, \theta') = \sum_{i=1}^{M} (\theta_i - \theta'_i)^2$ is the Hamming distance and $N \ge 2^{\frac{M}{8}}$.

The following result (proved in Section 5.6) provides lower bounds associated with Theorems 3.1 and 3.4, i.e., estimation and prediction errors, respectively in the commutative setting.

Theorem 4.3 (Lower bound–Commutative). Suppose Assumptions 1 and 3 hold. Then for $\alpha \geq \frac{1}{2}$, we have

$$\lim_{a \to 0} \lim_{n \to \infty} \inf_{\hat{\beta}} \sup_{\beta^* \in \mathcal{R}(T^{\alpha})} \mathbb{P}\left\{ \|\hat{\beta} - \beta^*\|_{\mathcal{H}} \ge an^{-\frac{t(\alpha - \frac{1}{2})}{1 + c + 2\alpha t}} \right\} = 1.$$

Further for $\alpha > 0$, we have

$$\lim_{a \to 0} \lim_{n \to \infty} \inf_{\hat{\beta}} \sup_{\beta^* \in \mathcal{R}(T^{\alpha})} \mathbb{P}\left\{ \|\hat{\beta} - \beta^*\|_{L^2} \ge an^{-\frac{\alpha t}{1 + c + 2\alpha t}} \right\} = 1$$

and

$$\lim_{a \to 0} \lim_{n \to \infty} \inf_{\hat{\beta}} \sup_{\beta^* \in \mathcal{R}(T^{\alpha})} \mathbb{P}\left\{ \mathbb{E} \langle \hat{\beta} - \beta^*, X \rangle_{L^2}^2 \ge a n^{-\frac{2\alpha t + c}{1 + c + 2\alpha t}} \right\} = 1,$$

where the infimum is taken over all possible estimators.

The following result (proved in Section 5.7) provides lower bounds on the estimation and prediction errors in the non-commutative setting. Combined with Theorems 3.3 and 3.5, Theorem 4.4 ensures that the convergence rates achieved in the non-commutative setting are optimal.

Theorem 4.4 (Lower bound–Non-commutative). Suppose Assumptions 2 and 4 hold. Then for any s > 0, we have

$$\lim_{a \to 0} \lim_{n \to \infty} \inf_{\hat{\beta}} \sup_{\beta^* \in \mathcal{R}(T^{\frac{1}{2}}\Lambda^s)} \mathbb{P}\left\{ \|\hat{\beta} - \beta^*\|_{\mathcal{H}} \ge an^{-\frac{sb}{1+b+2sb}} \right\} = 1$$

and

$$\lim_{a \to 0} \lim_{n \to \infty} \inf_{\hat{\beta}} \sup_{\beta^* \in \mathcal{R}(T^{\frac{1}{2}} \Lambda^s)} \mathbb{P}\left\{ \mathbb{E} \langle \hat{\beta} - \beta^*, X \rangle_{L^2}^2 \ge a n^{-\frac{b(2s+1)}{1+b+2sb}} \right\} = 1,$$

where the infimum is taken over all possible estimators.

5. Proofs

The proofs of the main results of Sections 3 and 4 are provided in this section.

5.1. **Proof of part (a) in Theorem 3.1.** The proof involves decomposing $\|\hat{\beta} - \beta^*\|_{L^2}$ into several terms and bounding each of the terms separately. We start by considering the error term

$$\begin{split} \|\hat{\beta} - \beta^*\|_{L^2} &= \|J\hat{\beta} - \beta^*\|_{L^2} = \|Jg_{\lambda}(J^*\hat{C}_n J)J^*\hat{R} - \beta^*\|_{L^2} \\ &= \|T^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_n)T^{\frac{1}{2}}\hat{R} - \beta^*\|_{L^2} \quad \text{(by Lemma A.1)} \\ &\leq \underbrace{\|T^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_n)(T^{\frac{1}{2}}\hat{R} - T^{\frac{1}{2}}\hat{C}_n\beta^*)\|_{L^2}}_{Term-1} + \underbrace{\|T^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_n)T^{\frac{1}{2}}\hat{C}_n\beta^* - \beta^*\|_{L^2}}_{Term-2}. \end{split}$$

Bounding Term-1:

$$\begin{split} \|T^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_{n})(T^{\frac{1}{2}}\hat{R}-T^{\frac{1}{2}}\hat{C}_{n}\beta^{*})\|_{L^{2}} \\ \leq \|T^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_{n})(\hat{\Lambda}_{n}+\lambda I)^{\frac{1}{2}}\|_{L^{2}\to L^{2}}\|(\hat{\Lambda}_{n}+\lambda I)^{-\frac{1}{2}}(\Lambda+\lambda I)^{\frac{1}{2}}\|_{L^{2}\to L^{2}} \\ \times \|(\Lambda+\lambda I)^{-\frac{1}{2}}(T^{\frac{1}{2}}\hat{R}-T^{\frac{1}{2}}\hat{C}_{n}\beta^{*})\|_{L^{2}} \\ \lesssim \|T^{\frac{1}{2}}(\Lambda+\lambda I)^{-\frac{1}{2}}\|_{L^{2}\to L^{2}}\|(\Lambda+\lambda I)^{\frac{1}{2}}(\hat{\Lambda}_{n}+\lambda I)^{-\frac{1}{2}}\|_{L^{2}\to L^{2}} \\ \times \|(\hat{\Lambda}_{n}+\lambda I)^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_{n})(\hat{\Lambda}_{n}+\lambda I)^{\frac{1}{2}}\|_{L^{2}\to L^{2}}\|(\Lambda+\lambda I)^{-\frac{1}{2}}(T^{\frac{1}{2}}\hat{R}-T^{\frac{1}{2}}\hat{C}_{n}\beta^{*})\|_{L^{2}} \\ \overset{(*)}{\lesssim}_{p}\|T^{\frac{1}{2}}(\Lambda+\lambda I)^{-\frac{1}{2}}\|_{L^{2}\to L^{2}}\|(\Lambda+\lambda I)^{-\frac{1}{2}}(T^{\frac{1}{2}}\hat{R}-T^{\frac{1}{2}}\hat{C}_{n}\beta^{*})\|_{L^{2}}, \end{split}$$

where we used Lemma A.6 and the regularization properties (2.3), (2.4) in (*). Next, we apply Lemma A.2 to conclude that

$$\begin{split} \|T^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_{n})(T^{\frac{1}{2}}\hat{R}-T^{\frac{1}{2}}\hat{C}_{n}\beta^{*})\|_{L^{2}} \lesssim_{p} \sqrt{\frac{\sigma^{2}\mathcal{N}(\lambda)}{n}}\|T^{\frac{1}{2}}(\Lambda+\lambda I)^{-\frac{1}{2}}\|_{L^{2}\to L^{2}}\\ \leq \lambda^{-\frac{c}{2(t+c)}}\sqrt{\frac{\sigma^{2}\mathcal{N}(\lambda)}{n}}, \end{split}$$

where the last inequality follows from the fact that

$$\|T^{\frac{1}{2}}(\Lambda + \lambda I)^{-\frac{1}{2}}\|_{L^{2} \to L^{2}} = \left(\sup_{i} \left|\frac{\mu_{i}}{\mu_{i}\xi_{i} + \lambda}\right|\right)^{\frac{1}{2}} \quad (T \text{ and } C \text{ commute})$$

$$\lesssim \left(\sup_{i} \left|\frac{i^{-t}}{i^{-(t+c)} + \lambda}\right|\right)^{\frac{1}{2}} \quad (\text{by Assumption 3})$$

$$\lesssim \lambda^{\frac{t-(t+c)}{2(t+c)}} = \lambda^{-\frac{c}{2(t+c)}} \quad (\text{by Lemma A.9}).$$
(5.1)

Bounding Term-2: As the bound on this term will depend on the source condition, the estimation has been considered under two cases over the range of α . Case-1: Suppose $\alpha \in (0, \frac{1}{2}]$. Consider

$$\begin{split} \|T^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_{n})T^{\frac{1}{2}}\hat{C}_{n}\beta^{*} - \beta^{*}\|_{L^{2}} \\ \leq \|T^{\frac{1}{2}}(g_{\lambda}(\hat{\Lambda}_{n}) - (\hat{\Lambda}_{n} + \lambda I)^{-1})T^{\frac{1}{2}}\hat{C}_{n}\beta^{*}\|_{L^{2}} + \|T^{\frac{1}{2}}(\hat{\Lambda}_{n} + \lambda I)^{-1}T^{\frac{1}{2}}\hat{C}_{n}\beta^{*} - \beta^{*}\|_{L^{2}} \\ \leq \underbrace{\|T^{\frac{1}{2}}((\hat{\Lambda}_{n} + \lambda I)^{-1}T^{\frac{1}{2}}\hat{C}_{n}\beta^{*} - (\Lambda + \lambda I)^{-1}T^{\frac{1}{2}}C\beta^{*})\|_{L^{2}}}_{Term-2a} \\ + \underbrace{\|T^{\frac{1}{2}}(g_{\lambda}(\hat{\Lambda}_{n}) - (\hat{\Lambda}_{n} + \lambda I)^{-1})T^{\frac{1}{2}}\hat{C}_{n}\beta^{*}\|_{L^{2}}}_{Term-2b} + \underbrace{\|T^{\frac{1}{2}}(\Lambda + \lambda I)^{-1}T^{\frac{1}{2}}C\beta^{*} - \beta^{*}\|_{L^{2}}}_{Term-2c} \end{split}$$

Bounding Term-2a: Define $\tilde{\beta} := (\Lambda + \lambda I)^{-1} T^{\frac{1}{2}} C \beta^*$ and $\Upsilon := \frac{t(1-2\alpha)}{2(t+c)}$. Then

$$\begin{split} \|T^{\frac{1}{2}}((\hat{\Lambda}_{n}+\lambda I)^{-1}T^{\frac{1}{2}}\hat{C}_{n}\beta^{*}-(\Lambda+\lambda I)^{-1}T^{\frac{1}{2}}C\beta^{*})\|_{L^{2}}\\ =\|T^{\frac{1}{2}}(\hat{\Lambda}_{n}+\lambda I)^{-1}(T^{\frac{1}{2}}\hat{C}_{n}\beta^{*}-\hat{\Lambda}_{n}\tilde{\beta}-\lambda\tilde{\beta})\|_{L^{2}}\\ =\|T^{\frac{1}{2}}(\hat{\Lambda}_{n}+\lambda I)^{-1}(T^{\frac{1}{2}}(C-\hat{C}_{n})(T^{\frac{1}{2}}\tilde{\beta}-\beta^{*}))\|_{L^{2}} \quad (\text{since }\lambda\tilde{\beta}=T^{\frac{1}{2}}C\beta^{*}-\Lambda\tilde{\beta})\\ \leq \|T^{\frac{1}{2}}(\hat{\Lambda}_{n}+\lambda I)^{-\frac{1}{2}}\|_{L^{2}\to L^{2}}\|(\hat{\Lambda}_{n}+\lambda I)^{-\frac{1}{2}}T^{\frac{1}{2}}(C-\hat{C}_{n})(T^{\frac{1}{2}}\tilde{\beta}-\beta^{*})\|_{L^{2}}. \end{split}$$

From equation (5.1) and the observation that $T^{\frac{1}{2}}\tilde{\beta} - \beta^* = -\lambda(\Lambda + \lambda I)^{-1}\beta^*$, we have

$$\begin{aligned} \|T^{\frac{1}{2}}((\hat{\Lambda}_{n}+\lambda I)^{-1}T^{\frac{1}{2}}\hat{C}_{n}\beta^{*}-(\Lambda+\lambda I)^{-1}T^{\frac{1}{2}}C\beta^{*})\|_{L^{2}} \\ \lesssim \lambda\lambda^{-\frac{c}{2(t+c)}}\|(\hat{\Lambda}_{n}+\lambda I)^{-\frac{1}{2}}T^{\frac{1}{2}}(C-\hat{C}_{n})(\Lambda+\lambda I)^{-1}\beta^{*}\|_{L^{2}} \\ \lesssim \sqrt{\lambda}\lambda^{-\frac{c}{2(t+c)}}\|(\Lambda+\lambda I)^{-\frac{1}{2}}T^{\frac{1}{2}}(C-\hat{C}_{n})(\Lambda+\lambda I)^{-\frac{1}{2}}T^{\alpha}\|_{L^{2}\to L^{2}}\|h\|_{L^{2}} \\ \lesssim \sqrt{\lambda}\lambda^{-\frac{c}{2(t+c)}}\|T^{\frac{1}{2}-\alpha}(\Lambda+\lambda I)^{-\Upsilon}T^{\alpha}(\Lambda+\lambda I)^{-(\frac{1}{2}-\Upsilon)}(C-\hat{C}_{n})(\Lambda+\lambda I)^{-\frac{1}{2}}T^{\alpha}\|_{L^{2}\to L^{2}} \\ \leq \sqrt{\lambda}\lambda^{-\frac{c}{2(t+c)}}\lambda^{-\Upsilon}\|T^{\alpha}(\Lambda+\lambda I)^{-(\frac{1}{2}-\Upsilon)}(C-\hat{C}_{n})(\Lambda+\lambda I)^{-(\frac{1}{2}-\Upsilon)}T^{\alpha}\|_{L^{2}\to L^{2}}. \end{aligned}$$
(5.2)

By using $p = (\frac{1}{2} - \Upsilon)$ in Lemma A.4, we get

$$\|T^{\frac{1}{2}}(\hat{\Lambda}_n + \lambda I)^{-1}T^{\frac{1}{2}}\hat{C}_n\beta^* - T^{\frac{1}{2}}(\Lambda + \lambda I)^{-1}T^{\frac{1}{2}}C\beta^*\|_{L^2} \lesssim_p \frac{\lambda^{-\frac{1-2\alpha t}{2(t+c)}}}{\sqrt{n}}.$$

Bound of Term-2b: Consider

$$\begin{split} \|T^{\frac{1}{2}}(g_{\lambda}(\hat{\Lambda}_{n}) - (\hat{\Lambda}_{n} + \lambda I)^{-1})T^{\frac{1}{2}}\hat{C}_{n}\beta^{*}\|_{L^{2}} \\ &= \|T^{\frac{1}{2}}(\hat{\Lambda}_{n} + \lambda I)^{-1}((\hat{\Lambda}_{n} + \lambda I)g_{\lambda}(\hat{\Lambda}_{n}) - I)T^{\frac{1}{2}}\hat{C}_{n}\beta^{*}\|_{L^{2}} \\ &\leq \|T^{\frac{1}{2}}(\hat{\Lambda}_{n} + \lambda I)^{-1}((\hat{\Lambda}_{n}g_{\lambda}(\hat{\Lambda}_{n}) - I) + \lambda g_{\lambda}(\hat{\Lambda}_{n}))T^{\frac{1}{2}}\hat{C}_{n}\beta^{*}\|_{L^{2}} \\ &\leq \|T^{\frac{1}{2}}(\Lambda + \lambda I)^{-\frac{1}{2}}\|_{L^{2} \to L^{2}}\|(\Lambda + \lambda I)^{\frac{1}{2}}(\hat{\Lambda}_{n} + \lambda I)^{-\frac{1}{2}}\|_{L^{2} \to L^{2}}\|(\hat{\Lambda}_{n} + \lambda I)^{-\frac{1}{2}}\|_{L^{2} \to L^{2}}\|(\hat{\Lambda}_{n} + \lambda I)^{-\frac{3}{2}}T^{\frac{1}{2}}\hat{C}_{n}\beta^{*}\|_{L^{2}} \\ &\times \|(\hat{\Lambda}_{n} + \lambda I)^{-\frac{1}{2}}(r_{\lambda}(\hat{\Lambda}_{n}) + \lambda g_{\lambda}(\hat{\Lambda}_{n}))(\hat{\Lambda}_{n} + \lambda I)^{\frac{3}{2}}\|_{L^{2} \to L^{2}}. \end{split}$$
(5.3)

Using equation (5.1), Lemma A.6 and the regularization properties (2.3), (2.4), (2.5) and (2.6) in (5.3) yields

$$\begin{aligned} \|T^{\frac{1}{2}}(g_{\lambda}(\hat{\Lambda}_{n}) - (\hat{\Lambda}_{n} + \lambda I)^{-1})T^{\frac{1}{2}}\hat{C}_{n}\beta^{*}\|_{L^{2}} \\ \lesssim_{p}\lambda\lambda^{-\frac{c}{2(t+c)}}\|(\hat{\Lambda}_{n} + \lambda I)^{-\frac{3}{2}}T^{\frac{1}{2}}\hat{C}_{n}\beta^{*}\|_{L^{2}} \\ \leq \lambda\lambda^{-\frac{c}{2(t+c)}}\|(\hat{\Lambda}_{n} + \lambda I)^{-\frac{1}{2}}(\Lambda + \lambda I)^{\frac{1}{2}}\|_{L^{2} \to L^{2}}(\|(\Lambda + \lambda I)^{-\frac{3}{2}}T^{\frac{1}{2}}C\beta^{*}\|_{L^{2}} \\ &+\|(\Lambda + \lambda I)^{-\frac{1}{2}}((\hat{\Lambda}_{n} + \lambda I)^{-1}T^{\frac{1}{2}}\hat{C}_{n}\beta^{*} - (\Lambda + \lambda I)^{-1}T^{\frac{1}{2}}C\beta^{*})\|_{L^{2}}). \end{aligned}$$

$$(5.4)$$

Similar to (5.1), we can see that

$$\begin{split} \|(\Lambda + \lambda I)^{-\frac{3}{2}} T^{\frac{1}{2}} C \beta^* \|_{L^2} &\leq \|(\Lambda + \lambda I)^{-\frac{3}{2}} T^{\frac{1}{2}} C T^{\alpha} \|_{L^2 \to L^2} \|h\|_{L^2} \\ &\lesssim \left(\sup_i \left| \frac{i^{-\frac{2}{3}(t(\alpha + \frac{1}{2}) + c)}}{i^{-(t+c)} + \lambda} \right| \right)^{\frac{3}{2}} \lesssim \lambda^{\frac{2\alpha t - 2t - c}{2(t+c)}}, \end{split}$$

thereby yielding

$$\|T^{\frac{1}{2}}(g_{\lambda}(\hat{\Lambda}_{n}) - (\hat{\Lambda}_{n} + \lambda I)^{-1})T^{\frac{1}{2}}\hat{C}_{n}\beta^{*}\|_{L^{2}}$$

$$\lesssim_{p}\sqrt{\lambda}\lambda^{-\frac{c}{2(t+c)}}\|(\hat{\Lambda}_{n} + \lambda I)^{-1}T^{\frac{1}{2}}\hat{C}_{n}\beta^{*} - (\Lambda + \lambda I)^{-1}T^{\frac{1}{2}}C\beta^{*}\|_{L^{2}} + \lambda^{\frac{ct}{t+c}}.$$
(5.5)

Using similar ideas from (5.2) to bound the norm term in the r.h.s. of (5.5), we get

$$\|T^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_{n})T^{\frac{1}{2}}\hat{C}_{n}\beta^{*} - T^{\frac{1}{2}}(\hat{\Lambda}_{n} + \lambda I)^{-1}T^{\frac{1}{2}}\hat{C}_{n}\beta^{*}\|_{L^{2}} \lesssim_{p} \frac{\lambda^{-\frac{1-2\alpha t}{2(t+c)}}}{\sqrt{n}} + \lambda^{\frac{\alpha t}{t+c}}.$$

Bounding Term-2c: Consider

$$\begin{split} \|T^{\frac{1}{2}}(\Lambda+\lambda I)^{-1}T^{\frac{1}{2}}C\beta^*-\beta^*\|_{L^2} &= \|T^{\frac{1}{2}}(\Lambda+\lambda I)^{-1}T^{\frac{1}{2}}CT^{\alpha}h-T^{\alpha}h\|_{L^2} \\ &= \left[\sum_i (\frac{\mu_i^{1+\alpha}\xi_i}{\mu_i\xi_i+\lambda}-\mu_i^{\alpha})^2\langle\phi_i,h\rangle^2\right]^{\frac{1}{2}} = \left[\sum_i (\frac{\lambda\mu_i^{\alpha}}{\mu_i\xi_i+\lambda})^2\langle\phi_i,h\rangle^2\right]^{\frac{1}{2}} \\ &\leq \lambda \sup_i \left[\frac{\mu_i^{\alpha}}{\mu_i\xi_i+\lambda}\right] \|h\|_{L^2} \lesssim \lambda \sup_i \left[\frac{i^{-\alpha t}}{i^{-(t+c)}+\lambda}\right] \|h\|_{L^2} \lesssim \lambda^{\frac{\alpha t}{t+c}}. \end{split}$$

Case-2 : Suppose $\alpha \geq \frac{1}{2}$. Let ν be the qualification of the regularization family and ν' be the greatest integer less than or equal to ν . Consider

$$\begin{split} \|T^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_{n})T^{\frac{1}{2}}\hat{C}_{n}\beta^{*} - \beta^{*}\|_{L^{2}} &= \|T^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_{n})T^{\frac{1}{2}}\hat{C}_{n}T^{\frac{1}{2}}T^{\alpha-\frac{1}{2}}h - T^{\frac{1}{2}}T^{\alpha-\frac{1}{2}}h\|_{L^{2}} \\ &= \|T^{\frac{1}{2}}r_{\lambda}(\hat{\Lambda}_{n})(\hat{\Lambda}_{n} + \lambda I)^{\nu}(\hat{\Lambda}_{n} + \lambda I)^{-(\nu-\nu')}(\Lambda + \lambda I)^{\nu-\nu'} \\ &\times (\Lambda + \lambda I)^{-(\nu-\nu')}(\hat{\Lambda}_{n} + \lambda I)^{-\nu'}T^{\alpha-\frac{1}{2}}h\|_{L^{2}} \\ &\leq \|T^{\frac{1}{2}}r_{\lambda}(\hat{\Lambda}_{n})(\hat{\Lambda}_{n} + \lambda I)^{\nu}(\hat{\Lambda}_{n} + \lambda I)^{-(\nu-\nu')}(\Lambda + \lambda I)^{\nu-\nu'} \\ &\times ((\Lambda + \lambda I)^{-(\nu-\nu')}(\hat{\Lambda}_{n} + \lambda I)^{-(\nu'-\nu')}(\Lambda + \lambda I)^{-\nu}T^{\alpha-\frac{1}{2}}h + (\Lambda + \lambda I)^{-\nu}T^{\alpha-\frac{1}{2}}h)\|_{L^{2}} \\ \lesssim_{p} \underbrace{\lambda^{\nu}\|(\Lambda + \lambda I)^{-(\nu-\nu')}(\hat{\Lambda}_{n} + \lambda I)^{-\nu'}T^{\alpha-\frac{1}{2}}h - (\Lambda + \lambda I)^{-\nu}T^{\alpha-\frac{1}{2}}h\|_{L^{2}}}_{Term-2d} \\ &+ \underbrace{\|T^{\frac{1}{2}}r_{\lambda}(\hat{\Lambda}_{n})(\hat{\Lambda}_{n} + \lambda I)^{\nu}(\hat{\Lambda}_{n} + \lambda I)^{-(\nu-\nu')}(\Lambda + \lambda I)^{\nu-\nu'}(\Lambda + \lambda I)^{-\nu}T^{\alpha-\frac{1}{2}}h\|_{L^{2}}}_{Term-2e}, \end{split}$$

where the last inequality is obtained by using Lemma A.6 and the qualification property (2.6) of the regularization family.

Bounding Term-2d: Note that

$$\lambda^{\nu} \| (\Lambda + \lambda I)^{-(\nu - \nu')} (\hat{\Lambda}_n + \lambda I)^{-\nu'} T^{\alpha - \frac{1}{2}} h - (\Lambda + \lambda I)^{-\nu} T^{\alpha - \frac{1}{2}} h \|_{L^2}$$

$$\leq \lambda^{\nu'} \| (\hat{\Lambda}_n + \lambda I)^{-\nu'} T^{\alpha - \frac{1}{2}} h - (\Lambda + \lambda I)^{-\nu'} T^{\alpha - \frac{1}{2}} h \|_{L^2}.$$
(5.6)

Using Lemma A.3, we get

$$\begin{split} &\|(\hat{\Lambda}_{n}+\lambda I)^{-\nu'}T^{\alpha-\frac{1}{2}}h-(\Lambda+\lambda I)^{-\nu'}T^{\alpha-\frac{1}{2}}h\|_{L^{2}} \\ \leq &\|(\hat{\Lambda}_{n}+\lambda I)^{-(\nu'-1)}[(\hat{\Lambda}_{n}+\lambda I)^{-1}T^{\alpha-\frac{1}{2}}h-(\Lambda+\lambda I)^{-1}T^{\alpha-\frac{1}{2}}h]\|_{L^{2}} \\ &+\|\sum_{i=1}^{\nu'-1}(\hat{\Lambda}_{n}+\lambda I)^{-i}(\Lambda-\hat{\Lambda}_{n})(\Lambda+\lambda I)^{-(\nu'+1-i)}T^{\alpha-\frac{1}{2}}h\|_{L^{2}} \\ \lesssim &\frac{1}{\lambda^{\nu'-1}}\|(\hat{\Lambda}_{n}+\lambda I)^{-1}T^{\alpha-\frac{1}{2}}-(\Lambda+\lambda I)^{-1}T^{\alpha-\frac{1}{2}}\|_{L^{2}\to L^{2}} \\ &+\sum_{i=1}^{\nu'-1}\frac{1}{\lambda^{i-\frac{1}{2}}}\frac{1}{\lambda^{\nu'+1-i-\frac{1}{2}}}\|(\hat{\Lambda}_{n}+\lambda I)^{-\frac{1}{2}}(\Lambda-\hat{\Lambda}_{n})(\Lambda+\lambda I)^{-\frac{1}{2}}T^{\alpha-\frac{1}{2}}\|_{L^{2}\to L^{2}} \\ \lesssim &\frac{1}{\lambda^{\nu'}}\|(\hat{\Lambda}_{n}+\lambda I)^{-\frac{1}{2}}(\Lambda-\hat{\Lambda}_{n})(\Lambda+\lambda I)^{-\frac{1}{2}}\|_{L^{2}\to L^{2}}. \end{split}$$

By using $p = \frac{1}{2}$ and $\alpha = \frac{1}{2}$ in Lemma A.5, we get

$$\|(\hat{\Lambda}_n + \lambda I)^{-\nu'} T^{\alpha - \frac{1}{2}} h - (\Lambda + \lambda I)^{-\nu'} T^{\alpha - \frac{1}{2}} h\|_{L^2} \lesssim_p \frac{1}{\lambda^{\nu'}} \frac{\lambda^{-\frac{1}{2(t+c)}}}{\sqrt{n}}.$$

Using this bound in (5.6) yields

$$\lambda^{\nu} \| (\Lambda + \lambda I)^{-(\nu - \nu')} (\hat{\Lambda}_n + \lambda I)^{-\nu'} T^{\alpha - \frac{1}{2}} h - (\Lambda + \lambda I)^{-\nu} T^{\alpha - \frac{1}{2}} h \|_{L^2} \lesssim_p \frac{\lambda^{-\frac{1}{2(t+c)}}}{\sqrt{n}}.$$
 (5.7)

Bounding Term-2e:

$$\begin{split} \|T^{\frac{1}{2}}r_{\lambda}(\hat{\Lambda}_{n})(\hat{\Lambda}_{n}+\lambda I)^{\nu}(\hat{\Lambda}_{n}+\lambda I)^{-(\nu-\nu')}(\Lambda+\lambda I)^{\nu-\nu'}(\Lambda+\lambda I)^{-\nu}T^{\alpha-\frac{1}{2}}h\|_{L^{2}} \\ \leq \|T^{\frac{1}{2}}(\Lambda+\lambda I)^{-\frac{1}{2}}\|_{L^{2}\to L^{2}}\|(\Lambda+\lambda I)^{\frac{1}{2}}(\hat{\Lambda}_{n}+\lambda I)^{-\frac{1}{2}}\|_{L^{2}\to L^{2}} \\ &\times \|(\hat{\Lambda}_{n}+\lambda I)^{\frac{1}{2}}r_{\lambda}(\hat{\Lambda}_{n})(\hat{\Lambda}_{n}+\lambda I)^{\nu-\frac{1}{2}}\|_{L^{2}\to L^{2}}\|(\Lambda+\lambda I)^{-(\nu-\frac{1}{2})}T^{\alpha-\frac{1}{2}}h\|_{L^{2}} \\ &\times \|(\hat{\Lambda}_{n}+\lambda I)^{\frac{1}{2}}(\hat{\Lambda}_{n}+\lambda I)^{-(\nu-\nu')}(\Lambda+\lambda I)^{\nu-\nu'}(\Lambda+\lambda I)^{-\frac{1}{2}}\|_{L^{2}\to L^{2}}. \end{split}$$

Using (5.1), Lemma A.6, Lemma A.2 and the qualification property (2.6) of regularization family, we get

$$\begin{aligned} \|T^{\frac{1}{2}}r_{\lambda}(\hat{\Lambda}_{n})(\hat{\Lambda}_{n}+\lambda I)^{\nu}(\hat{\Lambda}_{n}+\lambda I)^{-(\nu-\nu')}(\Lambda+\lambda I)^{\nu-\nu'}(\Lambda+\lambda I)^{-\nu}T^{\alpha-\frac{1}{2}}h\|_{L^{2}} \\ \lesssim_{p} \lambda^{-\frac{c}{2(t+c)}}\lambda^{\nu}\|(\Lambda+\lambda I)^{-(\nu-\frac{1}{2})}T^{\alpha-\frac{1}{2}}h\|_{L^{2}}, \end{aligned}$$

where

$$\begin{aligned} \|(\Lambda+\lambda I)^{-(\nu-\frac{1}{2})}T^{\alpha-\frac{1}{2}}h\|_{L^{2}} &\leq \|(\Lambda+\lambda I)^{-(\nu-\frac{1}{2})}T^{\alpha-\frac{1}{2}}\|_{L^{2}\to L^{2}}\|h\|_{L^{2}} \\ &\leq \sup_{i} \left|\frac{\mu_{i}^{\alpha-\frac{1}{2}}}{(\mu_{i}\xi_{i}+\lambda)^{(\nu-\frac{1}{2})}}\right| \|h\|_{L^{2}} \lesssim \sup_{i} \left|\frac{i^{-t(\alpha-\frac{1}{2})}}{(i^{-t-c}+\lambda)^{(\nu-\frac{1}{2})}}\right| \lesssim \lambda^{\frac{rt-\frac{t}{2}-(t+c)(\nu-\frac{1}{2})}{t+c}}, \end{aligned}$$

for $r = \min\{\alpha, \nu + \frac{c}{t}(\nu - \frac{1}{2})\}$, where the last inequality follows from Lemma A.9 with $(a, m, p, q, l) = (\frac{1}{2}, t, \alpha, t + c, \nu - \frac{1}{2})$, resulting in

$$\begin{aligned} \|T^{\frac{1}{2}}r_{\lambda}(\hat{\Lambda}_{n})(\hat{\Lambda}_{n}+\lambda I)^{\nu}(\hat{\Lambda}_{n}+\lambda I)^{-(\nu-\nu')}(\Lambda+\lambda I)^{\nu-\nu'}(\Lambda+\lambda I)^{-\nu}T^{\alpha-\frac{1}{2}}h\|_{L^{2}} \\ \lesssim_{p} \lambda^{\frac{rt}{t+c}}. \end{aligned}$$

Combining all the bounds we get

$$\|\beta^* - \hat{\beta}\|_{L^2} \lesssim_p \begin{cases} \lambda^{-\frac{c}{2(t+c)}} \sqrt{\frac{\sigma^2 \mathcal{N}(\lambda)}{n}} + \lambda^{\frac{\alpha t}{t+c}} + \frac{\lambda^{-\frac{1-2\alpha t}{2(t+c)}}}{\sqrt{n}} & \text{if } 0 < \alpha \leq \frac{1}{2} \\ \lambda^{-\frac{c}{2(t+c)}} \sqrt{\frac{\sigma^2 \mathcal{N}(\lambda)}{n}} + \lambda^{\frac{\alpha t}{t+c}} + \frac{\lambda^{-\frac{1}{2(t+c)}}}{\sqrt{n}} & \text{if } \frac{1}{2} \leq \alpha \leq \nu + \frac{c}{t}(\nu - \frac{1}{2}) \\ \lambda^{-\frac{c}{2(t+c)}} \sqrt{\frac{\sigma^2 \mathcal{N}(\lambda)}{n}} + \lambda^{\frac{(\nu + \frac{c}{t}(\nu - \frac{1}{2}))t}{t+c}} + \frac{\lambda^{-\frac{1}{2(t+c)}}}{\sqrt{n}} & \text{if } \alpha \geq \nu + \frac{c}{t}(\nu - \frac{1}{2}) \end{cases}$$

where the effective dimension $\mathcal{N}(\lambda)$ can be bounded as

$$\mathcal{N}(\lambda) = \operatorname{trace}(\Lambda(\Lambda + \lambda I)^{-1}) = \sum_{i} \frac{\mu_i \xi_i}{\mu_i \xi_i + \lambda} \lesssim \sum_{i} \frac{i^{-(t+c)}}{i^{-(t+c)} + \lambda} \lesssim \lambda^{-\frac{1}{t+c}}.$$

Therefore,

$$\|\hat{\beta} - \beta^*\|_{L^2} \lesssim_p \frac{\lambda^{-\frac{(c+1)}{2(t+c)}}}{\sqrt{n}} + \lambda^{\frac{rt}{t+c}} \lesssim n^{-\frac{rt}{1+c+2rt}}$$

14

for $\lambda = n^{-\frac{t+c}{1+c+2rt}}$ and the result follows.

5.2. **Proof of part (b) in Theorem 3.1.** We start by considering the error term in RKHS norm and applying Lemma A.1.

$$\begin{split} \|\hat{\beta} - \beta^*\|_{\mathcal{H}} &= \|J\hat{\beta} - \beta^*\|_{\mathcal{H}} = \|Jg_{\lambda}(J^*\hat{C}_n J)J^*\hat{R} - \beta^*\|_{\mathcal{H}} \\ &= \|T^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_n)T^{\frac{1}{2}}\hat{R} - \beta^*\|_{\mathcal{H}} = \|T^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_n)T^{\frac{1}{2}}\hat{R} - T^{\alpha}h\|_{\mathcal{H}} \\ &\leq \|T^{\frac{1}{2}}\|_{L^2 \to \mathcal{H}}\|g_{\lambda}(\hat{\Lambda}_n)T^{\frac{1}{2}}\hat{R} - T^{\alpha-\frac{1}{2}}h\|_{L^2} \quad (\|T^{\frac{1}{2}}\|_{L^2 \to \mathcal{H}} = 1) \\ &\leq \|g_{\lambda}(\hat{\Lambda}_n)(T^{\frac{1}{2}}\hat{R} - T^{\frac{1}{2}}\hat{C}_n\beta^*)\|_{L^2} + \|g_{\lambda}(\hat{\Lambda}_n)T^{\frac{1}{2}}\hat{C}_n\beta^* - T^{\alpha-\frac{1}{2}}h\|_{L^2} \\ &\leq (\|g_{\lambda}(\hat{\Lambda}_n)(\hat{\Lambda}_n + \lambda I)^{\frac{1}{2}}\|_{L^2 \to L^2}\|(\hat{\Lambda}_n + \lambda I)^{-\frac{1}{2}}(\Lambda + \lambda I)^{\frac{1}{2}}\|_{L^2 \to L^2} \\ &\times \|(\Lambda + \lambda I)^{-\frac{1}{2}}(T^{\frac{1}{2}}\hat{R} - T^{\frac{1}{2}}\hat{C}_n\beta^*)\|_{L^2}) + \|g_{\lambda}(\hat{\Lambda}_n)T^{\frac{1}{2}}\hat{C}_n\beta^* - T^{\alpha-\frac{1}{2}}h\|_{L^2}. \end{split}$$

From Lemma A.2, Lemma A.6 and regularization property (2.4), we get

$$\|\hat{\beta} - \beta^*\|_{\mathcal{H}} \lesssim_p \frac{1}{\sqrt{\lambda}} \frac{\lambda^{-\frac{1}{2(t+c)}}}{\sqrt{n}} + \|g_{\lambda}(\hat{\Lambda}_n)T^{\frac{1}{2}}\hat{C}_n\beta^* - T^{\alpha-\frac{1}{2}}h\|_{L^2}.$$

Now we bound the second term in the above inequality as

$$\begin{split} \|g_{\lambda}(\hat{\Lambda}_{n})T^{\frac{1}{2}}\hat{C}_{n}\beta^{*} - T^{\alpha-\frac{1}{2}}h\|_{L^{2}} &= \|r_{\lambda}(\hat{\Lambda}_{n})T^{\alpha-\frac{1}{2}}h\|_{L^{2}} \\ &= \|r_{\lambda}(\hat{\Lambda}_{n})(\hat{\Lambda}_{n} + \lambda I)^{\nu'}(\hat{\Lambda}_{n} + \lambda I)^{-\nu'}T^{\alpha-\frac{1}{2}}h\|_{L^{2}} \\ &= \|r_{\lambda}(\hat{\Lambda}_{n})(\hat{\Lambda}_{n} + \lambda I)^{\nu}(\hat{\Lambda}_{n} + \lambda I)^{-(\nu-\nu')}(\Lambda + \lambda I)^{\nu-\nu'} \\ &\times (\Lambda + \lambda I)^{-(\nu-\nu')}(\hat{\Lambda}_{n} + \lambda I)^{-\nu'}T^{\alpha-\frac{1}{2}}h\|_{L^{2}} \\ &= \|r_{\lambda}(\hat{\Lambda}_{n})(\hat{\Lambda}_{n} + \lambda I)^{\nu}(\hat{\Lambda}_{n} + \lambda I)^{-(\nu-\nu')}(\Lambda + \lambda I)^{\nu-\nu'} \\ &\times ((\Lambda + \lambda I)^{-(\nu-\nu')}(\hat{\Lambda}_{n} + \lambda I)^{-\nu'}T^{\alpha-\frac{1}{2}}h - (\Lambda + \lambda I)^{-\nu}T^{\alpha-\frac{1}{2}}h + (\Lambda + \lambda I)^{-\nu}T^{\alpha-\frac{1}{2}}h)\|_{L^{2}} \\ &\lesssim_{p}\lambda^{\nu}\underbrace{\|(\Lambda + \lambda I)^{-(\nu-\nu')}(\hat{\Lambda}_{n} + \lambda I)^{-\nu'}T^{\alpha-\frac{1}{2}}h - (\Lambda + \lambda I)^{-\nu}T^{\alpha-\frac{1}{2}}h\|_{L^{2}}}_{Term-1} \\ &+ \lambda^{\nu}\underbrace{\|(\Lambda + \lambda I)^{-\nu}T^{\alpha-\frac{1}{2}}h\|_{L^{2}}}_{Term-2}. \end{split}$$

The last inequality follows from Lemma A.6 and the qualification property (2.6) of the regularization family.

Bounding Term-1:

$$\|(\Lambda + \lambda I)^{-(\nu - \nu')} (\hat{\Lambda}_n + \lambda I)^{-\nu'} T^{\alpha - \frac{1}{2}} h - (\Lambda + \lambda I)^{-\nu} T^{\alpha - \frac{1}{2}} h \|_{L^2}$$

$$\leq \frac{1}{\lambda^{(\nu - \nu')}} \|(\hat{\Lambda}_n + \lambda I)^{-\nu'} T^{\alpha - \frac{1}{2}} h - (\Lambda + \lambda I)^{-\nu'} T^{\alpha - \frac{1}{2}} h \|_{L^2}.$$

From (5.7), we get

$$\|(\Lambda + \lambda I)^{-(\nu - \nu')} (\hat{\Lambda}_n + \lambda I)^{-\nu'} T^{\alpha - \frac{1}{2}} h - (\Lambda + \lambda I)^{-\nu} T^{\alpha - \frac{1}{2}} h\|_{L^2}$$

$$\lesssim_p \frac{1}{\lambda^{(\nu - \nu')}} \frac{1}{\lambda^{\nu'}} \frac{\lambda^{-\frac{1}{2(t+c)}}}{\sqrt{n}} = \lambda^{-\nu} \frac{\lambda^{-\frac{1}{2(t+c)}}}{\sqrt{n}}.$$

Bounding Term-2: We apply Lemma A.9 with $(a, m, p, q, l) = (\frac{1}{2}, t, \alpha, t + c, \nu)$ to get

$$\|(\Lambda + \lambda I)^{-\nu} T^{\alpha - \frac{1}{2}} h\|_{L^2} \lesssim \sup_i \left| \frac{\mu_i^{\alpha - \frac{1}{2}}}{(\mu_i \xi_i + \lambda)^{\nu}} \right| \lesssim \sup_i \left| \frac{i^{-t(\alpha - \frac{1}{2})}}{(i^{-(t+c)} + \lambda)^{\nu}} \right| \lesssim \lambda^{\frac{t(r - \frac{1}{2}) - (t+c)\nu}{t+c}},$$
ere $r = \min\{\alpha, \nu + \frac{c}{2}\nu + \frac{1}{2}\}$

where $r = \min\{\alpha, \nu + \frac{c}{t}\nu + \frac{1}{2}\}.$

Combining the bounds on Term-1 and Term-2, we have

$$\|g_{\lambda}(\hat{\Lambda}_{n})T^{\frac{1}{2}}\hat{C}_{n}\beta^{*} - T^{\alpha-\frac{1}{2}}h\|_{L^{2}} = \|r_{\lambda}(\hat{\Lambda}_{n})T^{\alpha-\frac{1}{2}}h\|_{L^{2}} \lesssim_{p} \frac{\lambda^{-\frac{1}{2(t+c)}}}{\sqrt{n}} + \lambda^{\frac{t(r-\frac{1}{2})}{t+c}}.$$

resulting in

$$\|\hat{\beta} - \beta^*\|_{\mathcal{H}} \lesssim_p \lambda^{\frac{t(r-\frac{1}{2})}{t+c}} + \frac{1}{\sqrt{\lambda}} \frac{\lambda^{-\frac{1}{2(t+c)}}}{\sqrt{n}}.$$

The result follows by choosing $\lambda = n^{-\frac{t+c}{1+c+2rt}}$.

5.3. Proof of Theorem 3.3. Consider

$$\begin{aligned} &\|\hat{\beta} - \beta^*\|_{\mathcal{H}} = \|J\hat{\beta} - \beta^*\|_{\mathcal{H}} = \|Jg_{\lambda}(J^*\hat{C}_n J)J^*\hat{R} - \beta^*\|_{\mathcal{H}} \\ &= \|T^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_n)T^{\frac{1}{2}}\hat{R} - \beta^*\|_{\mathcal{H}} \\ &\leq \|T^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_n)(T^{\frac{1}{2}}\hat{R} - T^{\frac{1}{2}}\hat{C}_n\beta^*)\|_{\mathcal{H}} + \|T^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_n)T^{\frac{1}{2}}\hat{C}_n\beta^* - \beta^*\|_{\mathcal{H}}. \end{aligned}$$

The first term on the r.h.s. can be bounded as

$$\begin{split} \|T^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_{n})(T^{\frac{1}{2}}\hat{R}-T^{\frac{1}{2}}\hat{C}_{n}\beta^{*})\|_{\mathcal{H}} \\ \leq \|T^{\frac{1}{2}}\|_{L^{2}\to\mathcal{H}}\|g_{\lambda}(\hat{\Lambda}_{n})(\hat{\Lambda}_{n}+\lambda I)^{\frac{1}{2}}\|_{L^{2}\to L^{2}}\|(\hat{\Lambda}_{n}+\lambda I)^{-\frac{1}{2}}(\Lambda+\lambda I)^{\frac{1}{2}}\|_{L^{2}\to L^{2}} \\ & \times \|(\Lambda+\lambda I)^{-\frac{1}{2}}(T^{\frac{1}{2}}\hat{R}-T^{\frac{1}{2}}\hat{C}_{n}\beta^{*})\|_{L^{2}}. \end{split}$$

Using regularization properties (2.4), Lemma A.6 and Lemma A.2, we get

$$\|T^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_n)(T^{\frac{1}{2}}\hat{R}-T^{\frac{1}{2}}\hat{C}_n\beta^*)\|_{\mathcal{H}} \lesssim_p \frac{1}{\sqrt{\lambda}}\sqrt{\frac{\sigma^2\mathcal{N}(\lambda)}{n}}.$$

Now we consider

$$\begin{split} \|T^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_{n})T^{\frac{1}{2}}\hat{C}_{n}\beta^{*} - \beta^{*}\|_{\mathcal{H}} &= \|T^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_{n})T^{\frac{1}{2}}\hat{C}_{n}T^{\frac{1}{2}}\Lambda^{s}h - T^{\frac{1}{2}}\Lambda^{s}h\|_{\mathcal{H}} \\ &= \|T^{\frac{1}{2}}(g_{\lambda}(\hat{\Lambda}_{n})\hat{\Lambda}_{n} - I)\Lambda^{s}h\|_{\mathcal{H}} = \|T^{\frac{1}{2}}r_{\lambda}(\hat{\Lambda}_{n})\Lambda^{s}h\|_{\mathcal{H}} \\ &= \|T^{\frac{1}{2}}r_{\lambda}(\hat{\Lambda}_{n})(\hat{\Lambda}_{n} + \lambda I)^{\nu}(\hat{\Lambda}_{n} + \lambda I)^{-(\nu-\nu')}(\hat{\Lambda}_{n} + \lambda I)^{-\nu'}\Lambda^{s}h\|_{\mathcal{H}} \\ &= \|T^{\frac{1}{2}}r_{\lambda}(\hat{\Lambda}_{n})(\hat{\Lambda}_{n} + \lambda I)^{\nu}(\hat{\Lambda}_{n} + \lambda I)^{-(\nu-\nu')}(\Lambda + \lambda I)^{(\nu-\nu')} \\ &\times [(\Lambda + \lambda I)^{-(\nu-\nu')}(\hat{\Lambda}_{n} + \lambda I)^{-\nu'}\Lambda^{s}h - (\Lambda + \lambda I)^{-\nu}\Lambda^{s}h + (\Lambda + \lambda I)^{-\nu}\Lambda^{s}h]\|_{\mathcal{H}} \\ \lesssim_{p}\underbrace{\lambda^{\nu}\|(\Lambda + \lambda I)^{-(\nu-\nu')}(\hat{\Lambda}_{n} + \lambda I)^{-\nu'}\Lambda^{s}h - (\Lambda + \lambda I)^{-\nu}\Lambda^{s}h\|_{L^{2}}}_{Term-3} \\ &+ \underbrace{\lambda^{\nu}\|(\Lambda + \lambda I)^{-\nu}\Lambda^{s}h\|_{L^{2}}}_{Term-4}, \end{split}$$

where we used Lemma A.6 and the qualification properties (2.6) of the regularization family in the last equality.

Bounding Term-3:

$$\lambda^{\nu} \| (\Lambda + \lambda I)^{-(\nu - \nu')} (\hat{\Lambda}_n + \lambda I)^{-\nu'} \Lambda^s h - (\Lambda + \lambda I)^{-\nu} \Lambda^s h \|_{L^2}$$

$$\leq \lambda^{\nu} \| (\Lambda + \lambda I)^{-(\nu - \nu')} ((\hat{\Lambda}_n + \lambda I)^{-\nu'} \Lambda^s h - (\Lambda + \lambda I)^{-\nu'} \Lambda^s h) \|_{L^2}$$

$$\leq \lambda^{\nu'} \| (\hat{\Lambda}_n + \lambda I)^{-\nu'} \Lambda^s h - (\Lambda + \lambda I)^{-\nu'} \Lambda^s h \|_{L^2}.$$

Using Lemma A.3, we get

$$\begin{split} &\|(\hat{\Lambda}_{n}+\lambda I)^{-\nu'}\Lambda^{s}h-(\Lambda+\lambda I)^{-\nu'}\Lambda^{s}h\|_{L^{2}} \\ \leq &\lambda^{-(\nu'-1)}\|(\hat{\Lambda}_{n}+\lambda I)^{-1}\Lambda^{s}h-(\Lambda+\lambda I)^{-1}\Lambda^{s}h\|_{L^{2}} \\ &+\left\|\sum_{i=1}^{\nu'-1}(\hat{\Lambda}_{n}+\lambda I)^{-i}(\Lambda-\hat{\Lambda}_{n})(\Lambda+\lambda I)^{-(\nu'+1-i)}\Lambda^{s}h\right\|_{L^{2}} \\ \lesssim &\lambda^{-(\nu'-1)}\|(\hat{\Lambda}_{n}+\lambda I)^{-1}(\Lambda-\hat{\Lambda}_{n})(\Lambda+\lambda I)^{-1}\Lambda^{s}\|_{L^{2}\to L^{2}} \\ &+\sum_{i=1}^{\nu'-1}\frac{1}{\lambda^{i-\frac{1}{2}}}\frac{1}{\lambda^{\nu'+1-i-\frac{1}{2}}}\|(\hat{\Lambda}_{n}+\lambda I)^{-\frac{1}{2}}(\Lambda-\hat{\Lambda}_{n})(\Lambda+\lambda I)^{-\frac{1}{2}}\|_{L^{2}\to L^{2}} \\ \lesssim &\lambda^{-\nu'}\|(\hat{\Lambda}_{n}+\lambda I)^{-\frac{1}{2}}(\Lambda-\hat{\Lambda}_{n})(\Lambda+\lambda I)^{-\frac{1}{2}}\|_{L^{2}\to L^{2}} \lesssim_{p}\lambda^{-\nu'}\frac{\lambda^{-\frac{1}{2b}}}{\sqrt{n}}, \end{split}$$

where the last inequality follows from Lemma A.5.

Bounding Term-4: We apply Lemma A.9 with $(a, m, p, q, l) = (0, b, s, b, \nu)$ to get

$$\lambda^{\nu} \| (\Lambda + \lambda I)^{-\nu} \Lambda^{s} h \|_{L^{2}} \leq \lambda^{\nu} \| (\Lambda + \lambda I)^{-\nu} \Lambda^{s} \|_{L^{2} \to L^{2}} \| h \|_{L^{2}}$$
$$\lesssim \lambda^{\nu} \sup_{i} \left| \frac{\tau_{i}^{s}}{(\tau_{i} + \lambda)^{\nu}} \right| \lesssim \lambda^{\nu} \sup_{i} \left| \frac{i^{-sb}}{(i^{-b} + \lambda)^{\nu}} \right| \leq \lambda^{\nu} \lambda^{\frac{rb-b\nu}{b}} = \lambda^{r},$$

where $r = \min\{s, \nu\}$.

Combing all the bounds, we get

$$\|\beta^* - \hat{\beta}\|_{\mathcal{H}} \lesssim_p \frac{1}{\sqrt{\lambda}} \sqrt{\frac{\sigma^2 \mathcal{N}(\lambda)}{n}} + \frac{\lambda^{-\frac{1}{2b}}}{\sqrt{n}} + \lambda^r,$$

where

$$\mathcal{N}(\lambda) = \operatorname{trace}(\Lambda(\Lambda + \lambda I)^{-1}) = \sum_{i} \left[\frac{\tau_i}{\tau_i + \lambda} \right] \lesssim \sum_{i} \left[\frac{i^{-b}}{i^{-b} + \lambda} \right] \lesssim \lambda^{-\frac{1}{b}}$$

resulting in

$$\|\hat{\beta} - \beta^*\|_{\mathcal{H}} \lesssim_p \frac{\lambda^{-(\frac{1}{2b} + \frac{1}{2})}}{\sqrt{n}} + \lambda^r.$$

The result follows for $\lambda = n^{-\frac{b}{1+b+2rb}}$.

5.4. Proof of Theorem 3.4. Consider

$$\|C^{\frac{1}{2}}(\hat{\beta}-\beta^*)\|_{L^2} = \|C^{\frac{1}{2}}(J\hat{\beta}-\beta^*)\|_{L^2} = \|C^{\frac{1}{2}}(Jg_{\lambda}(J^*\hat{C}_nJ)J^*\hat{R}-\beta^*)\|_{L^2}$$

Using Lemma A.1, we get

$$\begin{split} \|C^{\frac{1}{2}}(\hat{\beta}-\beta^{*})\|_{L^{2}} &= \|C^{\frac{1}{2}}(T^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_{n})T^{\frac{1}{2}}\hat{R}-\beta^{*})\|_{L^{2}} \\ &\leq \|C^{\frac{1}{2}}(T^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_{n})T^{\frac{1}{2}}\hat{R}-T^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_{n})T^{\frac{1}{2}}\hat{C}_{n}\beta^{*})\|_{L^{2}} \\ &\quad + \|C^{\frac{1}{2}}(T^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_{n})T^{\frac{1}{2}}\hat{C}_{n}\beta^{*}-\beta^{*})\|_{L^{2}} \\ &= \|C^{\frac{1}{2}}T^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_{n})(T^{\frac{1}{2}}\hat{R}-T^{\frac{1}{2}}\hat{C}_{n}\beta^{*})\|_{L^{2}} + \|C^{\frac{1}{2}}(T^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_{n})T^{\frac{1}{2}}\hat{C}_{n}\beta^{*}-\beta^{*})\|_{L^{2}} \\ &= \|\Lambda^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_{n})(T^{\frac{1}{2}}\hat{R}-T^{\frac{1}{2}}\hat{C}_{n}\beta^{*})\|_{L^{2}} + \|C^{\frac{1}{2}}(T^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_{n})T^{\frac{1}{2}}\hat{C}_{n}\beta^{*}-\beta^{*})\|_{L^{2}} \\ &\leq \|\Lambda^{\frac{1}{2}}(\Lambda+\lambda I)^{-\frac{1}{2}}\|_{L^{2}\to L^{2}}\|(\Lambda+\lambda I)^{\frac{1}{2}}(\hat{\Lambda}_{n}+\lambda I)^{-\frac{1}{2}}\|_{L^{2}\to L^{2}} \\ &\times \|(\hat{\Lambda}_{n}+\lambda I)^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_{n})(\hat{\Lambda}_{n}+\lambda I)^{\frac{1}{2}}\|_{L^{2}\to L^{2}} \\ &\times \|(\hat{\Lambda}_{n}+\lambda I)^{-\frac{1}{2}}(\Lambda+\lambda I)^{\frac{1}{2}}\|_{L^{2}\to L^{2}}\|(\Lambda+\lambda I)^{-\frac{1}{2}}(T^{\frac{1}{2}}\hat{R}-T^{\frac{1}{2}}\hat{C}_{n}\beta^{*})\|_{L^{2}} \\ &+ \|C^{\frac{1}{2}}(T^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_{n})T^{\frac{1}{2}}\hat{C}_{n}\beta^{*}-\beta^{*})\|_{L^{2}}. \end{split}$$

Using Lemma A.6, Lemma A.2 and regularization properties (2.3), (2.4), we get

$$\|C^{\frac{1}{2}}(\hat{\beta}-\beta^{*})\|_{L^{2}} \lesssim_{p} \frac{\lambda^{-\frac{1}{2(t+c)}}}{\sqrt{n}} + \|C^{\frac{1}{2}}(T^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_{n})T^{\frac{1}{2}}\hat{C}_{n}\beta^{*}-\beta^{*})\|_{L^{2}}.$$
(5.8)

Since β^* appearing in the second term on the r.h.s. of (5.8) depends on the source condition, we split the analysis into two cases depending on the value of α .

Case-1 : Let $\alpha \in (0, \frac{1}{2}]$.

$$\begin{split} \|C^{\frac{1}{2}}(T^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_{n})T^{\frac{1}{2}}\hat{C}_{n}\beta^{*}-\beta^{*})\|_{L^{2}} \\ \leq \underbrace{\|C^{\frac{1}{2}}T^{\frac{1}{2}}(g_{\lambda}(\hat{\Lambda}_{n})-(\hat{\Lambda}_{n}+\lambda I)^{-1})T^{\frac{1}{2}}\hat{C}_{n}\beta^{*}\|_{L^{2}}}_{Term-5} + \underbrace{\|C^{\frac{1}{2}}(T^{\frac{1}{2}}(\Lambda+\lambda I)^{-1}T^{\frac{1}{2}}C\beta^{*}-\beta^{*})\|_{L^{2}}}_{Term-6} \\ + \underbrace{\|C^{\frac{1}{2}}T^{\frac{1}{2}}((\hat{\Lambda}_{n}+\lambda I)^{-1}T^{\frac{1}{2}}\hat{C}_{n}\beta^{*}-(\Lambda+\lambda I)^{-1}T^{\frac{1}{2}}C\beta^{*})\|_{L^{2}}}_{Term-7}. \end{split}$$

Bounding Term-5:

$$\begin{split} \|C^{\frac{1}{2}}T^{\frac{1}{2}}(g_{\lambda}(\hat{\Lambda}_{n}) - (\hat{\Lambda}_{n} + \lambda I)^{-1})T^{\frac{1}{2}}\hat{C}_{n}\beta^{*}\|_{L^{2}} \\ \leq \|\Lambda^{\frac{1}{2}}(\hat{\Lambda}_{n} + \lambda I)^{-1}r_{\lambda}(\hat{\Lambda}_{n})T^{\frac{1}{2}}\hat{C}_{n}\beta^{*}\|_{L^{2}} + \lambda\|\Lambda^{\frac{1}{2}}(\hat{\Lambda}_{n} + \lambda I)^{-1}g_{\lambda}(\hat{\Lambda}_{n})T^{\frac{1}{2}}\hat{C}_{n}\beta^{*}\|_{L^{2}} \\ \lesssim \Big[\|(\hat{\Lambda}_{n} + \lambda I)^{-\frac{1}{2}}r_{\lambda}(\hat{\Lambda}_{n})(\hat{\Lambda}_{n} + \lambda I)^{\frac{3}{2}}\|_{L^{2} \to L^{2}} + \lambda\|(\hat{\Lambda}_{n} + \lambda I)^{-\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_{n})(\hat{\Lambda}_{n} + \lambda I)^{\frac{3}{2}}\|_{L^{2} \to L^{2}}\Big] \\ \times \|\Lambda^{\frac{1}{2}}(\Lambda + \lambda I)^{-\frac{1}{2}}\|_{L^{2} \to L^{2}}\|(\Lambda + \lambda I)^{\frac{1}{2}}(\hat{\Lambda}_{n} + \lambda I)^{-\frac{1}{2}}\|_{L^{2} \to L^{2}}\|(\hat{\Lambda}_{n} + \lambda I)^{-\frac{3}{2}}T^{\frac{1}{2}}\hat{C}_{n}\beta^{*}\|_{L^{2}}. \end{split}$$

From Lemma A.6 and regularization properties (2.3), (2.4) and (2.6), we get

$$\|C^{\frac{1}{2}}T^{\frac{1}{2}}(g_{\lambda}(\hat{\Lambda}_{n}) - (\hat{\Lambda}_{n} + \lambda I)^{-1})T^{\frac{1}{2}}\hat{C}_{n}\beta^{*}\|_{L^{2}} \lesssim_{p} \lambda \|(\hat{\Lambda}_{n} + \lambda I)^{-\frac{3}{2}}T^{\frac{1}{2}}\hat{C}_{n}\beta^{*}\|_{L^{2}}.$$

Using similar steps of (5.4), we get

$$\|C^{\frac{1}{2}}T^{\frac{1}{2}}(g_{\lambda}(\hat{\Lambda}_{n}) - (\hat{\Lambda}_{n} + \lambda I)^{-1})T^{\frac{1}{2}}\hat{C}_{n}\beta^{*}\|_{L^{2}} \lesssim_{p} \frac{\lambda^{-\frac{1-c-2t\alpha}{2(t+c)}}}{\sqrt{n}} + \lambda^{\frac{\alpha t + \frac{c}{2}}{t+c}}.$$

Bounding Term-6:

$$\begin{split} \|C^{\frac{1}{2}}(T^{\frac{1}{2}}(\Lambda+\lambda I)^{-1}T^{\frac{1}{2}}C\beta^{*}-\beta^{*})\|_{L^{2}} &= \|C^{\frac{1}{2}}(T^{\frac{1}{2}}(\Lambda+\lambda I)^{-1}T^{\frac{1}{2}}CT^{\alpha}h-T^{\alpha}h)\|_{L^{2}} \\ &= \left[\sum_{i}(\frac{\mu_{i}^{1+\alpha}\xi_{i}^{\frac{3}{2}}}{\mu_{i}\xi_{i}+\lambda}-\mu_{i}^{\alpha}\xi_{i}^{\frac{1}{2}})^{2}\langle\phi_{i},h\rangle^{2}\right]^{\frac{1}{2}} = \left[\sum_{i}(\frac{\lambda\mu_{i}^{\alpha}\xi_{i}^{\frac{1}{2}}}{\mu_{i}\xi_{i}+\lambda})^{2}\langle\phi_{i},h\rangle^{2}\right]^{\frac{1}{2}} \\ &\leq \lambda \sup_{i}\left[\frac{\mu_{i}^{\alpha}\xi_{i}^{\frac{1}{2}}}{\mu_{i}\xi_{i}+\lambda}\right]\|h\|_{L^{2}} \lesssim \lambda \sup_{i}\left[\frac{i^{-\alpha t-\frac{c}{2}}}{i^{-(t+c)}+\lambda}\right]\|h\|_{L^{2}} \lesssim \lambda^{\frac{\alpha t+\frac{c}{2}}{t+c}}. \end{split}$$

Bounding Term-7:

$$\|C^{\frac{1}{2}}T^{\frac{1}{2}}((\hat{\Lambda}_{n}+\lambda I)^{-1}T^{\frac{1}{2}}\hat{C}_{n}\beta^{*}-(\Lambda+\lambda I)^{-1}T^{\frac{1}{2}}C\beta^{*})\|_{L^{2}}$$

= $\|\Lambda^{\frac{1}{2}}((\hat{\Lambda}_{n}+\lambda I)^{-1}T^{\frac{1}{2}}\hat{C}_{n}\beta^{*}-(\Lambda+\lambda I)^{-1}T^{\frac{1}{2}}C\beta^{*})\|_{L^{2}}.$

Using the similar steps of (5.2), we get

$$\|C^{\frac{1}{2}}T^{\frac{1}{2}}((\hat{\Lambda}_{n}+\lambda I)^{-1}T^{\frac{1}{2}}\hat{C}_{n}\beta^{*}-(\Lambda+\lambda I)^{-1}T^{\frac{1}{2}}C\beta^{*})\|_{L^{2}} \lesssim_{p} \frac{\lambda^{-\frac{1-c-2t\alpha}{2(t+c)}}}{\sqrt{n}}.$$

Combining all the bounds, we get

$$\|C^{\frac{1}{2}}(T^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_{n})T^{\frac{1}{2}}\hat{C}_{n}\beta^{*}-\beta^{*})\|_{L^{2}} \lesssim_{p} \frac{\lambda^{-\frac{1-c-2t\alpha}{2(t+c)}}}{\sqrt{n}} + \lambda^{\frac{\alpha t+\frac{c}{2}}{t+c}}$$

Case-2: Suppose $\alpha \geq \frac{1}{2}$. Let ν be the qualification of the regularization family.

$$\begin{aligned} \|C^{\frac{1}{2}}(T^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_{n})T^{\frac{1}{2}}\hat{C}_{n}\beta^{*}-\beta^{*})\|_{L^{2}} \\ = \|C^{\frac{1}{2}}(T^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_{n})T^{\frac{1}{2}}\hat{C}_{n}T^{\frac{1}{2}}T^{\alpha-\frac{1}{2}}h-T^{\frac{1}{2}}T^{\alpha-\frac{1}{2}}h)\|_{L^{2}} \\ = \|C^{\frac{1}{2}}T^{\frac{1}{2}}r_{\lambda}(\hat{\Lambda}_{n})T^{\alpha-\frac{1}{2}}h\|_{L^{2}} = \|\Lambda^{\frac{1}{2}}r_{\lambda}(\hat{\Lambda}_{n})T^{\alpha-\frac{1}{2}}h\|_{L^{2}}.\end{aligned}$$

Following the steps of Case-2 in Theorem 3.1, we get

$$\|C^{\frac{1}{2}}(\hat{\beta}-\beta^*)\|_{L^2}^2 \lesssim_p \frac{\lambda^{-\frac{1}{t+c}}}{n} + \lambda^{\frac{2rt+c}{t+c}},$$

where $r = \min\{\alpha, \nu + \frac{1}{2} + \nu \frac{c}{t}\}$. Combining all the bounds, we get

$$\|C^{\frac{1}{2}}(\hat{\beta}-\beta^*)\|_{L^2}^2 \lesssim_p \frac{\lambda^{-\frac{1}{t+c}}}{n} + \lambda^{\frac{2rt+c}{t+c}}$$

and the result follows by choosing $\lambda = n^{-\frac{t+c}{1+c+2rt}}$.

5.5. Proof of Theorem 3.5. Consider

$$\|C^{\frac{1}{2}}(\hat{\beta}-\beta^*)\|_{L^2} = \|C^{\frac{1}{2}}(J\hat{\beta}-\beta^*)\|_{L^2} = \|C^{\frac{1}{2}}(Jg_{\lambda}(J^*\hat{C}_nJ)J^*\hat{R}-\beta^*)\|_{L^2}.$$

Following the steps of Theorem 3.4, we get

$$\begin{aligned} \|C^{\frac{1}{2}}(\hat{\beta}-\beta^{*})\|_{L^{2}} &\lesssim_{p} \|(\Lambda+\lambda I)^{-\frac{1}{2}}(T^{\frac{1}{2}}\hat{R}-T^{\frac{1}{2}}\hat{C}_{n}\beta^{*})\|_{L^{2}} \\ &+ \|C^{\frac{1}{2}}(T^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_{n})T^{\frac{1}{2}}\hat{C}_{n}\beta^{*}-\beta^{*})\|_{L^{2}}. \end{aligned}$$

Using Lemma A.2 yields

$$\|C^{\frac{1}{2}}(\hat{\beta}-\beta^{*})\|_{L^{2}} \lesssim_{p} \frac{\lambda^{-\frac{1}{2b}}}{\sqrt{n}} + \|C^{\frac{1}{2}}(T^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_{n})T^{\frac{1}{2}}\hat{C}_{n}\beta^{*}-\beta^{*})\|_{L^{2}}.$$

Now consider

$$\|C^{\frac{1}{2}}(T^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_{n})T^{\frac{1}{2}}\hat{C}_{n}\beta^{*}-\beta^{*})\|_{L^{2}} = \|\Lambda^{\frac{1}{2}}r_{\lambda}(\hat{\Lambda}_{n})\Lambda^{*}h\|_{L^{2}}.$$

Following steps similar to those of Theorem 3.3, we obtain

$$\begin{split} \|C^{\frac{1}{2}}(T^{\frac{1}{2}}g_{\lambda}(\hat{\Lambda}_{n})T^{\frac{1}{2}}\hat{C}_{n}\beta^{*}-\beta^{*})\|_{L^{2}} &\lesssim_{p}\frac{\lambda^{-\frac{1}{2b}}}{\sqrt{n}}+\lambda^{\nu}\|(\Lambda+\lambda I)^{-(\nu-\frac{1}{2})}\Lambda^{s}h\|_{L^{2}}\\ &\lesssim\frac{\lambda^{-\frac{1}{2b}}}{\sqrt{n}}+\lambda^{r+\frac{1}{2}}, \end{split}$$

where $r = \min\{s, \nu - \frac{1}{2}\}$. Combining all the bounds, we get

$$\|C^{\frac{1}{2}}(\hat{\beta} - \beta^*)\|_{L^2}^2 \lesssim_p \frac{\lambda^{-\frac{1}{b}}}{n} + \lambda^{2r+1}$$

and the result follows by choosing $\lambda = n^{-\frac{b}{1+b+2rb}}$.

5.6. **Proof of Theorem 4.3.** To establish the lower bound, without loss of generality, we assume $\epsilon \sim N(0, \sigma^2)$ since the lower bound for a special case guarantees the lower bound for the general case. Let M be the smallest integer greater than $c_0 n^{\frac{1}{q}}$ for some constant $c_0 > 0$ and q > 0, which will be specified later. For $\theta \in \{0, 1\}^M$, define

$$f_{\theta} := \sum_{k=1}^{M} \theta_k M^{-\frac{1}{2}} T^{\alpha} \phi_{k+M}.$$

Let P_{θ} denote the *n*-fold product of joint probability distributions for $\{(X_i, Y_i) : 1 \le i \le n\}$ with $\beta^* = f_{\theta}$. Then the Kullback-Leibler divergence between P_{θ} and $P_{\theta'}$ is

$$\mathcal{K}(P_{\theta}, P_{\theta'}) = \frac{n}{2\sigma^2} \|C^{\frac{1}{2}}(f_{\theta} - f_{\theta'})\|_{L_2}^2$$

for $\theta, \theta' \in \{0, 1\}^M$. Using the representation of f_{θ} and $f_{\theta'}$, we obtain

$$\begin{split} \mathcal{K}(P_{\theta}, P_{\theta'}) &= \frac{n}{2\sigma^2} \left\| \sum_{k=1}^{M} (\theta_k - \theta'_k) M^{-\frac{1}{2}} C^{\frac{1}{2}} T^{\alpha} \phi_{k+M} \right\|_{L_2}^2 \\ &= \frac{n}{2\sigma^2} \sum_{k=1}^{M} (\theta_k - \theta'_k)^2 M^{-1} \mu_{k+M}^{2\alpha} \xi_{k+M} \le \frac{n}{\sigma^2} M^{-1} \mu_M^{2\alpha} \xi_M H(\theta, \theta') \\ &\le \frac{d}{\sigma^2} n M^{-1} M^{-2\alpha t} M^{-c} M \le \frac{d}{\sigma^2 c_0^q} M^{q-2\alpha t-c}. \end{split}$$

In the last inequality, we used $c_0 n^{\frac{1}{q}} \leq M$. To get $\mathcal{K}(P_{\theta}, P_{\theta'}) \leq u \log(N)$ for some $0 < u < \frac{1}{8}$, we choose $q - 2\alpha t - c = 1$ and $c_0 = c'_0 u^{-\frac{1}{q}}$, where c'_0 is a large enough constant. So we have $q = 1 + 2\alpha t + c$.

Next, for $\alpha \geq \frac{1}{2}$, by considering the error term in the RKHS norm, we have

$$\begin{split} \|f_{\theta} - f_{\theta'}\|_{\mathcal{H}}^{2} &= \left\|\sum_{k=1}^{M} (\theta_{k} - \theta_{k}') M^{-\frac{1}{2}} T^{\alpha} \phi_{k+M} \right\|_{\mathcal{H}}^{2} &= \frac{1}{M} \left\|\sum_{k=1}^{M} (\theta_{k} - \theta_{k}') T^{\alpha - \frac{1}{2}} \phi_{k+M} \right\|_{L_{2}}^{2} \\ &= \frac{1}{M} \sum_{k=1}^{M} (\theta_{k} - \theta_{k}')^{2} \mu_{k+M}^{2(\alpha - \frac{1}{2})} \gtrsim M^{-1} H(\theta, \theta') \mu_{2M}^{2(\alpha - \frac{1}{2})} \overset{(*)}{\gtrsim} M^{-1} M M^{-2t(\alpha - \frac{1}{2})} \\ &= M^{-2t(\alpha - \frac{1}{2})} \overset{(\dagger)}{\gtrsim} u^{\frac{t(2\alpha - 1)}{q}} n^{-\frac{t(2\alpha - 1)}{q}}, \end{split}$$

where we used Lemma 4.2 in (*) and the fact that $\frac{M}{2} < c_0 n^{\frac{1}{q}}$ in (†). Hence,

$$\|f_{\theta} - f_{\theta'}\|_{\mathcal{H}} \gtrsim u^{\frac{t(\alpha - \frac{1}{2})}{1 + c + 2\alpha t}} n^{-\frac{t(\alpha - \frac{1}{2})}{1 + c + 2\alpha t}}.$$
(5.9)

Next, we consider the estimation error term in the L^2 norm as

$$\|f_{\theta} - f_{\theta'}\|_{L_{2}}^{2} = \left\| \sum_{k=1}^{M} (\theta_{k} - \theta_{k}') M^{-\frac{1}{2}} T^{\alpha} \phi_{k+M} \right\|_{L_{2}}^{2} = M^{-1} \sum_{k=1}^{M} (\theta_{k} - \theta_{k}')^{2} \mu_{k+M}^{2\alpha}$$

$$\gtrsim M^{-1} H(\theta, \theta') \mu_{2M}^{2\alpha} \gtrsim M^{-1} M M^{-2\alpha t} = M^{-2\alpha t} \gtrsim u^{\frac{2\alpha t}{q}} n^{-\frac{2\alpha t}{q}}.$$

Therefore, we get

$$\|f_{\theta} - f_{\theta'}\|_{L_2} \gtrsim u^{\frac{\alpha t}{1+c+2\alpha t}} n^{-\frac{\alpha t}{1+c+2\alpha t}}.$$
(5.10)

Next, we consider the error term for the prediction error:

$$\begin{split} \|C^{\frac{1}{2}}(f_{\theta} - f_{\theta'})\|_{L^{2}}^{2} &= \left\|\sum_{k=1}^{M} (\theta_{k} - \theta_{k}')M^{-\frac{1}{2}}C^{\frac{1}{2}}T^{\alpha}\phi_{k+M}\right\|_{L_{2}}^{2} \\ &= M^{-1}\sum_{k=1}^{M} (\theta_{k} - \theta_{k}')^{2}\mu_{k+M}^{2\alpha}\xi_{k+M} \gtrsim M^{-1}(2M)^{-2\alpha t}(2M)^{-c}H(\theta, \theta') \gtrsim M^{-2\alpha t-c} \\ &\geq u^{\frac{2\alpha t+c}{q}}n^{-\frac{2\alpha t+c}{q}}. \end{split}$$

Hence,

$$\|C^{\frac{1}{2}}(f_{\theta} - f_{\theta'})\|_{L^2}^2 \gtrsim u^{\frac{c+2\alpha t}{1+c+2\alpha t}} n^{-\frac{c+2\alpha t}{1+c+2\alpha t}}.$$
(5.11)

The desired results are obtained by applying Theorem 4.1 to equations (5.9), (5.10) and (5.11).

5.7. **Proof of Theorem 4.4.** Let M be the smallest integer greater than $c_0 n^{\frac{1}{q}}$ for some constant $c_0 > 0$ and q > 1 + b. For $\theta \in \{0, 1\}^M$, we define

$$f_{\theta} := \sum_{k=1}^{M} \theta_k M^{-\frac{1}{2}} T^{\frac{1}{2}} (T^{\frac{1}{2}} C T^{\frac{1}{2}})^s \phi_{k+M}.$$

Similar to the proof of Theorem 4.3, consider the Kullback-Leibler divergence between joint probability distributions P_{θ} and $P_{\theta'}$:

$$\begin{aligned} \mathcal{K}(P_{\theta}, P_{\theta'}) &= \frac{n}{2\sigma^2} \|C^{\frac{1}{2}}(f_{\theta} - f_{\theta'})\|_{L_2}^2 = \frac{n}{2\sigma^2} \left\|\sum_{k=1}^M (\theta_k - \theta'_k) M^{-\frac{1}{2}} C^{\frac{1}{2}} T^{\frac{1}{2}} \Lambda^s \phi_{k+M} \right\|_{L_2}^2 \\ &= \frac{nM^{-1}}{2\sigma^2} \sum_{k=1}^M (\theta_k - \theta'_k)^2 \tau_{k+M}^{1+2s} \le \frac{d'}{\sigma^2} nM^{-b(1+2s)} = \frac{d'}{\sigma^2} nM^{-b(1+2s)} \\ &\le \frac{d'}{\sigma^2 c_0^q} M^{-b(1+2s)+q}. \end{aligned}$$

In the last step, we used that $c_0 n^{\frac{1}{q}} \leq M$. To satisfy the condition on Kullback divergence, we take q = 1 + b(1 + 2s) and $c_0 = c'_0 u^{-\frac{1}{q}}$, where c'_0 is a large enough constant. Next, we consider the error term in RKHS norm:

$$\|f_{\theta} - f_{\theta'}\|_{\mathcal{H}}^{2} = M^{-1} \sum_{k=1}^{M} (\theta_{k} - \theta_{k}')^{2} \tau_{k+M}^{2s} \gtrsim M^{-1} \tau_{2M}^{2s} H(\theta, \theta')$$

$$\stackrel{(*)}{\gtrsim} M^{-1} M^{-b(2s)} M = M^{-b(2s)} \stackrel{(\dagger)}{\gtrsim} u^{\frac{2sb}{q}} n^{-\frac{2sb}{q}},$$

where we used Lemma 4.2 in (*) and the fact that $\frac{M}{2} < c_0 n^{\frac{1}{q}}$ in (†). Applying Theorem 4.1, we obtain

$$\lim_{a \to 0} \lim_{n \to \infty} \inf_{\hat{\beta}} \sup_{\beta^* \in \mathcal{R}(T^{\frac{1}{2}}(\Lambda)^s)} \mathbb{P}\{\|\hat{\beta} - \beta^*\|_{\mathcal{H}} \ge an^{-\frac{so}{1+b+2sb}}\} = 1.$$

The error term for prediction is given by

$$\begin{aligned} \|C^{\frac{1}{2}}(f_{\theta} - f_{\theta'})\|_{L^{2}}^{2} \gtrsim M^{-1}H(\theta, \theta')\tau_{2M}^{1+2s} \\ \gtrsim M^{-b(1+2s)} \gtrsim u^{\frac{b(1+2s)}{q}}n^{-\frac{b(1+2s)}{q}}, \end{aligned}$$

where applying Theorem 4.1 yields

$$\lim_{a \to 0} \lim_{n \to \infty} \inf_{\hat{\beta}} \sup_{\beta^* \in \mathcal{R}(T^{\frac{1}{2}}(\Lambda)^s)} \mathbb{P}\{\mathbb{E}\langle \hat{\beta} - \beta^*, X \rangle_{L^2}^2 \ge an^{-\frac{b(2s+1)}{1+b+2sb}}\} = 1$$

Acknowledgements

BKS is partially supported by the National Science Foundation (NSF) CAREER award DMS-1945396.

References

- [1] ARONSZAJN, N. Theory of reproducing kernels. Trans. Amer. Math. Soc. 68 (1950), 337-404.
- [2] BALASUBRAMANIAN, K., MÜLLER, H.-G., AND SRIPERUMBUDUR, B. K. Functional linear and singleindex models: A unified approach via Gaussian Stein identity. *Bernoulli. To appear. arXiv:2206.03975* (2024).
- BAUER, F., PEREVERZEV, S., AND ROSASCO, L. On regularization algorithms in learning theory. J. Complexity 23, 1 (2007), 52–72.
- [4] BENATIA, D., CARRASCO, M., AND FLORENS, J.-P. Functional linear regression with functional response. J. Econometrics 201, 2 (2017), 269–291.
- [5] CAI, T. T., AND HALL, P. Prediction in functional linear regression. Ann. Statist. 34, 5 (2006), 2159–2179.

- [6] CAI, T. T., AND YUAN, M. Minimax and adaptive prediction for functional linear regression. J. Amer. Statist. Assoc. 107, 499 (2012), 1201–1216.
- [7] CARDOT, H., FERRATY, F., AND SARDA, P. Functional linear model. Statist. Probab. Lett. 45, 1 (1999), 11–22.
- [8] CARDOT, H., FERRATY, F., AND SARDA, P. Spline estimators for the functional linear model. Statist. Sinica 13, 3 (2003), 571–591.
- CHEN, D., HALL, P., AND MÜLLER, H.-G. Single and multiple index functional regression models with nonparametric link. Ann. Statist. 39, 3 (2011), 1720–1747.
- [10] CORDES, H. O. Spectral Theory of Linear Differential Operators and Comparison Algebras, vol. 76 of London Mathematical Society Lecture Note Series. Cambridge University Press, Cambridge, 1987.
- [11] CUCKER, F., AND ZHOU, D.-X. Learning Theory: An Approximation Theory Viewpoint, vol. 24. Cambridge University Press, Cambridge, 2007.
- [12] CUEVAS, A., FEBRERO, M., AND FRAIMAN, R. Linear functional regression: The case of fixed design and functional response. *Canad. J. Statist.* 30, 2 (2002), 285–300.
- [13] ENGL, H. W., HANKE, M., AND NEUBAUER, A. Regularization of Inverse Problems, vol. 375 of Mathematics and its Applications. Kluwer Academic Publishers Group, Dordrecht, 1996.
- [14] HALL, P., AND HOROWITZ, J. L. Methodology and convergence rates for functional linear regression. Ann. Statist. 35, 1 (2007), 70–91.
- [15] HAPP, C., AND GREVEN, S. Multivariate functional principal component analysis for data observed on different (dimensional) domains. J. Amer. Statist. Assoc. 113, 522 (2018), 649–659.
- [16] HOLZLEITNER, M., AND PEREVERZYEV, S. V. On regularized polynomial functional regression. J. Complexity 83 (2024), Paper No. 101853, 21.
- [17] JAMES, G. M. Generalized linear models with functional predictors. J. R. Stat. Soc. Ser. B Stat. Methodol. 64, 3 (2002), 411–432.
- [18] JIANG, C.-R., AND WANG, J.-L. Functional single index models for longitudinal data. Ann. Statist. 39, 1 (2011), 362–388.
- [19] KOKOSZKA, P., AND REIMHERR, M. Introduction to Functional Data Analysis. Texts in Statistical Science Series. CRC Press, Boca Raton, FL, 2017.
- [20] LI, Y., AND HSING, T. On rates of convergence in functional linear regression. J. Multivariate Anal. 98, 9 (2007), 1782–1804.
- [21] LU, S., AND PEREVERZEV, S. V. Regularization Theory for Ill-posed Problems, vol. 58 of Inverse and Ill-posed Problems Series. De Gruyter, Berlin, 2013. Selected topics.
- [22] MOLLENHAUER, M., MÜCKE, N., AND SULLIVAN, T. Learning linear operators: Infinite-dimensional regression as a well-behaved non-compact inverse problem. arXiv preprint arXiv:2211.08875 (2022).
- [23] MORRIS, J. S. Functional regression. Annual Review of Statistics and Its Application 2 (2015), 321– 359.
- [24] MÜLLER, H.-G., AND STADTMÜLLER, U. Generalized functional linear models. Ann. Statist. 33, 2 (2005), 774–805.
- [25] PAULSEN, V. I., AND RAGHUPATHI, M. An Introduction to the Theory of Reproducing Kernel Hilbert Spaces, vol. 152 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 2016.
- [26] PEREVERZYEV, S. An Introduction to Artificial Intelligence Based on Reproducing Kernel Hilbert Spaces. Compact Textbooks in Mathematics. Birkhäuser, 2022.
- [27] RAMSAY, J. O. When the data are functions. Psychometrika 47, 4 (1982), 379–396.
- [28] RAMSAY, J. O., AND DALZELL, C. J. Some tools for functional data analysis. J. Roy. Statist. Soc. Ser. B 53, 3 (1991), 539–572.
- [29] RAMSAY, J. O., AND SILVERMAN, B. W. Applied Functional Data Analysis. Springer Series in Statistics. Springer-Verlag, New York, 2002. Methods and case studies.
- [30] REISS, P. T., GOLDSMITH, J., SHANG, H. L., AND OGDEN, R. T. Methods for scalar-on-function regression. Int. Stat. Rev. 85, 2 (2017), 228–249.
- [31] SHANG, Z., AND CHENG, G. Nonparametric inference in generalized functional linear models. Ann. Statist. 43, 4 (2015), 1742–1773.
- [32] STEINWART, I., AND CHRISTMANN, A. Support Vector Machines. Information Science and Statistics. Springer, New York, 2008.
- [33] TONG, H. Functional linear regression with Huber loss. J. Complexity 74 (2023), Paper No. 101696, 14.
- [34] TONG, H., AND NG, M. Analysis of regularized least squares for functional linear regression model. J. Complexity 49 (2018), 85–94.

- [35] TONG, H. Z., HU, L. F., AND NG, M. Non-asymptotic error bound for optimal prediction of functionon-function regression by RKHS approach. Acta Math. Sin. (Engl. Ser.) 38, 4 (2022), 777–796.
- [36] TSYBAKOV, A. B. Introduction to Nonparametric Estimation. Springer Series in Statistics.
- [37] WANG, J.-L., CHIOU, J.-M., AND MÜLLER, H.-G. Functional data analysis. Annual Review of Statistics and its Application 3 (2016), 257–295.
- [38] YUAN, M., AND CAI, T. T. A reproducing kernel Hilbert space approach to functional linear regression. Ann. Statist. 38, 6 (2010), 3412–3444.
- [39] ZHANG, F., ZHANG, W., LI, R., AND LIAN, H. Faster convergence rate for functional linear regression in reproducing kernel Hilbert spaces. *Statistics* 54, 1 (2020), 167–181.
- [40] ZHU, H., YAO, F., AND ZHANG, H. H. Structured functional additive regression in reproducing kernel Hilbert spaces. J. R. Stat. Soc. Ser. B. Stat. Methodol. 76, 3 (2014), 581–603.

APPENDIX A. SUPPLEMENTARY RESULTS

In this appendix, we present technical results that are used to prove the main results of the paper.

Lemma A.1. For any bounded linear positive operator $\mathcal{A}: L^2(S) \to L^2(S)$, we have

$$Jg_{\lambda}(J^*\mathcal{A}J)J^* = T^{1/2}g_{\lambda}(T^{\frac{1}{2}}\mathcal{A}T^{\frac{1}{2}})T^{1/2}.$$

Proof. Let $(m_i, \phi_i)_i$ be the eigensystem of the operator $J^* \mathcal{A} J : \mathcal{H} \to \mathcal{H}$. It is easy to verify that $(m_i, \frac{T^{1/2} \mathcal{A} J \phi_i}{m_i})_i$ is the eigensystem of $T^{1/2} \mathcal{A} T^{1/2}$ since

$$T^{1/2}\mathcal{A}T^{1/2}T^{1/2}\mathcal{A}J\phi_i = T^{1/2}\mathcal{A}T\mathcal{A}J\phi_i = T^{1/2}\mathcal{A}JJ^*\mathcal{A}J\phi_i = m_iT^{1/2}\mathcal{A}J\phi_i$$

and for any $i, j \in \mathbb{N}$,

$$\left\langle \frac{T^{1/2} \mathcal{A} J \phi_i}{m_i}, \frac{T^{1/2} \mathcal{A} J \phi_j}{m_j} \right\rangle_{L^2} = \left\langle \frac{\mathcal{A} J \phi_i}{m_i}, \frac{T \mathcal{A} J \phi_j}{m_j} \right\rangle_{L^2} = \left\langle \frac{\mathcal{A} J \phi_i}{m_i}, \frac{J J^* \mathcal{A} J \phi_j}{m_j} \right\rangle_{L^2}$$
$$= \left\langle \frac{J^* \mathcal{A} J \phi_i}{m_i}, \frac{J^* \mathcal{A} J \phi_j}{m_j} \right\rangle_{\mathcal{H}} = \left\langle \frac{m_i \phi_i}{m_i}, \frac{m_j \phi_j}{m_j} \right\rangle_{\mathcal{H}} = \langle \phi_i, \phi_j \rangle_{\mathcal{H}} = \delta_{ij}.$$

Therefore, using the spectral representation of $J^*\mathcal{A}J$, it follows that for all $f \in L^2(S)$,

$$T^{1/2}g_{\lambda}(T^{\frac{1}{2}}\mathcal{A}T^{\frac{1}{2}})T^{1/2}f = \sum_{i}g_{\lambda}(m_{i})\left\langle T^{1/2}f, \frac{T^{1/2}\mathcal{A}J\phi_{i}}{m_{i}}\right\rangle_{L^{2}}T^{1/2}\left(\frac{T^{1/2}\mathcal{A}J\phi_{i}}{m_{i}}\right)$$
$$= \sum_{i}g_{\lambda}(m_{i})\left\langle f, \frac{T\mathcal{A}J\phi_{i}}{m_{i}}\right\rangle_{L^{2}}\frac{T\mathcal{A}J\phi_{i}}{m_{i}} = \sum_{i}g_{\lambda}(m_{i})\left\langle f, \frac{JJ^{*}\mathcal{A}J\phi_{i}}{m_{i}}\right\rangle_{L^{2}}\frac{JJ^{*}\mathcal{A}J\phi_{i}}{m_{i}}$$
$$= \sum_{i}g_{\lambda}(m_{i})\langle f, J\phi_{i}\rangle_{L^{2}}J\phi_{i} = \sum_{i}g_{\lambda}(m_{i})\langle J^{*}f, \phi_{i}\rangle_{\mathcal{H}}J\phi_{i} = Jg_{\lambda}(J^{*}\mathcal{A}J)J^{*}f$$

and the result follows.

Lemma A.2. For any $\delta > 0$, with at least probability $1 - \delta$, we have that

$$\|(\Lambda+\lambda I)^{-\frac{1}{2}}T^{\frac{1}{2}}(\hat{R}-\hat{C}_n\beta^*)\|_{L^2} \leq \sqrt{\frac{\sigma^2\mathcal{N}(\lambda)}{n\delta}}.$$

Proof. For $i = 1, 2, \dots, n$, define $Z_i := (\Lambda + \lambda I)^{-\frac{1}{2}} T^{\frac{1}{2}} [Y_i X_i - (X_i \otimes X_i)\beta^*]$. Since the slope function β^* satisfies the operator equation $C\beta^* = \mathbb{E}[YX]$, the mean of the random variable Z_i is zero, i.e.,

$$\mathbb{E}[Z_i] = (\Lambda + \lambda I)^{-\frac{1}{2}} T^{\frac{1}{2}} [\mathbb{E}[YX] - C\beta^*] = 0.$$

By Markov's inequality, for any t > 0

$$\mathbb{P}\left(\left\|\frac{1}{n}\sum_{i=1}^{n}Z_{i}\right\|_{L^{2}}\geq t\right)\leq\frac{\mathbb{E}\|\frac{1}{n}\sum_{i=1}^{n}Z_{i}\|_{L^{2}}^{2}}{t^{2}}.$$

Note that

$$\mathbb{E}\left\|\frac{1}{n}\sum_{i=1}^{n}Z_{i}\right\|_{L^{2}}^{2} = \frac{1}{n^{2}}\sum_{i,j=1}^{n}\mathbb{E}\langle Z_{i}, Z_{j}\rangle_{L^{2}} = \frac{1}{n^{2}}\sum_{i\neq j}^{n}\mathbb{E}\langle Z_{i}, Z_{j}\rangle_{L^{2}} + \frac{1}{n^{2}}\sum_{i=1}^{n}\mathbb{E}\|Z_{i}\|_{L^{2}}^{2} = \frac{\mathbb{E}\|Z_{1}\|_{L^{2}}^{2}}{n}$$

and by taking $t = \sqrt{\frac{\mathbb{E} \|Z_1\|_{L^2}^2}{n\delta}}$, with at least probability $1 - \delta$, we have

$$\|(\Lambda + \lambda I)^{-\frac{1}{2}} T^{\frac{1}{2}} (\hat{R} - \hat{C}_n \beta^*)\|_{L^2} \le \sqrt{\frac{\mathbb{E} \|(\Lambda + \lambda I)^{-\frac{1}{2}} T^{\frac{1}{2}} (YX - (X \otimes X)\beta^*)\|_{L^2}^2}{n\delta}}.$$
 (A.1)

Consider

$$\mathbb{E}[\|(\Lambda + \lambda I)^{-\frac{1}{2}}T^{\frac{1}{2}}(YX - (X \otimes X)\beta^*)\|_{L^2}^2]$$

$$=\mathbb{E}[\|(\Lambda + \lambda I)^{-\frac{1}{2}}T^{\frac{1}{2}}(Y - \langle X, \beta^* \rangle_{L^2})X\|_{L^2}^2]$$

$$=\mathbb{E}[(Y - \langle X, \beta^* \rangle_{L^2})^2\|(\Lambda + \lambda I)^{-\frac{1}{2}}T^{\frac{1}{2}}X\|_{L^2}^2]$$

$$=\mathbb{E}[\epsilon^2 \langle (\Lambda + \lambda I)^{-\frac{1}{2}}T^{\frac{1}{2}}X, (\Lambda + \lambda I)^{-\frac{1}{2}}T^{\frac{1}{2}}X \rangle_{L^2}]$$

$$=\mathbb{E}[\epsilon^2 \operatorname{trace}((\Lambda + \lambda I)^{-1}T^{\frac{1}{2}}(X \otimes X)T^{\frac{1}{2}})]$$

$$=\mathbb{E}[\epsilon^2]\operatorname{trace}((\Lambda + \lambda I)^{-1}T^{\frac{1}{2}}CT^{\frac{1}{2}})$$

$$=\sigma^2\operatorname{trace}((\Lambda + \lambda I)^{-1}\Lambda) = \sigma^2\mathcal{N}(\lambda).$$

The result therefore follows from (A.1).

Lemma A.3. For any $n \ge 1$, we have

$$(\hat{\Lambda}_n + \lambda I)^{-n} - (\Lambda + \lambda I)^{-n} = (\hat{\Lambda}_n + \lambda I)^{-(n-1)} [(\hat{\Lambda}_n + \lambda I)^{-1} - (\Lambda + \lambda I)^{-1}] + \sum_{i=1}^{n-1} (\hat{\Lambda}_n + \lambda I)^{-i} (\Lambda - \hat{\Lambda}_n) (\Lambda + \lambda I)^{-(n+1-i)}.$$

Proof. Define

$$\Lambda_{\lambda} := (\Lambda + \lambda I)^{-1}$$
 and $\hat{\Lambda}_{\lambda} := (\hat{\Lambda}_n + \lambda I)^{-1}$.

Consider

$$\hat{\Lambda}^n_{\lambda} - \Lambda^n_{\lambda} = \hat{\Lambda}_{\lambda} (\hat{\Lambda}^{n-1}_{\lambda} - \hat{\Lambda}\Lambda^n_{\lambda} - \lambda\Lambda^n_{\lambda}) = \hat{\Lambda}_{\lambda} (\hat{\Lambda}^{n-1}_{\lambda} - \hat{\Lambda}\Lambda^n_{\lambda} - \Lambda^{n-1}_{\lambda} + \Lambda\Lambda^n_{\lambda}).$$

In the last step, we have used the fact that $\lambda \Lambda_{\lambda}^{n} = \Lambda_{\lambda}^{n-1} - \Lambda \Lambda_{\lambda}^{n}$, $\forall n \in \mathbb{N}$. Therefore,

$$\hat{\Lambda}^n_{\lambda} - \Lambda^n_{\lambda} = \hat{\Lambda}_{\lambda} (\Lambda - \hat{\Lambda}_n) \Lambda^n_{\lambda} + \hat{\Lambda}_{\lambda} (\hat{\Lambda}^{n-1}_{\lambda} - \Lambda^{n-1}_{\lambda}).$$

Doing similar step n-1 times, we get

$$\hat{\Lambda}^{n}_{\lambda} - \Lambda^{n}_{\lambda} = \hat{\Lambda}^{n-1}_{\lambda} [\hat{\Lambda}_{\lambda} - \Lambda_{\lambda}] + \sum_{i=1}^{n-1} \hat{\Lambda}^{i}_{\lambda} (\Lambda - \hat{\Lambda}_{n}) \Lambda^{n+1-i}_{\lambda}$$

and the result follows.

Lemma A.4. Suppose T and C commute and Assumptions 1, 3 and 5 hold. Then for all choices of p such that $2p(t+c) \ge 2\alpha t + c$, we have

$$\|(\Lambda + \lambda I)^{-p} T^{\alpha} (C - \hat{C}_n) T^{\alpha} (\Lambda + \lambda I)^{-p} \|_{L^2 \to L^2} \lesssim_p \frac{1}{\sqrt{n}} \lambda^{-\frac{1 + 4p(t+c) - (4\alpha t + 2c)}{2(t+c)}}.$$

Proof. Consider

$$\begin{aligned} &\|(\Lambda + \lambda I)^{-p} T^{\alpha} (C - \hat{C}_n) T^{\alpha} (\Lambda + \lambda I)^{-p} \|_{L^2 \to L^2} \\ &= \sup_{h: \|h\|_{L^2} = 1} \langle h, (\Lambda + \lambda I)^{-p} T^{\alpha} (C - \hat{C}_n) T^{\alpha} (\Lambda + \lambda I)^{-p} h \rangle_{L^2} \\ &= \sup_{h: \|h\|_{L^2} = 1} \langle T^{\alpha} (\Lambda + \lambda I)^{-p} h, (C - \hat{C}_n) T^{\alpha} (\Lambda + \lambda I)^{-p} h \rangle_{L^2}. \end{aligned}$$

Using the spectral representation of T and C, we get

$$\langle T^{\alpha}(\Lambda + \lambda I)^{-p}h, (C - \hat{C}_{n})T^{\alpha}(\Lambda + \lambda I)^{-p}h \rangle_{L^{2}}$$

$$= \left\langle \sum_{j \geq 1} \frac{\mu_{j}^{\alpha} \langle h, \phi_{j} \rangle_{L^{2}}}{(\mu_{j}\xi_{j} + \lambda)^{p}} \phi_{j}, \sum_{k \geq 1} \frac{\mu_{k}^{\alpha} \langle h, \phi_{k} \rangle_{L^{2}}}{(\mu_{k}\xi_{k} + \lambda)^{p}} (C - \hat{C}_{n})\phi_{k} \right\rangle_{L^{2}}$$

$$= \sum_{j,k \geq 1} \frac{\mu_{j}^{\alpha} \mu_{k}^{\alpha} \langle h, \phi_{j} \rangle_{L^{2}} \langle h, \phi_{k} \rangle_{L^{2}}}{(\mu_{j}\xi_{j} + \lambda)^{p} (\mu_{k}\xi_{k} + \lambda)^{p}} \langle \phi_{j}, (C - \hat{C}_{n})\phi_{k} \rangle_{L^{2}}$$

$$\leq \left(\sum_{j,k \geq 1} \frac{\mu_{j}^{2\alpha} \mu_{k}^{2\alpha}}{(\mu_{j}\xi_{j} + \lambda)^{2p} (\mu_{k}\xi_{k} + \lambda)^{2p}} \langle \phi_{j}, (C - \hat{C}_{n})\phi_{k} \rangle_{L^{2}}^{2} \right)^{\frac{1}{2}} \|h\|_{L^{2}},$$

where we applied Cauchy-Schwartz inequality in the last inequality. Therefore,

$$\mathbb{E} \| (\Lambda + \lambda I)^{-p} T^{\alpha} (C - \hat{C}_n) T^{\alpha} (\Lambda + \lambda I)^{-p} \|_{L^2 \to L^2}$$

$$\leq \left(\sum_{j,k \ge 1} \frac{\mu_j^{2\alpha} \mu_k^{2\alpha}}{(\mu_j \xi_j + \lambda)^{2p} (\mu_k \xi_k + \lambda)^{2p}} \mathbb{E} \langle \phi_j, (C - \hat{C}_n) \phi_k \rangle_{L^2}^2 \right)^{\frac{1}{2}},$$

which follows through an application of Jensen's inequality. Now we consider

$$\mathbb{E}\langle\phi_j, (C - \hat{C}_n)\phi_k\rangle_{L^2}^2 = \mathbb{E}\langle\phi_j, \frac{1}{n}\sum_{i=1}^n (C - X_i \otimes X_i)\phi_k\rangle_{L^2}^2$$
$$= \mathbb{E}\left[\frac{1}{n}\sum_{i=1}^n \langle\phi_j, (C - X_i \otimes X_i)\phi_k\rangle_{L^2}\right]^2 \le \frac{1}{n}\mathbb{E}\langle\phi_j, (X \otimes X)\phi_k\rangle_{L^2}^2$$
$$= \frac{1}{n}\mathbb{E}\langle X, (\phi_j \otimes \phi_k)X\rangle_{L^2}^2.$$

By Karhunen-Loève expansion, we have $X = \sum x_i \phi_i$, where $\mathbb{E}[x_i x_j] = \delta_{ij} \xi_i$. Let $\phi_j \otimes \phi_k =: \Gamma_{jk}$. Then we have

$$\mathbb{E}\langle X, (\phi_j \otimes \phi_k) X \rangle_{L^2}^2 = \mathbb{E}\langle X, \Gamma_{jk} X \rangle_{L^2}^2 = \mathbb{E}\left\langle \sum_{i \ge 1} x_i \phi_i, \sum_{l \ge 1} x_l \Gamma_{jk} \phi_l \right\rangle_{L^2}^2$$
$$= \sum_{i \ge 1} \mathbb{E}[x_i^4] \langle \phi_i, \Gamma_{jk} \phi_i \rangle_{L^2}^2 + \sum_{i,l \ge 1, i \ne l} \mathbb{E}[x_i^2] \mathbb{E}[x_l^2] \langle \phi_i, \Gamma_{jk} \phi_i \rangle_{L^2} \langle \phi_l, \Gamma_{jk} \phi_l \rangle_{L^2}$$
$$\lesssim \sum_{i \ge 1} (\mathbb{E}[x_i]^2)^2 \langle \phi_i, \Gamma_{jk} \phi_i \rangle_{L^2}^2 = \xi_j^2 \delta_{kj},$$

yielding

$$\mathbb{E}[\|(\Lambda + \lambda I)^{-p} T^{\alpha} (C - \hat{C}_{n}) T^{\alpha} (\Lambda + \lambda I)^{-p} \|_{L^{2} \to L^{2}}] \\ \lesssim \frac{1}{\sqrt{n}} \left(\sum_{j,k \ge 1} \frac{\mu_{j}^{2\alpha} \mu_{k}^{2\alpha} \xi_{j}^{2} \delta_{jk}}{(\mu_{j}\xi_{j} + \lambda)^{2p} (\mu_{k}\xi_{k} + \lambda)^{2p}} \right)^{\frac{1}{2}} \lesssim \frac{1}{\sqrt{n}} \left(\sum_{j \ge 1} \frac{j^{-4t\alpha} j^{-2c}}{(j^{-(t+c)} + \lambda)^{4p}} \right)^{\frac{1}{2}}$$

By Lemma A.10, for $2p(t+c) \ge 2\alpha t + c$, we get

$$\mathbb{E}[\|(\Lambda + \lambda I)^{-p} T^{\alpha} (C - \hat{C}_n) T^{\alpha} (\Lambda + \lambda I)^{-p}\|_{L^2 \to L^2}] \lesssim \frac{1}{\sqrt{n}} \lambda^{-\frac{1 + 4p(t+c) - (4\alpha t + 2c)}{2(t+c)}}.$$

The result follows from an application of Markov's inequality.

Lemma A.5. Suppose Assumptions 2, 4 and 5 hold. Then for $p \geq \frac{1}{2}$, we have

$$\|(\Lambda + \lambda I)^{-p} (\Lambda - \hat{\Lambda}_n) (\Lambda + \lambda I)^{-p} \|_{L^2 \to L^2} \lesssim_p \frac{1}{\sqrt{n}} \lambda^{-\frac{1 + 4bp - 2b}{2b}}.$$

Proof. Following similar steps as in Lemma A.4, we have

$$\mathbb{E} \| (\Lambda + \lambda I)^{-p} (\Lambda - \Lambda_n) (\Lambda + \lambda I)^{-p} \|_{L^2 \to L^2}$$

$$\lesssim \left(\sum_{j,k \ge 1} \frac{1}{(\tau_j + \lambda)^{2p} (\tau_k + \lambda)^{2p}} \mathbb{E} \langle e_j, (\Lambda - \hat{\Lambda}_n) e_k \rangle_{L^2}^2 \right)^{\frac{1}{2}},$$

where $(\tau_j, e_j)_j$ is the eigensystem of Λ . Note that

$$\mathbb{E}\langle e_j, (\Lambda - \hat{\Lambda}_n) e_k \rangle_{L^2}^2 = \frac{1}{n} \mathbb{E} \langle T^{\frac{1}{2}} X, (e_j \otimes e_k) T^{\frac{1}{2}} X \rangle_{L^2}^2.$$

Since $\mathbb{E}[T^{\frac{1}{2}}X \otimes T^{\frac{1}{2}}X] = \Lambda$, by Karhunen-Loève expansion, we have $T^{\frac{1}{2}}X = \sum x_i e_i$, where $\mathbb{E}[x_i x_j] = \delta_{ij} \tau_i$. Then using the similar ideas as in Lemma A.4, we get

$$\mathbb{E}\langle T^{\frac{1}{2}}X, (e_j \otimes e_k)T^{\frac{1}{2}}X\rangle_{L^2}^2 \lesssim \tau_j^2\delta_{jk}.$$

So we have that

$$\|(\Lambda + \lambda I)^{-p} (\Lambda - \hat{\Lambda}_n) (\Lambda + \lambda I)^{-p}\|_{L^2 \to L^2} \lesssim_p \frac{1}{\sqrt{n}} \left(\sum_j \frac{\tau_j^2}{(\tau_j + \lambda)^{4p}} \right)^{\frac{1}{2}}$$
$$\lesssim \frac{1}{\sqrt{n}} \left(\sum_j \frac{j^{-2b}}{(j^{-b} + \lambda)^{4p}} \right)^{\frac{1}{2}} \lesssim \frac{1}{\sqrt{n}} \lambda^{-\frac{1+4bp-2b}{2b}},$$

provided $p \geq \frac{1}{2}$.

Lemma A.6. Let $\delta \in (0,1)$ and $\lambda \ge c_{\delta}n^{-b}$ for some positive constant c_{δ} . Under Assumptions 4 and 5, with probability at least $1 - \delta$, the following holds:

- $\begin{array}{l} (a) \ \|(\hat{\Lambda}_n \Lambda)(\Lambda + \lambda I)^{-1}\|_{L^2 \to L^2} \leq \frac{1}{2}; \\ (b) \ \|(\Lambda + \lambda I)^l(\hat{\Lambda}_n + \lambda I)^{-l}\|_{L^2 \to L^2} \leq 2^l, \ \forall l \in [0, 1]; \\ (c) \ \|(\Lambda + \lambda I)^{-l}(\hat{\Lambda}_n + \lambda I)^l\|_{L^2 \to L^2} \leq (\frac{3}{2})^l, \ \forall l \in [0, 1]. \end{array}$

Proof. We start by considering

$$\begin{aligned} \|(\hat{\Lambda}_{n} - \Lambda)(\Lambda + \lambda I)^{-1}\|_{L^{2} \to L^{2}} &= \|(\Lambda + \lambda I)^{-1}(\hat{\Lambda}_{n} - \Lambda)\|_{L^{2} \to L^{2}} \\ &= \sup_{h,g: \|h\|_{L^{2}}, \|g\|_{L^{2}} = 1} |\langle (\Lambda + \lambda I)^{-\frac{1}{2}}g, (\Lambda + \lambda I)^{-\frac{1}{2}}(\hat{\Lambda}_{n} - \Lambda)h\rangle_{L^{2}}|. \end{aligned}$$

Using the fact that $h = \sum_{j \ge 1} \langle h, e_j \rangle_{L^2} e_j$ and $g = \sum_{j \ge 1} \langle g, e_j \rangle_{L^2} e_j$, we get

$$\langle (\Lambda + \lambda I)^{-\frac{1}{2}} g, (\Lambda + \lambda I)^{-\frac{1}{2}} (\hat{\Lambda}_n - \Lambda) h \rangle_{L^2}$$

$$= \left\langle \sum_{j \ge 1} \frac{\langle g, e_j \rangle_{L^2} e_j}{(\tau_j + \lambda)^{\frac{1}{2}}}, \sum_{k,l \ge 1} \frac{\langle h, e_k \rangle_{L^2} \langle e_l, (\hat{\Lambda}_n - \Lambda) e_k \rangle_{L^2} e_l}{(\tau_l + \lambda)^{\frac{1}{2}}} \right\rangle_{L^2}$$

$$= \sum_{j,k \ge 1} \frac{\langle g, e_j \rangle_{L^2} \langle h, e_k \rangle_{L^2}}{\tau_j + \lambda} \langle e_j, (\hat{\Lambda}_n - \Lambda) e_k \rangle_{L^2}.$$

Applying Cauchy–Schwartz inequality yields

$$\|(\Lambda+\lambda I)^{-1}(\hat{\Lambda}_n-\Lambda)\|_{L^2\to L^2} \le \left(\sum_{j,k\ge 1} \frac{1}{(\tau_j+\lambda)^2} \langle e_j, (\hat{\Lambda}_n-\Lambda)e_k \rangle_{L^2}^2\right)^{\frac{1}{2}}.$$

Now taking expectations on both sides and applying Jensen's Inequality, we have

$$\mathbb{E}\|(\Lambda+\lambda I)^{-1}(\hat{\Lambda}_n-\Lambda)\|_{L^2\to L^2} \le \left(\sum_{j,k\ge 1} \frac{1}{(\tau_j+\lambda)^2} \mathbb{E}\langle e_j, (\hat{\Lambda}_n-\Lambda)e_k\rangle_{L^2}^2\right)^{\frac{1}{2}}$$

From Lemma A.5, we can see that

$$\mathbb{E}\langle e_j, (\hat{\Lambda}_n - \Lambda) e_k \rangle_{L^2}^2 \lesssim \frac{1}{n} \tau_j^2 \delta_{jk}.$$

Therefore, an application of Markov's inequality yields

$$\|(\Lambda + \lambda I)^{-1}(\hat{\Lambda}_n - \Lambda)\|_{L^2 \to L^2} \lesssim_p \frac{1}{\sqrt{n}} \left(\sum_{j \ge 1} \frac{\tau_j^2}{(\tau_j + \lambda)^2}\right)^{\frac{1}{2}}$$
$$\lesssim \frac{1}{\sqrt{n}} \left(\sum_{j \ge 1} \frac{j^{-2b}}{(j^{-b} + \lambda)^2}\right)^{\frac{1}{2}} \lesssim \frac{1}{\sqrt{n}} \lambda^{-\frac{1}{2b}}$$

and the result in (a) follows under $\lambda \gtrsim n^{-b}$. As a consequence of part (a), we get

$$\begin{aligned} \| (\Lambda + \lambda I) (\hat{\Lambda}_n + \lambda I)^{-1} \|_{L^2 \to L^2} &= \| [(\hat{\Lambda}_n - \Lambda) (\Lambda + \lambda I)^{-1} + I]^{-1} \|_{L^2 \to L^2} \\ &\leq \frac{1}{1 - \| (\hat{\Lambda}_n - \Lambda) (\Lambda + \lambda I)^{-1} \|_{L^2 \to L^2}} \leq 2, \quad \forall \lambda \gtrsim n^{-b}, \end{aligned}$$

and part (b) follows from combining (a) with Lemma A.7 (Corde's inequality). For part (c), note that

$$\begin{aligned} \|(\Lambda + \lambda I)^{-1} (\hat{\Lambda}_n + \lambda I)\|_{L^2 \to L^2} &= \|I - (\Lambda - \hat{\Lambda}_n) (\Lambda + \lambda I)^{-1}\|_{L^2 \to L^2} \\ &\leq 1 + \|(\hat{\Lambda}_n - \Lambda) (\Lambda + \lambda I)^{-1}\|_{L^2 \to L^2} \\ &\leq \frac{3}{2}, \quad \forall \lambda \gtrsim n^{-b}, \end{aligned}$$

and applying Lemma A.7 (Corde's inequality) yields the result.

Lemma A.7. [10, Lemma 5.1] Suppose T_1 and T_2 are two positive bounded linear operators on a separable Hilbert space. Then

$$||T_1^p T_2^p|| \le ||T_1 T_2||^p$$
, when $0 \le p \le 1$.

Lemma A.8. [2, Lemma A.6] For any $0 < \alpha \leq \beta$,

$$\sup_{i\in\mathbb{N}}\left[\frac{i^{-\alpha}}{i^{-\beta}+\lambda}\right] \leq \lambda^{\frac{\alpha-\beta}{\beta}}, \ \forall \ \lambda > 0.$$

Lemma A.9. Let a, m, p, q and l be positive numbers. Then for $r = \min\{p, \frac{lq}{m} + a\}$ and a < p, we have

$$\sup_{i \in \mathbb{N}} \left[\frac{i^{-(p-a)m}}{(i^{-q}+\lambda)^l} \right] \le \lambda^{\frac{(r-a)m-ql}{q}}, \ \forall \ \lambda > 0.$$

Proof. Consider

$$\sup_{i\in\mathbb{N}}\left[\frac{i^{-(p-a)m}}{(i^{-q}+\lambda)^l}\right] = \left(\sup_{i\in\mathbb{N}}\left[\frac{i^{-(p-a)\frac{m}{l}}}{i^{-q}+\lambda}\right]\right)^l.$$

If $p \leq \frac{lq}{m} + a$, i.e., $(p-a)\frac{m}{l} \leq q$, then it follows from Lemma A.8 that

$$\sup_{i \in \mathbb{N}} \left[\frac{i^{-(p-a)m}}{(i^{-q} + \lambda)^l} \right] \le \lambda^{\frac{(p-a)m-ql}{q}}$$

If $p > \frac{lq}{m} + a$, i.e., $(p-a)\frac{m}{l} > q$, then

$$\sup_{i\in\mathbb{N}}\left[\frac{i^{-(p-a)m}}{(i^{-q}+\lambda)^l}\right] = \left(\sup_{i\in\mathbb{N}}\left[\frac{i^{-q}}{(i^{-q}+\lambda)}\right]\right)^l \left(\sup_{i\in\mathbb{N}}\left[i^{-\left(\frac{(p-a)m}{l}-q\right)}\right]\right)^l \le 1,$$

where the last step follows from Lemma A.8 and the fact that $\frac{(p-a)m}{l} - q > 0$. \Box Lemma A.10. For $\alpha > 1$, $\beta > 1$, and $q \ge \frac{\alpha}{\beta}$, we have

$$\sum_{i \in \mathbb{N}} \frac{i^{-\alpha}}{(i^{-\beta} + \lambda)^q} \lesssim \lambda^{-\frac{1+\beta q - \alpha}{\beta}}, \ \forall \ \lambda > 0.$$

Proof. Consider

$$\sum_{i\in\mathbb{N}}\frac{i^{-\alpha}}{(i^{-\beta}+\lambda)^q} = \sum_{i\in\mathbb{N}}\frac{i^{q\beta-\alpha}}{(1+\lambda i^\beta)^q} \lesssim \int_0^\infty \frac{x^{q\beta-\alpha}}{(1+\lambda x^\beta)^q} dx = \lambda^{-\frac{1+q\beta-\alpha}{\beta}} \int_0^\infty \frac{y^{q\beta-\alpha}}{(1+y^\beta)^q} dy.$$

Note that

$$\int_{0}^{\infty} \frac{y^{q\beta-\alpha}}{(1+y^{\beta})^{q}} dy = \int_{0}^{\infty} \frac{(y^{\beta})^{q-\frac{\alpha}{\beta}}}{(1+y^{\beta})^{q-\frac{\alpha}{\beta}}(1+y^{\beta})^{\frac{\alpha}{\beta}}} dy \le \int_{0}^{\infty} \frac{1}{(1+y^{\beta})^{\frac{\alpha}{\beta}}} dy = \int_{0}^{1} \frac{1}{(1+y^{\beta})^{\frac{\alpha}{\beta}}} dy + \int_{1}^{\infty} \frac{1}{(1+y^{\beta})^{\frac{\alpha}{\beta}}} dy.$$

For $y \ge 1$, $\frac{1}{1+y^{\beta}} \le \frac{1}{y^{\beta}}$ and therefore, $\int_{1}^{\infty} \frac{1}{(1+y^{\beta})^{\frac{\alpha}{\beta}}} dy \le \int_{1}^{\infty} y^{-\alpha} dy = \frac{1}{\alpha-1}$. On the other hand, $\int_{0}^{1} \frac{1}{(1+y^{\beta})^{\frac{\alpha}{\beta}}} dy \le 1$. The result therefore follows.

(N. Gupta) Department of Mathematics, Indian Institute of Technology Delhi, India $\mathit{Email}\ address:\ \texttt{ngupta.maths}@gmail.com$

(S. Sivananthan) Department of Mathematics, Indian Institute of Technology Delhi, India *Email address*: siva@maths.iitd.ac.in

(B. K. Sriperumbudur) Department of Statistics, Pennsylvania State University, USA $\mathit{Email}\ address:\ bks180psu.edu$