ON THE UNIQUENESS OF BEST APPROXIMATION IN ORLICZ SPACES

ANA BENAVENTE¹ O, JUAN COSTA PONCE², AND SERGIO FAVIER³ O

ABSTRACT. We study uniqueness of best approximation in Orlicz spaces L^{Φ} , for different types of convex functions Φ and for some finite dimensional approximation classes of functions, where Tchebycheff spaces, and more general approximation ones, are involved.

1. Introduction and notations

Let \Im be the class of all non decreasing functions φ defined for all real numbers $t \geq 0$, such that $\varphi(t) > 0$ if t > 0, and consider Ψ the class of all convex functions Φ defined by $\Phi(x) = \int_0^x \varphi(t) dt$, for $x \geq 0$, with $\varphi \in \Im$. We assume a Δ_2 condition for the functions Φ , which means that there exists a constant $\Lambda > 0$ such that $\Phi(2x) \leq \Lambda \Phi(x)$, for $x \geq 0$. We also denote by φ^- and φ^+ the left and right derivatives of Φ respectively.

Let $\Phi \in \Psi$ and let (Ω, \mathcal{A}, m) be the Lebesgue measure space where $\Omega \subset \mathbb{R}$ is a bounded set. We denote by $L^{\Phi} = L^{\Phi}(\Omega, \mathcal{A}, m)$ the Orlicz space given by the class of all \mathcal{A} -measurable functions f defined on Ω such that $\int_{\Omega} \Phi(|f|) dx < \infty$.

Given a set $S \subset L^{\Phi}$, an element $P \in S$ is called a best Φ -approximation of $f \in L^{\Phi}$ from the approximation class S if and only if

$$\int_{\Omega} \Phi(|f - P|) dx = \inf_{Q \in S} \int_{\Omega} \Phi(|f - Q|) dx$$

and, in this case, we write $P \in \mu_{\Phi}(f/S)$. The mapping $\mu_{\Phi} : L^{\Phi} \to S$ is called the *best* Φ -approximation operator given S.

Research partially supported CONICET, Universidad Nacional de San Luis.

 $^{2020\ \}textit{Mathematics Subject Classification}.\ \text{Primary 46E30}; \ \text{Secondary 41A10},\ 41\text{A50}.$

Key words and phrases. Orlicz Spaces, Best Approximation, Characterization of Best Approximation Operators.

Given a n-dimensional linear space $S_n \subseteq C([a,b])$, the existence of a best approximation is assured by Theorem 2.3 in [3]. Recall that S_n is called a Tchebycheff space if any non zero element in S_n has at most n-1 zeros in [a,b].

Uniqueness of best approximation was studied since 1924, for example in [7], and this problem was extensively developed in L^1 in the 70's. Conditions which assure uniqueness of best L^1 approximation have been studied, for instance for Galkin in [6] and Strauss [14] showed uniqueness in L^1 by polynomial splines. In [9] Micchelli considered best approximation by weak Tchebychev subspaces. For a very mentioned reference of this problem we also have to cite to DeVore [5], Kroo [8], Pinkus [10] and Strauss [14].

We point out that for Orlicz spaces $L^{\Phi}([a,b])$ it is easy to obtain uniqueness of best Φ - approximation for a given strictly convex function Φ . In this paper we present first a characterization theorem of best Φ - approximation which will be used in the sequel. Next, we get uniqueness of the best Φ - approximation, for a continuous function defined in a real interval [a,b], where we consider a general function Φ and where the approximation class is a Tchebycheff space which generalizes the classical result in L^1 , setted for example in [12]. Next we get uniqueness of the best Φ - approximation for more general approximation classes, considering suitable classes of functions Φ . We set also a sufficient and necessary condition to assure that the best approximation of a continuous function is unique, which is an extension, in some way, to Theorem 22 of [4], stablished in the L^1 space.

2. Characterization and properties of the best Φ -approximations.

In this section, we will set a characterization property of the best Φ - approximations, when the approximation class is a finite dimensional vector space S_n , which will be useful to get uniqueness of best Φ - approximation for continuous functions. On the other hand, it is well known that uniqueness results follows straight forward if Φ is a strictly convex function. We deal with a more general convex functions Φ , where the strictly convexity is not required.

At the end of this section, we will present a property for the not strictly convex functions that will be used in the study of the uniqueness for a suitable class of approximation functions.

The following characterization is a generalization of Theorem 2.2 in [1] for the case $\varphi^+(0) > 0$ and Theorem 2.2 in [2] for the non derivative case of Φ .

Theorem 2.1. Let $\Phi \in \Psi$ and $f \in L^{\Phi}(\Omega)$. Then $P \in \mu_{\Phi}(f/S_n)$ if and only if

$$\int_{\{Q>0, f>P\}\cup\{Q<0, f< P\}} \varphi^{-}(|f-P|)|Q| dx - \int_{\{Q<0, f>P\}\cup\{Q>0, f< P\}} \varphi^{+}(|f-P|)|Q| dx
\leq \varphi^{+}(0) \int_{\{f=P\}} |Q| dx,$$

for any $Q \in S_n$.

Proof. For $P \in \mu_{\Phi}(f/S_n)$, $Q \in S_n$ with $Q \neq P$ and $\epsilon \geq 0$, we define

$$F_Q(\epsilon) = \int_{\Omega} \Phi(|f - (P + \epsilon Q)|) dx.$$

Using the convexity of Φ , we have that F_Q is also a convex function on $[0, \infty)$, then

$$F_{Q}(a\epsilon_{1} + b\epsilon_{2}) = \int_{\Omega} \Phi\left(\left|(a+b)f - ((a+b)P + (a\epsilon_{1} + b\epsilon_{2})Q)\right|\right) dx$$

$$\leq \int_{\Omega} \Phi\left(a\left|f - (P + \epsilon_{1}Q)\right| + b\left|f - (P + \epsilon_{2}Q)\right|\right) dx$$

$$\leq a \int_{\Omega} \Phi\left(\left|f - (P + \epsilon_{1}Q)\right|\right) dx + b \int_{\Omega} \Phi\left(\left|f - (P + \epsilon_{2}Q)\right|\right) dx$$

$$= aF_{Q}\left(\epsilon_{1}\right) + bF_{Q}\left(\epsilon_{2}\right),$$

for ϵ_1 , $\epsilon_2 \geq 0$, and a+b=1. It follows that $F_Q(0)=\min_{[0,\infty)}F_Q(\epsilon)$, and the equality holds if, and only if, $0 \leq F_Q^+(0)$, where $F_Q^+(0)$ is the right derivative of F_Q in $\epsilon=0$. Now we compute the derivative.

$$F_Q^+(0) = \lim_{\epsilon \to 0^+} \frac{F_Q(0+\epsilon) - F_Q(0)}{\epsilon}$$

$$= \lim_{\epsilon \to 0^+} \frac{1}{\epsilon} \left\{ \int_{\Omega} \Phi(|f - (P + \epsilon Q)|) dx - \int_{\Omega} \Phi(|f - P|) dx \right\}$$

$$= \lim_{\epsilon \to 0^+} \int_{\Omega \cap \{Q \neq 0\}} \left[\frac{\Phi(|f - (P + \epsilon Q)|) - \Phi(|f - P|)}{\epsilon Q} \right] Q dx.$$

To analyze each absolute value in the argument of Φ , we split the set $\Omega \cap \{Q \neq 0\}$ in seven cases, which will yield the following seven integrals:

$$F_{Q}^{+}(0) = \lim_{\epsilon \to 0^{+}} \left\{ \int_{\{Q > 0, \ P < P + \epsilon Q < f\}} \left[\frac{\Phi(f - P - \epsilon Q) - \Phi(f - P)}{\epsilon Q} \right] Q \, dx \right.$$

$$+ \int_{\{Q < 0, \ P + \epsilon Q < P < f\}} \left[\frac{\Phi(f - P - \epsilon Q) - \Phi(f - P)}{\epsilon Q} \right] Q \, dx$$

$$+ \int_{\{Q < 0, \ P + \epsilon Q < f < P\}} \left[\frac{\Phi(f - P - \epsilon Q) - \Phi(P - f)}{\epsilon Q} \right] Q \, dx$$

$$+ \int_{\{Q < 0, \ P + \epsilon Q < f = P\}} \left[\frac{\Phi(f - P - \epsilon Q) - \Phi(P - f)}{\epsilon Q} \right] Q \, dx$$

$$+ \int_{\{Q > 0, \ f \le P < P + \epsilon Q\}} \left[\frac{\Phi(P + \epsilon Q - f) - \Phi(P - f)}{\epsilon Q} \right] Q \, dx$$

$$+ \int_{\{Q > 0, \ P < f \le P + \epsilon Q < P\}} \left[\frac{\Phi(P + \epsilon Q - f) - \Phi(P - f)}{\epsilon Q} \right] Q \, dx$$

$$+ \int_{\{Q < 0, \ f \le P + \epsilon Q < P\}} \left[\frac{\Phi(P + \epsilon Q - f) - \Phi(P - f)}{\epsilon Q} \right] Q \, dx$$

that is, $F_Q^+(0) = \lim_{\epsilon \to 0^+} \{ I_1 + I_2 + I_3 + I_4 + I_5 + I_6 + I_7 \}.$

By the convexity of Φ , for $0 < \epsilon \le 1$ we have:

(2)
$$\frac{|Q||\Phi(|f - (P + \epsilon Q)|) - \Phi(|f - P|)|}{\epsilon |Q|} \le |Q|(\varphi^{+}(|f - P| + |Q|) + \varphi^{+}(|f - P|)).$$

The function to the right of the inequality (2), is integrable in Ω because $\varphi^+ \in \Delta_2$, $|P| \leq ||P||_{\infty}$, $|Q| \leq ||Q||_{\infty}$ and $|\Omega| < \infty$. By the Dominated Convergence theorem we conclude that:

$$\lim_{\epsilon \to 0^+} I_1 = -\int_{\{Q > 0, \ f > P\}} \varphi^-(|f - P|) Q \, dx,$$

$$\lim_{\epsilon \to 0^+} I_2 = -\int_{\{Q < 0, f > P\}} \varphi^+(|f - P|) Q \, dx,$$

$$\lim_{\epsilon \to 0^{+}} I_{4} = \lim_{\epsilon \to 0^{+}} \int_{\{Q < 0, \ P + \epsilon Q < f = P\}} \left[\frac{\Phi(0) - \Phi(0 + \epsilon |Q|)}{\epsilon |Q|} \right] Q \, dx$$
$$= \varphi^{+}(0) \int_{\{Q < 0, \ f = P\}} |Q| \, dx,$$

$$\lim_{\epsilon \to 0^+} I_5 = \int_{\{Q > 0, \ f \leqslant P\}} \varphi^+(|f - P|)Q \, dx$$

$$= \int_{\{Q > 0, \ f < P\}} \varphi^+(|f - P|)Q \, dx + \varphi^+(0) \int_{\{Q > 0, \ f = P\}} |Q| dx$$

and

$$\lim_{\epsilon \to 0^+} I_7 = \int_{\{Q < 0, \ f < P\}} \varphi^-(|f - P|) Q \, dx.$$

For the remaining integrals, let us consider a decreasing and convergent sequence $\{\epsilon_n\} \searrow 0$, and the decreasing sets: $A_n := \{Q < 0, \ P + \epsilon_n Q < f < P\}$, then $\bigcap_{n=1}^{\infty} A_n = \emptyset$ and:

$$\lim_{n \to \infty} \int_{A_n} \left[\frac{\Phi(f - P - \epsilon_n Q) - \Phi(P - f)}{\epsilon_n Q} \right] Q dx \le \lim_{n \to \infty} \int_{A_n} \varphi^+(|f - P| + |Q|) + \varphi^+(|f - P|) dx$$

$$= \lim_{n \to \infty} \mu(A_n),$$

and
$$\lim_{n\to\infty} \mu(A_n) = \mu\left(\bigcap_{n=1}^{\infty} A_n\right) = \mu(\emptyset) = 0$$
. Then $\lim_{\epsilon\to 0^+} I_3 = 0$.

Now, let us consider the sets $B_n = \{Q > 0, P < f \leq P + \epsilon_n Q\}$, then $\bigcap_{n=1}^{\infty} B_n = \emptyset$ and again, by the Dominated Convergence Theorem we have $\lim_{\epsilon \to 0^+} I_6 = 0$.

Finally, replacing all the results in (1), we obtain

$$\begin{split} F_Q^+(0) &= -\int_{\{Q>0,\ f>P\}} \varphi^-(|f-P|)Q\,dx - \int_{\{Q<0,\ f>P\}} \varphi^+(|f-P|)Q\,dx, \\ &+ \int_{\{Q>0,\ f< P\}} \varphi^+(|f-P|)Q\,dx + \varphi^+(0) \int_{\{f=P\}} |Q|dx \\ &+ \int_{\{Q<0,\ f< P\}} \varphi^-(|f-P|)Q\,dx. \end{split}$$

As $F_Q^+(0) \geqslant 0$, we conclude the statement of the theorem.

For the next result we use the following notations. For a real number x we set $sgn(x) = \frac{x}{|x|}$, $x \neq 0$, and sign(0) = 0.

Remark 2.2. If $\Phi(x) = \int_0^x \varphi(t) dt$ is a derivable function then $\varphi^+ = \varphi^- = \varphi$, thus the last result becomes: $P \in \mu_{\Phi}(f/S_n)$ if and only if

$$\left| \int_{\Omega} \varphi(|f - P|) sgn(f - P) Q \, dx \right| \le \varphi(0^+) \int_{\{f = P\}} |Q| dx,$$

for each $Q \in S_n$.

The following result generalizes the Haar's L_1 -version theorem in [7].

Lemma 2.3. Let $\Phi \in \Psi$ and S_n be an n-dimensional linear space of continuous functions in almost every point in the interval [a,b]. If $P_1, P_2 \in \mu_{\Phi}(f/S_n)$ for $f \in L^{\Phi}$, then

$$[f(x) - P_1(x)][f(x) - P_2(x)] \ge 0$$

for almost every $x \in [a, b]$.

Proof. Let's define the functions $g_1(x) := f(x) - P_1(x)$, $g_2(x) := f(x) - P_2(x)$ and the number $\rho := \inf_{Q \in S_n} \int_a^b \Phi(|f - Q|) dx$. We observe that $\rho = \int_a^b \Phi(|g_1|) dx = \int_a^b \Phi(|g_2|) dx$. Using that Φ is a non-decreasing and convex function, we have:

$$\int_{a}^{b} \Phi(|f - \frac{P_1 + P_2}{2}| dx \le \frac{1}{2} \int_{a}^{b} \Phi(|g_1|) dx + \frac{1}{2} \int_{a}^{b} \Phi(|g_2|) dx = \rho.$$

Then $\frac{P_1+P_2}{2}$ is also a best Φ -approximation for f and

$$\int_{a}^{b} \Phi\left(\frac{|g_1 + g_2|}{2}\right) dx = \int_{a}^{b} \frac{\Phi(|g_1|)}{2} dx + \int_{a}^{b} \frac{\Phi(|g_2|)}{2} dx.$$

Which means that

(3)
$$\Phi\left(\frac{|g_1(x) + g_2(x)|}{2}\right) = \frac{\Phi(|g_1(x)|)}{2} + \frac{\Phi(|g_2(x)|)}{2}$$

in almost every $x \in [a, b]$.

That is because Φ is continuous and $\frac{\Phi(|g_1(x)|)}{2} + \frac{\Phi(|g_2(x)|)}{2} - \Phi\left(\frac{|g_1(x)+g_2(x)|}{2}\right) \geq 0$ in [a,b]. If for some set A with positive measure, $g_1(x)$ and $g_2(x)$ have different signs, then $|g_1(x)| + |g_2(x)| + |g_2(x)| + |g_2(x)|$ in A, and since Φ is a increasing and convex function, it satisfies $\Phi(\frac{|g_1(x)+g_2(x)|}{2}) < \Phi(\frac{|g_1(x)+|g_2(x)|}{2}) \leq \frac{\Phi(|g_1(x)|+\Phi(|g_2(x)|)}{2}$. But this last inequality contradicts (3), and the proof is then complete.

Recall that a function $\Phi:[0,\infty)\to\mathbb{R}$ is *convex* in an interval I if for all $x,y\in I$ and $0\leq\lambda\leq 1$, then $\Phi(\lambda x+(1-\lambda)y)\leq\lambda\Phi(x)+(1-\lambda)\Phi(y)$. And Φ is a *strictly convex* function in I if for all $x\neq y\in I$ and $0<\lambda<1$, then $\Phi(\lambda x+(1-\lambda)y)<\lambda\Phi(x)+(1-\lambda)\Phi(y)$. We observe that linear functions are convex, but not strictly convex. On the other hand, the following lemma shows that a convex, but not strictly convex function, is linear in some sub interval.

Lemma 2.4. Let $\Phi \in \Psi$. If Φ is not strictly convex in an interval I, then there exists an interval $J \subset I$ such that Φ is a straight line in J.

Proof. If Φ is not strictly convex, then there exists an interval $J = (x_1, x_3) \subset I$ and a convex combination of x_1 and x_3 called x_2 , such that the point $(x_2, \Phi(x_2))$ is in the line between the points $(x_1, \Phi(x_1))$ and $(x_3, \Phi(x_3))$, where the equation for this line is

(4)
$$\Phi(x) = \frac{\Phi(x_3) - \Phi(x_2)}{x_3 - x_2} (x - x_3) + \Phi(x_3).$$

If there is an $a \in (x_1, x_2)$ such that $\Phi(a) < \frac{\Phi(x_3) - \Phi(x_2)}{x_3 - x_2}(a - x_3) + \Phi(x_3)$, then $\frac{\Phi(x_3) - \Phi(a)}{x_3 - a} > \frac{\Phi(x_3) - \Phi(x_2)}{x_3 - x_2}$, and we have that $\Phi(x_2) > \frac{\Phi(x_3) - \Phi(a)}{x_3 - a}(x_2 - x_3) + \Phi(x_3)$. This inequality means that x_2 is a convex combination between a and x_3 such that $\Phi(x_2)$ is not in the segment between the points $(a, \Phi(a))$ and $(x_3, \Phi(x_3))$; which contradicts the convexity of Φ . A similar argument is used if $a \in (x_2, x_3)$. So (4) is satisfied for all x in y.

3. Uniqueness Results

Next we give uniqueness results of best Φ — approximation, for a suitable convex function Φ , which is non strictly convex and, according to Lemma 2.4, it is a straight line in some interval. In each case a convenient approximation class is considered.

Theorem 3.1. Let $\Phi \in \Psi$, $S_n \subseteq C([a,b])$ be a Tchebycheff space in [a,b] and f be a continuous function in [a,b]. Then there exists a unique best Φ -approximation of f in [a,b] from the class S_n .

Proof. Since the approximation class is finite dimensional, the existence of a best approximation, say P_1 , is assured by theorem 2.3 in [3]. It remains to prove the uniqueness of the best Φ - approximation function. Suppose that $f \notin S_n$ and for each $Q \in S_n$, consider the set $Z(f-Q) := \{x \in [a,b] : f(x) - Q(x) = 0\}$.

Suppose that $Z(f - P_1)$ has measure zero, then by Theorem 2.1 we conclude

$$\int_{\{Q>0, f>P_1\}} \varphi^-(|f-P_1|)sgn(f-P_1)Q \, dx + \int_{\{Q<0, f< P_1\}} \varphi^-(|f-P_1|)sgn(f-P_1)Q \, dx + \int_{\{Q<0, f> P_1\}} \varphi^+(|f-P_1|)sgn(f-P_1)Q \, dx + \int_{\{Q>0, f< P_1\}} \varphi^+(|f-P_1|)sgn(f-P_1|)Q \, dx + \int_{\{Q>0, f< P_1\}} \varphi^+(|f-P_1|)ggn(f-P_1|)Q \, dx + \int_{\{Q>0, f< P_1\}} \varphi^+(|f-P_1|)ggn(f-P_1|)Q \, dx + \int_{\{Q>0, f< P_1\}} \varphi^+(|f-P_1|)ggn(f-P_1|)Q \, dx + \int_{\{Q>0, f< P_1\}} \varphi^+(|f-$$

for every $Q \in S_n$.

Note that S_n has a basis $\{\delta_i\}_{i=1}^n$ such that for m < n, the set $\{\delta_i\}_{i=1}^m$ generate also a Tchebycheff space (see Theorem 2.29 in [13]).

We get that the function $sign[f(x) - P_1(x)]$ has at least n sign changes. In fact, suppose it has only m < n sign changes at the n points $x_1 < ... < x_m$ in (a, b). Then there exists a function $P_m \in S_n$, $P_m \neq 0$ that changes sign only in x_i , for i = 1, ..., m (see Proposition 2, page 195, in [11]). It follows that the function $P_m sign[f - P_1] \geq 0$ in [a, b].

Since φ^+ and φ^- are positive for x > 0, we have

$$0 = \int_{\{P_m > 0, f > P_1\}} \varphi^-(|f - P_1|) sgn(f - P_1) P_m dx$$

$$= \int_{\{P_m < 0, f < P_1\}} \varphi^-(|f - P_1|) sgn(f - P_1) P_m dx$$

$$= \int_{\{P_m < 0, f > P_1\}} \varphi^+(|f - P_1|) sgn(f - P_1) P_m dx$$

$$= \int_{\{P_m > 0, f < P_1\}} \varphi^+(|f - P_1|) sgn(f - P_1) P_m dx$$

Thus we conclude $f = P_1$ almost everywhere in [a, b], we get a contradiction.

Then, the function $sgn[f(x) - P_1(x)]$, has at least n sign changes. Now, suppose that there exists P_2 , another best Φ -approximation for f in [a, b]. By Lemma 2.3, $f(x) - P_2(x)$ is zero at the n-points where $f(x) - P_1(x)$ changes signs. Then $P_1(x) - P_2(x)$ has n-zeros, so $P_1 = P_2$.

Now, suppose that for any $P_1 \in \mu_{\Phi}(f/S_n)$ we have $m(Z(f-P_1)) > 0$ and consider another $P_2 \in \mu_{\Phi}(f/S_n)$. By the convexity of Φ , we have for $\lambda \in [0,1]$:

$$\int_{[a,b]} \Phi(|f - \lambda P_1 - (1 - \lambda)P_2|) dx \le \lambda \int_{[a,b]} \Phi(|f - P_1|) dx + (1 - \lambda) \int_{[a,b]} \Phi(|f - P_2|) dx$$

then $\lambda P_1 + (1-\lambda)P_2 \in \mu_{\Phi}(f/S_n)$. Suppose that for each $\lambda \in [0,1]$: $m(Z(f-\lambda P_1 - (1-\lambda)P_2)) > 0$. Then there exists $\lambda_1, \ \lambda_2 \in [0,1]$ such that the sets of zeros $Z(f-\lambda_1P_1 - (1-\lambda_1)P_2)$ and $Z(f-\lambda_2P_1 - (1-\lambda_2)P_2)$ have intersection with positive measure (see Lemma 4-7, page 109 in [12]). In particular, the intersection has n-points. Then $P_1 = P_2$ and this concludes the proof.

Next we consider wider approximation classes: the 1-space and the θ -space.

Definition 3.2. We say that the n-dimensional set $S_n \subset C([a,b])$ is a 1-space if there exists $h \in S_n$ such that h(x) > 0 for every $x \in [a,b]$, and if $P_1, P_2 \in S_n$ with $P_1 \neq P_2$ then $|\{P_1 = P_2\}| = 0$. If S_n satisfies only the last condition, we say that S_n is a 0-space.

Lemma 3.3. Let S_n a 1-space and $P \in \mu_{\Phi}(f/S_n)$. Then there exits $x \in [a,b]$ such that f(x) = P(x).

Proof. We assume P(x) - f(x) > 0, for every $x \in [a, b]$ and consider $h \in S_n$, h(x) > 0 for $x \in [a, b]$. Then, for $\varepsilon > 0$ such that $\varepsilon < \frac{\min_{x \in [a, b]} P(x) - f(x)}{\max_{x \in [a, b]} h(x)}$, we have

$$\varepsilon h(x) < P(x) - f(x), \ x \in [a, b].$$

Then

$$0 < \int_{[a,b]} \Phi(P - \varepsilon h - f) \, dx < \int_{[a,b]} \Phi(P - f) \, dx,$$

which is a contradiction since $P - \varepsilon h \in S_n$

Now we get the following uniqueness result when S_n is a 1-space.

Theorem 3.4. Let $\varphi \in \Im$ which is also a strictly increasing function in [0,b], b > 0, and $\Phi(x) = \int_0^x \varphi(t) dt$. If S_n is a 1-space, then for every $f \in C([a,b])$ the best approximation set $\mu_{\Phi}(f/S_n)$ is a singleton.

Proof. Suppose $P_1, P_2 \in \mu_{\Phi}(f/S_n)$, with $P_1 \neq P_2$. Now by Lemma 3.3 there exists an interval I such that $|f - P_1|(x) \leq b$, for every $x \in I$, and then the set $J = I \cap \{x \in [a, b] : P_1(x) \neq P_2(x)\}$ has positive measure. Then, from the strictly convexity of Φ in [0, b] and taking into account Lemma 2.3 we get

$$\int_{J} \Phi(|f - (\frac{P_1 + P_2}{2})|) \, dx < \frac{1}{2} \int_{J} \Phi(|f - P_1|) \, dx + \frac{1}{2} \int_{J} \Phi(|f - P_2|) \, dx$$

and then, since the convexity of Φ , we obtain

$$\int_{[a,b]} \Phi(|f - (\frac{P_1 + P_2}{2})|) \, dx < \frac{1}{2} \int_{[a,b]} \Phi(|f - P_1|) \, dx + \frac{1}{2} \int_{[a,b]} \Phi(|f - P_2|) \, dx,$$

which is a contradiction since $\frac{P_1+P_2}{2} \in S_n$.

For the next result we consider a n-dimensional 0-space $S_n \subseteq C([a,b])$ and we give a necessary and sufficient condition on the approximation class S_n which assures the uniqueness of $\mu_{\Phi}(f/S_n)$ for convex function Φ , which is a strictly convex function just for x grater than some positive real number. This result follows the same lines considered in [4] for L^1 and it is a generalization to Orlicz spaces.

Definition 3.5. The set $Z(f) := \{x \in [a,b] : f(x) = 0\}$ is called a γ -set if $0 \in \mu_{\Phi}(f/S_n)$.

Theorem 3.6. Consider the function Φ such that for k, c > 0 and $x \in [0, c]$, $\Phi(x) = kx$, and Φ is a strictly convex function for $x \in (c, \infty)$. Let S_n be a n-dimensional 0-space. Then, for every $f \in C([a, b]) \cap L^{\Phi}([a, b])$ the set $\mu_{\Phi}(f/S_n)$ has an unique element P_1 , if and only if

- a) The constant function 0 is the unique element of S_n which is 0 on a $\gamma-$ set or
- b) For $P_1 \in \mu_{\Phi}(f/S_n)$, the set $\{x \in [a,b] : |f(x) P_1(x)| > c\}$ has positive measure for every $f \in C([a,b]) \cap L^{\Phi}([a,b])$.

Proof. Suppose P_1, P_2 are two different elements in $\mu_{\Phi}(f/S_n)$. Due to the convexity of Φ , we have

$$\int_{[a,b]} \Phi(|f - \frac{P_1 + P_2}{2}|) \, dx = \frac{1}{2} \int_{[a,b]} \Phi(|f - P_1|) \, dx + \frac{1}{2} \int_{[a,b]} \Phi(|f - P_2|) \, dx.$$

Then we get

$$\Phi(|f(x) - \frac{P_1 + P_2}{2}(x)|) = \frac{1}{2}\Phi(|f(x) - P_1(x)|) + \frac{1}{2}\Phi(|f(x) - P_2(x)|), \quad x \in [a, b].$$

Thus for any $x \in Z(f - \frac{P_1 + P_2}{2})$ we have $f(x) - P_1(x) = f(x) - P_2(x)$, and then $P_1(x) - P_2(x) = 0$, and this is a contradiction of a).

Now assuming condition b) and suppose P_1 , $P_2 \in \mu_{\Phi}(f/S_n)$, $P_2 \neq P_1$, then we have m(I) > 0, for

$$I = \{x \in [a, b] : |f(x) - P_1(x)| > c\} \cap \{x \in [a, b] : P_1(x) \neq P_2(x)\},\$$

thus

$$\int_{I} \Phi(|f - (\frac{P_1 + P_2}{2})|) \, dx < \frac{1}{2} \int_{I} \Phi(|f - P_1|) \, dx + \frac{1}{2} \int_{I} \Phi(|f - P_2|) \, dx,$$

and then we have a contradiction.

On the other hand, suppose there exists $P_3 \in S_n$, $P_3 \neq 0$ and $0 \leq |P_3| \leq \frac{c}{2}$, such that $Z(f - P_1) \subseteq Z(P_3)$, $P_1 \in \mu_{\Phi}(f/S_n)$ and assume $0 \leq |f - P_1|(x) \leq c$, a. e. x.

Now, for $h(x) = |P_3(x)| sgn(f - P_1)(x)$, we have

$$\int_{[a,b]\cap\{f\neq P_1\}} \varphi(|h|)sgn(h)Q\,dx = \int_{[a,b]\cap\{f\neq P_1\}} \varphi(|h|)sgn(f-P_1)Q\,dx$$

and since $\varphi(|h|) = \varphi(|f - P_1|) = k$, we have

$$\int_{[a,b]\cap\{f\neq P_1\}} \varphi(|h|) sgn(h) Q \, dx = \int_{[a,b]\cap\{f\neq P_1\}} \varphi(|f-P_1|) sgn(f-P_1) Q \, dx,$$

and since $P_1 \in \mu_{\Phi}(f/S_n)$ and $Z(f-P_1) \subseteq Z(h)$, we get

$$\int_{[a,b]\cap\{h\neq 0\}} \varphi(|h|) \operatorname{sgn}(h) Q \, dx \le \varphi(0) \int_{Z(h)} |Q| \, dx,$$

which implies, since 2.2 that $0 \in \mu_{\Phi}(h/S_n)$.

Now

$$\int_{[a,b]} \Phi(|h - \varepsilon P_3|) \, dx = \int_{[a,b]} \Phi((h - \varepsilon P_3) \, sgn(h - \varepsilon P_3) \, dx,$$

and taking into account $\Phi(x) = kx$, for $x \leq c$ we get

$$\int_{[a,b]} \Phi(|h - \varepsilon P_3|) dx = \int_{[a,b]} k h \operatorname{sgn}(h - \varepsilon P_3) dx - \varepsilon \int_{[a,b]} k P_3 \operatorname{sgn}(h - \varepsilon P_3) dx,$$

and then, for $x \in [a, b]$, $h(x) - \varepsilon P_3(x) \neq 0$, it holds $sgn(h - \varepsilon P_3)(x) = sgn(f - P_1)(x)$, thus

$$\int_{[a,b]} \Phi(|h-\varepsilon P_3|) dx = \int_{[a,b]} k h \operatorname{sgn}(f-P_1) dx - \varepsilon \int_{[a,b]} k P_3 \operatorname{sgn}(f-P_1) dx.$$

Now, we use remark 2.2 to obtain

$$-\varepsilon \int_{[a,b]} k P_3 \operatorname{sgn}(f - P_1) dx \le \varepsilon \varphi(0) \int_{Z(f - P_1)} k |P_3| dx,$$

and then

$$\int_{[a,b]} \Phi(|h - \varepsilon P_3|) \, dx \le \int_{[a,b]} k \, h \, sgn(f - P_1) \, dx + \varepsilon \varphi(0) \int_{Z(f - P_1)} k \, |P_3| \, dx.$$

Thus, since $Z(f - P_1) \subseteq Z(P_3)$, we have $\int_{Z(f-P_1)} |P_3| dx = 0$, then

$$\int_{[a,b]} \Phi(|h - \varepsilon P_3|) \, dx \le \int_{[a,b]} k \, h \, sgn(f - P_1) \, dx = \int_{[a,b]} k \, h \, sgn(h) \, dx = \int_{[a,b]} \Phi(|h|) \, dx,$$

which implies that $\varepsilon P_3 \in \mu_{\Phi}(h/S_n)$, for every ε , $0 < \varepsilon < 1$.

Finally, we set an uniqueness result of $\mu_{\Phi}(f/S_n)$, where a specific convex function Φ and a suitable 1-space S_n are considered.

Theorem 3.7. Let $\Phi \in \Psi$, $\Phi(x) = \int_0^x \varphi(t) dt$, where $\varphi \in \Im$ is non continuous in a decreasing positive sequence a_n , which converges to 0. Let S_n be a 1-space such that every nonzero $P \in S_n$ has only a finite amount of zeros on [a,b]. Then for every $f \in C([a,b])$ the best approximation set $\mu_{\Phi}(f/S_n)$ is a singleton.

Proof. Suppose that $P_1, P_2 \in \mu_{\Phi}(f/S_n)$ with $P_1 \neq P_2$. Since $\int_a^b \Phi(|f-P_1|) - \Phi(|f-P_2|) dx = 0$, the function $\Phi(|f-P_1|) - \Phi(|f-P_2|)$ cannot be positive in [a,b]. Then, since $\Phi(0) = 0$ and Lemma 2.3, there exists $x_0 \in [a,b]$ such that $f(x) - P_1(x_0) = f(x) - P_2(x_0)$.

Actually, x_0 is a zero of $P_1 - P_2$, which is a nonzero element of S_n . Therefore, x_0 is an isolated point in [a,b] and there exists a positive real number $\delta_0 > 0$ such that for every $x \in (x_0, x_0 + \delta_0]$, we assume without loss of generality that $f(x) - P_1(x) > f(x) - P_2(x) > 0$.

Also there exists a real number $x_1 \in [x_0, x_0 + \delta_0]$ such that

$$(f - P_1)(x_1) = \max_{x \in [x_0, x_0 + \delta_0]} (f - P_1)(x)$$

Considering that $a_n \to 0$, there must be some n such that $a_n < (f - P_1)(x_1)$. Due to $f - P_1$ continuity, there is a real number $x_2 \in [x_0, x_0 + \delta_0]$ such that $(f - P_1)(x_2) = a_n$.

For this x_2 we can find some $\delta_1 > 0$ such that $(f - P_2)(x) < a_n < (f - P_1)(x)$ for every $x \in (x_2, x_2 + \delta_1) \subset [x_0, x_0 + \delta_0]$. If there were no such $\delta_1 > 0$, then $f - P_1$ would not reach up to its maximum at $(f - P_1)(x_1)$.

Finally, if we set $I := (x_2, x_2 + \delta_1)$, we can see that

$$\begin{split} \int_{[a,b]} \Phi(|f-(\frac{P_1+P_2}{2})|) \, dx &= \int_I \Phi(|f-(\frac{P_1+P_2}{2})|) \, dx + \int_{[a,b]-I} \Phi(|f-(\frac{P_1+P_2}{2})|) \, dx \\ &< \frac{1}{2} \int_I \Phi(|f-P_1|) \, dx + \frac{1}{2} \int_I \Phi(|f-P_2|) \, dx + \\ &+ \frac{1}{2} \int_{[a,b]-I} \Phi(|f-P_1|) \, dx + \frac{1}{2} \int_{[a,b]-I} \Phi(|f-P_2|) \, dx \\ &= \frac{1}{2} \int_{[a,b]} \Phi(|f-P_1|) \, dx + \frac{1}{2} \int_{[a,b]} \Phi(|f-P_2|) \, dx. \end{split}$$

This is a contradiction since $\frac{P_1+P_2}{2} \in S_n$.

References

- Acinas, S., Favier, S. and Lorenzo, R. (2024) Extension of the Best Polynomial Operator in Generalized Orlicz Spaces. arXiv:2402.17048 [math.CA]. https://doi.org/10.48550/arXiv.2402.17048
- Acinas, S. and Favier, S. (2016) Multivalued Extended Best Φ-Polynomial Approximation Operator, Numerical Functional Analysis and Optimization, 37 (11): 1339-1353. https://doi.org/10.1080/01630563.2016.1216445
- 3. Benavente, A., Favier, S. and Levis, F. (2017) Existence and characterization of best φ- approximations by linear subspaces, Adv. Pure Appl. Math. 8 (3): 209-217. https://doi.org/10.1515/apam-2015-0069
- Cheney, E.W. and Wulbert, D.E. (1969) The existence and unicity of best approximations, Math. Scand.,
 24: 113-140. https://doi.org/10.7146/math.scand.a-10925
- De Vore, R. (1968) One-sided approximation of functios, J. Approx. Theory, 1: 11-25. https://doi.org/10.1016/0021-9045(68)90054-3
- Galkin, R. V. (1974) The uniqueness of the element of best mean approximation to a continuous function using splines with fixed nodes, Math. Notes, 15: 3-8. https://doi.org/10.1007/BF01153536
- Jackson, D. (1924) A general class of problems in Approximation, Amer. J. Math., 46: 215-234. https://doi.org/10.2307/2370858

- 8. Kroo, A. (1985) On an L^1 -approximation problem, Proc. Amer. Math., **94**: 406-410. https://doi.org/10.1090/S0002-9939-1985-0787882-0
- Micchelli, C. A. (1977) Best L¹ approximation by weak Chebyshev systems and the uniqueness of interpolating perfect splines, J. Approx. Theory, 19: 1-14. https://doi.org/10.1016/0021-9045(77)90024-7
- Pinkus, A. (1986) Unicity subspaces in L¹ approximation, J. Approx. Theory, 48: 226-250. https://doi.org/10.1016/0021-9045(86)90007-9
- 11. Pinkus, A. (1989) On L1-Approximation, Cambridge University Press; i-vi.
- 12. Rice, J. (1964) The Approximation of Functions, Addison-Wesley Publishing Company, Inc., USA.
- 13. Schumaker, L. L. (1981) Spline Functions: Basic Theory. Wiley-Interscience.
- 14. Strauss, H. (1982) Unicity of best one-sided L^1 approximation, Numer. Math., **40**: 229-243. https://doi.org/10.1007/BF01400541
- ¹ Instituto de Matemática Aplicada San Luis, UNSL-CONICET and Departamento de Matemática, FCFMyN, UNSL, Av. Ejército de los Andes 950, 5700 San Luis, Argentina. Email address: abenaven@unsl.edu.ar
- ² Instituto de Matemática Aplicada San Luis, UNSL-CONICET and Departamento de Matemática, FCFMyN, UNSL, Av. Ejército de los Andes 950, 5700 San Luis, Argentina. Email address: costaponcejuan@gmail.com
- ³ INSTITUTO DE MATEMÁTICA APLICADA SAN LUIS, UNSL-CONICET AND DEPARTAMENTO DE MATEMÁTICA, FCFMyN, UNSL, Av. Ejército de los Andes 950, 5700 San Luis, Argentina Email address: sfavier@unsl.edu.ar