ON THE COMMUTATIVITY DEGREE OF A FINITE-DIMENSIONAL LIE ALGEBRA

AFSANEH SHAMSAKI, AHMAD ERFANIAN, AND MOHSEN PARVIZI

ABSTRACT. In this paper, we introduce the commutativity degree of a finitedimensional Lie algebra over a finite field and determine upper and lower bounds for it. Moreover, we study some relations between the notion of commutativity degree and known concepts in Lie algebras.

1. INTRODUCTION

In the past decades, the relation between probability theory and other branches of mathematics has attracted some authors. For instance, the relation between group theory and probability theory is considered and the commutativity degree of a finite group G denoted by d(G) defined as the probability that two randomly chosen elements of G commute. More precisely

$$d(G) = \frac{|\{(x,y) \in G \times G \mid xy = yx\}|}{|G|^2}$$

Gustafson [5] was the first one who obtained the upper bound $\frac{5}{8}$ for it. Later, Lescot [8] showed that if G is a group such that $d(G) > \frac{1}{2}$, then G is nilpotent. He studied the relation between the commutativity degree and the concept of isoclinism and proved that two isoclinic finite groups have the same commutativity degree, but the converse is not true in general. One of his results is the computation of the commutativity degree of D_{2n} and $Q_{2^{n+1}}$.

Recently, Nath [11] found the commutativity degree of finite *p*-groups with derived subgroups of order *p* such that $G' \subseteq Z(G)$ and also proved that for every positive integer *k* there exists a family of groups with the commutativity degree of $\frac{1}{k}$. In [12], by Haar measure the notion of commutativity degree was extended for infinite compact groups. Moreover, the commutativity degree of Moufang loops was defined and investigated by Ahmadidelir [1].

In spite of the fact that two concepts "Lie algebra" and "group" are totally different and they have different structures, but sometimes we may observe similar behaviours. In this paper, we are going to define the commutativity degree of a finite-dimensional Lie algebra over the field \mathbb{F}_q by following the same lines in defining it in group theory. As this notion originates in group theory, one may expect a considerable similarity between the notion of commutativity degree in groups and Lie algebras, but there are some important structural differences either in the definitions or in techniques. For instance, the role of conjugacy classes in computing the commutativity degree of a group is very important, but there is no such definition in Lie algebras.

Key words and phrases. Commutativity degree, Lie algebras.

2. Preliminaries

In the following, we recall the definition of a Lie algebra and state some necessary concepts and terminologies which are used in the rest of the paper.

Let \mathbb{F} be a field. A Lie algebra over the field \mathbb{F} is an \mathbb{F} -vector space L, together with a bilinear map called the Lie bracket $L \times L \to L$ with $(x, y) \mapsto [x, y]$, satisfying the following properties:

$$[x, x] = 0 \quad \text{for all} \quad x \in L, \tag{L1}$$

$$[x, [y, z]] + [z, [x, y]] + [y, [z, x]] = 0 \quad \text{for all} \quad x, y, z \in L.$$
 (L2)

The Lie bracket [x, y] is often referred to as the commutator of x and y. The condition L2 is known as the Jacobi identity. Since the Lie bracket [-, -] is bilinear, we have 0 = [x + y, x + y] = [x, x] + [x, y] + [y, x] + [y, y]. Hence the condition L1 implies [x, y] = -[y, x] for all $x, y \in L$. As an example of Lie algebras, if [x, y] = 0 for all $x, y \in L$, then L is called an abelian Lie algebra. In this paper, abelian n-dimensional Lie algebras are denoted by A(n).

A subalgebra H of L is a vector subspace of L provided that is closed under Lie bracket. By considering L as an n-dimensional Lie algebra over the field \mathbb{F}_q and $\{x_1, \ldots, x_n\}$ as a basis of L, we may display a as a linear combinations of x_1, \ldots, x_n for every element $a \in L$. In other words, we have $a = \alpha_1 x_1 + \cdots + \alpha_n x_n$, where $\alpha_i \in \mathbb{F}_q$ for all $1 \leq i \leq n$. Hence, there is q choices for every scaler and so the number of elements of L is equal to q^n and the number of elements of an mdimensional subalgebra H is equal to q^m such that m < n. An ideal of L is a vector subspace of L provided that $[I, L] \subset I$. Every ideal is always a subalgebra, but a subalgebra is not necessarily an ideal. For instance, let S be a subset of L. Then the centralizer of S in L is defined as $C_L(S) = \{x \in L \mid [x,s] = 0 \text{ for all } s \in S\}$ which is a subalgebra and not an ideal in general. There are several examples for an ideal of L. One important example of an ideal of L is the center of L defined as $Z(L) = \{z \in L \mid [z, x] = 0 \text{ for all } x \in L\}$. Note that L is abelian if and only if Z(L) = L. Assume that I and J are two ideals of L. Then one can see $[I,J] = \langle [x,y] \mid x \in I, y \in J \rangle$ is an ideal, which is named a product of ideals. By using it, we can construct another important example of an ideal of L. The derived subalgebra of L can be defined by putting I = J = L. It is common to be denoted by L^2 instead of [L, L]. One of known classes of Lie algebra is Heisenberg Lie algebras. From [2], a Lie algebra L is called Heisenberg if $L^2 = Z(L)$ and dim $L^2 = 1$, it is proved such Lie algebras are odd-dimensional and have the following presentation

$$L = H(m) = \langle x_i, y_i, z \mid [x_i, y_i] = z, \ 1 \le i \le m \rangle.$$

Suppose that L_1 and L_2 are two Lie algebras. The (external) direct sum of them is the vector space $L_1 \oplus L_2 = \{(x_1, x_2) \mid x_1 \in L_1, x_2 \in L_2\}$ with the following operation

$$[(x_1, x_2), (y_1, y_2)] = ([x_1, y_1], [x_2, y_2]).$$

Put $A = \{(x_1, 0) \mid x_1 \in L_1\}$ and $B = \{(0, x_2) \mid x_2 \in L_2\}$. It is easy to see that they are ideals of direct sum such that are isomorphic to L_1 and L_2 , respectively. Also, $A \cap B = 0$ and [A, B] = 0. Conversely, if I_1 and I_2 are two ideals of L such that $L = I_1 + I_2$ and $I_1 \cap I_2 = 0$, then the map $(x_1, x_2) \mapsto x_1 + x_2$ is an isomorphism of $I_1 \oplus I_2$ to L. So, L is internal direct sum of I_1 and I_2 . Note that $[I_1, I_2]$ is contained in $I_1 \cap I_2$ and so $[I_1, I_2] = 0$. The lower central series of a Lie algebra L is a series with terms

$$L_0 = L$$
 and $L_k = [L, L_{k-1}]$ for all $k \ge 1$.

Then the series $L_0 \supseteq L_1 \supseteq L_2 \supseteq \ldots$ is decreasing. The Lie algebra L is called nilpotent if $L_m = 0$ for some $m \ge 1$. For a nilpotent Lie algebra, the smallest nsuch that L has a central series of length n is called the nilpotency class of L. If the lower central series does not stop, then L is called non-nilpotent Lie algebra. There are various non-nilpotent Lie algebras. One of the non-nilpotent Lie algebras is $\langle x, y \mid [x, y] = x \rangle$ which plays an essential role in the next.

All notions and terminologies about Lie algebras are standard here and can be found in [3, 4].

3. The Commutativity degree

Let us start this section by definition of the commutativity degree of finitedimensional Lie algebra L over the field \mathbb{F}_q as the following.

Definition 3.1. Let *L* be a finite-dimensional Lie algebra over the field \mathbb{F}_q . The commutativity degree of *L* is defined as the ratio

$$d(L) = \frac{|\{(x,y) \in L \times L \mid [x,y] = 0\}|}{|L|^2}.$$

It is clear that d(L) = 1 if and only if L is abelian. So, we work on non-abelian Lie algebras. From now, all Lie algebras are considered finite-dimensional non-abelian Lie algebra over the field \mathbb{F}_q .

Put $B(L) = \{(x, y) \in L \times L \mid [x, y] = 0\}$. We observe that $|B(L)| = \sum_{x \in L} |C_L(x)|$, where $C_L(x) = \{y \in L \mid [x, y] = 0\}$. So, we can easily see that $d(L) = \frac{1}{|L|^2} \sum_{x \in L} |C_L(x)|$. For a Lie algebra L and every element $x \in L$, we denote ad_x as the map from L to L by the rule $ad_x(y) = [x, y]$ for all $y \in L$. It is obvious that ad_x is a linear map. The following Lemma shows that d(L) can be rewritten in term of $\operatorname{Im} ad_x$'s.

Lemma 3.2. Let L be a Lie algebra. Then

$$d(L) = \frac{1}{|L|} \sum_{x \in L} \frac{1}{|\operatorname{Im} ad_x|}.$$

Proof. Consider $ad_x : L \to L$. We have dim $L = \dim \operatorname{Im} ad_x + \dim \ker ad_x$. Since $\ker ad_x = C_L(x)$, so dim $L = \dim \operatorname{Im} ad_x + \dim C_L(x)$. Hence

(3.1)
$$|L| = q^{\dim L} = q^{\dim \operatorname{Im} ad_x + \dim C_L(x)} = |\operatorname{Im} ad_x||C_L(x)|.$$

Therefore, the result follows by using $d(L) = \frac{1}{|L|^2} \sum_{x \in L} |C_L(x)|$ and (3.1).

By Lemma 3.2, we may compute the commutativity degree of Heisenberg Lie algebras.

Example 3.3. Let *m* be a positive integer and $L = H(m) = \langle x_i, y_i, z \mid [x_i, y_i] = z$, $1 \leq i \leq m \rangle$. Since Im $ad_x \subseteq L^2$ and dim $L^2 = 1$, we have Im $ad_x = L^2$ for all $x \in L \setminus Z(L)$. Hence $|\operatorname{Im} ad_x| = q$ for all $x \in L \setminus Z(L)$ and

$$\begin{split} d(L) &= \frac{1}{|L|} \sum_{x \in L} \frac{1}{|\operatorname{Im} ad_x|} = \frac{|Z(L)|}{|L|} + \frac{1}{|L|} \sum_{x \in L \setminus Z(L)} \frac{1}{|\operatorname{Im} ad_x|} \\ &= \frac{|Z(L)|}{|L|} + \frac{1}{|L|} (|L| - |Z(L)|) \frac{1}{q} = \frac{q^{2m} + q - 1}{q^{2m+1}}. \end{split}$$

In the following example, we compute the commutativity degree of the unique non-nilpotent Lie algebra of dimension 2.

Example 3.4. Let $L = \langle x_1, y_1 | [x_1, y_1] = x_1 \rangle$. It is clear that Z(L) = 0 and so |Z(L)| = 1. If $x \neq 0$, then $\operatorname{Im} ad_x = L^2$, which implies $|\operatorname{Im} ad_x| = q$. Thus

$$\begin{split} d(L) &= \frac{1}{|L|} \sum_{x \in L} \frac{1}{|\operatorname{Im} ad_x|} = \frac{1}{|L|} + \frac{1}{|L|} \sum_{x \neq 0} \frac{1}{|\operatorname{Im} ad_x|} = \frac{1}{|L|} + \frac{1}{|L|} (|L| - |Z(L)|) \frac{1}{q} \\ &= \frac{q^2 + q - 1}{q^3}. \end{split}$$

The commutativity degree of a direct sum of two Lie algebras is given in the following.

Proposition 3.5. Let H and K be Lie algebras and $L = H \oplus K$. Then

$$d(L) = d(H)d(K).$$

Proof. Since there is a natural bijection between B(L) and $B(H) \times B(K)$, we have |B(L)| = |B(H)||B(K)|. So,

$$|L|^2 = q^{\dim H + \dim K} q^{\dim H + \dim K} = |H|^2 |K|^2.$$

Now, the result follows.

A Lie algebra L is called stem if $Z(L) \subseteq L^2$. From [6, Proposition 3.1], we know that every finite-dimensional Lie algebra L is isomorphic to $T \oplus A$, in which A is abelian and T is stem. Now, by Proposition 3.5 we have d(L) = d(T). Therefore, the commutativity degree of stem Lie algebras is important. Thus, without loss of generality we may always assume that all Lie algebras are stem.

The concept of isoclinism is a useful tool in computation of the commutativity degree of groups. As we know two groups G_1 and G_2 are called isoclinic provided that there are isomorphisms $\alpha : G_1/Z(G_1) \to G_2/Z(G_2)$ and $\beta : G'_1 \to G'_2$ such that the following diagram commutes:

$$\begin{array}{c} G_1/Z(G_1) \times G_1/Z(G_1) \longrightarrow G'_1 \\ & \downarrow^{\alpha \times \alpha} & \downarrow^{\beta} \\ G_2/Z(G_2) \times G_2/Z(G_2) \longrightarrow G'_2 \end{array}$$

The horizontal maps are defined by $(\overline{x}, \overline{y}) \mapsto [x, y]$. This notion is denoted by $G_1 \sim G_2$. Lescot [9] proved that two isoclinic groups have the same commutativity degree. Now, it is interesting to see what will happen for isoclinic Lie algebras. We know that the definition of isoclinism in Lie algebras the same as the group case by replacing G'_1 and G'_2 by L_1^2 and L_2^2 . In the following theorem, we show that two isoclinic Lie algebras L and M have the same commutativity degree similar to groups.

Theorem 3.6. Let L and M be the isoclinic Lie algebras. Then d(L) = d(M).

Proof. By using [6, Proposition 3.1], we have $L = T_1 \oplus A_1$ and $M = T_2 \oplus A_2$ in which T_1 and T_2 are stem and A_1 and A_2 are abelian. It is proved that [10, Lemma 1] $L \sim T_1$ and $M \sim T_2$ and hence $T_1 \sim T_2$. Now, by using [10, Theorem 11] $T_1 \cong T_2$ and so $d(T_1) = d(T_2)$. The result follows by using Proposition 3.5.

The concept of isoclinism enables us to compute the commutativity degree of Lie algebras more easily. The next example shows how it works.

Example 3.7. Consider the following Lie algebras

$$L = H(1) \oplus H(1) = \langle a_1, \dots, a_6 \mid [a_1, a_2] = a_5, [a_3, a_4] = a_6 \rangle \text{ and}$$
$$M = L_{6,7}^{(2)}(0) = \langle x_1, \dots, x_6 \mid [x_1, x_2] = x_5, [x_3, x_4] = x_5 + x_6, [x_1, x_3] = x_6 \rangle$$

over the field \mathbb{F}_2 . We claim that $d(L) = d(M) = \frac{25}{64}$. By using the identity homomorphism $\beta: L^2 \longrightarrow M^2$ such that $a_5 \longmapsto x_5$ and $a_6 \longmapsto x_5 + x_6$ and the isomorphism $\alpha: L/Z(L) \longrightarrow M/Z(M)$ given by

$$\overline{a}_1 \longmapsto \overline{x}_1 + \overline{x}_2 + \overline{x}_4, \quad \overline{a}_2 \longmapsto \overline{x}_2, \quad \overline{a}_3 \longmapsto \overline{x}_2 + \overline{x}_3 + \overline{x}_4, \quad \overline{a}_4 \longmapsto \overline{x}_4,$$

we can see $L \sim M$. Now, d(L) = d(M) by Theorem 3.6. On the other hand, $d(L) = d(H(1))d(H(1)) = \frac{25}{64}$ by Example 3.3 and Proposition 3.5. So, the result holds.

It is known that two groups with the same commutativity degree need not to be isoclinic (consider two groups D_8 and Q_8). Now, we have the same situation for Lie algebras.

Example 3.8. Take the following Lie algebras

$$L_{5,5} = \langle x_1, \dots, x_5 \mid [x_1, x_2] = x_3, [x_1, x_3] = x_5, [x_2, x_4] = x_5 \rangle \text{ and} \\ L_{5,7} = \langle x_1, \dots, x_5 \mid [x_1, x_2] = x_3, [x_1, x_3] = x_4, [x_1, x_4] = x_5 \rangle.$$

First, we show that $d(L_{5,5}) = d(L_{5,7}) = \frac{q^3 + q^2 - 1}{q^5}$. We know that $\operatorname{Im} ad_a \subseteq L^2$ for every $a \in L_{5,5}$ and $\dim L_{5,5}^2 = 2$ which implies $\dim \operatorname{Im} ad_a \leq 2$. Also, we can write $a = \alpha_1 x_1 + \alpha_2 x_2 + \alpha_3 x_3 + \alpha_4 x_4 + \alpha_5 x_5 + z$ for some $\alpha_i \in \mathbb{F}_q$ such that $1 \leq i \leq 5$ and $z \in Z(L)$. Then $\dim \operatorname{Im} ad_a = 2$ when $\alpha_1 \neq 0$. By counting the coefficients of a, the number of such elements is $(q-1)q^4$ when $\alpha_1 \neq 0$. If $\alpha_1 = 0$ and $\alpha_2 \neq 0$, then one can see $\dim \operatorname{Im} ad_a = 2$ by a similar method and the number of them is $(q-1)q^3$. Let $\alpha_1 = \alpha_2 = 0$ and $\alpha_3 \neq 0$. Then $\dim \operatorname{Im} ad_a = 1$ and their number is equal to $(q-1)q^2$. Finally, $\dim \operatorname{Im} ad_a = 1$ and the number of these elements is (q-1)q when $\alpha_1 = \alpha_2 = \alpha_3 = 0$ and $\alpha_4 \neq 0$. On the other hand, $|Z(L_{5,5})| = q$ thus by Lemma 3.2

$$d(L_{5,5}) = \frac{1}{|L|} \sum_{x \in L} \frac{1}{|\operatorname{Im} ad_x|} = \frac{|Z(L)|}{|L|} + \frac{1}{|L|} \sum_{x \in L \setminus Z(L)} \frac{1}{|\operatorname{Im} ad_x|}$$
$$= \frac{q}{q^5} + \frac{1}{q^5} \left(\frac{q^3 - q}{q} + \frac{q^5 - q^3}{q^2}\right) = \frac{q^3 + q^2 - 1}{q^5}.$$

Similarly, $d(L_{5,7}) = \frac{q^3 + q^2 - 1}{q^5}$ is obtained. Since $|L_{5,5}^2| \neq |L_{5,7}^2|$, they are not isoclinic.

4. Bounds on commutativity degree

In this section, we give an upper bound and a lower bound for the commutativity degree of a Lie algebra in term of q. As a result, we have the upper bound $\frac{5}{8}$ for the Lie algebra L (similar to the upper bound for the commutativity degree in groups). It is worth emphasizing that there is no Lie algebra with the commutativity degree in the interval $(\frac{5}{8}, 1)$.

We start with the upper bound for centerless Lie algebras, but before that we recall the breath of a Lie algebra used in the next theorems. For any Lie algebra L, the breath of L is denoted by b(L) defined as the set $b(L) = \max\{\dim \operatorname{Im} ad_x \mid x \in L\}$. This concept plays an essential role in classification of Lie algebras and other issues (see [7] for more details).

Theorem 4.1. Let L be an n-dimensional Lie algebra with Z(L) = 0. Then

$$d(L) \le \frac{q^n + q - 1}{q^{n+1}}.$$

The equality holds if and only if $\dim L^2 = 1$.

Proof. If $x \neq 0$, then $\operatorname{Im} ad_x$ is a non-zero and $\operatorname{Im} ad_x \subseteq L^2$ and so dim $\operatorname{Im} ad_x \geq 1$. Thus $\frac{1}{|\operatorname{Im} ad_x|} \leq \frac{1}{q}$ for all non-zero elements x. Hence by Lemma 3.2

$$d(L) = \frac{1}{|L|} \sum_{x \in L} \frac{1}{|\operatorname{Im} ad_x|} = \frac{1}{|L|} + \frac{1}{|L|} \sum_{x \neq 0} \frac{1}{|\operatorname{Im} ad_x|}$$
$$\leq \frac{1}{q^n} + \frac{1}{q^n} \frac{q^n - 1}{q} = \frac{q^n + q - 1}{q^{n+1}}.$$

Assume that dim $L^2 = 1$. Then b(L) = 1 by [7, Theorem 3.2.1] and so $|\operatorname{Im} ad_x| = q$ for all $x \in L \setminus Z(L)$. Therefore, $d(L) = \frac{q^n + q - 1}{q^{n+1}}$ by Lemma 3.2. Conversely, let the equality hold. We claim that dim $L^2 = 1$. By contrary, suppose that dim $L^2 \neq 1$. Since L is non-abelian, then dim $L^2 \geq 2$. So, there exists an element x such that $|\operatorname{Im} ad_x| > q$ by [7, Theorem 3.2.1]. Hence $d(L) < \frac{q^n + q - 1}{q^{n+1}}$, which is a contradiction.

The case that $Z(L) \neq 0$ is stated as the following.

Theorem 4.2. Let L be a Lie algebra with dim $Z(L) \ge 1$. Then

$$d(L) \le \frac{q^2 + q - 1}{q^3}.$$

Proof. By Lemma 3.2, we have

$$d(L) = \frac{1}{|L|} \sum_{x \in L} \frac{1}{|\operatorname{Im} ad_x|} = \frac{1}{|L|} \left(|Z(L)| + \sum_{x \in L \setminus Z(L)} \frac{1}{|\operatorname{Im} ad_x|} \right)$$
$$\leq \frac{|Z(L)|}{|L|} + \frac{1}{q} - \frac{|Z(L)|}{|L|q} = \frac{1}{q} + \frac{|Z(L)|}{|L|} (1 - \frac{1}{q}).$$

One the other hand, we know that $\dim L/Z(L) \ge 2$ for all non-abelian Lie algebras. Thus

$$d(L) \le \frac{q^2 + q - 1}{q^3}.$$

as required.

Corollary 4.3. Let L be a Lie algebra. Then

(4.1)
$$d(L) \le \frac{q^2 + q - 1}{q^3}.$$

Proof. Assume that dim L = n. If $Z(L) \ge 1$, then $d(L) \le \frac{q^2+q-1}{q^3}$ by Theorem 4.2. If Z(L) = 0, then $d(L) \le \frac{q^n+q-1}{q^{n+1}}$ by Theorem 4.1. It is enough to show that

$$\frac{q^n + q - 1}{q^{n+1}} \le \frac{q^2 + q - 1}{q^3}. \text{ We know } q^3(q - 1) \le q^{n+1}(q - 1) \text{ for all } n \ge 2, \text{ which implies } \frac{q - 1}{q^{n+1}} \le \frac{q - 1}{q^3}. \text{ Hence } \frac{q^n + q - 1}{q^{n+1}} \le \frac{q^2 + q - 1}{q^3}. \square$$

By the breath of a Lie algebra, we give a lower bound for the commutativity degree of Lie algebras. Let us state the following Lemma from [7] which can be used in the proof of Theorem 4.5.

Lemma 4.4. [7, Theorem 3.1.9] Let L be a Lie algebra such that b(L) = n, (n > 0). Then dim $(L/Z(L)) \ge n + 1$.

Theorem 4.5. Let L be a Lie algebra such that $\dim L/Z(L) = t$. Then

$$d(L) \ge \frac{1}{q^t} + \frac{1}{q^{t-1}} - \frac{1}{q^{2t-1}}$$

Proof. Let dim L/Z(L) = t. Theorem 4.4 shows that dim $L/Z(L) \ge \dim \operatorname{Im} ad_x + 1$ for all $x \in L$, so we have $\frac{1}{|\operatorname{Im} ad_x|} \ge \frac{1}{q^{t-1}}$. Hence by Lemma 3.2

$$d(L) = \frac{1}{|L|} \sum_{x \in L} \frac{1}{|\operatorname{Im} ad_x|} = \frac{1}{|L|} \left(|Z(L)| + \sum_{x \in L \setminus Z(L)} \frac{1}{|\operatorname{Im} ad_x|} \right)$$

$$\geq \frac{|Z(L)|}{|L|} + \frac{(|L| - |Z(L)|)}{|L|} \frac{1}{q^{t-1}} = \frac{1}{q^t} + \frac{1}{q^{t-1}} - \frac{1}{q^{2t-1}}.$$

The next corollary shows all central quotients of a Lie algebra L that d(L) attains the bound (4.1) has exactly 2-dimensional.

Corollary 4.6. Let L be a Lie algebra. Then dim L/Z(L) = 2 if and only if $d(L) = \frac{q^2+q-1}{a^3}$.

Proof. Since dim L/Z(L) = 2, then $\frac{q^2+q-1}{q^3} \leq d(L)$ by Theorem 4.5. On other hand, $d(L) \leq \frac{q^2+q-1}{q^3}$ by Corollary 4.1. So, $d(L) = \frac{q^2+q-1}{q^3}$. Conversely, let $x \notin Z(L)$. Then $|L|/|C_L(x)| \geq q$. Hence

$$\frac{q^2 + q - 1}{q^3} = d(L) = \frac{1}{|L|^2} \sum_{x \in L} |C_L(x)| \le \frac{1}{|L|^2} (|Z(L)||L| + (|L| - |Z(L)|) \frac{|L|}{q})$$
$$= \frac{1}{q} + \frac{|Z(L)|}{|L|} (1 - \frac{1}{q}).$$

Thus $\frac{|L|}{|Z(L)|} \leq q^2$. Since L is non-abelian, we have dim L/Z(L) = 2.

In the next theorem, we obtain the upper bound $\frac{5}{8}$ and show that it is just valid for the commutativity degree of all Lie algebras over the field \mathbb{F}_2 . Hence there is no Lie algebra L with $d(L) > \frac{5}{8}$.

Theorem 4.7. Let L be an n-dimensional Lie algebra. Then $d(L) \leq \frac{5}{8}$. Moreover, $d(L) = \frac{5}{8}$ if and only if L is isomorphic to one of the following Lie algebras over the field \mathbb{F}_2

$$\langle x, y, z \mid [x, y] = z \rangle = H(1) \oplus A(n-3)$$

or

$$\langle x, y \mid [x, y] = x \rangle \oplus A(n-2).$$

Proof. By Corollary 4.3, we have $d(L) \leq \frac{q^2+q-1}{q^3}$. Put $f(q) = \frac{q^2+q-1}{q^3}$. Since f(q) is a decreasing function for all $q \geq 2$, we have $d(L) \leq \frac{5}{8}$.

If L is isomorphic to one of the above Lie algebras, then $d(L) = \frac{5}{8}$ by Proposition 3.5, Examples 3.3 and 3.4 for q = 2. Conversely, let $d(L) = \frac{5}{8}$. Note that $d(L) = \frac{5}{8}$ when q = 2 by the first proof. Since $L = T \oplus A$ where T is stem and A is abelian by [6, Proposition 3.1], it is enough to determine the structure of T. Also, $\dim T/Z(T) = 2$ by Corollary 4.6 thus $\dim T^2 \leq 1$ by [10, Lemma 14]. On the other hand, T is non-abelian, so $\dim T^2 = 1$. Since $Z(T) \subseteq T^2$ and $\dim T^2 = 1$, we have $\dim Z(T) = 0$ or $\dim Z(T) = 1$. Therefore $\dim T = 2$ or 3 and so T is isomorphic to $\langle x, y, z \mid [x, y] = z \rangle \cong H(1)$ or $\langle x, y \mid [x, y] = x \rangle$ by [4, Theorem 3.1 and section 3.2.1], Examples 3.3 and 3.4. Hence L is isomorphic to one of the Lie algebras over the field \mathbb{F}_2

$$\langle x, y, z \mid [x, y] = z \rangle \cong H(1) \oplus A(n-3),$$

or

$$\langle x, y \mid [x, y] = x \rangle \oplus A(n-2).$$

Now, we are in a position to compare d(L) and d(H), where H is a subalgebra of L.

Lemma 4.8. Let L be a Lie algebra and H be a subalgebra of L. Then $\frac{|H|}{|C_H(x)|} \leq \frac{|L|}{|C_L(x)|}$ for all $x \in L$.

Proof. It is clear that $H + C_L(x) \subseteq L$ for all $x \in L$. Hence $|H + C_L(x)| \leq |L|$ and so $|H||C_L(x)|/|H \cap C_L(x)| \leq |L|$. Since $H \cap C_L(x) = C_H(x)$, we have $|H||C_L(x)|/|C_H(x)| \leq |L|$ as required.

The following proposition gives a comparison between d(H) and d(L) for every subalgebra H of L.

Proposition 4.9. Let L be a Lie algebra and H be a subalgebra of L. Then $\frac{|H|^2}{|L|^2}d(H) \leq d(L) \leq d(H)$. If L = H + Z(L), then d(L) = d(H).

Proof. By Lemma 4.8

(4.2)
$$\sum_{x \in L} |C_L(x)| \le \frac{|L|}{|H|} \sum_{x \in L} |C_H(x)|$$

If $x \in L$ and $y \in C_H(x)$, then [x, y] = 0. Thus $x \in C_L(y)$ and we can rewrite (4.2) as follows:

(4.3)
$$\sum_{x \in L} |C_L(x)| \le \frac{|L|}{|H|} \sum_{y \in H} |C_L(y)|.$$

By Lemma 4.8 and (4.3) we have

(4.4)
$$\sum_{x \in L} |C_L(x)| \le \frac{|L|^2}{|H|^2} \sum_{y \in H} |C_H(y)|$$

Consequently,

$$d(L) = \frac{1}{|L|^2} \sum_{x \in L} |C_L(x)| \le \frac{1}{|L|^2} \frac{|L|^2}{|H|^2} \sum_{y \in H} |C_H(y)| = \frac{1}{|H|^2} \sum_{y \in H} |C_H(y)| = d(H).$$

Since $|C_H(x)| \leq |C_L(x)|$, we have also

$$d(L) = \frac{1}{|L|^2} \sum_{x \in L} |C_L(x)| \ge \frac{1}{|L|^2} \sum_{x \in L} |C_H(x)| \ge \frac{1}{|L|^2} \sum_{x \in H} |C_H(x)| = \frac{|H|^2}{|H|^2 |L|^2} \sum_{x \in H} |C_H(x)|$$
$$= \frac{|H|^2}{|L|^2} d(H).$$

Therefore $\frac{|H|^2}{|L|^2}d(H) \leq d(L) \leq d(L)$. Let L = H + Z(L). Then $Z(L) = \bigcap_{x \in L} C_L(x)$ and $H + C_L(x) \subseteq L$ imply that $L = H + C_L(x)$. Hence $\frac{|H|}{|C_H(x)|} = \frac{|L|}{|C_L(x)|}$ for all $x \in L$ by the proof of Lemma 4.8. Therefore the inequality change into equality in (4.4) and so d(L) = d(H). \Box

Now, we compare d(L) with d(N) and d(L/N), where N is an ideal of L.

Proposition 4.10. Let L be a Lie algebra and N be an ideal of L. Then

$$d(L) \le d(\frac{L}{N})d(N).$$

If $N \cap L^2 = 0$, then the equality holds.

Proof. Let $y \in L$. By the second isomorphism theorem and $C_N(y) = C_L(y) \cap N$, we have $\frac{C_L(y)}{C_N(y)} \cong \frac{C_L(y)+N}{N} \subseteq C_{L/N}(y+N)$. Hence

$$\sum_{y \in L} |C_L(y)| = \sum_{S \in L/N} \sum_{y \in S} \frac{|C_L(y)|}{|C_N(y)|} |C_N(y)| = \sum_{S \in L/N} \sum_{y \in S} \frac{|C_L(y)|}{|C_L(y) \cap N|} |C_N(y)|$$

$$= \sum_{S \in L/N} \sum_{y \in S} \frac{|C_L(y) + N|}{|N|} |C_N(y)| \le \sum_{S \in L/N} \sum_{y \in S} |C_{L/N}(y + N)| |C_N(y)|$$

$$= \sum_{S \in L/N} |C_{L/N}(S)| \sum_{y \in S} |C_N(y)| = \sum_{S \in L/N} |C_{L/N}(S)| \sum_{y \in S} |\{x \in N \mid [x, y] = 0\}|$$

$$= \sum_{S \in L/N} |C_{L/N}(S)| \sum_{x \in N} |C_L(x) \cap S|.$$
(4.5)

If $C_L(x) \cap S = \emptyset$, then $|C_L(x) \cap S| < |C_N(x)|$. If $C_L(x) \cap S \neq \emptyset$, then there exist $x_0 \in C_L(x) \cap S$. So, we have $S = x_0 + N$

$$C_L(x) \cap S = C_L(x) \cap (x_0 + N) = (x_0 + C_L(x)) \cap (x_0 + N) = (C_L(x) \cap N) + x_0$$

= $C_N(x) + x_0$.

Hence $|C_L(x) \cap S| = |C_N(x) + x_0| = |C_N(x)|$. So, in any case (4.6) $|C_L(x) \cap S| \le |C_N(x)|$.

Finally, by using (4.5) and (4.6)

$$d(L) = \frac{1}{|L|^2} \sum_{S \in L/N} |C_{L/N}(S)| \sum_{x \in N} |C_L(x) \cap S|$$

$$\leq \frac{|N|^2}{|L|^2} \sum_{S \in L/N} |C_{L/N}(S)| \frac{1}{|N|^2} \sum_{x \in N} |C_N(x)| = d(L/N)d(N).$$

Assume that $x+N \in C_{L/N}(y+N)$. Thus $[x, y] \in N$. If $N \cap L^2 = 0$, then [x, y] = 0 and so $x+N \in \frac{C_L(y)+N}{N}$. Hence $C_{L/N}(y+N) \subseteq \frac{C_L(y)+N}{N}$. Since $\frac{C_L(y)+N}{N} \subseteq C_{L/N}(y+N)$, we have $\frac{C_L(y)+N}{N} = C_{L/N}(y+N)$. We can also see that $C_L(x) \cap S \neq \emptyset$ for all $x \in N$ and for $S \in \frac{L}{N}$. Therefore all inequalities change into equalities. \Box

Proposition 4.11. Let L be a Lie algebra and M, N be ideals of L such that $N \subseteq M$. Then $d(\frac{L}{N}) \leq d(\frac{L}{M})$.

Proof. Since $d(\frac{L}{M}) = d(L/N/M/N)$ by the third isomorphism theorem, we have

$$d(\frac{L}{M}) = d(\frac{L/N}{M/N}) \ge \frac{d(L/N)}{d(M/N)} \ge d(\frac{L}{N})$$

by Proposition 4.10 and the fact that $0 < d(M/N) \le 1$.

References

- [1] Ahmadidelir, Karim. On the commutativity degree in finite Moufang loops. Int. J. Group Theory 5 (2016), no. 3, 37–47.
- [2] Batten, Peggy; Stitzinger, Ernest. On covers of Lie algebras. Comm. Algebra 24 (1996), no. 14, 4301–4317.
- [3] Cicalò, Serena; de Graaf, Willem A.; Schneider, Csaba. Six-dimensional nilpotent Lie algebras. Linear Algebra Appl. 436 (2012), no. 1, 163–189.
- [4] Erdmann, Karin, and Mark J. Wildon. Introduction to Lie algebras. Vol. 122. London: Springer, 2006.
- [5] Gustafson, William H. What is the probability that two group elements commute? The American mathematical monthly 80.9 (1973), 1031-1034.
- [6] Johari, Farangis; Parvizi, Mohsen and Niroomand, Peyman. Capability and Schur multiplier of a pair of Lie algebras. J. Geom. Phys. 114 (2017), 184–196.
- [7] Khuhirun, Borworn. Classification of nilpotent Lie algebras with small breadth. North Carolina State University, 2014.
- [8] Lescot, Paul. Sur certains groupes finis. Rev. Math. Spéciales 8 (1987), 276-277.
- [9] Lescot, Paul. Isoclinism classes and commutativity degrees of finite groups. J. Algebra 177 (1995), no. 3, 847–869.
- [10] Moneyhun, Kay. Isoclinisms in Lie algebras. Algebras Groups Geom. 11 (1994), no. 1, 9–22.
- [11] Nath, Rajat Kanti. On asymptotic commutativity degree of finite groups. Proyecciones 38 (2019), no. 4, 829–835.
- [12] Rezaei, Rashid and Erfanian, Ahmad. On the commutativity degree of compact groups. Arch. Math. (Basel) 93 (2009), no. 4, 345–356.

DEPARTMENT OF MATHEMATICS, FACULTY OF MATHEMATICAL SCIENCES, FERDOWSI UNIVERSITY OF MASHHAD, MASHHAD, IRAN

Email address: Shamsaki.afsaneh@yahoo.com

Email address: erfanian@um.ac.ir

DEPARTMENT OF MATHEMATICS, FACULTY OF MATHEMATICAL SCIENCES, FERDOWSI UNIVERSITY OF MASHHAD, MASHHAD, IRAN

Email address: parvizi@um.ac.ir

DEPARTMENT OF MATHEMATICS, FACULTY OF MATHEMATICAL SCIENCES, FERDOWSI UNIVERSITY OF MASHHAD, MASHHAD, IRAN