# MARKOVIAN LIFTS OF STOCHASTIC VOLTERRA EQUATIONS IN SOBOLEV SPACES: SOLUTION THEORY, AN ITÔ FORMULA AND INVARIANT MEASURES

### FLORIAN HUBER

ABSTRACT. We investigate Markovian lifts of stochastic Volterra equations (SVEs) with completely monotone kernels and general coefficients within a class of weighted Sobolev spaces. Our primary focus is developing a comprehensive solution theory for a class of non-local stochastic evolution equations (SEEs) encompassing these Markovian lifts. This enables us to provide conditions for the existence of invariant measures for the lifted processes and the corresponding SVE. Another key contribution is an Itô-type formula for the stochastic Volterra equations under consideration.

# Contents

. introduction and main results	2
1.1. Notation	6
2. Preliminaries: Weighted spaces	7
3. The lifted equation	12
3.1. The operator and semigroup	13
3.2. Existence	18
3.3. Long term existence and invariant measures	35
4. Ito-formula for Volterra equations	39
5. Appendix	42
5.1. Proofs of technical results	42
5.2. Auxiliary Lemmata	47
References	18

1

<sup>2020</sup> Mathematics Subject Classification. 60H15,60H20,60G22,37A25.

 $Key\ words\ and\ phrases.$  stochastic partial differential equations, Volterra processes, , markovian lift, ergodic behaviour, ito formula.

The author gratefully acknowledges financial support through grant Y 1235 of the START-program during his employment at the University of Vienna, where parts of this paper were written. The author thanks Christa Cuchiero for bringing this topic to his attention.

## 1. INTRODUCTION AND MAIN RESULTS

Driven by the increasing prominence of rough volatility models in mathematical finance, as explored in works such as [36, 40] significant attention has been directed towards the study of stochastic Volterra equations (SVE) of the form:

(1.1) 
$$X_t = X_0 + \int_0^t k_b(t-s)b(s, X_s) ds + \int_0^t k_\sigma(t-s)\sigma(s, X_s) dW_s,$$

where W is a multidimensional Brownian motion and the coefficients  $b, \sigma$ , as well as convolution kernels  $k_b, k_\sigma$  meet certain integrability and regularity conditions. The introduction of such kernels is prevalent in applications to describe phenomena with memory that diminishes over time. Besides finance, these equations also find applications in the modeling of turbulence [12], mechanics [58, 53], biology [35, 62, 65], sociology [18, 29] and energy markets [11]. A key challenge posed by this type of equation is that solutions to stochastic Volterra equations are generally neither semimartingales nor Markov processes. Consequently, existing techniques for studying the existence, uniqueness, and qualitative features of SVEs need extension to accommodate these characteristics. Numerous researchers have contributed to this field (see for example [16, 17, 27, 67]). Recently, a comprehensive weak existence theory was developed in [3] (see also [44]), in the convolutional setting, and [60], in the non-convolutional setting. Equations with affine ( [6, 2, 19]) or polynomial coefficients ([48]) received particular attention, as their structure provides a convenient setting for the application of probabilistic methods.

The uniqueness of solutions to SVEs remains challenging in many cases. For kernels with singularities, pathwise uniqueness of SVEs without drift, under specific conditions on the coefficients, was first established in [57] via an infinite-dimensional lift, a strategy later extended to include a drift component in [59]. If the kernels are regular, in the sense that the kernels do not exhibit singularities, pathwise uniqueness was studied in [61] and [10] for Hölder continuous coefficients ( $\sigma \in C^{\alpha}$ ,  $\alpha \geq \frac{1}{2}$ ).

To investigate qualitative aspects of SVE solutions, one often resorts to an approach that restores the Markov property by suitably extending the state space of the process, at the cost of the new state space becoming infinite-dimensional. The resulting infinite-dimensional Markov process is called Markovian lift. This method was employed in [49, 15] to investigate the long-term behavior of SVEs. To our knowledge, the only direct study at the SVE level was conducted in [39]. There is no canonical way of obtaining such a lift and several forms have been used. The three most common ones are:

- Measure-valued lift: Originating in the context of fractional Brownian motion [26], it was later applied to affine SVEs in [31, 30]. This lifting notably inspired Markovian approximations for SVEs [1, 5]. [43] used the principle of the measure-valued lift and studied it in the spaces introduced in [26, 45, 7].
- Forward curve lift: Motivated by applications in mathematical finance, this lift was utilized in [4, 15].
- The superprocess-inspired lift, which enabled pathwise existence results for certain coefficients and the singular kernels  $k(t-s) \approx (t-s)^{-\alpha}$ . This lift was notably used in [57] to obtain a pathwise-uniqueness result (see also [59]) and conceptually related to catalytic superprocesses.

It should be noted that, on a purely formal level, the first and second lifts exhibit similar properties, as the principal operators in the lifted equations are Laplace transforms of each other, which is also reflected in the corresponding semigroups.

The main aspects of this work are the following:

- We study the (measure-valued) lift for a broad class of coefficients in weighted Sobolev spaces, which are more common in studying (S)PDEs and provide more flexibility compared to spaces used in prior studies [31, 30], or [43]. In particular, this choice gives us easy access to embedding theorems, extending existing results to more general coefficients.
- We present an Itô-type formula for the solution of stochastic Volterra equations which shares certain properties with the (functional) Itô formula from [66] (see also [20]), but is more convenient in its application.
- We extend the results regarding the existence of invariant measures from [49] to settings with general coefficients and a wider variety of kernels.

The primary tool will be the infinite-dimensional lift of (1.1), presented in [26], to a stochastic evolution equation (SEE) on of the form

(1.2) 
$$\mu_t = \mu_0 - \int_0^t x \mu_s \, \mathrm{d}s + \int_0^t \nu_b b(s, \langle \mu_s, 1 \rangle) \, \mathrm{d}s + \int_0^t \nu_\sigma \sigma(s, \langle \mu_s, 1 \rangle) \, \mathrm{d}W_s.$$

The tools, facilitating this lift are the following definition and theorem:

**Definition 1.** Let  $k: \mathbb{R}_+ \to \mathbb{R}_+$ . We say that k is a completely monotone function, if k is infinitely differentiable on  $(0,\infty)$  and satisfies  $(-1)^n \partial_t^n k(t) \geq 0$  for any  $n \in \mathbb{N} \cup \{0\}$  and  $t \in (0,\infty)$ .

**Theorem 2.** [63, Proposition 1.2 and Theorem 4.8]. The following three assertions are equivalent:

- (1) k is completely monotone on  $(0, \infty)$  (respectively on  $[0, \infty)$ );
- (2) k is represented as the Laplace transform of a unique Radon (respectively finite) measure  $\nu$  on  $[0,\infty)$ :

(1.3) 
$$k(\lambda) = \int_0^\infty e^{-\lambda x} \nu(dx), \quad \lambda > 0 (\text{ respectively } \lambda \ge 0)$$

(3) k is infinitely differentiable on  $(0,\infty)$  (respectively continuous on  $[0,\infty)$ , infinitely differentiable on  $(0,\infty)$ ) and satisfies  $(-1)^n k^{(n)} \geq 0$  for every  $n \in \mathbb{N}_0$ .

Let us briefly discuss the heuristics of the lift. Assuming complete monotonicity of the kernels in equation (1.1), and setting  $X_0 = 0$  for simplicity, Theorem 2 allows us to rewrite (1.1) as

(1.4) 
$$X_t = \int_0^t \int_{\mathbb{R}_+} e^{-x(t-s)} \nu_b b(s, X_s) \, \mathrm{d}s + \int_0^t \int_{\mathbb{R}_+} e^{-x(t-s)} \nu_\sigma \sigma(s, X_s) \, \mathrm{d}W_s.$$

Assuming further that X can be represented as the integral w.r.t. another measure  $\mu$  and formally using Fubini's theorem results in

$$(1.5) \qquad \langle \mu_t, 1 \rangle = \left\langle \int_0^t e^{-x(t-s)} \nu_b b(s, \langle \mu_s, 1 \rangle) \, \mathrm{d}s + \int_0^t e^{-x(t-s)} \nu_\sigma \sigma(s, \langle \mu_s, 1 \rangle) \, \mathrm{d}W_s, 1 \right\rangle.$$

We recognize that  $\mu$  should satisfy the equation

(1.6) 
$$\mu_t = \int_0^t e^{-x(t-s)} \nu_b b(s, \langle \mu_s, 1 \rangle) \, \mathrm{d}s + \int_0^t e^{-x(t-s)} \nu_\sigma \sigma(s, \langle \mu_s, 1 \rangle) \, \mathrm{d}W_s,$$

which corresponds to the mild formulation of the equation

(1.7) 
$$d\mu_t = -x\mu_t dt + \nu_b b(t, \langle \mu_t, 1 \rangle) dt + \nu_\sigma \sigma(t, \langle \mu_t, 1 \rangle) dW_t.$$

To make this formal observation rigorous, let us collect the paper's main results. The weighted Sobolev spaces used in the following results will be introduced in section 2 in detail. For now, just note that we will identify the dual space of  $W_w^{m,p}(\mathbb{R}_+)$  via an unweighted  $L^2(\mathbb{R}_+)$  duality with  $W_{\frac{1}{w}}^{-m,q}(\mathbb{R}_+)$ .

We will consider three different weighted spaces and weights  $\frac{1}{w_{-}} < \frac{1}{w_{-}} < \frac{1}{w_{+}}$  ( $w_{-} > w_{\sim} > w_{+}$ ) on  $\mathbb{R}_{+}$ , satisfying Assumption 24.

**Definition 3.** Let  $\theta_{\nu_k}, \theta_{\nu_{\sigma}}$  from Assumption (M 1) be given. For  $i \geq 0$ , we set

$$(w_{-})_{i}(x) := (1+x)^{2\eta_{-}-1+2i}$$

$$> (w_{\sim})_{i}(x) := (1+x)^{2\eta_{\sim}-1+2i}$$

$$> (w_{+})_{i}(x) := (1+x)^{2\eta_{+}-1+2i},$$

and impose the following conditions on these weights.

$$\eta_{+} = \begin{cases} -\varepsilon & \text{if } \theta_{\nu_{\sigma}} < \frac{1}{2}, \text{ where } 0 < \varepsilon < \frac{1}{2} - \theta_{\nu_{\sigma}}, \\ \theta_{\nu_{\sigma}} - \frac{1}{2} + \delta & \text{if } \theta_{\nu_{\sigma}} > \frac{1}{2}, \text{ where } 0 < \delta < \frac{1}{2}, \end{cases}$$

$$\eta_{-} > \max\{\theta_{\nu_{b}}, \theta_{\nu_{\sigma}}\},$$

$$\eta_{+} < \eta_{\sim} < \eta_{-}.$$

The ordering of the weights corresponds to the embedding of these spaces  $W_{\frac{1}{w_+}}^{-1,2} \hookrightarrow W_{\frac{1}{w_-}}^{-2,2} \hookrightarrow W_{\frac{1}{w_-}}^{-2,2}$ .

**Definition 4.** Let  $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{F}}, \tilde{\mathbb{P}})$ , where  $\tilde{\mathbb{F}} = (\tilde{\mathcal{F}}_s)_{s \in [0,T]}$  is a stochastic basis,  $\tilde{\mu}$  is a  $\mathbb{F}$ -progressively measurable process with laws supported on  $C([0,T],W_{\frac{1}{w_{\sim}}}^{-2,2})$  and  $\tilde{W}$  is a Wiener process. We call  $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{F}}, \tilde{\mathbb{P}}, \tilde{\mu}, \tilde{W})$  a probabilistically weak (or martingale) mild solution of

(1.8) 
$$d\mu_t = -x\mu_t dt + \nu_b b(t, \langle \mu_t, \psi \rangle) dt + \nu_\sigma \sigma(t, \langle \mu_t, \psi \rangle) dW_t,$$

with  $\mu_t \mid_{t=0} = \mu_0 \in W_{\frac{1}{w^{\sim}}}^{-1,2}, \ \psi \in W_{w^{\sim}}^{2,2}, \ if$ 

$$(1.9) \quad \widetilde{\mu}_t = e^{-xt}\widetilde{\mu}_0 + \int_0^t e^{-x(t-s)}\nu_b(x)b(s,\langle \widetilde{\mu}_s, \psi \rangle) \,\mathrm{d}s + \int_0^t e^{-x(t-s)}\nu_\sigma(x)\sigma(s,\langle \widetilde{\mu}_s, \psi \rangle) \,\mathrm{d}\widetilde{W}_s,$$

 $\widetilde{\mathbb{P}}$ -a.s. for each  $t \in [0, T]$ .

**Theorem 5.** Let Assumptions (Coeff 1), (Coeff 3) and (M 1) be satisfied and let p > 2. For  $\mu_0 \in L^p(\Omega, W_{\frac{1}{w_+}}^{-1,2})$ , equation (1.8) has a probabilistically weak, analytically mild solution  $\mu \in L^p(\Omega, C([0,T], W_{\frac{1}{w_+}}^{-2,2})) \cap L^2(\Omega, L^{\infty}([0,T], W_{\frac{1}{w_+}}^{-1,2}))$ . In the case when the weights are chosen such that one can set  $\psi = 1$ , we have the following equivalence between solutions of the SVE and the SEE.

**Theorem 6.** Let  $b: \mathbb{R}_+ \times \mathbb{R} \to \mathbb{R}$  and  $\sigma: \mathbb{R}_+ \times \mathbb{R} \to \mathbb{R}$  be continuous and satisfy the linear growth condition (Coeff 1). Further let  $\nu_b, \nu_\sigma$  be non-negative measures on  $\mathbb{R}_+$ , such that for every  $\lambda > 0$ ,  $\int_{\mathbb{R}_+} e^{-\lambda x} \nu_b(dx) = k_b(\lambda)$ ,  $\int_{\mathbb{R}_+} e^{-\lambda x} \nu_\sigma(dx) = k_\sigma(\lambda)$ , where  $k_b \in L^1(0,T)$ ,  $k_\sigma \in L^2(0,T)$  are completely monotone kernels. Let  $w, \frac{1}{w}$  be weight functions, such that Assumptions 46 hold.

(1) Assume there exists a solution X of

(1.10) 
$$X_t = X_0 + \int_0^t k_1(t-s)b(s,X_s) ds + \int_0^t k_2(t-s)\sigma(s,X_s) dW_s,$$

satisfying

(1.11) 
$$\int_0^t |b(s, X_s)| \, \mathrm{d}s < \infty, \qquad \int_0^t |\sigma(s, X_s)|^2 \, \mathrm{d}s < \infty.$$

Then the  $W_{\frac{1}{w}}^{-1,2}$ -valued process Y defined by

$$(1.12) Y_t = e^{-xt} x_0 \delta_0 + \int_0^t e^{-x(t-s)} \nu_b b(s, X_s) \, \mathrm{d}s + \int_0^t e^{-x(t-s)} \nu_\sigma \sigma(s, X_s) \, \mathrm{d}W_s,$$

is a mild solution of

$$(1.13) \mu_t = \mu_0 - \int_0^t x \mu_s \, \mathrm{d}s + \int_0^t \nu_b(x) b(s, \langle \mu_s, 1 \rangle) \, \mathrm{d}s + \int_0^t \nu_\sigma(x) \sigma(s, \langle \mu_s, 1 \rangle) \, \mathrm{d}W_s$$

with initial condition  $\mu_0 = x_0 \delta_0$  and it holds that

$$X_t = \langle \mu_t, 1 \rangle$$
, a.s. for a.e.  $t > 0$ .

(2) If  $\mu$  is a mild solution of the lifted stochastic evolution equation (1.13), according to Definition 42, with the initial condition  $x_0\delta_0$ , and let

(1.14) 
$$\int_0^t |b(s, \langle \mu_s, 1 \rangle)| \, \mathrm{d}s < \infty, \qquad \int_0^t |\sigma(s, \langle \mu_s, 1 \rangle)|^2 \, \mathrm{d}s < \infty,$$

then the  $\mathbb{R}^n$ -valued process X defined by

$$X_t := \langle \mu_t, 1 \rangle, t > 0,$$

is a solution of (1.10) with initial condition  $x_0$ . Furthermore, (1.12) holds a.s. for any  $t \ge 0$ .

In particular, the uniqueness holds for the SVE (1.10) with initial condition  $x_0$  if and only if the uniqueness holds for the lifted SEE (1.8) with the initial condition  $x_0\delta_0$ .

**Remark 7.** Unless otherwise stated, we will restrict ourselves to dimension d=1 to lighten the notation in later chapters. The vector-valued case can be obtained by slightly adapting the arguments presented in this work, and it does not pose much of a challenge as key results regarding our spaces have already been formulated in sufficient generality.

Combining the previous equivalence result and the following theorem will allow us to obtain results regarding the long-term behavior of solutions to the SVE.

**Theorem 8.** Let Assumptions (Coeff 1), (UL), (LT) be satisfied, then a mild solution  $\mu \in W_{\frac{1}{w_+}}^{-1,2}$  of equation (1.8) is a weak (and in particular generalized) Feller process with an invariant measure

**Theorem 9.** Let the kernels  $k_b \in L^1(0,\infty)$  and  $k_\sigma \in L^2(0,\infty)$  be both completely monotone, their associated lifted measures satisfy Assumption (LT), the lifted equation, with initial condition  $\mu_0 = \delta_0 x_0$ , satisfies Assumption (UL) and the coefficients satisfy Assumption (Coeff 1). Then there exists an invariant measure to the SVE (1.1) in the sense that there exists a measure  $\Pi$  and  $x_0 \sim \Pi$ ,  $X_t \sim \Pi$ .

A last consequence of the lift is the following Itô- type formula.

**Proposition 10.** Let  $k_b$ ,  $k_\sigma$  be completely monotone kernels such that their associated measures, given by (1.3), satisfy Assumptions (M 1) and (M 2). Let X denote the solution of the corresponding stochastic Volterra equation (1.1) and set

$$\Gamma_{st}(X) := X_t - \int_s^t k_b(t-r)b(r,X_r) dr - \int_s^t k_\sigma(t-r)\sigma(r,X_r) dW_r.$$

For  $f \in C^{1,2}(\mathbb{R}_+ \times \mathbb{R}, \mathbb{R})$ , the following Itô formula holds.

$$f(t, X_t) = f(t_0, \mathbb{E}\left[\Gamma_{t_0 t}(X) | \mathcal{F}_{t_0}^X\right]) + \int_{t_0}^t \partial_s f\left(s, \mathbb{E}\left[\Gamma_{s t}(X) | \mathcal{F}_s^X\right]\right) ds$$
$$+ \int_{t_0}^t \partial_x f\left(s, \mathbb{E}\left[\Gamma_{s t}(X) | \mathcal{F}_s^X\right]\right) k_{\sigma}(t-s) \sigma(s, X_s) dW_s$$
$$+ \frac{1}{2} \int_{t_0}^t \left(\partial_x^2 f\left(s, \mathbb{E}\left[\Gamma_{s t}(X) | \mathcal{F}_s^X\right]\right)\right) k_{\sigma}(t-s)^2 \sigma(s, X_s)^2 ds.$$

Note that  $\Gamma_{0t} = X_0$ .

1.0.1. Structure of the paper. In section 2 we introduce the mathematical framework for treating the lifted equation, which relies on weighted Sobolev spaces and we prove the most important embedding properties of these spaces. Section 3 is split into two parts. In the first part (subsection 3.1), we study the properties of certain operators related to our lift on the previously introduced spaces. In the second part (subsection 3.2), we obtain strong existence and uniqueness for solutions to the lifted equation with Lipschitz continuous coefficients. An approximation result, uniform estimates, and tightness give us a weak existence result for more general coefficients. In section 3.3, we use the weak (or generalized) Feller theory to obtain the existence of invariant measures for stochastic Volterra equations, under certain conditions. Section 4 is devoted to an Itô-type formula for stochastic Volterra equations and a condition for the global existence of solutions to such equations.

## 1.1. Notation.

- We usually write  $\mathbb{R}_+ := [0, \infty)$ .
- We denote by  $C_b^k(\mathbb{R})$  the space of bounded k-times continuously differentiable functions from  $\mathbb{R}$  to  $\mathbb{R}$ . If they additionally have compact support or vanish at  $\pm \infty$ , we write  $C_c^k(\mathbb{R})$  and  $C_0^k(\mathbb{R})$  respectively.

- To indicate a derivative, with respect to the x variable, we will use the letter D. Higher order derivatives will be denoted in the usual way by  $D^j = \underbrace{D \circ \cdots \circ D}_{k\text{-times}}$  for  $j \in \mathbb{N}$ .
- $\|\cdot\|_{\mathcal{L}(X,Y)}$  denotes the operator norm of an operator mapping from the Banach space X into the Banach space Y.
- For Banach spaces U, V, which are completely regular and  $\rho$  be a weight function on U with relatively compact sublevel sets, we define  $\mathcal{B}^{\rho}(U; V)$  as the closure of  $C_b(U; V)$  with respect to the norm  $||f||_{\rho} := \sup_{u \in U} \rho(u)^{-1} ||f(U)||_{V}$ .
- We follow the convention that C denotes a generic numerical constant that may change from line to line. We add the corresponding parameters as a subscript to highlight a particular dependence. If a parameter of interest does not appear in the subscript, the constant is uniform with respect to this parameter.

### 2. Preliminaries: Weighted spaces

We begin by introducing a class of weighted Lebesgue and Sobolev spaces which will serve as state spaces in our analysis.

**Definition 11.** We call a locally integrable function w on  $\mathbb{R}_+$ , such that w(x) > 0-a.e. a weight or weight function.

Every weight w induces a positive Borel-measure on  $\mathbb{R}_+$  via integration, i.e.  $\lambda_w(E) = \int_E w(x) dx$  for measurable sets  $E \subset \mathbb{R}_+$ .

**Definition 12.** Let w be a weight. For  $0 we define <math>L_w^p$  as the set of measurable functions u on  $\mathbb{R}_+$  such that

$$||u||_{L_w^p} = \left(\int_{\mathbb{R}_+} |u(x)|^p w(x) \, \mathrm{d}x\right)^{1/p} < \infty.$$

The following statements recall that elements of weighted spaces remain distributions on  $\mathbb{R}_+$ .

**Definition 13.** Let p > 1. We say that a weight function w satisfies the condition  $\mathcal{B}_p(\mathbb{R}_+)$  and write  $w \in \mathcal{B}_p(\mathbb{R}_+)$ , if

$$w^{-1/(p-1)} \in L^1_{loc}(\mathbb{R}_+).$$

**Theorem 14.** Let p > 1,  $w \in \mathcal{B}_p(\mathbb{R}_+)$  and K be a compact set such that  $K \subset \mathbb{R}_+$ . Then the embedding

$$L_w^p \hookrightarrow L^1(K)$$

is continuous.

*Proof.* The theorem directly results from Hölder's inequality.

Corollary 15. Under the assumptions of the previous theorem, we have

$$L_w^p \hookrightarrow L_{loc}^1(\mathbb{R}_+) \subset D'(\mathbb{R}_+),$$

where  $D'(\mathbb{R}_+)$  denotes the space of distributions.

Let  $m \in \mathbb{N}$  and  $1 \leq p < \infty$ . Let  $(w_0, \ldots, w_k)$  be a given weight functions. We introduce the norm

$$||u||_{W_w^{m,p}} = \left(\sum_{0 \le j \le m} \int_0^\infty |D^\alpha u|^p w_j dx\right)^{1/p}.$$

**Definition 16.** We denote by  $W_w^{m,p}$  the completion of  $\{u \in C_0^{\infty}(\mathbb{R}_+) : ||u||_{W_w^{m,p}} < \infty\}$  with respect to the norm  $||\cdot||_{W_w^{m,p}}$ .

If  $1 and <math>w \in \mathcal{B}_p(\mathbb{R}_+)$ , then  $W_w^{m,p}$  is a uniformly convex Banach space (see [51, section 4]).

Remark 17. Note that one could also introduce weighted Sobolev spaces  $\mathcal{W}_w^{m,p}$  by considering the set of all functions  $u \in L_w^p$  for which the weak derivatives  $D^j u$ , with  $j \leq m$ , belong to  $L_w^p$ . The weighted Sobolev space  $\mathcal{W}_w^{m,p}$  is a normed linear space if equipped with the norm  $\|\cdot\|_{W_w^{m,p}}$ . We have that  $W \subseteq \mathcal{W}$ . By definition, functions that are smooth in the interior of  $\mathbb{R}_+$  are dense in W, while the space W is known to contain all functions of finite well-defined "energy". If w is bounded from above and away from 0 from below  $0 < c_1 \leq w(x) \leq c_2 < \infty$ , the spaces W and W coincide on general domains, however if  $w \in L^2_{loc}(\mathbb{R}_+)$ , W = W does not need to hold.

Although  $w = (w_0, \dots, w_m)$  is not a weight function according to our definition, but a vector of weight functions, we will still call such a vector a weight function out of convenience.

For simplicity, we will write  $w \in \mathcal{B}_2(\mathbb{R}_+)$  or  $w \in L^{\infty}_{loc}(\mathbb{R}_+)$ . This notation will also be used for other component-wise properties.

**Remark 18.** An examples of such a weight which satisfies  $\mathcal{B}_2(\mathbb{R}_+)$  is given by w(x) with  $w_i(x) = |x|^a (1+|x|)^{(i+1)b}$  for  $0 \le a < 1$  and  $b \in \mathbb{R}$ .

**Theorem 19.** Let  $1 \le p < \infty$ ,  $w \in \mathcal{B}_2(\mathbb{R}_+)$ , then  $W_w^{m,p}$  is separable. If  $1 , <math>W_w^{m,p}$  is reflexive.

The proof follows the same lines as in the non-weighted setting (see [13, Theorem 1.3] and also [51, Section 2]).

We state the following embedding theorem on  $\mathbb{R}^d$  instead of  $\mathbb{R}_+$ , as this general setting is of interest on its own. For  $\mathbb{R}_+$ , the proof works analogously with the only difference that instead of balls of radius R in  $\mathbb{R}^d$ , denoted by  $B_R(0)$ , the intersections of balls in  $\mathbb{R}$  with  $\mathbb{R}_+$  is considered.

**Proposition 20.** Let w, w' and  $w_c$  be weight functions and let  $i \in (\mathbb{N} \cup \{0\})^d$  be a multi-index. We introduce the following conditions:

(1) There exists a constant c > 0, such that for every  $i \in (\mathbb{N} \cup \{0\})^d$  and  $x \in \mathbb{R}^d$ , assume that  $w_i(x) \geq c$ . Further there exists a  $0 < K \in \mathbb{R}$  such that for every  $\mathbb{R}_+ \ni R \geq K$ ,

$$\sup_{x \in B_R(0)} |w_i(x)| \le C_R.$$

(2) There exists a number K > 0, such that  $\frac{w_i(x)}{w'_i(x)}$  is monotonically decreasing, with respect to |x|, on  $B_K(0)^c$ .

(3) For  $i \in (\mathbb{N} \cup \{0\})^d$ ,

$$\lim_{R \to \infty} R^{i - \frac{d}{p}} \sup_{R < |x| < 2R} |w_c(x)| \left( \frac{1}{R} \int_R^{2R} |w_i^{-\frac{1}{p-1}}(x)| \, \mathrm{d}x \right)^{\frac{\rho - 1}{p\rho}} < \infty.$$

(4) For  $i \in (\mathbb{N} \cup \{0\})^d$ ,

$$\lim_{R \to \infty} R^{i - \frac{d}{p}} \left( \frac{1}{R} \int_{R}^{2R} |w_i^{-\frac{1}{p-1}}(x)| \, \mathrm{d}x \right)^{\frac{\rho - 1}{p\rho}} < \infty.$$

(5) For  $i \in (\mathbb{N} \cup \{0\})^d$ ,

$$\lim_{R \to \infty} R^{i - \frac{d}{p}} \sup_{R < |x| < 2R} |w_c(x)| \inf_{R \le |x| \le 2R} \left(\frac{1}{w_i(x)}\right)^{\frac{1}{p}} < \infty.$$

(6) For  $i \in (\mathbb{N} \cup \{0\})^d$ ,

$$\lim_{R \to \infty} R^{i - \frac{d}{p}} \inf_{R \le |x| \le 2R} \left( \frac{1}{w_i(x)} \right)^{\frac{1}{p}} < \infty.$$

- (7) Let  $w \in L^{\infty}_{loc}(\mathbb{R}^d)$ ,  $w \in \mathcal{B}_{\rho}$ ,  $\rho > 1$ , and for every  $i \geq 0$ . The following statements hold.
  - (1) Let conditions 1, 2 be satisfied. For 0 < d (s l)p < d and  $\frac{1}{p} \frac{s}{d} < \frac{1}{q} \frac{l}{d}$   $(1 \le q < \frac{dp}{(d (s l)p)})$ , the embedding

$$W^{s,p}_{\frac{1}{w}} \cap W^{l,q}_{\frac{1}{w'}} \hookrightarrow W^{l,q}_{\frac{1}{w}}$$

is compact.

In the case d = (s - l)p, the embedding is compact for every  $1 \le q < \infty$ .

(2) Let  $p \ge 1$ ,  $d \ge 1$ ,  $s, r \in \mathbb{N} \cup 0$  and let us define

$$V := C^r(\mathbb{R}^d) \cap C_b(\mathbb{R}^d) \qquad if \ p = 1 \ and \ s - r \ge d$$

$$V := C^{r,\eta}(\mathbb{R}^d) \cap C_b(\mathbb{R}^d) \qquad if \ (s - r) \ge \frac{d}{p} \ge (s - r) - 1 \ and \ 0 < \eta < (s - r) - \frac{d}{p}$$

$$V := C^{r,1}(\mathbb{R}^d) \cap C_b(\mathbb{R}^d) \qquad if \ d = (s - r) - 1 \ and \ p = 1$$

• Let Assumptions 3; 4 and 7 hold. If there exists an  $i \in (\mathbb{N} \cup \{0\})^d$ , such that  $w_i(0) = 0$ , then the embedding

$$W_w^{s,p\rho} \hookrightarrow V \cap C_{w_c}(\mathbb{R}^d),$$

is continuous.

• If Assumption 1, 5 and 6 hold, then the embedding

$$W_w^{s,p} \hookrightarrow V \cap C_{w_c}(\mathbb{R}^d)$$

is continuous.

The proof can be found in the Appendix, Section 5.1.

Corollary 21. Let d=1 and  $w, w_c$  be weight functions satisfying Assumption 1. Further, for each  $i \geq 0$  there exits a constant C>0, such that  $|\partial_x^i w_c(x)| \leq C_i w_c(x)$  for each  $x \in \mathbb{R}$ . w satisfies  $\lim_{R\to 0} R^{i-\frac{1}{p}} \inf_{R\leq |x|\leq 2R} \left(\frac{1}{w_i(x)}\right)^{\frac{1}{p}} < \infty$ . Then  $\|w_c \varphi\|_{L^{\infty}(\mathbb{R})} \leq C \|\varphi\|_{W_{\bar{w}}^{m,p}(\mathbb{R})}$ , where  $\widetilde{w}_i(x) \leq w_c(x)^p w_i(x)$ .

**Remark 22.** If we replace the finite limit in Assumptions 4, 6 with 0, we obtain that  $\lim_{x\to\infty} |u(x)| = 0$  uniformly.

Remark 23. Assumption 1 can be avoided, if one considers weights, which behave on  $B_R(0)$  like Muckenhoupt weights and used the established theory covering this type of weighted space (see e.g. [55]). This way one could allow for singularities at 0, e.g. for d = 1,  $x^{-a}(a+x)^{-b}$  with  $0 \le a < 1$  and  $b \in \mathbb{R}$ . The arguments are similar however, at later stages of the paper this would lead to further technicalities, which we want to avoid. We will hint at one reason in Remark 47 why switching to such weights would not significantly improve our estimates to warrant the additional technicalities.

Later on, we focus on a particular choice of weights, however, if we do not specify that our weight is of a specific form then we impose the following standing assumption for any weight in the remainder of this work.

**Assumption 24.** Let  $m \in \mathbb{N}$  be given. We assume that the weight function  $w = (w_0, \ldots, w_m)$  satisfies  $w_j \in \mathcal{B}_2(\mathbb{R}_+)$  as well as  $w_j \in L^{\infty}_{loc}$ , for every  $i \in \{0, \ldots, k\}$ . For a given weight function w, let  $\frac{1}{w}$  also be a weight function satisfying the above assumptions.

We identify the dual space of  $W_w^{m,p}$ , via the (unweighted)  $L^2$  duality with  $W_{\frac{1}{w}}^{-m,q}(\mathbb{R}_+)$ , whose norm is given by

$$||v||_{W_{\frac{1}{w}}^{-m,q}} = \sup_{u: ||u||_{W_{\frac{1}{w}}^{m,p}} = 1} |\langle v, u \rangle|.$$

In other words, we work on the following triple(s) of spaces

$$W_w^{m,p} \hookrightarrow L^2 \hookrightarrow W_{\frac{1}{w}}^{-m,q}$$

Remark 25. In particular  $(L_w^2)' = W_{\frac{1}{w}}^{0,2}$ , where

$$||u||_{L^{2}_{\frac{1}{2n}}}^{2} = \int_{0}^{\infty} \frac{|u(x)|^{2}}{w(x)} dx.$$

**Remark 26.** If  $w(x) = (1+x^2)^{b/2}$  for some  $b \in \mathbb{R}$ ,  $\nu$  is usually referred to as a tempered measure. We will slightly abuse the notation "tempered" and refer to measures  $\nu$ , satisfying  $\int_{\mathbb{R}} \frac{1}{w(x)} \nu(\,\mathrm{d}x) < \infty$ , as being (w-)tempered.

In the following Lemma, we identify a w-tempered measure with an element in the dual of certain weighted Sobolev spaces.

**Lemma 27.** Let  $w_c$ ,  $\widetilde{w}$  be a weight function satisfying the conditions of Corollary 21. Let  $d=1,\ p=2,\ i=1,2.$  If  $\nu$  is a non-negative, measure on  $\mathbb{R}_+$ , such that  $\int_{\mathbb{R}_+} \frac{1}{w_c(x)} \nu(\,\mathrm{d} x) < \infty$ , then  $\nu \in \left(W_{\widetilde{w}}^{1,2}\right)' = W_{\frac{1}{\widetilde{w}}}^{-1,2}$ .

*Proof.* Proposition 20 yields that  $\varphi \in W^{1,2}_{w'}$  is continuous and bounded on  $\mathbb{R}_+$  and in particular  $\varphi \in C_{w_c}(\mathbb{R}_+)$ . Hence, the dual pairing with a (w-tempered) Radon measure is well-defined.

$$\sup_{\|\varphi\|_{W^{1,2}_{\widetilde{\alpha}}=1}} |\langle \nu, \varphi \rangle| = \sup_{\|\varphi\|_{W^{1,2}_{\widetilde{\alpha}}=1}} \left| \int_{\mathbb{R}_+} \varphi(x) \nu(\,\mathrm{d}x) \right|$$

$$= \sup_{\|\varphi\|_{W_{\widetilde{w}}^{1,2}=1}} \left| \int_{\mathbb{R}_{+}} \varphi(x) \frac{w_{c}(x)}{w_{c}(x)} \nu(\,\mathrm{d}x) \right|$$

$$\leq \sup_{\|\varphi\|_{W_{\widetilde{w}}^{1,2}=1}} \|w_{c}\varphi\|_{L^{\infty}(\mathbb{R}_{+})} \left| \int_{\mathbb{R}_{+}} \frac{1}{w_{c}(x)} \nu(\,\mathrm{d}x) \right|$$

$$\leq C \sup_{\|\varphi\|_{W_{\widetilde{w}}^{1,2}=1}} \|\varphi\|_{W_{\widetilde{w}}^{1,2}} \left| \int_{\mathbb{R}_{+}} \frac{1}{w_{c}(x)} \nu(\,\mathrm{d}x) \right|.$$

**Remark 28.** In Proposition 20, as well as Lemma 27, it is possible to replace  $w_c$  and  $\frac{1}{w}$  by weights of the for  $\frac{1}{w_c}$  and w respectively, where  $w_c$  as well as w are increasing towards  $\infty$  and  $\partial_x \frac{1}{w} \leq C \frac{1}{w}$ , for some nonnegative constant C. The arguments are identical but are skipped for brevity.

We note that there might be many choices of weights, for which  $\int_{\mathbb{R}} \frac{1}{w(x)} \nu(\,\mathrm{d}x) < \infty$  is satisfied and some might be more canonical than others, given a particular measure  $\nu$  and application in mind. We will see that weights that behave like  $w(x) = (1+x)^{\beta}$  or  $w(x) = (1+x^2)^{\frac{\beta}{2}}$ , for  $\beta \in \mathbb{R}$  will appear naturally in our analysis. However, these weights will often appear in the form  $w(x) = (c_1 + c_2 x)^{\beta}$  with  $c_1, c_2 > 0$ . For this reason, we will introduce an equivalence relation on the family of such weights.

**Definition 29.** Let  $a, b, r, s, x \ge 0$ . We call two weights  $(a + bx)^{\beta}$  and  $(r + sx)^{\beta}$  equivalent, denoted by  $(a + bx)^{\beta} \cong (r + sx)^{\beta}$ , if there exist constants  $c_1, c_2 > 0$ , such that  $(c_1a + c_2bx)^{\beta} = (r + sx)^{\beta}$ .

It can be easily checked that this indeed defines an equivalence relation and the set of functions  $w(x) = (c_1 + c_2 x)^{\beta}$  with  $c_1, c_2 > 0$ . Next, we verify that the two equivalent weights, in terms of the relation specified above, give rise to equivalent weighted Sobolev norms, in the sense that they induce the same topology. This result will be used in later calculations where, rather than keeping track of constants appearing in the weight, we collect them via the inequality stated in the following Lemma in a constant in front of the norm.

**Lemma 30.** Let  $w_{1,i}(x) = (a_{1,i} + a_{2,i}x)^{\beta_i}$  and  $w_{2,i}(x) = (b_{1,i} + b_{2,i}x)^{\beta_i}$  be two equivalent weights with  $a_1, a_2, b_1, b_2 > 0$ . Then  $\|\cdot\|_{W^{m,p}_{w_1}}$  and  $\|\cdot\|_{W^{m,p}_{w_2}}$  are equivalent in the sense of norms, in the sense that there exist two constants  $c \leq C > 0$  which only depend on  $a_{1,i}, a_{2,i}, b_{1,i}, b_{2,i}$  and  $\beta_i$ , such that

$$c\|\cdot\|_{W^{m,p}_{w_2}} \le \|\cdot\|_{W^{m,p}_{w_1}} \le C\|\cdot\|_{W^{m,p}_{w_2}}.$$

Additionally  $\|\cdot\|_{W^{m,p}_{\frac{1}{w_1}}}$  and  $\|\cdot\|_{W^{m,p}_{\frac{1}{w_2}}}$  are equivalent

**Remark 31.** We want to keep the following trivial estimates in mind. For  $\eta > 0$  and  $x \geq 0$ ,

$$\frac{1+x}{1+\eta x} \le 1 \lor \frac{1}{\eta}, \quad \frac{1+x}{\eta+x} \le 1 \lor \frac{1}{\eta}.$$

For convenience, we want to state a Corollary of Proposition 20, specified to the settings where our weight components are of the form  $w_i(x) = (1+x)^{\beta_i}$ . We will need a simple lemma, telling us that the terms we encounter due to estimates on the annuli  $A_R$  can be equivalently expressed in terms of such weights.

**Lemma 32.** Let  $R \geq 1$ ,  $w = (w_1, ..., w_s)$  with  $w_i = (1+x)^{ip-d}$ . Recall the notation  $A_R := \{x : R \leq |x| \leq 2R\}$ . Then, there exist constants  $0 < \widetilde{c} \leq \widetilde{C} < \infty$ , such that for any  $l \geq 0$ ,  $1 \leq p < \infty$ ,

$$\widetilde{c}\|u\|_{W_w^{m,p}(A_R)} \le \left(\sum_{j=0}^m R^{jp-d}\|D^j u\|_{L^p(A_R)}^p\right)^{1/p} \le \widetilde{C}\|u\|_{W_w^{m,p}(A_R)}.$$

**Corollary 33** (of Proposition 20). Let the assumptions of Proposition 20 be satisfied and  $w = (w_1, \ldots, w_s)$  with  $w_i = (1+x)^{ip-d}$ , then the embedding  $W_w^{m,p} \hookrightarrow C_b$  is continuous.

Proof. Let  $R \geq 1, m \geq 0$  and  $w = (w_1, \dots, w_l)$ , with  $w_i = (1+x)^{ip-d}$ . On  $B_R(0)$ , the norms  $\|\cdot\|_{W^{m,p}}$  and  $\|\cdot\|_{W^{m,p}}$  are equivalent we obtain  $u \in C_b(B_R)$ . Lemma 32 combined with the arguments in the proof of Proposition 20 now yield that u is continuous everywhere and  $\|u\|_{L^{\infty}(A_R)} \leq C\|u\|_{W^{m,p}_w}$ .

**Lemma 34.** Let  $\nu$  be a non-negative radon measure and assume that there exists a  $\theta_{\nu} \in \mathbb{R}$ , such that  $\int_0^{\infty} \frac{1}{(1+x)^{\theta_{\nu}}} \nu(\,\mathrm{d}x) < \infty$ , then  $\nu \in W_{\frac{1}{n}}^{-1,2}$  with  $w_i(x) = (1+x)^{-1+2\theta_{\nu}+2i}$ .

*Proof.* Since  $\|(1+x)^{\theta_{\nu}}\varphi\|_{L^{\infty}(B_{R}(0))} \leq C(w,R,\theta_{\nu})\|\varphi\|_{W^{1,2}_{w}}$  for any weight w which is bounded from above and below on  $B_{R}(0)$ , it suffices to consider the estimates on the annuli  $A_{R}$ . We consider only the case  $\theta_{\nu} \geq 0$ , as the case  $\theta_{\nu} \leq 0$  is identical, but with a different constant.

$$||(1+x)^{\theta_{\nu}}\varphi||_{L^{\infty}(A_{R})} \leq C2^{\theta_{\nu}}(1+R)^{\theta_{\nu}} \left(\sum_{j=0}^{1} \int_{R}^{2R} |D^{j}u(x)|^{2} (1+x)^{j2-1} dx\right)^{1/2}$$

$$= C \left(\sum_{j=0}^{1} \int_{R}^{2R} |D^{j}u(x)|^{2} (1+R)^{2\theta_{\nu}} (1+x)^{j2-1} dx\right)^{1/2}$$

$$\leq C \left(\sum_{j=0}^{1} \int_{R}^{2R} |D^{j}u(x)|^{2} (1+x)^{2\theta_{\nu}} (1+x)^{j2-1} dx\right)^{1/2}$$

$$= C \left(\sum_{j=0}^{1} \int_{R}^{2R} |D^{j}u(x)|^{2} (1+x)^{2\theta_{\nu}} (1+x)^{j2-1} dx\right)^{1/2}.$$

## 3. The lifted equation

In this section, we obtain a solution theory for (1.8) under stricter conditions and obtain bounds on the solutions. Let us fix  $0 < T < \infty$  and unless explicitly stated otherwise, we will study our given problem on the finite time interval [0, T].

**Remark 35.** The multiplication of elements with negative Sobolev regularity and functions -x and  $e^{-tx}$  should be interpreted in the dual sense, i.e.  $\langle -x\mu, \varphi \rangle \coloneqq \langle \mu, -x\varphi \rangle$  and  $\langle e^{-(t-s)x}\nu, \varphi \rangle \coloneqq \langle \nu, e^{-(t-s)x}\varphi \rangle$ , for appropriate test functions  $\varphi$ .

3.1. **The operator and semigroup.** Before we proceed with the solution theory, we first want to understand better the operators and semigroup we will work with and how they act on our choice of spaces. This will be the content of this subsection.

Let us briefly recall the strong form on (1.8).

(3.1) 
$$\mu_t = \mu_0 - \int_0^t x \mu_s \, \mathrm{d}s + \int_0^t \nu_b b(\langle \mu_s, 1 \rangle) \, \mathrm{d}s + \int_0^t \nu_\sigma \sigma(\langle \mu_s, 1 \rangle) \, \mathrm{d}W_s.$$

The operator of primary interest will be the multiplication operator  $f \mapsto -xf$ .

**Lemma 36.** Let  $1 \le p < \infty$ . The family  $(S_t)_t := (e^{-t \cdot})_t$  of linear operators, where  $e^{-t \cdot}: f \mapsto (x \mapsto f(x)e^{-tx})$  is

(1) a strongly continuous contraction semigroup on  $W_w^{0,p}$ , a strongly continuous semigroup on  $W_w^{m,p}$   $(m \ge 0)$ . It has a densely defined generator, given by

$$(Af)(x) := -xf(x), \quad x \ge 0,$$
  
$$\mathcal{D}(A) := \left\{ f \in W_w^{m,p} \colon \| -xf \|_{W_w^{m,p}} < \infty \right\}.$$

(2) The adjoint semigroup  $(S_t^*)_t$  is a strongly continuous semigroup on  $W_{\frac{1}{w}}^{-m,q}$ ,

*Proof.* (1) Let  $m \ge 0$  and  $f \in W_w^{m,p}$ , then

$$||S_{t}f - f||_{W_{w}^{m,p}} = \left(\sum_{j=0}^{m} \int_{\mathbb{R}_{+}} |D^{j}e^{-tx}f(x) - D^{j}f(x)|^{p} w_{j}(x) dx\right)^{\frac{1}{p}}$$

$$= \left(\sum_{j=0}^{m} \int_{\mathbb{R}_{+}} |D^{j}(f(x)(e^{-tx} - 1))|^{p} ww_{j}(x) dx\right)^{\frac{1}{p}}$$

$$= \left(\sum_{j=0}^{m} \int_{\mathbb{R}_{+}} \left|\sum_{i=0}^{j} {j \choose i} D^{j-i}(e^{-tx} - 1) D^{i}f(x)\right|^{p} w_{j}(x) dx\right)^{\frac{1}{p}}$$

$$= \left(\sum_{j=0}^{m} \int_{\mathbb{R}_{+}} \left|(e^{-tx} - 1) D^{j}f(x) + \sum_{i=0}^{j-1} {j \choose i} (-t)^{j-i}e^{-tx}D^{i}f(x)\right|^{p} w_{j}(x) dx\right)^{\frac{1}{p}}.$$

By the dominated convergence theorem, we conclude that  $\lim_{t\to 0} \|S_t f - f\|_{W_w^{m,p}} = 0$ . We define  $A: \mathcal{D}(A) \to W_w^{m,p}$  by

$$(Af)(x) := -xf(x), \quad x \ge 0,$$
  
$$\mathcal{D}(A) := \left\{ f \in W_w^{m,p} \colon \| -xf \|_{W_w^{m,p}} < \infty \right\}.$$

Note that  $\mathcal{D}(A) \supset C_0^{\infty}$ , which is dense in  $W_w^{m,p}$  by definition.

$$\frac{1}{t} \|S_t f - f - tAf\|_{W_w^{m,p}} = \frac{1}{t} \left( \sum_{j=0}^m \int_{\mathbb{R}_+} \left| D^j \left( \left( e^{-tx} - 1 + x \right) f(x) \right) \right|^p w_j(x) \, \mathrm{d}x \right)^{\frac{1}{p}} \\
= \frac{1}{t} \left( \sum_{j=0}^m \int_{\mathbb{R}_+} \left| \sum_{i=0}^j \binom{j}{i} D^{j-i} \left( e^{-tx} - 1 + tx \right) D^i f(x) \right|^p w_j(x) \, \mathrm{d}x \right)^{\frac{1}{p}}$$

$$\begin{split} &= \frac{1}{t} \left( \sum_{j=0}^{m} \int_{\mathbb{R}_{+}} \left| \left( e^{-tx} - 1 + tx \right) D^{j} f(x) + j \left( -te^{-tx} + t \right) D^{j-1} f(x) \right. \\ &\quad + \left. \sum_{i=0}^{j-2} \binom{j}{i} \left( -t \right)^{j-i} e^{-tx} D^{i} f(x) \right|^{p} w_{j}(x) \, \mathrm{d}x \right)^{\frac{1}{p}} \\ &\leq C \left( \sum_{j=0}^{m} \int_{\mathbb{R}_{+}} \left| \frac{\left( e^{-tx} - 1 + tx \right)}{t} D^{j} f(x) + j \left( 1 - e^{-tx} \right) D^{j-1} f(x) \right|^{p} w_{j}(x) \, \mathrm{d}x \right)^{\frac{1}{p}} + O^{t \to 0}(t) \\ &\leq C \left( \sum_{j=0}^{m} \int_{\mathbb{R}_{+}} \left| \left| 1 - \frac{1}{t} \int_{0}^{t} e^{-sx} \, \mathrm{d}s \right| |D^{j} f(x)| + j \left| 1 - e^{-tx} \right| |D^{j-1} f(x)| \right|^{p} w_{j}(x) \, \mathrm{d}x \right)^{\frac{1}{p}} + O^{t \to 0}(t), \end{split}$$

where  $O^{t\to 0}(t)$  denotes terms of order t, as  $t\to 0$ . By the dominated convergence theorem, the right-hand side vanishes, as  $t\to 0$ . The contraction property, when k=0, can be seen immediately, since  $|e^{-tx}| \le 1$  for any  $0 \le t, x$ .

(2)

$$|\langle S_t^* \nu - \nu, f \rangle| = |\langle \nu, (S_t - 1) f \rangle|.$$

By the same arguments as above the semigroup is weakly continuous, i.e. for every  $f \in W_w^{m,p} \lim_{t\to 0} |\langle S_t^*\nu - \nu, f \rangle| = 0$ . By [37, Theorem 5.8], the semigroup  $S^*$  is even strongly continuous on  $W_{\frac{1}{w}}^{-m,p}$ .

Remark 37. We can define

$$A^*\nu := \operatorname{weak}^* \lim_{t \to 0} \frac{1}{t} \left( S_t^* \nu - \nu \right)$$

on the domain

$$\mathcal{D}\left(A^{*}\right) \coloneqq \left\{\nu \in W_{\frac{1}{w}}^{-m,p} \colon \operatorname{weak}^{*} \lim_{t \to 0} \frac{1}{t} \left(S_{t}^{*} \nu - \nu\right) \ \operatorname{exists} \ \right\}.$$

 $A^*$  is a weak\*-closed and weak\*-densely defined operator and coincides with the adjoint  $A^*$  of A (see [37, Definition B.8]), i.e.,

$$\mathcal{D}(A^*) := \left\{ \nu \in W_{\frac{1}{w}}^{-m,p} : \begin{array}{c} \text{there exists } \eta \in W_{\frac{1}{w}}^{-m,p} \text{ such that} \\ \langle f, \eta \rangle = \langle Af, \nu \rangle \text{ for all } f \in \mathcal{D}(A) \end{array} \right\},$$

and  $A^*$  is the adjoint of A. By [37, Corollary B. 12] it then follows that  $\sigma(A^*) = \sigma(A)$ .

Also note that in our case the adjoint semigroup  $(S_t^*)_t$  coincides with the so-called sun-dual semigroup of  $(S_t)_t$  (see [37, Section 2.6]).

From this point onwards, we set p = 2 and restrict our analysis to the Hilbert-space case. In the next lemma, we want to investigate certain mapping properties of the semigroup  $S^*$ .

## 3.1.1. Estimates on Hilbert spaces.

**Lemma 38.** Let 0 < t and let  $\eta \in W_{\frac{1}{w}}^{-m,2}$ , then  $S^* \eta \in W_{\frac{1}{w}}^{-m,2}$  with  $(1+x)^{-2\gamma} w_i(x) \leq \widetilde{w}_i(x)$ , for any  $\gamma \in [0,1]$ . If  $0 < t \leq T < \infty$ ,

$$||S_t^*\eta||_{W_{\frac{1}{w}}^{-m,2}} \le C||\eta||_{W_{\frac{1}{w}}^{-m,2}} \frac{(1 \lor T)^{\gamma}(1+t^m)}{t^{\gamma}}.$$

Proof.

$$\begin{split} \|S_t^*\eta\|_{W_{\overline{w}}^{-m,2}} &= \sup_{\psi \colon \|\psi\|_{W_{\overline{w}}^{m,2}=1}} |\langle \eta, S_t \psi \rangle| \\ &\leq \|\eta\|_{W_{\overline{w}}^{-m,2}} \sup_{\psi \colon \|\psi\|_{W_{\overline{w}}^{m,2}=1}} \sqrt{\sum_{j=0}^m \int_{\mathbb{R}_+} |D^j(e^{-tx}\psi(x))|^2 w_j(x) \, \mathrm{d}x} \\ &\leq \|\eta\|_{W_{\overline{w}}^{-m,2}} \sup_{\psi \colon \|\psi\|_{W_{\overline{w}}^{m,2}=1}} \sqrt{\int_{\mathbb{R}_+} \sum_{j=0}^m \left|\sum_{i=1}^j \binom{j}{i} \left(-t\right)^{j-i} e^{-tx} D^i \psi(x)\right|^2 w_j(x) \, \mathrm{d}x} \\ &\leq C \|\eta\|_{W_{\overline{w}}^{-m,2}} \sup_{\psi \colon \|\psi\|_{W_{\overline{w}}^{m,2}=1}} \sqrt{\int_{\mathbb{R}_+} (1+t^m)^2 \sum_{j=0}^m \int_{\mathbb{R}_+} |D^j \psi(x)|^2 \left(\frac{1}{1+tx}\right)^{2\gamma} w_j(x) \, \mathrm{d}x}, \end{split}$$

for any  $\gamma \in [0,1]$ . Hence,

$$\begin{split} \|S_t^*\eta\|_{W_{\frac{1}{w}}^{-m,2}} &\leq C\|\eta\|_{W_{\frac{1}{w}}^{-m,2}}(1+t^m) \sup_{\psi\colon \|\psi\|_{W_{\widetilde{w}}^{m,2}}=1} \sqrt{\int_{\mathbb{R}_+} \sum_{j=0}^m \int_{\mathbb{R}_+} |D^j\psi(x)|^2 \frac{1}{t^{2\gamma}} \left(\frac{(1\vee T)}{1+x}\right)^{2\gamma} w_j(x) \,\mathrm{d}x} \\ &\leq C\|\eta\|_{W_{\frac{1}{w}}^{-m,2}} \frac{(1+t^m)(1\vee T)^{\gamma}}{t^{\gamma}} \sup_{\psi\colon \|\psi\|_{W_{\widetilde{w}}^{m,2}}=1} \sqrt{\int_{\mathbb{R}_+} \sum_{j=0}^m \int_{\mathbb{R}_+} |D^j\psi(x)|^2 \left(\frac{1}{1+x}\right)^{2\gamma} w_j(x) \,\mathrm{d}x} \\ &\leq C\|\eta\|_{W_{\frac{1}{w}}^{-m,2}} \frac{(1+t^m)(1\vee T)^{\gamma}}{t^{\gamma}} \sup_{\psi\colon \|\psi\|_{W_{\widetilde{w}}^{m,2}}=1} \|\psi\|_{W_{\widetilde{w}}^{m,2}}, \end{split}$$

where 
$$\left(\frac{1}{1+x}\right)^{2\gamma}w(x) \leq \widetilde{w}$$
.

We will require one additional Lemma, which will be important in a later section.

**Lemma 39.** Let  $\eta \in W_{\frac{1}{w}}^{-m,p}$ ,  $\gamma \in [0,1]$  and  $\widetilde{w}$  such that  $(1+x)^{2\gamma}w_i(x) \leq \widetilde{w}_i(x)$ , for every  $i \geq 0$ . If  $|t-s| \leq 1$  then

$$||S_{t-s}^*S_s^*\eta - S_s^*\eta||_{W_{\frac{1}{\widetilde{w}}}^{-m,2}} \le C_\gamma ||\eta||_{W_{\frac{1}{w}}^{-m,2}} (1 \vee s)^m (t-s)^\gamma,$$

as well as

$$||S_{t-s}^*\eta - \eta||_{W_{\frac{1}{m}}^{-m,2}} \le C_{\gamma} ||\eta||_{W_{\frac{1}{m}}^{-m,2}} (t-s)^{\gamma}.$$

If 
$$|t-s| > 1$$

$$||S_{t-s}^* S_s^* \eta - S_s^* \eta||_{W_{\frac{1}{m}}^{-m,2}} \le C ||\eta||_{W_{\frac{1}{m}}^{-m,2}} (1 \vee s)^m (t-s)^m.$$

*Proof.* As in the previous lemma,

$$\|\varphi\|_{W_{\frac{1}{w}}^{m,2}} = \left(\sum_{i=0}^{m} \int_{\mathbb{R}_{+}} |\partial_{x}^{i}(e^{-sx}(e^{-(t-s)x} - 1)\varphi)|^{p} w_{i}(x) dx\right)^{\frac{1}{p}}$$

$$= \left(\sum_{i=0}^{m} \int_{\mathbb{R}_{+}} \left|\sum_{j_{1}+j_{2}+j_{3}=i} {i \choose j_{1}, j_{2}, j_{3}} \partial_{x}^{j_{1}}(e^{-sx}) \partial_{x}^{j_{2}}(e^{-(t-s)x} - 1) \partial_{x}^{j_{3}} \varphi\right|^{2} w_{i}(x) dx\right)^{\frac{1}{2}}.$$

For  $j \in \mathbb{N}$ ,

$$\partial_x^j(e^{-sx}) = (-1)^j s^j e^{-sx}$$
  
$$\partial_x^j(e^{-(t-s)x} - 1) = (-1)^j (t-s)^j e^{-(t-s)x}.$$

Also,

$$\begin{aligned} |\partial_x^{j_1}(e^{-sx})\partial_x^{j_2}(e^{-(t-s)x}-1)| &= \left| (-1)^{j_1}s^{j_1}e^{-sx}(-1)^{j_2}(t-s)^{j_2}e^{-(t-s)x} \right| \\ &\leq \left| s^{j_1}(t-s)^{j_2} \right|. \end{aligned}$$

If  $j_1 \in \mathbb{N}_0, j_2 = 0$ ,

$$|\partial_x^{j_1}(e^{-sx})(e^{-(t-s)x}-1)| \le C_{\gamma} |s^{j_1}(t-s)^{\gamma}(1+x)^{\gamma}|,$$

for any  $0 \le \gamma \le 1$ . Hence, if  $|t - s| \le 1$ ,

$$\left(\sum_{i=0}^{m} \int_{\mathbb{R}_{+}} \left| \sum_{j_{1}+j_{2}+j_{3}=i} {i \choose j_{1}, j_{2}, j_{3}} \partial_{x}^{j_{1}}(e^{-sx}) \partial_{x}^{j_{2}}(e^{-(t-s)x} - 1) \partial_{x}^{j_{3}} \varphi \right|^{2} w_{i}(x) dx \right)^{\frac{1}{2}}$$

$$\leq C_{\gamma} \left(\sum_{i=0}^{m} \int_{\mathbb{R}_{+}} (1 \vee s)^{2i} (t-s)^{2\gamma} (1+x)^{2\gamma} \left| \partial_{x}^{i} \varphi \right|^{2} w_{i}(x) dx \right)^{\frac{1}{2}}$$

$$\leq C_{\gamma} (1 \vee s)^{m} (t-s)^{\gamma} \left(\sum_{i=0}^{m} \int_{\mathbb{R}_{+}} \left| \partial_{x}^{i} \varphi \right|^{2} (1+x)^{2\gamma} w_{i}(x) dx \right)^{\frac{1}{2}}$$

To estimate  $\|S_{t-s}^*S_s^*\eta - S_s^*\eta\|_{W_{\frac{1}{w}}^{-m,2}}$ , we use that  $\eta \in W_{\frac{1}{w}}^{-m,2}$  and the previous estimates,

$$\begin{split} & \|S_{t-s}^* S_s^* \eta - S_s^* \eta\|_{W_{\frac{1}{w}}^{-m,2}} = \sup_{\|\varphi\|_{W_{\overline{w}}^{m,p}} = 1} |\langle \eta, S_s(S_{t-s} - 1)\varphi \rangle| \\ & \leq \sup_{\|\varphi\|_{W_{\overline{w}}^{m,2}} = 1} \|\eta\|_{W_{\frac{1}{w}}^{-m,2}}^2 \|S_s(S_{t-s} - 1)\varphi\|_{W_w^{m,2}} \\ & \leq C_\gamma \sup_{\|\varphi\|_{W_{\overline{w}}^{m,2}} = 1} \|\eta\|_{W_{\frac{1}{w}}^{-m,2}} (1 \vee s)^k (t-s)^\gamma \left( \sum_{i=0}^m \int_{\mathbb{R}_+} |\partial_x^i \varphi|^2 (1+x)^{2\gamma} w_i(x) \, \mathrm{d}x \right)^{\frac{1}{2}}. \end{split}$$

Hence, if  $(1+x)^{2\gamma}w_i(x) \leq \widetilde{w}_i(x)$ , for every  $i \geq 0$ , we obtain

$$||S_{t-s}^*S_s^*\eta - S_s^*\eta||_{W_{\frac{1}{w}}^{-m,2}} \le C_\gamma ||\eta||_{W_{\frac{1}{w}}^{-m,2}} (1 \lor s)^m (t-s)^\gamma.$$

The second inequality follows similarly:

$$||S_{t-s}\varphi - \varphi||_{W_{\frac{1}{w}}^{m,2}} = \left(\sum_{i=0}^{m} \int_{\mathbb{R}_{+}} |\partial_{x}^{i}((e^{-(t-s)x} - 1)\varphi)|^{2} w_{i}(x) dx\right)^{\frac{1}{2}}$$
$$= \left(\sum_{i=0}^{m} \int_{\mathbb{R}_{+}} \left|\sum_{j_{1}+j_{2}=i} {i \choose j_{1}, j_{2}} \partial_{x}^{j_{1}}(e^{-(t-s)x} - 1)\partial_{x}^{j_{2}}\varphi\right|^{2} w_{i}(x) dx\right)^{\frac{1}{2}}.$$

For  $j \in \mathbb{N}$ ,

$$\left|\partial_x^j (e^{-(t-s)x} - 1)\right| = \left|(-1)^j (t-s)^j e^{-(t-s)x}\right| \le \left|(t-s)^{j_2}\right|.$$

Hence, if  $|t - s| \le 1$ ,

$$\left(\sum_{i=0}^{m} \int_{\mathbb{R}_{+}} \left| \sum_{j_{1}+j_{2}=i} {i \choose j_{1}, j_{2}} \partial_{x}^{j_{1}} (e^{-(t-s)x} - 1) \partial_{x}^{j_{2}} \varphi \right|^{2} w_{i}(x) dx \right)^{\frac{1}{2}}$$

$$\leq C_{\gamma} \left( \sum_{i=0}^{m} \int_{\mathbb{R}_{+}} (t-s)^{2\gamma} (1+x)^{2\gamma} \left| \partial_{x}^{i} \varphi \right|^{2} w_{i}(x) dx \right)^{\frac{1}{2}}$$

$$\leq C_{\gamma} (t-s)^{\gamma} \left( \sum_{i=0}^{m} \int_{\mathbb{R}_{+}} \left| \partial_{x}^{i} \varphi \right|^{2} (1+x)^{2\gamma} w_{i}(x) dx \right)^{\frac{1}{2}}.$$

If |t-s| > 1, we can perform the same steps, but with  $|t-s|^k$ .

Considering the previous estimates and embeddings, the following lemma might seem redundant. However, it ties in more directly with Assumption (M 1) and gives more insight into how the weights in our spaces can be chosen, if we require  $||S_t^*\nu||_{W^{-1,2}_{\frac{1}{w}}} \in L^1(0,T)$  or  $||S_t^*\nu||_{W^{-1,2}_{\frac{1}{w}}} \in L^2(0,T)$ .

**Lemma 40.** Let  $\nu$  be a non-negative Radon measure on  $\mathbb{R}_+$ , such that there exists a  $0 \leq \theta_{\nu}$  for which  $\left| \int_{\mathbb{R}_+} \frac{1}{(1+x)^{\theta_{\nu}}} \nu(\,\mathrm{d}x) \right| < \infty$ . Let  $0 < t \leq T < \infty$  and consider the semigroup  $S_t^*$ , which was introduced in Lemma 36. Let  $\alpha \in \mathbb{R}$  and  $0 \leq \gamma < 1$  satisfy  $\theta_{\nu} \leq \gamma - \alpha$ . We set  $w_i = (1+x)^{-2\alpha-1+i2}$ , then

$$\|S_t^*\nu\|_{W_{\frac{1}{n}}^{-1,2}} \leq C \frac{1 \vee T^{\gamma}}{t^{\gamma}}.$$

Remark 41. We reiterated the crucial connection between the singularity of the kernel we lift  $(1 - \theta_{\nu})$ , the decay of the corresponding measure  $(\theta_{\nu})$ , the weighted space we work in (  $w_i = (1+x)^{-2\alpha-1+i2}$  with  $\theta_{\nu} - \gamma = \alpha$ ) and the time integrability of  $\|S_t^*\nu\|_{W^{-1,2}_{\frac{1}{w}}}$  (indicated by  $\gamma$ ).

*Proof.* Using the embedding from Lemma 34,

$$\begin{split} \|S_t^*\nu\|_{W_{\frac{1}{w}}^{-1,2}} &= \sup_{\varphi \colon \|\varphi\|_{W_w^{1,2}} = 1} |\langle \varphi, S_t^*\nu \rangle| \\ &= \sup_{\varphi \colon \|\varphi\|_{W_w^{1,2}} = 1} \left| \int_{\mathbb{R}_+} e^{-xt} \varphi(x) \nu(\,\mathrm{d}x) \right| \\ &\leq \sup_{\varphi \colon \|\varphi\|_{W_w^{1,2}} = 1} \|(1+x)^{-\alpha} \varphi\|_{\infty} \left| \int_{\mathbb{R}_+} e^{-xt} (1+x)^{\alpha} \nu(\,\mathrm{d}x) \right| \\ &\leq \sup_{\varphi \colon \|\varphi\|_{W_w^{1,2}} = 1} \|(1+x)^{-\alpha} \varphi\|_{\infty} \left| \int_{\mathbb{R}_+} e^{-xt} (1-\gamma) \frac{1}{t^{\gamma}} \left( \frac{1}{\frac{1}{t}+x} \right)^{\gamma} (1+x)^{\alpha} \nu(\,\mathrm{d}x) \right| \\ &\leq (1 \lor t)^{\gamma} \frac{1}{t^{\gamma}} \sup_{\varphi \colon \|\varphi\|_{W_w^{1,2}} = 1} \|\varphi\|_{W_w^{1,2}} \left| \int_{\mathbb{R}_+} e^{-xt(1-\gamma)} \left( \frac{1}{1+x} \right)^{\gamma} (1+x)^{\alpha} \nu(\,\mathrm{d}x) \right|, \end{split}$$

with  $w_i = (1+x)^{-2\alpha-1+i2}$ .

3.2. **Existence.** The goal of this section is to prove an existence result for equations of the type (3.2) with  $\psi \in W_w^{1,2}$ ,  $b, \sigma$  being uniformly continuous functions in the t and x-variable and  $\nu_i = \nu_i(\,\mathrm{d}x)$  (i = 1, 2) being a non-negative, (tempered) measures on  $\mathbb{R}_+$ , whose Laplace transforms satisfy a certain relation similar to (1.3). We first consider  $b, \sigma$  to be Lipschitz continuous in x and obtain the existence and uniqueness of solutions in a standard way. In the next step, we consider more general coefficients, which we approximate by Lipschitz functions, and use a tightness result to obtain existence in this general setting.

**Definition 42.** Given  $k > \frac{1}{2}$ ,  $\psi \in W_w^{m,2}$  and  $\mu_0 \in W_{\frac{1}{w}}^{-m,2}$ . A  $W_{\frac{1}{w}}^{-m,2}$ -valued predictable process  $\mu_t$ ,  $t \in [0,T]$  is called a mild solution of equation

(3.2) 
$$\mu_t = \mu_0 - \int_0^t x \mu_s \, \mathrm{d}s + \int_0^t \nu_b b(s, \langle \mu_s, \psi \rangle) \, \mathrm{d}s + \int_0^t \nu_\sigma \sigma(s, \langle \mu_s, \psi \rangle) \, \mathrm{d}W_s,$$
if

$$(3.3) \quad \mu_t = e^{-xt}\mu_0 + \int_0^t e^{-x(t-s)}\nu_b(x)b(s,\langle\mu_s,\psi\rangle)\,\mathrm{d}s + \int_0^t e^{-x(t-s)}\nu_\sigma(x)\sigma(s,\langle\mu_s,\psi\rangle)\,\mathrm{d}W_s,$$

 $\mathbb{P}$ -a.s. for each  $t \in [0,T]$ . If  $\mu$  has continuous sample paths, we will call it a continuous, mild solution.

**Assumption 43.** Let  $L_{T,b}, L_{T,\sigma}, C_{T,b}, C_{T,\sigma} \geq 0$ . The functions  $b: \mathbb{R}_+ \times \mathbb{R} \to \mathbb{R}$  and  $\sigma: \mathbb{R}_+ \times \mathbb{R} \to \mathbb{R}$  are continuous in both arguments and satisfy the following bounds for any  $t \in [0,T]$ , (Coeff 1) Linear growth:

$$|b(t,x)| \le C_{T,b}(1+|x|),$$
  $|\sigma(t,x)| \le C_{T\sigma}(1+|x|).$ 

(Coeff 2) Lipschitz continuity: For every  $t \in \mathbb{R}_+$ ,  $x, y \in \mathbb{R}$ ,

$$|b(t,x) - b(t,y)| \le L_{T,b}|x - y|, \qquad |\sigma(t,x) - \sigma(t,y)| \le L_{T,\sigma}|x - y|.$$

(Coeff 3) There exist mouli of continuity  $\rho_b, \rho_\sigma$ , such that for every  $t \in \mathbb{R}_+$ ,  $x, y \in \mathbb{R}$ ,  $|b(t, x) - b(t, y)| < \rho_b(|x - y|)$ ,  $|\sigma(t, x) - \sigma(t, y)| < \rho_\sigma(|x - y|)$ .

Remark 44. There is no issue in considering coefficients which, in addition to x, t, also depend on  $\omega \in \Omega$ , as long as the constants in the assumption are uniform in  $\omega$ . One can also allow constants  $C_t$  that exhibit an  $L^2([0,T])$ -singularity at 0. This case is a straightforward adaptation of the statements in this paper, however, we refrain from including this case in our arguments to not overload the proofs with additional parameters.

Notation 45. When we write  $C_{\text{Lip}_b}$ ,  $C_{\text{Lip}_\sigma}$ ,  $C_{\text{LG}_\sigma}$ ,  $C_{\text{LG}_\sigma}$ , we implicitly refer to the constants appearing in the previous assumption, without specifying the T dependence explicitly. If a constant without a second subscript appears, like  $C_{\text{Lip}}$ ,  $C_{\text{Lg}}$ , it is implied that  $\max\{C_{\text{Lip}_b}, C_{\text{Lip}_\sigma}\} \leq C_{\text{Lip}}$  and  $\max\{C_{\text{LG}_b}, C_{\text{LG}_\sigma}\} \leq C_{\text{LG}}$ .

We impose the following assumptions on  $\nu_b$  and  $\nu_{\sigma}$ . We will discuss their interpretation and alternative formulations afterward.

**Assumption 46.** (M 1)  $\nu_b$  and  $\nu_\sigma$  are non-negative measures on  $\mathbb{R}_+$  and there exist  $0 \le \theta_{\nu_b}, \theta_{\nu_\sigma} < 1$ , such that

(3.4) 
$$\frac{1}{(1+|x|)^{\theta_{\nu_b\setminus\sigma}}}\nu_{b\setminus\sigma}(\mathbb{R}_+) < \infty.$$

For the measures from Assumption (M 1), Lemma 40 implies that there exists a weight w and constants  $0 < a_b, a_\sigma \le 1$ , such that

(3.5) 
$$||S_{t-s}^* \nu_{b \setminus \sigma}||_{W_{\frac{1}{w}}^{-1,2}} \le \frac{C(T)}{(t-s)^{1-a_{b \setminus \sigma}}},$$

where the value of  $a_b, a_\sigma$  are connected to the choice of weight w. Hence, Lemma 40 allows us to identify a weight, for which

(3.6) 
$$||S_{t-s}^*\nu_b||_{W_{\frac{1}{w}}^{-1,2}} \in L^1(0,T), \qquad ||S_{t-s}^*\nu_\sigma||_{W_{\frac{1}{w}}^{-1,2}} \in L^2(0,T).$$

(M 2) We assume that this w satisfies Assumption 24, and the constant 1 function has finite  $W_w^{1,2}$  norm, i.e.  $\int_{\mathbb{R}_+} w_0(x) \, \mathrm{d}x < \infty$ .

Remark 47. Discussion of the assumptions.

- For the most part, it would suffice to impose (3.6) as a general Assumption and we will use the previous implication extensively.
- Assumption (M 2) is not strictly necessary for our solution theory. It only comes into play, when we want to relate an explicit SVE to the solution of the SEE. To "reconstruct" the correct kernel, we will want to test the mild solution of (1.8) with the constant 1 function. To obtain the SVE, the function in the non-local terms needs to coincide with the functions we are testing with, i.e.  $\psi = 1$ . This also reflects that an  $L^2$ -kernel is generally required to make sense of the stochastic integral. Intuitively, being able to set  $\psi = 1$  correlates with the corresponding SVE being well-posed in the sense of functions. (3.4) with  $\frac{1}{2} \leq \theta_{\nu_{\sigma}} < 1$  would correspond to a kernel with an  $L^1$  singularity inside the stochastic integral which could, for  $\sigma = 1$ , be interpreted as a generalized fractional Gaussian field with Hurst index < 0 (see [52]).
- It might seem reasonable to alter (M 1) by: There exist  $0 \le \theta_1^{\nu}, \theta_2^{\nu} < 1$ , such that  $\frac{1}{x^{\theta_1^{\nu}}(1+|x|)^{\theta_2^{\nu}}}\nu(dx)$  is a finite measure on  $\mathbb{R}_+$ . If we wanted to test (3.3) with functions, that vanish sufficiently fast at 0, we could consider weights, which are not  $L^1_{loc}$  around 0, which would indeed provide better estimates. Since, later on, we want Assumption

(M 2) to hold, we will always require a contribution of  $\theta_2^{\nu} > 0$ . Such a contribution is obtained via the semigroup and always results in a time dependence of the coefficient appearing in (3.5).

**Example 48.** We illustrate potential choices of weights for common examples of completely monotone kernels. In these examples, we verify the bound (3.5) via (3.4). Taking Assumption (M 2) into account, we notice that the higher order weights  $w_j$  for  $j \geq 1$  play no particular role. Hence these can be chosen freely to satisfy certain conditions related to the associated Sobolev spaces.

Let  $t < \infty$ .

(1) Let  $K_{\text{exp}}$  be a finite combination of exponential functions:

$$K_{\text{exp}}(t) := \sum_{i=1}^{k} c_i e^{-x_i t}, t > 0,$$

with  $k \in \mathbb{N}$ ,  $c_i > 0$  and  $x_i \in [0, \infty)$  for  $i = 1, \ldots, k$ . The corresponding Radon measure is

$$\nu_{\exp}(\,\mathrm{d}x) = \sum_{i=1}^{k} c_i \delta_{x_i}(\,\mathrm{d}x),$$

where  $\delta_{x_i}$  denotes the Dirac measure at point  $x_i$ . In this example, we set  $\theta_{\nu} = 0$  corresponding to the unweighted case. Hence, we will allow ourselves to perform the estimates in fractional Sobolev spaces. For estimate (3.5), we obtain

$$\sup_{\|\varphi\|_{W^{1/2+\varepsilon,2}=1}} \left| \int_0^\infty e^{-(t-s)x} \psi(x) \nu(\,\mathrm{d}x) \right| \leq \sup_{\|\varphi\|_{W^{1/2+\varepsilon,2}=1}} \sum_{i=1}^k c_i e^{-(t-s)x_i} \psi(x_i) \\
\leq C \sup_{\|\varphi\|_{W^{1/2+\varepsilon,2}=1}} \|\psi\|_{L^\infty} \left( 1 + \frac{1}{(1+\min_{i,x_i\neq 0} x_i(t-s))^{\theta}} \right) \\
\leq C,$$

for any  $\theta \in (0,1)$  and  $\varepsilon > 0$ . In this case, (3.5) is satisfied with a = 0.

Note that the additive constant inside the brackets only appears if there is an  $x_i$ , such that  $x_i = 0$ .

In the unweighted case, Assumption (M 2) might become an issue, but since the support of  $\nu_{\text{exp}}$  is contained in a ball of radius  $R = 2 \max\{x_1, \dots, x_k\}$ , it suffices to study the whole problem on the spaces  $W^{\alpha,p}(B_R(0))$  ( $0 \le \alpha, 1 \le p$ ).

(2) Let  $K_{frac}$  be the fractional kernel of order  $\alpha \in (0,1)$ :

$$K_{frac}(t) := \frac{1}{\Gamma(\alpha)} t^{\alpha - 1}, t > 0.$$

The corresponding Radon measure is

$$\nu_{\rm frac}(x) = \frac{1}{\Gamma(\alpha)\Gamma(1-\alpha)} x^{-\alpha} dx.$$

we will ignore the pre-factor  $\frac{1}{\Gamma(\alpha)\Gamma(1-\alpha)}$ . Let  $\gamma > 0$ .

$$\sup_{\|\varphi\|_{W^{1,2}^{1,2}}=1}\left|\int_{0}^{\infty}e^{-(t-s)x}\varphi(x)\nu(\,\mathrm{d}x)\right|\leq \sup_{\|\varphi\|_{W^{1,2}^{1,2}}=1}\|(1+x)^{\gamma}\varphi(x)\|_{\infty}\left|\int_{0}^{\infty}e^{-(t-s)x}\frac{1}{(1+x)^{\gamma}x^{\alpha}}\,\mathrm{d}x\right|$$

$$\leq \sup_{\|\varphi\|_{W_{w}^{1,2}}=1} \|(1+x)^{\gamma} \varphi(x)\|_{\infty} \left| \int_{0}^{\infty} e^{-(t-s)x(1-\beta)} \frac{1}{(t-s)^{\beta}} \frac{1}{\left(\frac{1}{(t-s)} + x\right)^{\beta} (1+x)^{\gamma} x^{\alpha}} dx \right|$$

$$\leq \frac{1}{(t-s)^{\beta}} \sup_{\|\varphi\|_{W^{1,2}}=1} \|\varphi(x)\|_{W_{w}^{1,2}} C(T).$$

(3.5) is satisfied with  $a = \beta$ . This illustrates that we require  $\alpha < 1$ ,  $\beta + \gamma + \alpha > 0$ . Assume, we wanted to square integrate the result, so  $\beta < \frac{1}{2}$ .

- First we consider the case  $\alpha \in (\frac{1}{2}, 1)$ : Let  $\widetilde{\varepsilon} > \varepsilon > 0$  be such that  $1 \alpha + \widetilde{\varepsilon} < 1$ . We set  $\gamma = -\varepsilon$ ,  $\beta = 1 - \alpha + \widetilde{\varepsilon} - \varepsilon$ . Let  $w = (w_0, w_1)$  with  $w_i = (1 + x)^{i2 - 1 - \varepsilon}$ , then also Assumption (M 2) is satisfied.
- For  $\alpha \in (0, \frac{1}{2})$ , we notice that we require additional decay, so we set  $\gamma = 1 + \varepsilon \beta \alpha$ . This results in a weight  $w = (w_0, w_1)$  with  $w_i = (1 + x)^{i2 1 + \gamma}$ . Hence we can not guarantee that Assumption  $(M \ 2)$  is satisfied.
- (3) A similar argument works for the kernel

$$k_{MVN}(t-s) = \frac{1}{\Gamma(\alpha)} \left( (t-s)^{\alpha-1} - (-s)_{+}^{\alpha-1} \right) .$$

(4) Let  $K_{gamma}$  be the Gamma kernel of the form

$$K_{gamma}(t) := \frac{1}{\Gamma(\alpha)} e^{-\beta t} t^{\alpha - 1}, t > 0,$$

for some  $\beta > 0$  and  $\alpha \in (\frac{1}{2}, 1)$ . The corresponding Radon measure is given by

$$\mu_{\text{gamma}}(dx) = \frac{1}{\Gamma(\alpha)\Gamma(1-\alpha)} (x-\beta)^{-\alpha} \mathbf{1}_{(\beta,\infty)} dx.$$

The measure  $\mu_{\text{gamma}}$  will be supported on  $[\beta, \infty)$ . Let  $\varepsilon > 0$  be such that  $1 - a + \varepsilon < 1$ . The only difference to the fractional kernel is that the weights will be shifted by  $\beta$ .

$$\left| \int_0^\infty e^{-(t-s)x} \varphi(x) \nu(\,\mathrm{d}x) \right| \leq \left\| (1+(x-\beta))^\gamma \varphi(x) \right\|_\infty \left| \int_\beta^\infty e^{-(t-s)x} \frac{1}{(1+(x-\beta))^\gamma (x-\beta)^\alpha} \,\mathrm{d}x \right|$$

$$\leq \left\| \frac{1}{(1+(x-\beta))^\varepsilon} \varphi(x) \right\|_\infty e^{-\beta(t-s)} \left| \int_0^\infty e^{-(t-s)x(1-\eta)} \frac{1}{\left(\frac{1}{(t-s)} + x\right)^\eta (1+x)^\gamma x^\alpha} \,\mathrm{d}x \right|$$

$$\leq \left\| \varphi(x) \right\|_{W_w^{1,2}} C(T) \frac{e^{-\beta(t-s)}}{(t-s)^\eta} \left| \int_0^1 \frac{1}{x^\alpha} \,\mathrm{d}x + \int_1^\infty \frac{1}{(1+x)^{\eta+\gamma+\alpha}} \,\mathrm{d}x \right|.$$

The choice of  $\alpha, \gamma \eta$  are identical to  $K_{frac}$ . (3.5) is again satisfied with  $a = \eta$ .

(5) Given any  $\beta > 0$  and let the kernel K satisfying Assumption (M 1), the exponentially damped kernel

$$K_{damp}(t) := e^{-\beta t} K(t), t > 0,$$

has the corresponding Radon measure

$$\nu_{\text{damp}}(dx) = \mathbf{1}_{[\beta,\infty)}(x)\nu_K(dx - \beta),$$

with support supp  $\nu_{\text{damp}} = \{x \in [\beta, \infty) \mid x - \beta \in \text{supp } \mu\}$ . This case, roughly, can be handled with the same specifications as for  $\nu_K$ , since

$$e^{-\beta t}K(t) = \int_0^\infty \mathbf{1}_{\operatorname{supp}(\nu_K)} e^{-\beta t} e^{-tx} \nu_K(\,\mathrm{d}x).$$

(6) Let  $\delta > 0$  and consider any completely monotone kernel K, satisfying Assumption (M 1), with the corresponding Radon measure  $\nu_K$ . The sifted kernel

$$K_{shift}(t) := K(t + \delta), t > 0,$$

possesses the corresponding Radon measure

$$\nu_{\text{shift}}(dx) = e^{-\delta x} \nu_K(dx),$$

with support supp( $\nu_{\text{shift}}$ ) = supp(supp  $\mu_{shift}$  = supp  $\mu$ ).

A direct calculation shows that the assumptions are satisfied with a bound of the form  $C(T)\frac{1}{(t-s+\delta)^{1-\alpha+\tilde{\varepsilon}}}$ .

Remark 49. On a purely formal level, measures  $\nu_{\text{frac}}$  with  $\alpha \in \left(0, \frac{1}{2}\right)$  correspond to singular kernels which are only  $L^1$  integrable in time. Integrated versions of SVEs with  $L^1$  kernels have been considered in [2]. The non-integrated SVE is not well posed in a strong\ pointwise sense, which is also reflected in the decay requirement of the test function in the lift. However, the formulation of the X process in [2] is considerably "better-behaved". A more in-depth study of this case is currently a work in progress.

We recall for  $\theta_{\nu_h}, \theta_{\nu_{\sigma}}$  from Assumption (M 1) the weights from Definition 3

$$(w_{-})_{i}(x) := (1+x)^{2\eta_{-}-1+2i}$$
  
>  $(w_{\sim})_{i}(x) := (1+x)^{2\eta_{\sim}-1+2i}$   
>  $(w_{+})_{i}(x) := (1+x)^{2\eta_{+}-1+2i}$ ,

with

$$\begin{split} \eta_{+} &= \begin{cases} -\varepsilon & \text{if } \theta_{\nu_{\sigma}} < \frac{1}{2}, \text{ where } 0 < \varepsilon < \frac{1}{2} - \theta_{\nu_{\sigma}}, \\ \theta_{\nu_{\sigma}} - \frac{1}{2} + \delta & \text{if } \theta_{\nu_{\sigma}} > \frac{1}{2}, \text{ where } 0 < \delta < \frac{1}{2}, \end{cases} \\ \eta_{-} &> \max\{\theta_{\nu_{b}}, \theta_{\nu_{\sigma}}\}, \\ \eta_{+} &< \eta_{\sim} < \eta_{-}. \end{split}$$

Remark 50. Let  $\mu \in W_{\frac{1}{w_+}}^{-1,2}$ ,  $\psi \in W_{w_+}^{1,2}$  and  $\nu \in W_{\frac{1}{w_-}}^{-1,2}$ . There is no ambiguity in the interpretation of the stochastic integral, since the driving noise W is one dimensional, and  $\mu \mapsto \nu_{\sigma}\sigma(\langle \mu, \psi \rangle)$  is a mapping from  $W_{\frac{1}{w_+}}^{-1,2}$  into the space of Hilbert-Schmidt operators from  $\mathbb{R}$  to  $W_{\frac{1}{w_-}}^{-1,2}$ , denoted by  $\mathcal{L}_2(\mathbb{R}, W_{\frac{1}{w_-}}^{-1,2})$ .

3.2.1. Step 1: The Lipschitz case.

**Theorem 51.** Let Assumptions (Coeff 1), (Coeff 2) and (M 1) be satisfied and T > 0 be fixed. Let  $W_{\frac{1}{w_+}}^{-1,2}$  correspond to the choice of space for which (3.6) holds. and  $\mu_0 \in L^2(0,T;W_{\frac{1}{w_+}}^{-1,2})$ 

be given. Then equation (1.8), with initial condition  $\mu_0$ , has a unique mild  $W_{\frac{1}{w_+}}^{-1,2}$  solution  $\mu \in L^2(\Omega, C([0,T], W_{\frac{1}{w_+}}^{-1,2})).$ 

Before we begin with the proof, we introduce the following useful Lemma.

**Lemma 52.** Let  $0 \le T < \infty$ ,  $f, g \in L^1(0,T)$  and  $g \ge 0$ , then

$$\sup_{t \le T} \int_0^t g(t - s) f(s) \, ds \le \int_0^T g(T - r) \sup_{0 \le u \le r} |f(u)| \, dr.$$

Proof.

$$\sup_{t \le T} \int_0^t g(t-s)f(s) \, \mathrm{d}s = \sup_{t \le T} \int_0^t g(T-(s-t+T))f(s) \, \mathrm{d}s 
= \sup_{t < T} \int_{T-t}^T g(T-r)f(r+t-T) \, \mathrm{d}r \le \sup_{t < T} \int_0^T g(T-r)\mathbf{1}_{\{T-t \le r \le T\}} |f(r+t-T)| \, \mathrm{d}r.$$

Since  $0 \le r + t - T \le r$ ,

$$\sup_{t \le T} \int_0^T g(T - r) \mathbf{1}_{T - t \le r \le T} |f(r + t - T)| dr \le \sup_{t \le T} \int_0^T g(T - r) \sup_{0 \le u \le r} |f(u)| dr 
\le \int_0^T g(T - r) \sup_{0 \le u \le r} |f(u)| dr.$$

Proof of Theorem 51. We define the operator  $\Gamma: L^2(\Omega, C([0,T], W_{\frac{1}{w_+}}^{-1,2})) \to L^2(\Omega, C([0,T], W_{\frac{1}{w_+}}^{-1,2}))$  by

$$\Gamma(\mu, \mu_0) := S_t^* \mu_0 + \int_0^t S_{t-s}^* \nu_b b(s, \langle \mu_s, \psi \rangle) \, \mathrm{d}s + \int_0^t S_{t-s}^* \nu_\sigma \sigma(s, \langle \mu_s, \psi \rangle) \, \mathrm{d}W_s.$$

The proof is a rather standard application of the Banach fixed-point theorem, verifying the self-mapping and contraction properties of  $\Gamma$ . We only verify the contraction property of the solution  $\Gamma$ , since the self-mapping property is verified analogously, using the estimate from Lemma 38 and a similar estimate will be shown in Lemma 57. In the following steps, we use that  $|\langle \mu_s, \psi \rangle| \leq \sup_{h \colon \|h\|_{W_{w_+}^{1,2}} = 1} |\langle \mu_s, h \rangle| = \|\mu_s\|_{W_{w_+}^{-1,2}}$ . We first consider  $\widetilde{T}$  to be fixed and "small".

$$\mathbb{E}\sup_{t\leq\widetilde{T}}\|\Gamma(\mu_{t}^{1},\mu_{0})-\Gamma(\mu_{t}^{2},\mu_{0})\|_{W_{-\frac{1}{w_{+}}}^{-1,2}}^{2}\leq C_{\operatorname{Lip}_{b}}\mathbb{E}\sup_{t\leq\widetilde{T}}\left(\int_{0}^{t}\|S_{t-s}^{*}\nu_{b}\|_{W_{-\frac{1}{w_{+}}}^{-1,2}}\|\mu_{s}^{1}-\mu_{s}^{2}\|_{W_{-\frac{1}{w_{+}}}^{-1,2}}\,\mathrm{d}s\right)^{2}$$

$$+C_{\operatorname{Lip}_{\sigma}}\mathbb{E}\sup_{t\leq\widetilde{T}}\left(\int_{0}^{t}\|S_{t-s}^{*}\nu_{\sigma}\|_{W_{-\frac{1}{w_{+}}}^{-1,2}}\|\mu_{s}^{1}-\mu_{s}^{2}\|_{W_{-\frac{1}{w_{+}}}^{-1,2}}^{2}\,\mathrm{d}s\right)$$

$$\leq C_{\operatorname{Lip}_{b}}\mathbb{E}\sup_{t\leq\widetilde{T}}\left(\int_{0}^{t}\|S_{t-s}^{*}\nu_{b}\|_{W_{-\frac{1}{w_{+}}}^{-1,2}}\|\mu_{s}^{1}-\mu_{s}^{2}\|_{W_{-\frac{1}{w_{+}}}^{-1,2}}\,\mathrm{d}s\right)^{2}$$

П

$$\begin{split} &+ C_{\operatorname{Lip}_{\sigma}} \mathbb{E} \left( \int_{0}^{\widetilde{T}} \left\| S_{\widetilde{T}-s}^{*} \nu_{\sigma} \right\|_{W_{\frac{1}{w_{+}}}^{-1,2}}^{2} \sup_{r \leq s} \left\| \mu_{r}^{1} - \mu_{r}^{2} \right\|_{W_{\frac{1}{w_{+}}}^{-1,2}}^{2} \, \mathrm{d}s \right) \\ &\leq C_{\operatorname{Lip}_{b}} \sup_{t \leq \widetilde{T}} \left( \int_{0}^{t} \left\| S_{t-s}^{*} \nu_{b} \right\|_{W_{\frac{1}{w_{+}}}^{-1,2}} \, \mathrm{d}s \right) \mathbb{E} \int_{0}^{\widetilde{T}} \left\| S_{\widetilde{T}-s}^{*} \nu_{b} \right\|_{W_{\frac{1}{w_{+}}}^{-1,2}} \sup_{r \leq s} \left\| \mu_{r}^{1} - \mu_{r}^{2} \right\|_{W_{\frac{1}{w_{+}}}^{-1,2}}^{2} \, \mathrm{d}s \\ &+ C_{\operatorname{Lip}_{\sigma}} \mathbb{E} \left( \int_{0}^{\widetilde{T}} \left\| S_{\widetilde{T}-s}^{*} \nu_{\sigma} \right\|_{W_{\frac{1}{w_{+}}}^{-1,2}}^{2} \sup_{r \leq s} \left\| \mu_{r}^{1} - \mu_{r}^{2} \right\|_{W_{\frac{1}{w_{+}}}^{-1,2}}^{2} \, \mathrm{d}s \right) \\ &\leq C_{\operatorname{Lip},\widetilde{T}} \mathbb{E} \sup_{s \leq \widetilde{T}} \left\| \mu_{s}^{1} - \mu_{s}^{2} \right\|_{W_{\frac{1}{w_{+}}}^{-1,2}}^{2}, \end{split}$$

Where  $C_{\widetilde{T}}$  depends on  $\widetilde{T}$  via the terms  $\int_0^{\widetilde{T}} \|S_{\widetilde{T}-s}^* \nu_b\|_{W_{\frac{1}{w_+}}^{-1,2}} ds$  and  $\int_0^{\widetilde{T}} \|S_{\widetilde{T}-s}^* \nu_\sigma\|_{W_{\frac{1}{w_+}}^{-1,2}}^2$ . By As-

sumption (M 1), both of these integrals can be controlled by a constant times  $\max\{\widetilde{T}^{2a_b}, \widetilde{T}^{2a_\sigma}\}$ . Hence, we obtain a contraction for  $\widetilde{T}$  small enough. Obtaining a solution on [0,T] now follows by standard arguments.

Corollary 53. The solution map  $\mu_0 \mapsto \mu_t$  is continuous for every  $0 < t \le T$ .

3.2.2. Step 2: General coefficients. Our strategy to show existence for the equation, when  $b, \sigma$  are merely uniformly continuous ( (Coeff 3)) and satisfy (Coeff 1) is to approximate general  $b, \sigma$  with Lipschitz continuous function and then pass to the limit. For precisely this limiting procedure, we will derive a-priori estimates on the mild solution of equation (1.8), which will be used in combination with certain tightness arguments.

A central tool to our approach is the next proposition, which allows us to approximate  $b, \sigma$  by a sequence of Lipschitz continuous functions  $b_n, \sigma_n$  with a uniform linear growth bound. By Theorem 51, for each  $n \in \mathbb{N}$ , the equation

(3.7) 
$$d\mu_n(t) = -x\mu_n(t) dt + \nu_b b_n(t, \langle \mu_n(t), \psi \rangle) dt + \nu_\sigma \sigma_n(t, \langle \mu_n(t), \psi \rangle) dW_t$$

has a solution, for each  $\psi \in W_{\frac{1}{w_+}}^{-1,2}$  in the sense of Definition 42 with values in  $L^2(\Omega, C([0,T], W_{\frac{1}{w_+}}^{-1,2}))$ .

**Proposition 54.** ([47, Proposition 1.1]) Suppose  $F: \mathbb{R}_+ \times \mathbb{R}^n \to \mathbb{R}^d$  is a Borel function of at most linear growth, i.e.

$$\exists L < \infty, \ \forall t > 0, \ \forall x \in \mathbb{R}^n, \ \|F(t,x)\| < L(1+\|x\|).$$

such that  $F(t,\cdot) \in C(\mathbb{R}^n,\mathbb{R}^d)$  for any  $t \in \mathbb{R}_+$ . Then there exists a sequence of Borel functions  $F_k \colon \mathbb{R}_+ \times \mathbb{R}^n \to \mathbb{R}^d$ ,  $k \ge 1$ , which have at most linear growth uniformly in k, namely

$$\forall k \in \mathbb{N}, \exists L < \infty, \forall t \ge 0, \forall x \in \mathbb{R}^n, ||F_k(t, x)|| \le L(1 + ||x||),$$

which are Lipschitz continuous in the second variable uniformly in the first one,

$$\forall k \in \mathbb{N}, \exists L_k < \infty, \forall t \geq 0, \forall x, y \in \mathbb{R}^n, ||F_k(t, x) - F_k(t, y)|| \leq L_k ||x - y||,$$

and which satisfy

$$\lim_{k \to \infty} F_k(t, \cdot) = F(t, \cdot) \quad \text{locally uniformly on } \mathbb{R}^n$$

for all  $t \geq 0$ .

**Remark 55.** Unlike the Lipschitz constant, the approximations  $F_k$  are share the same modulus of continuity as F.

3.2.3. A-priori estimates. We will derive a "spatial" and a "temporal" estimate to perform the limit  $n \to \infty$  in (3.7). The "spatial" estimate will be separated into two Lemmata since we will reuse Lemma 56 in Section 3.3.

**Lemma 56.** Let Assumptions (Coeff 1) be satisfied, 1 < p and let  $\mu$  be a mild solution of (1.8), with  $\mu_0 \in L^p(\Omega, W_{\frac{1}{m}}^{-1,2})$ .

Then there exists a constant  $C_{p,LG} > 0$ , which only depends on p and the linear growth condition (Coeff 1) and is independent of T, such

$$\mathbb{E} \sup_{t \leq T} \|\mu_{t}\|_{W_{\frac{1}{w_{+}}}^{-1,2}}^{p} \leq C_{p,\text{LG}} \mathbb{E} \sup_{t \leq T} \|S_{t}^{*}\mu_{0}\|_{W_{\frac{1}{w_{+}}}^{-1,2}}^{p} + C_{p,\text{LG}}h_{b,j}(T) \int_{0}^{T} \|S_{T-s}^{*}\nu_{b}\|_{W_{\frac{1}{w_{+}}}^{-1,2}} \mathbb{E} \sup_{r \leq s} \|\mu_{r}\|_{W_{\frac{1}{w_{+}}}^{-1,2}}^{p} \, ds$$

$$+ C_{p,\text{LG}}h_{\sigma,j}(T) \int_{0}^{T} \|S_{T-s}^{*}\nu_{\sigma}\|_{W_{\frac{1}{w_{+}}}^{-1,2}}^{2} \left( \mathbb{E} \sup_{r \leq s} \|\mu_{r}\|_{W_{\frac{1}{w_{+}}}^{-1,2}}^{2} \right)^{\frac{p}{2}} \, ds$$

$$+ C_{p,\text{LG}} \left( \int_{0}^{T} \|S_{T-s}^{*}\nu_{b}\|_{W_{\frac{1}{w_{+}}}^{-1,2}} \, ds \right)^{p} + C_{p,\text{LG}} \left( \int_{0}^{T} \|S_{T-s}^{*}\nu_{\sigma}\|_{W_{\frac{1}{w_{+}}}^{-1,2}}^{2} \, ds \right)^{\frac{p}{2}}.$$

where, j = 1, if  $2 \le p$  and j = 1, if  $1 < j \le 2$ . Depending on the value of p, we set

$$h_{b,1}(T) = h_{b,2}(T) = \left( \int_0^T \|S_{T-s}^* \nu_b\|_{W_{\frac{1}{w_+}}^{-1,2}} \, \mathrm{d}s \right)^{p-1}, \qquad h_{\sigma,1}(T) = \left( \int_0^T \|S_{T-s}^* \nu_\sigma\|_{W_{\frac{1}{w_+}}^{-1,2}}^2 \, \mathrm{d}s \right)^{\frac{p}{2}-1}$$

$$h_{\sigma,2}(T) = \left( \int_0^T \|S_{T-s}^* \nu_\sigma\|_{W_{\frac{1}{w_+}}^{-1,2}}^2 \right)^{p-1}.$$

*Proof.* Fix T > 0. We use the maximal inequality (see [46]), and Jensen's inequality, so that for  $p \ge 2$  we obtain

$$\begin{split} & \mathbb{E} \sup_{t \in [0,T]} \|\mu_t\|_{W^{-1,2}_{\frac{1}{w_+}}}^p \leq C_p \mathbb{E} \sup_{t \in [0,T]} \|S_t^* \mu_0\|_{W^{-1,2}_{\frac{1}{w_+}}}^p \\ & + C_{p,\text{LG}} \left( \int_0^T \|S_{T-s}^* \nu_b\|_{W^{-1,2}_{\frac{1}{w_+}}} \, \mathrm{d}s \right)^{p-1} \int_0^T \|S_{T-s}^* \nu_b\|_{W^{-1,2}_{\frac{1}{w_+}}} \mathbb{E} \sup_{t \leq s} \|\mu_t\|_{W^{-1,2}_{\frac{1}{w_+}}}^p \, \mathrm{d}s \\ & + C_{p,\text{LG}} \left( \int_0^T \|S_{T-s}^* \nu_\sigma\|_{W^{-1,2}_{\frac{1}{w_+}}}^2 \, \mathrm{d}s \right)^{\frac{p}{2}-1} \int_0^T \|S_{T-s}^* \nu_\sigma\|_{W^{-1,2}_{\frac{1}{w_+}}}^2 \mathbb{E} \sup_{t \leq s} \|\mu_t\|_{W^{-1,2}_{\frac{1}{w_+}}}^p \, \mathrm{d}s \\ & + C_{p,\text{LG}} \left( \int_0^T \|S_{T-s}^* \nu_b\|_{W^{-1,2}_{\frac{1}{w_+}}} \, \mathrm{d}s \right)^{\frac{p}{2}-1} \int_0^T \|S_{T-s}^* \nu_\sigma\|_{W^{-1,2}_{\frac{1}{w_+}}}^2 \mathbb{E} \sup_{t \leq s} \|\mu_t\|_{W^{-1,2}_{\frac{1}{w_+}}}^p \, \mathrm{d}s \end{split}$$

If  $1 \le p < 2$  and  $2\gamma_2 > p$ , then

$$\mathbb{E}\sup_{t\in[0,T]}\|\mu_t\|_{W^{-1,2}_{\frac{1}{w_+}}}^p \leq C_p\mathbb{E}\sup_{t\in[0,T]}\|S_t^*\mu_0\|_{W^{-1,2}_{\frac{1}{w_+}}}^p$$

$$\begin{split} &+C_{p,\mathrm{LG}}\left(\int_{0}^{T}\|S_{T-s}^{*}\nu_{b}\|_{W_{-\frac{1}{2}}^{-1,2}}\mathrm{d}s\right)^{p-1}\int_{0}^{T}\|S_{T-s}^{*}\nu_{b}\|_{W_{-\frac{1}{2}}^{-1,2}}\mathrm{E}\sup_{r\leq s}\|\mu_{r}\|_{W_{-\frac{1}{2}}^{-1,2}}^{p}\mathrm{d}s\\ &+C_{p,\mathrm{LG}}\mathbb{E}\left(\left(\int_{0}^{T}\|S_{T-s}^{*}\nu_{\sigma}\|_{W_{-\frac{1}{2}}^{-1,2}}^{2}\right)^{p-1}\int_{0}^{T}\|S_{T-s}^{*}\nu_{\sigma}\|_{W_{-\frac{1}{2}}^{-1,2}}^{2}\left(\sup_{r\leq s}\|\mu_{r}\|_{W_{-\frac{1}{2}}^{-1,2}}^{p}\sup\|\mu_{r}\|_{W_{-\frac{1}{2}}^{-1,2}}^{p}\right)\mathrm{d}s\right)^{1/2}\\ &+C_{p,\mathrm{LG}}\left(\int_{0}^{T}\|S_{T-s}^{*}\nu_{b}\|_{W_{-\frac{1}{2}}^{-1,2}}\mathrm{d}s\right)^{p}+C_{p,\mathrm{LG}}\left(\int_{0}^{T}\|S_{T-s}^{*}\nu_{\sigma}\|_{W_{-\frac{1}{2}}^{-1,2}}^{2}\mathrm{d}s\right)^{\frac{p}{2}}\\ &\leq C_{p}\mathbb{E}\sup_{t\in[0,T]}\|S_{t}^{*}\mu_{0}\|_{W_{-\frac{1}{2}}^{-1,2}}^{p}\mathrm{d}s\right)^{p-1}\int_{0}^{T}\|S_{T-s}^{*}\nu_{b}\|_{W_{-\frac{1}{2}}^{-1,2}}\mathbb{E}\sup_{r\leq s}\|\mu_{r}\|_{W_{-\frac{1}{2}}^{-1,2}}^{p}\mathrm{d}s\\ &+C_{p,\mathrm{LG}}\mathbb{E}\left(\sup_{r\leq T}\|\mu_{r}\|_{W_{-\frac{1}{2}}^{-1,2}}^{p}\mathrm{d}s\right)^{p-1}\int_{0}^{T}\|S_{T-s}^{*}\nu_{\sigma}\|_{W_{-\frac{1}{2}}^{-1,2}}^{2}\mathbb{E}\sup_{w_{+}}\|\mu_{r}\|_{W_{-\frac{1}{2}}^{-1,2}}^{p}\mathrm{d}s\\ &+C_{p,\mathrm{LG}}\mathbb{E}\left(\int_{0}^{T}\|S_{T-s}^{*}\nu_{b}\|_{W_{-\frac{1}{2}}^{-1,2}}\mathrm{d}s\right)^{p}+C_{p,\mathrm{LG}}\left(\int_{0}^{T}\|S_{T-s}^{*}\nu_{\sigma}\|_{W_{-\frac{1}{2}}^{-1,2}}^{2}\mathrm{d}s\right)^{\frac{p}{2}}\\ &\leq C_{p}\mathbb{E}\sup_{t\in[0,T]}\|S_{t}^{*}\mu_{0}\|_{W_{-\frac{1}{2}}^{-1,2}}^{p}\mathrm{d}s\right)^{p}+C_{p,\mathrm{LG}}\mathbb{E}\left(\int_{0}^{T}\|S_{T-s}^{*}\nu_{\sigma}\|_{W_{-\frac{1}{2}}^{-1,2}}^{p}\mathrm{d}s\right)^{p}\\ &+C_{p,\mathrm{LG}}\mathbb{E}\left(\int_{0}^{T}\|S_{T-s}^{*}\nu_{\sigma}\|_{W_{-\frac{1}{2}}^{-1,2}}^{p}\mathrm{d}s\right)^{p-1}\int_{0}^{T}\|S_{T-s}^{*}\nu_{\sigma}\|_{W_{-\frac{1}{2}}^{-1,2}}^{p}\mathrm{d}s\right)^{p}\\ &+C_{p,\mathrm{LG}}\mathbb{E}\left(\int_{0}^{T}\|S_{T-s}^{*}\nu_{\sigma}\|_{W_{-\frac{1}{2}}^{-1,2}}^{p}\mathrm{d}s\right)^{p-1}\int_{0}^{T}\|S_{T-s}^{*}\nu_{\sigma}\|_{W_{-\frac{1}{2}}^{-1,2}}^{p}\mathrm{d}s\right)^{p}\\ &+C_{p,\mathrm{LG}}\mathbb{E}\left(\int_{0}^{T}\|S_{T-s}^{*}\nu_{\sigma}\|_{W_{-\frac{1}{2}}^{-1,2}}^{p}\mathrm{d}s\right)^{p-1}\mathbb{E}\int_{0}^{T}\|S_{T-s}^{*}\nu_{\sigma}\|_{W_{-\frac{1}{2}}^{-1,2}}^{p}\mathrm{d}s\right)^{p}\\ &+C_{p,\mathrm{LG}}\left(\int_{0}^{T}\|S_{T-s}^{*}\nu_{\sigma}\|_{W_{-\frac{1}{2}}^{-1,2}}^{p}\mathrm{d}s\right)^{p}+C_{p,\mathrm{LG}}\left(\int_{0}^{T}\|S_{T-s}^{*}\nu_{\sigma}\|_{W_{-\frac{1}{2}}^{-1,2}}^{p}\mathrm{d}s\right)^{p}\\ &+C_{p,\mathrm{LG}}\left(\int_{0}^{T}\|S_{T-s}^{*}\nu_{\sigma}\|_{W_{-\frac{1}{2}}^{-1,2}}^{p}\mathrm{d}s\right)^{p}+C_{p,\mathrm{LG}}\left(\int_{0}^{T}\|S_{T-s}^{*}\nu_{\sigma}\|_{W_{-\frac{1}{2}}^{-1,2}}^{p}\mathrm{d}s\right)^{p}\\ &+C_{p,\mathrm{LG}}\left(\int$$

Choosing  $\varepsilon$  such that  $C_{p,\text{LG}}\varepsilon < 1$ , bringing  $C_{p,\text{LG}}\varepsilon\mathbb{E}\left(\sup_{r \leq T} \|\mu_r\|_{W^{-1,2}_{\frac{1}{w_+}}}^p\right)$  to the left hand side and dividing by  $(1 - C_{p,\text{LG}}\varepsilon)$  yields

$$\begin{split} \mathbb{E} \sup_{t \in [0,T]} \|\mu_t\|_{W^{-1,2}_{\frac{1}{w_+}}}^p &\leq C_{p,\mathrm{LG},\varepsilon} \mathbb{E} \sup_{t \in [0,T]} \|S_t^* \mu_0\|_{W^{-1,2}_{\frac{1}{w_+}}}^p \\ &+ C_{p,\mathrm{LG},\varepsilon} \left( \int_0^T \|S_{T-s}^* \nu_b\|_{W^{-1,2}_{\frac{1}{w_+}}} \, \mathrm{d}s \right)^{p-1} \int_0^T \|S_{T-s}^* \nu_b\|_{W^{-1,2}_{\frac{1}{w_+}}} \mathbb{E} \sup_{r \leq s} \|\mu_r\|_{W^{-1,2}_{\frac{1}{w_+}}}^p \, \mathrm{d}s \end{split}$$

$$+ C_{p,\text{LG},\varepsilon} \left( \int_{0}^{T} \|S_{T-s}^{*}\nu_{\sigma}\|_{W_{-\frac{1}{w_{+}}}^{-1,2}}^{2} \right)^{p-1} \int_{0}^{T} \|S_{T-s}^{*}\nu_{\sigma}\|_{W_{-\frac{1}{w_{+}}}^{-1,2}}^{2} \mathbb{E} \left( \sup_{r \leq s} \|\mu_{r}\|_{W_{\frac{1}{w_{+}}}^{-1,2}}^{p} \, \mathrm{d}s \right)$$

$$+ C_{p,\text{LG},\varepsilon} \left( \int_{0}^{T} \|S_{T-s}^{*}\nu_{b}\|_{W_{-\frac{1}{w_{+}}}^{-1,2}} \, \mathrm{d}s \right)^{p} + C_{p,\text{LG},\varepsilon} \left( \int_{0}^{T} \|S_{T-s}^{*}\nu_{\sigma}\|_{W_{-\frac{1}{w_{+}}}^{-1,2}}^{2} \, \mathrm{d}s \right)^{\frac{p}{2}}.$$

**Lemma 57.** Let the Assumptions (M 1), (Coeff 1) be satisfied, 1 < p and let  $\mu$  be a mild solution of (1.8), with  $\mu_0 \in L^p(\Omega, W_{\frac{1}{w_+}}^{-1,2})$ , where  $W_{\frac{1}{w_+}}^{-1,2}$  corresponds to the choice of space for which (3.6) holds. Then

(3.9) 
$$\mathbb{E} \sup_{t \le T} \|\mu_t\|_{W_{\frac{1}{n-1}}^{-1,2}}^p \le C_{p,b,\sigma,T,\mu_0}.$$

*Proof.* By Lemma 56 and Assumption (Coeff 1),

$$\mathbb{E} \sup_{t \leq T} \|\mu_t\|_{W_{\frac{1}{w_+}}^{-1,2}}^p \leq C_{p,\text{LG}} \mathbb{E} \sup_{t \leq T} \|S_t^* \mu_0\|_{W_{\frac{1}{w_+}}^{-1,2}}^p + C_{p,\text{LG}} h_{b,j}(T) \int_0^T \frac{C(T)}{(T-s)^{1-a_1}} \mathbb{E} \sup_{r \leq s} \|\mu_r\|_{W_{\frac{1}{w_+}}^{-1,2}}^2 ds$$

$$+ C_{p,\text{LG}} h_{\sigma,j}(T) \int_0^T \frac{C(T)}{(T-s)^{2-2a_2}} \mathbb{E} \sup_{r \leq s} \|\mu_r\|_{W_{\frac{1}{w_+}}^{-1,2}}^2 ds$$

$$+ C_{p,\text{LG}} T^{(p-1)a_1} + C_{p,\text{LG}} T^{(\frac{p}{2}-1)(2a_2-1)}.$$

$$h_{b,1}(T) = h_{b,2}(T) = C(T)T^{(p-1)a_1}, \qquad h_{\sigma,1}(T) = C(T)T^{(\frac{p}{2}-1)(2a_2-1)}, \qquad h_{\sigma,2}(T) = C(T)T^{(p-1)(2a_2-1)}.$$

Let u, A, G, F be non negative functions on  $\mathbb{R}_+$ . Since

$$u(t) \le A(t) + G(t) + F(t) \le A(t) + 2\max\{G(t), F(t)\},\$$

it suffices to estimate  $u(t) \leq A(t) + 2G(t)$  and  $u(t) \leq A(t) + 2F(t)$  separately. In our case,

$$\mathbb{E} \sup_{t \leq T} \|\mu_t\|_{W^{-1,2}_{\frac{1}{w_+}}}^p \leq C_{p,\text{LG}} \mathbb{E} \sup_{t \leq T} \|S_t^* \mu_0\|_{W^{-1,2}_{\frac{1}{w_+}}}^p + 2C_{p,b,\sigma,T} h_{b,j}(T) \int_0^T \frac{1}{(T-s)^{1-a_1}} \mathbb{E} \sup_{r \leq s} \|\mu_r\|_{W^{-1,2}_{\frac{1}{w_+}}}^2 ds + 2C_{p,\text{LG}} T^{(p-1)a_1},$$

$$\mathbb{E} \sup_{t \leq T} \|\mu_t\|_{W_{\frac{1}{w_+}}^{-1,2}}^p \leq C_{p,\text{LG}} \mathbb{E} \sup_{t \leq T} \|S_t^* \mu_0\|_{W_{\frac{1}{w_+}}^{-1,2}}^p + 2C_{p,b,\sigma,T} h_{\sigma,j}(T) \int_0^T \frac{1}{(T-s)^{2-2a_2}} \mathbb{E} \sup_{r \leq s} \|\mu_r\|_{W_{\frac{1}{w_+}}^{-1,2}}^2 ds + 2C_{p,\text{LG}} T^{(\frac{p}{2}-1)(2a_2-1)}.$$

We can include the terms  $b_1$  and  $b_{2,j}$  into the constant since it already depends on T. [68, Lemma 2.2] now yields

$$\mathbb{E} \sup_{t \le T} \|\mu_t\|_{W_{\frac{1}{w_+}}^{-1,2}}^p \le C_{p,\text{LG},T}.$$

Now we consider the time-regularity of  $t \mapsto \mu_t$ .

**Lemma 58.** Let Assumptions (Coeff 1) and (M 1) be satisfied,  $\gamma \in [0,1]$  be arbitrary,  $1 \leq p < \infty$ . Recall the weights from Definition 3 and let  $0 \leq \gamma_b, 2\gamma_\sigma \leq 1$  be such that  $\theta_{\nu_b} - \gamma_b \leq \eta_-$ ,  $\theta_{\nu_\sigma} - \gamma_\sigma \leq \eta_-$ . Assume  $\mu \in L^p(\Omega, L^\infty(0, T; W_{\frac{1}{w_+}}^{-1,2}))$  to be a mild solution to (1.8) with initial condition  $\mu_0 \in L^p(\Omega, W_{\frac{1}{w_+}}^{-1,2})$ , where  $W_{\frac{1}{w_+}}^{-1,2}$  corresponds to the choice of space for which (3.6) holds. Then for  $s \leq t$  with  $|t-s| \leq 1$ , the following estimate holds.

$$\mathbb{E}\|\mu_t - \mu_s\|_{W_{\frac{1}{w}}^{-2,2}}^p \leq \mathbb{E}\|\mu_t - \mu_s\|_{W_{\frac{1}{w}}^{-1,2}}^p \leq C_{p,\text{LG}}\left(|t-s|^{(1-\gamma_b)p} + |t-s|^{(1-2\gamma_\sigma)\frac{p}{2}} + |t-s|^{p(\eta_- - \eta_+ \wedge 1)}\right).$$

Proof of Lemma 58.

$$\mathbb{E}\|\mu_t - \mu_s\|_{W_{\frac{1}{w_-}}^{-1,2}}^p \le C\left(\mathbb{E}\|\mu_t - S_{t-s}^*\mu_s\|_{W_{\frac{1}{w_-}}^{-1,2}}^p + \mathbb{E}\|S_{t-s}^*\mu_s - \mu_s\|_{W_{\frac{1}{w_-}}^{-1,2}}^p\right).$$

By Lemma 34,  $\nu_b \in W_{\frac{1}{w_b}}^{-1,2}$ ,  $\nu_{\sigma} \in W_{\frac{1}{w_{\sigma}}}^{-1,2}$  with  $(w_b)_i = (1+x)^{2\theta_{\nu_b}-1+2i}$  and  $(w_{\sigma})_i = (1+x)^{2\theta_{\nu_{\sigma}}-1+2i}$ , for  $i \geq 0$  and  $x \in \mathbb{R}_+$ . Lemma 38 then implies

$$||S_t^* \nu_b||_{W_{\frac{1}{w_-}}^{-1,2}} \le C_T ||\nu_b||_{W_{\frac{1}{w_b}}^{-1,2}} |t|^{-\gamma_b} ||S_t^* \nu_\sigma||_{W_{\frac{1}{w_-}}^{-1,2}} \le C_T ||\nu_\sigma||_{W_{\frac{1}{w_\sigma}}^{-1,2}} |t|^{-\gamma_\sigma},$$

for any  $0 < t \le T$ . For simplicity, we will set

$$\theta_{\nu_b} - \gamma_b = \theta_{\nu_\sigma} - \gamma_\sigma = \eta_-.$$

$$\begin{split} \mathbb{E} \| \mu_{t} - S_{t-s}^{*} \mu_{s} \|_{W_{-\frac{1}{w}}^{-1,2}}^{p} &\leq C_{p} \mathbb{E} \| \mu_{s} - S_{s-s}^{*} \mu_{s} \|_{W_{-\frac{1}{w}}^{-1,2}}^{p} \\ &+ C_{p} \mathbb{E} \left( \int_{s}^{t} \| S_{r-s}^{*} \nu_{b} b(r, \langle \mu_{r}, \psi \rangle)) \|_{W_{-\frac{1}{w}}^{-1,2}} \, \mathrm{d}r \right)^{p} + C_{p} \mathbb{E} \left\| \int_{s}^{t} S_{r-s}^{*} \nu_{\sigma} \sigma(r, \langle \mu_{r}, \psi \rangle)) \, \mathrm{d}W_{r} \right\|_{W_{-\frac{1}{w}}^{-1,2}}^{p} \\ &\leq C_{p} \mathbb{E} \left( \int_{s}^{t} \| S_{r-s}^{*} \nu_{b} \|_{W_{-\frac{1}{w}}^{-1,2}} |b(r, \langle \mu_{r}, \psi \rangle)| \, \mathrm{d}r \right)^{p} + C_{p} \mathbb{E} \left( \int_{s}^{t} \| S_{r-s}^{*} \nu_{\sigma} \|_{W_{-\frac{1}{w}}^{-1,2}}^{2-1} |\sigma(r, \langle \mu_{r}, \psi \rangle)|^{2} \, \mathrm{d}r \right)^{p/2} \\ &\leq \mathbb{E} \left( \int_{s}^{t} \| S_{r-s}^{*} \nu_{b} \|_{W_{-\frac{1}{w}}^{-1,2}} |\sigma(r, \langle \mu_{r}, \psi \rangle)|^{2} \, \mathrm{d}r \right)^{p/2} \\ &+ \mathbb{E} \left( \int_{s}^{t} \| S_{r-s}^{*} \nu_{\sigma} \|_{W_{-\frac{1}{w}}^{-1,2}}^{2-1} |\sigma(r, \langle \mu_{r}, \psi \rangle)|^{2} \, \mathrm{d}r \right)^{p/2} \\ &\leq \mathbb{E} \left( \int_{s}^{t} \frac{1}{(r-s)^{\gamma_{b}}} \, \mathrm{d}r \right)^{p-1} \int_{s}^{t} \frac{1}{(r-s)^{\gamma_{b}}} |b(r, \langle \mu_{r}, \psi \rangle)|^{p} \, \mathrm{d}r \\ &+ \mathbb{E} \left( \int_{s}^{t} \frac{1}{(r-s)^{2\gamma_{\sigma}}} \, \mathrm{d}r \right)^{\frac{p}{2}-1} \left( \int_{s}^{t} \frac{1}{(r-s)^{2\gamma_{\sigma}}} |\sigma(r, \langle \mu_{r}, \psi \rangle)|^{2} \, \mathrm{d}r \right)^{p/2} \end{split}$$

$$\leq C_{p,\text{LG}} \mathbb{E}|t-s|^{(1-\gamma_b)(p-1)} \int_{s}^{t} \frac{1+\|\mu_r\|_{W^{-1,2}_{\frac{1}{w_+}}}^{p}}{(r-s)^{\gamma_b}} dr + \mathbb{E}|t-s|^{(2\gamma_{\sigma}\left(\frac{p}{2}-1\right))} \int_{s}^{t} \frac{1+\|\mu_r\|_{W^{-1,2}_{\frac{1}{w_+}}}^{p}}{(r-s)^{2\gamma_{\sigma}}} dr$$

$$\leq C_{p,\text{LG}} \left(|t-s|^{(1-\gamma_b)(p-1)} \int_{s}^{t} \frac{1}{(r-s)^{\gamma_b}} dr + |t-s|^{2\gamma_{\sigma}\left(\frac{p}{2}-1\right)} \int_{s}^{t} \frac{1}{(r-s)^{2\gamma_{\sigma}}} dr \right) \mathbb{E} \sup_{t\leq T} \left(1+\|\mu_r\|_{W^{-1,2}_{\frac{1}{w_+}}}^{p}\right)^{p}$$

$$\leq C_{p,\text{LG}} \left(|t-s|^{(1-\gamma_b)(p-1)+(1-\gamma_b)} + |t-s|^{(1-2\gamma_{\sigma})\left(\frac{p}{2}-1\right)+1-2\gamma_{\sigma}}\right) \mathbb{E} \sup_{t\leq T} \left(1+\|\mu_r\|_{W^{-1,2}_{\frac{1}{w_+}}}^{p}\right)$$

$$\leq C_{p,\text{LG}} \left(|t-s|^{(1-\gamma_b)p} + |t-s|^{(1-2\gamma_{\sigma})\frac{p}{2}}\right).$$

For the second estimate, recall that  $(w_+)_i(x) = (1+x)^{2\eta_+-1+2i}$  and use Lemma 39 to conclude that (since  $\eta_- > \eta_+$ )

$$\mathbb{E}\|S_{t-s}^*\mu_s - \mu_s\|_{W_{\frac{1}{w_-}}^{-1,2}} \le C\|\mu_s\|_{W_{\frac{1}{w_+}}^{-1,2}}^p |t-s|^{(\eta_- - \eta_+) \wedge 1}.$$

In summary, we obtain

$$\mathbb{E}\|\mu_t - \mu_s\|_{W_{\frac{1}{w}}^{-1,2}}^p \le C_{p,\mathrm{LG}}\left(|t-s|^{(1-\gamma_b)p} + |t-s|^{(1-2\gamma_\sigma)\frac{p}{2}} + |t-s|^{p((\eta_--\eta_+)\wedge 1)}\right).$$

**Corollary 59.** Let  $(\tau_n)_{n\in\mathbb{N}}$  be a sequence of stopping times such that  $0 \leq \tau_n \leq T$  and let  $\theta > 0$ , then

$$\mathbb{E}\|\mu_{\tau_n+\theta} - \mu_{\tau_n}\|_{W^{\frac{-2,2}{1}}_{\frac{1}{w_-}}}^p \le C_{p,\text{LG}}\left(|\theta|^{(1-\gamma_b)p} + |\theta|^{(1-2\gamma_\sigma)\frac{p}{2}} + |\theta|^{p(\eta_--\eta_+\wedge 1)}\right).$$

Equipped with these two estimates, we process to identify a convergent subsequence and potential limit.

**Definition 60.** For a separable Banach space V, we define  $C([0,T],V^{\text{weak}^*})$ , as the space of weakly continuous functions  $u:[0,T] \to V$ , equipped with the topology  $\mathcal{T}$ , such that for all h in the predual of V, denoted by V, the mapping

$$C([0,T],V^{\text{weak}^*}) \ni v \mapsto \langle v(\cdot),h \rangle_{V \times V} \in C([0,T],\mathbb{R})$$

is continuous. In particular  $v_n \to v$  in  $C([0,T];V^{\text{weak}^*})$  iff for all  $h \in V$ :

$$\lim_{n \to \infty} \sup_{t \in [0,T]} \left| \langle v_n(t) - v(t), h \rangle_{V \times V} \right| = 0.$$

We write  $C([0,T];V^{\text{weak}})$  for the same space but equipped with the weak topology.

Consider the ball

$$B_r := \{ y \in V; \quad ||y||_V \le r \}.$$

If V is separable, the weak\* topology induced on  $B_r$  is metrizable. Let q denote the metric compatible with the weak\* topology on  $B_r$ . Let us consider

$$C\left([0,T]; B_r^{\mathrm{weak}^*}\right) = \left\{u \in C\left([0,T]; V^{\mathrm{weak}^*}\right) : \sup_{t \in [0,T]} \|u(t)\|_V \leq r\right\},$$

which denotes the space of weakly continuous functions  $v:[0,T]\to V$  and such that  $\sup_{t\in[0,T]}\|v(t)\|_{V}\leq r$ . The space  $C\left([0,T];B_{r}^{\mathrm{weak}^{*}}\right)$  is metrizable with

$$\varrho(u,v) = \sup_{t \in [0,T]} q(u(t),v(t)).$$

**Remark 61.** The unit-ball on  $L^p(0,T;V)$  is weak\*-metrizable if and only if  $L^p(0,T;V)$  is separable, in which case compactness and sequential compactness coincide.

Since by the Banach-Alaoglu Theorem  $B_r^{\text{weak}^*}$  is compact,  $(C([0,T];B_r^{\text{weak}^*}),\varrho)$  is a complete metric space.

Let us consider  $V = W_{\frac{1}{w_{\perp}}}^{-1,2}$  and  $B_R$  the corresponding ball of radius R in  $W_{\frac{1}{w_{\perp}}}^{-1,2}$ .

**Remark 62.** Note that the weak topology on  $W_{\underline{1}}^{-1,2}$  coincides with the weakest topology for which the mappings  $v \mapsto \langle v(\cdot), \varphi \rangle_{W_{\frac{1}{w_{-}}}^{-1,2} \times W_{w_{+}}^{1,2}} \in C([0,T],\mathbb{R})$  are continuous.

**Lemma 63.** Let  $\mathcal{Z} := C([0,T], W_{\frac{1}{w_{\sim}}}^{-2,2}) \cap C\left([0,T]; (W_{\frac{1}{w_{+}}}^{-1,2})^{\text{weak}^*}\right)$ , equipped with the maximum of the two topologies. Then a set  $K \subset \mathcal{Z}$  is relatively compact in  $\mathcal{Z}$ , if the following conditions hold:

- $\begin{aligned} & (1) \ \sup_{u \in K} \sup_{t \in [0,T]} \|u\|_{W^{-1,2}_{\frac{1}{w_+}}} < \infty, \\ & (2) \ \lim_{\delta \to 0} \sup_{u \in K} \sup_{s,t \in [0,T], |t-s| \le \delta} \|u(t) u(s)\|_{W^{-2,2}_{\frac{1}{w_-}}} = 0. \end{aligned}$

*Proof.* Without loss of generality, we assume that K is closed in  $\mathcal{Z}$ . The first condition allows us to work on the metric subspace  $C\left([0,T]; B_R^{\text{weak}^*}\right) \subset C\left([0,T]; (W_{\frac{1}{w_+}}^{-1,2})^{\text{weak}^*}\right)$ , for some R > 0 large enough. Due to the compact embedding of  $W_{\frac{1}{w_+}}^{-1,2} \to W_{\frac{1}{w_+}}^{-2,2}$ , by Proposition 20, we can use standard arguments (see [64]) to obtain the compactness of K in  $C([0,T],W_{\underline{1}}^{-2,2})$ . By Lemma 84, any sequence  $(u_n)_n \subset C([0,T];B_R)$ , which converges in  $W_{\frac{1}{w_{n,i}}}^{-2,2}$ , also converges in  $C([0,T]; B_R^{\text{weak}^*})$ . This finishes the proof, since we found, for any bounded sequence in K, a convergent subsequence in  $\mathcal{Z}$ .

**Lemma 64.** Let  $\{\mu_n\}_{n\in\mathbb{N}}$  be a sequence of continuous,  $\mathbb{F}$ -adapted,  $W_{\frac{1}{n}}^{-2,2}$ -valued processes such that, for some 1 < p,

- (1)  $\sup_{u \in K} \mathbb{E} \sup_{t \in [0,T]} \|u\|_{W_{-1,2}}^p < \infty$ ,
- (2)  $\forall \varepsilon > 0 \quad \forall \eta > 0 \quad \exists \delta > 0 \text{ such that for every sequence } (\tau_n)_{n \in \mathbb{N}} \text{ of } \mathbb{F}\text{-stopping times}$ with  $\tau_n \leq T$  one has

$$\sup_{n\in\mathbb{N}}\sup_{0<\theta\leq\delta}\mathbb{P}\left(\left\|\mu_{n}\left(\tau_{n}+\theta\right)-\mu_{n}\left(\tau_{n}\right)\right\|_{W_{-\frac{1}{w_{-}}}^{-2,2}}\geq\eta\right)\leq\varepsilon.$$

Let  $\mathcal{L}_n$  denote the law of  $\mu_n$  on  $\mathcal{Z}$ . Then for every  $\varepsilon > 0$ , there exists a compact subset  $K_{\varepsilon} \subset \mathcal{Z}$ , such that

$$\sup_{n} \mathcal{L}_n(K_{\varepsilon}) \ge 1 - \varepsilon.$$

*Proof.* Let  $\varepsilon > 0$ . By the Markov inequality, we infer that for every  $n \in \mathbb{N}$  and R > 0,

$$\mathbb{P}\left(\sup_{t\in[0,T]}\|u\|_{W_{\frac{1}{w_{+}}}^{-1,2}}^{p}>R_{1}\right)\leq\frac{\mathbb{E}\sup_{t\in[0,T]}\|u\|_{W_{\frac{1}{w_{+}}}^{-1,2}}^{p}}{R_{1}}\leq\frac{C}{R}.$$

Choosing  $\frac{2C}{\varepsilon} \leq R$  yields

$$\sup_{n\in\mathbb{N}} \mathbb{P}\left(\sup_{t\in[0,T]} \|u\|_{W_{\frac{1}{w_{+}}}^{-1,2}}^{p} > R\right) \leq \frac{\varepsilon}{2}.$$

By [22, Lemma 3.6, Lemma 3.8], there exists a subset  $A_{\frac{\varepsilon}{2}} \subset C([0,T], W_{\frac{1}{w_{-}}}^{-2,2})$ , such that  $\mathcal{L}_{n}(A_{\frac{\varepsilon}{2}}) \geq 1 - \frac{\varepsilon}{2}$ . We can define  $K_{\varepsilon}$  as the closure of  $B_{R} \cap A_{\frac{\varepsilon}{2}}$  in  $\mathcal{Z}$ . By Lemma 63  $K_{\varepsilon}$  is compact in  $\mathcal{Z}$  and the claim follows.

Corollary 65. Let  $\mu_n$  denote a mild solution of (1.8), given by Theorem 51. The laws of  $\{\mu_n\}_{n\in\mathbb{N}}$  are tight on  $C([0,T],W_{\frac{1}{w_n}}^{-2,2})\cap C\left([0,T];B_R^{\mathrm{weak}^*}\right)$ .

By the Skorohod representation theorem (see e.g [23, Theorem A.1], or [50]), there exists a subsequence  $(n_k)_{k\in\mathbb{N}}$ , which is not relabeled, a probability space  $(\widetilde{\Omega}, \widetilde{\mathcal{F}}, \widetilde{\mathbb{P}})$  and, on this space,  $(C([0,T],W_{\frac{1}{w_{\sim}}}^{-2,2})\cap C\left([0,T];B_R^{\text{weak}^*}\right), C^0([0,T];\mathbb{R}))$ -valued random variables  $(\widetilde{\mu},\widetilde{W})$  and  $(\widetilde{\mu}_n,\widetilde{W}_n)$  such that  $(\widetilde{\mu}_n,\widetilde{W}_n)$  has the same law as  $(\mu_n,W)$  on  $\mathcal{B}(C([0,T],W_{\frac{1}{w_{\sim}}}^{-2,2})\times C^0([0,T];\mathbb{R}))$  and, as  $n\to\infty$ ,

$$(\widetilde{\mu}_n, \widetilde{W}^n) \to (\widetilde{\mu}, \widetilde{W})$$
 in  $C([0, T], W_{\frac{1}{m}}^{-2,2}) \cap C\left([0, T]; B_R^{\text{weak}^*}\right) \times C^0([0, T]; \mathbb{R})$   $\widetilde{\mathbb{P}}$ -a.s.

The first observation is that these new variables also satisfy (3.3)  $\widetilde{\mathbb{P}}$ -a.s., replacing  $(\mu_n, W)$  by  $(\widetilde{\mu}_n, \widetilde{W})$ . The proof is done via a regularization argument, similar to [21, Theorem 2.9.1] (or [14, 25]). It remains to verify that the limit still satisfies the proposed equation. For convenience, we will drop the tilde notation and agree to work on the new probability space for the remainder of this section. In the first lemma, we investigate the convergence of the individual terms of our approximate mild solution.

**Lemma 66.** For every  $t \in [0,T]$ , we have the following convergences

- (1)  $\lim_{n\to\infty} \mathbb{E} \|\widetilde{\mu}_n(t) \widetilde{\mu}(t)\|_{W^{-2,2}_{\frac{1}{n}}} = 0.$
- $(2) \lim_{n\to\infty} \mathbb{E} \left\| \int_0^t S_{t-s}^* \nu_b b_n(s, \langle \widetilde{\mu}_n(s), \psi \rangle) \, \mathrm{d}s \int_0^t S_{t-s}^* \nu_b b(s, \langle \widetilde{\mu}(s), \psi \rangle) \, \mathrm{d}s \right\|_{W_{\underline{1}}^{-2,2}} = 0.$
- (3)  $\lim_{n\to\infty} \mathbb{E} \left\| \int_0^t S_{t-s}^* \nu(\,\mathrm{d}x) \sigma_n(s, \langle \widetilde{\mu}_n(s), \psi \rangle) \,\mathrm{d}W_s^n \int_0^t S_{t-s}^* \nu(\,\mathrm{d}x) \sigma(s, \langle \widetilde{\mu}(s), \psi \rangle) \,\mathrm{d}W_s \right\|_{W_{\infty}^{-2,2}} = 0.$

Proof. The first claim already follows from the application of the Skorohod representation theorem. First note that

$$\|S_{t-s}^* \nu(dx)\|_{W_{\frac{1}{w_{\sim}}}^{-2,2}} \le C \|S_{t-s}^* \nu(dx)\|_{W_{\frac{1}{w_{+}}}^{-1,2}}$$

$$\left\| \int_{0}^{t} S_{t-s}^{*} \nu(\,\mathrm{d}x) b_{n}(\langle \widetilde{\mu}_{n}(s), \psi \rangle) \,\mathrm{d}s - \int_{0}^{t} S_{t-s}^{*} \nu(\,\mathrm{d}x) b(s, \langle \widetilde{\mu}(s), \psi \rangle) \,\mathrm{d}s \right\|_{W_{\frac{1}{w_{\infty}}}^{-2,2}}$$

$$\leq \left\| \int_{0}^{t} S_{t-s}^{*} \nu(\,\mathrm{d}x) b_{n}(s, \langle \widetilde{\mu}_{n}(s), \psi \rangle) \,\mathrm{d}s - \int_{0}^{t} S_{t-s}^{*} \nu(\,\mathrm{d}x) b_{n}(\langle \widetilde{\mu}(s), \psi \rangle) \,\mathrm{d}s \right\|_{W_{\frac{1}{w_{\infty}}}^{-2,2}}$$

$$+ \left\| \int_{0}^{t} S_{t-s}^{*} \nu(\,\mathrm{d}x) b_{n}(s, \langle \widetilde{\mu}(s), \psi \rangle) \,\mathrm{d}s - \int_{0}^{t} S_{t-s}^{*} \nu(\,\mathrm{d}x) b(s, \langle \widetilde{\mu}(s), \psi \rangle) \,\mathrm{d}s \right\|_{W_{\frac{1}{w_{\infty}}}^{-2,2}}$$

$$\leq \left\| \int_{0}^{t} S_{t-s}^{*} \nu(\,\mathrm{d}x) |b_{n}(s, \langle \widetilde{\mu}(s), \psi \rangle) - b_{n}(s, \langle \widetilde{\mu}(s), \psi \rangle) |\,\mathrm{d}s \right\|_{W_{\frac{1}{w_{\infty}}}^{-2,2}}$$

$$+ \left\| \int_{0}^{t} S_{t-s}^{*} \nu(\,\mathrm{d}x) |b_{n}(s, \langle \widetilde{\mu}(s), \psi \rangle) - b(s, \langle \widetilde{\mu}(s), \psi \rangle) |\,\mathrm{d}s \right\|_{W_{\frac{1}{w_{\infty}}}^{-2,2}}$$

$$\leq \left\| \int_{0}^{t} S_{t-s}^{*} \nu(\,\mathrm{d}x) |\rho_{b}(\langle \widetilde{\mu}_{n}(s), \psi \rangle - \langle \widetilde{\mu}(s), \psi \rangle) |\,\mathrm{d}s \right\|_{W_{\frac{1}{w_{\infty}}}^{-2,2}}$$

$$+ \left\| \int_{0}^{t} S_{t-s}^{*} \nu(\,\mathrm{d}x) |b_{n}(s, \langle \widetilde{\mu}(s), \psi \rangle) - b(s, \langle \widetilde{\mu}(s), \psi \rangle) |\,\mathrm{d}s \right\|_{W_{\frac{1}{w_{\infty}}}^{-2,2}}$$

$$+ \left\| \int_{0}^{t} S_{t-s}^{*} \nu(\,\mathrm{d}x) |b_{n}(s, \langle \widetilde{\mu}(s), \psi \rangle) - b(s, \langle \widetilde{\mu}(s), \psi \rangle) |\,\mathrm{d}s \right\|_{W_{\frac{1}{w_{\infty}}}^{-2,2}}$$

For  $D_1$ , We obtain

$$\mathbb{E} \left\| \int_{0}^{t} S_{t-s}^{*} \nu(\,\mathrm{d}x) |\rho_{b}(\langle \widetilde{\mu}_{n}(s), \psi \rangle - \langle \widetilde{\mu}_{s}, \psi \rangle)| \,\mathrm{d}s \right\|_{W_{\infty}^{-2,2}}$$

$$\leq \int_{0}^{t} \|S_{t-s}^{*} \nu(\,\mathrm{d}x)\|_{W_{\infty}^{-2,2}} \mathbb{E} |\rho_{b}(\langle \widetilde{\mu}_{n}(s), \psi \rangle - \langle \widetilde{\mu}(s), \psi \rangle)| \,\mathrm{d}s$$

$$\leq \sup_{s} \mathbb{E} |\rho_{b}(\langle \widetilde{\mu}_{n}(s), \psi \rangle - \langle \widetilde{\mu}(s), \psi \rangle)| \int_{0}^{t} \|S_{t-s}^{*} \nu(\,\mathrm{d}x)\|_{W_{\infty}^{-2,2}} \,\mathrm{d}s.$$

Since  $\widetilde{\mu}^n \to \widetilde{\mu}$  in  $C([0,T],W_{\frac{1}{w_{\sim}}}^{-2,2}) \cap C\left([0,T];B_R^{\text{weak}^*}\right)$  for almost every  $\omega \in \Omega$ ,  $\langle \widetilde{\mu}_n(s), \psi \rangle \to \langle \widetilde{\mu}(s), \psi \rangle$  in  $C([0,T],\mathbb{R})$  for almost every  $\omega \in \Omega$ . Recall that  $\rho_b$  is a continuous function from  $\mathbb{R} \to \mathbb{R}$ , and  $\lim \rho_b(x_n - x) \to 0$ , as  $x_n \to x$ . By the linear growth assumption on b and Lemma 56, we can use the dominated convergence theorem to conclude that

$$\mathbb{E}\lim_{n\to\infty} \left\| \int_0^t S_{t-s}^* \nu(\,\mathrm{d}x) |\rho_b(\langle \widetilde{\mu}_n(s), \psi \rangle - \langle \widetilde{\mu}_s, \psi \rangle)| \,\mathrm{d}s \right\|_{W^{-2,2}_{\frac{1}{2t-s}}} = 0.$$

 $D_2$  is handled in a similar way. Let us introduce the sequence of stopping times  $\tau_m := \inf\{t \ge 0 : \sup_t \|\widetilde{\mu}_n\|_{W^{-2,2}_{\frac{1}{w_{\infty}}}} > m\}.$ 

$$\mathbb{E} \left\| \int_0^{t \wedge \tau_m} S_{t-s}^* \nu(\,\mathrm{d}x) |b_n(\langle \widetilde{\mu}(s), \psi \rangle) - b(\langle \widetilde{\mu}(s), \psi \rangle) |\,\mathrm{d}s \right\|_{W_{\infty}^{-2,2}}$$

$$\leq \mathbb{E} \int_{0}^{t \wedge \tau_{m}} \|S_{t-s}^{*} \nu(\,\mathrm{d}x)\|_{W_{\frac{1}{w_{\sim}}}^{-2,2} \mathbf{1}_{\{\tau_{m}>0\}}} |b_{n}(\langle \widetilde{\mu}(s), \psi \rangle) - b(\langle \widetilde{\mu}(s), \psi \rangle)| \,\mathrm{d}s$$

$$\leq \mathbb{E} \int_{0}^{t} \|S_{t-s}^{*} \nu(\,\mathrm{d}x)\|_{W_{\frac{1}{w_{\sim}}}^{-2,2}} \sup_{z: \|z\|_{W_{\frac{1}{w_{\sim}}}^{-2,2} \leq m}} |b_{n}(\|z\|_{W_{\frac{1}{w_{\sim}}}^{-2,2}}) - b(\|z\|_{W_{\frac{1}{w_{\sim}}}^{-2,2}})| \,\mathrm{d}s.$$

By the linear growth assumption, the dominated convergence theorem and the locally uniform convergence,

$$\lim_{k} \mathbb{E} \int_{0}^{t} \|S_{t-s}^{*} \nu(\,\mathrm{d}x)\|_{W_{\frac{1}{w_{\infty}}}^{-2,2}} \sup_{z: \|z\|_{W_{\frac{1}{w_{\infty}}}^{-2,2}} \le m} |b_{n}(\|z\|_{W_{\frac{1}{w_{\infty}}}^{-2,2}}) - b(\|z\|_{W_{\frac{1}{w_{\infty}}}^{-2,2}}) |\,\mathrm{d}s = 0.$$

The stochastic terms are treated similarly. Note that

$$\mathbb{E}\left|\int_{0}^{t} \left\|S_{t-s}^{*}\nu(\,\mathrm{d}x)\sigma_{n}(\langle\widetilde{\mu}_{n}(s),\psi\rangle) - S_{t-s}^{*}\nu(\,\mathrm{d}x)\sigma(\langle\widetilde{\mu}(s),\psi\rangle)\right\|_{W^{-2,2}_{\frac{1}{w_{\sim}}}}^{2} \,\mathrm{d}s\right|$$

$$\leq C\mathbb{E}\left[\int_{0}^{t} \left\|S_{t-s}^{*}\nu(\,\mathrm{d}x)\right\|_{W^{-2,2}_{\frac{1}{w_{\sim}}}}^{2} \left(\sigma_{n}(\langle\widetilde{\mu}_{n}(s),\psi\rangle) - \sigma_{n}(\langle\widetilde{\mu}(s),\psi\rangle)\right)^{2} \,\mathrm{d}s\right]$$

$$+ C\mathbb{E}\left[\int_{0}^{t} \left\|S_{t-s}^{*}\nu(\,\mathrm{d}x)\right\|_{W^{-2,2}_{\frac{1}{w_{\sim}}}}^{2} \left(\sigma_{n}(\langle\widetilde{\mu}(s),\psi\rangle) - \sigma(\langle\widetilde{\mu}(s),\psi\rangle)\right)^{2} \,\mathrm{d}s\right] = S_{1} + S_{2}.$$

The convergence of both terms is very similar to  $D_1$  and  $D_2$ . The argument for the convergence of the stochastic integrals is a direct adaptation of [33, Lemma 2.1] or [21, Lemma 2.6.6, step 3] to the case of convergence in  $L^1(\Omega)$ . By standard arguments, see e.g. [24], we conclude that  $(\widetilde{\Omega}, \widetilde{\mathcal{F}}, \widetilde{\mathbb{P}})$  and  $(\widetilde{\mu}, \widetilde{W})$  satisfy the conditions of Definition 4 and are a probabilistically weak, mild solution to the SEE (1.8).  $\tau_m$  was merely a localizing sequence and we can let  $m \to \infty$  and obtain the claim.

Theorem 5 follows directly from the previous Lemmata.

Since a probabilistically weak solution usually connects to a martingale problem, we briefly argue that a mild solution, as obtained above, also satisfies a weak formulation of (1.8), which relates more directly to a martingale problem. We will drop the tilde notation when working with a probabilistically weak solution.

**Lemma 67.** Let  $\mu$  be a mild solution to (3.3) and assume that  $\nu_b, \nu_\sigma$  satisfy Assumption (M 1) and set  $\theta_{\nu} = \max\{\theta_{\nu_b}, \theta_{\nu_\sigma}\}$ . Let  $\varphi \in W_{\bar{w}}^{1,2}$ , with  $\bar{w}_i = (1+x)^{2+2\max\{\theta_{\nu_b}, \theta_{\nu_\sigma}\}-1+2i}$ ,  $(w_{\theta_{\nu}})_i = (1+x)^{2\max\{\theta_{\nu_b}, \theta_{\nu_\sigma}\}-1+2i}$ , then  $\mu$  satisfied the weak formulation

(3.10) 
$$\langle \mu_t, \varphi \rangle = \langle \mu_0, \varphi \rangle + \int_0^t \langle \mu_s, -x\varphi \rangle \, \mathrm{d}s$$

$$(3.11) + \int_0^t \langle \nu_b, \varphi \rangle b(s, \langle \mu_s, \psi \rangle) \, \mathrm{d}s + \int_0^t \langle \nu_\sigma, \varphi \rangle \sigma(s, \langle \mu_s, \psi \rangle) \, \mathrm{d}W_s.$$

*Proof.* The proof is standard, so we only sketch it. Without loss of generality, assume  $\mu_0 = 0$ . Otherwise we can consider the process  $\mu_t - S_t^* \mu_0$ . Fix  $\varphi$ , apply  $-x\varphi$  to (3.3) and integrate over time.

$$\int_0^t \langle \mu_s - S_s^* \mu_0, -x\varphi \rangle \, \mathrm{d}s = \int_0^t \int_0^s \langle S_{s-r}^* \nu_b, -x\varphi \rangle b(r, \langle \mu_r, \psi \rangle) \, \mathrm{d}r \, \mathrm{d}s$$

$$+ \int_0^t \int_0^s \langle S_{s-r}^* \nu_{\sigma}, -x\varphi \rangle \sigma(r, \langle \mu_r, \psi \rangle) \, dW_r \, ds$$

$$= \int_0^t \langle \nu_b, \int_r^t -S_{s-r} x\varphi \, ds \rangle b(r, \langle \mu_r, \psi \rangle) \, dr$$

$$+ \int_0^t \langle \nu_{\sigma}, \int_r^t -S_{s-r} x\varphi \, ds \rangle \sigma(r, \langle \mu_r, \psi \rangle) \, dW_r.$$

By Lemma 36  $S^*$  is a strongly continuous semigroup on  $W_{\frac{1}{u_{\theta_{-}}}}^{-1,2}$ . Hence

$$\begin{split} \int_0^t \langle \mu_s - S_s^* \mu_0, -x\varphi \rangle \, \mathrm{d}s &= \int_0^t \langle \nu_b, S_{t-r}\varphi \rangle b(r, \langle \mu_r, \psi \rangle) \, \mathrm{d}r - \int_0^t \langle \nu_b, \varphi \rangle b(r, \langle \mu_r, \psi \rangle) \, \mathrm{d}r \\ &+ \int_0^t \langle \nu_\sigma, S_{t-r}\varphi \rangle \sigma(r, \langle \mu_r, \psi \rangle) \, \mathrm{d}W_r - \int_0^t \langle \nu_\sigma, \varphi \rangle \sigma(r, \langle \mu_r, \psi \rangle) \, \mathrm{d}W_r \\ &= \langle \mu_t - S_t^* \mu_0, \varphi \rangle - \int_0^t \langle \nu_b, \varphi \rangle b(r, \langle \mu_r, \psi \rangle) \, \mathrm{d}r - \int_0^t \langle \nu_\sigma, \varphi \rangle \sigma(r, \langle \mu_r, \psi \rangle) \, \mathrm{d}W_r. \end{split}$$

In summary, we obtain

$$\langle \mu_t - S_t^* \mu_0, \varphi \rangle - \int_0^t \langle \nu_b, \varphi \rangle b(r, \langle \mu_r, \psi \rangle) \, \mathrm{d}r - \int_0^t \langle \nu_\sigma, \varphi \rangle \sigma(r, \langle \mu_r, \psi \rangle) \, \mathrm{d}W_r$$
$$= \int_0^t \langle \mu_s, -x\varphi \rangle \, \mathrm{d}s - \langle S_s^* \mu_0, \varphi \rangle + \langle \mu_0, \varphi \rangle.$$

The previous Lemma implies in particular, that for every  $\varphi \in W_{\frac{1}{2}}^{-1,2}$ , the process

$$M_{\varphi}(t,\mu) := \langle \mu_t, \varphi \rangle - \langle \mu_0, \varphi \rangle + \int_0^t \langle -x\mu_s, \varphi \rangle \, \mathrm{d}s - \int_0^t \langle \nu, \varphi \rangle b(s, \langle \mu_s, \psi \rangle) \, \mathrm{d}s,$$

is a square-integrable  $\mathbb{F}_t$  martingale with respect to the measure P, with quadratic variation

$$\int_0^t \langle \nu, \varphi \rangle^2 \sigma(s, \langle \mu_s, \psi \rangle)^2 \, \mathrm{d}s.$$

**Remark 68.** The equivalence with the weak formulation would also allow us to derive a dual formulation of the process  $\mu$  in the case  $\sigma(x) = x^{\gamma}$ , for  $\frac{1}{2} < \gamma < 1$ . Let us set  $\psi = 1$  for simplicity. Purely formal, by Itô's formula,

$$\mathbb{E}\left[e^{-\langle\mu(t),\varphi\rangle}\right] = \mathbb{E}\left[e^{-\langle\mu_0,\varphi\rangle}\right] - \mathbb{E}\left[\int_0^t e^{-\langle\mu(s),\varphi\rangle}\left(\langle -x\mu(s),\varphi\rangle + \langle\nu,\varphi\rangle\langle\mu(s),1\rangle - \frac{1}{2}\langle\nu,\varphi\rangle^2(\langle\mu(s),1\rangle)^{2\gamma}\right) ds\right].$$

This allows us to formulate an equation for the dual process. We consider the following stochastic evolution equation

$$Y(t) = Y(0) - \int_0^t xY(s) ds + \int_0^t \langle \nu, Y(s) \rangle ds + \int_0^t \langle \nu, Y(s) \rangle^{\frac{1}{\gamma}} dL,$$

where L is a non-negative Levy process with purely positive jumps, such that

$$\mathbb{E}\left[e^{-ixL(t)}\right] = e^{tc_L \int_{0_+}^{\infty} \left(e^{-ix\lambda} - 1 + (-ix\lambda)\right)\lambda^{-2\gamma - 1} d\lambda} = e^{-ti\frac{x}{2\gamma - 1} + tc_L \int_{0_+}^{\infty} \left(e^{-ix\lambda} - 1 - \mathbf{1}_{|\lambda| < 1} ix\lambda\right)\right)\lambda^{-2\gamma - 1} d\lambda}$$

with  $c_L = \frac{1}{2} \frac{3\gamma(2\gamma-1)}{\Gamma(2-2\gamma)}$  and truncation function  $h(\lambda) = \lambda \mathbf{1}_{[0,1]}(\lambda)$ . The corresponding characteristic triplet is given by  $\left(\frac{1}{2\gamma-1}, 0, \frac{1}{\lambda^{2\gamma+1}} d\lambda\right)$ . Hence, we have that

$$\mathbb{E}\left[e^{-\int_0^t\varphi\,\mathrm{d}L(s)}\right] = \mathbb{E}\left[e^{\int_0^t\varphi^{2\gamma}\,\mathrm{d}s}\right] = \mathbb{E}\left[e^{c_L\int_0^t\int_{0_+}^\infty\left(e^{-\varphi(s)\lambda}-1+\varphi(s)\lambda\right)\right)\lambda^{-2\gamma-1}\,\mathrm{d}\lambda\,\mathrm{d}s}\right].$$

Note that  $\int_{\mathbb{R}_+} |\lambda|^2 \wedge |\lambda| \nu(d\lambda) < \infty$ . To obtain the (mild) existence of a dual process, it suffices to solve the Volterra SDE

$$V_t = g(t) + \int_0^t k_b(t-s)V_s \,ds + \int_0^t k_\sigma(t-s)V_s^{\frac{1}{\gamma}} dL_s.$$

By [38, Theorem 4.4.11], the existence of a dual process would imply

$$\mathbb{E}\left[F(\mu_t, \varphi(0))\right] = \mathbb{E}\left[F(\mu_0, \varphi(t))\right],$$

and uniqueness in law for the SEE. The same idea was used in [56]. Since [3] requires a few technical modifications to apply to this setting, we will not follow this idea rigorously in this work.

3.3. Long term existence and invariant measures. In this section, we assume for simplicity that our solution  $\mu$  is unique in law. We refer to the introduction where we mentioned several uniqueness results which, due to Theorem 6, can be transferred from the SVE to the SEE. This assumption could be avoided by proving a Markov-selection result like in [41] and carefully performing the following arguments for the corresponding selection.

**Assumption 69.** (UL) We assume that the (mild) solution to (1.8) is unique in law. (UC) We assume that the coefficients  $b, \sigma$  only depend on x.

We Assume (UC) for the remainder of this chapter, which implies that the constants from Assumption (Coeff 1) can be chosen uniformly in T.

**Proposition 70.** Let  $\mu$  be a mild solution of equation (1.8), with initial condition  $\mu_0$ , which is unique in law (Assumption (UL)). Then the family  $\{\mu_t(\mu_0)\}_{t\geq 0, \mu_0\in W^{-1,2}_{\frac{1}{2}}}$  is a time-

homogeneous Markov process and in particular  $P_{t+s} = P_t P_s$ .

Proof.

$$\mu_{t_0+t} = e^{-(t+t_0)x}\mu_0 + \int_0^{t+t_0} e^{-(t+t_0-s)x}\nu_b b(\langle \mu_s, \psi \rangle) \, \mathrm{d}s + \int_0^{t+t_0} e^{-(t+t_0-s)x}\nu_\sigma \sigma(\langle \mu_s, \psi \rangle) \, \mathrm{d}W_s$$

$$= e^{-tx} \left( e^{-t_0x}\mu_0 + \int_0^{t_0} e^{-(t_0-s)x}\nu_b b(\langle \mu_s, \psi \rangle) \, \mathrm{d}s + \int_0^{t_0} e^{-(t_0-s)x}\nu_\sigma \sigma(\langle \mu_s, \psi \rangle) \, \mathrm{d}W_s \right)$$

$$+ \int_{t_0}^{t+t_0} e^{-(t+t_0-s)x}\nu_b b(\langle \mu_s, \psi \rangle) \, \mathrm{d}s + \int_{t_0}^{t+t_0} e^{-(t+t_0-s)x}\nu_\sigma \sigma(\langle \mu_s, \psi \rangle) \, \mathrm{d}W_s$$

$$= e^{-tx}\mu_{t_0}(\mu_0) + \int_{t_0}^{t+t_0} e^{-(t+t_0-s)x}\nu_b b(\langle \mu_s, \psi \rangle) \, \mathrm{d}s + \int_{t_0}^{t+t_0} e^{-(t+t_0-s)x}\nu_\sigma \sigma(\langle \mu_s, \psi \rangle) \, \mathrm{d}W_s$$

$$= e^{-tx}\mu_{t_0}(\mu_0) + \int_0^t e^{-(t-s)x}\nu_b b(\langle \mu_{s+t_0}, \psi \rangle) \, \mathrm{d}s + \int_0^t e^{-(t-s)x}\nu_\sigma \sigma(\langle \mu_{s+t_0}, \psi \rangle) \, \mathrm{d}W_s^{t_0}.$$

By uniqueness in law,  $P_t \varphi(\mu_0) = \mathbb{E}\left[\varphi(\mu_{t_0+t}(\mu_0))\right] = \mathbb{E}\left[\varphi(\mu_t(\mu_{t_0}(\mu_0)))\right]$ . In particular,  $\mathbb{E}\left[\varphi(\mu_{t_0+t}(\mu_0)) \mid \mathcal{F}_{t_0}\right] = \mathbb{E}\left[\varphi(\mu_t(\mu_{t_0}(\mu_0))) \mid \mathcal{F}_{t_0}\right] = P_t \varphi(\mu_{t_0}(\mu_0)).$ 

**Remark 71.** Recall that  $W_{\frac{1}{w_{-}}}^{-1,2}$  was defined as the dual space of  $W_{w_{+}}^{1,2}$  with respect to the unweighted  $L^2$  duality.

This section aims to study the weak and generalized Feller properties of solutions to (42). As a first step, we want to establish a weak sequential Feller property for the solution (1.8). Let the initial condition of equation (1.8) be denoted by  $\mu_0 \in W_{\frac{1}{w_+}}^{-1,2}$  and let the space of bounded, Borel measurable functions from  $W_{\frac{1}{w_+}}^{-1,2}$  to  $\mathbb{R}$  be denoted by  $\mathcal{B}_b(W_{\frac{1}{w_+}}^{-1,2},\mathbb{R})$ . For every  $\Phi \in \mathcal{B}_b(W^{-1,2}_{\frac{1}{w_{\perp}}}, \mathbb{R})$ , we define

$$P_t\Phi(\mu_0) := \mathbb{E}\left[\Phi(\mu_t(\mu_0))\right].$$

By continuity of the trajectories of  $\mu$ ,  $P_t$  forms a stochastically continuous semigroup on  $W_{\frac{1}{w_+}}^{-1,2}$ , i.e.

$$\lim_{t\to 0} P_t \Phi(\mu_0) = \Phi(\mu_0)$$

for every  $\Phi \in C_b(W_{\frac{1}{w}}^{-1,2})$ .

**Proposition 72.** Let  $\Phi: W_{\frac{1}{w_+}}^{-1,2} \to \mathbb{R}$  be a bounded and sequentially weakly continuous function and t > 0. Then  $P_t \Phi: W_{\frac{1}{w_+}}^{-1,2} \to \mathbb{R}$  is also a bounded sequentially weakly continuous function. In particular, if  $\mu_{0,n} \to \mu_0$  in  $W_{\frac{1}{w_+}}^{-1,2}$ , then for any  $t \ge 0$ ,  $P_t\Phi(\mu_{0,n}) \to P_t\Phi(\mu_0)$ , as  $n \to \infty$ .

**Remark 73.** Referring to the work [54], this theorem verifies that  $P_t(S_b((W_{\frac{1}{w_+}}^{-1,2})^{\text{weak *}})) \subseteq$  $S_b((W_{\frac{1}{w_+}}^{-1,2})^{\text{weak}*})$ , where  $S_b((W_{\frac{1}{w_+}}^{-1,2})^{\text{weak}*})$  denotes the set of all real-valued, bounded, weakly sequentially continuous functions on  $W_{\frac{1}{w_+}}^{-1,2}$ . This property is also referred to as the sequentially weak Feller property.

Proof. Let t > 0. We start on the filtered probability space  $(\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$ . Given a sequence  $(\mu_{0,n})_{n \in \mathbb{N}} \subseteq W_{\frac{1}{w_+}}^{-1,2}$ , converging weakly to  $\mu_0 \in W_{\frac{1}{w_+}}^{-1,2}$ , we need to verify that  $P_t\Phi(\mu_{0,n}) \to P_t\Phi(\mu_0)$ , where  $P_t\Phi(\mu_0) = \int_{W_{\frac{1}{w_+}}^{-1,2}} \Phi(\mu_t(\mu_0)) d\mathbb{P}^{\mu_0}$ . By Lemma 56,  $P_t\Phi$  is bounded from  $W_{\frac{1}{w_+}}^{-1,2}$ to  $\mathbb{R}$ . Theorem 5 yields the existence of a solution  $(\Omega^{\mu_{0,n}}, \mathcal{F}^{\mu_{0,n}}, \mathbb{F}^{\mu_{0,n}}, \mathbb{P}^{\mu_{0,n}}, \mu^{\mu_{0,n}}, W^{\mu_{0,n}})$  to equation (1.8) for each  $\mu_{0,n}$ . Hence  $P_t\Phi(\mu_{0,n})$  is well defined. Let  $\rho^{\mu_{0,n}}$  denote the joint law of  $(\mu^{\mu_0,n}, W^{\mu_0,n})$ . We already know by Theorem 5 that  $(\Omega^{\mu_0}, \mathcal{F}^{\mu_0}, \mathbb{F}^{\mu_0,n}, \mathbb{F}^{\mu_0}, \mu^{\mu_0}, W^{\mu_0})$  to equation (1.8) for  $\mu_0$ . By Lemma 56 and 58 (and since we assumed that  $(\mu_{0,n})_{n\in\mathbb{N}}\subseteq W_{\underline{1}}^{-1,2}$ 

was convergent), we conclude that the laws  $(\rho^{\mu_{0,n}})_n$  are tight on  $C\left([0,T];\left(W_{\frac{1}{w^{-1}}}^{-1,2}\right)^{\text{weak}-*}\right)\cap$  $C\left([0,T];W_{\frac{1}{1}}^{-2,2}\right).$ 

We set  $\mu_{\cdot}^n = \mu_{\cdot}(\mu_{0,n})$ . The Skorohod representation theorem (see [50]) yields the existence of a subsequence  $n_k$  which will not be relabelled, a new stochastic basis  $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{F}}, \tilde{\mathbb{P}})$ , where  $\tilde{\mathbb{F}} = (\tilde{\mathcal{F}}_s)_{s \in [0,T]}$  and  $\mathbb{F}$ -progressively measurable process  $\tilde{\mu}$ ,  $(\tilde{\mu}^n)_n$  with laws supported on  $W_{\frac{1}{w_n}}^{-2,2} \cap W_{\frac{1}{w_n}}^{-1,2}$  and a new Wiener process  $\tilde{W}$ , such that  $\tilde{\mu}_n$  has the same law as  $\mu^n$  and

$$\tilde{\mu}^n \to \tilde{\mu}$$
 on  $C([0,T], W_{\frac{1}{w_{\sim}}}^{-2,2}) \cap C\left([0,T], \left(W_{\frac{1}{w_{+}}}^{-1,2}\right)^{\text{weak}^*}\right)$   $\tilde{\mathbb{P}}$ -a.s. The system  $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{F}}, \tilde{\mathbb{P}}, \tilde{W}, \tilde{\mu})$  is a probabilistically weak solution of (1.8). In particular  $\tilde{\mu}_t^n \to \tilde{\mu}_t$  (weakly-\*) in  $W_{\frac{1}{w_{+}}}^{-1,2}$ . Since

 $\Phi$  was chosen as an element of the sequentially continuous bounded functions on  $W_{\frac{1}{w_+}}^{-1,2}$ ,  $\tilde{\mathbb{P}}$ -a.s.  $\Phi(\tilde{\mu}^n) \to \Phi(\tilde{\mu})$  in  $\mathbb{R}$ . The boundedness of  $\mu$  allows us to use Lebesgue's dominated

$$\lim_{n\to\infty} \tilde{\mathbb{E}}[\Phi(\tilde{\mu}_t^n)] = \tilde{\mathbb{E}}[\Phi(\tilde{\mu}_t)].$$

By the equality of laws of  $\tilde{\mu}^n$  and  $\mu^n$ , we obtain

convergence theorem to conclude that

$$\widetilde{\mathbb{E}}[\Phi(\widetilde{\mu}_t^n)] = \mathbb{E}^{\mu_{0,n}}[\Phi(\mu_t^n)] = P_t \Phi(\mu_{0,n}).$$

By Assumption (UL), we have that the solution to (1.8) is unique in law. Hence, it must hold that

$$\widetilde{\mathbb{E}}[\Phi(\widetilde{\mu}_t)] = \mathbb{E}[\Phi(\mu_t)] = P_t \Phi(\mu_0).$$

This yields

$$\lim_{n \to \infty} P_t \Phi(\mu_{0,n}) = \lim_{n \to \infty} \tilde{\mathbb{E}}[\Phi(\tilde{\mu}_t^n)] = \tilde{\mathbb{E}}[\Phi(\tilde{\mu}_t)] = \mathbb{E}[\Phi(\mu_t)] = P_t \Phi(\mu_0).$$

Corollary 74. Let  $\varrho(x) := 1 + \|x\|_{W_{\frac{1}{w_+}}^{-1,2}}$ , then  $P_t$  as defined above, is a generalized Feller semigroup.

*Proof.* Since we satisfy the necessary assumptions, [34, Theorem 5.3] implies that the weak(\*\*) Feller property of Proposition 72 implies the generalized Feller property.  $P_t$  is therefore a strongly continuous semigroup on  $\mathcal{B}^{\varrho}\left(\left(W_{\frac{1}{w_+}}^{-1,2}\right)^{\text{weak}^*}\right)$  (see section 1.1 regarding the notation).

To make use of the weak or generalized Feller property, we will derive additional estimates.

**Assumption 75.** (LT) Let  $\nu_i$ ,  $i \in \{b, \sigma\}$ , be such that for any  $0 \le s < t$ ,

(3.12) 
$$||S_{t-s}^*\nu_i||_{W_{\frac{1}{w_+}}^{-1,2}} \le CF_i(t-s),$$

with kernels  $F_i > 0$ ,  $F_b \in C(\mathbb{R}_+) \cap L^1(\mathbb{R}_+)$  and  $F_\sigma \in C(\mathbb{R}_+) \cap L^2(\mathbb{R}_+)$ .

**Lemma 76.** Let  $\mu$  be a mild solution of equation (1.8), let Assumption (LT) be satisfied and assume that  $\sup_{T>0} \mathbb{E} \sup_{t\in[0,T]} \|S_t^*\mu_0\|_{W_{\frac{1}{w}}^{-1,2}} < \infty$ . Then

$$\sup_{t \in [0,\infty)} \mathbb{E} \|\mu_t\|_{W^{-1,2}_{\frac{1}{w_+}}} \le \sup_{T>0} \mathbb{E} \sup_{t \in [0,T]} \|\mu_t\|_{W^{-1,2}_{\frac{1}{w_+}}} < \infty.$$

*Proof.* Let u, A, G, H be non negative functions on  $\mathbb{R}_+$ . Since

$$u(t) \le A(t) + G(t)HF(t) \le A(t) + 2\max\{G(t), H(t)\},\$$

it suffices to estimate  $u(t) \leq A(t) + 2G(t)$  and  $u(t) \leq A(t) + 2H(t)$  separately. Hence for  $i \in \{b, \sigma\}, j = 1, 2$ , we consider (3.14)

$$\mathbb{E} \sup_{t \leq T} \|\mu_t\|_{W_{\frac{1}{w_+}}^{-1,2}}^p \leq C \mathbb{E} \sup_{t \leq T} \|S_t^* \mu_0\|_{W_{\frac{1}{w_+}}^{-1,2}}^p + Ch_{i,j}(T) \int_0^T F_i(T-s) \left( \mathbb{E} \sup_{r \leq s} \|\mu_r\|_{W_{\frac{1}{w_+}}^{-1,2}}^2 \right)^{\widetilde{\gamma}_{i,j}} ds,$$

where, depending on p,

$$\widetilde{\gamma}_{b,1} = \widetilde{\gamma}_{b,2} = \gamma_b, \qquad \widetilde{\gamma}_{\sigma,1} = \gamma_{\sigma}, \qquad \widetilde{\gamma}_{\sigma,2} = 2\gamma_{\sigma} - 1,$$

$$h_{b,1}(T) = h_{b,2}(T) = \left( \int_0^T \|S_{T-s}^* \nu_b\|_{W_{\frac{1}{w_+}}^{-1,2}} \, \mathrm{d}s \right)^{p-1}, \qquad h_{\sigma,1}(T) = \left( \int_0^T \|S_{T-s}^* \nu_\sigma\|_{W_{\frac{1}{w_+}}^{-1,2}} \, \mathrm{d}s \right)^{\frac{p}{2}-1}$$

 $h_{\sigma,2}(T) = 1.$ 

Since  $\tilde{\gamma}_{i,j} \leq 1$ , we can use Young's inequality to obtain

$$\mathbb{E} \sup_{t \leq T} \|\mu_{t}\|_{W_{\frac{1}{w_{+}}}^{-1,2}}^{p} \leq C \mathbb{E} \sup_{t \leq T} \|S_{t}^{*}\mu_{0}\|_{W_{\frac{1}{w_{+}}}^{-1,2}}^{p} + Ch_{i,j}(T) \int_{0}^{T} F_{i}(T-s) \left(1 + \mathbb{E} \sup_{r \leq s} \|\mu_{r}\|_{W_{\frac{1}{w_{+}}}^{-1,2}}^{p}\right) ds$$

$$\leq C \left(\mathbb{E} \sup_{t \leq T} \|S_{t}^{*}\mu_{0}\|_{W_{\frac{1}{w_{+}}}^{-1,2}}^{p} + h_{i,j}(T) \int_{0}^{T} F_{i}(T-s) ds\right)$$

$$+ Ch_{i,j}(T) \int_{0}^{T} F_{i}(T-s) \mathbb{E} \sup_{r \leq s} \|\mu_{r}\|_{W_{\frac{1}{w_{+}}}^{-1,2}}^{p} ds.$$

Now, we use [68, Lemma 2.2], which allows us to bound the previous terms by the corresponding resolvents (of the second kind) of  $F_i$ , namely

$$f(t) \le a(t) + \int_0^t F(t-s)f(s) \,\mathrm{d}s,$$

implies

$$f(t) \le a(t) + \int_0^t R_F(t-s)a(s) \,\mathrm{d}s.$$

We will also use that [42, Theorem 1] implies that if the kernel  $F \in L^1(\mathbb{R}_+)$ , then its resolvent  $R_F \in L^1(\mathbb{R}_+)$ . By Assumption (LT)  $h_{i,j} \leq C$  and we obtain

$$\mathbb{E} \sup_{t \leq T} \|\mu_t\|_{W_{t}^{-1,2}}^2 \leq C \left( 1 + \mathbb{E} \sup_{t \leq T} \|S_t^* \mu_0\|_{W_{t}^{-1,2}}^p \right) + C \int_0^T R_{F_i}(T-s) \left( 1 + \mathbb{E} \sup_{t \leq T} \|S_t^* \mu_0\|_{W_{t}^{-1,2}}^p \right) ds.$$

Assumption (LT) and [42, Theorem 1] now yield that

$$\sup_{T} \mathbb{E} \sup_{t \le T} \|\mu_t\|_{W_{\frac{1}{w_+}}^{-1,2}}^p < \infty.$$

Let us verify (LT) for two examples. Recall that in the Example 48, the kernel obtained in the estimates for the Gamma-kernel was again a Gamma-kernel. The same property can be derived for the shifted fractional kernel. Hence, we will discuss the examples in this section in terms of the kernels  $F_i$ , appearing in Assumption (LT).

- The gamma-kernel:Let  $\delta > 0$ ,  $\beta \in (0, \frac{1}{2})$  and  $F_i(t) = e^{-t\delta}t^{\beta-1}$ . Note that  $\int_0^t e^{-(t-s)\delta}(t-s)^{\beta-1} ds = \delta^{-\beta} (\Gamma(\beta) \Gamma(\beta, \delta t))$ , which can be bounded by a constant. The resolvent (of the second kind) is given by  $R(t) = e^{-\delta t}t^{\beta-1}E_{\beta,\beta}(-t^{\beta})$ , where  $E_{\alpha,\beta}(z) = \sum_{n=0}^{\infty} \frac{z^n}{\Gamma(\alpha n+\beta)}$  denotes the Mittag-Leffler function. It can be easily verified that  $\sup_T \int_0^T R(s) ds < \infty$ .
- $\sup_T \int_0^T R(s) ds < \infty$ . • For fixed  $a, \varepsilon > 0$ , let  $F_i(t) = \frac{1}{(t+\varepsilon)^{1+\alpha}}$ . This kernel is clearly in  $L^1(\mathbb{R}_+)$ . Note that kernels of this form are completely monotone and fit in our previous analysis.

**Theorem 77.** Let  $\mu$  be a solution of (1.8) and let Assumptions (LT) hold. Then there exists at least one measure  $\mathcal{Q}$  on  $W_{\frac{1}{w_+}}^{-1,2}$ , such that  $P_t^*\mathcal{Q} = \mathcal{Q}$ .

*Proof.* The statement follows directly from [54, Proposition 3.1] and estimate (3.13).

Proof of Theorem 8. The theorem follows from Theorem 77, Proposition 72 and Corollary  $\overline{\phantom{a}}$ 

**Remark 78.** We could have obtained the result via the strategy used in [49]. Since  $(P_t)_{t\geq 0}$  be the generalised Feller semigroup associated to  $(\mu_t)_{t\geq 0}$ . Hence, for any  $(W_{\frac{1}{w_+}}^{-1,2})^{\text{weak}^*}$ -valued random variable  $\mu_0 \sim n$ . Pro  $(\mu_0) = \mathbb{E}_{w_+}[\rho(\mu_t)]$ 

random variable  $\mu_0 \sim \eta$ ,  $P_t \varrho (\mu_0) = \mathbb{E}_{\mu_0} [\varrho (\mu_t)]$ . We can use the weight  $\varrho(\mu) = 1 + \|\mu\|_{W^{-1,2}_{\frac{1}{w_+}}}^p$ , which is an admissible weight according to the

Definition [34, Definition 2.1].

$$\sup_{t\geq 0} \int_{W_{\frac{1}{w}}^{-1,2}} \mathbb{E}_{\mu_0} \left[ \varrho \left( \mu_t \right) \right] \, \mathrm{d}\eta \left( \mu_0 \right) = \sup_{t\geq 0} \mathbb{E} \left[ \varrho \left( \mu_t \right) \right] < \infty,$$

is enough for an application of [49, Lemma 3.1].

Proof of Theorem 9. Theorem 77 implies the existence of a probability measure  $\mathcal{Q}$  on  $W_{\frac{1}{w_+}}^{-1,2}$  such that, if  $\mu_0 = \delta_0 x_0 \sim \mathcal{Q}$ , for any  $t \geq 0$ ,  $\mathcal{Q}$  is the Law of  $\mu_t$ .  $(\langle \mu_t, 1 \rangle)_{t \geq s}$  is distributed according to the push-forward measure of  $\mathcal{Q}$  under the map  $\mu \mapsto \langle \mu, 1 \rangle$  and also strictly stationary.

### 4. Ito-formula for Volterra equations

Our last application of the lifting procedure is an Itô-type formula for SVEs. We will rely on the results from [32] (see also [28, 9]). For this, we consider the separable Hilbert spaces U, V, H, V' such that  $V \hookrightarrow H \hookrightarrow V'$  is continuous and dense. In this section, we impose Assumptions (M 2) and (M 1). Let  $\theta_{\nu} = \max\{\theta_{\nu_b}, \theta_{\nu_\sigma}\}$  and  $\varepsilon < 1 - \theta_{\nu}$ . We can simply set  $U = \mathbb{R}, \ V' = W_{\frac{1}{w_-}}^{-1,2}$ , with  $(w_-)_i(x) = (1+x)^{2\theta_{\nu}-1+2i}$ ,  $H = W_{\frac{1}{w_-}}^{-1,2}$ , with  $(w_-)_i(x) = (1+x)^{(\theta_{\nu}-\varepsilon)-1+2i}$  and  $V = W_{\frac{1}{w_+}}^{-1,2}$ , with  $(w_+)_i(x) = (1+x)^{-2\varepsilon-1+2i}$ . Let  $\Phi \in C^{1,2}([0,T] \times V, \mathbb{R})$ ,

then we denote the the partial Frechet derivatives of  $\Phi$  by

$$\partial_x \Phi \in C([0,T] \times V, L(V,\mathbb{R})), \partial_x \Phi = \frac{\partial \Phi}{\partial x}$$

$$\partial_x^2 \Phi \in C([0,T] \times V, L(V,L(V,\mathbb{R}))), \partial_x^2 \Phi = \frac{\partial^2 \Phi}{\partial x^2}$$

$$\partial_t \Phi \in C([0,T] \times V,\mathbb{R}), \partial_t \varphi = \frac{\partial \Phi}{\partial t}.$$

**Remark 79.** Our spaces changed slightly in this section since we no longer require the compactness of any embeddings.

We recall the following definition from [32].

**Definition 80.** (Mild Itô process). Let  $S^*: \Delta_t \to L(V',V)$  be a  $\mathcal{B}(\Delta_t)/\mathcal{S}(V',V)$ -measurable mapping satisfying  $S^*_{t_2,t_3}S^*_{t_1,t_2} = S^*_{t_1,t_3}$  for all  $t_1,t_2,t_3 \in [0,T]$  with  $t_1 < t_2 < t_3$ . Additionally, let  $Y: [0,T] \times \Omega \to V'$  and  $Z: [0,T] \times \Omega \to HS(U,V')$  be two predictable stochastic processes with  $\int_{\tau}^{t} \|S^*_{s,t}Y_s\|_{V} ds < \infty$ ,  $\mathbb{P}$ -a.s. and  $\int_{\tau}^{t} \|S^*_{s,t}Z_s\|_{HS(U,V)}^2 ds < \infty$   $\mathbb{P}$ -a.s. for all  $t \in [0,T]$ . Then a predictable stochastic process  $X: [0,T] \times \Omega \to H$  satisfying

$$X_{t} = S_{\tau,t}^{*} X_{\tau} + \int_{\tau}^{t} S_{s,t}^{*} Y_{s} \, \mathrm{d}s + \int_{\tau}^{t} S_{s,t}^{*} Z_{s} \, \mathrm{d}W_{s}$$

 $\mathbb{P}$ -a.s. for all  $t \in [0,T] \cap (\tau,\infty)$  is called a mild Itô process (with semigroup  $S^*$ , mild drift Y and mild diffusion Z).

By Lemma 36, 38,  $S^*$  satisfies the mapping properties mentioned in Definition 80.

**Theorem 81.** [32, Theorem 1] Let  $X : [0,T] \times \Omega \to H$  be a mild Itô process with semigroup  $S^* : \Delta_t \to L(V',V)$ , mild drift  $Y : [0,T] \times \Omega \to V'$  and mild diffusion  $Z : [0,T] \times \Omega \to HS(U,V')$ . Then

$$\mathbb{P}\left[\int_{t_{0}}^{t} \left\| (\partial_{2}\Phi) \left( s, S_{s,t}^{*}X_{s} \right) S_{s,t}^{*}Y_{s} \right\|_{V} + \left\| (\partial_{2}\Phi) \left( s, S_{s,t}^{*}X_{s} \right) S_{s,t}^{*}Z_{s} \right\|_{HS(U,V)}^{2} \, \mathrm{d}s < \infty \right] = 1$$

$$\mathbb{P}\left[\int_{t_{0}}^{t} \left\| (\partial_{1}\Phi) \left( s, X_{s} \right) \right\|_{V} + \left\| \left( \partial_{2}^{2}\Phi \right) \left( s, S_{s,t}^{*}X_{s} \right) \right\|_{L^{(2)}(V,V)} \left\| S_{s,t}^{*}Z_{s} \right\|_{HS(U,V)}^{2} \, \mathrm{d}s < \infty \right] = 1$$

and

$$\Phi(t, X_t) = \Phi(t_0, S_{t_0, t}^* X_{t_0}) + \int_{t_0}^t (\partial_1 \Phi) \left( s, S_{s, t}^* X_s \right) ds + \int_{t_0}^t (\partial_2 \Phi) \left( s, S_{s, t}^* X_s \right) S_{s, t}^* Y_s ds$$

$$+ \int_{t_0}^t (\partial_2 \Phi) \left( s, S_{s, t}^* X_s \right) S_{s, t}^* Z_s dW_s + \frac{1}{2} \sum_{j \in \mathbb{N}} \int_{t_0}^t (\partial_2^2 \Phi) \left( s, S_{s, t}^* X_s \right) \left( S_{s, t}^* Z_s e_j, S_{s, t}^* Z_s e_j \right) ds$$

 $\mathbb{P}$ -a.s. for all  $t_0, t \in [0, T]$  with  $t_0 < t$  and all  $\Phi \in C^{1,2}([0, T] \times V, \mathbb{R})$ .

Corollary 82. Let  $k_b$ ,  $k_\sigma$  be completely monotone kernels such that their associated measures, given by (1.3), satisfy Assumptions (M 1), (M 2). Let X denote the solution of the corresponding stochastic Volterra equation (1.1),  $S^*$  the semigroup from section 3.1 and  $f \in C^{1,2}$ . Then the following Itô formula holds.

$$f(t, X_t) = f(t_0, \langle S_{t_0, t}^* \mu_{t_0}, 1 \rangle) + \int_{t_0}^t \partial_s f(s, \langle S_{s, t}^* \mu_s, 1 \rangle) \, \mathrm{d}s + \int_{t_0}^t \partial_x f(s, \langle S_{s, t}^* \mu_s, 1 \rangle) k_b(t - s) b(s, X_s) \, \mathrm{d}s$$

$$+ \int_{t_0}^{t} \partial_x f(s, \langle S_{s,t}^* \mu_s, 1 \rangle) k_{\sigma}(t-s) \sigma(s, X_s) dW_s$$
$$+ \frac{1}{2} \int_{t_0}^{t} \left( \partial_x^2 f(s, \langle S_{s,t}^* \mu_s, 1 \rangle) \right) k_{\sigma}(t-s)^2 \sigma(s, X_s)^2 ds$$

*Proof.* We lift the SVE to the infinite-dimensional evolution equation (1.9). By Theorem 5,  $\mu \in C([0,T], W_{\frac{1}{w_+}}^{-1,2})$ . The required mapping property of the semigroup  $S^*$  is a consequence

of the inequality  $e^{-y} \leq \left(\frac{1}{1+y}\right)^{\theta}$ , for any  $\theta \in [0,1)$  and y > -1. Let  $\varphi(\mu) = f(t, \langle \mu, g \rangle)$ . The mild Itô formula now yields

$$\varphi(t,\mu_t) = f(t,\langle\mu_t,g\rangle) = f(t_0,\langle S_{t_0,t}^*\mu_0,g\rangle) + \int_{t_0}^t \partial_t f(s,\langle S_{s,t}^*\mu_s,g\rangle) \,\mathrm{d}s + \int_{t_0}^t \partial_x f(s,\langle S_{s,t}^*\mu_s,g\rangle) \langle S_{s,t}^*Y_s,g\rangle \,\mathrm{d}s \\ + \int_{t_0}^t \partial_x f(s,\langle S_{s,t}^*\mu_s,g\rangle) \langle S_{s,t}^*Z_s,g\rangle \,\mathrm{d}W_s + \frac{1}{2} \sum_{j \in \mathbb{N}} \int_{t_0}^t \left(\partial_x^2 f(s,\langle S_{s,t}^*\mu_s,g\rangle)\right) \left(S_{s,t}^*Z_s e_j, S_{s,t}^*Z_s e_j\right) \,\mathrm{d}s \\ = f(t_0,\langle S_{t_0,t}\mu_0,g\rangle) + \int_{t_0}^t \partial_t f(s,\langle S_{s,t}^*\mu_s,g\rangle) \,\mathrm{d}s \\ + \int_{t_0}^t \partial_x f(s,\langle S_{s,t}^*\mu_s,g\rangle) \langle S_{s,t}^*\nu b(s,\langle \mu_s,\varphi\rangle),g\rangle \,\mathrm{d}s \\ + \int_{t_0}^t \partial_x f(s,\langle S_{s,t}^*\mu_s,g\rangle) \langle S_{s,t}^*\nu \sigma(s,\langle \mu_s,\varphi\rangle),g\rangle \,\mathrm{d}W_s \\ + \frac{1}{2} \int_{t_0}^t \left(\partial_x^2 f(s,\langle S_{s,t}^*\mu_s,g\rangle)\right) \left(S_{s,t}^*\nu \sigma(s,\langle \mu_s,\varphi\rangle), S_{s,t}^*\nu \sigma(s,\langle \mu_s,\varphi\rangle)\right) (g \otimes g) \,\mathrm{d}s$$

Setting q = 1 yields

$$f(t, X_t) = f(t_0, \langle S_{t_0, t}^* \mu_0, 1 \rangle) + \int_{t_0}^t \partial_s f(s, \langle S_{s, t}^* \mu_s, 1 \rangle) \, \mathrm{d}s + \int_{t_0}^t \partial_x f(s, \langle S_{s, t}^* \mu_s, 1 \rangle) k(t - s) b(s, X_s) \, \mathrm{d}s$$

$$+ \int_{t_0}^t \partial_x f(s, \langle S_{s, t}^* \mu_s, 1 \rangle) k(t - s) \sigma(s, X_s) \, \mathrm{d}W_s$$

$$+ \frac{1}{2} \int_{t_0}^t \left( \partial_x^2 f(s, \langle S_{s, t}^* \mu_s, 1 \rangle) \right) k(t - s)^2 \sigma(s, X_s)^2 \, \mathrm{d}s$$

*Proof of Proposition* 10. The only part which is left to show is that

$$\mathbb{E}\left[X_t \mid \mathcal{F}_s^X\right] - \int_s^t k_b(t-r)\mathbb{E}\left[b(r,X_r) \mid \mathcal{F}_s^X\right] dr - \int_s^t k_b(t-r)\mathbb{E}\left[\sigma(r,X_r) \mid \mathcal{F}_s^X\right] dW_r$$
$$= \mathbb{E}\left[\langle e^{-x(t-s)}\mu_s, 1\rangle \mid \mathcal{F}_s^\mu\right] = \langle e^{-x(t-s)}\mu_s, 1\rangle.$$

This, however, follows directly from taking conditional expectations and noticing that the filtration generated by  $(X_t)_t$  is the same as the one generated by  $(\langle \mu_t, 1 \rangle)_t$ , since  $(X_t)_t$  is simply a finite-dimensional marginal of  $(\mu_t)_t$ . Hence the law of  $X_t$  conditioned on the filtration

 $\mathcal{F}^X$  coincides with the law of  $X_t$ , conditioned on the filtration generated by  $(\mu_t)_t$ . We obtain

$$\mathbb{E}\left[X_{t} \mid \mathcal{F}_{s}^{\mu}\right] = \mathbb{E}\left[\langle \mu_{t}, 1 \rangle \mid \mathcal{F}_{s}^{\mu}\right] = \mathbb{E}\left[\langle e^{-x(t-s)}\mu_{s}, 1 \rangle \mid \mathcal{F}_{s}^{\mu}\right] + \mathbb{E}\left[\int_{s}^{t} k_{b}(t-r)b(r, X_{r}) \, \mathrm{d}r + \int_{s}^{t} k_{\sigma}(t-r)\sigma(r, X_{r}) \, \mathrm{d}W_{r} \mid \mathcal{F}_{s}^{\mu}\right].$$

The next Corollary is now a direct consequence of Proposition 10.

**Corollary 83.** Let X be a solution of the stochastic Volterra equation (1.1). Suppose that there exists a  $C \geq 0$ , such that  $|X_0| \leq C$ , almost surely. If there exists a function  $V \in C^2(\mathbb{R}, \mathbb{R}_+)$ , constants  $0 < d, h, c_1, c_2 < \infty$  and  $1 \leq p$  which satisfy

(1) 
$$c_1|x|^p \leq V(x) \leq c_2|x|^p$$
 for all  $x \in \mathbb{R}$ ,

(2)

$$\mathcal{L}V(x,t,s) := V'(\Gamma_{st})k_b(t-s)b(X_s) + V''(\Gamma_{st})k_{\sigma}(t-s)^2\sigma(X_s)^2 \le hV(X_s) + d,$$

then the solution X is a global solution of the stochastic Volterra equation, in the sense that for each  $0 \le t < \infty$ ,  $\mathbb{E}[|X_t|] < \infty$ .

*Proof.* By the Itô formula for Volterra equations, for any t > 0,

$$V(X_t) = V(\mathbb{E}\left[\Gamma_{0t}|\mathcal{F}_0^X\right])$$

$$+ \int_0^t V'(\mathbb{E}\left[\Gamma_{st}|\mathcal{F}_s^X\right])k_b(t-s)b(X_s) + V''(\mathbb{E}\left[\Gamma_{st}|\mathcal{F}_s^X\right])k_\sigma(t-s)^2\sigma(X_s)^2 ds$$

$$+ \int_0^t V'(\Gamma_{st})k_\sigma(t-s)\sigma(X_s) dW_s$$

Taking the expectation, and using  $\mathcal{L}V(x,t,s) \leq hV(X_t) + d$ , we get

$$\mathbb{E}\left[V(X_t)\right] \le \mathbb{E}\left[V(X_0)\right] + h \int_0^t \mathbb{E}\left[V(X_s)\right] \, \mathrm{d}s + td.$$

Gronwall's inequality now yields

$$\mathbb{E}\left[V(X_t)\right] \le \left(\mathbb{E}\left[V(X_0)\right] + td\right)e^{ht}.$$

The statement follows by the assumption that  $V(x) \leq c_2|x|$ .

#### 5. Appendix

# 5.1. Proofs of technical results.

Of Theorem 6. (1) Let X be a solution of the SVE (1.10) with  $X_0 = x_0$ . Define

(5.1) 
$$Y_t = e^{-xt}Y_0 + \int_0^t e^{-x(t-s)}\nu_b(x)b(s, X_s) ds + \int_0^t e^{-x(t-s)}\nu_\sigma(x)\sigma(s, X_s) dW_s.$$

By the assumed integrability (1.11), Lemma 27 and Lemma 40, Y is a  $W_{\frac{1}{w}}^{-1,2}$ -valued adapted process and satisfies  $\int_0^T \|Y_t\|_{W_{\frac{1}{w}}}^2 dt < \infty$  a.s. for any  $t \geq 0$ , due to Young's convolution inequality. By the same estimates, performed in Section 3.2.2, Y is a

continuous  $W_{\frac{1}{w_{-}}}^{-2,2}$ -valued process, where  $w_{-}$  is given as in Definition 3. We note that the map  $f \mapsto \langle f, 1 \rangle$  from  $W_{\underline{1}}^{-1,2}$  to  $\mathbb R$  is linear. Hence,

$$\langle Y_t, 1 \rangle = \langle e^{-xt} Y_0, 1 \rangle + \int_0^t \langle e^{-x(t-s)} \nu_b(x), 1 \rangle b(s, X_s) \, \mathrm{d}s + \int_0^t \langle e^{-x(t-s)} \nu_\sigma(x), 1 \rangle \sigma(s, X_s) \, \mathrm{d}W_s$$
$$= x_0 + \int_0^t k_1(t-s) b(s, X_s) \, \mathrm{d}s + \int_0^t k_2(t-s) \sigma(s, X_s) \, \mathrm{d}W_s = X_t,$$

a.s. for a.e. t > 0. Therefore Y is a mild solution to (1.13) and  $X_t = \langle Y_t, 1 \rangle$  a.s. for a.e. t > 0.

(2) We now prove the converse part. Assume  $\mu$  is a mild solution of (1.13) and define the process  $X_t$  by

$$X_t := \langle \mu_t, 1 \rangle = \langle e^{-xt} \mu_0, 1 \rangle + \int_0^t \langle e^{-x(t-s)} \nu_b(x), 1 \rangle b(s, \langle \mu_s, 1 \rangle) \, \mathrm{d}s + \int_0^t \langle e^{-x(t-s)} \nu_\sigma(x), 1 \rangle \sigma(s, \langle \mu_s, 1 \rangle) \, \mathrm{d}W_s.$$

 $\mu$  is a  $W_{\frac{1}{w}}^{-1,2}$ -valued process and the map  $f \mapsto \langle f, 1 \rangle$  from  $W_{\frac{1}{w}}^{-1,2}$  to  $\mathbb{R}^n$  is continuous and hence Borel. Using assumptions and the properties of the measures  $\nu_b, \nu_\sigma$ , it can be shown that  $t \mapsto \langle \mu_t, 1 \rangle$  is continuous. This implies that X is a predictable  $\mathbb{R}$ -valued process. Applying the map  $f \mapsto \langle f, 1 \rangle$  to the mild formulation

$$\mu_t = e^{-xt}\mu_0, + \int_0^t e^{-x(t-s)}\nu_b(x)b(s, \langle \mu_s, 1 \rangle) \,\mathrm{d}s + \int_0^t e^{-x(t-s)}\nu_\sigma(x)\sigma(s, \langle \mu_s, 1 \rangle) \,\mathrm{d}W_s,$$

we see that

$$X_{t} = \left\langle e^{-t} \mu_{0}(\cdot) + \int_{0}^{t} e^{-(t-s)} \nu_{b}(\cdot) b\left(s, \langle \mu_{s}, 1 \rangle\right) \, \mathrm{d}s + \int_{0}^{t} e^{-(t-s)} \nu_{\sigma}(\cdot) \sigma\left(s, \langle \mu_{s}, 1 \rangle\right) \, \mathrm{d}W_{s}, 1 \right\rangle$$

$$= x_{0} + \int_{0}^{t} K(t-s) b\left(s, \langle \mu_{s}, 1 \rangle\right) \, \mathrm{d}s + \int_{0}^{t} K(t-s) \sigma\left(s, \langle \mu_{s}, 1 \rangle\right) \, \mathrm{d}W_{s}$$

$$= x_{0} + \int_{0}^{t} K(t-s) b\left(s, X_{s}\right) \, \mathrm{d}s + \int_{0}^{t} K(t-s) \sigma\left(s, X_{s}\right) \, \mathrm{d}W_{s},$$

a.s. for a.e. t > 0. Therefore, X is a solution of the SVE (1.10) with  $X_0 = x_0$ . We have

$$Y_{t} = e^{-t}Y_{0} + \int_{0}^{t} e^{-(t-s)} \nu_{b} b(s, X_{s}) ds + \int_{0}^{t} e^{-(t-s)} \nu_{\sigma} \sigma(s, X_{s}) dW_{s}$$

a.s. for any  $t \geq 0$ . This completes the proof.

Proof of Proposition 20. (1) Let  $\{u_n\}_n$  be a bounded sequence in  $W_w^{s,2}$  and let  $B_m(0)$  denote the ball with radius  $2^m$  in  $\mathbb{R}$ . First, we note that the restriction of  $W_w^{s,p}$  to  $W_w^{s,p}(B_m(0))$  is a continuous map. By Assumption 1,

$$\sum_{0 \le |j| \le s} \int_{B_m(0)} |D^j u|^p \, dx \le \sum_{0 \le |j| \le s} \int_{B_m(0)} |D^j u|^p w_j(x) \, dx \sup_{x \in B_m(0)} \left| \frac{1}{w_j(x)} \right|,$$

$$\sum_{0 \le |j| \le s} \int_{B_m(0)} |D^j u|^p w_j(x) \, dx \le C_R \sum_{0 \le |j| \le s} \int_{B_m(0)} |D^j u|^p \, dx.$$

Г

By our assumptions on the weight w, the sequence  $\{u_n\}_n$  is also bounded in  $W^{s,p}(B_m(0))$ . By Rellich's embedding theorem (see [8, Theorem 4.12]),  $W^{s,p}(B_m(0)) \hookrightarrow W^{l,q}(B_m(0))$ , when the parameters s, l, p, q satisfy the relation in the statement. This allows us to extract a subsequence  $\{u_{n_k}\}_k$  converging in  $W^{l,q}(B_m(0))$  and a further subsequence converging almost everywhere to a limit  $u^m$ . This sequence is still bounded in  $W^{s,p}_w$ . We can iterate the above procedure on a bigger ball with radius  $2^{m+1} > 2^m$ , yielding a further subsequence converging almost everywhere to a limit  $u^{m+1}$ . Due to the almost everywhere convergence,  $u^m = u^{m+1}$  on  $B_{m+1}(0)$ . On each consecutive ball, we can extract a new almost-surely convergent subsequence. In the following, we will not rename the extracted sub-sequences. This construction yields a diagonal sequence and a function u, such that  $u(x) = u^m(x)$  with m chosen such that for each  $x \in B_m(0)$ ,  $u_n^m(x) \stackrel{n \to \infty}{\to} u^m(x) = u(x)$  almost everywhere. According to Fatou's Lemma,

$$\sum_{0 \le |j| \le s} \int_{\mathbb{R}^d} \left| D^j u \right|^p w_j(x) \, \mathrm{d}x \le \liminf_{m \to \infty} \sum_{0 \le |j| \le s} \int_{\mathbb{R}^d} \left| D^j u_m^m \right|^p w_j(x) \, \mathrm{d}x \le C.$$

Let  $M \in \mathbb{N}$ ,  $M \geq K$ , where K was given in the assumptions. Recall that for every  $0 \leq |j| \leq l$ ,  $w'_j(x) > w_j(x)$  for all  $x \in \mathbb{R} \backslash B_M(0)$  and  $\frac{w_j(x)}{w'_j(x)} \to 0$  as  $|x| \to \infty$ . For  $u \in W^{l,q}_{w'}(\mathbb{R}^d \backslash B_M(0))$ ,

$$\sum_{j \leq l} \int_{\mathbb{R}^d \backslash B_M(0)} \left| D^j u \right|^q w_j(x) \, \mathrm{d}x \leq \sum_{j \leq l} \int_{\mathbb{R}^d \backslash B_M(0)} \left| D^j u \right|^q \frac{w_j(x)}{w_j'(x)} w_j'(x) \, \mathrm{d}x$$

$$\leq \sum_{j \leq l} \int_{\mathbb{R}^d \backslash B_M(0)} \left| D^j u \right|^q w_j'(x) \, \mathrm{d}x \frac{w_j(M)}{w_j'(M)}.$$

For M large enough,  $\frac{w_j(M)}{w_i'(M)} \leq \varepsilon$ , for any  $\varepsilon > 0$ . Hence,

$$\sum_{j \leq l} \int_{\mathbb{R}^d} \left| D^j u_m - D^j u \right|^q w_j(x) \, \mathrm{d}x = \sum_{j \leq l} \int_{B_M(0)} \left| D^j u_m - D^j u \right|^q w_j(x) \, \mathrm{d}x$$

$$+ \sum_{j \leq l} \int_{\mathbb{R}^d \setminus B_M(0)} \left| D^j u_m - D^j u \right|^q w_j(x) \, \mathrm{d}x$$

$$\leq \varepsilon + C \sum_{j \leq l} \frac{w_j(M)}{w_j(M)} \int_{\mathbb{R}^d \setminus B_M(0)} \left( \left| D^j u_m \right|^p + \left| D^j u \right|^p \right) w_j'(x) \, \mathrm{d}x$$

$$\leq (l+1)\varepsilon.$$

This finishes the proof of the first statement.

(2) Let  $B_R(0)$  be again the ball with radius R. By Assumption 1, there exists a constant such that

$$||u||_{L_w^{\infty}(B_R(0))} \le C_{w,R}||u||_{L^{\infty}(B_R(0))} \le C_{w,R}||u||_{W^{s,p}(B_R(0))}.$$

(a) Let there be an  $i \in \{0, ..., s\}$ , such that  $w_i(0) = 0$ . Since we assumed  $w_k \in \mathcal{B}^{\rho}(\mathbb{R}^d)$  for every k = 0, ..., s,

$$\left( \int_{B_R(0)} |D^j u(x)|^p \, \mathrm{d}x \right)^{\frac{1}{p}} = \left( \int_{B_R(0)} |D^j u(x)|^p \frac{w_j(x)^{1/\rho}}{w_j(x)^{1/\rho}} \, \mathrm{d}x \right)^{\frac{1}{p}}$$

$$\leq \left( \int_{B_R(0)} |D^j u(x)|^{p\rho} w_j(x) \, \mathrm{d}x \right)^{\frac{1}{p\rho}} \left( \int_{B_R(0)} \frac{1}{w_j(x)^{\frac{1}{\rho-1}}} \, \mathrm{d}x \right)^{\frac{\rho-1}{p\rho}} \\
\leq C \|D^j u\|_{L_w^{p\rho}(B_R(0))}.$$

By classical results, we know that  $u \in C(B_R(0))$  in each of the respective cases by [8, Theorem 4.12, Part II]. Since continuity and differentiability are local properties, and we assumed that  $w \in L^{\infty}_{loc}$ , we can shift the center of the ball and obtain the continuity and differentiability properties on the whole space. The statements including the boundedness and weights however require more work. Let  $A_R := B_{2R} \setminus \overline{B}_R$ . By [8, Theorem 4.12, Part I],  $W^{l,p}(B_{2R}) \hookrightarrow L^{\infty}(B_{2R})$ continuously. On  $A_R$ , with R > 1,

$$||u(\cdot)||_{L^{\infty}(A_R)} = ||u(R\cdot)||_{L^{\infty}(A_1)} \le C_{A_1} ||u(R\cdot)||_{W^{l,p}(A_1)} \le C_{A_1} \left( \sum_{j=0}^{l} R^{jp-d} ||D^j u||_{L^p(A_R)}^p \right)^{1/p}.$$

 $_{
m Hence,}$ 

$$\begin{split} \|u(\cdot)w_{c}(\cdot)\|_{L^{\infty}(A_{R})} &\leq \|w_{c}(\cdot)\|_{L^{\infty}(A_{R})} \|u(R\cdot)\|_{L^{\infty}(A_{1})} \\ &\leq C_{A_{1}} \|w_{c}(\cdot)\|_{L^{\infty}(A_{R})} \left( \sum_{j=0}^{l} R^{jp-d} \|D^{j}u\|_{L^{p}(A_{R})}^{p} \right)^{1/p} \\ &\leq C_{A_{1}} \|w_{c}(\cdot)\|_{L^{\infty}(A_{R})} \left( \sum_{j=0}^{l} R^{jp-d} \|w_{j}^{-\frac{1}{\rho-1}}(R\cdot)\|_{L^{1}(A_{1})}^{\frac{\rho-1}{\rho}} \|D^{k}u\|_{L^{p/\rho}_{w}(A_{R})}^{p/\rho} \right)^{1/p} \\ &\leq C_{A_{1}} \left( \sum_{j=0}^{l} R^{jp-d} \|w_{c}(\cdot)\|_{L^{\infty}(A_{R})}^{p} \|w_{j}^{-\frac{1}{\rho-1}}(R\cdot)\|_{L^{1}(A_{1})}^{\frac{\rho-1}{\rho}} \|D^{k}u\|_{L^{p/\rho}_{w}(A_{R})}^{p/\rho} \right)^{1/p} \\ &\leq C_{A_{1},l} \left( \sum_{j=0}^{l} R^{jp-d} \|w_{c}(\cdot)\|_{L^{\infty}(A_{R})}^{p} \|w_{j}^{-\frac{1}{\rho-1}}(R\cdot)\|_{L^{1}(A_{1})}^{\frac{\rho-1}{\rho}} \right)^{1/p} \|u\|_{W_{w}^{l,p\rho}} \\ &\leq C_{A_{1},l} \sum_{j=0}^{l} R^{j-\frac{d}{p}} \|w_{c}(\cdot)\|_{L^{\infty}(A_{R})} \|w_{j}^{-\frac{1}{\rho-1}}(R\cdot)\|_{L^{1}(A_{1})}^{\frac{\rho-1}{p}} \|u\|_{W_{w}^{l,p\rho}}. \end{split}$$

By Assumption, the terms on the right-hand side are bounded, which yields the claim. If s-l>1, the argument can be repeated with the derivatives of u. The boundedness of u follows from the identical steps, with  $w_c=1$ .

(b) The second case is even easier, since

$$\left( \int_{B_R(0)} |D^j u(x)|^p \, \mathrm{d}x \right)^{\frac{1}{p}} \le \frac{1}{c^{1/p}} \left( \int_{B_R(0)} |D^j u(x)|^p w_j(x) \, \mathrm{d}x \right)^{\frac{1}{p}}.$$

Considering again the annuli outside  $B_R(0)$ ,

$$||u(\cdot)w_c(\cdot)||_{L^{\infty}(A_R)} \le ||w_c(\cdot)||_{L^{\infty}(A_R)} ||u(R\cdot)||_{L^{\infty}(A_1)}$$

$$\leq C_{A_{1}} \|w_{c}(\cdot)\|_{L^{\infty}(A_{R})} \left( \sum_{j=0}^{l} R^{jp-d} \|D^{j}u\|_{L^{p}(A_{R})}^{p} \right)^{1/p}$$

$$\leq C_{A_{1}} \|w_{c}(\cdot)\|_{L^{\infty}(A_{R})} \left( \sum_{j=0}^{l} R^{jp-d} \inf_{y \in A_{R}} \frac{1}{w_{j}(y)} \|D^{k}u\|_{L^{p}_{w}(A_{R})}^{p} \right)^{1/p}$$

$$= C_{A_{1}} \left( \sum_{j=0}^{l} R^{jp-d} \sup_{y \in A_{R}} |w_{c}(y)|^{p} \inf_{y \in A_{R}} \frac{1}{w_{j}(y)} \|D^{k}u\|_{L^{p}_{w}(A_{R})}^{p} \right)^{1/p}$$

$$\leq C_{A_{1},l} \left( \sum_{j=0}^{l} R^{jp-d} \sup_{y \in A_{R}} |w_{c}(y)|^{p} \inf_{y \in A_{R}} \frac{1}{w_{j}(y)} \right)^{1/p} \|u\|_{W^{l,p}_{w}}$$

$$\leq C_{A_{1},l} \|u\|_{W^{l,p}_{w}} \sum_{j=0}^{l} R^{j-\frac{d}{p}} \sup_{y \in A_{R}} |w_{c}(y)| \inf_{y \in A_{R}} \left( \frac{1}{w_{j}(y)} \right)^{1/p}.$$

We conclude that

$$||u(\cdot)w_{c}(\cdot)||_{L^{\infty}} \leq ||u(\cdot)w_{c}(\cdot)||_{L^{\infty}(B_{R}(0))} + ||u(\cdot)w_{c}(\cdot)||_{L^{\infty}(B_{R}(0)^{c})}$$

$$\leq ||u(\cdot)w_{c}(\cdot)||_{L^{\infty}(B_{R}(0))} + \sup_{n \in \mathbb{N}} ||u(\cdot)w_{c}(\cdot)||_{L^{\infty}(A_{Rn})} \leq C||u||_{W_{w}^{l,p}}.$$

The boundedness of u follows from the identical steps, with  $w_c = 1$ .

Proof of Lemma 30.

$$\|\cdot\|_{W_{w_1}^{m,p}} = \left(\sum_{0 \le j \le k} \int_0^\infty |D^{\alpha}u|^p (a_{1,j} + a_{2,j}x)^{\beta_j} dx\right)^{1/p}$$

$$= \left(\sum_{0 \le j \le k} \int_0^\infty |D^{\alpha}u|^p \frac{(a_{1,j} + a_{2,j}x)^{\beta_j}}{(b_{1,j} + b_{2,j}x)^{\beta_j}} (b_{1,j} + b_{2,j}x)^{\beta_j} dx\right)^{1/p}.$$

Noting that  $\min_{j} \min \left\{ \left( \frac{a_{1,j}}{b_{1,j}} \right)^{\beta_{j}}, \left( \frac{a_{2,j}}{b_{2,j}} \right)^{\beta_{j}} \right\} \leq \frac{(a_{1,j} + a_{2,j} x)^{\beta_{j}}}{(b_{1,j} + b_{2,j} x)^{\beta_{j}}} \leq \max_{j} \max \left\{ \left( \frac{a_{1,j}}{b_{1,j}} \right)^{\beta_{j}}, \left( \frac{a_{2,j}}{b_{2,j}} \right)^{\beta_{j}} \right\},$  yields the result.

For the second case

$$\|\cdot\|_{W^{m,p}_{\frac{1}{w_1}}} = \left(\sum_{0 \le j \le k} \int_0^\infty |D^{\alpha} u|^p \frac{1}{(a_{1,j} + a_{2,j}x)^{\beta_j}} dx\right)^{1/p}$$

$$= \left(\sum_{0 \le j \le k} \int_0^\infty |D^{\alpha} u|^p \frac{(b_{1,j} + b_{2,j}x)^{\beta_j}}{(a_{1,j} + a_{2,j}x)^{\beta_j}} \frac{1}{(b_{1,j} + b_{2,j}x)^{\beta_j}} dx\right)^{1/p}.$$

Again, since  $\min_{j} \min \left\{ \left( \frac{b_{1,j}}{a_{1,j}} \right)^{\beta_{j}}, \left( \frac{b_{2,j}}{a_{2,j}} \right)^{\beta_{j}} \right\} \leq \frac{(b_{1,j} + b_{2,j} x)^{\beta_{j}}}{(a_{1,j} + a_{2,j} x)^{\beta_{j}}} \leq \max_{j} \max \left\{ \left( \frac{b_{1,j}}{a_{1,j}} \right)^{\beta_{j}}, \left( \frac{b_{2,j}}{a_{2,j}} \right)^{\beta_{j}} \right\},$  we obtain the result.

Proof of Lemma 32. For  $\beta > 0$  and  $x \in [R, 2R]$ ,

$$3^{-\beta}(1+x)^{\beta} \le 3^{-\beta}(R+2R)^{\beta} \le R^{\beta} \le (1+x)^{\beta}.$$

$$\frac{1}{(1+x)^{\beta}} \le \frac{1}{(1+R)^{\beta}} \le \frac{1}{R^{\beta}} \le \frac{1}{(\frac{1}{3} + \frac{2R}{3})^{\beta}} \le \frac{3^{\beta}}{(1+x)^{\beta}}.$$

Let  $C_{j,p} = \begin{cases} 1 \text{ if } jp-1 > 0 \\ 3 \text{ if } jp-1 < 0. \end{cases}$  . On the annuli  $A_R$ , we have

$$\left(\sum_{j=0}^{l} R^{jp-1} \|D^{j}u\|_{L^{p}(A_{R})}^{p}\right)^{1/p} = \left(\sum_{j=0}^{l} R^{jp-1} \int_{R}^{2R} |D^{j}u(x)|^{p} dx\right)^{1/p} \\
\leq \left(\sum_{j=0}^{l} C_{j,p,d} \int_{R}^{2R} |D^{j}u(x)|^{p} (1+x)^{jp-1} dx\right)^{1/p} \\
\leq \widetilde{C} \left(\sum_{j=0}^{l} \int_{R}^{2R} |D^{j}u(x)|^{p} (1+x)^{jp-1} dx\right)^{1/p} .$$

Let 
$$c_{j,p} = \begin{cases} \frac{1}{3} & \text{if } jp - 1 > 0\\ 1 & \text{if } jp - 1 < 0, \end{cases}$$

$$\left(\sum_{j=0}^{l} R^{jp-d} \|D^{j}u\|_{L^{p}(A_{R})}^{p}\right)^{1/p} = \left(\sum_{j=0}^{l} R^{jp-1} \int_{R}^{2R} |D^{j}u(x)|^{p} dx\right)^{1/p}$$

$$\geq \left(\sum_{j=0}^{l} c_{j,p}^{jp-1} \int_{R}^{2R} |D^{j}u(x)|^{p} (1+x)^{jp-1} dx\right)^{1/p}$$

$$\geq \widetilde{c} \left(\sum_{j=0}^{l} \int_{R}^{2R} |D^{j}u(x)|^{p} (1+x)^{jp-1} dx\right)^{1/p}.$$

# 5.2. Auxiliary Lemmata.

**Lemma 84.** Let 'U be a Banach space and 'V a separable Banach space. We denote their dual spaces by U, V. Further assume that the embeddings  $U \hookrightarrow V \hookrightarrow V \hookrightarrow U$  are continuous and  $U \subset V$  is dense. Let  $B_r^{\text{weak}^*}$  denote the ball of radius r in V, equipped with the weak-\* topology. Assume that the following conditions are satisfied,

- (1)  $u_n \to u$  in C([0,T];U),
- (2)  $\sup_{t \in [0,T]} ||u_n||_V \le r$ .

Then  $u, u_n \in C([0,T]; B_r^{\text{weak}^*})$  and  $u_n \to u$  in  $C([0,T]; B_r^{\text{weak}^*})$  as  $n \to \infty$ .

*Proof.* (1) We claim that

$$u_n \to u$$
 in  $C\left([0,T]; B_r^{\text{weak}^*}\right)$  as  $n \to \infty$ 

i.e. that for all  $\phi \in V$ 

$$\lim_{n \to \infty} \sup_{s \in [0,T]} \left| \langle u_n(s) - u(s), \phi \rangle_{V \times V} \right| = 0.$$

To verify this claim, we fix  $\phi \in V$  and  $\varepsilon > 0$ . Since U is dense in V, there exists  $\phi_{\varepsilon} \in U$  such that  $|\phi - \phi_{\varepsilon}|_{V} \leq \varepsilon$ . Using (2), we infer that for all  $s \in [0, T]$  the following estimates hold

$$\begin{split} \left| \langle u_n(s) - u(s), \phi \rangle_{V \times'V} \right| &\leq \left| \langle u_n(s) - u(s), \phi - \phi_\varepsilon \rangle_{V \times'V} \right| + \left| \langle u_n(s) - u(s), \phi_\varepsilon \rangle_{U \times'U} \right| \\ &\leq \left\| u_n(s) - u(s) \right\|_V \left\| \phi - \phi_\varepsilon \right\|_{V} + \left| \langle u_n(s) - u(s), \phi_\varepsilon \rangle_{U \times'U} \right| \\ &\leq \varepsilon \cdot \left\| u_n - u \right\|_{L^{\infty}(0,T;V)} + \left| \langle u_n(s) - u(s), \phi_\varepsilon \rangle_{U \times'U} \right| \\ &\leq 2\varepsilon \cdot \sup_{n \in \mathbb{N}} \left\| u_n \right\|_{L^{\infty}(0,T;V)} + \left| \langle u_n(s) - u(s), \phi_\varepsilon \rangle_{U \times'U} \right| \\ &\leq 2\varepsilon r + \sup_{s \in [0,T]} \left| \langle u_n(s) - u(s), \phi_\varepsilon \rangle_{U \times'U} \right| \\ &\leq 2\varepsilon r + \sup_{s \in [0,T]} \left\| u_n(s) - u(s) \right\|_U \left\| \phi_\varepsilon \right\|_{U}. \end{split}$$

Passing to the limit  $n \to \infty$ , we obtain

$$\limsup_{n \to \infty} \sup_{s \in [0,T]} \left| \langle u_n(s) - u(s), \phi \rangle_{V \times V} \right| \le 2r\varepsilon.$$

Since  $\varepsilon$  is arbitrary,

$$\lim_{n \to \infty} \sup_{s \in [0,T]} \left| \langle u_n(s) - u(s), \phi \rangle_{V \times V} \right| = 0,$$

for every  $\phi \in V$ . Since  $C([0,T]; B_r^{\text{weak}^*})$  is a complete metric space, we infer that  $u \in C([0,T]; B_r^{\text{weak}^*})$  as well. This completes the proof.

#### References

- [1] E. Abi Jaber. Lifting the Heston model. Quantitative finance, 19(12):1995–2013, 2019.
- [2] E. Abi Jaber. Weak existence and uniqueness for affine stochastic Volterra equations with L1-kernels. Bernoulli, 27(3):1583–1615, 2021.
- [3] E. Abi Jaber, C. Cuchiero, M. Larsson, and S. Pulido. A weak solution theory for stochastic Volterra equations of convolution type. *The Annals of Applied Probability*, 31(6):2924–2952, 2021.
- [4] E. Abi Jaber and O. El Euch. Markovian structure of the Volterra Heston model. Statistics & Probability Letters, 149:63–72, 2019.
- [5] E. Abi Jaber and O. El Euch. Multifactor approximation of rough volatility models. SIAM journal on financial mathematics, 10(2):309–349, 2019.
- [6] E. Abi Jaber, M. Larsson, and S. Pulido. Affine Volterra processes. The Annals of Applied Probability, 29(5):3155–3200, 2019.
- [7] E. Abi Jaber, E. Miller, and H. Pham. Linear-Quadratic control for a class of stochastic Volterra equations: solvability and approximation. *The Annals of Applied Probability*, 31(5):2244–2274, 2021.
- [8] R. A. Adams and J. J. Fournier. Sobolev spaces. Elsevier, 2003.

- [9] S. Albeverio, L. Gawarecki, V. Mandrekar, B. Rüdiger, and B. Sarkar. Itô formula for mild solutions of SPDEs with Gaussian and non-Gaussian noise and applications to stability properties. *Random Operators* and Stochastic Equations, 25(2):79–105, 2017.
- [10] A. Alfonsi and G. Szulda. On non-negative solutions of stochastic volterra equations with jumps and non-lipschitz coefficients. arXiv preprint arXiv:2402.19203, 2024.
- [11] O. E. Barndorff-Nielsen, F. E. Benth, and A. E. Veraart. Modelling energy spot prices by volatility modulated Lévy-driven Volterra processes. *Bernoulli*, 19(3):803–845, 2013.
- [12] O. E. Barndorff-Nielsen and J. Schmiegel. Time change, volatility, and turbulence. In *Mathematical control theory and finance*, pages 29–53. Springer, 2008.
- [13] V. Benci and D. Fortunato. Weighted Sobolev spaces and the nonlinear Dirichlet problem in unbounded domains. Annali di Matematica Pura ed Applicata, 121(1):319–336, 1979.
- [14] A. Bensoussan. Stochastic Navier-stokes Equations. Acta Applicandae Mathematicae, 38(3):267–304, 1995.
- [15] F. E. Benth, N. Detering, and P. Kruehner. Stochastic Volterra integral equations and a class of first-order stochastic partial differential equations. Stochastics, 94(7):1054–1076, 2022.
- [16] M. A. Berger and V. J. Mizel. Volterra equations with Itô integrals—I. The Journal of Integral Equations, pages 187–245, 1980.
- [17] M. A. Berger and V. J. Mizel. Volterra equations with itô integrals—II. The Journal of Integral Equations, pages 319–337, 1980.
- [18] C. Blundell, J. Beck, and K. A. Heller. Modelling reciprocating relationships with Hawkes processes. Advances in neural information processing systems, 25, 2012.
- [19] A. Bondi, G. Livieri, and S. Pulido. Affine Volterra processes with jumps. Stochastic Processes and their Applications, 168:104264, 2024.
- [20] O. Bonesini, A. Jacquier, and A. Pannier. Rough volatility, path-dependent PDEs and weak rates of convergence. arXiv preprint arXiv:2304.03042, 2023.
- [21] D. Breit, E. Feireisl, and M. Hofmanova. Stochastically Forced Compressible Fluid Flows. Walter de Gruyter, 2018.
- [22] Z. Brzeźniak and E. Motyl. Existence of a martingale solution of the stochastic Navier–Stokes equations in unbounded 2D and 3D domains. *Journal of Differential Equations*, 254(4):1627–1685, 2013.
- [23] Z. Brzeźniak and M. Ondreját. Stochastic geometric wave equations with values in compact Riemannian homogeneous spaces. *The Annals of Probability*, 41(3B):1938–1977, 2013.
- [24] Z. Brzeźniak, M. Ondreját, and J. Seidler. Invariant measures for stochastic nonlinear beam and wave equations. Journal of Differential Equations, 260(5):4157–4179, 2016.
- [25] Z. Brzeźniak, E. Hausenblas, and P. Razafimandimby. Stochastic Reaction-diffusion Equations Driven by Jump Processes. Potential analysis, 49(2):131–201, Sept. 2010.
- [26] L. Coutin and P. Carmona. Fractional Brownian motion and the Markov property. Electronic Communications in Probability, 3:12, 1998.
- [27] L. Coutin and L. Decreusefond. Stochastic Volterra equations with singular kernels. In *Stochastic analysis* and mathematical physics, pages 39–50. Springer, 2001.
- [28] S. Cox, A. Jentzen, R. Kurniawan, and P. Pušnik. On the mild Itô formula in Banach spaces. arXiv preprint arXiv:1612.03210, 2016.
- [29] R. Crane and D. Sornette. Robust dynamic classes revealed by measuring the response function of a social system. *Proceedings of the National Academy of Sciences*, 105(41):15649–15653, 2008.
- [30] C. Cuchiero and J. Teichmann. Markovian lifts of positive semidefinite affine Volterra-type processes. Decisions in Economics and Finance, 42:407–448, 2019.
- [31] C. Cuchiero and J. Teichmann. Generalized Feller processes and Markovian lifts of stochastic Volterra processes: the affine case. *Journal of evolution equations*, 20(4):1301–1348, 2020.
- [32] G. Da Prato, A. Jentzen, and M. Röckner. A mild Itô formula for SPDEs. *Transactions of the American Mathematical Society*, 372(6):3755–3807, 2019.
- [33] A. Debussche, N. Glatt-Holtz, and R. Temam. Local martingale and pathwise solutions for an abstract fluids model. *Physica D: Nonlinear Phenomena*, 240(14-15):1123–1144, 2011.
- [34] P. Dörsek and J. Teichmann. A semigroup point of view on splitting schemes for stochastic (partial) differential equations. arXiv preprint arXiv:1011.2651, 2010.
- [35] C. Duval, E. Luçon, and C. Pouzat. Interacting Hawkes processes with multiplicative inhibition. *Stochastic Processes and their Applications*, 148:180–226, 2022.

- [36] O. El Euch and M. Rosenbaum. The characteristic function of rough Heston models. Mathematical Finance, 29(1):3–38, 2019.
- [37] K.-J. Engel, R. Nagel, and S. Brendle. One-parameter semigroups for linear evolution equations, volume 194. Springer, 2000.
- [38] S. N. Ethier and T. G. Kurtz. Markov processes: characterization and convergence. John Wiley & Sons, 2009.
- [39] M. Friesen and P. Jin. Volterra square-root process: Stationarity and regularity of the law. The Annals of Applied Probability, 34(1A):318–356, 2024.
- [40] J. Gatheral, T. Jaisson, and M. Rosenbaum. Volatility is rough. In Commodities, pages 659–690. Chapman and Hall/CRC, 2022.
- [41] B. Goldys, M. Röckner, and X. Zhang. Martingale solutions and Markov selections for stochastic partial differential equations. Stochastic Processes and their Applications, 119(5):1725–1764, 2009.
- [42] G. Gripenberg. On the resolvents of Volterra equations with nonincreasing kernels. *Journal of Mathematical Analysis and Applications*, 76(1):134–145, 1980.
- [43] Y. Hamaguchi. Markovian lifting and asymptotic log-Harnack inequality for stochastic Volterra integral equations. arXiv preprint arXiv:2304.06683, 2023.
- [44] Y. Hamaguchi. Weak well-posedness of stochastic Volterra equations with completely monotone kernels and non-degenerate noise. arXiv preprint arXiv:2310.16030, 2023.
- [45] P. Harms and D. Stefanovits. Affine representations of fractional processes with applications in mathematical finance. Stochastic Processes and their Applications, 129(4):1185–1228, 2019.
- [46] E. Hausenblas and J. Seidler. A note on maximal inequality for stochastic convolutions. Czechoslovak Mathematical Journal, 51(4):785–790, 2001.
- [47] M. Hofmanová and J. Seidler. On weak solutions of stochastic differential equations. Stochastic analysis and applications, 30(1):100–121, 2012.
- [48] E. A. Jaber, C. Cuchiero, L. Pelizzari, S. Pulido, and S. Svaluto-Ferro. Polynomial Volterra processes. arXiv preprint arXiv:2403.14251, 2024.
- [49] A. Jacquier, A. Pannier, and K. Spiliopoulos. On the large-time behaviour of affine Volterra processes. arXiv preprint arXiv:2204.05270, 2022.
- [50] A. Jakubowski. The almost sure Skorokhod representation for subsequences in nonmetric spaces. Theory Probab. Appl., 42(1):209–216, 1997.
- [51] A. Kufner and B. Opic. How to define reasonably weighted Sobolev spaces. Commentationes Mathematicae Universitatis Carolinae, 25(3):537–554, 1984.
- [52] A. Lodhia, S. Sheffield, X. Sun, and S. Watson. Fractional gaussian fields: A survey. Probability Surveys, 13, 2016.
- [53] R. MacCamy. Approximations for a class of functional differential equations. SIAM Journal on Applied Mathematics, 23(1):70–83, 1972.
- [54] B. Maslowski and J. Seidler. On sequentially weakly Feller solutions to SPDE's. Atti della Accademia Nazionale dei Lincei. Classe di Scienze Fisiche, Matematiche e Naturali. Rendiconti Lincei. Matematica e Applicazioni, 10(2):69-78, 1999.
- [55] M. Meyries and M. Veraar. Characterization of a class of embeddings for function spaces with Muckenhoupt weights. *Archiv der Mathematik*, 103(5):435–449, 2014.
- [56] L. Mytnik. Weak uniqueness for the heat equation with noise. Annals of probability, pages 968–984, 1998.
- [57] L. Mytnik and T. S. Salisbury. Uniqueness for Volterra-type stochastic integral equations. arXiv preprint arXiv:1502.05513, 2015.
- [58] W. Noll, B. D. Coleman, and W. Noll. An approximation theorem for functionals, with applications in continuum mechanics. Springer, 1974.
- [59] D. J. Prömel and D. Scheffels. Pathwise uniqueness for singular stochastic Volterra equations with Hölder coefficients. arXiv preprint arXiv:2212.08029, 2022.
- [60] D. J. Prömel and D. Scheffels. On the existence of weak solutions to stochastic Volterra equations. Electronic Communications in Probability, 28:1–12, 2023.
- [61] D. J. Prömel and D. Scheffels. Stochastic Volterra equations with Hölder diffusion coefficients. Stochastic Processes and their Applications, 161:291–315, 2023.
- [62] P. Reynaud-Bouret and S. Schbath. Adaptive estimation for Hawkes processes; application to genome analysis. Annals of Statistics, 38(5):2781–2822, 2010.

- [63] R. L. Schilling, R. Song, and Z. Vondraček. Bernstein functions: theory and applications. Walter de Gruyter, 2009.
- [64] J. Simon. Compact sets in the space Lp (0, T; B). Annali di Matematica pura ed applicata, 146:65–96, 1986.
- [65] A. Verma, S. G. Jena, D. R. Isakov, K. Aoki, J. E. Toettcher, and B. E. Engelhardt. A self-exciting point process to study multicellular spatial signaling patterns. *Proceedings of the National Academy of Sciences*, 118(32):e2026123118, 2021.
- [66] F. Viens and J. Zhang. A martingale approach for fractional Brownian motions and related path dependent PDEs. *The Annals of Applied Probability*, 29(6):3489–3540, 2019.
- [67] Z. Wang. Existence and uniqueness of solutions to stochastic Volterra equations with singular kernels and non-Lipschitz coefficients. Statistics & probability letters, 78(9):1062–1071, 2008.
- [68] X. Zhang. Stochastic Volterra equations in Banach spaces and stochastic partial differential equation. Journal of Functional Analysis, 258(4):1361–1425, 2010.

ÉCOLE POLYTECHNIQUE FÉDÉRALE DE LAUSANNE, SWITZERLAND *Email address*: florian.huber@epfl.ch