GROMOV'S ELLIPTICITY OF PRINCIPAL \mathbb{G}_m -BUNDLES

SH. KALIMAN

ABSTRACT. We prove that every nontrivial principal \mathbb{G}_m -bundle over a complete uniformly rational variety is algebraically elliptic in the sense of Gromov.

1. INTRODUCTION

All varieties in this paper are smooth algebraic varieties defined over an algebraically closed field k of characteristic zero. This work is continuation of [\[AKZ23\]](#page-20-0) and [\[KZ23b\]](#page-21-0) and, thus, we adhere to the terminology and notations of these papers (see also [\[For17\]](#page-21-1) for the general theory of Oka manifolds and [\[For23\]](#page-21-2) for recent advances in this area). In particular, when we talk about Gromov's ellipticity (or, ellipticity for short) we always mean algebraic ellipticity leaving the holomorphic case aside. This enables us to use the following advantage - the notions of algebraic Gromov's ellipticity and subellipticity coincide [\[KZ23a\]](#page-21-3), while in the holomorphic case it is still an open problem. Recall that an algebraic variety is called uniformly rational if every point in it has a Zariski open neighborhood isomorphic to an open subset of \mathbb{A}^n and it is still unknown whether a smooth complete rational variety is uniformly rational (see, $[Gro89, 3.5.E''']$, $[BB14,$ Question 1.1] and [\[CPPZ21,](#page-21-6) p. 41]). The main results of [\[AKZ23,](#page-20-0) Theorem 3.3] are the following.

Theorem 1.1. Every complete uniformly rational variety is Gromov's elliptic.

Theorem 1.2. Let X be a complete uniformly rational variety and $\mathcal{D} \to X$ be an ample or anti-ample line bundle on X with zero section $Z_{\mathcal{D}}$. Then $\mathcal{D} \setminus Z_{\mathcal{D}}$ is elliptic.

However, the assumption that $\mathcal{D} \to X$ is ample or anti-ample is not necessary (e.g., see [\[AKZ23,](#page-20-0) Theorem 3.8]). What is necessary, of course, is the requirement that $\mathcal{D} \to X$ is a nontrivial line bundle. This leads to the following.

Question 1. Let X be a complete uniformly rational variety and $\mathcal{D} \to X$ be a nontrivial line bundle on X with zero section $Z_{\mathcal{D}}$. Is $\mathcal{D} \setminus Z_{\mathcal{D}}$ elliptic?

In this paper we show that the answer to this question is positive (Theorem 6.1). Let us briefly discuss our approach in the crucial case when X is a smooth complete rational surface (i.e., X is Gromov's elliptic by Theorem [1.1\)](#page-0-0). Given $v \in X$ we look for an open set $B \subset \mathbb{A}^1$ and a birational morphism $\varphi : B \times \mathbb{P}^1 \to X$ locally invertible at v. In particular, v is contained in $C = \varphi(b_0 \times \mathbb{P}^1)$ for some $b_0 \in B$. It follows from

Key words and phrases. Gromov ellipticity, spray, uniformly rational variety, complete surface. 2020 Mathematics Subject Classification Primary 14J60, 14M25, 14M27; Secondary 32Q56.

[\[KZ23b,](#page-21-0) Corollary 3.8] and [\[AKZ23,](#page-20-0) Corollary 2.7] that Y admits a family of rank 1 sprays dominating on the fiber over v if the restriction of $\mathcal D$ yields a nontrivial line bundle on C. If this is true for every $v \in X$, then one has subellipticity of Y. Thus, since subellipticity implies ellipticity [\[KZ23a\]](#page-21-3) we only need to check that $\mathcal{D}|_C$ is not trivial. Letting $\mathcal{D} = \mathcal{O}(D)$ for some SNC divisor D we observe that $\mathcal{D}|_C$ is not trivial if C meets only one irreducible component H of D (where $C \neq H$). Hence, our aim is to find such φ, D and H for any given $v \in X$. The realization of this plan is executed in Sections [3](#page-2-0)[-5](#page-8-0) (after preliminaries in Section [2\)](#page-1-0) which concludes the case of surfaces. Using induction by dimension we show in Section [6](#page-13-1) that the same idea works in the general case.

2. Preliminaries

The following basic definitions can be found in $\left[\frac{\text{Gro89}}{\text{Gro89}}\right]$ and $\left[\frac{\text{For17}}{\text{Gro89}}\right]$, Definition 6.1.1.

Definition 2.1. A spray of rank r over a smooth algebraic variety X is a triple (E, p, s) consisting of a vector bundle $p: E \to X$ of rank r and a morphism $s: E \to X$ such that $s|_Z = p|_Z$ where $Z \subset E$ stands for the zero section of p. This spray is *dominating* at $x \in X$ if the restriction $s|_{E_x}: E_x \to X$ to the fiber $E_x = p^{-1}(x)$ is dominant at the origin 0_x of the vector space E_x . The variety X is called *elliptic* if it admits a spray (E, p, s) which is dominating at each point $x \in X$. The variety X is called *subelliptic* if it admits a family of sprays (E_i, p_i, s_i) defined over X which is dominating at each point $x \in X$, that is,

$$
T_x X = \sum_{i=1}^n ds_i (T_{0_{i,x}} E_{i,x}) \quad \forall x \in X.
$$

Recall the following $[AKZ23,$ Definition 2.5] (see, also $[KZ23b,$ Definition 2.7]).

Definition 2.2. We say that a complete rational curve C on a smooth variety X verifies the *strengthened two-orbit property* at a smooth point $v \in C$ if there exists a pair of rank 1 sprays (E_i, p_i, s_i) $(i = 1, 2)$ on X such that C is covered by the onedimensional s_i -orbits $O_{i,v}$, where $s_i: p_i^{-1}(v) \to O_{i,v}$ is a birational morphism étale over v and (E_i, p_i, s_i) restricts to a spray on $O_{i,v}$ dominating at v.

If for any $v \in X$ there exists a curve $C = C_v$ as above, then we say that X verifies the strengthened curve-orbit property.

The reason why we need the curve-orbit property is the following [\[AKZ23,](#page-20-0) Corollary 2.7].

Proposition 2.3. Let X be a smooth variety, $\rho : \mathcal{D} \to X$ be a line bundle and Y = $\mathcal{D}\setminus Z_{\mathcal{D}}$. Suppose that

(i) X is elliptic and verifies the strengthened curve-orbit property with respect to a covering family F of projective rational curves $\{C_v\}$ on X that are smooth at v, and

(ii) $\rho: \mathcal{D} \to X$ restricts to a nontrivial line bundle on each member C_v of \mathcal{F} . Then Y is elliptic.

The first step in the proof of Proposition [2.3](#page-1-1) is to establish subellipticity using the strengthened curve-orbit property and then ellipticity follows by [\[KZ23a\]](#page-21-3).

Proposition 2.4. Let X be a smooth variety of dimension n and B be a smooth variety of dimension $n-1$. Assume that X admits a birational morphism $\varphi: B \times \mathbb{P}^1 \to X$ locally invertible over a neighborhood of a point $v \in X$. Let $C = \varphi(b \times \mathbb{P}^1)$, $b \in B$ be the curve containing v. Then C verifies the strengthened two-orbit property at v.

Proof. Let $v = \varphi(b, u_0)$ where $u_0 \in \mathbb{P}^1$. By [\[AKZ23,](#page-20-0) Lemma 2.4] for every point $u \in \mathbb{P}^1 \setminus \{u_0\}$ there is a rank 1 spray (E_u, p_u, s_u) on X whose restriction to C is also a spray such that $s_u|_{p_u^{-1}(v)} : p_u^{-1}(v) \to C$ is a birational morphism étale over v and the s_u -orbit is equal to $\varphi(b \times (\mathbb{P}^1 \setminus \{u\}))$. This yields the desired conclusion.

Propositions [2.3](#page-1-1) and [2.4](#page-2-1) imply now the following.

Corollary 2.5. Let X be a smooth complete elliptic variety, $\rho : \mathcal{D} \to X$ be a line bundle with the zero section Z, and $Y = \mathcal{D} \setminus Z_{\mathcal{D}}$. Suppose that for every $v \in X$ there is a morphism $\varphi_v : B_v \times \mathbb{P}^1 \to X$ as in Proposition [2.4](#page-2-1) and C_v be the curve $\varphi_v(b_v \times \mathbb{P}^1)$ containing v. Let the restriction of D to every curve C_v , $v \in X$ be a nontrivial line bundle. Then Y is elliptic.

3. CONTRACTIBLE GRAPHS

Let us recall first terminology and some standard facts about dual weighted graphs of curves. Let G be a simple normal crossing (SNC) curve in a smooth surface X , i.e., all irreducible components of G are smooth and each of its singularities is a point where exactly two components meet transversely. The *dual graph* of G is the graph whose vertices are the irreducible components of G and the edges between the vertices are the intersection points of these components. Assuming that G is complete one can consider the weighted dual graph Γ of G that is the dual graph of G with each vertex C equipped with the weight equal to the selfintersection number $C \cdot C$ of this component C in X. A vertex C is linear (resp. endpoint) if it has two (resp. one) neighbors in Γ . If C has at least three neighbors it is called a branch point. A non-circular connected graph without branch points will be called *linear*. Suppose that π : $\hat{X} \to X$ is the blowing up of a point $p \in G$ and $G' = \pi^{-1}(G)$. Then G' is again an SNC curve. If p is the intersection of two components of G with weights w_1 and w_2 , then the graph of G' is obtained from Γ by the following change

❝ w1 ❝ w2 =⇒ ❝ w¹ − 1 ❝ −1 ❝ w² − 1

,

while the rest of the dual graph remains the same (such change is called an *inner blowing up*). If p is a smooth point of G located in a component with weight w, then the dual graph of G' is obtained from Γ by the following change

The latter change is called an *outer blowing up* and unlike the inner blowing up it does not describe the curve G' uniquely since one needs to indicate the point of w at which the blowing up occurs. The contraction of a (-1) -component E in G also preserves SNC type, provided that E is not a branch point. In this case the image of G has the dual graph obtained from Γ by reversion of the above diagrams. A dual graph is called contractible if such contractions can reduce it to an empty graph.

Notation 3.1. From now on by $\mathbb{A}_{u_1,...,u_n}^n$ we denote an affine space \mathbb{A}^n equipped with a coordinate system (u_1, \ldots, u_n) .

Lemma 3.2. Let Γ be a contractible linear graph with a unique (-1)-vertex E and C be a neighbor of E in Γ . Then Γ is the dual graph of the minimal resolution $\pi: W \to \mathbb{A}^2_{x,y}$ of the indeterminacy points of a rational function y^m/x^k where k and $m \geq 1$ are relatively prime. Furthermore, there are relatively prime l and n and a neighborhood $U \simeq \mathbb{A}^2_{u,v}$ of $E \cap C$ in W such that $E \cap U$ and $C \cap U$ are coordinate axes and (u, v) coincides with $(y^m/x^k, x^n/y^l)$ or $(x^k/y^m, y^l/x^n)$.

Proof. We use induction on the number of vertices. Note that every weighted linear contractible graph with E as an endpoint must be of the form

This is exactly the resolution graph of indeterminacies of y^m/x^k when $m = 1$. Note that each point $C_i \cap C_{i+1}$ has a neighborhood in W isomorphic to \mathbb{A}^2 on which the functions x^{k+1-i}/y and y/x^{k-i} yield a coordinate system. For $k, m > 1$ denote now by C_1 and C_2 the neighbors of E. Denote their proper transforms in the image W' of W after the contraction of E by C'_1 and C'_2 and the new dual graph by Γ'. Note that one of them (say, C'_1) $'_{1}$) is a (-1)-vertex, while the weight of C'_{2} y_2 is at most -2 to guarantee further contraction. By the induction assumption one can suppose that Γ' is the minimal resolution $W' \to \mathbb{A}^2$ of the indeterminacy points of a function y^j/x^i . Furthermore, there is a neighborhood of the point $C_1' \cap C_2'$ y_2' in W' isomorphic to \mathbb{A}^2 equipped with the coordinate system $(u, v) = (y^j/x^i, x^n/y^l)$ (resp. $(u, v) = (x^i/y^j, y^l/x^n)$) with C'_1 given locally by $v = 0$ and the matrix

$$
\begin{bmatrix} j & l \\ i & n \end{bmatrix}
$$

having the determinant ± 1 (this assumption is obvious in the case $m = 1$). The blowing up of W' at $C'_1 \cap C'_2$ produces E and resolves indeterminacy of $u/v = y^{j+l}/x^{i+n}$. Note that the assumption on the matrix in Formula [\(1\)](#page-3-0) implies that $m = j + l$ and $k = i + n$ are relatively prime and the determinants of the matrices

$$
\begin{bmatrix}\nj+l & l \\
i+n & n\n\end{bmatrix}\n\text{ and }\n\begin{bmatrix}\nj+l & j \\
i+n & i\n\end{bmatrix}
$$

are ± 1 . Note also that $(u, v/u)$ (resp. $(v, u/v)$) is a local coordinate system in a neighborhood of $C_1 \cap E$ (resp. $C_2 \cap E$). This concludes the proof.

Remark 3.3. Let $1 \leq m \leq k$. Note that one of the matrices in Formula [\(2\)](#page-4-0) has determinant 1. Thus, one can always find a neighbor C of E in Γ and natural n and l such that $kl - mn = 1$ and Γ is obtained by inner blowing ups from the dual graph Γ_0 of the minimal resolution of the indeterminacy points of the function y^l/x^n with the proper transform of C being the only (-1) -vertex in Γ_0 . Actually, the same claim remains true in the case of $m = 1$ and $k \ge 1$ if one let $n = k - 1$ and $l = 1$ where in the case of $k = m = 1$ one has to use the proper transform of the y-axis instead of C.

Lemma 3.4. Let the notation of Remark [3.3](#page-4-1) hold. Then the functions x^k/y^m and y^{l}/x^{n} form a coordinate system in a neighborhood of the point $C \cap E$ with C and E being the coordinate axes.

Proof. As we stated in Lemma [3.2](#page-3-1) either (a) the functions x^k/y^m and y^l/x^n or (b) the functions y^m/x^k and x^n/y^l form a coordinate system in a neighborhood of $C \cap E$ with C and E being the coordinate axes. However, $(y^m/x^k)^l (x^n/y^l)^m = x^{-1}$. Since the lift of x to a neighborhood of $C \cap E$ is regular we see that (b) is impossible which yields the desired conclusion.

Lemma 3.5. Let Γ be a connected contractible (not necessarily linear) weighted graph such that it has only one (-1) -vertex E. Then Γ contains n–1 branch points E_1, \ldots, E_{n-1} such that

(1) each of them has exactly three neighbors;

(2) every E_i is contained in a linear subgraph Γ_i of Γ such that after changing the weight of E_i to -1 this subgraph becomes a weighted graph from Lemma [3.2](#page-3-1) with $m, k \geq$ 2;

(3) every E_i has a neighbor which is an endpoint of Γ_{i+1} where Γ_n is as in Lemma [3.2](#page-3-1) (in particular, it contains E).

Proof. We use induction by n. If $n = 1$, then Γ is linear and the conclusion follows from Lemma [3.2.](#page-3-1) Assume that the statement is true for $n-1$ and we want to prove it for n. To get Γ contracted one needs to contract Γ_n since otherwise the branch point E_{n-1} prevents further contraction. By Lemma [3.2](#page-3-1) Γ_n is of the desired form. Contracting Γ_n one changes the weight of E_{n-1} and this weight must become -1 to proceed with contraction. Hence, the induction assumption yields the desired conclusion. \Box

Proposition 3.6. Let $\psi: X \to X_0$ be a proper birational morphism of smooth complete rational surfaces and $H \subset X$ be a smooth irreducible curve such that $q_0 = \psi(H)$ is a singleton. Then $\psi = \mu \circ \theta$ where $\theta : X \to \tilde{X}$ is a morphism into a smooth complete rational surface \check{X} and $\mu : \check{X} \to X_0$ is a morphism such that the restriction of μ over $\breve{X} \setminus q_0$ is an isomorphism and the dual graph Γ of $G = \mu^{-1}(q_0)$ is of the same form

as in Lemma [3.5](#page-4-2) with $H = \theta(H)$ being the only (-1)-vertex. Furthermore, μ can be presented as a composition of morphisms $\mu_{i+1} : X'_{i+1} \to X'_{i}, i = 0, \ldots, n-1$ with $\breve{X} = X'_n$ and $X'_0 = X_0$ such that each μ_i has a connected exceptional divisor whose dual graph is Γ_i .

Proof. Note that X is obtained by a sequence of blowing up of a finite set in X_0 and infinitely near points. Choose in this sequence the smallest subsequence R which generates a proper transform H of H in a surface X over X_0 . That is, ψ factors through a proper birational morphism $\mu : \check{X} \to X_0$ with a connected contractible exceptional divisor G which contains H as an irreducible component. Now the first statement is the consequence of the following.

Claim. The dual graph Γ of G is a contractible graph with \tilde{H} being a unique (-1)vertex. Indeed, every (-1) -vertex in Γ appears in this sequence R of blowing ups only after the proper transforms of its neighbors have been already created (otherwise, the weight of this vertex cannot be -1). This implies that Γ cannot contain linear (-1) -vertices except for \tilde{H} since otherwise R is not the smallest subsequence. Hence, Γ satisfies the assumptions of Lemma [3.5](#page-4-2) and we have the first statement.

The second statement is automatic for $n = 1$. Hence, in the general case contracting the components corresponding to the vertices of Γ_n in \check{X} we get the desired conclusion by induction. \Box

4. Local coordinate systems

The following fact is well-known and the author does not know to whom its original proof has to be attributed.

Lemma 4.1. Let X_0 be a smooth complete rational surface and $q_0 \in X_0$. Then there is a neighborhood of q_0 in X isomorphic to the affine plane.

Proof. Recall that a variety Y belongs to class \mathcal{A}_0 if every point in Y has a neighborhood isomorphic to an affine space (see, $[For 06, Definition 2.3]$). Any variety obtained by blowing up of Y at a point is still in class \mathcal{A}_0 [\[Gro89,](#page-21-4) Section 3.5D]. In particular, every smooth complete rational surface X belongs to class \mathcal{A}_0 since such surface is obtained from \mathbb{P}^2 or a Hirzebruch surface by consequent blowing up of points \square

Lemma 4.2. Let π : $X \to X_0$ be a proper morphism of smooth projective rational surfaces such that its restriction is an isomorphism over $X_0 \setminus q_0$ for some $q_0 \in X_0$ and the weighted dual graph Γ_1 of $\pi^{-1}(q_0)$ is the graph of the minimal resolution of indeterminacy point of a rational function of the form x^k/y^m where m and k are relatively prime. Let C_n be the vertex in Γ_1 that is the proper transform of the exceptional divisor \tilde{C} of the blowing up $\tau : \tilde{X} \to X_0$ of X_0 at q_0 . Let $U_0 \simeq \mathbb{A}^2_{x_0,y_0}$ be an open subset of X_0 with the origin at q_0 and H be the proper transform in X of the x_0 -axis $H_0 \subset U_0$. Suppose that the graph Γ_1^+ of $\pi^{-1}(H_0)$ is of the form

$$
H \quad C_1 \quad C_2 \quad C_n
$$

where C_1, \ldots, C_n are the irreducible components of $\pi^{-1}(q_0)$. Then π coincides with the minimal resolution of indeterminacy of x_0^k/y_0^m at q_0 and $m \leq k$.

Proof. The graph of $\tau^{-1}(H_0)$ is a linear graph consisting of \tilde{C} and the proper transform \tilde{H} of H_0 . To obtain Γ_1^+ one has to use only inner blowing ups in this graph starting with the blowing up of the point $H \cap C$. This implies that π is uniquely determined by Γ_1 . Hence, π coincides either with the minimal resolution of indeterminacy of y_0^m/x_0^k or of y_0^k/x_0^m where $m \leq k$. However, in the latter case H meets C_n , whereas it is supposed to meet C_1 . Thus, $n = 1$ which implies $m = k = 1$ and concludes the proof.

Definition 4.3. (1) For $c \in \mathbf{k}$ and (x_0, y_0) as in Lemma [4.2](#page-5-0) a local coordinate system of the form $(x_0 + c, y_0)$ will be called a *bottom coordinate system* for $\pi : X \to X_0$.

(2) By Lemma [3.4](#page-4-3) for some relatively prime n and l the lift of x_0^k/y_0^m and y_0^l/x_0^n to X yields a coordinate system (x_1, y_1) on an open subset $U_1 \simeq \mathbb{A}^2$ of X such that y_1 vanishes on the only (−1)-vertex of Γ_1 . We call $(x_1 + c, y_1)$ a top coordinate system for $\pi: X \to X_0.$

Remark 4.4. (i) There is some freedom in the choice of a bottom coordinate system. Indeed, if one replaced y_0 by $y_0 + ax_0^l$ where $a \in \mathbf{k}$ and l is sufficiently large, then the proper transform of the new x_0 -axis will still meet C_1 . Similarly, one can replace x_0 by $x_0 + by_0, b \in \mathbf{k}$. Hence, we can assume that every given finite subset M of $U_0 \setminus \{q_0\}$ does not meet the coordinate axis.

(ii) A neighborhood $U_0 \simeq \mathbb{A}^2_{x_0,y_0}$ of q_0 as in Lemma [4.2](#page-5-0) always exists whenever k and m are relatively prime. Indeed, a neighborhood U_0 isomorphic to an affine plane always exists by Lemma [4.1.](#page-5-1) Consider now a smooth curve $H' \subset X$ meeting C_1 transversely. Then the proper transform $H'_0 \subset X_0$ of H contains q_0 and it is smooth since contractions preserve SNC type of curves. Choosing a coordinate system (x_0, y_0) on U_0 such that H_0 is tangent to H'_0 at q_0 with sufficiently high order we see that the graph of $\pi^{-1}(H_0)$ has the desired form. Using the similar argument we can alway find a smooth affine line $F_0 \subset X_0$ such that the graph of $\pi^{-1}(F_0)$ is linear with the proper transform of F_0 meeting C_n .

Lemma 4.5. Let π : $X \to X_0$ be a proper morphism of smooth projective rational surfaces such that its restriction is an isomorphism over $X_0 \setminus q_0$ for some $q_0 \in X_0$ and the dual graph Γ_1 of $\pi^{-1}(q_0)$ is the graph of a minimal resolution of indeterminacy point of a rational function of the form x^k/y^m where k and m are relatively prime. Let $U_0 \simeq \mathbb{A}^2_{x_0,y_0}$ be an open subset of X_0 with the origin at q_0 such that the graph of $\pi^{-1}(H_0)$ is linear where H_0 is the x_0 -axis. Then π can be presented as a composition of morphisms $\nu_i: X_i \to X_{i-1}, i = 1, ..., N$ with $X = X_N$ such that

(i) the restriction of ν_i is an isomorphism over $X_{i-1} \setminus \{q_{i-1}\}\$ for some $q_{i-1} \in X_{i-1}$ and the exceptional divisor $\nu_i^{-1}(q_{i-1})$ contains only one (-1) -curve H_i ;

(ii) there is an open subset $U_i \simeq \mathbb{A}^2_{x_i,y_i}$ of X_i such that $q_i \in H_i$ and y_i vanishes on $H_i \cap U_i$ for $i < N$;

(iii) (x_i, y_i) is a bottom coordinate system for ν_{i+1} and a top coordinate system ν_i ; i.e., ν_{i+1} is the minimal resolution of indeterminacy points of a function of the form

 $(x_i - c_i)^{k_i}/y_i^{m_i}$ where c_i is the x_i -coordinate of q_i and, furthermore, $m_i = k_i = 1$ for $i \leq N-2$, while $m_{N-1} \leq k_{N-1}$;

(iv) H_N coincides with the only irreducible component E of $\pi^{-1}(q_0)$ with selfinter $section -1$.

Proof. Let C_1, \ldots, C_n be as in Lemma [4.2.](#page-5-0) If the proper transform H of H_0 meets C_1 , then the conclusion follows from Lemma [4.2](#page-5-0) with $\pi = \nu_1$. In particular, this is true when the length n of Γ_1 is 1, i.e., $C_1 = C_n$ and π is the blowing up of X_0 at q_0 . Hence, we suppose that H meets C_n and use induction on n. Let $\nu_1 : X_1 \to X_0$ be the blowing up of q_0 , i.e., $H_1 = \nu_1^{-1}(q_0)$ is the proper transform of C_n . Then (x_0, y_0) is the bottom coordinate system for ν_1 and $\pi = \nu_1 \circ \pi_1$ where the restriction of $\pi_1 : X \to X_1$ is an isomorphism over $X_1 \setminus \{q_1\}$ for some $q_1 \in H_1$. Note that we have a top coordinate system (x_1, y_1) for H_1 with origin at q_1 (in particular, y_1 vanishes on H_1). Note also that the graph Γ' obtained by removing C_n and its adjacent edge from Γ_1 is a contractible graph which unique (-1) -vertex is still E. Furthermore, this graph is the dual graph of $\pi_1^{-1}(q_1)$. Since its length is $n-1$ we are done by the induction assumption.

Proposition 4.6. Let $\mu : \check{X} \to X_0$, $q_0 \in X_0$, and $\mu_{i+1} : X'_{i+1} \to X'_{i}$, $i = 1, ..., n-1$ be as in Proposition [3.6.](#page-4-4) Let (x_0, y_0) be a bottom coordinate system on $\mathbb{A}^2 \simeq U_0 \subset X_0$ for $\mu_1: X'_1 \to X_0$ with the origin at q_0 . Then μ can be presented as a composition of morphisms $\nu_i: X_i \to X_{i-1}, i = 1, ..., N$ with $\breve{X} = X_N$ such that

(i) the restriction of ν_i is an isomorphism over $X_{i-1} \setminus \{q_{i-1}\}\$ for some $q_{i-1} \in X_{i-1}$ and the exceptional divisor of ν_i contains only one (-1) -curve H_i ;

(ii) there is an open subset $U_i \simeq \mathbb{A}^2_{x_i,y_i}$ of X_i such that $q_i \in U_i \cap H_i$ and y_i vanishes on $H_i \cap U_i$ for $i < N$;

(iii) each ν_{i+1} is the minimal resolution of indeterminacy points of a function of the form $(x_i - c_i)^{k_i} / y_i^{m_i}$ where c_i is the x_i -coordinate of q_i and $m_i \leq k_i$.

Proof. If $n = 1$, then we are done by Lemma [4.5.](#page-6-0) Otherwise, let $\nu_1 = \mu_1$ and consider the top coordinate system (x_1, y_1) of μ_1 associated with (x_0, y_0) . In particular, y_1 vanishes on the (-1) -vertex H'_1 of the exceptional divisor of μ_1 . Let Γ and Γ_i be as in Proposition [3.6.](#page-4-4) The smallest subgraph of Γ consisting of the proper transform of H'_1 and all vertices of Γ_2 is linear. Thus, we can decompose μ_2 as in Lemma [4.5](#page-6-0) using (x_1, y_1) as the bottom coordinate system in the first factor of the decomposition. Now the induction yields the desired conclusion.

Lemma 4.7. Let the assumptions of Proposition [4.6](#page-7-0) hold, H_0 be the x_0 -coordinate in U_0 , and $\ddot{U}_0 \simeq \mathbb{A}^2_{\ddot{x}_0,\ddot{y}_0}$ be an open subset of X_0 with the origin at q_0 and \ddot{y}_0 vanishing on $H_0 \cap U_0$, i.e., (\ddot{x}_0, \ddot{y}_0) is another bottom coordinate system for $\nu_1 : X_1 \to X_0$. Let $\mathbb{A}^2_{\ddot{x}_i,\ddot{y}_i} \simeq \ddot{U}_i \subset X_i$ and \ddot{c}_i have the same meaning for (\ddot{x}_0,\ddot{y}_0) as U_i and c_i for (x_0, y_0) . Let α be an automorphism of X_0 such that $\alpha(q_0) = q_0$, $\alpha(U_0) = \ddot{U}_0$, and $\alpha^*((\ddot{x}_0, \ddot{y}_0)) = (x_0(1 + p(x_0, y_0)), y_0(1 + q(x_0, y_0)))$ where p and q are rational functions on $U_0 \simeq \mathbb{A}^2_{x_0,y_0}$ that are regular in a neighborhood of the origin and vanish at the origin. Then for every $i \geq 1$ one has $\ddot{c}_i = c_i$ and there exists an automorphism α_i of X_i such that $\alpha_i(U_i) = \ddot{U}_i$ and $\nu_i \circ \ldots \circ \nu_1 \circ \alpha_i = \alpha \circ \nu_i \circ \ldots \circ \nu_1$.

Proof. Recall that $x_1 = x_0^k/y_0^m$, $y_1 = y_0^l/x_0^n$, $\ddot{x}_1 = \ddot{x}_0^k/\ddot{y}_0^m$ and $\ddot{y}_1 = \ddot{y}_0^l/\ddot{x}_0^n$ where kl – $mn = 1$. In particular, $(x_0, y_0) = (x_1^l y_1^m, x_1^n y_1^k)$. Then

$$
\alpha^*(\ddot{x}_1) = x_1(1 + f_1(x_0, y_0)) = x_1(1 + f_1(x_1^l y_1^m, x_1^n y_1^k)) = x_1(1 + p_1(x_1, y_1))
$$

and

$$
\alpha^*(\ddot{y}_1) = y_1(1 + g_1(x_0, y_0)) = y_1(1 + g_1(x_1^l y_1^m, x_1^n y_1^k)) = y_1(1 + q_1(x_1, y_1))
$$

where f_1 and g_1 are rational functions regular at the origin and vanishing at it. Since $k, m > 0$ we see that p_1 and q_1 are rational functions regular on the x_1 -axis in $U_1 \simeq$ $\mathbb{A}^2_{x_1,y_1}$ and vanishing on it. Hence, $\ddot{c}_1 = c_1$ and for the lift $\alpha_1 : X_1 \to \ddot{X}_1$ of α one has $\alpha_1(\tilde{U}_1) = \tilde{U}_1$ and

$$
\alpha_1^*((x_1,y_1)) = (x_1(1+p_1(x_1,y_1)), y_1(1+q_1(x_1,y_1))).
$$

Let $(\tilde{x}_1, \tilde{y}_1) = (x_1 - c_1, y_1)$ and $(\tilde{\tilde{x}}_1, \tilde{\tilde{y}}_1) = (\tilde{x}_1 - c_1, \tilde{y}_1)$. Since both $p_1(x_1, y_1)$ and $q_1(x_1, y_1)$ vanish on the x_1 -axis we see that $\alpha^*(\tilde{\ddot{x}}_1) = \tilde{x}_1(1 + \tilde{p}_1(\tilde{x}_1, \tilde{y}_1))$ and $\alpha^*(\tilde{\ddot{y}}_1) =$ $\tilde{y}_1(1 + \tilde{q}_1(\tilde{x}_1, \tilde{y}_1))$ where \tilde{p}_1 and \tilde{q}_1 vanish at the origin. Thus, replacing (x_0, y_0) and (\ddot{x}_0, \ddot{y}_0) with $(\tilde{x}_1, \tilde{y}_1)$ and $(\tilde{\ddot{x}}_1, \tilde{\ddot{y}}_1)$, respectively, and using the induction on N we get the desired conclusion.

As an immediate consequence of Lemma [4.7](#page-7-1) we have the following.

Proposition 4.8. Let the assumptions of Lemma [4.7](#page-7-1) hold, R be the closure of $U_0 \setminus$ $\mu(U_N)$ and \ddot{R} be the closure of $\ddot{U}_0 \setminus \mu(\ddot{U}_N)$. Then $\alpha(R) = \ddot{R}$.

5. The case of surfaces

Let us fix notations for the rest of this section.

Notation 5.1. (1) Let $Q' = \mathbb{P}^1 \times \mathbb{P}^1$ be the quadric that is the natural completion of $\mathbb{A}^2_{x,y}$. Let $E'_b = \{y = b\}$ and $F'_a = \{x = a\}$ where $a, b \in \mathbb{P}^1$. Let $q'_b = F'_b \cap E'_0$ and $p'_b = F'_b \cap E'_\infty$. By $Q'_* \subset Q'$ we denote the union of $\mathbb{A}^2_{x,y}$ and the germ (Q', p'_∞) . We consider also the quadric Q'' that is the natural completion of $\mathbb{A}^2_{s,t}$ and suppose that E''_b, F''_a q''_a, Q''_*, q''_b b'' and p''_b have the same meaning for Q'' as E'_b , F'_a q'_{a}, Q'_{*}, q'_{b} η'_{b} and p'_{b} \int_b' for Q' .

(2) Let m and k be relatively prime natural numbers with $m \leq k$. We suppose that n and l are as in Remark [3.3.](#page-4-1) In particular, $kl - mn = 1$. By $\chi : Q'' \dashrightarrow Q'$ we denote the rational map that extends the morphism $\mathbb{A}^2_{s,t} \to \mathbb{A}^2_{x,y}$, $(s,t) \mapsto (s^l t^m, s^n t^k)$ (i.e., $\chi(s,t) = (st, t)$ for $m = k = 1$).

Lemma 5.2. Let $c \in \mathbf{k}$ and α be the automorphism of Q' whose restriction to $\mathbb{A}^2_{x,y}$ is given by $(x, y) \mapsto (x + c, y)$. Then $\alpha \circ \chi|_{Q''_*}$ yields a morphism $\chi_c : Q''_* \to Q'_*$.

Proof. Since the restriction of α yields an automorphism of Q'_{*} it suffices to consider the case $c = 0$. Note that p'_{∞} is contained a neighborhood isomorphic to $\mathbb{A}^2_{\hat{x},\hat{y}}$ where $(\hat{x}, \hat{y}) = (1/x, 1/y)$ and the union of this neighborhood with $\mathbb{A}^2_{x,y}$ contains $Q'_*,$. There is the similar neighborhood $\mathbb{A}^2_{\hat{s},\hat{t}}$ of p''_{∞} whose union with $\mathbb{A}^2_{s,t}$ contains Q''_* where (\hat{s},\hat{t}) = $(1/s, 1/t)$. The formula for χ in Notation [5.1](#page-8-1) implies that χ is regular on $\mathbb{A}^2_{\hat{s},\hat{t}}$ and

 $\chi(\mathbb{A}^2_{\hat{s},\hat{t}}) \subset \mathbb{A}^2_{\hat{x},\hat{y}}$ (more precisely, $\chi|_{\mathbb{A}^2_{\hat{s},\hat{t}}}$ is given by $(\hat{s},\hat{t}) \mapsto (\hat{s}^l\hat{t}^m,\hat{s}^n\hat{t}^k)$). Thus, χ is regular on Q''_* and $\chi(Q''_*) \subset Q'_*$ which concludes the proof.

Lemma 5.3. Let $\tau : \breve{Q} \to Q'_{*}$ be the minimal resolution of indeterminacy points of the function $(x-c)^k/y^m$ at q_c' c^{\prime} and \breve{E} be the only (-1)-component in $\tau^{-1}(q^{\prime}_{c})$ $_{c}^{\prime }).$

(1) Let $k > 1$. Then there is a morphism $\breve{\chi}_c : Q''_* \to \breve{Q}$ such that $\chi_c = \tau \circ \breve{\chi}_c$ and the restriction of χ_c to $\mathbb{A}^2_{s,t}$ is an embedding. Furthermore, $\chi_c(E_0'' \setminus \{q_\infty''\}) \subset E$ and there is an irreducible component C of $\tau^{-1}(q'_c)$ ζ_c') meeting \breve{E} such that $\breve{\chi}_c(F_0'')$ $\binom{m}{0} \setminus \{p''_0$ $'_{0}\})\subset C.$

(2) If $k = m = 1$, then the same conclusion holds if one replaces C with the proper transform of F_c' $_{c}^{\prime}$ in \breve{Q} .

Proof. Let $\check{\chi}_c$: $Q''_* \dashrightarrow \check{Q}$ be the lift of χ_c . Note that $s = \chi_c^*$ $c^*(x-c)^k/y^m$ and $t = \chi_c^*$ $c_c(y^l/(x-c)ⁿ)$. Hence, (1) follows from Lemma [3.4.](#page-4-3) The similar argument works for (2) with $s = \chi^*((x-c)/y)$ and $t = \chi^*(y)$. Since the restriction of τ over $Q'_* \setminus \{q'_c\}$ is an isomorphism we have the regularity of χ_c which concludes the proof.

Lemma 5.4. Let X be a smooth complete rational surface, $U_0 \simeq \mathbb{A}^2_{x_0,y_0}$ be an open subset of X, and $q = (c, 0) \in U_0$. Let $\nu : \hat{X} \to X$ be the minimal resolution of indeterminacy of the function $(x_0-c)^k/y_0^m$ at q, that is, (x_0, y_0) is a bottom coordinate system for v and we have a top coordinate system (x_1, y_1) on an open subset $U_1 \simeq \mathbb{A}^2_{x_1, y_1}$ of \hat{X} as in Definition [4.3.](#page-6-1) Suppose that $\varphi: Q'_{*} \to X$ is a morphism such that its restriction to $\mathbb{A}^2_{x,y}$ yields the isomorphism $\mathbb{A}^2_{x,y} \to \mathbb{A}^2_{x_0,y_0}$ for which $\varphi^*(x_0,y_0) = (x,y)$. Then there is a morphism $\hat{\varphi}: Q''_* \to \hat{X}$ such that its restriction yields an isomorphism $\mathbb{A}^2_{s,t} \to U_1$ with $\hat{\varphi}^*(x_1, y_1) = (s, t)$.

Proof. Let the notations of Lemmas [5.3](#page-9-0) hold. Then there is a commutative diagram

$$
Q''_* \xrightarrow{\check{X}c} \vec{Q} \xrightarrow{\kappa} \hat{X}
$$

$$
\downarrow_c \qquad \qquad \downarrow_r
$$

$$
Q'_* \xrightarrow{\varphi} X.
$$

Recall that $s = \chi_c^*$ $c^*(x-c)^k/y^m$ and $t=\chi_c^*$ $c^*(y^l/(x-c)^n)$. Hence, $s = (\varphi \circ \chi_c)^*((x_0 (c)^k/y_0^m$ and $t = (\varphi \circ \chi_c)^*(y_0^l/(x_0-c)^n)$. By Definition [4.3](#page-6-1) we can suppose that $(x_1, y_1) =$ $\nu^*((x_0-c)^k/y_0^m, y_0^l/(x_0-c)^n)$. Let $\hat{\varphi} := \kappa \circ \check{\chi}_c : Q''_* \to \hat{X}$. Then the commutativity of the diagram implies that $\hat{\varphi}^*(x_1, y_1) = (s, t)$ and we have the desired conclusion. \square

Proposition 5.5. Let the assumptions of Proposition [3.6](#page-4-4) hold with $X_0 = Q'$ and H being an irreducible component in a complete connected SNC curve $D \subset X$. Suppose that $\nu_i: X_i \to X_{i-1}$ and $U_i \simeq \mathbb{A}^2_{x_i,y_i}$, $i = 1,\ldots,N$ are as in Proposition [4.6,](#page-7-0) i.e., $X_N = \breve{X}$. Let $\psi(D)$ be a finite subset of U_0 and $q_0 = \psi(H)$ be the origin in $\mathbb{A}^2_{x,y} \subset Q'$. Suppose also that (x, y) coincides with (x_0, y_0) . Then replacing H, if necessary, by another irreducible component of D one can find a morphism $\varphi_N: Q''_* \to X_N$ such that its restriction yields an isomorphism $\mathbb{A}^2_{s,t} \to U_N \simeq \mathbb{A}^2_{x_N,y_N}$ for which $\varphi^*_N(x_N, y_N) = (s,t)$ and $\theta(D)$ is the union of the curve $H_N = \theta(H)$ and a finite subset.

Proof. Let $\mu^{-1}(q_0)$ contain the proper transform \tilde{H} of an irreducible component of D different from H. Contracting H_N and some other irreducible components in the weighted dual graph Γ of $\mu^{-1}(q_0)$ we obtain another contractible graph $\tilde{\Gamma}$ with the proper transform of H as the only (−1)-vertex. Note that we have a morphism $\tilde{\mu}$: $\tilde{X} \to X_0$ such that φ factors through $\tilde{\mu}$ and $\tilde{\Gamma}$ is the weighted dual graph of $\tilde{\mu}^{-1}(q_0)$. Since the replacement of H_N and μ by H and $\tilde{\mu}$ reduces the number of vertices in dual graphs we can assume from the beginning that H is the only irreducible component of D whose proper transform is contained in $\mu^{-1}(q_0)$. Let $\varphi_0: Q'_* \to X_0$ be the identity isomorphism, i.e., φ_0^* $\chi_0^*(x_0, y_0) = (x, y)$. By Lemma [5.4](#page-9-1) we have morphism $\varphi_1 : Q''_* \to X_1$ such that φ_1^* $f_1(x_1, y_1) = (s, t)$. Note that by our assumption unless $N = 1$ none of the irreducible components of $\nu_1^{-1}(q_0)$ is the proper transform of an irreducible component of D. Proceeding by induction we get a morphism $\varphi_N: Q'' \to X_N$ such that none of the irreducible components of $\mu^{-1}(q_0)$ but H_N is the proper transform of irreducible components of D and $\varphi_N^*(x_N, y_N) = (s, t)$ which is the desired conclusion.

Proposition 5.6. Let the assumptions of Proposition [5.5](#page-9-2) hold and μ be invertible over a neighborhood of a point $v_0 \in \mathbb{A}_{x_0}^* \times \mathbb{A}_{y_0}^* \subset U_0$. The one can suppose that

(i) $\varphi_N(p''_{\infty})$ is a general point of X_N ;

(ii) $v_N = \mu^{-1}(v_0)$ is contained in $\mathbb{A}_{x_N}^* \times \mathbb{A}_{y_N}^* \subset U_N$.

Proof. Let $\alpha: X_0 \to X_0$ be given by

$$
\alpha(x,y) = \left(\frac{ax}{a-x}, \frac{by}{b-y}\right) =: (\ddot{x}, \ddot{y})
$$

where a and b are general elements of **k**. Then $\mathbb{A}^2_{\ddot{x}_0,\ddot{y}_0} \simeq \ddot{U}_0$ is an open subset of X_0 containing v_0 and (\ddot{x}_0, \ddot{y}_0) is another bottom coordinate system for $\nu_1 : X_1 \to X_0$. Let $\mathbb{A}^2_{\ddot{x}_i,\ddot{y}_i} \simeq \ddot{U}_i$ have the same meaning as in Lemma [4.7,](#page-7-1) R be the closure of $U_0 \setminus \mu(U_N)$, and \ddot{R} be the closure of $\ddot{U}_0 \setminus \mu(\ddot{U}_N)$. By Proposition [4.8](#page-8-2) $\alpha^{-1}(\ddot{R}) = R$. Since a and b are general $\alpha^{-1}(v_0)$ is a general point of U_0 . In particular, $\alpha^{-1}(v_0)$ does not belong to R which implies (ii). Since $\alpha(p''_{\infty})$ is a general point of \ddot{U}_0 we have also (i) which concludes the proof. \Box

In the next lemmas we shall justify the assumptions of Propositions 5.5 and 5.6 .

Lemma 5.7. Let $\psi: X \to \mathbb{P}^n$ be a birational morphism of smooth complete varieties, $v \in X$, and ψ be invertible over a neighborhood of $v_0 = \psi(v)$. Let $\mathcal D$ be a nontrivial line bundle on X and D be a divisor in X such that $\mathcal{D} = \mathcal{O}_X(D)$. Suppose that the pushforward of D generates a nontrivial line bundle on \mathbb{P}^n . Then for an appropriate choice of D there is a birational morphism $\varphi : B \times \mathbb{P}^1 \to X$ such that

(i) φ is locally invertible over a neighborhood of v;

(ii) there is a general point o of some irreducible component of D such that each curve $\varphi(b \times \mathbb{P}^1)$, $b \in B$ meets D at o only and the intersection of this curve and D is transversal;

(iii) the restriction of D to $\varphi(b \times \mathbb{P}^1)$ is a nontrivial line bundle.

Proof. Let L be a hyperplane in \mathbb{P}^n that does not contain v_0 . One can choose a rational function f on \mathbb{P}^n with a divisor T such that $T + \psi_* D = mL$. Note that

 $\mathcal{O}(D) = \mathcal{O}(\psi^*(T) + D)$. Hence, we can suppose that $\psi_*D = mL$. By the assumption $m \neq 0$. Choose a general point *o* in L and a general hyperplane \mathbb{P}^{n-1} in \mathbb{P}^n . Then for \mathbb{A}^{n-1} ⊂ \mathbb{P}^{n-1} one can consider the set of lines each of which pass through o' and a point of \mathbb{A}^{n-1} . This set generates a morphism $\varphi_0: \mathbb{A}^{n-1} \times \mathbb{P}^1 \to \mathbb{P}^n$. Since o' is general we can suppose that ψ is invertible over a neighborhood of the line $\varphi_0(b_0 \times \mathbb{P}^1)$ containing v₀. Thus, ψ is invertible over $\varphi_0(B \times \mathbb{P}^1)$ for some open subset $B \subset \mathbb{A}^{n-1}$ containing b_0 . The lift of $\varphi_0|_{B\times\mathbb{P}^1}$ to X yields $\varphi: B\times\mathbb{P}^1\to X$ such that ever curve $\varphi(b\times\mathbb{P}^1)$ meets D only at a general point in the proper transform H of L . Hence, the restriction of D to $\varphi(b \times \mathbb{P}^1)$ is a nontrivial line bundle which concludes the proof.

Lemma 5.8. Let $\beta: Y \to \mathbb{P}^2$ be a birational morphism from a smooth complete surface Y such that it is invertible over a neighborhood of $u_0 \in \mathbb{P}^2$. Let T be a complete SNC curve in Y such that $u = \beta^{-1}(u_0) \notin T$ and $\beta(T)$ is finite. Then there is a commutative diagram

(3)
$$
\begin{array}{ccc}\nX & \stackrel{\lambda}{\to} & Y \\
\downarrow^{\psi} & & \downarrow^{\beta} \\
Q' & \stackrel{\gamma}{\dashrightarrow} & \mathbb{P}^2\n\end{array}
$$

where X is a smooth complete rational surface, the morphisms ψ and λ are birational and λ is locally invertible over a neighborhood of $\{u\} \cup T$. Furthermore, one can suppose that $\psi: X \to X_0 = Q'$ satisfies the assumptions of Propositions [5.6](#page-10-0) with $D = \lambda^{-1}(T)$, and $v_0 = \psi(\lambda^{-1}(u)).$

Proof. Let L be a general line in \mathbb{P}^2 . Then L does not contain u_0 and the set $M_0 \subset \mathbb{P}^2$ of points over which β is not invertible (note that $\beta(T) \subset M_0$). Thus, $U = \mathbb{P}^2 \setminus L$ is a neighborhood of u_0 isomorphic to $\mathbb{A}^2_{x_0,y_0}$. Let F be any irreducible component of T. Then β admits a decomposition as in Proposition [3.6](#page-4-4) with $\psi : X \to X_0$ and H replaced by $\beta: Y \to \mathbb{P}^2$ and F. Consequently, β admits a decomposition as in Proposition [4.6](#page-7-0) with (x_0, y_0) having the same meaning as the bottom coordinate system in Proposition [4.6.](#page-7-0) By Remark [4.4](#page-6-2) the coordinate system (x_0, y_0) can be chosen so that $u_0 \in \mathbb{A}_{x_0}^* \times \mathbb{A}_{y_0}^*$ $_{y_0}^*$. Choosing Q' as the natural completion of $\mathbb{A}^2_{x_0,y_0}$ we get the birational map $\gamma: Q^{\gamma} \dashrightarrow \mathbb{P}^2$ which gives rise to commutative diagram in (3) with λ being a rational map. However, resolving the indeterminacy points of λ (which are located in $\beta^{-1}(L)$) we can suppose that λ is a morphism. Let H be the proper transform of F in X. By construction all the assumption of Proposition [5.5](#page-9-2) for $\psi: X \to Q'$, H, D, v_0 , and (x_0, y_0) are valid. By Proposition [5.6](#page-10-0) we have the desired conclusion.

Proposition 5.9. Let the assumptions and conclusions of Lemma [5.8](#page-11-1) hold and M be a finite subset of X that does not contain $v = \lambda^{-1}(u)$. Then there is an irreducible component H in D and a birational morphism $\varphi : B \times \mathbb{P}^1 \to X$ such that

(i) φ is locally invertible over a neighborhood of v;

(ii) there is a general point $o \in H$ such that for every $b \in B$ the curve $\varphi(b \times \mathbb{P}^1)$ meets D at o only and the intersection of $\varphi(b \times \mathbb{P}^1)$ and H is transversal;

(iii) the image of φ does not meet M.

Proof. Recall that we have a decomposition $\psi = \mu \circ \theta$ as in Propositions [3.6](#page-4-4) and [4.6](#page-7-0) where $\theta: X \to \check{X} = X_N$ and $\mu: X_N \to Q'$. Furthermore, there is an irreducible component H of D such that $\theta(D)$ is contained in the union of the curve $H_N = \theta(H)$ and the set M_N of points over which θ is not invertible. By Lemma [5.8](#page-11-1) and Propositions [5.5-](#page-9-2)[5.6](#page-10-0) U_N contains $v_N = \mu^{-1}(v_0) = \theta(v)$ and we have a morphism $\varphi_N : Q''_* \to X_N$ invertible over U_N where U_N is the isomorphic image of $\mathbb{A}^2_{s,t}$ under φ_N . Since ψ is invertible over v_0 , θ is invertible over v_N and, thus, v_N does not belong to the finite set $M' = M_N \cup \theta(M)$. Let $M'' = \varphi_N^{-1}(M') \cap \mathbb{A}^2_{s,t}$ and $v_N = \varphi_N(s_0, t_0)$ where by Proposition [5.6](#page-10-0) $(s_0, t_0) \in \mathbb{A}_s^* \times \mathbb{A}_t^*$ *. Consider all a, b and $c \in \mathbf{k}$ for which the image of the morphism $\gamma_{a,b,c} : \mathbb{A}_z^1 \to Q''_*, z \mapsto (a + bz + cz^2, z)$ contains (s_0, t_0) . Note that for any point $(s_1, t_1) \neq (s_0, t_0)$ in $\mathbb{A}^2_{s,t}$ and $a \in \mathbf{k}$ one can choose b and c such that the image of $\gamma_{a,b,c}$ does not contain (s_1, t_1) . In particular, we can suppose that for a general a_0 and some pair (b_0, c_0) the image of γ_{a_0,b_0,c_0} does not meet M''. Hence, for an appropriate neighborhood $B \subset \mathbb{A}^1$ of b_0 we have a morphism $B \times \mathbb{A}^1_z \to \mathbb{A}^2_{s,t}$, $(b, z) \mapsto (a_0+bz+c_0z, z)$ invertible over (s_0, t_0) such that its image image does not meet M''. This map extends to a morphism $\gamma: B \times \mathbb{P}^1 \to Q''_*$ such that $\gamma(B \times \infty) = p''_{\infty}$. Note that $\varphi_N \circ \gamma(B \times \mathbb{P}^1)$ does not meet M' since the restriction of φ_N to $\mathbb{A}^2_{s,t}$ is an isomorphism onto U_N and $\varphi_N(p''_{\infty})$ is a general point of X_N by Proposition [5.6.](#page-10-0) Furthermore, every curve $\varphi_N \circ \gamma(b \times \mathbb{P}^1)$ meets H_N transversely and at one general point only since every curve $\gamma(b \times \mathbb{P}^1)$ meets the s-axis transversely and only at the point with s-coordinate a_0 . Since the image of $\varphi_N \circ \gamma$ does not meet M_N we see that $\varphi_N \circ \gamma$ admits a lift $\varphi: B \times \mathbb{P}^1 \to X$ such that (i)-(iii) hold. Hence, we are done.

Lemma 5.10. Let π : $\tilde{X} \to X$ and $\tilde{\varphi}$: $B \times \mathbb{P}^1 \to \tilde{X}$ be birational morphisms of smooth varieties such π is locally invertible over a neighborhood of $v \in X$, $\tilde{v} = \pi^{-1}(v)$ is contained in the image of $\tilde{\varphi}$ and $\tilde{\varphi}$ is locally invertible over a neighborhood of \tilde{v} . Let D be a line bundle on X and $\tilde{\mathcal{D}} = \pi^* \mathcal{D}$. Suppose that for $b \in B$ such that $\tilde{v} \in \hat{\varphi}(b \times \mathbb{P}^1)$ the restriction of $\tilde{\mathcal{D}}$ to $\tilde{\varphi}(b \times \mathbb{P}^1)$ is a nontrivial line bundle. Then there exists φ : $B \times \mathbb{P}^1 \to X$ locally invertible over a neighborhood of v such that the restriction of \mathcal{D} to $\varphi(b \times \mathbb{P}^1)$ is a nontrivial line bundle.

Proof. Put $\varphi = \pi \circ \tilde{\varphi}$. Note that if the restriction of \mathcal{D} to $\varphi(b \times \mathbb{P}^1)$ is a trivial line bundle, then so is the restriction of $\tilde{\mathcal{D}}$ to $\tilde{\varphi}(\tilde{s} \times \mathbb{P}^1)$. This yields the desired conclusion.

Theorem 5.11. Let X be a smooth complete rational surface and D be a nontrivial line bundle on X with zero section Z. Then $\mathcal{D} \setminus Z$ is elliptic.

Proof. Recall that X is of class \mathcal{A}_0 (e.g., see the proof of Lemma [4.1\)](#page-5-1) and, hence, X is elliptic since every variety of class \mathcal{A}_0 is elliptic [\[Gro89,](#page-21-4) Section 3.5]. Thus, by Corollary [2.5](#page-2-2) it suffices to show that for every $v \in X$ one can find a birational morphism $\varphi : B \times \mathbb{P}^1 \to X$ invertible over a neighborhood of v such that for $b_0 \in B$ with $v \in \varphi(b_0 \times \mathbb{P}^1)$ the restriction of \mathcal{D} to $\varphi(b_0 \times \mathbb{P}^1)$ is a nontrivial line bundle. Let $U \simeq \mathbb{A}^2$ be a neighborhood of v in X. Letting \mathbb{P}^2 be a completion of U we have a rational map $\beta : X \dashrightarrow \mathbb{P}^2$ which is regular and invertible over U. Resolving indeterminacy points of this map we get a morphism $\tilde{X} \to \mathbb{P}^2$ invertible over U. By Lemma [5.10](#page-12-0) we

can suppose that $X = \tilde{X}$, that is β is regular. By Lemma [5.7](#page-10-1) we can assume that β (supp D) is finite for some SNC divisor D such that $\mathcal{D} = \mathcal{O}(D)$. Hence, we have a morphism $\psi: X \to Q'$ as in Lemma [5.8.](#page-11-1) By Proposition [5.9](#page-11-2) there is a birational morphism $\varphi: B \times \mathbb{P}^1 \to X$ as in Proposition [5.9](#page-11-2) such that the curve $\varphi(b_0 \times \mathbb{P}^1)$ passing through v meets only one irreducible component H of D . Hence, the restriction of D to $\varphi(0 \times \mathbb{P}^1)$ is a nontrivial line bundle which yields the desired conclusion.

6. Main Theorem

The aim of this section is the following.

Theorem 6.1. Let X be a complete uniformly rational variety and $\mathcal{D} \to X$ be a nontrivial line bundle on X with zero section $Z_{\mathcal{D}}$. Then $Y = \mathcal{D} \setminus Z_{\mathcal{D}}$ is elliptic.

By Corollary [2.5](#page-2-2) it suffices to show that for every $v \in X$ one can find a smooth variety B and a birational morphism $\varphi : B \times \mathbb{P}^1 \to X$ invertible over a neighborhood of v such that for $b_0 \in B$ with $v \in \varphi(b_0 \times \mathbb{P}^1)$ the restriction of D to $\varphi(b_0 \times \mathbb{P}^1)$ is a nontrivial line bundle. To establish this fact we need to check the following property for every $v \in X$.

Definition 6.2. Let X be a smooth complete rational variety, v be a point in X with a neighborhood isomorphic to an open subset of \mathbb{A}^n (which is automatic in the case when X is uniformly rational), and $\mathcal D$ be a nontrivial line bundle on X. We say that the triple (X, \mathcal{D}, v) has P-property if there is an SNC divisor D' such that $\mathcal{D} = \mathcal{O}_X(D')$ and either

(a) there is a morphism of X onto \mathbb{P}^n such that its restriction to a neighborhood of v is an embedding and the pushforward of D' is not a principal divisor on \mathbb{P}^n , or, otherwise,

(b) there is a birational morphism $\varphi : B \times \mathbb{P}^1 \to X$ invertible over a neighborhood of v such that $\varphi(B \times \mathbb{P}^1)$ meets D' at a general point o of one irreducible component H of D' only and the intersection of every curve $\varphi(b \times \mathbb{P}^1)$, $b \in B$ and H at this point is transversal.

In fact, we do not need to establish this property for every triple as in Definition [6.2](#page-13-2) since by Lemma [5.10](#page-12-0) we can replace (X, \mathcal{D}, v) by $(\tilde{X}, \tilde{\mathcal{D}}, \tilde{v})$ where $\pi : \tilde{X} \to X$ is a proper birational morphism from a smooth variety \tilde{X} invertible over a neighborhood of $v, \tilde{\mathcal{D}} = \pi^* \mathcal{D}$, and $\tilde{v} = \pi^{-1}(v)$. Furthermore, by Lemma [5.7](#page-10-1) and Corollary [2.5,](#page-2-2) Theorem [6.1](#page-13-0) is valid under the assumption (a) of Definition [6.2.](#page-13-2) Thus, we suppose further that there is a birational morphism $\theta: X \to \mathbb{P}^n$ invertible over a neighborhood of $\theta(v)$ such that $\theta(\text{supp } D')$ is contained in the indeterminacy set of θ^{-1} .

Lemma 6.3. One can suppose that there is a \mathbb{P}^1 -bundle pr : $Q \to S$ over $S \simeq \mathbb{P}^{n-1}$ and a proper birational morphism $\alpha: X \to Q$ such that for the composition $\rho = \text{pr} \circ \alpha$: $X \rightarrow S$ one has the following

(i) $\alpha(\text{supp } D')$ is contained in the indeterminacy set P of α^{-1} and none of the fibers of pr is contained in P;

(ii) there is a neighborhood $W \subset S\pr(P)$ of $s_0 = \varrho(v)$ for which $\varrho|_{\varrho^{-1}(W)} : \varrho^{-1}(W) \to$ W is a \mathbb{P}^1 -bundle:

(iii) pr has two disjoint section S and S_0 .

Proof. Let P be the indeterminacy set of θ^{-1} . We consider a general line L through $v_0 = \theta(v)$ in \mathbb{P}^n (in particular, L does not meet P), a general point $p_0 \in L$, and a general hyperplane $S \simeq \mathbb{P}^{n-1}$ through v_0 in \mathbb{P}^n . Let $\gamma: Q \to \mathbb{P}^n$ be the blowing up of \mathbb{P}^n at p_0 . Then one has the natural morphism $pr: Q \to S$ which is a \mathbb{P}^1 -bundle. By Lemma [5.10](#page-12-0) we can replace X with its blowing up \breve{X} at $\theta^{-1}(p_0)$ (and we also replace v by its preimage in this blowing up). This yields the morphism $\alpha : \check{X} \to Q$. Note that the indeterminacy set \check{P} of α^{-1} coincides with $\gamma^{-1}(P)$. Since L does not meet P there is a neighborhood $W \subset S \setminus \mathrm{pr}(\check{P})$ of $\mathrm{pr}(v_0)$ such that $\mathrm{pr}^{-1}(W) \cap \check{P} = \emptyset$. Consequently, $\varrho|_{\varrho^{-1}(W)} : \varrho^{-1}(W) \to W$ is a \mathbb{P}^1 -bundle. Since $p_0 \notin P$ the exceptional divisor of γ does not meet \check{P} . Hence, none of the fibers of pr is contained in \check{P} because the exceptional divisor is a section S_0 of the \mathbb{P}^1 -bundle (the other section S is generated by S). This concludes the proof. \Box

Lemma 6.4. One can also suppose that there is a \mathbb{P}^1 -bundle $\text{pr}: Q \to S$ over a smooth complete rational variety S and a proper birational morphism $\alpha: X \to Q$ such that for the composition $\rho = \text{pr} \circ \alpha : X \to S$ one has the following

(i) $\alpha(\text{supp } D')$ is contained in the indeterminacy set P of α^{-1} , dim $\rho(H) = n - 2$ for every irreducible component H of $D = \alpha^{-1}(P)$, and none of the fibers of pr is contained in P;

(ii) the restriction of ϱ over a neighborhood $W \subset S$ of $s_0 = \varrho(v)$ is \mathbb{P}^1 -bundle, $\varrho^{-1}(W) \cap D = \emptyset$, and W is isomorphic to a subset of \mathbb{P}^{n-1} ;

(iii) pr has two disjoint section S and S_0 .

Proof. Let $\alpha: X \to Q$, $\varrho: X \to S$, pr : $Q \to S$, and P be as in Lemma [6.3.](#page-13-3) Let $\pi : \tilde{S} \to S$ be a proper birational morphism from a smooth variety $S, \tilde{X} = X \times_S \tilde{S}$, and $\tilde{Q} = Q \times_S \tilde{S}$. Let $\tilde{\varrho} : \tilde{X} \to \tilde{S}$, $\mathrm{pr}_X : \tilde{X} \to X$, $\mathrm{pr}_Q : \tilde{Q} \to Q$ be the natural projections and $\tilde{\alpha}$: $\tilde{X} \to \tilde{Q}$ be the birational morphism induced by α . By Raynaud's flattening theorem $\pi : \tilde{S} \to S$ can be chosen such that $\tilde{\varrho}$ is flat. Furthermore, since ϱ is flat over $s_0 = \rho(v)$ we can suppose that π is invertible over a neighborhood W of s_0 in S [\[Ray72,](#page-21-8) Chapter 4, Theorem 1]. Thus, the restriction of pr_X over W is invertible and, in particular it is invertible over neighborhood of $v \in \varrho^{-1}(W)$. Let \tilde{P} be the indeterminacy set of $\tilde{\alpha}^{-1}$, that is, $\tilde{P} \subset \text{pr}_Q^{-1}(P)$. Since \tilde{Q} is smooth every irreducible component \tilde{H} of $\tilde{D} = \tilde{\alpha}^{-1}(\tilde{P})$ is a hypersurface in \tilde{X} by van der Waerden's theorem (e.g., $\left[\text{DaSh94}, \text{Chap. II}, \text{Section 7.3}\right]$). Hence, $\tilde{\rho}(\tilde{H})$ must have dimension $n-2$ since each fiber of $\tilde{\varrho}$ is one-dimensional because of flatness. By Lemma [6.3](#page-13-3) (i) $\text{pr}^{-1}(s) \setminus P$ is a nonempty open subset of $pr^{-1}(s) \simeq \mathbb{P}^1$ for every $s \in S$. Hence, $\tilde{pr}^{-1}(s) \setminus \tilde{P}$ is a nonempty open subset of $\tilde{pr}^{-1}(s)$ for every $\tilde{s} \in \tilde{S}$. Thus, none of the fibers of \tilde{pr} is contained in \tilde{P} . By Lemma [5.10](#page-12-0) we can replace X by \tilde{X} (and v by $\mathrm{pr}_{X}^{-1}(v)$) which yields the desired conclusion, since (iii) follows from Lemma 6.3 (iii). **Lemma 6.5.** Let the notations of Lemma [6.4](#page-14-0) hold and F be an irreducible component of $T = \rho(D)$. Suppose that property $\mathcal P$ is true for the triple $(S, \mathcal O_S(F), s_0)$. Then there is a morphism $\varphi : B \times \mathbb{P}^1 \to S$ invertible over a neighborhood of s_0 such that the set $F \cap \varphi(B \times \mathbb{P}^1)$ consists of general points of F and at one of them o the intersection of every curve $\varphi(b \times \mathbb{P}^1)$, $b \in B$ and F is transversal.

Proof. Let Z be a proper closed subset of F. Recall that we have a morphism $\pi : S \to$ \mathbb{P}^{n-1} locally invertible over a neighborhood of $\pi(s_0)$. Let $Z' = \pi(Z)$. If $F' = \pi(F)$ is a hypersurface in \mathbb{P}^{n-1} , then consider a general point o' in $F' \setminus Z'$, a neighborhood B of $\pi(s_0)$ in a hyperplane $L \simeq \mathbb{P}^{n-2}$ of \mathbb{P}^{n-1} containing $\pi(s_0)$ but not o' , and the set of lines passing through o' and the points of B . One can choose B such that this set generates a morphism $\psi: B \times \mathbb{P}^1 \to \mathbb{P}^{n-1}$ and none of these lines meets Z'. Since o' is a general point of F' the morphism π is invertible over a neighborhood of the line through $\pi(s_0)$ and o' . Hence, for an appropriate B we can suppose that ψ admits a lift to S which is a desired morphism $\varphi : B \times \mathbb{P}^1 \to S$. If dim $F' \leq n-3$, then the desired conclusion follows from the assumption (b) of property P and we are done.

For a smooth irreducible subvariety C of codimension at least 2 in a smooth variety Y we denote by $\text{Bl}_C Y$ the blowing up of Y along C. If Y is a variety over V (that is, we have a morphism $\beta: Y \to V$ and F is a hypersurface in V such that $\beta(C) = F$ and $\beta|_C : C \to F$ is an unramified covering, then we call C a multisection over F.

Lemma 6.6. Let $\alpha: Y' \to Y_0' = V \times \mathbb{P}^1$ be a proper morphism of smooth n-dimensional varieties, pr : $Y_0' \to V$ be the natural projection, $\rho = \text{pr} \circ \alpha$, and F be a hypersurface in V such that the restriction of α over $V \setminus F$ is invertible. Then one can find an open subset V^* of V with a nonempty smooth $F^* = V^* \cap F$ such that $\alpha|_{\varrho^{-1}(V^*)} : \varrho^{-1}(V^*) \to$ ${\rm pr}^{-1}(V^*)$ is a composition of blowing ups $\nu_i: Y_i = \text{Bl}_{C_{i-1}} Y_{i-1} \to Y_{i-1}, i = 1, \ldots, N$ (where $Y_0 = \text{pr}^{-1}(V^*)$ and $Y_N = \varrho^{-1}(V^*)$) such that C_{i-1} is a multisection over F^* . Furthermore, $\varrho^{-1}(F^*)$ is an SNC divisor with every point in $\varrho^{-1}(F^*)$ contained in at most two irreducible components of $\varrho^{-1}(F^*)$.

Proof. Let F^* be any nonempty smooth affine open subset of F such that it admits a finite morphism onto an open subset W of \mathbb{A}^{n-2} . Choose an affine neighborhood $V^* \subset V$ of F^* such that the morphism $F^* \to W$ extends to a morphism $V^* \to W$, that is, $\varrho^{-1}(V^*)$ and $\operatorname{pr}^{-1}(V^*)$ are varieties over W. To simplify notations replace V and F by V^* and F^* , and morphisms over V by their restrictions over V^* . By [\[Har04,](#page-21-10) Chapter III, Corollary 10.7] we can suppose now that the natural morphisms $\varrho^{-1}(V) \to W$ and $\text{pr}^{-1}(V) \to W$ are smooth. Let **K** be the algebraic closure of the field of rational functions on W . For every algebraic variety X over W we denote by X^K the variety Spec K $\times_{\text{Spec } K} X$ and we also use upper index K for points in X^K and functions on it. Then we have proper morphisms $\alpha^{\mathbf{K}} : Y_N^{\mathbf{K}} \to Y_0^{\mathbf{K}} = (V \times \mathbb{P}^1)^{\mathbf{K}}$, ${\rm pr}^{\mathbf{K}}: Y_{0}^{\mathbf{K}} \to V^{\mathbf{K}}$ and $\varrho^{\mathbf{K}}: Y_{N}^{\mathbf{K}} \to V^{\mathbf{K}}$ where $Y_{N}^{\mathbf{K}}$ and $Y_{0}^{\mathbf{K}}$ $V_{0}^{\mathbf{K}}$ are smooth surfaces, $V^{\mathbf{K}}$ is a smooth curve and the restriction of $\alpha^{\mathbf{K}}$ is an isomorphism over $V^{\mathbf{K}} \setminus F^{\mathbf{K}}$. By [\[Har04,](#page-21-10) Chap. V, Theorem 5.3] α^{K} can be presented as a composition of monoidal transformations $\nu_i^{\mathbf{K}}$ $X_i^{\mathbf{K}}$: $Y_i^{\mathbf{K}} \rightarrow Y_{i-1}^{\mathbf{K}}$, $i = 1, ..., N$ with centers at points $q_{i-1}^{\mathbf{K}}$. The

SNC curve $(\varrho^{\mathbf{K}})^{-1}(F^{\mathbf{K}})$ consists of irreducible components $L_i^{\mathbf{K}}$ $\frac{\mathbf{k}}{i}$ such that the proper transform of $L_i^{\mathbf{K}}$ $\sum_{i}^{\mathbf{K}}$ in $Y_i^{\mathbf{K}}$ $\mathcal{I}_i^{\mathbf{K}}$ is the exceptional divisor of $\nu_i^{\mathbf{K}}$ ^K for $i \geq 1$, whereas $L_0^{\mathbf{K}}$ $_{0}^{\mathbf{K}}$ is the proper transform of $(pr^{K})^{-1}(F^{K})$. Each q_i^{K} i^k is either the intersection of the proper transforms in $Y_i^{\mathbf{K}}$ $L_i^{\mathbf{K}}$ of two components $L_j^{\mathbf{K}}$ \boldsymbol{K} and $L_l^{\mathbf{K}}$ \sum_{l}^{k} (where $j, l \leq i$) or it belongs to the proper transform of $L_i^{\mathbf{K}}$ **K** only. Equipping $L_j^{\mathbf{K}}$ with a coordinate function we can treat the coordinate of $q_i^{\mathbf{K}}$ i^k is a mutivalued function C_i over W and, therefore, over F. Taking a smaller $F = F^*$, if necessary, one can suppose that C_0 is a multisection over F located in $pr^{-1}(F) \simeq F \times \mathbb{P}^1$. Thus, we have $\nu_1 : Bl_{C_0} Y_0 = Y_1 \to Y_0$ and the field extension \mathbf{K}/\mathbf{k} transforms this morphism into $\nu_1^{\mathbf{K}}$ $Y_1^{\mathbf{K}}$: $Y_1^{\mathbf{K}} \rightarrow Y_0^{\mathbf{K}}$ $_{0}^{\prime}$. Assume that we have already constructed Y_i such that the preimage P_i of F in Y_i under the natural projection $Y_i \to V$ is a hypersurface of SNC type and it consists of irreducible components L_j , $j = 0, \ldots, i$ of the form $C \times \mathbb{P}^1$ each of which becomes the proper transform of $\tilde{L}_i^{\mathbf{K}}$ **K** in $Y_i^{\mathbf{K}}$ under the field extension \mathbf{K}/\mathbf{k} . Reducing the size of F we can suppose that C_i is a multisection over F such that either it is the intersection of L_j and L_l or it is contained in L_i only without meeting the other irreducible components of P_i . In particular, ν_{i+1} : $\text{Bl}_{C_i} Y_i = Y_{i+1} \to Y_i$ is the blowing up along a smooth center, $L_{i+1} \simeq C_i \times \mathbb{P}^1$, P_{i+1} is an SNC hypersurface, and the field extension \mathbf{K}/\mathbf{k} transforms this morphism into $\nu_{i+1}^{\mathbf{K}}$: $Y_{i+1}^{\mathbf{K}} \to Y_i^{\mathbf{K}}$ **EVA**. Thus, the induction concludes the proof. \square

Notation 6.7. (1) Let the notation of Lemmas [6.4](#page-14-0) and [6.5](#page-15-0) hold and $V^* \subset S$ be an open affine set such that $F^* = F \cap V^*$ contains *o* and admits a finite étale morphism onto an open subset of \mathbb{A}^{n-2} extendable to a morphism $V^* \to \mathbb{A}^{n-2}$. Furthermore, we suppose that V^* does not meet $T \setminus F$. For $Y = \varrho^{-1}(V^*)$ and $Y_0 = \text{pr}^{-1}(V^*)$ we have the morphisms $\alpha_V : Y \to Y_0$, $pr_V : Y_0 \to V^*$, and $\rho_V : Y \to V^*$ that are the restrictions of α , pr, and ϱ over V^* . Without loss of generality we can also suppose that $\varphi^{-1}(F^*) = \varphi^{-1}(o)$ coincides with $R = B \times p_0$ where $p_0 \in \mathbb{P}^1$.

(2) Let $\hat{Q} = (B \times \mathbb{P}^1) \times_{S,\varphi,\text{pr}} Q$, $\hat{X} = (B \times \mathbb{P}^1) \times_{S,\varphi,\varrho} X$, $\hat{D} = (B \times \mathbb{P}^1) \times_{S,\varphi,\varrho} D$ (where D is as in Lemma [6.4\)](#page-14-0), $\hat{pr}: \hat{Q} \to B \times \mathbb{P}^1$ and $\hat{\varrho}: \hat{X} \to B \times \mathbb{P}^1$ are the natural projections and the morphism $\hat{\alpha}$: $\hat{X} \to \hat{Q}$ is induced by α . For every $b \in B$ we let $\hat{Q}^b = \hat{\mathrm{pr}}^{-1}(b \times \mathbb{P}^1).$

(3) For every variety Z over V^* we let $\hat{Z} = Z \times_{V^*} \hat{V}^*$ where $\hat{V}^* = \varphi^{-1}(V^*)$, and for a morphism $\gamma: Z_1 \to Z_2$ of varieties above V^* we denote by $\hat{\gamma}: \hat{Z}_1 \to \hat{Z}_2$ the induced morphism. In particular, we have the morphisms $\hat{\alpha}_V : \hat{Y} \to \hat{Y}_0$, $\hat{\mathrm{pr}}_V : \hat{Y}_0 \to \hat{V}^*$, and $\hat{\varrho}_V : \hat{Y} \to \hat{V}^*$. For every $b \in B$ we consider $\hat{Y}^b_0 = \hat{\mathrm{pr}}_V^{-1}(\hat{V}^* \cap (b \times \mathbb{P}^1)) \subset \hat{Q}^b$ and $\hat{Y}^b = \hat{\varrho}_V^{-1}$ $_V^{-1}(\hat{V}^* \cap (b \times \mathbb{P}^1)).$

Lemma 6.8. One can suppose that there is a natural isomorphism $\mu: \hat{Q} \to B \times \hat{Q}^{b_0}$ over $B \times \mathbb{P}^1$ where b_0 is any point in B (say, $s_0 \in \varphi(b_0 \times \mathbb{P}^1)$).

Proof. By Lemma [6.4](#page-14-0) (iii) pr has two disjoint sections $\hat{\mathcal{S}}$ and $\hat{\mathcal{S}}_0$. Thus, $\hat{Q} \setminus \hat{\mathcal{S}}_0$ can be viewed as a line bundle $\mathcal L$ over $B\times \mathbb P^1$ and $\hat Q$ as the proectivization of $\mathcal L\oplus \mathcal E$ where $\mathcal E$ is the trivial line bundle. Let \mathcal{L}_b and \mathcal{E}_b be the restrictions of these bundles to $b \times \mathbb{P}^1$. Replacing B by an appropriate affine neighborhood of b_0 we can suppose that a divisor generating $\mathcal L$ is linearly equivalent to a divisor of the form $m(B \times q)$ where $q \in \mathbb{P}^1$ and $m \in \mathbb{Z}$. Thus, $\mathcal{L} = \kappa^* \mathcal{L}_{b_0}$ where $\kappa : B \times \mathbb{P}^1 \to \mathbb{P}^1 \simeq b_0 \times \mathbb{P}^1$ is the natural projection. Consequently, $\kappa^*(\mathcal{L}_{b_0} \oplus \mathcal{E}_{b_0}) = \mathcal{L} \oplus \mathcal{E}$ and \hat{Q} is isomorphic to $B \times \hat{Q}^{b_0}$ over $B \times \mathbb{P}^1$. \Box

Restricting the isomorphism μ from Lemma [6.8](#page-16-0) to $\hat{Y}_0 \subset \hat{Q}$ we get the following

Lemma 6.9. There is a rational map $\nu_0 : \hat{Y}_0 \dashrightarrow B \times \hat{Y}_0^{b_0}$ that is biregular over the neighborhood $U = \hat{V}^* \cap \hat{\text{pr}} \circ \mu^{-1}(B \times \hat{Y}^{b_0})$ of R.

Lemma 6.10. There is a rational map $\nu : \hat{Y} \dashrightarrow B \times \hat{Y}^{b_0}$ that is biregular over U and \hat{Y}^{b_0} is a smooth surface.

Proof. Let $\nu_{i+1}: Y_{i+1} \to Y_i$, $i = 1, ..., N$ be the decomposition of $\alpha_V: Y \to Y_0$ as in Lemma [6.6](#page-15-1) and let $\kappa_i : Y_i \to V^*$ be the natural projection. Then every an irreducible component L_j of $\kappa_i^{-1}(F^*)$ is of the form $C \times \mathbb{P}^1$ for some multisection C over F^* . Recall that $Y_{i+1} = \text{Bl}_{C_i} Y_i$ for a multisection C_i over F^* where either (a) $C_i = L_j \cap L_l$, $j, l < i$ or (b) C_i is contained in L_j only. Let $\hat{L}_j = \psi_i^{-1}(L_j)$ where $\psi_i : \hat{Y}_i = Y_i \times_{V^*} \hat{V}^* \to Y_i$ is the natural projection. Since $\varphi(R) = o$ we have $\hat{L}_j \simeq B \times (\kappa_i^{-1}(o) \cap L_j)$ which is the disjoint union of irreducible components isomorphic to $B \times \mathbb{P}^1$. Similarly, $\hat{C}_i = \psi_i^{-1}(C_i)$ is naturally isomorphic to $B \times K_i$ where $K_i = C_i \cap \kappa_i^{-1}(o)$ is a finite set. We can suppose that F^* is a principal divisor in V^* given by the zeros of a function f on V^* . We use the symbols f_i to denote the lifts of f to Y_i and we let $\hat{f}_i = \psi_i^* f_i$. The transversality of the intersection of F^* and the curves $\varphi(b \times \mathbb{P}^1)$ implies that \hat{f}_0 vanishes on \hat{L}_0 with multiplicity 1. This implies in turn that the scheme ψ_0^* $_{0}^{\ast}(C_{0})$ coincides with the reduced variety \hat{C}_0 . Hence, [\[Har04,](#page-21-10) Chapter II, Corollary 7.15] implies that $\hat{Y}_1 = \text{Bl}_{\psi_i^*(C_0)} \hat{Y}_0 = \text{Bl}_{\hat{C}_0} \hat{Y}_0$ where the latter variety admits a rational map $\text{Bl}_{\hat{C}_0}\hat{Y}_0 \dashrightarrow B \times \text{Bl}_{b_0 \times K_0} \hat{Y}_0^{b_0}$ over the rational map $\hat{Y}_0 \dashrightarrow B \times \hat{Y}_0^{b_0}$ that is biregular over U. Note also that $\hat{Y}_1^{b_0} = \text{Bl}_{b_0 \times K_0} \hat{Y}_0^{b_0}$ is a smooth surface. Assume that we constructed a rational map $\hat{Y}_i \dashrightarrow B \times Bl_{b_0 \times K_i} \hat{Y}_i^{b_0}$ which is biregular over U with $\hat{Y}_i^{b_0}$ being a smooth surface and that for every $j \leq i$ the zero multiplicity m_j of f_i on L_j coincides with the zero multiplicity of \hat{f}_i on \hat{L}_j . Hence, for a regular function g on Y_i vanishing on L_j with multiplcity 1 the function ψ_i^* i ^{*}_{*i*} g vanishes on \hat{L}_j with multiplicity 1. The latter implies that the scheme ψ_i^* $i(C_i)$ coincides with the reduced variety \hat{C}_i . This yields again a rational map $\hat{Y}_{i+1} \dashrightarrow B \times \text{Bl}_{b_0 \times K_i} \hat{Y}_i^{b_0}$ with the desired properties. Since C_i (resp. \hat{C}_i) is the transversal intersection of L_j and L_l (resp. \hat{L}_j and \hat{L}_l) in case (a) the zero multiplicities of f_{i+1} on L_{i+1} and \hat{f}_{i+1} on \hat{L}_{i+1} are both equal to $m_j + m_l$, whereas in case (b) they are equal to m_j . Thus, the equality of the multiplicities is preserved and we are done by induction.

The morphism $\alpha: X \to Q$ is invertible over $S \setminus T$ where T is as in Lemma [6.5.](#page-15-0) In particular, α is invertible over $V^* \setminus T = V^* \setminus F^*$. Since $\varphi^{-1}(V^* \setminus F^*) = \hat{V}^* \setminus R \supset U \setminus R$ we see that $\hat{\alpha}_V$ is invertible over $U \setminus R$ which implies an isomorphism $\hat{\varrho}_V^{-1}$ $_V^{-1}(U \setminus R) \simeq$ $\hat{\mathrm{pr}}^{-1}(U \setminus R)$ over $U \setminus R$.

Proposition 6.11. Let Z be the variety obtained by the natural gluing of $Z_1 = \hat{\varrho}_V^{-1}$ $V^{-1}(U)$ and $Z_2 = \hat{Q} \setminus \hat{\mathrm{pr}}^{-1}(R)$ via the isomorphism $\hat{\varrho}_V^{-1}$ $V_V^{-1}(U \setminus R) \simeq \hat{\mathrm{pr}}^{-1}(U \setminus R)$. Then there is a natural projection $\eta: Z \to B \times \mathbb{P}^1$ such that Z is isomorphic to $B \times Z^{b_0}$ over $\hat{Q} \simeq B \times \hat{Q}^{b_0}$ where $Z^b = \eta^{-1}(b \times \mathbb{P}^1)$.

Proof. Let η_1 be the restriction of $\hat{\varrho}_V$ to Z_1 and η_2 be the restriction of \hat{pr} to Z_2 . The isomorphism $\hat{\varrho}_V^{-1}$ $V^{-1}(U \setminus R) \simeq \hat{\mathrm{pr}}^{-1}(U \setminus R)$ implies that these maps agree over $U \setminus R$ and, thus, we have $\eta: Z \to B \times \mathbb{P}^1$. Note that $Z_1^{b_0} = Z^{b_0} \cap Z_1$ is naturally isomorphism to \hat{Y}^{b_0} , whereas $Z_2^{b_0} = Z^{b_0} \cap Z_2$ is naturally isomorphic to $\hat{Q}^{b_0} \setminus \hat{\mathrm{pr}}^{-1}(R)$. The rational maps from Lemmas [6.9](#page-17-0) and [6.10](#page-17-1) satisfy the following commutative diagram

$$
\hat{Y} \xrightarrow{\nu} B \times \hat{Y}^{b_0}
$$
\n
$$
\downarrow^{\hat{\alpha}_V} \qquad \qquad \downarrow^{(\mathrm{id}, \hat{\alpha}_V^{b_0})}
$$
\n
$$
\hat{Y}_0 \xrightarrow{\nu_0} B \times \hat{Y}_0^{b_0}
$$

where $\hat{\alpha}_V^{b_0}$ \hat{Y}^{b_0} : $\hat{Y}^{b_0} \to \hat{Y}_0^{b_0}$ is the restriction of $\hat{\alpha}_V$, the horizontal arrows are biregular over U, and the vertical arrows are biregular over $U \setminus R$. The restriction of ν_0 over $U \setminus R$ extends to an isomorphism between $Z_2 = \hat{Q} \setminus \hat{\text{pr}}^{-1}(R)$ and $B \times Z_2^{b_0}$ b_0 over $B \times \mathbb{P}^1$. The restriction of ν over U yields an embedding of Z_1 into $B \times \hat{Y}^{b_0} = B \times Z_1^{b_0}$ over $B \times \mathbb{P}^1$. The commutativity of the diagram implies now that this isomorphism and this embedding agree over $U \backslash R$. This yields the desired isomorphism $Z \simeq B \times (Z_1^{b_0} \cup Z_2^{b_0})$ $\binom{b_0}{2} = B \times Z^{b_0}.$ \Box

Let $\hat{X}, \hat{D}, \hat{\varrho}: \hat{X} \to B \times \mathbb{P}^1$, and $\hat{\alpha}: \hat{X} \to \hat{Q}$ be as in Notation [6.7.](#page-16-1) Since φ is invertible over a neighborhood of s_0 (see Lemma [6.5\)](#page-15-0) we see that the natural projection $\hat{X} \to X$ is invertible over v and, thus, the preimage of v is a singleton $\hat{v} \in \hat{X}$.

Lemma 6.12. Let the notations of Proposition [6.11](#page-17-2) hold. Then there is a rational map $\gamma : \hat{X} \dashrightarrow Z$ regular over Z for which

(i) the restriction of γ over U is biregular;

(ii) $M = \gamma(\hat{D} \setminus \hat{g}^{-1}(R))$ is a codimension 2 subvariety in Z such that for every $b \in B$ the set $M_b = Z^b \cap M$ is finite;

(iii) γ is biregular over $Z \setminus M$ (in particular, γ is biregular over a neighborhood of $\gamma(\hat{v})$).

Proof. Consider the restriction γ_1 of $\hat{\alpha}$ over $(B \times \mathbb{P}^1) \setminus R$ and the morphism γ_2 : $\hat{\varrho}^{-1}(U) \rightarrow \hat{\varrho}_V^{-1}$ $\overline{V}_V^{-1}(U)$ induced by the isomorphism $\varrho^{-1}(U) \to \varrho_V^{-1}$ $V_V^{-1}(U)$ (that is, γ_2 is the identity isomorphism). Since $\alpha^{-1}(S \setminus T)$ is naturally isomorphic to ${\rm pr}^{-1}(S \setminus T)$ we see that γ_1 and γ_2 agree over $U \setminus R$ which yields $\gamma : \hat{X} \dashrightarrow Z$ and statement (i). By Lemma [6.5](#page-15-0) φ is invertible over a neighborhood W of s_0 in S and by Lemma [6.4](#page-14-0) (ii) $\varrho^{-1}(W)$ does not meet D. Hence, \hat{D} does not meet $\hat{\varrho}^{-1}(\varphi^{-1}(W))$. Without loss of generality we can suppose that $\varphi(b \times \mathbb{P}^1)$ meets W for every $b \in B$. Then the set M_b is contained in the union of a finite number of fibers of the restriction $\eta_b: Z^b \to \mathbb{P}^1$ of $\eta: Z \to B \times \mathbb{P}^1$. Furthermore, by Lemma [6.4](#page-14-0) (i) M_b cannot contain a fiber of η_b . Hence, we have statement (ii). Since $\hat{\alpha}$ and γ coincide over $(B \times \mathbb{P}^1) \setminus R$ statement (i) implies that γ is biregular on the complement of $\hat{D} \setminus \hat{\varrho}^{-1}(R)$ in $\gamma^{-1}(Z)$. Since $v \notin D$ we have $\hat{v} \notin \hat{D}$ which concludes (iii) and the proof.

Proposition 6.13. Let D be a nontrivial line bundle on a smooth complete rational variety X and $v \in X$ be a point that possesses a neighborhood isomorphic to an open

subset of \mathbb{A}^n . Let $\pi : \tilde{X} \to X$ be a morphism from a smooth rational variety \tilde{X} locally invertible over a neighborhood of v, $\tilde{\mathcal{D}} = \pi^* \mathcal{D}$, and $\tilde{v} = \pi^{-1}(v)$. Suppose that for every triple (X, \mathcal{D}, v) as above the triple $(\tilde{X}, \tilde{\mathcal{D}}, \tilde{v})$ can be chosen so that P-property is valid for it whenever dim $X = n - 1$. Then the same remains true if dim $X = n$.

Proof. Since v has a neighborhood isomorphic to \mathbb{A}^n we can suppose that the conclu-sions of Lemma [6.4](#page-14-0) hold. Consider the triple (S, \mathcal{F}, s_0) where $\mathcal{F} = \mathcal{O}_S(F)$. By the assumption there exists a morphism $\pi : \tilde{S} \to S$ invertible over a neighborhood of s_0 such that P-property holds for the triple $(\tilde{S}, \tilde{\mathcal{F}}, \tilde{s}_0)$ where $\tilde{\mathcal{F}} = \pi^* \mathcal{F}$ and $\tilde{s}_0 = \pi^{-1}(s_0)$. Let $\tilde{X} = X \times_S \tilde{S}$, $\tilde{\mathcal{D}}$ be the lift of \mathcal{D} to \tilde{X} and \tilde{v} be the preimage of v in \tilde{X} . It suffices to prove P-property for $(X, \mathcal{D}, \tilde{v})$. Changing notations we can suppose that $(X, \mathcal{D}, v) = (\tilde{X}, \tilde{\mathcal{D}}, \tilde{v})$. Let $\varphi : B \times \mathbb{P}^1 \to S$ and $o \in F$ be as in Lemma [6.5,](#page-15-0) $b_0 \in B$ be such that $s_0 \in \varphi(b_0 \times \mathbb{P}^1)$ and, thus, $v \in \varrho^{-1}(\varphi(b_0 \times \mathbb{P}^1))$. Let the notations of Lemma [6.12](#page-18-0) hold and $\hat{D}' = \kappa^{-1}(\text{supp } D')$ where $\mathcal{D} = \mathcal{O}_X(D')$ and $\kappa : \hat{X} \to X$ is the natural projection. Then $\gamma(\hat{v})$ is contained in Z^{b_0} . By Proposition [5.9](#page-11-2) there is an irreducible component E_0 of $\gamma(\hat{D}') \cap Z^{b_0}$ for which one can find a birational morphism $\psi_0: B_1 \times \mathbb{P}^1 \to Z^{b_0}$ invertible over a neighborhood of $\gamma(\hat{v})$ in Z^{b_0} such that $\psi_0(B_1 \times \mathbb{P}^1) \cap M_{b_0} = \emptyset$, the set $\psi_0(B_1 \times \mathbb{P}^1) \cap \gamma(\hat{D}')$ is a general point o_1 of E_0 , and every curve $\psi_0(b_1 \times \mathbb{P}^1), b_1 \in B_1$ intersects E_0 transversely at o_1 . By Propo-sition [6.11](#page-17-2) there is an isomorphism $\tau : B \times Z^{b_0} \to Z$. Note that the morphism $\psi: B \times B_1 \times \mathbb{P}^1 \to Z$, $(b, b_1, t) \mapsto \tau(b, \psi_0(b_1, t))$ is locally invertible over a neighborhood of $\gamma(\hat{v})$ in Z. Let $b_1^0 \in B_1$ be such that $\gamma(\hat{v}) \in \psi(b_0 \times b_1^0 \times \mathbb{P}^1)$. Replacing $B \times B_1$ with an neighborhood \check{B} of (b_0, b_1^0) we can suppose that $\psi(\check{B} \times \mathbb{P}^1)$ does not meet M. Then every curve $\psi(\breve{b} \times \mathbb{P}^1) \subset Z^b$, $\breve{b} = (b, b_1) \in \breve{B}$ intersects $\gamma(\hat{D}')$ at the point $o_1^b = \tau((b, o_1))$ only with the intersection being transversal. Furthermore, by Lemma 6.12 (iii) one has a lift $\check{\psi}: \check{B} \times \mathbb{P}^1 \to \hat{X}$ of ψ invertible over \hat{v} such that every curve $\check{\psi}(\check{b} \times \mathbb{P}^1)$ intersects \hat{D}' at $\gamma^{-1}(o_1^b)$ only with the intersection being transversal and $\gamma^{-1}(o_1^b)$ contained in the irreducible component $\hat{H} = \gamma^{-1}(\tau(B \times E_0))$ of \hat{D}' . Note that $\hat{H} = \kappa^{-1}(H)$ where H is an irreducible component D' such that $\varrho(H) = F$ and $\breve{\sigma} = \kappa(\tau((b_0, o_1)))$ is a general point of H located above $o \in F$. The morphism $\chi = \kappa \circ \check{\psi} : \check{B} \times \mathbb{P}^1 \to X$ is locally invertible over $v, \chi(\check{B} \times \mathbb{P}^1) \cap \text{supp } D' = \check{o}$, and every curve $\chi(\hat{b} \times \mathbb{P}^1)$ intersects H at δ transversely. This concludes the proof.

Proof of Theorem [6.1.](#page-13-0) By Proposition [5.9](#page-11-2) and Lemma [5.7](#page-10-1) for a smooth projective surface X and a triple (X, \mathcal{D}, v) one can find a triple $(X, \mathcal{D}, \tilde{v})$ as in Proposition [6.13](#page-18-1) which satisfies the P -property. Proposition 6.13 and induction imply that the same is true when X is a complete uniformly rational variety of any dimension. Thus, we can suppose that there is a birational morphism $\varphi : B \times \mathbb{P}^1 \to X$ invertible over a neighborhood of v such that every curve $\varphi(b \times \mathbb{P}^1)$, $b \in B$ meets only one irreducible component of D' . Hence, the restriction of D to any of these curves is a nontrivial line bundle. Every complete uniformly rational variety is elliptic by Theorem [1.1.](#page-0-0) Now Corollary [2.5](#page-2-2) implies the desired conclusion.

While discussing principal \mathbb{G}_m -bundles we suppose a priori that they are locally trivial in the étale topology.

Corollary 6.14. Let X be a complete uniformly rational variety and $\beta: Y \to X$ be a nontrivial principal \mathbb{G}_m -bundle. Then Y is elliptic.

Proof. The morphism β extends to a proper morphism $\bar{Y} \to X$ which is a \mathbb{P}^1 -bundle such that the divisor $G = \overline{Y} \setminus Y$ is an unramified two-sheeted covering of X. If G consists of two sections, then Y can be viewed as the complement to a zero section of a nontrivial line bundle over X which implies ellipticity by Theorem [6.1.](#page-13-0) It remains to establish the following (certainly well-known) fact that G cannot be irreducible for a smooth complete rational variety X (that is, $\beta: Y \to X$ is locally trivial in the Zariski topology). Let us start first with the case when the ground field is C. Consider a birational map $\mathbb{P}^n \dashrightarrow X$. Resolving the indeterminacy set $P \subset \mathbb{P}^n$ of this map one gets a smooth complete rational variety W with birational morphisms $\eta: W \to \mathbb{P}^n$ and $\zeta: W \to X$. Every element $[\delta]$ of the fundamental group $\pi_1(W)$ can be presented by a differentiable map $\delta: S^1 \to W$ from a circle S^1 whose image does not meet $\eta^{-1}(P)$. Since $\pi_1(\mathbb{P}^n) = 0$ the map $\eta \circ \delta : S^1 \to \mathbb{P}^n$ can be extended to a differentiable map $\gamma : \Delta \to \mathbb{P}^n$ from a closed disc Δ with the boundary S^1 . By the transversality argument one can suppose that $\gamma(\Delta)$ does not meet P since the latter is of the complex codimension at least 2. Hence, γ admits a lift $\Delta \to W$ which shows that $[\delta]$ is the zero element of $\pi_1(W)$. Thus, $\pi_1(W) = 0$. Every element $[\sigma]$ of the fundamental group $\pi_1(X)$ can be presented by a differentiable map $\sigma : S^1 \to X$ whose image does not meet the indeterminacy set of ζ^{-1} . Hence, $[\sigma]$ belongs to the image of $\pi_1(W)$ and, thus, $\pi_1(X) = 0$. On the other hand, for G to be irreducible one needs a nontrivial $\pi_1(X)$ which concludes the case of complex varieties. Now the standard application of the Lefschetz principle implies that G cannot be irreducible in the case of a general ground field **k** which concludes the proof. \Box

M. Zaidenberg suggested the following strengthened version of Question 1 in Introducation to which the author does not know the answer.

Question 2 (Zaidenberg). Let X be a smooth complete elliptic variety that verifies the strengthened curve-orbit property and $\mathcal{D} \to X$ be a nontrivial line bundle with the zero section $Z_{\mathcal{D}}$. Is $Y = \mathcal{D} \setminus Z_{\mathcal{D}}$ elliptic?

It is worth mentioning that there are non-rational unirational varieties that satisfy the strengthened curve-orbit property (see $KZ24$).

Acknowledgements. The author is grateful to M. Zaidenberg for useful consultations.

REFERENCES

[AKZ23] I. V. Arzhantsev, Sh. Kaliman, and M. G. Zaidenberg, Varieties covered by affine spaces, uniformly rational varieties and their cones, Adv. Math. 437 (2024), Paper No. 109449, 18 pp.

- [BB14] F. Bogomolov and C. Böhning, On uniformly rational varieties. In: Topology, geometry, integrable systems, and mathematical physics, 33–48. Amer. Math. Soc. Transl. Ser. 2, 234. Amer. Math. Soc., 2014.
- [CPPZ21] I. Cheltsov, J. Park, Yu. Prokhorov, and M. Zaidenberg, Cylinders in Fano varieties, EMS Surv. Math. Sci. 8 (2021), 39–105, DOI 10.4171/EMSS/44.
- [DaSh94] V.I. Danilov, V.V. Shokurov, Algebraic Geometry I: Algebraic Curves, Algebraic Manifolds and Schemes, (Encyclopaedia of mathematical sciences, 23) Springer, 1994.
- [For06] F. Forstnerič, Holomorphic flexibility properties of complex manifolds. Amer. J. Math. 128:1 (2006), 239–270.
- [For17] F. Forstnerič, Stein Manifolds and Holomorphic Mapping. The Homotopy Principle in Complex Analysis, Springer-Verlag, Berlin-Heidelberg, 2011.
- [For23] F. Forstnerič, Recent developments on Oka manifolds, Indag. Math., New Ser. 34(2) (2023), 367–417.
- [Gro89] M. Gromov, Oka's principle for holomorphic sections of elliptic bundles, J. Amer. Math. Soc. 2 (1989), 851–897.
- [Har04] R. Hartshorne, Algebraic Geometry. Graduate Texts in Mathematics 52, Springer 2004.
- [KZ23a] S. Kaliman and M. Zaidenberg, Gromov ellipticity and subellipticity, Forum Math. 36 (2024), no. 2, 373-376.
- [KZ23b] S. Kaliman and M. Zaidenberg, Gromov ellipticity of cones over projective manifolds, arXiv:2303.02036 (2023).
- [KZ24] S. Kaliman and M. Zaidenberg, Algebraic Gromov's ellipticity of cubic hypersurfaces, arXiv:2402.04462 (2024).
- [Ray72] M. Raynaud, Flat modules in algebraic geometry, Compositio Math. 24 (1972), 11-31.

Department of Mathematics, University of Miami, Coral Gables, FL 33124, USA $\emph{Email address: }$ kaliman@math.miami.edu