THE CODEGREE ISOMORPHISM PROBLEM FOR FINITE SIMPLE GROUPS II

NGUYEN N. HUNG AND ALEXANDER MORETÓ

ABSTRACT. Let H be a nonabelian finite simple group. Huppert's conjecture asserts that if G is a finite group with the same set of complex character degrees as H, then $G \cong H \times A$ for some abelian group A. Over the past two decades, several specific cases of this conjecture have been addressed. Recently, attention has shifted to the analogous conjecture for character codegrees: if G has the same set of character codegrees as H, then $G \cong H$. Unfortunately, both problems have primarily been examined on a case-by-case basis. In this paper and the companion [HM22], we present a more unified approach to the codegree conjecture and confirm it for several families of simple groups.

1. INTRODUCTION

This paper is a sequel to our previous work [HM22], in which we investigated the codegree isomorphism problem for finite simple groups.

Let G be a finite group. The codegree of a character χ of G is defined as

$$\operatorname{cod}(\chi) := |G: \ker(\chi)|/\chi(1).$$

The set of all the codegrees of irreducible characters of G is denoted by cod(G), and the corresponding multiset of the codegrees with multiplicity is denoted by C(G). This C(G) is called the group pseudo-algebra of G. We proved in [HM22] that a finite group G and a simple group H are isomorphic if and only if they have the same group pseudo-algebra. In this paper, we drop the multiplicity and turn our attention to the so-called *codegree isomorphism conjecture* (see Problem 20.79 of the Kourovka Notebook [Kou20]), which says that if H is a nonabelian simple group and G is a

²⁰²⁰ Mathematics Subject Classification. Primary 20C15, 20C30, 20C33, 20D06.

Key words and phrases. Character codegrees, isomorphism problem, finite simple groups, Huppert's conjecture.

We are indebted to Pham Huu Tiep for several ideas in the proof of Theorem D for classical groups. We also thank him and Bob Guralnick for helpful comments on group cohomology and Eamonn O'Brien for checking some cases of Hypothesis 5.4. Part of this work was done while the first author was visiting the Vietnam Institute for Advanced Study in Mathematics (VIASM), and he thanks the VIASM for the financial support. The second author is supported by Ministerio de Ciencia e Innovación (Grant PID2022-137612NB-I00 funded by MCIN/AEI/10.13039/501100011033 and "ERDF A way of making Europe"). He also acknowledges support by Generalitat Valenciana CIAICO/2021/163.

finite group with cod(G) = cod(H) then $G \cong H$. In fact, we propose a stronger version.

Conjecture A. Let H be a finite nonabelian simple group and G a finite group. Then $cod(G) \subseteq cod(H)$ if and only if $G \cong H$.

The original conjecture has been confirmed for several sporadic simple groups, Suzuki groups, and linear groups in dimensions 2 and 3, see [BAK21, Aha22, GKL+22, LY23]. However, the approach in these papers has been primarily case-by-case, employing ad-hoc arguments for each specific group (in the case of sporadic groups) or family of groups (in the case of Suzuki and linear groups).

We pursue here (and in the companion paper [HM22]) a more uniform approach. In particular, we prove in Theorem 3.1 that, except possibly for odd-characteristic orthogonal groups in even dimension, Conjecture A follows from the following.

Conjecture B. Let H be a nonabelian simple group. Then, for every perfect group G with an elementary abelian normal p-subgroup N for some prime p dividing |H| such that G/N = H and N is a faithful irreducible H-module, there exists a faithful irreducible character χ of G such that $\chi(1)_p < |N|$.

This path allows us to confirm Conjecture A in several cases.

Theorem C. Both Conjectures A and B hold when H is an alternating group, a sporadic simple group, or a simple group of the following Lie types: ${}^{2}B_{2}$, ${}^{2}G_{2}$, ${}^{2}F_{4}$, G_{2} , ${}^{3}D_{4}$, F_{4} , ${}^{2}E_{6}$, $A_{\leq 3}$, ${}^{2}A_{\leq 6}$, C_{2} , C_{3} , and B_{3} .

When H is a simple group of Lie type, Conjecture B can be reduced further to the situation where the prime p is the defining characteristic of H, see Proposition 4.3.

Conjecture B is related to questions on group cohomology and p-groups that we will discuss in more detail in Section 5. This approach appears particularly promising and we hope that it will be studied further by experts in the field.

Our approach requires investigating the character codegrees of simple (and even quasisimple) groups. The first step is to demonstrate that if S and H are finite simple groups such that $\operatorname{cod}(S) \subseteq \operatorname{cod}(H)$, then $S \cong H$. This was achieved in [HM22], utilizing some deep results in the representation theory of finite simple groups, such as the classification of prime-power-degree representations, bounds for the largest degree of irreducible representations, and the existence of p-defect zero characters.

Another major step is to show that the codegree set of a nonabelian simple group is different from that of a nontrivial perfect central cover of the group. This turns out to be a daunting task. The following result might be helpful in future attempts to establish Huppert's original conjecture for character degrees in full generality. Several ideas in its proof for the case of classical groups was kindly given to us by Tiep.

Theorem D. Let S be a finite nonabelian simple group and G a quasisimple group with center $\mathbf{Z}(G)$ of prime order such that $G/\mathbf{Z}(G) \cong S$. Suppose that $S \not\cong P\Omega_{2n}^{\pm}(q)$ with $n \ge 4$ and q odd. Then there exists a faithful character $\chi \in Irr(G)$ such that $\chi(1)/|\mathbf{Z}(G)|$ is not a character degree of S. Consequently, $cod(G) \not\equiv cod(S)$.

We expect Theorem D to also hold for even-dimensional orthogonal groups, but the proof appears to be much more complicated, see Section 2 for further comment on this.

The proofs of Theorem D and C are respectively given in Sections 2 and 4. Section 3 shows that Conjecture A follows from Conjecture B. Finally, Section 5 discusses some reductions of Conjecture B to a problem on p-groups or a problem on group cohomology.

We conclude this Introduction by noting that we announced the main results of this paper in Remark 1 of [HM22]. Since then, there has been a number of preprints dealing with the codegree isomorphism conjecture for some families of the groups mentioned in Theorem C, using the results in [HM22]. Nevertheless, the approach in this paper remains original and independent of previous papers on the codegree isomorphism conjecture, except for [HM22]. Given that this paper strongly relies on the complex Theorem B of [HM22], we considered it appropriate to have that paper reviewed before publishing this one. We believe that the cohomological approach discussed in Section 5 is perhaps the best way to address most of the remaining cases.

2. Character degrees of quasisimple groups

The aim of this section is to prove Theorem D. That is, if S be a finite non-abelian simple group and G a quasi-simple group with center $\mathbf{Z}(G)$ of prime order such that $G/\mathbf{Z}(G) \cong S$, then there exists a faithful character $\chi \in \operatorname{Irr}(G)$ such that

$$\chi(1)/|\mathbf{Z}(G)| \notin \mathrm{cd}(S),\tag{1}$$

Here, as usual, cd(S) denotes the set of the degrees of irreducible characters of S. We begin with the easy case of alternating and sporadic groups.

Proposition 2.1. Theorem D holds when S is an alternating group or a sporadic simple group.

Proof. When S is a sporadic group or an alternating group of degree at most 9, the result can be verified using [Atl1].

Assume that $S = A_n$ for $n \ge 10$. In particular, S has a unique nontrivial perfect central cover, which is the Schur double cover, denoted by \widehat{A}_n . Assume, to the contrary, that $\chi(1)/|\mathbf{Z}(G)| \in \operatorname{cd}(S)$ for every faithful character $\chi \in \operatorname{Irr}(G)$.

Note that the so-called basic representation of \widehat{A}_n , the one labeled by the partition (n), has the degree $2^{\lfloor (n-2)/2 \rfloor}$ (see [KT12, p.1774], for instance). Therefore, we obtain that $2^{\lfloor (n-2)/2 \rfloor-1}$ would be a character degree of A_n . This is impossible, because the only possible prime-power character degree of A_n is n-1 (by [BBOO01, Theorem 5.1]), but the equation $2^{\lfloor (n-2)/2 \rfloor-1} = n-1$ returns no solutions.

The proof of Theorem D for groups of Lie type relies on Lusztig's classification of their irreducible characters. We briefly recall some basics that will be needed in the remaining of this section, and refer the reader to [Car85, DM91] for the full account of the theory.

Let $G = \mathbf{G}^F$ be the group of fixed points under a Frobenius endomorphism F of a connected reductive algebraic group \mathbf{G} over an algebraically closed field of characteristic p > 0. Let (\mathbf{G}^*, F^*) be the pair in duality with (\mathbf{G}, F) and let $G^* := \mathbf{G}^{*F^*}$. The set $\operatorname{Irr}(G)$ of irreducible characters of G is partitioned into subsets known as the (rational) Lusztig series $\mathcal{E}(G, s)$ associated to various G^* -conjugacy classes of semisimple elements $s \in G^*$. The series $\mathcal{E}(G, s)$ is defined to be the set of irreducible characters of G occurring in some Deligne-Lusztig character $R^{\mathbf{G}}_{\mathbf{T}}\theta$, where \mathbf{T} is an Fstable maximal torus of \mathbf{G} and $\theta \in \operatorname{Irr}(\mathbf{T}^F)$ such that the geometric conjugacy class of (\mathbf{T}, θ) corresponds to the G^* -conjugacy class containing s.

According to [DM91, Theorem 13.23 and Remark 13.24], for each semisimple element $s \in G^*$, there is a bijection $\chi \leftrightarrow \psi$ between $\mathcal{E}(G, s)$ and $\mathcal{E}(\mathbf{C}_{G^*}(s), 1)$ – the set of the unipotent characters of the centralizer $\mathbf{C}_{G^*}(s)$, such that

$$\chi(1) = |G^*: \mathbf{C}_{G^*}(s)|_{p'}\psi(1).$$
(2)

Remark that, when $\mathbf{C}_{G^*}(s)$ is disconnected, unipotent characters of $\mathbf{C}_{G^*}(s)$, by definition, are precisely those characters in $\operatorname{Irr}(\mathbf{C}_{G^*}(s))$ lying over unipotent characters of the connected component $(\mathbf{C}_{\mathbf{G}^*}(s)^0)^{F^*}$. Once a bijection is fixed, we will say that (s^{G^*}, ψ) is the Jordan decomposition of χ .

In certain cases, the desired character χ satisfying(1) can be chosen to be a semisimple character (see [GM20, Definition 2.6.9]). Every Lusztig series $\mathcal{E}(G, s)$ contains at least one semisimple character. Their degrees are the same and given by

$$|G^*: \mathbf{C}_{G^*}(s)|_{p'}.$$

We record the following well-known lemma that will be useful in producing faithful irreducible characters of finite groups of Lie type.

Lemma 2.2. Assume the above notation. Let $s \in G^*$ be a semisimple element. Assume that $|\mathbf{Z}(G)| = |G^*/(G^*)'|$. Then all characters in $\mathcal{E}(G, s)$ restricts trivially to $\mathbf{Z}(G)$ if and only if $s \in (G^*)'$. More precisely,

- (1) if $s \in (G^*)'$, then all characters in $\mathcal{E}(G, s)$ lie over the trivial character of $\mathbf{Z}(G)$, and
- (2) if $s \notin (G^*)'$, then all characters in $\mathcal{E}(G, s)$ lie over the same nontrivial character of $\mathbf{Z}(G)$.

Proof. See [NT13, Lemma 4.4(ii)] for the 'if' implication, and [Hun24, Lemma 5.8(ii)] for the other implication. The fact that all the characters in one Lusztig series lie over the same character of $\mathbf{Z}(G)$ is from [Mal07, Lemma 2.2].

Proposition 2.3. Theorem *D* holds when *S* is an exceptional group of Lie type.

Proof. We can ignore the case where S has trivial Schur multiplier. When S is one of the groups $\{G_2(3), G_2(4), {}^2E_6(2), Suz(8)\}$, the result can be checked using [Atl1]. It remains to consider

It remains to consider

$$S = E_6(q), {}^2E_6(q), \text{ or } E_7(q),$$

where

$$3 \mid (q-1), 3 \mid (q+1), \text{ or } 2 \mid (q-1),$$

respectively. The only possible nontrivial cover of S is then the universal cover, which turns out to be the corresponding finite reductive group of simply connected type. (Note that ${}^{2}E_{6}(2)$ is already treated above and excluded here as its Schur multiplier is exceptional and isomorphic to product of cyclic groups of order 2,2, and 3.) We denote these quasisimple groups by the usual notation

$$S_{sc} = E_6(q)_{sc}, {}^2E_6(q)_{sc}, \text{ or } E_7(q)_{sc}.$$

Note that the dual group of S_{sc} is the corresponding group of adjoint type, denoted by S_{ad} . By Lemma 2.2, if s is chosen to be outside the commutator subgroup S'_{ad} of S_{ad} , every character χ in the series $\mathcal{E}(S_{sc}, (s))$ lie over a nontrivial character of $\mathbf{Z}(S_{sc})$, and hence is faithful. Certainly, in our situation,

$$|S_{ad}: S'_{ad}| = |\mathbf{Z}(S_{sc})| \in \{2, 3\},\$$

and we may choose the desired semisimple element s of order $|\mathbf{Z}(S_{sc})|$. In such case, we can appeal to Lübeck's webpage [Lüb1] for the information on conjugacy classes of elements of order 2 or 3 in exceptional groups of Lie type and the structure of their centralizers. Once s is chosen, we compute the degree of a semisimple character $\chi_{(s)}$ in the series $\mathcal{E}(S_{sc}, s)$ (note that there may be more than one such a character) by the formula

$$\chi_{(s)}(1) = |S_{ad} : \mathbf{C}_{S_{ad}}(s)|_{p'},$$

where p is the defining characteristic of S. Using [Lüb2], one can check that both $\chi_{(s)}(1)$ and $\chi_{(s)}(1)/|\mathbf{Z}(S_{sc})|$ are not members of $\operatorname{cd}(S_{ad})$, which shows that the latter is not a member of $\operatorname{cd}(S)$, as wanted as in (1).

We provide complete arguments in the case $S = E_7(q)$, where $q = p^n$ is some odd prime power, and skip the details in the remaining cases. We will use cyclotomic polynomials $\Phi_i := \Phi_i(q)$ to express the orders of these centralizers. First assume that $q \equiv 1 \pmod{4}$. Then $E_7(q)_{ad}$ has one involution class (s) outside $E_7(q)$ with centralizer ${}^2A_7(q).2$ of order

$$|\mathbf{C}_{E_7(q)_{ad}}(s)| = 2q^{28}\Phi_1^4\Phi_2^7\Phi_3\Phi_4^2\Phi_6^2\Phi_8\Phi_{10}\Phi_{14}.$$

A semisimple character $\chi_{(s)}$ associated to (s) of $G = E_7(q)_{sc}$ then has the degree

$$\chi_{(s)}(1) = [E_7(q)_{ad} : \mathbf{C}_{E_7(q)_{ad}}(s)]_{p'} = \frac{1}{2} \Phi_1^3 \Phi_3^2 \Phi_5 \Phi_6 \Phi_7 \Phi_9 \Phi_{12} \Phi_{18}.$$

It can be checked that neither $\chi_{(s)}(1)$ nor $\chi_{(s)}(1)/2$ is a character degree of $E_7(q)_{ad}$, so that $\chi_{(s)}(1)/2 \notin \operatorname{cd}(E_7(q))$. Our desired character χ as in (1) now can be chosen to be $\chi_{(s)}$. For the other case $q \equiv -1 \pmod{4}$, $E_7(q)_{ad}$ has one involution class outside $E_7(q)$ with centralizer $A_7(q).2$ and similar arguments apply.

When $S = E_6(q)$ with $3 \mid (q-1)$, the character χ can be taken to be a semisimple character associated to a class $(s) \in E_6(q)_{ad} - E_6(q)$ of elements of order 3 with centralizer $A_2(q^3).3$. Finally, when $H = {}^2E_6(q)$ with q > 2 and $3 \mid (q+1), \chi$ can be chosen to be a semisimple character associated to a class of elements of order 3 with centralizer ${}^2A_2(q^3).3$.

In what follows, $\epsilon \in \{\pm\}$ or $\{\pm1\}$, depending on the context. Also, $PSL_n^+(q)$ stands for the linear groups and $PSU_n^-(q)$ stands for the unitary groups.

Proposition 2.4. Theorem D holds when $S = PSL_n^{\epsilon}(q)$ with $n \ge 2$ or $PSp_{2n}(q)$ with $n \ge 2$.

Proof. We recall that G is a quasisimple group such that $|\mathbf{Z}(G)|$ is of prime order, say r, and $G/\mathbf{Z}(G) \cong S$. We wish to find a faithful $\chi \in \operatorname{Irr}(G)$ such that $\chi(1)/r \notin \operatorname{cd}(S)$.

Consider $S = PSp_{2n}(q)$ with $n \ge 2$. We may assume that q is odd and $G = Sp_{2n}(q)$, so that r = 2. It is well known that G has a faithful Weil character χ of degree either $(q^n + 1)/2$ or $(q^n - 1)/2$. However, $\chi(1)/2 = (q^n \pm 1)/4$ is not a character degree of S, as the minimal nontrivial degree of a projective irreducible character of S is $(q^n - 1)/2$ (see [TZ96, Theorem 5.2]).

Consider $S = \text{PSL}_n^{\epsilon}(q)$. As the alternating groups have been treated in Proposition 2.1 and the cases $S = \text{PSL}_3(4)$, $\text{PSU}_4(2)$, $\text{PSU}_4(3)$, or $\text{PSU}_6(2)$ can be verified directly, we may assume that G is a quotient of $\text{SL}_n^{\epsilon}(q)$. In fact, if $|\mathbf{Z}(G)| = r$, then the cyclic group $\mathbf{Z}(\text{SL}_n^{\epsilon}(q))$, of order $\text{gcd}(n, q - \epsilon)$, has a unique subgroup C of order $\text{gcd}(n, q - \epsilon)/r$, and G must be isomorphic to $\text{SL}_n^{\epsilon}(q)/C$. Now $\text{SL}_n^{\epsilon}(q)$ has an irreducible Weil character χ of degree $\chi(1) = (q^n - \epsilon^n)/(q - \epsilon)$ whose kernel is precisely C, see [Tie15] and [Sze99] for the character formulas of the Weil character of G. An easy inspection from [TZ96, Table II] shows that $\chi(1)/2$ is smaller than the minimal nontrivial character degree of S, and so $\chi(1)/r \notin \text{cd}(S)$, as desired.

We set up some notation for the proof of Theorem D in the case where S is an odd-dimensional orthogonal group.

Let $\langle . | . \rangle$ be a non-degenerate symplectic form on the space $V := \mathbb{F}_q^{2n}$, where q is odd. The *conformal symplectic group* $\mathrm{CSp}_{2n}(q)$ is defined to be

$$\mathrm{CSp}_{2n}(q) := \{ g \in \mathrm{GL}(V) : \exists \tau(g) \in \mathbb{F}_q^*, \langle gu | gv \rangle = \tau(g) \langle u | v \rangle \forall u, v \in V \}.$$

We have $\mathbf{Z}(CSp_{2n}(q)) \cong C_{q-1}$ and the projective conformal symplectic group is

$$\operatorname{PCSp}_{2n}(q) := \operatorname{CSp}_{2n}(q) / \mathbf{Z}(\operatorname{CSp}_{2n}(q)).$$

This is indeed the type- C_n finite reductive group of adjoint type, see [Car85, p. 40]. Set

$$H := \operatorname{CSp}_{2n}(q)$$
 and $G^* := \operatorname{PCSp}_{2n}(q)$.

The group G^* is dual to

$$G := Spin_{2n+1}(q),$$

the type- B_n group of simply connected type, see [Car85, p.120]. Let \mathbf{G}^* and F^* be respectively a simple algebraic group and associated Frobenius map defining G^* , so that $\mathbf{G}^{*F^*} = G^*$.

Now, let s be a semisimple element in G^* . Consider a pre-image of s, which we will denote by the same s, in H. From the aforementioned Jordan decomposition, to control characters of G, we wish to understand the centralizer

$$C^* := \mathbf{C}_{G^*}(s).$$

Note that $\mathbf{C}_{\mathbf{G}^*}(s)$ is not always connected. Also, the F^* -fixed points of the connected component $\mathbf{C}_{\mathbf{G}^*}(s)^0$, which we denote by C^{*0} , is precisely the image of $\mathbf{C}_H(s)$ under the natural projection $\pi: H \to G^*$.

Lemma 2.5. Assume the above notation. Then the index of C^{*0} in C^* is either 1 or 2.

Proof. Let $g \in H$ be such that its image $\pi(g)$ under π belongs to C^* ; that is, $gsg^{-1} = \lambda s$ for some $\lambda \in \mathbb{F}_q^*$. Then, for every $u, v \in V$,

$$\lambda^{2}\tau(s)\langle u|v\rangle = \langle \lambda su|\lambda sv\rangle = \langle gsg^{-1}u|gsg^{-1}v\rangle$$
$$=\tau(g)\tau(s)\tau(g^{-1})\langle u|v\rangle = \tau(s)\langle u,v\rangle,$$

and so $\lambda \in \{\pm 1\}$. Thus $\mathbf{C}_H(s)$ has index 1 or 2 in $\pi^{-1}(C^*)$, proving the lemma. \Box

Remark 2.6. Note that, for certain semisimple element $s \in G^*$, the case $\lambda = -1$ in the proof of Lemma 2.5 does not occur. In such situation, $\mathbf{C}_{\mathbf{G}^*}(s)$ is connected and $\mathbf{C}_H(s)$ is the full pre-image of $C^* = C^{*0}$.

Lemma 2.7. Let q be a power of an odd prime p. Let G^* be the projective conformal symplectic group $\operatorname{PCSp}_{2n}(q)$ with $n \ge 3$ and s a semisimple element of G^* . Then the only possible degrees prime to p of a unipotent character of $\mathbf{C}_{G^*}(s)$ are 1 and 2.

Proof. As above, we denote a pre-image of s under the projection π by the same notation s. By [Ng10, Lemma 2.4], we have

$$\mathbf{C}_{\mathrm{Sp}_{2n}(q)}(s) \cong \mathrm{Sp}_{2k}(q) \times \mathrm{Sp}_{2(m-k)}(q) \times \prod_{i=1}^{t} \mathrm{GL}_{a_i}^{\alpha_i}(q^{k_i}),$$

where $0 \leq k \leq m \leq n$, $\alpha_i = \pm$, and $\sum_{i=1}^t k_i a_i = n - m$, if $\tau(s)$ is a square in \mathbb{F}_q , and

$$\mathbf{C}_{\mathrm{Sp}_{2n}(q)}(s) \cong \mathrm{Sp}_m(q^2) \times \prod_{i=1}^t \mathrm{GL}_{a_i}^{\alpha_i}(q^{k_i}),$$

where *m* is even, $\alpha_i = \pm$, and $\sum_{i=1}^{t} k_i a_i = n - m$, if $\tau(s)$ is not a square in \mathbb{F}_q . By [Mal07, Theorem 6.8] and the assumption on *q*, the only unipotent character of either Sp_(*q*) or SL[±]₋(*q*) of degree prime to *p* is the trivial character. It follows that, the only unipotent characters of $\mathbf{C}_{\text{Sp}_{2n}(q)}(s)$ of degree prime to *p* are linear characters.

Let $C^{*0} := \mathbf{C}_{\mathbf{G}^*}(s)^{0^{F^*}}$, as above. Using [DM91, Proposition 13.20] for the embedding $f : \mathbf{C}_{\mathrm{Sp}_{2n}(q)}(s) \hookrightarrow \mathbf{C}_{\mathrm{CSp}_{2n}(q)}(s)$ and the projection $\pi : \mathbf{C}_{\mathrm{CSp}_{2n}(q)}(s) \to C^{*0}$, we know that the degrees of unipotent characters of $\mathbf{C}_{\mathrm{Sp}_{2n}(q)}(s)$ and C^{*0} are the same. Therefore, the only unipotent characters of C^{*0} of degree prime to p are linear characters.

The result now follows from Lemma 2.5 that C^{*0} has index 1 or 2 in $\mathbf{C}_{G^*}(s)$. \Box

Proposition 2.8. Theorem *D* holds when $S = \Omega_{2n+1}(q)$ with $n \ge 3$.

Proof. The theorem can be easily checked for $\Omega_7(2)$ and $\Omega_7(3)$, those groups with an exceptional Schur multiplier. Also, $\Omega_{2n+1}(2^m) \cong \operatorname{Sp}_{2n}(2^m)$ is already considered in Proposition 2.4. We therefore may assume that $G = Spin_{2n+1}(q)$ with q odd, which is the (double) universal cover of S. Let $G^* := \operatorname{PCSp}_{2n}(q)$ and $H := \operatorname{CSp}_{2n}(q)$, as above. Also, let \mathbf{G}^* and F^* be respectively a simple algebraic group and associated Frobenius map defining G^* , so $\mathbf{G}^{*F^*} = G^*$.

Suppose that $q \equiv \epsilon \in \{\pm 1\} \pmod{4}$. According to [GLS97, Table 4.5.1], G^* contains an involution, say t_{ϵ} , that is not in $[G^*, G^*] = \operatorname{PSp}_{2n}(q)$. Furthermore, if $C^* := \mathbf{C}_{G^*}(t_{\epsilon})$ and $L^* := \mathbf{O}^{p'}(C^*)$ where p is the defining characteristic of S, then

$$L^* := \mathbf{O}^{p'}(C^*) = \mathrm{SL}_n^{\epsilon}(q) / C_{(n,2)} \text{ and } \mathbf{C}_{C^*}(L^*) = C_{q-\epsilon}.$$

Let $\mathbf{C} := \mathbf{C}_{\mathbf{G}^*}(t_{\epsilon})$ and \mathbf{C}^0 be its connected components containing the identity element. We have $\mathbf{C}/\mathbf{C}^0 \cong C_2$ and \mathbf{C}^0 is of type $A_{n-1}T_1$, according to [GLS97, Table 4.3.1]. The *F*-fixed-point group $C^{*0} := \mathbf{C}^{0^F}$ is therefore isomorphic to $A_{n-1}.C_{q-1}$ when $\epsilon = 1$ and ${}^2A_{n-1}.C_{q+1}$ when $\epsilon = -1$. Moreover, $C^*/C^{*0} \cong \mathbf{C}/\mathbf{C}^0 \cong C_2$ (see [GLS97, Theorem 4.2.2(i)]). All together, we have

$$|C^*| = 2q^{\frac{1}{2}n(n-1)} \prod_{i=1}^n (q^i - \epsilon^i).$$

Now, the Lusztig series defined by (t_{ϵ}) contains a (semisimple) character of degree D/2, where

$$D := 2[G^* : C^*]_{p'} = \frac{\prod_{i=1}^n (q^{2i} - 1)}{\prod_{i=1}^n (q^i - \epsilon^i)}$$

Note that, as t_{ϵ} does not belong to $[G^*, G^*]$, this character is faithful, again by Lemma 2.2.

We claim that D/4 is not a character degree of S, and this will complete the proof. Suppose otherwise, i.e., there is $\chi \in Irr(S)$ with $\chi(1) = D/4$. Let $s \in G^* = PCSp_{2n}(q)$ be a semisimple element labeling the Lusztig series containing χ . In fact,

$$s \in [G^*, G^*] = \operatorname{PSp}_{2n}(q),$$

by Lemma 2.2. Note that D/4 is coprime to p. Formula (2) implies that the degree of the unipotent character ψ of $\mathbf{C}_{G^*}(s)$ in Lusztig's Jordan decomposition is coprime to p as well. By Lemma 2.7, we have $\psi(1) = 1$ or 2. In summary, we obtain

$$\chi(1) = \frac{D}{4} = \alpha \cdot [G^* : \mathbf{C}_{G^*}(s)]_{p'}$$

for some $\alpha \in \{1, 2\}$. It follows that

$$|\mathbf{C}_{G^*}(s)|_{p'} = 4\alpha |\mathrm{GL}_n^{\pm}(q)|_{p'}.$$

Again, for notational convenience, we will identify s with a pre-image of s in $\operatorname{Sp}_{2n}(q)$. Observe that

$$|\mathbf{C}_{\mathrm{PCSp}_{2n}(q)}(s)| = \frac{\kappa}{q-1} |\mathbf{C}_{\mathrm{CSp}_{2n}(q)}(s)|$$

for some divisor κ of q-1. In fact, Lemma 2.5 shows that κ is a divisor of 2.

Now, $|\mathbf{C}_{CSp_{2n}(q)}(s)| = (q-1)|\mathbf{C}_{Sp_{2n}(q)}(s)|$, by [Ng10, Lemma 2.4]. It follows that,

$$|\mathbf{C}_{\mathrm{Sp}_{2n}(q)}(s)|_{p'} = \frac{4\alpha}{\kappa} |\mathrm{GL}_n^{\pm}(q)|_{p'}$$

and thus

$$|\mathbf{C}_{\mathrm{Sp}_{2n}(q)}(s)|_{p'} = \{2 \text{ or } 4 \text{ or } 8\} |\mathrm{GL}_n^{\pm}(q)|_{p'}.$$
(3)

Recall from [Ng10, Lemma 2.4] that

$$\mathbf{C}_{\mathrm{Sp}_{2n}(q)}(s) \cong \mathrm{Sp}_{2k}(q) \times \mathrm{Sp}_{2(m-k)}(q) \times \prod_{i=1}^{t} \mathrm{GL}_{a_i}^{\alpha_i}(q^{k_i}),$$

for some $0 \leq k \leq m \leq n$, $\alpha_i = \pm$, and $\sum_{i=1}^t k_i a_i = n - m$.

Suppose first that the group on the right-hand side of (3) is unitary. Suppose furthermore that n is odd. Zsigmondy's theorem then implies that t = 1, $k_1a_1 = n$, and in fact $\mathbf{C}_{\mathrm{Sp}_{2n}(q)}(s) \cong \mathrm{GU}_{a_1}(q^{k_1})$, which in turn implies that $|\mathbf{C}_{\mathrm{Sp}_{2n}(q)}(s)|_{p'} \leq$ $|\mathrm{GU}_n(q)|_{p'}$, violating (3). Let n be even. We see that $q^{n-1} + 1$ is a factor of $|\mathbf{C}_{\mathrm{Sp}_{2n}(q)}(s)|_{p'}$. Using a primitive prime divisor of $q^{2n-2} - 1$, we deduce that either $\mathrm{Sp}_{2n-2}(q)$ or $\mathrm{GU}_{a_i}(q^{k_i})$ for some relevant a_i and k_i with $k_ia_i = n-1$ must be a factor of $\mathbf{C}_{\mathrm{Sp}_{2n}(q)}(s)$. (Note that the case $\mathbf{C}_{\mathrm{Sp}_{2n}(q)}(s) \cong \mathrm{GU}_{a_1}(q^{k_1})$ is already eliminated.) In either case, there is only one other direct factor of $\mathbf{C}_{\mathrm{Sp}_{2n}(q)}(s)$, which must be either $\mathrm{Sp}_2(q)$ or $\mathrm{GL}_1^{\pm}(q)$. It is now straightforward to check that none of these cases can fulfill (3). The case of a linear group in (3) is entirely similar.

Proposition 2.9. Theorem *D* holds when $S = \Omega_{2n}^{\pm}(q)$ with $n \ge 4$ and q even.

Proof. Simple even-dimensional orthogonal groups in even characteristic have trivial Schur multiplier, unless the group is $\Omega_8^+(2)$. The Schur multiplier $\Omega_8^+(2)$ is the Klein four-group, and one can verify the result directly using [Atl1].

We have completed the proof of Theorem D.

The case of even-dimensional orthogonal groups in odd characteristic appears to be complicated. On one hand, the Schur multiplier of S is of order 4, and hence our group G (in Theorem D) is not the entire Schur cover of S. Lemma 2.2 is therefore not enough to produce a desired character. On the other hand, it is much more difficult than the odd-dimensional case to control the centralizer of a semisimple element in the dual group G^* , which in this case is the projective conformal orthogonal group $P(CO_{2n}^{\pm}(q)^0)$.

3. Conjecture **B** implies Conjecture A

In this short section we show that, if Conjecture B holds for a simple group H that is not an even-dimensional orthogonal group in odd characteristic, then Conjecture A holds for H.

Theorem 3.1. Let H be a non-abelian simple group. Suppose that $H \not\cong P\Omega_{2n}^{\pm}(q)$ with q odd and $n \ge 4$. If Conjecture B holds for H, then Conjecture A holds for H.

Proof. Let G be a minimal counterexample to Conjecture A. Since H is perfect, [HM22, Theorem 2.3] implies that G is perfect. Let N be a maximal normal subgroup of G. Since $\operatorname{cod}(G/N) \subseteq \operatorname{cod}(G) \subseteq \operatorname{cod}(H)$, and G/N is simple, it follows from [HM22, Theorem B] that $G/N \cong H$. Furthermore, by the minimality of G as a counterexample, we have that N is a minimal normal subgroup of G. Also, N is the unique minimal normal subgroup of G since, otherwise, $G = H \times H$, violating the assumption $\operatorname{cod}(G) \not \equiv \operatorname{cod}(H)$. In particular, every character in $\operatorname{Irr}(G|N)$ is faithful.

We claim that N is an elementary abelian p-group for some prime p that divides |H|. Suppose that N is not abelian. Then there exists a non-principal character in $\operatorname{Irr}(N)$ that extends to $\chi \in \operatorname{Irr}(G)$ (see [Mor23, Theorem 7.1] for instance). Clearly, $\chi(1) < |N|$. Hence $\operatorname{cod}(\chi) > |G/N| = |H|$, which contradicts $\operatorname{cod}(G) \subseteq \operatorname{cod}(H)$. We have shown that N is an elementary abelian p-group. Let $\pi(X)$ denote the set of prime divisors of |X|. Since $\pi(G) = \pi(\operatorname{cod}(G)) \subseteq \pi(\operatorname{cod}(H)) = \pi(H)$, we deduce that p divides |H|, as claimed.

Note that $\mathbf{C}_G(N) = N$ or $\mathbf{C}_G(N) = G$. Assume first that $\mathbf{C}_G(N) = G$. Then $N = \mathbf{Z}(G)$ and, as G is perfect and N is the unique minimal normal subgroup of G, G must be a quasisimple group with the center $\mathbf{Z}(G)$ of prime order. We are therefore done using Theorem D.

It remains to consider the case $\mathbf{C}_G(N) = N$. In particular, N is a faithful irreducible H-module. By Conjecture **B**, there exists $\chi \in \operatorname{Irr}(G)$ faithful such that $\chi(1)_p < |N|$. Therefore, $\operatorname{cod}(\chi)_p > |G|_p/|N| = |H|_p$. It follows that $\operatorname{cod}(\chi) \notin \operatorname{cod}(H)$, which contradicts the assumption $\operatorname{cod}(G) \subseteq \operatorname{cod}(H)$.

10

Note that if one could prove Theorem D for even-dimensional orthogonal groups, then the above argument also shows that Conjecture B implies Conjecture A for these groups.

4. Theorem C

We now work toward a proof of Conjecture B. The following simple observation will be useful.

Lemma 4.1. Let G be a finite group and $N \leq G$. Let p be a prime divisor of |G/N| and let P/N be a Sylow p-subgroup of G/N.

(1) Suppose that $\theta \in Irr(N)$ is G-invariant. Then there exists $\chi \in Irr(G|\theta)$ such that

$$(\chi(1)/\theta(1))_p \leq \sqrt{|G/N|_p}.$$

(2) Suppose that $\theta \in Irr(N)$ is *P*-invariant. Then there exists $\chi \in Irr(G|\theta)$ such that

$$(\chi(1)/\theta(1))_p \leqslant \sqrt{|G/N|_p}.$$

Proof. We first prove (1). Replacing (G, N, θ) by an isomorphic character triple (see Chapter 11 of [Isa76]), we may assume that N is central and θ is linear. Let $m_p(G)$ be the smallest p-part of the degrees of the characters in $\operatorname{Irr}(G|\theta)$. Since $|G/N| = \sum_{\chi \in \operatorname{Irr}(G|\theta)} \chi(1)^2$ is a multiple of $m_p(G)^2$, we deduce that $m_p(G) \leq \sqrt{|G/N|_p}$, as desired.

For (2), let $T = I_G(\theta)$ - the inertia subgroup of θ in G, so that $P \leq T$. By part (1), there exists $\psi \in \operatorname{Irr}(T|\theta)$ such that

$$(\psi(1)/\theta(1))_p \leqslant \sqrt{|T/N|_p} = \sqrt{|G/N|_p}.$$

Now, by Clifford's correspondence, $\chi = \psi^G \in \operatorname{Irr}(G|\theta)$. Notice that since |G:T| is a p'-number, $\chi(1)_p = \psi(1)_p$, and so

$$(\chi(1)/\theta(1))_p \leq \sqrt{|G/N|_p},$$

as desired.

The following result will allow us confirm Conjecture B in many cases.

Proposition 4.2. Conjecture *B* holds if $|N| > \sqrt{|H|_p}$. In particular, Conjecture *B* holds if $p^{2d} > |H|_p$, where *d* is the smallest degree of the non-trivial irreducible *p*-Brauer characters of *H*.

Proof. Consider the action of G on the p-group $\operatorname{Irr}(N)$. It is clear that there exists $\mathbf{1}_N \neq \lambda \in \operatorname{Irr}(N)$ in an orbit of p'-size. Set $T := I_G(\lambda)$, so that T contains a Sylow p-subgroup of G. By Lemma 4.1, there exists $\chi \in \operatorname{Irr}(G|\lambda)$ such that

$$\chi(1)_p \leqslant \sqrt{|G/N|_p} = \sqrt{|H|_p} < |N|,$$

as wanted.

In order to see the second part, note that since N is a faithful irreducible H-module, dim $N \ge d$. Hence $|N| \ge p^d > \sqrt{|H|_p}$.

The following completes the proof of Theorem C. We will write d(H) to denote the smallest degree of the nontrivial irreducible Brauer characters of H (the prime will be understood from the context).

Proposition 4.3. Let H be a nonabelian simple group. Let G be a perfect group with a normal elementary abelian p-subgroup N for some prime p dividing |H| such that G/N = H and N is a faithful irreducible H-module. Suppose that H belongs to one of the following:

- (1) sporadic groups,
- (2) alternating groups,
- (3) groups of Lie type in characteristic $\ell \neq p$,
- (4) exceptional group of Lie type in characteristic p: ${}^{2}B_{2}(q)$, ${}^{2}G_{2}(q)$, ${}^{2}F_{4}(q)$, $G_{2}(q)$, ${}^{2}D_{4}(q)$, $F_{4}(q)$, ${}^{2}E_{6}(q)$,
- (5) classical groups in characteristic p: $PSL_{\leq 4}(q)$, $PSU_{\leq 7}(q)$, $PSp_4(q)$, $PSp_6(q)$, $P\Omega_7(q)$, $P\Omega_8^{\pm}(q)$.

Then there exists $\chi \in Irr(G)$ faithful such that $\chi(1)_p < |N|$.

Proof. Let d(H) denote the smallest degree of nontrivial irreducible *p*-Brauer characters of *H*. By Proposition 4.2, the result follows for *H* if we are able to prove that

$$p^{2d(H)} > |H|_p.$$

(i) The smallest degrees of the nontrivial p-Brauer characters of the sporadic groups are collected in Table 1 of [Jan05]. It is routine to check that the desired inequality is satisfied.

(ii) We now consider the alternating groups. It is well-known that irreducible $\mathbb{F}_p S_n$ modules are in one-to-one bijection with *p*-regular partitions of *n*. Let λ be such a partition and D^{λ} denote its corresponding module. When the first part of λ is at least n - 4, an explicit lower bound for dim D^{λ} was obtained by James [Jam83]. On the other hand, it was shown by Müller in [Mül16, §6.2] that, for $n \ge 11$ and λ having the first part at most n - 4,

$$\dim D^{\lambda} \ge \min\left\{\frac{n^4 - 14n^3 + 47n^2 - 34n}{24}, g(n)\right\},\$$

where

$$g(n) := \begin{cases} 55 \cdot 2^{(n-11)/2} & \text{if } n \text{ is odd,} \\ 89 \cdot 2^{(n-12)/2} & \text{if } n \text{ is even.} \end{cases}$$

Note that $|\mathsf{A}_n|_p \leq p^{(n-1)/(p-1)}$ for p odd and $|\mathsf{A}_n|_2 \leq 2^{n-2}$. These bounds are sufficient to verify the inequality for alternating groups of degree, of course, $n \geq 11$. The case of smaller n can be checked using [Atl2].

(iii) Next, consider the case where H is a simple group of Lie type in characteristic $\ell \neq p$. A good bound for the smallest degree d(H) of the non-trivial irreducible p-Brauer characters of H was obtained by Landázuri, Seitz, and Zalesskii [LS74, SZ93] (see also [TZ96, Table I]). (Note that this bound does not depend on p.) From this it is easy to check that

$$p^{2d(H)} > |H|_{\ell}$$

for all H. Therefore, we are done if ℓ is a dominant prime in |H| (which means that $|H|_{\ell} \ge |H|_p$ for every p). If otherwise, by [KLST90, Theorem 3.3], H is either $PSL_2(\ell)$ with ℓ a Mersenne prime, $PSL_2(2^a)$ with $2^a + 1$ a Fermat prime, $PSL_2(8)$, or $PSU_3(3)$, in which cases the desired inequality can be easily verified.

(iv) Now, assume that H is a simple group of Lie type in characteristic p. Recall that, By Proposition 4.2, we are done if $|N| > \sqrt{|H|_p}$. Assume by contradiction that

$$|N| \leqslant \sqrt{|H|_p}.$$

Let $\lambda \in \operatorname{Irr}(N) \setminus \{\mathbf{1}_N\}$ and $T := I_G(\lambda)$. Since |G:T| is the number of G-conjugates of λ , we have |G:T| < |N|, and it follows that

$$|G:T| < \sqrt{|H|_p}.$$

If T = G then ker $(\lambda) \leq G$, and thus ker $(\lambda) = 1$, implying that N is cyclic, which cannot happen as H can be embedded into Aut(N). So T is a proper subgroup of G, and thus T/N is a proper subgroup of H = G/N as well. Let M be a maximal subgroup of G/N such that $T/N \subseteq M$. Then

$$|G:T| \ge |H:M|.$$

We now have $|H:M| < \sqrt{|H|_p}$, where M is a certain maximal subgroup of H and p is the defining characteristic of H. Note that H naturally permutes the right cosets of M, so [H:M] is the degree of a permutation representation of H. Minimal degrees of permutation representations of simple groups of Lie type have been worked out by various authors and are available, for example, in [GMPS15, Table 4]. The data shows that, for the listed groups in (iv) and (v), the minimal degree for H always exceeds $\sqrt{|H|_p}$. This contradiction completes the proof.

Theorem C follows from Theorem 3.1, Lemma 4.1, Proposition 4.2, and Proposition 4.3. Since we have proved Conjecture B for D_4 and 2D_4 in Proposition 4.3, Conjecture A for these groups would follow from a proof of Theorem D for them.

5. Further reductions

In this section we discuss some further possible reductions for Conjectures B and A.

The split-extension case of Conjecture B is elementary. Note that the conclusion of the following is much stronger than needed.

Proposition 5.1. Let G = HN, where H is simple non-abelian and N is a normal elementary abelian p-subgroup of G. Suppose that the (conjugation) action of H on N is faithful and irreducible. Then there exists a faithful character $\chi \in Irr(G)$ of degree prime to p.

Proof. Since H acts on the p-group $\operatorname{Irr}(N)$, there exists a non-principal character $\lambda \in \operatorname{Irr}(N)$ in an H-orbit of p'-size. Thus $T := I_G(\lambda)$ contains a Sylow p-subgroup of G. By [Isa76, Problem 6.18], the character λ extends to every Sylow subgroup of G. Now, [Isa76, Corollary 11.31] implies that there exists $\tilde{\lambda} \in \operatorname{Irr}(T|\lambda)$ that extends λ . Hence, by the Clifford correspondence, $\tilde{\lambda}^G$ is a faithful irreducible character of G of p'-degree, as wanted.

Given a simple group H of classical Lie type, we will refer to the natural module of the full cover \tilde{H} of H as the natural module of H. We will use the fact, following from [KL90, Proposition 5.4.11], that if φ is an irreducible Brauer character of Hwith $\varphi(1) > d$, where d is the dimension of the natural module for H, then $\varphi(1) > \log_p \sqrt{|H|_p}$.

Proposition 5.2. Let H be a classical simple group of Lie type over the field with p elements, where p is a prime. Suppose that, for every perfect group G with a normal subgroup N such that

- (1) G/N = H,
- (2) N is an elementary abelian p-group of order equal to the size of the natural module for H,
- (3) N is a faithful irreducible H-module, and
- (4) G is a non-split extension of N by H,

there exists a faithful character $\chi \in Irr(G)$ such that $\chi(1)_p < |N|$. Then Conjecture B holds for H.

Proof. By Proposition 5.1 and the hypothesis, we may assume that the extension is non-split and that |N| exceeds the size of the natural module for H. Since Nis a faithful irreducible \mathbb{F}_pH -module, Corollary 3F of [Won72] implies that N is absolutely irreducible. By hypothesis, we may assume that $\dim_{\mathbb{F}_p} N > d(H)$, which is the dimension of the natural module for H. But then [KL90, Proposition 5.4.11] implies that $|N| > \sqrt{|H|_p}$ and the result follows from Proposition 4.2.

We believe that it should be possible to remove the hypothesis on the order of the field in Proposition 5.2. As pointed out to us by Guralnick and Tiep, the second cohomology group $H^2(G, V)$, where G is a classical group of simply connected type and V is the natural module, is most of the time vanishing (see [Avr78, Dem73, Gri73], for instance). When $H^2(G, V)$ does vanish, Conjecture B would follow from Proposition 5.1. This applies, for example, when $H = PSL_n(2)$ for n > 5, as shown in [Dem73].

Theorem 5.3. Conjecture A holds when $H = PSL_n(2)$ for n > 5.

This provides the first family of groups of Lie type with arbitrarily large rank for which either of the original Huppert's conjecture on character degrees or the codegree counterpart is proved. We expect these ideas to work for other families of simple groups of Lie type with large rank.

We now present another reduction for Conjecture A in the case of classical groups to a problem on p-groups. We remark that, while it is common to reduce problems on finite groups to simple groups, it is unusual to reduce problems concerning groups that are close to simple to p-groups, which lie at the opposite end of the spectrum of finite groups.

Hypothesis 5.4. Let p be a prime and let H be a simple classical group over the field with $q = p^a$ elements. Suppose that Z is a normal (central) subgroup of order p of a p-group P and P/Z is a Sylow p-subgroup of H. Then there exists $\psi \in \operatorname{Irr}(P|Z)$ such that its degree is less than the size of the natural \mathbb{F}_q -module for H.

In the following, given a normal subgroup N of a group G, we write Irr(G|N) to denote the set of characters $\chi \in Irr(G)$ such that N is not contained in the kernel of χ .

Proposition 5.5. Let H be a simple classical group. Suppose that Hypothesis 5.4 holds for H. Then Conjecture B holds for H.

Proof. Recall the setup in Conjecture B that G is a perfect group having a normal elementary abelian p-group N for some prime p dividing |H| such that $G/N \cong H$. We wish to show that there exists a faithful character $\chi \in \operatorname{Irr}(G)$ such that $\chi(1)_p < |N|$. For this purpose, by Proposition 4.3, we may assume that the defining characteristic of H is p.

Let P/N be a Sylow *p*-subgroup of G/N. Let M be a maximal subgroup of N (so M is also elementary abelian whose rank is one less than that of N) such that M is normal in P. By hypothesis, there exists $\psi \in \operatorname{Irr}(P/M|N/M)$ such that $\psi(1)$ is less than the size of the natural \mathbb{F}_q -module V for H. Notice also that $|V| \leq |N|$, so $\psi(1) < |N|$.

Let $\lambda \in \operatorname{Irr}(N)$ be a (non-principal) character lying under ψ (here ψ is viewed as a character of P). Since $\psi^G(1)_p = \psi(1) < |N|$, there exists $\chi \in \operatorname{Irr}(G)$ lying over ψ such that $\chi(1)_p < |N|$. Since χ lies over ψ , χ also lies over λ . We conclude that $\chi \in \operatorname{Irr}(G|N)$, so χ is faithful, as wanted.

E. O'Brien has checked with computer that Hypothesis 5.4 holds when $H = \text{Sp}_8(2)$. In fact, every central extension of a Sylow 2-subgroup P of $\text{Sp}_8(2)$ has a character $\psi \in \text{Irr}(P|Z)$ with $\psi(1) \leq 2^6$. His computations also suggest that Hypothesis 5.4 holds when $H = \text{Sp}_{2n}(2)$ for $5 \leq n \leq 8$, but in these cases there are too many central extensions of the Sylow 2-subgroups. According to these computations, perhaps it is true that there always exists $\psi \in \text{Irr}(P|Z)$ such that its degree is at most 2^{d-2} , where d is the dimension of the natural module for H.

References

- [Aha22] N. Ahanjideh, Nondivisibility among irreducible character co-degrees, Bull. Aust. Math. Soc. 105 (2022), 68–74. 2
- [Avr78] G. S. Avrunin, A vanishing theorem for second degree cohomology, J. Algebra 53 (1978), 382–388. 15
- [BAK21] A. Bahri, Z. Akhlaghi, and B. Khosravi, An analogue of Huppert's conjecture for character codegrees, Bull. Aust. Math. Soc. 104 (2021), 278–286. 2
- [BBOO01] A. Balog, C. Bessenrodt, J. B. Olsson, and K. Ono, Prime power degree representations of the symmetric and alternating groups, J. London Math. Soc. 64 (2001), 344–356. 3
- [Car85] R. W. Carter, Finite groups of Lie type. Conjugacy classes and complex characters, Wiley and Sons, New York et al, 1985, 544 pp. 4, 7
- [Atl1] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker, and R. A. Wilson, Atlas of finite groups, Oxford University Press, London 1985. 3, 5, 10
- [Dem73] U. Dempwolff, On the second cohomology of GL(n, 2), J. Austral. Math. Soc. 16 (1973), 207–209. 15
- [DM91] F. Digne and J. Michel, Representations of finite groups of Lie type, London Mathematical Society Student Texts 21, 1991, 159 pp. 4, 8
- [GM20] M. Geck and G. Malle, *The character theory of finite groups of Lie type: a guided tour*, Cambridge studies in advanced mathematics **187**, 2020, 394 pp. **4**
- [GKL+22] M. Gintz, M. Kortje, M. Laurence, Y. Liu, Z. Wang, and Y. Yang, On the characterization of some non-abelian simple groups with few codegrees, *Comm. Algebra* 50 (2022), 3932–3939. 2
- [GLS97] D. Gorenstein, R. Lyons, and R. Solomon, The classification of the finite simple groups, Number 3, Part 1, American Mathematical Society, 1997. 8
- [Gri73] R. L. Griess, Automorphisms of extra special groups and nonvanishing degree 2 cohomology *Pacific J. Math.* 48 (1973), 403–422. 15
- [GMPS15] S. Guest, J. Morris, C. E. Praeger, and P. Spiga, On the maximum orders of elements of finite almost simple groups and primitive permutation groups, *Trans. Amer. Math.* Soc. 367 (2015), 7665–7694. 13
- [Hun24] N. N. Hung, The continuity of *p*-rationality and a lower bound for *p'*-degree irreducible characters of finite groups, *Trans. Amer. Math. Soc.* **377** (2024), 323–344. 4
- [HM22] N. N. Hung and A. Moretó, The codegree isomorphism problem for finite simple groups, Quarterly J. Math., to appear, arXiv:2301.00446. 1, 2, 3, 10

- [Isa76] I. M. Isaacs, Character theory of finite groups, Academic Press, 1976. 11, 14
- [Jam83] G. D. James, On the minimal dimensions of irreducible representations of symmetric groups, Math. Proc. Cambridge Philos. Soc. 94 (1983), 417–424. 12
- [Jan05] C. Jansen, The minimal degrees of faithful representations of the sporadic simple groups and their covering groups, LMS J. of Computation and Mathematics 8 (2005), 122–144. 12
- [Atl2] C. Jansen, K. Lux, R. Parker, and R. Wilson, An atlas of Brauer characters. Appendix 2 by T. Breuer and S. Norton. London Mathematical Society Monographs 11, Oxford Science Publications, The Clarendon Press, Oxford University Press, New York, 1995, 327 pp. 13
- [KLST90] W. Kimmerle, R. Lyons, R. Sandling, D. N. Teague, Composition factors from the group ring and Artin's theorem on orders of simple groups, *Proc. London Math. Soc.* 60 (1990), 89–122. 13
- [KL90] P. Kleidman and M. Liebeck, The subgroups structure of the finite classical groups. Cambridge University Press, Cambridge, 1990. 14, 15
- [KT12] A. S. Kleshchev and P. H. Tiep, Small-dimensional projective representations of symmetric and alternating groups, Algebra Number Theory 6 (2012), 1773–1816. 3
- [Kou20] E. I. Khukhro and V. D. Mazurov, The Kourovka notebook: Unsolved problems in group theory, **20**th edition. https://kourovka-notebook.org. **1**
- [LS74] V. Landázuri and G. Seitz, On the minimal degrees of projective representations of the finite Chevalley groups, J. Algebra 32 (1974), 418–443. 13
- [LY23] Y. Liu and Y. Yang, Huppert's analogue conjecture for PSL(3, q) and PSU(3, q), Results Math. 78, article number 7, (2023). 2
- [Lüb1] F. Lübeck, Elements of order 2 and 3 in exceptional groups of Lie type, available at http://www.math.rwth-aachen.de/ Frank.Luebeck/ chev/23elts.html. 5
- [Lüb2] F. Lübeck, Character Degrees and their Multiplicities for some Groups of Lie Type of Rank < 9, available at http://www.math.rwthaachen.de/ Frank.Luebeck/chev/DegMult/index.html?LANG=en. 5
- [Mal07] G. Malle, Height 0 characters of finite groups of Lie type, *Represent. Theory* **11** (2007), 192–220. **4**, **8**
- [Mor23] A. Moretó, Methods and questions in character degrees of finite groups, Vietnam J. Math. 51 (2023), 685–701. 10
- [Mül16] J. Müller, On low-degree representations of the symmetric group, J. Algebra 465 (2016), 356–398. 12
- [NT13] G. Navarro and P. H. Tiep, Characters of relative p'-degree over normal subgroups, Ann. of Math. 178 (2013), 1135–1171. 4
- [Ng10] H. N. Nguyen, Low-dimensional complex characters of the symplectic and orthogonal groups, Comm. Algebra 38 (2010), 1157–1197. 7, 9
- [SZ93] G. M. Seitz and A. E. Zalesskii, On the minimal degrees of projective representations of the finite Chevalley groups, II, J. Algebra 158 (1993), 233–243. 13
- [Sze99] F. Szechtman, Weil representations of unitary groups, J. Algebra 221 (1999), 161–187.
 6
- [Tie15] P.H. Tiep, Weil representations of finite general linear groups and finite special linear groups, *Pacific J. Math.* 279 (2015), 481–498. 6
- [TZ96] P. H. Tiep and A. E. Zalesskii, Minimal characters of the finite classical groups, Comm. Algebra 24 (1996), 2093–2167. 6, 13
- [Won72] W. Wong, Representations of Chevalley groups in characteristic p, Nagoya J. Math. 45 (1972), 39–78 14

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF AKRON, AKRON, OH 44325, USA $\mathit{Email}\ address:\ hungnguyen@uakron.edu$

Departamento de Matemáticas, Universidad de Valencia, 46100 Burjassot, Valencia, Spain

Email address: alexander.moreto@uv.es

18