

P. JONES' INTERPOLATION THEOREM FOR NONCOMMUTATIVE MARTINGALE HARDY SPACES II

NARCISSE RANDRIANANTOANINA

ABSTRACT. Let \mathcal{M} be a semifinite von Neumann algebra equipped with an increasing filtration $(\mathcal{M}_n)_{n \geq 1}$ of (semifinite) von Neumann subalgebras of \mathcal{M} . For $1 \leq p \leq \infty$, let $\mathcal{H}_p^c(\mathcal{M})$ denote the noncommutative column martingale Hardy space constructed from column square functions associated with the filtration $(\mathcal{M}_n)_{n \geq 1}$ and the index p . We prove the following real interpolation identity: if $0 < \theta < 1$ and $1/p = 1 - \theta$, then

$$(\mathcal{H}_1^c(\mathcal{M}), \mathcal{H}_\infty^c(\mathcal{M}))_{\theta, p} = \mathcal{H}_p^c(\mathcal{M}).$$

This is new even for classical martingale Hardy spaces as it is previously known only under the assumption that the filtration is regular. We also obtain analogous result for noncommutative column martingale Orlicz-Hardy spaces.

1. INTRODUCTION

This article is a continuation of our previous work [37] on the investigation of interpolation spaces of compatible couples of various types of noncommutative martingale Hardy spaces and related spaces. The study of interpolations of classical martingale Hardy spaces has a long history which we refer to [14, 39, 40, 43]. Recall that the first class of noncommutative martingale Hardy spaces were introduced by Pisier and Xu in the seminal paper [32]. These Hardy spaces were constructed from column/row square functions and the column versions are commonly denoted by \mathcal{H}_p^c for $0 < p \leq \infty$. Later, another class of noncommutative martingale Hardy spaces were considered by Junge and Xu in [19] which are based on conditioned column/row square functions. The column versions of these second type of Hardy space are denoted by h_p^c for $0 < p \leq \infty$. Both classes of Hardy spaces are instrumental in the development of noncommutative martingale theory the last two decades. The topic of interpolations of these classes of Hardy spaces turns out to be at the forefront of these developments.

Let us recall some background on interpolations of noncommutative martingale Hardy spaces. We refer to [19, 32] for the classes of noncommutative BMO-spaces discussed below. The study of interpolations of noncommutative martingale Hardy spaces was initiated by Musat in [29] where the complex interpolation of the compatible couple $(\mathcal{H}_1^c, \mathcal{BMO}^c)$ was given. Later, Bekjan *et al.* established in [2] that the analogue of Musat's result is valid for couple of column/row conditioned Hardy spaces. More precisely, they obtained the corresponding result for the compatible couple (h_1^c, \mathbf{bmo}^c) . A common theme in these two earlier articles is that all couples considered have one of the endpoints consisting of appropriate types of noncommutative martingale BMO-spaces. We should also mention the articles [3, 4] for related interpolation results in this general direction.

In the recent article [37], we obtained a Peter Jones type result for real interpolations of the couple (h_p^c, h_∞^c) for $0 < p < \infty$. We refer to [5, 16, 17, 31] for background concerning Peter Jones's interpolation result. It is a natural question if the results from [37] remains valid for other classes of Hardy spaces.

2020 *Mathematics Subject Classification.* Primary: 46L52, 46L53, 46B70. Secondary: 46E30, 60G42, 60G48.

Key words and phrases. Noncommutative martingale; Hardy spaces; real interpolation.

The primary objective of the present article is to investigate the compatible couple $(\mathcal{H}_1^c, \mathcal{H}_\infty^c)$. Curiously, general interpolation results for this type of couples of Hardy spaces from classical martingale theory do not appear to be available in the literature. In fact, all existing results to date in this direction require that either the filtration of σ -algebras involved is regular or the Hardy spaces considered were of special type such as those restricted to martingales whose sequences of quadratic variations are dominated by predictable sequence of random variables. Our findings remedy this situation for both commutative and noncommutative settings. More precisely, we establish a general result which states that as in the conditioned case, the family of Hardy spaces $\{\mathcal{H}_p^c\}_{1 \leq p \leq \infty}$ forms a real interpolation scale. In other words, we obtain that if $0 < \theta < 1$ and $1/p = 1 - \theta$, then (with equivalent norms)

$$(1.1) \quad (\mathcal{H}_1^c, \mathcal{H}_\infty^c)_{\theta, p} = \mathcal{H}_p^c$$

where $(\cdot, \cdot)_{\theta, q}$ denotes the real interpolation method. We should point out that, as is already known from the classical case, the identity (1.1) does not extend to the compatible couple $(\mathcal{H}_p^c, \mathcal{H}_\infty^c)$ for $0 < p < 1$. This is in strong contrast with the conditioned case from [37] and thus highlighting that the two situations can be quite different. However, our method of proof is based on insights from techniques used in [37]. Indeed, the decisive step in our argument is an estimate from above of the K -functional of the couple $(\mathcal{H}_2^c, \mathcal{H}_\infty^c)$. We refer to Theorem 3.1 below for details. The main feature of the estimate is our use of the dual Cesàro operator. Although this particular estimate does not imply that the couple $(\mathcal{H}_2^c, \mathcal{H}_\infty^c)$ is K -closed in some appropriate noncommutative couple (L_2, L_∞) as in the case of conditioned spaces treated in [37], it is sufficient to deduce real interpolations of the couple $(\mathcal{H}_2^c, \mathcal{H}_\infty^c)$. The case of the couple $(\mathcal{H}_1^c, \mathcal{H}_\infty^c)$ is then deduced from the couple $(\mathcal{H}_2^c, \mathcal{H}_\infty^c)$ via standard use of Wolff's interpolation theorem.

Motivated by the recent interest on martingale Hardy spaces associated with Orlicz function spaces in the classical setting, we also consider the case of noncommutative Orlicz-Hardy spaces. We refer to recent articles [26, 27] for more perspective and background on real interpolations of compatible couples involving martingale Orlicz-Hardy spaces in the classical setting. We show as an extension of our techniques that (1.1) remains valid for noncommutative martingale Hardy spaces associated with Orlicz functions satisfying some natural conditions.

The paper is organized as follows. In Section 2, we give a brief introduction of noncommutative spaces and review the construction of noncommutative martingale Hardy spaces associated with symmetric spaces of measurable operators. This section also contains relevant discussions on some concepts from interpolation theory. More specifically, the real interpolation method is discussed. It also includes some background concerning the Cesàro operator and its dual operator that play key parts in the estimate of the K -functional discussed above. Section 3 is where we provide the formulation and proof of our primary result together with extensions and applications. The section also contains a paragraph dealing with noncommutative martingale Hardy spaces associated with general symmetric spaces. More specifically, we consider real interpolations of couples involving classes of noncommutative martingale Hardy spaces associated with Orlicz function spaces.

2. DEFINITIONS AND PRELIMINARY RESULTS

Throughout, we use c_{abs} to denote some absolute constant whose value may change from one statement to the next. We write $A \lesssim B$ if there is some absolute constant c_{abs} such that $A \leq c_{abs}B$. We say that A is equivalent to B if $A \lesssim B$ and $B \lesssim A$. In this case, we write $A \approx B$.

2.1. Generalized singular value functions and noncommutative spaces. In what follows, H is a separable Hilbert space and $\mathcal{M} \subseteq \mathcal{B}(H)$ denotes a semifinite von Neumann algebra equipped

with a faithful normal semifinite trace τ . The identity in \mathcal{M} will be denoted by $\mathbf{1}$. A closed and densely defined operator a on H is said to be *affiliated* with \mathcal{M} if $u^*au = a$ for each unitary operator u in the commutant \mathcal{M}' of \mathcal{M} . An operator x is called τ -*measurable* if x is affiliated with \mathcal{M} and for every $\varepsilon > 0$, there exists a projection $p \in \mathcal{M}$ such that $p(H) \subset \text{dom}(x)$ and $\tau(\mathbf{1}-p) < \varepsilon$. The set of all τ -measurable operators will be denoted by $L_0(\mathcal{M})$. Given a self-adjoint operator $x \in L_0(\mathcal{M})$ and a Borel set $B \subset \mathbb{R}$, we denote by $\chi_B(x)$ its spectral projection. The *distribution function* of x is defined by

$$\lambda_s(x) = \tau(\chi_{(s,\infty)}(x)), \quad s \in \mathbb{R}.$$

For $x \in L_0(\mathcal{M})$, the *generalized singular value function* of x is defined by

$$\mu_t(x) = \inf \{s > 0 : \lambda_s(|x|) \leq t\}, \quad t > 0.$$

The function $t \mapsto \mu_t(x)$ is decreasing and right-continuous. In the case that \mathcal{M} is the abelian von Neumann algebra $L_\infty(0, \alpha)$ ($0 < \alpha \leq \infty$) with the trace given by the integration with respect to the Lebesgue measure, $L_0(\mathcal{M})$ is the space of all measurable functions, with non-trivial distribution, and $\mu(f)$ is the decreasing rearrangement of the measurable function f (see [25]). In the abelian case, we write $L_0(0, \alpha)$ instead of $L_0(L_\infty(0, \alpha))$ ($0 < \alpha \leq \infty$). For more discussions on generalized singular value functions, we refer the reader to [10].

Let $0 < \alpha \leq \infty$. A (quasi) Banach function space $(E, \|\cdot\|_E)$ on the interval $(0, \alpha)$ is called *symmetric* if for every $g \in E$ and for every measurable function f with $\mu(f) \leq \mu(g)$, we have $f \in E$ and $\|f\|_E \leq \|g\|_E$.

Given a symmetric quasi-Banach function space E on $(0, \alpha)$, we define the corresponding noncommutative space of operators by setting:

$$E(\mathcal{M}, \tau) = \left\{ x \in L_0(\mathcal{M}) : \mu(x) \in E \right\}.$$

Equipped with the quasi-norm $\|x\|_{E(\mathcal{M}, \tau)} := \|\mu(x)\|_E$, the linear space $E(\mathcal{M}, \tau)$ becomes a complex quasi-Banach space ([22, 38, 42]) and is usually referred to as the *noncommutative symmetric space* associated with (\mathcal{M}, τ) corresponding to $(E, \|\cdot\|_E)$. We remark that if $0 < p < \infty$ and $E = L_p$, then $E(\mathcal{M}, \tau)$ is exactly the usual noncommutative L_p -space $L_p(\mathcal{M}, \tau)$ associated with the pair (\mathcal{M}, τ) . In the sequel, $E(\mathcal{M}, \tau)$ will be abbreviated to $E(\mathcal{M})$.

Beside L_p -spaces, we also make extensive use of two classes of symmetric spaces. Namely, Lorentz spaces and Orlicz spaces. We begin with the former.

• *Lorentz spaces*: Let $0 < p, q \leq \infty$. The *Lorentz space* $L_{p,q}(0, \infty)$ is the space of all $f \in L_0(0, \infty)$ for which $\|f\|_{p,q} < \infty$ where

$$\|f\|_{p,q} = \begin{cases} \left(\int_0^\infty \mu_t^q(f) d(t^{q/p}) \right)^{1/q}, & 0 < q < \infty; \\ \sup_{t>0} t^{1/p} \mu_t(f), & q = \infty. \end{cases}$$

If $1 \leq q \leq p < \infty$ or $p = q = \infty$, then $L_{p,q}(0, \infty)$ is a symmetric Banach function space. If $1 < p < \infty$ and $p \leq q \leq \infty$, then $L_{p,q}(0, \infty)$ can be equivalently renormed to become a symmetric Banach function space ([5, Theorem 4.6]). In general, $L_{p,q}(0, \infty)$ is only a symmetric quasi-Banach function space.

We will also use a different type of Lorentz spaces which we briefly describe. Let $\phi : [0, \infty) \rightarrow [0, \infty)$ be an increasing concave continuous function such that $\phi(0) = 0$ and $\lim_{t \rightarrow \infty} \phi(t) = \infty$. The Lorentz space $\Lambda_\phi(0, \infty)$ is defined by setting:

$$\Lambda_\phi(0, \infty) := \left\{ f \in L_0(0, \infty) : \int_0^\infty \mu_s(f) d\phi(s) < \infty \right\}$$

equipped with the norm

$$\|f\|_{\Lambda_\phi(0,\infty)} := \int_0^\infty \mu_s(f) d\phi(s).$$

Clearly, $\Lambda_\phi(0, \infty)$ is a symmetric Banach function space. In the case $\phi(t) = \log(1+t)$, we use the notation $\Lambda_{\log}(0, \infty)$ for $\Lambda_\phi(0, \infty)$.

• *Orlicz spaces:* By an *Orlicz function* Φ on $[0, \infty)$, we mean a continuous increasing function satisfying $\Phi(0) = 0$ and $\lim_{t \rightarrow \infty} \Phi(t) = \infty$. An Orlicz function Φ is said to be *p-convex* if the function $t \mapsto \Phi(t^{1/p})$ is convex, and to be *q-concave* if the function $t \mapsto \Phi(t^{1/q})$ is concave. For a given Orlicz function Φ that is *p-convex* and *q-concave* for $0 < p \leq q < \infty$, the associated *Orlicz space* $L_\Phi(0, \infty)$ is defined by setting

$$L_\Phi(0, \infty) := \left\{ f \in L_0(0, \infty) : \int_0^\infty \Phi\left(\frac{|f(s)|}{\lambda}\right) ds < \infty \text{ for some } \lambda > 0 \right\}$$

equipped with the quasi-norm

$$\|f\|_\Phi := \inf \left\{ \lambda > 0 : \int_0^\infty \Phi\left(\frac{|f(s)|}{\lambda}\right) ds \leq 1 \right\}.$$

The Orlicz space $L_\Phi(0, \infty)$ is a symmetric quasi-Banach function space. We refer to [24, 28] for more details on Orlicz functions and Orlicz spaces. We will also make use of the following more general space: for $0 < r \leq \infty$, the space $L_{\Phi,r}(0, \infty)$ is the collection of all $f \in L_0(0, \infty)$ for which $\|f\|_{\Phi,r} < \infty$ where

$$\|f\|_{\Phi,r} := \begin{cases} \left(r \int_0^\infty (t \|\chi_{\{|f|>t\}}\|_\Phi)^r \frac{dt}{t} \right)^{1/r}, & 0 < r < \infty; \\ \sup_{t>0} t \|\chi_{\{|f|>t\}}\|_\Phi, & r = \infty. \end{cases}$$

The space $L_{\Phi,r}(0, \infty)$ was introduced in [11] and was called Orlicz-Lorentz space there. Note that if $\Phi(t) = t^p$, then $L_{\Phi,r}(0, \infty)$ coincides with the Lorentz space $L_{p,r}(0, \infty)$. The space $L_{\Phi,\infty}(0, \infty)$ is also known as the weak Orlicz space.

We conclude this subsection by recalling the notion of *submajorization*. Given $x, y \in L_0(\mathcal{M})$, we say that y is *submajorized* in the sense of Hardy-Littlewood-Pólya by x (written $y \prec\prec x$) if

$$\int_0^t \mu_s(y) ds \leq \int_0^t \mu_s(x) ds, \quad t > 0.$$

In the sequel, we will frequently use the submajorization inequality

$$(2.1) \quad \mu(x+y) \prec\prec \mu(x) + \mu(y), \quad x, y \in L_0(\mathcal{M}).$$

Another fact that is important below is that if $T : L_1(\mathcal{M}) + \mathcal{M} \rightarrow L_1(\mathcal{M}) + \mathcal{M}$ satisfies $\max\{\|T : L_1(\mathcal{M}) \rightarrow L_1(\mathcal{M})\|; \|T : \mathcal{M} \rightarrow \mathcal{M}\|\} \leq 1$ then for every $x \in L_1(\mathcal{M}) + \mathcal{M}$, $Tx \prec\prec x$. This fact can be found in [9, Proposition 4.1]. In particular, if $x \in L_1(\mathcal{M}) + \mathcal{M}$ and $(p_k)_{k \geq 1}$ is a sequence of mutually disjoint projections from \mathcal{M} then,

$$(2.2) \quad \sum_{k \geq 1} p_k x p_k \prec\prec x.$$

2.2. Noncommutative martingale Hardy spaces. By a filtration $(\mathcal{M}_n)_{n \geq 1}$, we mean an increasing sequence of von Neumann subalgebras of \mathcal{M} whose union is w^* -dense in \mathcal{M} . Throughout, we will work with a fixed filtration $(\mathcal{M}_n)_{n \geq 1}$. For every $n \geq 1$, we assume further that there is a trace preserving conditional expectation \mathcal{E}_n from \mathcal{M} onto \mathcal{M}_n . This is the case if for every $n \geq 1$, the restriction of the trace τ on \mathcal{M}_n is semifinite. It is well-known that for $1 \leq p < \infty$, the \mathcal{E}_n 's extend to be contractive projections from $L_p(\mathcal{M}, \tau)$ onto $L_p(\mathcal{M}_n, \tau|_{\mathcal{M}_n})$. In particular, they are well-defined on $L_1(\mathcal{M}) + \mathcal{M}$.

Definition 2.1. A sequence $x = (x_n)_{n \geq 1}$ in $L_1(\mathcal{M}) + \mathcal{M}$ is called a *noncommutative martingale* with respect to the filtration $(\mathcal{M}_n)_{n \geq 1}$ if for every $n \geq 1$,

$$\mathcal{E}_n(x_{n+1}) = x_n.$$

Let E be a symmetric quasi-Banach function space and $x = (x_n)_{n \geq 1}$ be a martingale. If for every $n \geq 1$, $x_n \in E(\mathcal{M}_n)$, then we say that $(x_n)_{n \geq 1}$ is an $E(\mathcal{M})$ -martingale. In this case, we set

$$\|x\|_{E(\mathcal{M})} = \sup_{n \geq 1} \|x_n\|_{E(\mathcal{M})}.$$

If $\|x\|_{E(\mathcal{M})} < \infty$, then x will be called a bounded $E(\mathcal{M})$ -martingale.

For a martingale $x = (x_n)_{n \geq 1}$, we set $dx_n = x_n - x_{n-1}$ for $n \geq 1$ with the usual convention that $x_0 = 0$. The sequence $dx = (dx_n)_{n \geq 1}$ is called the *martingale difference sequence* of x . A martingale x is called a *finite martingale* if there exists $N \in \mathbb{N}$ such that $dx_n = 0$ for all $n \geq N$.

Let us now review some basic definitions related to martingale Hardy spaces associated to noncommutative symmetric spaces.

Following [32], we define the *column square functions* of a given martingale $x = (x_k)_{k \geq 1}$ by setting:

$$S_{c,n}(x) = \left(\sum_{k=1}^n |dx_k|^2 \right)^{1/2}, \quad S_c(x) = \left(\sum_{k=1}^{\infty} |dx_k|^2 \right)^{1/2}.$$

For convenience, we will use the notation

$$S_{c,n}(a) = \left(\sum_{k=1}^n |a_k|^2 \right)^{1/2}, \quad S_c(a) = \left(\sum_{k=1}^{\infty} |a_k|^2 \right)^{1/2}$$

for sequences $a = (a_k)_{k \geq 1}$ in $L_1(\mathcal{M}) + \mathcal{M}$ that are not necessarily martingale difference sequences. It is worth pointing out that the infinite sums of positive operators stated above may not always make sense as operators. However, if the sequence $(S_{c,n}(x))_{n \geq 1}$ is order bounded, then it admits a supremum. In that case, $S_c(x)$ may be taken to be the limit of the sequence $(S_{c,n}(x))_{n \geq 1}$ for the measure topology. Similar remark applies to the sequence $(S_{c,n}(a))_{n \geq 1}$.

We will now describe noncommutative martingale Hardy spaces associated with symmetric Banach function spaces. In this paper, we will only work with Hardy spaces built from square functions. We refer to [36, 37] for other types of noncommutative martingale Hardy spaces.

Assume that E is a symmetric Banach function space on $(0, \infty)$. We denote by \mathcal{F}_E the collection of all finite martingales in $E(\mathcal{M}) \cap \mathcal{M}$. For $x = (x_k)_{k \geq 1} \in \mathcal{F}_E$, we set:

$$\|x\|_{\mathcal{H}_E^c} = \|S_c(x)\|_{E(\mathcal{M})}.$$

Then $(\mathcal{F}_E, \|\cdot\|_{\mathcal{H}_E^c})$ is a normed space. If we denote by $(e_{i,j})_{i,j \geq 1}$ the family of unit matrices in $\mathcal{B}(\ell_2)$, then the correspondence $x \mapsto \sum_{k \geq 1} dx_k \otimes e_{k,1}$ maps \mathcal{F}_E isometrically into a (not necessarily closed) linear subspace of $E(\mathcal{M} \overline{\otimes} \mathcal{B}(\ell_2))$.

We define the *column martingale Hardy space* $\mathcal{H}_E^c(\mathcal{M})$ to be the completion of $(\mathcal{F}_E, \|\cdot\|_{\mathcal{H}_E^c})$. It then follows that $\mathcal{H}_E^c(\mathcal{M})$ embeds isometrically into a closed subspace of the quasi-Banach space $E(\mathcal{M} \otimes \mathcal{B}(\ell_2))$.

We remark that using the above definition with $L_p(0, \infty)$ where $1 \leq p < \infty$, we recover the definition of $\mathcal{H}_p^c(\mathcal{M})$ as defined in [32]. However, the case $p = \infty$ is not covered by the above description since it requires separability. We define $\mathcal{H}_\infty^c(\mathcal{M})$ as the collection of all martingales in \mathcal{M} for which the column square functions exists in \mathcal{M} . The norm in $\mathcal{H}_\infty^c(\mathcal{M})$ is defined by:

$$\|x\|_{\mathcal{H}_\infty^c} = \|\mathcal{S}_c(x)\|_\infty, \quad x \in \mathcal{H}_\infty^c(\mathcal{M}).$$

In the sequel, we will also make use of the more general column space $E(\mathcal{M}; \ell_2^c)$ which is defined as the set of all sequences $a = (a_k)_{k \geq 1}$ in $E(\mathcal{M})$ for which $\mathcal{S}_c(a)$ exists in $E(\mathcal{M})$. In this case, we set

$$\|a\|_{E(\mathcal{M}; \ell_2^c)} = \|\mathcal{S}_c(a)\|_{E(\mathcal{M})}.$$

Under the above quasi-norm, one can easily see that $E(\mathcal{M}; \ell_2^c)$ is a quasi-Banach space. The closed subspace of $E(\mathcal{M}; \ell_2^c)$ consisting of adapted sequences will be denoted by $E^{\text{ad}}(\mathcal{M}; \ell_2^c)$. That is,

$$E^{\text{ad}}(\mathcal{M}; \ell_2^c) = \left\{ (a_n)_{n \geq 1} \in E(\mathcal{M}; \ell_2^c) : \forall n \geq 1, a_n \in E(\mathcal{M}_n) \right\}.$$

Below, we use the notation $\mathcal{H}_{p,q}^c(\mathcal{M})$ for the noncommutative column martingale Hardy space associated with the Lorentz space $L_{p,q}(0, \infty)$. Similarly, $\mathcal{H}_\Phi^c(\mathcal{M})$ and $\mathcal{H}_{\Phi,r}^c(\mathcal{M})$ are used for noncommutative column Hardy spaces associated with the function spaces $L_\Phi(0, \infty)$ and $L_{\Phi,r}(0, \infty)$ respectively.

Note that for $1 < p < \infty$, it follows from the noncommutative Stein inequality that $L_p^{\text{ad}}(\mathcal{M}; \ell_2^c)$ is a complemented subspace of $L_p(\mathcal{M}; \ell_2^c)$. Similarly, for $1 \leq p < \infty$, $\mathcal{H}_p^c(\mathcal{M})$ is a complemented subspace of $L_p^{\text{ad}}(\mathcal{M}; \ell_2^c)$. The case $p = 1$ is a consequence of the noncommutative Lépingle-Yor inequality ([34]). However, in general, $\mathcal{H}_1^c(\mathcal{M})$ is not a complemented subspace of $L_1(\mathcal{M}; \ell_2^c)$. Likewise, $\mathcal{H}_\infty^c(\mathcal{M})$ is not a complemented subspace of $L_\infty(\mathcal{M}; \ell_2^c)$.

2.3. Basics of interpolations. Let (A_0, A_1) be a compatible couple of quasi-Banach spaces in the sense that both A_0 and A_1 embed continuously into some topological vector space \mathcal{Z} . This allows us to define the spaces $A_0 \cap A_1$ and $A_0 + A_1$. These are quasi-Banach spaces when equipped with quasi-norms:

$$\|x\|_{A_0 \cap A_1} = \max \left\{ \|x\|_{A_0}, \|x\|_{A_1} \right\}$$

and

$$\|x\|_{A_0 + A_1} = \inf \left\{ \|x_0\|_{A_0} + \|x_1\|_{A_1} : x = x_0 + x_1, x_0 \in A_0, x_1 \in A_1 \right\},$$

respectively.

Definition 2.2. A (quasi) Banach space A is called an *interpolation space* for the couple (A_0, A_1) if $A_0 \cap A_1 \subseteq A \subseteq A_0 + A_1$ and whenever a bounded linear operator $T : A_0 + A_1 \rightarrow A_0 + A_1$ is such that $T(A_0) \subseteq A_0$ and $T(A_1) \subseteq A_1$, we have $T(A) \subseteq A$ and

$$\|T : A \rightarrow A\| \leq c \max \left\{ \|T : A_0 \rightarrow A_0\|, \|T : A_1 \rightarrow A_1\| \right\}$$

for some constant c .

If A is an interpolation space for the couple (A_0, A_1) , we write $A \in \text{Int}(A_0, A_1)$. Below, we are primarily interested in an interpolation method generally referred to as *the real interpolation method* which we now briefly review.

A fundamental notion for the construction of real interpolation spaces is the *K-functional*. For $x \in A_0 + A_1$, we define the *K-functional* by setting for $t > 0$,

$$K(x, t) = K(x, t; A_0, A_1) = \inf \left\{ \|x_0\|_{A_0} + t\|x_1\|_{A_1} : x = x_0 + x_1, x_0 \in A_0, x_1 \in A_1 \right\}.$$

Note that for each $t > 0$, $x \mapsto K(x, t)$ gives an equivalent quasi-norm on $A_0 + A_1$.

If $0 < \theta < 1$ and $1 \leq \gamma < \infty$, the real interpolation space $A_{\theta, \gamma} = (A_0, A_1)_{\theta, \gamma}$ is defined by $x \in A_{\theta, \gamma}$ if and only if

$$\|x\|_{(A_0, A_1)_{\theta, \gamma}} = \left(\int_0^\infty (t^{-\theta} K(x, t; A_0, A_1))^\gamma \frac{dt}{t} \right)^{1/\gamma} < \infty.$$

If $\gamma = \infty$, we define $x \in A_{\theta, \infty}$ if and only if

$$\|x\|_{(A_0, A_1)_{\theta, \infty}} = \sup_{t > 0} t^{-\theta} K(x, t; A_0, A_1) < \infty.$$

For $0 < \theta < 1$ and $0 < \gamma \leq \infty$, the functional $\|\cdot\|_{\theta, \gamma}$ is a quasi-norm. In the case where A_0 and A_1 are Banach spaces and $1 \leq \gamma \leq \infty$, $(A_{\theta, \gamma}, \|\cdot\|_{\theta, \gamma})$ can be renormed to be a Banach space. Moreover, the space $A_{\theta, \gamma}$ is an interpolation space for the couple (A_0, A_1) in the sense of Definition 2.2. There is also an equivalent description of $A_{\theta, \gamma}$ using a dual notion called *J-functionals* but this will not be needed for our purpose below. Our main references for interpolations are the books [5] and [6].

It is worth mentioning that the real interpolation method is well understood for the couple (L_{p_0}, L_{p_1}) for both the classical case and the noncommutative case. We record here that Lorentz spaces can be realized as real interpolation spaces for the couple (L_{p_0}, L_{p_1}) . More precisely, if \mathcal{N} is a semifinite von Neumann algebra, $1 \leq p_0 < p_1 \leq \infty$, $0 < \theta < 1$, and $1 \leq q \leq \infty$ then, up to equivalent quasi-norms (independent of \mathcal{N}),

$$(L_{p_0}(\mathcal{N}), L_{p_1}(\mathcal{N}))_{\theta, q} = L_{p, q}(\mathcal{N})$$

where $1/p = (1 - \theta)/p_0 + \theta/p_1$. By reiteration, if $1 \leq \lambda, \gamma \leq \infty$, we also have

$$(2.3) \quad (L_{p_0, \lambda}(\mathcal{N}), L_{p_1, \gamma}(\mathcal{N}))_{\theta, q} = L_{p, q}(\mathcal{N})$$

with equivalent quasi-norms when $1 \leq p_0 < p_1 \leq \infty$, $0 < \theta < 1$, $1 \leq q \leq \infty$, and $1/p = (1 - \theta)/p_0 + \theta/p_1$. These facts can be found in [33] and will be used repeatedly throughout.

Wolff's interpolation theorem will be needed in the next subsection. We record it here for convenience.

Theorem 2.3 ([41, Theorem 1]). *Let B_i ($i = 1, 2, 3, 4$) be quasi-Banach spaces such that $B_1 \cap B_4$ is dense in B_j ($j = 2, 3$) and satisfy:*

$$B_2 = (B_1, B_3)_{\phi, r} \quad \text{and} \quad B_3 = (B_2, B_4)_{\theta, q}$$

for $0 < \phi, \theta < 1$ and $0 < r, q \leq \infty$. Then

$$B_2 = (B_1, B_4)_{\xi, r} \quad \text{and} \quad B_3 = (B_1, B_4)_{\zeta, q}$$

where $\xi = \frac{\phi\theta}{1 - \phi + \phi\theta}$ and $\zeta = \frac{\theta}{1 - \phi + \phi\theta}$.

It is more convenient to apply the Wolff's interpolation theorem using intervals.

Definition 2.4. A family of quasi-Banach spaces $(A_{p,\gamma})_{p,\gamma \in (0,\infty]}$ is said to form a *real interpolation scale* on an interval $I \subseteq \mathbb{R} \cup \{\infty\}$ if for every $p, q \in I$, $0 < \gamma_1, \gamma_2, \gamma \leq \infty$, $0 < \theta < 1$, and $1/r = (1 - \theta)/p + \theta/q$,

$$A_{r,\gamma} = (A_{p,\gamma_1}, A_{q,\gamma_2})_{\theta,\gamma}$$

with equivalent quasi-norms.

The next lemma is a version of Wolff's interpolation theorem at the level of family of real interpolation scale.

Lemma 2.5 ([36, Lemma 3.4]). *Assume that a family of quasi-Banach spaces $\mathfrak{F} = (A_{p,\gamma})_{p,\gamma \in (0,\infty]}$ forms a real interpolation scale on two different intervals I and J . If $|J \cap I| > 1$, then \mathfrak{F} forms a real interpolation scale on the interval $I \cup J$.*

A more general real interpolation type spaces will be essential in our consideration below. Recall that a Banach function space \mathcal{F} has a monotone norm if whenever $f, g \in \mathcal{F}$, $|f| \leq |g| \implies \|f\|_{\mathcal{F}} \leq \|g\|_{\mathcal{F}}$.

Definition 2.6. An interpolation space E for a Banach couple (E_0, E_1) is said to be *given by a K -method* if there exists a Banach function space \mathcal{F} with monotone norm such that $x \in E$ if and only if $t \mapsto K(x, t; E_0, E_1) \in \mathcal{F}$ and there exists a constant $C_E > 0$ such that

$$C_E^{-1} \|t \mapsto K(x, t; E_0, E_1)\|_{\mathcal{F}} \leq \|x\|_E \leq C_E \|t \mapsto K(x, t; E_0, E_1)\|_{\mathcal{F}}.$$

In this case, we write $E = (E_0, E_1)_{\mathcal{F}; K}$.

The following fact will be used in the sequel. This is known as a result of Brudnyi and Krugliak (see [21, Theorem 6.3]).

Proposition 2.7. *Let $1 \leq p < q \leq \infty$. Every interpolation space $E \in \text{Int}(L_p, L_q)$ is given by a K -method.*

In the next section, we will sparingly use the concept of K -closed couple which we refer to [23, 31, 37] for formal definition and relevant properties.

We end this subsection with a discussion on an extrapolation result that will be needed in the next section. It involves the notion of Cesàro operator together with its formal dual. Recall that the *Cesàro operator* $C : (L_1 + L_\infty)(0, \infty) \rightarrow (L_{1,\infty} + L_\infty)(0, \infty)$ is defined by the formula

$$(Cf)(t) := \frac{1}{t} \int_0^t f(s) ds, \quad f \in (L_1 + L_\infty)(0, \infty).$$

The operator C^* which we will refer to as *the dual Cesàro operator* is defined by setting

$$(C^*f)(t) := \int_t^\infty \frac{f(s)}{s} ds, \quad f \in \Lambda_{\log}(0, \infty).$$

It is known that C^* is bounded from $\Lambda_{\log}(0, \infty)$ into $(L_1 + L_\infty)(0, \infty)$. A proof of this fact can be found in [15, Fact 2.9]. We refer to C^* as the (formal) *dual of the Cesàro operator* due to the fact that

$$\langle Cf, g \rangle = \langle f, C^*g \rangle, \quad f, g \in L_2(0, \infty).$$

We will need the following well known results on boundedness of Cesàro operators (see [12, Theorem 327] for the detailed proof).

Lemma 2.8. *The operators C and C^* satisfy the following properties:*

$$\|C\|_{L_p \rightarrow L_p} = p', \quad 1 < p \leq \infty,$$

and

$$\|C^*\|_{L_p \rightarrow L_p} = p, \quad 1 \leq p < \infty,$$

where p' is the conjugate index of p .

We now consider an extrapolation result from [15]. Assume that (\mathcal{N}_1, ν_1) and (\mathcal{N}_2, ν_2) are semifinite von Neumann algebras and suppose that $T : L_2(\mathcal{N}_1, \nu_1) \rightarrow L_2(\mathcal{N}_2, \nu_2)$ is a linear operator that admits bounded extensions from $L_p(\mathcal{N}_1, \nu_1)$ to $L_p(\mathcal{N}_2, \nu_2)$ for all $2 < p < \infty$ and satisfies further that

$$\|T\|_{L_p(\mathcal{N}_1) \rightarrow L_p(\mathcal{N}_2)} \leq p, \quad 2 < p < \infty.$$

Such assumption is referred to in [15] as the *First Extrapolation Condition*. The following extrapolation will be used to deduce Proposition 2.10 below.

Theorem 2.9 ([15, Theorem 3.2]). *Suppose that T satisfies the First Extrapolation condition. Then T admits a bounded linear extension from $(\Lambda_{\log} \cap (L_2 + L_\infty))(\mathcal{N}_1, \nu_1)$ into $(L_2 + L_\infty)(\mathcal{N}_2, \nu_2)$. Moreover, we have*

$$\mu^2(Tx) \prec\prec c_{\text{abs}}(C^* \mu(x))^2, \quad x \in (\Lambda_{\log} \cap (L_2 + L_\infty))(\mathcal{N}_1, \nu_1).$$

The following variant of the dual Doob inequality is essential in the our proof in next section.

Proposition 2.10. *Let $(e_k)_{k \geq 1}$ be a mutually disjoint sequence of projections in \mathcal{M} . If a is a positive operator so that $a^{1/2} \in (\Lambda_{\log} \cap (L_2 + L_\infty))(\mathcal{M})$, then*

$$\mu\left(\sum_{k \geq 1} \mathcal{E}_{k-1}(e_k a e_k)\right) \prec\prec c_{\text{abs}}\left(C^*[\mu^{1/2}(a)]\right)^2.$$

In preparation of the proof, we recall a crucial result established by Junge in [18, Proposition 2.8] (see also [7] for the nonseparable case).

Proposition 2.11. *Let \mathcal{N} be a semifinite von Neuman subalgebra of \mathcal{M} . If $\mathcal{E} : \mathcal{M} \rightarrow \mathcal{N}$ is the trace preserving conditional expectation, then there exists an \mathcal{N} -module linear isometry $u : L_2(\mathcal{M}) \rightarrow L_2(\mathcal{N} \overline{\otimes} \mathcal{B}(\ell_2))$ such that*

$$u(x)^* u(y) = \mathcal{E}(x^* y) \otimes e_{1,1}, \quad x, y \in L_2(\mathcal{M}).$$

Proof of Proposition 2.10. Let $\Theta : L_2(\mathcal{M}) \rightarrow L_2(\mathcal{M} \overline{\otimes} \mathcal{B}(\ell^2))$ be defined by

$$\Theta(x) = \sum_{k \geq 1} x e_k \otimes e_{k,1}.$$

Then, for $2 \leq p < \infty$ and $x \in L_p(\mathcal{M})$, we have

$$\begin{aligned} \|\Theta(x)\|_p^p &= \left\| \left(\sum_{k \geq 1} e_k |x|^2 e_k \right)^{1/2} \right\|_p^p \\ &= \left\| \sum_{k \geq 1} e_k |x|^2 e_k \right\|_{p/2}^{p/2} \\ &\leq \| |x|^2 \|_{p/2}^{p/2} = \|x\|_p^p. \end{aligned}$$

That is, $\|\Theta\|_{L_p \rightarrow L_p} \leq 1$. Next, for $k \geq 1$, let $u_k : L_2(\mathcal{M}) \rightarrow L_2(\mathcal{M}_k \overline{\otimes} \mathcal{B}(\ell_2))$ be the \mathcal{M}_k -module map as in Proposition 2.11 that satisfies

$$|u_k(x)|^2 = \mathcal{E}_k(x^*x) \otimes e_{1,1}, \quad x \in L_2(\mathcal{M}).$$

Consider the mapping T formally defined by the formula

$$Ta = \sum_{k \geq 1} u_{k-1}(a_{k1}) \otimes e_{k,1}, \quad a = (a_{kj})_{k,j \geq 1} \in L_0(\mathcal{M} \overline{\otimes} \mathcal{B}(\ell_2)).$$

One can easily see that $|T(a)|^2 = \sum_{k \geq 1} \mathcal{E}_{k-1}(|a_{k1}|^2) \otimes e_{1,1} \otimes e_{1,1}$. By the dual Doob inequality ([18, 20]), we have for $2 \leq p < \infty$,

$$\begin{aligned} \|Ta\|_p^p &= \| |Ta|^2 \|_{p/2}^{p/2} \\ &\leq (c_{abs}(p/2)^2)^{p/2} \left\| \sum_{k \geq 1} |a_{k1}|^2 \right\|_{p/2}^{p/2} \\ &\leq c_{abs}^{p/2} p^p \|a\|_p^p. \end{aligned}$$

We obtain that $\|T\|_{L_p \rightarrow L_p} \leq c_{abs}^{1/2} p$. This shows in particular that the operator

$$T \circ \Theta : L_2(\mathcal{M}) \rightarrow L_2(\mathcal{M} \overline{\otimes} \mathcal{B}(\ell_2) \overline{\otimes} \mathcal{B}(\ell_2))$$

satisfies the First Extrapolation Condition. Applying Theorem 2.9, we have for every $x \in (\Lambda_{\log} \cap (L_2 + L_\infty))(\mathcal{M})$,

$$\mu^2(T \circ \Theta(x)) \prec\prec c_{abs}(C^* \mu(x))^2.$$

Note that $\mu^2(T \circ \Theta(x)) = \mu(|T \circ \Theta(x)|^2)$ and $|T \circ \Theta(x)|^2 = \sum_{k \geq 1} \mathcal{E}_{k-1}(e_k |x|^2 e_k) \otimes e_{1,1} \otimes e_{1,1}$. The desired conclusion follows by applying the above submajorization to $x = a^{1/2}$. \square

3. INTERPOLATIONS OF MARTINGALE HARDY SPACES

3.1. An estimate of the K -functional of the couple $(\mathcal{H}_2^c, \mathcal{H}_\infty^c)$. The following is the primary result of the paper.

Theorem 3.1. *For every $x \in \mathcal{H}_{\Lambda_{\log}}^c(\mathcal{M}) \cap (\mathcal{H}_2^c(\mathcal{M}) + \mathcal{H}_\infty^c(\mathcal{M}))$ and $t > 0$, the following inequality holds:*

$$K(x, t; \mathcal{H}_2^c(\mathcal{M}), \mathcal{H}_\infty^c(\mathcal{M})) \leq C_{abs} \left(\int_0^{t^2} \left(\mu_u(S_c(x)) + C^*[\mu(S_c(x))](u) \right)^2 du \right)^{1/2}$$

where C^* is the dual Cesàro operator.

Proof. The argument below follows the strategy used for the conditioned case in [37, Proposition 3.2]. The main new idea is the incorporation of the dual Cesàro operator C^* . We include all crucial details for completeness.

Assume that $x \in \mathcal{H}_{\Lambda_{\log}}^c(\mathcal{M}) \cap (\mathcal{H}_2^c(\mathcal{M}) + \mathcal{H}_\infty^c(\mathcal{M}))$. This is equivalent to the condition that $S_c(x) \in \Lambda_{\log}(\mathcal{M}) \cap (L_2 + L_\infty)(\mathcal{M})$ or $\mu(S_c(x)) \in \Lambda_{\log}(0, \infty) \cap (L_2 + L_\infty)(0, \infty)$. In particular, $C^*[\mu(S_c(x))]$ is well-defined and belongs to $(L_2 + L_\infty)(0, \infty)$. We will construct a concrete decomposition of x that will provide the desired estimate on the K -functional.

Fix $t > 0$ and set a parameter

$$(3.1) \quad \lambda := \frac{3}{t} \left(\int_0^{t^2} \mu_u(S_c^2(x)) du + \kappa \int_0^{t^2} (C^*[\mu(S_c(x))](u))^2 du \right)^{1/2}$$

where κ denotes the absolute constant from Proposition 2.10. Although at first this choice of λ seems artificial, it will become more transparent during the course of the proof.

As in [37], the argument requires two steps.

• Step 1.

We apply the construction of Cuculescu's projections to the submartingale $(S_{c,k}^2(x))_{k \geq 1}$ and the parameter λ^2 . That is, we start with $q_0 = \mathbf{1}$ and for $k \geq 1$, we set

$$q_k := q_{k-1} \chi_{[0, \lambda^2]}(q_{k-1} S_{c,k}^2(x) q_{k-1}) = \chi_{[0, \lambda^2]}(q_{k-1} S_{c,k}^2(x) q_{k-1}) q_{k-1}.$$

Then $(q_k)_{k \geq 1}$ is a decreasing sequence of projections in \mathcal{M} satisfying the following properties:

- (1) $q_k \in \mathcal{M}_k$ for every $k \geq 1$;
- (2) q_k commutes with $q_{k-1} S_{c,k}^2(x) q_{k-1}$ for all $k \geq 1$;
- (3) $q_k S_{c,k}^2(x) q_k \leq \lambda^2 q_k$ for all $k \geq 1$;
- (4) if we set $q = \bigwedge_{k \geq 1} q_k$, then $\lambda^2(\mathbf{1} - q) \leq \sum_{k \geq 1} (q_{k-1} - q_k) S_{c,k}^2(x) (q_{k-1} - q_k)$.

Verifications of these facts concerning Cuculescu's projections can be found in [7, 8, 30, 35].

Using the sequence $(q_k)_{k \geq 0}$, we consider the following adapted sequence $\alpha = (\alpha_k)_{k \geq 1}$ by setting:

$$(3.2) \quad \alpha_k = dx_k q_k, \quad k \geq 1.$$

Next, we modify α into a martingale difference sequence by setting:

$$(3.3) \quad d\beta_k = \alpha_k - \mathcal{E}_{k-1}(\alpha_k), \quad k \geq 1.$$

We denote the corresponding martingale by $\beta = (\beta_k)_{k \geq 1}$. The main difference with the conditioned case is that $(q_k)_{k \geq 0}$ is only an adapted sequence as oppose to being predictable in [37] and thus the adjustment taken in the definition of β .

We record the next lemma for further use.

Lemma 3.2. *We have the following properties:*

- (i) $\sup_{k \geq 1} \|\alpha_k\|_\infty \leq \lambda$;
- (ii) $\sup_{k \geq 1} \|d\beta_k\|_\infty \leq 2\lambda$;
- (iii) $\mathcal{S}_c^2(\alpha) \prec\prec 4\mathcal{S}_c^2(x)$;
- (iv) $\mu(\mathcal{S}_c^2(\beta)) \prec\prec 8\mu(\mathcal{S}_c^2(x)) + 2\kappa \left(C^* [\mu(\mathcal{S}_c(x))] \right)^2$.

Proof. The first item follows easily from the construction. Indeed, given $k \geq 1$,

$$\begin{aligned} |\alpha_k|^2 &= q_k |dx_k|^2 q_k \\ &= q_k [S_{c,k}^2(x) - S_{c,k-1}^2(x)] q_k \\ &\leq q_k S_{c,k}^2(x) q_k \leq \lambda^2 q_k. \end{aligned}$$

The second item clearly follows from the fact that conditional expectations are contractive projections on \mathcal{M} . For the third item, we consider first the martingale difference sequence

$$d\gamma_k = dx_k q_{k-1}, \quad k \geq 1.$$

We have from the definition of $d\gamma$ that for $m \geq 1$ (with $S_{c,0}(x) = 0$),

$$\begin{aligned} S_{c,m}^2(\gamma) &= \sum_{k=1}^m q_{k-1} |dx_k|^2 q_{k-1} \\ &= \sum_{k=1}^m q_{k-1} [S_{c,k}^2(x) - S_{c,k-1}^2(x)] q_{k-1} \\ &= \sum_{k=1}^m q_{k-1} S_{c,k}^2(x) q_{k-1} - \sum_{k=1}^m q_{k-1} S_{c,k-1}^2(x) q_{k-1}. \end{aligned}$$

Performing some indexing shift, we obtain that

$$S_{c,m}^2(\gamma) = q_{m-1} S_{c,m}^2(x) q_{m-1} + \sum_{k=1}^{m-1} (q_{k-1} S_{c,k}^2(x) q_{k-1} - q_k S_{c,k}^2(x) q_k).$$

From the fact that q_k commutes with $q_{k-1} S_{c,k}^2(x) q_{k-1}$, we deduce that

$$\begin{aligned} S_{c,m}^2(\gamma) &= q_{m-1} S_{c,m}^2(x) q_{m-1} + \sum_{k=1}^{m-1} (q_{k-1} - q_k) S_{c,k}^2(x) (q_{k-1} - q_k) \\ &\leq q_{m-1} S_c^2(x) q_{m-1} + \sum_{k=1}^{m-1} (q_{k-1} - q_k) S_c^2(x) (q_{k-1} - q_k). \end{aligned}$$

Note that the finite family of projections $\{q_{k-1} - q_k : 1 \leq k \leq m-1\} \cup \{q_{m-1}\}$ is mutually disjoint. We may deduce from (2.2) that for every $m \geq 1$, $S_{c,m}^2(\gamma) \prec\prec S_c^2(x)$. Next, for every $w > 0$, the monotone convergence theorem gives:

$$\int_0^w \mu_u(S_c^2(\gamma)) du = \lim_{m \rightarrow \infty} \int_0^w \mu_u(S_{c,m}^2(\gamma)) du \leq \int_0^w \mu_u(S_c^2(x)) du.$$

That is, $S_c^2(\gamma) \prec\prec S_c^2(x)$.

On the other hand, a simple computation gives:

$$\begin{aligned} S_c^2(d\gamma - \alpha) &= \sum_{k \geq 1} (q_{k-1} - q_k) [S_{c,k}^2(x) - S_{c,k-1}^2(x)] (q_{k-1} - q_k) \\ &\leq \sum_{k \geq 1} (q_{k-1} - q_k) S_c^2(x) (q_{k-1} - q_k) \end{aligned}$$

and therefore, we also have from (2.2) that $S_c^2(d\gamma - \alpha) \prec\prec S_c^2(x)$. Using the elementary inequality $|a + b|^2 \leq 2|a|^2 + 2|b|^2$ for operators a and b , we can conclude that

$$S_c^2(\alpha) \leq 2S_c^2(d\gamma - \alpha) + 2S_c^2(\gamma) \prec\prec 4S_c^2(x)$$

which is item (iii).

For the last item, we begin with the simple fact that

$$S_c^2(\beta) \leq 2S_c^2(\alpha) + 2 \sum_{k \geq 1} |\mathcal{E}_{k-1}(\alpha_k)|^2.$$

Recall that $\alpha_k = dx_k q_k = dx_k(q_k - q_{k-1}) + dx_k q_{k-1}$. Since $(dx_k q_{k-1})_{k \geq 1}$ is a martingale difference sequence, we have $\mathcal{E}_{k-1}(\alpha_k) = \mathcal{E}_{k-1}(dx_k(q_k - q_{k-1}))$. Applying Kadison's inequality $|\mathcal{E}_{k-1}(a)|^2 \leq$

$\mathcal{E}_{k-1}(|a|^2)$ for any operator a and $k \geq 1$, we further obtain that

$$\begin{aligned} S_c^2(\beta) &\leq 2S_c^2(\alpha) + 2 \sum_{k \geq 1} \mathcal{E}_{k-1}((q_{k-1} - q_k)|dx_k|^2(q_{k-1} - q_k)) \\ &\leq 2S_c^2(\alpha) + 2 \sum_{k \geq 1} \mathcal{E}_{k-1}[(q_{k-1} - q_k)S_c^2(x)(q_{k-1} - q_k)]. \end{aligned}$$

It follows from (2.1) and Item (iii) that

$$\begin{aligned} \mu(S_c^2(\beta)) &\prec\prec 2\mu(S_c^2(\alpha)) + 2\mu\left(\sum_{k \geq 1} \mathcal{E}_{k-1}[(q_{k-1} - q_k)S_c^2(x)(q_{k-1} - q_k)]\right) \\ &\prec\prec 8\mu(S_c^2(x)) + 2\mu\left(\sum_{k \geq 1} \mathcal{E}_{k-1}[(q_{k-1} - q_k)S_c^2(x)(q_{k-1} - q_k)]\right). \end{aligned}$$

Next, we apply Proposition 2.10 to the mutually disjoint sequence $\{(q_{k-1} - q_k)\}$ and the operator $S_c^2(x)$. Note that $S_c(x) \in (\Lambda_{\log} \cap (L_2 + L_\infty))(\mathcal{M})$ and therefore it satisfies the assumption used in Proposition 2.10. We obtain that

$$\mu(S_c^2(\beta)) \prec\prec 8\mu(S_c^2(x)) + 2\kappa\left(C^*[\mu(S_c(x))]\right)^2$$

which is the desired submajorization. \square

• Step 2. Construction of a decomposition that will provide the stated estimate on the K -functional.

As in the first step, we apply the construction of Cuculescu's projections to the submartingale $(S_{c,k}^2(\beta))_{k \geq 1}$ with the parameter λ^2 where β is the martingale from (3.3). That is, setting $\pi_0 = \mathbf{1}$ and for $k \geq 1$, we define:

$$\pi_k := \pi_{k-1} \chi_{[0, \lambda^2]}(\pi_{k-1} S_{c,k}^2(\beta) \pi_{k-1}) = \chi_{[0, \lambda^2]}(\pi_{k-1} S_{c,k}^2(\beta) \pi_{k-1}) \pi_{k-1}.$$

Then, $(\pi_k)_{k \geq 1}$ is a decreasing sequence of projections in \mathcal{M} . As before, it satisfies the following properties:

- (1) $\pi_k \in \mathcal{M}_k$ for every $k \geq 1$;
- (2) π_k commutes with $\pi_{k-1} S_{c,k}^2(\beta) \pi_{k-1}$ for all $k \geq 1$;
- (3) $\pi_k S_{c,k}^2(\beta) \pi_k \leq \lambda^2 \pi_k$ for all $k \geq 1$;
- (4) if we set $\pi = \bigwedge_{k \geq 1} \pi_k$, then $\lambda^2(\mathbf{1} - \pi) \leq \sum_{k \geq 1} (\pi_{k-1} - \pi_k) S_{c,k}^2(\beta) (\pi_{k-1} - \pi_k)$.

Next, we define two martingales y and z by setting:

$$(3.4) \quad z = \sum_{k \geq 1} d\beta_k \pi_{k-1} = \sum_{k \geq 1} [dx_k q_k - \mathcal{E}_{k-1}(dx_k q_k)] \pi_{k-1} \quad \text{and} \quad y = x - z.$$

We will show that this decomposition provides the desired estimate on the K -functional. We consider first the martingale z . We claim that $z \in \mathcal{H}_\infty^c(\mathcal{M})$ with

$$(3.5) \quad \|z\|_{\mathcal{H}_\infty^c} \leq \sqrt{5}\lambda.$$

To verify (3.5), we first fix $m \geq 1$ and estimate $S_{c,m}(z)$. From the definition of z , we have:

$$\begin{aligned}
S_{c,m}^2(z) &= \sum_{k=1}^m \pi_{k-1} |d\beta_k|^2 \pi_{k-1} \\
&= \sum_{k=1}^m (\pi_{k-1} S_{c,k}^2(\beta) \pi_{k-1} - \pi_{k-1} S_{c,k-1}^2(\beta) \pi_{k-1}) \\
&= \sum_{k=1}^m \pi_{k-1} S_{c,k}^2(\beta) \pi_{k-1} - \sum_{k=1}^{m-1} \pi_k S_{c,k}^2(\beta) \pi_k \\
&= \pi_{m-1} S_{c,m}^2(\beta) \pi_{m-1} + \sum_{k=1}^{m-1} (\pi_{k-1} - \pi_k) S_{c,k}^2(\beta) (\pi_{k-1} - \pi_k),
\end{aligned}$$

where the last equality follows from the commutativity between π_k and $\pi_{k-1} S_{c,k}^2(\beta) \pi_{k-1}$. Recall from Lemma 3.2 (ii) that $\|d\beta_k\|_\infty \leq 2\lambda$. Using this fact, we have

$$\begin{aligned}
\pi_{k-1} S_{c,k}^2(\beta) \pi_{k-1} &= \pi_{k-1} [S_{c,k-1}^2(\beta) + |d\beta_k|^2] \pi_{k-1} \\
&\leq 5\lambda^2 \pi_{k-1}.
\end{aligned}$$

We can deduce that for every $m \geq 1$,

$$S_{c,m}^2(z) \leq \lambda^2 \pi_{m-1} + 5\lambda^2 \sum_{k=1}^{m-1} (\pi_{k-1} - \pi_k) \leq 5\lambda^2 \mathbf{1}.$$

Since this holds for arbitrary $m \geq 1$, we have $S_c^2(z) \leq 5\lambda^2 \mathbf{1}$ which shows that $\|z\|_{\mathcal{H}_\infty^c} \leq \sqrt{5}\lambda$ and thus proving inequality (3.5).

We now deal with the martingale y . We will estimate the norm of y in $\mathcal{H}_2^c(\mathcal{M})$. The verification of the next lemma is the most delicate part of the argument. We take the opportunity to point out that the proof below together with the submajorization in Lemma 3.2(iv) motivated the choice of λ taken in (3.1).

Lemma 3.3. *The projections q and π satisfy the following property:*

$$\max \{ \tau(\mathbf{1} - q), \tau(\mathbf{1} - \pi) \} \leq t^2.$$

Proof. Fix $w > t^2$. We will show that $\mu_w(\mathbf{1} - q) = \mu_w(\mathbf{1} - \pi) = 0$. We provide the argument for $\mathbf{1} - \pi$. The case of $\mathbf{1} - q$ is simpler since it does not depend on the second step.

Assume the opposite, i.e, $\mu_w(\mathbf{1} - \pi) = 1$. We start with the fact that

$$(3.6) \quad \lambda^2(\mathbf{1} - \pi) \leq \sum_{k \geq 1} (\pi_{k-1} - \pi_k) S_c^2(\beta) (\pi_{k-1} - \pi_k).$$

Taking generalized singular values and integrals, inequality (3.6) gives

$$\lambda^2 \int_0^w \mu_u(\mathbf{1} - \pi) du \leq \int_0^w \mu_u \left(\sum_{k \geq 1} (\pi_{k-1} - \pi_k) S_c^2(\beta) (\pi_{k-1} - \pi_k) \right) du$$

By submajorization and the fact that $\mu(\mathbf{1} - \pi)$ is a characteristic function and therefore is identically equal to 1 on the interval $[0, w]$ by assumption, we have

$$\lambda^2 w \leq \int_0^w \mu_u(S_c^2(\beta)) du \leq 8 \int_0^w \mu_u(S_c^2(x)) du + 2\kappa \int_0^w \left(C^*[\mu(S_c(x))](u) \right)^2 du$$

where the second inequality comes from the submajorization in Lemma 3.2(iv). Using the specific value of λ in (3.1), this leads to

$$\begin{aligned} 9w \int_0^{t^2} \mu_u(S_c^2(x)) \, du + 9w\kappa \int_0^{t^2} \left(C^*[\mu(S_c(x))](u) \right)^2 \, du \\ \leq 8t^2 \int_0^w \mu_u(S_c^2(x)) \, du + 2t^2\kappa \int_0^w \left(C^*[\mu(S_c(x))](u) \right)^2 \, du \\ = I + II. \end{aligned}$$

Next, we estimate I and II separately. For I , we have:

$$\begin{aligned} I &= 8t^2 \int_0^{t^2} \mu_u(S_c^2(x)) \, du + 8t^2 \int_{t^2}^w \mu_u(S_c^2(x)) \, du \\ &\leq 9t^2 \int_0^{t^2} \mu_u(S_c^2(x)) \, du + 8t^2(w - t^2)\mu_{t^2}(S_c^2(x)). \end{aligned}$$

Similarly, II can be estimated as follows:

$$\begin{aligned} II &= 2t^2\kappa \int_0^{t^2} \left(C^*[\mu(S_c(x))](u) \right)^2 \, du + 2t^2\kappa \int_{t^2}^w \left(C^*[\mu(S_c(x))](u) \right)^2 \, du \\ &\leq 9t^2\kappa \int_0^{t^2} \left(C^*[\mu(S_c(x))](u) \right)^2 \, du + 2t^2\kappa \int_{t^2}^w \left(C^*[\mu(S_c(x))](u) \right)^2 \, du \\ &\leq 9t^2\kappa \int_0^{t^2} \left(C^*[\mu(S_c(x))](u) \right)^2 \, du + 2t^2\kappa(w - t^2) \left(C^*[\mu(S_c(x))](t^2) \right)^2 \end{aligned}$$

where in the last estimate we have used the fact that the function $C^*[\mu(S_c(x))]$ is decreasing. Using these estimates on I and II , we obtain after rearrangement and division by $w - t^2$ that

$$9 \int_0^{t^2} \mu_u(S_c^2(x)) \, du + 9\kappa \int_0^{t^2} \left(C^*[\mu(S_c(x))](u) \right)^2 \, du \leq 8t^2\mu_{t^2}(S_c^2(x)) + 2\kappa t^2 \left(C^*[\mu(S_c(x))](t^2) \right)^2.$$

But the left hand side of the preceding inequality is larger than the quantity $9t^2\mu_{t^2}(S_c^2(x)) + 9\kappa t^2 \left(C^*[\mu(S_c(x))](t^2) \right)^2$ which is a contradiction. Thus, we may conclude that $\mu_w(\mathbf{1} - \pi) = 0$. This shows in particular that $\tau(\mathbf{1} - \pi) \leq t^2$. The proof for $\tau(\mathbf{1} - q)$ is identical. \square

Now we proceed with the estimation of the \mathcal{H}_2^c -norm of y . We begin by observing that for $k \geq 1$, it follows from the definitions z and y that dy_k can be split into three separate parts:

$$\begin{aligned} dy_k &= dx_k - d\beta_k\pi_{k-1} \\ &= dx_k(\mathbf{1} - \pi_{k-1}) + dx_k(\mathbf{1} - q_k)\pi_{k-1} + \mathcal{E}_{k-1}(dx_k q_k)\pi_{k-1} \\ &= dx_k(\mathbf{1} - \pi_{k-1}) + dx_k(\mathbf{1} - q_k)\pi_{k-1} + \mathcal{E}_{k-1}[dx_k(q_k - q_{k-1})]\pi_{k-1}. \end{aligned}$$

Using the facts that conditional expectations are contractive projections on L_2 and the functional $\|\cdot\|_2^2$ is convex, we have for each $k \geq 1$,

$$\begin{aligned} \|dy_k\|_2^2 &\leq 4\|dx_k(\mathbf{1} - \pi_{k-1})\|_2^2 + 4\|dx_k(\mathbf{1} - q_k)\pi_{k-1}\|_2^2 + 2\|\mathcal{E}_{k-1}[dx_k(q_k - q_{k-1})]\pi_{k-1}\|_2^2 \\ &\leq 4\|dx_k(\mathbf{1} - \pi_{k-1})\|_2^2 + 4\|dx_k(\mathbf{1} - q_k)\|_2^2 + 2\|dx_k(q_{k-1} - q_k)\|_2^2 \\ &\leq 4\|dx_k(\mathbf{1} - \pi)\|_2^2 + 6\|dx_k(\mathbf{1} - q)\|_2^2. \end{aligned}$$

Taking summation over k , we deduce the following estimate:

$$\begin{aligned}
\|y\|_{\mathcal{H}_2^c}^2 &= \sum_{k \geq 1} \|dy_k\|_2^2 \\
&\leq 4 \sum_{k \geq 1} \|dx_k(\mathbf{1} - \pi)\|_2^2 + 6 \sum_{k \geq 1} \|dx_k(\mathbf{1} - q)\|_2^2 \\
&= 4 \sum_{k \geq 1} \tau[(\mathbf{1} - \pi)|dx_k|^2(\mathbf{1} - \pi)] + 6 \sum_{k \geq 1} \tau[(\mathbf{1} - q)|dx_k|^2(\mathbf{1} - q)] \\
&= 4\tau[(\mathbf{1} - \pi)S_c^2(x)(\mathbf{1} - \pi)] + 6\tau[(\mathbf{1} - q)S_c^2(x)(\mathbf{1} - q)].
\end{aligned}$$

Moreover, it follows from Lemma 3.3 and properties of generalized singular values that

$$\begin{aligned}
\|y\|_{\mathcal{H}_2^c}^2 &\leq 4 \int_0^\infty \mu_u((\mathbf{1} - \pi)S_c^2(x)(\mathbf{1} - \pi)) du + 6 \int_0^\infty \mu_u((\mathbf{1} - q)S_c^2(x)(\mathbf{1} - q)) du \\
&= 4 \int_0^{t^2} \mu_u((\mathbf{1} - \pi)S_c^2(x)(\mathbf{1} - \pi)) du + 6 \int_0^{t^2} \mu_u((\mathbf{1} - q)S_c^2(x)(\mathbf{1} - q)) du \\
&\leq 10 \int_0^{t^2} \mu_u(S_c^2(x)) du.
\end{aligned}$$

Thus, we arrive at the inequality,

$$(3.7) \quad \|y\|_{\mathcal{H}_2^c} \leq \sqrt{10} \left(\int_0^{t^2} \mu_u(S_c^2(x)) du \right)^{1/2}.$$

We now estimate the K -functional using the decomposition $x = y + z$. By combining (3.5) and (3.7), we have

$$\begin{aligned}
K(x, t; \mathcal{H}_2^c(\mathcal{M}), \mathcal{H}_\infty^c(\mathcal{M})) &\leq \|y\|_{\mathcal{H}_2^c} + t\|z\|_{\mathcal{H}_\infty^c} \\
&\leq \sqrt{10} \left(\int_0^{t^2} \mu_u(S_c^2(x)) du \right)^{1/2} + 3\sqrt{5} \left(\int_0^{t^2} \mu_u(S_c^2(x)) du + \kappa \int_0^{t^2} (C^*[\mu(S_c(x))](u))^2 du \right)^{1/2}.
\end{aligned}$$

We can now conclude that

$$K(x, t; \mathcal{H}_2^c(\mathcal{M}), \mathcal{H}_\infty^c(\mathcal{M})) \leq [\sqrt{10} + 3\sqrt{5}(1 + \kappa)^{1/2}] \left(\int_0^{t^2} (\mu_u(S_c(x)) + C^*[\mu(S_c(x))](u))^2 du \right)^{1/2}.$$

The proof is complete. \square

For application purposes, it is important to view the statement of Theorem 3.1 as a comparison between two different K -functionals. We recall that for $f \in L_2(0, \infty) + L_\infty(0, \infty)$ and $t > 0$, we have from [13] the following equivalence:

$$K(f, t; L_2, L_\infty) \approx \left(\int_0^{t^2} (\mu_u(f))^2 du \right)^{1/2}.$$

With this connection, we have the following reformulation of Theorem 3.1.

Remark 3.4. For every $x \in \mathcal{H}_{\Lambda_{\log}}^c(\mathcal{M}) \cap (\mathcal{H}_2^c(\mathcal{M}) + \mathcal{H}_\infty^c(\mathcal{M}))$ and $t > 0$, the following inequality holds:

$$K(x, t; \mathcal{H}_2^c(\mathcal{M}), \mathcal{H}_\infty^c(\mathcal{M})) \leq C_{abs} K(\mu(S_c(x)) + C^*[\mu(S_c(x))], t; L_2, L_\infty).$$

At the time of this writing, we do not know if the use of the operator C^* in the estimate in Theorem 3.1 can be avoided. That is, it is unclear if for all $x \in \mathcal{H}_2^c(\mathcal{M}) + \mathcal{H}_\infty^c(\mathcal{M})$, the equivalence $K(x, t; \mathcal{H}_2^c(\mathcal{M}), \mathcal{H}_\infty^c(\mathcal{M})) \approx K(S_c(x), t; L_2(\mathcal{M}), \mathcal{M})$ holds as in the conditioned case treated in [37]. In other words, it is still an open question if the couple $(\mathcal{H}_2^c(\mathcal{M}), \mathcal{H}_\infty^c(\mathcal{M}))$ is K -closed in the larger couple $(L_2(\mathcal{M} \overline{\otimes} \mathcal{B}(\ell_2)), \mathcal{M} \overline{\otimes} \mathcal{B}(\ell_2))$. Nevertheless, the estimate given in Theorem 3.1 and Remark 3.4 is sufficient to deduce satisfactory results concerning interpolations of the couple $(\mathcal{H}_1^c(\mathcal{M}), \mathcal{H}_\infty^c(\mathcal{M}))$ as we will explore in the next subsection.

3.2. Applications of Theorem 3.1.

3.2.1. *A Peter Jones type interpolation theorem.* Our result in this part constitutes the initial motivation for the paper. It fully resolved the real interpolations for the couple $(\mathcal{H}_1^c(\mathcal{M}), \mathcal{H}_\infty^c(\mathcal{M}))$. It will be deduced from Theorem 3.1 and Wolff's interpolation theorem.

Theorem 3.5. *If $0 < \theta < 1$, $1/p = 1 - \theta$, and $1 \leq \gamma \leq \infty$, then*

$$(\mathcal{H}_1^c(\mathcal{M}), \mathcal{H}_\infty^c(\mathcal{M}))_{\theta, \gamma} = \mathcal{H}_{p, \gamma}^c(\mathcal{M})$$

with equivalent norms.

Proof. We need two steps.

• *Step 1.* This concerns the couple $(\mathcal{H}_2^c(\mathcal{M}), \mathcal{H}_\infty^c(\mathcal{M}))$. Let $0 < \theta < 1$ and $1/q = (1 - \theta)/2$. Fix $y \in (\mathcal{H}_2^c(\mathcal{M}), \mathcal{H}_\infty^c(\mathcal{M}))_{\theta, \gamma}$. We verify first that

$$(3.8) \quad \|y\|_{\mathcal{H}_{q, \gamma}^c} \lesssim \|y\|_{(\mathcal{H}_2^c(\mathcal{M}), \mathcal{H}_\infty^c(\mathcal{M}))_{\theta, \gamma}}.$$

Indeed, since for every symmetric function space E on $(0, \infty)$, the space $\mathcal{H}_E^c(\mathcal{M})$ embeds isometrically into $E(\mathcal{M}; \ell_2^c)$ by the map $x \mapsto (dx_n)_{n \geq 1}$, we have

$$\begin{aligned} \|y\|_{\mathcal{H}_{q, \gamma}^c} &= \|(dy_n)\|_{L_{q, \gamma}(\mathcal{M}; \ell_2^c)} \\ &\approx \|(dy_n)\|_{(L_2(\mathcal{M}; \ell_2^c), L_\infty(\mathcal{M}; \ell_2^c))_{\theta, \gamma}} \\ &\leq \|y\|_{(\mathcal{H}_2^c(\mathcal{M}), \mathcal{H}_\infty^c(\mathcal{M}))_{\theta, \gamma}}. \end{aligned}$$

For the reverse inequality, assume first that $x \in \mathcal{H}_2^c(\mathcal{M}) + \mathcal{H}_\infty^c(\mathcal{M})$ with $S_c(x) \in \Lambda_{\log}(\mathcal{M})$. By Theorem 3.1 (see also Remark 3.4), we have for $1 \leq \gamma \leq \infty$,

$$\begin{aligned} \|x\|_{(\mathcal{H}_2^c(\mathcal{M}), \mathcal{H}_\infty^c(\mathcal{M}))_{\theta, \gamma}} &\lesssim \|\mu(S_c(x)) + C^*[\mu(S_c(x))]\|_{(L_2, L_\infty)_{\theta, \gamma}} \\ &\approx_{q, \gamma} \|\mu(S_c(x)) + C^*[\mu(S_c(x))]\|_{q, \gamma} \\ &\lesssim_{p, \gamma} \|\mu(S_c(x))\|_{q, \gamma} + \|C^*[\mu(S_c(x))]\|_{q, \gamma}. \end{aligned}$$

The important fact here is that since $2 < q < \infty$, we have from Lemma 2.8 and interpolation that $C^* : L_{q, \gamma} \rightarrow L_{q, \gamma}$ is bounded. Therefore, we obtain further that

$$(3.9) \quad \|x\|_{(\mathcal{H}_2^c(\mathcal{M}), \mathcal{H}_\infty^c(\mathcal{M}))_{\theta, \gamma}} \lesssim_{q, \gamma} \|\mu(S_c(x))\|_{q, \gamma} = \|x\|_{\mathcal{H}_{q, \gamma}^c}.$$

Now we remove the extra assumption that $S_c(x) \in \Lambda_{\log}(\mathcal{M})$. Consider an arbitrary $y \in \mathcal{H}_{q, \gamma}^c(\mathcal{M})$. Since \mathcal{M}_1 is semifinite, we may fix an increasing sequence of projections $\{e_j\}$ in \mathcal{M}_1 with $e_j \uparrow \mathbf{1}$ and so that for every $j \geq 1$, $\tau(e_j) < \infty$. For $j \geq 1$, we define the martingale

$$y^{(j)} = (y_n e_j)_{n \geq 1}.$$

The sequence of martingales $(y^{(j)})_{j \geq 1}$ satisfies the following properties:

- (i) for every $j \geq 1$, $y^{(j)} \in \mathcal{H}_{\Lambda_{\log}}^c(\mathcal{M}) \cap \mathcal{H}_{q, \gamma}^c(\mathcal{M})$;

(ii) $\lim_{j \rightarrow \infty} \|y^{(j)} - y\|_{\mathcal{H}_{q,\gamma}^c} = 0$.

We verify first that for every $j \geq 1$, $y^{(j)} \in \mathcal{H}_{q,\gamma}^c(\mathcal{M})$. Indeed, one easily sees that $S_c(y^{(j)}) = (e_j S_c^2(y) e_j)^{1/2} = |S_c(y) e_j|$. It then follows that

$$\|y^{(j)}\|_{\mathcal{H}_{q,\gamma}^c} = \|S_c(y^{(j)})\|_{q,\gamma} \leq \|S_c(y)\|_{q,\gamma} = \|y\|_{\mathcal{H}_{q,\gamma}^c} < \infty.$$

For the Λ_{\log} -case, we have from the definition that for any given $j \geq 1$,

$$\begin{aligned} \|y^{(j)}\|_{\mathcal{H}_{\Lambda_{\log}}^c} &= \int_0^\infty \frac{\mu_t(S_c(y) e_j)}{1+t} dt \\ &= \int_0^{\tau(e_j)} \frac{\mu_t(S_c(y) e_j)}{1+t} dt \\ &\leq \int_0^{\tau(e_j)} \mu_t(S_c(y)) dt \\ &\leq \|y\|_{\mathcal{H}_{q,\gamma}^c} \|\chi_{[0,\tau(e_j)]}(\cdot)\|_{q',\gamma'} < \infty \end{aligned}$$

where the next to last inequality comes from the fact that $L_{q,\gamma}$ is the Köthe dual of $L_{q',\gamma'}$. This verifies the first item. As a consequence, inequality (3.9) applies to $y^{(j)}$ for every $j \geq 1$.

The second item follows at once from $S_c(y^{(j)} - y) = |S_c(y)(\mathbf{1} - e_j)|$. Now, using these two properties and (3.9), we get that $(y^{(j)})_{j \geq 1}$ is a Cauchy sequence in $(\mathcal{H}_2^c(\mathcal{M}), \mathcal{H}_\infty^c(\mathcal{M}))_{\theta,\gamma}$. On the other hand, (3.8) and the second item imply that the limit of the Cauchy sequence $(y^{(j)})_{j \geq 1}$ in $(\mathcal{H}_2^c(\mathcal{M}), \mathcal{H}_\infty^c(\mathcal{M}))_{\theta,\gamma}$ must be y . Thus, by taking limits, we may conclude that:

$$\begin{aligned} \|y\|_{(\mathcal{H}_2^c(\mathcal{M}), \mathcal{H}_\infty^c(\mathcal{M}))_{\theta,\gamma}} &= \lim_{j \rightarrow \infty} \|y^{(j)}\|_{(\mathcal{H}_2^c(\mathcal{M}), \mathcal{H}_\infty^c(\mathcal{M}))_{\theta,\gamma}} \\ &\lesssim_{q,\gamma} \lim_{j \rightarrow \infty} \|y^{(j)}\|_{\mathcal{H}_{q,\gamma}^c} \\ &= \|y\|_{\mathcal{H}_{q,\gamma}^c}. \end{aligned}$$

This shows that (3.9) is valid for any $x \in \mathcal{H}_{q,\gamma}^c(\mathcal{M})$ and combining with (3.8), we conclude that

$$(\mathcal{H}_2^c(\mathcal{M}), \mathcal{H}_\infty^c(\mathcal{M}))_{\theta,\gamma} = \mathcal{H}_{q,\gamma}^c(\mathcal{M}).$$

By reiteration, we may also state the slightly more general conclusion that if $2 < r < \infty$, $0 < v, \gamma \leq \infty$, $0 < \theta < 1$, and $1/q = (1 - \theta)/r$, then

$$(\mathcal{H}_{r,v}^c(\mathcal{M}), \mathcal{H}_\infty^c(\mathcal{M}))_{\theta,\gamma} = \mathcal{H}_{q,\gamma}^c(\mathcal{M}).$$

• *Step 2.* It is already known that the result holds if both endpoints consist of Hardy spaces with finite indices ([29]). Therefore, it suffices to apply Wolff interpolation theorem. More specifically, we use Lemma 2.5. For $1 \leq \gamma \leq \infty$, set $A_{p,\gamma} := \mathcal{H}_{p,\gamma}^c(\mathcal{M})$ when $1 \leq p < \infty$ and $A_{\infty,\gamma} := \mathcal{H}_\infty^c(\mathcal{M})$.

By Step 1, the family $\{A_{p,\gamma}\}_{p,\gamma \in [1,\infty]}$ forms a real-interpolation scale on the interval $I = (2, \infty]$. On the other hand, from the finite indices, the family $\{A_{p,\gamma}\}_{p,\gamma \in [1,\infty]}$ forms a real-interpolation scale on the interval $J = [1, \infty)$. Clearly, $|I \cap J| > 1$. By Lemma 2.5, we conclude that the family $\{A_{p,\gamma}\}_{p,\gamma \in [1,\infty]}$ forms a real-interpolation scale on $I \cup J = [1, \infty]$. This completes the proof. \square

Remark 3.6. The argument used in Step 1 of the proof above can be easily adapted to provide the following more general statement: if $E = (L_2, L_\infty)_{\mathcal{F};K}$ where \mathcal{F} is a Banach function space with monotone norm and the operator C^* is bounded on E , then

$$(3.10) \quad \mathcal{H}_E^c(\mathcal{M}) = (\mathcal{H}_2^c(\mathcal{M}), \mathcal{H}_\infty^c(\mathcal{M}))_{\mathcal{F};K}$$

with equivalent norms.

3.2.2. Extensions to Hardy spaces associated with general symmetric spaces. We have the following result for Hardy spaces associated with symmetric spaces.

Theorem 3.7. *Assume that $E \in \text{Int}(L_2, L_\infty)$ and $C^* : E \rightarrow E$ is bounded. Let \mathcal{G} be a Banach function space with monotone norm. If $F = (E, L_\infty)_{\mathcal{G};K}$ and F is r -concave for some $r < \infty$, then*

$$\mathcal{H}_F^c(\mathcal{M}) = (\mathcal{H}_E^c(\mathcal{M}), \mathcal{H}_\infty^c(\mathcal{M}))_{\mathcal{G};K}$$

with equivalent norms.

The proof is based on the following general reiteration for K -functionals.

Proposition 3.8 ([1]). *Let (A_1, A_0) be a couple of quasi-Banach spaces and \mathcal{F} be a quasi-Banach function space with monotone quasi-norm. If $X = (A_0, A_1)_{\mathcal{F};K}$ and $a \in X + A_1$, then for every $t > 0$,*

$$K(a, \rho(t); X, A_1) \approx I(t, a) + \frac{\rho(t)}{t} K(a, t; A_0, A_1),$$

where $I(t, a) = \|\chi_{(0,t)}(\cdot)K(a, \cdot; A_0, A_1)\|_{\mathcal{F}}$ and $\rho(t) \approx t\|\chi_{(t,\infty)}(\cdot)\|_{\mathcal{F}} + \|u \mapsto u\chi_{(0,t)}(u)\|_{\mathcal{F}}$.

Proof of Theorem 3.7. We begin by verifying that Theorem 3.1 extends to the present situation. That is, if $x \in \mathcal{H}_E^c(\mathcal{M}) + \mathcal{H}_\infty^c(\mathcal{M})$ is such that $S_c(x) \in \Lambda_{\log}(\mathcal{M})$, then for every $s > 0$, the following holds:

$$(3.11) \quad K(x, s; \mathcal{H}_E^c(\mathcal{M}), \mathcal{H}_\infty^c(\mathcal{M})) \leq C_{abs} K(\mu(S_c(x)) + C^*[\mu(S_c(x))], s; E, L_\infty).$$

By assumption, we have from Remark 3.6 that $\mathcal{H}_E^c(\mathcal{M}) = (\mathcal{H}_2^c(\mathcal{M}), \mathcal{H}_\infty^c(\mathcal{M}))_{\mathcal{F};K}$. Therefore by Proposition 3.8, we have for $t > 0$,

$$K(x, \rho(t); \mathcal{H}_E^c(\mathcal{M}), \mathcal{H}_\infty^c(\mathcal{M})) \approx I(t, x) + \frac{\rho(t)}{t} K(x, t; \mathcal{H}_2^c(\mathcal{M}), \mathcal{H}_\infty^c(\mathcal{M}))$$

where $I(t, x) = \|\chi_{(0,t)}(\cdot)K(x, \cdot; \mathcal{H}_2^c(\mathcal{M}), \mathcal{H}_\infty^c(\mathcal{M}))\|_{\mathcal{F}}$. It then follows from Theorem 3.1 and the monotonicity of $\|\cdot\|_{\mathcal{F}}$ that:

$$I(t, x) \lesssim \|\chi_{(0,t)}(\cdot)K(\mu(S_c(x)) + C^*[\mu(S_c(x))], \cdot; L_2, L_\infty)\|_{\mathcal{F}} := \widehat{I}(t, \mu(S_c(x)) + C^*[\mu(S_c(x))]).$$

We can then deduce that

$$\begin{aligned} K(x, \rho(t); \mathcal{H}_E^c(\mathcal{M}), \mathcal{H}_\infty^c(\mathcal{M})) &\lesssim \widehat{I}(t, \mu(S_c(x)) + C^*[\mu(S_c(x))]) + \frac{\rho(t)}{t} K(\mu(S_c(x)) + C^*[\mu(S_c(x))], t; L_2, L_\infty) \\ &\approx K(\mu(S_c(x)) + C^*[\mu(S_c(x))], \rho(t); E, L_\infty). \end{aligned}$$

One can verify as in the proof of [37, Theorem 3.13] that the range of $\rho(\cdot)$ is $[0, \infty)$ which proves (3.11).

In turn, the estimate (3.11) implies that if $x \in \mathcal{H}_E^c(\mathcal{M}) + \mathcal{H}_\infty^c(\mathcal{M})$ is such that $S_c(x) \in \Lambda_{\log}(\mathcal{M})$, then

$$\begin{aligned} \|x\|_{(\mathcal{H}_E^c(\mathcal{M}), \mathcal{H}_\infty^c(\mathcal{M}))_{\mathcal{G};K}} &\lesssim \|\mu(S_c(x)) + C^*[\mu(S_c(x))]\|_{(E, L_\infty)_{\mathcal{G};K}} \\ &\approx \|\mu(S_c(x)) + C^*[\mu(S_c(x))]\|_F. \end{aligned}$$

Next, since F is r -concave, we have $F \in \text{Int}(L_2, L_r)$. It follows from Lemma 2.8 and interpolation that C^* is a bounded operator on F . This implies further that

$$\|x\|_{(\mathcal{H}_E^c(\mathcal{M}), \mathcal{H}_\infty^c(\mathcal{M}))_{\mathcal{G};K}} \leq C_E \|Id + C^* : F \rightarrow F\| \cdot \|x\|_{\mathcal{H}_F^c}.$$

As above, the extra assumption that $x \in \mathcal{H}_{\Lambda_{\log}}^c(\mathcal{M})$ can be removed by approximations for which we omit the details. \square

Our argument above is clearly handicapped by the fact that we only have estimate on the K -functional for the couple $(\mathcal{H}_2^c(\mathcal{M}), \mathcal{H}_\infty^c(\mathcal{M}))$. We suspect that this extra assumption is not necessary. We leave as an open problem that Theorem 3.7 can be improved to cover all spaces $E \in \text{Int}(L_1, L_\infty)$.

As an illustration, we treat the case of martingale Orlicz-Hardy spaces. In this special situation, the restriction in Theorem 3.7 is not needed. We start from recalling that at the level of function spaces, the following result holds:

Proposition 3.9 ([26, Proposition 3.3]). *Let Φ be an Orlicz function, $1 \leq \gamma \leq \infty$, and $0 < \theta < 1$. If $\Phi_0^{-1}(t) = [\Phi^{-1}(t)]^{1-\theta}$, then*

$$(L_\Phi, L_\infty)_{\theta, \gamma} = L_{\Phi_0, \gamma}.$$

By reiteration, we also deduce the following: assume that $0 < \theta, \eta < 1$ and $1 \leq \lambda, \gamma \leq \infty$. Set Ψ_1 and Ψ_2 such that $\Psi_1^{-1}(t) = [\Phi^{-1}(t)]^{1-\theta}$ and $\Psi_2^{-1}(t) = [\Phi^{-1}(t)]^{1-\theta\eta}$. Then

$$(L_\Phi, L_{\Psi_1, \lambda})_{\eta, \gamma} = L_{\Psi_2, \gamma}.$$

Next, we recall that $\mathcal{H}_p^c(\mathcal{M})$ is complemented in $L_p^{\text{ad}}(\mathcal{M}; \ell_2^c)$ for $1 \leq p < \infty$. Since the above identity transfers to the corresponding spaces of adapted sequences (see [37]), it follows that if Φ and Ψ_1 are convex Orlicz functions that are also q -concave for some $1 \leq q < \infty$, then the following holds:

$$(3.12) \quad (\mathcal{H}_\Phi^c(\mathcal{M}), \mathcal{H}_{\Psi_1, \lambda}^c(\mathcal{M}))_{\eta, \gamma} = \mathcal{H}_{\Psi_2, \gamma}^c(\mathcal{M}).$$

We may view this as the Orlicz extension of having finite indices.

The aim of the next result is to show that as in conditioned case, the full equivalence in Proposition 3.9 transfers to martingale Hardy spaces. This provides an extension of Theorem 3.5 to the case of martingale Orlicz-Hardy spaces.

Theorem 3.10. *Let $0 < \theta < 1$ and $1 \leq \gamma \leq \infty$. If Φ is a convex Orlicz function that is q -concave for $1 \leq q < \infty$, then for $\Phi_0^{-1}(t) = [\Phi^{-1}(t)]^{1-\theta}$, the following holds:*

$$(\mathcal{H}_\Phi^c(\mathcal{M}), \mathcal{H}_\infty^c(\mathcal{M}))_{\theta, \gamma} = \mathcal{H}_{\Phi_0, \gamma}^c(\mathcal{M}).$$

Proof. We will consider three cases.

- Case 1. Assume first that Φ is p -convex and q -concave for $2 < p \leq q < \infty$. In this case, $L_\Phi \in \text{Int}(L_2, L_q)$ and the statement follows immediately from Theorem 3.7 and Proposition 3.9. In fact, we have the following slightly more general statement:

$$(\mathcal{H}_{\Phi, \lambda}^c(\mathcal{M}), \mathcal{H}_\infty^c(\mathcal{M}))_{\theta, \gamma} = \mathcal{H}_{\Phi_0, \gamma}^c(\mathcal{M}).$$

- Case 2. Assume that $1/2 < \theta < 1$. One can easily see that Φ_0 is $(1-\theta)^{-1}$ -convex and $q(1-\theta)^{-1}$ -concave. Moreover, $(1-\theta)^{-1} > 2$. Fix $1/2 < \psi < \theta$ and define Φ_1 so that

$$\Phi_1^{-1}(t) = [\Phi^{-1}(t)]^{1-\psi}, \quad t > 0.$$

We note that Φ_1 is $(1-\psi)^{-1}$ -convex with $(1-\psi)^{-1} > 2$ and $q(1-\psi)^{-1}$ -concave. Moreover, $\Phi_0^{-1}(t) = [\Phi_1^{-1}(t)]^{1-\theta_0}$ for $1-\theta_0 = \frac{1-\theta}{1-\psi}$. Applying Case 1 to Φ_1 , we have

$$(\mathcal{H}_{\Phi_1, \lambda}^c(\mathcal{M}), \mathcal{H}_\infty^c(\mathcal{M}))_{\theta_0, \gamma} = \mathcal{H}_{\Phi_0, \gamma}^c(\mathcal{M}).$$

On the other hand, we also have from (3.12) that

$$(\mathcal{H}_{\Phi}^c(\mathcal{M}), \mathcal{H}_{\Phi_0, \gamma}^c(\mathcal{M}))_{\theta_1, \lambda} = \mathcal{H}_{\Phi_1, \lambda}^c(\mathcal{M})$$

where $\theta_1 = \psi/\theta$. By Wolff's interpolation theorem stated in Theorem 2.3 with $B_1 = \mathcal{H}_{\Phi}^c(\mathcal{M})$, $B_2 = \mathcal{H}_{\Phi_1, \lambda}^c(\mathcal{M})$, $B_3 = \mathcal{H}_{\Phi_0, \gamma}^c(\mathcal{M})$, and $B_4 = \mathcal{H}_{\infty}^c(\mathcal{M})$, we conclude that

$$(\mathcal{H}_{\Phi}^c(\mathcal{M}), \mathcal{H}_{\infty}^c(\mathcal{M}))_{\xi, \lambda} = \mathcal{H}_{\Phi_0, \lambda}^c(\mathcal{M})$$

where $\xi = \frac{\theta_0}{1 - \theta_1 + \theta_1 \theta_0}$. A simple calculation shows that $\xi = \theta$.

• Case 3. Assume that $0 < \theta \leq 1/2$. Fix ψ so that $0 < 1 - \psi < \theta \leq 1/2$. Set $\tilde{\Phi}$ satisfying $\tilde{\Phi}^{-1}(t) = [\Phi_0^{-1}(t)]^{1-\psi}$ for $t > 0$. Note that $1/2 < \psi < 1$. Using Case 2 with Φ_0 in place of Φ and $\tilde{\Phi}$ in place of Φ_0 , we get

$$(\mathcal{H}_{\Phi_0, \lambda}^c(\mathcal{M}), \mathcal{H}_{\infty}^c(\mathcal{M}))_{\psi, \lambda} = \mathcal{H}_{\tilde{\Phi}, \lambda}^c(\mathcal{M}).$$

Next, since for every $t > 0$, $\tilde{\Phi}(t)^{-1} = [\Phi^{-1}(t)]^{(1-\theta)(1-\psi)} = [\Phi^{-1}(t)]^{(1-\tilde{\theta})}$, it follows (3.12) that

$$(\mathcal{H}_{\tilde{\Phi}}^c(\mathcal{M}), \mathcal{H}_{\tilde{\Phi}, \lambda}^c(\mathcal{M}))_{\eta, \lambda} = \mathcal{H}_{\Phi_0, \lambda}^c(\mathcal{M})$$

where $\eta = \frac{\theta}{\tilde{\theta}} = \frac{\theta}{\theta + \psi - \psi\theta}$. We use Wolff's interpolation theorem with $B_1 = \mathcal{H}_{\tilde{\Phi}}^c(\mathcal{M})$, $B_2 = \mathcal{H}_{\Phi_0, \lambda}^c(\mathcal{M})$, $B_3 = \mathcal{H}_{\tilde{\Phi}, \lambda}^c(\mathcal{M})$, and $B_4 = \mathcal{H}_{\infty}^c(\mathcal{M})$ to conclude that

$$(\mathcal{H}_{\tilde{\Phi}}^c(\mathcal{M}), \mathcal{H}_{\infty}^c(\mathcal{M}))_{v, \lambda} = \mathcal{H}_{\Phi_0, \lambda}^c(\mathcal{M})$$

where $v = \frac{\eta\psi}{1 - \eta + \eta\psi}$. One can easily verify that $v = \theta$. The proof is complete. \square

We conclude this section with the corresponding result for BMO-spaces. This may be viewed as an Orlicz generalization of the real interpolation form of Musat's result ([29]). To the best of our knowledge, the only available result in the literature is for classical martingale Hardy spaces associated with regular filtration (see [27, Corollary 4.9]). The proof outlined below is based on interpolation of spaces of adapted sequences.

Theorem 3.11. *Let $0 < \theta < 1$ and $1 \leq \gamma \leq \infty$. If Φ is a convex Orlicz function that is q -concave for $1 \leq q < \infty$, then for $\Phi_0^{-1}(t) = [\Phi^{-1}(t)]^{1-\theta}$, the following holds:*

$$(\mathcal{H}_{\Phi}^c(\mathcal{M}), \mathcal{BMO}^c(\mathcal{M}))_{\theta, \gamma} = \mathcal{H}_{\Phi_0, \gamma}^c(\mathcal{M}).$$

Sketch of the proof. Assume first that Φ is p -convex and q -concave for $1 < p < q < \infty$. Denote by Φ^* (resp. Φ_0^*) the Orlicz function complementary to the convex function Φ (resp. Φ_0). Then Φ^* is q' -convex and p' -concave where p' and q' denote the conjugate indices of p and q respectively. In this case, $L_{\Phi^*} \in \text{Int}(L_{q'}, L_{p'})$. A fortiori, $L_{\Phi^*} \in \text{Int}(L_{p'}, L_1)$. Let \mathcal{F} be a Banach function space with monotone norm so that $L_{\Phi^*} = (L_{p'}, L_1)_{\mathcal{F}, K}$. The existence of such \mathcal{F} is given by Proposition 2.7. It follows that $L_{\Phi^*}(\mathcal{M}; \ell_2^c) = (L_{p'}(\mathcal{M}; \ell_2^c), L_1(\mathcal{M}; \ell_2^c))_{\mathcal{F}, K}$. Similarly, $L_{\Phi_0^*}^{\text{ad}}(\mathcal{M}; \ell_2^c) = (L_{p'}^{\text{ad}}(\mathcal{M}; \ell_2^c), L_1^{\text{ad}}(\mathcal{M}; \ell_2^c))_{\mathcal{F}, K}$.

By Proposition 3.8, one can express the K -functionals of the couple $(L_{\Phi^*}(\mathcal{M}; \ell_2^c), L_1(\mathcal{M}; \ell_2^c))$ (resp. $L_{\Phi_0^*}^{\text{ad}}(\mathcal{M}; \ell_2^c), L_1^{\text{ad}}(\mathcal{M}; \ell_2^c)$) in terms of those of the couple $(L_{p'}(\mathcal{M}; \ell_2^c), L_1(\mathcal{M}; \ell_2^c))$ (resp. $(L_{p'}^{\text{ad}}(\mathcal{M}; \ell_2^c), L_1^{\text{ad}}(\mathcal{M}; \ell_2^c))$).

The important fact here is that the couple $(L_{p'}^{\text{ad}}(\mathcal{M}; \ell_2^c), L_1^{\text{ad}}(\mathcal{M}; \ell_2^c))$ is K -closed in the couple $(L_{p'}(\mathcal{M}; \ell_2^c), L_1(\mathcal{M}; \ell_2^c))$ (see [37, Proposition 3.19]). Therefore, we also have that the couple $(L_{\Phi_0^*}^{\text{ad}}(\mathcal{M}; \ell_2^c), L_1^{\text{ad}}(\mathcal{M}; \ell_2^c))$ is K -closed in the larger couple $(L_{\Phi^*}(\mathcal{M}; \ell_2^c), L_1(\mathcal{M}; \ell_2^c))$.

From K -closedness, we deduce that if $(L_{\Phi^*}, L_1)_{\theta, \gamma'} = E$, then

$$(L_{\Phi^*}^{\text{ad}}(\mathcal{M}; \ell_2^c), L_1^{\text{ad}}(\mathcal{M}; \ell_2^c))_{\theta, \gamma'} = E^{\text{ad}}(\mathcal{M}; \ell_2^c).$$

By complementation, we obtain further that

$$(\mathcal{H}_{\Phi^*}^c(\mathcal{M}), \mathcal{H}_1^c(\mathcal{M}))_{\theta, \gamma'} = \mathcal{H}_E^c(\mathcal{M}).$$

With the facts that $\mathcal{H}_{\Phi^*}^c(\mathcal{M}) = (\mathcal{H}_{\Phi}^c(\mathcal{M}))^*$, $\mathcal{H}_{E^*}^c(\mathcal{M}) = (\mathcal{H}_E^c(\mathcal{M}))^*$, and $(\mathcal{H}_1^c(\mathcal{M}))^* = \mathcal{BMO}^c(\mathcal{M})$, we may apply duality (see [6, Theorem 3.7.1]) to deduce that

$$(\mathcal{H}_{\Phi}^c(\mathcal{M}), \mathcal{BMO}^c(\mathcal{M}))_{\theta, \gamma} = \mathcal{H}_{E^*}^c(\mathcal{M}).$$

Observe that $E^* = (L_{\Phi}, L_{\infty})_{\theta, \gamma} = L_{\Phi_0, \gamma}$. This proves the theorem for the case Φ being p -convex with $p > 1$.

For general convex function Φ as stated, one can repeat the argument used in Case 2 of the proof of Theorem 3.10 (but without the restriction $1/2 < \theta < 1$). We leave the details to the interested reader. \square

4. CONCLUDING REMARKS

By taking adjoints, all results from the previous section are valid for noncommutative martingale row Hardy spaces.

Recall that the mixed martingale Hardy spaces are defined as follows: for $E \in \text{Int}(L_p, L_2)$ with $0 < p < 2$,

$$\mathcal{H}_E(\mathcal{M}) = \mathcal{H}_E^c(\mathcal{M}) + \mathcal{H}_E^r(\mathcal{M})$$

while for $F \in \text{Int}(L_2, L_q)$ with $2 < q \leq \infty$,

$$\mathcal{H}_F(\mathcal{M}) = \mathcal{H}_F^c(\mathcal{M}) \cap \mathcal{H}_F^r(\mathcal{M}).$$

Using similar argument as in the proof of [2, Theorem 4.5], we may also deduce the corresponding interpolation result for mixed Hardy spaces which reads as follows:

Theorem 4.1. *If $0 < \theta < 1$, $1/p = 1 - \theta$, and $1 \leq \gamma \leq \infty$, then*

$$(\mathcal{H}_1(\mathcal{M}), \mathcal{H}_{\infty}(\mathcal{M}))_{\theta, \gamma} = \mathcal{H}_{p, \gamma}(\mathcal{M})$$

with equivalent norms.

Motivated by results from the previous section and the Musat's result on the complex interpolation of $(\mathcal{H}_1^c(\mathcal{M}), \mathcal{BMO}^c(\mathcal{M}))$, a natural direction of interest is the complex interpolation method for the couple $(\mathcal{H}_1^c(\mathcal{M}), \mathcal{H}_{\infty}^c(\mathcal{M}))$ (or the couple $(\mathcal{H}_1(\mathcal{M}), \mathcal{H}_{\infty}(\mathcal{M}))$). For a given compatible couple of Banach spaces (A_0, A_1) and $0 < \theta < 1$, let $[A_0, A_1]_{\theta}$ denote the complex interpolation space of exponent θ as defined in [6]. The following question remains unresolved.

Problem 4.2. Assume that $1 < \theta < 1$ and $1/p = 1 - \theta$. Does one have $[\mathcal{H}_1^c(\mathcal{M}), \mathcal{H}_{\infty}^c(\mathcal{M})]_{\theta} = \mathcal{H}_p^c(\mathcal{M})$?

We should point out that the corresponding problem for the conditioned case is still open (see [2, Problem 5]).

Acknowledgments. I am very grateful to the anonymous referee for a careful reading of the paper and for providing valuable suggestions that improved the presentation of the paper.

REFERENCES

- [1] I. Ahmed, G. E. Karadzhov, and A. Raza, *General Holmstedt's formulae for the K -functional*, J. Funct. Spaces (2017), Art. ID 4958073, 9pp. MR 3614335
- [2] T. Bekjan, Z. Chen, M. Perrin, and Z. Yin, *Atomic decomposition and interpolation for Hardy spaces of noncommutative martingales*, J. Funct. Anal. **258** (2010), no. 7, 2483–2505. MR 2584751 (2011d:46131)
- [3] T. N. Bekjan, *Interpolation of noncommutative symmetric martingale spaces*, J. Operator Theory. **77** (2017), no. 2, 245–259. MR 3634506
- [4] T. N. Bekjan, Z. Chen, M. Raikhan, and M. Sun, *Interpolation and the John-Nirenberg inequality on symmetric spaces of noncommutative martingales*, Studia Math. **262** (2022), no. 3, 241–273. MR 4358459
- [5] C. Bennett and R. Sharpley, *Interpolation of operators*, Academic Press Inc., Boston, MA, 1988. MR 89e:46001
- [6] J. Bergh and J. Löfström, *Interpolation spaces. An introduction*, Springer-Verlag, Berlin, 1976, Grundlehren der Mathematischen Wissenschaften, No. 223. MR MR0482275 (58 #2349)
- [7] Z. Chen, N. Randrianantoanina, and Q. Xu, *Atomic decompositions for noncommutative martingales*, J. Funct. Anal. **284** (2023), no. 9, Paper No. 109877, 47pp.
- [8] I. Cuculescu, *Martingales on von Neumann algebras*, J. Multivariate Anal. **1** (1971), 17–27. MR 45 #4464
- [9] P. G. Dodds, T. K. Dodds, and B. de Pagter, *Noncommutative Köthe duality*, Trans. Amer. Math. Soc. **339** (1993), 717–750. MR 94a:46093
- [10] T. Fack and H. Kosaki, *Generalized s -numbers of τ -measurable operators*, Pacific J. Math. **123** (1986), 269–300. MR 87h:46122
- [11] Z. Hao and L. Li, *Orlicz-Lorentz Hardy martingale spaces*, J. Math. Anal. Appl. **482** (2020), no. 1, 123520, pp 27. MR 4013836
- [12] G. H. Hardy, J. E. Littlewood, and G. Pólya, *Inequalities*, Cambridge Mathematical Library, Cambridge University Press, Cambridge, 1988, Reprint of the 1952 edition.
- [13] T. Holmstedt, *Interpolation of quasi-normed spaces*, Math. Scand. **26** (1970), 177–199. MR 54 #3440
- [14] S. Janson and P. W. Jones, *Interpolation between H^p spaces: the complex method*, J. Funct. Anal. **48** (1982), no. 1, 58–80. MR 671315
- [15] Y. Jiao, F. Sukochev, L. Wu, and D. Zanin, *Distributional inequalities for noncommutative martingales*, J. Funct. Anal. **284** (2023), no. 5, Paper No. 109798, 55pp. MR 4521734
- [16] P. W. Jones, *L^∞ estimates for the $\bar{\partial}$ problem in a half-plane*, Acta Math. **150** (1983), no. 1-2, 137–152. MR 697611
- [17] ———, *On interpolation between H^1 and H^∞* , Interpolation spaces and allied topics in analysis (Lund, 1983), Lecture Notes in Math., vol. 1070, Springer, Berlin, 1984, pp. 143–151. MR 760480
- [18] M. Junge, *Doob's inequality for non-commutative martingales*, J. Reine Angew. Math. **549** (2002), 149–190. MR 2003k:46097
- [19] M. Junge and Q. Xu, *Noncommutative Burkholder/Rosenthal inequalities*, Ann. Probab. **31** (2003), no. 2, 948–995. MR 2004f:46078
- [20] ———, *On the best constants in some non-commutative martingale inequalities*, Bull. London Math. Soc. **37** (2005), no. 2, 243–253. MR 2005k:46170
- [21] N. Kalton and S. Montgomery-Smith, *Interpolation of Banach spaces*, Handbook of the geometry of Banach spaces, Vol. 2, North-Holland, Amsterdam, 2003, pp. 1131–1175. MR 1999193
- [22] N. J. Kalton and F. A. Sukochev, *Symmetric norms and spaces of operators*, J. Reine Angew. Math. **621** (2008), 81–121. MR 2431251 (2009i:46118)
- [23] S. V. Kislyakov and Q. Xu, *Real interpolation and singular integrals*, Algebra i Analiz **8** (1996), no. 4, 75–109, translation in St. Petersburg Math. J. **8** (1997), no. 4, 593–615. MR 1418256
- [24] M. A. Krasnosel'skiĭ and J. B. Rutickiĭ, *Convex functions and Orlicz spaces*, Translated from the first Russian edition by Leo F. Boron, P. Noordhoff Ltd., Groningen, 1961. MR 0126722 (23 #A4016)
- [25] S. G. Kreĭn, Yu. Ī. Petunin, and E. M. Semĕnov, *Interpolation of linear operators*, Translations of Mathematical Monographs, vol. 54, American Mathematical Society, Providence, R.I., 1982, Translated from the Russian by J. SzHucs. MR 649411
- [26] L. Long, H. Tian, and D. Zhou, *Interpolation of martingale Orlicz-Hardy spaces*, Acta Math. Hungar. **163** (2021), no. 1, 276–294. MR 4217969
- [27] L. Long, F. Weisz, and G. Xie, *Real interpolation of martingale Orlicz Hardy spaces and BMO spaces*, J. Math. Anal. Appl. **505** (2022), no. 2, Paper No. 125565, 23. MR 4300991

- [28] L. Maligranda, *Orlicz spaces and interpolation*, Seminários de Matemática [Seminars in Mathematics], vol. 5, Universidade Estadual de Campinas, Departamento de Matemática, Campinas, 1989. MR 2264389 (2007e:46025)
- [29] M. Musat, *Interpolation between non-commutative BMO and non-commutative L_p -spaces*, J. Funct. Anal. **202** (2003), no. 1, 195–225. MR 1994770
- [30] J. Parcet and N. Randrianantoanina, *Gundy's decomposition for non-commutative martingales and applications*, Proc. London Math. Soc. (3) **93** (2006), no. 1, 227–252. MR MR2235948
- [31] G. Pisier, *Interpolation between H^p spaces and noncommutative generalizations. I*, Pacific J. Math. **155** (1992), no. 2, 341–368. MR 1178030
- [32] G. Pisier and Q. Xu, *Non-commutative martingale inequalities*, Comm. Math. Phys. **189** (1997), 667–698. MR 98m:46079
- [33] ———, *Non-commutative L^p -spaces*, Handbook of the geometry of Banach spaces, Vol. 2, North-Holland, Amsterdam, 2003, pp. 1459–1517. MR 2004i:46095
- [34] Y. Qiu, *A non-commutative version of Lépingle-Yor martingale inequality*, Statist. Probab. Lett. **91** (2014), 52–54. MR 3208115
- [35] N. Randrianantoanina, *Non-commutative martingale transforms*, J. Funct. Anal. **194** (2002), 181–212. MR 2003m:46098
- [36] ———, *Interpolation between noncommutative martingale Hardy and BMO spaces: the case $0 < p < 1$* , Canad. J. Math. **74** (2022), no. 6, 1700–1744. MR 4520665
- [37] ———, *P. Jones' interpolation theorem for noncommutative martingale Hardy spaces*, Trans. Amer. Math. Soc. **376** (2023), no. 3, 2089–2124. MR 4549700
- [38] F. Sukochev, *Completeness of quasi-normed symmetric operator spaces*, Indag. Math. (N.S.) **25** (2014), no. 2, 376–388. MR 3151823
- [39] F. Weisz, *Interpolation between martingale Hardy and BMO spaces, the real method*, Bull. Sci. Math. **116** (1992), no. 2, 145–158. MR 1168308
- [40] ———, *Martingale Hardy spaces and their applications in Fourier analysis*, Lecture Notes in Mathematics, vol. 1568, Springer-Verlag, Berlin, 1994. MR MR1320508 (96m:60108)
- [41] T. H. Wolff, *A note on interpolation spaces*, Harmonic analysis (Minneapolis, Minn., 1981), Lecture Notes in Math., vol. 908, Springer, Berlin-New York, 1982, pp. 199–204. MR 654187
- [42] Q. Xu, *Analytic functions with values in lattices and symmetric spaces of measurable operators*, Math. Proc. Cambridge Philos. Soc. **109** (1991), 541–563. MR 92g:46036
- [43] ———, *Some results related to interpolation on Hardy spaces of regular martingales*, Israel J. Math. **91** (1995), no. 1-3, 173–187. MR 1348311

DEPARTMENT OF MATHEMATICS, MIAMI UNIVERSITY, OXFORD, OHIO 45056, USA

Email address: randrin@miamioh.edu