OVER-MAHONIAN NUMBERS: BASIC PROPERTIES AND UNIMODALITY

ALI KESSOURI, MOUSSA AHMIA, AND SALIM MESBAHI

ABSTRACT. In this paper, we propose the number of permutations of length n having k overlined inversions, which we call over-Mahonian number. We study useful properties and some combinatorial interpretations by lattice paths/overpartitions and tilings. Furthermore, we prove combinatorially that these numbers form a log-concave sequence and therefore unimodal.

1. INTRODUCTION

Let [n] denote the set $\{1, 2, ..., n\}$. Let $\sigma = \sigma_1 \sigma_2 \cdots \sigma_n$ be a permutation of length n, i.e., a permutation of [n]. The group of all the permutations of [n] is the symmetric group S_n . The identity element of S_n is the permutation $\iota = \iota_1 \iota_2 \cdots \iota_n$ where $\iota_i = i$ for all $1 \le i \le n$.

Definition 1.1. A pair (σ_i, σ_j) is called an **inversion** of the permutation σ if i < j and $\sigma_i > \sigma_j$.

Example 1.2. The permutation 2431 has 4 inversions: (2, 1), (4, 1), (4, 3) and (3, 1).

Remark 1.3. The only permutation with no inversion is the identity element ι , and the permutation of length n with the maximum number of inversions is the permutation $\alpha = \alpha_1 \alpha_2 \cdots \alpha_n$ where $\alpha_i = n + 1 - i$, which has $\binom{n}{2}$ inversions. Moreover the number of inversions of a permutation and its inverse are equal.

The backward permutation of σ is the permutation σ' where $\sigma'_i = \sigma_{n+1-i}$ for all $1 \le i \le n$.

A pair (σ_i, σ_j) is called a backward inversion of the permutation σ if i < j and $\sigma_i < \sigma_j$. If a permutation of length n has k inversions then it has $\binom{n}{2} - k$ backward inversions [10].

The major index of a permutation σ of length n is the sum of indices i that satisfy $\sigma_i > \sigma_{i+1}$. For instance the permutation 4231 has major index 1 + 3 = 4.

The enumeration of permutations of length n according to their number of inversions, and the study of numbers i(n, k) of permutations of length n having k inversions, is a

Date: June 18, 2024.

²⁰¹⁰ Mathematics Subject Classification. 05A05, 05A15.

Key words and phrases. Over-Mahonian numbers, inversions, permutations, lattice paths, overpartitions, tilings.

classic area of combinatorics. The best-known result is the following [12]:

(1)
$$\sum_{k=0}^{\binom{n}{2}} i(n,k)x^k = (1+x)\cdots(1+x+\cdots+x^{n-1}),$$

where i(n, k) is known as the Mahonian number. MacMahon [11] proved that this number is equal to the number of permutations of length n with major index equal to k.

Our paper consists of three sections. In the first section, we define the over-Mahonian numbers, and we give the combinatorial prove of some recurrence relations of these numbers and some identities. Furthermore, we give the generating function of the over-Mahonian numbers and we deduce an important identity in terms of the double factorial. In the second section, we give combinatorial interpretations for the over-Mahonian numbers using lattice paths/overpartitions and tilings. Finally, in the last section, we prove combinatorially that these numbers form a log-concave sequence and therefore unimodal, by constructing the appropriate injection.

2. Basic properties of over-Mahonian numbers

First of all, we propose the following definition.

Definition 2.1. Let (σ_i, σ_j) be an inversion of the permutation σ . We say (σ_i, σ_j) is an **overlined inversion** if the entry σ_i is overlined or not.

According to this definition, we propose the definition of the following number.

Definition 2.2. Let $\overline{\binom{n}{k}}^{\uparrow}$ be the number of permutations of length n with k overlined inversions. We call this number **over-Mahonian number**.

Example 2.3. The permutations of 3 with 2 overlined inversions are:

$$312, \overline{3}12, 231, \overline{2}31, 2\overline{3}1, \overline{2}\overline{3}1.$$

So
$$\overline{\begin{bmatrix} 3\\2 \end{bmatrix}}^{\uparrow} = 6.$$

From Definition 2.2, we can establish that the over-Mahonian number $\overline{\binom{n}{k}}^{\uparrow}$ satisfies the following recurrence relation.

Theorem 2.4. For positive integers n and k,

(2)
$$\overline{\begin{bmatrix}n\\k\end{bmatrix}}^{\uparrow} = \overline{\begin{bmatrix}n-1\\k\end{bmatrix}}^{\uparrow} + 2\sum_{j=1}^{n-1} \overline{\begin{bmatrix}n-1\\k-j\end{bmatrix}}^{\uparrow},$$

where $\overline{\begin{bmatrix}n\\0\end{bmatrix}}^{\uparrow} = 1$ and $\overline{\begin{bmatrix}n\\k\end{bmatrix}}^{\uparrow} = 0$ unless $0 \le k \le {\binom{n}{2}}$.

For any permutation $\sigma = \sigma_1 \cdots \sigma_n$ of length n having k overlined inversions. We remove σ_1 form σ we have either a permutation of length n-1 having k overlined inversions if the entry σ_1 does not form an overlined inversion with the others which corresponds to $\overline{\binom{n-1}{k}}^{\uparrow}$, or a permutation of length n-1 having k-j overlined inversions if the entry σ_1 is not overlined and form j overlined inversions, for $1 \leq j \leq n-1$, with the others or the entry σ_1 is overlined and form j overlined inversions with the others, then we obtain for the both $\overline{\binom{n-1}{k-j}}^{\uparrow}$, that gives us $2\sum_{j=1}^{n-1} \overline{\binom{n-1}{k-j}}^{\uparrow}$, and therefore the equality (2).

These numbers build a triangle which we call "over-Mahonian triangle", see Table 1.

$n \backslash k$	0	1	2	3	4	5	6	7	8	9	10
1	1										
2	1	2									
3	1	4	6	4							
4	1	6	16	26	28	20	8				
5	1	8	30	72	126	172	188	164	112	56	16

Table 1: The over-Mahonian triangle.

From Theorem 2.4, we establish that the over-Mahonian numbers $\overline{\binom{n}{k}}^{\uparrow}$ satisfy the following recurrence relation.

Proposition 2.5. For $0 \le k \le {n \choose 2}$, we have

(3)
$$\overline{\binom{n}{k}}^{\uparrow} = \overline{\binom{n}{k-1}}^{\uparrow} + \overline{\binom{n-1}{k}}^{\uparrow} + \overline{\binom{n-1}{k-1}}^{\uparrow} - 2\overline{\binom{n-1}{k-n}}^{\uparrow}.$$

Proof. By relation (2), we have

(4)
$$\overline{\binom{n}{k}}^{\uparrow} = \overline{\binom{n-1}{k}}^{\uparrow} + 2\overline{\binom{n-1}{k-1}}^{\uparrow} + \dots + 2\overline{\binom{n-1}{k-n+1}}^{\uparrow}$$

and

(5)
$$\overline{\binom{n}{k-1}}^{\uparrow} = \overline{\binom{n-1}{k-1}}^{\uparrow} + 2\overline{\binom{n-1}{k-2}}^{\uparrow} + \dots + 2\overline{\binom{n-1}{k-n}}^{\uparrow}.$$

It follows from relation (5) that

$$\overline{\begin{bmatrix}n\\k\end{bmatrix}}^{\uparrow} = \overline{\begin{bmatrix}n-1\\k\end{bmatrix}}^{\uparrow} + \overline{\begin{bmatrix}n-1\\k-1\end{bmatrix}}^{\uparrow} + \left(\overline{\begin{bmatrix}n-1\\k-1\end{bmatrix}}^{\uparrow} + 2\overline{\begin{bmatrix}n-1\\k-2\end{bmatrix}}^{\uparrow} + \dots + 2\overline{\begin{bmatrix}n-1\\k-n\end{bmatrix}}^{\uparrow}\right) - 2\overline{\begin{bmatrix}n-1\\k-n\end{bmatrix}}^{\uparrow}$$
$$= \overline{\begin{bmatrix}n-1\\k\end{bmatrix}}^{\uparrow} + \overline{\begin{bmatrix}n-1\\k-1\end{bmatrix}}^{\uparrow} + \overline{\begin{bmatrix}n-1\\k-1\end{bmatrix}}^{\uparrow} + \overline{\begin{bmatrix}n-1\\k-1\end{bmatrix}}^{\uparrow} - 2\overline{\begin{bmatrix}n-1\\k-n\end{bmatrix}}^{\uparrow}$$
$$= \overline{\begin{bmatrix}n\\k-1\end{bmatrix}}^{\uparrow} + \overline{\begin{bmatrix}n-1\\k\end{bmatrix}}^{\uparrow} + \overline{\begin{bmatrix}n-1\\k-1\end{bmatrix}}^{\uparrow} - 2\overline{\begin{bmatrix}n-1\\k-n\end{bmatrix}}^{\uparrow}.$$
As desired.

As desired.

From Definition 2.2, we have also the following identities.

Proposition 2.6. For any integer $n \ge 1$, we have

(1)
$$-\overline{\begin{bmatrix} n\\ n\\ 2 \end{bmatrix}}^{\dagger} = 2^{n-1}.$$

(2) $-\overline{\begin{bmatrix} n\\ 1 \end{bmatrix}}^{\dagger} = 2 \times (n-1).$
(3) $-\overline{\begin{bmatrix} n\\ k \end{bmatrix}}^{\dagger}$ is even if $n \ge 2$ and $k \ge 1$

(1) $-\overline{\binom{n}{\binom{n}{2}}}^{\uparrow}$ counts the number of permutations of length *n* having $\binom{n}{2}$ overlined Proof. inversions, this means that $\sigma_1 > \cdots > \sigma_{n-1} > \sigma_n$ such that the entries $\sigma_1, \ldots, \sigma_{n-1}$ are overlined or not, that gives us 2^{n-1} possible permutations.

- (2) $-\overline{\begin{bmatrix}n\\1\end{bmatrix}}^{\uparrow}$ counts the number of permutations of length *n* having one overlined inversion, this means that $\sigma_1 < \cdots < \sigma_j > \sigma_{j+1} < \cdots < \sigma_n$ such that σ_j is overlined or not, for $1 \leq j \leq (n-1)$, that gives us $2 \times (n-1)$ possible permutations.
- (3) If $n \ge 2$ and $k \ge 1$, we can necessarily have overlined inversions, and for each inversion (σ_i, σ_j) we have two possibilities: either the entry σ_i is overlined or not overlined, then the total number of permutations of length n having k overlined inversions is even.

The over-Mahonian numbers $\overline{\binom{n}{k}}^{\uparrow}$ have the following row generating function.

Theorem 2.7. For every positive integer n,

$$\sum_{k=0}^{\binom{n}{2}} \overline{\binom{n}{k}}^{\uparrow} z^k = \begin{cases} 1, & \text{if } n = 1, \\ (1+2z)\cdots(1+2z+\cdots+2z^{n-1}), & \text{if } n > 1. \end{cases}$$

Proof. Let

$$f_n(z) = \sum_{k=0}^{\binom{n}{2}} \overline{\binom{n}{k}}^{\uparrow} z^k$$

be the row generating function of the over-Mahonian number.

The case of n = 1 is trivial. For $n \ge 2$, using (2) we obtain

$$\begin{split} f_n(z) &= \sum_{k=0}^{\binom{n}{2}} \overline{\binom{n}{k}}^{\uparrow} z^k = \sum_{k=0}^{\binom{n}{2}} \left(\overline{\binom{n-1}{k}}^{\uparrow} + 2\sum_{j=1}^{n-1} \overline{\binom{n-1}{k-j}}^{\uparrow} \right) z^k \\ &= \sum_{k=0}^{\binom{n}{2}} \overline{\binom{n-1}{k}}^{\uparrow} z^k + 2\sum_{k=0}^{\binom{n}{2}} \sum_{j=1}^{n-1} \overline{\binom{n-1}{k-j}}^{\uparrow} z^k \\ &= f_{n-1}(z) + 2\sum_{j=1}^{n-1} z^j \sum_{k=j}^{\binom{n}{2}-j} \overline{\binom{n-1}{k-j}}^{\uparrow} z^{k-j} \\ &= f_{n-1}(z) + 2\sum_{j=1}^{n-1} z^j \sum_{k=0}^{\binom{n-1}{2}} \overline{\binom{n-1}{k}}^{\uparrow} z^k \\ &= f_{n-1}(z) \left(1 + 2\sum_{j=1}^{n-1} z^j\right). \end{split}$$

Iterating, we obtain

$$f_n(z) = (1+2z)\cdots(1+2z+\cdots+2z^{n-1}).$$

By setting z = 1 in Theorem 2.7, we obtain the following interested result.

Corollary 2.8. For positive integer n, the over-Mahonian numbers satisfy the following identity:

$$\sum_{k=0}^{\binom{n}{2}} \overline{\begin{bmatrix} n \\ k \end{bmatrix}}^{\uparrow} = (2n-1)!!,$$

where $(2n-1)!! = 1 \times 3 \times \cdots \times (2n-1)$.

3. Combinatorial interpretations of the over-Mahonian numbers

In this section, we give combinatorial interpretations for the over-Mahonian numbers using lattice paths/overpartitions and tilings.

3.1. A lattice path interpretation. Ghemit and Ahmia in [8] established that the Mahonian number i(n, k) counts the number of lattice paths from $u_1 = (0, 0)$ to $v_1 = (n-1, k)$ taking at most j North steps at the level j.

By the same approach, we give in this subsection a combinatorial interpretation for the over-Mahonian numbers $\frac{\left[n\right]^{\uparrow}}{\left[k\right]^{\uparrow}}$ by lattice path.

Let $\mathcal{P}_{n,k}^{\uparrow}$ denote the set of lattice paths from the point (0,0) to the point (n-1,k), with only North steps (vertical steps = (0,1)), East steps (horizontal steps = (1,0)) and

North-East steps (diagonal steps = (1, 1)), such that the number of North steps in each level j is at most j if there is no a diagonal step before and at most j - 1 North steps after the diagonal step, where the levels associated to vertical lines are from 0 to n - 1 as shown in the example of Figure 1.




FIGURE 1. A path P in $\mathcal{P}_{4.6}^{\uparrow}$.

Theorem 3.1. The over-Mahonian number $\overline{{\binom{n}{k}}}^{\uparrow}$ counts the number of lattice paths from $u_1 = (0,0)$ to $v_1 = (n-1,k)$ taking at most j North steps at the level j if there is no a diagonal step before, and at most (j-1) North steps at the level j after a diagonal step. That is

$$\overline{\begin{bmatrix}n\\k\end{bmatrix}}^{\uparrow} = \mid \mathcal{P}_{n,k}^{\uparrow} \mid$$

Proof. Since $\overline{\binom{n}{k}}^{\uparrow}$ counts the number of permutations of length n having k overlined inversions, here it suffices to prove that there exists a bijection between these permutations and the lattice paths of $\mathcal{P}_{n,k}^{\uparrow}$. So we do this as follows:

For each path $P \in \mathcal{P}_{n,k}^{\uparrow}$, we can easily find the permutation associated to P, this permutation is obtained as follows : we associate to the point (0,0) the entry 1, the first step is necessarily an East step or a North-East step. If this step is an East step, we move to the point (1,0) and we add to this point the entry 2 on the right of 1 (i.e., 12) and if this step is a North-East step, we move to the point (1,1) and we add in this point the entry $\overline{2}$ to the left of 1 (i.e., $\overline{2}1$, which gives us an inversion). At the point (1,0), we have three cases: if the next step of P is an East step, we add the entry $\overline{3}$ to the right of 2 (i.e., 123), if the next step of P is an North-East step, we add the entry $\overline{3}$ to the right of 2 and we move the entry $\overline{3}$ one position to the left of 2 (i.e., 1 $\overline{3}2$, which gives us an inversion), and if the next step of P is an North step, here we move the entry 2 one position to the left (i.e., 21, which gives us an inversion). At the point (1,1), here we have two cases: if the next step of P is an North step, here we move the entry 2 one position to the left (i.e., 21, which gives us an inversion). At the point (1,1), here we have two cases: if the

next step of P is an East step, we add the entry 3 to the right of 1 (i.e., $\overline{2}13$), **if** the next step of P is an North-East step, here we add the entry $\overline{3}$ to the right of 1 and we move the entry $\overline{3}$ one position to the left of 1 (i.e., $\overline{2}\overline{3}1$, which gives us an inversion). We proceed the same operations with next points until we arrive to the point (n-1,k), that gives us the desired permutation. An example is shown in Figure 2.




FIGURE 2. The path associated to the permutation $\sigma = \overline{4321}$.

3.2. An overpartition interpretation. In this subsection, we give a combinatorial interpretations for the over-Mahonian numbers by overpartitions.

First of all, we start by the following two definitions.

Definition 3.2. A partition $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ of *n* is a non-increasing sequence of positive integers (i.e., $\lambda_1 \ge \lambda_2 \ge \dots \ge \lambda_k$) whose sum is *n*, and p(n) denotes the number of partitions of *n*. For convenience, define p(0) = 1.

For $1 \leq i \leq k$, λ_i is called a part of λ . Although we considered each part in a partition is positive, we sometimes allow "zero" as a part.

The length of λ denoted $l(\lambda)$ is the number of parts of λ ; and the weight of λ denoted $|\lambda|$ is the sum of the parts of λ .

Definition 3.3. [7, Corteel and Lovejoy] An **overpartition** of n is a non-increasing sequence of natural numbers whose sum is n in which the first occurrence of a number may be overlined and $\overline{p}(n)$ denotes the number of overpartitions of n. For convenience, define $\overline{p}(0) = 1$.

For example, there are 8 overpartitions of 3 enumerated by:


 $3,\overline{3},2+1,\overline{2}+1,2+\overline{1},\overline{2}+\overline{1},1+1+1,\overline{1}+1+1.$

Since the bijection between lattice paths and tiling is weight-preserving, we obtain the following overpartition interpretation of the over-Mahonian numbers.

Theorem 3.4. The over-Mahonian number $\overline{\binom{n}{k}}^{\uparrow}$ counts the number of overpartitions into k parts in which each part j must be used at most j times if the part j is not overlined before and at most j-1 times if the part j is overlined before, and the largest part $\leq n-1$.

Proof. Since $\overline{\binom{n}{k}}^{\uparrow}$ counts the number of lattice paths from (0,0) to (n-1,k) taking at most j North steps at the level j if there is no a diagonal step before, and at most j-1 North steps at the level j after a diagonal step. So, if we take the parts (resp. the overlined parts) as the number of the cases above each North step (resp. North-East step) in the paths associated to $\overline{\binom{n}{k}}^{\uparrow}$, then we can easily see that the over-Mahonian number $\overline{\binom{n}{k}}^{\uparrow}$ counts the number of overpartition into k parts in which each part j must be used at most j times if the part j is not overlined before and at most j-1 times if the part j is overlined before, and the largest part $\leq n-1$.

For example, Figure 3 shows the 16 overpartitions into k = 2 parts and the largest part is n - 1 = 3.






FIGURE 3. All lattice paths/overpartitions for n = 4 and k = 2.

3.3. A tiling interpretation. Ghemit and Ahmia [9] proved also that the Mahonian number i(n, k) counts the number of ways to tile a board of length (n + k - 1) using k red squares and n - 1 blue squares with at most j red squares successively if there are j blue squares before. Motivated by their work, we give in this subsection a combinatorial interpretation for the over-Mahonian numbers using the same combinatorial objects.

Let $\mathcal{T}_{n,k}^{\uparrow}$ be the set of all tilings of a board of length (n + k - 1) using only at most n - 1 blue squares, at most k red squares and black rectangles with an area of two squares (square pair), where the number of successive red squares is at most j if there is j blue squares before, and the number of successive red squares is at most j - 1 after black rectangle, and it is necessary that the number of red squares plus the number of black rectangle is equal to k, and the number of blue squares plus the number of black rectangle is equal to n - 1. The black rectangle is considered as blue square and red square in that order.

Remark 3.5. There is a bijection between the sets $\mathcal{P}_{n,k}^{\uparrow}$ and $\mathcal{T}_{n,k}^{\uparrow}$ indeed, each East step is replaced by blue square and North step is replaced by red square and diagonal step is replaced by black rectangle. See Figure 4.




FIGURE 4. The tiling/lattice path associated to the permutation $\sigma = 2\overline{3}1$.

According to this remark, we can easily obtain the following tiling interpretation for the over-Mahonian numbers.

Corollary 3.6. The over-Mahonian number $\overline{\binom{n}{k}}^{\uparrow}$ is equal to the cardinal of the set $\mathcal{T}_{n,k}^{\uparrow}$. That is

$$\overline{\begin{bmatrix}n\\k\end{bmatrix}}^{\uparrow} = \mid \mathcal{T}_{n,k}^{\uparrow} \mid .$$

Example 3.7. For n = 3 and k = 2, we obtain $\begin{bmatrix} 3 \\ 2 \end{bmatrix}^{\uparrow}$ tilings with 2 blue squares and 2 red squares and black rectangle, as shown in Figure 5.




FIGURE 5. The tiling interpretation associated to $\begin{bmatrix} 3\\2 \end{bmatrix}^{\top}$.

4. Combinatorial proof of the log-Concavity and unimodality of the over-Mahonian numbers

Log-concave and unimodal sequences arise often in combinatorics, geometry and algebra. The reader is referred to [6, 14] for surveys of the diverse techniques employed to sequences and polynomials for which are log-concave or unimodal.

A sequence of nonnegative numbers $\{x_k\}_k$ is log-concave if $x_{i-1}x_{i+1} \leq x_i^2$ for all i > 0, which is equivalent to (for relevant results one can see [5, 14]):

$$x_{i-1}x_{j+1} \le x_i x_j \quad for \quad j \ge i \ge 1.$$

A finite sequence of real numbers a_0, \ldots, a_m is said to be unimodal if there exist an index $0 \le m^* \le m$, called the mode of the sequence, such that a_k increases up to $k = m^*$ and decreases from then on, that is, $a_0 \le a_1 \le \cdots \le a_{m^*}$ and $a_{m^*} \ge a_{m^*+1} \ge \cdots \ge a_m$. It is easy to see that if a sequence is log-concave then it is unimodal [5].

A polynomial is called log-concave (resp. unimodal) if its coefficients form a log-concave (resp. an unimodal) sequence. As examples, it is well-known that Gaussian polynomials and q-multinomial coefficients [15, Theorem 3.11] are unimodal. And it is a classic result (see for instance [4] for a proof) that the product of log-concave (resp. unimodal) polynomials is log-concave (resp. unimodal). For example $\sum_{k=0}^{\binom{n}{2}} i(n,k)x^k$ is log-concave (resp. unimodal) polynomial, that is, the sequence $i(n,0),\ldots,i(n,\binom{n}{2})$ is log-concave (resp. unimodal).

The first non-generating function proof of the log-concavity of the Mahonian numbers i(n, k) in k is proposed by Bóna [3]. He used an injection property and the induction

hypothesis over n, but this injection is non-constructive. For this, Ghemit and Ahmia proposed in their paper [9] a constructive proof to replace the non-constructive injection of Bóna.

The generating function proof of the log-concavity (resp. the unimodality) of the over-Mahonian numbers $\left\{\overline{\binom{n}{k}}^{\uparrow}\right\}_{k}$ in k is easy to establish from Theorem 2.7, because $\sum_{k=0}^{\binom{n}{2}} \overline{\binom{n}{k}}^{\uparrow} x^{k}$ is the product of log-concave (resp. unimodal) polynomials. But in this section, motivated by work of Ghemit and Ahmia [9], we show that the over-Mahonian numbers $\left\{\overline{\binom{n}{k}}^{\uparrow}\right\}_{k}$ form a log-concave sequence in k and therefore unimodal by constructing the appropriate injection.

First of all, we propose the following key definition.

Definition 4.1. Let σ be a permutation of length n. For $0 \leq i \leq n-1$, let m_i^{σ} be the number of appearances of the entry i+1 as the first element of the overlined inversions of σ . The total number of overlined inversions of σ is exactly $\sum_{i=0}^{n-1} m_i^{\sigma}$.

Theorem 4.2. The sequence of over-Mahonian numbers $\left\{\overline{\begin{bmatrix} n\\k \end{bmatrix}}^{\uparrow}\right\}_k$ is log-concave in k, that is :

$$\left(\overline{\begin{bmatrix}n\\k\end{bmatrix}}^{\uparrow}\right)^2 - \overline{\begin{bmatrix}n\\k-1\end{bmatrix}}^{\uparrow}\overline{\begin{bmatrix}n\\k+1\end{bmatrix}}^{\uparrow} \ge 0,$$

for all $0 < k \leq \binom{n}{2}$.

Proof. Let $\overline{P}(n,k)$ denote the set of permutations of length n with k overlined inversions, and $\overline{\left[\frac{n}{k}\right]}^{\uparrow} = |\overline{P}(n,k)|$ the number of permutations of length n with k overlined inversions.

To prove the log concavity of the sequence of over-Mahonian numbers $\{\overline{\begin{bmatrix} n\\k \end{bmatrix}}^{\uparrow}\}_k$, it is equivalent to find an injection $f_{n,k,k}$ from $\overline{P}(n,k+1) \times \overline{P}(n,k-1)$ to $\overline{P}(n,k) \times \overline{P}(n,k)$, such that, $f_{n,k,k}(\sigma,\tau) = (\theta,\pi)$ with $(\sigma,\tau) \in \overline{P}(n,k+1) \times \overline{P}(n,k-1)$ and $(\theta,\pi) \in \overline{P}(n,k) \times \overline{P}(n,k)$.

Let $(\sigma, \tau) \in \overline{P}(n, k+1) \times \overline{P}(n, k-1)$. We will define I as the largest integer which satisfies the following three conditions:

(6)
$$\sum_{j\geq I} m_j^{\sigma} \geq \sum_{j\geq I+1} m_j^{\tau} + 1$$
, if the entry $(j+1) = (I+1)$ of the σ not overlined,

(7)
$$\sum_{j\geq I} m_j^{\sigma} > \sum_{j\geq I+1} m_j^{\tau} + 1, \quad \text{if the entry } (j+1) = (I+1) \quad \text{of the } \sigma \quad \text{is overlined}$$

and

(8)
$$\sum_{j \ge I} m_j^{\tau} + 1 - \sum_{j \ge I+1} m_j^{\sigma} \le I, \quad \text{if } \sum_{j \ge I} m_j^{\sigma} < \sum_{j \ge I} m_j^{\tau} + 1.$$

Now, we define the map $f_{n,k,k}$ as follows:

$$f_{n,k,k}(\sigma,\tau) = (\theta,\pi),$$

where :

- θ is obtained by the following modifications on τ :
 - For the entry (j+1), from (I+2) to n in ascending order: if the entry (j+1) from σ is overlined, then overlined the entry (j+1) from τ , else not overlined the entry (j+1) from τ .
 - For the entry (j + 1), from n to (I + 2) in decreasing order: the entry j + 1 moves m_j^{τ} positions to the right and overlined, if the entry j + 1 is overlined. Else, the entry j + 1 only moves m_j^{τ} positions to the right.
 - For the entry j + 1 = I + 1: the entry j + 1 moves $\left(\sum_{i=1}^{I} m_i^{\sigma} \sum_{i=1}^{I} m_i^{\tau} 1\right)$ positions to the left and overlined, if the entry j + 1 is overlined. Else, the entry j + 1 only moves $\left(\sum_{i=1}^{I} m_i^{\sigma} \sum_{i=1}^{I} m_i^{\tau} 1\right)$ positions to the left.
 - For the entry (j + 1), from (I + 2) to n in ascending order: the entry j + 1 moves m_j^{σ} positions to the left and overlined, if the entry j + 1 is overlined. Else, the entry j + 1 only moves m_j^{σ} positions to the left.
- π is obtained by the following modifications on σ :
 - For the entry (j+1), from (I+2) to n in ascending order: if the entry (j+1) from τ is overlined, then overlined the entry (j+1) from σ , else not overlined the entry (j+1) from σ .
 - For the entry (j + 1), from n to (I + 2) in decreasing order: the entry j + 1 moves m_j^{σ} positions to the right and overlined, if the entry j + 1 is overlined. Else, the entry j + 1 only moves m_j^{σ} positions to the right.
 - For the entry j + 1 = I + 1: the entry j + 1 moves $\left(\sum_{i=1}^{I} m_i^{\sigma} \sum_{i=1}^{I} m_i^{\tau} 1\right)$ positions to the right and overlined, if the entry j + 1 is overlined. Else, the entry j + 1 only moves $\left(\sum_{i=1}^{I} m_i^{\sigma} \sum_{i=1}^{I} m_i^{\tau} 1\right)$ positions to the right.
 - For the entry (j + 1), from (I + 2) to n in ascending order: the entry j + 1 moves m_j^{τ} positions to the left and overlined, if the entry j + 1 is overlined. Else, the entry j + 1 only moves m_j^{τ} positions to the left.

To prove that $f_{n,k,k}$ is well defined we only need to check that:

(9)
$$0 \le \sum_{j=1}^{I} m_{j}^{\sigma} - \sum_{j=1}^{I} m_{j}^{\tau} - 1 \le m_{I}^{\sigma}$$

and

(10)
$$m_I^{\tau} + \sum_{j=1}^I m_j^{\sigma} - \sum_{j=1}^I m_j^{\tau} - 1 \le I.$$

Now, we simplify the inequality (9) as follows:

$$\sum_{j=1}^{I} m_{j}^{\sigma} - \sum_{j=1}^{I} m_{j}^{\tau} - 1 = \sum_{j=1}^{n-1} m_{j}^{\sigma} - \sum_{j=I+1}^{n-1} m_{j}^{\sigma} - \sum_{j=1}^{n-1} m_{j}^{\tau} + \sum_{j=I+1}^{n-1} m_{j}^{\tau} - 1$$
$$= \left(k + 1 - \sum_{j=I+1}^{n-1} m_{j}^{\sigma}\right) - \left(k - 1 - \sum_{j=I+1}^{n-1} m_{j}^{\tau}\right) - 1$$
$$= 1 - \sum_{j=I+1}^{n-1} m_{j}^{\sigma} + \sum_{j=I+1}^{n-1} m_{j}^{\tau}.$$

Then inequality (10) becomes:

$$m_{I}^{\tau} + \sum_{j=1}^{I} m_{j}^{\sigma} - \sum_{j=1}^{I} m_{j}^{\tau} - 1 = m_{I}^{\tau} + 1 - \sum_{j=I+1}^{n-1} m_{j}^{\sigma} + \sum_{j=I+1}^{n-1} m_{j}^{\tau}$$
$$= 1 - \sum_{j=I+1}^{n-1} m_{j}^{\sigma} + \sum_{j=I}^{n-1} m_{j}^{\tau}.$$

Now, we show inequality (9), i.e.

$$0 \le \sum_{j=1}^{I} m_j^{\sigma} - \sum_{j=1}^{I} m_j^{\tau} - 1 \le m_I^{\sigma}.$$

Here, we have two cases : **First case :** If $\sum_{j\geq I} m_j^{\sigma} \geq \sum_{j\geq I+1} m_j^{\tau} + 1$, we have:

$$\begin{split} 1 - \sum_{j \ge I} m_j^\sigma + \sum_{j \ge I+1} m_j^\tau \le 0 \Leftrightarrow 1 - \sum_{j \ge I+1} m_j^\sigma - m_I^\sigma + \sum_{j \ge I+1} m_j^\tau \le 0 \\ \Leftrightarrow 1 - \sum_{j \ge I+1} m_j^\sigma + \sum_{j \ge I+1} m_j^\tau \le m_I^\sigma \\ \Leftrightarrow \sum_{j=1}^I m_j^\sigma - \sum_{j=1}^I m_j^\tau - 1 \le m_I^\sigma. \end{split}$$

Second case : If $\sum_{j \ge I} m_j^{\sigma} > \sum_{j \ge I+1} m_j^{\tau} + 1$, we have :

$$\begin{split} 1 - \sum_{j \ge I} m_j^\sigma + \sum_{j \ge I+1} m_j^\tau < 0 \Leftrightarrow 1 - \sum_{j \ge I+1} m_j^\sigma - m_I^\sigma + \sum_{j \ge I+1} m_j^\tau < 0 \\ \Leftrightarrow 1 - \sum_{j \ge I+1} m_j^\sigma + \sum_{j \ge I+1} m_j^\tau < m_I^\sigma \\ \Leftrightarrow \sum_{j=1}^I m_j^\sigma - \sum_{j=1}^I m_j^\tau - 1 < m_I^\sigma. \end{split}$$

It remains to show that :

$$\sum_{j=1}^{I} m_j^{\sigma} - \sum_{j=1}^{I} m_j^{\tau} - 1 \ge 0.$$

For this, we have two cases: **First case :** If $\sum_{j=I+1}^{n-1} m_j^{\sigma} < \sum_{j=I+2}^{n-1} m_j^{\tau} + 1$, we have :

$$\begin{split} -\sum_{j=I+1}^{n-1} m_j^{\sigma} &> -\sum_{j=I+2}^{n-1} m_j^{\tau} - 1 \Leftrightarrow 1 - \sum_{j=I+1}^{n-1} m_j^{\sigma} > -\sum_{j=I+2}^{n-1} m_j^{\tau} \\ &\Leftrightarrow 1 - \sum_{j=I+1}^{n-1} m_j^{\sigma} + \sum_{j=I+2}^{n-1} m_j^{\tau} > 0 \\ &\Leftrightarrow 1 - \sum_{j=I+1}^{n-1} m_j^{\sigma} + \sum_{j=I+1}^{n-1} m_j^{\tau} > m_{I+1}^{\tau} \ge 0 \\ &\Leftrightarrow 1 - \sum_{j=I+1}^{n-1} m_j^{\sigma} + \sum_{j=I+1}^{n-1} m_j^{\tau} > 0 \\ &\Leftrightarrow \sum_{j=1}^{I} m_j^{\sigma} - \sum_{j=1}^{I} m_j^{\tau} - 1 > 0. \end{split}$$

Second case : If $\sum_{j=I+1}^{n-1} m_j^{\sigma} \ge \sum_{j=I+2}^{n-1} m_j^{\tau} + 1$ and $\sum_{j=I+1}^{n-1} m_j^{\sigma} < \sum_{j=I+1}^{n-1} m_j^{\tau} + 1$, we have

$$1 - \sum_{j=I+1}^{n-1} m_j^{\sigma} + \sum_{j=I+1}^{n-1} m_j^{\tau} > 0 \Leftrightarrow \sum_{j=1}^{I} m_j^{\sigma} - \sum_{j=1}^{I} m_j^{\tau} - 1 > 0.$$

We show the inequality (10): $m_I^{\tau} + \sum_{j=1}^I m_j^{\sigma} - \sum_{j=1}^I m_j^{\tau} - 1 \le I$. Here, we have two cases: **First case :** If $\sum_{j \ge I} m_j^{\sigma} < \sum_{j \ge I} m_j^{\tau} + 1$, using the condition (8), we obtain:

$$\sum_{j \ge I} m_j^\tau + 1 - \sum_{j \ge I+1} m_j^\sigma \le I \Leftrightarrow m_I^\tau + \sum_{j=1}^I m_j^\sigma - \sum_{j=1}^I m_j^\tau - 1 \le I.$$

Second case : If $\sum_{j\geq I} m_j^{\sigma} \geq \sum_{j\geq I} m_j^{\tau} + 1$, we have :

$$\begin{split} 1 - \sum_{j \ge I} m_j^{\sigma} + \sum_{j \ge I} m_j^{\tau} &\le 0 \Leftrightarrow 1 - \sum_{j \ge I+1} m_j^{\sigma} + \sum_{j \ge I} m_j^{\tau} \le m_I^{\sigma} \le I \\ &\Leftrightarrow 1 - \sum_{j \ge I+1} m_j^{\sigma} + \sum_{j \ge I} m_j^{\tau} \le I \\ &\Leftrightarrow m_I^{\tau} + \sum_{j=1}^I m_j^{\sigma} - \sum_{j=1}^I m_j^{\tau} - 1 \le I. \end{split}$$

Now, we must check that $(\theta, \tau) \in \overline{P}(n,k) \times \overline{P}(n,k)$ which means that the number of inversions in θ is k and the number of inversions in π is k.

The number of inversions in θ is calculated as follows

$$\sum_{j=1}^{I} m_j^{\tau} + \left(\sum_{j=1}^{I} m_j^{\sigma} - \sum_{j=1}^{I} m_j^{\tau} - 1\right) + \sum_{j=I+1}^{n-1} m_j^{\sigma} = \sum_{j=1}^{I} m_j^{\sigma} + \sum_{j=I+1}^{n-1} m_j^{\sigma} - 1$$
$$= \sum_{j=1}^{n-1} m_j^{\sigma} - 1$$
$$= (k+1) - 1 = k.$$

The number of inversions of π is calculated as follows

$$\sum_{j=1}^{I} m_j^{\sigma} - \left(\sum_{j=1}^{I} m_j^{\sigma} - \sum_{j=1}^{I} m_j^{\tau} - 1\right) + \sum_{j=I+1}^{n-1} m_j^{\tau} = \sum_{j=1}^{n-1} m_j^{\tau} + 1$$
$$= (k-1) + 1 = k.$$

Then $(\theta, \tau) \in \overline{P}(n,k) \times \overline{P}(n,k)$. Furthermore, $f_{n,k,k}^{-1} = f_{n,k+1,k-1}$, which gives us the injectivity of $f_{n,k,k}$ and completes the proof.

Here, we give an illustrative example.

Example 4.3. Let $\sigma = \overline{3}\overline{2}\overline{4}\overline{5}1$ a permutation with 5 overlined inversions and let $\tau = 12\overline{5}\overline{4}3$ a permutation with 3 overlined inversions.

$m_1^{\sigma} = 1$	$m_1^{\tau} = 0$
$m_2^{\sigma} = 2$	$m_2^{\tau} = 0$
$m_3^\sigma = 1$	$m_3^{\tau} = 1$
$m_4^\sigma = 1$	$m_4^{\tau} = 2$

First, we must find the index I, we have :

For I = 4, $I + 1 = \overline{5}$, so $m_4^{\sigma} = 1 = 0 + 1$ then I = 4 does not satisfy the second condition (7).

For I = 3, $I + 1 = \overline{4}$, so $m_3^{\sigma} + m_4^{\sigma} = 2 < m_4^{\tau} + 1 = 3$ then I = 3 does not satisfy the second condition (7).

For I = 2, $I + 1 = \overline{2}$, so $m_2^{\sigma} + m_3^{\sigma} + m_4^{\sigma} = 4 = m_3^{\tau} + m_4^{\tau} + 1 = 4$ then I = 2 does not satisfy the second condition (7).

For I = 1, $I + 1 = \overline{3}$, so $m_1^{\sigma} + m_2^{\sigma} + m_3^{\sigma} + m_4^{\sigma} = 5 > m_2^{\tau} + m_3^{\tau} + m_4^{\tau} + 1 = 4$ then I = 1 satisfies the second condition (7). Thus, I = 1 and our application gives :

$$\overline{32451} \rightarrow 3\overline{2451} \rightarrow 3\overline{2415} \rightarrow 3\overline{2145} \rightarrow \overline{21345} \rightarrow \overline{21435} \rightarrow \overline{21543}$$
$$12\overline{543} \rightarrow 12\overline{543} \rightarrow 12\overline{435} \rightarrow 12\overline{435} \rightarrow 12\overline{345} \rightarrow \overline{31245} \rightarrow \overline{31425} \rightarrow \overline{31452}$$

Hence, $f_{5,4,4}(\overline{3}\,\overline{2}\,\overline{4}\,\overline{5}1,12\overline{5}\,\overline{4}3) = (\overline{2}1\overline{5}\,\overline{4}3,\overline{3}1\overline{4}\,\overline{5}2)$, where $\theta = \overline{2}1\overline{5}\,\overline{4}3$ has 4 overlined inversions, and $\pi = \overline{3}1\overline{4}\,\overline{5}2$ has 4 overlined inversions.

Since the log-concave sequence is unimodal [5], we have immediately from the previous theorem the following result.

Corollary 4.4. The sequence of over-Mahonian numbers $\left\{ \overline{\begin{bmatrix} n \\ k \end{bmatrix}}^{\uparrow} \right\}_k$ is unimodal in k.

Remark 4.5. Since there is a bijection between the permutations with overlined inversions and the paths as shown in Subsection 3.1, then the previous theorem can be proved using Sagan's paths approach [13] by adding a new condition to his involution as follows. (See also the approaches proposed in [8, 2]).

Let $P_1, P_2 \in P_{n,k}^{\uparrow}$. And let $u \xrightarrow{P} v$ denote P has initial vertex u and final vertex v.

Definition 4.6. Given $u_1 \xrightarrow{P_1} v_1$ and $u_2 \xrightarrow{P_2} v_2$. Then define the involution $\varphi_I(P_1, P_2) = (P'_1, P'_2)$ where :

$$P_1^{'} = u_1 \xrightarrow{P_1} v_0 \xrightarrow{P_2} v_2 \text{ and } P_2^{'} = u_2 \xrightarrow{P_2} v_0 \xrightarrow{P_1} v_1$$

i.e., switches the portions of P_1 and P_2 after v_0 (see, Figure 6), where v_0 is the last vertex of $P_1 \cap P_2$.

Acknowledgment

The authors would like to thank the referees for many valuable remarks and suggestions to improve the original manuscript. This work was supported by DG-RSDT (Algeria), PRFU Project, No. C00L03UN180120220002.




FIGURE 6. The involution φ_I .

References

- G. E. Andrews, The theory of partitions, Cambridge Mathematical Library. Cambridge University Press, Cambridge, 1998. Reprint of the 1976 original.
- [2] A. Bazeniar, M. Ahmia, A. Bouchair, Log-concave sequences of bi^snomial coefficients with their analogs and symmetric function, *Indian J. Pure Appl. Math.* 53, 127-137 (2022).
- [3] M. Bóna, A Combinatorial Proof of the Log-concavity of a famous sequence counting permutations, Electron. J. Combin. 11(2), #N2 (2005).
- [4] M. Bóna, Combinatorics of Permutations. CRC Press, (2004).
- [5] F. Brenti, Unimodal, log-concave and Polya frequency sequences in combinatorics, Mem. Amer. Math. Soc. 413 (1989).
- [6] F. Brenti, Log-concave and unimodal sequence in algebra, combinatorics and geometry: an update, *Elec. Contemp. Math.* 178, 71–84 (1997).
- [7] S. Corteel, J. Lovejoy, Overpartitions, Trans. Amer. Math. Soc. 356(4), 1623–1635 (2004).
- [8] Y. Ghemit, M. Ahmia, Overpartition analogues of q-bi^snomial coefficients: basic properties and logconcavity, *The Ramanujan Journal.* 62, 431-455 (2023).
- Y. Ghemit, M. Ahmia, An analogue of Mahonian numbers and log-concavity, Annals of Combinatorics. 27, 895-916 (2023).
- [10] D. J. L. Indong, G. R. Peralta, Inversions of permutations in symmetric, alternating, and dihedral groups, J. Integer Seq. 11, (2008).
- [11] P. A. MacMahon, The indices of permutations and the derivation therefrom of functions of a single variable associated with the permutations of any assemblage of objects, Amer. J. Math. 35, 281-322 (1913).
- [12] G. Rodrigues, Note sur les inversions, ou dérangements produits dans les permutations, J. Math. Pures Appl. 4, 236-240 (1839).
- [13] B. E. Sagan, Log concave sequences of symmetric functions and analogs of the Jacobi-Trudi determinants, Trans. Amer. Math. Soc. 329(2), 795–811 (1992).
- [14] R. P. Stanley, Log-concave and unimodal sequences inalgebra, combinatorics and geometry, Ann. N.Y. Acad. Sci. 576, 500–534 (1989).
- [15] J. J. Sylvester, Proof of the hitherto undemonstrated fundamental theorem of invariants, *Philos. Mag.* 5, 178-188 (1878).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF FERHAT ABBAS SETIF 1, ALGERIA Email address: ali.kessouri@ensta.edu.dz or ali.kessouri@univ-setif.dz

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MOHAMED SEDDIK BENYAHIA, LMAM LABORA-TORY, BP 98 OULED AISSA JIJEL 18000, ALGERIA *Email address*: moussa.ahmia@univ-jijel.dz or ahmiamoussa@gmail.com

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF FERHAT ABBAS SETIF 1, ALGERIA *Email address:* salim.mesbahi@univ-setif.dz

18