Note on higher-group structure in 6d self-dual gauge theory

Tatsuki Nakajima¹, Kikyo Nakamura^{*1}, and Tadakatsu Sakai^{2,1}

¹Department of Physics, Nagoya University

²Kobayashi-Maskawa Institute for the Origin of Particles and the Universe,

Nagoya University

Abstract

We analyze higher-group structure of a 6d model coupled with a self-dual 2-form gauge field. This model is defined from 6d axion-electrodynamics with a 1-form Chern-Weil(CW) symmetry gauged dynamically. The gauging leads to a Green-Schwarz-West-Sagnotti(GSWS) term, which gives rise to an anomaly through a GSWS transformation acting on the 2form gauge field. We cancel this anomaly by gauging a 3-form CW symmetry in 6d axionelectrodynamics. We find out the global symmetries in the resultant model and derive the gauge invariant action in the presence of the background gauge fields. It is argued that a discrete 1-form symmetry is anomalous because turning on the associated background gauge field causes quantum inconsistency due to an operator-valued ambiguity. Higher-group structure in this model that is manifested as a Green-Schwarz-like transformation for CW background gauge fields is discussed.

Contents

1 Introduction

2 Model

 $\mathbf{2}$

3

^{*}present address: Meitetsucom Co., Ltd.

3	EoM-based current and global symmetry	8
	3.1 Dualizing the 4-form gauge field	14
4	Gauging the EoM-based global symmetry	15
	4.1 Operator-valued ambiguity	18
5	Discussions	18
Α	Dimensional reduction to 5d	20

1 Introduction

The paper [1] discusses higher-group structure in 6d axion-electrodynamics by extending interesting papers [2,3], which explore 3-group structure encoded in 4d axion-electrodynamics. It is seen that the 6d model possesses much richer structure of higher-group compared with the 4d case, because the 6d model allows Chern-Weil(CW) currents of higher rank to exist. The higher-group structure is manifested as a Green-Schwarz(GS)-like transformation law [4] for the background gauge fields associated with the CW symmetry, although the mathematical framework for it is difficult to formulate. For recent studies of higher-group structure, see also [5–35].

In this paper, we discuss higher-group structure in a 6d model with a self-dual gauge field. This model is obtained by promoting a CW 2-form background gauge field in the 6d axionelectrodynamics to a dynamical gauge field. One of the motivations for this paper is to discuss how the gauging of the CW symmetry affects the higher-group structure that is worked out in [1] in the hope that a new higher-group structure emerges. The gauging of the CW symmetry leads to a coupling between the 2-form self-dual gauge field b and the CW 4-form current $\frac{1}{4\pi^2} da \wedge da$ with abeing a photon field. This is equal to a Green-Schwarz-West-Sagnotti(GSWP) term [36,37], which enables us to cancel 6d reducible anomalies by using a GS-like transformation law obeyed by b. A typical example is given by a U(1) gauge theory coupled with chiral fermions, where the box $U(1)^4$ anomaly can be canceled by using the GSWP mechanism. In this paper, we focus on an anomaly free model that is constructed by gauging another CW symmetry in the 6d axion-electrodynamics with no chiral fermions coupled. As discussed in [1], the 6d axion-electrodynamics admits the CW current $da/2\pi$. We gauge this symmetry by introducing a dynamical 4-form gauge field coupled with the CW current. The anomaly free model is obtained by requiring that the 4-form gauge field make an appropriate GS-like transformation under the U(1) gauge transformation as well.

We next discuss the global symmetries realized in the resultant 6d model. It is seen that a discrete 1-form symmetry found in [1] that acts on the photon is broken explicitly, but gets recovered by modifying it so that it acts on the self-dual gauge field and 4-form gauge field together with the photon. We argue that the 1-form symmetry is anomalous, however, because turning on the corresponding background field results in an operator-valued ambiguity, which states that the theory in the presence of the background gauge field has a quantum Chern-Simons(CS) action that is ambiguous in lifting it to 7d. It is found that higher-group structure in this model is less interesting than that in 6d axion-electrodynamics generically, because the GS-like transformation is induced only for a 1-form CW background gauge field for generic cases.

The organization of this paper is as follows. In section 2, we construct the 6d model by gauging dynamically CW symmetries in 6d axion-electrodynamics. Section 3 works out the global symmetries of the model by studying the equations of motion in detail. In section 4, we turn on the background gauge fields for the global symmetry to show that an operator-values ambiguity is caused by that for the discrete 1-form symmetry. Higher-group structure in this model is discussed in section 5. The appendix A discusses dimensional reduction of the 6d model to 5d for the purpose of evading the subtlety of the self-dual 2-form gauge field in obtaining the global higher-form symmetries in 6d.

2 Model

The action of the 6d axion electrodynamics is given by

$$S_{\rm aE}[\phi,a] = -\int_{\mathcal{M}_6} \left(\frac{1}{2}|d\phi|^2 + \frac{1}{2}|da|^2 - \frac{N}{48\pi^3}\phi(da)^3\right).$$
(2.1)

Here, $|\chi_p|^2 = (1/p!) \chi_p \wedge \star \chi_p$ for the *p*-form χ_p with \star being the Hodge star. We take \mathcal{M}_6 to be a spin manifold so that $N \in \mathbb{Z}$. We denote $(da)^3 = da \wedge da \wedge da$ for simplicity. This action admits discrete \mathbb{Z}_N symmetries that follow from the equation of motion(EoM) for ϕ and *a*. *a* is normalized as $\int da \in 2\pi\mathbb{Z}$. Turning on the corresponding background field gives rise to an operator-valued ambiguity. This is canceled by gauging the CW symmetry and requiring that the associated background gauge field make an appropriate GS transformation. The CW current reads¹

$$j_{\rm CW}^{[1]} = \frac{1}{4\pi^2} (da)^2 , \qquad j_{\rm CW}^{[2]} = \frac{1}{4\pi^2} \, d\phi \wedge da , \qquad j_{\rm CW}^{[3]} = \frac{1}{2\pi} \, da , \qquad j_{\rm CW}^{[4]} = \frac{1}{2\pi} \, d\phi .$$

Let B_2^{CW} , C_3^{CW} , D_4^{CW} and E_5^{CW} be the background gauge fields that couple minimally with the CW currents $j_{\text{CW}}^{[1]}$, $j_{\text{CW}}^{[2]}$, $j_{\text{CW}}^{[3]}$ and $j_{\text{CW}}^{[4]}$ respectively. It is shown that the gauge invariant field strength for the CW gauge field should be defined by

$$G_{3} = dB_{2}^{CW} - \frac{N}{2\pi} A_{1}B_{2} ,$$

$$H_{4} = dC_{3}^{CW} + \frac{N}{4\pi} (B_{2})^{2} ,$$

$$I_{5} = dD_{4}^{CW} - \frac{1}{2\pi} A_{1} dC_{3}^{CW} + \frac{1}{2\pi} B_{2} dB_{2}^{CW} - \frac{N}{4\pi^{2}} A_{1} (B_{2})^{2} ,$$

$$J_{6} = dE_{5}^{CW} + \frac{N}{2\pi} B_{2} dC_{3}^{CW} + \frac{N}{12\pi^{2}} (B_{2})^{3} .$$
(2.2)

Here, (A_1, A_0) with $NA_1 = dA_0$ and (B_2, B_1) with $NB_2 = dB_1$ are the background \mathbb{Z}_N gauge fields for the EoM-based 0-form and 1-form \mathbb{Z}_N symmetries, respectively. Hereafter, we denote a symmetry group G of rank p as $G^{[p]}$. For instance, the U(1) gauge symmetry acting on the photon field is a 0-form symmetry and is referred to as $U(1)^{[0]}$ with the gauge transformation given by $\delta^{[0]}a = d\lambda_0$.

Now we promote C_2^{CW} to a dynamical, self-dual 2-form gauge field b. The action becomes

$$S_{aEb}[\phi, a, b] = \int_{\mathcal{M}_6} \left(-\frac{1}{2} |d\phi|^2 - \frac{1}{2} |da|^2 - \frac{1}{8\pi} \left| db - \frac{M}{2\pi} \omega_3(a) \right|^2 + \frac{N}{48\pi^3} \phi(da)^3 + \frac{M}{8\pi^2} b \wedge (da)^2 \right) ,$$
(2.3)

with $\omega_3(a) = a \wedge da$ being the CS 3-form. *b* is normalized as $\int db \in 2\pi\mathbb{Z}$ and $M \in \mathbb{Z}$. The self-duality condition for *b* is given by

$$db - \frac{M}{2\pi}\omega_3(a) = \star \left(db - \frac{M}{2\pi}\omega_3(a)\right) . \tag{2.4}$$

As a consistency check, we note that the EoM reads

$$\frac{1}{4\pi}d \star \left(db - \frac{M}{2\pi}\omega_3(a)\right) + \frac{1}{8\pi^2}(da)^2 = 0 ,$$

¹For the normalization of the CW currents, we follow that of [18], where the integral of the CW current over a closed cycle in \mathcal{M}_6 is Z-valued.

on which imposing (2.4) leads to the Bianchi identity for the field strength $db - (M/2\pi) \omega_3(a)$. This field strength is left unchanged under the $U(1)^{[0]}$ gauge transformation if b makes a GS-like transformation as

$$\delta^{[0]}b = \frac{M}{2\pi}\lambda_0 da \; .$$

The GSWS term

$$\frac{M}{8\pi^2}b\wedge (da)^2$$

is utilized for canceling reducible anomalies in 6d [36, 37].

As an example for the GSWS mechanism, we consider n right-handed fermions of $U(1)^{[0]}$ charge p. They lead to the one-loop box anomaly

$$2\pi I_6 = -n \, p^4 \frac{1}{4! (2\pi)^3} \lambda_0 \, (da)^3 \,, \tag{2.5}$$

which descends from the anomaly polynomial

$$I_8 = -n p^4 \frac{1}{4! (2\pi)^4} (da)^4$$

By imposing

$$2\pi I_6 + \delta^{[0]} S = 0 ,$$

with

$$\delta^{[0]}S = \frac{M}{8\pi^2} \delta^{[0]}b \wedge (da)^2 = \frac{M^2}{2(2\pi)^3} \lambda_0 (da)^3 ,$$

we find

$$M = \left(\frac{n \, p^4}{12}\right)^{1/2} \,. \tag{2.6}$$

This is consistent if the RHS is integer.

The axion electrodynamics coupled with the right-handed fermions is obtained by the Peccei-Quinn(PQ) mechanics [38]. To see this, consider the model

$$U(1)_{L} \times U(1)_{R}$$

$$n_{L} \times \psi_{L} \qquad (p,0)$$

$$(n_{L}+n) \times \psi_{R} \qquad (0,p)$$

$$\Phi \qquad (p,-p)$$

$$b \qquad (0,0)$$

$$(2.7)$$

 $\psi_{L,R}$ are left- and right-handed chiral fermions, respectively. The vector-like subgroup of $U(1)_L \times U(1)_R$ is dynamically gauged with the dynamical gauge field given by a. We note that the one-loop $U(1)^4$ anomaly from ψ_L and ψ_R is given by (2.5). This is canceled by the GSWS mechanics by setting M equal to (2.6). Φ is a complex scalar field that is neutral under $U(1)^{[0]}$. They are coupled with the Yukawa coupling as

$$\mathcal{L}_{\mathrm{Y}} = \Phi \, \psi_L^{i\dagger} \psi_{Ri} + \mathrm{c.}c. \; .$$

with $i = 1, 2, \dots, n_L$. The rest of ψ_{Ri} with $i = n_L + 1, \dots, n$ has no Yukawa coupling with ψ_L .

The PQ mechanism occurs by turning on the vev of Φ :

$$\langle \Phi \rangle = v$$

This breaks $U(1)_A$ while the $U(1)^{[0]}$ gauge symmetry is left unbroken. The fluctuations about this vacuum are given by

$$\Phi = (v + \rho)e^{i\phi}$$

 ρ is massive and thus can be integrated out. The Yukawa term becomes

$$\mathcal{L}_{\mathrm{Y}} = v e^{i\phi} \psi_L^{\dagger} \psi_R + \cdots$$

The $U(1)_A$ transformation

$$\psi_L \to e^{ip\alpha} \psi_L , \quad \psi_R \to e^{-ip\alpha} \psi_R ,$$

with

$$\phi - 2p\alpha = 0$$

removes ϕ from the Yukawa coupling. Then, n_L pairs of the chiral fermions ψ_L and ψ_R gain a non-vanishing, ϕ -independent mass term and thus can be integrated out. The $U(1)_A$ rotation gives rise to the axion coupling via $U(1)_A$ anomaly

$$2 \times n_L p^4 \alpha \frac{(da)^3}{3!(2\pi)^3} = n_L p^3 \phi \frac{(da)^3}{3!(2\pi)^3} .$$

This allows us to identify

$$N = n_L p^3 . (2.8)$$

Our main focus in this paper is on a model defined by gauging the 3-form CW symmetry. This is done by promoting the CW background gauge field D_4^{CW} to a dynamical 4-form gauge field u, which couples minimally with the corresponding CW current as

$$Ku \wedge j_{\rm CW}^{[3]} = \frac{K}{2\pi} u \wedge da \; .$$

Here, $K \in \mathbb{Z}$ because of the normalization $\int du \in 2\pi\mathbb{Z}$. Then, this model is anomaly free by requiring that u make a GS-like transformation under $U(1)^{[0]}$ as

$$\delta^{[0]}u = -\frac{M^2}{8\pi^2 K} \lambda_0 \wedge (da)^2 \; .$$

This implies that the gauge invariant field strength of u is given by

$$i_5 = du + \frac{M^2}{8\pi^2 K} a \wedge (da)^2 = du + \frac{M^2}{8\pi^2 K} \omega_5(a) .$$
(2.9)

To summarize, the gauge symmetry and the gauge transformation of the present model reads

• $U(1)^{[0]}$

$$\delta^{[0]}a = d\lambda_0 , \quad \delta^{[0]}b = \frac{M}{2\pi}\lambda_0 da , \quad \delta^{[0]}u = -\frac{M^2}{8\pi^2 K}\lambda_0 (da)^2 . \tag{2.10}$$
$$U(1)^{[1]}$$

 $\delta^{[1]}b = d\lambda_1 \ .$

• $U(1)^{[3]}$

$$\delta^{[3]}u = d\lambda_3$$
 .

The consistency of (2.10) with the quantization condition $\int du \in 2\pi\mathbb{Z}$ requires

$$\frac{M^2}{8\pi^2 K} (2\pi)^3 \in 2\pi\mathbb{Z} \ .$$

This is equivalent to stating that there exists an integer r_1 such that

$$M^2 = 2Kr_1 \; .$$

This shows that M is even. Let $m_1 = \text{gcd}(K, r_1)$. Then,

$$K = m_1 \widetilde{K} , \ r_1 = m_1 \widetilde{r}_1 , \qquad (2.11)$$

with $\widetilde{K}, \widetilde{r}_1 \in \mathbb{Z}$ being relatively prime. These satisfy

$$M^2 = 2m_1^2 \widetilde{K} \widetilde{r}_1 \; .$$

This implies that there exists an integer L_1 such that

$$2\widetilde{K}\,\widetilde{r}_1 = L_1^2\,,\qquad(2.12)$$

which requires that L_1 be even. It follows that M is written in terms of m_1 and L_1 as

$$M = m_1 L_1 . (2.13)$$

3 EoM-based current and global symmetry

Start with the action

$$S = -\int_{\mathcal{M}_6} \left(\frac{1}{2} |d\phi|^2 + \frac{1}{2} |da|^2 + \frac{1}{8\pi} |db - \frac{M}{2\pi} \omega_3(a)|^2 + \frac{1}{2} |du + \frac{M^2}{8\pi^2 K} \omega_5(a)|^2 \right) + S_{\rm CS} .$$
(3.1)

Here

$$S_{\rm CS} = \int_{\mathcal{M}_6} \left(\frac{N}{48\pi^3} \phi(da)^3 + \frac{M}{8\pi^2} (da)^2 \wedge b + \frac{K}{2\pi} da \wedge u \right) \\ = \int_{\Omega_{\mathcal{M}_6}} \left(\frac{N}{48\pi^3} d\phi \wedge (da)^3 + \frac{M}{8\pi^2} (da)^2 \wedge db + \frac{K}{2\pi} da \wedge du \right) ,$$

where $\Omega_{\mathcal{M}_6}$ is a seven-manifold with $\partial \Omega_{\mathcal{M}_6} = \mathcal{M}_6$. The EoM for ϕ reads

$$d^{\star}d\phi + \frac{N}{48\pi^3} \, (da)^3 = 0 \; .$$

This defines the conserved current

$$j_{\rm EoM}^{[0]} = -\star d\phi - \frac{N}{48\pi^3}\,\omega_5 \;.$$

As discussed in [1], this leads to a 0-form global \mathbb{Z}_N symmetry, which acts as

$$\delta_{\Lambda_0}\phi = \Lambda_0 \; , \;$$

with $\Lambda_0 = 2\pi/N$.

The EoM for u reads

$$d \star \left(du + \frac{M^2}{8\pi^2 K} \omega_5 \right) + \frac{K}{2\pi} da = 0 . \qquad (3.2)$$

This defines the conserved current

$$j_{\rm EoM}^{[4]} = - \star \left(du + \frac{M^2}{8\pi^2 K} \omega_5 \right) - \frac{K}{2\pi} a \, .$$

Noting that $i_5 = du + \frac{M^2}{8\pi^2 K} \omega_5$ is gauge invariant, the symmetry generator associated with $j_{\text{EoM}}^{[4]}$ is gauge invariant if it generates a 4-form \mathbb{Z}_K global symmetry. This acts as

$$\delta_{\Lambda_4} u = \Lambda_4 \; ,$$

with

$$\int \Lambda_4 \in \frac{2\pi}{K} \mathbb{Z} \; .$$

The EoM for b reads

$$\frac{1}{4\pi}d \star \left(db - \frac{M}{2\pi}\omega_3\right) + \frac{M}{8\pi^2}(da)^2 = 0.$$
(3.3)

The EoM-based conserved current is given by

$$j_{\rm EoM}^{[2]} = -\frac{1}{4\pi} \star \left(db - \frac{M}{2\pi} \omega_3 \right) - \frac{M}{8\pi^2} \omega_3 = -\frac{1}{4\pi} db \; .$$

Identification of the global symmetry generated by this current is tricky because of the self-duality condition of b. One way to do that is to dimensionally reduce the 6d action (3.1) to 5d, where breduces to a U(1) gauge field. For details, see the appendix A. As discussed there, the resultant action has a 1-form global \mathbb{Z}_M symmetry, which is the manifestation of the 2-form global \mathbb{Z}_M symmetry in 6d after dimensional reduction. This acts as

$$\delta_{\Lambda_2} b = \Lambda_2 \; ,$$

with

$$\int \Lambda_2 \in \frac{2\pi}{M} \mathbb{Z} \ .$$

The EoM for a reads

$$d^{\star}da - \frac{2M}{16\pi^2} \left[da \wedge \star \left(db - \frac{M}{2\pi} \omega_3 \right) + d \left\{ a \wedge \star \left(db - \frac{M}{2\pi} \omega_3 \right) \right\} \right]$$
$$+ \frac{M^2}{8\pi^2 K} \left[(da)^2 \wedge \star \left(du + \frac{M^2}{8\pi^2 K} \omega_5 \right) + 2 d \left\{ \omega_3 \wedge \star \left(du + \frac{M^2}{8\pi^2 K} \omega_5 \right) \right\} \right]$$
$$- \frac{N}{48\pi^3} 3 d\phi \wedge (da)^2 - \frac{M}{8\pi^2} 2 db \wedge da - \frac{K}{2\pi} du = 0 . \tag{3.4}$$

This can be written in a gauge invariant form as

$$d^{\star}da - \frac{M}{2\pi^{2}}da \wedge \left(db - \frac{M}{2\pi}\omega_{3}\right) + \frac{3M^{2}}{8\pi^{2}K}(da)^{2} \wedge \star \left(du + \frac{M^{2}}{8\pi^{2}K}\omega_{5}\right) - \frac{K}{2\pi}\left(du + \frac{M^{2}}{8\pi^{2}K}\omega_{5}\right) - \frac{N}{16\pi^{3}}d\phi \wedge (da)^{2} = 0.$$
(3.5)

It can be verified that the LHS of (3.5) is an exact form leading to an EoM-based conserved current. To see this, we note that (3.2) is integrated as

$$\star \left(du + \frac{M^2}{8\pi^2 K} \omega_5 \right) = -\frac{K}{2\pi} a + \frac{1}{2\pi} dv_0 \; .$$

Here, v_0 is a 0-form field, which is required to transform as

$$\delta^{[0]}v_0 = K\lambda_0 , \quad \delta^{[1]}v_0 = \delta^{[3]}v_0 = 0 .$$
(3.6)

It then follows that

$$du + \frac{M^2}{8\pi^2 K} \omega_5 = -\frac{K}{2\pi} \star \left(a - \frac{1}{K} \, dv_0\right) \,. \tag{3.7}$$

Inserting this into (3.5) gives

$$d \star da - \frac{M}{2\pi^2} da \wedge db + \frac{3M^2}{16\pi^3 K} (da)^2 \wedge dv_0 - \frac{K}{2\pi} du - \frac{N}{16\pi^3} d\phi \wedge (da)^2 = 0 .$$

This can be integrated as

$$j_4 \equiv \star da - \frac{M}{2\pi^2} da \wedge b + \frac{3M^2}{16\pi^3 K} (da)^2 v_0 - \frac{K}{2\pi} u - \frac{N}{16\pi^3} \phi (da)^2 = -\frac{1}{2\pi} dv_3 , \qquad (3.8)$$

with v_3 being a 3-form gauge field. j_4 defines an EoM-based conserved current. It is easy to show that v_3 transforms as

$$\delta^{[0]}v_3 = 0 , \quad \delta^{[1]}v_3 = \frac{M}{\pi} \lambda_1 \wedge da , \quad \delta^{[3]}v_3 = K\lambda_3 .$$
(3.9)

As clear from (3.6) and (3.9), v_0 and v_3 can be regarded as NG bosons associated with the spontaneous symmetry breaking for the gauge group $U(1)^{[0]}$ and $U(1)^{[3]}$, respectively. This is a manifestation of the Stückelberg mechanism, which gives rise to a mass for the gauge field a and u via the coupling $(K/2\pi) da \wedge u$. We normalize v_0 and v_3 as

$$\int dv_0 \in 2\pi\mathbb{Z} \ , \quad \int dv_3 \in 2\pi\mathbb{Z}$$

Then, the Stückelberg mechanism is equivalent to Higgsing the gauge groups $U(1)^{[0]}$ and $U(1)^{[3]}$ by the vev of operators of K units of charges, so that the gauge groups are broken to $\mathbb{Z}_{K}^{[0]}$ and $\mathbb{Z}_{K}^{[3]}$, respectively [39].

We can improve $j_{\text{EoM}}^{[1]}$ by adding a total derivative term to eliminate v_3 :

$$j_{\rm EoM}^{\prime[1]} = j_{\rm EoM}^{[1]} + d\Omega_3$$

= $\star da - \frac{M}{8\pi^2} da \wedge b - \frac{3M}{4\pi K} \left(db - \frac{M}{2\pi} \omega_3 \right) \wedge \star \left(du + \frac{M^2}{8\pi^2 K} \omega_5 \right) - \frac{K}{2\pi} u - \frac{N}{16\pi^3} \phi \left(da \right)^2 ,$
(3.10)

where

$$\Omega_3 = \frac{3M}{8\pi^2 K} \left(db - \frac{M}{2\pi} \,\omega_3 \right) v_0 + \frac{3M}{8\pi^2} \,a \wedge b \;.$$

We now show that the transformation laws generated by the EoM-based current are given by

$$\delta_{\Lambda_1} a = \Lambda_1 , \quad \delta_{\Lambda_1} b = -\frac{M}{2\pi} \Lambda_1 \wedge a , \quad \delta_{\Lambda_1} u = \Lambda_1 \wedge \left(\frac{M^2}{8\pi^2 K} \omega_3 - \frac{3M}{4\pi K} db\right) . \tag{3.11}$$

For this purpose, we gauge Λ_1 with $d\Lambda_1 \neq 0$. Using

$$\delta_{\Lambda_1} \left(db - \frac{M}{2\pi} \omega_3 \right) = -\frac{M}{\pi} d\Lambda_1 \wedge a ,$$

$$\delta_{\Lambda_1} \left(du + \frac{M^2}{8\pi^2 K} \omega_3 \right) = -\frac{3M}{4\pi K} d\Lambda_1 \wedge \left(db - \frac{M}{2\pi} \omega_3 \right) ,$$

it is found that

$$\delta_{\Lambda_1} \mathcal{L} = d\Lambda_1 \wedge \left[-\star da + \frac{M}{4\pi^2} a \wedge \star \left(db - \frac{M}{2\pi} \omega_3 \right) + \frac{3M}{4\pi K} \left(db - \frac{M}{2\pi} \omega_3 \right) \wedge \star \left(du + \frac{M^2}{8\pi^2 K} \omega_5 \right) \right. \\ \left. + \frac{N}{16\pi^3} \phi (da)^3 + \frac{M}{4\pi^2} da \wedge b + \frac{K}{2\pi} u \right] - \frac{3M}{8\pi^2} \Lambda_1 \wedge da \wedge db \; .$$

Using

$$\Lambda_1 \wedge da \wedge db = d\Lambda_1 \wedge (a \wedge db + t \cdot d(a \wedge b)) - d \left[\Lambda_1 \wedge (a \wedge db + t \cdot d(a \wedge b))\right] ,$$

with t being any real number, it is found that

$$\delta_{\Lambda_1} \mathcal{L} = d\Lambda_1 \wedge \left[-\star da + \frac{M}{4\pi^2} a \wedge \star \left(db - \frac{M}{2\pi} \omega_3 \right) + \frac{3M}{4\pi K} \left(db - \frac{M}{2\pi} \omega_3 \right) \wedge \star \left(du + \frac{M^2}{8\pi^2 K} \omega_5 \right) \right. \\ \left. + \frac{N}{16\pi^3} \phi(da)^3 + \frac{K}{2\pi} u + \frac{M}{4\pi^2} \left(1 - \frac{3t}{2} \right) da \wedge b + \frac{3M}{8\pi^2} \left(-1 + t \right) a \wedge db \right] \\ \left. + \frac{3M}{8\pi^2} d \left[\Lambda_1 \wedge \left(a \wedge db + t \cdot d(a \wedge b) \right) \right] \,.$$

Imposing the self-duality condition (2.4) reduces it to

$$\delta_{\Lambda_1} \mathcal{L} = d\Lambda_1 \wedge \left[-\star da + \frac{3M}{4\pi K} \left(db - \frac{M}{2\pi} \omega_3 \right) \wedge \star \left(du + \frac{M^2}{8\pi^2 K} \omega_5 \right) + \frac{N}{16\pi^3} \phi (da)^3 + \frac{K}{2\pi} u + \frac{M}{4\pi^2} \left(1 - \frac{3t}{2} \right) da \wedge b + \frac{M}{8\pi^2} \left(-1 + 3t \right) a \wedge db \right] + \frac{3M}{8\pi^2} d \left[\Lambda_1 \wedge (a \wedge db + t \cdot d(a \wedge b)) \right] .$$

Setting t = 1/3 and restoring Λ_1 to a global transformation parameter, we obtain the conservation law for $j'_{\text{EoM}}^{[1]}$.

Let

$$U^{[1]}(\alpha, \mathcal{M}_4) = e^{i\alpha \int_{\mathcal{M}_4} j_{\text{EoM}}^{\prime [1]}}$$

be the symmetry generator defined by $j_{\text{EoM}}^{\prime[1]}$. The transformation parameter α is written in terms of Λ_1 as

$$\alpha = \int \Lambda_1 \; ,$$

and constrained from the gauge invariance of the symmetry generator. The conserved charge $\int_{\mathcal{M}_4} j_{\text{EoM}}^{\prime[1]}$ is rewritten in a gauge invariant form by lifting it to a five-dimensional manifold. The gauge invariance of $U^{[1]}$ amounts to requiring that it be independent of how to lift it. We are led to impose the condition

$$\exp i\alpha \int_{\mathcal{M}_5} \left(-\frac{M}{8\pi^2} da \wedge db - \frac{K}{2\pi} du - \frac{N}{16\pi^3} d\phi \wedge (da)^2 \right) = 1 , \qquad (3.12)$$

for any closed manifold \mathcal{M}_5 . By using

$$\frac{1}{2\pi} \int_{\mathcal{M}_5} du = n_1 , \quad (n_1 \in \mathbb{Z})$$
$$\frac{1}{8\pi^2} \int_{\mathcal{M}_5} da \wedge db = \frac{n_2}{2} , \quad (n_2 \in \mathbb{Z})$$
$$\frac{1}{16\pi^3} \int_{\mathcal{M}_5} d\phi \wedge (da)^2 = \frac{n_3}{2} , \quad (n_3 \in \mathbb{Z})$$

we obtain

$$\left(n_1K + \frac{n_2}{2}M + \frac{n_3}{2}N\right)\alpha \in 2\pi\mathbb{Z} .$$

Using (2.11) and (2.13), we find

$$n_1K + \frac{n_2}{2}M + \frac{n_3}{2}N = m_1\left(n_1\tilde{K} + n_2\frac{L_1}{2}\right) + \frac{N}{2}n_3$$
.

Note that $L_1/2 \in \mathbb{Z}$. If there exists an nontrivial integer q such that

$$q = \gcd(m_1, \frac{N}{2}) , \qquad (3.13)$$

the symmetry generator is gauge invariant by setting

$$\alpha \in \frac{2\pi}{q}\mathbb{Z} .$$

This shows that the EoM-based current leads to the global 1-form \mathbb{Z}_q symmetry. It follows from (3.13) that m_1 and N are written as

$$m_1 = q\widetilde{m}_1 , \quad N = q\widetilde{N} , \qquad (3.14)$$

with $\widetilde{m}_1, \ \widetilde{N}/2 \in \mathbb{Z}$ relatively prime with each other.

As a consistency check for the existence of $\mathbb{Z}_q^{[1]}$, we show that (3.11) is consistent with the normalization condition of b and u. It is found that

$$\delta_{\Lambda_1} \int db = -\frac{M}{2\pi} \int \Lambda_1 \wedge da \in \frac{2\pi M}{q} \mathbb{Z} ,$$

$$\delta_{\Lambda_1} \int du = \frac{M^2}{8\pi^2 K} \int \Lambda_1 \wedge (da)^2 \in \frac{2\pi M^2}{Kq} \mathbb{Z}$$

Using (2.13) and (3.14) gives

$$\frac{M}{q} = \widetilde{m}_1 L_1 \; ,$$

guaranteeing that the normalization condition for b is consistent with the global $\mathbb{Z}_q^{[1]}$ symmetry transformation. Furthermore, it is verified that

$$\frac{M^2}{Kq} = \frac{M}{K} \widetilde{m}_1 L_1 = \frac{\widetilde{m}_1 L_1^2}{\widetilde{K}} = 2\widetilde{m}_1 \widetilde{r}_1 \ .$$

3.1 Dualizing the 4-form gauge field

It is interesting to rewrite the Lagrangian by dualizing u. The gauge invariant field strength of u is given by (2.9), which obeys the Bianchi identity

$$di_5 = \frac{M^2}{8\pi^2 K} (da)^3 \; .$$

Instead of u, we regard i_5 as an independent variable. This is achieved by using a Lagrange multiplier ξ that imposes the Bianchi identity as a constraint. We consider

$$S' = S + \frac{1}{2\pi} \int \xi \left(di_5 - \frac{M^2}{8\pi^2 K} (da)^3 \right)$$

=
$$\int_{\mathcal{M}_6} \left\{ -\frac{1}{2} |d\phi|^2 - \frac{1}{2} |da|^2 - \frac{1}{8\pi} |db - \frac{M}{2\pi} \omega_3|^2 + \frac{N}{48\pi^3} \phi (da)^3 + \frac{M}{8\pi^2} (da)^2 b - \frac{1}{2} |i_5|^2 + \frac{1}{2\pi} i_5 \wedge (d\xi - Ka) - \frac{M^2}{16\pi^3 K} \xi (da)^3 \right\}.$$

Here, ξ is a compact scalar with $\xi \sim \xi + 2\pi$. The EoM of i_5 reads

$$\star \, i_5 = \frac{1}{2\pi} \left(d\xi - Ka \right) \; . \tag{3.15}$$

The $U(1)^{[0]}$ gauge invariance of i_5 requires that ξ transform under it as

$$\delta^{[0]}\xi = K\lambda_0 \ . \tag{3.16}$$

It follows that ξ is identified with an NG boson associated with the spontaneous breaking of $U(1)^{[0]}$ to $\mathbb{Z}_{K}^{[0]}$. This is consistent with those obtained in the previous section, where the NG boson associated with the symmetry breaking $U(1)^{[0]} \to \mathbb{Z}_{K}^{[0]}$ is given by v_{0} .

By using (3.15), the action becomes

$$S' = \int_{\mathcal{M}_6} \left\{ -\frac{1}{2} |d\phi|^2 - \frac{1}{2} |da|^2 - \frac{1}{8\pi^2} |d\xi - Ka|^2 - \frac{1}{8\pi} |db - \frac{M}{2\pi} \omega_3|^2 + \frac{N}{48\pi^3} \phi (da)^3 - \frac{M^2}{16\pi^3 K} \xi (da)^3 + \frac{M}{8\pi^2} (da)^2 \wedge b \right\}.$$
(3.17)

As a consistency check, we see that the axionic coupling between ξ and $(da)^3$ is gauge invariant by showing that it is properly normalized:

$$\frac{M^2}{16\pi^2 K} = \frac{1}{48\pi^3} \frac{3M^2}{K} = \frac{6r_1}{48\pi^3} ,$$

with $r_1 \in \mathbb{Z}$. Furthermore, we note that the anomaly from the GSWS transformation $\delta^{[0]}b$ is canceled by that from $\delta^{[0]}\xi$ thanks to the axionic coupling.

4 Gauging the EoM-based global symmetry

We gauge the EoM-based global symmetry by promoting $\lambda_{0,1,2,4}$ to non-closed, gauge transformation functions. The gauge invariant action can be obtained by activating the corresponding background gauge fields.

For the 0-form \mathbb{Z}_N symmetry, we consider

$$(A_1, A_0)$$
, $NA_1 = dA_0$.

The field strength for ϕ is defined as $d\phi - A_1$.

For the 2-form \mathbb{Z}_M symmetry, we consider

$$(V_3, V_2)$$
, $MV_3 = dV_2$.

The field strength for b is given by $db - V_3$.

For the 4-form \mathbb{Z}_K symmetry, we consider

$$(W_5, W_4)$$
, $KW_5 = dW_4$,

leading to the field strength $du - W_5$ for u.

We discuss the gauging of the 1-form \mathbb{Z}_q symmetry in some detail. We turn on the associated background gauge field

$$(B_2, B_1)$$
, $qB_2 = dB_1$.

It follows from (3.11) that the field strength for a is given simply by $da - B_2$. For the purpose of defining the field strength of b and u for the \mathbb{Z}_q gauge symmetry, we first modify the transformation law for b in (3.11) so that the RHS becomes \mathbb{Z}_q -invariant:

$$\delta_{\Lambda_1} b = -\frac{M}{2\pi} \Lambda_1 \wedge \left(a - \frac{1}{q} B_1 \right) \; .$$

By noting $\Lambda_1 = \frac{1}{q} \delta_{\Lambda_1} B_1$, this is rewritten as

$$\delta_{\Lambda_1}\overline{b} = 0$$
, $\overline{b} \equiv b + \frac{M}{2\pi q}B_1 \wedge \left(a - \frac{1}{q}B_1\right)$.

The field strength for b is defined naturally as

$$d\overline{b} = db - \frac{M}{2\pi q} da \wedge B_1 + \frac{M}{2\pi} a \wedge B_2 .$$

We also modify the transformation law of u in (3.11) so that the RHS becomes \mathbb{Z}_q -invariant:

$$\delta_{\Lambda_1} u = \Lambda_1 \wedge \left(\frac{M^2}{8\pi^2 K} \,\omega_3(a - \frac{1}{q}B_1) - \frac{3M}{4\pi K} \left(db - \frac{M}{2\pi q} da \wedge B_1 + \frac{M}{2\pi} a \wedge B_2 \right) \right)$$

It then follows that

$$\delta_{\Lambda_1}\overline{u} = 0 , \quad \overline{u} \equiv u - \frac{1}{q}B_1 \wedge \left(\frac{M^2}{8\pi^2 K}\omega_3(a - \frac{1}{q}B_1) - \frac{3M}{4\pi K}d\overline{b}\right) .$$

The field strength for u reads

$$d\overline{u} = du - B_2 \wedge \left(\frac{M^2}{8\pi^2 K}\omega_3(a - \frac{1}{q}B_1) - \frac{3M}{4\pi K}d\overline{b}\right) + \frac{M^2}{8\pi^2 Kq}B_1 \wedge (da - B_2)^2 .$$

Now we examine the combined gauge transformation of $U(1)^{[0]}$ and $\mathbb{Z}_q^{[1]}$ for b and u to define the corresponding field strengths. We first assume that it is given by the linear combonation of $\delta^{[0]}$ and δ_{Λ_1} with the gauge field a appearing in $\delta^{[0]}$ replaced by $a - \frac{1}{q}B_1$. Then, it follows that

$$b \stackrel{?}{\to} b + \frac{M}{2\pi}\lambda_0(da - B_2) - \frac{M}{2\pi}\Lambda_1 \wedge \left(a - \frac{1}{q}B_1\right) .$$

Then, \overline{b} transforms as

$$\overline{b} \stackrel{?}{\to} \overline{b} + \frac{M}{2\pi} \lambda_0 (da - B_2) - \frac{M}{2\pi q} d\lambda_0 \wedge (B_1 + q\Lambda_1)$$
$$= \overline{b} + \frac{M}{2\pi} \lambda_0 (da - B_2) - d\left(\frac{M}{2\pi q} \lambda_0 (B_1 + q\Lambda_1)\right) + \frac{M}{2\pi} \lambda_0 (B_2 + d\Lambda_1)$$

The 4th term makes it impossible to define a gauge invariant field strength even using a GS-like transformation. This problem is resolved by modifying the gauge transformation as

$$b \to b + \frac{M}{2\pi}\lambda_0(da - B_2) - \frac{M}{2\pi}\lambda_0(B_2 + d\Lambda_1) - \frac{M}{2\pi}\Lambda_1 \wedge \left(a - \frac{1}{q}B_1\right) ,$$

from which we obtain

$$\overline{b} \to \overline{b} + \frac{M}{2\pi} \lambda_0 (da - B_2) - d\left(\frac{M}{2\pi q} \lambda_0 \left(B_1 + q\Lambda_1\right)\right)$$

We are thus led to define the field strength of b for the combined gauge transformation:

$$Db = d\overline{b} - \frac{M}{2\pi}\omega_3(a - \frac{1}{q}B_1) \; .$$

It is natural to work with \overline{b} rather than b as an independent, dynamical field. To see this, we note that the combined gauge transformation of db is given by

$$db \to db + \frac{M}{2\pi} d\lambda_0 \wedge (da - B_2) - \frac{M}{2\pi} d\lambda_0 \wedge (B_2 + d\Lambda_1) - \frac{M}{2\pi} d\left(\Lambda_1 \wedge \left(a - \frac{1}{q} B_1\right)\right) \quad .$$

This shows that the gauge transformation spoils the normalization condition $\int db \in 2\pi\mathbb{Z}$. On the contrary, by using

$$d\overline{b} \to d\overline{b} + \frac{M}{2\pi} d\lambda_0 \wedge (da - B_2) ,$$

the gauge transformation acts on $\int d\overline{b}$ as a $2\pi \frac{M}{q}\mathbb{Z}$ shift. This is consistent because $\int d\overline{b}$ is not quantized.

The gauge transformation and the field strength of u for the combined gauge transformation can be derived in the same manner. We assume that the gauge transformation should be modified from the linear combination $(\delta^{[0]} + \delta_{\Lambda_1})u$ by adding an additional transformation:

$$u \to u - \frac{M^2}{8\pi^2 K} \lambda_0 (da - B_2)^2 + \Lambda_1 \wedge \left(\frac{M^2}{8\pi^2 K} \omega_3 (a - \frac{1}{q}B_1) - \frac{3M}{4\pi K} d\overline{b}\right) + \delta' u$$

from which

$$\overline{u} \to \overline{u} - \frac{M^2}{8\pi^2 K} \lambda_0 (da - B_2)^2 - d\left(\frac{M^2}{4\pi^2 Kq} \lambda_0 \left(B_1 + q\Lambda_1\right) \wedge (da - B_2)\right) \\ + \frac{M^2}{4\pi^2 K} \lambda_0 \left(B_2 + d\Lambda_1\right) \wedge (da - B_2) + \delta' u \;.$$

We set

$$\delta' u = -\frac{M^2}{4\pi^2 K} \lambda_0 \left(B_2 + d\Lambda_1 \right) \wedge \left(da - B_2 \right) \,.$$

It follows that the field strength for u reads

$$Du = d\overline{u} + \frac{M^2}{8\pi^2 K} \omega_5 \left(a - \frac{1}{q}B_1\right) \,.$$

As in the case for b, we should treat \overline{u} as an independent, dynamical field rather than u.

We now derive the action in the presence of the background gauge fields. This is obtained by replacing the derivative of the dynamical fields in (3.1) with the field strengths we have constructed:

$$S = -\int_{\mathcal{M}_6} \left(\frac{1}{2} |d\phi - A_1|^2 + \frac{1}{2} |da - B_2|^2 + \frac{1}{8\pi} |Db - V_3|^2 + \frac{1}{2} |Du - W_5|^2 \right) + S_{\rm CS} , \qquad (4.1)$$

where

$$S_{\rm CS} = \int_{\Omega_{\mathcal{M}_6}} \left(\frac{N}{48\pi^3} (d\phi - A_1) \wedge (da - B_2)^3 + \frac{M}{8\pi^2} (da - B_2)^2 \wedge (Db - V_3) + \frac{K}{2\pi} (da - B_2) \wedge (Du - W_5) \right)$$

=
$$\int_{\Omega_{\mathcal{M}_6}} \left(\frac{N}{48\pi^3} (d\phi - A_1) \wedge (da - B_2)^3 + \frac{M}{8\pi^2} (da - B_2)^2 \wedge (d\overline{b} - V_3) + \frac{K}{2\pi} (da - B_2) \wedge (d\overline{u} - W_5) \right)$$

(4.2)

As a consistency check, we note that the CS action is left unchanged manifestly under the small gauge transformations. Furthermore, the integrand of the CS action is a closed 7-form so that it is regarded as a 6d action.

4.1 Operator-valued ambiguity

We now examine if the resultant CS action (4.2) suffers an operator-valued ambiguity, which is a quantum inconsistency stating that the quantum parts of the CS action written in terms of the dynamical fields depend on how to lift them to a seven-manifold. It is found that there arise the operator-valued ambiguities as far as the background gauge field (B_2, B_1) is turned on. To see this, focus on the term

$$\int_{\Omega_{\mathcal{M}_6}} \frac{K}{2\pi} da \wedge d\overline{u}$$

This depends on the choice of the seven-manifold $\Omega_{\mathcal{M}_6}$ even if K is an integer because the integral $\int d\overline{b}$ over any closed 3-cycle is not quantized in the presence of (B_2, B_1) . This is a manifestation of the quantum anomaly of the EoM-based $\mathbb{Z}_q^{[1]}$ symmetry via the operator-valued ambiguity. This result is consistent with those obtained in [40].² This paper discusses a discrete 1-form symmetry in $\mathcal{N} = (1,0)$ 6d theories coupled with tensor and vector multiplets. It is shown that turning on the background gauge field associated with a 1-form symmetry spoils the Dirac quantization condition for BPS string charges, which gives rise to massive excitation modes that break the 1-form symmetry explicitly.

In order to remove the operator-valued ambiguity, we have to turn off (B_2, B_1) . Then, the dynamical field \overline{b} and \overline{u} reduces to b and u, respectively. The resultant 6d model has no operator-valued ambiguity, because the flux integral of b and u over any closed cycles is quantized.

5 Discussions

As discussed in [1], the 6d axion-electrodynamics has rich higher-group structure that is manifested as GS-like transformation laws for the background gauge fields associated with the CW symmetries. A key role is played by the background gauge field for a discrete 1-form symmetry, (B_2, B_1) , as clear from the gauge invariant field strengths given in (2.2). In this paper, we have investigated

²See also [41] for a related work.

the model that is obtained by promoting the CW background gauge field B_2^{CW} and D_4^{CW} to the dynamical field *b* and *u*, respectively. It is shown that the background gauge field for the EoMbased $\mathbb{Z}_q^{[1]}$ symmetry, which is denoted by (B_2, B_1) as well, must be turned off because it would cause an operator-valued ambiguity. We examine whether there exists a nontrivial higher-group structure even in the absence of (B_2, B_1) .

The CW currents in this model are written in terms of the wedge products of the gauge invariant closed forms

$$du + \frac{M}{4\pi K} da \wedge db \ , \ \ da \ , \ \ d\phi \ .$$

The gauge invariance of the CW 5-form is guaranteed because

$$du + \frac{M}{4\pi K} da \wedge db = du + \frac{M^2}{8\pi^2 K} \omega_5(a) + \frac{M}{4\pi K} da \wedge \left(db - \frac{M}{2\pi} \omega_3(a) \right)$$

We first compute the integral of the 5-form over a closed 5-cycle. Using (2.11) and (2.13) gives

$$\frac{M}{2K} = \frac{L_1}{2\tilde{K}}$$

Set

$$s = \gcd(\frac{L_1}{2}, \widetilde{K}) ,$$

or equivalently

$$\frac{L_1}{2} = sL' , \quad \widetilde{K} = sK' ,$$

with L' and K' relatively prime. It is found from these and (2.12) that there exists an odd integer s' such that

$$2s = s'K'$$

with K' even. It then follows that

$$\int \left(du + \frac{M}{4\pi K} da \wedge db \right) \in \frac{2\pi}{K'} \mathbb{Z} \; .$$

It is thus natural to normalize the CW 5-form as

$$K'\left(du + \frac{M}{4\pi K}da \wedge db\right) = K'du + \frac{L'}{2\pi}da \wedge db$$

We consider the CW 6-form current

$$d\phi \wedge \left(K'du + \frac{L'}{2\pi} da \wedge db \right)$$

Let ϑ_0^{CW} be the CW 0-form background gauge field coupled with this current with the normalization condition given by $\int d\vartheta_0^{\text{CW}} \in 2\pi\mathbb{Z}$. We propose that the interaction term in the presence of (A_1, A_0) , (V_3, V_2) and (W_5, W_4) takes the form

$$S_{\rm CW} = \frac{2m_1 sQ}{4\pi^2} \int_{\mathcal{M}_6} \vartheta_0^{\rm CW} \left(d\phi - A_1 \right) \wedge \left[K'(du - W_5) + \frac{L'}{2\pi} da \wedge (db - V_3) \right] ,$$

with $Q \in \mathbb{Z}$. It is found that the operator-valued ambiguity comes from the term

$$-\frac{2m_1sQ}{4\pi^2}\int_{\mathcal{M}_6}\vartheta_0^{\mathrm{CW}}A_1\wedge\left(K'du+\frac{L'}{2\pi}da\wedge(db-V_3)\right)$$

This is canceled by adding the local counterterm

$$\Delta S = \frac{2m_1s}{2\pi} \int_{\mathcal{M}_6} \left(\eta_1^{\mathrm{CW}} + \frac{Q}{2\pi} \vartheta_0^{\mathrm{CW}} A_1 \right) \wedge \left(K'(du - W_5) + \frac{L'}{2\pi} da \wedge (db - V_3) \right) \; .$$

Here, η_1^{CW} is the background gauge field associated with the CW current $K'du + \frac{L'}{2\pi}da \wedge db$ with $\int d\eta_1^{\text{CW}} \in 2\pi\mathbb{Z}$. It is easy to see that there appears no operator-valued ambiguity in $S_{\text{CW}} + \Delta S$. The gauge invariance of ΔS induces a GS-like transformation for the 1-form CW gauge field η_1^{CW} that is charactorized by the integer Q.

So far, we have considered the cases with a nontrivial $K \in \mathbb{Z}$. When restricting to K = 1, we have q = 1 so that $\mathbb{Z}_q^{[1]}$ symmetry is trivial. It is seen that this case admits much richer higher-group structure. To explore it in more detail deserves a further study.

A Dimensional reduction to 5d

Start with part of the action (3.1):

$$\hat{S} = -\int_{\mathcal{M}_6} \left(\frac{1}{2} |\hat{d}\hat{\phi}|^2 + \frac{1}{2} |\hat{d}\hat{a}|^2 + \frac{1}{8\pi} |\hat{d}\hat{b} - \frac{M}{2\pi} \hat{\omega}_3|^2 + \frac{1}{2} \left| \hat{d}\hat{u} + \frac{M^2}{8\pi^2 K} \hat{\omega}_5 \right|^2 - \frac{N}{48\pi^3} \hat{\phi} (\hat{d}\hat{a})^3 - \frac{M}{8\pi^2} (\hat{d}\hat{a})^2 \hat{b} - \frac{K}{2\pi} \hat{d}\hat{a} \hat{u} \right)$$

The fields with a hat are defined in 6d before dimensional reduction. Let

$$M, N = 0, 1, \dots, 5$$
, (6d spacetime indices)
 $\mu, \nu = 0, 1, \dots, 4$, (5d spacetime indices)

with $X_5 \sim X_5 + 1$. The space-time metric is given by

$$\eta_{MN} = \text{diag}(-+++++)$$
.

Define the 5d gauge fields as

$$\hat{b} = b_2 + b_1 \wedge dX_5$$
, $\hat{a} = a + \zeta dX_5$, $\hat{u} = u + u_3 \wedge dX_5$.

This choice is natural because the fluxes of the resultant 5d gauge fields are normalized as $2\pi\mathbb{Z}$. Then,

$$\hat{db} - \frac{M}{2\pi}\hat{\omega}_3 = db_2 - \frac{M}{2\pi}\omega_3 + \left(db_1 - \frac{M}{2\pi}\left(\zeta da - d\zeta \wedge a\right)\right) \wedge dX_5 \equiv G_3 + \mathcal{F}_2 \wedge dX_5 \ .$$

Let $\hat{\star}$ and \star be the Hodge star operation in 6d and 5d, respectively. Using the formulae

$$\hat{\star} (dX^{\mu_1} \wedge \dots \wedge dX^{\mu_p}) = \star (dX^{\mu_1} \wedge \dots \wedge dX^{\mu_p}) \wedge dX_5 ,$$
$$\hat{\star} (dX^{\mu_1} \wedge \dots \wedge dX^{\mu_q} \wedge dX_5) = (-)^{q+1} \star (dX^{\mu_1} \wedge \dots \wedge dX^{\mu_q}) ,$$

we obtain

$$\hat{\star} \left(\hat{d}\hat{b} - \frac{M}{2\pi} \hat{\omega}_3 \right) = \star G_3 \wedge dX_5 - \star \mathcal{F}_2 \; .$$

The self-duality condition for b is rewritten as

$$\star G_3 = \mathcal{F}_2 , \quad \star \mathcal{F}_2 = -G_3 . \tag{A.1}$$

The field strength of \hat{u}_4 is reduced as

$$\hat{i}_5 = \hat{d}\hat{u}_4 + \frac{M^2}{8\pi^2 K}\hat{\omega}_5 = i_5 + \mathcal{F}_4 \wedge dX_5 ,$$

where

$$i_5 = du_4 + \frac{M^2}{8\pi^2 K} \omega_5 , \quad \mathcal{F}_4 = du_3 + \frac{M^2}{8\pi^2 K} \left(\zeta (da)^2 - 2d\zeta \wedge \omega_3 \right)$$

Then,

$$\hat{\star}\left(\hat{d}\hat{u}_4 + \frac{M^2}{8\pi^2 K}\hat{\omega}_5\right) = (\star i_5) \wedge dX_5 - \star \mathcal{F}_4 \; .$$

The 0-form gauge symmetry transformation acts on \hat{a} as

$$\hat{\delta}^{[0]}\hat{a} = \hat{d}\hat{\lambda}_0 . \tag{A.2}$$

•

Reducing $\hat{\lambda}_0$ to 5d leads to

$$\delta^{[0]}a = d\lambda_0 \; ,$$

with ζ left unchanged. ζ is a compact boson of period 2π . This is seen from the 0-form gauge transformation with $\hat{\lambda}_0 = 2\pi X_5$.

The 1-form gauge symmetry transformation acts on \hat{b} as

$$\hat{\delta}^{[1]}\hat{b} = \hat{d}\hat{\lambda}_1$$
.

Reducing $\hat{\lambda}_1$ to 5d as

$$\hat{\lambda}_1 = \lambda_1 + \lambda_0 \, dX_5 \; ,$$

it acts as

$$\delta^{[1]}b_2 = d\lambda_1 , \quad \delta^{[1]}b_1 = d\lambda_0 .$$

The 3-form gauge symmetry transformation acts on \hat{u}_4 as

$$\hat{\delta}^{[3]}\hat{u}_4 = \hat{d}\hat{\lambda}_3$$
.

Reducing $\hat{\lambda}_3$ to 5d as

$$\hat{\lambda}_3 = \lambda_3 + \lambda_2 \wedge dX_5 \; ,$$

it acts as

$$\delta^{[3]}u_4 = d\lambda_3 , \quad \delta^{[3]}u_3 = d\lambda_2$$

The GS laws for \hat{b} and \hat{u}_4 under $U(1)^{[0]}$ are given in (2.10), which is reduced to 5d as

$$\delta^{[0]}b_2 = \frac{M}{2\pi}\lambda_0 da , \quad \delta^{[0]}b_1 = \frac{M}{2\pi}\lambda_0 d\zeta ,$$

and

$$\delta^{[0]}u_4 = -\frac{M^2}{8\pi^2 K} \lambda_0 (da)^2 , \quad \delta^{[0]}u_3 = -\frac{M^2}{4\pi^2 K} \lambda_0 d\zeta \wedge da ,$$

respectively. Note that these are consistent with the normalization condition for the 5d gauge fields.

Upon dimensional reduction to 5d, the action becomes

$$S = \int \left[-\frac{1}{8\pi} |G_3|^2 - \frac{1}{8\pi} |\mathcal{F}_2|^2 + \frac{M}{4\pi^2} b_2 \wedge d\zeta \wedge da + \frac{M}{8\pi^2} b_1 \wedge (da)^2 - \frac{1}{2} |i_5|^2 - \frac{1}{2} |\mathcal{F}_4|^2 + \frac{K}{2\pi} u_4 \wedge d\zeta + \frac{K}{2\pi} u_3 \wedge da - \frac{1}{2} |d\phi|^2 + \frac{N}{16\pi^3} \phi \, d\zeta \wedge (da)^2 - \frac{1}{2} |da|^2 - \frac{1}{2} |d\zeta|^2 \right].$$

Here, we used $\int dX_5 = 1$. As a check, the equations of motion for b_2 and b_1 read

$$d \star G_3 = -\frac{M}{\pi} d\zeta \wedge da , \quad d \star \mathcal{F}_2 = \frac{M}{2\pi} (da)^2 .$$
 (A.3)

These are consistent with the self-duality consistion (A.1).

So far, we regard b_2 and b_1 as independent variables and impose the self-duality condition (A.1) on the equations of motion. Instead, we attempt to reformulate the action so that G_3 becomes an independent variable rather than b_2 . For this purpose, we note that the third term in the 5d action is rewritten as

$$\frac{M}{4\pi^2}b_2 \wedge d\zeta \wedge da = \Delta \mathcal{L} + \frac{1}{4\pi}G_3 \wedge \mathcal{F}_2 - \frac{M^2}{16\pi^3}\zeta \,\omega_5 + \frac{M}{8\pi^2}b_1 \wedge (da)^2 ,$$

where

$$\Delta \mathcal{L} = -\frac{1}{4\pi} d \left[b_2 \wedge \mathcal{F}_2 - \frac{M}{2\pi} b_1 \wedge \omega_3 \right] \; .$$

Now we define

$$\begin{split} \bar{S} &= S - \int \Delta \mathcal{L} \\ &= \int \left[-\frac{1}{8\pi} |G_3|^2 - \frac{1}{8\pi} |\mathcal{F}_2|^2 + \frac{1}{4\pi} G_3 \wedge \mathcal{F}_2 + \frac{M}{4\pi^2} b_1 \wedge (da)^2 - \frac{M^2}{16\pi^3} \zeta \, \omega_5 \right. \\ &\quad \left. -\frac{1}{2} |i_5|^2 - \frac{1}{2} |\mathcal{F}_4|^2 + \frac{K}{2\pi} u_4 \wedge d\zeta + \frac{K}{2\pi} u_3 \wedge da - \frac{1}{2} |d\phi|^2 + \frac{N}{16\pi^3} \phi \, d\zeta \wedge (da)^2 \right. \\ &\quad \left. -\frac{1}{2} |da|^2 - \frac{1}{2} |d\zeta|^2 \right] \,. \end{split}$$

This technique is utilized in [42].

Regarding G_3 as an independent variable, the EoM for G_3 reads

$$\star G_3 = \mathcal{F}_2 \; ,$$

reproducing the self-duality condition (A.1). We note that the Gauss law constraint for G_3 follows from the EoM for b_1 . This guarantees that G_3 is regarded as an independent variable instead of b_2 . Integrating out G_3 leads to

$$\bar{S} = \int \left[-\frac{1}{2} |da|^2 - \frac{1}{2} |d\zeta|^2 - \frac{1}{4\pi} |\mathcal{F}_2|^2 - \frac{1}{2} |i_5|^2 - \frac{1}{2} |\mathcal{F}_4|^2 - \frac{1}{2} |d\phi|^2 + \frac{M}{4\pi^2} b_1 \wedge (da)^2 - \frac{M^2}{16\pi^3} \zeta \,\omega_5 + \frac{K}{2\pi} u_4 \wedge d\zeta + \frac{K}{2\pi} u_3 \wedge da + \frac{N}{16\pi^3} \phi \,d\zeta \wedge (da)^2 \right].$$
(A.4)

The EoM for b_1 is

$$d \star \mathcal{F}_2 = \frac{M}{2\pi} (da)^2 . \tag{A.5}$$

The EoM for u_4 is

$$d \star i_5 = -\frac{K}{2\pi} d\zeta \ . \tag{A.6}$$

The EoM for u_3 is

$$d \star \mathcal{F}_4 = \frac{K}{2\pi} da \ . \tag{A.7}$$

The EoM for ϕ is

$$d \star d\phi = -\frac{N}{16\pi^3} d\zeta \wedge (da)^2$$
.

A bit lengthy computation shows that the EoMs for ζ and a read

$$d \star d\zeta + \frac{M}{2\pi^2} da \wedge \star \mathcal{F}_2 - \frac{K}{2\pi} i_5 - \frac{3M^2}{8\pi^2 K} (da)^2 \wedge \star \mathcal{F}_4 - \frac{N}{16\pi^3} d\phi \wedge (da)^2 = 0 ,$$

and

$$\begin{aligned} -d \star da &+ \frac{M}{2\pi^2} d\zeta \wedge \star \mathcal{F}_2 + \frac{M}{2\pi^2} da \wedge \mathcal{F}_2 - \frac{3M^2}{8\pi^2 K} (da)^2 \wedge \star i_5 - \frac{3M^2}{4\pi^2 K} d\zeta \wedge da \wedge \star \mathcal{F}_4 \\ &+ \frac{K}{2\pi} \mathcal{F}_4 + \frac{N}{8\pi^3} d\phi \wedge d\zeta \wedge da = 0 , \end{aligned}$$

respectively. Note that all the EoMs are gauge invariant manifestly.

We discuss the symmetries realized in (A.4) and their relation to the 6d theory. (A.5) gives a conserved current

$$j_{\rm EoM}^{[1]} = \frac{1}{2\pi} \star \mathcal{F}_2 - \frac{M}{4\pi^2} \omega_3 \; .$$

This defines a 1-form symmetry generator. It is easy to see that the gauge invariance requires it to generate $\mathbb{Z}_M^{[1]}$. This is a manifestation of the 2-form \mathbb{Z}_M symmetry in 6d after dimensional reduction to 5d.

The EoM-based conserved current for ζ and a can be most easily obtained by dimensionally reducing the EoM-based conserved current for \hat{a}

$$\hat{j}_{\rm EoM}^{\prime[1]} = \hat{\star}\hat{d}\hat{a} - \frac{M}{8\pi^2}\hat{d}\hat{a} \wedge \hat{b} - \frac{3M}{4\pi K}\left(\hat{d}\hat{b} - \frac{M}{2\pi}\hat{\omega}_3\right) \wedge \hat{\star}\left(\hat{d}\hat{u}_4 + \frac{M^2}{8\pi^2 K}\hat{\omega}_5\right) - \frac{K}{2\pi}\hat{u}_4 - \frac{N}{16\pi^3}\hat{\phi}\left(\hat{d}\hat{a}\right)^2.$$

to 5d:

$$\hat{j}_{\text{EoM}}^{\prime[1]} = j_{\text{EoM}}^{\prime[0]} + j_{\text{EoM}}^{\prime[1]} \wedge dX_5 \;.$$

We find

$$j_{\rm EoM}^{\prime [0]} = \star d\zeta - \frac{M}{8\pi^2} da \wedge b_2 - \frac{3M}{4\pi K} \star \mathcal{F}_2 \wedge \star \mathcal{F}_4 - \frac{K}{2\pi} u_4 - \frac{N}{16\pi^3} \phi(da)^2 ,$$

$$j_{\rm EoM}^{\prime [1]} = \star da - \frac{M}{8\pi^2} \left(da \wedge b_1 + d\zeta \wedge b_2 \right) + \frac{3M}{4\pi K} \star \mathcal{F}_2 \wedge \star i_5 - \frac{3M}{4\pi K} \mathcal{F}_2 \wedge \star \mathcal{F}_4 - \frac{K}{2\pi} u_3 - \frac{N}{8\pi^3} \phi d\zeta \wedge da .$$

It is easy to show that $dj'_{\rm EoM}^{[0]} = dj'_{\rm EoM}^{[1]} = 0$ by using the EoMs in 5d. Note that both depend on b_2 , which is eliminated nonlocally by solving (A.1).

Now we examine the gauge invariance condition of the symmetry generators associated with these currents. That for $j'_{EoM}^{[0]}$ requires that the corresponding symmetry generator define \mathbb{Z}_q with q given in (3.13), because the CS-like terms in $j'_{EoM}^{[0]}$ take exactly the same form as in $\hat{j}'_{EoM}^{[1]}$. Furthermore, it is seen that the symmetry generator for $j'_{EoM}^{[1]}$ is required to generate \mathbb{Z}_q as well by examining the difference of two ways of lifting the CS-like terms in $j'_{EoM}^{[1]}$ to a higher-dimensions.

References

- [1] Tatsuki Nakajima, Tadakatsu Sakai, and Ryo Yokokura. Higher-group structure in 2n-dimensional axion-electrodynamics. JHEP, 01:150, 2023. arXiv:2211.13861, doi:10.1007/JHEP01(2023)150.
- Yoshimasa Hidaka, Muneto Nitta, and Ryo Yokokura. Higher-form symmetries and 3group in axion electrodynamics. *Phys. Lett. B*, 808:135672, 2020. arXiv:2006.12532, doi:10.1016/j.physletb.2020.135672.

- [3] Yoshimasa Hidaka, Muneto Nitta, and Ryo Yokokura. Global 3-group symmetry and 't Hooft anomalies in axion electrodynamics. JHEP, 01:173, 2021. arXiv:2009.14368, doi:10.1007/JHEP01(2021)173.
- [4] Michael B. Green and John H. Schwarz. Superstring Field Theory. Nucl. Phys. B, 243:475–536, 1984. doi:10.1016/0550-3213(84)90488-7.
- [5] Anton Kapustin and Ryan Thorngren. Higher Symmetry and Gapped Phases of Gauge Theories. Prog. Math., 324:177-202, 2017. arXiv:1309.4721, doi:10.1007/978-3-319-59939-7_5.
- [6] Anton Kapustin and Ryan Thorngren. Topological Field Theory on a Lattice, Discrete Theta-Angles and Confinement. Adv. Theor. Math. Phys., 18(5):1233-1247, 2014. arXiv:1308.2926, doi:10.4310/ATMP.2014.v18.n5.a4.
- [7] Eric Sharpe. Notes on generalized global symmetries in QFT. Fortsch. Phys., 63:659-682, 2015. arXiv:1508.04770, doi:10.1002/prop.201500048.
- [8] Lakshya Bhardwaj, Davide Gaiotto, and Anton Kapustin. State sum constructions of spin-TFTs and string net constructions of fermionic phases of matter. JHEP, 04:096, 2017. arXiv:1605.01640, doi:10.1007/JHEP04(2017)096.
- [9] Anton Kapustin and Ryan Thorngren. Fermionic SPT phases in higher dimensions and bosonization. JHEP, 10:080, 2017. arXiv:1701.08264, doi:10.1007/JHEP10(2017)080.
- [10] Yuji Tachikawa. On gauging finite subgroups. SciPost Phys., 8(1):015, 2020.
 arXiv:1712.09542, doi:10.21468/SciPostPhys.8.1.015.
- [11] R. Costa de Almeida, J. P. Ibieta-Jimenez, J. Lorca Espiro, and P. Teotonio-Sobrinho. Topological Order from a Cohomological and Higher Gauge Theory perspective. 11 2017. arXiv:1711.04186.
- [12] Francesco Benini, Clay Córdova, and Po-Shen Hsin. On 2-Group Global Symmetries and their Anomalies. JHEP, 03:118, 2019. arXiv:1803.09336, doi:10.1007/JHEP03(2019)118.
- [13] Clay Córdova, Thomas T. Dumitrescu, and Kenneth Intriligator. Exploring 2-Group Global Symmetries. JHEP, 02:184, 2019. arXiv:1802.04790, doi:10.1007/JHEP02(2019)184.

- [14] Clement Delcamp and Apoorv Tiwari. From gauge to higher gauge models of topological phases. JHEP, 10:049, 2018. arXiv:1802.10104, doi:10.1007/JHEP10(2018)049.
- [15] Xiao-Gang Wen. Emergent anomalous higher symmetries from topological order and from dynamical electromagnetic field in condensed matter systems. *Phys. Rev. B*, 99(20):205139, 2019. arXiv:1812.02517, doi:10.1103/PhysRevB.99.205139.
- [16] Clement Delcamp and Apoorv Tiwari. On 2-form gauge models of topological phases. JHEP, 05:064, 2019. arXiv:1901.02249, doi:10.1007/JHEP05(2019)064.
- [17] Ryan Thorngren. Topological quantum field theory, symmetry breaking, and finite gauge theory in 3+1D. *Phys. Rev. B*, 101(24):245160, 2020. arXiv:2001.11938, doi:10.1103/PhysRevB.101.245160.
- [18] Clay Cordova, Thomas T. Dumitrescu, and Kenneth Intriligator. 2-Group Global Symmetries and Anomalies in Six-Dimensional Quantum Field Theories. JHEP, 04:252, 2021. arXiv:2009.00138, doi:10.1007/JHEP04(2021)252.
- [19] Po-Shen Hsin and Alex Turzillo. Symmetry-enriched quantum spin liquids in (3 + 1)d. JHEP, 09:022, 2020. arXiv:1904.11550, doi:10.1007/JHEP09(2020)022.
- [20] Po-Shen Hsin and Ho Tat Lam. Discrete theta angles, symmetries and anomalies. SciPost Phys., 10(2):032, 2021. arXiv:2007.05915, doi:10.21468/SciPostPhys.10.2.032.
- [21] Sergei Gukov, Po-Shen Hsin, and Du Pei. Generalized global symmetries of T[M] theories.
 Part I. JHEP, 04:232, 2021. arXiv:2010.15890, doi:10.1007/JHEP04(2021)232.
- [22] Nabil Iqbal and Napat Poovuttikul. 2-group global symmetries, hydrodynamics and holography. SciPost Phys., 15(2):063, 2023. arXiv:2010.00320, doi:10.21468/SciPostPhys.15.2.063.
- [23] Tomáš Brauner. Field theories with higher-group symmetry from composite currents. JHEP, 04:045, 2021. arXiv:2012.00051, doi:10.1007/JHEP04(2021)045.
- [24] Oliver DeWolfe and Kenneth Higginbotham. Generalized symmetries and 2-groups via electromagnetic duality in AdS/CFT. Phys. Rev. D, 103(2):026011, 2021. arXiv:2010.06594, doi:10.1103/PhysRevD.103.026011.

- [25] Ben Heidenreich, Jacob McNamara, Miguel Montero, Matthew Reece, Tom Rudelius, and Irene Valenzuela. Chern-Weil global symmetries and how quantum gravity avoids them. JHEP, 11:053, 2021. arXiv:2012.00009, doi:10.1007/JHEP11(2021)053.
- [26] Fabio Apruzzi, Sakura Schafer-Nameki, Lakshya Bhardwaj, and Jihwan Oh. The Global Form of Flavor Symmetries and 2-Group Symmetries in 5d SCFTs. *SciPost Phys.*, 13(2):024, 2022. arXiv:2105.08724, doi:10.21468/SciPostPhys.13.2.024.
- [27] Lakshya Bhardwaj. 2-Group symmetries in class S. SciPost Phys., 12(5):152, 2022.
 arXiv:2107.06816, doi:10.21468/SciPostPhys.12.5.152.
- [28] Yoshimasa Hidaka, Muneto Nitta, and Ryo Yokokura. Topological axion electrodynamics and 4-group symmetry. *Phys. Lett. B*, 823:136762, 2021. arXiv:2107.08753, doi:10.1016/j.physletb.2021.136762.
- [29] Yoshimasa Hidaka, Muneto Nitta, and Ryo Yokokura. Global 4-group symmetry and 't Hooft anomalies in topological axion electrodynamics. *PTEP*, 2022(4):04A109, 2022.
 arXiv:2108.12564, doi:10.1093/ptep/ptab150.
- [30] Fabio Apruzzi, Lakshya Bhardwaj, Dewi S. W. Gould, and Sakura Schafer-Nameki. 2-Group Symmetries and their Classification in 6d. 10 2021. arXiv:2110.14647.
- [31] Michele Del Zotto, Iñaki García Etxebarria, and Sakura Schafer-Nameki. 2-Group Symmetries and M-Theory. SciPost Phys., 13:105, 2022. arXiv:2203.10097, doi:10.21468/SciPostPhys.13.5.105.
- [32] Mirjam Cvetič, Jonathan J. Heckman, Max Hübner, and Ethan Torres. 0-form, 1-form, and 2-group symmetries via cutting and gluing of orbifolds. *Phys. Rev. D*, 106(10):106003, 2022. arXiv:2203.10102, doi:10.1103/PhysRevD.106.106003.
- [33] Tatsuki Nakajima, Tadakatsu Sakai, and Ryo Yokokura. BCF anomaly and highergroup structure in the low energy effective theories of mesons. JHEP, 01:175, 2023. arXiv:2212.12987, doi:10.1007/JHEP01(2023)175.
- [34] Naoto Kan, Okuto Morikawa, Yuta Nagoya, and Hiroki Wada. Higher-group structure in lattice Abelian gauge theory under instanton-sum modification. *Eur. Phys.*

J. C, 83(6):481, 2023. [Erratum: Eur.Phys.J.C 84, 22 (2024)]. arXiv:2302.13466, doi:10.1140/epjc/s10052-023-11616-6.

- [35] Yu Nakayama. Higher group Weyl symmetry. Phys. Lett. B, 848:138405, 2024. arXiv:2306.12662, doi:10.1016/j.physletb.2023.138405.
- [36] Michael B. Green, John H. Schwarz, and Peter C. West. Anomaly Free Chiral Theories in Six-Dimensions. Nucl. Phys. B, 254:327–348, 1985. doi:10.1016/0550-3213(85)90222-6.
- [37] Augusto Sagnotti. A Note on the Green-Schwarz mechanism in open string theories. Phys. Lett. B, 294:196-203, 1992. arXiv:hep-th/9210127, doi:10.1016/0370-2693(92)90682-T.
- [38] R. D. Peccei and Helen R. Quinn. CP Conservation in the Presence of Instantons. Phys. Rev. Lett., 38:1440-1443, 1977. doi:10.1103/PhysRevLett.38.1440.
- [39] Tom Banks and Nathan Seiberg. Symmetries and Strings in Field Theory and Gravity. Phys. Rev. D, 83:084019, 2011. arXiv:1011.5120, doi:10.1103/PhysRevD.83.084019.
- [40] Fabio Apruzzi, Markus Dierigl, and Ling Lin. The Fate of Discrete 1-Form Symmetries in 6d. SciPost Phys., 12:047, 2022. arXiv:2008.09117, doi:10.21468/SciPostPhys.12.2.047.
- [41] Pietro Benetti Genolini and Luigi Tizzano. Instantons, symmetries and anomalies in five dimensions. JHEP, 04:188, 2021. arXiv:2009.07873, doi:10.1007/JHEP04(2021)188.
- [42] Federico Bonetti and Thomas W. Grimm. Six-dimensional (1,0) effective action of Ftheory via M-theory on Calabi-Yau threefolds. JHEP, 05:019, 2012. arXiv:1112.1082, doi:10.1007/JHEP05(2012)019.