

# The universal $\mathfrak{gl}$ -weight system and the chromatic polynomial

M. Kazarian<sup>\*</sup>; N. Kodaneva<sup>†</sup>; S. Lando<sup>‡</sup>

June 18, 2024

---

<sup>\*</sup>Higher School of Economics, Skolkovo Institute of Science and Technology

<sup>†</sup>Higher School of Economics

<sup>‡</sup>Higher School of Economics, Skolkovo Institute of Science and Technology

$\mathfrak{gl}(N)$	Lie algebra of all $N \times N$ matrices
$E_{i,j} \in \mathfrak{gl}(N)$	matrix unit, $1 \leq i, j \leq N$
$w_{\mathfrak{gl}}$	weight system associated to the series $\mathfrak{gl}$ of Lie algebras
$U\mathfrak{gl}(\infty)$	universal enveloping algebra of the limit Lie algebra $\mathfrak{gl}(\infty)$
$ZU\mathfrak{gl}(N)$	center of $U\mathfrak{gl}(N)$
$C_1, C_2, \dots$	Casimir elements generating the center of $U\mathfrak{gl}(\infty)$
$\alpha \in \mathbb{S}_m$	permutation, which determines the hyperedges of the hypermap
$\mathbb{S}_m$	group of permutations of $m$ elements
$G(\alpha)$	the digraph of a permutation $\alpha$
$f(\alpha)$	number of faces of $\alpha$
$c(\alpha)$	number of cycles in $\alpha$
$V(\alpha)$	set of cycles of $\alpha$
$v_1, v_2, \dots \in V(\alpha)$	cycles of $\alpha$
$a(\alpha)$	number of ascents of $\alpha$
$\alpha _U$	restriction of $\alpha$ to $U \subset \{1, 2, \dots, m\}$
$\gamma(\alpha)$	the intersection graph of $\alpha$
$\sigma = (1, 2, \dots, m) \in \mathbb{S}_m$	standard long cycle, the hypervertex of a hypermap
$\varphi \in \mathbb{S}_m$	permutation, which determines the hypofaces of the hypermap
$\mathcal{A}_m$	the vector space spanned by equivalence classes of permutations of $m$ elements
$[\alpha]$	equivalence class of $\alpha$
$\mathcal{A}$	the rotational Hopf algebra of permutations
$\alpha_1 \# \alpha_2$	concatenation product of permutations $\alpha_1$ and $\alpha_2$
$\mu(\alpha)$	coproduct of $\alpha$
$\bar{\alpha}$	canonical form of $\alpha$
$P$	subspace of primitive elements of a Hopf algebra
$\pi$	projection to the subspace of primitive elements
$\mathcal{A}_m^+$	the vector space spanned by equivalence classes of positive permutations of $m$ elements
$\mathcal{A}^+$	the rotational Hopf algebra of positive permutations
$\mathcal{B}_m$	the vector space spanned by equivalence classes of involutions of $m$ elements without fixed points
$\mathcal{B}$	the rotational Hopf algebra of chord diagrams
$\mathcal{G}_n$	the vector space spanned by graphs with $n$ vertices
$\mathcal{G}$	the Hopf algebra of graphs
$G$	simple graph
$K_n$	complete graph with $n$ vertices
$V(G)$	the set of vertices of $G$
$G _U$	the subgraph of $G$ induced by $U \subset V(G)$
$\chi_G(x)$	chromatic polynomial of $G$
$X(\alpha)(N; p_1, p_2, \dots)$	prechromatic substitution
$X_0(\alpha)$	top coefficient of $X_\alpha$

# 1 Introduction

Weight systems are functions on chord diagrams satisfying so-called Vassiliev's 4-term relations. They are closely related to finite type knot invariants, see [17, 12].

Certain weight systems can be derived from graph invariants, see a recent account in [9]. One of the first examples of such an invariant was the chromatic polynomial [4].

Another main source of weight systems are Lie algebras, the construction due to D. Bar-Natan [1] and M. Kontsevich [12]. In recent papers [18, 9], the weight systems associated to the Lie algebras  $\mathfrak{gl}(N)$ ,  $N = 1, 2, 3, \dots$ , were unified in a universal  $\mathfrak{gl}$ -weight system, which takes values in the ring  $\mathbb{C}[N, C_1, C_2, C_3, \dots]$  of polynomials in infinitely many variables. Note that this weight system is associated to the HOMFLYPT polynomial, which is an important and powerful knot invariant. The unification has been achieved by extending the  $\mathfrak{gl}(N)$ -weight systems from chord diagrams, which can be considered as involutions without fixed points modulo cyclic shifts, to arbitrary permutations<sup>1</sup>. The main goal of the extension was to produce an efficient way to compute explicitly the values of the  $\mathfrak{gl}(N)$ -weight systems, and M. Kazarian suggested a recurrence relation for such a computation, which works for permutations rather than just for chord diagrams.

A natural question then arises, namely, which already known weight systems can be obtained from the universal  $\mathfrak{gl}$  weight system. In addition to understanding the internal relationship between weight systems, knowing that a given weight system can be induced from the  $\mathfrak{gl}$ -weight system would immediately lead to extending the former to arbitrary permutations.

To each chord diagram, one can associate a graph, called the intersection graph of the chord diagram. Certain weight systems are completely determined by the intersection graphs; this is true, for example, for the  $\mathfrak{sl}(2)$ - and for the  $\mathfrak{gl}(1|1)$ -weight systems [3]. This is true no longer for more complicated Lie algebras, say, for  $\mathfrak{sl}(3)$ . In general, the relationship between Lie algebra weight systems and polynomial graph invariants looks rather mysterious.

In our recent paper [11], we suggested a substitution that makes the universal  $\mathfrak{gl}$  weight system into the interlace polynomial. The present paper is devoted to investigation of the relationship between the universal  $\mathfrak{gl}$  weight system and the chromatic polynomial.

In this case, the situation is less straightforward. Namely, it is easy to show that there does not exist a substitution transforming the universal  $\mathfrak{gl}$  weight system into the chromatic polynomial. However, we suggest a substitution, from which the chromatic polynomial emerges as the *leading term* of the result, as  $N$ , the rank of the Lie algebras  $\mathfrak{gl}(N)$ , tends to infinity. This substitution provides an interesting function on permutations, which takes values in the ring of polynomials in two variables. We start to study properties of this function. In particular, we prove that the leading term in  $N$  coincides with the

---

<sup>1</sup>The authors are grateful to M. Karev for pointing out that the extension can be considered as a special case of a construction in [6].

$$f \left( \text{diagram 1} \right) - f \left( \text{diagram 2} \right) + f \left( \text{diagram 3} \right) - f \left( \text{diagram 4} \right) = 0$$

Figure 1: 4-term relation

chromatic polynomial of the intersection graph of the permutation not only for chord diagrams, but for a wider class of permutations, which we call positive permutations.

The existence of a relationship between the chromatic polynomial weight system and the series  $\mathfrak{gl}(N)$  of Lie algebras was established earlier in [15]. Our construction seems to be much simpler and more straightforward than the one in [15].

The results of the present paper lead to numerous new questions. For example, it would be interesting to know whether the chromatic polynomial can be naturally extended to hypermaps with arbitrarily many hypervertices (a permutation considered modulo cyclic shifts can be treated as a hypermap with a single vertex). Also, there are universal weight systems for permutations constructed through other series of Lie algebras and Lie superalgebras [10]; their specializations also can lead to already known, as well as new, weight systems and their extensions to permutations.

### 1.1 4-term relations and weight systems

In V. Vassiliev's theory [17] of finite type knot invariants, a function on chord diagrams with  $n$  chords is associated to every knot invariant of order not greater than  $n$ . He showed that any function constructed in this way satisfies so-called 4-term relations. Then in [12], M. Kontsevich showed that every function on chord diagrams satisfying the four-term relations defines a knot invariant of finite type.

Below, we give necessary definitions and state the properties of the objects we require. A complete account can be found in [14, 2].

A *chord diagram* of order  $n$  is an oriented circle with  $2n$  points on it, which are split into  $n$  disjoint pairs, each pair connected by a chord, considered up to orientation preserving diffeomorphisms of the circle. A function  $f$  on chord diagrams is called a *weight system* if it satisfies the 4-term relation shown in Fig. 1. The vector space spanned by chord diagrams modulo 4-term relations is endowed with a natural multiplication: modulo 4-term relations, the concatenation of two chord diagrams is well defined.

### 1.2 Graphs, embedded graphs and hypermaps

Weight systems are closely related to invariants of graphs, embedded (= ribbon, topological) graphs and delta-matroids, see a detailed account in [9]. Here we recall briefly certain notions that will be used in the present paper.

A chord diagram can be treated as an orientable embedded graph (a combinatorial map) with a single vertex: the vertex is the outer circle, while the edges (ribbons) are obtained by thickening the chords.

Permutations considered up to standard cyclic shift can be treated as orientable hypermaps with a single vertex. A *hypermap* is a triple of permutations  $(\sigma, \alpha, \varphi)$  in  $\mathbb{S}_m$  whose product is the identity permutation,  $\sigma \circ \alpha \circ \varphi = \text{id}$ , considered up to simultaneous conjugation by one and the same permutation. A hypermap is a map if the permutation  $\alpha$  is an involution without fixed points. Cycles in the decomposition of  $\sigma$  (resp.,  $\alpha, \varphi$ ) into a product of disjoint cycles are *hypervertices* (resp., *hyperedges*, *hyperfaces*) of the hypermap.

Below, we will always assume that the first permutation  $\sigma$  is the standard cycle,  $\sigma = (1, 2, \dots, m)$ , and since  $\varphi = \alpha^{-1} \circ \sigma^{-1}$  is uniquely reconstructed from  $\alpha$ , we will speak about  $\alpha$  as about a hypermap with a single vertex.

The number  $f(\alpha)$  of (hyper)faces for a given permutation  $\alpha$  is the number of disjoint cycles in the permutation  $\varphi$ . If  $\alpha$  is an involution without fixed points (an arc diagram), then the number of its hyperfaces coincides with the number of ordinary faces.

To each hypermap, an orientable two-dimensional surface with boundary is naturally associated. The hyperfaces of the hypermap are in a natural one-to-one correspondence with the connected components of the boundary of the surface. For a single vertex, the surface is constructed by attaching the hyperedges (which are disks corresponding to the cycles in  $\alpha$ ) to the vertex, which is the disk, which corresponds to the unique cycle in  $\sigma$ . The rule of gluing is prescribed by  $\alpha$ .

### 1.3 $\mathfrak{gl}$ -weight system

The  $\mathfrak{gl}$ -weight system  $w_{\mathfrak{gl}}$  for permutations is defined in [18], see also [9]. This function on permutations takes values in the algebra  $\mathbb{C}[N][C_1, C_2, \dots]$  of polynomials in infinitely many variables  $C_1, C_2, \dots$  whose coefficients depend on the additional variable  $N = C_0$ . The variables  $C_i$  should be treated as the standard *Casimir elements*, which generate the center of the universal enveloping algebra  $U\mathfrak{gl}(\infty)$  of the limit Lie algebra  $\mathfrak{gl}(\infty)$ . The definition proceeds as follows.

The value of the  $\mathfrak{gl}(N)$ -weight system on a permutation  $\alpha \in \mathbb{S}_m$  is defined as the element

$$w_{\mathfrak{gl}(N)}(\alpha) = \sum_{i_1, \dots, i_m=1}^N E_{i_1, i_{\alpha(1)}} \cdots E_{i_m, i_{\alpha(m)}}$$

of the universal enveloping algebra  $U\mathfrak{gl}(N)$ , where  $E_{i,j} \in \mathfrak{gl}(N)$  are the matrix units,  $1 \leq i, j \leq N$ .

An arc diagram can be interpreted as an involution without fixed points on the set of the arc ends. The  $\mathfrak{gl}$ -weight system for permutations extends the  $\mathfrak{gl}(N)$ -weight system for chord diagrams to arbitrary permutations. This extension provides a way for recursive computation of the  $\mathfrak{gl}(N)$ -weight systems,

as well as unifies the latter weight systems in a universal one, containing all the specific weight systems, for a given  $N$ .

In order to introduce the recurrence, we will need the notion of the digraph of a permutation. A permutation can be represented as an oriented graph. The  $m$  vertices of the graph correspond to the permuted elements. They are placed on the horizontal line, and numbered from left to right in the increasing order. The arc arrows show the action of the permutation (so that each vertex is incident with exactly one incoming and one outgoing arc edge). The digraph  $G(\alpha)$  of a permutation  $\alpha \in \mathbb{S}_m$  consists of these  $m$  vertices and  $m$  oriented edges, for example:

$$G((1\ n+1)(2\ n+2)\cdots(n\ 2n)) = \begin{array}{c} \text{---} \bullet \text{---} \bullet \text{---} \cdots \text{---} \bullet \text{---} \bullet \text{---} \text{---} \\ \text{1} \quad \text{2} \quad \cdots \quad \text{n} \quad \text{n+1} \quad \text{n+2} \quad \cdots \quad \text{2n} \\ \text{---} \end{array}$$

Chord diagrams are permutations of special kind, namely, involutions without fixed points. For them, the initial definition coincides with the one above.

Define *Casimir elements*  $C_m \in U\mathfrak{gl}(N)$ ,  $m = 1, 2, \dots$ :

$$C_m = w_{\mathfrak{gl}(N)}((1, 2, \dots, m)) = \sum_{i_1, i_2, \dots, i_m=1}^N E_{i_1, i_2} E_{i_2, i_3} \cdots E_{i_m, i_1};$$

associated to the standard cycles  $1 \mapsto 2 \mapsto 3 \mapsto \cdots \mapsto m \mapsto 1$ .

**Theorem 1.1** *The center  $ZU\mathfrak{gl}(N)$  of the universal enveloping algebra  $U\mathfrak{gl}(N)$  of  $\mathfrak{gl}(N)$  is identified with the polynomial ring  $\mathbb{C}[C_1, \dots, C_N]$ .*

**Theorem 1.2 (Zhuoke Yang, [18])** *The  $w_{\mathfrak{gl}(N)}$  invariant of permutations possesses the following properties:*

- for the empty permutation, the value of  $w_{\mathfrak{gl}(N)}$  is equal to 1;
- $w_{\mathfrak{gl}(N)}$  is multiplicative with respect to concatenation of permutations;
- (**Recurrence Rule**) For the graph of an arbitrary permutation  $\alpha$  in  $\mathbb{S}_m$ , and for any two neighboring elements  $l, l+1$ , of the permuted set  $\{1, 2, \dots, m\}$ , we have for the values of the  $w_{\mathfrak{gl}(N)}$  weight system

$$\begin{array}{c} \text{---} \bullet \text{---} \bullet \text{---} \text{---} \\ \text{---} \end{array} = \begin{array}{c} \text{---} \bullet \text{---} \bullet \text{---} \text{---} \\ \text{---} \end{array} + \begin{array}{c} \text{---} \bullet \text{---} \bullet \text{---} \text{---} \\ \text{---} \end{array} - \begin{array}{c} \text{---} \bullet \text{---} \bullet \text{---} \text{---} \\ \text{---} \end{array} \quad (1)$$

For the special case  $\alpha(l+1) = l$ , the recurrence looks like follows:

$$\begin{array}{c} \text{---} \bullet \text{---} \bullet \text{---} \text{---} \\ \text{---} \end{array} = \begin{array}{c} \text{---} \bullet \text{---} \bullet \text{---} \text{---} \\ \text{---} \end{array} + C_1 \times \begin{array}{c} \text{---} \bullet \text{---} \bullet \text{---} \text{---} \\ \text{---} \end{array} - N \times \begin{array}{c} \text{---} \bullet \text{---} \bullet \text{---} \text{---} \\ \text{---} \end{array} \quad (2)$$

Below, we will write the four terms in the recurrence relation for a permutation as  $\alpha, \alpha', \beta_1, \beta_2$ , so that the (first) relation looks like

$$w_{\mathfrak{gl}}(\alpha) = w_{\mathfrak{gl}}(\alpha') + w_{\mathfrak{gl}}(\beta_1) - w_{\mathfrak{gl}}(\beta_2).$$

We do not specify the place where the two consecutive elements are exchanged, but we will do this in each specific case.

**Example 1.3** Let us compute the value of  $w_{\mathfrak{gl}}$  on the cyclic permutation  $(1\ 3\ 2)$  by switching the places of the nodes 2 and 3:

$$\begin{array}{c} \begin{array}{c} \text{---} \text{---} \text{---} \\ \text{1} \quad \text{2} \quad \text{3} \\ \text{(1 3 2)} \end{array} = \begin{array}{c} \text{---} \text{---} \text{---} \\ \text{1} \quad \text{2} \quad \text{3} \\ \text{(1 2 3)} \end{array} + C_1 \times \begin{array}{c} \text{---} \text{---} \text{---} \\ \text{1} \\ \text{(1)} \end{array} - N \times \begin{array}{c} \text{---} \text{---} \text{---} \\ \text{1} \quad \text{2} \\ \text{(1 2)} \end{array} \end{array}$$

$$\begin{aligned} w_{\mathfrak{gl}(N)}((1\ 3\ 2)) &= w_{\mathfrak{gl}(N)}((1\ 2\ 3)) + C_1 \cdot w_{\mathfrak{gl}(N)}((1)) - N \cdot w_{\mathfrak{gl}(N)}((1\ 2)) \\ &= C_3 + C_1^2 - NC_2 \end{aligned}$$

**Corollary 1.4** *The  $\mathfrak{gl}(N)$ -weight systems, for  $N = 1, 2, \dots$ , are combined into a universal  $\mathfrak{gl}$ -weight system taking values in the ring of polynomials in infinitely many variables  $\mathbb{C}[N; C_1, C_2, \dots]$ .*

*After substituting a given value of  $N$  and an expression of higher Casimirs  $C_{N+1}, C_{N+2}, \dots$  in terms of the lower ones  $C_1, C_2, \dots, C_N$ , this weight system specifies into the  $\mathfrak{gl}(N)$ -weight system.*

We say that a function on permutations is of  $\mathfrak{gl}$ -type if it is invariant with respect to the standard shift and satisfies the same recurrence relation as the  $\mathfrak{gl}$ -weight system.


**Example 1.5** *The number  $f(\alpha)$  of faces of a permutation is a function of  $\mathfrak{gl}$ -type. In fact, it is easy to see that the number of faces is preserved by replacing the first permutation on the left hand side of the recurrence with the first one on the right hand side, as well as replacing the second permutation on the left with the second one on the right.*

## 2 Rotational Hopf algebras of chord diagrams and permutations

In this section, we introduce three Hopf algebras, the rotational Hopf algebra of chord diagrams, the rotational Hopf algebra of permutations, and the rotational Hopf algebra of positive permutations. They have similar structural properties, and form a chain of graded Hopf algebras with respect to inclusion. Our further goal will be to associate to the  $\mathfrak{gl}$ -weight system a Hopf algebra homomorphism

of each of these three Hopf algebras to the Hopf algebra of polynomials in infinitely many variables.

We start with the rotational Hopf algebra of permutations. By the *concatenation product of two permutations*  $\alpha_1 \in \mathbb{S}_{m_1}$  and  $\alpha_2 \in \mathbb{S}_{m_2}$  we mean the permutation  $\alpha_1 \# \alpha_2 \in \mathbb{S}_{m_1+m_2}$  whose directed graph is obtained by attaching the base line of the directed graph of  $\alpha_2$  to the right of that of the directed graph of  $\alpha_1$ , followed by the shift of the numbering of permuted elements:



Let  $\mathcal{A}_m$  be the vector space (for definiteness, all the vector spaces we consider are over the ground field  $\mathbb{C}$  of complex numbers) freely spanned by equivalence classes of permutations of  $m$  elements modulo the following equivalence relation:

- any cyclic shift of a permutation is equivalent to it,  $\alpha \sim \sigma^{-1}\alpha\sigma$ ;
- the concatenation product of any two permutations is equivalent to the concatenation product of any pair of permutations cyclically equivalent to the factors.

Denote the equivalence class of a permutation  $\alpha$  by  $[\alpha]$ . Let

$$\mathcal{A} = \mathcal{A}_0 \oplus \mathcal{A}_1 \oplus \mathcal{A}_2 \oplus \dots$$

be the infinite dimensional vector space, which is the direct sum of the finite dimensional vector spaces  $\mathcal{A}_m$ ,  $m = 0, 1, 2, \dots$ , the vector space  $\mathcal{A}_m$  being freely spanned by the equivalence classes of permutations of  $m$  elements. The concatenation product of permutations makes  $\mathcal{A}$  into an infinite dimensional commutative graded algebra. We introduce the comultiplication  $\mu : \mathcal{A} \rightarrow \mathcal{A} \otimes \mathcal{A}$  in the following way: for a generator  $[\alpha] \in \mathcal{A}_m$  take a permutation  $\alpha \in \mathbb{S}_m$  representing this generator and set

$$\mu([\alpha]) = [\mu(\alpha)] = \sum_{I \sqcup J = V(\alpha)} [\alpha|_I] \otimes [\alpha|_J],$$

where the summation is carried over all the ways to represent the set  $V(\alpha)$  of disjoint cycles of  $\alpha$  into a disjoint union of two subsets  $I, J$ , and  $\alpha|_I$  denotes the restriction of  $\alpha$  to the subset of elements contained in the cycles from  $I$ . Here, following [11], we define the *restriction*  $\alpha|_U \in \mathbb{S}_{|U|}$  of a permutation  $\alpha \in \mathbb{S}_m$  to a subset  $U \subset \{1, 2, \dots, m\}$  as a permutation of the elements of  $U$  preserving their relative cyclic order inside each disjoint cycle in  $\alpha$ , composed with the subsequent renumbering to  $\{1, 2, \dots, |U|\}$ .

The standard argument shows that the vector space  $\mathcal{A}$  endowed with the multiplication  $\#$  and the comultiplication  $\mu$  is a graded connected commutative cocommutative Hopf algebra. We call it the *rotational Hopf algebra of permutations*.



**Definition 2.1** An *ascent* of a permutation  $\alpha \in \mathbb{S}_m$  is a number  $i$ ,  $0 < i < m$  such that  $\alpha(i) > i$ . We denote the number of ascents in  $\alpha$  by  $a(\alpha)$ .

A permutation  $\alpha \in \mathbb{S}_m$  is *positive* if  $a(\alpha) = m - c(\alpha)$  where  $c(\alpha)$  is the number of cycles in  $\alpha$ .

Obviously, the number of ascents of a permutation is preserved under its cyclic shifts, whence is well-defined for the equivalence class of a permutation.

The standard cycle  $\sigma = (1, 2, 3, \dots, m)$  is the only positive permutation consisting of a single cycle. Its number of ascents is  $a(\sigma) = m - 1$ . Its inverse  $\sigma^{-1} = (m, m - 1, m - 2, \dots, 1)$  has the minimal possible number of ascents 1. The numbers of ascents of other cyclic permutations are strictly between these two extremal values.

A permutation is positive if and only if its restriction to each disjoint cycle is a standard cycle. Positive permutations in  $\mathbb{S}_m$  are in one-to-one correspondence with partitions of the set  $\{1, 2, \dots, m\}$ : each part of the partition specifies a single positive disjoint cycle. In particular, any arc diagram, and, more generally, any involution is a positive permutation. Remark that a cyclic shift of a positive permutation also is a positive permutation.

Equivalence classes of positive permutations generate a graded Hopf subalgebra  $\mathcal{A}^+ \subset \mathcal{A}$ ,

$$\mathcal{A}^+ = \mathcal{A}_0^+ \oplus \mathcal{A}_1^+ \oplus \mathcal{A}_2^+ \oplus \dots,$$

$\mathcal{A}_m^+ \subset \mathcal{A}_m$ ,  $m = 0, 1, 2, \dots$ , which we call the *rotational Hopf algebra of positive permutations*.

By definition, the Hopf subalgebra

$$\mathcal{B} = \mathcal{B}_0 \oplus \mathcal{B}_1 \oplus \mathcal{B}_2 \oplus \dots,$$

where  $\mathcal{B}_m \subset \mathcal{A}_m^+ \subset \mathcal{A}_m$ ,  $m = 0, 1, 2, \dots$  are the vector subspaces spanned by equivalence classes of involutions without fixed points, is the *rotational Hopf algebra of chord diagrams*. The definition means, in particular, that all the vector spaces  $\mathcal{B}_m$  with odd indices  $m$  are zero.

Note that the rotational Hopf algebra of chord diagrams  $\mathcal{B}$  is different from the conventional Hopf algebra of chord diagrams [12, 2], since, in contrast to the usual case, we do not impose the equivalence modulo 4-term relations. The latter can also be extended to arbitrary permutations, which leads to a corresponding new Hopf algebra structure, but we do not require it in the present paper.

An equivalence class of a permutation  $[\alpha]$  is said to be *connected* if it cannot be represented as the concatenation product of two equivalence classes of smaller degree. Each of the Hopf algebras  $\mathcal{A}$ ,  $\mathcal{A}^+$ , and  $\mathcal{B}$  is generated by connected equivalence classes, and there are no further linear relations. This means that the dimension of the vector space  $P(\mathcal{A}_m)$  of primitive elements of degree  $m$  is equal to the number of connected equivalence classes of permutations in  $\mathbb{S}_m$ .

Two more Hopf algebras we will need are the Hopf algebra of polynomials in infinitely many variables  $\mathbb{C}[p_1, p_2, p_3, \dots]$  and the Hopf algebra of graphs  $\mathcal{G}$ . The comultiplication in the Hopf algebra of polynomials is defined on the generators by  $\mu : p_m \mapsto 1 \otimes p_m + p_m \otimes 1$ ,  $m = 1, 2, 3, \dots$ . Since the Hopf algebra

homomorphisms we are going to construct will be just filtered, not graded, we define a filtration rather than grading on this Hopf algebra by setting the degree of a variable  $p_m$  equal to  $m$ ,  $m = 1, 2, 3, \dots$ .

The Hopf algebra of graphs  $\mathcal{G}$  is defined as an infinite direct sum

$$\mathcal{G} = \mathcal{G}_0 \oplus \mathcal{G}_1 \oplus \mathcal{G}_2 \oplus \dots,$$

where the vector space  $\mathcal{G}_m$ ,  $m = 0, 1, 2, \dots$ , is freely spanned by isomorphism classes of graphs on  $m$  vertices. The multiplication in it is induced by the disjoint union, and the comultiplication  $\mu : \mathcal{G} \rightarrow \mathcal{G} \otimes \mathcal{G}$  is defined on the generators by

$$\mu : G \rightarrow \sum_{I \sqcup J = V(G)} G|_I \otimes G|_J,$$

where the summation is carried over all (ordered) partitions of the set  $V(G)$  of vertices of  $G$  into two disjoint subsets and  $G|_I$  denotes the subgraph of  $G$  induced by the subset  $I \subset V(G)$  of its vertices. Once again, we treat the Hopf algebra of graphs  $\mathcal{G}$  as filtered by the subspaces

$$\mathcal{G}_0 \subset \mathcal{G}_0 \oplus \mathcal{G}_1 \subset \mathcal{G}_0 \oplus \mathcal{G}_1 \oplus \mathcal{G}_2 \subset \dots$$

rather than a graded one.

The notion of intersection graph of a chord diagram [4] admits a natural extension to arbitrary permutations. Namely, for a permutation  $\alpha$  of the set  $\{1, 2, \dots, m\}$ , we say that two its disjoint cycles *interlace* if there are two elements in the first cycle and two elements in the second one following the base line in the alternating order. The *intersection graph*  $\gamma(\alpha)$  of a permutation  $\alpha$  is the abstract graph whose vertices are in one-to-one correspondence with the disjoint cycles of  $\alpha$ , and two vertices are connected by an edge iff the corresponding cycles interlace. Equivalent permutations have isomorphic intersection graphs. An equivalence class is connected if and only if its intersection graph is connected.

The mapping  $[\alpha] \rightarrow \gamma(\alpha)$  taking an equivalence class of permutations to its intersection graph extends by linearity to a Hopf algebra homomorphism  $\gamma : \mathcal{A} \rightarrow \mathcal{G}$  from the rotational Hopf algebra of permutations to the Hopf algebra of graphs. Note, however, that this homomorphism is filtered rather than graded: the grading in the Hopf algebra of graphs is given by the number of vertices, which corresponds to the number of cycles in a permutation rather than the number of permuted elements.

Since all the Hopf algebras we consider are generated by their primitive elements, their algebra homomorphism is a homomorphism of Hopf algebras iff it takes any primitive element in the source to a primitive element in the target.

### 3 Chromatic substitution

Our goal in the present section is to define a filtered Hopf algebra homomorphism from the rotational Hopf algebra of permutations  $\mathcal{A}$  to the Hopf algebra of

polynomials in infinitely many variables. This homomorphism, when specialized to a single variable and restricted to the rotational Hopf algebras of positive permutations  $\mathcal{A}^+$  and of chord diagrams  $\mathcal{B}$ , provides the chromatic polynomial of the intersection graph of a positive permutation or a chord diagram, respectively.

For a permutation  $\alpha$ , we set

$$X(\alpha) = N^{c(\alpha)-m} w_{\text{gl}}(\alpha) |_{C_k := p_k N^{k-1}, k=1,2,\dots}.$$

The value  $X(\alpha)$  is a Laurent polynomial in  $N$  and a polynomial in the variables  $p_1, p_2, \dots$ , which we call the *prechromatic invariant*.

The recursion for  $w_{\text{gl}}$ , when rewritten in terms of  $X$ , acquires the following form:

- the invariant  $X$  takes equal values on all permutations in one equivalence class;
- it is multiplicative with respect to concatenation product of permutations;
- the value of  $X$  on the standard (positive)  $m$ -cycle is equal to  $p_m$ ;
- if  $\ell$  and  $\ell + 1$  belong to different cycles of  $\alpha$ , we have

$$X \left( \begin{array}{c} \text{Diagram 1} \\ \hline \ell \quad \ell+1 \end{array} \right) = X \left( \begin{array}{c} \text{Diagram 2} \\ \hline \bullet \quad \bullet \end{array} \right) + X \left( \begin{array}{c} \text{Diagram 3} \\ \hline \bullet \end{array} \right) - X \left( \begin{array}{c} \text{Diagram 4} \\ \hline \bullet \end{array} \right) \quad (3)$$

- if  $\ell$  and  $\ell + 1$  belong to one cycle of  $\alpha$  and  $\ell + 1 \neq \alpha^{\pm 1}(\ell)$ , we have

$$X \left( \begin{array}{c} \text{Diagram 1} \\ \hline k \quad k+1 \end{array} \right) = X \left( \begin{array}{c} \text{Diagram 2} \\ \hline \bullet \quad \bullet \end{array} \right) + N^{-2} \cdot \left( X \left( \begin{array}{c} \text{Diagram 3} \\ \hline \bullet \end{array} \right) - X \left( \begin{array}{c} \text{Diagram 4} \\ \hline \bullet \end{array} \right) \right) \quad (4)$$

- in the case  $\alpha(\ell + 1) = \ell$ , we have

$$X \left( \begin{array}{c} \text{Diagram 1} \\ \hline k \quad k+1 \end{array} \right) = X \left( \begin{array}{c} \text{Diagram 2} \\ \hline \bullet \quad \bullet \end{array} \right) + p_1 N^{-2} \cdot X \left( \begin{array}{c} \text{Diagram 3} \\ \hline \bullet \end{array} \right) - X \left( \begin{array}{c} \text{Diagram 4} \\ \hline \bullet \end{array} \right) \quad (5)$$

As an immediate corollary of these relations we obtain

**Lemma 3.1** *The prechromatic invariant takes values in the ring of polynomials in  $N^{-2}, p_1, p_2, \dots$ . In other words, its value on any permutation  $\alpha$  can be written as a finite sum*

$$X(\alpha) = X_0(\alpha) + N^{-2} X_1(\alpha) + N^{-4} X_2(\alpha) + \dots$$

where each  $X_k(\alpha)$  is a polynomial in the variables  $p_1, p_2, \dots$

We are mostly interested in the leading term  $X_0$  of this expansion. The following assertion is the main result of the present paper.

**Theorem 3.2** *The value  $X_0(\alpha)$  on a permutation  $\alpha \in \mathbb{S}_m$  extends by linearity to a filtered Hopf algebra homomorphism  $X_0 : \mathcal{A} \rightarrow \mathbb{C}[p_1, p_2, \dots]$ .*

**Corollary 3.3** *The restriction of  $X_0$  to the rotational Hopf algebras of positive permutations  $\mathcal{A}^+ \subset \mathcal{A}$  and of chord diagrams  $\mathcal{B} \subset \mathcal{A}$  is a filtered Hopf algebra homomorphism. Under the substitution  $p_m := x$ ,  $m = 1, 2, \dots$  its value on an equivalence class of a positive permutation or a chord diagram becomes the chromatic polynomial of the corresponding intersection graph.*

We call the substitution  $p_m := x$  to  $X$  the *chromatic substitution*.

**Example 3.4** *The positive permutation  $(1, 3, 5, 8, 10)(2, 4, 7)(6, 9, 11)$  consists of three disjoint cycles. Any two of these cycles interlace, whence its intersection graph is the complete graph  $K_3$ . The chromatic substitution on this permutation is*

$$\begin{aligned} & X((1, 3, 5, 8, 10)(2, 4, 7)(6, 9, 11))(N; x, x, \dots) \\ &= x(x-1)(x-2) \\ &+ x(x-1)(3x-5)N^{-2} \\ &+ 2x(x-1)(3x-2)N^{-4} \\ &+ x(x-1)(5x-1)N^{-6} \\ &+ x^2(2x-1)N^{-8}. \end{aligned}$$

The coefficient of the leading term in  $N$  is the chromatic polynomial  $\chi_{K_3}(x)$ .

**Example 3.5** *For the value of the  $\mathfrak{gl}$ -weight system on the chord diagram of order 5 whose intersection graph is the complete graph  $K_5$ , we have*

$$\begin{aligned} w_{\mathfrak{gl}}(K_5) &= 24C_2N^4 + (24C_3 - 50C_2^2 - 24C_1^2)N^3 \\ &- (24C_4 + 10C_2C_3 - 35C_2^3 - 70C_1^2C_2 + 72C_1C_2 - 32C_2)N^2 \\ &+ (10C_2C_4 + 96C_1C_3 - 10C_2^4 - 50C_1^2C_2^2 + 30C_1C_2^2 - 82C_2^2 - 20C_1^4 + 48C_1^3 - 32C_1^2)N \\ &- 40C_1C_2C_3 + C_2^5 + 10C_1^2C_2^3 + 30C_2^3 + 15C_1^4C_2 - 20C_1^3C_2 + 10C_1^2C_2, \end{aligned}$$

which, under the chromatic substitution, becomes

$$\begin{aligned} X(K_5)(N; x, x, \dots) &= x(x-1)(x-2)(x-3)(x-4) \\ &+ 2x(x-1)(5x^3 - 20x^2 + 25x - 16)N^{-2} \\ &+ x^2(15x^3 - 40x^2 + 58x - 32)N^{-4}. \end{aligned}$$

It would be interesting to know the combinatorial meaning of the coefficients of the chromatic substitution at negative powers of  $N$ . In particular, whether they depend on the intersection graph of a chord diagram (or a positive permutation) only.

**Theorem 3.6** *The value of the chromatic substitution on a nonpositive permutation  $\alpha \in \mathbb{S}_m$  is a polynomial in  $N^{-2}$  with zero free term.*

**Example 3.7** *For the inverse standard cycle (132), the chromatic substitution to the value of the  $\mathfrak{gl}$ -weight system yields*

$$X_{\lrcorner}(132)(N; x, x, \dots) = N^{-2}(C_3 + C_1^2 - NC_2)|_{C_k := xN^{k-1}} = x^2N^{-2}.$$

*Its free term is 0.*

The following further specializations of the chromatic substitution are useful. It is shown in [11] that for the substitution  $C_k := N^{k-1}$ ,  $k = 1, 2, \dots$ , the universal  $\mathfrak{gl}$ -weight system takes a permutation  $\alpha$  to  $N^{f(\alpha)-1}$ , where  $f(\alpha)$  is the number of faces of  $\alpha$ ,  $w_{\mathfrak{gl}}(\alpha)|_{C_k := N^{k-1}} = N^{f(\alpha)-1}$ .

**Lemma 3.8** *We have the following specializations of the universal  $\mathfrak{gl}$ -weight system:*

- *for the substitution  $C_k := N^{k+1}$ ,  $k = 1, 2, \dots$ , the universal  $\mathfrak{gl}$ -weight system takes a permutation  $\alpha \in \mathbb{S}_m$  to the monomial  $N^{m+c(\alpha)}$ ,  $w_{\mathfrak{gl}}(\alpha)|_{C_k := N^{k+1}} = N^{m+c(\alpha)}$ ;*
- *for the substitution  $N := 1, C_k := x$ ,  $k = 1, 2, \dots$ , the universal  $\mathfrak{gl}$ -weight system takes a permutation  $\alpha \in \mathbb{S}_m$  to the monomial  $x^{c(\alpha)}$ ,  $w_{\mathfrak{gl}}(\alpha)|_{N:=1, C_k:=x} = x^{c(\alpha)}$ .*

*Recall that  $c(\alpha)$  denotes the number of cycles in  $\alpha$ .*

Indeed, the first assertion of the lemma is true for the standard cycles. Moreover, the function  $\alpha \mapsto N^{m+c(\alpha)}$  is a function of  $\mathfrak{gl}$ -type, that is, it satisfies the  $\mathfrak{gl}$  recurrence relation (1): the value  $m + c(\alpha)$  is the same for both  $\alpha$  and  $\alpha'$ , as well as for both  $\beta_1$  and  $\beta_2$ .

For the second assertion, the argument is the same.

## 4 Proof of the main results

In this section we give a proof of the main results of the paper. We start with defining a linear order on the set of equivalence classes of permutations in  $\mathbb{S}_m$ , that is, on the set of basic vectors in the spaces  $\mathcal{A}_m$ . Then we argue by a double induction with respect to  $m$  and this linear ordering.

### 4.1 A linear ordering on the set of equivalence classes of permutations

Associate to the equivalence class  $[\alpha]$  of a permutation  $\alpha$  its canonical form in the following way. Write each permutation in the equivalence class as a product of disjoint cycles, where for each cycle we chose the lexicographically

minimal presentation (the one with the smallest first element), and write the cycles in the lexicographic order (that is, the order in which their first elements increase). Now, among all the permutations in the equivalence class, choose the permutation with the lexicographically smallest canonical form. This will be the *canonical form*  $\bar{\alpha}$  of  $[\alpha]$ . The lexicographic ordering of canonical forms defines a linear ordering on the basic vectors in  $A_m$ .

## 4.2 Proof of Theorem 3.2

We argue by induction over  $m$ , the number of permuted elements, and then by the linear ordering of the basic elements of  $\mathcal{A}_m$  defined above.

Suppose we have proved already the assertion for all permutations of less than  $m$  elements, so that we know that for each permutation  $\alpha$  of at most  $m-1$  elements the leading term  $X_0(\alpha)$  in  $N$  is a linear polynomial in  $p_1, p_2, \dots, p_i$ .

The comultiplicativity of the mapping  $X_0 : \mathcal{A} \rightarrow \mathbb{C}[p_1, p_2, \dots]$  can be reformulated in the following way. Consider the mapping

$$(X_0 \otimes X_0) \circ \mu : \mathcal{A} \rightarrow \mathbb{C}[p_1, p_2, \dots] \otimes \mathbb{C}[p_1, p_2, \dots],$$

which we denote by  $\Xi$ . For a given permutation  $\alpha$ , it looks like

$$\Xi : \alpha \mapsto \sum_{I \sqcup J = V(\alpha)} X_0(\alpha|_I) \otimes X_0(\alpha|_J),$$

where the summation is carried over all ordered partitions of the set  $V(\alpha)$  of disjoint cycles of  $\alpha$  into two disjoint subsets.

**Lemma 4.1** *The mapping  $\Xi$  behaves as follows with respect to the recurrence relations applied to a permutation  $\alpha$ :*

- under (3), with  $X_0(\alpha) = X_0(\alpha') + X_0(\beta_1) - X_0(\beta_2)$ , we have

$$\Xi(\alpha) = \Xi(\alpha') + \Xi(\beta_1) - \Xi(\beta_2);$$

- under (4), with  $X_0(\alpha) = X_0(\alpha') + N^{-2}(X_0(\beta_1) - X_0(\beta_2))$ , we have

$$\Xi(\alpha) = \Xi(\alpha');$$

- under (5), with  $X_0(\alpha) = X_0(\alpha') + N^{-2}X_0(\beta_1) - X_0(\beta_2)$ , we have

$$\Xi(\alpha) = \Xi(\alpha') - \Xi(\beta_2).$$

Before proving Lemma 4.1 let us deduce Theorem 3.2 from it. The comultiplicativity of  $X_0$  is equivalent to the identity

$$\Xi(\alpha) = X_0(\alpha)|_{p_k := p_k \otimes 1 + 1 \otimes p_k}.$$

Since  $X_0(C_k) = p_k$ ,  $k = 1, 2, 3, \dots$ , the comultiplicativity property is equivalent to the assertion that the image of the mapping

$$\Xi : \mathcal{A} \rightarrow \mathbb{C}[p_1, p_2, \dots] \otimes \mathbb{C}[p_1, p_2, \dots]$$

coincides with the subring generated by the elements  $p_k \otimes 1 + 1 \otimes p_k$ .

Suppose  $m$  is the minimal number for which the assertion of Theorem 3.2 is not proved yet. Let  $[\alpha]$  be the lexicographically minimal basic vector in  $A_m$  possessing this property, and let  $\bar{\alpha}$  be the corresponding canonical form. Obviously,  $\bar{\alpha}$  is connected and contains at least 2 disjoint cycles. Take the minimal number  $k$ ,  $1 \leq k < m$  such that  $\bar{\alpha}(k) > k + 1$ . Applying the recurrence relation to the pair  $\bar{\alpha}(k) - 1, \bar{\alpha}(k)$  of permuted elements, we fall into one of the three cases of Lemma 4.1. Since in each of the cases each of the permutations  $\bar{\alpha}', \beta_1, \beta_2$  are smaller than  $\bar{\alpha}$ , the induction argument proves the Theorem.

**Proof of Lemma 4.1.**

- If the pair  $\ell - 1, \ell$  is like in recursion (3), so that  $\ell - 1$  and  $\ell$  belong to two different disjoint cycles of  $\alpha$ , then let  $v_1 \in V(\alpha)$  be the cycle containing  $\ell$ , and  $v_2 \in V(\alpha)$  be the cycle containing  $\ell - 1$ . The disjoint cycles  $V(\alpha')$  naturally correspond one-to-one to the disjoint cycles  $V(\alpha)$ , while each of the permutations  $\beta_1, \beta_2$  has one disjoint cycle less: the pair of cycles  $v_1, v_2$  is replaced by a single cycle. Now, we have

$$\begin{aligned} \Xi(\alpha) - \Xi(\alpha') - \Xi(\beta_1) + \Xi(\beta_2) = & \\ & \sum_{\substack{I \sqcup J = V(\alpha) \\ \{v_1, v_2\} \subset I}} X_0(\alpha|_I) \otimes (X_0(\alpha|_J) - X_0(\alpha'|_J) - X_0(\beta_1|_J) + X_0(\beta_2|_J)) \\ & \sum_{\substack{I \sqcup J = V(\alpha) \\ \{v_1, v_2\} \subset J}} (X_0(\alpha|_I) - X_0(\alpha'|_I) - X_0(\beta_1|_I) + X_0(\beta_2|_I)) \otimes X_0(\alpha|_J) \\ & \sum_{\substack{I \sqcup J = V(\alpha) \\ v_1 \in I, v_2 \in J \text{ or } v_2 \in I, v_1 \in J}} (X_0(\alpha|_I) \otimes X_0(\alpha|_J) - X_0(\alpha'|_I) \otimes X_0(\alpha'|_J)). \end{aligned}$$

Each of the three summands on the right is zero: for the first two sums the expression in brackets is 0 due to recurrence (3) applied to the permutations  $\alpha|_J$  and  $\alpha|_I$ , respectively, while for the third sum, it vanishes because  $\alpha|_I = \alpha'|_I, \alpha|_J = \alpha'|_J$ .

- If the pair  $\ell - 1, \ell$  is like in recurrence (4), so that  $\ell - 1$  and  $\ell$  belong to the same disjoint cycle of  $\alpha$ , then let  $v \in V(\alpha)$  be the cycle containing this pair of elements. The disjoint cycles  $V(\alpha')$  naturally correspond one-to-one to the disjoint cycles  $V(\alpha)$ . We claim that for an arbitrary subset  $I \subset V(\alpha)$  the values  $X_0(\alpha|_I)$  and  $X_0(\alpha'|_I)$  coincide. Indeed, if  $I$  does not contain  $v$ , then  $\alpha|_I = \alpha'|_I$ . If  $I$  contains  $v$ , then the two restrictions coincide no longer. However, in this case the permutations  $\alpha|_I$  and  $\alpha'|_I$  are related by the same recurrence (4), and the equality follows from the induction hypothesis.

- If the pair  $\ell - 1, \ell$  is like in recursion (5), so that  $\ell - 1$  and  $\ell$  are two successive elements in the same disjoint cycle, then let  $v \in V(\alpha)$  be this cycle. The disjoint cycles  $V(\alpha')$  and  $V(\beta_2)$  naturally correspond one-to-one to the disjoint cycles  $V(\alpha)$ , while the permutation  $\beta_1$  has one disjoint cycle more: the cycle  $v$  is replaced by two cycles. Thus, for any subset  $I \subset V(\alpha)$  not containing  $v$ , we have  $\alpha|_I = \alpha'|_I = \beta_2|_I$ . As a consequence,

$$\begin{aligned} \Xi(\alpha) - \Xi(\alpha') + \Xi(\beta_2) &= \\ &= \sum_{\substack{I \sqcup J = V(\alpha) \\ v \in I}} (X_0(\alpha|_I) - X_0(\alpha'|_I) + X_0(\beta_2|_I)) \otimes X_0(\alpha|_J) \\ &+ \sum_{\substack{I \sqcup J = V(\alpha) \\ v \in J}} X_0(\alpha|_J) \otimes (X_0(\alpha|_I) - X_0(\alpha'|_I) + X_0(\beta_2|_I)). \end{aligned}$$

Each of the two summands on the right vanishes: the expression in brackets is 0 due to recurrence (5) applied to the permutations  $\alpha|_J$  and  $\alpha|_I$ .

Lemma 4.1 is proved, which completes the proof of Theorem 3.2.

### 4.3 Values of $X_0$ on cyclic permutations

In this section we compute explicitly the value of  $X_0$  on cyclic permutations.

**Proposition 4.2** *If  $\alpha \in \mathbb{S}_m$  is a cyclic permutation, that is,  $c(\alpha) = 1$ , then the value  $X_\alpha$  of the prechromatic invariant on it has the form*

$$X(\alpha) = X_0(p_1, p_2, \dots) + X_1(p_1, p_2, \dots)N^{-2} + \dots,$$

where  $X_0$  is a linear polynomial in  $p_1, \dots, p_m$ ,  $X_1$  is at most a quadratic polynomial in  $p_1, \dots, p_{m-2}$ ,  $X_2$  is at most a cubic polynomial in  $p_1, \dots, p_{m-4}$ ,  $\dots$ . For a cyclic permutation  $\alpha \in \mathbb{S}_m$  having  $m - k$  ascents,  $a(\alpha) = m - k$ , the coefficient  $X_0$  is

$$X_0(\alpha) = p_m - \binom{k-1}{1} p_{m-1} + \binom{k-1}{2} p_{m-2} - \dots + (-1)^{k-1} \binom{k-1}{k-1} p_{m-k+1}.$$

**Remark 4.3** *In particular, for  $m > 1$  the coefficient of  $p_1$  in the linear polynomial  $L_1$  is 0, so that it depends only on  $p_2, \dots, p_m$ ; for  $k > 1$ , under the substitution  $p_k = x$  the polynomial  $X_0(\alpha)$  vanishes.*

Let  $[\alpha]$  be the minimal cyclic permutation for which the assertion of the Proposition is not proved yet. Take its canonical form  $\bar{\alpha} \in \mathbb{S}_m$  and let  $\ell$ ,  $1 \leq \ell < m - 1$  be the minimal element such that  $\bar{\alpha}(\ell) > \ell + 1$ . (If such an element  $\ell$  does not exist, then  $\bar{\alpha}$  is a standard cycle, for which the assertion of the Proposition is obvious.) There are two possibilities:

- either  $\bar{\alpha}(\bar{\alpha}(\ell)) = \bar{\alpha}(\ell) - 1$ ;
- or  $\bar{\alpha}(\bar{\alpha}(\ell)) \neq \bar{\alpha}(\ell) - 1$ .



In the first case, recurrence (5) is applicable to the points  $\bar{\alpha}(\ell) - 1, \bar{\alpha}(\ell)$ . The number of ascents in the permutation  $\bar{\alpha}'$ , the first permutation on the right, is one more than  $a(\bar{\alpha})$ , and, by the induction hypothesis, the polynomial  $X_0(\bar{\alpha}')$  for it is

$$p_m - \binom{m - a(\bar{\alpha}) - 2}{1} p_{m-1} + \binom{m - a(\bar{\alpha}) - 2}{2} p_{m-2} - \dots$$

The second permutation  $\beta_1$  on the right-hand side is a permutation of  $m - 2$  elements; by the induction hypothesis, after multiplication by  $p_1 N^{-2}$  the value of  $X$  on it does not contribute to the coefficient of  $N^0$ . The second element  $\beta_2$  is a cyclic permutation of  $m - 1$  elements, with  $a(\bar{\alpha})$  ascents. By the induction hypothesis, the value of  $X_0$  on it is

$$p_{m-1} - \binom{m - a(\bar{\alpha}) - 2}{1} p_{m-2} + \binom{m - a(\bar{\alpha}) - 2}{2} p_{m-3} - \dots$$

to the coefficient of  $N^{m-1}$ . After subtracting this value from the contribution of  $\bar{\alpha}'$ , we deduce that the value  $X_0(\bar{\alpha})$  is as required.

In the second case, apply recurrence (4) to the points  $\bar{\alpha}(\ell) - 1, \bar{\alpha}(\ell)$ . Then  $\bar{\alpha}'$ , the first term on the right, is a cyclic permutation of  $m$  elements whose number of ascents coincides with  $a(\bar{\alpha})$ . The two other permutations on the right  $\beta_1, \beta_2$  are permutations of  $m - 1$  elements, each consisting of two disjoint cycles. Therefore, the polynomial  $X$  on none of them contributes to free term in  $N$ , hence the assertion of the lemma is true for  $[\alpha]$  as well.

The property that  $X_k$  is a polynomial of degree at most  $k + 1$  in  $p_i$  also follows immediately from the induction argument above. The Proposition is proved.

#### 4.4 Proof of Corollary 3.3

The proof of the fact that the leading term in  $N$  of the chromatic substitution on positive permutations coincides with the chromatic polynomial of their intersection graphs splits into two steps. First, we need to prove that this leading term depends on the intersection graph only. We show this by proving that it satisfies the deletion-contraction relation. In addition, we prove that for a nonpositive permutation  $\alpha$ , the free term in the chromatic substitution is 0.

The proof proceeds by induction on the number  $m$  of permuted elements, and the ordering on the equivalence classes  $[\alpha]$  of permutations defined above. Suppose we have already proved Corollary 3.3 for all positive permutations of at most  $m - 1$  elements. For the cyclic positive permutation  $\sigma \in \mathbb{S}_m$ , the assertions are also valid.

Let  $[\alpha], \alpha \in \mathbb{S}_m$  be the minimal equivalence class of a positive permutation, for which Corollary 3.3 is not proved yet. It contains at least two disjoint cycles. Take its canonical representative  $\bar{\alpha}$ , let  $v_1$  be the disjoint cycle containing 1, and let  $\ell, 2 \leq \ell \leq m - 1$  be the minimal element of the cycle  $v_1$  such that  $\ell - 1$  does not belong to this cycle. Let us denote the cycle of  $\bar{\alpha}$  which contains  $\ell - 1$  by  $v_2$ .

Apply recurrence (1) to the pair of points  $\ell - 1, \ell$  and then apply the chromatic substitution to all its four terms. There are two possibilities:

- the intersection graph  $\gamma(\bar{\alpha})$  of the permutation  $\bar{\alpha}$  and the intersection graph  $\gamma(\bar{\alpha}')$  of the permutation  $\bar{\alpha}'$  are isomorphic to one another;
- these two intersection graphs are distinct.

In the second case, the intersection graph  $\gamma(\bar{\alpha}')$  is obtained from  $\gamma(\bar{\alpha})$  by either deleting or adding the edge connecting  $v_1$  and  $v_2$ .

Both permutations  $\beta_1, \beta_2$  on the right-hand side of recurrence (1) are permutations of  $m - 1$  elements containing  $c(\alpha) - 1$  disjoint cycles. If the intersection graphs  $\gamma(\bar{\alpha})$  and  $\gamma(\bar{\alpha}')$  are isomorphic, then, by the induction hypothesis, the contribution of the chromatic substitution to the coefficient of  $N^0$  on both  $\beta_1$  and  $\beta_2$  is 0. Indeed, each of these two permutations contains a nonpositive disjoint cycle, the one obtained by merging the cycles  $v_1, v_2$  of  $\bar{\alpha}$ .

Now suppose the graphs  $\gamma(\bar{\alpha})$  and  $\gamma(\bar{\alpha}')$  are not isomorphic. Then if  $v_1, v_2$  are connected in  $\gamma(\bar{\alpha})$  and not in  $\gamma(\bar{\alpha}')$ , then the permutation  $\beta_1$  is not positive, while  $\beta_2$  is. In this case the contribution of the chromatic substitution on  $\beta_1$  to the coefficient of  $N^0$  is 0, while the intersection graph  $\gamma(\beta_2)$  of the permutation  $\beta_2$  is the result of contracting the edge  $(v_1, v_2)$  in the intersection graph  $\gamma(\bar{\alpha})$ . The formula for the coefficient of  $N^0$  in the chromatic substitution for  $\bar{\alpha}$  then becomes the contraction-deletion relation for the chromatic polynomial, which proves that this coefficient is, indeed, the chromatic polynomial of  $\gamma(\bar{\alpha})$ . If  $v_1, v_2$  are not connected in  $\gamma(\bar{\alpha})$  and are connected in  $\gamma(\bar{\alpha}')$ , the argument is the same, with the exchange of the roles of the permutations  $\beta_1, \beta_2$ .

In the case the permutation  $\bar{\alpha}$  is not positive, all the three permutations  $\bar{\alpha}', \beta_1, \beta_2$  are not positive as well. Hence, by the induction hypothesis, none of the three permutations contributes to the coefficient of  $N^{m-c(\alpha)}$  in the chromatic substitution, whence the same is true for the permutation  $\bar{\alpha}$ . This completes the proof of Corollary 3.3.

It is worth to recall that the chromatic polynomial can be considered as a filtered Hopf algebra homomorphism  $\chi : \mathcal{G} \rightarrow \mathbb{C}[x]$ , see, e.g. [9]. In our current setting, it can be considered also as a filtered Hopf algebra homomorphism  $\chi : \mathcal{A}^+ \rightarrow \mathbb{C}[x]$  from the rotational Hopf algebra of positive permutations to the Hopf algebra of polynomials in a single variable.

## 5 Further results

In this section, we present several results about the chromatic substitution, which both clarify its nature and simplify its computation.

**Theorem 5.1** *If  $\alpha$  is a nonpositive long cycle of length  $m$ , then the degree in  $N$  of the polynomial  $X(\alpha)(N; x, x, \dots)$  is less than  $-1$ . In particular, for cyclic permutations with  $m - 2$  ascents, this polynomial has degree  $-2$  in  $N$ .*

**Remark 5.2** *The chromatic substitution on cyclic permutations with less than  $m-2$  ascents also can provide polynomials of degree  $-2$  in  $N$ . For example, the permutation  $(1, 2, 3, 6, 5, 4) \in \mathbb{S}_6$  has  $m-3=3$  ascents, while the chromatic substitution on it yields*

$$X((1, 2, 3, 6, 5, 4))(N; x, x, \dots) = x^2 N^{-2}.$$

**Example 5.3** *The long cycle  $\alpha = (1, 3, 5, 2, 4)$  is nonpositive; the number of faces in it is  $f(\alpha) = 1$ . The chromatic substitution on this permutation is*

$$X_\alpha(N; x, x, \dots) = x(x-1)N^{-2} + x^2 N^{-4}.$$

**Lemma 5.4** *The value of the chromatic substitution on a cyclic permutation in  $\mathbb{S}_m$  with  $m-2$  ascents is  $x^2 N^{m-3}$ .*

**Proof.** We proceed by induction on  $m$ . Suppose the assertion of the lemma is true for all cyclic permutations of up to  $m-1$  elements. Each cyclic permutation of positivity  $m-4$  can be obtained from the positive cyclic permutation by a composition of exchanging the pairs of elements  $(m-1, m)$ ,  $(m-2, m-1)$ ,  $\dots$ . Applying the recurrence rule for  $\mathfrak{gl}$  to the first of this exchanges and using the induction assumption, we conclude that the value of the chromatic substitution on the cyclic permutation  $(1, 2, \dots, m-2, m, m-1)$  indeed is  $x^2 N^{m-3}$ . Further exchanges preserve this value since for them the right hand side of the recurrence relation vanishes.  $\square$

**Theorem 5.5** *All the coefficients but one in  $X(\alpha)(N; x, x, \dots)$ , for an arbitrary permutation  $\alpha$  of at least one element, are divisible by  $x(x-1)$ . The only exception is the coefficient of  $N^{f(\alpha)-m}$ ; it is divisible by  $x$  and its value at  $x=1$  is 1.*

Indeed, we know that the composition of the chromatic substitution with  $x := 1$  makes its value on a permutation  $\alpha$  into  $N^{f(\alpha)-m}$ , while substitution  $x := 0$  makes it into 0 for all permutations but the empty one.

**Theorem 5.6** *Let  $\alpha \in \mathbb{S}_m$  be a permutation such that  $\alpha(1) = 2$ . Then*

$$X(\alpha)(N; x, x, \dots) = X(\alpha|_{\{2,3,\dots,m\}})(N; x, x, \dots).$$

Note that since permutations are considered up to cyclic shifts, a similar assertion is valid for any permutation  $\alpha \in \mathbb{S}_m$  such that  $\alpha(k) \equiv k+1 \pmod{m}$  for some  $k \in \{1, \dots, m\}$ . This means that if a permutation takes an element to the next one in the cyclic order, then we can replace this pair of consecutive elements by a single one, the value of the chromatic substitution being preserved. The proof repeats that for the main theorem for permutations.

Associate to a permutation  $\alpha$  the *weighted intersection graph*  $\gamma^w(\alpha)$ , which is a simple graph with  $\mathbb{N}$ -weighted vertices, in the following way: the underlying graph of  $\gamma^w(\alpha)$  coincides with  $\gamma(\alpha)$ , and the weight of a vertex is the length of the cycle in  $\alpha$ , which corresponds to this cycle.

## References

- [1] D. Bar-Natan, *On the Vassiliev knot invariants*, *Topology*, 1995, **34**, 423–472
- [2] S. Chmutov, S. Duzhin, and J. Mostovoy, *Introduction to Vassiliev Knot Invariants*, Cambridge University Press, 2012
- [3] S. Chmutov and S. Lando, *Mutant knots and intersection graphs*, *Algebraic & Geometric Topology*, 2007, **7**, 1579–1598
- [4] S.V.Chmutov, S.V.Duzhin, S.K.Lando *Vassiliev knot invariants I. Introduction*, *Advances in Soviet Mathematics*, vol. 21 (1994), pp. 117–126
- [5] S. Chmutov, M. Kazarian, and S. Lando, *Polynomial graph invariants and the KP hierarchy*, *Selecta Math. New Series* (2020) 26:34
- [6] S. Duzhin, M. Karev, *Detecting the orientation of string links by finite type invariants* *Functional Analysis and Its Applications*, 41(3), 208–216 (2007)
- [7] J. A. Ellis-Monaghan, I. Moffatt, *Graphs on surfaces: dualities, polynomials, and knots*, 2013, Springer
- [8] M. Kazarian, S. Lando, *Combinatorial solutions to integrable hierarchies*, *Russ. Math. Surveys*, **70** (3) 453–482 (2015).
- [9] M. Kazarian, S. Lando, *Weight systems and invariants of graphs and embedded graphs*, *Russ. Math. Surveys*, **77** (5) 893–942 (2022).
- [10] M.Kazarian, Zhuoke Yang, *Universal polynomial so weight system*, in preparation
- [11] N. Kodaneva, S. Lando *Polynomial graph invariants induced from the  $\mathfrak{gl}$ -weight system*, arXiv:2312.17519
- [12] M. Kontsevich. *Vassiliev knot invariants*, in: *Advances in Soviet Math.*, 16(2):137–150, 1993.
- [13] S. Lando, *On primitive elements in the bialgebra of chord diagrams*, *Translations of the American Mathematical Society-Series 2*, **180**, 167–174, 1997, Providence [etc.] American Mathematical Society, 1949-
- [14] S. Lando and A. Zvonkin, *Graphs on Surfaces and Their Applications*, 2003
- [15] J. Lieberum, *Chromatic weight systems and the corresponding knot invariants*, *Math. Ann.*, Volume 317, pages 459–482 (2000)
- [16] W. R. Schmitt, *Incidence Hopf algebras*, *Journal of Pure and Applied Algebra* **96** 299–330 (1994).
- [17] V. Vassiliev, *Cohomology of knot spaces* in: *Advance in Soviet Math.v. 1*, 1990, 23–69

- [18] Zhuoke Yang, *New approaches to  $\mathfrak{gl}(N)$  weight system*, Izvestiya Mathematics, 2023, vol. 77:6, 150–166; arXiv:2202.12225 (2022)
- [19] Zhuoke Yang, *On the Lie superalgebra  $\mathfrak{gl}(m|n)$  weight system*, Journal of Geometry and Physics Vol. 187 Article 104808 (2022)