Strong convergence rates for long-time approximations of SDEs with non-globally Lipschitz continuous coefficients [†]

Xiaoming Wu^{a,b}, Xiaojie Wang^b *

^a Department of Mathematics , Southern University of Science and Technology, Shenzhen, Guangdong, P. R. China
 ^b School of Mathematics and Statistics, HNP-LAMA, Central South University, Changsha, Hunan, P. R. China

June 18, 2024

Abstract

This paper is concerned with long-time strong approximations of SDEs with non-globally Lipschitz coefficients. Under certain non-globally Lipschitz conditions, a long-time version of fundamental strong convergence theorem is established for general one-step time discretization schemes. With the aid of the fundamental strong convergence theorem, we prove the expected strong convergence rate over infinite time for two types of schemes such as the backward Euler method and the projected Euler method in non-globally Lipschitz settings. Numerical examples are finally reported to confirm our findings.

Keywords: SDEs with non-globally Lipschitz coefficients, backward Euler method, projectd Euler method, strong convergence

1 Introduction

Stochastic differential equations (SDEs) find prominent applications in engineering, physics, chemistry, finance and many other branches of science. Nevertheless, nonlinear SDEs rarely have closed-form solution available and one often resorts to their numerical approximations. To numerically study SDEs, the globally Lipschitz conditions are imposed on the coefficient functions of SDEs [21,32]. Under the restrictive assumptions, fundamental mean-square convergence theorems were established for one-step approximation schemes of SDEs on the finite time horizon [32] and the infinite time horizon [22]. However, the coefficients of most nonlinear SDEs from applications violate the traditional but restrictive assumptions. What happens when the commonly used schemes in the global Lipschitz settings are applied to SDEs with non-globally Lipschitz coefficients? As asserted by [15], the most popular Euler-Maruyama method produces divergent numerical approximations when used to solve SDEs with super-linearly growing coefficients over finite interval [0, T]. In the literature, a large amount of attention has been attracted to construct

[†]This work was supported by Natural Science Foundation of China (12071488, 12371417, 11971488). The authors want to thank Prof. Zhihui Liu and Yajie She for helpful comments.

^{*}Corresponding author.

E-mail addresses: 12331004@mail.sustech.edu.cn,x.j.wang7@csu.edu.cn.

and analyze convergent numerical approximations of SDEs with super-linearly growing coefficients, see, e.g., [1-3,8-11,13,14,16,17,19,20,27-29,33,35,39-42]. Indeed, a majority of existing works are devoted to developing and analyzing schemes for strong approximations of SDEs with non-globally Lipschitz conditions over the finite time interval [0, T]. In the strong convergence analysis over the finite time interval [0, T]. In the strong convergence theorem due to [32] in the globally Lipschitz setting and its counterpart in a non-globally Lipschitz setting [39].

However, the long-time approximations of SDEs also plays an important role in many scientific areas such as high-dimensional sampling, Bayesian inference, statistical physics and machine learning [7, 12, 36, 43]. In the last decades, there are some important works devoted to long-time approximations of SDEs with globally Lipschitz continuous coefficients, see, e.g., [31, 37, 38] and references therein. More recently, some researchers examined long-time approximations of SDEs in non-globally Lipschitz setting [4–6, 8, 22–25, 30, 34], to just mention a few.

To the best of our knowledge, the fundamental strong convergence theorem for long-time approximations of SDEs with non-globally Lipschitz continuous coefficients is still absent in the literature. In this paper, we attempt to fill the gap by establishing a long-time version of fundamental strong convergence theorem for general one-step time discretization schemes applied to SDEs with non-globally Lipschitz coefficients (Theorem 2.5). As applications of the fundamental strong convergence theorem, we prove the expected strong convergence rate of order $\frac{1}{2}$ over infinite time for two types of time-stepping schemes such as the backward Euler method and the projected Euler method in non-globally Lipschitz settings (Theorem 3.9 and Theorem 4.7).

The outline of the paper is as follows. In the next section, we present some standard notations and assumptions that will be used in our proofs and obtain the long-time fundamental strong convergence theorem. In sections 3 and 4, we consider the long-time strong convergence analysis of two well-known methods, e.g., the backward Euler scheme and the projected Euler scheme in the non-globally Lipschitz settings. Numerical experiments are presented in section 5 to verify our theoretical findings.

2 The long-time fundamental strong convergence theorem

Notation: Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a complete probability space and $\{\mathcal{F}_t^w\}_{0 \le t < \infty}$ be an increasing family subalgebras induced by w(t) for $0 \le t < \infty$, where $(w(t), \mathcal{F}_t^w) = ((w_1(t), w_2(t), \cdots, w_m(t))^T, \mathcal{F}_t^w)$ is an *m*-dimensional Wiener process. Let $|\cdot|$ and $\langle \cdot, \cdot \rangle$ be the Euclidean norm and the inner product of vectors in \mathbb{R}^d , respectively. By M^T we denote the transpose of a vector or matrix M. For a given matrix M, we use $||M|| := \sqrt{trace(M^T M)}$ to denote the trace norm of M. On the probability space $(\Omega, \mathcal{F}, \mathbb{P})$, we use \mathbb{E} to denote the expectation and $L^q(\Omega; \mathbb{R}^{d \times m}), q \in \mathbb{N}$, to denote the family of $\mathbb{R}^{d \times m}$ -valued variables with the norm defined by $\|\xi\|_{L^q(\Omega; \mathbb{R}^{d \times m})} = (\mathbb{E}[\|\xi\|^q])^{\frac{1}{q}} < \infty$. And |x| denotes the integer part of x.

Consider the autonomous SDEs in the Itô form of

$$\begin{cases} dX(t) = f(X(t)) dt + g(X(t)) dW(t), & t \ge 0, \\ X(0) = X_0 \in \mathbb{R}^d, \end{cases}$$
(2.1)

where $f = (f^1, f^2, \dots, f^d)^T$ is the drift coefficient function and $g = (g^{ij})_{d \times m} = (g^1, g^2, \dots, g^m) = (g_1^T, g_2^T, \dots, g_d^T)^T$ is the diffusion coefficient function. It is worth noting that f^i is a real-valued

function, g^i is a d-column vector function and g_j is a m-row vector function. In addition, $\{W_t\}_{t\geq 0}$ is an *m*-dimensional Wiener process and the initial data X_0 is independent of w.

We aim to establish a fundamental strong convergence theorem for general one-step approximation schemes, used to approximate SDEs (2.1), with uniform step size $h = \frac{1}{N}, N \ge 1$. Here we introduce a new notation X(t, x; s) for $t \leq s < \infty$, which denotes the solution of (2.1) with the initial condition X(t, x; t) = x. When we write $X(t), t \ge 0$, we mean a solution of SDEs (2.1) with the initial value $X(0) = X_0$. In addition, we introduce one-step approximation Z(t, x; t+h)for X(t, x; t + h), where $t \ge 0, 0 < h < 1$, which is defined as follows:

$$Z(t, x; t+h) = x + \Psi(t, x, h; \xi_t),$$
(2.2)

where Ψ is a function from $[0,\infty) \times \mathbb{R}^d \times (0,T] \times \mathbb{R}^m$ to \mathbb{R}^m , ξ_t is a random variable with sufficiently high-order moments. Moreover, we define Z(t, x; t+h) as an approximation of the solution at t+hwith initial value Z(t, x; t) = x. Then, a numerical approximation $\{Z_k\}_{k>0}$ can be constructed on the uniform mesh grid $\{t_k = kh, k \ge 0\}$, given by

$$Z_0 = X_0, Z_{k+1} = Z_k + \Psi(t_k, Z_k, h; \xi_k), k \ge 0,$$
(2.3)

where ξ_k for $k \ge 0$ are independent of $Z_0, Z_1, \cdots, Z_{k-1}, \xi_0, \xi_1, \cdots, \xi_{k-1}$. Alternatively, one can write

$$Z_0 = X_0, Z_{k+1} = Z(t_k, Z_k; t_{k+1}) = Z(t_0, Z_0; t_{k+1}), k \ge 0.$$
(2.4)

To facilitate the strong analysis, we impose the following assumptions.

Assumption 2.1. Suppose that the drift and diffusion coefficients of SDEs given by (2.1) satisfy a contractive monotone condition. Additionally, there exists a non-integer constant $p^* \in [1,\infty)$ and $\alpha_1 \in (0, \infty)$ such that

$$\langle x - y, f(x) - f(y) \rangle + \frac{2p^* - 1}{2} \|g(x) - g(y)\|^2 \le -\alpha_1 |x - y|^2.$$
 (2.5)

Assume that f satisfies the polynomial growth Lipschitz condition, more accurately, there exist positive constants $\kappa \geq 1$ and $c_1 > 0$ such that

$$|f(x) - f(y)|^2 \le c_1 \left(1 + |x|^{2\kappa - 2} + |y|^{2\kappa - 2} \right) |x - y|^2.$$
(2.6)

In addition, assume the initial data X_0 satisfies

$$\mathbb{E}\big[|X_0|^{2p^*}\big] < \infty. \tag{2.7}$$

According to (2.5), for any $q \in (0, |p^*|]$, there exists a sufficient small coefficient $0 < \epsilon_1 < \alpha_1$ such that

$$\langle x, f(x) \rangle + \frac{2q-1}{2} \|g(x)\|^2 \le -(\alpha_1 - \epsilon_1)|x|^2 + c_0,$$
 (2.8)

where $c_0 = \frac{|f(0)|^2}{4\epsilon_1} + \frac{(2p^*-1)^2 ||g(0)||^2}{4(p^*-q)} - \frac{2p^*-1}{2} ||g(0)||^2$. Furthermore, (2.6) immediately implies

$$|f(x)|^2 \le c_2 |x|^{2\kappa} + c_3, \tag{2.9}$$

where $c_2 = \frac{2c_1(\kappa+1)}{\kappa}$ and $c_3 = 2|f(0)|^2 + \frac{2c_1(\kappa-1)}{\kappa}$. To show the 2*p*-th moment $(p \ge 1)$ boundedness of X(t), we first establish the following lemma, which is a slight modification of [18, Lemma 8.1].

Lemma 2.2. Let $r : [0, \infty) \to [0, \infty)$ and $\xi : [0, \infty) \to [0, \infty)$ be nonnegative continuous functions. If there exists a positive constant β such that

$$r(t) - r(s) \le -\beta \int_{s}^{t} r(u) \, du + \int_{s}^{t} \xi(u) \, du$$
 (2.10)

for any $0 \leq s < t < \infty$, then

$$r(t) \le r(0) + \int_0^t e^{-\beta(t-u)} \xi(u) \, du.$$
 (2.11)

Equipped with Assumption 2.1 and Lemma 2.2, we shall give the moment boundedness of the solution over infinite time, which is an interesting result independently.

Theorem 2.3. Suppose Assumption 2.1 hold. There exists a positive constant $C_1 := C_1(p, \alpha_1, \epsilon_1, c_0)$ independent of t, such that

$$\mathbb{E}[|X(t)|^{2p}] \le C_1 \mathbb{E}[(1+|X_0|^{2p})], \qquad 1 \le p < p^*.$$
(2.12)

Proof of Theorem 2.3. By the Itô formula, one sees

$$(1 + |X(t)|^2)^p = (1 + |X_0|^2)^p + 2p \int_0^t (1 + |X(s)|^2)^{p-1} \langle X(s), f(X(s)) \rangle ds + 2p \int_0^t (1 + |X(s)|^2)^{p-1} \langle X(s), g(X(s)) dW(s) \rangle + p \int_0^t (1 + |X(s)|^2)^{p-1} ||g(X(s))||^2 ds + 2p(p-1) \int_0^t (1 + |X(s)|^2)^{p-2} |X(s)^T g(X(s))|^2 ds,$$

$$(2.13)$$

which straightforwardly implies

$$(1 + |X(t)|^2)^p \le (1 + |X_0|^2)^p + 2p \int_0^t (1 + |X(s)|^2)^{p-1} \langle X(s), f(X(s)) \rangle \,\mathrm{d}s + 2p \int_0^t (1 + |X(s)|^2)^{p-1} \langle X(s), g(X(s)) \,\mathrm{d}W(s) \rangle + p(2p-1) \int_0^t (1 + |X(s)|^2)^{p-1} \|g(X(s))\|^2 \,\mathrm{d}s.$$
 (2.14)

According to (2.8), we have

$$(1+|X(t)|^2)^p \leq (1+|X_0|^2)^p - 2p(\alpha_1 - \epsilon_1) \int_0^t (1+|X(s)|^2)^p \,\mathrm{d}s + 2p(c_0 + \alpha_1 - \epsilon_1) \int_0^t (1+|X(s)|^2)^{p-1} \,\mathrm{d}s + 2p \int_0^t (1+|X(s)|^2)^{p-1} \langle X(s), g(X(s)) \,\mathrm{d}W(s) \rangle \,.$$
 (2.15)

Then by the Young inequality, we obtain

$$(1+|X(t)|^{2})^{p} \leq (1+|X_{0}|^{2})^{p} - p(\alpha_{1}-\epsilon_{1}) \int_{0}^{t} (1+|X(s)|^{2})^{p} ds + \int_{0}^{t} \frac{(2(c_{0}+\alpha_{1}-\epsilon_{1}))^{p}}{(\alpha_{1}-\epsilon_{1})^{p-1}} ds + 2p \int_{0}^{t} (1+|X(s)|^{2})^{p-1} \langle X(s), g(X(s)) dW(s) \rangle.$$
(2.16)

For every integer $n \ge 1$, define the stopping time:

$$\tau_n := \inf\{s \in [0, t] : |X(s)| \ge n\}.$$
(2.17)

Clearly, $\tau_n \uparrow t$ a.s.. Moreover, it follows from (2.16) and the property of the Itô integral that

$$\mathbb{E}\Big[\left(1+|X(t\wedge\tau_{n})|^{2}\right)^{p}\Big]+p(\alpha_{1}-\epsilon_{1})\mathbb{E}\Big[\int_{0}^{t\wedge\tau_{n}}\left(1+|X(s)|^{2}\right)^{p}\,\mathrm{d}s\Big] \\
\leq \mathbb{E}\Big[\left(1+|X_{0}|^{2}\right)^{p}\Big]+\int_{0}^{t}\frac{(2(c_{0}+\alpha_{1}-\epsilon_{1}))^{p}}{(\alpha_{1}-\epsilon_{1})^{p-1}}\,\mathrm{d}s.$$
(2.18)

Letting $n \to \infty$ and by the Fatou lemma, we get

$$\mathbb{E}\Big[\left(1+|X(t)|^{2}\right)^{p}\Big]+p(\alpha_{1}-\epsilon_{1})\int_{0}^{t}\mathbb{E}\Big[\left(1+|X(s)|^{2}\right)^{p}\Big]\,\mathrm{d}s \\
\leq \mathbb{E}\Big[\left(1+|X_{0}|^{2}\right)^{p}\Big]+\int_{0}^{t}\frac{(2(c_{0}+\alpha_{1}-\epsilon_{1}))^{p}}{(\alpha_{1}-\epsilon_{1})^{p-1}}\,\mathrm{d}s.$$
(2.19)

Taking $r(t) = \mathbb{E}\left[\left(1 + |X(t)|^2\right)^p\right]$, $\beta = p(\alpha_1 - \epsilon_1)$ and $\xi = \frac{(2(c_0 + \alpha_1 - \epsilon_1))^p}{(\alpha_1 - \epsilon_1)^{p-1}}$ in Lemma 2.2 shows

$$\mathbb{E}\Big[\left(1+|X(t)|^{2}\right)^{p}\Big] \leq \mathbb{E}\Big[\left(1+|X_{0}|^{2}\right)^{p}\Big] + \int_{0}^{t} e^{-p(\alpha_{1}-\epsilon_{1})(t-u)} \frac{(2(c_{0}+\alpha_{1}-\epsilon_{1}))^{p}}{(\alpha_{1}-\epsilon_{1})^{p-1}} du
\leq \mathbb{E}\Big[\left(1+|X_{0}|^{2}\right)^{p}\Big] + \frac{(2(c_{0}+\alpha_{1}-\epsilon_{1}))^{p}}{p(\alpha_{1}-\epsilon_{1})^{p}} (1-e^{-p(\alpha_{1}-\epsilon_{1})t})
\leq C_{1}\mathbb{E}\Big[\left(1+|X_{0}|^{2p}\right)\Big].$$
(2.20)

Thus the proof is finished. \Box

To establish a strong convergence theorem over an infinite time, we also rely on the following result.

Theorem 2.4. Let Assumption 2.1 hold. For the representation

$$X(t,x;t+\theta) - X(t,y;t+\theta) = x - y + \mathcal{R}_{t,x,y}(t+\theta), \qquad (2.21)$$

we have for $1 \leq p < \frac{p^*}{\kappa}$,

$$\mathbb{E}\left[|X(t,x;t+h) - X(t,y;t+h)|^{2p}\right] \le |x-y|^{2p} \exp(-2p\alpha_1 h),$$
(2.22)

$$\mathbb{E}\left[|\mathcal{R}_{t,x,y}(t+h)|^{2p}\right] \le C_2 \left(1+|x|^{2\kappa-2}+|y|^{2\kappa-2}\right)^{\frac{p}{2}}|x-y|^{2p}h^p, \tag{2.23}$$

where $C_2 > 0$ and do not depend on t, h.

Proof of Theorem 2.4. For $s \ge t$, we introduce the process

$$\mathcal{D}_{t,x,y}(s) := X(t,x;s) - X(t,y;s)$$

and note that

$$\mathcal{R}_{t,x,y}(s) := \mathcal{D}_{t,x,y}(s) - (x - y).$$

Clearly,

$$d\mathcal{D}_{t,x,y}(s) = \left(f(X(t,x;s)) - f(X(t,y;s))\right) ds + \left(g(X(t,x;s)) - g(X(t,y;s))\right) dW(s).$$
(2.24)

By applying the Itô formula, we obtain the following for any $t \le s \le t + h$,

$$e^{2\alpha_{1}p(s-t)}|\mathcal{D}_{t,x,y}(s)|^{2p} - |\mathcal{D}_{t,x,y}(t)|^{2p}$$

$$\leq \int_{t}^{s} 2\alpha_{1}pe^{2\alpha_{1}p(\theta-t)}|\mathcal{D}_{t,x,y}(\theta)|^{2p} d\theta$$

$$+ \int_{t}^{s} 2pe^{2\alpha_{1}p(\theta-t)}|\mathcal{D}_{t,x,y}(\theta)|^{2p-2} \langle \mathcal{D}_{t,x,y}(\theta), f(X(t,x;\theta)) - f(X(t,y;\theta)) \rangle d\theta \qquad (2.25)$$

$$+ \frac{2p(2p-1)}{2} \int_{t}^{s} e^{2\alpha_{1}p(\theta-t)}|\mathcal{D}_{t,x,y}(\theta)|^{2p-2} ||g(X(t,x;\theta)) - g(X(t,y;\theta))||^{2} d\theta$$

$$+ \int_{t}^{s} 2pe^{2\alpha_{1}p(\theta-t)}|\mathcal{D}_{t,x,y}(\theta)|^{2p-2} \langle \mathcal{D}_{t,x,y}(\theta), (g(X(t,x;\theta)) - g(X(t,y;\theta))) \rangle dW(\theta) \rangle.$$

Further, utilizing (2.5) and taking expectations, we have

$$\mathbb{E}\left[e^{2\alpha_1 p(s-t)} |\mathcal{D}_{t,x,y}(s)|^{2p}\right] \le \mathbb{E}\left[|\mathcal{D}_{t,x,y}(t)|^{2p}\right].$$
(2.26)

The estimate (2.22) is straightforward. Now we prove (2.23). By using the Itô formula and (2.5),

we obtain for $\theta \geq 0$,

$$\begin{aligned} |\mathcal{R}_{t,x,y}(t+\theta)|^{2p} &\leq 2p \int_{t}^{t+\theta} |\mathcal{R}_{t,x,y}(s)|^{2p-2} \left\langle \mathcal{R}_{t,x,y}(s), f(X(t,x;s)) - f(X(t,y;s)) \right\rangle \,\mathrm{d}s \\ &+ \frac{2p(2p-1)}{2} \int_{t}^{t+\theta} |\mathcal{R}_{t,x,y}(s)|^{2p-2} \left\| g(X(t,x;s)) - g(X(t,y;s)) \right\|^{2} \,\mathrm{d}s \\ &+ 2p \int_{t}^{t+\theta} |\mathcal{R}_{t,x,y}(s)|^{2p-2} \left\langle \mathcal{R}_{t,x,y}(s), \left(g(X(t,x;s)) - g(X(t,y;s)) \right) \,\mathrm{d}W(s) \right\rangle \right\rangle \\ &\leq 2p \int_{t}^{t+\theta} |\mathcal{R}_{t,x,y}(s)|^{2p-2} \left(\left\langle \mathcal{D}_{t,x,y}(s), f(X(t,x;s)) - f(X(t,y;s)) \right\rangle \right. \\ &+ \frac{2p-1}{2} \left\| g(X(t,x;s)) - g(X(t,y;s)) \right\|^{2} \right) \,\mathrm{d}s \\ &- 2p \int_{t}^{t+\theta} |\mathcal{R}_{t,x,y}(s)|^{2p-2} \left\langle \mathcal{R}_{t,x,y}(s), \left(g(X(t,x;s)) - g(X(t,y;s)) \right) \right\rangle \,\mathrm{d}W(s) \right\rangle \\ &\leq 2p \int_{t}^{t+\theta} |\mathcal{R}_{t,x,y}(s)|^{2p-2} \left\langle \mathcal{R}_{t,x,y}(s), \left(g(X(t,x;s)) - g(X(t,y;s)) \right) \right\rangle \,\mathrm{d}W(s) \right\rangle \\ &\leq 2p \int_{t}^{t+\theta} |\mathcal{R}_{t,x,y}(s)|^{2p-2} \left\langle \mathcal{R}_{t,x,y}(s), \left(g(X(t,x;s)) - g(X(t,y;s)) \right) \right\rangle \,\mathrm{d}W(s) \right\rangle \\ &\leq 2p \int_{t}^{t+\theta} |\mathcal{R}_{t,x,y}(s)|^{2p-2} \left| f(X(t,x;s)) - f(X(t,y;s)) \right| |x-y| \,\mathrm{d}s \\ &+ 2p \int_{t}^{t+\theta} |\mathcal{R}_{t,x,y}(s)|^{2p-2} \left\langle \mathcal{R}_{t,x,y}(s), \left(g(X(t,x;s)) - g(X(t,y;s)) \right) \,\mathrm{d}W(s) \right\rangle . \end{aligned}$$

According to the Cauchy-Bunyakovsy-Schwarz inequality (twice), (2.6), (2.12) and (2.22), we get

$$\mathbb{E}\left[\left|\mathcal{R}_{t,x,y}(t+\theta)\right|^{2p}\right] \leq 2p|x-y|\int_{t}^{t+\theta} \left(\mathbb{E}\left[\left|f(X(t,x;s)) - f(X(t,y;s))\right|^{p}\right]\right)^{\frac{2}{2p}} \left(\mathbb{E}\left[\left|\mathcal{R}_{t,x,y}(s)\right|^{2p}\right]\right)^{\frac{2p-2}{2p}} ds \\ \leq 2p\sqrt{c_{1}}|x-y|\int_{t}^{t+\theta} \left(\mathbb{E}\left[\left(1+|X(t,x;s)|^{2\kappa-2}+|X(t,y;s)|^{2\kappa-2}\right)^{\frac{p}{2}}\right] \\ \left|X(t,x;s) - X(t,y;s)\right|^{p}\right]\right)^{\frac{1}{p}} \left(\mathbb{E}\left[\left|\mathcal{R}_{t,x,y}(s)\right|^{2p}\right]\right)^{1-\frac{1}{p}} ds \\ \leq 2p\sqrt{c_{1}}|x-y|\int_{t}^{t+\theta} \left(\mathbb{E}\left[\left(1+|X(t,x;s)|^{2\kappa-2}+|X(t,y;s)|^{2\kappa-2}\right)^{p}\right]\right)^{\frac{1}{2p}} \\ \left(\mathbb{E}\left[|X(t,x;s) - X(t,y;s)|^{2p}\right]\right)^{\frac{1}{2p}} \left(\mathbb{E}\left[|\mathcal{R}_{t,x,y}(s)|^{2p}\right]\right)^{1-\frac{1}{p}} ds \\ \leq C'|x-y|^{2}\left(1+|x|^{2\kappa-2}+|y|^{2\kappa-2}\right)^{\frac{1}{2}}\int_{t}^{t+\theta} \left(\mathbb{E}\left[|\mathcal{R}_{t,x,y}(s)|^{2p}\right]\right)^{1-\frac{1}{p}} ds.$$

By employing the Gronwall inequality [26, p.46, Theorem 8.3], we obtain

$$\mathbb{E}\left[|\mathcal{R}(t+\theta)|^{2p}\right] \le C_2 \left(1+|x|^{2\kappa-2}+|y|^{2\kappa-2}\right)^{\frac{p}{2}} |x-y|^{2p} \theta^p.$$
(2.29)

The proof of Lemma 2.4 is thus completed. \Box

Now we present the long-time fundamental strong convergence theorem as follows.

Theorem 2.5. (A long-time fundamental strong convergence theorem) Suppose

- (H_1) Assumption 2.1 hold.
- (H₂) The one-step approximation $Z(t_0, X_0; h)$ is given by (2.2) has the following orders of accuracy: for some $p \ge 1$ there are $\eta_1 \ge 1$, $\eta_2 \ge 1$ and $0 \le h_0 \le 1$, $q_1 > 1$, $\frac{1}{2} < q_2 \le q_1 \frac{1}{2}$ such that for $0 < h \le h_0$, the numerical method has, respectively, local weak and strong errors of order q_1 and q_2 , defined as

$$\left| \mathbb{E} \left[X(t_0, X_0; h) - Z(t_0, X_0; h) \right] \right| \le C_3 \mathbb{E} \left[\left(1 + |X_0|^{\eta_1} \right) \right] h^{q_1},$$
(2.30)

$$\left(\mathbb{E}\left[|X(t_0, X_0; h) - Z(t_0, X_0; h)|^{2p}\right]\right)^{\frac{1}{2p}} \le C_4 \left(\mathbb{E}\left[\left(1 + |X_0|^{2p\eta_2}\right)\right]\right)^{\frac{1}{2p}} h^{q_2},\tag{2.31}$$

where $C_3, C_4 > 0$ and independent of h, t.

(H₃) The approximation Z_k is given by (2.3) has finite moments, i.e., for some $p \ge 1$ there exist $\eta_3 \ge 1$, $h_0 > 0$ and $C_5 > 0$ such that for all $0 < h \le h_0$ and $k \ge 0$,

$$\mathbb{E}\left[|Z_k|^{2p}\right] \le C_5 \mathbb{E}\left[\left(1 + |X_0|^{2p\eta_3}\right)\right],\tag{2.32}$$

where C_5 not depend on h, t, k.

Then there exists a constant $\lambda := \max\{\eta_1\eta_3, (\frac{\kappa-1}{2}+\eta_2)\eta_3\}$, for $h \leq h_1 := \min\{\frac{1}{p\alpha_1}, h_0\}$, the global error is bounded as follows:

$$\left(\mathbb{E}\left[|X_k - Z_k|^{2p}\right]\right)^{\frac{1}{2p}} \le C\left(\mathbb{E}\left[\left(1 + |X_0|^{2\lambda p}\right)\right]\right)^{\frac{1}{2p}} h^{q_2 - \frac{1}{2}},\tag{2.33}$$

where C is a constant and do not depend on h, t, k.

The proof is given in Appendix.

3 Applications: Strong Convergence Rate of the Backward Euler Method over Infinite Time

In this section, we will utilize the previously obtained strong convergence theorem to derive the strong convergence rate of the backward Euler scheme under certain conditions. The backward Euler method applied to SDEs (2.1) takes the following form:

$$Z_k = Z_{k-1} + f(Z_k)h + g(Z_{k-1})\Delta W_{k-1}, \qquad (3.1)$$

where $\Delta W_{k-1} := W(t_k) - W(t_{k-1}), k \ge 1$. Then the one-step approximation of (3.1) reads:

$$Z(t,x;t+h) = x + \int_{t}^{t+h} f(Z(t,x;t+h)) \,\mathrm{d}s + \int_{t}^{t+h} g(x) \,\mathrm{d}W(s).$$
(3.2)

To apply the strong convergence theorem, the first step is to ensure the 2*p*-th moment $(p \ge 1)$ boundedness of Z_k . To this end, we make the following assumptions.

Assumption 3.1. Suppose that the diffusion coefficient of SDEs (2.1) satisfies the global Lipschitz condition. Namely, there exists a constant $\beta_1 > 0$ such that

$$||g(x) - g(y)||^2 \le \beta_1 |x - y|^2.$$
(3.3)

According to Assumptions 2.1, 3.1, there exist some positive constants α_2, β_2 and $2\alpha_2 > (2p^* - 1)\beta_2$, such that

$$\langle x, f(x) \rangle \le -\alpha_2 |x|^2 + c_4, \tag{3.4}$$

$$\|g(x)\|^{2} \le \beta_{2}|x|^{2} + c_{5}, \qquad (3.5)$$

where $c_4, c_5 > 0$ are independent of x.

Lemma 3.2. If there exist some positive constants μ , ν such that $1 - \mu h > 0$ and the sequence $\{Y_k\}_{k\in\mathbb{N}}$ satisfies

$$|Y_k|^{2p} \le (1 - \mu h)|Y_{k-1}|^{2p} + \nu h|Y_{k-1}|^{2p-2}$$
(3.6)

for any $p \ge 1$ and $k \in \mathbb{N}$, then we obtain

$$|Y_k|^{2p} \le C(1+|Y_0|^{2p}),\tag{3.7}$$

where C only depends on p, μ, ν and Y_0 is the initial value.

Proof of Lemma 3.2. When p = 1, by (3.6) we obtain

$$\begin{aligned} |Y_k|^2 &\leq (1 - \mu h)^k |Y_0|^2 + \left(1 + (1 - \mu h) + \dots + (1 - \mu h)^{k-1}\right) \nu h \\ &= (1 - \mu h)^k |Y_0|^2 + \frac{1 - (1 - \mu h)^k}{\mu h} \nu h \\ &\leq C \left(1 + |Y_0|^2\right). \end{aligned}$$
(3.8)

For p > 1, we can utilize the Young inequality to obtain:

$$|Y_k|^{2p} \le \left(1 - \frac{\mu}{2}h\right)|Y_{k-1}|^{2p} + \nu^p \left(\frac{2p-2}{\mu p}\right)^{p-1}h.$$
(3.9)

Following the same procedure as in (3.8), one can straightforwardly derive the inequality (3.7).

Theorem 3.3. Let Assumptions 2.1 and 3.1 hold. For $p \in [1, \lfloor p^* \rfloor]$ and $k \in \mathbb{N}$, there exists a constant $K := K(\alpha_2, \beta_2, c_4, c_5, p) > 0$ such that

$$\mathbb{E}\left[|Z_k|^{2p}\right] \le K \mathbb{E}\left[\left(1 + |X_0|^{2p}\right)\right].$$
(3.10)

In the following proof and throughout the rest of the paper, the symbol K will represent a generic constant that may vary from line to line. Specifically, the factors upon which K depends may differ, but it remains independent of t, h, k.

Proof of Theorem 3.3. From (3.1) and (3.4), we have

$$(1+2\alpha_2 h)^p |Z_k|^{2p} \le \left(|Z_{k-1} + g(Z_{k-1})\Delta W_{k-1}|^2 + Kh \right)^p.$$
(3.11)

Let us first consider the case p = 1. Taking expectations of (3.11) and employing (3.5), we get

$$(1+2\alpha_2 h)\mathbb{E}[|Z_k|^2] \le (1+\beta_2 h)\mathbb{E}[|Z_{k-1}|^2] + Kh.$$
(3.12)

Since $2\alpha_2 > \beta_2$, we have $\frac{1+\beta_2h}{1+2\alpha_2h} = 1 - \gamma h$, where $\gamma = \frac{2\alpha_2 - \beta_2}{1+2\alpha_2h} > 0$. By (3.12) and Lemma 3.2, we derive that $\mathbb{E}[|Z_k|^2] \le (1 - \gamma h)\mathbb{E}[|Z_{k-1}|^2] + Kh$ $\le K\mathbb{E}[(1 + |X_0|^2)]. \tag{3.13}$

For integer p > 1, we note that:

$$J_{1} := \mathbb{E}\left[\left(|Z_{k-1} + g(Z_{k-1})\Delta W_{k-1}|^{2} + Kh\right)^{p} |\mathcal{F}_{t_{k-1}}\right]$$

$$\leq \mathbb{E}\left[|Z_{k-1} + g(Z_{k-1})\Delta W_{k-1}|^{2p} |\mathcal{F}_{t_{k-1}}\right] + Kh\left(1 + |Z_{k-1}|^{2(p-1)}\right)$$

$$= \mathbb{E}\left[\sum_{l=0}^{2p} C_{2p}^{l} |g(Z_{k-1})\Delta W_{k-1}|^{l} |Z_{k-1}|^{2p-l} |\mathcal{F}_{t_{k-1}}\right] + Kh\left(1 + |Z_{k-1}|^{2(p-1)}\right)$$

$$\leq (1 + c(p, \beta_{2}, h)) |Z_{k-1}|^{2p} + Kh\left(1 + |Z_{k-1}|^{2(p-1)}\right),$$
(3.14)

where $c(p, \beta_2, h) := \sum_{l=1}^{p} C_{2p}^{2l} h^l \beta_2^l (2l-1)!!$. By taking the conditional expectations of (3.12) and then applying (3.14), we see

$$\mathbb{E}[|Z_k|^{2p}] \le (1 - \gamma h) \mathbb{E}[|Z_{k-1}|^{2p}] + Kh \mathbb{E}[(1 + |Z_{k-1}|^{2(p-1)})], \qquad (3.15)$$

where $\gamma = \frac{(1+2\alpha_2 h)^p - (1+c(p,\beta_2,h))}{h(1+2\alpha_2 h)^p}$, Verifying $\gamma > 0$ is straightforward. Therefore, by applying Lemma 3.2 and Theorem 3.3, we have established the result for integer p. For non-integer p, we can derive the result using the Young inequality. Thus the proof is finished. \Box

In order to analyze the strong convergence rate, we also introduce an auxiliary one-step approximation,

$$Z_E(t,x;t+h) := x + \int_t^{t+h} f(x) \,\mathrm{d}s + \int_t^{t+h} g(x) \,\mathrm{d}W(s), \qquad (3.16)$$

which can be regarded as a one-step approximation of the Euler-Maruyama method. Subtracting (3.16) from (3.2) yields

$$Z(t,x;t+h) := Z_E(t,x;t+h) + \underbrace{\int_t^{t+h} f(Z(t,x;t+h)) - f(x) \,\mathrm{d}s}_{:=I_f}.$$
(3.17)

To derive the one-step error of the backward Euler scheme (3.2), we require several lemmas.

Lemma 3.4. Suppose Assumptions 2.1 and 3.1 hold. For any $1 \le p < \frac{p^*}{\kappa}$, one gets that

$$\mathbb{E}[|X(h) - X_0|^{2p}] \le C_6 \mathbb{E}[(1 + |X_0|^{2\kappa p})]h^p, \qquad (3.18)$$

where C_6 does not depend on h, t.

Proof of Lemma 3.4. By applying (2.9), (3.5) and (2.12), and then combining the Hölder

inequality with the moment inequality [26, Theorem 7.1], we can obtain that

$$\mathbb{E}\left[|X(h) - X_{0}|^{2p}\right] = \mathbb{E}\left[\left|\int_{0}^{h} f(X(s)) \,\mathrm{d}s + \int_{0}^{h} g(X(s)) \,\mathrm{d}W(s)\right|^{2p}\right] \\
\leq C_{p}h^{2p-1} \int_{0}^{h} \mathbb{E}\left[|f(X(s))|^{2p}\right] \,\mathrm{d}s + C_{p}h^{p-1} \int_{0}^{h} \mathbb{E}\left[\left\|g(X(s))\|^{2p}\right] \,\mathrm{d}s \\
\leq C_{p}h^{2p-1} \int_{0}^{h} \mathbb{E}\left[\left(c_{3} + c_{2}|X(s)|^{2\kappa}\right)^{p}\right] \,\mathrm{d}s + C_{p}h^{p-1} \int_{0}^{h} \mathbb{E}\left[\left(c_{5} + \beta_{2}|X(s)|^{2}\right)^{p}\right] \,\mathrm{d}s \\
\leq C_{6}\mathbb{E}\left[\left(1 + |X_{0}|^{2\kappa p}\right)\right]h^{p}.$$
(3.19)

Here and in the following, the letter C_p is used to denote a generic positive constant, which only depend on p and may vary for each appearance. \Box

Lemma 3.5. Let Assumptions 2.1 and 3.1 be satisfied. For any $1 \le p \le \frac{\lfloor p^* \rfloor}{\kappa}$, one can see that

$$\mathbb{E}[|Z(t,x;t+h) - x|^{2p}] \le C_7 (1+|x|^{2\kappa p}) h^p, \qquad (3.20)$$

where C_7 is independent of h, t.

Proof of Lemma 3.5. Applying (2.9), (3.5), (3.10), the Hölder inequality and the moment inequality, yields

$$\mathbb{E}\left[\left|Z(t,x;t+h)-x\right|^{2p}\right] = \mathbb{E}\left[\left|\int_{t}^{t+h} f(Z(t,x;t+h))\,\mathrm{d}s + \int_{t}^{t+h} g(x)\,\mathrm{d}W(s)\right|^{2p}\right] \\ \leq C_{p}h^{2p-1}\int_{t}^{t+h} \mathbb{E}\left[\left|f(Z(t,x;t+h))\right|^{2p}\right]\,\mathrm{d}s + C_{p}h^{p-1}\int_{t}^{t+h} \mathbb{E}\left[\left\|g(x)\right\|^{2p}\right]\,\mathrm{d}s \\ \leq C_{7}\left(1+|x|^{2\kappa p}\right)h^{p}.$$
(3.21)

Thus the proof is finished. \Box

Lemma 3.6. Let Assumptions 2.1 and 3.1 be fulfilled. Then

$$\begin{aligned} \left| \mathbb{E}[X(t,x;t+h) - Z_E(t,x;t+h)] \right| &\leq K \left(1 + |x|^{2\kappa-1} \right) h^{\frac{3}{2}}, \\ \left| \mathbb{E}[X(t,x;t+h) - x] \right| &\leq K \left(1 + |x|^{\kappa} \right) h, \\ \left| \mathbb{E}[Z_E(t,x;t+h) - x] \right| &\leq K \left(1 + |x|^{\kappa} \right) h. \end{aligned}$$
(3.22)

Proof of Lemma 3.6. For the first item in (3.22), we can utilize a variant of (2.6), (2.12) and

(3.18), as well as the Hölder inequality to acquire that

$$\begin{aligned} \left| \mathbb{E}[X(t,x;t+h) - Z_{E}(t,x;t+h)] \right| &= \left| \mathbb{E} \Big[\int_{t}^{t+h} f(X(s)) - f(x) \, \mathrm{d}s + \int_{t}^{t+h} g(X(s)) - g(x) \, \mathrm{d}W(s) \Big] \right| \\ &\leq \sqrt{c_{1}} \int_{t}^{t+h} \mathbb{E} \Big[(1 + |X(s)|^{2\kappa-2} + |x|^{2\kappa-2})^{\frac{1}{2}} |X(s) - x| \Big] \, \mathrm{d}s \\ &\leq \sqrt{c_{1}} \int_{t}^{t+h} \left(\mathbb{E} \Big[(1 + |X(s)|^{2\kappa-2} + |x|^{2\kappa-2})^{\frac{2\kappa-1}{2\kappa-2}} \Big] \right)^{\frac{\kappa-1}{2\kappa-1}} \\ &\qquad \left(\mathbb{E} \Big[|X(s) - x|^{\frac{2\kappa-1}{\kappa}} \Big] \right)^{\frac{\kappa}{2\kappa-1}} \, \mathrm{d}s \\ &\leq K \int_{t}^{t+h} (1 + |x|^{\kappa-1}) (1 + |x|^{\kappa}) (s - t)^{\frac{1}{2}} \, \mathrm{d}s \\ &\leq K (1 + |x|^{2\kappa-1}) h^{\frac{3}{2}}. \end{aligned}$$

$$(3.23)$$

Furthermore, utilizing (2.9) and (2.12), the remaining terms in (3.22) can be analogously validated. Thus the proof is accomplished. \Box

Lemma 3.7. Under Assumptions 2.1 and 3.1, the one-step backward Euler scheme (3.2) obeys

$$\begin{aligned} \left| \mathbb{E}[X(t,x;t+h) - Z(t,x;t+h)] \right| &\leq K \left(1 + |x|^{2\kappa - 1} \right) h^{\frac{3}{2}}, \\ \left| \mathbb{E}[Z(t,x;t+h)] \right| &\leq K \left(1 + |x|^{\kappa} \right) h. \end{aligned}$$
(3.24)

Proof of Lemma 3.7. We will begin by discussing the following inequality:

$$\begin{aligned} \left| \mathbb{E}[X(t,x;t+h) - Z(t,x;t+h)] \right| &= \left| \mathbb{E}[X(t,x;t+h) - Z_E(t,x;t+h) - I_f] \right| \\ &\leq \left| \mathbb{E}[X(t,x;t+h) - Z_E(t,x;t+h)] \right| + \left| \mathbb{E}[I_f] \right|. \end{aligned}$$
(3.25)

The first term on the right-hand side of (3.25) is estimated in Lemma 3.6, yielding:

$$\left| \mathbb{E}[X(t,x;t+h) - Z_E(t,x;t+h)] \right| \le K \left(1 + |x|^{2\kappa - 1} \right) h^{\frac{3}{2}}.$$
(3.26)

For the second item on the right-hand side of (3.25), by using (2.6), along with Theorem 3.3 and Lemma 3.5, as well as applying the Hölder inequality, one can obtain

$$\begin{aligned} |\mathbb{E}[I_{f}]| &= \left| \mathbb{E} \Big[\int_{t}^{t+h} f(Z(t,x;t+h)) - f(x) \, \mathrm{d}s \Big] \right| \\ &\leq \int_{t}^{t+h} \mathbb{E} \Big[|f(Z(t,x;t+h)) - f(x)| \Big] \, \mathrm{d}s \\ &\leq \sqrt{c_{1}} \int_{t}^{t+h} \mathbb{E} \Big[(1 + |Z(t,x;t+h)|^{2\kappa-2} + |x|^{2\kappa-2})^{\frac{1}{2}} |Z(t,x;t+h) - x| \Big] \, \mathrm{d}s \end{aligned} \tag{3.27} \\ &\leq \sqrt{c_{1}} \int_{t}^{t+h} \Big(\mathbb{E} \Big[(1 + |Z(t,x;t+h)|^{2\kappa-2} + |x|^{2\kappa-2})^{\frac{2\kappa-1}{2\kappa-2}} \Big] \Big)^{\frac{\kappa-1}{2\kappa-1}} \\ &\qquad \left(\mathbb{E} \Big[|Z(t,x;t+h) - x|^{\frac{2\kappa-1}{\kappa}} \Big] \Big)^{\frac{\kappa}{2\kappa-1}} \, \mathrm{d}s \\ &\leq K \Big(1 + |x|^{2\kappa-1} \Big) h^{\frac{3}{2}}, \end{aligned}$$

where I_f is given by (3.17). Combining this with (3.26), we obtain the first inequality in (3.24). Thanks to (2.9) and Theorem 3.3, the second item in (3.24) becomes apparent. \Box

Lemma 3.8. Under Assumptions 2.1, 3.1, for any $1 \le p \le \frac{\lfloor p^* \rfloor}{2\kappa - 1}$, we obtain

$$\left(\mathbb{E}\left[|I_f|^{2p}\right]\right)^{\frac{1}{2p}} \le K\left(1+|x|^{(4\kappa-2)p}\right)^{\frac{1}{2p}}h^{\frac{3}{2}},$$

$$\left(\mathbb{E}\left[|X(t,x;t+h)-Z(t,x;t+h)|^{2p}\right]\right)^{\frac{1}{2p}} \le K\left(1+|x|^{(4\kappa-2)p}\right)^{\frac{1}{2p}}h,$$
(3.28)

where I_f is given by (3.17).

Proof of Lemma 3.8. To establish the first inequality in (3.28), we employ (2.6), the Hölder inequality, Theorem 3.3, as well as Lemma 3.5 to deduce

$$\mathbb{E}\left[|I_{f}|^{2p}\right] = \mathbb{E}\left[\left|\int_{t}^{t+h} f(Z(t,x;t+h)) - f(x) \,\mathrm{d}s\right|^{2p}\right] \\
\leq c_{1}^{p}h^{2p-1} \int_{t}^{t+h} \mathbb{E}\left[\left(1 + |Z(t,x;t+h)|^{2\kappa-2} + |x|^{2\kappa-2}\right)^{p} |Z(t,x;t+h) - x|^{2p}\right] \,\mathrm{d}s \\
\leq c_{1}^{p}h^{2p-1} \int_{t}^{t+h} \left(\mathbb{E}\left[\left(1 + |Z(t,x;t+h)|^{2\kappa-2} + |x|^{2\kappa-2}\right)^{\frac{(2\kappa-1)p}{\kappa-1}}\right]\right)^{\frac{\kappa-1}{2\kappa-1}} \\
\left(\mathbb{E}\left[|Z(t,x;t+h) - x|^{\frac{2(2\kappa-1)p}{\kappa}}\right]\right)^{\frac{\kappa}{2\kappa-1}} \,\mathrm{d}s \\
\leq K\left(1 + |x|^{(4\kappa-2)p}\right)h^{3p}.$$
(3.29)

By utilizing (2.6), (3.3) and (2.12), along with (3.29), we can apply the Hölder inequality, the moment inequality and Lemma 3.4 to estimate the other inequality in (3.28) as follows:

$$\begin{split} & \mathbb{E}\left[|X(t,x;t+h) - Z(t,x;t+h)|^{2p}\right] \\ &= \mathbb{E}\left[\left|\int_{t}^{t+h} f(X(s)) - f(Z(t,x;t+h)) \,\mathrm{d}s + \int_{t}^{t+h} g(X(s)) - g(x) \,\mathrm{d}W(s)\right|^{2p}\right] \\ &= \mathbb{E}\left[\left|\int_{t}^{t+h} f(X(s)) - f(x) \,\mathrm{d}s - I_{f} + \int_{t}^{t+h} g(X(s)) - g(x) \,\mathrm{d}W(s)\right|^{2p}\right] \\ &\leq C_{p}h^{2p-1} \int_{t}^{t+h} \mathbb{E}\left[|f(X(s)) - f(x)|^{2p}\right] \,\mathrm{d}s + C_{p}\mathbb{E}\left[|(I_{f})|^{2p}\right] \\ &+ C_{p}h^{p-1} \int_{t}^{t+h} \mathbb{E}\left[\left||g(X(s)) - g(x)|\right|^{2p}\right] \,\mathrm{d}s \\ &\leq c_{1}^{p}C_{p}h^{2p-1} \int_{t}^{t+h} \mathbb{E}\left[\left(1 + |X(s)|^{2\kappa-2} + |x|^{2\kappa-2}\right)^{p}|X(s) - x|^{2p}\right] \,\mathrm{d}s \\ &+ K\left(1 + |x|^{(4\kappa-2)p}\right)h^{3p} + C_{p}h^{p-1}\beta_{1}^{p} \int_{t}^{t+h} \mathbb{E}\left[|X(s) - x|^{2p}\right] \,\mathrm{d}s \\ &\leq Kh^{2p-1} \int_{t}^{t+h} \left(\mathbb{E}\left[\left(1 + |X(s)|^{2\kappa-2} + |x|^{2\kappa-2}\right)^{\frac{(2\kappa-1)p}{\kappa-1}}\right]\right)^{\frac{\kappa-1}{2\kappa-1}} \\ &\left(\mathbb{E}\left[|X(s) - x|^{\frac{2(2\kappa-1)p}{\kappa}}\right]\right)^{\frac{\kappa}{2\kappa-1}} \,\mathrm{d}s + K\left(1 + |x|^{(4\kappa-2)p}\right)h^{3p} + K\left(1 + |x|^{2\kappa p}\right)h^{2p} \\ &\leq K\left(1 + |x|^{(4\kappa-2)p}\right)h^{2p}. \end{split}$$

Thus the proof is finished. \Box

Theorem 3.9. Under Assumptions 2.1, 3.1, for any $1 \le p \le \frac{\lfloor p^* \rfloor}{2\kappa - 1}$ and $0 < h \le h_1$, the backward Euler method (3.2) has a strong convergence rate of order $\frac{1}{2}$ over infinite time, namely,

$$\left(\mathbb{E}\left[|X_k - Z_k|^{2p}\right]\right)^{\frac{1}{2p}} \le K\left(\mathbb{E}\left[\left(1 + |X_0|^{(5\kappa - 3)p}\right)\right]\right)^{\frac{1}{2p}} h^{\frac{1}{2}}.$$
(3.31)

Proof of Theorem 3.9. With the aid of Theorem 3.3, Lemmas 3.7, 3.8 and according to Theorem 2.5, one can straightforwardly obtain the long-time strong convergence rate of the backward Euler method (3.2). \Box

4 Applications: Strong Convergence Rate of the Projected Euler Method over Infinite Time

In this section, let us consider another application of the strong convergence theorem and analyze the strong convergence rate of a kind of projected Euler scheme [2] under certain assumptions. The projected Euler method of SDEs (2.1) proposed here is given as follows:

$$\begin{cases} \bar{Z}_k := \Phi(Z_k), \\ Z_{k+1} := \bar{Z}_k + hf(\bar{Z}_k) + g(\bar{Z}_k)\Delta W_k, \quad Z_0 = X_0, \end{cases}$$
(4.1)

where $\Delta W_k := W(t_{k+1}) - W(t_k), k \ge 1$ and the projection operator $\Phi \colon \mathbb{R}^d \to \mathbb{R}^d$ is assumed to satisfy the following assumptions.

Assumption 4.1. Let the mapping $\Phi \colon \mathbb{R}^d \to \mathbb{R}^d$ obey

$$\begin{aligned} |\Phi(x)| &\le h^{-\frac{1}{2(\kappa+1)}}, \\ |\Phi(x) - \Phi(y)| &\le |x - y| \end{aligned}$$
(4.2)

and $\Phi(0) = 0 \in \mathbb{R}^d$, where κ is given by (2.6).

Taking
$$y = 0$$
 in (4.2) yields

$$|\Phi(x)| \le |x|. \tag{4.3}$$

Below we divide the analysis of the strong convergence rate of the scheme (4.1) into three parts. The first part is to establish the 2p-th $(p \ge 1)$ moment boundedness of the numerical solution.

4.1 Bounded moments of the projected Euler method

This part is to establish the boundedness of the high-order moments of the projected Euler method. Based on Assumption 2.1, we can conclude that the diffusion coefficient of SDEs (2.1) satisfies the following inequality:

$$||g(x) - g(y)||^2 \le \beta_3 (1 + |x|^{\kappa - 1} + |y|^{\kappa - 1}) |x - y|^2,$$
(4.4)

where β_3 is a constant. As a consequence, we have

$$\|g(x)\|^{2} \le \beta_{4}|x|^{\kappa+1} + c_{6}, \qquad (4.5)$$

where $\beta_4 = \frac{2\beta_3(\kappa+3)}{\kappa+1}$, $c_6 = 2 ||g(0)||^2 + \frac{2\beta_3(\kappa-1)}{\kappa+1}$. In addition, employing (2.9), (4.5) and the fact that $|\Phi(Z_{k-1})| \leq h^{-\frac{1}{2(\kappa+1)}}$ yields

$$|f(\bar{Z}_{k-1})|^2 \le c_2 |\bar{Z}_{k-1}|^{2\kappa} + c_3 \le c_2 h^{-1} + c_3$$
(4.6)

and

$$\left\|g(\bar{Z}_{k-1})\right\|^{2} \leq \beta_{4} |\bar{Z}_{k-1}|^{\kappa+1} + c_{6} \leq \beta_{4} h^{-\frac{1}{2}} + c_{6}.$$
(4.7)

Next, we shall establish the moment bounds for the projected Euler scheme.

Theorem 4.2. Let Assumptions 2.1,4.1 hold. For any $1 \le p \le \lfloor p^* \rfloor$ and $0 < h \le h_2 := \min\{h_1, \frac{1}{2p(\alpha_1 - \epsilon_1)}\}$, one has

$$\mathbb{E}\left[|Z_k|^{2p}\right] \le K \mathbb{E}\left[\left(1 + |X_0|^{2p}\right)\right].$$
(4.8)

Proof of Theorem 4.2. From (4.1), one can derive

$$1 + |Z_{k}|^{2} = 1 + |\bar{Z}_{k-1} + f(\bar{Z}_{k-1})h + g(\bar{Z}_{k-1})\Delta W_{k-1}|^{2}$$

= 1 + $|\bar{Z}_{k-1}|^{2} + 2\langle \bar{Z}_{k-1}, f(\bar{Z}_{k-1})h \rangle + |g(\bar{Z}_{k-1})\Delta W_{k-1}|^{2}$
+ 2 $\langle \bar{Z}_{k-1} + f(\bar{Z}_{k-1})h, g(\bar{Z}_{k-1})\Delta W_{k-1} \rangle + |f(\bar{Z}_{k-1})h|^{2}.$ (4.9)

Combining this with (4.6) shows

$$\mathbb{E}[(1+|Z_k|^2)^p | \mathcal{F}_{t_{k-1}}] \le \mathbb{E}[(1+\xi)^p | \mathcal{F}_{t_{k-1}}](1+|\bar{Z}_{k-1}|^2)^p + Kh(1+|\bar{Z}_{k-1}|^2)^{p-1},$$
(4.10)

where we denote

$$\xi := \frac{2\langle \bar{Z}_{k-1}, f(\bar{Z}_{k-1})h \rangle + |g(\bar{Z}_{k-1})\Delta W_{k-1}|^2 + 2\langle \bar{Z}_{k-1} + f(\bar{Z}_{k-1})h, g(\bar{Z}_{k-1})\Delta W_{k-1} \rangle}{1 + |\bar{Z}_{k-1}|^2} := \xi_1 + \xi_2 + \xi_3.$$

In what follows four cases are considered.

Case 1. When p = 1, one obtains that

$$\mathbb{E}\left[\left(1+|Z_{k}|^{2}\right)|\mathcal{F}_{t_{k-1}}\right] \leq \mathbb{E}\left[(1+\xi)|\mathcal{F}_{t_{k-1}}\right]\left(1+|\bar{Z}_{k-1}|^{2}\right)+Kh \\
= \mathbb{E}\left[\left(1+\xi_{1}+\xi_{2}+\xi_{3}\right)|\mathcal{F}_{t_{k-1}}\right]\left(1+|\bar{Z}_{k-1}|^{2}\right)+Kh.$$
(4.11)

Utilizing the properties of Brownian motion, we have the following:

$$\mathbb{E}[\xi_{1} + \xi_{2} | \mathcal{F}_{t_{k-1}}] = \mathbb{E}\left[\frac{2\left\langle \bar{Z}_{k-1}, f(\bar{Z}_{k-1})h\right\rangle + |g(\bar{Z}_{k-1})\Delta W_{k-1}|^{2}}{1 + |\bar{Z}_{k-1}|^{2}}\Big| \mathcal{F}_{t_{k-1}}\right]$$

$$= \frac{2\left\langle \bar{Z}_{k-1}, f(\bar{Z}_{k-1})h\right\rangle + \left\|g(\bar{Z}_{k-1})\right\|^{2}h}{1 + |\bar{Z}_{k-1}|^{2}}$$
(4.12)

and

$$\mathbb{E}[\xi_3|\mathcal{F}_{t_{k-1}}] = \mathbb{E}\left[\frac{2\left\langle \bar{Z}_{k-1} + f(\bar{Z}_{k-1})h, g(\bar{Z}_{k-1})\Delta W_{k-1} \right\rangle}{1 + |\bar{Z}_{k-1}|^2} \middle| \mathcal{F}_{t_{k-1}}\right] = 0.$$
(4.13)

Taking these two estimates into account, we can derive from (4.11) that

$$\mathbb{E}[\xi|\mathcal{F}_{t_{k-1}}] = \frac{2\left\langle \bar{Z}_{k-1}, f(\bar{Z}_{k-1})h \right\rangle + \left\| g(\bar{Z}_{k-1}) \right\|^2 h}{1 + |\bar{Z}_{k-1}|^2}.$$
(4.14)

By (2.8) and (4.2), we acquire

$$\mathbb{E}\left[\left(1+|Z_k|^2\right)\right] \le (1-2(\alpha_1-\epsilon_1)h)\mathbb{E}\left[\left(1+|Z_{k-1}|^2\right)\right] + Kh.$$
(4.15)

Then from Lemma 3.2 we can see that the scheme (4.8) is evident when p = 1.

Case 2. When p = 2, one sees that

$$\mathbb{E}\Big[\left(1+|Z_{k}|^{2}\right)^{2}\big|\mathcal{F}_{t_{k-1}}\Big] \leq \mathbb{E}\Big[\left(1+\xi\right)^{2}|\mathcal{F}_{t_{k-1}}\Big]\left(1+|\bar{Z}_{k-1}|^{2}\right)^{2}+Kh\left(1+|\bar{Z}_{k-1}|^{2}\right) \\ = \mathbb{E}\Big[\left(1+2\xi+\xi^{2}\right)|\mathcal{F}_{t_{k-1}}\Big]\left(1+|\bar{Z}_{k-1}|^{2}\right)^{2}+Kh\left(1+|\bar{Z}_{k-1}|^{2}\right) \tag{4.16}$$

and

$$\mathbb{E}\left[\xi^{2}|\mathcal{F}_{t_{k-1}}\right] = \mathbb{E}\left[\left(\xi_{1}+\xi_{2}+\xi_{3}\right)^{2}\middle|\mathcal{F}_{t_{k-1}}\right] \\
= \mathbb{E}\left[\left(\xi_{1}^{2}+\xi_{2}^{2}+\xi_{3}^{2}+2\xi_{1}\xi_{2}+2\xi_{1}\xi_{3}+2\xi_{2}\xi_{3}\right)\middle|\mathcal{F}_{t_{k-1}}\right].$$
(4.17)

Next, we will provide estimates for each term in (4.17). By (4.6), the first item on the right-hand side of (4.17) is bounded as:

$$\mathbb{E}\left[\xi_{1}^{2}|\mathcal{F}_{t_{k-1}}\right] \leq \mathbb{E}\left[\frac{4|\bar{Z}_{k-1}|^{2}|f(\bar{Z}_{k-1})h|^{2}}{\left(1+|\bar{Z}_{k-1}|^{2}\right)^{2}}\Big|\mathcal{F}_{t_{k-1}}\right] \leq \frac{4|\bar{Z}_{k-1}|^{2}\left(c_{2}h+c_{3}h^{2}\right)}{\left(1+|\bar{Z}_{k-1}|^{2}\right)^{2}} \leq \frac{Kh}{1+|\bar{Z}_{k-1}|^{2}}.$$
(4.18)

For the second item on the right-hand side of (4.17), using (4.7) gives

$$\mathbb{E}\left[\xi_{2}^{2}|\mathcal{F}_{t_{k-1}}\right]\mathbb{E}\left[\frac{|g(\bar{Z}_{k-1})\Delta W_{k-1}|^{4}}{\left(1+|\bar{Z}_{k-1}|^{2}\right)^{2}}\Big|\mathcal{F}_{t_{k-1}}\right] \leq \frac{3\left(\beta_{4}h^{-\frac{1}{2}}+c_{6}\right)^{2}h^{2}}{\left(1+|\bar{Z}_{k-1}|^{2}\right)^{2}} \leq \frac{Kh}{1+|\bar{Z}_{k-1}|^{2}}.$$
(4.19)

Employing (2.8) and (3.5) to bound the third and fourth items on the right-hand side of (4.17) as follows: $5 \pm \overline{5} = 5 \pm \overline{5} = 5$

$$\mathbb{E}\left[\xi_{3}^{2}|\mathcal{F}_{t_{k-1}}\right] \leq \mathbb{E}\left[\frac{4|\bar{Z}_{k-1} + f(\bar{Z}_{k-1})h|^{2}|g(\bar{Z}_{k-1})\Delta W_{k-1}|^{2}}{\left(1 + |\bar{Z}_{k-1}|^{2}\right)^{2}}\Big|\mathcal{F}_{t_{k-1}}\right] \\
\leq \frac{4h\left\|g(\bar{Z}_{k-1})\right\|^{2}}{1 + |\bar{Z}_{k-1}|^{2}} + \frac{Kh}{1 + |\bar{Z}_{k-1}|^{2}} \tag{4.20}$$

and

$$\mathbb{E}\left[2\xi_{1}\xi_{2}|\mathcal{F}_{t_{k-1}}\right] = \mathbb{E}\left[\frac{4\left\langle\bar{Z}_{k-1}, f(\bar{Z}_{k-1})h\right\rangle|g(\bar{Z}_{k-1})\Delta W_{k-1}|^{2}}{\left(1+|\bar{Z}_{k-1}|^{2}\right)^{2}}\Big|\mathcal{F}_{t_{k-1}}\right] \le \frac{Kh}{1+|\bar{Z}_{k-1}|^{2}}.$$
(4.21)

Due to the properties of Brownian motion, we know

$$\mathbb{E}\big[2\xi_1\xi_3|\mathcal{F}_{t_{k-1}}\big] = \mathbb{E}\big[2\xi_2\xi_3|\mathcal{F}_{t_{k-1}}\big] = 0.$$

Combining (4.18), (4.19), (4.20) with (4.21), we obtain

$$\mathbb{E}\left[\xi^{2} \big| \mathcal{F}_{t_{k-1}}\right] \leq \frac{4h \left\| g(\bar{Z}_{k-1}) \right\|^{2}}{1 + |\bar{Z}_{k-1}|^{2}} + \frac{Kh}{1 + |\bar{Z}_{k-1}|^{2}}.$$
(4.22)

Next, by (2.8) and (4.2), we have

$$\mathbb{E}\Big[\left(1+|Z_{k}|^{2}\right)^{2}\left|\mathcal{F}_{t_{k-1}}\right] \leq \left(1+\frac{4\left\langle\bar{Z}_{k-1},f(\bar{Z}_{k-1})h\right\rangle+6h\left\|g(\bar{Z}_{k-1})\right\|^{2}}{1+|\bar{Z}_{k-1}|^{2}}\right)\left(1+|\bar{Z}_{k-1}|^{2}\right)^{2}+Kh\left(1+|\bar{Z}_{k-1}|^{2}\right) \\ \leq \left(1-4(\alpha_{1}-\epsilon_{1})h\right)\left(1+|Z_{k-1}|^{2}\right)^{2}+Kh\left(1+|Z_{k-1}|^{2}\right). \tag{4.23}$$

Thanks to Lemma 3.2, the inequality (4.8) holds when p = 2.

Case 3. For p = 3, we make a further decomposition as follows:

$$\mathbb{E}\Big[\left(1+|Z_{k}|^{2}\right)^{3}\big|\mathcal{F}_{t_{k-1}}\Big] \leq \mathbb{E}\Big[\left(1+\xi\right)^{3}\big|\mathcal{F}_{t_{k-1}}\Big]\left(1+|\bar{Z}_{k-1}|^{2}\right)^{3}+Kh\left(1+|\bar{Z}_{k-1}|^{2}\right)^{2} \\ = \mathbb{E}\Big[\left(1+3\xi+3\xi^{2}+\xi^{3}\right)\big|\mathcal{F}_{t_{k-1}}\Big]\left(1+|\bar{Z}_{k-1}|^{2}\right)^{3}+Kh\left(1+|\bar{Z}_{k-1}|^{2}\right)^{2} \tag{4.24}$$

and

$$\mathbb{E}\left[\xi^{3}\big|\mathcal{F}_{t_{k-1}}\right] = \mathbb{E}\left[\left(\xi_{1} + \xi_{2} + \xi_{3}\right)^{3}\big|\mathcal{F}_{t_{k-1}}\right] \\ = \mathbb{E}\left[\left(\xi_{2}^{3} + 3\xi_{1}^{2}\xi_{2} + 3\xi_{2}\xi_{3}^{2} + \xi_{1}^{3} + 3\xi_{1}\xi_{2}^{2} + 3\xi_{1}\xi_{3}^{2} + 3(\xi_{1} + \xi_{2})^{2}\xi_{3} + \xi_{3}^{3}\right)\big|\mathcal{F}_{t_{k-1}}\right].$$

$$(4.25)$$

Next, we will bound these terms separately. By applying (4.7), we obtain

$$\mathbb{E}\left[\xi_{2}^{3}\big|\mathcal{F}_{t_{k-1}}\right] = \mathbb{E}\left[\frac{|g(\bar{Z}_{k-1})\Delta W_{k-1}|^{6}}{\left(1+|\bar{Z}_{k-1}|^{2}\right)^{3}}\Big|\mathcal{F}_{t_{k-1}}\right] \le \frac{15\left(\beta_{4}h^{-\frac{1}{2}}+c_{6}\right)^{3}h^{3}}{\left(1+|\bar{Z}_{k-1}|^{2}\right)^{3}} \le \frac{Kh}{1+|\bar{Z}_{k-1}|^{2}}.$$
(4.26)

With the help of (4.6) and (4.7), the second and third items on the right-hand side of (4.25) can be estimated as

$$\mathbb{E}\left[3\xi_{1}^{2}\xi_{2}\big|\mathcal{F}_{t_{k-1}}\right] = \mathbb{E}\left[\frac{12|\bar{Z}_{k-1}|^{2}|f(\bar{Z}_{k-1})h|^{2}|g(\bar{Z}_{k-1})\Delta W_{k-1}|^{2}}{\left(1+|\bar{Z}_{k-1}|^{2}\right)^{3}}\Big|\mathcal{F}_{t_{k-1}}\right] \\
\leq \frac{12|\bar{Z}_{k-1}|^{2}\left(c_{2}h^{-1}+c_{3}\right)\left(\beta_{4}h^{-\frac{1}{2}}+c_{6}\right)h^{3}}{\left(1+|\bar{Z}_{k-1}|^{2}\right)^{3}} \\
\leq \frac{Kh}{1+|\bar{Z}_{k-1}|^{2}}$$
(4.27)

and

$$\mathbb{E}\left[3\xi_{2}\xi_{3}^{2}\big|\mathcal{F}_{t_{k-1}}\right] \leq \mathbb{E}\left[\frac{12|\bar{Z}_{k-1} + f(\bar{Z}_{k-1})h|^{2}|g(\bar{Z}_{k-1})\Delta W_{k-1}|^{4}}{\left(1 + |\bar{Z}_{k-1}|^{2}\right)^{3}}\Big|\mathcal{F}_{t_{k-1}}\right] \leq \frac{Kh}{1 + |\bar{Z}_{k-1}|^{2}}.$$
(4.28)

In view of (2.8), it is not difficult to observe that

$$\mathbb{E}\left[\left(\xi_1^3 + 3\xi_1\xi_2^2 + 3\xi_1\xi_3^2\right) \middle| \mathcal{F}_{t_{k-1}}\right] \le \frac{Kh}{1 + |\bar{Z}_{k-1}|^2}.$$
(4.29)

By applying properties of the Brownian motion, we know

$$\mathbb{E}\left[\xi_{3}^{3}\big|\mathcal{F}_{t_{k-1}}\right] = \mathbb{E}\left[3(\xi_{1}+\xi_{2})^{2}\xi_{3}|\mathcal{F}_{t_{k-1}}\right] = 0.$$

Then, using (4.26)-(4.29), we are able to obtain

$$\mathbb{E}\left[\xi^3 \middle| \mathcal{F}_{t_{k-1}}\right] \le \frac{Kh}{1+|\bar{Z}_{k-1}|^2}.$$
(4.30)

By (2.8) and (4.2), we see

$$\mathbb{E}\Big[\left(1+|Z_{k}|^{2}\right)^{3}\big|\mathcal{F}_{t_{k-1}}\Big] \leq \left(1+\frac{6\left\langle\bar{Z}_{k-1},f(\bar{Z}_{k-1})h\right\rangle+15h\left\|g(\bar{Z}_{k-1})\right\|^{2}}{1+|\bar{Z}_{k-1}|^{2}}\right)\left(1+|\bar{Z}_{k-1}|^{2}\right)^{3} + Kh\left(1+|\bar{Z}_{k-1}|^{2}\right)^{2} \leq (1-6(\alpha_{1}-\epsilon_{1})h)\left(1+|Z_{k-1}|^{2}\right)^{3} + Kh\left(1+|Z_{k-1}|^{2}\right)^{2}.$$
(4.31)

By applying Lemma 3.2, the proof of the inequality (4.8) is thus completed for the case when p = 3.

Case 4. For $p \ge 4$, based on the previous analysis, we have

$$\mathbb{E}\left[\left(1+|Z_{k}|^{2}\right)^{p}\left|\mathcal{F}_{t_{k-1}}\right] \leq \mathbb{E}\left[\left(1+p\xi+\frac{p(p-1)}{2}\xi^{2}+\frac{p(p-1)(p-2)}{6}\xi^{3}\right)\left|\mathcal{F}_{t_{k-1}}\right]\right] \\ \left(1+|\bar{Z}_{k-1}|^{2}\right)^{p}+Kh\left(1+|\bar{Z}_{k-1}|^{2}\right)^{p-1}.$$
(4.32)

Taking (4.14), (4.22) and (4.30) into account and using (2.8), (4.2) we derive from (4.32) that

$$\mathbb{E}\left[\left(1+|Z_{k}|^{2}\right)^{p}\left|\mathcal{F}_{t_{k-1}}\right] \leq \left(1+\frac{2p\left\langle\bar{Z}_{k-1},f(\bar{Z}_{k-1})h\right\rangle+p(2p-1)\left\|g(\bar{Z}_{k-1})\right\|^{2}}{1+|\bar{Z}_{k-1}|^{2}}\right) \times \left(1+|\bar{Z}_{k-1}|^{2}\right)^{p}+Kh\left(1+|\bar{Z}_{k-1}|^{2}\right)^{p-1} \leq (1-2p(\alpha_{1}-\epsilon_{1})h)\left(1+|Z_{k-1}|^{2}\right)^{p}+Kh\left(1+|Z_{k-1}|^{2}\right)^{p-1}.$$
(4.33)

Therefore, (4.8) is evident for integer p by applying Lemma 3.2. For non-integer p, we can bound these terms using the Young inequality, which completes the proof. \Box

4.2 Strong convergence rate of the projected Euler method

In light of the fundamental strong convergence theorem, Theorem 2.5, we need to verify the local weak and strong errors in (2.30) and (2.31). For the purpose of the strong convergence analysis, we will start from the following auxiliary lemmas.

Lemma 4.3. Suppose Assumption 2.1 hold. For $1 \le p < \frac{p^*}{\kappa}$, one obtains that

$$\mathbb{E}[|X(h) - X_0|^{2p}] \le C_8 \mathbb{E}[(1 + |X_0|^{2\kappa p})]h^p.$$
(4.34)

where C_8 do not depend on h, t.

Proof of Lemma 4.3. Based on (2.9) and (4.7), we can use the Hölder inequality, the moment inequality and (2.12) to obtain

$$\mathbb{E}\left[|X(h) - X_{0}|^{2p}\right] = \mathbb{E}\left[\left|\int_{0}^{h} f(X(s)) \,\mathrm{d}s + \int_{0}^{h} g(X(s)) \,\mathrm{d}W(s)\right|^{2p}\right]$$

$$\leq C_{p}h^{2p-1} \int_{0}^{h} \mathbb{E}\left[\left(c_{3} + c_{2}|X(s)|^{2\kappa}\right)^{p}\right] \,\mathrm{d}s + C_{p}h^{p-1} \int_{0}^{h} \mathbb{E}\left[\left(c_{6} + \beta_{4}|X(s)|^{\kappa+1}\right)^{p}\right] \,\mathrm{d}s$$

$$\leq C_{8}\mathbb{E}\left[\left(1 + |X_{0}|^{2\kappa p}\right)\right]h^{p}.$$
(4.35)

Thus the proof is finished. \Box

Lemma 4.4. For $h \in (0,1]$ and $q \ge 1$, consider the mapping $\mathbb{R}^d \ni x \mapsto \Phi(x) \in \mathbb{R}^d$ which satisfies

$$|x - \Phi(x)| \le 2\left(1 + |x|^{q+1}\right) h^{\frac{q}{2(\kappa+1)}}.$$
(4.36)

Proof of Lemma 4.4. We will divide the analysis into two cases: $|x| \leq h^{-\frac{1}{2(\kappa+1)}}$ and $|x| > h^{-\frac{1}{2(\kappa+1)}}$ $h^{-\frac{1}{2(\kappa+1)}}$.

For $|x| \leq h^{-\frac{1}{2(\kappa+1)}}$, we have $x = \Phi(x)$, thus (4.36) is evident. For $|x| > h^{-\frac{1}{2(\kappa+1)}}$, by using the elementary inequality, we can easily obtain

$$\left|x - \Phi(x)\right| = \left|x - h^{-\frac{1}{2(\kappa+1)}} \frac{x}{|x|}\right| \le |x| + \left|h^{-\frac{1}{2(\kappa+1)}}\right| \le 2\left(1 + |x|^{q+1}\right)h^{\frac{q}{2(\kappa+1)}}.$$
(4.37)

Thus the proof is accomplished. \Box

Lemma 4.5. Under the Assumptions 2.1 and 4.1, we obtain

$$\left| \mathbb{E} \left[Z_E(t, x; t+h) - Z(t, x; t+h) \right] \right| \le K \left(1 + |x|^{3\kappa+4} \right) h^{\frac{3}{2}},$$

$$\left| \mathbb{E} \left[X(t, x; t+h) - Z(t, x; t+h) \right] \right| \le K \left(1 + |x|^{3\kappa+4} \right) h^{\frac{3}{2}}.$$
(4.38)

Proof of Lemma 4.5. We can employ (2.6), the Hölder inequality and Lemma 4.4 for the first item in (4.38), to infer

$$\begin{aligned} & \left| \mathbb{E} \left[Z_E(t, x; t+h) - Z(t, x; t+h) \right] \right| \\ &= \left| \mathbb{E} \left[x - \Phi(x) + \int_t^{t+h} f(x) - f(\Phi(x)) \, \mathrm{d}s + \int_t^{t+h} g(x) - g(\Phi(x)) \, \mathrm{d}W(s) \right] \right| \\ &\leq K \left(1 + |x|^{3\kappa+4} \right) h^{\frac{3}{2}}. \end{aligned}$$
(4.39)

Finally, using Lemma 3.6 and (4.39), we have

$$\begin{aligned} & \left| \mathbb{E} \big[X(t,x;t+h) - Z(t,x;t+h) \big] \big| \\ &= \left| \mathbb{E} \big[(X(t,x;t+h) - Z_E(t,x;t+h) + Z_E(t,x;t+h) - Z(t,x;t+h)) \big] \big| \\ &\leq \left| \mathbb{E} \big[(X(t,x;t+h) - Z_E(t,x;t+h)) \big] \big| + \left| \mathbb{E} \big[(Z_E(t,x;t+h) - Z(t,x;t+h)) \big] \big| \\ &\leq K \big(1 + |x|^{3\kappa + 4} \big) h^{\frac{3}{2}}. \end{aligned}$$

$$(4.40)$$

Thus the proof is completed. \Box

Lemma 4.6. Under Assumptions 2.1, 4.1, for any $1 \le p \le \frac{\lfloor p^* \rfloor}{2\kappa - 1}$, we have

$$\left(\mathbb{E}\left[|X(t,x;t+h) - Z_E(t,x;t+h)|^{2p}\right]\right)^{\frac{1}{2p}} \leq K\left(1 + |x|^{(4\kappa-2)p}\right)^{\frac{1}{2p}}h, \\
\left(\mathbb{E}\left[|Z_E(t,x;t+h) - Z(t,x;t+h)|^{2p}\right]\right)^{\frac{1}{2p}} \leq K\left(1 + |x|^{4(\kappa+1)p+1}\right)^{\frac{1}{2p}}h, \\
\left(\mathbb{E}\left[|X(t,x;t+h) - Z(t,x;t+h)|^{2p}\right]\right)^{\frac{1}{2p}} \leq K\left(1 + |x|^{4(\kappa+1)p+1}\right)^{\frac{1}{2p}}h.$$
(4.41)

Proof of Lemma 4.6. First, let us consider the first term in (4.41). By using (2.6), (4.4), (2.12), the moment inequality, the Hölder inequality and Lemma 4.3, one can get

$$\mathbb{E}\left[|X(t,x;t+h) - Z_{E}(t,x;t+h)|^{2p}\right] \\
= \mathbb{E}\left[\left|\int_{t}^{t+h} f(X(s)) - f(x) \, \mathrm{d}s + \int_{t}^{t+h} g(X(s)) - g(x) \, \mathrm{d}W(s)\right|^{2p}\right] \\
\leq Kh^{2p-1} \int_{t}^{t+h} \mathbb{E}\left[\left(1 + |X(s)|^{2\kappa-2} + |x|^{2\kappa-2}\right)^{p} |X(s) - x|^{2p}\right] \, \mathrm{d}s \\
+ Kh^{p-1} \int_{t}^{t+h} \mathbb{E}\left[\left(1 + |X(s)|^{\kappa-1} + |x|^{\kappa-1}\right)^{p} |X(s) - x|^{2p}\right] \, \mathrm{d}s \\
\leq Kh^{p-1} \int_{t}^{t+h} \left(\mathbb{E}\left[\left(1 + |X(s)|^{2\kappa-2} + |x|^{2\kappa-2}\right)^{\frac{(2\kappa-1)p}{\kappa-1}}\right]\right)^{\frac{\kappa-1}{2\kappa-1}} \\
\left(\mathbb{E}\left[|X(s) - x|^{\frac{2(2\kappa-1)p}{\kappa}}\right]\right)^{\frac{\kappa}{2\kappa-1}} \, \mathrm{d}s \\
\leq K\left(1 + |x|^{(2\kappa-2)p}\right)(1 + |x|^{2\kappa p})h^{p-1} \int_{t}^{t+h} s^{p} \, \mathrm{d}s \\
\leq K\left(1 + |x|^{(4\kappa-2)p}\right)h^{2p}.$$
(4.42)

Next, utilizing (2.6), (4.4), (4.36) and the Hölder inequality, the second item of the inequality (4.41) can be estimated as follows:

$$\mathbb{E}\left[|(Z_E(t,x;t+h) - Z(t,x;t+h))|^{2p} \right]
= \mathbb{E}\left[\left| x - \Phi(x) + \int_t^{t+h} f(x) - f(\Phi(x)) \, \mathrm{d}s + \int_t^{t+h} g(x) - g(\Phi(x)) \, \mathrm{d}W(s) \right|^{2p} \right]
\leq C_p \mathbb{E}\left[|x - \Phi(x)|^{2p} \right] + C_p h^{2p-1} \int_t^{t+h} \mathbb{E}\left[|f(x) - f(\Phi(x))|^{2p} \right] \, \mathrm{d}s
+ C_p h^{p-1} \int_t^{t+h} \mathbb{E}\left[|g(x) - g(\Phi(x))|^{2p} \right] \, \mathrm{d}s
\leq K \left(1 + |x|^{4(\kappa+1)p+1} \right) h^{2p}.$$
(4.43)

Further, in view of (4.42) and (4.43), we obtain

$$\mathbb{E}\left[|(X(t,x;t+h) - Z(t,x;t+h))|^{2p}\right]
= \mathbb{E}\left[|(X(t,x;t+h) - Z_E(t,x;t+h) + Z_E(t,x;t+h) - Z(t,x;t+h))|^{2p}\right]
\leq \mathbb{E}\left[|((X(t,x;t+h) - Z_E(t,x;t+h))|^{2p}\right] + \mathbb{E}\left[|(Z_E(t,x;t+h) - Z(t,x;t+h))|^{2p}\right]
\leq K\left(1 + |x|^{4(\kappa+1)p+1}\right)h^{2p}.$$
(4.44)

Thus the proof is finished. \Box

Armed with Theorem 4.2, Lemmas 4.5, 4.6 and due to Theorem 2.5, one can straightforwardly obtain the long-time strong convergence rate of the scheme, presented as follows.

Theorem 4.7. Let Assumptions 2.1, 4.1 hold. For any $1 \le p \le \frac{\lfloor p^* \rfloor}{2\kappa - 1}$ and $0 < h \le h_2$, the projected Euler method (4.1) has a strong convergence rate of order $\frac{1}{2}$ over infinite time, namely,

$$\left(\mathbb{E}\left[|X_k - Z_k|^{2p}\right]\right)^{\frac{1}{2p}} \le K\left(\mathbb{E}\left[\left(1 + |X_0|^{(6\kappa+8)p}\right)\right]\right)^{\frac{1}{2p}} h^{\frac{1}{2}}.$$
(4.45)

5 Numerical Experiments

In this section, we will test the previous findings by performing numerical simulations of some examples of nonlinear SDEs.

Example 5.1. Let us consider the stochastic Ginzburg-Landau (GL) equation [15, 21], in the form

$$\begin{cases} dX(t) = \left[(\eta + \frac{1}{2}\sigma^2)X(t) - \vartheta X^3(t) \right] dt + \sigma X(t) \, dW(t), \quad t \in [0, T], \\ X(0) = X_0 > 0, \end{cases}$$
(5.1)

where $\sigma, \vartheta > 0$ and $W : [0, T] \times \Omega \to \mathbb{R}$ is the real-valued standard Brownian motions.

The coefficients are set as: $\eta = -3/2, \sigma = 1$ and $\vartheta = 1$. It is easy to see that the coefficients satisfy Assumptions 2.1 and 3.1, therefore Theorems 3.9 and 4.7 are applicable here. Next, we will consider the error caused by the temporal discretization of the problem (5.1), using both the backward Euler and the projected Euler methods. In Figure 1 and Figure 2, we plot the strong approximation errors of these two numerical schemes for the SDE (5.1). Set T = 16 and use the following time step sizes: $h \in \{2^{-7}, 2^{-6}, 2^{-5}, 2^{-4}, 2^{-3}\}$, where $h = 2^{-12}$ is considered as the exact solution. We will use M = 10000 sample paths to simulate the expectation.



Figure 1: Strong convergence rate of the backward Euler method for (5.1).

From Figure 1 and Figure 2, the expected strong convergence rate of order $\frac{1}{2}$ for both the backward Euler method and the projected Euler method is numerically confirmed.

Example 5.2. Consider the following semi-linear stochastic partial differential equation (SPDE):

$$\begin{cases} du(t,x) = \left[\frac{\partial^2}{\partial x^2}u(t,x) + u(t,x) - u(t,x)^3\right] dt + g(u(t,x)) dW_t, & t \in (0,T], & x \in (0,1) \\ u(t,0) = u(t,1) = 0, & (5.2) \\ u(0,x) = u_0(x), & \end{cases}$$

where $g: \mathbb{R} \to \mathbb{R}$ and $W: [0, T] \times \Omega \to \mathbb{R}$ is the real-valued standard Brownian motions.



Figure 2: Strong convergence rate of the projected Euler method for (5.1).

Such an SPDE is usually termed as the stochastic Allen-Cahn equation. We begin by introducing a spatial discretization with a step size $\Delta x := \frac{1}{N}$ on the interval [0, 1] and denoting the discrete spatial points as $x_i = i\Delta x, i = 1, 2, \dots, N-1$. The discretization yields an SDE system:

$$dX_t = [\mathbb{A}X_t + \mathbb{F}(X_t)] dt + \mathbb{G}(X_t) dW_t, \quad t \in (0, T], \quad X_0 = x_0,$$
(5.3)

where $X_t = (X_{1,t}, X_{2,t}, \cdots, X_{N-1,t})^T := (u(t, x_1), u(t, x_2), \cdots, u(t, x_{N-1}))^T$, $\mathbb{A} \in \mathbb{R}^{(N-1) \times (N-1)}$, $x_0 = (u_0(x_1), u_0(x_2), \cdots, u_0(x_{N-1}))^T$ and

$$\mathbb{A} = K^{2} \begin{pmatrix} -2 & 1 & 0 & \cdots & 0 & 0 \\ 1 & -2 & 1 & \cdots & 0 & 0 \\ 0 & 1 & -2 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & -2 & 1 \\ 0 & 0 & 0 & \cdots & 1 & -2 \end{pmatrix},$$
$$\mathbb{F}(X) = \begin{pmatrix} f(X_{1}) \\ f(X_{2}) \\ \vdots \\ f(X_{N-1}) \end{pmatrix}, \quad \mathbb{G}(X) = \begin{pmatrix} g(X_{1}) \\ g(X_{2}) \\ \vdots \\ g(X_{N-1}) \end{pmatrix}$$

Now we turn our attention to the temporal discretization of the SDE system (5.3), using the backward Euler and projected Euler methods. For the following numerical tests, we fix $g(u) = \sin u + 1$, T = 30 and and initialize the system with $u_0(x) = 1$.

In Figure 3 and Figure 4, we plot the strong approximation errors of the two time-stepping schemes for the SDE system (5.3) with K = 4. We use the following time step-sizes: $h \in \{\frac{15}{2^{10}}, \frac{15}{2^9}, \frac{15}{2^8}, \frac{15}{2^7}, \frac{15}{2^6}\}$ and the numerical approximation with $h_{\text{exact}} = \frac{15}{2^{12}}$ is identified as the exact solution. Moreover, M = 5000 sample paths are used to approximate the expectation. From Figure 3 and Figure 4, one can tell the expected strong convergence rate of order $\frac{1}{2}$ for both the backward Euler method and the projected Euler method, which confirms the theoretical findings.



Figure 3: Strong convergence rate of the backward Euler method for (5.3).



Figure 4: Strong convergence rate of the projected Euler method for (5.3).

References

- Adam Andersson and Raphael Kruse. Mean-square convergence of the BDF2-Maruyama and backward Euler schemes for SDE satisfying a global monotonicity condition. *BIT Numerical Mathematics*, 57(1):21–53, 2017.
- [2] Wolf-Jürgen Beyn, Elena Isaak, and Raphael Kruse. Stochastic C-stability and B-consistency of explicit and implicit Euler-type schemes. *Journal of Scientific Computing*, 67(3):955–987, 2016.
- [3] Wolf-Jürgen Beyn, Elena Isaak, and Raphael Kruse. Stochastic C-stability and B-consistency of explicit and implicit Milstein-type schemes. *Journal of Scientific Computing*, 70:1042–1077, 2017.
- [4] Charles-Edouard Brehier. Approximation of the invariant distribution for a class of ergodic sdes with one-sided lipschitz continuous drift coefficient using an explicit tamed euler scheme. ESAIM: Probability and Statistics, 27:841–866, 2023.
- [5] Nicolas Brosse, Alain Durmus, Éric Moulines, and Sotirios Sabanis. The tamed unadjusted Langevin algorithm. *Stochastic Processes and their Applications*, 129(10):3638–3663, 2019.
- [6] Chuchu Chen, Jialin Hong, and Yulan Lu. Stochastic differential equation with piecewise continuous arguments: Markov property, invariant measure and numerical approximation. *Discrete and Continuous Dynamical Systems-B*, 28(1):765–807, 2022.
- [7] Arnak S Dalalyan. Theoretical guarantees for approximate sampling from smooth and logconcave densities. Journal of the Royal Statistical Society Series B: Statistical Methodology, 79(3):651–676, 2017.
- [8] Wei Fang and Michael B Giles. Adaptive Euler–Maruyama method for SDEs with nonglobally Lipschitz drift. *The Annals of Applied Probability*, 30(2):526–560, 2020.
- [9] Siqing Gan, Youzi He, and Xiaojie Wang. Tamed Runge-Kutta methods for SDEs with superlinearly growing drift and diffusion coefficients. *Applied Numerical Mathematics*, 152:379–402, 2020.
- [10] Qian Guo, Wei Liu, Xuerong Mao, and Rongxian Yue. The truncated Milstein method for stochastic differential equations with commutative noise. *Journal of Computational and Applied Mathematics*, 338:298–310, 2018.
- [11] Desmond J Higham, Xuerong Mao, and Andrew M Stuart. Strong convergence of Euler-type methods for nonlinear stochastic differential equations. SIAM journal on numerical analysis, 40(3):1041–1063, 2002.
- [12] Jialin Hong and Xu Wang. Invariant measures for stochastic nonlinear Schrödinger equations. Springer, 2019.
- [13] Martin Hutzenthaler and Arnulf Jentzen. Numerical approximations of stochastic differential equations with non-globally Lipschitz continuous coefficients. American Mathematical Soc., 2015.

- [14] Martin Hutzenthaler and Arnulf Jentzen. On a perturbation theory and on strong convergence rates for stochastic ordinary and partial differential equations with nonglobally monotone coefficients. The Annals of Probability, 48(1):53–93, 2020.
- [15] Martin Hutzenthaler, Arnulf Jentzen, and Peter E Kloeden. Strong and weak divergence in finite time of Euler's method for stochastic differential equations with non-globally Lipschitz continuous coefficients. Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences, 467(2130):1563–1576, 2011.
- [16] Martin Hutzenthaler, Arnulf Jentzen, and Peter E Kloeden. Strong convergence of an explicit numerical method for SDEs with nonglobally Lipschitz continuous coefficients. *The Annals* of Applied Probability, 22(4):1611–1641, 2012.
- [17] Martin Hutzenthaler, Arnulf Jentzen, and Xiaojie Wang. Exponential integrability properties of numerical approximation processes for nonlinear stochastic differential equations. *Mathematics of Computation*, 87(311):1353–1413, 2018.
- [18] K Itô and M Nisio. On stationary solutions of a stochastic differential equation. J. Math. Kyoto Univ, 4(1):1–75, 1964.
- [19] Cónall Kelly and Gabriel J Lord. Adaptive time-stepping strategies for nonlinear stochastic systems. IMA Journal of Numerical Analysis, 38(3):1523–1549, 2018.
- [20] Cónall Kelly, Gabriel J Lord, and Fandi Sun. Strong convergence of an adaptive time-stepping Milstein method for SDEs with monotone coefficients. *BIT Numerical Mathematics*, 63(2):33, 2023.
- [21] Peter E Kloeden and Eckhard Platen. Numerical Solution of Stochastic differential equations. Springer, 1992.
- [22] Ruilin Li, Hongyuan Zha, and Molei Tao. Sqrt (d) dimension dependence of Langevin Monte Carlo. ICLR, 2022.
- [23] Xiaoyue Li, Xuerong Mao, and George Yin. Explicit numerical approximations for stochastic differential equations in finite and infinite horizons: truncation methods, convergence in pth moment and stability. *IMA Journal of Numerical Analysis*, 39(2):847–892, 2019.
- [24] Wei Liu, Xuerong Mao, and Yue Wu. The backward Euler-Maruyama method for invariant measures of stochastic differential equations with super-linear coefficients. *Applied Numerical Mathematics*, 184:137–150, 2023.
- [25] Wei Liu and Yudong Wang. Strong convergence in the infinite horizon of numerical methods for stochastic differential equations. arXiv preprint arXiv:2307.05039, 2023.
- [26] Xuerong Mao. Stochastic differential equations and applications. Elsevier, 2007.
- [27] Xuerong Mao. The truncated Euler-Maruyama method for stochastic differential equations. Journal of Computational and Applied Mathematics, 290:370–384, 2015.

- [28] Xuerong Mao and Lukasz Szpruch. Strong convergence and stability of implicit numerical methods for stochastic differential equations with non-globally Lipschitz continuous coefficients. Journal of Computational and Applied Mathematics, 238:14–28, 2013.
- [29] Xuerong Mao and Lukasz Szpruch. Strong convergence rates for backward Euler-Maruyama method for non-linear dissipative-type stochastic differential equations with super-linear diffusion coefficients. Stochastics An International Journal of Probability and Stochastic Processes, 85(1):144–171, 2013.
- [30] Jonathan C Mattingly, Andrew M Stuart, and Desmond J Higham. Ergodicity for SDEs and approximations: locally Lipschitz vector fields and degenerate noise. *Stochastic processes and their applications*, 101(2):185–232, 2002.
- [31] Jonathan C Mattingly, Andrew M Stuart, and Michael V Tretyakov. Convergence of numerical time-averaging and stationary measures via Poisson equations. SIAM Journal on Numerical Analysis, 48(2):552–577, 2010.
- [32] Grigori N Milstein and Michael V Tretyakov. Stochastic numerics for mathematical physics, volume 39. Springer, 2004.
- [33] Andreas Neuenkirch and Lukasz Szpruch. First order strong approximations of scalar sdes defined in a domain. *Numerische Mathematik*, 128:103–136, 2014.
- [34] Chenxu Pang, Xiaojie Wang, and Yue Wu. Linear implicit approximations of invariant measures of semi-linear sdes with non-globally lipschitz coefficients. *Journal of Complexity*, 83:101842, 2024.
- [35] Sotirios Sabanis. Euler approximations with varying coefficients: the case of superlinearly growing diffusion coefficients. Annals of Applied Probability, 26(4):2083–2105, 2016.
- [36] Yang Song, Jascha Sohl-Dickstein, Diederik P Kingma, Abhishek Kumar, Stefano Ermon, and Ben Poole. Score-based generative modeling through stochastic differential equations. In the 9th International Conference on Learning Representations (ICLR), 2021.
- [37] Denis Talay. Second-order discretization schemes of stochastic differential systems for the computation of the invariant law. *Stochastics: An International Journal of Probability and Stochastic Processes*, 29(1):13–36, 1990.
- [38] Denis Talay. Stochastic Hamiltonian systems: exponential convergence to the invariant measure, and discretization by the implicit Euler scheme. *Markov Process. Related Fields*, 8(2):163–198, 2002.
- [39] Michael V Tretyakov and Zhongqiang Zhang. A fundamental mean-square convergence theorem for SDEs with locally Lipschitz coefficients and its applications. SIAM Journal on Numerical Analysis, 51(6):3135–3162, 2013.
- [40] Xiaojie Wang. Mean-square convergence rates of implicit Milstein type methods for SDEs with non-Lipschitz coefficients. Advances in Computational Mathematics, 49(3):37, 2023.

- [41] Xiaojie Wang and Siqing Gan. The tamed Milstein method for commutative stochastic differential equations with non-globally Lipschitz continuous coefficients. *Journal of Difference Equations and Applications*, 19(3):466–490, 2013.
- [42] Xiaojie Wang, Jiayi Wu, and Bozhang Dong. Mean-square convergence rates of stochastic theta methods for SDEs under a coupled monotonicity condition. *BIT Numerical Mathematics*, 60(3):759–790, 2020.
- [43] Max Welling and Yee W Teh. Bayesian learning via stochastic gradient Langevin dynamics. In Proceedings of the 28th international conference on machine learning (ICML-11), pages 681–688. Citeseer, 2011.

A Appendix

Proof of Theorem 2.5. To highlight the dependence on initialization, we denote the solution of an SDE as $X(t_0, X_0; t_0+t)$. Now, consider the error of the method $Z(t_0, X_0; t_{k+1})$ at the (k+1)-step

$$\rho_{k+1} := X(t_0, X_0; t_{k+1}) - Z(t_0, X_0; t_{k+1})
= X(t_k, X_k; t_{k+1}) - Z(t_k, Z_k; t_{k+1})
= X(t_k, X_k; t_{k+1}) - X(t_k, Z_k; t_{k+1}) + X(t_k, Z_k; t_{k+1}) - Z(t_k, Z_k; t_{k+1}).$$
(A.1)

It is apparent that the primary distinction on the right-hand side of equation (A.1) arises from the difference on the initial data at time t_k , resulting in errors in the solution at the (k + 1)th step. This can be reformulated as:

$$\mathcal{D}_{t_k, X_k, Z_k}(t_{k+1}) := X(t_k, X_k; t_{k+1}) - X(t_k, Z_k; t_{k+1})$$

$$:= \rho_k + \mathcal{R}_{t_k, X_k, Z_k}(t_{k+1})$$
(A.2)

where \mathcal{R} is given by (2.21). The second difference in equality (A.1) is the one-step error at the (k+1)-step and we denote it as

$$\mathcal{V}_{k+1} := X(t_k, Z_k; t_{k+1}) - Z(t_k, Z_k; t_{k+1}).$$
(A.3)

Let $p \ge 1$ be an integer. We have

$$\mathbb{E}\left[|\rho_{k+1}|^{2p}\right] = \mathbb{E}\left[|\mathcal{D}_{t_k,X_k,Z_k}(t_{k+1}) + \mathcal{V}_{k+1}|^{2p}\right] \\
= \mathbb{E}\left[|\mathcal{D}_{t_k,X_k,Z_k}(t_{k+1})|^2 + 2\left\langle \mathcal{D}_{t_k,X_k,Z_k}(t_{k+1}), V_{k+1}\right\rangle + |\mathcal{V}_{k+1}|^2\right]^p \\
\leq \mathbb{E}\left[|\mathcal{D}_{t_k,X_k,Z_k}(t_{k+1})|^{2p}\right] + 2p\mathbb{E}\left[|\mathcal{D}_{t_k,X_k,Z_k}(t_{k+1})|^{2p-2}\left\langle \rho_k + \mathcal{R}_{t_k,X_k,Z_k}(t_{k+1}), \mathcal{V}_{k+1}\right\rangle \right] \\
+ \widetilde{K}_1 \sum_{l=2}^{2p} \mathbb{E}\left[|\mathcal{D}_{t_k,X_k,Z_k}(t_{k+1})|^{2p-l}|\mathcal{V}_{k+1}|^l\right],$$
(A.4)

where $\widetilde{K}_1 > 0$ only depends on p. For the first term on the right-hand side of (A.4), by (2.22) we have

$$\mathbb{E}\left[|\mathcal{D}_{t_k,X_k,Z_k}(t_{k+1})|^{2p}\right] \le \mathbb{E}\left[|\rho_k|^{2p}\right] \exp(-2p\alpha_1 h).$$
(A.5)

Next, we perform a further decomposition of the second term on the right-hand side of (A.4) as follows:

$$2p\mathbb{E}[|\mathcal{D}_{t_{k},X_{k},Z_{k}}(t_{k+1})|^{2p-2}\langle\rho_{k}+\mathcal{R}_{t_{k},X_{k},Z_{k}}(t_{k+1}),\mathcal{V}_{k+1}\rangle] = 2p\mathbb{E}[|\rho_{k}|^{2p-2}\langle\rho_{k},\mathcal{V}_{k+1}\rangle] + 2p\mathbb{E}[(|\mathcal{D}_{t_{k},X_{k},Z_{k}}(t_{k+1})|^{2p-2} - |\rho_{k}|^{2p-2})\langle\rho_{k},\mathcal{V}_{k+1}\rangle] + 2p\mathbb{E}[|\mathcal{D}_{t_{k},X_{k},Z_{k}}(t_{k+1})|^{2p-2}\langle\mathcal{R}_{t_{k},X_{k},Z_{k}}(t_{k+1}),\mathcal{V}_{k+1}\rangle].$$
(A.6)

Due to the \mathcal{F}_{t_k} -measurability of ρ_k and the conditional variant of (2.30), we obtain for the first term on the right-hand side of (A.6),

$$2p\mathbb{E}[|\rho_k|^{2p-2} \langle \rho_k, \mathcal{V}_{k+1} \rangle] \le 2pC_3 \mathbb{E}[|\rho_k|^{2p-1} (1+|Z_k|^{\eta_1})] h^{q_1}.$$
(A.7)

For the second term on the right-hand side of (A.6), noting that it equals zero when p = 1, we have for $p \ge 2$,

$$2p\mathbb{E}\left[\left(|\mathcal{D}_{t_{k},X_{k},Z_{k}}(t_{k+1})|^{2p-2}-|\rho_{k}|^{2p-2}\right)\langle\rho_{k},\mathcal{V}_{k+1}\rangle\right] \\ \leq 2p\widetilde{K}_{2}\mathbb{E}\left[|\mathcal{R}_{t_{k},X_{k},Z_{k}}(t_{k+1})||\rho_{k}||\mathcal{V}_{k+1}|\sum_{l=0}^{2p-3}|\mathcal{D}_{t_{k},X_{k},Z_{k}}(t_{k+1})|^{2p-3-l}|\rho_{k}|^{l}\right],$$
(A.8)

where $\widetilde{K}_2 > 0$ only depends on p. Additionally, one can utilize \mathcal{F}_{t_k} -measurability of ρ_k and the conditional variants of (2.31), (2.22) and (2.23), along with the Hölder inequality, to derive for $p \geq 2$,

$$2p\mathbb{E}\left[\left(|\mathcal{D}_{t_{k},X_{k},Z_{k}}(t_{k+1})|^{2p-2}-|\rho_{k}|^{2p-2}\right)\langle\rho_{k},\mathcal{V}_{k+1}\rangle\right]$$

$$\leq 2p\widetilde{K}_{2}C_{2}C_{4}\mathbb{E}\left[|\rho_{k}|^{2p-1}\left(1+|X_{k}|^{2\kappa-2}+|Z_{k}|^{2\kappa-2}\right)^{\frac{1}{4}}\left(1+|Z_{k}|^{2p\eta_{2}}\right)^{\frac{1}{2p}}\right]$$

$$h^{q_{2}+\frac{1}{2}}\sum_{l=0}^{2p-3}\exp(-(2p-3-l)\alpha_{1}h)$$

$$\leq 2p\widetilde{K}_{3}\mathbb{E}\left[|\rho_{k}|^{2p-1}\left(1+|X_{k}|^{2\kappa-2}+|Z_{k}|^{2\kappa-2}\right)^{\frac{1}{4}}\left(1+|Z_{k}|^{2p\eta_{2}}\right)^{\frac{1}{2p}}\right]h^{q_{2}+\frac{1}{2}},$$
(A.9)

where $\widetilde{K}_3 > 0$ depend on \widetilde{K}_2 , C_2 and C_4 .

Moreover, one can bound the third term on the right-hand side of (A.6), by employing the conditional variants of (2.31), (2.22), (2.23), along with applying the Hölder inequality (twice):

$$2p\mathbb{E}\left[|\mathcal{D}_{t_{k},X_{k},Z_{k}}(t_{k+1})|^{2p-2} \langle \mathcal{R}_{t_{k},X_{k},Z_{k}}(t_{k+1}),\mathcal{V}_{k+1}\rangle\right]$$

$$\leq 2p\mathbb{E}\left[\left(\mathbb{E}\left[|\mathcal{D}_{t_{k},X_{k},Z_{k}}(t_{k+1})|^{2p}|\mathcal{F}_{t_{k}}\right]\right)^{\frac{2p-2}{2p}} \left(\mathbb{E}\left[\langle \mathcal{R}_{t_{k},X_{k},Z_{k}}(t_{k+1}),V_{k+1}\rangle^{p}|\mathcal{F}_{t_{k}}\right]\right)^{\frac{2}{2p}}\right]$$

$$\leq 2p\mathbb{E}\left[\left(\mathbb{E}\left[|\mathcal{D}_{t_{k},X_{k},Z_{k}}(t_{k+1})|^{2p}|\mathcal{F}_{t_{k}}\right]\right)^{\frac{2p-2}{2p}} \left(\mathbb{E}\left[|\mathcal{R}_{t_{k},X_{k},Z_{k}}(t_{k+1})|^{2p}|\mathcal{F}_{t_{k}}\right]\right)^{\frac{1}{2p}} \left(\mathbb{E}\left[|\mathcal{V}_{k+1}|^{2p}|\mathcal{F}_{t_{k}}\right]\right)^{\frac{1}{2p}}\right]$$

$$\leq 2pC_{2}C_{4}\mathbb{E}\left[|\rho_{k}|^{2p-1}\left(1+|X_{k}|^{2\kappa-2}+|Z_{k}|^{2\kappa-2}\right)^{\frac{1}{4}}\left(1+|Z_{k}|^{2p\eta_{2}}\right)^{\frac{1}{2p}}\right]h^{q_{2}+\frac{1}{2}}\exp\left(-(2p-2)\alpha_{1}h\right)$$

$$\leq 2p\widetilde{K}_{4}\mathbb{E}\left[|\rho_{k}|^{2p-1}\left(1+|X_{k}|^{2\kappa-2}+|Z_{k}|^{2\kappa-2}\right)^{\frac{1}{4}}\left(1+|Z_{k}|^{2p\eta_{2}}\right)^{\frac{1}{2p}}\right]h^{q_{2}+\frac{1}{2}},$$
(A.10)

where $\widetilde{K}_4 > 0$ depends on C_2, C_4 .

Following the conditional versions of (2.31) and (2.22) as well as the Hölder inequality, one can bound the third term on the right-hand side of the inequality (A.4) as follows:

$$\widetilde{K}_{1} \sum_{l=2}^{2p} \mathbb{E} \left[|\mathcal{D}_{t_{k},X_{k},Z_{k}}(t_{k+1})|^{2p-l} |\mathcal{V}_{k+1}|^{l} \right] \\
\leq \widetilde{K}_{1} \sum_{l=2}^{2p} \mathbb{E} \left[\left(\mathbb{E} \left[|\mathcal{D}_{t_{k},X_{k},Z_{k}}(t_{k+1})|^{2p} |\mathcal{F}_{t_{k}} \right] \right)^{\frac{2p-l}{2p}} \left(\mathbb{E} \left[|\mathcal{V}_{k+1}|^{2p} |\mathcal{F}_{t_{k}} \right] \right)^{\frac{l}{2p}} \right] \\
\leq \widetilde{K}_{1} \sum_{l=2}^{2p} (C_{4})^{l} \mathbb{E} \left[|\rho_{k}|^{2p-l} \left(1 + |Z_{k}|^{2p\eta_{2}} \right)^{\frac{l}{2p}} \right] h^{lq_{2}} \exp(-(2p-l)\alpha_{1}h) \\
\leq \widetilde{K}_{5} \sum_{l=2}^{2p} \mathbb{E} \left[|\rho_{k}|^{2p-l} \left(1 + |Z_{k}|^{2p\eta_{2}} \right)^{\frac{l}{2p}} \right] h^{lq_{2}},$$
(A.11)

where $\widetilde{K}_5 > 0$ depends on \widetilde{K}_1, C_4 . By Substituting the inequalities (A.5) to (A.11) into (A.4) and recalling that $q_1 \ge q_2 + \frac{1}{2}$, one can obtain

$$\mathbb{E}\left[|\rho_{k+1}|^{2p}\right] \leq \mathbb{E}\left[|\rho_{k}|^{2p}\right] \exp\left(-2p\alpha_{1}h\right) + 2pC_{3}\mathbb{E}\left[|\rho_{k}|^{2p-1}\left(1+|Z_{k}|^{\eta_{1}}\right)\right]h^{q_{1}} \\
+ 2p\widetilde{K}_{3}\mathbb{E}\left[|\rho_{k}|^{2p-1}\left(1+|X_{k}|^{2\kappa-2}+|Z_{k}|^{2\kappa-2}\right)^{\frac{1}{4}}h^{q_{2}+\frac{1}{2}}\left(1+|Z_{k}|^{2p\eta_{2}}\right)^{\frac{1}{2p}}\right] \\
+ 2p\widetilde{K}_{4}\mathbb{E}\left[|\rho_{k}|^{2p-1}\left(1+|X_{k}|^{2\kappa-2}+|Z_{k}|^{2\kappa-2}\right)^{\frac{1}{4}}h^{q_{2}+\frac{1}{2}}\left(1+|Z_{k}|^{2p\eta_{2}}\right)^{\frac{1}{2p}}\right] \\
+ \widetilde{K}_{5}\sum_{l=2}^{2p}\mathbb{E}\left[|\rho_{k}|^{2p-l}h^{lq_{2}}\left(1+|Z_{k}|^{2p\eta_{2}}\right)^{\frac{l}{2p}}\right].$$
(A.12)

Then using the Young inequality for (A.12), we have

$$\mathbb{E}\left[|\rho_{k+1}|^{2p}\right] \leq \mathbb{E}\left[|\rho_{k}|^{2p}\right] \exp(-2p\alpha_{1}h) + \frac{p\alpha_{1}h}{8} \mathbb{E}\left[|\rho_{k}|^{2p}\right] + \widetilde{K}\mathbb{E}\left[\left(1 + |Z_{k}|^{\eta_{1}}\right)^{2p}\right]h^{2p(q_{2}-\frac{1}{2})+1} \\
+ \frac{p\alpha_{1}h}{8}\mathbb{E}\left[|\rho_{k}|^{2p}\right] + \widetilde{K}\mathbb{E}\left[\left(1 + |X_{k}|^{2\kappa-2} + |Z_{k}|^{2\kappa-2}\right)^{\frac{p}{2}}\left(1 + |Z_{k}|^{2p\eta_{2}}\right)\right]h^{2p(q_{2}-\frac{1}{2})+1} \\
+ \frac{\sum_{l=2}^{2p}\frac{2p-l}{2p}}{8}\alpha_{1}h\mathbb{E}\left[|\rho_{k}|^{2p}\right] + \widetilde{K}\mathbb{E}\left[\left(1 + |Z_{k}|^{2p\eta_{2}}\right)\right]h^{2p(q_{2}-\frac{1}{2})+1},$$
(A.13)

where $\widetilde{K} > 0$ is independent of h, t.

Given the assumption that $0 < h \le \frac{1}{p\alpha_1}$ and the inequality $e^{-x} \le 1 - x + \frac{x^2}{2}$ for 0 < x < 1, we can proceed by utilizing the inequalities $\sum_{l=2}^{2p} \frac{2p-l}{2p} \le p$, (2.12) and (2.32) and obtain

$$\mathbb{E}\left[|\rho_{k+1}|^{2p}\right] \leq \left(1 - \frac{p}{2}\alpha_{1}h + \frac{2p}{8}\alpha_{1}h + \frac{p}{8}\alpha_{1}h\right)\mathbb{E}\left[|\rho_{k}|^{2p}\right] + \widetilde{K}\mathbb{E}\left[\left(1 + |Z_{k}|^{2p\eta_{1}}\right)\right]h^{2p(q_{2}-\frac{1}{2})+1} \\
+ \widetilde{K}\mathbb{E}\left[\left(1 + |X_{k}|^{2\kappa-2} + |Z_{k}|^{2\kappa-2}\right)^{\frac{p}{2}}\left(1 + |Z_{k}|^{2p\eta_{2}}\right)\right]h^{2p(q_{2}-\frac{1}{2})+1} \\
\leq \left(1 - \frac{p}{8}\alpha_{1}h\right)\mathbb{E}\left[|\rho_{k}|^{2p}\right] + \widetilde{K}\mathbb{E}\left[\left(1 + |X_{0}|^{2\lambda p}\right)\right]h^{2p(q_{2}-\frac{1}{2})+1},$$
(A.14)

where $\lambda = \max\{\eta_1\eta_3, (\frac{\kappa-1}{2} + \eta_2)\eta_3\}$. Equipped with the above inequality, one sees

$$\mathbb{E}[|\rho_k|^{2p}] \le C \mathbb{E}[(1+|X_0|^{2\lambda p})] h^{2p(q_2-\frac{1}{2})},$$
(A.15)

where the constant C does not depend on h, t, k. Namely,

$$\left(\mathbb{E}\left[|X(t_0, X_0; t_k) - Z(t_0, X_0; t_k)|^{2p}\right]\right)^{\frac{1}{2p}} \le C\left(\mathbb{E}\left[\left(1 + |X_0|^{2\lambda p}\right)\right]\right)^{\frac{1}{2p}} h^{q_2 - \frac{1}{2}}.$$
(A.16)

To conclude, Theorem 2.5 is proved for integer p. For non-integer p, the conclusion can be derived using the Young inequality. Therefore, the proof is completed. \Box