# Isometric Euclidean submanifolds with isometric Gauss maps

M. Dajczer, M. I. Jimenez and Th. Vlachos

#### Abstract

We investigate isometric immersions  $f: M^n \to \mathbb{R}^{n+2}, n \geq 3$ , of Riemannian manifolds into Euclidean space with codimension two that admit isometric deformations that preserve the metric of the Gauss map. In precise terms, the preservation of the third fundamental form of the submanifold must be ensured throughout the deformation. For minimal isometric deformations of minimal submanifolds this is always the case. Our main result is of a local nature and states that if f is neither minimal nor reducible, then it is a hypersurface of an isometrically deformable hypersurface  $F: \tilde{M}^{n+1} \to \mathbb{R}^{n+2}$  such that the deformations of F induce those of f. Moreover, for a particular class of such submanifolds, a complete local parametric description is provided.

To what extent is a submanifold of Euclidean space locally determined by the properties of its Gauss map? To lend clarity to this query, let  $f: M^n \to \mathbb{R}^{n+p}$  be an isometric immersion of a Riemannian manifold of dimension  $n \geq 2$  into Euclidean space with codimension p. The Gauss map of f assigns to each point  $x \in M^n$  its tangent space  $f_*T_xM$  seen as a vector subspace of  $\mathbb{R}^{n+p}$ . Thus it is a map  $\phi_f: M^n \to Gr_n(\mathbb{R}^{n+p})$  into the Grassmannian of non-oriented n-dimensional vector subspaces of  $\mathbb{R}^{n+p}$ . Now suppose that there exists a non-congruent isometric immersion  $g: M^n \to \mathbb{R}^{n+p}$  such that the Gauss maps  $\phi_f$  and  $\phi_g$  of f and g are constrained by some predetermined metric condition. To address the initial question, one must determine the extent to which f is governed by the requested metric demand.

Certainly, the simplest requirement is to ask the Gauss maps  $\phi_f$  and  $\phi_g$  to be congruent. In simpler terms, this means assuming that f and g have the same Gauss map. A complete answer to this question was given by Dajczer and Gromoll in [4]. Roughly speaking, it was shown that f has to be a minimal non-holomorphic isometric immersion of a Kaehler manifold and that g must be any element in its one-parameter associated family of minimal isometric immersions.

Key words and phrases. Euclidean submanifold, isometric Gauss map.

Mathematics Subject Classification: 53B25, 53C24, 53C42

Let  $Gr_n(\mathbb{R}^{n+p})$  be endowed with the standard metric as a symmetric space. We recall from [10] that the pullback of that metric is given by the third fundamental form  $III_f$  of f. In terms of the second fundamental form  $\alpha_f \colon T_x M \times T_x M \to N_f M(x)$  of fat  $x \in M^n$  and the corresponding shape operators we have

$$III_f(X,Y)(x) = \sum_{i=1}^n \langle \alpha_f(X,X_i), \alpha_f(Y,X_i) \rangle = \sum_{j=1}^p \langle A_{\xi_j}^2 X, Y \rangle, \tag{1}$$

where  $\{X_i\}_{1 \le i \le n}$  and  $\{\xi_j\}_{1 \le j \le p}$  are orthonormal basis of  $T_x M$  and  $N_f M(x)$ .

A quite weaker condition is to require the submanifolds  $f, g: M^n \to \mathbb{R}^{n+p}$  to have isometric Gauss maps. That means that the graphs  $M^n \to M^n \times Gr_n(\mathbb{R}^{n+p})$  of  $\phi_f$  and  $\phi_g$  are isometric. That is, the submanifolds share identical third fundamental forms. This paper is devoted to examining the scenario that emerges under this condition.

In the case of hypersurfaces, namely, for codimension p = 1, a complete answer was obtained in [5]. It was shown that the submanifold has to be a minimal Sbrana-Cartan hypersurface. We recall that Sbrana-Cartan hypersurfaces are those of rank two, that is, with precisely two non-zero principal curvatures that allow isometric deformations. For the parametric classification of these hypersurfaces in space forms, we refer to [3]as well as to Chapter 11 of [7] for additional information relevant to this paper. The case of surfaces with codimension p = 2 was investigated by Vlachos [11]. This paper is dedicated to exploring the case of submanifolds of higher dimension than two but still in codimension p = 2. As already evidenced by the findings for surfaces in [11], preserving the metric of the Gauss map is quite a weak assumption. Thus, if the goal is to achieve parametric classifications, it becomes quite imperative to introduce additional assumptions. On the other hand, there are plenty of known examples. For instance, it is quite easy to demonstrate that any minimal submanifold allowing minimal isometric deformations, of which there are many, possesses this property. Of course, cylinders constructed over such submanifolds are also examples. Other quite obvious examples go as follows: Let  $f_1: L^p \to \mathbb{R}^{p+1}$  be a minimal Sbrana-Cartan hypersurface. Given any hypersurface  $h: N^{n-p} \to \mathbb{R}^{n-p+1}$  let  $f = h \times f_1: N^{n-p} \times L^p \to \mathbb{R}^{n+2}$  be the extrinsic product of immersions. The isometric deformations to be considered are given by the ones of the Sbrana-Cartan hypersurface.

Before discussing a more elaborate class of examples, we recall that the *relative nullity* vector subspace  $\Delta_0(x) \subset T_x M$  at  $x \in M^n$  of an isometric immersion  $f: M^n \to \mathbb{R}^N$  is the kernel of its second fundamental form at that point, that is,

$$\Delta_0(x) = \{ X \in T_x M \colon \alpha_f(X, Y) = 0 \text{ for all } Y \in T_x M \}$$

Also recall that on any open subset of  $M^n$  where  $\nu_f(x) = \dim \Delta_0(x)$  is constant, then  $x \in M^n \to \Delta_0(x)$  is an integrable distribution whose leaves are totally geodesic submanifolds in  $M^n$  that are mapped by f onto open subsets of  $\nu_f$ -dimensional affine vector subspaces of  $\mathbb{R}^N$ .

Let  $F: \tilde{M}^{n+1} \to \mathbb{R}^{n+2}$  be a minimal Sbrana-Cartan hypersurface and then let  $G: \tilde{M}^{n+1} \to \mathbb{R}^{n+2}$  be an element of its associated one-parameter family of isometric minimal deformations. Let  $j: M^n \to \tilde{M}^{n+1}$  be an isometric embedding whose unit normal vector field  $\eta \in \Gamma(N_j M)$  at any point lies in the common relative nullity distribution of F and G. It is not difficult to verify that the submanifolds  $f = F \circ j: M^n \to \mathbb{R}^{n+2}$  and  $g = G \circ j: M^n \to \mathbb{R}^{n+2}$  have isometric Gauss maps. The following family of examples, fitting into this scenario, are particularly relevant for this paper.

**Examples**: Let  $f_1: L^p \to \mathbb{S}^{p+1} \subset \mathbb{R}^{p+2}$  be a minimal hypersurface of rank two in the unit sphere. Then let  $h: N^{n-p} \to \mathbb{R}^{n-p+1}$ ,  $2 \leq p \leq n-1$ , be an immersion such that  $h = (h_0, h_1)$  with  $h_1 > 0$  for a given orthogonal splitting  $\mathbb{R}^{n-p+1} = \mathbb{R}^{n-p} \oplus \mathbb{R}$ . Let  $M^n$  be the warped product manifold  $M^n = N^{n-p} \times_{h_1} L^p$  and let the isometric immersion  $f: M^n \to \mathbb{R}^{n-p} \oplus \mathbb{R}^{p+2} = \mathbb{R}^{n+2}$  be the warped product of h and  $\iota \circ f_1$ , where  $\iota: \mathbb{S}^{p+1} \to \mathbb{R}^{p+2}$  denotes the inclusion. That is, we have that  $f = (h_0, h_1 f_1)$ . Given an isometric deformation  $g_1: L^p \to \mathbb{S}^{p+1}$  of  $f_1$  then the submanifold  $g = (h_0, h_1 g_1)$  is an isometric deformation of f such that both submanifolds have isometric Gauss maps.

If  $(\eta_0, \eta_1)$  is a unit normal vector field of h then  $\eta = (\eta_0, \eta_1 f_1)$  is a normal vector field to f. Moreover, if  $\xi$  is a unit normal vector field to the minimal immersion  $f_1$ , then seen in  $\mathbb{R}^{n+2}$  it is also normal to f. Now consider the hypersurface  $F: M^n \times (-\epsilon, \epsilon) \to \mathbb{R}^{n+2}$ , for some  $\epsilon > 0$ , defined by  $F(x,t) = f(x) + t\eta(x)$ . Then F is a minimal hypersurface of constant rank two having  $\xi(x,t) = \xi(x)$  as Gauss map. Moreover, the similarly defined minimal hypersurface  $G: M^n \times (-\epsilon, \epsilon) \to \mathbb{R}^{n+2}$  is isometric to F. Thus F and G belong to a one-parameter family of isometric Sbrana-Cartan hypersurfaces.

Before stating the result of this paper, we recall some facts from the Sbrana-Cartan classification theory of isometrically deformable hypersurfaces.

A hypersurface  $F: N^m \to \mathbb{R}^{m+1}$ ,  $m \geq 3$ , is called *surface-like* if it is a cylinder over either a surface in  $\mathbb{R}^3$  or the cone of a surface in  $\mathbb{S}^3 \subset \mathbb{R}^4$ . According to the Sbrana-Cartan theory, any deformation of F is given by an isometric deformation of the surface in either  $\mathbb{R}^3$  or  $\mathbb{S}^3$ , depending on the case.

The Sbrana-Cartan hypersurface  $F: N^m \to \mathbb{R}^{m+1}$ ,  $m \geq 3$ , is called of *elliptic type* if it is not surface-like and the associated tensor  $J \in \Gamma(\text{End}(\Delta^{\perp}))$  satisfies  $J^2 = -I$ ; for details see [3] or [7]. According to the Sbrana-Cartan theory, these hypersurfaces can belong either to the discrete class or to the continuous class. The submanifolds in the former admit a single isometric deformation whereas the ones in the latter a smooth one parameter family of isometric deformations.

In the following result, that  $f: M^n \to \mathbb{R}^{n+2}$  is *locally extrinsically irreducible* means that there is no open subset  $U \subset M^n$  splitting as a Riemannian product of manifolds  $U = U_1 \times U_2$  such that  $f|_U$  splits extrinsically as  $f|_U = f_1 \times f_2$  where  $f_i: U_i \to \mathbb{R}^{n_i}$ with  $n+2 = n_1 + n_2$ . Note that this assumption excludes cylinders. Moreover, that fand g extend isometrically means that there is an isometric embedding  $j: M^n \to N^{n+1}$  into a Riemannian manifold  $N^{n+1}$  and two isometric immersions  $F: N^{n+1} \to \mathbb{R}^{n+2}$  and  $G: N^{n+1} \to \mathbb{R}^{n+2}$  such that  $f = F \circ j$  and  $g = G \circ j$ .

**Theorem 1.** Let  $f: M^n \to \mathbb{R}^{n+2}$ ,  $n \geq 3$ , be a locally extrinsically irreducible and nowhere minimal isometric immersion of a Riemannian manifold free of flat points. Let  $g: M^n \to \mathbb{R}^{n+2}$  be an isometric immersion, not congruent to f when restricted to any open subset of  $M^n$ , such that f and g have isometric Gauss maps. Then along any connected component of an open and dense subset of  $M^n$ , we have the following:

(i) The submanifolds f, g extend uniquely to isometric non-congruent Sbrana-Cartan hypersurfaces  $F, G: \tilde{M}^{n+1} \to \mathbb{R}^{n+2}$  that are either surface-like or are of elliptic type. In the latter case, either g is unique or is any element within the one-parameter family of isometric immersions determined by the isometric deformations of an elliptic Sbrana-Cartan hypersurface of continuous class.

(ii) If the extension  $F: \tilde{M}^{n+1} \to \mathbb{R}^{n+2}$  is a minimal hypersurface, then either

- (a) f is any submanifold given in Examples that satisfies the above assumptions. Therefore, it is the extrinsic warped product immersion of  $h: N^{n-p} \to \mathbb{R}^{n-p+1}$ and  $\iota \circ f_1: L^p \to \mathbb{R}^{p+2}$  where  $f_1: L^p \to \mathbb{S}^{p+1}$  is rank two minimal hypersurface, or
- (b)  $f = \iota \circ f_1$  where  $f_1 \colon M^n \to \mathbb{S}^{n+1}(r)$  is a minimal hypersurface of rank two,

and similarly for g in both cases. Moreover, the hypersurface F is the cylinder over the cone of  $f_1: L^p \to \mathbb{S}^{p+1}$  or just the cone, and similar for G.

Note that the above result shows that only some Sbrana-Cartan hypersurfaces carry a hypersurface whose Gauss map remains isometric throughout the deformations.

The submanifolds in part (ii) can be characterized as the ones having parallel normalized mean curvature vector fields. By normalized we mean dividing the mean curvature vector field by its norm to render it a unit vector field.

In the terminology introduced in [1] the isometric deformations considered in the theorem are not genuine. This means that f and g indeed possess isometric extensions, and these extensions dictate the deformations; see [7] for additional information. The conclusion of the theorem does not hold for minimal immersions. For instance, genuine deformations are exemplified in [6] and [8] whereas for non-genuine examples we refer to Theorem 5 in [2]. This explains why these submanifolds have been excluded from the theorem's statement.

## 1 The proof

Let  $f, g: M^n \to \mathbb{R}^{n+p}$  be isometric immersions. The Gauss equation

$$\langle R(X,Y)Z,W\rangle = \langle \alpha_f(X,W), \alpha_f(Y,Z)\rangle - \langle \alpha_f(X,Z), \alpha_f(Y,W)\rangle \text{ for } X,Y,Z,W \in \mathfrak{X}(M)$$

for f together with the one for g give that

$$n\langle \alpha_f(X,Y), H_f \rangle - III_f(X,Y) = \operatorname{Ric}(X,Y) = n\langle \alpha_g(X,Y), H_g \rangle - III_g(X,Y).$$
(2)

Here Ric stands for the Ricci curvature tensor of  $M^n$  whereas  $H_f$  and  $H_g$  denote the mean curvature vector fields of f and g, respectively.

Assume that  $f, g: M^n \to \mathbb{R}^{n+p}$  have isometric Gauss maps. Then (2) gives

$$\langle \alpha_f(X,Y), H_f \rangle = \langle \alpha_g(X,Y), H_g \rangle \text{ for any } X, Y \in \mathfrak{X}(M).$$
 (3)

In particular, the mean curvature vector fields satisfy  $||H_f|| = ||H_g||$ .

In what follows, we restrict ourselves to isometric immersions  $f, g: M^n \to \mathbb{R}^{n+2}$ with isometric Gauss maps. We have from (2) that this condition holds true when the submanifolds are both minimal. Therefore, we proceed further under the assumption that they are non-minimal at any point.

In the sequel, let  $\{\eta, \xi\} \subset \Gamma(N_f M)$  and  $\{\bar{\eta}, \bar{\xi}\} \subset \Gamma(N_g M)$  be local orthonormal normal frames where  $\eta \in \Gamma(N_f M)$  and  $\bar{\eta} \in \Gamma(N_g M)$  are vector fields collinear with  $H_f$ and  $H_g$ , respectively. It follows from (3) that the shape operators for f and g satisfy

$$A_{\eta} = \bar{A}_{\bar{\eta}} \tag{4}$$

and

$$\operatorname{tr} A_{\xi} = \operatorname{tr} \bar{A}_{\bar{\xi}} = 0. \tag{5}$$

Moreover, from (1) and (4) it follows that

$$A_{\xi}^2 = \bar{A}_{\bar{\xi}}^2. \tag{6}$$

To establish the validity of the theorem, we will require the lemmas presented next.

**Lemma 2.** Let  $f: M^n \to \mathbb{R}^{n+2}$  be a nowhere minimal isometric immersion free of flat points, and let  $g: M^n \to \mathbb{R}^{n+2}$  be an isometric immersion such that f and g have isometric Gauss maps. Assume that either  $A_{\xi} = \bar{A}_{\bar{\xi}} = 0$  or that  $0 \neq A_{\xi} = \pm \bar{A}_{\bar{\xi}}$  at any point of  $M^n$ . Then f and g are congruent submanifolds.

*Proof:* If  $A_{\xi} = \bar{A}_{\bar{\xi}} = 0$ , and considering that  $A_{\eta} = \bar{A}_{\bar{\eta}}$  with rank  $A_{\eta} \ge 2$  by assumption, then the result follows from Corollary 2.2, Proposition 2.9 in [7] and the so called Fundamental theorem of submanifolds; for the latter see Theorem 1.10 in [7].

Assume that  $0 \neq A_{\xi} = \pm \bar{A}_{\bar{\xi}}$ . Then the map  $\varphi \colon N_f M \to N_g M$  defined by  $\varphi \eta = \bar{\eta}$  and  $\varphi \xi = \pm \bar{\xi}$  is a vector bundle isometry that preserves the second fundamental form. It is then an elementary fact (cf. Lemma 4.16 in [7]) that  $\varphi$  is parallel, that is, it preserves the normal connections, and the result follows again from the Fundamental theorem of submanifolds.

From the Gauss equation and (4) we have that

$$A_{\xi}X \wedge A_{\xi}Y = \bar{A}_{\bar{\xi}}X \wedge \bar{A}_{\bar{\xi}}Y \text{ for any } X, Y \in \mathfrak{X}(M).$$
(7)

Equivalently, the symmetric bilinear map  $\beta \colon T_x M \times T_x M \to \mathbb{R}^{1,1}$  defined by

$$\beta(X,Y) = (\langle A_{\xi}X,Y \rangle, \langle \bar{A}_{\bar{\xi}}X,Y \rangle) \text{ for any } X,Y \in \mathfrak{X}(M)$$

is a flat bilinear form. Here  $\mathbb{R}^{1,1}$  is  $\mathbb{R}^2$  endowed with the Lorentzian inner product

$$\langle \langle (a,b), (c,d) \rangle \rangle = ac - bd$$

and  $\beta$  being flat means that

$$\langle\langle \beta(X,W), \beta(Y,Z) \rangle\rangle = \langle\langle \beta(X,Z), \beta(Y,W) \rangle\rangle$$
 for all  $X, Y, Z, W \in \mathfrak{X}(M)$ 

The bilinear map  $\beta$  is said to be null if

$$\langle \langle \beta(X, W), \beta(Y, Z) \rangle \rangle = 0$$
 for all  $X, Y, Z, W \in \mathfrak{X}(M)$ .

It is simple to check that  $\beta$  is null if and only if  $A_{\xi} = \pm \bar{A}_{\xi}$ .

**Lemma 3.** Assume that  $\beta$  at  $x \in M^n$  is not null. Then the vector subspace

$$\mathcal{N}(\beta)(x) = \{ X \in T_x M \colon \beta(X, Y) = 0 \text{ for any } Y \in T_x M \}$$

satisfies  $\dim(\mathcal{N}(\beta)(x)) \ge n-2$ .

*Proof:* Since  $\beta$  is not null, we have that span{ $\beta(X, Y): X, Y \in \mathfrak{X}(M)$ } =  $\mathbb{R}^{1,1}$ . Then the claim follows from Lemma 4.20 in [7].

**Lemma 4.** Let  $f, g: M^n \to \mathbb{R}^{n+2}$  be nowhere minimal isometric immersions with isometric Gauss maps. Assume that they are not congruent when restricted to any open subset of  $M^n$ . If  $M^n$  is free of flat points then either  $A_{\xi} = \bar{A}_{\bar{\xi}} = 0$  or  $\Delta = \ker A_{\xi} = \ker \bar{A}_{\bar{\xi}}$ has dimension n-2.

*Proof:* By Lemma 2 the flat bilinear form  $\beta$  is not null. Then Lemma 3 yields that  $\dim(\mathcal{N}(\beta)) \geq n-2$ . Hence  $\mathcal{N}(\beta) \subset \ker A_{\xi} \cap \ker \bar{A}_{\bar{\xi}}$ . Thus (5) and (6) give for  $A_{\xi}$  and  $\bar{A}_{\bar{\xi}}$  that either both vanish or that the common kernel  $\Delta$  has dimension n-2.

Proof of Theorem 1: Let the connection forms  $\psi, \bar{\psi} \in \Gamma(TM^*)$  for f and g be given by

$$\nabla_X^{\perp}\eta = \psi(X)\xi$$
 and  $\bar{\nabla}_X^{\perp}\bar{\eta} = \bar{\psi}(X)\bar{\xi}$ ,

where  $\nabla^{\perp}$  and  $\bar{\nabla}^{\perp}$  are the respective normal connections. The Codazzi equation

$$(\nabla_X A_\eta)Y - A_{\nabla_X^{\perp}\eta}Y = (\nabla_Y A_\eta)X - A_{\nabla_X^{\perp}\eta}X$$

and (4) yield that

$$\psi(X)A_{\xi}Y - \psi(Y)A_{\xi}X = \bar{\psi}(X)\bar{A}_{\bar{\xi}}Y - \bar{\psi}(Y)\bar{A}_{\bar{\xi}}X \text{ for any } X, Y \in \mathfrak{X}(M).$$
(8)

According to Lemma 4 we consider two cases. First suppose that  $A_{\xi} = \bar{A}_{\bar{\xi}} = 0$  along an open connected subset  $U \subset M^n$ . Since  $A_{\eta} = \bar{A}_{\bar{\eta}}$  and rank  $A_{\eta} \geq 2$ , following the arguments used in the proof of Lemma 2, it follows that the submanifolds  $f|_U$  and  $g|_U$ are contained in affine hyperplanes of  $\mathbb{R}^{n+2}$ , and thus are congruent in  $\mathbb{R}^{n+2}$ . But this has been ruled out.

In the sequel, we assume that f and g are restricted to a connected component of the open and dense subset where rank  $A_{\xi} = \operatorname{rank} \bar{A}_{\bar{\xi}} = 2$ , with  $\Delta = \ker A_{\xi} = \ker \bar{A}_{\xi}$ , and that one of the following cases of consideration holds.

Case  $\psi = 0$  and  $\bar{\psi} \neq 0$ . This case is not possible. In fact, from (8) we have

$$\bar{\psi}(X)\bar{A}_{\bar{\xi}}Y = \bar{\psi}(Y)\bar{A}_{\bar{\xi}}X$$
 for any  $X, Y \in \mathfrak{X}(M)$ .

But then rank  $\bar{A}_{\bar{\xi}} < 2$ , which is a contradiction.

Case  $\psi \neq 0 \neq \overline{\psi}$ . First suppose that ker  $\psi = \ker \overline{\psi}$ . If  $X \in \Gamma((\ker \psi)^{\perp})$  and  $Y \in \Gamma(\Delta^{\perp})$  are orthogonal then (8) yields

$$\psi(X)A_{\xi}Y = \bar{\psi}(X)\bar{A}_{\bar{\xi}}Y.$$
(9)

By (7) we have that  $A_{\xi}$  and  $\bar{A}_{\bar{\xi}}$  restricted to  $\Delta^{\perp}$  have the same determinant. This together with (5) and (9) give that  $A_{\xi} = \pm \bar{A}_{\bar{\xi}}$ , and Lemma 2 yields a contradiction.

Therefore, we have that  $\ker \psi \neq \ker \overline{\psi}$ . Then (8) yields  $\Delta = \ker \psi \cap \ker \overline{\psi}$ . Hence, from (5) and (7) it follows that

$$\bar{A}_{\bar{\xi}} = A_{\xi} \circ R,\tag{10}$$

where R is an isometry of TM acting as the identity on  $\Delta$ .

For simplicity, henceforth we denote the restriction of any tensor to  $\Delta^{\perp}$  in the same manner. Replacing (10) in (8) gives

$$\psi(X)A_{\xi}Y - \psi(Y)A_{\xi}X = \bar{\psi}(X)A_{\xi}RY - \bar{\psi}(Y)A_{\xi}RX \text{ for any } X, Y \in \mathfrak{X}(M).$$
(11)

Let  $\{X, Y\}$  be an orthonormal frame of eigenvectors of  $A_{\xi}$  in  $\Gamma(\Delta^{\perp})$  associated to the eigenvalues  $\pm \lambda$ , respectively. With respect to this frame set

$$R = \begin{bmatrix} \cos \gamma & -\sin \gamma \\ \sin \gamma & \cos \gamma \end{bmatrix} \text{ where } \gamma \in C^{\infty}(M).$$
(12)

Then we obtain from (11) and (12) that

$$\bar{\psi} = \psi \circ R. \tag{13}$$

Set  $N_f M = L \oplus P$  where  $L = \operatorname{span}\{\xi\}$  and  $P = \operatorname{span}\{\eta\}$ . Then let

$$\Omega(x) = \operatorname{span}\{(\tilde{\nabla}_X \delta)_{f_*TM \oplus P} \colon X \in T_x M \text{ and } \delta \in \Gamma(L)\}.$$
(14)

Since the vector field  $\xi$  is constant along  $\Delta$ , then  $\Omega$  is a smooth vector subbundle of  $f_*\Delta^{\perp} \oplus P$  of rank two. Let  $\Lambda$  be the line vector bundle defined by the orthogonal splitting  $f_*\Delta^{\perp} \oplus P = \Omega \oplus \Lambda$ . Then let  $0 \neq X_0 \in \Gamma(\Delta^{\perp})$  be given by  $f_*X_0 + \eta \in \Gamma(\Lambda)$ , that is, it satisfies that

$$0 = \langle \tilde{\nabla}_X (f_* X_0 + \eta), \xi \rangle = \langle A_\xi X, X_0 \rangle + \psi(X) \text{ for any } X \in \Gamma(\Delta^{\perp}).$$
(15)

Using (10), (13) and (15) we obtain that  $g_*X_0 + \bar{\eta}$  is orthogonal to  $\tilde{\nabla}_X \bar{\xi}$  for any  $X \in \Delta^{\perp}$ . Hence also

$$\langle \bar{A}_{\bar{\xi}}X, X_0 \rangle + \bar{\psi}(X) = 0 \text{ for any } X \in \Gamma(\Delta^{\perp}).$$
 (16)

The maps  $F, G \colon \tilde{M}^{n+1} = M^n \times (-\epsilon, \epsilon) \to \mathbb{R}^{n+2}$  given by

$$F(x,t) = f(x) + t(f_*X_0 + \eta) \text{ and } G(x,t) = g(x) + t(g_*X_0 + \bar{\eta})$$
(17)

define for small  $\epsilon > 0$  non-congruent isometric hypersurfaces of rank two. The fact that they are isometric follows using (15) and (16). Note that the unit vector fields  $\xi(x,t) = \xi(x)$  and  $\overline{\xi}(x,t) = \overline{\xi}(x)$  are the Gauss maps of F and G, respectively. Then  $\Delta \oplus \text{span}\{\partial_t\}$  are the relative nullity subspaces of both immersions.

The subsequent computations are done along f and g, that is, for t = 0. First we construct orthonormal frames of  $\Omega$  and  $\overline{\Omega}$  which are similarly defined. Let  $Y_1 \in \Gamma(\Delta^{\perp})$  be a unit norm vector field orthogonal to  $X_0$ . If  $\rho \in C^{\infty}(M)$  is given by  $\rho^2 = 1 + 1/||X_0||^2$ , then the vector field  $Z_1 = (1/\rho)((1-\rho^2)X_0 + \eta)$  together with  $Y_1$  form an orthonormal basis of  $\Omega$ . Also  $\{Y_1, \overline{Z}_1\} \in \overline{\Omega}$  is an orthonormal frame, where  $\overline{Z}_1$  is similarly defined in terms of  $\overline{\eta}$ . If  $A^F$  and  $A^G$  denote the shape operators of F and G, respectively, then

$$A^{F}X = A_{\xi}X + \psi(X)\eta \text{ and } A^{G}X = \bar{A}_{\bar{\xi}}X + \bar{\psi}(X)\bar{\eta} \text{ for any } X \in \Gamma(\Delta^{\perp}).$$
(18)

Setting  $X_1 = X_0 / ||X_0||$ , we obtain from (15) and (16) that

$$\langle A_{\xi}X_1, X_1 \rangle = -\|X_0\|^{-1}\psi(X_1) \text{ and } \langle \bar{A}_{\bar{\xi}}X_1, X_1 \rangle = -\|X_0\|^{-1}\bar{\psi}(X_1).$$
 (19)

Then (5) gives that

$$\langle A_{\xi}Y_1, Y_1 \rangle = \|X_0\|^{-1}\psi(X_1) \text{ and } \langle \bar{A}_{\bar{\xi}}Y_1, Y_1 \rangle = \|X_0\|^{-1}\bar{\psi}(X_1).$$
 (20)

Using (15), (16) and (18) it follows that

$$\langle A^F Y_1, Z_1 \rangle = \rho \psi(Y_1) \text{ and } \langle A^G Y_1, \overline{Z}_1 \rangle = \rho \overline{\psi}(Y_1).$$
 (21)

Since  $X_0 + \eta$  is contained in the common relative nullity subspaces of F and G, that is,  $A^F(X_0 + \eta) = 0 = A^G(X_0 + \eta)$ , we obtain from (15), (16) and (18) that

$$\langle A^F X, \eta \rangle = \psi(X) \text{ and } \langle A^G X, \bar{\eta} \rangle = \bar{\psi}(X) \text{ for any } X \in \Gamma(\Delta^{\perp}).$$
 (22)

Moreover, we have

$$\langle A^F Z_1, Z_1 \rangle = \frac{1}{\rho^2} \langle A^F ((1 - \rho^2) X_0 + \eta), (1 - \rho^2) X_0 + \eta \rangle$$
  
=  $-\frac{1}{\rho^2} \langle \rho^2 A^F X_0, (1 - \rho^2) X_0 + \eta \rangle$ 

and similarly for  $A^G$ . It follows using (19) and (22) that

$$\langle A^F Z_1, Z_1 \rangle = -\frac{\rho^2}{\sqrt{\rho^2 - 1}} \psi(X_1) \text{ and } \langle A^G \bar{Z}_1, \bar{Z}_1 \rangle = -\frac{\rho^2}{\sqrt{\rho^2 - 1}} \bar{\psi}(X_1).$$
 (23)

The frames  $\{Y_1, Z_1\}$  and  $\{Y_1, \overline{Z}_1\}$  coincide after identifying  $f_*TM \oplus \operatorname{span}\{\eta\}$  and  $g_*TM \oplus \operatorname{span}\{\overline{\eta}\}$  with the tangent space of  $M^n \times (\epsilon, \epsilon)$  at corresponding points. Then we have from (20), (21) and (23) that the shape operators have the expressions

$$A^{F} = \rho \begin{bmatrix} \phi \psi(X_{1}) & \psi(Y_{1}) \\ \psi(Y_{1}) & -\frac{1}{\phi} \psi(X_{1}) \end{bmatrix}, \quad A^{G} = \rho \begin{bmatrix} \phi \bar{\psi}(X_{1}) & \bar{\psi}(Y_{1}) \\ \bar{\psi}(Y_{1}) & -\frac{1}{\phi} \bar{\psi}(X_{1}) \end{bmatrix}$$
(24)

where  $\phi = \sqrt{\rho^2 - 1}/\rho$ . Since we can choose orientation of the frames such that R has the expression (12), we obtain using (13) that

$$A^G = A^F(\cos\gamma I + \sin\gamma J), \tag{25}$$

where J with respect to  $\{Y_1, Z_1\}$  is given by

$$J = \begin{bmatrix} 0 & -1/\phi \\ \phi & 0 \end{bmatrix}.$$

Since  $J^2 = -I$ , then either F and G are both surface-like or are isometric elliptic Sbrana-Cartan hypersurfaces.

Note that the second fundamental form of the surfaces involved in the surface-like case are as in (24) and satisfy the condition (25). In particular, they have negative extrinsic Gauss curvature.

Case  $\psi = \overline{\psi} = 0$ . The submanifolds have flat normal bundle and the shape operators associated to the vector fields  $\xi$  and  $\eta$  are Codazzi tensors, namely, they satisfy

$$(\nabla_X A_{\xi})Y = (\nabla_Y A_{\xi})X \text{ for any } X, Y \in \mathfrak{X}(M)$$
(26)

and

$$(\nabla_X A_\eta) Y = (\nabla_Y A_\eta) X \text{ for any } X, Y \in \mathfrak{X}(M).$$
(27)

Taking the  $\Delta$ -component of (26) for  $X \in \Gamma(\Delta^{\perp})$  and identity  $T \in \Gamma(\Delta)$  yields that  $\Delta$  is a totally geodesic distribution. The Ricci equation

$$\langle R^{\perp}(X,Y)\eta,\xi\rangle = \langle [A_{\eta},A_{\xi}]X,Y\rangle$$
 for any  $X,Y \in \mathfrak{X}(M)$ 

gives that  $A_{\eta}$  and  $A_{\xi}$  commute. Similarly, we have using (4) that also  $A_{\eta}$  and  $\bar{A}_{\bar{\xi}}$  commute. It follows from (5) and (6) pointwise that  $\bar{A}_{\bar{\xi}} = A_{\xi} \circ R$  being R a rotation of angle  $\theta$  on  $\Delta^{\perp}$  whereas the identity on  $\Delta$ . Moreover, it follows from (26) that  $\theta$  is constant and that R is not the identity, up to sign, from Lemma 2. Since  $A_{\eta}$  commutes with R, then

$$A_n|_{\Delta^\perp} = bI$$
 where  $b \in C^\infty(M)$ .

Set  $D = \ker(A_{\eta} - bI)$  and notice that  $\Delta^{\perp} \subset D$ . Hence dim  $D \geq 2$  everywhere. We assume further that we are restricted to on an open subset of  $M^n$  where D possesses constant dimension. Equation (27) gives that

$$\nabla_T A_\eta Z - A_\eta \nabla_T Z = Z(b)T + b \nabla_Z T - A_\eta \nabla_Z T$$

for any  $T \in \Gamma(D)$  and  $Z \in \Gamma(D^{\perp})$ . Then the S-component for any  $S \in \Gamma(D)$  yields

$$\langle (A_{\eta} - bI)Z, \nabla_T S \rangle = -Z(b) \langle T, S \rangle.$$

It follows that the distribution D is umbilical. Let  $\delta$  be the corresponding umbilical vector field. It is easily seen that the distribution D is spherical, that is,  $(\nabla_T \delta)_{D^{\perp}} = 0$  for any  $T \in \Gamma(D)$ . For instance, see Exercise 1.17 in [7].

Let  $U \subset M^n$  be an open subset where the totally geodesic distribution  $\Delta$  has constant dimension being U the saturation by maximal leaves of some cross section of the foliation. Then the quotient space of leaves is Hausdorff and hence a two dimensional manifold  $L^2$ . Since the vector field  $\xi$  is constant along the leaves of  $\Delta$ , then it determines a surface  $h: L^2 \to \mathbb{S}^{n+1}$  in the unit sphere. From now on, we restrict ourselves to open subsets where this situation holds. Note that these subsets form an open and dense subset of  $M^n$ .

Assume that the surface  $h: L^2 \to \mathbb{S}^{n+1}$  is substantial in  $\mathbb{S}^{p+1} \subset \mathbb{S}^{n+1}$  for some  $2 \leq p \leq n-1$ . This means that p+1 is the least dimension of the totally geodesic sphere  $\mathbb{S}^{p+1}$  that contains the surface. We argue that the Euclidean vector subspace  $\mathbb{R}^{p+2}$  that contains  $\mathbb{S}^{p+1}$  is spanned by the derivatives of all order of  $\xi$  with respect to the vectors fields in  $\Gamma(\Delta^{\perp})$ . In fact, the Euclidean space  $\mathbb{R}^{p+2}$  is spanned by the derivatives of all order of  $\xi$  in the ambient space by vectors fields of  $\Gamma(TM)$ . On one hand, we have  $\tilde{\nabla}_Z \xi = 0$  for  $Z \in \Gamma(\Delta)$ . On the other hand  $\tilde{\nabla}_X \xi = -f_* A_\xi X \in \Gamma(\Delta^{\perp})$  for

 $X \in \Gamma(\Delta^{\perp})$ . Then  $\tilde{\nabla}_Z \tilde{\nabla}_X \xi \in \Gamma(\Delta^{\perp})$  since the distribution  $\Delta$  is totally geodesic. Given that the ambient space is flat, then

$$\tilde{\nabla}_Z \tilde{\nabla}_{X_2} \tilde{\nabla}_{X_1} \xi = \tilde{\nabla}_{X_2} \tilde{\nabla}_Z \tilde{\nabla}_{X_1} \xi + \tilde{\nabla}_{[Z,X_2]} \tilde{\nabla}_{X_1} \xi$$

for any  $X_1, X_2 \in \Gamma(\Delta^{\perp})$ . Hence the left hand side is spanned by derivatives of  $\xi$  along  $\Gamma(\Delta^{\perp})$ . Similarly, we reach the same conclusion from

$$\tilde{\nabla}_Z \tilde{\nabla}_{X_k} \dots \tilde{\nabla}_{X_1} \xi = \tilde{\nabla}_{X_k} \tilde{\nabla}_Z \dots \tilde{\nabla}_{X_1} \xi + \tilde{\nabla}_{[Z,X_k]} \tilde{\nabla}_{X_{k-1}} \dots \tilde{\nabla}_{X_1} \xi.$$

for any k and  $X_i \in \Gamma(\Delta^{\perp}), 1 \leq i \leq k$ .

Assume that h is substantial in  $\mathbb{S}^{p+1} \subset \mathbb{R}^{p+2} \subset \mathbb{R}^{n+2}$  for some  $1 \leq p \leq n-1$ . We have that  $\tilde{\nabla}_X \xi = -f_* A_{\xi} X \in \Gamma(\Delta^{\perp})$  for  $X \in \Gamma(\Delta^{\perp})$ . Then the derivative with respect to  $Y \in \Gamma(\Delta^{\perp})$  gives

$$\tilde{\nabla}_Y \tilde{\nabla}_X \xi = -f_* (\nabla_Y A_\xi X)_D - \langle A_\xi X, A_\xi Y \rangle \xi - \langle A_\xi X, Y \rangle (b\eta + f_* \delta).$$
(28)

We analyze the nature of the derivatives along vectors in  $\Delta^{\perp}$  of the terms on the right hand side of (28). Since D is an umbilical distribution, we have that  $\tilde{\nabla}_X f_* Z \in$  $\Gamma(D)$  for any  $X \in \Gamma(\Delta^{\perp})$  and  $Z \in \Gamma(\Delta \cap D)$ . Therefore, the covariant derivative of the term  $f_*(\nabla_Y A_{\xi}X)_D$  along vector fields in  $\Gamma(\Delta^{\perp})$  belongs to the vector subbundle  $f_*D \oplus \text{span}\{\xi, b\eta + f_*\delta\}$ . We have from (27) that b is constant along D. Then, using that D is spherical, that  $\Delta^{\perp} \subset D$  and that b is constant along  $\Delta^{\perp}$ , it follows that consecutive derivatives of  $b\eta + f_*\delta$  with respect to vectors fields in  $\Gamma(\Delta^{\perp})$  also belong to  $f_*D \oplus \text{span}\{\xi, b\eta + f_*\delta\}$ . Since the surface h is substantial in  $\mathbb{S}^{p+1} \subset \mathbb{R}^{p+2}$ , we have seen that  $\mathbb{R}^{p+2}$  is spanned by the derivatives of all orders of  $\xi$  with respect to the vector fields in  $\Gamma(\Delta^{\perp})$ . It follows that

$$\mathbb{R}^{p+2} \subset f_* D(x) \oplus \operatorname{span}\{\xi(x), b\eta(x) + f_*\delta(x)\} \text{ at any } x \in M^n.$$
(29)

Decompose orthogonally  $\mathbb{R}^{n+2} = \mathbb{R}^{n-p} \oplus \mathbb{R}^{p+2}$  and any  $V \in \mathbb{R}^{n+2}$  as  $V = V_1 + V_2$ accordingly. Then  $\eta = \eta_1 + \eta_2$ . Suppose that  $\eta_1 = 0$  on an open subset  $M^n$ . Then  $N_f M(x) \subset \mathbb{R}^{p+2}$  and thus  $\mathbb{R}^{n-p} \subset f_* T_x M$  for any point in the open subset. Then  $\mathbb{R}^{n-p}$  conforms a tangent totally geodesic distribution contained in the relative nullity subspaces of the submanifold. But then f is a cylinder with an (n-p)-dimensional Euclidean factor on an open subset of  $M^n$ , which is ruled out by assumption.

Suppose that  $\eta_2 = 0$  on an open subset  $M^n$ . Since  $\eta$  belongs to  $\mathbb{R}^{n-p}$  this is also the case of  $\tilde{\nabla}_X \eta = -f_*A_\eta X = -bf_*X$  for any  $X \in \Gamma(\Delta^{\perp})$ . But since  $f_*\Delta^{\perp}(x) \subset \mathbb{R}^{p+2}$ , then b = 0 and hence  $\delta = 0$ . Having that  $\xi \in \mathbb{R}^{p+2}$  it now follows from (29) that  $\mathbb{R}^{p+2} =$  $f_*D_2(x) \oplus \text{span}\{\xi(x)\}$  where  $f_*D_2 = f_*TM \cap \mathbb{R}^{p+2}$  and that  $\mathbb{R}^{n-p} = f_*K(x) \oplus \text{span}\{\eta(x)\}$ where  $f_*K = f_*TM \cap \mathbb{R}^{n-p}$ . Then  $TM = D_2 \oplus K$  where  $D_2$  and K are orthogonal totally geodesic distributions. Since  $D_2 \subset D$  from (29), then  $\alpha_f(T, W) = 0$  for any  $T \in \Gamma(D_2)$ and  $W \in \Gamma(K)$ . Now we have from Theorem 8.4 in [7] that the immersion is an extrinsic product of a hypersurface  $f_1: M_1 \to \mathbb{R}^{n-p}$  and a minimal hypersurface  $f_2: M_2 \to \mathbb{R}^{p+2}$  of rank two, but this also is ruled out by assumption.

In view of the above, let  $U \subset M^n$  be a connected component of the open and dense subset where  $\eta_1 \neq 0 \neq \eta_2$ . Setting  $f_*D_2 = f_*TM \cap \mathbb{R}^{p+2}$  we have that dim  $D_2 \geq p$ . Since b = 0 implies  $\delta = 0$ , it then follows from (29) that  $D_2 \subset D$ . Since  $\eta_2$  is orthogonal to  $D_2$  then

$$\mathbb{R}^{p+2} = f_* D_2(x) \oplus \operatorname{span}\{\xi(x), \eta_2(x)\} \text{ at any } x \in U$$
(30)

and thus dim  $D_2 = p$  where  $p \ge 2$  since  $\Delta^{\perp} \subset D_2$ .

We claim that the distribution  $D_2$  is spherical. On one hand, we have from (30) that  $\tilde{\nabla}_S f_*T \in \mathbb{R}^{p+2}$  for any  $S, T \in \Gamma(D_2)$ . On the other hand, we know that

$$\tilde{\nabla}_S f_*T = f_*(\nabla_S T)_D + \langle A_{\xi}S, T \rangle \xi + \langle S, T \rangle (b\eta + f_*\delta), \text{ for any } S, T \in \Gamma(D_2).$$

Since  $\xi(x) \in \mathbb{R}^{p+2}$  then  $f_*(\nabla_S T)_D + \langle S, T \rangle (b\eta + f_*\delta) \in \mathbb{R}^{p+2}$ . Thus

$$(f_*(\nabla_S T)_D)_1 + \langle S, T \rangle (b\eta_1 + (f_*\delta)_1) = 0 \text{ for any } S, T \in \Gamma(D_2).$$
(31)

Using that

$$0 = \langle f_*(\nabla_S T)_D, \eta \rangle = \langle f_*(\nabla_S T)_D, \eta_1 \rangle + \langle f_*(\nabla_S T)_D, \eta_2 \rangle$$

we obtain from (31) that

$$\langle f_*(\nabla_S T)_D, \eta_2 \rangle = \langle S, T \rangle \langle b\eta_1 + (f_*\delta)_1, \eta_1 \rangle \text{ for any } S, T \in \Gamma(D_2).$$
 (32)

We have from (30) and (32) that

$$(f_*(\nabla_S T)_D)_2 = f_*(\nabla_S T)_{D_2} + \|\eta_2\|^{-2} \langle f_*(\nabla_S T)_D, \eta_2 \rangle \eta_2$$
  
=  $f_*(\nabla_S T)_{D_2} + \langle S, T \rangle a \eta_2$ 

where  $a = \|\eta_2\|^{-2} \langle b\eta_1 + (f_*\delta)_1, \eta_1 \rangle$ . Then using (31) it follows that

$$f_*(\nabla_S T)_D - f_*(\nabla_S T)_{D_2} = \langle S, T \rangle f_*\sigma. \text{ for any } S, T \in \Gamma(D_2),$$
(33)

where  $f_*\sigma = a\eta_2 - b\eta_1 - (f_*\delta)_1 \in \Gamma(D)$ . Being the distribution D spherical with mean curvature vector field  $\delta$ , we have from (33) that the distribution  $D_2$  is umbilical with mean curvature vector field  $\tilde{\delta} = \delta + \sigma$ .

Since  $\tilde{\nabla}_T \eta = -f_* A_\eta T = -bf_* T \in \mathbb{R}^{p+2}$  for any  $T \in \Gamma(D_2)$  then  $\tilde{\nabla}_T \eta_1 = 0$ . Hence,  $\eta_1$  is parallel in  $\mathbb{R}^{n-p}$  along  $D_2$  and then  $\|\eta_2\|$  is constant along  $D_2$ .

We have that

$$\nabla_T f_* \delta = f_* \nabla_T \delta + \langle A_\xi \delta, T \rangle \xi + \langle A_\eta T, \delta \rangle \eta$$
 for any  $T \in D_2$ ,

where  $\nabla_T \delta$  is collinear with T since  $D_2 \subset D$  and D is a spherical distribution. This gives that  $(f_*\delta)_1$  is parallel in  $\mathbb{R}^{n-p}$  along  $D_2$  and then  $||f_*\delta_2||$  is constant along  $D_2$ , where we used that  $A_{\xi}\delta = 0$  since  $\delta \in \Gamma(\Delta)$  and that  $\langle A_{\eta}T, \delta \rangle = \langle bT, \delta \rangle = 0$ . It follows from

$$0 = \langle f_*\sigma, \eta \rangle = a \|\eta_2\|^2 - b \|\eta_1\|^2 - \langle (f_*\delta)_1, \eta_1 \rangle$$

that the function a is constant along  $D_2$ . Since  $\tilde{\nabla}_T \eta = -bT$  and  $\eta_1$  is constant along  $D_2$  then  $\tilde{\nabla}_T f_* \sigma$  is a multiple of T. Having that also  $\nabla_T \delta$  is a multiple of T it follows that  $\nabla_T \tilde{\delta}$  is a multiple of T for any  $T \in \Gamma(D_2)$ . Thus the distribution  $D_2$  is spherical as claimed.

We have that  $f_*K = f_*TM \cap \mathbb{R}^{n-p}$  satisfies dim K = n-p-1. Also  $\tilde{\nabla}_T f_*W \in \mathbb{R}^{n-p}$ for any  $T \in \Gamma(D_2)$  and  $W \in \Gamma(K)$ . Thus  $\langle \nabla_T S, W \rangle = -\langle \nabla_T W, S \rangle = 0$  for  $T, S \in \Gamma(D_2)$ . Since dim D = p hence  $D_2^{\perp} = K \oplus \text{span}\{\tilde{\delta}\}$  at any point of U.

We claim that the distribution  $D_2^{\perp}$  is totally geodesic. It holds that

$$\langle \hat{\nabla}_Z \eta, f_*T \rangle = -\langle A_\eta Z, T \rangle = -b\langle Z, T \rangle = 0$$

if  $Z \in \Gamma(D_2^{\perp})$  and  $T \in \Gamma(D_2)$ . Then  $\langle \tilde{\nabla}_Z \eta_2, f_*T \rangle = 0$  for any  $Z \in \Gamma(D_2^{\perp})$  and  $T \in \Gamma(D_2)$ since  $\langle \tilde{\nabla}_Z \eta_1, f_*T \rangle = 0$  because  $\tilde{\nabla}_Z \eta_1 \in \mathbb{R}^{n-p}$ . Since  $\delta$  is orthogonal to  $D_2$  and  $\xi$ , we have from (30) that  $(f_*\delta)_2$  and  $\eta_2$  are collinear. Hence  $\langle \tilde{\nabla}_Z(f_*\delta)_2, f_*T \rangle = 0$  for any  $Z \in \Gamma(D_2^{\perp})$ and  $T \in \Gamma(D_2)$ . Since  $\tilde{\delta} = \delta + \sigma$  where  $f_*\sigma = a\eta_2 - b\eta_1 - (f_*\delta)_1$  then  $f_*\tilde{\delta} = c\eta_2 - b\eta_1$  for  $c \in C^{\infty}(U)$ . Hence we have that  $\langle \tilde{\nabla}_Z f_*\tilde{\delta}, f_*T \rangle = 0$  for any  $Z \in \Gamma(D_2^{\perp})$ . On the other hand, it is trivial that  $\langle \nabla_Y W, T \rangle = 0$  if  $W \in \Gamma(K), T \in \Gamma(D_2)$  and  $Y \in \mathfrak{X}(U)$ . We conclude that  $(\nabla_Z W)_{D_2} = 0$  for any  $Z, W \in \Gamma(D_2^{\perp})$ , and the claim has been proved.

Finally, since  $\alpha_f(T, Z) = 0$  for any  $T \in \Gamma(D_2)$  and  $Z \in \Gamma(D_2^{\perp})$ , then Theorem 10.4 in [7] and a result due to Nölker [9], which is also Theorem 10.21 in [7], gives that  $f|_U$ is part of a warped product of immersions as required.

Now assume that the surface  $h: L^2 = V/\Delta \to \mathbb{S}^{n+1}$  is substantial for an open subset  $V \subset M^n$ . From (29) we have

$$\mathbb{R}^{n+2} = D(x) \oplus \operatorname{span}\{\xi(x), b\eta(x) + \delta(x)\}$$

at any point  $x \in V$ . Then dim D = n, thus  $\delta(x) = 0$  and therefore  $A_{\eta} = bI$ . Note that  $b(x) \neq 0$  for any  $x \in V$  since f is nowhere minimal. Being  $\eta$  parallel in the normal connection, we have that f(V) is contained in an umbilical hypersurface  $\mathbb{S}^{n+1}(r) \subset \mathbb{R}^{n+2}$ . Note that  $\xi$  is a unit normal vector field to  $f|_V$  in  $\mathbb{S}^{n+1}(r)$ . It follows from (5) that  $f|_V$ is a rank two minimal hypersurface of  $\mathbb{S}^{n+1}(r)$ .

Next we argue that the submanifold g is of the same type as f. The argument is for when f is a warped product since the other case is trivial. We have seen that Uis intrinsically part of a Riemannian warped product determined by the distributions  $D_2$  and  $D_2^{\perp}$ , where the former is spherical and the latter is totally geodesic. Since  $\Delta^{\perp} \subset D_2$  and  $\Delta$  is the common kernel of  $A_{\xi}$  and  $\bar{A}_{\bar{\xi}}$ , it then follows from (4) that the second fundamental form of g also satisfies that  $\alpha_g(X, Y) = 0$  for any  $X \in \Gamma(D_2)$  and  $Y \in \Gamma(D_2^{\perp})$ . Then also g is an extrinsic warped product of immersions.

For the sake of simplicity, we continue under the assumption that f and g satisfy the requested conditions on all of  $M^n$ , rather than on a connected component within an open and dense subset. The maps  $F, G: \tilde{M}^{n+1} = M^n \times (-\epsilon, \epsilon) \to \mathbb{R}^{n+2}$  given by

$$F(x,t) = f(x) + t\eta \text{ and } G(x,t) = g(x) + t\bar{\eta}$$
(34)

parametrize for some  $\epsilon > 0$  isometric hypersurfaces. We know that f in Examples is a warped product of a hypersurface  $h: N^{n-p} \to \mathbb{R}^{n-p+1}$  with a minimal hypersurface  $f_1: L^p \to \mathbb{S}^{p+1}$  of rank two and that the Gauss map of F is determined by the Gauss map  $\xi$  of  $f_1$ . Since the image of  $\xi$  lies in  $\mathbb{S}^{p+1}$  then  $\mathbb{R}^{n-p} \subset F_*T_{(x,t)}\tilde{M}$  at any  $(x,t) \in \tilde{M}$ . Thus F is a cylinder over a minimal hypersurface  $F_1: \tilde{L}^{p+1} \to \mathbb{R}^{p+2}$ . From (34) we have

$$\langle F(x,t),\xi(x,t)\rangle = 0$$
 at any  $(x,t) \in \hat{M}$ ,

that is, the support function of F vanishes. Hence  $F_1$  is the cone over  $f_1$ . Thus F is a cylinder over the cone of a rank two minimal hypersurface  $f_1: L^p \to \mathbb{S}^{p+1}, 2 \leq p \leq n-1$ . If f is a rank two minimal hypersurface of a sphere and since in this situation the support function of F also vanishes, then F is the cone over f. The same assertions hold for G. In both cases, the extensions F and G are rank two minimal hypersurfaces. In particular, they are either surface-like or elliptic Sbrana-Cartan hypersurfaces.

Next we show the uniqueness of the extensions by means of an argument that does not rely on an assumption regarding the normal connection forms.

We consider orthonormal frames  $\{\eta, \xi\} \subset \Gamma(N_f M)$  and  $\{\bar{\eta}, \xi\} \subset \Gamma(N_g M)$  such that  $\eta$  and  $\bar{\eta}$  lie in the direction of the corresponding mean curvature vector fields. Thus (4) and (5) hold. We claim that, up to signs,  $\eta$  and  $\bar{\eta}$  are the unique normal vector fields whose corresponding shape operators coincide. In fact, let  $A_{\eta_2} = \bar{A}_{\bar{\eta}_2}$  where  $\eta_2 = a\eta + b\xi \in \Gamma(N_f M)$  and  $\bar{\eta}_2 = c\bar{\eta} + d\bar{\xi} \in \Gamma(N_g M)$  are unit vector fields. Then (4) yields

$$(a-c)A_{\eta} = d\bar{A}_{\bar{\xi}} - bA_{\xi}.$$

If  $a - c \neq 0$  it follows from (5) that tr  $A_{\eta} = 0$ . But then f would be minimal, which has been excluded. Thus a = c and it follows from Lemma 2 that b = d = 0, proving the claim.

Let  $\bar{F}, \bar{G}: N^{n+1} \to \mathbb{R}^{n+2}$  be non-congruent isometric extensions of f and g. Let  $\eta_1$  be a vector field normal to  $M^n$  in  $N^{n+1}$  of unit norm. Then  $\bar{F}_*\eta_1$  and  $\bar{G}_*\eta_1$  are normal to f and g, respectively, and the corresponding shape operators satisfy  $A_{\bar{F}_*\eta_1} = \bar{A}_{\bar{G}_*\eta_1}$ . Hence, up to sign, we have that  $\bar{F}_*\eta_1 = \eta$  and  $\bar{G}_*\eta_1 = \bar{\eta}$ . Thus the vector fields  $\eta$  and  $\bar{\eta}$  are tangent along f and g to  $\bar{F}(N)$  and  $\bar{G}(N)$ , respectively. Hence, along f and g the vector fields  $\xi$  and  $\bar{\xi}$  are normal to  $\bar{F}(N)$  and  $\bar{G}(N)$ . Since rank  $A_{\xi} = \operatorname{rank} \bar{A}_{\bar{\xi}} = 2$  and  $\bar{F}$  and  $\bar{G}$  are non-congruent, then these hypersurfaces have rank two. Therefore, there is  $X_0 \in \Gamma(\Delta^{\perp})$ , possible zero, such that  $f_*X_0 + \eta$  lies in the relative nullity of  $\bar{F}$ , because otherwise,  $\bar{F}$  would have rank larger than two. Since the leaves of relative nullity of an immersion are mapped to open subsets of affine subspaces, then the segments  $t(f_*X_0 + \eta)$  are contained in  $\bar{F}(N)$  for small t. Having that the same holds for  $\bar{G}$ , then these extensions coincide with F and G given either by (17) or (34), according to the case.

To conclude the proof of the theorem, we argue regarding the statement there about the set of deformations with isometric Gauss maps.

Let  $g_1: M^n \to \mathbb{R}^{n+2}$  be an isometric deformation of f, other than g, with an isometric Gauss map. Then f and  $g_1$  extend uniquely to Sbrana-Cartan hypersurfaces  $F_1$  and  $G_1$ . We have seen that  $X_0 \in \Gamma(\Delta^{\perp})$  given by

$$\langle A_{\xi}X, X_0 \rangle + \psi(X) = 0$$
 for any  $X \in \Gamma(\Delta^{\perp})$ 

is such that  $f_*X_0 + \eta$  is contained in the relative nullity vector subspaces of  $F_1$ . This shows that  $F_1$  coincides with F given by (17) or (34), and thus that  $G_1$  is an isometric deformation of F.

If F and G are the Sbrana-Cartan extensions of f and g, respectively, then there is an isometric embedding  $j: M^n \to \tilde{M}^{n+1}$  such that  $f = F \circ j$  and  $g = G \circ j$ . The second fundamental forms  $A^F$  and  $A^G$  of F and G are related by (25). If F lies in the continuous class, then it admits a one-parameter family of deformations, say  $G_s$ , with shape operators

$$A^{G_s} = A^F(\cos\gamma_s I + \sin\gamma_s J) \text{ where } \gamma_s \in C^{\infty}(\tilde{M}).$$
(35)

Let  $\xi_s$  be the Gauss map of  $G_s$ . Calling  $g_s = G_s \circ j$ , then the shape operator of  $g_s$  associated to  $\xi_s$  is given by  $j_*\bar{A}_{\xi_s} = (A^{G_s}|_{j_*TM})_{j_*TM}$ . We have from (5) that

$$\operatorname{tr}\left((A^F \circ J|_{j_*TM})_{j_*TM}\right) = 0.$$

Therefore it follows from (35) that tr  $\bar{A}_{\xi_s} = 0$ . If  $\{\eta_s, \xi_s\} \subset \Gamma(N_{g_s}M)$  is an orthonormal frame, then we have that  $\eta_s$  lies in the direction of the mean curvature vector field of  $g_s$ . Note that  $\eta_s$  is normal to j in  $\tilde{M}^{n+1}$ . Therefore it follows from (2) that f and  $g_s$  have the same third fundamental form. Since we have seen that any isometric deformation of f with isometric Gauss map extends to an isometric deformation of F, then the immersions  $g_s$  are all the possible isometric deformations of f with isometric Gauss maps. If F is of the discrete class, that is, if it admits a single isometric deformation, then also f admits a unique isometric deformation g such that f and g have isometric Gauss maps.

#### 2 Some comments

(1) There is some additional information for the submanifolds in part (ii) of Theorem 1. In fact, it was established that the submanifold is either a warped product of immersions with a minimal factor or just a minimal hypersurface of a sphere. Then the isometric deformations in the former case are given in terms of the deformations of the minimal factor, whereas in the latter case, by the deformation in the sphere of the submanifold itself.

(2) To illustrate the difficulty in constructing an example that belongs to part (i) of Theorem 1 but not to part (ii), an example we currently lack, we make the following observation.

**Proposition 5.** Let  $F: \overline{M}^{n+1} \to \mathbb{R}^{n+2}$ ,  $n \geq 3$ , be an elliptic Sbrana-Cartan hypersurface and let  $G: \overline{M}^{n+1} \to \mathbb{R}^{n+2}$  be an isometric deformation Let  $j: M^n \to \overline{M}^{n+1}$  be an isometric immersion satisfying at any point of  $M^n$  that

$$tr((A^{F}|_{j_{*}TM})_{j_{*}TM}) = tr((A^{G}|_{j_{*}TM})_{j_{*}TM}) = 0.$$
(36)

Then  $f = F \circ j \colon M^n \to \mathbb{R}^{n+2}$  and  $g = G \circ j \colon M^n \to \mathbb{R}^{n+2}$  have isometric Gauss maps.

Proof: If  $\eta$  is a unit vector field normal to j(M) in  $\overline{M}^{n+1}$  and  $\eta_0 = F_*\eta$ ,  $\overline{\eta}_0 = G_*\eta$ , then  $A_{\eta_0} = \overline{A}_{\overline{\eta}_0}$ . Let  $\xi$  and  $\overline{\xi}$  be unit vector fields normal to F and G along j, respectively. From (36) we have tr  $A_{\xi} = \operatorname{tr} \overline{A}_{\overline{\xi}} = 0$ . Then from (2) and (36) it follows that f and g have the same third fundamental form.

(3) A very similar result to Theorem 1 holds if, instead of Euclidean space, we consider submanifolds with isometric Gauss maps in the round sphere. This similarity is expected since their cones are examples in Euclidean space.

Marcos Dajczer is partially supported by the grant PID2021-124157NB-I00 funded by MCIN/AEI/10.13039/501100011033/ 'ERDF A way of making Europe', Spain, and are also supported by Comunidad Autónoma de la Región de Murcia, Spain, within the framework of the Regional Programme in Promotion of the Scientific and Technical Research (Action Plan 2022), by Fundación Séneca, Regional Agency of Science and Technology, REF, 21899/PI/22.

Miguel I. Jimenez is supported by FAPESP with the grants 2022/05321-9 and 2023/06762-1.

Miguel I. Jimenez expresses gratitude to the Mathematics department of the University of Ioannina for their kind hospitality during the development of this work.

## References

 M. Dajczer and L. Florit, Genuine deformations of submanifolds, Comm. Anal. Geom. 12 (2004), 1105–1129.

- [2] M. Dajczer and L. Florit, Genuine rigidity of Euclidean submanifolds in codimension two, Geom. Dedicata 106 (2004), 195–210.
- [3] M. Dajczer, L. Florit and R. Tojeiro, On deformable hypersurfaces in space forms, Ann. Mat. Pura Appl. 174 (1998), 361–390.
- [4] M. Dajczer and D. Gromoll, Real Kaehler submanifolds and uniqueness of the Gauss map, J. Differential Geom. 22 (1985), 13–28.
- [5] M. Dajczer and D. Gromoll, Euclidean hypersurfaces with isometric Gauss maps, Math. Z. 191 (1986), 201–205.
- [6] M. Dajczer and D. Gromoll, The Weierstrass representation for complete real Kaehler submanifolds of codimension two, Invent. Math. 119 (1995), 235–242.
- [7] M. Dajczer and R. Tojeiro, "Submanifold theory. Beyond an introduction". Universitext. Springer, New York, 2019.
- [8] M. Dajczer and Th. Vlachos, A class of complete minimal submanifolds and their associated families of genuine deformations, Comm. Anal. Geom. 26 (2018), 699– 721.
- [9] S. Nölker, Isometric immersions of warped products, Differ. Geom. Appl. 6 (1996), 1–30.
- [10] M. Obata, The Gauss map of immersions of Riemannian manifolds in spaces of constant curvature, J. Differential Geom. 2 (1968), 217–223.
- [11] Th. Vlachos, Isometric deformations of surfaces preserving the third fundamental form, Ann. Mat. Pura Appl. 187 (2008), 137–155.

Marcos Dajczer Departamento de Matemáticas Universidad de Murcia, E-30100 Espinardo, Murcia – Spain e-mail: marcos@impa.br

Miguel Ibieta Jimenez Instituto de Ciências Matemáticas e de Computação Universidade de São Paulo São Carlos SP 13566-590 – Brazil e-mail: mibieta@icmc.usp.br Theodoros Vlachos University of Ioannina Department of Mathematics Ioannina – Greece e-mail: tvlachos@uoi.gr